


Environmental microbiology.
v. 17, no. 8 (Aug. 2015)
General Collection
W1 EN984UM
2015-09-16 06:29:15

Environmental microbiology



PROPERTY OF THE
NATIONAL
LIBRARY OF
MEDICINE

Special Issue on Fungal Ecology

Bacteria in truffles contribute to aroma

Virulence traits of human pathogen *A. fumigatus* protect against amoeba

Disruption of photoautotrophic intertidal mats by filamentous fungi



WILEY Blackwell

Motif Exhibit 1064, Page 1 of 17

Discover this journal online at
Wiley Online Library
wileyonlinelibrary.com

environmental microbiology

http://www.env-micro.com

Editors

- Kenneth N. Timmis.** Institute of Microbiology, Technical University Braunschweig, Spielmannstrasse 7, D-38106 Braunschweig, Germany. E-mail: kntimmi@gmail.com
- Martin Polz.** Department of Civil and Environmental Engineering, Massachusetts Institute of Technology, 77 Massachusetts Avenue, 48-417, Cambridge, MA 02139, USA. Tel (617) 253 7128; Fax (617) 253 8850; E-mail: mpolz@mit.edu
- Michael Wagner.** Department of Microbial Ecology, University of Vienna, Althanstr. 14, A-1090 Vienna, Austria. Tel +43 1 4277 54390; Fax +43 1 4277 54389; E-mail: wagner@microbial-ecology.net
- Mike Jetten.** Department of Microbiology, Huygens Bldg Room 02.406, IWWR-FNWI, Radboud University Nijmegen, Toernooiveld 1, 6525 ED Nijmegen, The Netherlands. Tel +31 (24) 365 2940/2941; Fax +31 (24) 365 2830; E-mail: m.jetten@science.ru.nl
- Michael Galperin.** National Center for Biotechnology Information, NLM, National Institutes of Health, Bldg 38A, Bethesda, MD 20894, USA. Tel +1 301 435 5910; Fax +1 301 435 7793; E-mail: galperin@ncbi.nlm.nih.gov
- Victoria Orphan.** Argonne National Laboratory, Building 202, Room B373, 9700, South Cass Avenue, Lemont, IL, 60439, USA. Tel: +1 630 252 7489; E-mail: vorphan@gps.caltech.edu
- Paola Bonfante.** Dipartimento di Scienza della Vita e Biologia dei Sistemi, Dept. of Life Science and Systems Biology, Viale Mattioli 25 – 10125 Torino, Italy. Tel. +39 011 670 5965 (office); +39 011 650 2927 (lab.); E-mail: paola.bonfante@unito.it; p.bonfante@ipp.cnr.it
- Jack A. Gilbert.** Argonne National Laboratory, Building 202, Room B373, 9700, South Cass Avenue, Lemont, IL, 60439, USA. Tel: +1 630 252 7489; E-mail: gilbertjack@ANL.gov
- Rachel Whitaker.** Department of Microbiology, University of Illinois, 601 S. Goodwin Ave., Urbana, IL 61801, USA. Tel: +1 217 244 8420; E-mail: rwhitaker@life.illinois.edu
- Juan Luis Ramos Martin (Editor, and Reviews and Special Issues Editor).** Abengoa Research, Campus Palmas Altas, C/ Energía Solar nº 1, Palmas Altas, Seville, Spain. Tel +34 648102090; Fax +34 955413371; E-mail: juan.ramos@research.abengoa.com

Editorial Board

- Martin Ackermann *ETH Zurich and Eawag, Switzerland*
- Sonja-Vereina Albers *Marburg, Germany*
- Mads Albertsen *Aalborg, Denmark*
- Rudolf Amann *Bremen, Germany*
- Ricardo Amils *Madrid, Spain*
- John Archibald *Nova Scotia, Canada*
- Adam P. Arkin *Berkeley, CA, USA*
- Manfred Auer *Berkeley, CA, USA*
- Mark Bailey *Oxon, UK*
- Nitin Baliga *Seattle, WA, USA*
- Doug Bartlett *Jolla, CA, USA*
- Fernando Baquero *Madrid, Spain*
- David Berry *Vienna, Austria*
- Stefan Bertilsson *Uppsala, Sweden*
- Antje Boetius *Bremen, Germany*
- David Bourne *Queensland, Australia*
- Erhard Bremer *Marburg, Germany*
- Andreas Brune *Marburg, Germany*
- Carmen Buchrieser *Paris, France*
- Dan Buckley *Ithaca, NY, USA*
- Don Canfield *Odense, Denmark*
- David A. Caron *Los Angeles, CA, USA*
- Colleen Cavanaugh *Cambridge, MA, USA*
- Rick Cavicchioli *Sydney, Australia*
- Haiyan Chu *Nanjing, China*
- Rita Colwell *College Park, MD, USA*
- Ralf Conrad *Marburg, Germany*
- Pierre Cornelis *Brussels, Belgium*
- Milton da Costa *Coimbra, Portugal*
- Tom Curtis *Newcastle-upon-Tyne, UK*
- Daniele Daffonchio *Milan, Italy*
- Jaap Sinninghe Damsté *Den Burg, The Netherlands*
- Antoine Danchin *Paris, France*
- Roberto Danovaro *Ancona, Italy*
- Seana Davidson *Seattle, WA, USA*
- Eduardo Diaz *Madrid, Spain*
- Ulrich Dobrindt *Münster, Germany*
- Xiuzhu Dong *Beijing, China*
- Harold L. Drake *Bayreuth, Germany*
- Arnold Driessen *Groningen, The Netherlands*
- Nicole Dubilier *Bremen, Germany*
- Leo Eberl *Zürich, Switzerland*
- Katrina Edwards *Los Angeles, CA, USA*
- Ed J. Feil *Bath, UK*
- Julia Foght *Edmonton, Canada*
- Patrick Forterre *Paris, France*
- Jim K. Fredrickson *Richland, WA, USA*
- Michael W. Friedrich *Bremen, Germany*
- Jed Fuhrman *Los Angeles, CA, USA*
- Roger Garrett *Copenhagen, Denmark*
- Christopher van der Gast *Wallingford, UK*
- Steve Giovannoni *Corvallis, OR, USA*
- Lone Gram *Kgs. Lyngby, Denmark*
- Robert Gunsalus *Los Angeles, CA, USA*
- Martin W. Hahn *Mondsee, Austria*
- John E. Hallsworth *Belfast, Northern Ireland*
- Jo Handelsman *New Haven, CT, USA*
- Shige Harayama *Tokyo, Japan*
- John Hayes *Berkeley, CA, USA*
- Ian Head *Newcastle-upon-Tyne, UK*
- Michael Hecker *Greifswald, Germany*
- Tori M. Hoehler *Moffett Field, CA, USA*
- Andrew Holmes *Sydney, Australia*
- Steve D'Hondt *Narragansett, RI, USA*
- M. Claire Homer-Devine *Seattle, WA, USA*
- Phil Hugenholz *Queensland, Australia*
- Dieter Jahn *Braunschweig, Germany*
- Janet Jansson *Berkeley, CA, USA*
- Bo Barker Jørgensen *Aarhus, Denmark*
- Gary M. King *Baton Rouge, LA, USA*
- Staffan Kjelleberg *Sydney, Australia*
- Martin Klotz *Charlotte, NC, USA*
- Rob Knight *Boulder, CO, USA*
- George Kowalchuk *Heteren, The Netherlands*
- Michael Kühl *Helsingør, Denmark*
- Nikos K. Kypides *Walnut Creek, CA, USA*
- Rup Lal *Delhi, India*
- Jared R. ("Tuck") Leadbetter *Pasadena, CA, USA*
- Werner Liesack *Marburg, Germany*
- Debbie Lindell *Haifa, Israel*
- Steven Lindow *Berkeley, CA, USA*
- Frank Löffler *Knoxville, TN, USA*
- Victor de Lorenzo *Madrid, Spain*
- Derek Lovley *Amherst, MA, USA*
- Alexander Loy *Wien, Austria*
- Tillmann Lueders *Neuherberg, Germany*
- Rod Mackie *Urbana, IL, USA*
- Mike Madigan *Carbondale, IL, USA*
- Ramon Massana *Barcelona, Spain*
- Timothy R. McDermott *Bozemann, MT, USA*
- Terry McGenity *Colchester, UK*
- Mike McInerney *Norman, OK, USA*
- Trina McMahon *Madison, WI, USA*
- Rainer U. Meckenstock *Neuherberg, Germany*
- Jan Roelof van der Meer *Lausanne, Switzerland*
- Gabriella Molinari *Braunschweig, Germany*
- Mary Ann Moran *Athens, GA, USA*
- Francois Morel *Princeton, NJ, USA*
- Andrés Moya *Valencia, Spain*
- Volker Müller *Frankfurt, Germany*
- Colin Murrell *Coventry, UK*
- Ken Nealson *Los Angeles, CA, USA*
- Doug Nelson *Davis, CA, USA*
- Josh Neufeld *Ontario, Canada*
- Graeme Nicol *Aberdeen, UK*
- Per Halkjaer Nielsen *Aalborg, Denmark*
- Balbina Nogales *Palma de Mallorca, Spain*
- Daniel Noguera *Madison, WI, USA*
- Howard Ochman *Austin, TX, USA*
- Maarja Opik *Tartu, Estonia*
- Huub Op den Camp *Nijmegen, The Netherlands*
- Jörg Overmann *Braunschweig, Germany*
- Norman R. Pace *Boulder, CO, USA*
- Hans Paerl *Morehead City, NC, USA*
- R. John Parkes *Cardiff, UK*
- Ann Pearson *Cambridge, MA, USA*
- Jakob Pernthaler *Kilchberg, Switzerland*
- Jörn Piel *Zürich, Switzerland*
- David Prangishvili *Paris, France*
- Jim Prosser *Aberdeen, UK*
- Carla Pruzzo *Genova, Italy*
- Ralf Rabus *Oldenburg, Germany*
- Paul Rainey *Auckland, New Zealand*
- David Reiman *Palo Alto, CA, USA*
- Niels Peter Revsbech *Aarhus, Denmark*
- Katharina Riedel *Greifswald, Germany*
- Ute Römling *Stockholm, Sweden*
- Forest Rohwer *San Diego, CA, USA*
- Fernando Rojo *Madrid, Spain*
- Eugene Rosenberg *Ramat Aviv, Israel*
- Edward G. Ruby *Madison, WI, USA*
- Mak Saito *Woods Hole, MA, USA*
- Bernhard Schink *Konstanz, Germany*
- Christa Schlieper *Vienna, Austria*
- Michael Schloter *Oberschleissheim, Germany*
- Tom Schmidt *Ann Arbor, MI, USA*
- Andreas Schramm *Aarhus, Denmark*
- Barth F. Smets *Kgs. Lyngby, Denmark*
- Hauke Smidt *Wageningen, The Netherlands*
- Mitch Sogin *Woods Hole, MA, USA*
- John Spear *Golden, CO, USA*
- Alfred Spormann *Stanford, CA, USA*
- Lisa Y. Stein *Alberta, Canada*
- Ramunas Stepanauskas *East Boothbay, ME, USA*
- Thorsten Stoeck *Kaiserslautern, Germany*
- Marc Strous *Calgary, Canada*
- Matthew B Sullivan *Tucson, AZ, USA*
- Mike Taylor *Auckland, New Zealand*
- Andreas Teske *Chapel Hill, NC, USA*
- Burkhard Tümmler *Hannover, Germany*
- Gene Tyson *Brisbane, Australia*
- Tim Ulrich *Vienna, Austria*
- Gerrit Voodouw *Calgary, Canada*
- Julia A. Vorholt *Zürich, Switzerland*
- Willem M. de Vos *Wageningen, The Netherlands*
- Larry Wackett *St. Paul, MN, USA*
- Judy Wall *Columbia, MO, USA*
- Nicole Webster *Queensland, Australia*
- Friedrich Widdel *Bremen, Germany*
- Paul Williams *Nottingham, UK*
- Willie Wilson *Plymouth, UK*
- K. Eric Wommack *Newark, DE, USA*
- Michail M. Yakimov *Messina, Italy*
- Jonathan P. Zehr *Santa Cruz, CA, USA*
- Liping Zhao *Shanghai, China*
- Steve Zinder *Ithaca NY, USA*
- Erik Zinser *Knoxville, TN, USA*

Genomics Updates: Michael Galperin. National Center for Biotechnology Information, NLM, National Institutes of Health, Bldg 38A, Bethesda, MD 20894, USA. Tel (+1) 301 435 5910; Fax (+1) 301 435 7793; E-mail galperin@ncbi.nlm.nih.gov

Production Editor: Jonathan R. Rapay, John Wiley & Sons Philippines, SPI Bldg., Pascor Drive, Sto. Nino, Paranaque City 1700, Philippines. Fax: +632 325 0768; E-mail: jrapay@wiley.com

Cover Illustration: (A) Mature *Macrotermes natalensis* colony in a South African landscape (photo by D. K. Aanen). (B) The underground fungus comb, in which workers maintain *Termitomyces* (photo by D. K. Aanen). (C) The royal chamber in which the queen and king reside. (D) A *Macrotermes natalensis* royal pair after the ceiling of the royal chamber has been removed. (E) Fruiting structures of *Termitomyces reticulatus* emerging from a colony of *Odontotermes badius* or *Odontotermes transvaalensis*. For further details, the readers are referred to the article by Michael Poulsen on pp. 2562–2572 of this issue.

Information for Subscribers

Environmental Microbiology is published in 12 issues per year. Institutional subscription prices for the *Environmental Microbiology Package* for 2015 are: Print & Online: US\$6575 (US), US\$7672 (Rest of World) €4522 (Europe), £3560 (UK). Prices are exclusive of tax. Asia-Pacific GST, Canadian GST and European VAT will be applied at the appropriate rates. For more information on current tax rates, please go to wileyonlinelibrary.com/tax-vat. The price includes online access to the current and all online back files to January 1st 2007, where available. For other pricing options, including access information and terms and conditions, please visit wileyonlinelibrary.com/access.

*Includes *Environmental Microbiology* and *Environmental Microbiology Reports*

Delivery Terms and Legal Title

Where the subscription price includes print issues and delivery is to the recipient's address, delivery terms are Delivered at Place (DAP); the recipient is responsible for paying any import duty or taxes. Title to all issues transfers FOB our shipping point, freight prepaid. We will endeavour to fulfil claims for missing or damaged copies within six months of publication, within our reasonable discretion and subject to availability.

This journal is available at Wiley Online Library. Visit <http://wileyonlinelibrary.com> to search the articles and register for table of contents e-mail alerts.

Back Issues

Single issues from current and prior year volumes are available at the current single issue price from cs-journals@wiley.com. Earlier issues may be obtained from Periodicals Service Company, 11 Main Street, Germantown, NY 12526, USA. Tel: +1 518 537 4700, Fax: +1 518 537 5899, E-mail: psc@periodicals.com

Periodical ID Statement A

ENVIRONMENTAL MICROBIOLOGY (ISSN: 1462-2912) is published monthly. US mailing agent: Mercury Media Processing, LLC 1850 Elizabeth Avenue, Suite #C, Rahway, NJ 07065 USA. Periodical postage paid at Rahway, NJ.

POSTMASTER: Send all address changes to ENVIRONMENTAL MICROBIOLOGY, John Wiley & Sons Inc., C/O The Sheridan Press, PO Box 465, Hanover, PA 17331.

For submission instructions, subscription and all other information, visit: www.env-micro.com

Printed in the UK by Hobbs the Printers Ltd
Typeset by Toppan Best-set Premedia Limited

ISSN 1462-2912 (Print)
ISSN 1462-2920 (Online)

Contents

Highlight

- 2559 Could in situ DNA-hybridization chain reaction enable simple and effective detection of identity and function in whole cell hybridizations?
D. R. Noguera

Minireviews

- 2562 Towards an integrated understanding of the consequences of fungus domestication on the fungus-growing termite gut microbiota
M. Poulsen
- 2573 Microbial players involved in the decline of filamentous and colonial cyanobacterial blooms with a focus on fungal parasitism
M. Gerphagnon, D. J. MacArthur, D. Latour, C. M. M. Gachon, F. Van Ogtrop, F. H. Gleason & T. Sime-Ngando

Research articles

- 2588 New tricks of an old enemy: isolates of *Fusarium graminearum* produce a type A trichothecene mycotoxin
E. Varga, G. Wiesenberger, C. Hametner, T. J. Ward, Y. Dong, D. Schöffbeck, S. McCormick, K. Broz, R. Stückler, R. Schumacher, R. Krska, H. C. Kistler, F. Berthiller & G. Adam
- 2601 Characterization of protein kinase *PsSRPKL*, a novel pathogenicity factor in the wheat stripe rust fungus
Y. Cheng, X. Wang, J. Yao, R. T. Voegelé, Y. Zhang, W. Wang, L. Huang & Z. Kang
- 2618 Fungal lysis by a soil bacterium fermenting cellulose
A. C. Tolonen, T. Cerisy, H. El-Sayed, M. Boutard, M. Salanoubat & G. M. Church
- 2628 Production of trichodiene by *Trichoderma harzianum* alters the perception of this biocontrol strain by plants and antagonized fungi
M. G. Malmierca, S. P. McCormick, R. E. Cardozo, N. J. Alexander, E. Monte & S. Gutiérrez
- 2647 Bacteria associated with truffle-fruiting bodies contribute to truffle aroma
R. Spivak, A. Deveau, N. Valdez, N. Kirchhoff, P. Frey-Klett & P. Karlovsky
- 2661 Protein kinase FgSch9 serves as a mediator of the target of rapamycin and high osmolarity glycerol pathways and regulates multiple stress responses and secondary metabolism in *Fusarium graminearum*
Q. Gu, C. Zhang, F. Yu, Y. Yin, W.-B. Shim & Z. Ma
- 2677 Network analysis reveals that bacteria and fungi form modules that correlate independently with soil parameters
A. B. de Menezes, M. T. Prendergast-Miller, A. E. Richardson, P. Toscas, M. Farrell, L. M. MacDonald, G. Baker, T. Wark & P. H. Thrall
- 2690 The global regulator FtsG1 is required for expression of secondary metabolite gene clusters but not for pathogenicity in *Fusarium fujikuroi*
C. B. Michielse, L. Stüttgen, S. Janeska, C. M. K. Sieber, B. Arndt, J. J. Espino, H.-U. Humpf, U. Güldener & B. Tuszynski
- 2709 The composition of arbuscular mycorrhizal fungal communities in the roots of a ruderal forb is not related to the forest fragmentation process
G. Grilli, C. Urcelay, L. Galetto, J. Davison, M. Vassar, Ü. Saks, T. Jauris & M. Öpik
- 2721 Loss of Dfg5 glycosylphosphatidylinositol-anchored membrane protein confers enhanced heat tolerance in *Saccharomyces cerevisiae*
O. Nasution, J. Lee, K. Srinivasa, I.-G. Choi, Y. M. Lee, E. Kim, W. Choi & W. Kim
- 2735 A small molecule species specifically inhibits *Fusarium* myosin I
C. Zhang, Y. Chen, Y. Yin, H.-H. Ji, W.-B. Shim, Y. Hou, M. Zhou, X. Li & Z. Ma
- 2747 Whose truffle is this? Distribution patterns of ectomycorrhizal fungal diversity in *Tuber melanosporum* brûlés developed in multi-host Mediterranean plant communities
E. Taschen, M. Sauve, A. Taudière, J. Parlane, M.-A. Selosse & F. Richard
- 2762 The MAD5-box transcription factor FgMcm1 regulates cell identity and fungal development in *Fusarium graminearum*
C. Yang, H. Liu, G. Li, M. Liu, Y. Yun, C. Wang, Z. Ma & J.-R. Xu
- 2777 Primary succession of *Bistorta vivipara* (L.) Delabre (Polygonaceae) root-associated fungi mirrors plant succession in two glacial chronosequences
M. Davey, R. Blaaild, U. Vik, T. Carlsen, H. Kausrud & P. B. Eldesen
- 2791 Community composition and population genetics of insect pathogenic fungi in the genus *Metarhizium* from soils of a long-term agricultural research system
R. M. Kepler, T. A. Ugine, J. E. Maul, M. A. Cavigelli & S. A. Rehner
- 2805 Fungicide efflux and the MgMFS1 transporter contribute to the multidrug resistance phenotype in *Zymoseptoria tritici* field isolates
S. Omrane, H. Sghyer, C. Audéon, C. Lanen, C. Duplaix, A.-S. Walker & S. Fillingier
- 2824 Globally invading populations of the fungal plant pathogen *Verticillium dahliae* are dominated by multiple divergent lineages
D. P. G. Short, S. Gurung, P. Gladieux, P. Inderbitzin, Z. K. Atallah, F. Nigro, G. Li, S. Benlioglu & K. V. Subbarao
- 2841 Contribution of arbuscular mycorrhizal fungi of sedges to soil aggregation along an altitudinal alpine grassland gradient on the Tibetan Plateau
X. Li, J. Zhang, J. Gai, X. Cai, P. Christie & X. Li
- 2858 Virulence determinants of the human pathogenic fungus *Aspergillus fumigatus* protect against soil amoeba predation
F. Hillmann, S. Novohradská, D. J. Mattern, T. Forberger, T. Heinekamp, M. Westermann, T. Winckler & A. A. Brakhage
- 2870 Genetic variability and bottleneck detection of four *Tricholoma matsutake* populations from northeastern and southwestern China
D.-F. Zeng & B. Chen
- 2882 The composition of arbuscular mycorrhizal fungal communities differs among the roots, spores and extraradical mycelia associated with five Mediterranean plant species
S. Varela-Cervero, M. Vassar, J. Davison, J. M. Barea, M. Öpik & C. Azcón-Aguilar
- 2896 pH dependency of sclerotial development and pathogenicity revealed by using genetically defined oxalate-minus mutants of *Sclerotinia sclerotiorum*
L. Xu, M. Xiang, D. White & W. Chen
- 2910 Disruption of photoautotrophic intertidal mats by filamentous fungi
C. Carreira, M. Staal, D. Falkowski, R. P. de Vries, M. Middelboe & C. P. D. Brussaard
- 2922 Understanding fungal functional biodiversity during the mitigation of environmentally dispersed pentachlorophenol in cork oak forest soils
A. Varela, C. Martins, O. Núñez, I. Martins, J. A. M. P. Houbaker, T. M. Martins, M. C. Leitão, I. McLellan, W. Vetter, M. T. Galceran, R. A. Samson, A. Hursthouse & C. Silva Pereira
- 2935 *De novo* biosynthesis of cytokinins in the biotrophic fungus *Claviceps purpurea*
J. Hinsch, J. Vrabka, B. Oeser, O. Novák, P. Galuszka & P. Tuszynski
- 2952 Metagenome sequence of *Elaphomyces granulatus* from sporocarp tissue reveals Ascomycota ectomycorrhizal fingerprints of genome expansion and a Proteobacteria-rich microbiome
C. A. Quandt, A. Köhler, C. N. Hesse, T. J. Sharpton, F. Martin & J. W. Spatafora
- 2969 Activation of the signalling mucin MoMsb2 and its functional relationship with Cbp1 in *Magnaporthe oryzae*
G. Wang, G. Li, S. Zhang, C. Jiang, J. Qin & J.-R. Xu
- 2982 Surviving the heat: heterogeneity of response in *Saccharomyces cerevisiae* provides insight into thermal damage to the membrane
S. Guyot, P. Gervais, M. Young, P. Winckler, J. Dumont & H. M. Davey
- 2993 Biosynthesis of the antifungal haterulalide, cycodylin A, in *Serratia*, and its regulation by quorum sensing, RpoS and Hfq
M. A. Matilla, F. J. Leeper & G. P. C. Salmond
- 3009 Compartmentalized and contrasted response of ectomycorrhizal and soil fungal communities of Scots pine forests along elevation gradients in France and Spain
A. Rincón, B. Santamaría-Pérez, S. G. Rabasa, A. Conce, B. Marçais & M. Buée
- 3025 Early rhizosphere microbiome composition is related to the growth and Zn uptake of willows introduced to a former landfill
T. H. Bell, B. Cloutier-Hurteau, F. Al-Otaibi, M.-C. Turmel, E. Yergeau, F. Courchesne & M. St-Arnaud
- 3039 Fine-scale spatial genetic structure analysis of the black truffle *Tuber aestivum* and its link to aroma variability
V. Molinier, C. Murat, H. Frochet, D. Wipf & R. Spivak
- 3051 Inner Mongolian steppe arbuscular mycorrhizal fungal communities respond more strongly to water availability than to nitrogen fertilization
X. Li, T. Zhu, F. Peng, Q. Chen, S. Lin, P. Christie & J. Zhang
- 3069 Hsp12p and PAU genes are involved in ecological interactions between natural yeast strains
D. Rivero, L. Berná, I. Stefanini, E. Baruffini, A. Bergerat, A. Csikász-Nagy, C. De Filippo & D. Cavalieri
- 3082 CaCO₃ and SrCO₃ bioprecipitation by fungi isolated from calcareous soil
Q. Li, L. Csetenyi, G. I. Paton & G. M. Gadd
- 3098 Uncovering the abilities of *Agaricus bisporus* to degrade plant biomass throughout its life cycle
A. Patyshakulyeva, H. Post, M. Zhou, E. Jurak, A. J. R. Heck, K. S. Hildén, M. A. Kabel, M. R. Mäkelä, M. A. F. Attelaar & R. P. de Vries

Web alert

- 3110 Fungal toxins and diseases
L. P. Wackett

Forthcoming papers in *Environmental Microbiology*

In-plant sporulation phenotype: a major life history trait to understand the evolution of *Alnus*-infective *Frankia* strains
A. C. Pozzi, H. H. Bautista-Guerrero, I. Nouiou, L. Cotin-Galvan, R. Pepin, P. Fournier, F. Menu, M. P. Fernandez & A. Herrera-Belaroussi

An improved method to set significance thresholds for β diversity testing in microbial community comparisons
A. Gülay & B. F. Smets

The MqsR/MqsA toxin/antitoxin system protects *Escherichia coli* during bile acid stress
B. W. Kwan, D. M. Lörd, W. Peti, R. Page, M. J. Benedik & T. K. Wood

Diversity and distribution of unicellular opisthokonts along the European coast analysed using high-throughput sequencing
J. del Campo, D. Mallo, R. Massana, C. de Vargas, T. A. Richards & I. Ruiz-Trillo

Soil bacterial communities are shaped by temporal and environmental filtering: evidence from a long-term chronosequence
Z. Freedman & D. R. Zak

Methane-derived carbon flow through microbial communities in arctic lake sediments
R. He, M. J. Woolter, J. W. Pohman, J. M. Tiedje & M. B. Leigh

Differential transcriptional response to antibiotics by *Pseudomonas putida* DOT-T1E
C. Molina-Santiago, A. Daddaoua, M. Gómez-Lozano, Z. Udaondo, S. Molin & J.-L. Ramos

'Ménage à trois': a selfish genetic element uses a virus to propagate within *Thermotogales*
J. Lossouarn, C. L. Nesbo, C. Mercier, O. Zhaxybayeva, M. S. Johnson, R. Charchuck, J. Farasin, N. Bienvenu, A.-C. Baudoux, G. Michoud, M. Jebbar & C. Geslin

Enzymes of the benzoyl-coenzyme A degradation pathway in the hyperthermophilic archaeon *Ferroplasma acidiphilum*
G. Schmid, S. B. René & M. Boll

Hydrogen sulfide can inhibit and enhance oxygenic photosynthesis in a cyanobacterium from sulfidic springs
J. M. Klatt, S. Haas, P. Yilmaz, D. de Beer & L. Polerecky



1888

AMS/NATL LIB OF MEDICINE
ORDER #: 12752412
35320657249026001/MB
DC-60350-AB 12752412
ILS: SR43252

EBSCO

ENVIRONMENTAL MICROBIOLOGY / FOR AMERICAS/
2015 VOLUME :17 ISSUE :8

V E
SISAC

Bacteria associated with truffle-fruited bodies contribute to truffle aroma

Richard Splivallo,^{1,2*} Aurélie Deveau,³ Nayuf Valdez,¹ Nina Kirchhoff,¹ Pascale Frey-Klett³ and Petr Karlovsky¹

¹Molecular Phytopathology and Mycotoxin Research, Georg-August University of Göttingen, Göttingen, Germany.

²Institute for Molecular Biosciences, Biozentrum/Campus Riedberg, Frankfurt am Main, Germany.

³INRA, UMR 1136 INRA Université de Lorraine 'Interactions Arbres/Micro-organismes', Champenoux, France.

Summary

Truffles, symbiotic fungi renowned for the captivating aroma of their fruited bodies, are colonized by a complex bacterial community of unknown function. We characterized the bacterial community of the white truffle *Tuber borchii* and tested the involvement of its microbiome in the production of sulphur-containing volatiles. We found that sulphur-containing volatiles such as thiophene derivatives, characteristic of *T. borchii* fruited bodies, resulted from the biotransformation of non-volatile precursor(s) into volatile compounds by bacteria. The bacterial community of *T. borchii* was dominated by α - and β -*Proteobacteria*. Interestingly, all bacteria phyla/classes tested in this study were able to produce thiophene volatiles from *T. borchii* fruited body extract, irrespective of their isolation source (truffle or other sources). This indicates that the ability to produce thiophene volatiles might be widespread among bacteria and possibly linked to primary metabolism. Treatment of fruited bodies with antibacterial agents fully suppressed the production of thiophene volatiles while fungicides had no inhibitory effect. This suggests that during the

sexual stage of truffles, thiophene volatiles are exclusively synthesized by bacteria and not by the truffle. At this stage, the origin of thiophenes precursor in *T. borchii* remains elusive and the involvement of yeasts or other bacteria cannot be excluded.

Introduction

Truffles are symbiotic fungi that develop underground in association with plant roots, forming ectomycorrhizas (Mello *et al.*, 2006). Ectomycorrhizal associations, dominant in boreal and temperate forests, are of high ecological relevance since they improve plant nutrition and health (Read, 1991; Buscot *et al.*, 2000; Martin *et al.*, 2001). About 180 truffle species (*Tuber spp*) associate with angiosperm and gymnosperms and naturally occur in Europe, North-America and Asia (Bonito *et al.*, 2010).

The genome sequencing of the Périgord black truffle *Tuber melanosporum* expanded the status of truffles from a food delicacy to a scientific model valuable for the study of complex symbiotic interactions (Martin *et al.*, 2010). Truffles ectomycorrhizas and fruited bodies harbour a diverse microbial community including bacteria, yeasts and filamentous fungi (Barbieri *et al.*, 2005; 2007; Buzzini *et al.*, 2005; Pacioni *et al.*, 2007). Of these microbes, only bacteria have been extensively studied in truffles. Complex bacterial communities have been reported to establish in different truffle species: *T. aestivum* (Gryndler *et al.*, 2013), *T. magnatum* (Barbieri *et al.*, 2007), *T. melanosporum* (Antony-Babu *et al.*, 2013) and *T. borchii* (Sbrana *et al.*, 2000; Barbieri *et al.*, 2005). Bacteria colonize both the external (peridium) and internal part (gleba) of truffles and seem to be selected from the soil communities during the early stage of truffle formation (Antony-Babu *et al.*, 2013). Although bacterial communities differ depending on truffle species analysed, a core microbiome composed of α -*Proteobacteria* from the family of *Bradyrhizobiaceae* seems common to all species studied so far (Barbieri *et al.*, 2005; 2007; Antony-Babu *et al.*, 2013). Factors responsible for the selection of these bacteria remain mysterious. However, these bacteria could have a role in the development, growth and nutrition of truffle-fruited bodies (Sbrana *et al.*, 2000; 2002; Barbieri *et al.*, 2007; 2010; Antony-Babu *et al.*, 2013; Pavić *et al.*, 2013).

Truffle-fruited bodies emit intense aromas (Splivallo and Maier, 2011; Splivallo *et al.*, 2011). Sulphur-

Received 24 April, 2014; revised 26 May, 2014; accepted 28 May, 2014. *For correspondence. E-mail richard.splivallo@a3.epfl.ch; Tel. +49 69 798 42 193; Fax +49 69 798 29 527. Authors' contributions: RS and AD drafted the manuscript with input from all other co-authors. RS designed the experiments, performed the statistics and also performed the bioassays with truffle mycelium/fruited bodies/bacteria/antibacterial agents. N.V. isolated bacteria and performed the bacterial sequence analysis. N.K., P.K. and R.S. performed the volatile profiling and volatile data analysis on fruited bodies. A.D., P.F.-K. and R.S. designed the FISH analysis, and R.S. and A.D. analysed the FISH data. All authors read and approved the final manuscript.

containing volatiles have a central role in truffle aroma because they serve as attractants to mammals and contribute to truffle aroma sensed by humans (Talou *et al.*, 1990; Culleré *et al.*, 2010; Splivallo and Maier, 2011; Splivallo *et al.*, 2011). The origin of sulphur-containing volatiles in truffles is unclear as they might be derived from the truffle fungus itself but also from the microbial community inhabiting truffle-fruiting bodies (Buzzini *et al.*, 2005; Splivallo and Maier, 2011; Splivallo *et al.*, 2011). Analysis of the Black truffle's genome *T. melanosporum*, however, suggested that Black truffle may produce its volatiles without the involvement of bacteria (Martin *et al.*, 2010; Maxmen, 2010). Whether other truffle species can produce their volatiles by themselves or require the participation of associated bacteria remain unknown.

Here, we investigated the role of truffle-associated bacteria in the formation of the aroma of a white truffle species, *Tuber borchii*. *T. borchii* is a cultivated truffle species naturally occurring in Europe that has been recently introduced in New Zealand (Bonito *et al.*, 2010). *T. borchii* has been for long used in laboratories as a model organism to study truffles because of the relative faster growth of its mycelium compared with other truffle species. *T. borchii*-fruiting bodies emit some volatiles [thiophene derivatives such as 3-methylthiophene, hereafter referred to as (1), and 3-methyl-4,5-dihydrothiophene, referred to as (2)], which are species-specific and might be partially responsible for their characteristic aroma (Bellesia *et al.*, 2001; Mauriello *et al.*, 2004; Zeppa *et al.*, 2004; Splivallo *et al.*, 2007). The concentrations of thiophene volatiles in *T. borchii* have been reported to occur solely in fully mature (71–100% maturity) truffles (Zeppa *et al.*, 2004) and to increase upon storage at room temperature (Bellesia *et al.*, 2001), questioning the role of truffle-associated bacteria in the production of the aroma. To test this hypothesis, we analysed the evolution of the composition of the bacterial communities along storage as well as the production of thiophene derivatives. Then we tested the ability of bacterial isolates from *T. borchii* to produce thiophene volatiles. Overall, our results demonstrated that thiophene volatiles characteristic of *T. borchii*-fruiting bodies were produced by the microbiome inhabiting truffle-fruiting bodies.

Results

Composition of T. borchii bacterial communities evolves during storage

As a first step, we characterized the composition of bacterial communities associated to *T. borchii* during storage. Six *T. borchii* truffles were subsampled after 0, 2, 4 and 6 days of storage at room temperature. The bacterial communities were characterized and quantified by fluorescent *in situ* hybridization (FISH) using the eubacterial universal probe EUB338 as well as probes specifically targeting

Firmicutes (LGC354A), *Actinobacteria* (HGC69a), *Bacteroidetes* (CF319), α - (ALF1b), β - (BET42a), γ -*Proteobacteria* (GAM42a) and *Enterobacteriales* (EntB and EntD). Because bacterial distribution within fruiting bodies can be highly non-homogenous (Antony-Babu *et al.*, 2013), quantification of the bacterial communities was done by homogenizing gleba samples through grinding. *T. borchii* bacterial community composition was comparable with the one previously described in *T. borchii* (Barbieri *et al.*, 2007), with an overall majority of *Proteobacteria* (Fig. 1) among which α - and β -*Proteobacteria* were dominant. γ -*Proteobacteria*, including *Enterobacteriales*, and bacteria belonging to *Bacteroidetes* were also detected but at a much lower level. No *Firmicutes* or *Actinobacteria* could be detected at day 0 (Fig. 1). The overall bacterial community evolved upon storage with a shift of community composition with time. The quantity of bacteria first tended to increase at 2 days and was then reduced at day 6. A similar reduction in community size over time was observed in the α - and the β -*Proteobacteria*. The *Bacteroidetes*, the *Enterobacteriales*, the *Actinobacteria* and the γ -*Proteobacteria* populations remained quite low and stable during storage. Last, the *Firmicutes* population size slightly increased during the last days of storage.

Additionally to address bacterial cell distribution within undisrupted truffle tissues, thin sections of *T. borchii* were hybridized with FISH probes against α - and β -*Proteobacteria*, the two main bacterial groups identified in ground samples. α - and β -*Proteobacteria* could be observed in both the peridium and the gleba by confocal microscopy imaging (Fig. 2). Dense colonies of α - and β -*Proteobacteria* were observed in the peridium. However, this colonization was patchy with vast area without visible bacterial cells (Fig. 2A and B). Bacterial cells were present in between fungal cells but not inside the fungal cells in both gleba and peridium. Colonies of α -*Proteobacteria* did not appear to contain bacteria from other phyla in the peridium as illustrated by the complete overlay of *Eubacteria* and α probes (Fig. 2A). In contrast, β -*Proteobacteria* were found in mixed population with other bacteria (Fig. 2B) as demonstrated by the absence of overlay between eubacterial probe (FITC, green) and β -*Proteobacteria* (cy3, red). In the gleba, α -*Proteobacteria* were found as isolated cells as well as dense colonies (Fig. 2C) while β -*Proteobacteria* were more evenly distributed (Fig. 2D).

Bacterial community composition differ between peridium and gleba

Antony-Babu and colleagues (2013) recently demonstrated that bacterial communities from *Tuber melanosporum* strongly differed between the peridium and the gleba. To determine if a similar pattern was also present




Fig. 1. Change in the bacterial population inside *T. borchii* fruiting bodies as a function of storage time. The panels show bacterial population dynamics ($\pm$ SE) of all bacteria [*Eubacteria* expressed as bacterial cell count per milligram of truffle-fruiting body (dry weight)] and specific classes/groups (expressed as percentage of the total *Eubacteria*). The community was dominated by α - and β -*Proteobacteria*. No statistical difference was detected among storage days within *Eubacteria* or bacterial classes/groups ($n=6$ truffle-fruiting bodies, for all ANOVA: $0.03 < F < 1.62$; $0.22 < P < 0.99$). *Enterobacteriales* data from FISH probe EntD (probe EntB gave comparable results – not shown).

in *T. borchii*, we compared by FISH bacterial community composition in gleba and peridium samples at day 0. Only the population size of *Bacteroidetes* significantly differed between gleba and peridium. The population size of the latter phylum was five times higher in the peridium compared with the gleba (gleba: 137 ± 122 ; peridium: 697 ± 101 ; unit: cell count/mg dry weight; P -value: 0.026, Mann–Whitney U -test). No statistical difference between peridium and gleba was observed for the *Firmicutes* (gleba: 0 ± 0 ; peridium: 10 ± 10), *Actinobacteria* (gleba: 0 ± 0 ; peridium: 45 ± 45), α -*Proteobacteria* (gleba: 1370 ± 499 ; peridium: 947 ± 365), β -*Proteobacteria* (gleba: 1356 ± 453 ; peridium: 1336 ± 483), γ -*Proteobacteria* (gleba: 115 ± 82 ; peridium: 137 ± 43) and *Enterobacteriales* (gleba: 240 ± 210 ; peridium: 6 ± 3).

Concentration of thiophene volatiles in *T. borchii* gleba correlates with the bacterial abundance

Concentrations of thiophene volatiles have been reported to increase with the storage of truffle-fruiting bodies at room temperature (Bellesia *et al.*, 2001), suggesting a potential role of bacteria in the production of these volatile compounds. We hypothesized that the concentrations of

thiophene volatiles correlated with bacterial community composition and abundance within truffle-fruiting bodies. To test this hypothesis, thiophenes produced from the same gleba samples characterized by FISH were quantified by solid-phase microextraction–gas chromatography/mass spectrometry (SPME-GC/MS). Thiophene volatiles (1) and (2) were emitted all along the time course. The total density of *Eubacteria* were significantly correlated with the levels of thiophene volatile (2) only [volatile (1): Pearson $R^2 = 0.008$, $P = 0.686$; volatile (2): $R^2 = 0.220$, $P = 0.021$]. A significant correlation for volatiles (1) and/or (2) was also observed for the dominant α -*Proteobacteria* and a minor group representing *Bacteroidetes* (for α -*Proteobacteria*, volatile (1): $R^2 = 0.171$, $P = 0.045$; volatile (2): $R^2 = 0.172$, $P = 0.044$; for *Bacteroidetes*, volatile (1): $R^2 = 0.002$, $P = 0.821$; volatile (2): $R^2 = 0.303$, $P = 0.005$).

Bacteria isolated from fruiting bodies, but not *T. borchii* mycelia, are able to generate thiophene volatiles from *T. borchii*-fruiting bodies

The correlation between the evolution of bacterial communities during storage and the concentration of




Fig. 2. Localization of α - and β -*Proteobacteria* in thin sections of *T. borchii* peridium and gleba.

A,B. Hybridization of 30 μ m sections of *T. borchii* of peridium.

C,D. Hybridization of gleba (C,D) with the universal eubacteria Eub338 mix probe coupled to FITC (A, B, C) or cy3 (D) and with α -*Proteobacteria* specific probe coupled to cy3 (A, C) or β -*Proteobacteria* specific probe BET42A coupled to cy3 (B) or FITC (D) as observed by confocal microscopy.

White bars represent 10 μ m. asc., asci; b, bacterial cluster; myc., fungal cell; sp., spore. Each picture is representative of observations.

thiophene volatiles led us to hypothesize that some bacteria could be involved in the production of these compounds. To test this hypothesis, we tested the ability of bacteria isolated from *T. borchii*-fruiting bodies to produce thiophene compounds. Bacteria were isolated from soil adhering to the peridium and from the gleba of *T. borchii*-fruiting bodies on tryptic soy agar (TSA 3%) plates. Based on colony PCR and 16S rRNA sequences, the isolated strains belonged to β - and γ -*Proteobacteria*, *Bacteroidetes*, *Firmicutes* and *Actinobacteria* (Table 1). We were not able to isolate strains from the α -*Proteobacteria* phylum despite their high abundance in the fruiting body. Since those bacteria could also be involved in *T. borchii* aroma, we tested two *Rhizobiales* strains obtained from the German Collection of Microorganisms and Cell Cultures (Table 1).

A bioassay approach was used to test the ability of single bacterial strains to transform ethyl acetate extracts of truffle-fruiting bodies into thiophene volatiles (Supporting Information Fig. S2). Surprisingly, thiophene volatiles (1) and (2) were produced by all bacterial isolates but only in the presence of fruiting body extract (Fig. 3A and B). The amount of thiophene compounds produced varied between bacterial strains, *Proteobacteria* and more especially α -, β -*Proteobacteria* and *Enterobacteriales* among the γ -*Proteobacteria* being the most efficient producers (Fig. 3B). Surprisingly thiophene volatiles (1) and (2) were not only produced by bacteria isolated from *T. borchii*-fruiting bodies, but also by bacteria isolated from other sources (i.e. soil, plants, Table 1). The quantities of thiophene volatiles (1) and (2) released in our bioassays were similar to ~10 times lower to those detected from

Table 1. Bacteria tested in this study for their ability to produce thiophene volatiles. List of bacteria isolated from *T. borchii* (from TartufLangue, Cuneo, Italy, and collected 03/2008) and elsewhere.

Isolate reference in the present manuscript	Isolated from	Gleba or peridium/ adhering soil	GenBank Accession Number 16S rRNA	Bacterial phylum/class or order	Genus
A	<i>T. borchii</i>	Gleba	KO618433	Actinobacteria	<i>Microbacterium</i>
B	<i>T. borchii</i>	Gleba	KO618434	Bacteroidetes/Flavobacteria	<i>Chryseobacterium</i>
C	<i>T. borchii</i>	Peridium/adhering soil	KO618435	Bacteroidetes/Spingobacteria	<i>Spingobacterium</i>
D	<i>T. borchii</i>	Gleba	KO618436	Bacteroidetes/Spingobacteria	<i>Spingobacterium</i>
E	<i>T. borchii</i>	Peridium/adhering soil	KO618437	Firmicutes/Bacillales	<i>Brochothrix</i>
F	<i>T. borchii</i>	Peridium/adhering soil	KO618438	Firmicutes/Bacillales	<i>Brochothrix</i>
G	<i>T. borchii</i>	Peridium/adhering soil	KO618439	β -proteobacteria/Burkholderiales	<i>Comamonas</i>
H	<i>T. borchii</i>	Gleba	KO618440	β -proteobacteria/Burkholderiales	<i>Comamonas</i>
I	<i>T. borchii</i>	Peridium/adhering soil	KO618441	γ -proteobacteria/Enterobacteriales	<i>Serratia</i>
J (<i>Serratia plymuthica</i> IC14 ^b)	Soil (melon plantation)	–	–	γ -proteobacteria/Enterobacteriales	<i>Serratia</i>
K (<i>Serratia marcescens</i> MG1 ^a)	Cucumber (fruit)	–	–	γ -proteobacteria/Enterobacteriales	<i>Serratia</i>
L (<i>Serratia proteamaculans</i> B5a ^b)	Salmon	–	–	γ -proteobacteria/Enterobacteriales	<i>Serratia</i>
M	<i>T. borchii</i>	Peridium/adhering soil	KO618442	γ -proteobacteria/Pseudomonadales	<i>Pseudomonas</i>
N	<i>T. borchii</i>	Gleba	KO618443	γ -proteobacteria/Pseudomonadales	<i>Pseudomonas</i>
O	<i>T. borchii</i>	Peridium/adhering soil	KO618444	γ -proteobacteria/Xanthomonadales	<i>Stenotrophomonas</i>
P	<i>T. borchii</i>	Gleba	KO618445	γ -proteobacteria/Xanthomonadales	<i>Stenotrophomonas</i>
Q (<i>Ensifer adhaerens</i> strain LFG19a; DSMZ ref. 18131 ^a)	Not determined	–	–	α -proteobacteria/Rhizobiales	<i>Ensifer</i>
R (<i>Bosea thiooxidans</i> strain BI-42; DSMZ ref. 9653 ^a)	Agricultural Soil	–	AJ250796	α -proteobacteria/Rhizobiales	<i>Bosea</i>

a. Available from the German Collection of Microorganisms and Cell Cultures (www.dsmz.de).

b. Isolates of *Serratia* tested in an earlier report for their plant-growth-promoting properties (Blom *et al.*, 2011) (kindly provided by Dr. Laure Weisskopf, University of Zürich, Switzerland).




Fig. 3. Ability of bacteria and truffle mycelium to produce thiophene volatiles.

A. Chromatograms [total ion current (TIC) and extracted m/z 97] illustrating that bacteria but not truffle mycelium have the ability to transform truffle extract into two volatiles characteristic of *T. borchii*-fruiting bodies; volatile (1) is 3-methylthiophene and volatile (2) is 3-methyl-4,5(2H)thiophene. Chromatograms are shown with normalized intensities.

B. Levels ($\pm$ SE) of thiophene volatiles (1) and (2) (log scale normalized to growth) and corresponding growth ($\pm$ SE) (OD600) of different bacterial classes/groups ($n = 3$) (isolates are coded from A to R – See Table 1). Different letters denote values that differ significantly from each other (Fischer LSD post-hoc test $P < 0.05$) – ANOVA for 3-methylthiophene: $F = 15.94$, $P = 0.000000$; ANOVA for 3-methyl-4,5(2H)thiophene: $F = 6.26$, $P < 0.001$.

C. Quantities of thiophenes (1) and (2) released per SPME vial by *T. borchii* gleba ($n = 6$) and in the bioassays with bacteria ($n = 3$ per strain) described in Supporting Information Fig. S1. To make the values comparable between bioassays and gleba, thiophene quantities (extracted masses m/z 97 and m/z100) have been normalized to the TIC of the chromatograms.

T. borchii-fruiting bodies (Fig. 3C). Killed bacteria by autoclaving did not produce thiophene volatiles indicating that active metabolism was required (Fig. 3A).

To test whether the fungus was also able to produce thiophene volatiles by itself, volatile profiles of *in vitro* pure culture of *T. borchii* strain 43Bo isolated in 1996 were analysed. No thiophene volatiles were produced by *T. borchii* mycelial cultures *in vitro* (Fig. 3A). To make sure that the inability of *T. borchii* mycelial culture to produce thiophene volatiles was not a consequence of prolonged storage in pure culture of strain 43Bo, mycelium was freshly isolated from a fruiting body producing thiophene volatiles. The newly isolated strain IG2 was not able to produce any thiophene derivatives, confirming the previous results. Altogether these data indicate that bacteria but not truffle mycelium are able to biotransform the non-volatile precursors of *T. borchii* gleba into volatile thiophenes.

The precursor of thiophene volatiles is specific of the sexual stage of T. borchii and of no other truffle species

In a second step, we tested whether bacteria could produce thiophene volatiles when grown on *T. borchii* mycelium. Most surprisingly, even the most efficient thiophene-producer under our assay's conditions, an *Enterobacteriales* of the *Serratia* genus (Strain I, Table 1), failed to produce the thiophene volatiles typical of *T. borchii*-fruiting bodies when grown on sterilized *T. borchii* mycelium (strain 43Bo) that had been pre-grown in pure culture. Live mycelial cultures of *T. borchii* strain 43Bo also failed to produce thiophene volatiles on malt extract with or without addition of L-methionine, which had been suggested to be the precursor of thiophene volatiles in *T. borchii* (Zeppa *et al.*, 2004). Methionine, however, did induce in live mycelial cultures (strain 43Bo) the synthesis of numerous sulphur (non-thiophene) volatiles such as dimethyl sulphide and dimethyl disulphide (data not shown). These results suggest that mycelium of *T. borchii* does not produce the precursors of thiophenes *in vitro*. Furthermore, the *Serratia* isolate I (Table 1) produced the thiophene volatiles (1) and (2) on *T. borchii* gleba but not when grown on the gleba of other truffle species, including the white truffle *T. magnatum* or the black truffles *T. aestivum*, *T. mesentericum* and *T. brumale*.

Overall, these results indicate that thiophene production is linked to the sexual stage (fruiting body) of *T. borchii*. They also demonstrate that the aroma precursor of thiophene volatiles is/are unique to *T. borchii*.

Antibacterial but not antifungal agents inhibit the production of thiophene volatiles in T. borchii-fruiting bodies

To further investigate which organisms are responsible for the production of specific truffle-fruiting body volatiles, we

used antimicrobial agents targeting either fungi or bacteria. A homogenate of *T. borchii*-fruiting bodies was treated with aqueous solutions containing either the broad-spectrum antibacterial agent streptomycin, or one of the antifungal agents clotrimazole or amphotericin B, or pure water as a control. The concentrations of volatiles (1) and (2) were measured at the beginning of the experiment and after 48 h (Fig. 4). Production of thiophene volatiles (1) and (2) was only suppressed by the bactericide streptomycin (Fig. 4). The concentration of thiophenes (1) and (2) in samples treated with the fungicides (48 h, Fig. 4) increased to the same level as in the water control (Fig. 4), indicating that fungal metabolism was not necessary for the production of thiophene volatiles. Furthermore, the concentration of thiophene volatiles (1) and (2) induced by the different treatments correlated with the bacterial population density in the gleba samples (cultivable fraction of bacteria – Fig. 4).

Bacteria fed with methionine produce a volatile related in structure to the thiophenes characteristic of T. borchii-fruiting bodies

Interaction of fungi with bacteria might activate fungal biosynthetic pathways that are otherwise suppressed in pure mycelial cultures (Scherlach and Hertweck, 2009; Schroeckh *et al.*, 2009). We therefore co-cultured our *Serratia* bacterial isolate I (Table 1) with mycelia of white and black truffles, with or without an excess of L-methionine. Regardless of L-methionine addition, neither volatile (1) and (2) was detected. Surprisingly, another thiophene volatile, 2-methyltetrahydrothiophen-3-one [designated (3)], was detected in the presence of excess methionine not only from co-cultures of white or black truffle mycelia with *Serratia* but also from the single bacterial cultures without mycelium (Fig. 5). Cultures of truffle mycelia supplemented with extra methionine failed to produce volatile (3) when *Serratia* was absent (Fig. 5), confirming our hypothesis that the formation of thiophene derivatives by truffle-fruiting bodies depends on bacteria generating sulfur-containing heterocycles from linear precursor(s) such as L-methionine.

Discussion

Some fungi are hotspot for bacteria (Warmink *et al.*, 2009). Truffle-fruiting bodies do not derogate to this observation and are intensively colonized by bacteria that can reach densities up to 10^8 cells per gram of truffle (Barbieri *et al.*, 2005; 2007; Antony-Babu *et al.*, 2013). Similar to what has been previously described (Barbieri *et al.*, 2005), samples of *T. borchii* analysed here were intensively colonized by bacteria. In line with previous reports about *T. borchii* (Barbieri *et al.*, 2005),




Fig. 4. Effect of antimicrobial treatments on truffle volatiles. Mean levels ($\pm$ SD – log scale) of thiophene volatiles (1) and (2) and fraction of cultivable bacteria in fruiting body homogenate of *T. borchii* upon treatment with: A. H₂O/control or aqueous solutions of the antifungal agents clotrimazole B. Amphotericin B. C. Antibacterial agent streptomycin $n = 9$ replicates per treatment. Compared with the initial concentrations of thiophene volatiles released at the beginning of the experiment (H₂O/Control, 0h), synthesis of thiophene volatiles was only blocked by the antibacterial agent streptomycin (C). The increase in thiophene concentrations observed in the samples treated with the antifungal agents (A, B) for 48 h were equivalent to the H₂O/Control 48 h, demonstrating that truffle fungi are not involved in the production of thiophene volatiles. Different letters indicate significant differences among treatments ($P < 0.05$, ANOVA and Fischer LSD *post-hoc* test).

T. magnatum (Barbieri *et al.*, 2007) and *T. melanosporum* (Antony-Babu *et al.*, 2013), the bacterial community was dominated by α -*Proteobacteria*. These similarities between studies and between truffles species suggest that truffle-fruited bodies, regardless the species, provide a specific habitat for some bacteria. This core microbiome of truffle-fruited bodies might be supplemented with additional species depending on the fungal species, the maturation stage or the environment as suggested by the divergence observed for some classes. For example the abundance of β -*Proteobacteria* (Fig. 1) was much higher here compared with earlier reports on white truffles *T. borchii* (Barbieri *et al.*, 2005) and *T. magnatum* (Barbieri *et al.*, 2007).

α -*Proteobacteria* are very versatile in adapting to diverse environments ranging from the oceans (volcanoes on the ocean floor, surface waters) to soil (Ettema and Andersson, 2009). They also have an intriguing ability to interact with plants in either pathogenic or non-pathogenic mutualist/commensal relations (i.e. *Rhizobium*, *Azospirillum*). In fungi, the α -*Proteobacteria* phylum is




Fig. 5. Methionine induces thiophene derivative 2-methyltetrahydrothiophen-3-one in *Serratia*. A. Volatile (3) was not detected from *T. borchii* fruiting bodies. B. It was detected whenever the bacterium was present. C. It was not detected when the bacterium was absent. Chromatograms (extracted mass m/z 116) are shown with normalized intensities.

dominant not only in truffles (Barbieri *et al.*, 2005; 2007; Antony-Babu *et al.*, 2013), but also in the ectomycorrhizal fungus *Laccaria bicolor* (Bertaux *et al.*, 2005), suggesting the existence of some adaptative or competitive mechanism over other bacterial phyla. What creates the competitive advantage for α -*Proteobacteria* in truffles is unknown, but theoretically it might be driven by an increased ability to compete for specific resources (niche effect) or to degrade/produce toxic secondary metabolites (Hibbing *et al.*, 2010). A study on the genome of 92 α -*Proteobacteria* revealed major genetic differences between those whose lifestyles were associated to plants (symbiotic and non-symbiotic) compared with those that were not plant-associated (Pini *et al.*, 2011). By similarity, genetic adaptation might also exist in α -*Proteobacteria* associated to fungi and might possibly result from co-evolution between bacteria and their fungal hosts. In the specific case of the Périgord truffle *T. melanosporum*, the abundance of α -*Proteobacteria* was shown to increase in the truffle gleba over the period from a collection season (Antony-Babu *et al.*, 2013). Taken together with the information that nitrogen fixation actively takes place in truffle-fruited bodies of *T. magnatum* (Barbieri *et al.*, 2010), and that some members of the α -*Proteobacteria* are well known for their ability to fix nitrogen, this suggests that the dominance of the latter bacterial phylum in truffles might be driven by special nutritional requirements. Another possibility might be that α -*Proteobacteria* might be better at degrading antibacterial agents (maybe the precursor of thiophene derivatives) than other bacterial phyla. Both theories, only speculative at this stage, will need to be tested experimentally.

In addition, our work suggests the existence of a structuration of the bacterial communities within the fruiting bodies of *T. borchii*. As observed in *T. melanosporum* (Antony-Babu *et al.*, 2013), the composition of bacterial communities in the peridium diverged from the one of the gleba with an enrichment of bacteria belonging to the *Bacteroidetes* phylum in the peridium of both truffle species. Interestingly, our data also suggest that bacterial taxa could behave differently within the truffle with α -*Proteobacteria* tending to form 'pure' colonies, while β -*Proteobacteria* being present in mixed communities. These patterns suggest the existence of regulated network of interactions between bacteria in the truffle-fruited bodies. Mechanisms regulating these networks and their consequence on the community function remain to be explored.

If the presence of bacteria in truffle-fruited bodies is known since decades, the nature of the interactions occurring between bacterial communities and truffle fungi remain unclear. Do bacteria only consume truffle nutrients or do mutualistic interactions occur between the fungal host and the bacterial community? One of our aims was to

investigate if thiophene derivatives, volatiles characteristic of *T. borchii*'s aroma, were actually derived from the truffle fungus itself or from the microbiome inhabiting truffle-fruited bodies. By following an approach similar to Koch's postulate in microbiology, we demonstrated that the bacterial community inhabiting *T. borchii*-fruiting bodies was responsible for the production of thiophene derivatives (1) and (2) characteristic of the fruiting body aroma of the latter species. Such intimate interactions between bacteria and fungi have been documented earlier, for example in the case of the rice fungal pathogen *Rhizopus* which toxins are derived from endobacteria of the *Burkholderia* genus (Partida-Martinez and Hertweck, 2005). Unlike in the *Rhizopus/Burkholderia* case, the interaction in truffles is not limited to a single bacterial genus, and indeed all bacteria tested in this study were able to produce thiophene volatiles from *T. borchii*-fruiting body extract, irrespective of whether they had been isolated from truffles or from other sources (i.e. plant, soil). This indicates that the ability to produce thiophene volatiles is widespread among bacteria and might possibly be linked to primary metabolism. This also raises the question of the ecological role of thiophene volatiles in nature.

Methionine was suggested to be the precursor of thiophene volatiles in *T. borchii* (Zeppa *et al.*, 2004). Methionine did induce in our assays with *Serratia* volatile (3), a thiophene of related structure to volatiles (1) and (2), but not the latter two volatiles. There is no data in literature on the pathways or the genes leading to the synthesis of thiophene volatiles (1) and (2). However, data exist on the synthesis of volatile (3), both in yeast (Howell *et al.*, 2005) and in bacteria (Nawrath *et al.*, 2010). In the yeast *Saccharomyces cerevisiae*, volatile (3) might be derived from the cyclization and reorganization of 4-mercapto-4-methylpentan-2-one (4-MMP), itself released from the cleavage from a cystein moiety (Cys-4-MMP). Volatile (3) is furthermore strongly induced by the deletion of YAL012W, a putative carbon-sulfur lyase, suggesting an enzymatic biosynthesis route in yeast (Howell *et al.*, 2005). A BLAST search of YAL012W against 3271 expressed sequence tags of *T. borchii* (Lazzari *et al.*, 2007) did not reveal any match in the white truffle; this suggests that the latter gene might either not be expressed or might be absent in *T. borchii*. In the bacterium *Chitinophaga Fx7914*, volatile (3) is derived from two primary metabolites, homocysteine and pyruvate (Nawrath *et al.*, 2010). By analogy to the intermediates and pathways of the latter two examples, thiophene volatiles (1) and (2) might be derived in truffles from primary metabolites synthesized or transformed by truffles, but also by yeasts (Buzzini *et al.*, 2005) and bacteria.s

The production of volatiles (1) and (2) by bacteria from *T. borchii* gleba (or gleba extract) but not from other truffle species supports the inference that *T. borchii* is involved

in the synthesis of at least one intermediate leading to thiophene volatiles **(1)** and **(2)**. Additionally, the upregulation in sulfur metabolism in the fruiting bodies of *T. borchii* (Zeppa *et al.*, 2010) and *T. melanosporum* (Martin *et al.*, 2010; Splivallo and Maier, 2011; Splivallo *et al.*, 2011) provides indirect evidence that truffles are actively involved in the production of sulfur metabolites during their sexual stage. Indeed the upregulation in homocysteine synthase (*tbhos*) and putative sulfate transporter (*tbsul1*) in mature *T. borchii*-fruiting bodies suggests that these genes might be indirectly involved in the synthesis of volatiles (Zeppa *et al.*, 2010).

Overall, this strongly suggests that thiophene volatiles are derived from bacteria and further suggests that the non-volatile precursors of thiophene compounds might be produced by the intimate interactions of truffles, yeasts and bacteria. In contrast, other non-thiophene volatiles might be completely synthesized by the fungus itself. The complex blend of 20 to 50 molecules creating the aroma of *T. borchii* would thus originate from the mixed activity of the fungus and its microbiome.

Our results demonstrate that bacteria are central players in the production of thiophene volatiles **(1)** and **(2)** in *T. borchii*. The formation of other volatiles or the contribution of bacteria to aroma formation in other truffle species remains to be investigated. Similarly, the full biosynthetic pathway leading to thiophene volatiles and the exact contribution of yeasts and truffle remains to be elucidated. Our results open new horizons in the biotechnological production of fungal aromas suggesting that microbes might play a central role in the production of key fungal odorants (Splivallo and Maier, 2011). They also raise new ecological and mechanistic questions about the multitrophic interactions among truffles, their plant symbionts, mammals, insects and the bacterial community associated with truffle-fruiting bodies. By participating in the elaboration of sulphur volatile compounds that attract mammals, bacteria could indirectly participate in the dissemination of truffle spores and thus play a key role in the life cycle of the fungus.

Experimental procedures

Truffle-fruiting bodies

Truffle-fruiting bodies of *T. borchii*, *T. magnatum*, *T. aestivum*, *T. mesentericum* and *T. brumale* were purchased (Supporting Information Table S1) and were identified based on spore morphology (Ceruti *et al.*, 2003).

Truffle mycelial cultures

Mycelia of *T. borchii* isolated more than 15 years ago (strain 1Bo = ATCC 96540 and 43Bo) (Bonuso *et al.*, 2009) as well as a strain (strain IG2 – GenBank Accession KF414978)

freshly isolated from a *T. borchii*-fruiting body which produced volatiles **(1)** and **(2)** and *T. melanosporum* (strain TN1) were grown as described earlier (Splivallo *et al.*, 2012). For the time series and feeding experiment with methionine, mycelial cultures and culture-negative controls (no mycelium) in malt extract broth were homogenized with a blender and transferred to SPME vials (aliquots of 4.5 ml per vial). Vials were supplemented with 0.5 ml of an aqueous solution containing either L-methionine (final concentration in SPME vials 5 mM) or water (control). Samples were either incubated in the dark or under 16 h photoperiods (to reflect natural conditions in the soil) at 23°C. Volatiles were measured by SPME-GC/MS after 1, 2, 3, and 5 days of incubation as described hereafter.

Selection of bacterial strains

Bacteria were isolated from the surface (peridium and adhering soil) and the inner part (gleba) of six fresh mature *T. borchii*-fruiting bodies (purchased in 2008 from Tartuflangue, Cuneo, Italy). Unwashed truffles were shortly vortexed in a sterile 0.85% NaCl solution, and a ~300 mg piece of gleba was subsequently excised and homogenized in 0.85% NaCl. Bacteria were isolated from the NaCl solutions by dilution plating on TSA 3% (Barbieri *et al.*, 2005). A total of 29 strains were selected based on their colony morphology and colour (not shown). The full 16S rRNA was amplified by colony PCR using the universal primers 'UP-Forward' and 'UP-Reverse' (Barbieri *et al.*, 2000; 2005). Sequencing was performed from both ends by Macrogen Europe (Amsterdam, the Netherlands).

The ability of bacteria to produce thiophene volatiles was tested on the following isolates: (i) A sub-selection of 13 bacterial isolates from our 29 strains was done reflecting the typical bacterial diversity in *T. borchii* (Barbieri *et al.*, 2005) and included two strains of β -*Proteobacteria*, five strains of γ -*Proteobacteria*, three strains of the *Bacteroidetes*, two strains of the *Firmicutes* and one strain of the *Actinobacteria* (GenBank accession numbers are listed in Table 1). (ii) Five bacterial strains isolated from sources different from truffle-fruiting bodies (i.e. plant, soil) were also tested for their ability to produce thiophene volatiles. They included two strains of α -*Proteobacteria* and three strains of *Serratia* (*Enterobacteriales*) (Table 1). All bacterial isolates were stored at -70°C in 1:1 glycerol (85%) : tryptic soy broth (TSB). Before bioassays were performed, bacterial glycerol stocks were plated on TSA (3%) to check for purity.

Analysis of aroma profiles by SPME-GC/MS

Aroma of truffle gleba (300 mg), bacterial and mycelial cultures were profiled by SPME-GC/MS as described earlier (Splivallo *et al.*, 2012). The occurrence of thiophene volatiles 3-methylthiophene **(1)** and 3-methyl-4,5-dihydrothiophene **(2)** was confirmed in *T. borchii* samples collected independently in Piedmont ($n = 20$ samples collected in 2006, 2008 – See Supporting Information Table S1). The identity of all volatiles listed in this study was confirmed through their MS fragmentation patterns, Kovats indices and when available with synthetic standards (i.e. 3-methylthiophene and see also Supporting Information Table S2).

Bioassays with fruiting body ethyl acetate extract

To test the ability of bacteria isolated from *T. borchii*-fruiting bodies and of the other bacterial isolates from soil and plants (Table 1), as well as the ability of truffle mycelium to generate thiophene volatiles, 490 g *T. borchii*-fruiting bodies were freeze dried, homogenized in a mortar and extracted overnight with 2 × 400 ml hexane followed by 2 × 400 ml ethyl acetate. The ethyl acetate fractions (800 ml) were pooled, concentrated till dryness in a vacuum evaporator at 50°C and resuspended in 450 ml malt extract broth (1%, pH 7.0). A negative control containing only dried ethyl acetate was prepared in the same way. The malt extract solution was sterile filtered (0.22 µm) and transferred (900 µl) to sterile SPME vials. SPME vials were either inoculated with (i) 100 µl of actively growing bacterial cultures (single strains in TSB 30 g l⁻¹), with (ii) TSB only (negative control for the bacterial cultures) or with (iii) a pellet of 100 µl of *Serratia* Isolate I previously killed by autoclaving (another negative control for the bacterial cultures to make sure that active bacterial growth was required to produce thiophene volatiles) – see Supporting Information Fig. S1 for experimental design – or with (iv) *T. borchii* mycelium strain 43Bo or IG2, or with (v) malt extract broth (control for the mycelial cultures described in *Truffle mycelial cultures*). SPME vials containing bacterial cultures (and the malt extract control) were incubated under gentle vertical shaking at 15°C, reflecting spring soil temperatures typical of Italian regions where *T. borchii* truffles are collected. Furthermore, emission of thiophene volatiles and bacterial growth were determined after 25 h of incubation, which corresponds to the late exponential growth phase for our *Serratia* isolate I strain and the maximum emission of thiophene volatiles (1) and (2). SPME vials containing homogenized truffle mycelium were incubated for 25 h in the dark at 23°C (optimal growth condition for *T. borchii*) before generating volatile fingerprints.

Determination of bacterial growth

Bacterial growth was measured at 600 nm (OD600) after blank subtraction (100 µl of bacterial culture or malt extract broth per well) in an Epoch 96-well microtiter plate spectrophotometer (BioTeck Instruments, VT, USA).

Bioassays with dried fruiting body and mycelium homogenates

Truffle-fruiting bodies (1–3 g per species) of *T. borchii*, *T. magnatum*, *T. aestivum* (syn. *T. uncinatum*), *T. mesentericum*, *T. brumale* (Supporting Information Table S1) or mycelial cultures of *T. borchii* (strains 1Bo and 43Bo) (Bonuso *et al.*, 2009), and *T. melanosporum* (strain TN1) and negative culture controls (prepared as described in *Truffle mycelial cultures*), were freeze dried and each sample was homogenized separately to a fine powder in a mortar. The homogenates were sterilized with 10 ml CH₂Cl₂ (overnight incubation), after which CH₂Cl₂ was removed from the homogenates in a two-step process (vacuum evaporation at 50°C for 6 h followed by heating to 80°C for 3 h). A quantity of 50 mg of sterile fruiting body homogenate or mycelium homogenate was transferred to a sterilized 20 ml SPME vial,

which was then inoculated with either 100 µl bacterial glycerol stock (1:1 bacterium in TSB : 85% glycerol), or with 100 µl 1:1 TSB : 85% glycerol as control or with truffle mycelium following the same scheme described in *Bioassays with fruiting body ethyl acetate extract*. The detailed scheme of the growth conditions and volatile sampling is described in Supporting Information Fig. S2.

Bioassays with mixed cultures truffle mycelium/bacteria

A quantity of 2.0 g of truffle mycelial cultures (*T. borchii* strains 1Bo and 43Bo, and *T. melanosporum* strain TN1) and culture-negative controls (without mycelium), prepared as described earlier (Splivallo *et al.*, 2012), were transferred to 20 ml SPME vials and inoculated with 100 µl of *Serratia* sp. isolate I (identified as *Serratia* sp. based on 16S rRNA sequencing) glycerol stock (1:1 bacterium grown in TSB:85% glycerol) or with 100 µl 1:1 TSB : 85% glycerol as control. Samples in SPME vials were either incubated as such or supplemented with excess L-methionine (3 mg per SPME vials supplied in 100 µl H₂O) in order to check if excess methionine induced the production of thiophene volatiles. Samples were incubated for 48 h at 23°C in the dark before generating the aroma fingerprinting by SPME-GC/MS.

Treatment of fruiting bodies with antibacterial and antifungal agents

Two *T. borchii* truffles (~40 g in total) were homogenized in a mortar until obtaining a paste which was then transferred to 2.0 ml Eppendorf tubes containing each 300 mg truffle homogenate. Sterile water was added to the samples: 1.0 ml per tube of either pure water (control) or water containing the broad-spectrum antibacterial agent streptomycin (400 µg ml⁻¹) or one of the antifungal agents clotrimazole (400 µg ml⁻¹) or amphotericin B (250 µg ml⁻¹). Tubes were closed and incubated at 20°C on a rotary shaker (160 r.p.m.). After 48 h incubation, samples were centrifuged (2 min at 12 000 g), the aqueous phase was removed from the gleba homogenate, and the latter was then transferred to 20 ml SPME vials to quantify thiophene volatiles (1) and (2). Additionally, volatile fingerprinting was performed on 300 mg gleba homogenate at the beginning of the experiment (0 h) to measure the initial quantities of thiophene volatiles (1) and (2).

Fluorescence in situ hybridization (FISH)

Six fruiting bodies of *T. borchii* were washed with tap water, dried and cut using a sterile scalpel to generate four subsamples of 500 mg each per truffle-fruiting body. Samples were stored in 2.0 ml Eppendorf tubes and allowed to age at room temperature for 0 (samples processed immediately), 2, 4 and 6 days. Each subsample was divided in two parts, one 300 mg sample for volatile fingerprinting by SPME-GC/MS and one 200 mg sample which was fixed for subsequent FISH analysis. Fixation was performed in 2.0 ml Eppendorf tubes by incubating samples overnight at 4°C in 1200 µl of a 2.25% paraformaldehyde solution in phosphate saline buffer (PBS) – PBS = 8 g of NaCl, 0.2 g of KCl, 1.44 g of Na₂HPO₄, 0.24 g of KH₂PO₄ in 100 ml distilled H₂O, pH adjusted to 7.4.

Samples were washed with 3×1.0 ml ice-cold PBS after fixation and stored in 1:1 EtOH96% : PBS at -20°C until FISH analysis. FISH was performed on fruiting body homogenates (gleba samples at days 0, 2, 4 and 6 and peridium samples at day 0) or on thin sections of gleba and peridium samples ($30\ \mu\text{m}$) as described for *T. magnatum* (Barbieri *et al.*, 2007) with the probes listed in Supporting Information Table S3. Each sample was co-hybridized with the universal probe mix Eub338 coupled to FITC to quantify bacterial population and a phylum specific probe coupled to cy3 (Supporting Information Table S3). Signals representing all bacterial cells and specific phyla were quantified visually on homogenized samples after image acquisition under an epifluorescence microscope (BX41, Olympus). Each homogenized sample was mounted and hybridized independently two times and three images were acquired for each replicate (6 images in total). The mean bacterial cell number was normalized to the sample biomass and expressed as [cell count/mg dry weight of gleba]. Fluorescent signals representing *Firmicutes*, α -, β -, γ -*Proteobacteria*, *Bacteroidetes* and *Enterobacteriales* (Supporting Information Table S3) were counted directly under the fluorescence microscope and expressed as the percentage of *Eubacteria* (mean of $n \geq 3$ observation from the same sample). Values expressed as %*Eubacteria* were then transformed to [cell count / mg dry weight of gleba] after multiplication by the total bacterial cell count in the sample.

Thin sections of *T. borchii*-fruiting bodies ($30\ \mu\text{m}$) were hybridized with the probes specific for α - and β -*Proteobacteria* listed in Supporting Information Table S3 and observed with a Radiance 2100 Rainbow confocal microscope (Bio-Rad).

Data processing

All bioassays in SPME vials performed to check the capacity of bacteria or truffle mycelia to produce thiophene volatiles comprised at least three independent replicates. Treatment of *T. borchii* fruiting body with antimicrobial agents was performed on four replicates per treatment. Six *T. borchii*-fruiting bodies were used for FISH (subsamples at 0, 2, 4 and 6 days of aging).

Volatiles fingerprints were processed as described earlier (Splivallo *et al.*, 2012). Specific *m/z* typical of thiophene volatiles were furthermore extracted from the total ion chromatograms for data visualization and statistics (*m/z* 97 occurs both in compounds (1) and (2), while *m/z* 100 occurs in (2) only and *m/z* 106 in 2-methyltetrahydrothiophen-3-one (3) only – MS fragmentation patterns are shown in Supporting Information Fig. S3).

Acknowledgements

We would like to thank the following people for their support and for material provided: Prof. Ivo Feussner and Dr. Cornelia Herfurt for access to the GC/MS and Dr. Laure Weisskopf for providing the *Serratia* strains MG1, B5a and IC14 (Supporting Information Table S1); Prof. Alessandra Zambonelli for *T. borchii* mycelia strains 1Bo and 43Bo, and Dr. Gérard Chevalier for *T. melanosporum* mycelium strain TN1. Financial support was provided by the Deutsche Forschungsgemeinschaft – DFG (<http://www.dfg.de>, DFG

Grants SPL1191/2-1 and SPL1191/4-1 to RS). AD lab is supported by the Laboratory of Excellence ARBRE (ANR-11-LABX-0002-01).

Conflict of interest statement

RS declares that a patent has been filed regarding the production of truffle aroma using truffle-associated microbes (Splivallo and Maier, 2011). The other author(s) declare that they have no competing interests.

References

- Antony-Babu, S., Deveau, A., Van Nostrand, J.D., Zhou, J., Le Tacon, F., Robin, C., *et al.* (2013) Black truffle-associated bacterial communities during the development and maturation of *Tuber melanosporum* ascocarps and putative functional roles. *Environ Microbiol.* doi:10.1111/1462-2920.12294.
- Barbieri, E., Potenza, L., Rossi, I., Sisti, D., Giomaro, G., Rossetti, S., *et al.* (2000) Phylogenetic characterization and in situ detection of a Cytophaga-Flexibacter-Bacteroides phylogroup bacterium in *Tuber borchii* Vittad. Ectomycorrhizal mycelium. *Appl Environ Microbiol* **66**: 5035–5042.
- Barbieri, E., Bertini, L., Rossi, I., Ceccaroli, P., Saltarelli, R., Guidi, C., *et al.* (2005) New evidence for bacterial diversity in the ascoma of the ectomycorrhizal fungus *Tuber borchii* Vittad. *FEMS Microbiol Lett* **247**: 23–35.
- Barbieri, E., Guidi, C., Bertaux, J., Frey-Klett, P., Garbaye, J., Ceccaroli, P., *et al.* (2007) Occurrence and diversity of bacterial communities in *Tuber magnatum* during truffle maturation. *Environ Microbiol* **9**: 2234–2246.
- Barbieri, E., Ceccaroli, P., Saltarelli, R., Guidi, C., Potenza, L., Basaglia, M., *et al.* (2010) New evidence for nitrogen fixation within the Italian white truffle *Tuber magnatum*. *Fungal Biol* **114**: 936–942.
- Bellesia, F., Pinetti, A., Tirillini, B., and Bianchi, A. (2001) Temperature-dependent evolution of volatile organic compounds in *Tuber borchii* from Italy. *Flavour Fragrance J* **16**: 1–6.
- Bertaux, J., Schmid, M., Hutzler, P., Hartmann, A., Garbaye, J., and Frey-Klett, P. (2005) Occurrence and distribution of endobacteria in the plant-associated mycelium of the ectomycorrhizal fungus *Laccaria bicolor* S238N. *Environ Microbiol* **7**: 1786–1795.
- Blom, D., Fabbri, C., Connor, E.C., Schiestl, F.P., Klausner, D.R., Boller, T., *et al.* (2011) Production of plant growth modulating volatiles is widespread among rhizosphere bacteria and strongly depends on culture conditions. *Environ Microbiol* **13**: 3047–3058.
- Bonito, G.M., Gryganskyi, A.P., Trappe, J.M., and Vilgalys, R. (2010) A global meta-analysis of *Tuber* ITS rDNA sequences: species diversity, host associations and long-distance dispersal. *Mol Ecol* **19**: 4994–5008.
- Bonuso, E., Zambonelli, A., Bergemann, S.E., Iotti, M., and Garbelotto, M. (2009) Multilocus phylogenetic and coalescent analyses identify two cryptic species in the Italian bianchetto truffle, *Tuber borchii* Vittad. *Conserv Genet* **11**: 1453–1466.

- Buscot, F., Munch, J.C., Charcosset, J.Y., Gardes, M., Nehls, U., and Hampp, R. (2000) Recent advances in exploring physiology and biodiversity of ectomycorrhizas highlight the functioning of these symbioses in ecosystems. *FEMS Microbiol Rev* **24**: 601–614.
- Buzzini, P., Gasparetti, C., Turchetti, B., Cramarossa, M.R., Vaughan-Martini, A., Martini, A., *et al.* (2005) Production of volatile organic compounds (VOCs) by yeasts isolated from the ascocarps of black (*Tuber melanosporum* Vitt.) and white (*Tuber magnatum* Pico) truffles. *Arch Microbiol* **184**: 187–193.
- Ceruti, A., Fontana, A., and Nosenzo, C. (2003) *Le specie europee del genere Tuber: Una revisione storica* (Torino: Museo regionale di scienze naturali di Torino).
- Culleré, L., Ferreira, V., Chevrete, B., Venturini, M.E., Sánchez-Gimeno, A.C., and Blanco, D. (2010) Characterisation of aroma active compounds in black truffles (*Tuber melanosporum*) and summer truffles (*Tuber aestivum*) by gas chromatography–olfactometry. *Food Chem* **122**: 300–306.
- Ettema, T.J.G., and Andersson, S.G.E. (2009) The alpha-proteobacteria: the Darwin finches of the bacterial world. *Biol Lett* **5**: 429–432.
- Gryndler, M., Soukupová, L., Hřelová, H., Gryndlerová, H., Borovička, J., Streiblová, E., and Jansa, J. (2013) A quest for indigenous truffle helper prokaryotes. *Environ Microbiol Rep* **5**: 346–352.
- Hibbing, M.E., Fuqua, C., Parsek, M.R., and Peterson, S.B. (2010) Bacterial competition: surviving and thriving in the microbial jungle. *Nat Rev Microbiol* **8**: 15–25.
- Howell, K.S., Klein, M., Swiegers, J.H., Hayasaka, Y., Elsey, G.M., Fleet, G.H., *et al.* (2005) Genetic determinants of volatile-thiol release by *Saccharomyces cerevisiae* during wine fermentation. *Appl Environ Microbiol* **71**: 5420–5426.
- Lazzari, B., Caprera, A., Cosentino, C., Stella, A., Milanese, L., and Viotti, A. (2007) ESTuber db: an online database for *Tuber borchii* EST sequences. *BMC Bioinformatics* **8** (Suppl. 1): S13.
- Martin, F., Duplessis, S., Ditengou, F., Lagrange, H., Voiblet, C., and Lapeyrie, F. (2001) Developmental cross talking in the ectomycorrhizal symbiosis: signals and communication genes. *New Phytol* **151**: 145–154.
- Martin, F., Kohler, A., Murat, C., Balestrini, R., Coutinho, P.M., Jaillon, O., *et al.* (2010) Perigord black truffle genome uncovers evolutionary origins and mechanisms of symbiosis. *Nature* **464**: 1033–1038.
- Mauriello, G., Marino, R., DAuria, M., Cerone, G., and Rana, G.L. (2004) Determination of volatile organic compounds from truffles via SPME-GC-MS. *J Chromatogr Sci* **42**: 299–305.
- Maxmen, A. (2010) Truffle's savoury secret revealed. *Nat News*. doi:10.1038/news.2010.156.
- Mello, A., Murat, C., and Bonfante, P. (2006) Truffles: much more than a prized and local fungal delicacy. *FEMS Microbiol Lett* **260**: 1–8.
- Nawrath, T., Gerth, K., Müller, R., and Schulz, S. (2010) The Biosynthesis of the Aroma Volatile 2-Methyltetrahydrothiophen-3-one in the Bacterium *Chitinophaga Fx7914*. *Chembiochem* **11**: 1914–1919.
- Pacioni, G., Leonardi, M., Aimola, P., Ragnelli, A.M., Rubini, A., and Paolocci, F. (2007) Isolation and characterization of some mycelia inhabiting *Tuber ascomata*. *Mycol Res* **111**: 1450–1460.
- Partida-Martinez, L.P., and Hertweck, C. (2005) Pathogenic fungus harbours endosymbiotic bacteria for toxin production. *Nature* **437**: 884–888.
- Pavić, A., Stanković, S., Saljnikov, E., Krüger, D., Buscot, F., Tarkka, M., and Marjanović, Ž. (2013) Actinobacteria may influence white truffle (*Tuber magnatum* Pico) nutrition, ascocarp degradation and interactions with other soil fungi. *Fungal Ecol* **6**: 527–538.
- Pini, F., Galardini, M., Bazzicalupo, M., and Mengoni, A. (2011) Plant-bacteria association and symbiosis: are there common genomic traits in alphaproteobacteria? *Genes* **2**: 1017–1032.
- Read, D.J. (1991) Mycorrhizas in ecosystems. *Experientia* **47**: 376–391.
- Sbrana, C., Bagnoli, G., Bedini, S., Filippi, C., Giovannetti, M., and Nuti, M.P. (2000) Adhesion to hyphal matrix and antifungal activity of *Pseudomonas* strains isolated from *Tuber borchii* ascocarps. *Can J Microbiol* **46**: 259–268.
- Sbrana, C., Agnolucci, M., Bedini, S., Lepera, A., Toffanin, A., Giovannetti, M., and Nuti, M.P. (2002) Diversity of culturable bacterial populations associated to *Tuber borchii* ectomycorrhizas and their activity on *T. borchii* mycelial growth. *FEMS Microbiol Lett* **211**: 195–201.
- Scherlach, K., and Hertweck, C. (2009) Triggering cryptic natural product biosynthesis in microorganisms. *Org Biomol Chem* **7**: 1753–1760.
- Schroeckh, V., Scherlach, K., Nützmann, H.-W., Shelest, E., Schmidt-Heck, W., Schuemann, J., *et al.* (2009) Intimate bacterial-fungal interaction triggers biosynthesis of archetypal polyketides in *Aspergillus nidulans*. *Proc Natl Acad Sci USA* **106**: 14558–14563.
- Splivallo, R., and Maier, C. (2011) Production of natural truffle flavours from truffle mycelium (in French). Patent publication nb PCT/IB2010/052913.
- Splivallo, R., Bossi, S., Maffei, M., and Bonfante, P. (2007) Discrimination of truffle-fruited body versus mycelial aromas by stir bar sorptive extraction. *Phytochemistry* **68**: 2584–2598.
- Splivallo, R., Ottonello, S., Mello, A., and Karlovsky, P. (2011) Truffle volatiles: from chemical ecology to aroma biosynthesis. *New Phytol* **189**: 688–699.
- Splivallo, R., Valdez, N., Kirchhoff, N., Ona, M.C., Schmidt, J.-P., Feussner, I., and Karlovsky, P. (2012) Intraspecific genotypic variability determines concentrations of key truffle volatiles. *New Phytol* **194**: 823–835.
- Talou, T., Gaset, A., Delmas, M., Kulifaj, M., and Montant, C. (1990) Dimethyl sulphide: the secret for black truffle hunting by animals? *Mycol Res* **94**: 277–278.
- Warmink, J.A., Nazir, R., and van Elsas, J.D. (2009) Universal and species-specific bacterial ‘fungiphiles’ in the mycospheres of different basidiomycetous fungi. *Environ Microbiol* **11**: 300–312.
- Zeppa, S., Gioacchini, A.M., Guidi, C., Guescini, M., Pierleoni, R., Zambonelli, A., and Stocchi, V. (2004) Determination of specific volatile organic compounds synthesised during *Tuber borchii* fruit body development by

2660 R. Splivallo et al.

solid-phase microextraction and gas chromatography/mass spectrometry. *Rapid Commun Mass Spectrom* **18**: 199–205.

Zeppa, S., Marchionni, C., Saltarelli, R., Guidi, C., Ceccaroli, P., Pierleoni, R., *et al.* (2010) Sulfate metabolism in *Tuber borchii*: characterization of a putative sulfate transporter and the homocysteine synthase genes. *Curr Genet* **56**: 109–119.

Supporting information

Additional Supporting Information may be found in the online version of this article at the publisher's web-site:

Supporting Information Fig. S1. Schematic representation of the bioassay used to test the ability of bacteria to transform truffle-fruited body extract into thiophene volatiles.

Supporting Information Fig. S2. Schematic representation of the bioassay used to test the ability of bacteria to transform truffle-fruited body homogenate into thiophene volatiles.

Supporting Information Fig. S3. MS fragmentation pattern of thiophene volatiles.

Supporting Information Table S1. Countries/regions of origin and species of truffle-fruited bodies used in this study.

Supporting Information Table S2. Volatiles listed in this study and mode of identification.

Supporting Information Table S3. List of FISH probes used in this study.