GENE

LINDA HALL
AUG 15 2007
LIBRARY

ALSO AVAILABLE ONLINE:

http://www.elsevier.com/locate/gene

Full guide for authors available at this site

GENE

Volume 398 (2007)

Editor-in-Chief
Giorgio Bernardi
Naples, Italy

Managing Editor
Alberto Bernardi
Gif-sur-Yvette, France

ELSEVIER

AMSTERDAM – BOSTON – JENA – LONDON – NEW YORK – OXFORD – PARIS – PHILADELPHIA – SAN DIEGO – ST. LOUIS

GENE

AIMS AND SCOPE

The journal will publish papers on all structural, functional and evolutionary aspects of genes, chromatin, chromosomes and genomes.

EDITORIAL POLICY

The overriding criteria for publication are originality, high scientific quality, and interest to a wide audience of readers. Papers outside the scope of the journal, not having the requested scientific criteria of interest or quality, or not having been prepared according to the instructions outlined below, will be sent back to the authors without delay.

The articles published in *Gene* represent the opinions and scientific findings of the authors. Whilst the Publisher and Editors of *Gene* make every effort to ensure the accuracy of all published matter, they can accept no responsibility or liability, collectively or individually, for any erroneous, misleading or unintentionally damaging statements which may appear in the journal. Submission of a paper to Gene implies that it is not being published or submitted elsewhere (including symposia volumes or other types of publication).

The Editors assume that all new isolates and mutants described in papers published in *Gene* will be made freely available upon written request to all qualified members of the scientific community. If delays in strain or vector distribution are anticipated or if these are available from sources other than the authors, this should be indicated.

Gene requires all new nucleotide and amino acid sequences to be deposited. Obtain and include an accession number with all sequences. The most convenient method for submitting sequence data is by World Wide Web:

EMBL via Webin: http://www.ebi.ac.uk.embl/Submission/webin.html
GenBank via Bankit: http://www.ncbi.nlm.nih.gov/Bankit
DDBJ via Sakura: http://sakura.ddbj.nig.ac.jp/

Alternatively, the stand-alone submission tool 'Sequin' is available from the EBI at http://www3.ebi.ac.uk/Services/Sequin and from NCBI at http://www.ncbi.nlm.nih.gov/Sequin/

If requested the database will withhold release of data until publication.

Publication information. *Gene* (ISSN 0378-1119). For 2007, volumes 386–406 are scheduled for publication. Subscription prices are available upon request from the Publisher or from the Regional Sales Office nearest you or from this journal's web site (http://www.elsevier.com/locate/gene). Further information is available on this journal and other Elsevier products through Elsevier's website: (http://www.elsevier.com). Subscriptions are accepted on a prepaid basis only and are entered on a calendar year basis. Issues are sent by standard mail (surface within Europe, air delivery outside Europe). Priority rates are available upon request. Claims for missing issues should be made within six months of the date of dispatch.

Enquiries concerning manuscripts and proofs: Visit Elsevier's Author Gateway (http://authors.elsevier.com) for the facility to track accepted articles and set up e-mail alerts to inform you of when an article's status has changed. The Author Gateway also provides detailed artwork guidelines, copyright information, frequently asked questions and more.
Contact details for questions arising after acceptance of an article, especially those relating to proofs, are provided after registration of an article for publication.

Language Polishing. Authors who require information about language editing and copyediting services pre- and post-submission please visit http://www.elsevier.com/wps/find/authorshome.authors/languagepolishing or contact authorsupport@elsevier.com for more information. Please note Elsevier neither endorses nor takes responsibility for any products, goods or services offered by outside vendors through our services or in any advertising. For more information please refer to our Terms & Conditions http://www.elsevier.com/wps/find/termsconditions.cws_home/termsconditions.

ALSO AVAILABLE ONLINE:

http://www.elsevier.com/locate/gene

Full guide for authors available at this site

Gene is cited in: Biological Abstracts; Chemical Abstracts; Elsevier BIOBASE/Current Awareness in Biological Sciences; Current Contents (Life Sciences); EMBASE/Excerpta Medica; Index Medicus; Pascal M; Science Citation Index; Reference Update.

Protein on cover: space filling model of a *Narcissus pseudonarcissus* lectin monomer. Image reproduced with permission from CCLRC Daresbury Laboratory, United Kingdom. Structure originally published in J. Mol. Biol. 290 (1999) 185–199.

USA mailing notice: *Gene* (ISSN 0378-1119) is published twice a month by Elsevier B.V. (P.O. Box 211, 1000 AE Amsterdam, The Netherlands). Annual subscription price in the USA US$ 10,053.00 (valid in North, Central and South America), including air speed delivery. Periodical postage rate paid at Jamaica, NY 11431.
USA POSTMASTER: Send address changes to *Gene*, Elsevier, 6277 Sea Harbor Drive, Orlando, FL 32887-4800.
AIRFREIGHT AND MAILING in the USA by Mercury International Limited, 365 Blair Road, Avenel, NJ 07001.

Printed in the United Kingdom by Henry Ling Limited, at the Dorset Press, Dorchester, DT1 1HD

FOUNDED IN 1976 BY WACLAW SZYBALSKI

Case No.: IPR2023-00206
U.S. Patent No. 9,943,096

Vol. 398 Issues 1–2

GENE

15 August 2007

CONTENTS

Special Issue

Devoted to the XIVth International Conference on Dioxygen Binding and Sensing Proteins, Naples, Italy, 3–7 September 2006

Organizers and Editors: Luc Moens, Martino Bolognesi, Guido di Prisco and Cinzia Verde

Moens, L., Bolognesi, M., di Prisco, G. and Verde, C. 1
Editor's comment

Part 1: New structures in prokaryotic and eukaryotic hemoglobins

Nardini, M., Pesce, A., Milani, M. and Bolognesi, M. 2–11
Protein fold and structure in the truncated (2/2) globin family

Landfried, D.A., Vuletich, D.A., Pond, M.P. and Lecomte, J.T.J. 12–28
Structural and thermodynamic consequences of *b* heme binding for monomeric apoglobins and other apoproteins

Kuwada, T., Hasegawa, T., Sato, S., Sato, I., Ishikawa, K., Takagi, T. and Shishikura, F. 29–34
Crystal structures of two hemoglobin components from the midge larva *Propsilocerus akamusi* (Orthocladiinae, Diptera)

Marden, M.C., Dewilde, S., Kiger, L., Hamdane, D., Uzan, J., Burmester, T., Hankeln, T., Moens, L., Baudin-Creuza, V., Celier, C. and Wajcman, H. 35–41
Exploiting a list of protein sequences

Part 2: Bacterial hemoglobins and flavohemoglobins

Ascenzi, P., Bolognesi, M., Milani, M., Guertin, M. and Visca, P. 42–51
Mycobacterial truncated hemoglobins: From genes to functions

Bonamore, A., Attili, A., Arenghi, F., Catacchio, B., Chiancone, E., Morea, V. and Boffi, A. 52–61
A novel chimera: The "truncated hemoglobin-antibiotic monooxygenase" from *Streptomyces avermitilis*

Stevanin, T.M., Read, R.C. and Poole, R.K. 62–68
The *hmp* gene encoding the NO-inducible flavohaemoglobin in *Escherichia coli* confers a protective advantage in resisting killing within macrophages, but not *in vitro*: Links with swarming motility

Giordano, D., Parrilli, E., Dettaï, A., Russo, R., Barbiero, G., Marino, G., Lecointre, G., di Prisco, G., Tutino, L. and Verde, C. 69–77
The truncated hemoglobins in the Antarctic psychrophilic bacterium *Pseudoalteromonas haloplanktis* TAC125

Part 3: Plant and non-vertebrate hemoglobins

Garrocho-Villegas, V., Gopalasubramaniam, S.K. and Arredondo-Peter, R. 78–85
Plant hemoglobins: What we know six decades after their discovery

Hebelstrup, K.H., Igamberdiev, A.U. and Hill, R.D. 86–93
Metabolic effects of hemoglobin gene expression in plants

Part 4: Function of new globins in vertebrates (neuro- and cytoglobins)

Xi, Y., Obara, M., Ishida, Y., Ikeda, S. and Yoshizato, K. 94–102
Gene expression and tissue distribution of cytoglobin and myoglobin in the Amphibia and Reptilia: Possible compensation of myoglobin with cytoglobin in skeletal muscle cells of anurans that lack the myoglobin gene

Trandafir, F., Hoogewijs, D., Altieri, F., Rivetti di Val Cervo, P., Ramser, K., Van Doorslaer, S., Vanfleteren, J.R., Moens, L. and Dewilde, S. 103–113
Neuroglobin and cytoglobin as potential enzyme or substrate

Fordel, E., Thijs, L., Martinet, W., Schrijvers, D., Moens, L. and Dewilde, S. 114–122
Anoxia or oxygen and glucose deprivation in SH-SY5Y cells: A step closer to the unraveling of neuroglobin and cytoglobin functions

contents continued on last page of this issue

0378-1119(20070815)398:1/2;1-W

Available online at
ScienceDirect
www.sciencedirect.com

Contents (continued)

Part 5: Globin physiology and adaptation: evolution of oxygen transporters

Remington, N., Stevens, R.D., Wells, R.S., Holn, A., Dhungana, S., Taboy, C.H., Crumbliss, A.L., Henkens, R. and Bonaventura, C. 123–131
Genetic diversity of coastal bottlenose dolphins revealed by structurally and functionally diverse hemoglobins

Vinogradov, S.N., Hoogewijs, D., Bailly, X., Mizuguchi, K., Dewilde, S., Moens, L. and Vanfleteren, J.R. 132–142
A model of globin evolution

di Prisco, G., Eastman, J.T., Giordano, D., Parisi, E. and Verde, C. 143–155
Biogeography and adaptation of Notothenioid fish: Hemoglobin function and globin–gene evolution

Wittenberg, J.B. 156–161
On optima: The case of myoglobin-facilitated oxygen diffusion

Part 6: Globins and NO

De Rosa, M.C., Alinovi, C.C., Galtieri, A., Scatena, R. and Giardina, B. 162–171
The plasma membrane of erythrocytes plays a fundamental role in the transport of oxygen, carbon dioxide and nitric oxide and in the maintenance of the reduced state of the heme iron

Khan, A.A., Sun, Y., Jin, K., Mao, X.O., Chen, S., Ellerby, L.M. and Greenberg, D.A. 172–176
A neuroglobin-overexpressing transgenic mouse

Part 7: Structure, function and evolution of hemocyanin and hemerythrin

Paoli, M., Giomi, F., Hellmann, N., Jaenicke, E., Decker, H., Di Muro, P. and Beltramini, M. 177–182
The molecular heterogeneity of hemocyanin: Structural and functional properties of the 4 × 6-meric protein of *Upogebia pusilla* (Crustacea)

Decker, H., Schweikardt, T., Nillius, D., Salzbrunn, U., Jaenicke, E. and Tuczek, F. 183–191
Similar enzyme activation and catalysis in hemocyanins and tyrosinases

Giomi, F. and Beltramini, M. 192–201
The molecular heterogeneity of hemocyanin: Its role in the adaptive plasticity of Crustacea

Part 8: Ligand binding and dynamics in globins

Ronda, L., Faggiano, S., Bettati, S., Hellmann, N., Decker, H., Weidenbach, T. and Mozzarelli, A. 202–207
Hemocyanin from *E. californicum* encapsulated in silica gels: Oxygen binding and conformational states

Deng, P., Nienhaus, K., Palladino, P., Olson, J.S., Blouin, G., Moens, L., Dewilde, S., Geuens, E. and Nienhaus, G.U. 208–223
Transient ligand docking sites in *Cerebratulus lacteus* mini-hemoglobin

Bruno, S., Faggiano, S., Spyrakis, F., Mozzarelli, A., Cacciatori, E., Dominici, P., Grandi, E., Abbruzzetti, S. and Viappiani, C. 224–233
Different roles of protein dynamics and ligand migration in non-symbiotic hemoglobins AHb1 and AHb2 from *Arabidopsis thaliana*

Samuni, U., Dantsker, D., Roche, C.J. and Friedman, J.M. 234–248
Ligand recombination and a hierarchy of solvent slaved dynamics: The origin of kinetic phases in hemeproteins

This material may be protected by Copyright law (Title 17 U.S. Code)

ELSEVIER

Available online at www.sciencedirect.com
ScienceDirect

Gene 398 (2007) 78–85

GENE

www.elsevier.com/locate/gene

Review

Plant hemoglobins: What we know six decades after their discovery ☆

Verónica Garrocho-Villegas, Sabarinathan Kuttalingam Gopalasubramaniam, Raúl Arredondo-Peter *

Laboratorio de Biofísica y Biología Molecular, Facultad de Ciencias, Universidad Autónoma del Estado de Morelos, Ave. Universidad 1001, Col. Chamilpa, 62210 Cuernavaca, Morelos, México

Received 29 November 2006; received in revised form 30 January 2007; accepted 31 January 2007
Available online 25 April 2007

Abstract

This review describes contributions to the study of plant hemoglobins (Hbs) from a historical perspective with emphasis on non-symbiotic Hbs (nsHbs). Plant Hbs were first identified in soybean root nodules, are known as leghemoglobins (Lbs) and have been characterized in detail. It is widely accepted that a function of Lbs in nodules is to facilitate the diffusion of O_2 to bacteroids. For many years Hbs could not be identified in plants other than N_2-fixing legumes, however in the 1980s a Hb was isolated from the nodules of the non-legume dicot plant *Parasponia*, a *hb* gene was cloned from the non-nodulating *Trema*, and Hbs were detected in nodules of actinorhizal plants. Gene expression analysis showed that *Trema* Hb transcripts exist in non-symbiotic roots. In the 1990s nsHb sequences were also identified in monocot and primitive (bryophyte) plants. In addition to Lbs and nsHbs, Hb sequences that are similar to microbial truncated (2/2) Hbs were also detected in plants. Plant nsHbs have been characterized in detail. These proteins have very high O_2-affinities because of an extremely low O_2-dissociation constant. Analysis of rice Hb1 showed that distal His coordinates heme Fe and stabilizes bound O_2; this means that O_2 is not released easily from oxygenated nsHbs. Non-symbiotic *hb* genes are expressed in specific plant tissues, and overexpress in organs of stressed plants. These observations suggest that nsHbs have functions additional to O_2-transport, such as to modulate levels of ATP and NO.
© 2007 Elsevier B.V. All rights reserved.

Keywords: History; Leghemoglobin; Non-symbiotic; Truncated

1. Introduction

Hemoglobins (Hbs) are hemeproteins that reversibly bind O_2. The most common function of Hbs is associated with the transport of O_2, however Hbs also bind other gaseous ligands, such as NO (Cooper, 1999; Poole and Hughes, 2000; Moller and Skibsted, 2002; Dordas et al., 2003b; Milani et al., 2003; Frey and Kalio, 2005), and organic molecules (Ollesch et al., 1999; Bonamore et al., 2003; D'Angelo et al., 2004; Rinaldi et al., 2006), which suggests that they are multifunctional proteins in living organisms. Phylogenomic analysis showed that Hbs are widespread in organisms, as they have been identified in genomes of archaeobacteria, eubacteria and eukaryotes, including plants (Vinogradov et al., 2005, 2006). In plants three types of Hbs have been identified: symbiotic, non-symbiotic (nsHb) and truncated (2/2) Hbs (tHbs) (Ross et al., 2002). This review describes contributions to the study of plant Hbs from a historical perspective with emphasis on nsHbs. The authors refer interested readers to reviews by Appleby (1992), Fuchsman (1992), Arredondo-Peter et al. (1998), Hill (1998), Ross et al. (2002) and Kundu et al. (2003) for detailed discussion and relevant references on the structure, function and molecular biology of plant Hbs.

2. The discovery of plant Hbs

Plant Hbs were first identified by Kubo (1939). This author analyzed the red pigment of soybean root nodules. Nodules form in legumes after root infection by soil bacteria of the genera *Rhizobium*, *Bradyrhizobium*, *Azorhizobium*, *Mesorhizobium* and *Sinorhizobium* (collectively know as rhizobia). Nodules

Abbreviations: Hb, hemoglobin; HGT, horizontal gene transfer; Lb, leghemoglobin; Mb, myoglobin; nsHb, non-symbiotic hemoglobin; NO, nitric oxide; tHb, truncated (2/2) hemoglobin.

☆ Plenary lecture presented by RAP at the XIVth. International Conference on "Dioxygen Binding and Sensing Proteins", Naples, Italy, September 3–7, 2006.

* Corresponding author. Tel.: +52 777 329 7020; fax: +52 777 329 7040.
E-mail address: ra@buzon.uaem.mx (R. Arredondo-Peter).

0378-1119/$ - see front matter © 2007 Elsevier B.V. All rights reserved.
doi:10.1016/j.gene.2007.01.035

Case No.: IPR2023-00206
U.S. Patent No. 9,943,096

are red in color and are the plant organs where nitrogen fixation (by symbiotic rhizobia) occurs. Spectral examination of red pigment isolated by Kubo showed absorption bands that are characteristic of deoxy-ferrous, O_2-ferrous, CO-ferrous, and ferric Hbs. Kubo also crystallized hemin from this pigment and showed that crystals are identical to those of hemin from horse Hb. From this evidence Kubo concluded that the red pigment isolated from soybean root nodules was a hemeprotein.

To identify the physiological role of this hemeprotein in nodules, Kubo incubated nodule bacteria with it and observed stimulation of bacterial O_2-consumption. From this observation he concluded that the physiological role of the hemeprotein in nodules is to stimulate the assimilation and transport of O_2. The nodule hemeprotein discovered by Kubo has been characterized in detail by others. Biochemical, biophysical and molecular biology analyses have shown that Kubo's hemeprotein is a plant Hb with similar (*i.e.* structural) properties to animal Hbs (Fig. 1). Kubo's plant Hb was named as leghemoglobin (Lb) by Virtanen and Laine (1946) and is also known as plant symbiotic Hb.

3. The discovery of the function of Lb in root nodules

The role of Lb in N_2-fixing nodules was elucidated by Wittenberg, Bergersen, Appleby and Turner in 1974 (Wittenberg et al., 1974). These authors observed that the addition of Lb to suspensions of bacteroids enhanced the rate of O_2-uptake and nitrogenase activity (measured as the reduction of acetylene to ethylene), and promoted the formation of ethylene with increasing air O_2 concentrations. Ferric Lb, ferric horseradish peroxidase and ferric cytochrome *c* peroxidase were ineffective, however twelve O_2-binding proteins other than Lb also increased O_2-uptake and acetylene reduction by bacteroids. From these observations the authors concluded that oxyLb and other O_2-binding proteins facilitated the diffusion of O_2 to bacteroids in suspension, and that the function of Lb in nodules is to facilitate the diffusion of O_2 to bacteroids at an internal concentration too low to inhibit or destroy their O_2-sensitive nitrogenase. This concept of high O_2-flux at low free O_2 in the bacteroid vicinity is now generally accepted.



Fig. 1. Tertiary structure of Kubo's nodule hemeprotein (soybean leghemoglobin *a*) (gray) overlaid on the structure of spermwhale myoglobin (Mb) (black). Coordinates for the Lb*a* and Mb were obtained from the Brookhaven Protein Database using the identification numbers **1BIN** and **2MYC**, respectively, and structures were displayed using the PyMol program (http://www.pymol.org/funding.html). Helices (and CD loop in Lb*a*) are shown with the A–H letters.

4. The hypothesis for the origin of Lbs by horizontal gene transfer

For many years Hbs were identified in N_2-fixing legumes, but not elsewhere in the plant kingdom. Early attempts to detect Hbs in non-legumes using biochemical and molecular biology approaches were unsuccessful, so plant Hbs were apparently restricted to N_2-fixing legumes, and thus its origin was a mystery. In 1982 Jeffreys hypothesized that plant Hbs originated by a unique act of horizontal gene transfer (HGT) from a phytophagous insect to a primitive legume *via* a viral vector (Jeffreys, 1982). For this hypothesis to be correct the gene structure of insect *hb* and plant *lb* genes should be identical. Leghemoglobins are encoded by genes that are interrupted by three introns (Jensen et al., 1981; Hyldig-Nielsen et al., 1982). After Jeffreys' HGT hypothesis it was revealed that *hb* gene from the insect *Chironomus* contains no introns (in Appleby et al., 1988), thus the HGT hypothesis for the origin of plant Hbs was discarded, and the vertical evolution hypothesis of plant Hbs was favored. It is now known that other insect *hb* genes do contain introns, some in the exact positions found for legume *lb*s (Kloek et al., 1993; Hankeln et al., 2002; Hoogewijs et al., 2004); even so, other evidence in favor of vertical rather than horizontal evolution of plant *hb* genes soon emerged (see Section 5). An implication of the vertical evolution hypothesis is that Hbs might exist in all (legume and non-legume) land plants.

5. The vertical evolution of plant Hbs: discovery of plant non-symbiotic Hbs

A discovery that supported the vertical evolution hypothesis of plant Hbs was the identification of a Hb in the N_2-fixing root nodules of the non-legume dicot plant *Parasponia* (Appleby et al., 1983). *Parasponia* is a member of the Ulmaceae family, and the only non-legume known to be nodulated by *Rhizobium* strains. *Parasponia* Hb was extracted under anaerobic conditions and final purification was achieved by preparative isoelectric focusing, which produced a single major component of oxyHb (with pI of 6.28), and small amount of ferric Hb (with pI of 6.67). The absorption spectra of (deoxy-ferrous, O_2-ferrous and CO-ferrous) *Parasponia* Hb are similar to those of Lbs and animal Hbs and kinetic constants (O_2-affinity and O_2-association and -dissociation constants) are also similar in *Parasponia* Hb and soybean and other legume Lbs (Gibson et al., 1989). This evidence suggested that the function of Hb in *Parasponia* nodules is similar to function of Lb in legume nodules, *i.e.* to facilitate the diffusion of O_2 to bacteriods.

Motif Exhibit 1030, Page 7 of 13

Case No.: IPR2023-00206
U.S. Patent No. 9,943,096

During the 1980s a cDNA probe for *Parasponia* Hb was used to detect *hb* sequences in *Trema* and *Celtis*. *Trema* and *Celtis* are non-nodulating (ulmacean) plants closely related to *Parasponia*. Southern blot analysis revealed the existence of Hb-like sequences in *Trema* and (possibly) *Celtis* DNAs (Appleby et al., 1988; Boguz et al., 1988). DNA cloning and sequencing showed that *Trema* sequences have all the characteristics of a functional plant *hb* gene: the *Trema hb* gene is interrupted by three introns and exon sequences are 93% similar to *Parasponia hb*. Also, Northern blot analysis using the *Parasponia* Hb cDNA probe showed that Hb transcripts exist in *Trema* roots and in *Parasponia* roots and nodules, but not in *Trema* and *Parasponia* leaves. Western blot analysis using anti-*Parasponia* Hb antibodies showed that mRNA is translated into monomeric and dimeric Hbs in *Trema* roots and *Parasponia* roots and nodules, respectively (Boguz et al., 1988).

During the same decade Tjepkema detected high concentrations of Hb-like proteins in nodule extracts of actinorhizal plants (which are nodulated by the actinomycete *Frankia*), such as *Casuarina cunninghamiana* and *Myrica gale*, and low concentrations in nodules of *Comptonia peregrina*, *Alnus rubra* and *Eleagnus angustifolia* (Tjepkema, 1983). A few years latter after Tjepkema's report Hbs were purified and characterized from the actinorhizal *Casuarina glauca* (Fleming et al., 1987), *Alnus glutinosa* (Suharjo and Tjepkema, 1995), and *M. gale* (Pathirana and Tjepkema, 1995) root nodules. The spectral properties of actinorhizal Hbs are similar to those of Lbs and non-plant Hbs.

The detection of Hbs in *Parasponia*, *Trema* and actinorhizal plants suggested that plant Hbs are widespread in land plants and that plant *hb* genes vertically evolved from a common ancestor. Also, the detection of Hb transcripts in *Parasponia* and *Trema* roots showed that *hb* genes express and Hbs probably function in non-symbiotic plant organs.

cDNAs coding for monocot (barley and rice) nsHbs were cloned and sequenced in the 1990s. The first monocot Hb cDNA was cloned from barley (Taylor et al., 1994). The deduced amino acid sequence of barley Hb is 71% similar to *Parasponia* Hb. Southern blot analysis using the barley Hb cDNA probe showed that a single copy (or a very low copy number) of the *hb* gene exists in barley and that *hb*-like sequences exist in other monocot species, such as rye, maize and wheat. In 1997 Arredondo-Peter et al. reported the cloning and characterization of *hb1* and *2* genes from rice (Arredondo-Peter et al., 1997). Southern blot analysis showed that at least three copies of the *hb* gene exist in rice, however recent work showed that five copies of the *hb* gene exist in the rice genome (Lira-Ruan et al., 2002; Bustos-Rivera et al. manuscript in preparation). Sequence comparison showed that the predicted Hb1 and 2 proteins are 93% similar to each other, and that the rice Hbs are 68 to 82% similar to other non-symbiotic Hbs and about 50% similar to symbiotic Hbs (Arredondo-Peter et al., 1997).

Recently, Hb sequences were identified in very many evolved and primitive plants (Table 1), mostly because of numerous plant genome sequencing programs. For example, nsHb sequences have been identified in monocots such as maize and teosinte (Aréchaga-Ocampo et al., 2001) and wheat (Larsen, 2003), dicots such as soybean (Andersson et al., 1996), *Arabidopsis* (Trevaskis et al., 1997), chicory (Hendriks et al., 1998) and tomato (Wang et al., 2003), and bryophytes such as the moss *Physcomitrella patens* (Arredondo-Peter et al., 2000). The existence of nsHbs in the genome of primitive bryophytes suggests that the ancestor of land plant Hbs existed when the land was colonized by bryophytes, about 550 million years ago, and the existence of Hbs in primitive and evolved plants (Table 1) shows that Hbs are widespread in land plants. Moreover, *nshb* genes from the bryophytes *Marchantia polymorpha*, *Ceratodon purpureus* and *Physcomitrella patens* (GenBank accession numbers AY026341, EF028054 and EF028055, respectively) are interrupted by three introns in identical positions to introns of known plant *nshb* and *lb* genes. This evidence shows that *nshb* and *lb* genes evolved from an ancestral plant *hb* gene that was interrupted by three introns.

In addition to Lbs and nsHbs, Hb sequences that are similar to those of microbial truncated (2/2) Hbs (the reader is referred to reviews by Pesce et al. (2000) and Wittenberg et al. (2002) for a detailed description of the "truncated (2/2) folding" of globins) were detected in primitive and evolved plants (Table 1) (Watts et al., 2001). Sequence alignment showed that plant 2/2-like Hbs are highly conserved (not shown), and phylogenetic analysis showed that plant 2/2-like Hbs and Lbs and nsHbs evolved through different lineages (Fig. 2). Phylogenomic analysis by Vinogradov et al. (2006) revealed that plant 2/2-like Hbs are more similar to bacterial 2/2 Hbs than to plant (symbiotic and non-symbiotic) Hbs, which suggested that plant 2/2-like Hbs originated by a HGT event from a bacterial (*2/2 hb* gene) donor. Expression analysis showed that 2/2-like *hb* genes express in vegetative and embryonic organs and in organs from stressed plants (Hunt et al., 2001; Watts et al., 2001), however the function of 2/2-like Hbs in plant organs is not yet known, although kinetic properties of a recombinant *Arabidopsis* 2/2-like Hb suggest that these proteins may function as O_2-carriers (Watts et al., 2001).

6. Non-symbiotic (class 1) Hbs: hexacoordinate proteins with very high O_2-affinity

A distinctive characteristic of plant (class 1) nsHbs is that the affinity of these proteins for O_2 is very high (Duff et al., 1997; Arredondo-Peter et al., 1997; Trevaskis et al., 1997). For example, the O_2-affinity of rice Hb1 is 78-times higher than that from soybean Lb*a* (Arredondo-Peter et al., 1997). This high O_2-affinity of rice Hb1 is mainly due to a very low O_2-dissociation constant (k_{off}=0.038 s^{-1}), which suggests that O_2 is stabilized after binding to Hb1. Spectral analysis of deoxy-ferrous rice Hb1 showed that a distal ligand coordinates to Fe: the unliganded ferrous state of Hb1 exhibits peaks at 526 and 556 nm, which is similar to the absorption spectrum of (ferrous) cytochrome *b* where the heme Fe is hexacoordinate (Smith, 1978; Weiss and Ziganke, 1978). Histidine was identified as the distal ligand to heme Fe in wild type rice Hb1 by comparison with its mutant Hb1 H74L (in which distal His H74 was replaced by Leu). Absorbance spectra of mutant Hb1 showed no

Motif Exhibit 1030, Page 8 of 13

Case No.: IPR2023-00206
U.S. Patent No. 9,943,096

Table 1
Hemoglobin sequences identified in land plants

Organism	Hb[a] (accession number[b])
I. Seed plants	
a) Angiosperms	
Monocotyledonous	
Hordeum vulgare (barley)	nsHb-1 (U94968), tHb (AF376063)
Oryza sativa (rice)	nsHb1–1 (U76029), nsHb2–1 (U76031), nsHb3–1 (AF335504), nsHb4–1 (AF335504), tHb (BAD32857)
Saccharum spp. (sugarcane)	nsHb-1 (C0373680)
Triticum aestivum (wheat)	nsHb-1 (AY151390), tHb (AY151391)
Zea mays ssp. *mays* (maize)	nsHb-1 (AY005818), nsHb-2 (DQ171946)
Z. mays ssp. *parviglumis* (teosinte)	nsHb-1 (AF291052)
Dicotyledonous (legumes)	
Astragalus sinicus	symHb (DQ199647)
Canavalia lineata (bay bean)	symHb (UD9671)
Glycine max (soybean)	symHb (V00453), nsHb-1 (U47143), tHb (AA548191)
Lotus japonicus	symHb (AB238218), nsHb-1 (AB238220)
Lupinus luteus (yellow lupine)	symHb (Y00401)
Phaseolus vulgaris (kidney bean)	symHb (P02234)
Pisum sativum (pea)	symHb (P02233)
Psophocarpus tetragonolobus (winged bean)	symHb (X65874)
Medicago sativa (alfalfa)	symHb (M32883), nsHb-1 (AF172172)
M. trunculata (barrel medic)	symHb (X57733), tHb (ABE93362)
Sesbania rostrata	symHb (M23313)
Vicia faba (broad bean)	symHb (Z54159)
V. sativa (common vetch)	symHb (YO0229)
Vigna unguiculata (cowpea)	symHb I (U33206), symHb II (U33207)
Dicotyledonous (no-legumes)	
Alnus firma	nsHb-1 (AB221344)
Arabidopsis thaliana	nsHb-1 (NM_127165), nsHb-2 (NM_111887), tHb (NM_119421)
Beta vulgaris (sugar beet)	nsHb-2 (BE590299)
Brassica napus (canola)	nsHb-2 (AY026337)
Casuarina glauca (swamp oak)	nsHb-2 (X77694), nsHb-1(X53950)
Cichorium intybus × *C. endivia* (chicory)	nsHb1–2 (CAA07547), nsHb2–2 (AJ277797)
Citrus unshiu (satsuma orange)	nsHb-1 (AY026338)
Datisca glomerata (durango root)	tHb (CAD33536)
Euryale ferox (fox nut)	nsHb1–2 (AY281295), nsHb2–2 (AY281296)
Gossypium hirsutum (cotton)	nsHb-1 (AY899302), nsHb-2 (AY026340)
Lycopersicon esculentum (tomato)	nsHb-1 (AY026343), nsHb-2 (AY026344)
Malus domestica (apple)	nsHb-1 (AY224132)
Myrica gale	nsHb-1 (EF405885)
Nicotiana tabacum (tobacco)	nsHb-1 (BQ842804)

Table 1 (*continued*)

Organism	Hb[a] (accession number[b])
I. Seed plants	
a) Angiosperms	
Dicotyledonous (no-legumes)	
Parasponia andersonii	NsHb-1 (U27194)
P. rigida	nsHb-1 (P68169)
Pyrus communis (pear)	nsHb-1(AAY224133)
Raphanus sativus (horseradish)	nsHb-1 (AY286331)
Solanum tuberosum (potato)	nsHb2–1 (AY151389)
Trema orientalis	nsHb-1 (AF027215)
T. tomentosa	nsHb-1 (P07803)
T. virgata	nsHb-1 (AJ131349)
b) Gymnosperms	
Pinus tadea (pine)	tHb (BQ291271)
II. Bryophytes	
a) Mosses	
Physcomitrella patens	nsHb (AF218049)
Ceratodon purpureus	nsHb (AF309562)
b) Liverworts	
Marchantia polymorpha (liverwort)	nsHb (AY026341)

[a]nsHb, non-symbiotic Hb; nsHb-1, non-symbiotic Hb (class 1) with high O_2-affinity; nsHb-2, non-symbiotic Hb (class 2) with moderate O_2-affinity and/or sequence similarity to symbiotic Hbs; symHb, symbiotic Hb (including Lbs) with moderate O_2-affinity, which is specifically localized in nodules of N_2-fixing plants; tHb, 2/2-like plant Hb.
[b]Accession numbers in the GenBank database.

evidence of His coordination. Also, the substitution of H74 by L in Hb1 resulted in an increase of O_2-association and -dissociation constants. A conclusion from these observations was that distal His hexacoordinates to Fe and stabilizes bound O_2 in oxygenated Hb1 (Arredondo-Peter et al., 1997; Hargrove et al., 2000; Goodman and Hargrove, 2001).

7. Localization of nsHbs and expression of *nshb* genes in plant organs

Use of anti-rice Hb1 antibodies and Western blot showed that Hbs are localized in embryonic (coleoptiles, seminal roots and embryos) and (young and mature) vegetative (leaves and roots) rice organs. These observations showed that nsHbs are synthesized in different rice organs and during plant development (Lira-Ruan et al., 2001). However, analysis by confocal microscopy revealed that nsHbs are localized in the cytoplasm of differentiated and differentiating cell types of the developing rice seedling, such as the aleurone, scutellum, root cap cells, sclerenchyma, and tracheary elements. The immunolocalization of nsHbs in the vasculature of root, mesocotyl and leaf tissues, specifically in the xylem, suggests that nsHbs play roles in the metabolism accompanying the differentiation of the xylem (Ross et al., 2001).

Also, high levels of nsHb transcripts and proteins have been detected in organs and tissues of stressed plants. For example, *nshb* genes overexpress in barley aleurones subjected to microaerobiosis (<5% O_2) or sustained anoxia (up to 12 h) and in roots of flooded barley plants (Taylor et al., 1994). In

Motif Exhibit 1030, Page 9 of 13

Case No.: IPR2023-00206
U.S. Patent No. 9,943,096



Fig. 2. Cluster (phenetic) analysis of selected plant Hbs. Full Hb sequences (partial or incomplete Hb sequences were not included in this analysis) were obtained from the GenBank database using the accession numbers shown in Table 1. Hb sequences were aligned and phenogram was constructed using the Clustal W program (Thompson et al., 1997). The possible origin of plant 2/2-like Hbs by HGT is indicated according to Vinogradov et al. (2006). Abbreviations are the same as in Table 1. *Parasponia* Hbs were classified as nsHb-1 because they cluster with high O_2-affinity (class 1) nsHbs, however *Parasponia* Hbs are intermediate because they are localized in nodules and non-symbiotic roots and their O_2-affinity is similar to that of symHbs (Lbs).

etiolated rice levels of Hbs increased 2.4-fold compared to control plants grown under a normal light/dark regimen (Lira-Ruan et al., 2001), and levels of nsHb transcripts increased in microaerobic (0.1% O_2) and osmotically (1% sucrose) stressed *Arabidopsis* roots (Trevaskis et al., 1997). These observations suggest that nsHbs take part of the plant response to some stress conditions.

8. Multiple functions of plant nsHbs?

The kinetic constants (see Section 6) and localization of nsHbs in metabolically active tissues as well as the over-expression of *nshb* genes in stressed plants (see Section 7) suggest that nsHbs have functions other than or additional to O_2 transport. In 1998 it was proposed that nsHbs may play three

Case No.: IPR2023-00206
U.S. Patent No. 9,943,096

major roles in plant metabolism (Arredondo-Peter et al., 1998): 1) bind gaseous ligands, such as O_2, NO and CO, to modulate cell metabolism through interaction with other cell molecules, which could modify the rate constants for ligand binding; 2) bind small organic molecules, such as fatty acids, for transport and synthesis processes; and 3) scavenge O_2 based on the extremely tight binding of O_2 or interact with flavoproteins (for example with a ferric Hb reductase) for electron transfer.

Work aimed at the elucidation of nsHb function suggests that these proteins play roles in plant cells by maintaining levels of ATP and modulating levels of NO. Experiments with maize cells transformed with sense and antisense barley Hb cDNA (Hb^+ and Hb^- cells, respectively) showed that ATP levels decreased about 5, 40 and 65% when Hb^+, native (untransformed) and Hb^- maize cells were subjected to microaerobiosis, respectively (Sowa et al., 1998). Thus, it was concluded that a function of nsHbs is to maintain the energy *status* of cells under low O_2-concentrations. Also, work from the same laboratory showed that substantial amounts of NO were produced in alfalfa root cultures maintained under hypoxic (3% O_2) conditions, and that the amount of NO accumulated in alfalfa cells underexpressing Hb (Hb^- lines) was 2.5-fold higher than that in overexpressing barley Hb (Hb^+ lines) (Dordas et al., 2003a). It has been claimed that *Arabidopsis* nsHb1 scavenges NO and that it reduces levels of NO under hypoxic stress (Perazzolli et al., 2004). From these and others' observations it was concluded that a function of nsHbs is to modulate levels of NO (Dordas et al., 2003a; Perazzolli et al., 2004; Igamberdiev et al., 2005; Perazzolli et al., 2006) and, directly or indirectly, regulate a number of NO-dependent procceses in the plant cell.

9. Concluding remarks and future directions

Plant Hbs were described for the first time more than six decades ago. Since then substantial amounts of information about their structure, function and evolution have accumulated. However, there is still plenty of work to be done for a complete characterization and understanding of these plant proteins. For example, it is widely accepted that a function of Lb in nodules is to facilitate the diffusion of O_2 to the respiring bacteroids, however the mechanism for O_2-delivery from oxygenated Lb to the peribacteroid membrane is not yet known. It is well documented that Hbs interact with proteins (Low, 1986; Nagata et al., 1995; Park et al., 2002) and membrane lipids (Bonamore et al., 2003; Wakasugi et al., 2004; Rinaldi et al., 2006), thus it is of interest to evaluate whether or not Lb interacts with peribacteroidal membranes and if this interaction promotes O_2-delivery from oxygenated Lb. Also, primitive *nshb* genes could be selected within a particular plant family to help develop primitive symbiosis. Therefore, it is of interest to analyze primitive legumes to identify the ancestral *nshb/lb* that evolved into a functional symbiotic Hb.

Non-symbiotic Hbs have been characterized in detail. For example, *in vitro* analysis showed that O_2-affinity of (class 1) nsHbs is very high, mostly due to a very low O_2-dissociation constant. Thus, it is of interest to evaluate whether or not the O_2-affinity of nsHbs is also high *in vivo*, otherwise it is possible that nsHbs function as O_2-carriers or -sensors in plant tissues (Appleby et al., 1988). It is known that *nshb* genes (over)express in specific plant tissues (including organs from plants subjected to stress conditions), however details about the regulation of *nshb* gene expression are still not known. This information is essential to understand the role that nsHbs play in the physiology of a plant cell. A number of potential promoters have been identified upstream of the rice *hb1–4* genes (Lira-Ruan et al., 2002), thus it is likely that the regulation of *nshb* genes is complex. However, it has been shown that the expression of an *Arabidopsis nshb* gene is enhanced by the addition of the cytokinin 2iP (6-[γ-γ-dimethylallylamino]purine) (Hunt et al., 2001) and that a cytokinin-regulated element modulates the expression of rice *hb2* (Ross et al., 2004). This data shows that the expression of *nshb* genes is mediated by plant hormones, thus it is of interest to identify plant hormones that affect the synthesis of nsHbs and the role that these proteins play in the physiological/metabolic processes that are hormonally modulated.

Finally, compared to Lbs and nsHbs, little is known about the function of plant 2/2-like Hbs. However, it was postulated that these proteins originated (by HGT) from a bacterial tHb (Vinogradov et al., 2006), so it is likely that properties of plant 2/2-like Hbs and some bacterial tHbs are similar. Several lines of evidence suggest that a function of microbial tHbs is to detoxify NO (Ouellet et al., 2002; Milani et al., 2003; Giangiacomo et al., 2005; Sarma et al., 2005), thus it is of interest to evaluate whether or not plant 2/2-like Hbs participate in the metabolism of NO in plant cells. For example, (reduced) plant 2/2-like Hbs may bind O_2 and NO, dioxygenate NO and generate NO_3^- and ferric Hb. For this process, it is of interest to identify mechanisms that reduce ferric to ferrous plant 2/2-like Hb (which will subsequently bind O_2 for dioxygenation of NO), as has been documented for the reduction of ferric to ferrous Lbs (Lee and Klucas, 1984; Saari and Klucas, 1984, 1987; Klucas et al., 1988; Becana and Klucas, 1990; Ji et al., 1991).

In summary, the diversity of interactions of plant Hbs with molecules suggests that Hbs are multifunctional proteins that are essential for the functioning of plant cells. If this is true, Hbs have played an important role during the evolution and adaptation of plants to land environment.

Acknowledgements

Authors are grateful to an anonymous reviewer for corrections made to improve the English language of this article. Work in the authors' laboratory has been funded during the last decade by CoNaCyT (project nos. 25229N and 42873Q), Promep and DGAPA/PAPIIT-UNAM, México. V.G.V. and S.K.G. are postdoctoral fellows financed by CoNaCyT (IdAP 9272 and 8816, respectively), México.

References

Andersson, C.R., Jensen, E.O., Llewellyn, D.J., Dennis, E.S., Peacock, W.J., 1996. A new hemoglobin gene from soybean: a role for hemoglobin in all plants. Proc. Natl. Acad. Sci. U.S.A. 93, 5682–5687.

Appleby, C.A., 1992. The origin and functions of haemoglobin in plants. Sci. Prog. 76, 365–398.

Appleby, C.A., Bogusz, D., Dennis, E.S., Peacock, W.J., 1988. A role for hemoglobin in all plant roots? Plant Cell Environ. 11, 359–367.

Case No.: IPR2023-00206
U.S. Patent No. 9,943,096

Appleby, C.A., Tjepkema, J.D., Trinick, M.J., 1983. Hemoglobin in a nonleguminous plant *Parasponia*: possible genetic origin and function in nitrogen fixation. Science 220, 951–953.

Aréchaga-Ocampo, E., Sáenz-Rivera, J., Sarath, G., Klucas, R.V., Arredondo-Peter, R., 2001. Cloning and expression analysis of hemoglobin genes from maize (*Zea mays* ssp. *mays*) and teosinte (*Zea mays* ssp. *parviglumis*). Biochim. Biophys. Acta N, Gene Struct. Expr. 1522, 1–8.

Arredondo-Peter, R., Hargrove, M.S., Moran, J.F., Sarath, G., Klucas, R.V., 1998. Plant hemoglobins. Plant Physiol. 118, 1121–1126.

Arredondo-Peter, R., et al., 1997. Rice hemoglobins: gene cloning, analysis and oxygen-binding kinetics of a recombinant protein synthesized in *Escherichia coli*. Plant Physiol. 115, 1259–1266.

Arredondo-Peter, R, Ramírez, M, Sarath, G, Klucas, RV, 2000. Sequence analysis of an ancient hemoglobin cDNA isolated from the moss *Physcomitrella patens* (Accession No. AF218049). Plant Physiol. 122, 1457.

Becana, M., Klucas, R.V., 1990. Enzymatic and nonenzymatic mechanisms for ferric leghemoglobin reduction in legume root nodules. Proc. Natl. Acad. Sci. U.S.A. 87, 7295–7299.

Boguz, D., Appleby, C.A., Dennis, E.S., Trinick, W.J., Trinick, M., 1988. Functioning haemoglobin genes in non-nodulating plants. Nature 331, 178–180.

Bonamore, A., Farina, A., Gattoni, M., Schinina, M.E., Bellelli, A., Boffi, A., 2003. Interaction with membrane lipids and heme ligand binding properties of *Escherichia coli* flavohemoglobin. Biochemistry 42, 5792–5801.

Cooper, C.E., 1999. Nitric oxide and iron proteins. Biochim. Biophys. Acta 1411, 290–309.

Dordas, C., Hasinoff, B.B., Igamberdiev, A.U., Manasc'h, N., Rivoal, J., Hill, R.D., 2003a. Expression of a stress-induced hemoglobin affects NO levels produced by alfalfa root cultures under hypoxic stress. Plant J. 35, 763–770.

Dordas, C., Rivoal, J., Hill, R.D., 2003b. Plant hemoglobins, nitric oxide and hypoxic stress. Ann. Bot. 91, 173–178.

Duff, S.M.G., Wittenberg, J.B., Hill, R.D., 1997. Expression, purification and properties of recombinant barley (*Hordeum* sp.) hemoglobin: optical spectra and reactions with gaseous ligands. J. Biol. Chem. 272, 16746–16752.

D'Angelo, P., et al., 2004. Unusual heme iron-lipid acyl chain coordination in *Escherichia coli* flavohemoglobin. Biophys. J. 86, 3882–3892.

Fleming, A.I., Wittenberg, J.B., Wittenberg, B.A., Dudman, W.F., Appleby, C.A., 1987. The purification, characterization and ligand-binding kinetics of hemoglobin from root nodules of the non-leguminous *Casuarina glauca*-*Frankia* symbiosis. Biochim. Biophys. Acta 911, 209–220.

Frey, A.D., Kalio, P.T., 2005. Nitric oxide detoxification — a new era for bacterial globins in biotechnology? Trends Biotechnol. 23, 69–73.

Fuchsman, W., 1992. Plant hemoglobins. Adv. Comp. Environ. Physiol. 13, 23–58.

Giangiacomo, L., Ilari, A., Boffi, A., Morea, V., Chiancone, E., 2005. The truncated oxygen-avid hemoglobin from *Bacillus subtilis*. J. Biol. Chem. 280, 9192–9202.

Gibson, Q.H., Wittenberg, J.B., Wittenberg, B.A., Bogusz, D., Appleby, C.A., 1989. The kinetics of ligand binding to plant hemoglobins. J. Biol. Chem. 264, 100–107.

Goodman, M.D., Hargrove, M.S., 2001. Quaternary structure of rice nonsymbiotic hemoglobin. J. Biol. Chem. 276, 6834–6839.

Hankeln, T., et al., 2002. Characterization of *Drosophila* hemoglobin. J. Biol. Chem. 277, 29012–29017.

Hargrove, M., et al., 2000. Crystal structure of a non-symbiotic hemoglobin. Structure 8, 1005–1014.

Hendriks, T., et al., 1998. A nonsymbiotic hemoglobin gene is expressed during somatic embryogenesis in *Cichorium*. Biochim. Biophys. Acta 1443, 193–197.

Hill, R.D., 1998. What are hemoglobins doing in plants? Can. J. Microbiol. 76, 707–712.

Hoogewijs, D., et al., 2004. Genome-wide analysis of the globin gene family of *Caenorhabditis elegans*. IUBMB Life 56, 697–702.

Hunt, P.W., et al., 2001. Expression and evolution of functionally distinct haemoglobin genes in plants. Plant Mol. Biol. 47, 677–692.

Hyldig-Nielsen, J.J., et al., 1982. The primary structures of two leghemoglobin genes from soybean. Nucleic Acids Res. 10, 689–701.

Igamberdiev, A.U., Baron, K., Manac'h-Little, N., Stoimenova, M., Hill, R.D., 2005. The haemoglobin/nitric oxide cycle: involvement in flooding stress and effects on hormone signaling. Ann. Bot. 96, 557–564.

Jeffreys, A.J., 1982. Evolution of globin genes. In: Dover, G.A., Flavell, R.B. (Eds.), Genome Evolution. Academic Press, New York, pp. 157–176.

Jensen, E.O., Paludan, K., Hyldig-Nielsen, J.J., Jorgensen, P., Marcker, K.A., 1981. The structure of a chromosomal leghaemoglobin gene from soybean. Nature 291, 677–679.

Ji, L., Wood, S., Becana, M., Klucas, R.V., 1991. Purification and characterization of soybean root nodule ferric leghemoglobin reductase. Plant Physiol. 96, 32–37.

Kloek, A.P., Sherman, D.R., Goldberg, D.E., 1993. Novel gene structure and evolutionary context of *Caenorhabditis elegans* globin. Gene 129, 215–221.

Klucas, R., Ramesh, V., Shearman, L., Saari, L., 1988. Ferric leghemoglobin reductase from soybean root nodules. In: Bothe, H., deBrujin, F.J., Newton, W.E. (Eds.), Nitrogen Fixation: Hundred Years After. Gustav Fischer Verlag, Stuttgart, p. 637.

Kubo, H., 1939. Über hämoprotein aus den wurzelknöllchen von leguminosen. Acta Phytochim. (Tokyo) 11, 195–200.

Kundu, S., Trent III, J.T., Hargrove, M.S., 2003. Plants, humans and hemoglobins. Trends Plant Sci. 8, 387–393.

Larsen, K., 2003. Molecular cloning and characterization of cDNAs encoding hemoglobin from wheat (*Triticum aestivum*) and potato (*Solanum tuberosum*). Biochim. Biophys. Acta 1621, 299–305.

Lee, K.K., Klucas, R.V., 1984. Reduction of ferric leghemoglobin in soybean root nodules. Plant Physiol. 74, 984–988.

Lira-Ruan, V., Ross, E., Sarath, G., Klucas, R.V., Arredondo-Peter, R., 2002. Mapping and analysis of a hemoglobin gene family from rice (*Oryza sativa*). Plant Physiol. Biochem. 40, 199–202.

Lira-Ruan, V., Sarath, G., Klucas, R.V., Arredondo-Peter, R., 2001. Synthesis of hemoglobins in rice (*Oryza sativa* var. Jackson) plants growing in normal and stress conditions. Plant Sci. 161, 279–287.

Low, P.S., 1986. Structure and function of the cytoplasmic domain of band 3: center of erythrocyte membrane–peripheral protein interactions. Biochim. Biophys. Acta 864, 145–167.

Milani, M., Pesce, A., Ouellet, H., Guertin, M., Bolognesi, M., 2003. Truncated hemoglobins and nitric oxide action. IUBMB Life 55, 623–627.

Moller, J.K.S., Skibsted, L.H., 2002. Nitric oxide and myoglobins. Chem. Rev. 102, 1167–1178.

Nagata, K., Okano, Y., Nozawa, Y., 1995. Possible interaction of haemoglobin with a low Mr GTP-binding protein, ram p25. Biochem. Mol. Biol. Int. 35, 507–515.

Ollesch, G., Kaunzinger, A., Juchelka, D., Schubert-Zsilavecz, M., Ermler, U., 1999. Phospholipid bound to the flavohemoprotein from *Alcaligenes eutrophus*. Eur. J. Biochem. 262, 396–405.

Ouellet, H., et al., 2002. Truncated hemoglobin HbN protects *Mycobacterium bovis* from nitric oxide. Proc. Natl. Acad. Sci. U.S.A. 99, 5902–5907.

Park, K.W., Kim, K.J., Howard, A.J., Stark, B.C., Webster, D.A., 2002. *Vitreoscilla* hemoglobin binds to subunit I of cytochrome *bo* ubiquinol oxidases. J. Biol. Chem. 277, 33334–33337.

Pathirana, S.M., Tjepkema, J.D., 1995. Purification of hemoglobin from the actinorhizal root of *Myrica gale* L. Plant Physiol. 107, 827–831.

Perazzolli, M., et al., 2004. *Arabidopsis* nonsymbiotic hemoglobin AHb1 modulates nitric oxide bioactivity. Plant Cell 16, 1–10.

Perazzolli, M., Romero-Puertas, M.C., Delledonne, M., 2006. Modulation of nitric oxide bioactivity by plant haemoglobins. J. Exp. Bot. 57, 479–488.

Pesce, A., et al., 2000. A novel two-over-two α-helical sandwich fold is characteristic of the truncated hemoglobin family. EMBO J. 19, 2424–2434.

Poole, R.K., Hughes, M.N., 2000. New functions for the ancient globin family: bacterial responses to nitric oxide and nitrosative stress. Mol. Microbiol. 36, 775–783.

Rinaldi, A.C., Bonamore, A., Macone, A., Boffi, A., Bozzi, A., Giulio, A.D., 2006. Interaction of *Vitreoscilla* hemoglobin with membrane lipids. Biochemistry 45, 4069–4076.

Ross, E.J.H., Lira-Ruan, V., Arredondo-Peter, R., Klucas, R.V., Sarath, G., 2002. Recent insights into plant hemoglobins. Rev. Plant Biochem. Biotechnol. 1, 173–189.

Ross, E.J.H., et al., 2001. Non-symbiotic hemoglobins are synthesized during germination and in differentiating cell types. Protoplasma 218, 125–133.

Ross, E.J.H., Stone, J.M., Elowsky, C.G., Arredondo-Peter, R., Klucas, R.V., Sarath, G., 2004. Activation of the *Oryza sativa* non-symbiotic

Motif Exhibit 1030, Page 12 of 13

Case No.: IPR2023-00206
U.S. Patent No. 9,943,096

haemoglobin-2 promoter by the cytokinin-regulated transcription factor, ARR1. J. Exp. Bot. 55, 1721–1731.

Saari, L.L., Klucas, R.V., 1984. Ferric leghemoglobin reductase from soybean root nodules. Arch. Biochem. Biophys. 231, 102–113.

Saari, L.L., Klucas, R.V., 1987. Nonenzymatic reduction of ferric leghemoglobin. Biochim. Biophys. Acta 912, 198–202.

Sarma, H., Sharma, B.K., Tiwari, S.C., Mishra, A.K., 2005. Truncated hemoglobins: a single structural motif with versatile functions in bacteria, plants and unicellular eukaryotes. Symbiosis 39, 151–158.

Smith, L., 1978. Bacterial cytochromes and their spectral characterization. Methods Enzymol. 53, 202–212.

Sowa, A.W., Duff, S.M.G., Guy, P.A., Hill, R.D., 1998. Altering hemoglobin levels changes energy status in maize cells under hypoxia. Proc. Natl. Acad. Sci. U.S.A. 95, 10317–10321.

Suharjo, U.K.J., Tjepkema, J.D., 1995. Occurrence of hemoglobin in the nitrogen-fixing root nodules of *Alnus glutinosa*. Physiol. Plant. 95, 247–252.

Taylor, E.R., Nie, X.Z., MacGregor, A.W., Hill, R.D., 1994. A cereal haemoglobin gene is expressed in seed and root tissues under anaerobic conditions. Plant Mol. Biol. 24, 853–862.

Thompson, J.D., Gibson, T.J., Plewniak, F., Jeanmougin, F., Higgins, D.G., 1997. The clustal X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. Nucleic Acids Res. 24, 4876–4882.

Tjepkema, J.D., 1983. Hemoglobins in the nitrogen-fixing root nodules of actinorhizal plants. Can. J. Bot. 61, 2924–2929.

Trevaskis, B., et al., 1997. Two hemoglobin genes in *Arabidopsis thaliana*: the evolutionary origins of leghemoglobins. Proc. Natl. Acad. Sci. U.S.A. 94, 12230–12234.

Vinogradov, S.N., et al., 2006. A phylogenomic profile of globins. BMC Evol. Biol. 6, 31–47.

Vinogradov, S.N., et al., 2005. Three globin lineages belonging to two structural classes in genomes from the three kingdoms of life. Proc. Natl. Acad. Sci. U.S.A. 102, 11385–11389.

Virtanen, A.I., Laine, T., 1946. Red, brown and green pigments in leguminous root nodules. Nature 1157, 25–26.

Wakasugi, K., Nakano, T., Kitatsuji, C., Morishima, I., 2004. Human neuroglobin interacts with flotillin-1, a lipid raft microdomain-associated protein. Biochem. Biophys. Res. Commun. 318, 453–460.

Wang, Y.H., Kochian, L.V., Doyle, J.J., Garvin, D.F., 2003. Two tomato non-symbiotic haemoglobin genes are differentially expressed in response to diverse changes in mineral nutrient status. Plant Cell Environ. 26, 673–680.

Watts, R.A., Hunt, P.W., Hvitved, A.N., Hargrove, M.S., Peacock, W.J., Dennis, E.S., 2001. A hemoglobin from plants homologous to truncated hemoglobins of microorganisms. Proc. Natl. Acad. Sci. U.S.A. 98, 10119–10124.

Weiss, H., Ziganke, B., 1978. Purification of cytochrome *b* from *Neurospora crassa* and other sources. Methods Enzymol. 53, 212–221.

Wittenberg, J.B., Bergersen, F.J., Appleby, C.A., Turner, G.L., 1974. Facilitated oxygen diffusion: the role of leghemoglobin in nitrogen fixation by bacteroids isolated from soybean root nodules. J. Biol. Chem. 249, 4057–4066.

Wittenberg, J.B., Bolognesi, M., Wittenberg, B.A., Guertin, M., 2002. Truncated hemoglobins: a new family of hemoglobins widely distributed in bacteria, unicellular eukaryotes, and plants. J. Biol. Chem. 277, 871–874.