
VOLUME 46 JANUARY 1998 NUMBER 1
JOURNAL OF AGRICULTURAL AND FOOD CHEMISTRY
PUBLISHED BY THE AMERICAN CHEMICAL SOCIETY
AVAILABLE ON THE WEB
http://pubs.acs.org

JOURNAL OF AGRICULTURAL AND FOOD CHEMISTRY

EDITOR

IRVIN E. LIENER

Department of Biochemistry, College of Biological Sciences
University of Minnesota, 1479 Gortner Avenue
St. Paul, Minnesota 55108
(612) 625-8244; Fax (612) 624-5358
E-mail jafc@gold.tc.umn.edu

ASSOCIATE EDITORS

G. Wayne Ivie
U.S. Department of Agriculture

Marshall Phillips
Strategic Biocon

ADVISORY BOARD

Terry E. Acree
Cornell University

Eric Block
State University of New York at Albany

Daniel R. Caldwell
University of Wyoming

John W. Finley
Monsanto Company

Paul Hedin
U.S. Department of Agriculture

Chi-Tang Ho
Rutgers University

Robert J. McGorrin
Kraft General Foods

Steven Nagy
Florida Department of Citrus

Sara Risch
Science by Design

Russell Rouseff
Florida Department of Citrus

Richard Scanlan
Oregon State University

Walter Wolf
U.S. Department of Agriculture

STEENBOCK
MEMORIAL LIBRARY
FEB 0 4 1998
U.W.- MADISON

WEB EDITIONS FROM



ACS PUBLICATIONS
Essential Resources for the Chemical Sciences
http://pubs.acs.org

Proven, acknowledged excellence in science publishing combines with the convenience and advantages of the Web.

- ***New features and functions available only with Web Editions***
  ACS Web Editions deliver the full content of the journal right to the desktop on the cover date of publication. Users can search in titles, abstracts, or full text, or browse the author index or table of contents. Web Editions can be viewed and printed in two formats, as text (HTML) or as complete page images (PDF). Users can also link to nationally recognized databases, and have faster access to information through Articles ASAP (As Soon As Publishable), where articles are posted on the Web as soon as they have completed the ACS Publications quality control process. Enhanced graphics and interactive video features allow users to interpret visual materials as never before.
- ***A benefit for individual and institutional subscribers — pricing options and flexible coverage***
  Web Editions can be purchased separately or with print subscriptions at a discounted combination rate. Institutions can purchase just the amount of distribution required for a Web Edition, for the library, lab, department, or other setting. Site licenses are also available.

AMERICAN CHEMICAL SOCIETY
MEMBER & SUBSCRIBER SERVICES
P.O. BOX 3337, COLUMBUS, OH 43210 USA
1-800-333-9511 ■ 614-447-3776
Fax: 614-447-3671
E-mail: service@acs.org

JOURNAL OF AGRICULTURAL AND FOOD CHEMISTRY

Journal of Agricultural and Food Chemistry (ISSN 0021-8561) is published monthly by the American Chemical Society at 1155 16th St., N.W., Washington, DC 20036. Periodicals postage paid at Washington, DC, and additional mailing offices. POSTMASTER: Send address changes to *Journal of Agricultural and Food Chemistry*, Member & Subscriber Services, P.O. Box 3337, Columbus, OH 43210.

American Chemical Society
1155 16th St., N.W.
Washington, DC 20036
(202) 872-4600
TDD (202) 452-2120
Fax (202) 872-8264

Journal Publishing Operations
American Chemical Society
2540 Olentangy River Road
P.O. Box 3330, Columbus, OH 43210
(614) 447-3665
Fax (614) 447-3745; Telex 6842086
E-mail acsproof@acs.org

Member & Subscriber Services
American Chemical Society
P.O. Box 3337, Columbus, OH 43210
(614) 447-3776; (800) 333-9511
Fax (614) 447-3671
E-mail service@acs.org

Advertising Office
Centcom, Ltd.
676 East Swedesford Road
Suite 202, Wayne, PA 19087-1612
(610) 964-8061

Publications Division
Robert D. Bovenschulte, Director

Journal Publishing Operations
Mary E. Scanlan, Director

Editorial Office
Anne C. O'Melia, Manager; Kathleen E. Duffy, Journals Editing Manager; Eileen M. Richardson, Associate Editor

Circulation Office
Paula Commodore, Manager

Canadian GST Reg. No. 127571347
Printed in the USA

Copyright Permission: Reprographic copying beyond that permitted by Section 107 or 108 of the U.S. Copyright Act is allowed for *internal use only*, provided that $15.00 per article is paid to the Copyright Clearance Center, 222 Rosewood Drive, Danvers, MA 01923. Republication or reproduction for sale of articles or abstracts in this journal is permitted only under license from the ACS. Direct these and other permission requests to the Copyright Office, ACS, Washington, DC. Tel: (202) 872-4367. Fax: (202) 872-6060. E-mail: copyright@acs.org.

Subscription Donations: Members may donate/share their personal subscriptions to/with libraries but only after 5 years from publication.

Editorial Information

Instructions for Authors and Copyright Status Form: See the first issue of each volume or visit the Publications Division Web site (http://pubs.acs.org). Please conform to these instructions when submitting manuscripts.

Manuscript Submission: Send manuscripts to the Editor, Irvin E. Liener, Department of Biochemistry, College of Biological Sciences, University of Minnesota, 1479 Gortner Ave., St. Paul, MN 55108.

Accepted Papers and Proofs: Direct correspondence to Journal Publishing Operations, Columbus, OH. Tel: (614) 447-3665. Fax: (614) 447-3745. E-mail: acsproof@acs.org.

Page Charges: A $40.00 per page fee may be paid for papers published in this journal. Payment does not affect acceptance or scheduling of papers.

Journal Policies: The American Chemical Society and its Editors assume no responsibility for the statements and opinions advanced by contributors. Registered names and trademarks, etc., used in this publication, even without specific indication thereof, are not to be considered unprotected by law.

Document Number: At the end of each document is a 9-character code that serves as a link between the printed and electronic (CJACS Plus Images) products and facilitates retrieval of the document in electronic form.

Publisher Item Identifier (PII): A PII has been adopted by the ACS and several other publishers to provide unique identification of individual published documents. The PII appears at the bottom of the first page. Include the PII in all document delivery requests for copies of the document. See the Publications Division Web site (http://pubs.acs.org) for more details about the PII.

1998 Subscription and Ordering Information

		USA	Canada, Mexico	Europe*	All Other Countries*
Printed Edition	Members	$ 40.00	$ 65.00	$ 95.00	$114.00
	Student Members	$ 30.00	$ 55.00	$ 85.00	$104.00
	Nonmembers	$170.00	$195.00	$225.00	$244.00
	Institutional	$575.00	$600.00	$630.00	$649.00
Microfiche	Members	$ 40.00	$ 52.00	$ 52.00	$ 52.00
	Institutional	$575.00	$587.00	$587.00	$587.00
Supporting Information on Microfiche		$ 25.00	$ 31.00	$ 31.00	$ 31.00

* Air service included.

Web Edition: This journal is also available to subscribers via the Internet. Contact Member & Subscriber Services [Tel: (614) 447-3776 or (800) 333-9511. Fax: (614) 447-3671. E-mail: service@acs.org] or visit the Publications Division Web site (http://pubs.acs.org) for additional details.

New and Renewal Subscriptions: Send with payment to American Chemical Society, Department L-0011, Columbus, OH 43268-0011. Institutional subscribers in Japan must enter subscription orders with Maruzen Company Ltd., 3-10 Nihonbashi 2-chome, Chuo-ku, Tokyo 103, Japan. Tel: (03) 272-7211.

Change of Address: Notify Member & Subscriber Services, ACS, Columbus, OH. Tel: (614) 447-3776 or (800) 333-9511. Fax: (614) 447-3671. E-mail: service@acs.org. Include both old and new addresses and a mailing label from a recent issue.

Microfilm, Microfiche, Back Issue, and Printed Edition Single Issue Orders: Send requests to Publications Support Services, ACS, Washington, DC. Tel: (202) 872-4376. Fax: (202) 872-6325. Printed edition not available prior to 1994.

Bulk Reprint Orders: For quotes and information, contact Publications Support Services, ACS, Washington, DC. Tel: (202) 872-4376. Fax: (202) 872-6325.

Claims for Issues Not Received: Claims will be honored only if submitted within 90 days of the issue date (subscribers in North America) or within 180 days of the issue date (all other subscribers). Send requests to Member & Subscriber Services, ACS, Columbus, OH. Tel: (614) 447-3776 or (800) 333-9511. Fax: (614) 447-3671. E-mail: service@acs.org.

Supporting Information (SI): SI is available in print, electronic-only, and combination print/electronic formats. See table of contents for format availability. SI in print is available as photocopy, as 24× microfiche, or via the Internet (free to subscribers). SI except structure factors is also included in the microfilm edition. For photocopy or microfiche orders, see SI paragraph at end of journal article for number of pages. For further information, contact Publications Support Services, ACS, Washington, DC. Tel: (202) 872-4376. Fax: (202) 872-6325. SI in electronic-only format is available via the Internet (http://pubs.acs.org). For additional information on electronic access, send e-mail to si-help@acs.org or phone (202) 872-6333.

© Copyright 1998 American Chemical Society

JAFCAU 46(1) 1−360 (1998)
ISSN 0021-8561

Registered in U.S. Patent and Trademark Office
Copyright 1998 by the American Chemical Society

JANUARY 1998

JOURNAL OF AGRICULTURAL AND FOOD CHEMISTRY®

EDITORIAL

1 Editorial

RAPID COMMUNICATIONS

2 Distribution of Aflatoxin in Pistachios. 5. Sampling and Testing U.S. Pistachios for Aflatoxin
Thomas F. Schatzki

5 Enzymatic Activity in Root Exudates of Dahurian Wild Rye (*Elymus dauricus*) That Degrades 2-Chlorobenzoic Acid
S. D. Siciliano, H. Goldie, and J. J. Germida*

FOOD CHEMISTRY/BIOCHEMISTRY

8 Growth-Inhibiting Effects of *Cinnamomum cassia* Bark-Derived Materials on Human Intestinal Bacteria
Hoi-Seon Lee and Young-Joon Ahn*

13 Dearomatization of Antioxidant Rosemary Extracts by Treatment with Supercritical Carbon Dioxide
Sara López-Sebastián, Elena Ramos, Elena Ibáñez, Jose M. Bueno, Lorenzo Ballester, Javier Tabera, and Guillermo Reglero*

20 Differences in Chlorophyll Loss at 13 °C for Two Broccoli (*Brassica oleracea* L.) Cultivars Associated with Antioxidant Enzyme Activities
P. M. A. Toivonen* and Mark Sweeney

25 Antioxidant Activity of Berry and Fruit Wines and Liquors
I. Marina Heinonen,* Pekka J. Lehtonen, and Anu I. Hopia

32 Phenolic Compounds in Different Olive Varieties
Marco Esti,* Luciano Cinquanta, and Ennio La Notte

36 Adsorption Behavior of Egg Yolk Low-Density Lipoproteins in Oil-in-Water Emulsions
Yoshinori Mine

42 Relation between Fatty Acid Content and Its Evolution during Fermentation and Utilization of Free Amino Acids in Vacuum-Filtered Viura Must
Belén Ayestarán, Julián Garrido, and Carmen Ancín*

49 Antioxidative Properties of Histidine-Containing Peptides Designed from Peptide Fragments Found in the Digests of a Soybean Protein
Hua-Ming Chen, Koji Muramoto,* Fumio Yamauchi, Kenshiro Fujimoto, and Kiyoshi Nokihara

Case No.: IPR2023-00206
U.S. Patent No. 9,943,096

54 Multidimensional Capillary GC–GC for the Analysis of Real Complex Samples. 3. Enantiomeric Distribution of Monoterpene Hydrocarbons and Monoterpene Alcohols of Mandarin Oils
Luigi Mondello,* Maurizio Catalfamo, Anna Rita Proteggente, Ivana Bonaccorsi, and Giovanni Dugo

62 Mobility of "Unfreezable" and "Freezable" Water in Waxy Corn Starch by ^{2}H and ^{1}H NMR
S. Li, L. C. Dickinson, and P. Chinachoti*

72 Influence of Protein Interfacial Composition on Salt Stability of Mixed Casein Emulsions
Herley Casanova and Eric Dickinson*

77 Isolation and Identification of β-Casein A^1-4P and β-Casein A^2-4P in Commercial Caseinates
Loren S. Ward and Eric D. Bastian*

84 Destabilization of Oil-in-Water Emulsions Formed Using Highly Hydrolyzed Whey Proteins
Samson O. Agboola, Harjinder Singh,* Peter A. Munro, Douglas G. Dalgleish, and Anne M. Singh

91 Viscoelastic Properties of Protein-Stabilized Emulsions: Effect of Protein–Surfactant Interactions
Jianshe Chen and Eric Dickinson*

98 Effect of Crystallinity on the Glass Transition Temperature of Starch
Akinori Mizuno,* Masata Mitsuiki, and Masao Motoki

104 New Melanoidin-like Maillard Polymers from 2-Deoxypentoses
Roland Tressl,* Georg T. Wondrak, Ralph-Peter Krüger, and Dieter Rewicki*

111 NMR Analytical Approach To Clarify the Antioxidative Molecular Mechanism of Catechins Using 1,1-Diphenyl-2-picrylhydrazyl
Yusuke Sawai* and Kanzo Sakata

115 Copper(II) Complexes of Opiate-like Food Peptides
Elzbieta Lodyga-Chruscinska, Giovanni Micera,* Ewa Szajdzinska-Piêtek, and Daniele Sanna

COMPOSITION OF FOODS/FEEDS

119 Flesh Lipid and Carotenoid Composition of Scottish Farmed Atlantic Salmon (*Salmo salar*)
J. Gordon Bell,* John McEvoy, John L. Webster, Fiona McGhee, Rose Mary Millar, and John R. Sargent

128 Effects of Ripening, Cultivar Differences, and Processing on the Carotenoid Composition of Mango
Adriana Z. Mercadante and Delia B. Rodriguez-Amaya*

131 Alfalfa Saponins and Their Implication in Animal Nutrition
Sucharita Sen, Harinder P. S. Makkar, and Klaus Becker*

ANALYTICAL METHODS

141 Enzymatic Determination of Citric Acid in Honey by Using Polyvinylpolypyrrolidone Clarification
Inés Mato, José F. Huidobro,* Verónica Cendón, Soledad Muniategui, Miguel A. Fernández-Muiño, and M. Teresa Sancho

145 Application of an Electronic Aroma Sensing System to Cork Stopper Quality Control
S. Rocha, I. Delgadillo,* A. J. Ferrer Correia, A. Barros, and P. Wells

152 Development of an Enzyme-Linked Immunosorbent Assay for the Detection of Metosulam
J. S. Parnell and J. Christopher Hall*

157 Colorimetric Assay of Alditols in Complex Biological Samples
Juan Sánchez

161 Analytical Method for the Determination of Atrazine and Its Dealkylated Chlorotriazine Metabolites in Urine by Gas Chromatography/Mass Selective Detection
Liakat H. Bodalbhai, Robert A. Yokley,* and Max W. Cheung

168 Determination of SO_2 in Wines Using a Flow Injection Analysis System with Potentiometric Detection
Alberto N. Araújo, Cristina M. C. M. Couto, José L. F. C. Lima, and Maria C. B. S. M. Montenegro*

Case No.: IPR2023-00206
U.S. Patent No. 9,943,096

173 Sample Preparation for Arsenic, Copper, Iron, and Lead Determination in Sugar
Jale Leblebici* and Mürvet Volkan

NUTRITION/TOXICOLOGY

178 Metabolic Study of Isoproturon in Goats Following a Single Oral Administration: Toxicokinetics and Recovery
Sanis Juliet, Tapan K. Mandal, Biswanath Mal, Ashim Chowdhury, Anjan Bhattacharyya, and Animesh K. Chakraborty*

184 Dose-Response Effect of Dietary Docosahexaenoic Acid on Fatty Acid Profiles of Serum and Tissue Lipids in Rats
Morio Saito,* Megumi Ueno, Kazuhiro Kubo, and Michio Yamaguchi

194 Unidentified Factors in Jojoba Meal Prevent Oviduct Development in Broiler Breeder Females
Sabien Vermaut,* Okanlawon Onagbesan, Veerle Bruggeman, Guido Verhoeven, Luc Berghman, Gerda Flo, Marnix Cokelaere, and Eddy Decuypere

202 Characterization of *N*-(Nitrosomethyl)urea in Nitrosated Fermented Fish Products
Dajun Deng,* Tong Li, Hong Ma, Ruming Wang, Liankun Gu, and Jing Zhou

206 Effects of Microwave Heating on the Loss of Vitamin B_{12} in Foods
Fumio Watanabe,* Katsuo Abe, Tomoyuki Fujita, Mashahiro Goto, Miki Hiemori, and Yoshihisa Nakano

211 Organic Acids in Selected Foods Inhibit Intestinal Brush Border Pteroylpolyglutamate Hydrolase in Vitro: Potential Mechanism Affecting the Bioavailability of Dietary Polyglutamyl Folate
Ming-Min Wei and Jesse F. Gregory, III*

FLAVORS AND AROMAS

220 Thermal Degradation of Allyl Isothiocyanate in Aqueous Solution
Chung-Wen Chen and Chi-Tang Ho*

224 Mechanistic Studies on Thiazolidine Formation in Aldehyde/Cysteamine Model Systems
Tzou-Chi Huang,* Lee-Zen Huang, and Chi-Tang Ho

228 Evaluation of Solid-Phase Microextraction for Analysis of Volatile Metabolites Produced by Staphylococci
L. Vergnais, F. Masson, M. C. Montel, J. L. Berdagué, and R. Talon*

235 Quantitative Model Studies on the Effectiveness of Different Precursor Systems in the Formation of the Intense Food Odorants 2-Furfurylthiol and 2-Methyl-3-furanthiol
T. Hofmann and P. Schieberle*

242 Determination of Key Flavor Components in Methylene Chloride Extracts from Processed Grapefruit Juice
P. Jella, R. Rouseff,* K. Goodner, and W. Widmer

248 Natural Abundance ^{2}H Nuclear Magnetic Resonance Study of the Origin of Raspberry Ketone
Giovanni Fronza, Claudio Fuganti,* Claude Guillou, Fabiano Reniero, and Daniel Joulain

CHEMICAL CHANGES DURING PROCESSING/STORAGE

255 Carbon Monoxide, Nitric Oxide, and Nitrogen Dioxide Levels in Gas Ovens Related to Surface Pinking of Cooked Beef and Turkey
D. P. Cornforth,* J. K. Rabovitser, S. Ahuja, J. C. Wagner, R. Hanson, B. Cummings, and Y. Chudnovsky

262 Changes in the Dietary Fiber (Resistant Starch and Nonstarch Polysaccharides) Content of Cooked Flours Prepared from Three Chinese Indigenous Legume Seeds
Peter C.-K. Cheung* and Chi-Fai Chau

266 High-Pressure and Temperature Effects on Enzyme Inactivation in Tomato Puree
Almudena Hernández and M. Pilar Cano*

271 Effects of Drying Pretreatments on the Cell Wall Composition of Grape Tissues
A. Femenia, E. S. Sánchez, S. Simal, and C. Rosselló*

Motif Exhibit 1049, Page 6 of 14

Case No.: IPR2023-00206
U.S. Patent No. 9,943,096

277 Changes in the Amino Acid Composition of Dehydrated Orange Juice during Accelerated Nonenzymatic Browning
M. Dolores del Castillo, Nieves Corzo,* M. Carmen Polo, Encarnación Pueyo, and Agustín Olano

281 Inactivation Kinetics Study of the Kunitz Soybean Trypsin Inhibitor and the Bowman–Birk Inhibitor
Rob van den Hout,* Marieke Pouw, Harry Gruppen, and Klaas van't Riet

CHEMISTRY OF CROP AND ANIMAL PROTECTION

286 Antifeedant *Delphinium* Diterpenoid Alkaloids. Structure–Activity Relationships
Azucena González-Coloma,* Ana Guadaño, Carmen Gutiérrez, Raimundo Cabrera, Eduardo de la Peña, Gabriel de la Fuente, and Matías Reina

291 Field Persistence Studies on Pendimethalin Residues in Onions and Soil after Herbicide Postemergence Application in Onion Cultivation
Nicholas G. Tsiropoulos* and George E. Miliadis

296 Effect of Long-Chained Esters on the Insecticidal Properties of L-Canavanine
Gerald A. Rosenthal,* Palesa Nkomo, and Douglas L. Dahlman

300 Biotransformation of (+)- and (−)-Limonene by the Larvae of Common Cutworm (*Spodoptera litura*)
Mitsuo Miyazawa,* Takashi Wada, and Hiromu Kameoka

ENVIRONMENTAL CHEMISTRY

304 Accumulation, Metabolism, and Depuration of Organotin Compounds in the Marine Mussels *Mytilus graynus* and *Mytilus edulis* under Natural Conditions
Takashi Suzuki,* Isao Yamamoto, Hisashi Yamada, Nahoko Kaniwa, Kazunari Kondo, and Mitsunori Murayama

314 Formation of *N*-Nitrosoterbuthylazine and *N*-Nitrosoterbutryn in a Model System of Soil Water
Marco Trevisan, Elena Graviani, Attilio A. M. Del Re, Anna Arnoldi,* Angela Bassoli, Dario Cova, and Andrea Rossignoli

318 Formation of Sterilized Edible Films Based on Caseinates: Effects of Calcium and Plasticizers
Emna Mezgheni, Giuseppe D'Aprano, and Monique Lacroix*

325 Investigation of the Heterogeneously Catalyzed Hydrolysis of Organophosphorus Pesticides
Andrea Dannenberg and Simo O. Pehkonen*

335 Soil Adsorption Coefficients of *s*-Triazines Estimated with a New Gradient HPLC Method
A. Kaune,* R. Brüggemann, M. Sharma, and A. Kettrup

BIOTECHNOLOGY

344 Molecular Cloning of a cDNA Encoding Copper/Zinc Superoxide Dismutase from Papaya Fruit and Overexpression in *Escherichia coli*
Ming-Tse Lin, Ti-Jung Kuo, and Chi-Tsai Lin*

349 Purification and Characterization of a Tripeptidase from *Lactobacillus sake*
Yolanda Sanz, Francis Mulholland, and Fidel Toldrá*

354 Heterologous Expression of a *Candida molischiana* Anthocyanin-β-glucosidase in a Wine Yeast Strain
Paloma Sánchez-Torres, Luis González-Candelas, and Daniel Ramón*

There is no Supporting Information for this issue.

* In papers with more than one author, the asterisk indicates the name of the author to whom inquiries about the paper should be addressed.

Case No.: IPR2023-00206
U.S. Patent No. 9,943,096

This material may be protected by Copyright law (Title 17 U.S. Code)

Quantitative Model Studies on the Effectiveness of Different Precursor Systems in the Formation of the Intense Food Odorants 2-Furfurylthiol and 2-Methyl-3-furanthiol

T. Hofmann and P. Schieberle*

Deutsche Forschungsanstalt für Lebensmittelchemie, Lichtenbergstrasse 4, D-85748 Garching, Germany

The yields of the two intense food odorants 2-furfurylthiol (FFT) and 2-methyl-3-furanthiol (MFT) obtained by heating mixtures of possible precursors in model systems varying in temperature, pH value, or water content were determined by using stable isotope dilution assays. Although pentoses generated much higher amounts of FFT and MFT than hexoses when heated in the presence of cysteine, glucose and rhamnose also gave significant yields. Studies on several intermediates indicated the highest yields for MFT (1.4 mol %) when hydroxyacetaldehyde and mercapto-2-propanone were reacted for 6 min at 180 °C in the absence of water. Both intermediates also generated significant amounts of FFT (0.05 mol %). However, the system furan-2-aldehyde/H_2S showed a 10 times higher efficiency in generating FFT. Thiamin and norfuraneol/cysteine were less effective precursors of MFT. The results imply that different formation pathways may run in parallel during food processing and may account for the different amounts of the two odorants present in the respective food.

Keywords: *2-Furfurylthiol; 2-methyl-3-furanthiol; 4-hydroxy-5-methyl-3(2H)-furanone; furan-2-aldehyde; flavor precursors; mercapto-2-propanone; 3-deoxyribosulose*

INTRODUCTION

2-Furfurylthiol (FFT) and 2-methyl-3-furanthiol (MFT) exhibit intense roasted, coffee-like, and meatlike odor notes at the extremely low odor threshold of 0.0025 ng/L in air (Hofmann and Schieberle, 1995). FFT was reported for the first time as a food constituent in a study on roasted coffee (Reichstein and Staudinger, 1926) and, later, among the volatiles of cooked meat (Garbusov et al., 1976). MFT was identified for the first time among the food volatiles in an investigation on the constituents of heated canned tuna fish (Withycombe and Mussinan, 1988).

Using aroma extract dilution analysis (AEDA), both FFT and MFT have recently been established as key aroma contributors to several thermally processed foods, such as cooked beef (Gasser and Grosch, 1988) and roasted coffee (Blank et al., 1992). Furthermore, FFT was elucidated as an important odorant in freshly popped corn (Schieberle, 1991), roasted white sesame seeds (Schieberle, 1996), and, very recently, in heated commercial as well as in self-prepared baker's yeast extracts (Münch et al., 1997).

A great number of investigations have been performed in the past 30 years to clarify the precursors of FFT and MFT in foods. Model studies have elucidated pentoses and cysteine as precursors of either FFT [e.g. Ledl and Severin (1973), Mottram (1987), Whitfield et al. (1988), Farmer and Mottram (1990a), Zeiler-Hilgart (1994), and Mottram and Whitfield (1995)] or MFT [e.g. Mottram (1987), Whitfield et al. (1988), Martin (1988), Farmer and Mottram (1990a,b), Meynier and Mottram (1995), and Mottram and Whitfield (1994)]. In addition, FFT has been reported to be formed by heating glucose in the presence of either hydrogen sulfide and ammonia (Shibamoto and Russel, 1976, 1977) or cysteine (Yeo and Shibamoto, 1991).

Very recently, by application of the AEDA, FFT and MFT have also been established as key contributors to the overall odors of processed flavors manufactured by heating ribose (Hofmann and Schieberle, 1995) and glucose or rhamnose (Hofmann and Schieberle, 1997) in the presence of cysteine.

Shibamoto and Russell (1977) and Silwar and Tressl (1989) reported the formation of FFT from furan-2-aldehyde, a major dehydration product of pentoses, in the presence of sulfur sources, such as cysteine or hydrogen sulfide. Quantitative studies performed by using stable isotope dilution assays (Münch et al., 1997) recently confirmed that FFT is indeed formed in high concentrations during thermal treatment of furan-2-aldehyde in the presence of cysteine. Further data (Münch et al., 1997) suggested the importance of retro-Aldol reactions in FFT formation.

Whitfield et al. (1993) and Mottram and Whitfield (1994) reported that MFT is formed by reacting 4-hydroxy-5-methyl-3(2*H*)-furanone [norfuraneol (NF)], another major degradation product of pentoses, with cysteine or hydrogen sulfide, respectively. However, van den Ouweland and Peer (1975) were unable to identify MFT in such reaction systems. Quantitative studies (Zeiler-Hilgart, 1994) confirmed these results showing that the norfuraneol/cysteine system is indeed not very effective in generating MFT.

The thermal degradation of thiamin has been discussed as an alternative reaction pathway in MFT formation during food processing (van der Linde et al., 1979; Hartman et al., 1984; Reineccius and Liardon,

* Author to whom correspondence should be addressed (e-mail Lebchem.Schieberle@lrz.tu-muenchen.de; fax +49-89-289 141 83; telephone +49-89-289 141 70).

S0021-8561(97)00598-0 CCC: $15.00 © 1998 American Chemical Society
Published on Web 01/19/1998

Case No.: IPR2023-00206
U.S. Patent No. 9,943,096

1985; Gasser, 1990; Güntert et al., 1990; Werkhoff et al., 1990; Hincelin et al., 1992; Zeiler-Hilgart, 1994).

To reveal the effectiveness of the precursor systems discussed above in generating both odorants, exact quantitative data are a prerequisite. However, the instability of FFT, and especially that of MFT, may cause severe losses during enrichment procedures needed to quantify these trace odorants (Schieberle and Hofmann, 1997). Stable isotope dilution analysis (SIDA) is the best method to overcome these problems [cf. Schieberle (1995)]. The following quantitative model studies were, therefore, undertaken to gain more detailed insight into the efficiency of possible precursors and, also, in reaction mechanisms governing the formation of FFT as well as that of MFT during food processing.

EXPERIMENTAL PROCEDURES

Chemicals. 2-Furfurylthiol, 2-methyl-3-furanthiol, 2-oxopropanal (30% solution in water), furan-2-aldehyde, D-ribose, D-xylose, D-glucose, L-rhamnose, D-glucose-6-phosphate, maltose, L-cysteine, thiamin, and sodium hydrosulfide hydrate (NaHS) were obtained from Aldrich (Steinheim, Germany), and hydroxyacetaldehyde (glycolaldehyde) was from Roth (Germany).

Syntheses. The following compounds were synthesized following closely the procedures given in parentheses: mercapto-2-propanone (Hofmann and Schieberle, 1995); [$^{13}C_4$]-mercapto-2-butanone (Hofmann and Schieberle, unpublished data); [$^2H_2$]furfurylthiol (Sen and Grosch, 1991); [$^2H_3$]-2-methyl-3-furanthiol (Sen and Grosch, 1991); [$^{13}C_2$]-4-hydroxy-2,5-dimethyl-3(2*H*)-furanone (Sen et al., 1991).

Preparation of 3-Deoxyribosulose. The preparation was performed following the method described for 3-deoxyxylulose (Madson and Feather, 1981) with some modifications. A stirred solution of ribose (55 mmol) and *p*-toluidine (55 mmol) in ethanol/acetic acid (236 mL; 22:1 by volume) was refluxed for 30 min. After addition of benzoyl hydrazine (120 mmol), refluxing was continued for another 420 min. The reaction mixture was then cooled to room temperature and kept at −30 °C overnight. Separated 3-deoxyribosulose bis(benzylhydrazone) was isolated by filtration over a Büchner funnel and washed with ice-cooled ethanol (total volume = 100 mL; yield = 24 mmol, 43.6% of theoretical yield). The hydrazone (24 mmol) and benzaldehyde (16 mL) were then refluxed in a mixture of ethanol (300 mL), water (500 mL), and acetic acid (22 mL). After 2 h, the mixture was cooled to room temperature and the ethanol was evaporated in vacuo, while water (500 mL) was added within 30 min. The mixture was then cooled in an ice bath and the separated benzaldehyde−benzoylhydrazone was removed by filtration over a Büchner funnel. Mixed-bed H^+/OH^--ion-exchange resin (Serdolyt MB3; 60 g; Serva, Heidelberg, Germany) was added to the solution, and after being stirred for 15 min in the dark, the suspension was filtered. The filtrate was concentrated in vacuo to ~100 mL, then washed with diethyl ether (6 × 50 mL), and finally concentrated in vacuo to ~5 mL. The residue was taken up in a mixture of ethanol (50 mL) and water (5 mL), mixed-bed H^+/OH^--ion-exchange resin (Serdolyt MB3; 10 g) was added, and the suspension was stirred for a further 15 min in the dark. After filtration, the solution was concentrated in vacuo to ~5 mL. Freeze-drying yielded 3-deoxyribosulose as an amorphous powder (10 mmol; 42%).

Preparation of NF. Xylose (400 mmol), *n*-propylamine (200 mmol), and KH_2PO_4 (400 mmol) were dissolved in water (400 mL), and the pH was adjusted to 7.0 with aqueous NaOH (15%). After refluxing for 30 min, the mixture was cooled to room temperature, water (150 mL) was added, and the pH was adjusted to 4.0 with hydrochloric acid (1 mol/L). After filtration, the aqueous solution was extracted with ethyl acetate (8 × 200 mL), and the organic layer was dried over Na_2SO_4 and finally concentrated in vacuo to ~5 mL. The target compound was isolated from the mixture by column chromatography (30 × 4 cm) on silica gel (G60; 7% water). After flushing with ethyl acetate/hexane (3:1, by volume), an aliquot of the reaction mixture was applied onto the top of the column. Elution was performed with ethyl acetate/hexane (3:1, by volume), affording the target compound within 400 mL of the eluate. After recrystallization of the crude product from methanol (2 mL), NF (12 mmol; 3%) was obtained as a white solid: MS(EI), *m/z* (%) 43 (100), 114 (52), 55 (9), 58 (4), 71 (3); ^1H NMR (360 MHz in $CDCl_3$) (ppm; multiplicity; hydrogen at relevant carbon atom) δ 2.27 (s, 3H, CH_3), δ 4.52 (s, 2H, CH_2); ^{13}C NMR (360 MHz in [$^2H_4$]CH_3OH) (ppm; multiplicity; relevant carbon atom) δ 11.7 (CH_3), δ 72.6 (CH_2), δ 134.3 (C), δ 175.3 (COH), δ 196.6 (CO).

Model Experiments. In a first series of experiments, the reactants (cf. amounts detailed in the tables) were dissolved in phosphate buffer (0.5 mol/L) at pH 3.0, 5.0, or 7.0 and thermally treated for 20 min at 145 °C in a laboratory autoclave (200 mL; Type II; Roth, Germany). In a second series of experiments, the reactants were intimately mixed with silica gel (3.0 g containing 300 μL of the respective phosphate buffer; 2 mol/L) and reacted for 6 min at 180 °C.

Quantitation of FFT and MFT by SIDA. After cooling of the reaction mixtures to 20 °C, the internal standards [$^2H_2$]-2-furfurylthiol and [$^2H_3$]-2-methyl-3-furanthiol (10 μg each dissolved in 0.5 mL of pentane) were added to the aqueous reaction mixtures and, after equilibration for 10 min, the analyte and the labeled standard were isolated by extraction with diethyl ether (three times, total volume = 60 mL). After drying over Na_2SO_4, quantitation of FFT and MFT was performed by mass chromatography using the fragment ions and response factors as described recently (Guth et al., 1995).

Quantitation of Furan-2-aldehyde. After cooling, certain amounts of 5-methylfurfural were added to the reaction mixtures as the internal standard. After extraction of the analyte and the standard with diethyl ether (four times, total volume = 50 mL) followed by drying over Na_2SO_4, quantitation of furfural was performed by mass chromatography using the ion trap detector. An MS response factor of 0.95 was calculated from mixtures containing known amounts of furan-2-aldehyde and 5-methylfurfural by using the molecular ions obtained in the chemical ionization mode.

Quantitation of Mercapto-2-propanone and NF. After cooling to room temperature, certain amounts of the labeled homologues [$^{13}C_4$]mercapto-2-butanone and [$^{13}C_2$]-4-hydroxy-2,5-dimethyl-3(2*H*)-furanone were added to the reaction mixtures. After extraction with diethyl ether (four times, total volume = 50 mL) and drying over Na_2SO_4, quantitation was performed by mass chromatography using a gas chromatograph coupled to an ion trap detector. MS response factors for mercapto-2-propanone (1.08) and NF (0.98) were calculated from mixtures containing known amounts of the analyte and the labeled standard.

High-Resolution Gas Chromatography/Mass Spectrometry (HRGC/MS). HRGC analyses were performed by means of a gas chromatograph Type 5300 (Fisons, Mainz, Germany) and by using an FFAP capillary column (30 m × 0.32 mm; 0.5 μm; J&W Scientific, Fisons Instruments, Mainz, Germany). The samples (1 μL) were applied by on-column injection at 40 °C. After 2 min, the oven temperature was raised at 40 °C/min to 60 °C, held for 2 min isothermally, then raised at 6 °C/min to 230 °C, and held for 5 min. Quantitations were performed using an Finnigan 800 ion trap detector (Bremen, Germany), running in the chemical ionization mode with methanol as the reagent gas.

NMR Spectroscopy. ^1H and ^{13}C NMR spectra were recorded in [$^2H_4$]methanol (MSD isotopes, Montreal, Canada) by means of an AM 360 (Bruker, Karlsruhe, Germany). The ^1H signals were assigned according to tetramethylsilane (TMS) as the internal standard.

RESULTS AND DISCUSSION

Carbohydrates and Cysteine as Precursors of FFT and MFT. To investigate their effectiveness in

Case No.: IPR2023-00206
U.S. Patent No. 9,943,096

Table 1. Amounts of FFT and MFT Generated from Cysteine and Various Carbohydrates[a]

carbohydrate	amount[b] (μg) FFT	MFT
ribose	12.1	19.8
xylose	9.6	14.3
fructose	3.2	2.5
glucose	2.8	1.9
glucose-6-phosphate	0.9	0.6
rhamnose	0.8	0.8
maltose	0.6	0.3

[a] A solution (100 mL) of cysteine (3.3 mmol) and the corresponding carbohydrate (10.0 mmol) was reacted at pH 5.0 in a laboratory autoclave. [b] Data are mean values of triplicates and differed by not more than 10%.

Table 2. Influence of pH on the Amounts of FFT and MFT Generated from Cysteine and Carbohydrates[a]

carbohydrate	FFT[b] (μg) at pH 3.0	5.0	7.0	5.0[c]	MFT[b] (μg) at pH 3.0	5.0	7.0	5.0[c]
ribose	22.9	12.1	1.2	97.2	55.3	19.8	2.5	25.1
glucose	0.7	2.8	0.6	1.4	0.3	1.9	0.4	4.2
rhamnose	0.2	0.8	0.1	0.4	0.1	0.8	0.1	3.1

[a] A solution (100 mL) of cysteine (3.3 mmol) and the corresponding carbohydrate (10.0 mmol) was reacted in a laboratory autoclave. [b] See footnote *b* in Table 1. [c] A mixture of cysteine (3.3 mmol) and the corresponding carbohydrate (10.0 mmol) was dry-heated for 6 min at 180 °C.

generating FFT and MFT, aqueous solutions of several monosaccharides, as well as glucose-6-phosphate and the disaccharide maltose, were reacted in the presence of cysteine. The amounts of both odorants were determined by SIDA. The results established the pentoses ribose and xylose as the most effective precursors in the generation of both odorants (Table 1). Fructose and glucose were more efficient than glucose-6-phosphate, rhamnose, or maltose, which yielded only small amounts of FFT and MFT when heated in the presence of cysteine.

To gain a more detailed insight into factors governing the formation of FFT and MFT from pentoses and hexoses, the influence of the pH on odorant formation from ribose, glucose, and rhamnose was studied. The formation of both odorants from pentoses and hexoses was influenced by pH in a different way (Table 2). From ribose, the liberation of FFT and MFT increased significantly with lowering the pH from 7.0 to 3.0. Although comparatively lower amounts were formed, the generation of both odorants in the presence of glucose or rhamnose went through a maximum at pH 5.0.

To examine the influence of water and the reaction temperature, the three carbohydrates were heated under dry-heating conditions in the presence of cysteine. A small amount of concentrated buffer was added to maintain pH 5.0. Under these conditions, ribose generated much higher concentrations of FFT than under aqueous conditions, whereas MFT was only slightly increased. In contrast, FFT especially was lower when hexoses were reacted under dry-heating conditions.

In summary, the highest amounts of FFT were obtained by dry-heating the ribose/cysteine mixture (Table 2), whereas MFT was preferentially formed from this model mixture under aqueous conditions at pH 3.0. Substitution of ribose by hexoses generally gave different results. Besides lower yields, the formation of FFT from both hexose/cysteine systems was highest under aqueous conditions at pH 5.0, whereas the MFT was formed from these mixtures preferentially under dry-heating conditions. The data suggested that the formation of FFT and MFT from pentoses and hexoses follows different reaction pathways and prompted us to look into several possible intermediates of FFT and MFT formation in more detail.

Table 3. Generation of FFT from 5-Carbon Precursors in the Presence of Hydrogen Sulfide[a]

precursor	FFT μg[b]	%
ribose	9.2	0.008
3-deoxyribosulose	78.6	0.08
furan-2-aldehyde	550.8	0.48

[a] An aqueous solution (50 mL) of hydrogen sulfide (1 mmol) and the precursor (1 mmol) was reacted at pH 5.0 in a laboratory autoclave. [b] See footnote *b* in Table 1.

Table 4. Generation of MFT from Ribose or NF in the Presence of Hydrogen Sulfide[a]

precursor (1 mmol)	MFT μg[b]	%
ribose	15.1	0.01
NF	211.2	0.19

[a] An aqueous solution (50 mL) of hydrogen sulfide (1 mmol) and the precursor was heated at pH 5.0 in a laboratory autoclave. [b] See footnote *b* in Table 1.

Intermediates in FFT and MFT Formation from Pentoses. Dehydration of pentoses may result in the formation of either the 1- or the 3-deoxypentosone. Their subsequent cyclization and elimination of water lead to the formation of either NF or furan-2-aldehyde as the major products. However, it is yet unknown whether both deoxyosones are formed in equal amounts.

The reaction of furan-2-aldehyde with hydrogen sulfide, which can be liberated from cysteine, was recently shown to generate the FFT very effectively (Münch et al., 1997). It might, therefore, be assumed that also the 3-deoxypentosone could act as a precursor of this flavor compound.

To compare their efficiencies as FFT precursors, furan-2-aldehyde and 3-deoxypentosone were each heated in the presence of hydrogen sulfide, and the amounts of FFT formed were compared with those formed from ribose. More FFT was liberated from the furan-2-aldehyde, compared to ribose and the 3-deoxyribosulose, by factors of 60 or 7, respectively (Table 3). A reaction mechanism explaining FFT formation via a reductive sulfhydrylation of furan-2-aldehyde with hydrogen sulfide has recently been proposed (Münch et al., 1997).

NF belongs to the major degradation product of pentoses and is assumed to be formed via the 1-deoxypentosone. When heated in the presence of cysteine, NF has been found to generate significant amounts of MFT (Whitfield et al., 1993). To compare its effectiveness with that of ribose, NF was reacted with hydrogen sulfide and the amount of MFT liberated was quantified. In agreement with literature data (Whitfield et al., 1993), the odorant was generated very efficiently from the furanone (Table 4). Compared to ribose, by a factor of ≈14 higher amounts of MFT were formed from NF. NF is generated in very high yields from ribose/cysteine (Table 5). This result therefore explains the preferential generation of MFT from the pentose via NF as the intermediate.

A hypothetical reaction sequence leading to MFT from NF and hydrogen sulfide as the intermediates is dis-

Case No.: IPR2023-00206
U.S. Patent No. 9,943,096



Figure 1. Hypothetical reaction sequence leading from NF and H_2S to MFT.

Table 5. Amounts (Micrograms)[a] of the Pairs Furan-2-aldehyde (FA)/FFT and NF/MFT Generated from Carbohydrates in the Presence of Cysteine[b]

carbohydrate	FA	FFT	NF	MFT
ribose	67.5	12.1	54530.0	19.8
glucose	6.9	2.8	13.5	1.9
rhamnose	6.2	0.8	19.1	0.8

[a] See footnote *b* in Table 1. [b] An aqueous solution (100 mL) of cysteine (3.3 mmol) and the corresponding carbohydrate (10.0 mmol) was reacted at pH 5.0 in a laboratory autoclave.

played in Figure 1. Incorporation of a thiol group at carbon 3 yields 4-mercapto-5-methyl-3(2*H*)-furanone in a first reaction step, which in turn might be reduced by a reductone or by a further molecule of hydrogen sulfide. Elimination of water from the intermediate 3-hydroxy-4-mercapto-5-methyl-1,2-dihydrofuran would give rise to MFT.

Intermediates in FFT and MFT Formation from Hexoses. Because MFT and FFT were found to be generated also from hexoses (cf. Table 2), it might be speculated that NF and furan-2-aldehyde are also the key intermediates in the formation of both odorants. However, to form these precursors from hexoses, a cleavage of the carbohydrate skeleton must occur. This is possible by α-dicarbonyl cleavage and vinylogous retro-Aldol cleavage, leading to several cleavage products (C-5/C-1, C-4/C-2, or C-3/C-3). Such short-chain carbonyl compounds, e.g., hydroxyacetaldehyde and 2-oxopropanal, have been reported to be formed from hexoses (Hayashi and Namiki, 1986; Nedvidek et al., 1992). A hypothesis on the formation of NF and furan-2-aldehyde starting from 2-oxopropanal and hydroxyacetaldehyde is shown in Figure 2. Depending on the side of the nucleophilic attack, an Aldol reaction of the short-chain compounds might result in either furan-2-aldehyde or NF. Nucleophilic attack of the methyl group of 2-oxopropanal at the hydroxyacetaldehyde results in the formation of 3-deoxypentosone (**Ib**; Figure 2), which upon 2,5-cyclization and dehydration gives rise to furan-2-aldehyde (FA; **IIb**). However, the reaction of the methylene group of hydroxyacetaldehyde with the aldehyde function of the 2-oxopropanal might result in 1-deoxypentosone (**Ia**; Figure 2), which upon 2,5-cyclization and water elimination yields NF (**IIa**).

To check this hypothesis, hydroxyacetaldehyde and



Figure 2. Formation of furan-2-aldehyde (FA) and NF by an Aldol reaction of 2-oxopropanal and hydroxyacetaldehyde.

2-oxopropanal were reacted in aqueous solution at different pH values and the amounts of FA and NF formed were quantified. Both compounds were formed

Case No.: IPR2023-00206
U.S. Patent No. 9,943,096

Table 6. Amounts of Furan-2-aldehyde (FA) and NF Generated from Hydroxyacetaldehyde and 2-Oxopropanal[a]

	FA		NF	
pH	μg[b]	%	μg[b]	%
3.0	101.2	0.11	153.1	0.13
5.0	364.5	0.40	885.1	0.78
7.0	1443.1	1.60	3610.5	3.18

[a] An aqueous solution (50 mL) of hydroxyacetaldehyde (1 mmol) and 2-oxopropanal (1 mmol) was reacted in a laboratory autoclave. [b] See footnote *b* in Table 1.

in significant yields, with neutral conditions (pH 7.0) favoring their formation (Table 6). It is interesting to note that in all reaction mixtures more NF than FA had been formed. Referring to the mechanism shown in Figure 2, these data indicate hydroxyacetaldehyde is the better nucleophile.

Assuming furan-2-aldehyde and NF to be the key intermediates in MFT and FFT formation, there should, consequently, be a correlation between the amounts of FA and FFT as well as between those of NF and MFT formed in processed carbohydrate/cysteine mixtures. As shown in Table 5, nearly 10 times more furan-2-aldehyde was generated in the ribose compared to the glucose system and simultaneously more FFT was generated. However, these data also corroborate the key role of furan-2-aldehyde in FFT generation from hexoses. In contrast, the amounts of NF were not well correlated with those of MFT. From ribose, by a factor of 3000 more NF was formed compared to MFT, whereas in the glucose system only by a factor of 10 more NF was present compared to MFT. This discrepancy implied that the MFT formation might not run exclusively via NF as the key intermediate.

We, therefore, thought of an alternative route for the formation of MFT starting from the possible precusors mercapto-2-propanone and hydroxyacetaldehyde. As proposed in Figure 3, depending on the side of the nucleophilic attack, the Aldol reaction of mercapto-2-propanone and hydroxyacetaldehyde might result in the formation of either MFT or FFT. Reaction of the methylene group of the mercapto-2-propanone gives the intermediate 4,5-dihydroxy-3-mercapto-2-pentanone (**Ia** in Figure 3), which upon cyclization and dehydration should give MFT (**Ib**). On the other hand, a nucleophilic attack of the methyl group of the mercapto-2-propanone on hydroxyacetaldehyde leads to the formation of 4,5-dihydroxy-1-mercapto-2-pentanone (**IIa**), which, after cyclization and elimination of water, might result in FFT (**IIb**). To elucidate this hypothesis, first, the formation of mercapto-2-propanone from 2-oxopropanal and H_2S was studied. The data indicated that high amounts of mercapto-2-propanone were generated from H_2S and the oxoaldehyde (Table 7). Because higher H_2S concentrations increased the formation of the mercaptoketone (Table 7), it might be assumed that H_2S is the limiting factor in this reaction.

In a next step, hydroxyacetaldehyde was reacted with mercapto-2-propanone in aqueous solution and the amounts of FFT and MFT were determined (Table 8). At pH 3.0, low concentrations of both odorants were formed. Increasing the pH to 5.0 significantly favored the formation of the MFT, whereas the increase in FFT concentration was low (Table 8). To study the influence of the reaction temperature and the water content on the formation of FFT and MFT, both intermediates were



Figure 3. Reaction pathway leading from hydroxyacetaldehyde and mercapto-2-propanone to FFT and MFT.

Table 7. Generation of Mercapto-2-propanone from 2-Oxopropanal and Hydrogen Sulfide[a]

reaction system		mercapto-2-propanone	
2-oxopropanal (mmol)	H_2S (mmol)	μg[b]	%
1	1	1650	1.8
1	2	3600	4.0

[a] An aqueous solution (50 mL) of hydrogen sulfide and 2-oxopropanal was reacted at pH 5.0 in a laboratory autoclave. [b] See footnote *b* in Table 1.

dry-heated and the odorant formation was monitored (experiment 4; Table 8). The results showed that compared to an aqueous system (cf. experiments 2 and 4; Table 8), the FFT formation increased slightly during dry-heating. However, the MFT generation was enhanced by a factor of nearly 6. The significant increase of MFT during dry-heating was well in line with the influence of the heating conditions on its generation from glucose or rhamnose (cf. Table 2).

The results indicate an important role of mercapto-2-propanone and hydroxyacetaldehyde in the formation of MFT from hexoses. To establish that this reaction is involved in MFT and FFT formation, the concentration of mercapto-2-propanone generated in thermally treated carbohydrate/cysteine mixtures was measured.

Case No.: IPR2023-00206
U.S. Patent No. 9,943,096

Table 8. Amounts of FFT and MFT Generated from Hydroxyacetaldehyde and Mercapto-2-propanone[a]

		FFT		MFT	
expt	pH	μg[b]	%	μg[b]	%
1	3.0	26.1	0.02	15.5	0.01
2	5.0	40.5	0.04	268.1	0.23
3	7.0	58.2	0.05	311.5	0.27
4	5.0[c]	51.3	0.05	1553.9	1.39

[a] An aqueous solution (50 mL) of hydroxyacetaldehyde (1 mmol) and mercapto-2-propanone (1 mmol) was reacted in a laboratory autoclave. [b] See footnote *b* in Table 1. [c] A mixture of hydroxyacetaldehyde (1 mmol) and mercapto-2-propanone (1 mmol) was intimately mixed with silica gel (3.0 g containing 300 μL of phosphate buffer, pH 5.0) and was dry-heated for 6 min at 180 °C.

Table 9. Formation of Mercapto-2-propanone from Glucose and Cysteine[a]

pH	amount[b] (μg)
3.0	11.4
5.0	59.6
7.0	26.5

[a] A solution (100 mL) of glucose (10.0 mmol) and cysteine (3.3 mmol) was reacted in a laboratory autoclave. [b] See footnote *b* in Table 1.

Table 10. Amounts of MFT Liberated from Various Precursors or Precursor Systems[a]

	MFT	
precursor (1.0 mmol) each	μg[b]	%
hydroxyacetaldehyde/mercapto-2-propanone	268.1	0.24
NF/H_2S	211.2	0.19
NF/cysteine	50.8	0.05
thiamin	8.2	0.01

[a] An aqueous solution (50 mL) of the precursors was reacted at pH 5.0 in a laboratory autoclave. [b] See footnote *b* in Table 1.

The data (Table 9) confirmed that the mercapto-2-propanone is indeed formed when carbohydrates are reacted in the presence of cysteine, with its formation going through a maximum at pH 5.0. This result is consistent with the preferential generation of both odorants at this pH value, e.g., in the glucose/cysteine mixture (Table 2).

Besides carbohydrates, thiamin especially has been reported as an important precursor of MFT. To study the effectiveness of MFT formation from thiamin, MFT was quantified in a heated aqueous solution of thiamin and the yields were compared with those obtained in the various precursor systems described above. The data established the binary mixtures hydroxacetaldehyde/mercapto-2-propanone followed by NF/H_2S as the most effective precursors systems (Table 10). Thiamin was less efficient, because by factors of 30 or 25 less MFT was generated compared to the two binary precursor mixtures.

Because, on the one hand, the efficiency of thiamin in MFT generation is very low and, on the other hand, thiamin is only a minor compound in foods, the liberation of MFT from thiamine during food processing should not contribute much to food flavors. Furthermore, since the carbohydrate fraction of foods contains usually more hexoses than pentoses, it seems likely that mercapto-2-propanone and hydroxyacetaldehyde are important intermediates involved in MFT generation during food processing.

LITERATURE CITED

Blank, I.; Sen, A.; Grosch, W. Potent odorants of roasted powder and brew of arabica coffee. *Z. Lebensm. Unters. Forsch.* **1992**, *195*, 239–245.

Farmer, L. J.; Mottram, D. S. Recent studies on the formation of meat-like aroma compounds. In *Flavour Science and Technology*; Bessier, Y., Thomas, A. F., Eds.; Wiley: Chichester, U.K., 1990a; pp 113–116.

Farmer, L. J.; Mottram, D. S. Interaction of lipid in the Maillard reaction between cysteine ad ribose: the effect of a triglyceride and three phospholipids on the volatile products. *J. Sci. Food Agric.* **1990b**, *53*, 505–525.

Garbusov, W.; Rehefeld, G.; Wölm, G.; Golvna, R. V.; Rothe, M. Volatile sulfur compounds contributing to meat flavour. Part I. Components identified in boiled meat. *Nahrung* **1976**, *20*, 235–238.

Gasser, U. Volatile aroma compounds in cooked beef and hen meat, meat flavorings and meat extract. Ph.D. Dissertation, Technical University of Munich, 1990.

Gasser, U.; Grosch, W. Identification of volatile flavour compounds with high aroma values from cooked beef. *Z. Lebensm. Unters. Forsch.* **1988**, *186*, 489–494.

Güntert, M.; Brüning, J.; Emberger, R.; Köpsel, M.; Kuhn, W.; Thielmann, T.; Werkhoff, P. Identification and formation of some selected sulfur-containing flavour compounds in various meat model systems. *J. Agric. Food Chem.* **1990**, *38*, 2027–2041.

Guth, H.; Hofmann, T.; Schieberle, P.; Grosch, W. Model reactions on the stability of disulfides in heated foods. *J. Agric. Food Chem.* **1995**, *43*, 2199–2203.

Hartmann, G. J.; Carlin, J. T.; Scheide, J. D.; Ho., C. T. Volatile products formed from the thermal degradation of thiamin at high and low moisture levels. *J. Agric. Food Chem.* **1984**, *32*, 1014–1018.

Hayashi, T.; Namiki, M. Role of sugar fragmentation in an early stage browning of amino-carbonyl reaction of sugar with amino acid. *Agric. Biol. Chem.* **1986**, *50*, 1965–1970.

Hincelin, O.; Ames, J. M.; Apriyanto, A.; Elmore, J. S. The effect of xylose on the generation of volatiles from heated thiamine. *Food Chem.* **1992**, *44*, 381–389.

Hofmann, T.; Schieberle, P. Evaluation of the key odorants in a thermally treated solution of ribose and cysteine by aroma extract dilution techniques. *J. Agric. Food Chem.* **1995**, *43*, 2187–2194.

Hofmann, T.; Schieberle, P. Identification of potent aroma compounds in thermally treated mixtures of glucose/cysteine and rhamnose/cysteine using aroma extract dilution techniques. *J. Agric Food Chem.* **1997**, *45*, 898–906.

Ledl, F.; Severin, T. Thermal degradation of cysteine and xylose in tributyrin. *Chem. Microbiol. Technol. Lebensm.* **1973**, *2*, 155–160.

Madson, M. A.; Feather, M. S. An improved preparation of 3-deoxy-D-erythro-hexos-2-ulose via the bis(benzoylhydrazone) and some related constitutional studies. *Carbohydr. Res.* **1981**, *94*, 183–191.

Martin, M. On the Maillard reaction of cysteine and methionine with reducing sugars. Ph.D. Dissertation, Technical University of Berlin, 1988.

Meynier, A.; Mottram, D. S. The effect of pH on the formation of volatiles compounds in meat-releated model systems. *Food Chem.* **1995**, *52*, 361–366.

Mottram, D. S. The effect of lipid on the formation of volatile heterocyclic compounds in the Maillard reaction. In *Flavour Science and Technology*; Martens, M., Dalen, G. A., Russwurm, H., Eds.; Wiley: Chichester, U.K., 1987; pp 29–34.

Mottram, D. S.; Whitfield, F. B. Aroma volatiles from meat-like Maillard systems. In *Thermally Generated Flavors: Maillard, Microwave and Extrusion Processes*; Parliment, T. H., Morello, M. J., McGorrin, R. J., Eds.; ACS Symposium Series 543; American Chemical Society: Washington, DC, 1994; pp 180–191.

Mottram, D. S.; Whitfield, F. B. Volatile compounds from the reaction of cysteine, ribose, and phospholipid in low moisture systems. *J. Agric. Food Chem.* **1995**, *43*, 984–988.

Case No.: IPR2023-00206
U.S. Patent No. 9,943,096

Münch, P.; Hofmann, T.; Schieberle, P. Comparison of key odorants generated by thermal treatment of commercial and self-prepared yeast extracts: influence of the amino acid composition on the odorant formation. *J. Agric. Food Chem.* **1997**, *45*, 1338−1344.

Nedvidek, W.; Ledl, F.; Fischer, P. Detection of 5-hydroxymethyl-2-methyl-3(2H)-furanone and of α-dicarbonyl compounds in reaction mixtures of hexoses and pentoses with different amines. *Z. Lebensm. Unters. Forsch.* **1992**, *194*, 222−228.

Reichstein, T.; Staudinger, H. GB Pat. 260, 1926, 960.

Reineccius, G. A.; Liardon, R. The use of characoal traps and microwave desorption for the analysis of headspace volatiles above heated thiamine solutions. In *Topics in Flavor Research*; Berger, R. G., Nitz, S., Schreier, P., Eds.; Eichhorn: Hangenham, Germany, 1985; pp 125−136.

Schieberle, P. Primary odorants in popcorn. *J. Agric. Food Chem.* **1991**, *39*, 1141−1144.

Schieberle, P. New developments in methods for analysis of volatile flavor compounds and their precursors. In *Characterization of Food: Emerging Methods*; Goankar, A. G., Ed.; Elsevier: Amsterdam, 1995; pp 403−431.

Schieberle, P. Odour-active compounds in moderately roasted sesame. *Food Chem.* **1996**, *55*, 145−152.

Schieberle, P.; Hofmann, T. Identification of the key odorants in processed ribose-cysteine Maillard mixtures by instrumental analysis and sensory studies. In *Flavour Science*; Taylor, A. J., Mottram, D. S., Eds.; The Royal Society of Chemistry, London, 1997; pp 175−181.

Sen, A.; Grosch, W. Synthesis of six deuterated sulfur containing odorants to be used as internal standards in quantification assays. *Z. Lebensm. Unters. Forsch.* **1991**, *192*, 541−547.

Sen, A.; Schieberle, P.; Grosch, W. Quantitative determination of 2,5-dimethyl-4-hydroxy-3(2H)-furanone and its methyl ether using a stable isotope dilution assay. *Lebensm. Wiss. Technol.* **1991**, *24*, 364−369.

Shibamoto, T.; Russel, G. F. Study of meat volatiles associated with aroma generated in a D-glucose−hydrogen sulfide−ammonia model system. *J. Agric. Food Chem.* **1976**, *24*, 843−846.

Shibamoto, T.; Russel, G. F. A study of the volatiles isolated from a D-glucose−hydrogen sulfide−ammonia model system. *J. Agric. Food Chem.* **1977**, *25*, 109−112.

Silwar, R.; Tressl, R. GC/MS analysis of odorants formed in a cysteine/methionine, furfural model system. *Z. Lebensm. Unters. Forsch.* **1989**, *189*, 205−211.

Van den Ouweland, G. A. M.; Peer, H. G. Components contributing to beef flavor. Volatile compounds produced by the reaction of 4-hydroxy-5-methyl-3(2H)-furanone and its thio analog with hydrogen sulfide. *J. Agric. Food Chem.* **1975**, *23*, 501−505.

Van der Linde, L. M.; van Dort, J. M.; de Valois, P.; Boelens, H.; de Rijke, D. Volatile components from thermally degraded thiamin. In *Progress in Flavor Research*; Land, D. G., Nursten, E. H., Eds.; Applied Science Publishers: London, 1979; pp 219−224.

Werkhoff, P.; Brüning, J.; Emberger, R.; Güntert, M.; Köpsel, M.; Kuhn, W.; Surburg, H. Isolation and characterization of volatile sulfur-containing meat flavor components in model systems. *J. Agric. Food Chem.* **1990**, *38*, 777−791.

Whitfield, F. B.; Mottram, D. S.; Brock, S.; Puckey, D. J.; Salter, L. J. Effect of phospholipid on the formation of volatile heterocyclic compounds in heated aqueous solutions of amino acids and ribose. *J. Sci. Food Chem.* **1988**, *42*, 261−272.

Whitfield, F. B.; Mottram, D. S.; Shaw, K. J. The formation of some novel thiols and disulfides in model systems containing 4-hydroxy-5-methyl-3(2H)-furanone. In *Progress in Flavour Precursor Studies*; Schreier, P., Winterhalter, P., Eds.; Allured Publishing: Carol Stream, IL, 1993; pp 395−400.

Withycombe, D. A.; Mussinan, D. Identification of 2-methyl-3-furanthiol in steam distillate from canned tuna fish. *J. Food Sci.* **1988**, *53*, 658.

Yeo, H. C. H.; Shibamoto, T. Microwave-induced volatiles of the Maillard model system under different pH conditions. *J. Agric. Food Chem.* **1991**, *39*, 370−373.

Zeiler-Hilgart, G. Model experiments on the formation of 2-methyl-3-furanethiol and 2-furfurylthiol in cooked meat. Ph.D. Dissertation, Technical University of Munich, 1994.

Received for review July 11, 1997. Revised manuscript received October 14, 1997. Accepted October 15, 1997.[⊗]

JF9705983

[⊗] Abstract published in *Advance ACS Abstracts,* November 15, 1997.

Motif Exhibit 1049, Page 14 of 14