

AGRICULTURAL AND BIOLOGICAL CHEMISTRY

4

Vol. 54, APRIL, 1990

ANALYTICAL CHEMISTRY

ORGANIC CHEMISTRY

BIOLOGICAL CHEMISTRY

FOOD & NUTRITION

MICROBIOLOGY & FERMENTATION INDUSTRY

PESTICIDE CHEMISTRY

RECEIVED

JUN 18 1990

UNIV. WIS. LIBRARY

ABCHA 6 54 (4) 855-1108 (1990)

ISSN 0002-1369

Published by

JAPAN SOCIETY FOR BIOSCIENCE, BIOTECHNOLOGY, AND AGROCHEMISTRY

(Formerly THE AGRICULTURAL CHEMICAL SOCIETY OF JAPAN)

JAPAN SOCIETY FOR BIOSCIENCE, BIOTECHNOLOGY, AND AGROCHEMISTRY (NIPPON NŌGEIKAGAKU KAI)

President

Hideo CHIBA

Vice Presidents

Hiuga SAITO Akira KUNINAKA

EDITORIAL BOARD

Chairman: Toshimasa YANO

Tadashi ASAH
Etsushiro DOI
Kiyoshi ISONO
Tohru KOMANO
Junya MIZUTANI
Mitsugi SENDA

Teruhiko BEPPU
Akira ENDO
Kazuo IZAKI
Shingo MARUMO
Takahisa OHTA
Kenji SODA
Akira YOSHIDA

Seiya CHIBA
Morifusa ETO
Akio KOBAYASHI
Shigeki MIZUNO
Yasuji OSHIMA
Shozo TODA

Advisory Editor

Kin-ichirō SAKAGUCHI

Managing Editors

Takeshi KITAHARA Kazuo NAGAI

Associate Editors

Yasumasa KUWAHARA Akihiro OKITANI

Agricultural and Biological Chemistry publishes original papers on agricultural and biological chemistry. Manuscripts submitted for publication and inquiries regarding them should be sent to the Editorial Board, the Japan Society for Bioscience, Biotechnology, and Agrochemistry, Japan Academic Societies Center Building, 2-4-16, Yayoi, Bunkyo-ku, Tokyo 113, Japan. Authors are earnestly required to consult the latest "Instructions to Authors", which are published in Vol. 54, No. 1.

Agricultural and Biological Chemistry is published monthly. The annual overseas subscription rates are: members, U.S. \$ 70.00 and non-members, \$ 180.00.

The Japan Society for Bioscience, Biotechnology, and Agrochemistry also publishes another journal in Japanese, *Nippon Nōgeikagaku Kaishi*.

AGENCIES

JAPAN PUBLICATIONS TRADING Co., Ltd.

P. O. Box 5030, Tokyo International, Tokyo, Japan

MARUZEN Co., Ltd.

P. O. Box 5050, Tokyo International, Tokyo, Japan

©Copyright, 1990, by the Japan Society for Bioscience, Biotechnology, and Agrochemistry.

The publication of this Journal was supported in part by a Grant-in-Aid for Publication of Scientific Research Result from the Ministry of Education, Science and Culture, which is gratefully acknowledged.

CONTENTS

Vol. 54, No. 4

Analytical Chemistry

- Volatile Components of TO-YO-RAN Flowers (*Cymbidium faberi* and *Cymbidium virens*)
.....A. OMATA, S. NAKAMURA, K. YOMOGIDA, K. MORIAI, Y. ICHIKAWA and I. WATANABE... 1029
- (N) Separation of Asn-linked Sialyloligosaccharides Labeled with *p*-Aminobenzoic Acid Ethyl Ester by Highperformance Liquid Chromatography
.....M. OHTA, M. KOBATAKE, A. MATSUMURA and F. MATSUURA.... 1045

Organic Chemistry

- (N) Evidence for Brassinosteroids in Strobilus of *Equisetum arvense* L.... S. TAKATSUTO, H. ABE and K. GAMOH.... 1057
- (N) Suppression of Tumor Promoter-induced Inflammation of Mouse Ear by Ursolic Acid and 4,4-Dimethylcholestane Derivatives.... M. HIROTA, T. MORI, M. YOSHIDA and R. IRIYE.... 1073
- (N) (–)- ϵ -Viniferin and Related Oligostilbenes from *Carex pumila* Thunb. (Cyperaceae)
.....H. KURIHARA, J. KAWABATA, S. ICHIKAWA and J. MIZUTANI.... 1097
- (SC) Novel C₁₀-Kaurane Type of Diterpene (Oryzalde A), a New Antimicrobial Compound Isolated from Healthy Leaves of a Bacterial Leaf Blight-resistant Cultivar of Rice Plant
.....M. WATANABE, Y. SAKAI, T. TERAOKA, H. ABE, Y. KONO, J. UZAWA, K. KOBAYASHI, Y. SUZUKI and A. SAKURAI.... 1103
- (SC) Brassinolide Is Biosynthesized from Castasterone in *Catharanthus roseus* Crown Gall Cells
.....T. YOKOTA, Y. OGINO, N. TAKAHASHI, H. SAIMOTO, S. FUJIOKA and A. SAKURAI.... 1107

Biological Chemistry

- Action Pattern of *Aeromonas hydrophila* Chitinase on Partially *N*-Acetylated Chitosan
.....M. MITSUTOMI, A. OHTAKARA, T. FUKAMIZO and S. GOTO.... 871
- Purification and Some Properties of a Protopectin-solubilizing Enzyme that Has Potent Activity on Sugar Beet Protopectin.....T. SAKAI and T. SAKAMOTO.... 879
- Succinate Degradation through the Citric Acid Cycle in *Bradyrhizobium japonicum* J501 Bacteroids under Low Oxygen Concentrations
.....S. TAJIMA, I. KIMURA, K. KOUZAI and T. KASAI.... 891
- Purification, Properties, and Transglycosylation Reaction of β -*N*-Acetylhexosaminidase from *Nocardia orientalis*.....F. NANJO, M. ISHIKAWA, R. KATSUMI and K. SAKAI.... 899
- Purification and Some Properties of β -Fructofuranosidase I from *Arthrobacter* sp. K-1
.....K. FUJITA, K. HARA, H. HASHIMOTO and S. KITAHATA.... 913
- Purification and Some Properties of Endo-1,3- β -D-xylanase from *Pseudomonas* sp. PT-5
.....I. YAMAURA, T. MATSUMOTO, M. FUNATSU and E. MUKAI.... 921
- Changes in Dolichyl Fatty Ester Compositions during the Germination of Soybean Embryos
.....M. ISHINAGA, K. MUKAI and C. TANIBE.... 943
- Purification and Some Properties of Chitinase from the Stomach of Japanese Eel, *Anguilla japonica*
.....M. KONO, T. MATSUI, C. SHIMIZU and D. KOGA.... 973
- Cholesterol Biosynthesis Inhibitory Activities of Novel Triterpenoid Acids from *Kadsura heteroclita* and *K. longipedunculata*
.....K. KANGOURI, T. MIYOSHI, A. IKEDA, S. OMURA, L. LI and H. XUE.... 993
- Purification and Some Properties of Dipeptidyl Carboxypeptidase from *Bacillus pumilus*
.....Y. NAGAMORI, N. FUJISHIMA and S. OKADA.... 999
- (N) The Presence of Protein Disulfide Isomerase in the Yeast *Saccharomyces cerevisiae*
.....Y. KATAKURA, T. MIZUNAGA, T. MIURA and Y. MARUYAMA.... 1043
- (N) Ozone Tolerance and Glutathione Reductase in Tobacco Cultivars
.....K. TANAKA, T. MACHIDA and T. SUGIMOTO.... 1061
- (N) Enzymatic Deglycation of Fructosyl-lysine
.....N. WATANABE, M. OHTSUKA, S. TAKAHASHI, Y. SAKANO and D. FUJIMOTO.... 1063
- (N) Some Properties and Action of Poly(glutamic acid) Hydrolase II from *Micromonospora melanosporea* IFO 12515.....T. MURO, Y. NAGAMORI, S. OKADA and Y. TOMINAGA.... 1065
- (N) Purification and Characterization of Alliin Lyase from Welsh Onion, *Allium fistulosum* L.
.....M. FUJITA, M. ENDO and M. SANO.... 1077
- (N) A Simple Method for Elucidating Structures of Galactomanno-oligosaccharides by Sequential Actions of β -Mannosidase and α -Galactosidase
.....I. KUSAKABE, R. KANEKO, N. TAKADA, A. F. ZAMORA, W. L. FERNANDEZ and K. MURAKAMI.... 1081
- (N) Further Studies on the Histidine Residue Located at the High Affinity Saccharide-binding Site of Ricin E.....T. HATAKEYAMA, N. YAMASAKI and G. FUNATSU.... 1085

Continued on inside back cover

Continued from back cover

- (N) Fatty Acid Composition of Phospholipids from Rat Granulomas Induced by Soybean and Coconut Oils Containing Tocopherol Acetate
.....M. ISHINAGA, M. TANIMOTO, Y. TANIGUCHI, K. MUKAI and K. YOKORO....1091
- (N) Inhibition of Angiotensin-Converting Enzyme by Synthetic Peptide Fragments of Various β -Caseins
.....M. KOHMURA, N. NIO and Y. ARIYOSHI....1101

Food & Nutrition

- Characterization of Zinc Chelating Compounds in Instant Coffee
.....T. ASAKURA, Y. NAKAMURA, N. INOUE, M. MURATA and S. HOMMA.... 855
- Kinetic Analysis of Soy-protein Denaturation by a Temperature-programmed Heat-denaturation Technique.....H. YOSHII, T. FURUTA, S. NOMA and T. NODA.... 863
- Relationship between the Emulsion Stability and Phospholipid Distribution in the Aqueous Phases Inside and Outside of an Emulsion Droplet.....K. CHIBA and M. TADA.... 907✓
- Effect of Aging on the Rheological Properties of Gluten Gel.....T. MITA.... 927
- Curie-point Pyrolysis of Cellulose in the Presence of Potassium Malate
.....K. IZAWA, M. MATSUKURA and Y. ISHIZU.... 957
- Effects of Dietary Palm Oil and Tocotrienol Concentrate on Plasma Lipids, Eicosanoid Productions and Tissue Fatty Acid Compositions in Rats
.....K. IMAIZUMI, J. NAGATA, M. SUGANO, H. MAEDA and Y. HASHIMOTO.... 965
- Molecular Size Distribution of Glutenin from Wheat Flour and the Polymerization of Proteins during Purification.....G. DANNO, S. R. XU and M. NATAKE.... 985
- Studies on the Precursors of Monoterpene Alcohols in Tea Leaves
.....M. YANO, K. OKADA, K. KUBOTA and A. KOBAYASHI....1023
- Chemical Composition of Lipids, Especially Triacylglycerol, in Grape Seeds
.....M. OHNISHI, S. HIROSE, M. KAWAGUCHI, S. ITO and Y. FUJINO....1035
- (N) Major Antioxidative Substances in Leaves of Atsumi-kabu (Red Turnip, *Brassica campestris* L.)
.....K. IGARASHI, M. ITOH and T. HARADA....1053
- (N) Esterification of Ovalbumin Depresses the Nutritive Values
.....Y. FURUICHI, H. UMEKAWA and T. TAKAHASHI....1069
- (N) Effect of Immunoglobulin Production Stimulating Factors in Foodstuffs on Immunoglobulin Production of Human Lymphocytes
.....K. YAMADA, I. IKEDA, M. MAEDA, S. SHIRAHATA and H. MURAKAMI....1087
- (N) Effects of Phenyl Compounds on Proliferation and IgM Production of Human-Human Hybridoma HB4C5 Cells Cultured in Serum-free Medium
.....M. MAEDA, K. YAMADA, I. IKEDA, H. NAKAJIMA, M. TAJIMA and H. MURAKAMI....1093

Microbiology & Fermentation Industry

- Protoplast Transfection of *Lactobacillus casei* by Phage PL-1 DNA
.....K. WATANABE, Y. KAKITA, Y. NAKASHIMA and T. SASAKI.... 937
- Expression of Recombinant Bovine β -Lactoglobulin in *Escherichia coli*
.....C. A. BATT, L. D. RABSON, D. W. S. WONG and J. E. KINSELLA.... 949
- Cloning and Nucleotide Sequence of the *KHR* Killer Gene of *Saccharomyces cerevisiae*
.....K. GOTO, Y. IWATUKI, K. KITANO, T. OBATA and S. HARA.... 979
- A New Depsipeptide Antibiotic, Citropeptin....Y. HAYAKAWA, M. NAKAGAWA, Y. TODA and H. SETO....1007
- (N) Growth and Levels of Isocitrate Lyase in *Rhodospseudomonas* sp. No. 7 Incubated in a Medium Containing Ethanol and Dicarboxylic Acid
.....T. FUJII, T. WATANABE, N. OOMURA, H. TAHAMA and A. ANDO....1049

Pesticide Chemistry

- Effect of Salithion Enantiomers on Larval Growth, Carbohydrases, Acetylcholinesterase, Adenylate Cyclase Activities and Cyclic Adenosine 3',5'-monophosphate Level of *Musca domestica* and *Tribolium castaneum*
.....A. HIRASHIMA, R. UENO, K. OYAMA, H. KOGA and M. ETO....1013
- (N) Syntheses of 3-Nitro-2,4,6-trihydroxy-benzamide and Thiobenzamide Derivatives as Highly Potent Inhibitors of Photosynthetic Electron Transport
.....I. HONDA, K. YONEYAMA, M. KONNAI, N. TAKAHASHI and S. YOSHIDA....1071

Abstracts

NIPPON NŌGEIKAGAKU KAISHI Vol. 54, No. 4..... A5

((N), Note; (RP), Rapid Paper; (SC), Short Communication)

Expression of Recombinant Bovine β -Lactoglobulin in *Escherichia coli*

Carl A. BATT,* Laurel D. RABSON,[†] Dominic W. S. WONG,^{††}
and John E. KINSELLA

Department of Food Science, Cornell University,
Ithaca, New York 14853, U.S.A.

Received September 20, 1989

Bovine β -lactoglobulin A was expressed in *Escherichia coli* in its mature form. The gene was constructed using a cDNA clone which coded for amino acid residues Leu-11 to Ile-162 and a synthetic oligonucleotide coding for the initial 10 amino acids preceded by a translational start. The met- β -lactoglobulin was expressed using a tac promoter vector, pTTQ18, and accounted for approximately 15% of the total cellular protein. The recombinant met- β -lactoglobulin migrated with the same molecular weight as native β -lactoglobulin A on SDS-PAGE. The majority of the met- β -lactoglobulin produced was found in an insoluble form but could be solubilized using guanidine-HCl. The renatured preparation was >80% pure and migrated similarly to purified β -lactoglobulin A under nondenaturing conditions.

β -Lactoglobulin is a whey protein, found in the milk of animals including bovine species, sheep, deer, dogs, and pigs, but not in humans.¹⁾ It occurs as several different variants due to genetic differences in each species. It is a single polypeptide of 18 kDa which corresponds to 162 amino acid residues. The complete amino acid sequence of bovine β -lactoglobulin has been reported and genetic variations in the amino acid sequence identified.²⁾

The function of β -lactoglobulin is not clear, especially since the milk of certain mammalian species appear to be devoid of it. β -Lactoglobulin has a significant degree of amino acid sequence homology with retinol binding protein,¹⁾ which functions *in vivo* to bind and transport the nonpolar retinol molecule through the aqueous medium of the blood serum from the liver to the target cells. Crystallographic studies of both proteins have demonstrated that they share the same conformational structure as well; both structures

consist of 8 strands of antiparallel beta sheets twisted into a calyx-shaped beta-barrel, with the interior cavity of the calyx being predominantly hydrophobic.³⁾ Both proteins contain, at the bottom of the interior cavity, a tryptophan residue which has been identified as important for the binding of retinol to retinol binding protein. β -Lactoglobulin has been shown to bind retinol more strongly than retinol binding protein,⁴⁾ as well as binding other nonpolar molecules insoluble in an aqueous medium, such as nonanone. The binding coefficient for retinol is approximately 2×10^{-8} M.⁵⁾ Furthermore, it has been demonstrated that retinol bound in this manner is protected from chemical oxidation while being transported.

The gene coding for ovine β -lactoglobulin was first isolated from a cDNA library by Gaye *et al.*⁶⁾ Subsequently we isolated a gene coding for the bovine β -lactoglobulin A from a cDNA bank constructed from poly(A⁺)

* Present address: School of Medicine, SUNY-Stony Brook, NY, U.S.A.

†† Present address: USDA-WRRC, Albany, CA, U.S.A.

* Corresponding author: Carl A. BATT, 107 Stocking Hall, Cornell University, Ithaca, NY 14853, U.S.A.

Abbreviations: [], denotes plasmid carrying state; IPTG, isopropyl- β -D-thiogalactopyranoside.

mRNA isolated from bovine mammary gland.⁷⁾ The gene was isolated using rabbit polyclonal antibodies raised against bovine β -lactoglobulin. Its nucleotide sequence codes for β -lactoglobulin A, from amino acid residues Ile-12 to Ile-162, as based on the amino acid sequence reported by Braunitzer *et al.*⁸⁾ In addition to the 455-bp coding sequence, our clone, pB β L4-10, contains a 3' nontranslated region of approximately 270 bp. High level expression of β -lactoglobulin was required as a prelude to studying the structure/function relationships through the use of site-directed mutagenesis. This study describes the expression and purification of a met- β -lactoglobulin in *E. coli*. It also represents the first report to our knowledge of the expression of a milk protein in its mature form using *E. coli*.

Materials and Methods

Bacterial strains and plasmids. *E. coli* TG1 was used as a host strain for all experiments. pTTQ18 was purchased from Amersham Co., it is a *tac* promoter vector with a *lacI^s* gene.⁹⁾ The cDNA β -lactoglobulin gene was carried on pB β lac4-10.⁷⁾ It is a pBR322 derivative carrying a 700-bp G-C tailed fragment in the *Pst*I site of the β -lactamase gene and includes the 3' nontranslated region with the polyA⁺ tail.

DNA manipulations. Restriction enzymes were purchased from either Amersham Co. or Boehringer-Mannheim and used according to their instructions. Plasmid DNA was isolated using the large scale procedure of Maniatis *et al.*¹⁰⁾ For small scale isolation the alkaline lysis method of Maniatis *et al.*¹⁰⁾ was used. The oligonucleotides were synthesized using an Applied Biosystems Model 70 synthesizer, and were not purified before use.

Recombinant met- β -lactoglobulin production. *E. coli* TG1 [pTTQ18 β lac.7.7.1] was grown in LB medium at 37°C with shaking to an OD₆₀₀ of 0.5. Isopropyl- β -D-thiogalactopyranoside (IPTG) was added to a final concentration of 2 mM and the culture incubated for an additional 4 hr at 37°C. The cells were harvested by centrifugation and lysed by the method of Schein and Noteborn¹¹⁾ or by passage through a French pressure cell. The cell debris and inclusion bodies containing the met- β -lactoglobulin were collected by centrifugation.

Purification of recombinant met- β -lactoglobulin. The insoluble form of β -lactoglobulin contained in inclusion

bodies was solubilized by treatment with 8 M guanidine-HCl with 1 mM dithiothreitol. The solution was held at 4°C for 30 min and any insoluble material removed by centrifugation. The 8 M guanidine-HCl was diluted 1:10 in 10 mM potassium phosphate buffer (pH 6.2), 1 mM DTT, and 1 mM PMSF and then dialyzed against 100X volume of 10 mM phosphate buffer with 1 mM DTT, overnight at 4°C. Any insoluble material remaining after dialysis was removed by centrifugation and the supernatant stored at -20°C.

PAGE and Western analysis. Western analysis was done using a 12.5% SDS PAGE gel and electroblotted onto nitrocellulose membranes. For non-denaturing PAGE the samples were prepared without the addition of SDS in the sample buffer and SDS was omitted from both the stacking and resolving gels as well as from the tank buffer. β -Lactoglobulin forms A and B could be clearly separated on a non-denaturing PAGE under these conditions (data not shown). Immunologically reacting protein was detected using polyclonal antibodies raised in Flemish Giant rabbits.⁷⁾ Primary antibody/antigen complexes were detected using goat anti-rabbit antiserum conjugated to horseradish peroxidase and developed using the BioRad HRP substrate.

Results

Construction of met- β -lactoglobulin gene

The cDNA coding for the bovine β -lactoglobulin gene, pB β lac4-10, was missing the first ten amino acids from the mature protein.⁷⁾ Curiously, the initial cDNA clone for the ovine β -lactoglobulin gene was also truncated for the first 10 amino acids from the amino terminus of the mature protein. Attempts to identify additional cDNA clones containing these missing amino acids using pB β lac4-10 as a probe from the same bovine mammary cDNA bank were not successful. Therefore an oligonucleotide coding for the initial ten amino acids of mature β -lactoglobulin, an ATG translation start codon and a ribosome binding site (RBS) spaced 10 bp upstream from the translational start was synthesized (Fig. 1). The codons used for the first ten amino acids were chosen based upon an *E. coli* codon bias using a reverse translation program which specifically selects codons using a prescribed codon bias.¹²⁾ Both the top and bottom strands were synthesized. Other salient features of this oligonucleotide were the addition of a *Hpa*I site between the




Fig. 1. Schematic for Construction of the Met-r- β -lactoglobulin and Its Expression in pTTQ18.

pBβlac4-10 is a pBR322 derivative carrying the 0.7 kb cDNA coding for bovine β -lactoglobulin from Ile-12 to Ile-162 flanked by an *EcoRV* and an *XbaI* site. This fragment was cloned into the complimentary sites of the Bluescript and Amp^r transformants selected. A transformant carrying the appropriate insert was selected by restriction analysis and a 52-bp oligonucleotide inserted into the *HindIII/EcoRV* sites 5' to the cDNA. Amp^r transformants were selected and screened for the presence of the oligonucleotide using a unique *HpaI* site designed into the oligonucleotide. Finally a 525bp *KpnI/XbaI* fragment was removed from Bluescript and inserted into the complimentary sites of pTTQ18 for expression via the *tac* promoter. Amp^r transformants were screened for the r-met- β -lactoglobulin insert by restriction analysis.

RBS and the translational start in addition to a *Hind*III site at the 5' end.

The coding sequence for β -lactoglobulin on pB β lac4-10 was fortuitously defined by an *Eco*RV site on the 5' end and an *Xba*I site on the 3' end. The entire coding region was excised as an *Eco*RV/*Xba*I fragment and cloned into the same restriction sites into the multiple cloning site of Bluescript KS (Stratagene). The sole motivation for using this vector is the presence of an *Eco*RV site within the multiple cloning site. Recombinants were selected and confirmed by restriction analysis. Both strands of the synthetic oligonucleotide coding for the 5' end of the mature β -lactoglobulin gene were annealed, kinased and ligated into *Hind*III/*Eco*RV digested Bluescript/B β lac4-10. Recombinant clones carrying the oligonucleotide were selected based on the presence of the unique *Hpa*I site in the oligonucleotide. A 510 bp *Hind*III/*Xba*I fragment carrying the oligo- β -lactoglobulin construct was subcloned into M13mp19 and sequenced to confirm the alignment of the oligonucleotide-cDNA junction. All clones investigated contained the ex-

pected nucleotide sequence for the oligonucleotide, in frame with the coding region of the β -lactoglobulin gene.

The met- β -lactoglobulin gene was removed from Bluescript on a *Kpn*I/*Xba*I fragment and ligated into *Kpn*I/*Xba*I-digested pTTQ18. pTTQ18 is a *tac* promoter vector which carries a *lac*I^q gene insuring a low level of uninduced expression.⁹⁾ Recombinants carrying the met- β -lactoglobulin gene in the appropriate orientation were selected by restriction analysis. The final construct was designated pTTQ18- β lac.7.7.1.

Expression and purification of met- β -lactoglobulin

E. coli TG1 [pTTQ18 β lac.7.7.1] was grown in LB medium and at an OD₆₀₀ of 0.5, induced with 2 mM IPTG. Cells were grown for an additional 4 hrs and then harvested by centrifugation. The production of met- β -lactoglobulin in whole cells was demonstrated by Western analysis using rabbit anti- β -lactoglobulin serum and is presented in Fig. 2. No met- β -lactoglobulin was observed when the




Fig. 2. Western Analysis of *E. coli* TG-1 [pTTQ18 β lac.7.7.1] Lysates Separated on a 12.5% SDS-PAGE. Lane A, β -lactoglobulin (Sigma); lane B, no IPTG added; lane C, induced with 2 mM IPTG; lane D, 12,000 $\times$ g pellet from culture induced with IPTG. M > indicates the position of β -lactoglobulin A.




Fig. 3. SDS-PAGE Analysis of *E. coli* TG-1 [pTTQ18 β lac.7.7.1] Lysates.

Lane A, β -lactoglobulin (Sigma); lane B, Insoluble fraction after guanidine hydrochloride treatment and dialysis; lane C, soluble fraction after guanidine hydrochloride treatment and dialysis; lane D, 12,000 $\times$ g pellet from whole cell lysate. (M) > indicates the position of β -lactoglobulin A.

cultures were not induced with IPTG. Densitometric scanning of the lysate from an IPTG induced culture separated on a SDS-PAGE stained with Coomassie blue reveals that β -lactoglobulin accounts for approximately 15% of the total cellular protein (Fig. 3).

Many heterologous proteins expressed at high levels in *E. coli* form inclusion bodies.¹²⁾ Met- β -lactoglobulin is no exception and insoluble complexes of it were observed in cell lysates. Attempts to produce a soluble form of met- β -lactoglobulin by growing IPTG induced cultures at 30°C¹¹⁾ (data not shown) were not successful. The majority of the met- β -lactoglobulin formed was insoluble, with the overall production at 30°C less than that observed at 37°C (data not shown). Furthermore the insoluble met- β -lactoglobulin could not be

solubilized using fast-flow ion exchange chromatography as demonstrated for an SV40 T antigen peptide, Th, interleukin 2, and the v-myb oncoprotein by Hoess *et al.*¹⁴⁾

Inclusion bodies containing met- β -lactoglobulin could be solubilized using 8M guanidine-HCl and the met- β -lactoglobulin remained soluble after dialysis to in the presence of dithiothreitol. The solubilized met- β -lactoglobulin migrated with the same molecular weight as authentic bovine β -lactoglobulin on denaturing PAGE (Fig. 3). Under nondenaturing PAGE conditions which allow the separation of the A and the B forms of β -lactoglobulin, the met- β -lactoglobulin migrated similarly to bovine β -lactoglobulin A (Fig. 4). A trace of a higher molecular weight band, barely detectable, may represent the dimer in both the bovine and met- β -lactoglobulin, again suggesting that the recombinant form has a similar conformation to the native β -lactoglobulin.

Discussion

cDNA clones for a number of bovine milk proteins have been reported including those for α_{s1} casein, β casein, κ casein,¹⁵⁻¹⁷⁾ α -lactalbumin,¹⁸⁾ and β -lactoglobulin.^{7,19)} The α_{s1} casein has been expressed in *E. coli* using the *tac* promoter and an insoluble, immunologically reactive protein was observed with an estimated yield of 16 mg/l.²⁰⁾ We estimate that *E. coli* TG1 [pTTQ18 β lac.7.7.1] produces β -lactoglobulin in excess of 50 mg/l culture medium.

The high level of recombinant met- β -lactoglobulin produced in *E. coli* [pTTQ18 β lac.7.7.1] could be the result of a variety of features in our construct. First, the proper spatial alignment of the RBS and the translational start of the protein was provided in the design of the oligonucleotide. The effect of varying the distance between the RBS and translational start have been elegantly demonstrated, for example with the expression of chymosin.²¹⁾ A change in the spacing between the RBS and translational start from 8 bp to




Fig. 4. Western Analysis of Purified and Recombinant β -Lactoglobulin Separated on a 12.5% Non-denaturing-PAGE.

Lane A, β -lactoglobulin A (Sigma); lane B, soluble fraction after guanidine hydrochloride treatment and dialysis from lysate of *E. coli* TG-1 [pTTQ18 β lac.7.7.1] induced with IPTG. (M)>indicates position of monomer; (D)>indicates position of dimer.

6 bp resulted in a 20-fold decrease in the level of chymosin expressed. In our construct, the spacing between the RBS and the translational start was 8 bp, although we have not conducted an exhaustive study to demonstrate that this distance is optimal for expression. Second, we used an *E. coli* codon bias for the first ten amino acids of met- β -lactoglobulin, to insure efficient translation.²²⁾ Last, although the physiological role of β -lactoglobulin is not known, it has no demonstrated enzymatic activity and does not bind divalent cations. It should therefore not present any toxicological problems to *E. coli* even at high intracellular concentrations. pTTQ18 is tightly regulated by virtue of a *lacI*^q on the plasmid; expression is only obtained upon the addition of IPTG or another suitable inducer. Therefore the potential deleterious effects (if any) of β -lactoglobulin on growth are minimized until the culture is induced by the addition of IPTG.

Our current interest in β -lactoglobulin is focused on elucidating and manipulating its structure and hence its function. It has recently been suggested by Sawyer that both molecules belong to a larger class of evolutionarily related proteins, including others such as bilin-binding proteins from various insects, which function to transport insoluble or chemically fragile molecules, and he has pointed out the possibility of using this basic biological architecture to design artificial transport proteins for industrial applications.²³⁾ Continued work will serve as a model for the rational design of proteins to serve as receptors for small molecules and in general examine the molecular recognition of small molecules by proteins and designing proteins for specific food and industrial applications.

Acknowledgments. This work was supported by a grant from the Cornell Biotechnology Program which is

sponsored by the New York State Science and Technology Foundation, a consortium of industries, the U.S. Army Research Office, and the National Science Foundation. The authors wish to thank Lindsay Sawyer and John Brady for their helpful suggestions.

References

- 1) S. Pervaiz and K. Brew, *Science*, **228**, 335 (1985).
- 2) L. Creamer, D. Parry and G. Malcolm, *Arch. Biochem. Biophys.*, **227**, 98 (1983).
- 3) M. Papiz, L. Sawyer, E. Eliopoulos, A. North, J. Findlay, R. Sivaprasadarao, T. Jones, M. Newcomer and P. Kraulis, *Nature*, **324**, 383 (1986).
- 4) R. Hemley, B. E. Kohler and P. Siriski, *Biophys. J.*, **28**, 447 (1979).
- 5) R. D. Fugate and P.-S. Song, *Biochim. Biophys. Acta*, **625**, 28 (1980).
- 6) P. Gaye, D. Hue-Delahaie, J.-C. Mercier, S. Soulier, J. Vilotte and J.-P. Furet, *Biochimie*, **68**, 1097 (1986).
- 7) A. C. Jamieson, M. A. Vandeyar, Y. C. Kang, J. E. Kinsella and C. A. Batt, *Gene*, **61**, 85 (1987).
- 8) G. Braunitzer, R. Chen, B. Schrank and A. Stangl, *Hoppe-Seyler's Z. Physiol. Chem.*, **354**, 867 (1973).
- 9) M. J. R. Stark, *Gene*, **51**, 255 (1987).
- 10) T. Maniatis, E. F. Fritsch and J. Sambrook, "Molecular Cloning: A Laboratory Manual," Cold Spring Harbor Laboratory, Cold Spring Harbor.

New York, 1982.

- 11) C. H. Schein and M. H. M. Noteborn, *Bio/Technol.*, **6**, 291 (1988).
- 12) M. P. Weiner and H. A. Scheraga, *CABIOS*, **5**, 191 (1989).
- 13) J. K. Kane and D. J. Hartley, *Trends Biotechnol.*, **6**, 95 (1988).
- 14) A. Hoess, A. K. Arthur, G. Wanner and E. Fanning, *Bio/Technol.*, **6**, 1214 (1988).
- 15) A. Stewart, I. Willis and A. Mackinlay, *Nucl. Acids Res.*, **12**, 3895 (1984); J.-C. Mercier, P. Gaye, S. Soulier, D. Hue-Delahaie and J.-L. Vilotte, *Biochimie*, **67**, 959 (1985).
- 16) R. Jimenez-Flores, Y. Kang and T. Richardson, *Biochem. Biophys. Res. Commun.*, **142**, 617 (1987).
- 17) Y. C. Kang and T. Richardson, *J. Dairy Sci.*, **71**, 29 (1987).
- 18) W. L. Hurley and L. A. Schuler, *Gene*, **61**, 119 (1987).
- 19) V. N. Ivanov, E. S. Judinkova and S. I. Gorodetsky, *Biol. Chem. Hoppe-Seyler*, **369**, 425 (1988).
- 20) M. Nagao, Y. Nakagawa, A. Ishii, R. Sasaki, H. Tanaka and H. Chiba, *Agric. Biol. Chem.*, **52**, 191 (1988).
- 21) Y. Kawaguchi, N. Yanagida, T. Uozumi and T. Beppu, *Agric. Biol. Chem.*, **50**, 499 (1986).
- 22) B. A. Springer and S. G. Sligar, *Proc. Natl. Acad. Sci. U.S.A.*, **84**, 8961 (1987).
- 23) L. Sawyer, *Nature*, **327**, 659 (1987).