SDSU LIBRARY - BROOKINGS, SD
3 1574 50493 4784

Supplement to
The American Journal of
CLINICAL NUTRITION
Official Journal of The American Society for Clinical Nutrition, Inc
SEPTEMBER 2003 • VOLUME 78 • NUMBER 3(S)
ISSN 0002-9165

Proceedings of
a symposium held in
Loma Linda, CA

April 8–11, 2002

# Fourth International Congress on Vegetarian Nutrition

Guest
Scientific
Editors

Joan Sabaté
Sujatha Rajaram

612.3905
Am 3
vol. 78
no. 3 Supp.

# The American Journal of CLINICAL NUTRITION

*Official Publication of The American Society for Clinical Nutrition, Inc*

THE CLINICAL DIVISION OF THE AMERICAN SOCIETY FOR NUTRITIONAL SCIENCES

*The American Journal of Clinical Nutrition* (ISSN 0002-9165) is the official Journal of and is published monthly, with two volumes per year, by The American Society for Clinical Nutrition, Inc, 9650 Rockville Pike, Bethesda, MD 20814-3998 USA. The primary focus of the Journal is the publication of basic and clinical studies relevant to human nutrition. The Information for Authors appears in the January issue each year and is also available on the Internet at **WWW.AJCN.ORG**.


All rights reserved. © 2003 by The American Society for Clinical Nutrition. The contents of this publication may not be reproduced in whole or part without consent of the copyright owner. When an abstract appears at the beginning of an article, however, it may be reproduced in any form or translated without specific permission provided the original citation is included. The table of contents and cover also may be reproduced in any form or translated without specific permission.

Statements made and opinions expressed in Original Research Communications, Letters to the Editor, Editorials, Presidential Addresses, Book Reviews, and certain other special articles appearing in this Journal represent the views of the authors and do not necessarily reflect the position of the Journal or The American Society for Clinical Nutrition.

**Subscription Services:** Contact Dues and Subscriptions, *The American Journal of Clinical Nutrition*, 9650 Rockville Pike, L-2310, Bethesda, MD 20814-3998. Phone: (301) 530-7029. Fax: (301) 571-5728. E-mail: staff@dues.faseb.org. Print + online subscription rates 2003: members $80 per year; non-member individuals $155; institutions $245; certified students $60 (online only). Agents are allowed a 5% discount. Contact this office for international rates, changes of address, and back issue and supplement sales. Visa, MasterCard, and American Express accepted. Claims must be made within 90 days of mail date. Periodicals postage paid at Bethesda and additional mailing offices. Canadian IPM 0059811 and GST 134892330.

**Advertising Representative:** FASEB AdNet, 9650 Rockville Pike, Bethesda, MD 20814-3998. Phone: (800) 43-FASEB, ext 7157. Fax: (301) 571-0683. The American Society for Clinical Nutrition does not endorse any commercial enterprise.

**Redactory and Printing Services:** Cadmus Professional Communications, 1991 Northampton Street, Easton, PA 18042-9111 and its affiliate ScholarOne, Inc, 375 Greenbrier Drive, Suite 200, Charlottesville, VA 22901-1618. The Journal is printed in Easton, PA, by Cadmus Professional Communications.

**Postmaster:** Send address changes to *The American Journal of Clinical Nutrition*, 9650 Rockville Pike, L-2310, Bethesda, MD 20814-3998. Be sure to include a copy of the old address label.


**Editor-in-Chief**
Charles H Halsted, MD

**Manuscripts Editor**
D'Ann Finley, PhD

**Administrative Assistant**
Sharon Lovejoy
Zann Gates

**Associate Editors**
Richard J Havel, MD
Janet C King, PhD
Robert Rucker, PhD
Robert M Russell, MD

**Editorial Office: Send manuscripts to** Editorial Office, *The American Journal of Clinical Nutrition*, 3247 Meyer Hall, University of California, One Shields Avenue, Davis, CA 95616-8790. Phone: (530) 752-8363. Fax: (530) 752-8371. E-mail: ajcn@ucdavis.edu.

**Consulting Editors**

Steven A Abrams, MD
Stephanie A Atkinson, PhD, RD
Lars F Berglund, MD, PhD
Barbara A Bowman, PhD
Raymond Burk, MD
Benjamin Caballero, MD, PhD
Carroll E Cross, MD
Robert Eckel, MD
Kent Erickson, PhD
Balz Frei, PhD
Michael I Goran, PhD
K Michael Hambidge, MD, ScD
David Heber, MD, PhD
William C Heird, MD
Marc Hellerstein, MD, PhD
Steven B Heymsfield, MD
James Hill, MD
Sheila M Innis, PhD
S Jill James, PhD
Khursheed Jeejeebhoy, PhD
Peter Jones, PhD
Martijn B Katan, PhD
Ronald M Krauss, MD
Nancy Krebs, MD
Allen Levine, PhD
Bo Lonnerdal, PhD
Mary Frances Picciano, PhD
Helga M Refsum, MD
Susan B Roberts, PhD
Hamid Said, PhD, PharmD
Charles B Stephensen, PhD
Christine A Swanson, PhD
Elizabeth C Theil, PhD
Connie M Weaver, PhD
Bruce M Wolfe, MD
Regina G Ziegler, PhD
Thomas R Ziegler, MD

**Production Office:** For inquiries regarding copyright, single article sales, and reprints, contact *The American Journal of Clinical Nutrition*, 9650 Rockville Pike, Bethesda, MD 20814-3998. Phone: (301) 530-7038. Fax: (301) 571-8303. E-mail: journal@ascn.faseb.org. Internet: WWW.AJCN.ORG.

**Managing Editor**
Elizabeth C Horowitz

**Senior Technical Editors**
Brenda Allen Milot
Jennifer L Scales

**Technical Editors**
Christina G Chase
Darren T Early
Carolinda Hales

## The American Society for Clinical Nutrition, Inc

**SUSTAINING MEMBERS**

The American Society for Clinical Nutrition, Inc, is pleased to acknowledge the generous support of these organizations.

Abbott Laboratories, Ross Products Division
Campbell Soup Company
The Coca-Cola Company
The Dannon Company, Inc
Gerber Products Company
GlaxoSmithKline
The Kellogg Company
Kraft Foods
McNeil Nutritionals
Mead Johnson Nutritionals
Monsanto Company
National Dairy Council
Nestlé Nutrition Institute of Nestlé USA
The Procter & Gamble Company
Roche Vitamins Inc
Slim-Fast Foods Company
The Sugar Association, Inc
Unilever Bestfoods
Wyeth Consumer Healthcare
Wyeth Nutrition

**OFFICERS 2002–2003**

**President**
Robert M Russell, MD

**Vice President**
Dale A Schoeller, PhD

**Vice President-Elect**

**Past President**
Steven B Heymsfield, MD

**Secretary**
Janet C King, PhD

**Treasurer**

**Treasurer-Elect**
Richard J Deckelbaum, MD

**Councilors**
Naomi K Fukagawa, MD, PhD
Benjamin Caballero, MD, PhD
Ronenn Roubenoff, MD, MHS

**ASNS Representative**
Sharon M Donovan, PhD, RD

**Editor-in-Chief**
Charles H Halsted, MD

**Executive Officer**
Sandra A Schlicker, PhD

Supplement to
The American Journal of

# CLINICAL NUTRITION

VOLUME 78 NUMBER 3(S) SEPTEMBER 2003

## Fourth International Congress on Vegetarian Nutrition

501S Preface. *J Sabaté and S Rajaram*

### Nutrient, food, and dietary patterns

502S The contribution of vegetarian diets to health and disease: a paradigm shift? *J Sabaté*

508S Nutrients, foods, and dietary patterns as exposures in research: a framework for food synergy. *DR Jacobs Jr and LM Steffen*

514S Transcending reductionism in nutrition research. *I Hoffmann*

517S Health benefits of fruit and vegetables are from additive and synergistic combinations of phytochemicals. *RH Liu*

521S A search for truth in dietary epidemiology. *GE Fraser*

### Epidemiologic studies of vegetarians

526S Does low meat consumption increase life expectancy in humans? *PN Singh, J Sabaté, and GE Fraser*

533S Mortality in British vegetarians: review and preliminary results from EPIC-Oxford. *TJ Key, PN Appleby, GK Davey, NE Allen, EA Spencer, and RC Travis*

539S Lessons from dietary studies in Adventists and questions for the future. *W Willett*

### Vegetarian diets and risk of selected chronic diseases

544S Plant-based foods and prevention of cardiovascular disease: an overview. *FB Hu*

552S The effect of vegetarian diet, plant foods, and phytochemicals on hemostasis and thrombosis. *S Rajaram*

559S Epidemiologic evidence of the protective effect of fruit and vegetables on cancer risk. *E Riboli and T Norat*

570S Fruits and vegetables in the prevention of cellular oxidative damage. *RL Prior*

579S Spicing up a vegetarian diet: chemopreventive effects of phytochemicals. *JW Lampe*

584S Dietary protein, calcium metabolism, and skeletal homeostasis revisited. *JE Kerstetter, KO O'Brien, and KL Insogna*

593S Dietary phytoestrogens and their effect on bone: evidence from in vitro and in vivo, human observational, and dietary intervention studies. *KDR Setchell and E Lydeking-Olsen*

610S Type 2 diabetes and the vegetarian diet. *DJA Jenkins, CWC Kendall, A Marchie, AL Jenkins, LSA Augustin, DS Ludwig, ND Barnard, and JW Anderson*

617S Dietary *cis*-monounsaturated fatty acids and metabolic control in type 2 diabetes. *E Ros*

### Nutritional adequacy and public health issues

626S What do vegetarians in the United States eat? *EH Haddad and JS Tanzman*

(Continued)

633S Bioavailability of iron, zinc, and other trace minerals from vegetarian diets. *JR Hunt*

640S Achieving optimal essential fatty acid status in vegetarians: current knowledge and practical implications. *BC Davis and PM Kris-Etherton*

647S Nut consumption and body weight. *J Sabaté*

651S Variant Creutzfeldt-Jakob disease: pathology, epidemiology, and public health implications. *CR Trevitt and PN Singh*

**Environmental impact of vegetarian diets**

657S Nutrition ecology: the contribution of vegetarian diets. *C Leitzmann*

660S Sustainability of meat-based and plant-based diets and the environment. *D Pimentel and M Pimentel*

664S Quantification of the environmental impact of different dietary protein choices. *L Reijnders and S Soret*

This material may be protected by Copyright law (Title 17 U.S. Code)

# Sustainability of meat-based and plant-based diets and the environment[1–3]

*David Pimentel and Marcia Pimentel*

**ABSTRACT** Worldwide, an estimated 2 billion people live primarily on a meat-based diet, while an estimated 4 billion live primarily on a plant-based diet. The US food production system uses about 50% of the total US land area, 80% of the fresh water, and 17% of the fossil energy used in the country. The heavy dependence on fossil energy suggests that the US food system, whether meat-based or plant-based, is not sustainable. The use of land and energy resources devoted to an average meat-based diet compared with a lactoovovegetarian (plant-based) diet is analyzed in this report. In both diets, the daily quantity of calories consumed are kept constant at about 3533 kcal per person. The meat-based food system requires more energy, land, and water resources than the lactoovovegetarian diet. In this limited sense, the lactoovovegetarian diet is more sustainable than the average American meat-based diet. *Am J Clin Nutr* 2003;78(suppl):660S–3S.

**KEY WORDS** Meat-based diet, plant-based diet, environment, natural resources, fossil, energy, fuel

## INTRODUCTION

Worldwide, an estimated 2 billion people live primarily on a meat-based diet, while an estimated 4 billion live primarily on a plant-based diet. The shortages of cropland, fresh water, and energy resources require most of the 4 billion people to live on a plant-based diet. The World Health Organization recently reported that more than 3 billion people are malnourished (1, 2). This is the largest number and proportion of malnourished people ever recorded in history. In large measure, the food shortage and malnourishment problem is primarily related to rapid population growth in the world plus the declining per capita availability of land, water, and energy resources (3).

Like the world population, the US population continues to grow rapidly. The US population doubled in the past 60 y and is projected to double again in the next 70 y (4) (**Figure 1**). The US food production system uses about 50% of the total US land area, approximately 80% of the fresh water, and 17% of the fossil energy used in the country (3). The heavy dependence on fossil energy suggests that the US food system, whether meat-based or plant-based, is not sustainable. The use of land and energy resources devoted to an average meat-based diet compared with a lactoovovegetarian (plant-based) diet is analyzed in this report. In both diets, the daily quantity of calories consumed was kept constant at about 3533 kcal per person.

## LACTOOVOVEGETARIAN DIET

The lactoovovegetarian diet was selected for this analysis because most vegetarians are on this or some modified version of this diet. In addition, the American Heart Association reported that the lactoovovegetarian diet enables individuals to meet basic nutrient needs (5).

A comparison of the calorie and food consumption of a lactoovovegetarian diet and a meat-based diet is provided in **Table 1**. In the lactoovovegetarian diet, the meat and fish calories were replaced by proportionately increasing most other foods consumed in Table 1 in the vegetarian diet except sugar and sweeteners, fats, and vegetable oils. The total weight of food consumed was slightly higher (1002 kg per year) in the lactoovovegetarian diet than in the meat-based diet (995 kg per year). The most food calories consumed in both diets were associated with food grains, and the second largest amount of calories consumed was from sugar and sweeteners.

The amount of feed grains used to produce the animal products (milk and eggs) consumed in the lactoovovegetarian diet was about half (450 kg) the amount of feed grains fed to the livestock (816 kg) to produce the animal products consumed in the meat-based diet (Table 1). This is expected because of the relatively large amount of animal products consumed in the meat-based diet (7). Less than 0.4 ha of cropland was used to produce the food for the vegetarian-based diet, whereas about 0.5 ha of cropland was used in the meat-based diet (8). This reflects the larger amount of land needed to produce the meat-based diet (Table 1).

The major fossil energy inputs for grain, vegetable, and forage production include fertilizers, agricultural machinery, fuel, irrigation, and pesticides (8, 9). The energy inputs vary according to the crops being grown (10). When these inputs are balanced against their energy and protein content, grains and some legumes, such as soybeans, are produced more efficiently in terms of energy inputs than vegetables, fruits, and animal products (8). In the United States, the average protein yield from a grain crop such as corn is 720 kg/ha (10). To produce 1 kcal of plant protein requires an input of about 2.2 kcal of fossil energy (10).

## MEAT-BASED DIET

The meat-based diet differs from the vegetarian diet in that 124 kg of meat and 20.3 kg of fish are consumed per year (Table 1). Note that the number of calories is the same for both diets because the vegetarian foods consumed were proportionately

[1] From the Department of Ecology and Evolutionary Biology, Cornell University, Ithaca, NY.

[2] Presented at the Fourth International Congress on Vegetarian Nutrition, held in Loma Linda, CA, April 8–11, 2002. Published proceedings edited by Joan Sabaté and Sujatha Rajaram, Loma Linda University, Loma Linda, CA.

[3] Address reprint requests to D Pimentel, Department of Ecology and Evolutionary Biology, Cornell University, 5126 Comstock Hall, Ithaca, NY 14853. E-mail: dp18@cornell.edu.



*Am J Clin Nutr* 2003;78(suppl):660S–3S. Printed in USA. © 2003 American Society for Clinical Nutrition

Case No.: IPR2023-00208
U.S. Patent No. 11,013,250






**FIGURE 1.** Projection of US population growth in the next 70 y (4).

increased to make sure that both diets contained the same number of calories. The total calories in the meat and fish consumed per day was 480 kcal. The foods in the meat-based diet providing the most calories were food grains and sugar and sweeteners—similar to the lactoovovegetarian diet.

In the United States, more than 9 billion livestock are maintained to supply the animal protein consumed each year (11). This livestock population on average outweighs the US human population by about 5 times. Some livestock, such as poultry and hogs, consume only grains, whereas dairy cattle, beef cattle, and lambs consume both grains and forage. At present, the US livestock population consumes more than 7 times as much grain as is consumed directly by the entire American population (11). The amount of grains fed to US livestock is sufficient to feed about 840 million people who follow a plant-based diet (7). From the US livestock population, a total of about 8 million tons (metric) of animal protein is produced annually. With an average distribution assumed, this protein is sufficient to supply about 77 g of animal protein daily per American. With the addition of about 35 g of available plant protein consumed per person, a total of 112 g of protein is available per capita in the United States per day (11). Note that the recommended daily allowance (RDA) for adults per day is 56 g of protein from a mixed diet. Therefore, based on these data, each American consumes about twice the RDA for protein. Americans on average are eating too much and are consuming about 1000 kcal in excess per day per capita (12, 13). The protein consumed per day on the lactoovovegetarian diet is 89 g per day. This is significantly lower than the 112 g for the meat-based diet but still much higher than the RDA of 56 g per day.

About 124 kg of meat is eaten per American per year (6). Of the meat eaten, beef amounts to 44 kg, pork 31 kg, poultry 48 kg, and other meats 1 kg. Additional animal protein is obtained from the consumption of milk, eggs, and fish. For every 1 kg of high-quality animal protein produced, livestock are fed about 6 kg of plant protein. In the conversion of plant protein to animal protein, there are 2 principal inputs or costs: 1) the direct costs of production of the harvest animal, including its feed; and 2) the indirect costs for maintaining the breeding herds.

Fossil energy is expended in livestock production systems (**Table 2**). For example, broiler chicken production is the most efficient, with an input of 4 kcal of fossil energy for each 1 kcal of broiler protein produced. The broiler system is primarily dependent on grain. Turkey, also a grain-fed system, is next in efficiency, with a ratio of 10:1. Milk production, based on a mixture of two-thirds grain and one-third forage, is relatively efficient, with a ratio of 14:1. Both pork and egg production also depend on grain. Pork production has a ratio of 14:1, whereas egg production has a 39:1 ratio.

The 2 livestock systems depending most heavily on forage but also using significant amounts of grain are the beef and lamb production systems (**Table 3**). The beef system has a ratio of 40:1, while the lamb has the highest, with a ratio of 57:1 (Table 2). If these animals were fed on only good-quality pasture, the energy inputs could be reduced by about half.

The average fossil energy input for all the animal protein production systems studied is 25 kcal fossil energy input per 1 kcal of protein produced (Table 2). This energy input is more than 11 times

**TABLE 1**
Per capita food consumption, energy, and protein of foods of a meat-based compared with a lactoovovegetarian diet in the United States

| Food | Meat-based diet[1] | Energy | Protein | Lactoovovegetarian diet[2] | Energy | Protein |
|---|---|---|---|---|---|---|
| | *kg* | *kcal* | *g* | *kg* | *kcal* | *g* |
| Food grain | 114 | 849 | 24.9 | 152 | 1132 | 33.2 |
| Pulses (legumes) | 4.3 | 40 | 2.0 | 7.5 | 70 | 4.5 |
| Vegetables | 239 | 147 | 6.6 | 286 | 155 | 8.8 |
| Oil crops | 6 | 71 | 3.0 | 8 | 95 | 4.0 |
| Fruit | 109 | 122 | 1.4 | 112 | 122 | 1.9 |
| Meat | 124 | 452 | 41.1 | 0 | 0 | 0 |
| Fish | 20.3 | 28 | 4.7 | 0 | 0 | 0 |
| Dairy products | 256 | 385 | 22.5 | 307.1 | 473 | 30.0 |
| Eggs | 14.5 | 55 | 4.2 | 19.2 | 73 | 5.6 |
| Vegetable oils | 24 | 548 | 0.2 | 25 | 570 | 0.2 |
| Animal fats | 6.7 | 127 | 0.1 | 6.7 | 127 | 0.1 |
| Sugar and sweeteners | 74 | 686 | 0.2 | 74 | 686 | 0.2 |
| Nuts | 3.1 | 23 | 0.6 | 4.0 | 30 | 0.8 |
| Total | 994.9 | 3533 | 111.5 | 1001.5 | 3533 | 89.3 |
| Feed grains[3] | 816.0 | — | — | 450.0 | — | — |

[1] Data from FAOSTAT (6).
[2] Estimated.
[3] Feed grains are cereal grains fed to livestock.

Motif Exhibit 1036, Page 6 of 8

Case No.: IPR2023-00208
U.S. Patent No. 11,013,250



**TABLE 2**
Animal production in the United States and the fossil energy required to produce 1 kcal of animal protein

| Livestock and animal products | Production volume[1] | Ratio of energy input to protein output[2] |
|---|---|---|
| | *× 10⁶* | *kcal* |
| Lamb | 7 | 57:1 |
| Beef cattle | 74 | 40:1 |
| Eggs | 77 000 | 39:1 |
| Swine | 60 | 14:1 |
| Dairy (milk) | 13 | 14:1 |
| Turkeys | 273 | 10:1 |
| Broilers | 8000 | 4:1 |

[1] Data from US Department of Agriculture (11).
[2] Data from Pimentel (9).

greater than that for grain protein production, which is about 2.2 kcal of fossil energy input per 1 kcal of plant protein produced (**Table 4**). This is for corn and assumes 9% protein in the corn. Animal protein is a complete protein based on its amino acid profile and has about 1.4 times the biological value of grain protein (8).

## LAND RESOURCES

More than 99.2% of US food is produced on land, while <0.8% comes from oceans and other aquatic ecosystems. The continued use and productivity of the land is a growing concern because of the rapid rate of soil erosion and degradation throughout the United States and the world. Each year about 90% of US cropland loses soil at a rate 13 times above the sustainable rate of 1 ton/ha/y (28). Also, US pastures and rangelands are losing soil at an average of 6 tons/ha/y. About 60% of United States pastureland is being overgrazed and is subject to accelerated erosion.

The concern about high rates of soil erosion in the United States and the world is evident when it is understood that it takes approximately 500 y to replace 25 mm (1 in) of lost soil (28). Clearly, a farmer cannot wait for the replacement of 25 mm of soil. Commercial fertilizers can replace some nutrient loss resulting from soil erosion, but this requires large inputs of fossil energy.

## WATER RESOURCES

Agricultural production, including livestock production, consumes more fresh water than any other activity in the United States. Western

**TABLE 3**
Grain and forage inputs per kilogram of animal product produced

| Livestock | Grain[1] | Forage[2] |
|---|---|---|
| | *kg* | *kg* |
| Lamb | 21 | 30 |
| Beef cattle | 13 | 30 |
| Eggs | 11 | — |
| Swine | 5.9 | — |
| Turkeys | 3.8 | — |
| Broilers | 2.3 | — |
| Dairy (milk) | 0.7 | 1 |

[1] Data from US Department of Agriculture (11).
[2] Data from Morrison (14) and Heitschmidt et al (15).

**TABLE 4**
Energy inputs and costs of corn production per hectare in the United States

| Inputs | Quantity | Energy | Cost |
|---|---|---|---|
| | | *kcal × 1000* | *$* |
| Labor (h)[1] | 11.4 (16)[2] | 462 | 114.00[3] |
| Machinery (kg) | 55 (8) | 1018 (17) | 103.21 (18) |
| Diesel (L) | 42.2 (19, 20) | 481 (17) | 8.87 (21) |
| Gasoline (L) | 32.4 (19, 20) | 328 (17) | 9.40 (21) |
| Nitrogen (kg) | 144.6 (22) | 2688 (23) | 89.65 (21) |
| Phosphorus (kg) | 62.8 (22) | 260 (23) | 34.54 (21) |
| Potassium (kg) | 54.9 (22) | 179 (23) | 17.02 (21) |
| Lime (kg) | 699 (22) | 220 (17) | 139.80 (16) |
| Seeds (kg) | 21 (8) | 520 (17) | 74.81 (24) |
| Irrigation (cm) | 33.7 (25) | 320 (17) | 123.00 |
| Herbicides (kg) | 3.2 (22) | 320 (17) | 64.00[4] |
| Insecticides (kg) | 0.92 (22) | 92 (17) | 18.40[4] |
| Electricity (kWh) | 13.2 (19, 20) | 34 (17) | 2.38[5] |
| Transportation (kg)[6] | 151 | 125 (17) | 45.30[7] |
| Total (kg yield) | 7965 (27) | 7047[8] | 844.38 |

[1] It is assumed that a person works 2000 h/y and uses an average of 8100 L oil equivalents/y.
[2] Reference.
[3] It is assumed that farm labor is paid $10/h.
[4] It is assumed that herbicide and insecticide prices are $20/kg.
[5] The price of electricity is $0.07/kWh (26).
[6] Goods transported include machinery, fuels, and seeds that were shipped an estimated 1000 km.
[7] Transport was estimated to cost $0.30/kg.
[8] Ratio of kcal input to output = 1:4.07.

agricultural irrigation accounts for 85% of the fresh water consumed (29). The water required to produce various foods and forage crops ranges from 500 to 2000 L of water per kilogram of crop produced. For instance, a hectare of US corn transpires more than 5 million L of water during the 3-mo growing season. If irrigation is required, more than 10 million L of water must be applied. Even with 800–1000 mm of annual rainfall in the US Corn Belt, corn usually suffers from lack of water in late July, when the corn is growing the most.

Producing 1 kg of animal protein requires about 100 times more water than producing 1 kg of grain protein (8). Livestock directly uses only 1.3% of the total water used in agriculture. However, when the water required for forage and grain production is included, the water requirements for livestock production dramatically increase. For example, producing 1 kg of fresh beef may require about 13 kg of grain and 30 kg of hay (17). This much forage and grain requires about 100 000 L of water to produce the 100 kg of hay, and 5400 L for the 4 kg of grain. On rangeland for forage production, more than 200 000 L of water are needed to produce 1 kg of beef (30). Animals vary in the amounts of water required for their production. In contrast to beef, 1 kg of broiler can be produced with about 2.3 kg of grain requiring approximately 3500 L of water.

## CONCLUSION

Both the meat-based average American diet and the lactoovovegetarian diet require significant quantities of nonrenewable fossil energy to produce. Thus, both food systems are not sustainable in the long term based on heavy fossil energy requirements. However, the meat-based diet requires more energy, land, and water resources than the lactoovovegetarian diet. In this

Motif Exhibit 1036, Page 7 of 8

Case No.: IPR2023-00208
U.S. Patent No. 11,013,250



limited sense, the lactoovovegetarian diet is more sustainable than the average American meat-based diet.

The major threat to future survival and to US natural resources is rapid population growth. The US population of 285 million is projected to double to 570 million in the next 70 y, which will place greater stress on the already-limited supply of energy, land, and water resources. These vital resources will have to be divided among ever greater numbers of people.

## REFERENCES

1. World Health Organization. Micronutrient malnutrition—half of the world's population affected. World Health Organization 1996;78:1–4.
2. World Health Organization. Malnutrition worldwide. 2000. Internet: http://www.who.int/nut/malnutrition_worldwide.htm (accessed 27 July 2000).
3. Pimentel D, Pimentel M. World population, food, natural resources, and survival. World Futures 2003;59:145–67.
4. US Bureau of the Census. Statistical abstract of the United States. Washington, DC: Government Printing Office, 2001.
5. American Heart Association. American Heart Association: home page. 2001. Internet: http://www.americanheart.org (accessed 22 December 2001).
6. FAOSTAT. Food balance sheets. Internet: http://armanncorn:98ivysub@faostat.fao.org/lim…ap.pl?FoodBalanceSheet&Domain (accessed 22 December 2001).
7. Pimentel D. Livestock production and energy use. In: Cleveland CJ, ed. Encyclopedia of energy (in press).
8. Pimentel D, Pimentel M. Food, energy and society. Niwot, CO: Colorado University Press, 1996.
9. Pimentel D. Livestock production: energy inputs and the environment. In: Scott SL, Zhao X, eds. Canadian Society of Animal Science, proceedings. Vol 47. Montreal, Canada: Canadian Society of Animal Science, 1997:17–26.
10. Pimentel D, Doughty R, Carothers C, Lamberson S, Bora N, Lee K. Energy use in developing and developed crop production. In: Lal R, Hansen D, Uphoff N, Slack S, eds. Food security and environmental quality in the developing world. Boca Raton, FL: CRC Press, 2002:129–51.
11. US Department of Agriculture. Agricultural statistics. Washington, DC: US Department of Agriculture, 2001.
12. Centers for Disease Control and Prevention. Obesity and overweight. 2002. Internet: http://www.gov/nccdphp/dnpa/obesity/index.htm (accessed 22 January 2002).
13. Surgeon General. The virtual office of the Surgeon General. 2002. Internet: http://www.google.com/search?q=cache:oQexukpqAwC:www.surgeongeneral.gov/ (accessed 22 January 2002).
14. Morrison FB. Feeds and feeding. Ithaca, NY: Morrison Publishing Company, 1956.
15. Heitschmidt RK, Short RE, Grings EE. Ecosystems, sustainability, and animal agriculture. J Anim Sci 1996;74:1395–405.
16. US Department of Agriculture, National Agricultural Statistics Service. Agricultural prices, 1998 summary. Washington, DC: US Department of Agriculture, 1999.
17. Pimentel D. Handbook of energy utilization in agriculture. Boca Raton, FL: CRC Press, 1980.
18. Hoffman TR, Warnock WD, Hinman HR. Crop enterprise budgets, Timothy-legume and alfalfa hay, Sudan grass, sweet corn and spring wheat under rill irrigation. Farm Business Reports EB 1173, Kittitas County, Washington. Pullman, WA: Washington State University, 1994.
19. National Agricultural Statistics Service. Farm labor. Internet: http//usda.mannlib.cornell.edu. (accessed 22 December 1999).
20. US Department of Agriculture, Economic Research Service, Economics and Statistics System. Corn-state: costs of production. Washington, DC: US Department of Agriculture, 1991. (Stock #94018.)
21. Hinman H, Pelter G, Kulp E, Sorensen E, Ford W. Enterprise budgets for fall potatoes, winter wheat, dry beans, and seed peas under rill irrigation. Farm Business Management Reports. Pullman, WA: Washington State University, 1992.
22. US Department of Agriculture. National Agricultural Statistics Service. Washington, DC: US Department of Agriculture, Economic Research Service, 1997.
23. Food and Agricultural Organization. Agricultural statistics. 1999. Internet: http://apps.fao.org/cgi-bin/nph-db.pl?subset-agriculture (accessed 22 November 1999).
24. US Department of Agriculture. Farm business briefing room, 1998. Washington, DC: US Department of Agriculture, 1998.
25. McGuckin JT, Gollehon N, Ghosh S. Water conservation in irrigated agriculture: a stochastic production frontier model. Water Resour Res 1992;28:305–12.
26. US Bureau of the Census. Statistical abstract of the United States, 2000. Washington, DC: Government Printing Office, 1998.
27. US Department of Agriculture. Agricultural statistics. Washington, DC: US Department of Agriculture, 1998.
28. Pimentel D, Kounang N. Ecology of soil erosion in ecosystems. Ecosystems 1998;1:416–26.
29. Pimentel D, Houser J, Preiss E, et al. Water resources: agriculture, the environment, and Society. BioScience 1997;47:97–106.
30. Thomas GW. Water: critical and evasive resource on semi-arid lands. In: Jordan WR, ed. Water and water policy in world food supplies. College Station, TX: Texas A&M University Press, 1987:83–90.

Motif Exhibit 1036, Page 8 of 8