

AUGUST 1995
VOLUME 43
NUMBER 8

Journal of Agricultural and Food Chemistry

RECEIVED
SEP - 8 1995



Published by the
American Chemical Society

Associate Editors: G. Wayne Ivie
Marshall Phillips

Manuscript Office: Marie Crunkleton
Wiggins, Manager; Bryan D. Tweedy,
Senior Associate Editor; Sharon A.
Dixon, Editorial Assistant

ADVISORY BOARD

Terry E. Acree	Steven Nagy
Eric Block	R. Petroski
John W. Finley	Russ Rousseff
Paul Hedin	Richard Scanlan
Chi-Tang Ho	Allan E. Smith
R. McGorin	Walter Wolf

Journal of Agricultural and Food Chemistry (ISSN 0021-8561) is published monthly by the American Chemical Society at 1155 16th St., N.W., Washington, DC 20036. Second-class postage paid at Washington, DC, and additional mailing offices. POSTMASTER: Send address changes to *Journal of Agricultural and Food Chemistry*, Member & Subscriber Services, P.O. Box 3337, Columbus, OH 43210. Canadian GST Reg. No. 127571347. Printed in the USA.

PUBLICATIONS DIVISION

Robert H. Marks, Director

Journal Publishing Operations: Charles R. Bertsch, Director

Editorial Office: Mary E. Scanlan, Manager; Anne C. O'Melia, Journals Editing Manager; Eileen M. Richardson, Associate Editor

Advertising Office: Centcom, Ltd., 1599 Post Road East, P.O. Box 231, Westport, CT 06881.

© Copyright 1995 American Chemical Society. **Copyright permission:** Reprographic copying beyond that permitted by Section 107 or 108 of the U.S. Copyright Act is allowed, provided that the appropriate per-copy fee is paid directly to the Copyright Clearance Center, Inc., 222 Rosewood Dr., Danvers, MA 01923. Anyone may excerpt up to 200 words per article for noncommercial, scholarly purposes with appropriate credit to the source. Direct other reprint permission requests to ACS Copyright Office, Publications Division, at the ACS Washington address. For quotes and information on ordering bulk reprints, please call (202) 872-4539 or write to the Product Services Department at the ACS Washington address.

Editorial Information

Guide for Authors and copyright transfer form appear in the first issue of each volume.

Manuscripts for publication should be submitted (six copies) to Manuscript Office, *Journal of Agricultural and Food Chemistry*, at the ACS Washington address.

American Chemical Society
1155 16th St., N.W.
Washington, DC 20036
(202) 872-4600
TDD (202) 872-8733
FAX (202) 872-4615

Correspondence regarding **accepted papers and proofs** should be directed to Journal Publishing Operations, at the address below.

Page charges of \$40.00 per page may be paid for papers published in this journal. Payment does not affect acceptance or scheduling of papers.

The American Chemical Society and its Editors assume no responsibility for the statements and opinions advanced by contributors.

Registered names and trademarks, etc., used in this publication, even without specific indication thereof, are not to be considered unprotected by law.

A document number appears after the bibliographic information for each paper. It is a nine character code unique to each manuscript and is assigned during editorial production. It serves as a link between the documents in the printed product and the electronic products (CJACS Plus Images and journals on CD-ROM) and makes it easy to retrieve the full text of the exact paper.

1995 Subscription and Ordering Information

Members may share/donate their personal subscriptions with/to libraries and the like but only after 5 years from publication.

	U.S.	Canada and Mexico	Europe*	All Other Countries*
Printed				
Members—1 yr	\$ 35	\$ 54	\$ 77	\$ 91
Nonmembers	\$420	\$439	\$462	\$476
Microfiche				
Members—1 yr	\$ 35	\$ 35	\$ 46	\$ 46
Nonmembers	\$420	\$420	\$431	\$431
Supporting information (microfiche)	\$ 15	\$ 21	\$ 21	\$ 21

* Air service included.

Nonmember subscribers in Japan must enter subscription orders with Maruzen Company Ltd., 3-10 Nihonbashi 2-chome, Chuo-ku, Tokyo 103, Japan. Tel: (03) 272-7211.

Microform and Back Issue orders should be sent to the Microforms & Back Issues Office at the ACS Washington address.

New and renewal subscriptions should be sent with payment to American Chemical Society, Department L-0011, Columbus, OH 43268-0011. Subscriptions are available only on a calendar year basis.

Journal Publishing Operations
American Chemical Society
2540 Olentangy River Road
P.O. Box 3330
Columbus, OH 43210
(614) 447-3600, Ext. 3171
TELEX 6842086; FAX (614) 447-3745

Journal of Agricultural and Food Chemistry

Claims for missing numbers will not be allowed if loss was due to failure of notice of change of address to be received in the time specified; if claim is dated (a) North America—more than 90 days beyond issue date, (b) all other foreign—more than 180 days beyond issue date. Claims are handled by Member & Subscriber Services.

Supporting Information is noted in the table of contents with a ■. It is available as photocopy (\$12.00 for up to 3 pages and \$1.50 per page for additional pages, plus \$2.00 for foreign postage) or as 24× microfiche (\$12.00, plus \$1.00 for foreign postage). Canadian residents should add 7% GST. See Supporting Information notice at the end of journal article for number of pages. Orders *must* give complete title of article, names of authors, journal, issue date, and page numbers. Prepayment is required and prices are subject to change. In 1995, electronic Supporting Information is available to current journal subscribers via the Internet using either gopher or World Wide Web protocols. Most often, the material is available as PDF files, which may be viewed using Adobe's Acrobat Reader, a program which is freely available on the Internet. However, some articles may include computer programs, PostScript files, word processing files, experimental data in a standard format (e.g., crystallographic parameters in CIF format), etc. In order to download the Supporting Information, users will need to enter their journal subscriber number, which can be found on the mailing label. Detailed instructions for using this service can be found on the Internet. With gopher, connect to pubs.acs.org, go to the "ACS Publications" selection, then to the "Supporting Information" selection. Read the README file in this directory for detailed instructions. When using a WWW client (e.g., Mosaic, Netscape), connect to the URL "<http://pubs.acs.org/>", select the "Supporting Info. for Journals" link. For further information on electronic access, send electronic mail to gopher@acsinfo.acs.org, or phone (202) 872-4434. For information on microforms, contact Microforms & Back Issues at the ACS Washington address or phone (202) 872-4554. Supporting information except structure factors also appears in the microfiche edition.

Member & Subscriber Services
American Chemical Society
P.O. Box 3337
Columbus, OH 43210
(614) 447-3776; FAX (614) 447-3671
(800) 333-9511

Journal of Agricultural and Food Chemistry®

JAFCAU 43(8) 1963-2316 (1995)
ISSN 0021-8561

Registered in U.S. Patent and Trademark Office
Copyright 1995 by the American Chemical Society

AUGUST 1995

STEENBOCK LIBRARY
SEP - 8 1995

SHORT COMMUNICATIONS

- 1963 Analytical Method for Quantifying Butyric Acid, Malathion, and Diazinon in Recycled Poly(ethylene terephthalate)
Vanee Komolprasert,* Anna R. Lawson, and Walter A. Hargraves
- 1966 Trace Determination of Naringenin and Hesperitin by Tandem Mass Spectrometry
■ **Randy A. Weintraub,* Barbara Ameer, Jodie V. Johnson, and Richard A. Yost**

FOOD CHEMISTRY/BIOCHEMISTRY

- 1969 Discovery and Synthesis of a New Series of High-Potency
L-Aspartyl-D- α -aminoalkanoyl-(S)- α -alkylbenzylamide Sweeteners
James G. Sweeny,* Lihong L. D'Angelo, Edith A. Ricks, and Guillermo A. Iacobucci
- 1977 ϵ -(γ -Glutamyl)lysine: Hydrolysis by γ -Glutamyltransferase of Different Origins, When Free or Protein Bound
Katsuya Seguro,* Yoshiyuki Kumazawa, Tomoko Ohtsuka, Hiroyuki Ide, Noriki Nio, Masao Motoki, and Koji Kubota
- 1982 Production of Peptic Hemoglobin Hydrolysates: Bitterness Demonstration and Characterization
Isabelle Aubes-Dufau, Jean-Louis Seris, and Didier Combes*
- 1989 Chemical Transformation of Hydrolyzed Fumonisin B₁ to Hydrolyzed Fumonisin B₂
Farid A. Badria, Hamed K. Abbas, and W. Thomas Shier*
- 1993 Effects of Dissociation, Deamidation, and Reducing Treatment on Structural and Surface Active Properties of Soy Glycinin
Jorge R. Wagner and Jacques Guéguen*
- 2001 Partial Reduction of Soy Protein Isolate Disulfide Bonds
Silvana Petruccielli and María Cristina Añón*
- 2007 Functional Changes of Carboxymethyl Potato Starch by Conjugation with Whey Proteins
Makoto Hattori, Wenhong Yang, and Koji Takahashi*
- 2012 Quantitative and Qualitative Variations in Resin Content and Guayulins (A and B) among Different Guayule Cultivars
Om P. Sidhu, Neelima Ratti, and Hari M. Behl*
- 2016 Phenolic Acids in Flaxseed
B. Dave Oomah,* Edward O. Kenaschuk, and Giuseppe Mazza
- 2020 Studies on the Constituents of *Chenopodium pallidicaule* (Cañihua) Seeds. Isolation and Characterization of Two New Flavonol Glycosides
Luca Rastrelli, Paola Saturnino, Oreste Schettino, and Antonio Dini*

- 2025 Immunoreactivity of Goat's Milk Casein Fractionated by Ion-Exchange Chromatography
Ana I. Haza, Paloma Morales, Rosario Martín, Teresa García, Gonzalo Anguita, Isabel González, Bernabé Sanz, and Pablo E. Hernández*
- 2030 Effect of Benzylaminopurine on the Flavanones Hesperidin, Hesperetin 7-O-Glucoside, and Prunin in Tangelo Nova Fruits
J. A. Del Río,* M. D. Fuster, F. Sabater, I. Porras, A. García-Lidón, and A. Ortuño
- 2035 Purification and Characterization of a Novel Cysteiny-Specific Peptidase from Tainong 57 Sweetpotato Roots
Tsong-Rong Yan,* Chun-Yi Liao, and Yaw-Huei Lin
- 2040 Identification of Conjugated Triene Oxidation Products of α -Farnesene in Apple Skin
Daryl D. Rowan,* John M. Allen, Simon Fielder, Julie A. Spicer, and Margaret A. Brimble
- 2046 Multivariate Evaluation of Lipid Hydrolysis and Oxidation Data from Light and Dark Muscle of Frozen Stored Rainbow Trout (*Oncorhynchus mykiss*)
Torbjörn Ingemansson, Peter Kaufmann, and Bo Ekstrand*

COMPOSITION OF FOODS/FEEDS

- 2053 Characterization of Citrus Honey (*Citrus* Spp.) Produced in Spain
J. Serra Bonvehí* and F. Ventura Coll
- 2058 Changes in Ascorbic Acid Content of Green Asparagus during the Harvesting Period and Storage
María J. Esteve, Rosaura Farré,* Ana Frigola, and Gonzalo Clemente
- 2062 Quality Characteristics of Yellow-Seeded *Brassica* Seed Meals: Protein, Carbohydrates, and Dietary Fiber Components
Joseph Simbaya, Bogdan A. Slominski,* Gerhard Rakow, Lloyd D. Campbell, R. Keith Downey, and J. Milton Bell
- 2067 Effect of Soil Temperature on the Seed Composition of Three Spanish Cultivars of Groundnut (*Arachis hypogaea* L.)
Sabine D. Golombek,* Rudravarapu Sridhar, and Umaid Singh
- 2071 Chemical and Biochemical Aspects of Slaughterhouse Sludge Intended for Feed Purposes
Nicoline G. Fransen,* Bert A. P. Urlings, Ben G. F. Kessels, and Peter G. H. Bijker

ANALYTICAL METHODS

- 2077 Optimizing Nitrogen-Phosphorus Detector Gas Chromatography for Pesticide Analysis
William M. Draper
- 2083 Development of an Enzyme-Linked Immunosorbent Assay for the Detection of the Triazole Fungicide Myclobutanil
András Székács and Bruce D. Hammock*
- 2092 Improved Methods for Conjugating Selected Mycotoxins to Carrier Proteins and Dextran for Immunoassays
H. Xiao, J. R. Clarke, R. R. Marquardt,* and A. A. Frohlich
- 2098 Synthesis and Rearrangement Reactions of Ester-Linked Lignin-Carbohydrate Model Compounds
Kaichang Li and Richard F. Helm*
- 2104 HPLC/MS Application to Anthocyanins of *Vitis vinifera* L.
Alessandro Baldi, Annalisa Romani, Nadia Mulinacci, Franco F. Vincieri,* and Bruno Casetta
- 2110 Determination of Tebuthiuron and Metabolites in Bovine Milk by GC-MSD
Bonnie S. Rutherford,* Darcy D. Shackelford, and Jerry J. Ulmer
- 2117 Development and Validation of a Liquid Chromatographic Method for the Determination of Furosemide, a Diuretic, in Bovine Milk
Badar Shaikh*

- 2122 Determination of Spectinomycin Residues in Bovine Milk Using Liquid Chromatography with Electrochemical Detection
Patricia G. Schermerhorn,* Pak-Sin Chu, and Philip James Kijak
- 2126 Determination of Ethylene Oxide in Spices Using Headspace Gas Chromatography
James E. Woodrow, Michael M. McChesney, and James N. Seiber*
- 2130 Retention of Radiolytic CO Gas in Irradiated Pepper Grains and Irradiation Detection of Spices and Dry Grains with the Level of Stocked CO Gas
Masakazu Furuta,* Takaaki Dohmaru, Tadashi Katayama, Hirokazu Toratani, and Atsuhiko Takeda
- 2134 Method for the Production and Characterization of Tomato Cutin Oligomers
Stanley F. Osman,* Herve C. Gerard, William F. Fett, Robert A. Moreau, and Robert L. Dudley
- 2138 Headspace Solid-Phase Microextraction Analysis of Volatile Organic Sulfur Compounds in Black and White Truffle Aroma
Fabio Pelusio, Torben Nilsson,* Luca Montanarella, Roberto Tilio, Bo Larsen, Sergio Facchetti, and Jørgen Ø. Madsen

NUTRITION/TOXICOLOGY

- 2144 Efficacy of β -Ionone in the Chemoprevention of Rat Mammary Carcinogenesis
Suzanne G. Yu, Peter J. Anderson, and Charles E. Elson*
- 2148 Resistance of Sorghum α -, β -, and γ -Kafirins to Pepsin Digestion
Maria P. Oria, Bruce R. Hamaker,* and Jeannette M. Shull
- 2154 Isolation and Characterization of Vasoactive Lipids from the Needles of *Pinus ponderosa*
M. S. Al-Mahmoud, S. P. Ford, R. E. Short, D. B. Farley, Lane Christenson, and J. P. N. Rosazza*
- 2162 Nutritional Assessment of *Vernonia amygdalina* Leaves in Growing Mice
G. O. Igile, W. Oleszek,* S. Burda, and M. Jurzysta
- 2167 Levels of the Cholesterol-Elevating Diterpenes Cafestol and Kahweol in Various Coffee Brews
Rob Urgert, Guido van der Weg, Truus G. Kosmeijer-Schuil, Peter van de Bovenkamp, Robert Hovenier, and Martijn B. Katan*

FLAVORS AND AROMAS

- 2173 Biosynthesis of 13-Hydroperoxylinoleate, 10-Oxo-8-decenoic Acid, and 1-Octen-3-ol from Linoleic Acid by a Mycelial-Pellet Homogenate of *Pleurotus pulmonarius*
Smadar Assaf, Yizhak Hadar, and Carlos G. Dosoretz*
- 2179 Simulation of Retronasal Aroma Using a Modified Headspace Technique: Investigating the Effects of Saliva, Temperature, Shearing, and Oil on Flavor Release
Deborah D. Roberts and Terry E. Acree*
- 2187 Evaluation of the Key Odorants in a Thermally Treated Solution of Ribose and Cysteine by Aroma Extract Dilution Techniques
Thomas Hofmann and Peter Schieberle*
- 2195 Determination of the Chemical Structure of the Intense Roasty, Popcorn-like Odorant 5-Acetyl-2,3-dihydro-1,4-thiazine
Thomas Hofmann, Rainer Hässner, and Peter Schieberle*
- 2199 Model Reactions on the Stability of Disulfides in Heated Foods
Helmut Guth, Thomas Hofmann, Peter Schieberle, and Werner Grosch*
- 2204 Nonequilibrium Partition Model for Predicting Flavor Retention in Microwave and Convection Heated Foods
Kris B. de Roos* and Ernst Graf
- 2212 Volatile Chemicals Formed in the Headpace of a Heated D-Glucose/L-Cysteine Maillard Model System
Katsumi Umano, Yukio Hagi, Kazuaki Nakahara, Akihiro Shyoji, and Takayuki Shibamoto*

CHEMICAL CHANGES DURING PROCESSING/STORAGE

- 2219 Fourier-Transform Infrared and Raman Spectroscopic Study of Biochemical and Chemical Treatments of Oak Wood (*Quercus rubra*) and Barley (*Hordeum vulgare*) Straw
D. Stewart,* H. M. Wilson, P. J. Hendra, and I. M. Morrison
- 2226 Model Reactions on Roast Aroma Formation. 15. Investigations on the Formation of Pyrido[3,4-*d*]imidazoles during the Maillard Reaction
Un-Sook Gi and Werner Baltes*
- 2231 Evolution of Trypsin Inhibitor Activity during Germination of Lentils
Juana Frias, Concepcion Diaz-Pollan, Cliff L. Hedley, and Concepcion Vidal-Valverde*
- 2235 Protein Hydrolysis during Soybean Tempe Fermentation with *Rhizopus oligosporus*
Johan C. de Reu,* R. Marcel ten Wolde, Jolan de Groot, M. J. Robert Nout, Frank M. Rombouts, and Harry Gruppen
- 2240 Changes in Texture and Cell Wall Polysaccharides of Olive Fruit during "Spanish Green Olive" Processing
Ana Jiménez, Rafael Guillén, Coral Sánchez, Juan Fernández-Bolaños, and Antonia Heredia*
- 2247 Pyrolysis of Cherry Red Tobacco and 1-Deoxy-1-[(*S*)-2-(3-pyridyl)-1-pyrrolidinyl]- β -D-fructose (Pyranose and Furanose Isomers) Amadori Products of Cherry Red Tobacco
W. W. Weeks,* Marcos P. Campos, and Serban Moldoveanu
- 2254 Stability of Polyunsaturated Fatty Acids in Egg Powder Processed and Stored under Various Conditions
Francesc Guardiola, Rafael Codony,* Albert Manich, Magda Rafecas, and Josep Boatella

CHEMISTRY OF CROP AND ANIMAL PROTECTION

- 2260 Fungicidal Activity of Phenyl *N*-(4-Substituted-phenyl)thionocarbamates
Martha Albores-Velasco,* John Thorne, and Ralph L. Wain
- 2262 Chemical Composition and Fungitoxic Properties to Phytopathogenic Fungi of Essential Oils of Selected Aromatic Plants Growing Wild in Turkey
Frank Müller-Riebau,* Bernhard Berger, and Oktay Yegen
- 2267 Diversity of Sesquiterpenes in 46 Potato Cultivars and Breeding Selections
Anne E. Desjardins,* Susan P. McCormick, and Dennis L. Corsini
- 2273 Polyphenolic Pattern in Apple Tree Leaves in Relation to Scab Resistance. A Preliminary Study
Anna Picinelli,* Enrique Dapena, and Juan José Mangas
- 2279 Persistence of Pirimicarb in Peaches and Nectarines
Paolo Cabras,* Marinella Melis, Franco Cabitza, Mario Cubeddu, and Lorenzo Spanedda
- 2283 Prangolarin Is a Chemical Facilitator for the Enhanced Development of the Infection Process in the Epicarp of *Citrus limon* by *Penicillium digitatum*
Yutaka Arimoto, Fumio Sugawara,* Shigeo Yoshida, and Isamu Yamaguchi
- 2286 Synthesis and Biological Evaluation of New Pyrethroids Having Halogen, Keto, or Nitro Group-Containing Substituents
Ming-Kuan Chyan and Scott J. Norton*
- 2291 Mineralization of the Herbicide Atrazine in Soil Inoculated with a *Pseudomonas* Strain
Clotaire Yanze Kontchou and Nikolaus Gschwind*

ENVIRONMENTAL CHEMISTRY

- 2295 Clomazone Sorption in Soil: Incubation Time, Temperature, and Soil Moisture Effects
Todd L. Mervosh, Gerald K. Sims,* Edward W. Stoller, and Timothy R. Ellsworth

BIOTECHNOLOGY

- 2301 Development of Monoclonal Antibodies for Red Snapper (*Lutjanus campechanus*) Identification Using Enzyme-Linked Immunosorbent Assay
Tung-shi Huang, Maurice R. Marshall, Kuo-jang Kao, W. Steven Otwell, and Cheng-i Wei*
- 2308 Fiber Optic Immunosensor for Polychlorinated Biphenyls
Christopher Q. Zhao, Nabil A. Anis, Kim R. Rogers, Richard H. Kline, Jr., Jeremy Wright, Amira T. Eldefrawi, and Mohyee E. Eldefrawi*

CORRECTION

- 2316 Effect of Coatings and Prolonged Storage Conditions on Fresh Orange Flavor Volatiles, Degrees Brix, and Ascorbic Acid Levels
Elizabeth A. Baldwin,* Myrna Nisperos-Carriedo, Philip E. Shaw, and Jacqueline K. Burns

■ Supporting information for this paper is available separately (consult a current masthead page for ordering information). Supporting information is available to subscribers electronically via the Internet at <http://pubs.acs.org> (WWW) and pubs.acs.org (Gopher). All supporting information except for structure factor tables will also appear following the paper in the microfilm edition of the journal.

* In papers with more than one author, the asterisk indicates the name of the author to whom inquiries about the paper should be addressed.

Evaluation of the Key Odorants in a Thermally Treated Solution of Ribose and Cysteine by Aroma Extract Dilution Techniques

Thomas Hofmann and Peter Schieberle*

Deutsche Forschungsanstalt für Lebensmittelchemie and Institut für Lebensmittelchemie der Technischen Universität München, Lichtenbergstrasse 4, D-85748 Garching, Germany

Application of the aroma extract dilution analysis on a solvent extract isolated from a thermally treated solution (145 °C; 20 min) of cysteine/ribose led to the identification of 2-furfurylthiol, 3-mercapto-2-pentanone, 2-methyl-3-furanthiol, 5-acetyl-2,3-dihydro-1,4-thiazine, 3-mercapto-2-butanone, and bis(2-methyl-3-furyl) disulfide showing the highest flavor dilution factors among the 29 odor-active volatiles. HRGC/olfactometry of decreasing headspace volumes established especially 2-furfurylthiol and 2-methyl-3-furanthiol as important odorants and revealed 2-thenyl mercaptan and ethyl mercaptan as further key contributors to the overall roasty, meatlike, sulfury odor of the model mixture. 5-Acetyl-2,3-dihydro-1,4-thiazine, identified for the first time among the volatiles of Maillard model reactions or foods, exhibited an intense roasty, popcorn-like odor at the low odor threshold of 0.06 ng/L of air, which was of the same order of magnitude as those reported in the literature for the roasty-smelling odorants 2-acetyl-1-pyrroline and 2-acetyl-2-thiazoline.

Keywords: *Aroma extract dilution analysis; nonenzymatic browning; Maillard reaction; ribose; cysteine; flavor; 5-acetyl-2,3-dihydro-1,4-thiazine*

INTRODUCTION

The Maillard reaction between reducing sugars and amino acids is known to generate flavors similar to those of certain processed foods [cf. Lane and Nursten (1983)] and, in particular, thermally treated model mixtures containing cysteine elicit odors resembling meat flavor (Morton et al., 1960). Therefore, precursor mixtures used in the industrial production of meat flavorings (reaction flavors, processed flavors) usually contain cysteine (van den Ouweland et al., 1978; MacLeod and Seyyedain-Ardebili, 1981; MacLeod, 1986).

In recent studies on the volatile fractions of model systems containing cysteine, an increasing number of reaction products have been identified (Martin, 1988; Whitfield et al., 1988; Tressl et al., 1989; Farmer et al., 1989; Güntert et al., 1990; Werkhoff et al., 1991; Silwar, 1992), giving insights into the complex reaction pathways governing the degradation of cysteine in the presence of carbohydrates, phospholipids, or thiamin. Sulfur-containing compounds especially predominate and were suggested as important odorants, e.g., in a reaction system containing thiamin and cysteine (Güntert et al., 1993).

Although very odor-active compounds have been identified, until now, a correlation between the overall odor of a reaction mixture containing exclusively cysteine and ribose and the compounds responsible for the flavor notes has not been established. However, a systematic evaluation of the odorants rendering the typical odor to such Maillard reaction mixtures would be helpful, e.g., in optimizing the processing conditions in the manufacturing of reaction flavors.

Application of aroma extract dilution techniques allows the detection and evaluation of key odorants in food flavor extracts [cf. reviews by Acree (1993) and Grosch (1993)], and recently such techniques have also been successfully applied to extracts obtained from

thermally treated solutions of proline/2-oxopropanal (Schieberle, 1990), proline/glucose (Roberts and Acree, 1994), and thiamin/cysteine (Güntert et al., 1993).

The purpose of this investigation was, therefore, to elucidate the most odor-active compounds in a thermally treated cysteine/ribose mixture by application of aroma extract dilution techniques.

EXPERIMENTAL PROCEDURES

Chemicals. 1-Chloro-2-propanone, 2-(hydroxymethyl)thiophene (2-thienylmethanol), methyl mercaptan, and ethyl mercaptan were from Aldrich (Steinheim, Germany). 2,3,5,6-Tetramethyl-1,4-dioxane-2,5-diol was from Lancaster (Mühlheim, Germany). L-Cysteine and D-(–)-ribose were from Sigma (Munich, Germany). Compounds **2**, **5**, **8**, **9**, **12**, **15**, **17**, **21**, **23**, **25**, and **30** (Table 1) were also from Aldrich, and **6** was from Lancaster.

Syntheses. *2-Methyl-3-thiophenethiol.* To a stirred solution of diisopropylamide (10 mmol) in dry tetrahydrofuran (10 mL) maintained in an atmosphere of pure argon was added dropwise *n*-butyl lithium (10 mmol) in *n*-hexane (2.5 mL) at –78 °C within 10 min. After the addition of 3-bromothiophene (10 mmol); the mixture was stirred for a further 180 min at –78 °C and then methyl iodide (10.5 mmol), dissolved in dry tetrahydrofuran, was dropwise added. After stirring for a further 30 min, the mixture was allowed to react at room temperature for 16 h. After the addition of water (20 mL), the 3-bromo-2-methylthiophene formed (yield 69%) was extracted with diethyl ether (total volume 90 mL) and the solution dried over Na₂SO₄. After evaporation of the solvent, the crude 3-bromo-2-methylthiophene (6.9 mmol) was dissolved in dry diethyl ether (10 mL), which was stirred at –78 °C in an argon atmosphere. After the dropwise addition of *n*-butyl lithium (7.3 mmol in *n*-hexane), the mixture was allowed to react for a further 20 min and, after the addition of powdered sulfur (6.9 mmol), reacted for 60 min with rising temperature to –10 °C. The solution was then poured onto ice (30 g) and, after the pH was adjusted to 3.0, the target compound was extracted with diethyl ether (total volume 100 mL). The 2-methyl-3-thiophenethiol was purified by microdistillation (yield 58%) and characterized by its mass spectrum.

MS/EI: *m/z* (%) 130 (100), 97 (72), 45 (68), 129 (55), 69 (34), 59 (32), 85 (20). The data agreed with those previously reported by van den Ouweland and Peer (1975).

* Author to whom correspondence should be addressed [telephone (089) 3209-4170; fax (089) 3209-4183].




Figure 1. Mass spectrum of 2-thenyl mercaptan.

2-Thenyl Mercaptan. 2-(Hydroxymethyl)thiophene (3 mmol) was added to a solution of thiourea (4 mmol) in hydrochloric acid (4 mL; 0.3% by weight), which was maintained under argon atmosphere. After stirring for 4 h at room temperature, aqueous sodium hydroxide (3 mL; 5% by weight) followed by diethyl ether (10 mL) was added and the mixture stirred in argon atmosphere for a further 30 min. The organic layer was discarded and the aqueous layer extracted twice with diethyl ether (total volume 50 mL) to remove the unreacted (hydroxymethyl)thiophene and byproducts. After cooling with ice water, the aqueous phase was acidified to pH 3.0 (hydrochloric acid, 10% by weight) and the target compound was extracted with diethyl ether (total volume 80 mL). After drying over Na_2SO_4 , the extract was concentrated to 40 mL by using a Vigreux column and then purified by sublimation in vacuo (yield 39%) using the apparatus reported earlier (Sen et al., 1991). The compound was characterized by its ^1H -NMR data in CD_2Cl_2 (assignment of the atoms bearing the H atoms refers to Figure 1): δ 2.08 (t, 1H, $J_{1-2} = 7.63$ Hz, H at 1), 3.98 (d, 2H, $J_{2-1} = 7.63$ Hz, H at C2), 6.93 (dd, 1H, $J_{3-4} = 3.27$ Hz, H at 3), 6.96 (dd, 1H, $J_{4-3} = 3.27$ Hz, $J_{4-5} = 1.09$ Hz, H at 4), 7.22 (dd, 1H, $J_{5-4} = 1.09$ Hz, H at C5) and its mass spectrum (Figure 1).

3-Mercapto-2-butanone. Acetoin (1 mmol) was liberated from its dimer 2,3,5,6-tetramethyl-1,4-dioxane-2,5-diol by weak acid treatment (SiO_2 in dichloromethane/water; 10 + 1 by vol) and subsequently dissolved in ethanol/water (6 mL; 5 + 1 by vol). The solution was saturated with H_2S at 0 °C and then stirred for 12 h at 80 °C in a closed vessel. After the addition of an aqueous sodium hydroxide solution (30 mL; 2 mol/L), the unreacted acetoin was extracted with diethyl ether (total volume 45 mL). The aqueous layer was then acidified to pH 3 with hydrochloric acid (1 mol/L), and the 3-mercapto-2-butanone was isolated by extraction with diethyl ether (three times; total volume 60 mL; yield 10%).

MS(EI): m/z (%) 43 (100), 61 (58), 104 (28), 60 (20), 72 (6), 71 (4), 106 (2). The spectrum was in good agreement with data published by Werkhoff et al. (1989).

1-Mercapto-2-propanone (Asinger et al., 1964). Under a constant stream of H_2S , 1-chloro-2-propanone (10 mmol) was added dropwise within 10 min to an ice-cooled stirred solution of sodium hydroxide (1.3 g) in water (6 mL), which had been presaturated with H_2S . After stirring for a further 60 min, an aqueous solution of sodium hydroxide was added (30 mL; 2 mol/L) and the excess of 1-chloro-2-propanone extracted with diethyl ether (45 mL). The aqueous phase was acidified to pH 3 with hydrochloric acid (1 mol/L), and the title compound (71% yield) was extracted with diethyl ether (total volume 60 mL) and characterized by mass spectrometry.

MS(EI): m/z (%) 43 (100), 90 (36), 47 (23), 45 (12), 73 (6), 58 (4), 71 (4), 92 (3).

Mixed Disulfides (20, 22, and 27–29 in Table 1). The mixed disulfides were synthesized according to the following general procedure: The corresponding thiols (0.1 mmol each) were dissolved in diethyl ether (15 mL) and, after the addition of an aqueous solution of copper(II) sulfate (3 mol/L; 15 mL), shaken for 35 min at room temperature. After complexation of the excess copper ions by the addition of ammonium acetate

(2 g), the ethereal layer was separated, washed three times with brine (total volume 60 mL), and then dried over Na_2SO_4 . The mixed disulfide was separated from the two bis-substituted disulfides formed in the reaction mixtures by HPLC on a RP-18 stationary phase as described for the isolation of bis-(2-mercaptoethyl)furan (Hofmann and Schieberle, unpublished results). The mass spectral data of the synthesized compounds are shown in Figure 2. The spectra reported for 2-methyl-3-furyl 2-oxo-3-butyl disulfide and 2-methyl-3-furyl 2-oxo-3-pentyl disulfide were in good agreement with those published recently by Ruther and Baltes (1994).

The following odorants were synthesized as reported in the literature: bis(2-methyl-3-furyl) disulfide (Evers et al., 1976); 3-mercapto-2-pentanone (Asinger et al., 1964); dimethyl trisulfide (Milligan et al., 1963); 5-acetyl-2,3-dihydro-1,4-thiazine and 5-propionyl-2,3-dihydro-1,4-thiazine (Hofmann and Schieberle, 1995); 2-acetyl-2-thiazoline (Cerny and Grosch, 1992, 1993); 3-hydroxy-2-pyranone [Wiley and Jarboe, 1956; mass spectra and ^1H -NMR data agreed with those published by Normington et al. (1986)]; and 4-hydroxy-5-methyl-3(2H)-furanone (Peer et al., 1968).

Model Reaction. L-Cysteine (3.3 mmol) and D-ribose (10 mmol) in phosphate buffer (100 mL, 0.5 mol/L, pH 5.0) were allowed to react in a laboratory autoclave (Model II; Roth, Karlsruhe, Germany) by raising the temperature within 20 min from 20 to 145 °C.

Isolation of the Volatile Reaction Products. After cooling, the dark yellow reaction mixture was extracted five times with diethyl ether (total volume 300 mL). The combined organic phases were treated three times with an aqueous sodium bicarbonate solution (0.5 mol/L; total volume 100 mL), and the ethereal solution containing the neutral/basic compounds (fraction NB) was washed twice with brine (total volume 100 mL) and then dried over Na_2SO_4 . The bicarbonate solution containing the acidic compounds was adjusted to pH 3.0 with hydrochloric acid (1 mol/L), and the acidic volatiles were then extracted with diethyl ether (total volume 150 mL; fraction AF). After washing with brine, the ethereal solution was dried over Na_2SO_4 . Each fraction was distilled from the nonvolatile material in the apparatus described recently (Sen et al., 1991).

Column Chromatography. Fraction NB was concentrated to 1 mL by distilling off the solvent on a Vigreux column (60 cm $\times$ 1 cm). The solution was placed on the top of a water-cooled (10–12 °C) column, designed for flash chromatography (20 cm $\times$ 2 cm i.d.; J. T. Baker BV, Deventer, The Netherlands, no. 7022-01) and equipped with a reservoir (250 mL) and a pressure regulator. The column was then filled with a slurry of silica gel (30–60 μm , Baker analyzed reagent) in *n*-pentane. Chromatography was performed under nitrogen pressure maintaining a constant flow rate of 5 mL/min and by using the following solvents: *n*-pentane (150 mL; NB I), *n*-pentane/diethyl ether (150 mL; 95 + 5 by vol; NB II), *n*-pentane/diethyl ether (150 mL; 8 + 2 by vol; NB III), *n*-pentane/diethyl ether (150 mL; 1 + 1 by vol; NB IV), diethyl ether/methanol (150 mL; 99.5 + 0.5 by vol; NB V), and finally diethyl ether/methanol (150 mL; 9 + 1 by vol; NB VI).

Enrichment of Thiols by Affinity Chromatography (Full and Schreier, 1994). Fraction AF also containing the thiols was concentrated to 1 mL and, after the addition of *n*-pentane (1 mL), placed on the top of a water-cooled (10–12 °C) glass column (5 cm $\times$ 1 cm i.d.) filled with a slurry of Affi-Gel 501 (Bio-Rad, Munich, Germany) in 2-propanol (30 mL). Nonbound compounds were first eluted with *n*-pentane/dichloromethane (60 mL; 2 + 1 by vol; AF I), and then the complexed compounds were isolated by elution with *n*-pentane/dichloromethane (30 mL; 2 + 1 by vol) containing 1,4-dithiothreitol (10 mmol; AF II). To separate the excess of the dithiol, the volatile material and the solvent were distilled off by using the apparatus described earlier (Sen et al., 1991).

High-Resolution Gas Chromatography (HRGC)/Mass Spectrometry (MS). HRGC was performed with a Type 5160 gas chromatograph (Fisons Instruments, Mainz, Germany) by using the following capillaries: FFAP (30 m $\times$ 0.32 mm fused silica capillary, free fatty acid phase, 0.25 μm ; J&W Scientific, Fisons Instruments, Mainz, Germany) and SE-54 (30 m $\times$ 0.32

Table 1. Intense Odorants (FD $\geq$ 4) Formed by Heating of an Aqueous Ribose/Cysteine Solution^a

no.	odorant ^b	fraction ^c	RI on		FD factor ^d		reported as volatile compound ^e
			FFAP	SE-54	I	II	
1	3-mercapto-2-butanone	AF II	1282	820	128	<1	5, 7
2	2-methyl-3-furanthiol	AF II	1300	869	256	<1	4, 6
3	3-mercapto-2-pentanone ^f	AF II	1347	907	512	<1	6, 7
4	1-mercapto-2-propanone	AF II	1357	<800	16	<1	5
5	2-furfurylthiol	AF II	1431	909	1024	<1	3-7
6	2-methyltetrahydrothiophen-3-one	AF I/NB III	1512	1017	8	16	3-7
7	2-methyl-3-thiophenethiol	AF II	1559	1059	16	<1	2, 6
8	2-acetylthiazole	NB III	1615	1142	<1	8	1, 3-6, 11
9	2-formylthiophene	NB III	1669	1000	<1	64	4-6
10	2-thenyl mercaptan	AF II	1682	1092	8	<1	4, 5, 12
11	2-acetyl-2-thiazoline	NB IV	1720	1111	<1	64	8
12	acetylthiophene	NB IV	1740	1090	<1	16	4-6, 11
13	unknown		1784		<1	16	
14	unknown		1800		<1	8	
15	2-(hydroxymethyl)thiophene	AF I/NB IV	1917	1030	2	<1	5
16	3-hydroxy-2H-pyran-2-one	AF II	1965	989	16	<1	5
17	4-hydroxy-2,5-dimethyl-3(2H)-furanone	AF II	2016	1100	32	<1	5, 11
18	bis(2-methyl-3-furyl) disulfide	AF I/NB I	2100	1526	128	16	7, 8, 10
19	4-hydroxy-5-methyl-3(2H)-furanone (norfuraneol)	AF II	2105	1044	64	<1	5, 6
20	2-methyl-3-furyl 2-oxo-3-butyl disulfide	AF I	2127	1491	4	<1	9
21	2-mercaptopropanoic acid	AF II	2130	1057	4	<1	5
22	2-methyl-3-furyl 2-oxo-3-pentyl disulfide	AF I	2145	1561	8	<1	9
23	3-hydroxy-4,5-dimethyl-2(5H)-furanone (sotolon) ^g	AF II	2153	1112	16	<1	
24	5-acetyl-2,3-dihydro-1,4-thiazine	NB VI	2177	1374	<1	256	
25	3-mercaptopropanoic acid	AF II	2193	1057	16	<1	5
26	5-propionyl-2,3-dihydro-1,4-thiazine	NB VI	2235	1456	<1	8	
27	2-furfuryl 2-methyl-3-furyl disulfide	AF I/NB I	2323	1624	16	4	10
28	2-furfuryl 2-oxo-3-butyl disulfide	AF I	2352	1577	4	<1	
29	2-furfuryl 2-oxo-3-pentyl disulfide	AF I	2385	1649	4	<1	
30	bis(2-furfuryl) disulfide	AF I/NB I	2465	1673	16	4	10

^a A mixture of ribose (10 mmol) and cysteine (3.3 mmol) was reacted for 20 min in phosphate buffer (100 mL; 0.5 mol/L; pH 5.0). ^b The compound was identified by comparing it with the reference substance on the basis of the following criteria: retention index (RI) on two capillary columns given in the table, mass spectra obtained by MS (EI) and MS (CI), and odor quality perceived at the sniffing port. ^c Fraction in which most of the compound appeared after chromatography on silica gel or Affi-Gel, respectively. ^d Flavor dilution (FD) factor determined in extracts containing the acidic (I) or the neutral/basic volatiles (II). Analyses were performed by two assessors in duplicates. The data differed to not more than two FD factors. ^e Reported in the literature as volatile reaction product of cysteine, cystine, or cysteamine reacted in the presence of carbohydrates or carbohydrate degradation products. (1) Mulders (1973); (2) van den Ouweland and Peer (1975); (3) Mussinan and Katz (1973); (4) Whitfield et al. (1988); (5) Martin (1988); (6) Farmer et al. (1989); (7) Güntert et al. (1990); (8) Sakaguchi and Shibamoto (1978); (9) Whitfield et al. (1993); (10) Farmer and Patterson (1991); (11) Silwar (1992); (12) Güntert et al. (1993). ^f 3-Mercapto-2-pentanone contained small amounts of its isomer 2-mercapto-3-pentanone. ^g The MS signals were too weak for an unequivocal interpretation. The compound was identified on the basis of the remaining criteria given in footnote b.

mm fused silica capillary DB-5; 0.25 μ m; J&W Scientific, Fisons Instruments). The samples were applied by the on-column injection technique at 40 °C. After 2 min, the temperature of the oven was raised at 40 °C/min to 50 °C (SE-54) or 60 °C (FFAP), respectively, held for 5 min isothermally, then raised at 6 °C/min to 230 °C, and held for 15 min. The flow of the carrier gas helium was 2.5 mL/min. At the end of the capillary, the effluent was split 1:1 (by vol) into an FID and a sniffing port using deactivated but uncoated fused silica capillaries (50 cm $\times$ 0.32 mm). The FID and the sniffing port were held at 180 °C. Linear retention indices (RI) of the compounds were calculated from the retention times of *n*-alkanes by using a computer program (Halang et al., 1978). MS analysis was performed with an MS-8230 (Finnigan, Bremen, Germany) in tandem with the capillaries described above. Mass spectra in the electron impact mode (MS/EI) were generated at 70 eV and in the chemical ionization mode (MS/CI) at 115 eV with isobutane as reactant gas.

Aroma Extract Dilution Analysis (AEDA); Approximation of Odor Threshold Values. The flavor dilution (FD) factors of the odor-active compounds were determined by AEDA (Ullrich and Grosch, 1987; Schieberle and Grosch, 1987) of the following dilution series: The original extract (100 μ L) of either fraction NB or fraction AF obtained from 100 mL of the model reaction was stepwise diluted with diethyl ether (1 + 1). HRGC/olfactometry was performed with aliquots (1.0 μ L) using capillary FFAP (fraction AF) and capillary DB-5 (fraction NB), respectively.

Static Headspace Analysis/Olfactometry (SHO). The reaction mixture (100 mL) was pipetted into a septum-sealed, thermostated vessel (240 mL total volume) and equilibrated for 1 h at 40 °C. Decreasing headspace volumes (20 to 0.02

mL) were taken off by means of a gastight syringe and then analyzed by SHO on the FFAP column as detailed by Guth and Grosch (1993). The perceived odors were attributed to the odorants identified in the preceding experiments using solvent extracts by means of odor quality, odor intensity, and retention index. The FD factors given in Table 2 were calculated by dividing the largest volume analyzed (20 mL; FD = 1 by definition) by the lowest volume in which the respective odorant was still detectable.

RESULTS

In preliminary experiments, first, the influence of the pH (5.0 and 7.0) and of the ribose/cysteine ratio (1:1 to 10:1) on the overall odor of the thermally treated reaction mixture was studied. The results of the sensory evaluation (data not shown) revealed that sulfury, meatlike odors were generally more prominent in reaction mixtures treated in phosphate buffer at pH 5.0, whereas at pH 7.0 caramel, burnt notes predominated. Furthermore, heating of several ribose (100 mmol/L)/cysteine mixtures at pH 5.0 with lowered concentrations of cysteine (100 to 10 mmol/L) showed significant differences in the overall odors, which were described as rubbery, sulfury, and H₂S-like in a 1:1 mixture and burnt and caramel-like in the 10:1 mixture. The most prominent roasty, meatlike odor was detectable in a mixture containing 100 mmol/L ribose and 33 mmol/L cysteine. Therefore, this reaction mixture was chosen for the evaluation by aroma extract dilution analyses.




Figure 2. Mass spectra of mixed disulfides: (A) 2-methyl-3-furyl 2-oxo-3-butyl disulfide; (B) 2-methyl-3-furyl 2-oxo-3-pentyl disulfide; (C) 2-furfuryl 2-methyl-3-furyl disulfide; (D) 2-furfuryl 2-oxo-3-butyl disulfide; (E) 2-furfuryl 2-oxo-3-pentyl disulfide.

Acidic Volatile Fraction. To avoid interferences with the gas chromatographic separations, the acidic volatiles from the reaction/mixture (fraction AF) were separated by treating the flavor extract with sodium bicarbonate. Application of the AEDA on fraction AF, eliciting a strong sulfury note, revealed 21 odor-active compounds in the FD factor range 4–1024 (column I in Table 1). On the basis of their high FD factors five compounds (1–3, 5, and 18), exhibiting sulfury, roasty, meatlike odors, were the main contributors to the overall odor of this fraction.

To get enough material for identification, 25 samples of the model reaction were combined (total volume 2.5 L); the acidic volatiles were isolated by treatment of the extract with sodium bicarbonate and then separated by affinity chromatography with phenylmercury chloride. This method has been recently developed (Full and Schreier, 1994) to enrich thiols from flavor extracts. By application of HRGC/olfactometry and HRGC/MS (Table 1; footnote *b*) to fraction AF II containing the compounds, which had been bound as mercury adducts, four of the five most important odorants in fraction AF could be identified (Table 1): **5**, 2-furfurylthiol (roasty, coffee-like); **3**, 3-mercapto-2-pentanone (sulfury, catty); **2**, 2-methyl-3-furanthiol (meatlike, sulfury); and **1**, 3-mer-

capto-2-butanone (sulfury, rotten). In addition, **19** [4-hydroxy-5-methyl-3(2*H*)-furanone (norfuranol; caramel-like)], **17** [4-hydroxy-2,5-dimethyl-3(2*H*)-furanone (HDF, caramel-like)], **23** [3-hydroxy-4,5-dimethyl-2(5*H*)-furanone (sotolon, seasoning-like)], **16** [3-hydroxy-2*H*-pyran-2-one (seasoning-like)], **7** [2-methyl-3-thiophenethiol (meatlike, sulfury)], **4** [1-mercapto-2-propanone (sulfury)], **10** [2-thenyl mercaptan (roasty)], and **21** and **25** [2- and 3-mercaptopropanoic acid (sulfury, seasoning-like)], contributing with lower FD factors to the overall odor, were identified in fraction AF II (Table 1). It is interesting to note that the affinity chromatography with phenylmercury chloride seems to be useful also in the enrichment of enoloxo compounds, such as norfuranol, HDF, 3-hydroxy-2*H*-pyran-2-one, or sotolon (**19**, **17**, **16**, and **23**; Table 1).

The fifth key odorant, **18**, was identified in fraction AF I as bis(2-methyl-3-furyl) disulfide.


Neutral Basic Fraction. Application of the AEDA on the neutral/basic volatile fraction (fraction NB) exhibiting an intense roasty, popcorn-like odor revealed 12 odorants (column II in Table 1) of which four compounds showing moderate FD factors [**6**, 2-methyl-tetrahydrothiophen-3-one; **18**, bis(2-methyl-3-furyl) disulfide; **27**, 2-furfuryl 2-methyl-3-furyl disulfide; and **30**,

Table 2. Results of Static Headspace Analysis/Olfactometry (SHO) of the Cysteine/Ribose Model Solution

odorant ^a	RI on FFAP	vol ^b (mL)	FD factor ^c
hydrogen sulfide ^d	<900	5	4
methyl mercaptan	<900	2.5	8
ethyl mercaptan	<900	0.16	125
2,3-butanedione (diacetyl)	968	20	1
3-mercapto-2-butanone	1282	2.5	8
2-methyl-3-furanthiol	1300	0.08	250
3-mercapto-2-pentanone ^d	1347	0.64	32
dimethyl trisulfide	1356	20	1
2-furfurylthiol	1431	0.02	1000
unknown (sulfury)	1438	10	2
unknown (sulfury)	1511	5	4
2-methyl-3-thiophenethiol	1559	0.64	32
2-thenyl mercaptan	1682	0.16	125
2-acetyl-2-thiazoline ^e	1720	20	1
5-acetyl-2,3-dihydro-1,4-thiazine ^e	2177	20	1

^a The compound was identified by comparing its mass spectra, retention index, odor quality, and odor activity (displayed by the odor threshold in air) with the reference compound (cf. Table 1).

^b Lowest headspace volume required to perceive the odorant at the sniffing port. ^c Calculated by dividing the largest volume analyzed (20 mL) by the lowest headspace volume required to perceive the odorant at the sniffing port. ^d 3-Mercapto-2-pentanone contained a smaller proportion of its isomer 2-mercapto-3-pentanone. ^e MS signals were too weak for an unequivocal interpretation.

**Figure 3.** Mass spectrum of peak 24: 5-acetyl-2,3-dihydro-1,4-thiazine.

bis(2-furfuryl) disulfide] had already been identified in the acidic fraction (column I, Table 1). Among the remaining eight compounds (8, 9, 11–14, 24, and 26) odorant 24, exhibiting an intense roasty, popcorn-like odor, showed the highest odor activity in the neutral/basic fraction.

After separation of the neutral/basic volatiles from 2.5 L of the reaction mixture by flash chromatography on silica gel, the mass spectrum shown in Figure 3 was obtained for compound 24 in fraction NB VI. As indicated in the figure, purification by preparative GLC and application of high-resolution mass spectrometry as well as various NMR techniques led to its structure being identified as 5-acetyl-2,3-dihydro-1,4-thiazine. The details of the NMR, MS, and synthetic experiments demonstrating the chemical identity of compound 24 will be published shortly (Hofmann et al., 1995a).

On the basis of their FD factors, compounds 9 and 11, also exhibiting roasty odors, followed next in rank to 5-acetyl-2,3-dihydro-1,4-thiazine. The identification experiments (Table 1, footnote b) revealed that 2-formylthiophene (9) and 2-acetyl-2-thiazoline (11) evoked the roasty odors detected by AEDA in the corresponding odor-active regions.

Compound 26, contributing with a lower FD factor to the odor of the neutral/basic fraction, also exhibited a popcorn-like, roasty odor. On the basis of mass spectral data [MS/EI m/z (%) 157 (100), 100 (64), 72 (42), 45 (40), 57 (34), 101 (22)] and NMR experiments followed by synthesis, compound 26 was identified as 5-propionyl-2,3-dihydro-1,4-thiazine. The analytical details will be reported shortly (Hofmann et al., 1995a).

Eight disulfides (18, 20, 22, and 27–30), known as oxidation products of the corresponding thiols (Farmer and Patterson, 1991; Whitfield et al., 1993), were additionally identified which, except for bis(2-methyl-3-furyl) disulfide (18; Table 1), contributed with low FD factors to the overall odor of the reaction mixture. Unexpectedly, the FD factors and, therefore, the concentrations of these neutral compounds were higher in the acidic than in the neutral/basic fraction (cf. columns I and II, Table 1), suggesting that the disulfides were formed as artifacts during the workup procedure. This assumption was very recently corroborated by our findings (Hofmann et al., 1995b) that after 10 days storage of, e.g., 2-methyl-3-furanthiol in diethyl ether at 6 °C, more than 50% of the thiol had been converted into the corresponding disulfide.

Static Headspace Analysis/Olfactometry (SHO).

The results of the static headspace analysis are summarized in Table 2. In a 20 mL sample of air above the model solution, 15 odor-active compounds were detectable by GC/Olfactometry, of which 12 compounds could be identified (Table 2). Eight of these odorants had already been identified in the AEDA of the solvent extract (1–3, 5, 7, 10, 11, and 24, Table 1), whereas hydrogen sulfide, methyl mercaptan, ethyl mercaptan, diacetyl, and dimethyl trisulfide had not. This is undoubtedly due to the high volatility of these four odorants causing losses during concentration of the solvent extracts. On the other hand, possibly due to their lower vapor pressure, compounds sensorially detected in the solvent extract (4, 6, 8, 9, 12, 15–23, and 25–30, Table 1) were not sensorially detectable in the headspace analysis.

A stepwise reduction of the headspace volume elicited the most odor-active compounds in the headspace of the reaction mixture. 2-Furfurylthiol followed by 2-methyl-3-furanthiol, 2-thenyl mercaptan and ethyl mercaptan were detected in the lowest headspace volumes analyzed (0.02–0.16 mL; Table 2). The two first mentioned thiols were also detected with the highest FD factors in the solvent extract, whereas ethyl mercaptan was sensorially not detectable and the 2-thenyl mercaptan occurred in the extract with a much lower FD factor (cf. 10, Table 1). Presumably, the last thiol is relatively labile, which would lead to significant losses of this odorant during concentration of the extract.

Interestingly, no disulfide was detected in the headspace analysis, corroborating the assumption that a significant proportion of these flavor compounds detected by AEDA result as artifacts during the workup procedure.

5-Acetyl-2,3-dihydro-1,4-thiazine (ADT), which was detected among the key odorants of the solvent extract, only showed a low FD factor in the headspace analysis. Since low odor thresholds in air and water were determined for ADT (Table 3), a significant retronasal flavor contribution of this odorant is to be assumed. However, this needs to be proved by further quantitative experiments.

Table 3. Odor Thresholds of Selected Sulfur-Containing Odorants Identified in the Cysteine/Ribose System

odorant	odor threshold ^a (ng/L of air)
2-methyl-3-thiophenethiol	0.0032–0.0128
2-thenyl mercaptan	0.003–0.012
5-acetyl-2,3-dihydro-1,4-thiazine ^c	0.02–0.08
2-acetyl-2-thiazoline	0.02–0.08
3-mercapto-2-pentanone	0.05–0.2
3-mercapto-2-butanone	0.2–0.8
1-mercapto-2-propanone	1.7–6.8
2-furfurylthiol	0.0025–0.01 ^b
2-methyl-3-furanthiol	0.0025–0.01 ^b
bis(2-methyl-3-furyl) disulfide	0.0006–0.0024 ^b
bis(2-furfuryl) disulfide	0.00015–0.0006 ^b
2-furfuryl 2-methyl-3-furyl disulfide	0.0004–0.0016
2-methyl-3-furyl 3-oxo-2-butyl disulfide	0.01–0.04
2-furfuryl 3-oxo-2-butyl disulfide	0.004–0.016

^a Determined by HRGCO/olfactometry as recently described (Ullrich and Grosch, 1987). (*E*)-2-Decenal (odor threshold = 2.7 ng/L of air) instead of hexanal was used as the reference odorant.

^b Adapted from the literature (Gasser and Grosch, 1990b). ^c Threshold in water = 0.6 µg/L (retronasally) and 1.7 µg/L (nasally) (unpublished results).

Odor Threshold Values. The odor threshold values of selected sulfur-containing odorants in the cysteine/ribose system were approximated by a HRGC/O method. With the exception of 1-mercapto-2-propanone, all compounds showed very low odor thresholds below 1 ng/L of air (Table 3). The lowest value was found for bis(2-furfuryl) disulfide, which was in very good agreement with data reported earlier by Gasser and Grosch (1990a). The odor threshold of the newly identified flavor compound 5-acetyl-2,3-dihydro-1,4-thiazine was of the same order of magnitude as the odor threshold of 2-acetyl-2-thiazoline, recently established as a potent odorant in roast beef (Cerny and Grosch, 1993).


DISCUSSION

The results indicated 2-furfurylthiol, 2-methyl-3-furanthiol, 2-thenyl mercaptan, and ethyl mercaptan to be as the main contributors to the overall roasty, meatlike odor of a thermally treated aqueous cysteine/ribose solution and suggested 3-mercapto-2-pentanone, 5-acetyl-2,3-dihydro-1,4-thiazine, 3-mercapto-2-butanone, and bis(2-methyl-3-furyl) disulfide as further key flavor compounds in the reaction mixture.

3-Mercapto-2-pentanone, 2-methyl-3-furanthiol, and thenyl mercaptan have also been reported among the most odor-active compounds of a thermally treated thiamin/cysteine mixture containing no carbohydrate (Güntert et al., 1993), and quantitative investigation (Grosch and Zeiler-Hilgart, 1993) has established that thiamin/cysteine is a much more effective precursor mixture in the formation of 2-methyl-3-furanthiol than, e.g., cysteine/ribose. However, 2-thenyl mercaptan was not detected among the key odorants when cysteine was omitted from the reaction mixture (Güntert et al., 1993), suggesting that cysteine is the main precursor of this odorant.

On the basis of AEDA results 2-furfurylthiol, 2-methyl-3-furanthiol, 3-mercapto-2-pentanone, and bis(2-methyl-3-furyl) disulfide have also been detected as key odorants in a commercial meat flavoring (flavoring D; Gasser and Grosch, 1990a). Furthermore, 2-methyl-3-furanthiol and 2-furfurylthiol were established as key odorants in cooked beef and chicken broth (Gasser, 1990; Gasser and Grosch, 1990b). In addition, 3-mercapto-2-pentanone significantly contributed to the overall odor

Scheme 1. Hypothetical Pathway Leading from Cysteine and 2,3-Butandione to 5-Acetyl-2,3-dihydro-1,4-thiazine



of chicken broth but did not much contribute to the flavor of cooked beef, whereas the reverse was found for bis-(2-methyl-3-furyl)-disulfide (Gasser, 1990).

2-Furfurylthiol and 2-methyl-3-furanthiol have also been established as key odorants in roasted coffee (Holscher et al., 1990; Blank et al., 1992), suggesting cysteine and/or thiamin as the precursors in the coffee beans. However, in freshly prepared popcorn (Schieberle, 1991) and roasted sesame seeds (Schieberle, 1993) 2-furfurylthiol but not 2-methyl-3-furanthiol was identified as key odorant, indicating a different formation pathway for 2-furfurylthiol here.

5-Acetyl-2,3-dihydro-1,4-thiazine has not been reported previously among the volatiles of Maillard model systems or of food flavors, respectively. The odorant could be formed from a reaction of cysteine and diacetyl, a well-known degradation product of sugars, as detailed in Scheme 1. Formation of an enamine and subsequent decarboxylation would yield the key intermediate I (Scheme 1). Tautomerization and subsequent cyclization of I by a Michael-type nucleophilic attack of the thiol group at the activated methyl carbon atom would yield 5-(2-hydroxyethenyl)-2,3,6-trihydro-1,4-thiazine. Oxidation of this enaminol, either by a second molecule of diacetyl or by a further dicarbonyl compound present in the reaction system, would then result in 5-acetyl-2,3,5-trihydro-1,4-thiazine, which due to the electronegativity of the sulfur atom tautomerizes into the more stable 5-acetyl-2,3-dihydro-1,4-thiazine.

A similar reaction of 2,3-pentanedione would result in 5-propionyl-2,3-dihydro-1,4-thiazine, which was also detected among the odorants of the reaction mixture.

LITERATURE CITED

- Acree, T. E. Bioassays for flavor. In *Flavor Science*; Acree, T. E.; Teranishi, R., Eds.; American Chemical Society: Washington, DC, 1993; pp 1–20.
- Asinger, F.; Schäfer, W.; Herkelmann, G.; Römgens, M.; Reintges, B. D.; Scharein, G.; Wegerhoff, A. Effect of sulfur and ammonia on tridecanon-(7) and pentanone-(2). *Justus Liebig's Ann. Chem.* **1964**, 672, 156–178.
- Blank, I.; Sen, A.; Grosch, W. Potent odorants of the roasted powder and brew of Arabica coffee. *Z. Lebensm. Unters. Forsch.* **1992**, 195, 239–245.
- Cerny, C.; Grosch, W. Evaluation of potent odorants in roasted beef by aroma extract dilution analysis. *Z. Lebensm. Unters. Forsch.* **1992**, 194, 322–325.
- Cerny, C.; Grosch, W. Quantification of character-impact odour compounds of roasted beef. *Z. Lebensm. Unters. Forsch.* **1993**, 196, 417–422.

- Evers, W. J.; Heinsohn, H. H.; Mayers, B. J.; Sanderson, A. Furans substituted at the three position with sulfur. In *Phenolic, Sulfur and Nitrogen Compounds in Food Flavors*; Charalambous, G., Katz, I., Eds.; American Chemical Society: Washington, DC, 1976; pp 184–193.
- Farmer, L. J.; Patterson, R. L. W. Compounds contributing to meat flavour. *Food Chem.* **1991**, *40*, 201–205.
- Farmer, L. J.; Mottram, D. S.; Whitfield, F. B. Volatile compounds produced in Maillard reactions involving cysteine, ribose and phospholipids. *J. Sci. Food Agric.* **1989**, *49*, 347–368.
- Full, G.; Schreier, P. Covalent chromatography. A valuable method for the aroma analysis of thiols at trace levels. *Lebensmittelchemie* **1994**, *48*, 1–5.
- Gasser, U. Flavor volatiles of beef and chicken broths. Ph.D. Dissertation, Technical University of Munich, 1990.
- Gasser, U.; Grosch, W. Aroma extract dilution analysis of commercial meat flavorings. *Z. Lebensm. Unters. Forsch.* **1990a**, *190*, 511–515.
- Gasser, U.; Grosch, W. Primary odorants of chicken broth. *Z. Lebensm. Unters. Forsch.* **1990b**, *190*, 3–8.
- Grosch, W. Detection of potent odorants in foods by aroma extract dilution analysis. *Trends Food Sci. Technol.* **1993**, *4*, 68–73.
- Grosch, W.; Zeiler-Hilgart, G. Formation of meat-like flavor compounds. In *Flavor Precursors*; Teranishi, R., Takeoka, G. R., Güntert, M., Eds.; American Chemical Society: Washington, DC, 1993; pp 183–192.
- Güntert, M.; Brüning, J.; Emberger, R.; Köpsel, M.; Kuhn, W.; Thielmann, T.; Werkhoff, P. Identification and formation of some selected sulfur-containing flavor compounds in various meat model systems. *J. Agric. Food Chem.* **1990**, *38*, 2027–2041.
- Güntert, M.; Bertram, J. H.; Hopp, R.; Silberzahn, W.; Sommer, H.; Werkhoff, P. Thermal generation of flavor compounds from thiamin and various amino acids. In *Recent Developments in Flavor and Fragrance Chemistry*; Hopp, R., Mori, K., Eds.; Verlag Chemie (VCH): Weinheim, Germany, 1993; pp 215–240.
- Guth, H.; Grosch, W. Identification of potent odorants in static headspace samples of green and black tea powders on the basis of aroma extract dilution analysis (AEDA). *Flavour Fragrance J.* **1993**, *8*, 173–178.
- Halang, W. A.; Langlais, R.; Kugler, E. Cubic spline interpolation for the calculation of retention indices in temperature-programmed gas liquid chromatography. *Anal. Chem.* **1978**, *50*, 1829–1832.
- Hofmann, T.; Hässner, T.; Schieberle, P. Determination of the chemical structure of the intense roasty, popcorn-like odorant 5-acetyl-2,3-dihydro-1,4-thiazine. *J. Agric. Food Chem.* **1995a**, *43*, 2195–2198.
- Hofmann, T.; Schieberle, P.; Grosch, W. Model studies on the oxidative stability of selected thiols occurring in food flavors. *J. Agric. Food Chem.* **1995b**, submitted for publication.
- Holscher, W.; Vitzthum, O. G.; Steinhart, H. Identification and sensorial evaluation of impact compounds in roasted Colombian coffee. *Cafe, Cacao, The.* **1990**, *34*, 205–212.
- Kato, S.; Kurata, T.; Fujimaki, M. Volatile compounds produced by the reaction of L-cysteine or L-cystine with carbonyl compounds. *Agric. Biol. Chem.* **1973**, *37*, 539–544.
- Lane, M. J.; Nursten, H. E. The variety of odors produced in Maillard model systems and how they are influenced by the reaction conditions. In *The Maillard Reaction in Foods and Nutrition*; ACS Symposium Series 215; American Chemical Society: Washington, DC, 1983; pp 141–158.
- MacLeod, G. The scientific and technological basis of meat flavors. In *Developments in Food Flavors*; Birch, G. G., Lindley, M. G., Eds.; Elsevier Applied Science: London, 1986; pp 191–223.
- MacLeod, G.; Seyyedain-Ardebili, M. Natural and simulated meat flavors (with particular reference to beef). *Crit. Rev. Food Sci. Nutr.* **1981**, *14*, 309–437.
- Martin, N. Model studies on the reaction of cystine/cysteine and methionine/S-methylcysteine with reducing carbohydrates. Ph.D. Dissertation, University of Berlin, Germany (Nov 1988).
- Milligan, B.; Saville, B.; Swan, J. M. Trisulphides and tetrasulphides from bunte salts. *J. Chem. Soc.* **1963**, 3608–3614.
- Morton, I. D.; Akroyd, P.; May, C. G. Flavouring substance. U.K. Pat. 836694, 1960.
- Mulders, E. J. Volatile compounds from the non-enzymatic browning reaction of the cysteine/cystine-ribose system. *Z. Lebensm. Unters. Forsch.* **1973**, *152*, 193–201.
- Mussinan, C. J.; Katz, J. Isolation and identification of some sulfur chemicals present in two model systems approximating cooked meat. *J. Agric. Food Chem.* **1973**, *21*, 43–45.
- Normington, K. W.; Baker, J.; Molina, M.; Wishnok, J. S.; Tannenbaum, S. R.; Puju, S. Characterization of a nitrite scavenger, 3-hydroxy-2-pyranone, from Chinese wild plum juice. *J. Agric. Food Chem.* **1986**, *34*, 215–217.
- Peer, H. G.; van den Ouweland, G. A. M.; de Groot, C. N. The reaction of aldopentoses and secondary amine salts, a convenient method of preparing 4-hydroxy-5-methyl-2,3-dihydrofuran-3-one. *Recl. Trav. Chim. Pays-Bas* **1968**, *87*, 1011–1016.
- Roberts, D. D.; Acree, T. E. Gas chromatography-olfactometry of glucose-proline Maillard reaction products. *Thermally Generated Flavors*; ACS Symposium Series 543; American Chemical Society: Washington, DC, 1994; pp 71–79.
- Ruther, J.; Baltes, W. Sulfur-containing furans in commercial meat flavorings. *J. Agric. Food Chem.* **1994**, *42*, 2254–2259.
- Sakaguchi, M.; Shibamoto, T. Formation of heterocyclic compounds from the reaction of cysteamine and D-glucose, acetaldehyde or glyoxal. *J. Agric. Food Chem.* **1978**, *26*, 1179–1183.
- Schieberle, P. Studies on bakers yeast as source of Maillard-type bread flavour compounds. In *The Maillard Reaction in Food Processing, Human Nutrition and Physiology*; Finot, P. A., Aeschbacher, H. U., Hurrell, R. F., Liardon, R., Eds.; Birkhäuser: Basel, 1990; pp 187–196.
- Schieberle, P. Primary odorants in popcorn. *J. Agric. Food Chem.* **1991**, *39*, 1141–1143.
- Schieberle, P. Studies on the flavour of roasted white sesame seeds. In *Progress in Flavour Precursor Studies*; Schreier, P., Winterhalter, P., Eds.; Allured Publishing: Carol Stream, IL, 1993; pp 343–360.
- Schieberle, P.; Grosch, W. Evaluation of the flavour of wheat and rye bread crusts by aroma extract dilution analysis. *Z. Lebensm. Unters. Forsch.* **1987**, *185*, 111–113.
- Sen, A.; Schieberle, P.; Grosch, W. Quantitative determination of (E)- β -damascenone in foods using a stable isotope dilution assay. *J. Agric. Food Chem.* **1991**, *39*, 757–759.
- Silwar, R. Gas chromatography mass spectrometry investigation of aroma compounds from the reaction of cysteine and rhamnose under roasting conditions. *Z. Lebensm. Unters. Forsch.* **1992**, *195*, 112–119.
- Tressl, R.; Helak, B.; Martin, N.; Kersten, E. Formation of amino acid specific Maillard products and their contribution to thermally generated aromas. *Thermal Generation of Aromas*; ACS Symposium Series 409; American Chemical Society: Washington, DC, 1989; pp 156–171.
- Ullrich, F.; Grosch, W. Identification of the most intense volatile flavour compounds formed during autooxidation of linoleic acid. *Z. Lebensm. Unters. Forsch.* **1987**, *184*, 277–282.
- van den Ouweland, G. A. M.; Peer, H. G. Components contributing to beef flavour. Volatile components produced by the reaction of 4-hydroxy-5-methyl-3(2H)-furanone and its thio analog with hydrogen sulfide. *J. Agric. Food Chem.* **1975**, *23*, 501–505.
- van den Ouweland, G. A. M.; Olsman, H.; Peer, H. G. Challenges in meat flavor research. In *Agricultural and Food Chemistry: Past, Present, Future*; Teranishi, R., Ed.; AVI: Westport, CT, 1978; pp 292–314.
- Werkhoff, P.; Brüning, J.; Emberger, R.; Güntert, M.; Köpsel, M.; Kuhn, W. Studies on volatile sulfur-containing flavour components in yeast extract. In *Proceedings of the 11th International Congress of Essential Oils, Fragrances and Flavours*; Vol. 4, Chemistry, Analysis and Structure; Bhat-tacharyya, S. C., Sen, N., Sethi, K. L., Eds.; Oxford & IBH Publishing: New Delhi, 1989; pp 215–243.

- Werkhoff, P.; Bretschneider, W.; Güntert, M.; Hopp, R.; Köpsel, M.; Surburg, H. Identification and formation of novel sulfur compounds in a meat flavour model system. *Chem. Mikrobiol. Technol. Lebensm.* **1991**, *13*, 111-124.
- Whitfield, F. B.; Mottram, D. S.; Brock, S.; Puckey, D. J.; Salter, L. J. Effect of phospholipid on the formation of volatile heterocyclic compounds in heated aqueous solution of amino acids and ribose. *J. Sci. Food Agric.* **1988**, *42*, 261-272.
- Whitfield, F. B.; Mottram, D. S.; Shaw, K. J. The formation of some novel thiols and disulphides in model systems containing 4-hydroxy-5-methyl-3(2H)-furanone. In *Progress in Flavour Precursor Studies*; Schreier, P., Winterhalter, P., Eds.; Allured Publishing: Carol Stream, IL, 1993; pp 395-400.

Wiley, R. H.; Jarboe, C. H. 2-Pyrones. XIX. 3-Hydroxy-2-pyrone and 4-arylhydrazono-2,3-pyranones. *J. Am. Chem. Soc.* **1956**, *78*, 2398-2401.

Received for review October 20, 1994. Revised manuscript received April 4, 1995. Accepted May 26, 1995.*

JF9405968

* Abstract published in *Advance ACS Abstracts*, July 1, 1995.