

VOLUME 17 NUMBER 3 • MARCH 1999

nature biotechnology


<http://biotech.nature.com>

**Cancer-targeting
cytotoxic RNases**

**Synthetic promoters
flex their muscles**

Splicing up gene therapy

Iron-fortified rice



***** 5-DIGIT 53706
53706UNIV 0000 DEC99 02000

BI

00210

UNIV OF WISCONSIN
STEENBOCK MEMORIAL LIBR
550 BABCOCK DR
MADISON WI 53706-1201

nature biotechnology



Structure of an engineered dimeric human ribonuclease (foreground) which has been altered to both facilitate targeting to cancer cells and to render it insensitive to its natural cytosolic inhibitor (shown in background). For more details, see Suzuki et al. p 265, and Wagner and Lipovsek, p 227. Illustration by Ken Eward (BioGrafx, Mount Vernon, Ohio).

OPINION

Editorial

- Killer potatoes: Where's the data? 207
More gods and monsters 207

Commentary

- Feeding the world, harmoniously 208
Tom Crompton

- Keeping the biotechnology of antisense in context 209
C.A. Stein

Correspondence

- Spin science 210
Ronald W. Eastman

- Shorter is better 210
M. Raafat El-Gewely

IN BRIEF

Business and Regulatory News

- Warner Lambert acquires speculative Agouron pipeline. . . IPO's increase in Europe. . . Cortech relists on Nasdaq. . . India sequences chickpea. . . Shaman switches gears. . . Marimastat fails again. . . Japan easy on gene therapy. . . First US-German start-up. . . Entremed's stock seesaws. . . New plant to desulfurize oil. . . Research collaborations 211

Research News

- Controlled release microchip. . . Muscling in on bone formation. . . Reprogramming neural stem cells. . . Sizing up DNA. . . Priming axon regeneration 213

This month in *Nature Biotechnology*

- Antitumor vaccines with punch. . . Building better promoters. . . Invading diagnostics. . . A synthetic IL-10 mimic. . . Stealthy RNases. . . Making antibodies more human. . . Reprogramming messages 214



Nature Biotechnology (ISSN 1087-0156) is published monthly by Nature America Inc., headquartered at 345 Park Avenue South, New York, NY 10010-1707. Editorial Office: 345 Park Avenue South, New York, NY 10010-1707. Tel: (212) 726 9200, Fax: (212) 696 9635. North American Advertising: Nature Biotechnology, 345 Park Avenue South, New York, NY 10010-1707. Tel: (212) 726-9200. Fax: (212) 696-9006. European Advertising: Nature Biotechnology, Porters South, Crinan Street, London N1 9XW. Tel: 44 171 833 4000. Fax: 44 171 843 4996. New subscriptions, renewals, changes of address, back issues, and all customer service questions in the Americas should be addressed to: Nature Biotechnology Subscription Department, P.O. Box 5054, Brentwood, TN 37024-5054. Tel: (800) 524-0384, Direct Dial (615) 377 3322, Fax: (615) 377 0525. Outside the Americas: Nature Biotechnology, Macmillan Magazines Ltd., Houndsmill, Brunel Road, Basingstoke, RG21 6XS, U.K.. Tel: +44-(0)1256-329242. Fax: +44-(0)1256 812358. Email: subscriptions@nature.com. Annual subscription rates: U.S./Canada/Mexico: U.S. \$595, Canada add 7% for GST (institutional/corporate), U.S. \$99, Canada add 7% for GST (individual making personal payment BN: 14091 1595 RT); U.K./Europe: £275 (institutional/corporate), £62 (individual making personal payment); Rest of world (excluding Japan): £320 (institutional/corporate), £83 (individual making personal payment); Japan: Contact Japan Publications Trading Co. Ltd., 2-1 Sarugaku-c'ho 1 chome, Chiyoda-ku, Tokyo 101, Japan, Tel: 03 292-3755. Back issues: U.S./Canada, \$45, Canada add 7% for GST; Rest of world: surface U.S. \$43, air mail U.S. \$45. Reader inquiry service: Nature Biotechnology, P.O. Box 5464, Pittsfield, MA 01203-9332. Periodicals postage paid at New York, NY 10010, and additional mailing offices. POSTMASTER: Send address changes to Nature Biotechnology Subscription Department, P.O. Box 5054, Brentwood, TN 37024-5054. Executive Officers of Nature America Inc.: Nicholas Byam Shaw, Chairman of the Board; Mary Waltham, President; Edward Vais, Secretary-Treasurer. Printed by Publishers Press, Inc., Shepherdsville, KY, US. Copyright ©1999 Nature America Inc.

CONTENTS

ANALYSIS

Business and Regulatory News	215
Stem cell transformations proliferate.....	215
UK government supports biotech	217
USDA appeases organic lobby.....	217
Synergen lineage may finally pay off.....	219
Overview of FY 2000 US federal budget request	220
Cancer Pathways' target not validated by clinical results	220
Pharma strategies extend drug lives	222
Return of the Neuer Markt	222
Research News and Views	223
A new routine in molecular gymnastics	223
Ryszard Kole	SEE ALSO 246
Piecing together more efficient gene expression.....	224
Nikunj V. Somia, Tal Kafri, and Inder M. Verma	SEE ALSO 241
Extending genetic vaccines with chemokines.....	226
Thomas Kipps and Robert Mendoza	SEE ALSO 253
Smartbombs and cloaking devices	227
Richard W. Wagner and Dasa Lipovsek	SEE ALSO 265
DREB takes the stress out of growing up.....	229
Nicholas Smirnoff and John A. Bryant	SEE ALSO 287
Pumping iron.....	230
Michael R. Sussman	SEE ALSO 282

FEATURES

Proteomics: translating genomics into products?	233
Alan Dove	
Drug approval in Europe	237
Gary Walsh	


RESEARCH ARTICLES

Synthetic muscle promoters: activities exceeding naturally occurring regulatory sequences.....	241
Xuyang Li, Eric M. Eastman, Robert J. Schwartz, and Ruxandra Draghia-Akli	SEE ALSO 224
Spliceosome-mediated RNA <i>trans</i> -splicing as a tool for gene therapy	246
M. Puttaraju, Sharon F. Jamison, S. Gary Mansfield, Mariano A. Garcia-Blanco, and Lloyd G. Mitchell	SEE ALSO 223
Genetic fusion of chemokines to a self tumor antigen induces protective, T-cell dependent antitumor immunity	253
Arya Biragyn, Kenji Tani, Michael C. Grimm, Steven Weeks, and Larry W. Kwak	SEE ALSO 226
Directed evolution of thymidine kinase for AZT phosphorylation using DNA family shuffling.....	259
Fred C. Christians, Leonardo Scapozza, Andreas Cramer, Gerd Folkers, and Willem P.C. Stemmer	

page 255

CONTENTS

Engineering receptor-mediated cytotoxicity into human ribonucleases by steric blockade of inhibitor interaction	265
Motoshi Suzuki, Shailendra K. Saxena, Ester Boix, Robert J. Prill, Veena M. Vasandani, Jane E. Ladner, Cynthia Sung, and Richard J. Youle SEE ALSO 227	
A synthetic mimic of a discontinuous binding site on interleukin-10	271
Ulrich Reineke, Robert Sabat, Rolf Misselwitz, Heinz Welfle, Hans-Dieter Volk, and Jens Schneider-Mergener	
A recombinant, fully human monoclonal antibody with antitumor activity constructed from phage-displayed antibody fragments	276
Gerwin A. Huls, Ingmar A.F.M. Heijnen, Maria E. Cuomo, Jacob C. Koningsberger, Luus Wiegman, Edwin Boel, Anne-Reneé van der Vuurst de Vries, Sabine A.J. Loyson, Wijnand Helfrich, Gerard P. van Berge Henegouwen, Marja van Meijer, John de Kruif, and Ton Logtenberg	
Iron fortification of rice seed by the soybean ferritin gene	282
Fumiyuki Goto, Toshihiro Yoshihara, Naoki Shigemoto, Seiichi Toki, and Fumio Takaiwa SEE ALSO 230	
Improving plant drought, salt, and freezing tolerance by gene transfer of a single stress-inducible transcription factor	287
Mie Kasuga, Qiang Liu, Setsuko Miura, Kazuko Yamaguchi-Shinozaki, and Kazuo Shinozaki SEE ALSO 229	
Polymorphism identification and quantitative detection of genomic DNA by invasive cleavage of oligonucleotide probes	292
Victor Lyamichev, Andrea L. Mast, Jeff G. Hall, James R. Prudent, Michael W. Kaiser, Tsetska Takova, Robert W. Kwiatkowski, Tamara J. Sander, Monika de Arruda, David A. Arco, Bruce P. Neri, and Mary Ann D. Brow	



page 277

RESOURCES



page 297

Patents	
Have you bargained away patent rights?	297
Breffni Baggot	
Recent antibody-related patents	299
In the Laboratory	
Displacement chromatography for biopolymer separation	300
Ruth Freitag	
Books	
Electric genes	303
Terry Gaasterland	
Internet	
Web PCR	304
Stacey McCann	
People	
Promotions, advancements	305
New Products	
Genomics, antibodies	306
Industry Trends	
Mental disorder drug discovery	307
Aris Persidis and Rachel M. Copen	

Iron fortification of rice seed by the soybean ferritin gene

Fumiyuki Goto¹, Toshihiro Yoshihara^{1*}, Naoki Shigemoto^{2,3}, Seiichi Toki², Fumio Takaiwa^{2*}

¹Department of Bio-Science, Central Research Institute Electric Power Industry, 1646 Abiko, Chiba 270-1194, Japan. ²Department of Biotechnology, National Institute of Agrobiological Resources, Tsukuba, Ibaraki 305-8602, Japan. ³Institute of Biotechnology, Hiroshima Pref. Agricultural Research Center, Higashi-Hiroshima, Hiroshima 739-0151, Japan. *Corresponding author (e-mail: yoshihara@criepi.denken.or.jp; takaiwa@ss.abr.affrc.go.jp).

Received 2 April 1998; accepted 19 January 1999

To improve the iron content of rice, we have transferred the entire coding sequence of the soybean ferritin gene into *Oryza sativa* (L. cv. Kita-ake) by *Agrobacterium*-mediated transformation. The rice seed-storage protein glutelin promoter, *GluB-1*, was used to drive expression of the soybean gene specifically in developing, self-pollinated seeds (T1 seeds) of transgenic plants, as confirmed by reverse transcription PCR analysis. Stable accumulation of the ferritin subunit in the rice seed was demonstrated by western blot analysis, and its specific accumulation in the endosperm by immunologic tissue printing. The iron content of T1 seeds was as much as threefold greater than that of their untransformed counterparts.

Keywords: nutrition, iron content, genetic improvement, glutelin promoter, *Oryza sativa*

Iron deficiency is a serious nutritional problem that afflicts an estimated 30% of the world population¹, especially where vegetable-based diets are the primary food source². Although supplements added to food or taken in tablet form are effective in preventing and controlling iron deficiency, such treatments are difficult to implement in developing countries because of the associated costs and the small number of primary health care programs³. Some crops (e.g., spinach and leguminous plants) are known for their high iron content⁴. However, these plants also contain oxalic acid and phytate-like substances that decrease their bioavailability⁵. The iron content of crops has been improved by increasing the iron concentration of the hydroponic culture media or soil. However, this method is costly and cannot be used to target iron accumulation to a desirable part of the plant. In addition, the excess iron or the chelator in the media suppresses plant growth and can also reduce commercial value and productivity⁶.

Ferritin is an iron-storage protein found in animals, plants, and bacteria⁷. Comprising 24 homologous or heterologous subunits, it assembles to form a large complex (450 kDa) that stores up to 4500 iron atoms in its central cavity^{7,8}. The ferritin gene has been isolated and sequenced in plants, including soybean⁹, French bean¹⁰, pea¹¹, and maize¹². Although the amino acid sequence homology between plant and mammalian ferritin is low (39–49% within the transit and extension peptides⁸; 56–66% without the additional peptide⁹), its molecular conformation and iron-storage capacity is conserved.

Ferritin is thought to play two main roles in living cells, both through modulation of iron content¹³. One is to provide iron for the synthesis of iron proteins such as ferredoxin and cytochromes. The other is to prevent damage from free radicals produced by iron/dioxygen interactions. In animals, ferritin synthesis is regulated at the translational, not the transcriptional, level¹³. Translation of the ferritin mRNA increases in response to high levels of iron, thus producing more ferritin for storage, and conversely, translation is decreased by low levels of iron. In plants, however, iron-dependent regulation of phytoferritin synthesis occurs at the transcriptional level^{14,15} and is also responsive to developmental cues^{16,17}.

Recent studies show that ferritin is used by both plants and animals as the storage form of iron¹⁸ and that, orally administered, it

can provide a source of iron for treatment of rat anemia¹⁹. These findings suggest that increasing the ferritin content of cereals by genetic engineering may help to solve the problem of dietary iron deficiency. We have explored that possibility by introducing the soybean ferritin gene into rice plants (*Oryza sativa* L. cv. Kita-ake) by *Agrobacterium*-mediated transformation. The introduced ferritin gene was expressed under the control of a rice seed-storage protein glutelin promoter, *GluB-1* (-1302/+18)²⁰, to mediate the accumulation of iron specifically in the grain. Our results indicated that the soybean ferritin could be highly expressed and accumulated in the endosperm tissue of heterologous rice plants. Notably, the transgenic seeds stored up to three times more iron than the normal seeds.

Results

Expression of the soybean ferritin in rice seeds. Thirty independent transgenic rice lines (T0 plants) were obtained by *Agrobacterium*-mediated transformation, and the insertion of the ferritin cDNA in their chromosomes was confirmed by PCR analysis of leaves from T0 plants (data not shown). All lines produced self-pollinated seeds (T1 and T2 seeds), and heterozygous lines were determined by screening with bialaphos and/or PCR, except in experiments to determine iron content of seeds. Southern blot analysis of total DNA isolated from T2 leaves of several lines (Fk1, 4, 7, 11, 16, 22, and 25) showed that independent transformants contained one to several copies of the introduced gene per haploid genome (Fig. 1A and B).

To examine the expression of the ferritin gene, total RNA isolated from T2 immature seeds of three lines (Fk1, 11, and 22) was analyzed by reverse transcriptase PCR (RT-PCR) analysis using primers specific for soybean ferritin. Figure 1C shows that a 566 bp band corresponding to a portion of the soybean ferritin cDNA could be amplified from cDNA prepared from seeds, but not leaves, of transformed plants. This suggested that the expression of the ferritin gene was tissue-specific. The specificity of the RT-PCR reaction for the soybean ferritin gene was confirmed by the absence of an amplification product from cDNA prepared from seeds of nontransformed plants (N4).

Total protein homogenates isolated from T1 seeds (Fk1, 4, 7, 11, 16, 22, and 25) were analyzed by western blotting. A polyclonal anti-

RESEARCH

body directed against the soybean ferritin gene specifically immunodecorated a 28 kDa band, confirming that the soybean ferritin protein accumulates in seeds (Fig. 2A and B). The 28 kDa band specific to soybean ferritin was detected in protein homogenates from soybean, but not from untransformed rice and tobacco. Though ferritin levels varied between different individual lines, the protein could be detected in almost all the transformants assayed (Fig. 2B), and the amount was estimated to be between 0.01–0.3% of the total seed

protein (data not shown). The 28-kDa band is very similar in size to that observed in the native soybean cotyledon and transgenic tobacco leaf containing the ferritin gene (Fig. 2A). The absence of proteolytic products indicated that the soybean ferritin was stable in the heterogeneous rice seed. In addition, another specific hybridized band about 32 kDa in molecular size was observed in *Escherichia coli* and native soybean, but was barely detectable in transgenic rice and tobacco plants (Fig. 2A). This band may correspond to a precursor




Figure 1. (A) Construction of *GluB-1* promoter—soybean ferritin chimeric gene in the pGPTV-35S-bar binary vector. (a) Deduced hybridization position of digoxigenin-labeled ferritin cDNA probe. (b) Annealing sites of RT-PCR primers, which amplified a fragment of 566 bp, for detecting the introduced soybean ferritin gene expression. (B) Southern blot hybridization of DNA prepared from individual T2 and a nontransformed control plants. Total DNA (10 µg) was digested with *Hind*III and hybridized with digoxigenin-labeled soybean ferritin cDNA probe. Lane 1 and 2, two and one copy (per a haploid genome) markers; lane 3, nontransformant (N4); lane 4, Fk1; lane 5, Fk1n (the ferritin cDNA was not detected by PCR method); lane 6, Fk4; lane 7, Fk7; lane 8, Fk11; lane 9, Fk16; lane 10, Fk22; and lane 11, Fk25. (C) RT-PCR analysis of total RNA from transgenic rice expressing the soybean ferritin gene. NTr, nontransformant (N4); Tr, transformant (1, Fk1; 2, Fk11; and 3, Fk22); M, ϕ X174/*Hae*III; N, no template; P, pGPTV-bar/Fer; S, seed; L, leaf.




Figure 2. Western blot analysis of soybean ferritin protein in the transgenic rice seeds. (A) Total protein extracts from *E. coli*, rice (T1-Fk1), and tobacco (lanes 1, 2, and 4) transformed with the soybean ferritin cDNA, and from nontransformed rice, tobacco, and soybean (lanes 3, 5, and 6) were separated by SDS-PAGE and immunoblotted with the soybean ferritin rabbit polyclonal antibody. Twenty micrograms were loaded in each lane, except for the *E. coli* extract, which used 28 µg. (B) Variability in soybean ferritin expression in different transgenic rice lines. 3.75 µg of total protein from one matured seed of each transformant (T1-Fk1, 4, 7, 11, 16, 22, and 25) and nontransformant (N; N4) was analyzed by western blot analysis with soybean ferritin antibody. Soybean ferritin subunit produced by *E. coli* (4.75 µg) was used as the positive control (P).




Figure 3. Immunological tissue printing of a seed from transgenic rice expressing soybean ferritin cDNA. Em, embryo; En, endosperm; SA, subaleurone layer. Each of three grains of three T1 transgenic lines (Fk1, 11, and 22) were analyzed by tissue printing using the soybean ferritin polyclonal antibody. Representative staining from a typical line (Fk1) is shown, with tissues that accumulate soybean ferritin staining dark brown.

RESEARCH




Figure 4. Comparison of iron content in transgenic rice seeds expressing soybean ferritin. Each dot represents the iron content of one seed. Ten T1 seeds were independently measured in each experimental line (11 transformed lines: 1, Fk19; 2, Fk4; 3, Fk17; 4, Fk13; 5, Fk13; 6, Fk25; 7, Fk2; 8, Fk7; 9, Fk22; 10, Fk11; and 11, Fk1; and four nontransformed rice lines: 1, N1; 2, N3; 3, N2; 4, N4). Bars indicate an average value of each experimental line. Asterisks indicate significant differences between the pooled mean values of all nontransformants and that of each independent transgenic lines at $p < 0.05$ (t -test).

Table 1. Iron content in some tissues of a transgenic rice line expressing soybean ferritin cDNA.

	Seed	Leaf ($\mu\text{g-Fe/g-DW}$)	Stem	Root
Transformants				
(Fk 1)	38.1 ± 4.5^a	104.8 ± 7.9^b	162.8 ± 8.1^b	962.1 ± 65.6^b
(Fk 11)	35.9 ± 7.7^a	98.8 ± 25.8^b	175.5 ± 17.2^b	966.8 ± 105.7^b
Nontransformants				
(N4)	14.3 ± 3.0	119.3 ± 8.9	170.0 ± 5.3	956.0 ± 37.4

Numerical status shows means of iron concentration in transformants (Fk1 and Fk11) and nontransformants (N4) from three (nonseed tissues) or ten (seed) independent replications. ^aSignificant or ^bnonsignificant differences between the mean values were calculated by t -test ($p < 0.05$).

form of the protein (see Discussion). Bands of greater than 32 kDa in transformed and nontransformed tobacco (Fig. 2A, lanes 4 and 5) may be due to the nonspecific reaction.

Immunological tissue printing was carried out to visualize the distribution of soybean ferritin in T1 seed (Fk1). Although a little background staining could be seen in the nontransgenic seed, darker brown staining was visible in the transgenic rice endosperm tissue (Fig. 3). The subaleurone layer of the endosperm stained most intensely, indicating a high accumulation of the ferritin in this tissue.

Tissue-specific enrichment of the iron content. Iron levels in the 11 mature whole T1 seeds varied from 13.3 ± 2.8 to 38.1 ± 4.5 $\mu\text{g/g}$ dry weight (DW) (Fig. 4). Given that the iron contents in seeds of nontransformants ranged from 8.6 ± 3.3 to 14.3 ± 3.0 $\mu\text{g/g}$ DW, we estimated a threefold increase in iron content of seeds from the highest-expressing transgenic rice lines. When the pooled mean values between all transformants (22.5 ± 1.3 $\mu\text{g/g}$ DW) and all nontransformants (11.2 ± 0.9 $\mu\text{g/g}$ DW) were compared by t -test, there was a significant difference ($p = 1.53 \times 10^{-11}$). Iron content was similar in non-seed tissues (i.e., leaf, stem and root) from T1 transformants and nontransformants (Table 1).




Figure 5. Distribution of iron in the rice seeds transformed with soybean ferritin cDNA. Iron content of embryo (open rectangular) and endosperm (closed rectangular) are shown for 10 T2 seeds of a transformed (Fk1) and nontransformed rice line (N4) are shown. Mean values are indicated $\pm$ standard error ($n=3$). Differences between transformants and nontransformants were calculated to be significant, with t -test value of $p = 0.02$.

To determine whether the transformation affected the iron content of seed tissue, we separately determined iron content in the embryo and endosperm of T2 seeds. The average iron content in the endosperm of a transformant (Fk1) was 3.4 ± 0.5 $\mu\text{g/10 grains}$, and that of a nontransformant (N4) was 1.6 ± 0.1 $\mu\text{g/10 grains}$ (Fig. 5). In contrast, the iron content of the embryo of the transformant and nontransformant were 1.2 ± 0.1 $\mu\text{g/10 grains}$ and 1.3 ± 0.2 $\mu\text{g/10 grains}$, respectively. Statistical analysis using the t -test showed a significant difference in the averaged iron content of endosperm between the transformant and nontransformant ($p = 0.02$), but not in that of embryo ($p = 0.64$). These results demonstrated that under the regulation of the glutelin promoter, the increase in iron content is endosperm specific. Expression of the soybean ferritin gene did not noticeably affect the appearance of the rice grain, as seeds from the transformants were approximately the same weight as those of nontransformants (approximately 18 mg DW). Moreover, the seeds from transformants were unaltered in their external appearance and did not exhibit signs of chlorosis, a symptom of iron deficiency.

Discussion

Soybean ferritin is synthesized as a 28 kDa protein in transgenic rice seeds, corresponding to the expected size of the native matured subunit²¹. From comparisons of ferritin amino acid sequences, including soybean ferritin, Ragland et al.⁹ proposed that the plant ferritin subunits could be generated from a single precursor by posttranslational processing. The processing yields three types of subunits of different molecular masses: a 32 kDa precursor, a 28 kDa matured subunit, and a 26.5 kDa partially degraded subunit. The proper assembly of the protein is required for import of the subunit into the plastid, as well as for iron storage function. In fact, a 32 kDa immunoreactive band corresponding to the ferritin precursor was observed in native soybean and was the predominant form in *E. coli* expressing the ferritin gene. This is consistent with the results of Kimata and Theil²² and Van Wuytswinkel et al.^{23,24}, who observed that the prokaryotic and in vitro translation systems were incapable of processing the precursor into the mature subunit. In contrast to the usual susceptibility of foreign proteins to proteolytic degradation²⁵⁻²⁷, the 28 kDa mature subunit appeared to be stable, suggesting that the endosperm tissue is particularly appropriate for accumulation of foreign ferritin.

The rice glutelin promoter, *GluB-1*, which is used to express the soybean ferritin cDNA in the transformation, has two distinct advantages in this experimental system other than the fact that the

RESEARCH

gene originated from rice. One is that it conferred endosperm-specific expression of the ferritin gene, which led to specific accumulation of the protein within that tissue. This accumulation was consistent with the β -glucuronidase staining pattern directed by the *GluB-1* promoter^{28,29}. The other advantage was the ability of the promoter to induce a high level of expression. Takaiwa et al.²⁷ indicated that glycinins expressed under the control of *GluB-1* promoter accumulated at the level of 0.02–4% of total proteins in dry seed of tobacco. This level was more than 40-fold higher than that achieved with a constitutive expression system using the cauliflower mosaic virus (CaMV) 35S promoter³⁰. Our results (approximately 0.01–0.3% of total protein) are comparable to these earlier findings. The varying expression levels were independent of the copy numbers in the transformants and may possibly arise from the different sites of transgene integration in the rice chromosome³¹.

The iron content of the rice seed was significantly increased by expression of the soybean ferritin gene under control of the endosperm-specific promoter, *GluB-1*. Specifically, the seeds from the transgenic rice accumulated up to threefold more iron ($38.1 \pm 4.5 \mu\text{g/g DW}$) than normal seeds ($11.2 \pm 0.9 \mu\text{g/g DW}$). This achievement suggests that it may be feasible to produce "ferritin rice" as an iron supplement in the human diet. The iron content in a meal-size portion of "ferritin rice" (approximately 5.7 mg-Fe/150 g-DW) would be sufficient to provide 30–50% of the daily adult iron requirement (approximately 13–15 mg Fe [ref. 4]). We recently demonstrated that expression of the soybean ferritin gene in tobacco under the control of the CaMV 35S-regulated promoter resulted in a maximum iron content in transformed leaves that was about 1.3 times that of nontransformed leaves ($109.3 \mu\text{g/g DW}$ compared with $84.0 \mu\text{g/g DW}$, respectively)³². This increase seems to be low compared with the rice system. However, it is interesting to note that absolute amounts of iron increase in these two systems are very similar ($24.3 \mu\text{g/g DW}$ in tobacco versus $26.9 \mu\text{g/g DW}$ in rice). Although a positive correlation can be observed between exogenous ferritin accumulation and iron content³², it is possible that the amount of iron accumulation is restricted by transport of iron to the ferritin molecule, rather than simply by levels of ferritin protein. Thus, it may be possible to store larger amounts of iron in the exogenous ferritin molecule by cointegrating the ferritin gene and the iron reductase-like iron transporter gene³³.

Experimental protocol

Construction and transformation of chimeric gene and plant cultivation. Ferritin cDNA was isolated from soybean cotyledons (Glycine max L. cv. Kitano-shiki) a few days after germination. polyCAT mRNA was isolated from 1 g fresh weight (FW) of the cotyledons and the ferritin cDNA (0.84 kb) was amplified by RT-PCR. The primers for the reverse transcription were random primers (p[dN]6), and those for the PCR corresponded to the ferritin cDNA sequence reported by Lescure et al.¹³ Forward primer: 5'-ATGGCTCTTGCTCCATCCAAAGTT; Reverse primer: 5'-TTGATCAAAGTGCCAAACACCGTG. The ferritin cDNA fragment was then cloned into the pCR II vector (Invitrogen, San Diego, CA) and the sequence was confirmed. Alignment of the fragment sequence with the known soybean ferritin sequence¹⁵ showed 95% identity in the conserved regions (data not shown). The divergence may have resulted from the use of different soybean cultivars. Next, the ferritin cDNA was inserted into a modified binary vector pGPTV-35S-bar, replacing the *uidA* gene. This modified vector was reconstructed from the pGPTV-bar binary vector³⁴ by exchange of the *Nos* with the CaMV 35S as the promoter of the bar gene. The ferritin cDNA was introduced between the restriction enzyme site of *Bam*HI and *Sac*I under control of the rice seed-storage protein glutelin promoter, *GluB-1* (-1302/+18)²⁰. It was flanked at the 3' end by the *Nos* polyadenylation signal (Fig. 1A). The plasmid pGPTV-35S-bar, containing the soybean ferritin gene, was transferred into *Agrobacterium tumefaciens* strain C58 or EH105 by electroporation. This *Agrobacterium* was used to infect 3-week-old calli derived from a mature rice seed (*Oryza sativa* L. cv. Kita-ake) for three days. Next the calli were successively cultured in N6 selection medium³⁵ (supplemented with 3% sucrose, 0.4% gelrite, 100 mg/l myoinositol, 300 mg/l

casamino acids, 2878 mg/l proline, 2 mg/l 2,4-D, 500 mg/l Claforan and 5 mg/l bialaphos) for 6 weeks, and MS regeneration medium³⁶ (supplemented with 3% sucrose, 3% sorbitol, 0.8% agar, 2 g/l casamino acids, 0.2 mg/l NAA, 2 mg/l kinetin, 250 mg/l Claforan, and 5 mg/l bialaphos) for 5 weeks at $28 \pm 1^\circ\text{C}$, under continuous light condition. Each line was designated Fk (Ferritin/Kita-ake) followed by an ordered number from 1 through 30, respectively. Total DNA was isolated from leaves of regenerated plantlets and tested for the integration of the soybean ferritin gene by the PCR method (using the same primer set as described for gene construction). These transformants (T0 plants) were subsequently acclimated in the MS medium without hormones and antibiotics for 3 weeks, transplanted to soil, and grown in a greenhouse ($30 \pm 3^\circ\text{C}$, 12 hours light/12 hours dark) until harvesting. T0 plants and the self-pollinated heterozygous progenies (T1 and T2) were used for the experiments described below.

Southern blot analysis. Total DNA was prepared from leaves of T2 progenies of transformants, according to the method of Shure et al.³⁷ *Hind*III-digested DNA (10 μg) was separated by agarose gel electrophoresis, blotted to a nylon membrane, and hybridized with a ferritin cDNA probe labeled with digoxigenin (Boehringer Mannheim, Mannheim, Germany). Hybridization and detection were carried out according to Engler-Blum et al.³⁸

RNA isolation from the transformants and characterization. Total RNA was extracted from immature T2 seeds collected about 14–21 days after pollination and from T1 leaves (harvested with the seeds) using a commercial kit based on guanidinium thiocyanate and a column filtration system (RNAqueous kits; Ambion, Austin, TX). After purification by lithium chloride precipitation and DNase digestion, expression of the introduced soybean ferritin gene was analyzed by the RT-PCR method, using the total RNAs and a set of primers as follows: Forward primer: 5'-CAGT-GTTGGGGATGCTCTGAA; Reverse primer: 5'-GGTCATTGTTGCGATCT-GCCA. These primers were designed to amplify a 566 bp portion of the soybean ferritin cDNA (Fig. 1A).

Protein extraction and immunologic detection. One T1 immature seed (about 14–21 days after pollination) from each of the transformants was separately homogenized with mortar and pestle in extraction buffer (80 mM Tris-HCl, pH 7.0, 2% SDS, 2% 2-mercaptoethanol, and 20% glycerol). After 3 min boiling, the resulting extracts were centrifuged at 12,000 rpm for 20 min, and the supernatant fractions were collected. Proteins in these extracts were measured according to the method of Bradford³⁹, separated by SDS-PAGE, and transferred to PVDF membrane (Immobilon-P⁵⁰; Millipore, Tokyo) using a buffer system consisting of 25 mM Tris-H₃BO₃, pH 9.5, and 20% methanol. The production of the soybean ferritin was confirmed by western blot analysis based on the ordinary method⁴⁰, except for using TBS buffer (Tris-HCl, pH 7.5, 0.15 M NaCl). For the western blotting procedure, 1/2000 diluted polyclonal antiserum diluted in TBS buffer containing 0.05% Tween 20 and 5% nonfat dried milk. The antiserum was raised in a male rabbit against the soybean ferritin subunit³², which was produced in *E. coli* harboring the ferritin gene in expression vector pQE30 (Qiagen, Hilden, Germany). Protein extracts from nontransformed soybean cotyledons and transgenic *E. coli* and tobacco³² harboring the ferritin gene were processed in parallel as positive controls.

In situ localization of the ferritin in transgenic rice was observed by the immunologic tissue printing method of Kamachi et al.⁴¹ An immature seed (about 14–18 days after pollination, T1) from transgenic rice was cut in half at the vertical section and pressed against a nitrocellulose membrane. Then the membrane was treated with the soybean ferritin polyclonal antiserum used for the western blotting, diluted for 1/2000 with TBS buffer. For detection, a biotinylated goat antirabbit IgG solution and ABC reagent containing avidin DH and biotinylated alkaline phosphatase H or horseradish peroxidase (Vectastain ABC Elite or ABC-AP kit; Vector Laboratories, Burlingame, CA) was used according to the protocol recommended by the manufacturer. Substrate kit II (Vectastain ABC-AP kit; Vector Laboratories) or Immunostain HRP-1000 (Konica, Tokyo) were used for visualizing the signal.

Iron content determination. Iron content was determined from 11 independent transformed lines and four nontransformed lines. Ten dry mature T1 seeds from each transformant and nontransformant were individually weighed and wet-ashed with HNO₃ and H₂O₂ solution at 110°C for 2 days. In addition, 10 T2 seeds each from a transformant and a nontransformant were separated into the endosperm, and embryo tissues and wet-ashed as described above. The dried precipitate was consequently dissolved in 2 ml of 1N HCl solution. The iron content was then measured by absorbance of Fe-O-phenanthroline at 510 nm, and calculated against the standard of FeSO₄ (ref. 41).

RESEARCH

Acknowledgments

We wish to express our gratitude to J. Guerin, CSIRO (Commonwealth Scientific and Industrial Research Organization) for critically reading this paper and D. Becker (Max-Planck Institute) for providing the binary vector pGPTV-bar. We also thank Ms. Kusaka, Ms. Inoue, Ms. Miyake, and Ms. Tsunokawa (CRIEPI) for their technical assistance. This research was supported, in part, by research grants from the Enhancement Center-of-Excellence, the Special Coordination Funds for promoting the Science and Technology and from Bio-oriented Technology Research Advancement Institute (PRO-BRAIN) to F.T.

- World Health Organization (WHO). *National Strategies for Overcoming Micronutrient Malnutrition*. Document A45/3. (WHO, Geneva, Switzerland, 1992).
- Craig, W.J. Iron status of vegetarians. *Am. J. Clin. Nutr.* **59**, 1233S–1237S (1994).
- Yip, R. Iron deficiency, contemporary scientific issues and international programmatic approaches. *J. Nutr.* **124**, 1479S–1490S (1994).
- Anonymous. In *Standard Tables of Food Composition* (ed. Investigation Committee of Food Resources, Science and Technology Agency, Japan, 1992).
- Gillooly, M. et al. The effect of organic acids, phytates and polyphenols on the absorption of iron from vegetables. *Br. J. Nutr.* **49**, 331–342 (1983).
- Inoue, K., Tashima, K., Sanada, K. & Yokota, H. Production of iron-enriched leaf vegetables using a soaking method. *Jap. J. Soil Sci. Plant Nutr.* **66**, 527–534 (1995).
- Theil, E.C. Ferritin: structure, gene regulation, and cellular function in animals, plants and microorganisms. *Annu. Rev. Biochem.* **56**, 289–315 (1987).
- Andrews, S. et al. Structure, function and evolution of ferritins. *J. Inorg. Chem.* **47**, 161–174 (1992).
- Ragland, M. et al. Evidence for a conservation of ferritin sequences among plants and animals and for a transit peptide in soybean. *J. Biol. Chem.* **265**, 18339–18344 (1990).
- Spence, M.J., Henzl, M.T. & Lammers, P.J. The structure of a *Phaseolus vulgaris* cDNA encoding the iron storage protein ferritin. *Plant Mol. Biol.* **17**, 499–504 (1991).
- Lobréaux, S., Yewdall, S., Briat, J.F. & Harrison, P.M. Amino-acid sequence and predicted three-dimensional structure of pea seed (*Pisum sativum*) ferritin. *Biochem. J.* **288**, 931–939 (1992).
- Lobréaux, S., Massenet, O. & Briat, J.F. Iron induces ferritin synthesis in maize plantlets. *Plant Mol. Biol.* **19**, 563–575 (1992).
- Theil, E.C. Regulation of ferritin and transferrin receptor mRNAs. *J. Biol. Chem.* **265**, 4771–4774 (1990).
- Van der Mark, F., Bienfait, F. & van den Ende, H. Variable amounts of translatable ferritin mRNA in bean leaves with various iron contents. *Biochem. Biophys. Res. Commun.* **115**, 463–469 (1983).
- Lescure, A.M. et al. Ferritin gene transcription is regulated by iron in soybean cell cultures. *Proc. Natl. Acad. Sci. USA* **88**, 8222–8226 (1991).
- Lobréaux, S. & Briat, J.F. Ferritin accumulation and degradation in different organs of pea (*Pisum sativum*) during development. *Biochem. J.* **274**, 601–606 (1991).
- Ragland, M. & Theil, E.C. Ferritin (mRNA, protein) and iron concentration during soybean nodule development. *Plant. Mol. Biol.* **21**, 555–560 (1993).
- Theil, E.C., Burton, J.W. & Beard J.L. A sustainable solution for dietary iron deficiency through plant biotechnology and breeding to increase seed ferritin control. *Eur. J. Clin. Nutr.* **51**, S28–S31 (1997).
- Beard, J.L., Burton, J.W. & Theil E.C. Purified ferritin and soybean meal can be sources of iron for treating iron deficiency in rats. *J. Nutr.* **126**, 154–160 (1996).
- Takaiwa, F., Oono, K., Wing, D. & Kato, A. Sequences of three members and expression of a new major subfamily of glutelin gene from rice. *Plant Mol. Biol.* **17**, 875–885 (1991).
- Proudhon, D., Briat, J.F. & Lescure, A.M. Iron induction of ferritin synthesis in soybean cell suspensions. *Plant Physiol.* **90**, 586–590 (1989).
- Kimata, Y. & Theil, E.C. Posttranscriptional regulation of ferritin during nodule development in soybean. *Plant Physiol.* **104**, 263–270 (1994).
- Van Wuytswinkel, O., Savino, G. & Briat, J.F. Purification and characterization of recombinant pea-seed ferritins expressed in *Escherichia coli*: influence of N-terminus deletions on protein solubility and core formation in vitro. *Biochem. J.* **305**, 253–261 (1995).
- Van Wuytswinkel, O. & Briat, J.F. Conformational changes and in vitro core-formation modifications induced by site-directed mutagenesis of the specific N-terminus of pea seed ferritin. *Biochem. J.* **305**, 959–965 (1995).
- Sturm, A., Voelker, T.A., Herman, E.M. & Chrispeels, M.J. Correct glycosylation, golgi-processing, and targeting to protein bodies of the vacuolar protein phyto-nemagglutinin in transgenic tobacco. *Planta*. **175**, 170–183 (1988).
- Ohtani, T., Wallace, J.C., Thompson, G.A., Gallili, G. & Larkins, B.A. Normal and lysine containing zeins are unstable in transgenic tobaccoseeds. *Plant Mol. Biol.* **16**, 117–128 (1990).
- Takaiwa, F. et al. High level accumulation of soybean glycinin in vacuole-derived protein bodies in the endosperm tissue of transgenic tobacco seed. *Plant Sci.* **111**, 39–49 (1995).
- Takaiwa, F. et al. Characterization of common cis-regulatory elements responsible for the endosperm-specific expression of members of the rice glutelin multi-gene family. *Plant Mol. Biol.* **30**, 1207–1221 (1996).
- Wu, C.-Y., Suzuki, A., Washida, H. & Takaiwa, F. The GCN4 motif in a rice glutelin gene is essential for endosperm-specific expression and is activated by Opaque-2 in transgenic rice plants. *Plant J.* **14**, 673–683 (1998).
- Utsumi, S. et al. Synthesis, processing and accumulation of modified glycinins of soybean in the seeds, leaves and stems of transgenic tobacco. *Plant Sci.* **92**, 191–202 (1993).
- Shirsat, A.H., Wilford, N. & Croy, R.R.D. 1989. Gene copy number and levels of expression in transgenic plants of a seed specific gene. *Plant Sci.* **61**, 75–80.
- Goto, F., Yoshihara, T. & Saiki, H. Iron accumulation in tobacco plants expressing soybean ferritin gene. *Transgenic Res.* **7**, 173–180 (1998).
- Delhaize, E. A metal-accumulator mutant of *Arabidopsis thaliana*. *Plant Physiol.* **111**, 849–855 (1996).
- Becker, D., Kemper, E., Schell, J. & Masterson, R. New plant binary vectors with selectable markers located proximal to the left T-DNA border. *Plant Mol. Biol.* **20**, 1195–1197 (1992).
- Chu, C., Wang, C. & Sun, C. Establishment of an efficient medium for anther culture of rice through comparative experiment on the nitrogen sources. *Scientia Sinica* **18**: 659–668 (1975).
- Murashige, T. & Skoog, F. A revised medium for rapid growth and bio assays with tobacco tissue culture. *Physiol. Plantarum*. **15**, 473–497 (1962).
- Shure, M., Wessler, S. & Fedoroff, N. Molecular identification and isolation of *Waxy* locus in maize. *Cell* **35**, 225–233 (1983).
- Enger-Blum, G., Meier, M., Frank, J. & Müller, G.A. Reduction of background problems in non-radioactive northern and Southern blot analyses enables higher sensitivity than ³²P-based hybridizations. *Anal. Biochem.* **210**, 235–244 (1993).
- Bradford, M.M. A rapid and sensitive method for the quantification of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* **72**, 248–254 (1976).
- Sambrook, J., Fritsch, E.F. & Maniatis, T. In *Molecular Cloning* 2nd ed. (eds. Ford, N. & Nolan, C.) 18.67–75. (Cold Spring Harbor Laboratory Press, New York, 1989).
- Kamachi, K., Yamaya, T., Hayakawa, T., Mae, T. & Ojima, K. Vascular bundle-specific localization of cytosolic glutamine synthetase in rice leaves. *Plant Physiol.* **99**, 1481–1486 (1992).
- Anonymous. In *Standard Methods for the Examination of Water and Wastewater*. 18th edn (eds Greenberg, A.E., Clesceri, L.S., & Eaton, A.D.) 3-66–3-68 (American Public Health Association, Washington, DC, 1992).