JAFCAU
Case No.: IPR2023-00208
U.S. Patent No. 11,013,250

JANUARY 1996
VOLUME 44
NUMBER 1

Journal of Agricultural and Food Chemistry

Published by the
American Chemical Society

Motif Exhibit 1051, Page 1 of 12

EDITOR: IRVIN E. LIENER

Associate Editors: G. Wayne Ivie
Marshall Phillips

Manuscript Office: Marie Crunkleton Wiggins, Manager; Bryan D. Tweedy, Senior Associate Editor

ADVISORY BOARD

Terry E. Acree
Eric Block
Daniel R. Caldwell
John W. Finley
Paul Hedin
Chi-Tang Ho
R. McGorrin
Steven Nagy
R. Petroski
Russ Rousseff
Richard Scanlan
Walter Wolf

Journal of Agricultural and Food Chemistry (ISSN 0021-8561) is published monthly by the American Chemical Society at 1155 16th St., N.W., Washington, DC 20036. Second-class postage paid at Washington, DC, and additional mailing offices. POSTMASTER: Send address changes to *Journal of Agricultural and Food Chemistry*, Member & Subscriber Services, P.O. Box 3337, Columbus, OH 43210. Canadian GST Reg. No. 127571347. Printed in the USA.

PUBLICATIONS DIVISION

Robert H. Marks, Director

Journal Publishing Operations: Charles R. Bertsch, Director

Editorial Office: Mary E. Scanlan, Manager; Anne C. O'Melia, Journals Editing Manager; Eileen M. Richardson, Associate Editor

Advertising Office: Centcom, Ltd., 1599 Post Road East, P.O. Box 231, Westport, CT 06881.

© Copyright 1996 American Chemical Society.
Copyright permission: Reprographic copying beyond that permitted by Section 107 or 108 of the U.S. Copyright Act is allowed *for internal use only*, provided $12.00 per article is paid directly to the Copyright Clearance Center (CCC), 222 Rosewood Dr., Danvers, MA 01923, USA. Republication or reproduction for sale of articles or abstracts in this journal is permitted only under license from ACS. Direct these and other permission requests to the ACS Copyright Office at the ACS Washington address. For quotes and information on ordering bulk reprints, please call (202) 872-4539 or write to Customer Service and Sales at the ACS Washington address.

Journal of Agricultural and Food Chemistry

Editorial Information

Guide for Authors and copyright status form appear in the first issue of each volume. These instructions are available via the World Wide Web by selecting Instructions for Authors and Editors at URL http://pubs.acs.org.

Manuscripts for publication should be submitted (six copies) to Manuscript Office, *Journal of Agricultural and Food Chemistry*, at the ACS Washington address.

Correspondence regarding **accepted papers and proofs** should be directed to Journal Publishing Operations, at the address below.

Page charges of $40.00 per page may be paid for papers published in this journal. Payment does not affect acceptance or scheduling of papers.

The American Chemical Society and its Editors assume no responsibility for the statements and opinions advanced by contributors.

Registered names and trademarks, etc., used in this publication, even without specific indication thereof, are not to be considered unprotected by law.

At the end of each document is a unique 9-character document number that serves as a link between the printed and electronic (CJACS Plus Images) products and facilitates the retrieval of the document in electronic form.

1996 Subscription and Ordering Information

Members may share/donate their personal subscriptions with/to libraries and the like but only after 5 years from publication.

	U.S.	Canada and Mexico	Europe*	All Other Countries*
Printed				
Members—1 yr	$ 37	$ 57	$ 85	$ 96
Nonmembers	$483	$503	$531	$542
Microfiche				
Members—1 yr	$ 37	$ 37	$ 49	$ 49
Nonmembers	$483	$483	$495	$495
Supporting information (microfiche)	$ 17	$ 23	$ 23	$ 23

* Air service included.

New and renewal subscriptions should be sent with payment to American Chemical Society, Department L-0011, Columbus, OH 43268-0011. Subscriptions are available only on a calendar year basis.

Nonmember subscribers in Japan must enter subscription orders with Maruzen Company Ltd., 3-10 Nihonbashi 2-chome, Chuo-ku, Tokyo 103, Japan. Tel: (03) 272-7211.

Microform and Back Issue orders should be sent to the Microforms & Back Issues Office at the ACS Washington address.

Claims for issues not received will be honored only if submitted within 90 days of the issue date (subscribers in North America) or within 180 days of the issue date (all other subscribers). Claims are handled by Member & Subscriber Services.

Supporting Information is noted on the table of contents with a ■. It is available as photocopy ($12.00 for up to 3 pages and $1.50 per page for each additional page, plus $2.00 for foreign postage) or as 24× microfiche ($12.00, plus $1.00 for foreign postage). Canadian residents should add 7% GST. See Supporting Information notice at the end of journal article for number of pages. Orders *must* give complete title of article, names of authors, journal, issue date, and page numbers. Prepayment is required and prices are subject to change. For information on microforms, contact the Microforms & Back Issues Office at the ACS Washington address or phone (202) 872-4554. Supporting information except structure factors also appears in the microfilm edition.

Supporting Information is also available via the Internet using either World Wide Web or gopher protocols. WWW users should select Electronic Supporting Information at URL http://pubs.acs.org. A README document is available for gopher users in the directory Supporting Information under ACS Publications at pubs.acs.org. Detailed instructions for using this service, along with a description of the file formats, are available at these sites. To download the Supporting Information, enter the journal subscription number from your mailing label. For additional information on electronic access, send electronic mail to gopher@acsinfo.acs.org or phone (202) 872-4434.

American Chemical Society
1155 16th St., N.W.
Washington, DC 20036
(202) 872-4600
TDD (202) 872-8733
FAX (202) 872-4615

Journal Publishing Operations
American Chemical Society
2540 Olentangy River Road
P.O. Box 3330
Columbus, OH 43210
(614) 447-3600, Ext. 3171
TELEX 6842086; FAX (614) 447-3745

Member & Subscriber Services
American Chemical Society
P.O. Box 3337
Columbus, OH 43210
(614) 447-3776; FAX (614) 447-3671
(800) 333-9511

Case No.: IPR2023-00208
U.S. Patent No. 11,013,250

STEENBOCK LIBRARY
FEB 06 1996

JAFCAU 44(1) 1–394 (1996)
ISSN 0021-8561

Registered in U.S. Patent and Trademark Office
Copyright 1996 by the American Chemical Society

JANUARY 1996

Journal of Agricultural and Food Chemistry®

SHORT COMMUNICATIONS

1 Tetrodotoxin Concentrations in Cultured Puffer Fish, *Fugu rubripes*
Kendo Matsumura

3 Nuisance Proteins of Wine Are Grape Pathogenesis-Related Proteins
Elizabeth J. Waters,* Neil J. Shirley, and Patrick J. Williams

REVIEWS

6 Nutritional Value of Proteins from Different Food Sources. A Review
Mendel Friedman

FOOD CHEMISTRY/BIOCHEMISTRY

30 Effects of Postharvest Dips in Calcium Chloride on Strawberry
José M. García,* Salvador Hérrera, and Ana Morilla

34 Identification of Isorhamnetin 4′-Glucoside in Onions
Yang-Kyun Park and Chang Y. Lee*

37 Varietal Differences in the Phenolic Content and Superoxide Radical Scavenging Potential of Wines from Different Sources
Michikatsu Sato, Narasimhan Ramarathnam,* Yumiko Suzuki, Tetsuro Ohkubo, Masao Takeuchi, and Hirotomo Ochi

42 Caseinolytic Specificity of Cardosin, an Aspartic Protease from the Cardoon *Cynara cardunculus* L.: Action on Bovine α_s- and β-Casein and Comparison with Chymosin
I. Queiroz Macedo,* Carlos J. Faro, and Euclides M. Pires

48 Comparison of Hydration Behavior of Bovine and Caprine Caseins As Determined by Oxygen-17 Nuclear Magnetic Resonance: Temperature Dependence of Colloidal Stability
Adela Mora-Gutierrez,* Harold M. Farrell, Jr., and Thomas F. Kumosinski

54 Rheological Properties of Thiolated and Succinylated Caseins
E. D. Strange,* V. H. Holsinger, and D. H. Kleyn

59 Competitive Adsorption between Dioleoylphosphatidylcholine and Sodium Caseinate on Oil–Water Interfaces
Yuan Fang and Douglas G. Dalgleish*

65 Studies on Vitamin E and Meat Quality. 1. Effect of Feeding High Vitamin E Levels on Time-Related Pork Quality
P. Dirinck,* A. De Winne, M. Casteels, and M. Frigg

Case No.: IPR2023-00208
U.S. Patent No. 11,013,250

69 Enzymatic Phosphorylation by a Casein Kinase II of Native and Succinylated Soy Storage Proteins Glycinin and β-Conglycinin
Marie-Christine Ralet,* Dominique Fouques, Thierry Chardot, and Jean-Claude Meunier

76 Fractionation of Peptides in Protease Digests of Proteins by Preparative Isoelectric Focusing in the Absence of Added Ampholyte: A Biocompatible and Low-Cost Approach Referred to as Autofocusing
Maki Yata, Kenji Sato,* Kozo Ohtsuki, and Makoto Kawabata

80 Nature of Polyphenol–Protein Interactions
Karl J. Siebert,* Nataliia V. Troukhanova, and Penelope Y. Lynn

86 Rheology and Microstructure of β-Lactoglobulin/Sodium Polypectate Gels
Embola E. Ndi, Barry G. Swanson, Gustavo V. Barbosa-Canovas, and Lloyd O. Luedecke*

93 Denaturant-Induced Conformations of Globulins
Shela Gorinstein,* Marina Zemser, Michael Friedman, Nora L. Vasco-Méndez, and Octavio Paredes-López

100 Structural Stability of Globulins
Shela Gorinstein,* Marina Zemser, and Octavio Paredes-López

106 Mechanism for Stabilization of Fish Actomyosin by Sodium Lactate
Grant A. MacDonald, Tyre C. Lanier,* Harold E. Swaisgood, and Donald D. Hamann

113 Interaction of Sucrose and Zinc for Cryoprotection of Surimi
Grant A. MacDonald, Tyre C. Lanier,* and Francis G. Giesbrecht

119 Inhibition of Protein and Lipid Oxidation in Beef Heart Surimi-like Material by Antioxidants and Combinations of pH, NaCl, and Buffer Type in the Washing Media
Subramanian Srinivasan, Youling L. Xiong,* and Eric A. Decker

126 Wild Rice Hull Antioxidants
Abdulwahab M. Asamarai, Paul B. Addis,* Richard J. Epley, and Thomas P. Krick

131 Antioxidant Activity of a Rosemary Extract and Its Constituents, Carnosic Acid, Carnosol, and Rosmarinic Acid, in Bulk Oil and Oil-in-Water Emulsion
Edwin N. Frankel,* Shu-Wen Huang, Robert Aeschbach, and Elizabeth Prior

COMPOSITION OF FOODS/FEEDS

136 Neutral Sugar Contents of Corn Gluten Meal and Corn Gluten Feed
Y. Victor Wu

139 Characterization of Citrus Honey by Deuterium NMR
Pinhas Lindner,* Elisha Bermann, and Biana Gamarnik

ANALYTICAL METHODS

141 Selective Recognition of the Herbicide Atrazine by Noncovalent Molecularly Imprinted Polymers
Martin Siemann, Lars I. Andersson, and Klaus Mosbach*

146 HPLC Method for Detection of Ergotamine, Ergosine, and Ergine after Intravenous Injection of a Single Dose
Ali S. Moubarak,* Edgar L. Piper, Zelpha B. Johnson, and Miroslav Flieger

149 Purification and Quantitative Analysis of Veratridine and Cevadine by HPLC
J. Daniel Hare

153 High-Performance Liquid Chromatographic Method for the Determination of Blasticidin S in Formulated Products with Photodiode Array Detection
Chi-Chu Lo,* Yuh-Jyuan Lee, and Chai-Ju Chang

159 Determination of Fumonisins in Corn: Evaluation of Competitive Immunoassay and HPLC Techniques
Eric W. Sydenham,* Gordon S. Shephard, Pieter G. Thiel, Charles Bird, and Brinton M. Miller

165 Flow Injection System with Multisite Detection for Spectrophotometric Determination of Calcium and Magnesium in Soil Extracts and Natural Waters
Ana Rita A. Nogueira, Sandra M. B. Brienza, Elias A. G. Zagatto, José L. F. C. Lima,* and Alberto N. Araújo

170 Discrimination of *Arabica* and *Robusta* in Instant Coffee by Fourier Transform Infrared Spectroscopy and Chemometrics
Romain Briandet, E. Katherine Kemsley,* and Reginald H. Wilson

175 Detection of Adulteration in Orange Juices by a New Screening Method Using Proton NMR Spectroscopy in Combination with Pattern Recognition Techniques
Jack T. W. E. Vogels,* Loes Terwel, Albert C. Tas, Frans van den Berg, Fred Dukel, and Jan van der Greef

181 Determination of Formetanate Hydrochloride in Strawberries
Renwei Hu, Valérie Petay, and Josette Fournier*

185 Determination of Safranal from Saffron (*Crocus sativus* L.) by Thermal Desorption−Gas Chromatography
Gonzalo L. Alonso, M. Rosario Salinas,* Francisco J. Esteban-Infantes, and M. Angeles Sánchez-Fernández

189 Determination of Zinc Phosphide Residues in the California Ground Squirrel *(Spermophilus beecheyi)* by Gas Chromatography−Flame Photometric Detection
Richard E. Mauldin,* David A. Goldade, Richard M. Engeman, Margaret J. Goodall, Rori K. Craver, and John J. Johnston

NUTRITION/TOXICOLOGY

195 Preparative LC Isolation and Purification of Fumonisin B_1 from Rice Culture
Filmore I. Meredith,* Charles W. Bacon, Ronald D. Plattner, and William P. Norred

199 Effect of a Daily Supplementation of Polyethylene Glycol on Intake and Digestion of Tannin-Containing Leaves (*Quercus calliprinos*, *Pistacia lentiscus*, and *Ceratonia siliqua*) by Goats
N. Silanikove,* N. Gilboa, I. Nir, A. Perevolotsky, and Z. Nitsan

FLAVORS AND AROMAS

206 Volatile Constituents from the Flowers of Japanese Honeysuckle (*Lonicera japonica*)
William S. Schlotzhauer,* Sam D. Pair, and Robert J. Horvat

210 Chemical Changes during the Development and Ripening of the Fruit of *Cucumis melo* (Cv. Makdimon)
Youming Wang, S. Grant Wyllie,* and David N. Leach

217 Sesquiterpenic Epoxides and Alcohols Derived from Hydrocarbons of Vetiver Essential Oil
Isabelle Bombarda, Emile M. Gaydou,* Jacqueline Smadja, and Robert Faure

223 Pyrolysis/GC/MS Analysis of *N*-(1-Deoxy-D-fructos-1-yl)-L-phenylalanine: Identification of Novel Pyridine and Naphthalene Derivatives
Anahita Keyhani and Varoujan A. Yaylayan*

230 Isolation and Identification of Precursor of 4-Hydroxy-2(or 5)-ethyl-5(or 2)-methyl-3(2*H*)-furanone from Isolated Soybean Protein and Shoyu
Masaoki Sasaki

236 DETBA Value and Hexanal Production with the Combination of Unsaturated Fatty Acids and Extracts Prepared from Soybean Seeds Lacking Two or Three Lipoxygenase Isozymes
Shu Furuta,* Yoichi Nishiba, Makita Hajika, Kazunori Igita, and Ikuo Suda

240 Mechanistic Studies of Tetramethylpyrazine Formation under Weak Acidic Conditions and High Hydrostatic Pressure
Tzou-Chi Huang, Hui-Yin Fu, and Chi-Tang Ho*

247 Volatile Compounds Produced by the Reaction of Leucine and Valine with Glucose in Propylene Glycol
Mohd Wahid Samsudin,* Sun Rongtao, and Ikram M. Said

251 Model Studies on the Oxidative Stability of Odor-Active Thiols Occurring in Food Flavors
Thomas Hofmann, Peter Schieberle, and Werner Grosch*

CHEMICAL CHANGES DURING PROCESSING/STORAGE

256 Potent Bacterial Mutagens Produced by Chlorination of Simulated Poultry Chiller Water
William F. Haddon,* Ronald G. Binder, Rosalind Y. Wong, Leslie A. Harden, Robert E. Wilson, Mabry Benson, and Kenneth L. Stevens

Case No.: IPR2023-00208
U.S. Patent No. 11,013,250

264 Influence of Storage Temperature on Fruit Ripening and Olive Oil Quality
José M. García,* Francisca Gutiérrez, José M. Castellano, Salud Perdiguero, Ana Morilla, and Miguel A. Albi

268 Volatiles in Distillates of Cider Aged in American Oak Wood
Juan Mangas,* Roberto Rodríguez, Javier Moreno, and Domingo Blanco

274 A New Benzochromanone Derivative from Cape Aloe
Giovanna Speranza,* Antonella Di Meo, Paolo Manitto, Diego Monti, and Gabriele Fontana

278 Effect of Long-Term Storage on Cell Wall Neutral Sugars and Galacturonic Acid of Two Sweetpotato Cultivars
William M. Walter, Jr.,* and Cheryl S. Palma

282 On the Role of 2,3-Dihydro-3,5-dihydroxy-6-methyl-4(*H*)-pyran-4-one in the Maillard Reaction
Myong-Ock Kim and Werner Baltes*

CHEMISTRY OF CROP AND ANIMAL PROTECTION

290 Electrochemical Properties of the Herbicide Cacalol and Its Derivatives in Protic and Aprotic Solvents by Using Cyclic Voltammetry. Correlation With Hill's Reaction Activities
Martha Aguilar-Martínez, Manuel Jiménez-Estrada, Norma A. Macías-Ruvalcaba, and Blas Lotina-Hennsen*

296 Insect Antifeedant Ryanodane Diterpenes from *Persea indica*
Azucena González-Coloma, David Terrero, Aurea Perales, Pierre Escoubas, and Braulio M. Fraga*

301 Antifungal Activity against Food-Borne Fungi of *Aspidistra elatior* Blume
Mamoru Koketsu,* Mujo Kim, and Takehiko Yamamoto

304 Isolation and Identification of Residues of 4″-(*epi*-Methylamino)-4″-deoxyavermectin B_{1a} Benzoate from the Surface of Cabbage
Christopher L. Wrzesinski,* Byron H. Arison, Jack Smith, Deborah L. Zink, William J. A. VandenHeuvel, and Louis S. Crouch

313 Microbial Transformations of Clomazone
Song Yu Liu, Mark Shocken, and John P. N. Rosazza*

320 Synthesis and Cytokinin Activity of Two Fluoro Derivatives of N^6-Isopentenyladenine
Denis Clemenceau, Jack Cousseau, Valérie Martin, Huguette Molines, Claude Wakselman, René Mornet,* Fabien Nogue, and Michel Laloue

ENVIRONMENTAL CHEMISTRY

324 Products and Kinetics of Cloransulam-methyl Aerobic Soil Metabolism
Jeffrey D. Wolt,* Joelene K. Smith, Jerry K. Sims, and Dennis O. Duebelbeis

333 Hydrolysis Kinetics of Thifensulfuron Methyl in Aqueous Buffer Solutions
Jean-Pierre Cambon* and Jean Bastide

BIOTECHNOLOGY

338 Synthesis of Oleic Acid Esters Catalyzed by Immobilized Lipase
Maja Habulin, Vlasta Krmelj, and Željko Knez*

343 Quantitative Analysis of 2,4-Dichlorophenoxyacetic Acid in Water Samples by Two Immunosensing Methods
Christine Wittmann,* Frank F. Bier, Sergei A. Eremin, and Rolf D. Schmid

351 Enzymatic Approach to Waste Minimization in a Cattle Dipping Operation: Economic Analysis
Kenneth J. Grice, Gregory F. Payne,* and Jeffrey S. Karns

358 Study of Putative Glycosylation Sites in Bovine β-Casein Introduced by PCR-Based Site-Directed Mutagenesis
Byung-Kwon Choi and Rafael Jiménez-Flores*

365 The Composition of Insect-Protected Cottonseed Is Equivalent to That of Conventional Cottonseed
Sharon A. Berberich,* Joel E. Ream, Teri L. Jackson, Randall Wood, Robert Stipanovic, Patricia Harvey, Shari Patzer, and Roy L. Fuchs

372 Production of a Sensitive Monoclonal Antibody to Sterigmatocystin and Its Application to ELISA of Wheat
Suzhen Li, Pu Yan Chen, Ronald R. Marquardt,* Zhengkang Han, and James R. Clarke

376 Development and Application of a Rapid Immunoassay for Difenzoquat in Wheat and Barley Products
Jupiter M. Yeung,* Richard D. Mortimer, and Peter G. Collins

381 Purification and Biochemical Comparison of 1-Aminocyclopropane-1-carboxylic Acid Deaminase Proteins Expressed in Delayed Ripening Tomato and *Escherichia coli*: Studies for a Food Safety Assessment
Rory F. Finn, Richard M. Leimgruber, Denis M. Boyle, Michael G. Jennings, Nancy M. Kimack, Christine E. Smith, Bruce F. Bishop, Ronald B. Frazier, Kimberly M. Magin, Roy L. Fuchs, and Andrew J. Reed*

388 Safety Assessment of 1-Aminocyclopropane-1-carboxylic Acid Deaminase Protein Expressed in Delayed Ripening Tomatoes
Andrew J. Reed,* Keith A. Kretzmer, Mark W. Naylor, Rory F. Finn, Kimberly M. Magin, Bruce G. Hammond, Richard M. Leimgruber, Stephen G. Rogers, and Roy L. Fuchs

There is no supporting information for this issue.

* In papers with more than one author, the asterisk indicates the name of the author to whom inquiries about the paper should be addressed.

This material may be protected by Copyright law (Title 17 U.S. Code)

J. Agric. Food Chem. 1996, 44, 251–255

Model Studies on the Oxidative Stability of Odor-Active Thiols Occurring in Food Flavors

Thomas Hofmann,† Peter Schieberle,‡ and Werner Grosch*,†

Deutsche Forschungsanstalt für Lebensmittelchemie and Institut für Lebensmittelchemie der Technischen Universität München, Lichtenbergstrasse 4, D-85748 Garching, Germany

Dilute ethereal solutions of mercaptoacetone (MA), 2-mercapto-3-butanone (MB), 3-mercapto-2-pentanone (MP), 2-methyl-3-furanthiol (MFT), and 2-furfurylthiol (FFT), either alone or in pairs, were stored at 6 °C. During a period of 10 days the oxidation of the thiols to the corresponding disulfides or mixed disulfides was followed by high-resolution gas chromatography/mass spectrometry. In diethyl ether MFT showed the highest oxidation rate, which decreased in the order MFT ≫ MA > MB = MP = FFT. After 1 day, mixed disulfides (e.g. MFT–FFT, FFT–MB) were detectable when pairs of the five thiols dissolved in diethyl ether were stored at 6 °C. MFT oxidized more slowly in dichloromethane than in diethyl ether, and it was stable in pentane. However, a heat treatment increased the oxidation rate of MFT dissolved in dichloromethane or pentane.

Keywords: *2-Methyl-3-furanthiol; 2-furfurylthiol; disulfide formation*

INTRODUCTION

The thiols shown in Figure 1 are formed by the Maillard reaction of cysteine and might contribute to the flavors of heated foods and beverages, due to their low odor thresholds (Whitfield et al., 1992). Actually, 2-methyl-3-furanthiol (MFT) and 2-furfurylthiol (FFT) have been detected as character impact odorants of roasted coffee (Tressl, 1981, 1989, 1990; Tressl et al., 1983; Holscher et al., 1990; Blank et al., 1992), cooked meat (Gasser and Grosch, 1988, 1990a; Farmer and Patterson, 1991; Grosch and Zeiler-Hilgart, 1992), and wheat bread (Baltes and Song, 1994).

It is well-known that thiols oxidize easily to disulfides. For this reason it is not surprising that the 15 disulfides or mixed disulfides (Table 1) resulting from the oxidation of the 5 thiols listed in Figure 1 have been identified in commercial meat flavorings (Gasser and Grosch, 1990b; Ruther and Baltes, 1994a,b) as well as in model Maillard reaction systems (Tressl et al., 1983, 1990; Hartmann et al., 1984; Reineccius and Liardon, 1985; Silwar and Tressl, 1989; Farmer and Mottram, 1990; Güntert et al., 1990, 1992a,b; Werkhoff et al., 1990; Zhang and Wo, 1991; Farmer and Patterson, 1991; Whitfield et al., 1992; Mottram and Whitfield, 1994; Hofmann and Schieberle, 1995). FFT–FFT and MFT–MFT (nomenclature in Table 1) were found in cooked meat (Gasser and Grosch, 1988, 1990a; Farmer and Patterson, 1991; Grosch and Zeiler-Hilgart, 1992), and FFT–FFT was found in roasted coffee (Tressl, 1981) as well as in wheat bread (Baltes and Song, 1994). The contribution of MFT–MFT to the flavor of roasted coffee (Blank et al., 1992) and black tea (Guth and Grosch, 1993) has been suggested on the basis of aroma extract dilution analysis.

Furthermore, it has been revealed recently (Eiserich and Shibamoto, 1994) that the rapid oxidation of MFT to its disulfide and to other products contributes to the antioxidative activity of Maillard reaction products.

* Author to whom correspondence should be addressed (fax +49-89-3209-4183).

† Deutsche Forschungsanstalt für Lebensmittelchemie.

‡ Institut für Lebensmittelchemie.



Figure 1. Chemical structures of the five thiols used in the model experiments: mercaptoacetone (MA), 2-mercapto-3-butanone (MB), 3-mercapto-2-pentanone (MP), 2-methyl-3-furanthiol (MFT), and 2-furfurylthiol (FFT).

Table 1. Nomenclature of the Disulfides and Mixed Disulfides

disulfides	
MA–MA	bis(2-oxopropyl) disulfide
MB–MB	bis(3-oxo-2-butyl) disulfide
MP–MP	bis(2-oxo-3-pentyl) disulfide
MFT–MFT	bis(2-methyl-3-furyl) disulfide
FFT–FFT	bis(2-furfuryl) disulfide
mixed disulfides	
MA–MB	2-oxopropyl 3-oxo-2-butyl disulfide
MA–MP	2-oxopropyl 2-oxo-3-pentyl disulfide
MP–MB	2-oxo-3-pentyl 3-oxo-2-butyl disulfide
MFT–MA	2-methyl-3-furyl 2-oxo-propyl disulfide
MFT–MB	2-methyl-3-furyl 3-oxo-2-butyl disulfide
MFT–MP	2-methyl-3-furyl 2-oxo-3-pentyl disulfide
FFT–MA	2-furfuryl 2-oxo-propyl disulfide
FFT–MB	2-furfuryl 3-oxo-2-butyl disulfide
FFT–MP	2-furfuryl 2-oxo-3-pentyl disulfide
FFT–MFT	2-furfuryl 2-methyl-3-furyl disulfide

In the studies mentioned above, the volatiles have been extracted with a solvent either in the course of simultaneous distillation and extraction or from an adsorbent in the course of dynamic headspace analysis. On the basis of the sensitivity of the thiols toward oxidation, the question arises whether the disulfides identified might be partially formed during storage and/or preparation of concentrated extracts for GC/MS analysis.

To check this source of artifact formation, the thiols shown in Figure 1 were dissolved in diethyl ether; MFT was also dissolved in dichloromethane and *n*-pentane, and their oxidation to the corresponding disulfides or

0021-8561/96/1444-0251$12.00/0 © 1996 American Chemical Society

Table 2. Mass Spectral Data (MS-EI) and Retention Indices (RI) of Disulfides and Mixed Disulfides

compound[a]	retention indices FFAP	retention indices SE-54	m/z (relative abundance in %) mol ion	predominant fragment ions
MA–MA	2347	1405	178 (10)	43 (100), 57 (9), 135 (6), 121 (4), 90 (3)
MB–MB	2204	1460	206 (25)	43 (100), 61 (28), 59 (24), 72 (20), 71 (19), 103 (19), 104 (18), 163 (16), 119 (14)
MP–MP	2247	1594	234 (25)	43 (100), 85 (37), 41 (30), 57 (24), 73 (24), 55 (21), 71 (18), 105 (13)
MFT–MFT	2100	1526	226 (100)	113 (95), 114 (40), 51 (25), 85 (18), 45 (18), 61 (16), 72 (16)
FFT–FFT	2465	1673	226 (20)	81 (100), 53 (80), 45 (20), 82 (20), 51 (15), 85 (12)
MA–MB	2280	1436	192 (11)	43 (100), 71 (9), 192 (9), 149 (8), 57 (8), 121 (2)
MA–MP	2308	1506	206 (20)	43 (100), 73 (26), 72 (16), 57 (14), 74 (12), 163 (11), 55 (11), 133 (10), 105 (10)
MP–MB	2227	1532	220 (64)	43 (100), 73 (56), 71 (42), 41 (41), 59 (35), 177 (32), 85 (32), 75 (30), 45 (24)
MFT–MA	2200	1462	202 (29)	43 (100), 113 (65), 114 (31), 44 (22), 51 (10), 81 (8), 144 (5)
MFT–MB	2127	1491	216 (50)	43 (100), 113 (74), 114 (52), 59 (52), 85 (24), 173 (20), 51 (20), 86 (18)
MFT–MP	2145	1561	230 (61)	43 (100), 113 (90), 114 (50), 45 (32), 145 (29), 73 (24), 85 (21), 187 (19)
FFT–MA	2400	1540	202 (18)	53 (100), 81 (92), 43 (79), 45 (37), 82 (33), 51 (19), 85 (16)
FFT–MB	2352	1577	216 (5)	81 (100), 43 (61), 53 (48), 45 (19), 82 (15), 59 (12), 85 (9)
FFT–MP	2385	1649	230 (4)	81 (100), 43 (22), 53 (19), 82 (8), 45 (8)
MFT–FFT	2323	1624	226 (19)	81 (100), 113 (18), 53 (14), 45 (11), 114 (10), 51 (9), 43 (6)

[a] For abbreviations see Table 1.

Table 3. Oxidation of Thiols to the Corresponding Disulfides[a]

storage period (days at 6 °C)	amount in μg MFT	MFT–MFT	MA	MA–MA	MB	MB–MB	MP	MP–MP	FFT	FFT–FFT
0	80.1	0	79.3	0	80.0	0	80.2	0	79.9	0
1	63.4	16.7	73.9	5.4	79.1	1.7	78.6	1.6	78.2	1.7
2	57.8	22.2	73.4	5.9	78.4	2.4	77.9	2.3	77.7	2.2
3	53.9	26.2	72.5	6.7	77.7	3.0	77.5	2.7	77.3	2.6
6	44.1	35.1	70.3	8.9	76.3	4.4	76.4	3.6	76.3	3.6
10	37.4	42.6	69.0	10.2	75.0	5.8	75.7	4.5	75.5	4.2

[a] The thiol, dissolved in diethyl ether (500 μL), was stored at 6 °C. The thiols and disulfides are abbreviated as reported in the legend of Figure 1 and in Table 1, respectively.

mixed disulfides was measured. The solvents selected for these model experiments are frequently used in flavor analysis (cf. references cited above).

EXPERIMENTAL PROCEDURES

Chemicals. The following compounds were obtained commercially: 2-acetylpyrrole, 2-furfurylthiol, and 2-methyl-3-furanthiol (Aldrich, Steinheim, Germany).

Synthesis. The following thiols were synthesized as reported in the literature given: 1-mercapto-2-propanone and 3-mercapto-2-butanone (Hofmann and Schieberle, 1995); 3-mercapto-2-pentanone (Asinger et al., 1964).

Immediately before the start of a model experiment, each thiol was freed from disulfides: the thiol (ca. 150 μg) was dissolved in the corresponding solvent (either diethyl ether, dichloromethane, or *n*-pentane; 20 mL each) and extracted with an aqueous solution of sodium hydroxide (10 mL, 1 mol/L). The aqueous phase was then acidified to pH 3 with hydrochloric acid (1 mol/L), and the thiol was isolated by extraction with the respective solvent (total volume 15 mL). The ice-cooled solution of the thiol was concentrated to 10 mL by evaporating the solvent with a stream of nitrogen. The amount of each thiol was determined gas chromatographically with 2-acetylpyrrole as an internal standard.

Disulfides. The corresponding thiol (0.2 mmol) was dissolved in diethyl ether (15 mL) and, after addition of an aqueous solution of copper(II) sulfate (3 mol/L; 15 mL), shaken for 35 min at room temperature. After addition of ammonium acetate (2 g) and shaking, the ethereal layer was separated, washed three times with brine (total volume 60 mL), and then dried over Na_2SO_4. The mixed disulfides were synthesized by oxidation of a mixture of the corresponding two thiols (Hofmann and Schieberle, 1995). MS-EI data of the disulfides and mixed disulfides are summarized in Table 2.

Storage of the Thiols. Solutions (500 μL each) containing the thiols in the amounts given in Tables 3–5 and 2-acetylpyrrole (25 μg) as an internal standard were pipetted into brown glass vials. After the vials were sealed with a septum, they were either stored at 6 °C in a refrigerator or refluxed for 1 h.

Table 4. Oxidation of MFT to MFT–MFT during Storage in Different Solvents[a]

expt	solvent	days of storage 0	1	2	9
		MFT (μg)			
1	diethyl ether	81.5	69.7	59.5	40.4
2	diethyl ether[b]	81.5	66.0	59.3	40.6
3	dichloromethane	80.5	78.1	77.5	75.4
4	*n*-pentane	79.6	78.9	78.5	77.7
		MFT–MFT (μg)			
1	diethyl ether	0.0	21.8	27.0	40.7
2	diethyl ether[b]	0.0	15.3	21.9	40.3
3	dichloromethane	0.0	2.9	3.4	5.0
4	*n*-pentane	0.0	0.8	1.1	1.7

[a] MFT, dissolved in the respective solvent (500 μL), was stored at 6 °C. [b] Butylhydroxyanisole (22 μg) was added prior to storage.

At the times given in Tables 3–5 samples of 1.0 μL were drawn with a syringe and then analyzed by HRGC/MS.

High-Resolution Capillary Gas Chromatography (HRGC)/Mass Spectrometry (MS). HRGC was performed with a Fisons gas chromatograph (Type 5300) by using the following capillaries: FFAP (30 m × 0.32 mm; fused silica capillary, free fatty acid phase, 0.25 μm; J&W Scientific, Fisons Instruments, Mainz, Germany) and SE-54 (30 m × 0.32 mm; fused silica capillary, DB-5, 0.25 μm, J&W Scientific, Fisons Instruments). The samples (1.0 μL) were applied by the on-column injection technique at 40 °C. After 2 min, the temperature of the oven was raised at 40 °C/min to 60 °C (FFAP) or 50 °C (SE-54), held isothermally for 5 min, then raised at 6 °C/min to 230 °C, and held for 15 min. Linear retention indices (RI) of the disulfides on the capillaries FFAP and SE-54 were calculated from the retention times of *n*-alkanes (cf. Table 2). MS analysis was performed with an INCOS XL (Finnigan, Bremen, Germany). Mass spectra were generated in the electron impact mode (MS-EI) at 70 eV.

RESULTS AND DISCUSSION

In a first set of experiments, each of the five thiols shown in Figure 1 was dissolved in diethyl ether and

Case No.: IPR2023-00208
U.S. Patent No. 11,013,250

Table 5. Oxidation of Pairs of Thiols to Their Disulfides and Mixed Disulfides, Respectively

expt	reaction system[a]	amount (μg) after 1 day of storage at 6 °C		
		thiol	disulfide	mixed disulfide
1	MFT plus FFT	MFT (64.3), FFT (78.5)	MFT−MFT (14.9), FFT−FFT (0.5)	MFT−FFT (1.9)
2	MFT plus MA	MFT (68.9), MA (74.3)	MFT−MFT (8.9), MA−MA (3.4)	MFT−MA (4.5)
3	MFT plus MB	MFT (64.9), MB (78.5)	MFT−MFT (14.2), MB−MB (0.6)	MFT−MB (1.2)
4	MFT plus MP	MFT (64,6), MP (78.7)	MFT−MFT (14.5), MP−MP (0.5)	MFT−MP (2.0)
5	FFT plus MA	FFT (78.8), MA (74.6)	FFT−FFT (0.5), MA−MA (4.8)	FFA−MA (1.2)
6	FFT plus MB	FFT (78.7), MB (78.8)	FFT−FFT (0.8), MB−MB (0.8)	FFT−MB (0.8)
7	FFT plus MP	FFT (78.8), MP (78.9)	FFT−FFT (0.8), MP−MP (0.8)	FFT−MP (0.8)
8	MA plus MB	MA (74.2), MB (78.4)	MA−MA (4.8), MB−MB (0.6)	MA−MB (2.0)
9	MA plus MP	MA (74.5), MP (79.3)	MA−MA (4.9), MP−MP (0.5)	MA−MP (1.3)
10	MB plus MP	MB (78.8), MP (78.9)	MB−MB (0.8), MP−MP (0.7)	MB−MP (0.7)

[a] Pairs of thiols (80 μg each) were dissolved in diethyl ether (500 μL). The abbreviations of the thiols and disulfides refer to Figure 1 and Table 1, respectively.

then stored at 6 °C. The decrease of each thiol and the increase of the corresponding disulfide were measured over a period of 10 days.

After 10 days, a portion of each thiol had been oxidized to the correspondig disulfide (Table 3), which was identified by comparison of the MS-EI with that obtained for the synthesized reference compound.

According to the data in Table 3, MFT oxidized very rapidly: 20% of the thiol was converted into MFT−MFT after 1 day of storage and more than 50% after 10 days of storage. Compared to MFT, the four other thiols were more stable. After 10 days of storage (Table 3), their losses amounted to not more than 5 (MP) or 13% (MA).

The high oxidation rate is in agreement with the high antioxidative activity of MFT (Eiserich and Shibamoto, 1994). This activity is based on the easy abstraction of a hydrogen atom from the thiol group of MFT with the intermediate formation of a thiyl radical being stabilized by electron delocalization. As this stabilization is lacking in FFT, MA, MB, and MP, the formation of a thiyl radical is less favorable and consequently these thiols oxidize much more slowly than MFT.

A comparison of experiments 1 and 2 in Table 4 indicates that the oxidation of MFT dissolved in diethyl ether was not inhibited by the antioxidant BHT.

The influence of the solvent on the oxidation of MFT at 6 °C was studied in further experiments (Table 4). Compared to diethyl ether, storage in dichloromethane significantly lowered the oxidation rate during the storage period of 9 days. While in diethyl ether about 50% of the MFT had been oxidized to the disulfide, the decrease of MFT in dichloromethane amounted only to 6% (Table 4). In *n*-pentane MFT was most stable as no significant oxidation was observed. After that, solutions of MFT in dichloromethane and pentane were heated for 1 h under the conditions used for the simultaneous distillation and extraction according to the procedure of Nickerson and Likens (1966). The results indicated that MFT is oxidized under these conditions. In dichloromethane as well as in *n*-pentane 7.5 or 2.1 μg of MFT−MFT was formed from 80 μg of MFT.

In a third set of experiments pairs of different thiols were dissolved in diethyl ether in a weight ratio of 1:1. After storage at 6 °C, the mixed disulfides, formed by the oxidation of the two thiols present in each reaction system, were identified by comparison of the mass spectral data with those of the reference compounds (Table 2). As shown in Table 5, the mixed disulfides were present in the reaction systems in addition to the disulfides that had already been detected in the first set of experiments (Table 3).

The consumption of the thiols and the formation of the corresponding disulfides and mixed disulfides were



Figure 2. Time course of the oxidation of a mixture of MFT and FFT in diethyl ether. MFT and FFT (80 μg each)—free from disulfides—were stored in diethyl ether (500 μL) at 6 °C for 10 days. The consumption of MFT (●) and FFT (○) and the production of the mixed disulfide MFT−FFT (▼) and of the corresponding disulfides MFT−MFT (▽) and FFT−FFT (□) were determined.



Figure 3. Time course of the oxidation of a mixture of MFT and MA in diethyl ether MFT and MA (80 μg each)—free from disulfides—were dissolved in diethyl ether (500 μL) and then stored at 6 °C for 10 days. The consumption of MFT (●) and MA (○) and the production of the mixed disulfide MFT−MA (▼) and of the corresponding disulfides MFT−MFT (▽) and MA−MA (□) were determined.

quantified during the storage period of 10 days in the reaction systems MFT plus FFT (Figure 2) and MFT plus MA (Figure 3). In both mixtures the unstable MFT was oxidized to MFT−MFT. However, in the presence of the stable FFT the rate was higher than in the presence of the more reactive MA, which to some extent protected MFT like an antioxidant. The time courses shown in Figures 2 and 3 confirm the results of the first set of experiments (Table 3), indicating that the oxidation rate of the thiols was highest at the first day.

In Table 5 the concentrations of the disulfides, formed after 1 day of storage in the 10 reaction systems investigated, are compared with the amounts of the

Case No.: IPR2023-00208
U.S. Patent No. 11,013,250

unreacted thiols left. In experiments 1–4 (Table 5) the most reactive thiol, MFT, was combined with each of the resting thiols shown in Figure 1. In the presence of the relatively stable thiols FFT, MB, and MP (experiments 1, 3, and 4) the consumption of MFT at 1 day was of the same order of magnitude as in the reaction system containing MFT alone (Table 3). However, in the presence of the more reactive MA (experiment 2; Table 5) MFT was partially protected because only 14% of the MFT was oxidized to the disulfides MFT–MFT and MFT–MA.

The consumption of MA, which was next reactive compared to MFT, amounted to 6% (experiments 2, 5, 8, and 9), while the more stable thiols FFT, MB, and MP were oxidized to only 2% (Table 5).

Mixed disulfides were identified in addition to the two disulfides resulting from one thiol (Table 5). After 1 day, the largest amount of a mixed disulfide (MFT–MA; 4.5 μg) was formed in the reaction system containing MFT and MA (experiment 2; Table 5). As expected in this experiment as well as in further experiments that contained thiols differing in their reactivities (e.g. experiments 1, 3, and 4; Table 5), the amount of the mixed disulfide lay between the amounts of the disulfide of the most reactive thiol (MFT–MFT in experiments 1–4) and those of the more stable thiols (FFT–FFT, MA–MA, MB–MB, and MP–MP in experiments 1–4).

CONCLUSION

The results clearly demonstrate that during handling and storage in organic solvents oxidation of thiols may generate significant amounts of the corresponding disulfides. The data imply that during concentration and/or storage of volatile food fractions prior to HRGC analysis, especially in diethyl ether, certain amounts of disulfides are formed as artifacts. In consequence, heated foods and model systems, in which the disulfides listed in Table 2 have been identified, have to be reexamined to clarify whether these odorants actually contribute to the food flavors or whether the main portion is formed in the extract after isolation from the food. To prevent possible artifact formation during the work up procedure, the thiols can be protected by conversion into derivatives immediately after isolation, e.g. by reaction with 4-vinylpyridine (Cavins and Friedman, 1970; Guth et al., 1995).

LITERATURE CITED

Asinger, F.; Schäfer, W.; Herkelman, G.; Römings, M.; Reintges, B. D.; Scharein, G.; Wegerhoff, A. *Justus Liebigs Ann. Chem.* **1964**, *672*, 156–178 (in German).

Baltes, W.; Song, C. In *Thermally Generated Flavors: Maillard, Microwave and Extrusion Processes*; Parliment, T. H., Morello, M. J., McGorrin, R. J., Eds.; ACS Symposium Series 543; American Chemical Society: Washington, DC, 1994; pp 192–205.

Blank, I.; Sen, A.; Grosch, W. Potent odorants of the roasted powder and brew of Arabica coffee. *Z. Lebensm. Unters. Forsch.* **1992**, *195*, 239–245.

Cavins, J. F.; Friedman, M. An internal standard for amino acid analysis: *S*-β-(4-pyridylethyl)-L-cysteine. *Anal. Biochem.* **1970**, *35*, 489–493.

Eiserich, J. P.; Shibamoto, T. Antioxidative activity of volatile heterocyclic compounds. *J. Agric. Food Chem.* **1994**, *42*, 1060–1063.

Evers, W. J.; Heinsohn, H. H.; Mayers, B. J.; Sanderson, A. In *Phenolic, Sulfur and Nitrogen Compounds in Food Flavors*; Charalambous, G., Katz, I., Eds.; ACS Symposium Series 26; American Chemical Society: Washington, DC, 1976; pp 184–193.

Farmer, L. J.; Mottram, D. S. In *Flavour Science and Technology*; Bessiere, Y., Thomas, A. F., Eds.; Wiley: New York, 1990; pp 113–116.

Farmer, L. J.; Patterson, R. L. S. Compounds contributing to meat flavour. *Food Chem.* **1991**, *40*, 201–205.

Gasser, U.; Grosch, W. Identification of volatile flavour compounds with high aroma values from cooked beef. *Z. Lebensm. Unters. Forsch.* **1988**, *186*, 489–494.

Gasser, U.; Grosch, W. Primary odorants of chicken broth. *Z. Lebensm. Unters. Forsch.* **1990a**, *190*, 3–8.

Gasser, U.; Grosch, W. *Z. Lebensm. Unters. Forsch.* **1990b**, *190*, 511–515 (in German).

Grosch, W.; Zeiler-Hilgart, G. In *Flavor Precursors, Thermal and Enzymatic Conversions*; Teranishi, R., Takeoka, G. R., Güntert, M., Eds.; American Chemical Society: New York, 1992; pp 183–192.

Güntert, M.; Brüning, J.; Emberger, R.; Köpsel, M.; Kuhn, W.; Thielmann, Th.; Werkhoff, P. Identification and formation of some selected sulfur containing flavor compounds in various meat model systems. *J. Agric. Food Chem.* **1990**, *38*, 2027–2041.

Güntert, M.; Bertram, H.-J.; Hopp, R.; Sommer, H.; Werkhoff, P. In *Progress in Flavour Precursor Studies*; Schreier, P., Winterhalter, P., Eds.; Allured Publishing: Carol Stream, IL, 1992a; pp 361–378.

Güntert, M.; Brüning, J.; Emberger, R.; Hopp, R.; Köpsel, M.; Surburg, H.; Werkhoff, P. In *Flavor Precursors, Thermal and Enzymatic Conversions*; Teranishi, R., Takeoka, G. R., Güntert, M., Eds.; American Chemical Society: New York, 1992b; pp 140–163.

Guth, H.; Grosch, W. Identification of potent odourants in static headspace samples of green and black tea powders on the basis of aroma extract dilution analysis (AEDA). *Flavour Fragrance J.* **1993**, *8*, 173–178.

Guth, H.; Hofmann, T.; Schieberle, P.; Grosch, W. Model reactions on the stability of disulfides in heated foods. *J. Agric. Food Chem.* **1995**, *43*, 2199–2203.

Hartmann, G. J.; Carlin, J. T.; Scheide, J. D.; Ho, C.-T. Volatile products formed by the thermal degradation of thiamin at high and low moisture levels. *J. Agric. Food Chem.* **1984**, *32*, 1015–1018.

Hofmann, T.; Schieberle, P. Evaluation of the key odorants in a thermally treated solution of ribose and cysteine by aroma extract dilution techniques. *J. Agric. Food Chem.* **1995**, *43*, 2187–2194.

Holscher, W.; Vitzthum, O. G.; Steinhart, H. Identification and sensorial evaluation of aroma-impact-compounds in roasted columbian coffee. *Cafe Cacao The* **1990**, *34*, 205–212.

Mottram, D. S.; Whitfield, F. B. In *Thermally Generated Flavors: Maillard, Microwave and Extrusion Processes*; Parliment, T. H., Morello, M. J., McGorrin, R. J., Eds.; ACS Symposium Series 543; American Chemical Society: Washington, DC, 1994; pp 180–192.

Nickerson, G. B.; Likens, S. T. Gas chromatographic evidence for the occurrence of hop oil components in beer. *J. Chromatogr.* **1966**, *21*, 1–5.

Reineccius, G. A.; Liardon, R. In *Topics in Flavour Research*; Berger, R. G., Nitz, S., Schreier, R., Eds.; Eichhorn: Hangenham, Germany, 1985; pp 125–136.

Ruther, J.; Baltes, W. Investigations on the composition of commercial meat flavorings. *Lebensmittelchemie* **1994a**, *48*, 27–34 (in German).

Ruther, J.; Baltes, W. Sulfur-containing furans in commercial meat flavorings. *J. Agric. Food Chem.* **1994b**, *42*, 2254–2259.

Silwar, R.; Tressl, R. *Z. Lebensm. Unters. Forsch.* **1989**, *189*, 205–211 (in German).

Case No.: IPR2023-00208
U.S. Patent No. 11,013,250

Tressl, R. In *Thermal Generation of Aromas*; Parliment, T. H., McGorrin, R. J., Ho, C.-T., Eds.; ACS Symposium Series 409; American Chemical Society: Washington DC, 1989; pp 285–301.

Tressl, R. In *Flavour Science and Technology*; Bessiere, Y., Thomas, A. F., Eds.; Wiley: New York, 1990; pp 87–104.

Tressl, R.; Silwar, R. Investigation of sulfur-containing components in roasted coffee. *J. Agric. Food Chem.* **1981**, *29*, 1078–1082.

Tressl, R.; Helak, B.; Grünewald, K.-G.; Silwar, R. In *International Symposium on Food Flavors, Paris*; Adda, J., Richard, H., Eds.; Technique et Documentation-Lavoisier: Paris, 1983; pp 207–230.

Werkhoff, P.; Brüning, J.; Emberger, R.; Güntert, M.; Köpsel, M.; Kuhn, W.; Surburg, H. Isolation and characterization of volatile sulfur-containing meat flavor compounds in model systems. *J. Agric. Food Chem.* **1990**, *38*, 777–791.

Whitfield, F. B.; Mottram, D. S.; Shaw, K. J. In *Progress in Flavor Precursor Studies*; Schreier, P., Winterhalter, P., Eds.; Allured Publishing: Carol Stream, IL, 1993; pp 395–400.

Zhang, Y.; Ho, C.-T. Formation of meatlike aroma compounds from thermal reaction of ionosine 5′-monophosphate with cysteine and gluthatione. *J. Agric. Food Chem.* **1991**, *39*, 1145–1148.

Received for review February 1, 1995. Revised manuscript received July 25, 1995. Accepted September 27, 1995.[⊗]

JF9500703

[⊗] Abstract published in *Advance ACS Abstracts,* December 1, 1995.