


Volume 266
Number 5
14 March 1997

JMIB

JOURNAL OF MOLECULAR BIOLOGY



ACADEMIC PRESS



JMB

Editor-in-Chief

Peter Wright

Department of Molecular Biology, The Scripps Research Institute, 10550 N. Torrey Pines Road,
La Jolla, CA 92037, U.S.A.

Executive Editor

Jonathan Karn

MRC Laboratory of Molecular Biology, Hills Road, Cambridge CB2 2QH, U.K.

Founding Editor

Sir John Kendrew

Consulting Editor

Sydney Brenner

Advisory Editors

Fred Cohen
Alan R. Fersht
Max Gottesman

Robert Huber
Aaron Klug
Moshe Yaniv

Board of Editors

Wolfgang Baumeister, Max-Planck-Institut für Biochemie, Am Klopferspitz 18a, D-82152 Martinsried bei München, Germany
Nam-Hai Chua, Laboratory of Plant Molecular Biology, The Rockefeller University, 1230 York Avenue, New York, NY 10021, U.S.A.

Fred Cohen, Dept of Cellular and Molecular Pharmacology, University of California, San Francisco, CA 94143-0450, U.S.A.
David E. Draper, Dept of Chemistry, Johns Hopkins University, 3400 North Charles Street, Baltimore, MD 21218, U.S.A.
Richard H. Ebright, The State University of New Jersey, Rutgers, Waksman Institute, PO Box 759, Piscataway, NJ 08855-0759, U.S.A.

Alan R. Fersht, University Chemical Laboratory, Cambridge University, Lensfield Road, Cambridge, CB2 1EW, U.K.
Max Gottesman, Institute of Cancer Research, College of Physicians and Surgeons of Columbia University, 701 W. 168th Street, New York, NY 10032, U.S.A.

Barry Holland, Institut de Genetique et Microbiologie, Batiment 409, Universite de Paris XI, 91405 Orsay Cedex 05, France
Barry Honig, Dept of Biochemistry & Molecular Biophysics, College of Physicians & Surgeons of Columbia University, 630 W. 168th Street, NY 10032, U.S.A.

Robert Huber, Max Planck Institute für Biochemie, 82152 Martinsried bei München, Germany
Aaron Klug, MRC Laboratory of Molecular Biology, Hills Road, Cambridge CB2 2QH, U.K.

Jeffrey H. Miller, Dept of Microbiology, University of California, 405 Hilgard Avenue, Los Angeles, CA 90024, U.S.A.

Michael F. Moody, School of Pharmacy, University of London, 29-39 Brunswick Square, London WC1N 1AX, U.K.

Kiyoshi Nagai, Laboratory of Molecular Biology, Hills Road, Cambridge CB2 2QH, U.K.

Steven Reed, Dept of Molecular Biology, The Scripps Research Institute, 10666 N. Torrey Pines Road, La Jolla, CA 92037, U.S.A.

Douglas Rees, Division of Chemistry and Chemical Engineering, California Institute of Technology, 147-75CH, Pasadena, CA 91125, U.S.A.

Timothy J. Richmond, Institut für Molekularbiologie und Biophysik Eidgenössische Technische Hochschule, Hönggerberg, CH 8093 Zürich, Switzerland

Gerald Smith, Fred Hutchinson Cancer Research Center, Basic Sciences Division - A1-162, 1124 Columbia Street, Seattle, WA 98104-2092, U.S.A.

Janet M. Thornton, Biomolecular Structure and Modelling Unit, Dept of Biochemistry and Molecular Biology, University College of London, Room 628, Gower Street, London WC1E 6BT, U.K.

Ignacio Tinoco, Dept of Chemistry, University of California, Berkeley, CA 94720-1460, U.S.A.

Gunnar Von Heijne, Dept. of Biochemistry, University of Stockholm, S-106 91 Stockholm, Sweden

James Wells, Protein Engineering, Genentech Inc, 460 Point San Bruno Blvd, South San Francisco, CA 94080, U.S.A.

Ian Wilson, Dept of Molecular Biology, The Scripps Research Institute, 10666 N. Torrey Pines Road, La Jolla, CA 92037, U.S.A.

Keith R. Yamamoto, Dept of Cellular and Molecular Pharmacology, University of California, San Francisco, CA 94143-0450, U.S.A.

Mitsuhiro Yanagida, Dept of Biophysics, Faculty of Science, Kyoto University, Sakyo-Ku, Kyoto 606, Japan

Moshe Yaniv, Department of Biotechnology, Pasteur Institute, 25 rue de Dr Roux, 75724 Paris Cedex 15, France

Editorial Office

Journal of Molecular Biology, Academic Press Editorial Office, 10d St Edward's Passage, Cambridge CB2 3PJ, U.K.

JOURNAL OF MOLECULAR BIOLOGY: ISSN 0022-2836. Volumes 265-274, 1997, is published weekly (except the last two weeks in December; 50 issues in total) by Academic Press at 24-28 Oval Road, London NW1 7DX, England. Annual worldwide subscription price including postage: £2280. Personal subscription rates are available from the publisher. Subscription orders should be sent to Academic Press, Foots Cray High Street, Sidcup, Kent DA14 5HP, U.K. (Tel: 0181-300 3322). Send notices of changes of address to the publisher at least 6-8 weeks in advance, including both old and new addresses.

Academic Press journals do not levy page charges.

Buyers in Canada should add GST at the current rate of 7%. Periodicals postage paid at Jamaica, NY 11431, U.S.A.

Air freight and mailing in the U.S.A. by Publications Expediting Inc., 200 Meacham Avenue, Elmont, NY 11003, U.S.A.

U.S.A. POSTMASTERS: send change of addresses to JOURNAL OF MOLECULAR BIOLOGY, c/o Publications Expediting, Inc., 200 Meacham Avenue, Elmont, NY 11003, U.S.A.

Printed in U.K., on acid free paper.

Listed in Biological Abstracts, Science Citation Index, Chemical Abstracts, Reference Update, Excerpta Medica database EMBASE, Current Contents, SCISEARCH, Research Alert, Biotechnology Citation Index, Biochemistry and Biophysics Citation Index, Index Medicus, Genetics Abstracts, Biological & Agricultural Index.

Volume 266, Number 5

Contents

Communication

Solution Structure of the Albumin-binding GA Module: A Versatile Bacterial Protein Domain (mb960856)	M. U. Johansson, M. de Chateau M. Wikström, S. Forsen T. Drakenberg and L. Björck	859–865
--	---	---------

Articles

Design of CytR Regulated, cAMP-CRP Dependent Class II Promoters in <i>Escherichia coli</i> : RNA Polymerase-Promoter Interactions Modulate the Efficiency of CytR Repression (mb960852)	H.-H. Kristensen, P. Valentin-Hansen and L. Søgaard-Andersen	866–876
Analysis of Interactions Between the Codon–Anticodon Duplexes within the Ribosome: Their Role in Translation (mb960802)	V. I. Lim	877–890
Protein P7 of Phage $\phi 6$ RNA Polymerase Complex, Acquiring of RNA Packaging Activity by <i>In Vitro</i> Assembly of the Purified Protein onto Deficient Particles (mb960817)	J. T. Juuti and D. H. Bamford	891–900
The Bacteriophage $\phi 29$ Packaging Proteins Supercoil the DNA Ends (mb960843)	S. Grimes and D. Anderson	901–914
RNA-DNA Hybrid Formation at a Bacteriophage T4 Replication Origin (mb960844)	K. Carles-Kinch and K. N. Kreuzer	915–926
Single-stranded DNA Binding Properties of the <i>uvsY</i> Recombination Protein of Bacteriophage T4 (mb960829)	M. A. Sweezy and S. W. Morrical	927–938
A Survey of Polypeptide Deformylase Function Throughout the Eubacterial Lineage (mb960835)	D. Mazel, E. Coïc, S. Blanchard W. Saurin and P. Marlière	939–949
A Dynamic <i>in Vivo</i> View of the HIV-I Rev-RRE Interaction (mb960858)	B. Charpentier, F. Stutz and M. Rosbash	950–962
Crystal Structure of the Gramicidin/Potassium Thiocyanate Complex (mb960837)	D. A. Doyle and B. A. Wallace	963–977
Module Swaps between Related Translocator Proteins pIV^{fl} , pIV^{IKe} and PuID: Identification of a Specificity Domain (mb960866)	S. Daefler, M. Russel and P. Model	978–992

Traffic Control of Completely Assembled MHC Class I Molecules Beyond the Endoplasmic Reticulum (mb960822)	S. Joyce	993-1001
Three-dimensional Reconstruction of <i>Limulus polyphemus</i> Hemocyanin from Cryoelectron Microscopy (mb960840)	J.-C. Taveau, N. Boisset, J. Lamy, O. Lambert and J. N. Lamy	1002-1015
Crystal Structures of <i>Escherichia coli</i> and <i>Salmonella typhimurium</i> 3-Isopropylmalate Dehydrogenase and Comparison with their Thermophilic Counterpart from <i>Thermus thermophilus</i> (mb960797)	G. Wallon, G. Kryger, S. T. Lovett T. Oshima, D. Ringe, G. A. Petsko	1016-1031
Characterization of Recombinant Soybean Leghemoglobin <i>a</i> and Apolar Distal Histidine Mutants (mb960833)	M. S. Hargrove, J. K. Barry E. A. Brucker, M. B. Berry G. N. Phillips Jr, J. S. Olson R. Arredondo-Peter, J. M. Dean R. V. Klucas and G. Sarath	1032-1042
Structure-based Identification of a Ricin Inhibitor (mb960865)	X. Yan, T. Hollis, M. Svinth, P. Day A. F. Monzingo, G. W. A. Milne and J. D. Robertus	1043-1049
Energetics of Target Peptide Recognition by Calmodulin: A Calorimetric Study (mb960785)	P. L. Wintrode and P. L. Privalov	1050-1062
Author Index		1063-1064

The home page for the Journal of Molecular Biology can be found at

<http://www.hbuk.co.uk/ap/journals/mb.htm>

IDEAL, Academic Press' online scientific journal library, where guest users can freely search and browse journal abstracts, can be found at:


<http://www.idealibrary.com/> (San Jose, USA) and

<http://www.europe.idealibrary.com/> (Bath, UK)

The main Academic Press home page is at <http://www.apnet.com/> where you will find general information about Academic Press and the AP Online Book Catalog.

Copyright © 1997 Academic Press Limited. The appearance of the code at the bottom of the first page of a paper in this Journal indicates the copyright owner's consent that copies of the paper may be made for personal or internal use, or for the personal or internal use of specific clients in the U.S.A. This consent is given on the condition, within the U.S.A., that the copier pay the stated per-copy fee through the Copyright Clearance Center, Inc., 27 Congress Street, Salem, MA 01970, for copying beyond that permitted by Section 107 or 108 of the U.S. Copyright Law. This consent does not extend to other kinds of copying, such as copying for general distribution, for advertising or promotional purposes, for creating new collective works, for resale or for copying or distributing copies outside the U.S.A.

Cover Illustration: Solution structure of the albumin-binding GA molecule—a versatile bacterial protein domain. See paper by Johansson *et al.* *J. Mol. Biol.* **266**, 859-865.



Typeset by BPC Digital Techset Ltd; printed and bound by BPC Wheatons Ltd
Both members of The British Printing Company Ltd
This journal is printed on acid-free paper

JMB



Characterization of Recombinant Soybean Leghemoglobin *a* and Apolar Distal Histidine Mutants

Mark S. Hargrove^{1*}, Jennifer K. Barry¹, Eric Allen Brucker¹
Michael B. Berry¹, George N. Phillips Jr¹, John S. Olson¹
Raúl Arredondo-Peter², Jeanenne M. Dean², Robert V. Klucas²
and Gautam Sarath²

¹Department of Biochemistry
and Cell Biology, Rice
University, Houston, TX
77005-1892, USA

²Department of Biochemistry
University of Nebraska-Lincoln
Lincoln, NE 68588-0664, USA

The cDNA for soybean leghemoglobin *a* (Lba) was cloned from a root nodule cDNA library and expressed in *Escherichia coli*. The crystal structure of the ferric acetate complex of recombinant wild-type Lba was determined at a resolution of 2.2 Å. Rate constants for O₂, CO and NO binding to recombinant Lba are identical with those of native soybean Lba. Rate constants for hemin dissociation and auto-oxidation of wild-type Lba were compared with those of sperm whale myoglobin. At 37°C and pH 7, soybean Lba is much less stable than sperm whale myoglobin due both to a fourfold higher rate of auto-oxidation and to a ~600-fold lower affinity for hemin. The role of His61(E7) in regulating oxygen binding was examined by site-directed mutagenesis. Replacement of His(E7) with Ala, Val or Leu causes little change in the equilibrium constant for O₂ binding to soybean Lba, whereas the same mutations in sperm whale myoglobin cause 50 to 100-fold decreases in K_{O₂}. These results show that, at neutral pH, hydrogen bonding with His(E7) is much less important in regulating O₂ binding to the soybean protein. The His(E7) to Phe mutation does cause a significant decrease in K_{O₂} for Lba, apparently due to steric hindrance of the bound ligand. The rate constants for O₂ dissociation from wild-type and native Lba decrease significantly with decreasing pH. In contrast, the O₂ dissociation rate constants for mutants with apolar E7 residues are independent of pH, suggesting that hydrogen bonding to the distal histidine residue in the native protein is enhanced under acid conditions. All of these results support the hypothesis that the high affinity of Lba for oxygen and other ligands is determined primarily by enhanced accessibility and reactivity of the heme group.

© 1997 Academic Press Limited

Keywords: soybean leghemoglobin; ligand binding; site-directed mutagenesis

*Corresponding author

Introduction

Leghemoglobins (Lb) are small monomeric heme proteins found in root nodules of leguminous plants. These proteins have unusually high affinities for oxygen compared to vertebrate myoglobins and hemoglobins. The association rate constants for O₂ binding are very large (1×10^8 to 3×10^8 M⁻¹ s⁻¹), whereas the dissociation rate constants are in the "normal" range for hemoglobins (5 to 30 s⁻¹; Gibson *et al.*, 1989). High oxygen affi-

nity is required to prevent inhibition of nitrogen fixation, and a moderate dissociation rate constant is necessary for facilitating diffusion of oxygen to a high-affinity bacteroid oxidase (Appleby, 1992).

The tertiary folding patterns of lupin and soybean Lb are very similar to those of mammalian myoglobins, and the α and β subunits of human hemoglobin. This pattern consists of seven α -helices (A to H, minus the D helix found in most mammalian myoglobins) wrapped around an iron-protoporphyrin IX prosthetic group that is coordinated to the histidine side-chain of the eighth residue in the F helix. Leghemoglobins also contain a "distal" histidine residue at position E7. Polar interactions involving His-(E7) in mammalian myoglobins have

Abbreviations used: Lb, leghemoglobin; EPR, electron paramagnetic resonance; Mb, myoglobin.

been shown to play a key role in regulating ligand binding (Springer *et al.*, 1994). The evidence for hydrogen bonding between bound O₂ and the distal histidine residue in leghemoglobin is less clear.

Comparisons of the ligand binding properties of leghemoglobins with those from a wide variety of animal myoglobins and hemoglobins have suggested that the small rates of O₂ dissociation from soybean and lupin Lb are due to hydrogen bonding with His(E7) (Appleby *et al.*, 1983; Mims *et al.*, 1983; Stetzkowski *et al.*, 1979; Rohlf *et al.*, 1988). Initial electron paramagnetic resonance (EPR) experiments with cobalt-containing soybean Lb indicated weak hydrogen bonding at neutral pH and a significant strengthening of this interaction at acid pH (Ikeda-Saito *et al.*, 1981). More recent spin echo EPR studies confirmed this observation and further suggested that hydrogen bonding with the distal histidine residue may occur only at low pH (Lee *et al.*, 1993). Fuchsman & Appleby (1979) observed a single narrow IR band at $\nu_{\text{C-O}} = 1947 \text{ cm}^{-1}$ for LbaCO at pH 7. This peak corresponds to the main band observed in sperm whale MbCO ($\nu_{\text{C-O}} = 1945 \text{ cm}^{-1}$) and indicates a similar electrostatic environment around the bound ligand in both proteins. More recently, Harutyunyan *et al.* (1995) reported a distance of 2.7 Å between the second ligand atom and N^δ of His(E7) in lupin oxyleghemoglobin II, indicating a hydrogen bond between the imidazole group and bound O₂. Thus, hydrogen bonding is thought to occur in leghemoglobins, especially at low pH. However, the strength of this interaction and its exact role in governing the kinetics and equilibria of ligand binding to plant hemoglobins have not been established directly by site-directed mutagenesis.

Like all oxygen binding proteins, leghemoglobins must be in the Fe(II) oxidation state to bind oxygen. Within the root nodule, the protein is kept in the ferrous state by a reductase that catalyses the reduction of metleghemoglobin by NADH (Saari & Klucas, 1984; Ji *et al.*, 1994a,b). However, little is known about the molecular recognition and protein-protein interaction between the enzyme and its substrate. Soybean root nodules contain four leghemoglobin gene products, termed Lba, Lbc1, Lbc2 and Lbc3, and four variants of these four proteins in which the N termini are acetylated (Appleby, 1984). The amino acid sequences and ligand binding properties of the four gene products are very similar (Martin *et al.*, 1990). Several studies have shown changes in the relative concentrations of the individual leghemoglobins during plant growth, but the physiological consequences of differential expression and turnover are not well understood (Appleby, 1984).

The expression of recombinant hemoglobins and myoglobins has allowed systematic studies of the structural factors that regulate ligand binding to these proteins (Nagai *et al.*, 1987; Mathews *et al.*, 1991; Kloeck *et al.*, 1993; Springer *et al.*, 1994). In order to carry out similar mechanistic studies, we

have cloned Lba from a soybean cDNA library, inserted it into two expression vectors, and used both constitutive and inducible systems to express large quantities of protein. The crystal structure of recombinant wild-type Lba was determined to 2.2 Å and compared with that of the native protein. The ligand binding, spectral, auto-oxidation and hemin dissociation properties of native and recombinant Lba were also compared in detail. His61(E7) to Ala, Val, Leu and Phe mutations were constructed to examine the role of the distal histidine residue in governing O₂ binding. Finally, Tyr30(B10) to Phe and Leu65(E11) to Val mutations were made to examine the structural origin of differences between the rates of O₂ binding to lupin LbII and soybean Lba.

Results

Properties of wild-type Lba

As described in Materials and Methods, the N-terminal methionine residue of recombinant Lba is removed post-translationally during constitutive expression in *Escherichia coli* TB1 cells, making the wild type recombinant Lba chemically identical with native Lba (Figure 1). The absorbance spectra of wild-type metLb, deoxyLb, LbO₂ and LbCO are identical with those of the native protein (Table 1). The kinetic parameters for O₂, CO and NO binding, hemin loss and auto-oxidation are also the same for the wild-type and native proteins (Tables 2, 4, and 5). These data show that the protein obtained from *E. coli* TB1 cells is functionally and spectrally the same as native soybean leghemoglobin *a*. Lbs expressed in the inducible pET28a/BL21DE3 system are only partially processed, yielding a mixed sample of N-met and "native-like" protein. However, the samples prepared from the pET28a/BL21DE3 system are kinetically and spectrally identical with those prepared in TB1. Furthermore, it has been observed that addition of an entire N-terminal S-tag (Novagen) domain has no effect on the kinetic or spectral properties of the Lbs examined here (unpublished results).

Crystal structure of wild type acetomet-Lba

The crystallization system described by Vainshtein *et al.* (1975) was followed starting with aquomet wild-type Lba. Since 0.5 M ammonium acetate is present in the solution, the acetomet form of the protein is produced in the crystals. The backbone and side-chain positions in the acetate complex of wild-type metLba are very similar to those of the nicotinate complex of native metLba (Ollis *et al.*, 1983; Ellis *et al.*, 1997). An overlay of the backbone atoms plus the side-chains of the proximal and distal histidine residues of the two structures is shown in Figure 2. The native nicotinate structure is shown in yellow and molecule *a* of the two molecules in the asymmetric unit of the wild-type structure is shown in blue. The ligands have been




Figure 1. The cDNA and amino acid sequence of recombinant soybean Lba in the pEMBL19⁺ plasmid pLba. The ribosome binding site lies between the 5' *KpnI* and *NcoI* restriction sites. The flanking 5' *NcoI* and 3' *EcoRI* restriction sites can be used to move the gene independently of the promoter region. The gene sequence corresponds with that observed for soybean Lba by Hyldig-Nielsen *et al.* (1982).

removed for clarity. These data show that the protein obtained from *E. coli* is structurally the same as native soybean Lba. The most significant differences between the two structures are in the positions of the heme group, distal histidine residue, portions of the CD and FG corners, and part of the F helix including His(F8). These differences appear to be due to the size of the coordinating ligand. In

the native structure, the large nicotinate ring forces the distal histidine residue out toward solvent and pushes the heme group downward. The proximal histidine residue, F helix and FG corner rotate in order to compensate for this movement of the porphyrin ring. The CD corner also appears to move away from the bound ligand, but this region is either mobile or disordered in both structures.

Table 1. Comparison of the wavelength maxima for the Soret absorbance bands of soybean Lb and sperm whale Mb at pH 7, 25°C

Protein	Wavelength peak (nm)				
	Fe(II)	Fe(III)	Fe(II)O ₂	Fe(II)CO	Fe(II)NO
Native Lb	428	403	409	416	418
Wild-type Lb	428	403	409	416	418
H61L Lb	425	395	414	419	418
Wild-type Mb	434	409	415	424	421
H64L Mb	434	393	415	425	421

Figure 3 shows electron density around the heme binding site of wild-type Lba. The view is looking into the pocket from solvent with the distal portion of the heme pocket located in the upper half of the drawing. The heme is shown in white, the acetate ligand in yellow and the protein chain in red. Well-resolved density for His61(E7) is seen next to the bound acetate group. Other prominent residues are (counter-clockwise): Leu65(E11), located just behind His61(E7); and Phe44(CD1), located to the left of His61(E7), just above the heme plane.

The distal histidine residue in the structure of native soybean metLba appears to be hydrogen bonding with the carbonyl oxygen atom of the nicotinate ligand (Ollis *et al.*, 1983). In the wild-type metLba structure, N^ε of His61(E7) is 2.64 Å and 3.08 Å away from the carbonyl oxygen atom of the acetate group in the two protein molecules found in the asymmetric unit. This proximity suggests that a hydrogen bond is also formed between bound acetate and the distal histidine residue in the wild-type protein. Harutyunyan *et al.* (1995) have shown that a hydrogen bond occurs between the distal histidine residue and the second coordinated ligand atom in the crystal structure of lupin LbO₂. Thus, in leghemoglobins, the distal histidine

residue appears to be quite flexible and is capable of interacting electrostatically with ligands of varying size.

Role of His61(E7) in O₂ binding

Hydrogen bonding between bound oxygen and the distal histidine residue in sperm whale myoglobin has been established by X-ray crystallography, and its effect on oxygen affinity has been determined in site-directed mutagenesis studies (Springer *et al.*, 1994). Following the strategy of Springer *et al.* (1989), we replaced His61(E7) with Ala, Val, Leu and Phe to examine quantitatively how oxygen binding is affected by loss of hydrogen-bonding potential at this position in Lba. Table 2 presents a comparison of the effects of these E7 mutations on the rate and equilibrium constants for O₂ binding to soybean Lba with those observed previously for O₂ binding to sperm whale myoglobin.

In general, the aliphatic replacements have a much smaller effect on O₂ binding to leghemoglobin than to myoglobin. These differences are particularly striking when the oxygen dissociation rate constants are examined. Only the Phe-E7 mutation in




Figure 2. Crystal structure of the acetate complex of recombinant soybean metLba. A stereo overlay of the backbone atoms of the nicotinate complex of native metLba (yellow) and wild-type metLba (blue) including the proximal and distal histidine side-chains. The nicotinate and acetate ligands have been removed for clarity.




Figure 3. Electron density of the heme pocket in wild-type metLba-acetate. The heme prosthetic group is shown in white, the acetate ligand in yellow and the protein in red. The distal histidine residue is within hydrogen-bonding distance of the carbonyl oxygen atom of the bound acetate group.

Lba produces a large increase in k_{O_2} , whereas in myoglobin all of the mutations cause at least 100-fold increases in k_{O_2} . These results clearly show that the hydrogen bond between oxygen and His-E7 in leghemoglobin is very weak at neutral pH and not the main determinant of the high affinity of this protein for O_2 , in agreement with the EPR results of Ikeda-Saito *et al.* (1981) and Lee *et al.* (1993). The three- to fourfold increases in k_{O_2} observed for the Val(E7) and Leu(E7) Lba mutants are balanced by roughly equal increases in k_{O_2} so there is little net change in overall oxygen affinity. The Ala(E7) and Phe(E7) mutations produce an increase and decrease, respectively, in K_{O_2} with respect to the wild-type protein. These latter changes seem to imply that the E7 residue may sterically hinder the bound ligand. However, an exact interpretation of these results will require high-resolution crystal structures of the proteins. Regardless, it is clear that the polarity of His(E7) plays a much smaller role in regulating ligand binding to leghe-moglobin than to myoglobin.

The effects of pH on oxygen binding to native, wild-type and E7 mutants of soybean leghemoglobin are presented in Table 3. As shown by Appleby *et al.* (1983), when the pH is decreased from 7 to 5, the rate constant for oxygen dissociation from Lba decreases twofold with no change in the association rate constant. The resultant increase in affinity at low pH has been attributed to a strengthening of the hydrogen bond between His61 and bound ligand. The apolar mutants of Lba show no pH dependence (Table 3). These mutagenesis results support the view that His61 plays a role in the pH dependence of O_2 binding to native leghemoglobin and that hydrogen bonding to the bound ligand is enhanced under acidic conditions.

CO and NO binding

The rate and equilibrium constants for carbon monoxide (CO) and nitric oxide (NO) binding to Lba, sperm whale myoglobin and the correspond-

Table 2. The effects of apolar substitutions at position E7 on the rate and equilibrium constants of O_2 binding to soybean Lba and sperm whale Mb at pH 7, 20°C

Residue at 61(E7)	k'_{O_2} $\mu M^{-1} s^{-1}$		k_{O_2} s^{-1}		K_{O_2} μM^{-1}	
	Lb	Mb	Lb	Mb	Lb	Mb
His(wild-type)	130	17	5.6	15	23	1.1
Ala	290	53	3.1	2300	94	0.023
Val	280	110	27	10000	10	0.011
Leu	400	98	24	4100	17	0.023
Phe	130	75	280	10000	0.45	0.0075

Data for sperm whale myoglobin were taken from Springer *et al.* (1994).

Table 3. Dependence of O₂ binding parameters on pH for native, wild-type, and Leu-E7 soybean Lba at 20°C.

Soybean Lb	pH 7			pH 5		
	k'_{O_2} ($\mu\text{M}^{-1}\text{s}^{-1}$)	k_{O_2} (s^{-1})	K_{O_2} (μM^{-1})	k'_{O_2} ($\mu\text{M}^{-1}\text{s}^{-1}$)	k_{O_2} (s^{-1})	K_{O_2} (μM^{-1})
Native	130	5.6	23	140	2.7	51
Wild-type	130	5.6	23	130	2.2	60
H61A	290	3.1	94	250	2.7	92
H61V	280	27	10	280	28	10
H61L	400	24	17	470	25	19
H61F	130	280	0.45	120	230	0.54

ing Leu(E7) mutants are presented in Table 4. Replacement of the distal histidine residue with Leu increases the association rate constants for CO and NO binding to both proteins, but again the effects are much greater in myoglobin. The rate constants for CO and NO dissociation are largely unaffected by the Leu(E7) replacement. For both ligands, wild-type Lba has larger association rate constants and smaller dissociation rate constants than wild-type myoglobin. These results point out that with or without a distal histidine residue, leghemoglobin is inherently much more reactive than myoglobin.

Autooxidation and hemin loss

A comparison of resistances to auto-oxidation and hemin dissociation is given in Table 5. Brantley *et al.* (1993) have shown that the rate constants for auto-oxidation of myoglobin mutants correlate inversely with O₂ affinity. Thus, we expected Lba to auto-oxidize slowly due to its high O₂ affinity. However, the opposite effect was observed; k_{autoox} for Lba is about fourfold greater than that of wild-type sperm whale Mb. This result does correlate with a much weaker hydrogen bond between bound O₂ and His(E7) in leghemoglobin. Brantley *et al.* (1993) showed that this polar interaction inhibits protonation of the Fe-O₂ complex and its subsequent dismutation into ferric iron and the neutral superoxide radical. Both myoglobin and leghemoglobin show a marked increase in k_{autoox} with decreasing pH, presumably due to protonation of bound O₂.

The rate constant for hemin dissociation from native and wild-type Lba is $\sim 6\text{ h}^{-1}$ at pH 7.0, 37°C. This rate is 600 times faster than that for myoglobin under the same conditions and, in combination with the increased auto-oxidation rate, results in a much less stable holoprotein (Hargrove & Olson, 1996). Lba also has a much larger temperature dependence for hemin dissociation than myoglobin ($E_a \approx 33\text{ kcal/mol}$ for aquometLba versus $E_a \approx 8\text{ kcal/mol}$ for sperm whale aquometMb at pH 7; Hargrove *et al.*, 1994; and unpublished results). As shown in Table 5, k_{-H} for met Lba is $\sim 0.8\text{ h}^{-1}$ at 20°C, which is roughly tenfold less than the rate observed at 37°C. The rate constant for hemin loss from myoglobin decreases by only a factor of ~ 2 when the temperature is decreased from 37 to 20°C at both pH 5.0 and pH 7.0 (Hargrove *et al.*, 1994; and unpublished results).

The rates of hemin loss from the distal histidine mutants of Lba are too fast to be measured conveniently at 37°C. As a result, comparisons between mutant and wild-type leghemoglobins were measured at 20°C. The distal histidine residue of Lba clearly stabilizes bound hemin, as shown by the H61L substitution, which increases the rate constant for hemin dissociation ~ 50 -fold. A similar effect occurs when His(E7) is replaced by Leu in myoglobin. A large part of this effect is probably due to loss of the coordinated water molecule from the ferric H61L protein. Both Leu-E7 metMb and metLba show a broad absorbance band centered at 395 nm, which is characteristic of 5-coordinated ferric proteins (Table 1; Quillin *et al.*, 1993). The pH dependence of hemin dissociation from Lba is

Table 4. Comparison of the CO and NO binding parameters of soybean Lba and sperm whale Mb at pH 7, 20°C

Protein	k'_{CO} ($\mu\text{M}^{-1}\text{s}^{-1}$)	k_{CO} (s^{-1})	K_{CO} (μM^{-1})	k'_{NO} ($\mu\text{M}^{-1}\text{s}^{-1}$)	k_{NO} (s^{-1})	K_{NO} (μM^{-1})
Native Lb	17	0.0078	2200	170	0.00002	9,000,000
Wild-type Lb	15	0.0084	1800	190	0.00002	9,000,000
H61L Lb	170	0.0024	72,000	320	0.00002	16,000,000
Wild-type Mb	0.51	0.019	27	22	0.00010	220,000
H64L Mb	26	0.024	1100	190	0.00013	1,500,000

Data for sperm whale myoglobin were taken from Springer *et al.* (1994).

Table 5. Comparison of the rate constants for hemin dissociation (k_{-H}) from and auto-oxidation (k_{autoox}) of soybean Lba and sperm whale Mb

Protein	k_{autoox} (h^{-1}) at 37°C		k_{-H} (h^{-1}) ^b at 20°C	
	pH 7	pH 5	pH 7	pH 5
Native Lb	0.2	1.6	0.7	1.3
Wild-type Lb	0.2	3.0	0.9	3.0
H61L Lb	0.3	6.6	47	45
Wild-type Mb	0.055	2.4	0.005 ^c (0.01)	0.4(1.0)
H64L Mb	10	200 ^a	0.094 ^c (0.20)	5.2 ^c (11)

The k_{autoox} values for MbO₂ were taken from Brantley *et al.* (1993) and the values of k_{-H} for metMb from Hargrove *et al.* (1996b). Rates of auto-oxidation were measured in the presence of 1mM EDTA, catalase, and superoxide dismutase as described by Brantley *et al.* (1993). The k_{-H} values in parentheses for metMb were measured at 37°C by Hargrove *et al.* (1996b) and Hargrove *et al.* (1994).

^a The value of k_{autoox} for Leu(E7) Mb at pH 5 was estimated assuming the same pH dependence as is observed for the Val(E7) Mb mutant by Brantley *et al.* (1993).

^b The rates of hemin loss from leghemoglobin were measured at 20°C to allow comparison with mutant Lbs, which lose heme too rapidly at 37°C.

^c The rates of hemin loss at 20°C were computed using the k_{-H} value at 37°C and an activation energy of 8.0 kcal/mol. The values in parentheses were measured at 37°C (Hargrove *et al.*, 1996b).

much smaller than that of myoglobin. The resistances of both proteins to auto-oxidation and hemin loss are similar at pH 5, whereas at pH 7 sperm whale myoglobin is much more stable.

Comparison with lupin LbII

As shown in Table 6, the association and dissociation rate constants for O₂ binding to lupin LbII are three- to fivefold greater than the corresponding parameters for soybean Lba. This implies more rapid rates of ligand entry into and exit from the distal pocket of the lupin protein. Both proteins contain His(E7) and Phe(CD1). In lupin LbII the B10 and E11 residues are smaller, Phe and Val, respectively, *versus* Tyr and Leu in soybean leghemoglobin. However, this decrease in size does not appear to account for the ligand binding differences. Mutation of Leu65(E11) to Val has little effect on O₂ binding to soybean Lba, and the Tyr30(B10) to Phe substitution causes decreases, not increases, in k'_{O_2} and k_{O_2} (Table 6). As suggested by Gibson *et al.* (1989), the differences in kinetic behavior among leghemoglobins appear to be due to amino acid changes remote from the ligand binding site. The increase in affinity due to the Y30F substitution is striking and suggests that an in-

depth study of the role of this residue in ligand binding will be rewarding.

Discussion

The crystal structure of recombinant Lba and the data reported in Tables 1 and 4 show that the protein expressed in *E. coli* TB1 cells is functionally and structurally identical with that of native Lba. It is unclear why the N-terminal methionine residue is removed from this protein when wild-type myoglobin grown under the same conditions is expressed with the N-Met still attached (Springer & Sligar, 1987; Phillips *et al.*, 1990). The N-terminal amino acid sequences of the two proteins are similar (Lba, VAFTE-; Mb, VLSEG-). There is no clear explanation for the difference in post-translational processing during constitutive expression.

Leghemoglobin is much less stable than myoglobin due to a fourfold higher rate of auto-oxidation and a ~600-fold higher rate of hemin loss at pH 7, 37°C. The rate constant for hemin loss of the H61L Lba mutant, ~50 h⁻¹, is in the range of that for H93G sperm whale myoglobin, which lacks a covalent bond between the heme and globin (Hargrove *et al.*, 1996a). The increase in k_{-H} caused by the H61L mutation suggests that the distal histidine residue of Lb stabilizes bound heme by hydrogen bonding to a coordinated water molecule as is observed in myoglobin. The decreased stability of Lba probably accounts for the lower yields in the constitutive expression (TB1) system relative to that of myoglobin. Purification of leghemoglobin in the presence of CO or other distal ligands prevents heme loss and greatly increases protein yields in both the constitutive and inducible expression systems.

The oxygen association rate constants for leghemoglobins are very fast, of the order of 10⁸ M⁻¹ s⁻¹, and are close to the rate expected for a diffusion-limited association reaction with an unhindered

Table 6. Comparison of the rate constants for O₂ binding to soybean Lba and Lupin LbII at pH 7, 20°C

Protein	k'_{O_2} ($\mu\text{M}^{-1}\text{s}^{-1}$)	k_{O_2} (s^{-1})	K_{O_2} (μM^{-1})
Lupin LbII	320	25	13
Soybean Lba	130	5.6	23
Soybean Lba (L65V)	180	5.8	30
Soybean Lba (Y30F)	79	0.75	100

The rate constants for lupin LbII were taken from Gibson *et al.* (1989).

model heme ($k'_{\text{diffusion}} \approx 5 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$, Reisberg & Olson, 1980). Gibson *et al.* (1989) also suggested that the Lb rates may be the fastest possible for ligand association to a myoglobin-like protein. The high rates of ligand binding to Lb result in part from a very reactive heme iron atom as has been demonstrated with laser photolysis experiments (Stetzkowski *et al.*, 1985; Gibson *et al.*, 1989). Harutyunyan *et al.* (1995) have shown that the proximal histidine residue in lupin Lb is much more free to rotate and move toward the heme plane than in myoglobin. These results provide a structural explanation for the high reactivity of leghemoglobins. The heme pocket of Lba is also more flexible and kinetically accessible than those of myoglobin and other heme proteins (Rohlf *et al.*, 1988). This has been demonstrated by many techniques and is exemplified by the ability of Lba to bind large ligands such as imidazole, nicotinate, and long-chain alkyl isocyanides, which do not bind tightly to myoglobin or hemoglobin (Stetzkowski *et al.*, 1979; Kong *et al.*, 1983; Mims *et al.*, 1983).

Even though the bimolecular rate constants are large, His(E7) still appears to offer some resistance to ligand association. The distal heme pocket of the deoxy forms of myoglobin and lupin leghemoglobin both contain a non-coordinated water molecule that hydrogen bonds to His(E7) (Quillin *et al.*, 1993; Harutyunyan *et al.*, 1995). This water molecule must be displaced before other ligands can bind. The increased association rate constants for oxygen binding to the H61A, H61V, and H61L mutants of Lba are probably due to the loss of water in the deoxy forms of these proteins. The same mutations produce even larger increases in k'_{O_2} when constructed in sperm whale myoglobin (Quillin *et al.*, 1993; Table 2).

The functional evidence for a hydrogen bond between bound O_2 and His(E7) in leghemoglobin is equivocal at pH 7.0. The Ala, Val and Leu(E7) substitutions do not produce large decreases in K_{O_2} as is observed in myoglobin. However, the pH dependence of O_2 dissociation from leghemoglobin does appear to involve the distal histidine residue. Decreasing the pH from 7 to 5 causes a twofold increase in O_2 affinity when the E7 residue is histidine but has no effect when the E7 residue is apolar. This result suggests that hydrogen bonding occurs under acid conditions, but even at low pH, the strength of this interaction is still much less than that observed in myoglobin.

The β subunit of R-state human hemoglobin shows high rates of ligand association, and replacement of the distal histidine residue with either Gly or Phe has little effect on O_2 binding (Mathews *et al.*, 1989). Thus both soybean Lba and R-state human β subunits have evolved mechanisms for stabilizing bound O_2 and inhibiting auto-oxidation that do not involve a strong hydrogen bond with His(E7). In contrast, strong favorable polar interactions occur in the distal pockets of all mammalian myoglobins and the α subunit of human hemoglobin.

However, the distal histidine residue is very important for retaining heme in leghemoglobin and human β subunits. Replacing the distal histidine residue with smaller apolar residues causes ≥ 50 -fold increases in the rate of heme loss from met-Lba and ferric β subunits, explaining the high degree of conservation of His(E7) in these proteins (Table 5; Hargrove *et al.*, 1994).

Materials and Methods

A full-length cDNA for Lba was isolated from a soybean nodule cDNA library using anti-Lba antibodies as a probe (Ji *et al.*, 1994a; Sambrook *et al.*, 1989). Phage (λ -GT11) containing the soybean Lba insert were placed in *E. coli*, harvested, and the DNA isolated. Lba cDNA was excised and ligated into pBluescript K+ (Invitrogen). To obtain the coding region, polymerase chain reaction (PCR) primers containing a *Nco*I site at the starting ATG codon and an *Eco*RI restriction site downstream of the TAA stop codon were used to amplify the Lba gene from the pBluescript-Lba vector. PCR-amplified products were purified by agarose gel electrophoresis and ligated into a pCRII vector (Invitrogen). After transformation, an *E. coli* colony containing the pCRII-Lba construct was used to obtain purified plasmid with the *Nco*I-Lba-*Eco*RI sequence. This gene fragment was removed from pCRII, purified from an agarose gel, and ligated into pACYC184. A duplex oligonucleotide of the constitutive *lac* promoter used by Springer & Sligar (1987) was constructed with the following sequence of restriction sites: 5' *Bsu*36I-*Kpn*I-*lac* promoter-*Nco*I 3'. The pACYC184-Lba plasmid was cut with *Nco*I and *Bsu*36I, and the oligonucleotide containing the *lac* promoter was ligated to the Lba cDNA. This *lac* promoter-Lba construct was removed from pACYC184 and ligated into pEMBL19+ using the flanking *Kpn*I/*Eco*RI sites.

The initial cDNA product cloned in this manner contained two base-pair differences compared to the Lba sequence reported by Hyldig-Nielsen *et al.* (1982). These changes were probably introduced during the PCR reaction used to amplify the cDNA. The gene was corrected to the native sequence by two rounds of site-directed mutagenesis using the Kunkel (1985) method. The resulting plasmid, pLba, contains the native Lba cDNA sequence in pEMBL19+ under control of the *lac* promoter (Figure 1). The Kunkel method of mutagenesis was also used to generate the B10, E7 and E11 mutants starting from the "corrected" wild-type Lba gene.

Expression of wild-type and mutant Lbas

E. coli TB-1 cells were transformed with pLba and grown in LB medium (with ampicillin) at 37°C. The final yield of purified leghemoglobin ($\sim 0.6 \text{ mg/l}$) was roughly one-seventh of that for sperm whale myoglobin expressed under the same conditions ($\sim 4 \text{ mg/l}$). H61F, H61V and H61A proteins showed no appreciable expression in the *lac* promoter-TB1 system. Consequently, the *Nco*I-Lba-*Eco*RI fragment was cloned into pET28a (Novagen) for high-level, inducible expression in *E. coli* BL21DE3 cells. Shake flasks of BL21DE3 cells transformed with the pET28a-Lba vector were grown to an absorbance of ~ 0.6 in $2 \times \text{YT}$ medium at 33°C, induced with 1 mM isopropyl- β -D-thiogalactopyranoside and incubated for an additional four hours. Apo-Lba was found in the supernatant and pellet of the cell lysis mixture. Protein was re-

covered from the lysis pellet by dissolution in 6 M guanidinium chloride followed by dialysis into 20 mM Tris (pH 8.0). Holoprotein was reconstituted by the addition of CO-heme or imidazole-hemin to the soluble and pellet fractions of the lysis mixture. The final yield of wild-type or mutant Lba was ~10 mg/l of medium. Reconstituted holo-Lba was precipitated with 90% saturated ammonium sulfate for further purification.

Regardless of the method of expression, recombinant Lba was purified as described by Jun *et al.* (1994) with the following modifications. The 90% saturated ammonium sulfate Lba pellet was resuspended in 1.8 M ammonium sulfate and loaded directly onto a phenyl Sepharose column, and the protein was not oxidized prior to G-75 chromatography. Purified Lba was in the CO or met-imidazole form (depending on the method of reconstitution) and had a Soret to 280 nm absorbance ratio greater than 4. Native Lba was purified from soybean root nodules using the method of Jun *et al.* (1994). N-terminal sequencing of the purified Lbs was performed at the UNL-Protein Core Facility on an ABI-494 Procise sequencer using a manufacturer-supplied protocol.

Crystallography of wild-type Lba

Crystals of the acetate complex of wild-type metLba were grown at room temperature to 0.5 mm × 0.05 mm × 0.05 mm in three to five days using the hanging drop method. The starting concentration of metLba was 1 mM in 0.1 M sodium phosphate (pH 6.0). The precipitant was 2.4 M ammonium sulfate, 0.5 M ammonium acetate at pH 6.8 (Vainshtein *et al.*, 1975). X-ray diffraction data were collected at room temperature on an R-axis IIc imaging plate system using copper K α radiation from a Siemens rotating anode operating at 50 mV and 90 mA. Crystals were in the $P2_12_12_1$ space group. The data were 90% complete at the limiting resolution of 2.2 Å, with an R_{merge} of 10.3%. Phases were determined by molecular replacement using XPLOR (Brünger, 1987) starting with the coordinates for the structure of Lba-nicotinate provided by Dr Paul Ellis (PDB file 1fsl; Ollis *et al.*, 1983; Ellis *et al.*, 1997). Two molecules related by a 2-fold non-crystallographic symmetry axis were present in the asymmetric unit. After several cycles of refinement, manual fitting and solvent addition, the crystallographic R-factor converged to 19.8% (R_{free} = 29.7%) with root-mean-square deviations of 0.018 Å in bond lengths, 1.93° in bond angles, 21.36° in dihedral angles and 2.29° in improper angles. Molecule a of the two molecules in the asymmetric unit was used to produce Figures 2 and 3. The coordinates of the structure of wild-type metLba-acetate have been assigned number 1bin in the Protein Data Bank at the Brookhaven National Laboratory.

Spectroscopic and kinetic characterization of wild-type and mutant Lba

Absorbance spectra were measured using a SLM 3000 diode array spectrophotometer. LbCO was prepared by diluting metLb into a cuvette containing 20 mM Tris (pH 8.0) buffer equilibrated with 1 atm CO and an excess of sodium dithionite. MetLb was formed by the reaction of ferrous Lb with potassium ferricyanide followed by chromatography on a Sephadex G-25 column equilibrated in 20 mM Tris (pH 8.0). LbO₂ was produced by the reduction of metLb with sodium dithionite followed by G-25 chromatography in the presence of oxygen.

DeoxyLba was produced by the addition of several granules of sodium dithionite to a cuvette containing metLb in anaerobic buffer.

The rate constants for O₂, CO and NO binding were measured using established rapid mixing and laser photolysis methods (Olson, 1981; Quillin *et al.*, 1995; Moore & Gibson, 1976). O₂ and CO association rate constants at pH 5.0 were measured in 100 mM sodium acetate (pH 5.0). O₂ dissociation at pH 5 was measured by mixing a LbO₂ sample in 10 mM potassium phosphate (pH 7.0) with 100 mM sodium acetate (pH 5.0), equilibrated with 1 atm CO (Appleby *et al.*, 1983).

Rate constants for hemin dissociation and auto-oxidation were measured as described by Hargrove *et al.* (1994) and Brantley *et al.* (1993), respectively. Hemin dissociation from H61L Lba was measured in a Gibson-Dionex stopped-flow apparatus as described by Hargrove *et al.* (1996a). Mes (2-(N-morpholino)ethanesulfonic acid) was used as the buffer at pH 5 for measurement of hemin dissociation because acetate coordination to metLb markedly inhibits this reaction.

Acknowledgments

This work was supported by United States Public Health Service grants GM35649 and HL-47020, grant C-612 from the Robert A. Welch Foundation, and the W. M. Keck Foundation (J.S.O.), grant 003604-025 from the State of Texas Advanced Technology Program (G.N.P./J.S.O.), a National Institutes of Health Postdoctoral Fellowship ARO8355 (E.A.B.), NSF grant OSR-92552255 (R.V.K.), USDA-CREES 95-37305-2441 (R.V.K./G.S.), and the Nebraska Research Initiative (G.S.). R.A.P. was partially supported by DGAPA (Dirección General de Asuntos del Personal Académico)-UNAM and CoNaCyT (Consejo Nacional de Ciencia y Tecnología), Mexico. We also thank M. Ikeda-Saito, C. Appleby and A. Wilkinson for reading the original manuscript and making helpful suggestions.

References

- Appleby, C. A. (1984). Leghemoglobins and *Rhizobium* respiration. *Annu. Rev. Plant Physiol.* **35**, 443–478.
- Appleby, C. A. (1992). The origin and functions of haemoglobin in plants. *Sci. Prog.* **76**, 365–398.
- Appleby, C. A., Bradbury, J. H., Morris, R. J., Wittenberg, B. A., Wittenberg, J. B. & Wright, P. E. (1983). Leghemoglobin. *J. Biol. Chem.* **258**, 2254–2259.
- Brantley, R. E., Smerdon, S. J., Wilkinson, A. J., Singleton, E. W. & Olson, J. S. (1993). The mechanism of autooxidation of myoglobin. *J. Biol. Chem.* **268**, 6995–7010.
- Brünger, A. (1987). Crystallographic R factor refinement by molecular dynamics. *Science*, **235**, 458–460.
- Ellis, P. J., Appleby, C. A., Guss, J. M., Hunter, W. N., Ollis, D. L. & Freeman, H. C. (1997). The crystal structure of ferric soybean leghemoglobin a nicotinate at 2.3 Å resolution. *Acta. Crystallog. sect. D*, in the press.
- Fuchsman, W. H. & Appleby, C. A. (1979). CO and O₂ complexes of soybean leghemoglobins: pH effects upon infrared and visible spectra. Comparisons with CO and O₂ complexes of myoglobin and hemoglobin. *Biochemistry*, **18**, 1309–1321.

- Gibson, Q. H., Wittenberg, J. B., Wittenberg, B. A., Bogusz, D. & Appleby, C. A. (1989). The kinetics of ligand binding to plant hemoglobins. *J. Biol. Chem.* **264**, 100–107.
- Hargrove, M. S. & Olson, J. S. (1996). The stability of holomyoglobin is determined by heme affinity. *Biochemistry*, **35**, 11310–11318.
- Hargrove, M. S., Singleton, E. W., Quillin, M. L., Ortiz, L. A., Phillips, G. N., Jr, Mathews, A. J. & Olson, J. S. (1994). His(E7)Tyr as a reagent for measurement of heme loss. *J. Biol. Chem.* **269**, 4207–4214.
- Hargrove, M. S., Barrick, D. & Olson, J. S. (1996a). The association rate constant for heme binding to globin is independent of protein structure. *Biochemistry*, **35**, 11293–11299.
- Hargrove, M. S., Wilkinson, A. J. & Olson, J. S. (1996b). Structural factors governing heme dissociation from metmyoglobin. *Biochemistry*, **35**, 11300–11309.
- Harutyunyan, E. H., Safonova, T. N., Kuranova, I. P., Popov, A. N., Teplyakov, A. V., Obmolova, G. V., Rusakov, A. A., Vainshtein, B. K., Dodson, G. G., Wilson, J. C. & Perutz, M. F. (1995). The structure of deoxy- and oxy-leghaemoglobin from lupin. *J. Mol. Biol.* **251**, 104–115.
- Hyldig-Nielsen, J. J., Jensen, E. O., Paludan, K., Wiborg, O., Garret, R., Jorgensen, P. & Marcker, K. A. (1982). The primary structures of two leghemoglobin genes from soybean. *Nucl. Acids Res.* **10**, 689–695.
- Ikeda-Saito, M., Hori, H., Inubushi, T. & Yonetani, T. (1981). Studies on cobalt myoglobins and hemoglobins. *J. Biol. Chem.* **256**, 10267–10271.
- Ji, L., Becana, M., Sarath, G. & Klucas, R. V. (1994a). Cloning and sequence analysis of a cDNA encoding ferric leghemoglobin reductase from soybean nodules. *Plant Physiol.* **104**, 453–459.
- Ji, L., Becana, M., Sarath, G., Shearman, L. & Klucas, R. V. (1994b). Over-expression in *Escherichia coli* and characterization of a soybean ferric leghemoglobin reductase. *Plant Physiol.* **106**, 203–209.
- Jun, H.-K., Sarath, G. & Wagner, F. W. (1994). Detection and purification of modified leghemoglobins from soybean root nodules. *Plant Sci.* **100**, 31–40.
- Kloek, A. P., Yang, J., Mathews, F. S., Frieden, C. & Goldberg, D. E. (1993). The tyrosine B10 hydroxyl is crucial for oxygen avidity of *Ascaris* hemoglobin. *J. Biol. Chem.* **269**, 2377–2379.
- Kong, S. B., Cutnell, J. D. & La Mar, G. N. (1983). Proton nuclear magnetic resonance study of the dynamic stability of the heme pocket of soybean leghemoglobin. *J. Biol. Chem.* **258**, 3843–3849.
- Kunkel, T. A. (1985). Rapid and efficient site-specific mutagenesis without phenotypic selection. *Proc. Natl Acad. Sci. USA*, **82**, 488–492.
- Lee, H. C., Wittenberg, J. B. & Peisach, J. (1993). Role of hydrogen bonding to bound dioxygen in soybean leghemoglobin. *Biochemistry*, **32**, 11500–11506.
- Martin, K. D., Saari, L., Guang-Xin, W., Wang, T., Parkhurst, L. J. & Klucas, R. V. (1990). Kinetics and thermodynamics of oxygen, CO, and azide binding by the subcomponents of soybean leghemoglobin. *J. Biol. Chem.* **265**, 19588–19593.
- Mathews, A. J., Rohlfs, R. J., Olson, J. S., Tame, J., Renaud, J. P. & Nagai, K. (1989). The effects of E7 and E11 mutations on the kinetics of ligand binding to R state human hemoglobin. *J. Biol. Chem.* **264**, 16573–16583.
- Mathews, A. J., Olson, J. S., Renaud, J.-P., Tame, J. & Nagai, K. (1991). The assignment of carbon monoxide association rate constants to the α and β subunits in native and mutant human deoxyhemoglobin tetramers. *J. Biol. Chem.* **266**, 21631–21693.
- Mims, M. P., Porras, A. G., Olson, J. S., Noble, R. W. & Peterson, J. A. (1983). Ligand binding to heme proteins. *J. Biol. Chem.* **258**, 14219–14232.
- Moore, E. G. & Gibson, Q. H. (1976). Cooperativity in the dissociation of nitric oxide from hemoglobin. *J. Biol. Chem.* **251**, 2788–2794.
- Nagai, K., Luisi, B., Shih, D., Miyazaki, G., Imai, K., Poyart, C., De Young, A., Kwiatkowski, L., Noble, R. W., Lin, S.-H. & Yu, N.-T. (1987). Distal residues in the oxygen binding site of haemoglobin studied by protein engineering. *Nature*, **329**, 858–860.
- Ollis, D. L., Appleby, C. A., Colman, P. M., Cutten, A. E., Guss, J. M., Venkatappa, M. P. & Freeman, H. C. (1983). Crystal structure of soybean ferric leghaemoglobin *a* nicotinate at a resolution of 3.3 Å. *Aust. J. Chem.* **36**, 451–468.
- Olson, J. S. (1981). Stopped-flow, rapid mixing measurements of ligand binding to hemoglobin and red cells. *Methods Enzymol.* **76**, 631–652.
- Olson, J. S. & Phillips, G. N., Jr (1996). Kinetic pathways and barriers for ligand binding to myoglobin. *J. Biol. Chem.* **271**, 17593–17596.
- Phillips, G. N., Jr, Arduini, R. M., Springer, B. A. & Sligar, S. G. (1990). Crystal structure of myoglobin from a synthetic gene. *Proteins: Struct. Funct. Genet.* **7**, 358–365.
- Quillin, M. L., Arduini, R. M., Olson, J. S. & Phillips, G. N., Jr (1993). High-resolution crystal structures of distal histidine mutants of sperm whale myoglobin. *J. Mol. Biol.* **234**, 140–155.
- Quillin, M. L., Li, T., Olson, J. S., Phillips, G. N., Jr, Dou, Y., Ikeda-Saito, M., Regan, R., Carlson, M., Gibson, Q. H., Li, H. & Elber, R. (1995). Structural and functional effects of apolar mutations of the distal valine in myoglobin. *J. Mol. Biol.* **245**, 416–436.
- Reisberg, P. I. & Olson, J. S. (1980). Rates of isonitrile binding to the isolated α and β subunits of human hemoglobin. *J. Biol. Chem.* **255**, 4151–4158.
- Rohlfs, R. J., Olson, J. S. & Gibson, Q. H. (1988). A comparison of the geminate recombination kinetics of several monomeric heme proteins. *J. Biol. Chem.* **263**, 100–107.
- Saari, L. L. & Klucas, R. V. (1984). Ferric leghemoglobin reductase from soybean root nodules. *Arch. Biochem. Biophys.* **231**, 102–113.
- Sambrook, J., Fritsch, E. F. & Maniatis, T. (1989) *Molecular Cloning: A Laboratory Manual*, 2nd edit, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Springer, B. A. & Sligar, S. G. (1987). High-level expression of sperm whale myoglobin in *Escherichia coli*. *Proc. Natl Acad. Sci. USA*, **84**, 8961–8965.
- Springer, B. A., Egeberg, K. D., Sligar, S. G., Rohlfs, R. J., Mathews, A. J. & Olson, J. S. (1989). Discrimination between oxygen and carbon monoxide and inhibition of autooxidation by myoglobin. *J. Biol. Chem.* **264**, 3057–3060.
- Springer, B. A., Sligar, S. G., Olson, J. S. & Phillips, G. N., Jr (1994). Mechanisms of ligand recognition in myoglobin. *Chem. Rev.* **94**, 699–714.
- Stetzkowski, F., Cassoly, R. & Banerjee, R. (1979). Binding of alkylisocyanides with soybean leghemoglobin. *J. Biol. Chem.* **254**, 11351–11356.
- Stetzkowski, F., Banerjee, R., Marden, M. C., Beece, D. K., Bowne, S. F., Doster, W., Eisenstein, L., Frauenfelder, H., Reinisch, L., Shyamsunder, E. & Jung, C. (1985). Dynamics of dioxygen and carbon monoxide binding to soybean leghemoglobin. *J. Biol. Chem.* **260**, 8803–8809.

Vainshtein, B. K., Arutyunyan, E. G., Kuranova, I. P.,
Borisov, V. V., Sosfenov, N. I., Pavlovskii, A. G.,
Grebenko, A. I. & Konareva, N. V. (1975). X-ray

study of leghemoglobin. I. Purification, crystalliza-
tion, and preparation of derivatives containing
heavy atoms. *Sov. Phys. Crystallog.* **19**, 598–601.

Edited by K. Nagai

(Received 9 September 1996; received in revised form 27 November 1996; accepted 4 December 1996)