



Balanced globin protein expression and heme biosynthesis improve production of human hemoglobin in *Saccharomyces cerevisiae*

Lifang Liu^a, José L. Martínez^a, Zihe Liu^a, Dina Petranovic^a, Jens Nielsen^{a,b,*}

^a Novo Nordisk Foundation Center for Biosustainability, Department of Chemical and Biological Engineering, Chalmers University of Technology, Kemivägen 10, SE-41296 Göteborg, Sweden

^b Novo Nordisk Foundation Center for Biosustainability, Technical University of Denmark, Fremtidsvej 3, DK-2970 Hørsholm, Denmark

ARTICLE INFO

Article history:

Received 25 October 2012

Received in revised form

19 August 2013

Accepted 4 October 2013

Available online 2 November 2013

Keywords:

Human hemoglobin

Heme biosynthesis

Prosthetic group

Protein production

Saccharomyces cerevisiae

ABSTRACT

Due to limitations associated with whole blood for transfusions (antigen compatibility, transmission of infections, supply and storage), the use of cell-free hemoglobin as an oxygen carrier substitute has been in the center of research interest for decades. Human hemoglobin has previously been synthesized in yeast, however the challenge is to balance the expression of the two different globin subunits, as well as the supply of the prosthetic heme required for obtaining the active hemoglobin ($\alpha_2\beta_2$). In this work we evaluated the expression of different combinations of α and β peptides and combined this with metabolic engineering of the heme biosynthetic pathway. Through evaluation of several different strategies we showed that engineering the biosynthesis pathway can substantially increase the heme level in yeast cells, and this resulted in a significant enhancement of human hemoglobin production. Besides demonstration of improved hemoglobin production our work demonstrates a novel strategy for improving the production of complex proteins, especially multimers with a prosthetic group.

Crown Copyright © 2013 Published by Elsevier Inc. on behalf of International Metabolic Engineering Society. All rights reserved.

1. Introduction

Hemoglobin is the major oxygen transport protein in blood and is found in erythrocytes. The protein is a tetrameric metalloprotein consisting of four globin subunits ($\alpha_2\beta_2$) with each subunit containing a prosthetic heme group. One heme molecule is comprised of a protoporphyrin IX and a ferrous iron which has high affinity for binding oxygen (Belcher et al., 2010; Benesch and Benesch, 1963). Due to the concerns with whole blood for transfusion, e.g. limited resources, cross-typing requirements and transfusion transmitted disease, substantial efforts have been put into the development of safe and effective human blood substitutes (Looker et al., 1992; Wagenbach et al., 1991). Aside from synthetic perfluorocarbon (PFC) polymers (Biro, 1993) and erythrocyte products (which are purified hemoglobin from a variety of mammalian sources), cell-free hemoglobin-based oxygen carrier (HBOC) has been extensively studied for its high efficiency in oxygen transfer and its contribution to the maintenance of osmotic pressure (Lu et al., 2011). The main limitation with this product is protein instability due to tetramer dissociation into dimers, resulting in short retention time and renal toxicity, and high

oxygen affinity in the body caused by absence of a co-factor 2, 3-bis (phospho) glycerate (2,3-BPG) (Kroeger and Kundrot, 1997; Looker et al., 1992; Wagenbach et al., 1991). A series of advances have been achieved including chemical modifications of natural hemoglobin to extend its half-life in circulation (Harris and Palmer, 2008) which have greatly stimulated further studies on HBOC (Lu et al., 2011).

With the development of recombinant DNA technology, it became possible to engineer hemoglobin with genetic engineering, which enabled designed modification to improve its biological properties as well as to enhance its production. A variety of strategies have been applied in different organisms since the late 80s ranging from bacteria (Hoffman et al., 1990) to higher organisms, including plants (Dieryck et al., 1997) and animals (Behringer et al., 1989; Martinez et al., 2012). The main advances include (i) expression of α and β globin chains simultaneously and refold together, with addition of exogenous heme *in vitro* (Hoffman et al., 1990) (ii) Co-expression of the methionine amino-peptidase (Met-AP) gene with the globin genes to reduce the N-terminal methionine in modified globins produced in *Escherichia coli* (Shen et al., 1993) (iii) Site-directed mutations in β globin chain in order to reduce the extreme oxygen affinity due to the lack of 2,3-BPG allosteric regulation (Hernan et al., 1992) (iv) Tandem fusion of two α globins to prevent tetramer disassociation into $\alpha\beta$ dimers (Looker et al., 1992) (v) Co-expression of bacterial heme membrane transporter systems to increase heme uptake and further

* Corresponding author at: Novo Nordisk Foundation Center for Biosustainability, Department of Chemical and Biological Engineering, Chalmers University of Technology, Kemivägen 10, SE-41296 Göteborg, Sweden. Fax: +46 31 772 3801.
E-mail address: nielsenj@chalmers.se (J. Nielsen).

enhancement of hemoglobin production (Graves et al., 2008). In addition to this, several recent studies showed that α -hemoglobin stabilizing protein (AHSP) could prevent α globin precipitation and hereby have a positive effect on hemoglobin production (Feng et al., 2004; Feng et al., 2005; Vasseur-Godbillon et al., 2006).

The yeast *S. cerevisiae* is a well-known cell factory for production of fuels (Lee et al., 2008), chemicals (Chemler et al., 2006; Nevoigt, 2008) and heterologous proteins, especially industrial enzymes and pharmaceuticals (Hou et al., 2012; Martinez et al., 2012). There are several reasons why yeast can potentially be a highly competitive expression host for production of human hemoglobin: it is general regarded as safe (GRAS organism) which avoids the risk of pathogen contaminations from human or animal resources, it is able to perform post-translational modifications, including correct processing the globin termini, folding and subunit assembling (Porro et al., 2005), being the prerequisite for proper reconstitution of a soluble heme-containing tetramer in vivo. Furthermore, being a long-term eukaryal model organism, numerous studies have been carried out on *S. cerevisiae* from which extensive genome and physiological information have been generated (Petranovic and Nielsen, 2008). For these reasons several recombinant proteins such as human insulin (Novo Nordisk), human bovine serum albumin (Novozymes) and hepatitis vaccines (GlaxoSmithKline, Merck) among others, have been industrially produced and approved for medical applications in *S. cerevisiae* (Martinez et al., 2012; Walsh, 2010).

Hemoglobin synthesis is highly conserved through evolution and the non-globin part, the prosthetic group protoporphyrin IX, is common among divergent species such as human and yeast, so one can utilize the endogenous yeast heme pathway for the production of human recombinant hemoglobin (Kumar, 1995). Studies suggest that recombinant hemoglobin produced in yeast remains fully functional and shows identical rate of dissociation as the native human hemoglobin (Coghlan et al., 1992). However, with improved globin production it is questionable whether endogenous heme production is sufficient. Thus, it has been found that supplementation with heme can accelerate hemoglobin synthesis in immature cultured erythroid cells, which suggests that heme availability is a rate-controlling step for hemoglobin production (Fibach et al., 1995). Additionally, heme association to globins was suggested to occur co-translationally and could therefore facilitate the formation of the proper tertiary structure on the ribosome (Komar et al., 1997). A coordinated optimization of endogenous heme level is therefore expected to improve heterologous hemoglobin production in yeast cells.

The heme biosynthesis pathway in yeast (Fig. 1) contains eight steps starting from 5-aminolevulinic acid (ALA) synthesis from succinyl-CoA and glycine (catalyzed by Hem1p) in mitochondria. After being transported to the cytosol to form porphobilinogen (PBG) (catalyzed by Hem2p), the intermediate is further converted to hydroxymethylbilane (catalyzed by Hem3p) and then uroporphyrinogen III (by Hem4p) and finally into coproporphyrinogen III (catalyzed by Hem12p). The coproporphyrinogen III is transported back to the mitochondria and converted to protoporphyrinogen IX by Hem13p following synthesis of protoporphyrin IX (by Hem14p). The last step is the incorporation of an iron atom into protoporphyrin IX to form the heme molecule catalyzed by Hem15p.

In this study, we evaluated the effect of metabolic engineering the heme biosynthetic pathway on heterologous hemoglobin production (Fig. 1): we overexpressed, individually or in different combinations (with promoters of different strengths), three genes coding for enzymes that represent confirmed (*HEM2* and *HEM3*) or suspicious (*HEM12*) rate-limiting steps in the heme biosynthetic pathway (Hoffman et al., 2003), and additionally co-expressed α and β globin genes in different ratios. Using this approach we could evaluate a large number of combinations with engineering the heme

biosynthesis and globin subunit co-expression and hereby identify the optimal combination for high level human hemoglobin production.

2. Materials and methods

2.1. Plasmids and strains

S. cerevisiae CEN.PK 113-11C (*MATa his3 Δ 1 ura3–52 MAL2-8c SUC2*) (Entian and Kötter, 1998) was used as the host strain for all the experiments performed in this study. Plasmid constructions were performed following standard molecular biology techniques using *E. coli* DH5 α (Bethesda Research Laboratories). All yeast transformations were performed following standard lithium acetate method (Daniel Gietz and Woods, 2002) and selected on synthetic dextrose media SD-His (heme overexpression strains), SD-Ura (globin overexpression strains), and SD-His-Ura (heme and globin double overexpression strains) respectively. For cultivation, yeast strains were grown in liquid SD-His, SD-Ura and SD-His-Ura accordingly.

All the heme overexpression plasmids were constructed using a bidirectional expression plasmid pLYC04 (Chen et al., 2013). The heme genes were amplified from the chromosomal DNA of CEN.PK 113-11C (Table S1). Plasmids H2, H3 and H12 were constructed by inserting the corresponding genes into the expression cassette of pLYC04 under the *TEF1* promoter. Plasmids H2H3 and H2H12 were constructed by inserting *HEM3* and *HEM12* respectively into H2 under the *PGK1* promoter. H3H2H12 was built by inserting a *HEM3* expression cassette amplified from H3 into H2H12. The globin overexpression plasmids were constructed from a different bidirectional expression plasmid pSP-GM1 (Chen et al., 2012). The DNA sequences (with restriction sites at both ends), namely *HBA* (coding α), *HBB* (coding β) and *HBAA* (coding di- α) were synthesized by GeneScript, Inc., and codon optimized to facilitate their expression in *S. cerevisiae*. The *HBAA* sequence coding di- α was made by linking two *HBA* gene sequences (first one without stop codon) with one glycine linker. The *HBB* fragment was digested with *Sall* and *KpnI* and ligated under the *PGK1* promoter of pSP-GM1 first. Then the *HBA* and *HBAA* fragments were digested by *NotI* and *SacI* and ligated under the *TEF1* promoter of pSP-GM1 to make B/A and B/AA respectively. Plasmid B/A/A was made by inserting the *HBA* expression cassette amplified from B/A into plasmid B/A between *PstI* and *MfeI*. Plasmid B/AA/B was constructed by inserting the *HBB* expression cassette amplified from B/A into the *MfeI* site of plasmid B/AA. All the primers and the synthesized globin gene sequences are listed in the supplementary file (Table S1 and S2).

CEN.PK 113-11C was transformed with the heme and globin overexpression plasmids, resulting in different recombinant host strains (Fig. 1B, Table 1).

2.2. Coproporphyrin assay

Coproporphyrin levels were measured by HPLC using the HPLC Porphyrin Kit from ALPCO Diagnostics Inc. Strains transformed with the heme overexpression plasmids were cultured in SD-His (containing 6.7 g/L of yeast nitrogen base w/o amino acids (Difco Laboratories, Sparks, MD), 0.77 g/L of complete supplement mixture (CSM, w/o histone) (MP Biomedicals, Solon, OH) and 2% glucose) medium at 30 °C with constant shaking at 180 rpm during 96 h before being harvested. Cell pellets collected from 35 ml of culture were resuspended into 1 mL of 0.6% HCl after being washed twice with ddH₂O. The suspended cell pellets were kept in dark at 4 °C before being loaded onto HPLC (PU-2089 Plus pump, AS-2057 Plus auto sampler, and FP-920 fluorescence




Fig. 1. Engineering strategy for optimizing the production of human hemoglobin. (A) Strategy overview: Yeast genes *HEM2*, *HEM3* and *HEM12* (highlighted in red) in heme biosynthesis pathway were overexpressed for an improved level of heme in cell cytosol. Human α and β globins were overproduced on plasmid in cytosol. Correct folding and assembling resulted in completed hemoglobin, failure in assembling resulted in degradation. (B) Strain construction: CEN.PK 113-11C as the expression host being transformed with i) heme overexpression plasmids: gene *HEM2*, *HEM3* and *HEM12* were inserted into the expression cassette *TEF1-ADH1*, either individually or in different combinations in the cassettes *TEF1-ADH1* and *PGK1-CYC1* on pLYC04. ii) globin overexpression plasmids: *HBA*, *HBB*, and *HBAA* were inserted into pSP-GM1 in the cassettes of *TEF1-ADH1* and *PGK1-CYC1* in different combinations. iii) heme and globin overexpression plasmids: The top three best heme/coproporphyrin producing plasmids H3, H12 and H3H2H12 were transformed in different combinations together with the globin overexpression plasmids B/A, B/AA, B/A/A and B/AA/B. Hemoglobin structure was modified from (Lu et al., 2011). Heme biosynthesis pathway was modified from (Hoffman et al., 2003). (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

detector, JASCO). The eluents, standard, and reversed phased C_{18} -column were included in the kit and the running conditions were referred to the kit.

2.3. Heme/porphyrin determination assay

The intracellular heme concentration was measured using a protocol derived from (Sassa, 1976) and slightly modified from (Michener et al., 2012). Cells were grown to a desired density and harvested at $OD_{600} \times mL = 8$ (e.g. if $OD_{600} = 2$, 4 mL of culture is harvested). The cell pellets were resuspended in 500 μ L of 20 mM oxalic acid and then kept in dark at 4 $^{\circ}$ C overnight. 500 μ L of preheated 2 M oxalic acid was added and half of the mixture was taken into an amber tube and heated up to 95–98 $^{\circ}$ C for 30 min during which the other half mixture was kept at room temperature. 200 μ L of clear supernatant was transferred into a black 96

plate and fluorescence with excitation at $\lambda = 400$ nm and emission at $\lambda = 600$ nm was monitored. The total porphyrin level (bound heme and free porphyrins) corresponds to the heated sample, and the free porphyrins level corresponds to the non-heated sample.

2.4. SDS-PAGE assay

For protein extraction, yeast cell samples were taken at the indicated times and cells were harvested by centrifugation. Pellets were resuspended in 2 ml of NaOH containing 7% 2-mercaptoethanol plus 2 ml of trichloroacetic acid (TCA) and incubated 2 min at room temperature. After incubation, cells were washed with 1 M Tris-HCl (pH 7.5) and centrifuged. The cell pellet was resuspended in loading buffer (SDS 3%, Glycerol 10%, 2-mercaptoethanol 5%, Tris pH 6.8 0.0625 M, 0.01 mg bromophenol blue) and heated up to 95 $^{\circ}$ C during 5 min. Cells were discarded by centrifugation and the remaining protein

Table 1

Plasmids and strains used in this study.

Plasmid/Strain	Description	Marker	Reference
Heme over-expression plasmids			
pIYC04	Empty plasmid with bidirectional gene expression cassettes (<i>TEF1-ADH1</i> and <i>PGK1-CYC1</i> , 2 μ)	HIS3	(Chen et al., 2013)
H2	pIYC04 inserted with <i>HEM2</i> under <i>TEF1</i> promoter	HIS3	This study
H3	pIYC04 inserted with <i>HEM3</i> under <i>TEF1</i> promoter	HIS3	This study
H12	pIYC04 inserted with <i>HEM12</i> under <i>TEF1</i> promoter	HIS3	This study
H2H3	H2 inserted with <i>HEM3</i> under <i>PGK1</i> promoter	HIS3	This study
H2H12	H2 inserted with <i>HEM12</i> under <i>PGK1</i> promoter	HIS3	This study
H3H2H12	H2H12 inserted with an <i>HEM3</i> expression cassette under <i>TEF1</i> promoter	HIS3	This study
Globin over-expression plasmids			
pSP-GM1	Empty plasmid with bidirectional gene expression cassettes (<i>TEF1-ADH1</i> and <i>PGK1-CYC1</i> , 2 μ)	URA3	(Chen et al., 2012)
B/A	pSP-GM1 inserted with <i>HBA</i> under <i>TEF1</i> and <i>HBB</i> under <i>PGK1</i>	URA3	This study
B/AA	pSP-GM1 inserted with <i>HBA-Gly-HBA</i> under <i>TEF1</i> and <i>HBB</i> under <i>PGK1</i>	URA3	This study
B/A/A	B/A inserted with a <i>HBA</i> over-expression cassette under <i>TEF1</i>	URA3	This study
B/AA/B	B/AA inserted with a <i>HBB</i> over-expression cassette under <i>PGK1</i>	URA3	This study
Engineered strains for heterologous hemoglobin production			
H3 + B/A	CEN.PK 113-11C transformed with plasmids H3 and B/A		This study
H3 + B/AA	CEN.PK 113-11C transformed with plasmids H3 and B/AA		This study
H3 + B/AA/B	CEN.PK 113-11C transformed with plasmids H3 and B/AA/B		This study
H3 + B/A/A	CEN.PK 113-11C transformed with plasmids H3 and B/A/A		This study
H3 + pSP-GM1	CEN.PK 113-11C transformed with plasmids H3 and pSP-GM1		This study
H12 + B/A/A	CEN.PK 113-11C transformed with plasmids H12 and B/A/A		This study
pIYC04 + B/A/A	CEN.PK 113-11C transformed with plasmids pIYC04 and B/A/A		This study
pIYC04 + pSP-GM1	CEN.PK 113-11C transformed with plasmids pIYC04 and pSP-GM1		This study

Note: slash separates the globin overexpression cassettes.

suspension was used for loading into gradient (4–12%) polyacrylamide bis–tris gels (Novex, Invitrogen) at the selected protein concentrations. The marker ladder used was Novex Sharp Pre-stained Protein Standard from Invitrogen. Software GelPro Analyzer (Media Cybernetics) was used for quantification of band intensities.

2.5. Assay for active hemoglobin

Washed cell pellets were resuspended into 500 μ l of 100 mM of PBS buffer supplemented with 1 $\times$ protease inhibitor (Halt Protease Inhibitor Cocktail, Thermo), pH 7.0. The cell suspension was loaded into a FastPrep™ Lysing tube with an equal volume of glass beads (425–600 μ m diameter, Sigma). Cells were broken by applying 8 pulses of 20 s with the speed of 6 m/s using a FastPrep 24 instrument (MP, Biomedicals, LLC). The sample tubes were placed on ice for 3 min between pulses. Clear supernatants were harvested by centrifugation at full speed for active hemoglobin assay and protein quantification. Lipid removal agent Cleanascite (X2555-100, BIOTECH SUPPORT GROUP) was added according to product description in case of unclear suspensions in the supernatant. Protein concentration was measured by BCA protein assay kit (Cat 23,225, Pierce). The active hemoglobin was measured using the QuantiChrom™ Hemoglobin Assay Kit (DIHEMOGLOBIN-250, Clonagen), based on an improved Triton/NaOH method, in which the hemoglobin is converted into a uniform colored end product that can be measured by spectrophotometer.

2.6. Relative gene transcription levels by qPCR

Yeast cells were cultured at 30 °C with shaking at 180 rpm in shake flask containing 50 ml of SD-His. Total RNA was isolated from 10 ml of cells with OD₆₀₀ around 1. The cells were disrupted by Fast-Prep 24 (MP-Biomedicals) using glass beads (Lysing matrix C, MP-Biomedicals). Extracted RNA was purified using the RNeasy kit (Qiagen). 200 ng of total RNA was used to synthesize cDNA from which 2 μ l of the synthesized cDNA was used as the template for the qPCR reaction to a final reaction volume of 20 μ l, using the Brilliant III Ultra Fast SYBRGreen QPCR Master mix (Stratagene) in

Mx3005P QPCR System (Agilent Technologies). Cycle thresholds (Ct) were normalized to the Ct value of *S. cerevisiae* ACT1.

2.7. Batch fermentations

Fresh colonies were picked to inoculate into 50 mL of SD- His-Ura media (containing 6.7 g/L of yeast nitrogen base w/o amino acids (Difco Laboratories, Sparks, MD), 0.75 g/L of complete supplement mixture (CSM, w/o histone and uracil) (MP Biomedicals, Solon, OH) and 2% glucose) to make seed cultures. After growing for 24 h at 30 °C at 180 rpm, the seed cultures were inoculated into the bioreactors with an initial OD₆₀₀ of 0.1. Fermentations were performed in DasGip 1.0-l stirrer-pro vessels (DASGIP GmbH, Germany) with a working volume of 600 mL in SD-His-Ura media. The temperature was controlled at 30 °C with agitation at 600 rpm and aeration at 1 vvm. Dissolved oxygen was controlled above 30% throughout the experiment. The impeller speed was automatically adjusted in response to the sensor. The pH was maintained at 6.0 using 2 M KOH in response to the pH sensor (Mettler Toledo, Switzerland). All fermentations were performed in biological duplicates. Dry cell weight was measured by filtering 5 mL of cell culture through a 0.45 μ m filter membrane and calculating the increased weight of the dried filter. Supernatants filtered through 0.45 μ m membrane were loaded to HPX-87G column (BIORAD, USA) on a Dionex Ultimate 3000 HPLC (Dionex Softron GmbH, Germany) to measure the concentrations of glucose, ethanol, and glycerol in the cultures. The samples were running with a flow rate of 0.6 mL/min at 65 °C using 5 mM H₂SO₄ as mobile phase.

3. Results and discussion

To investigate the impact of increased endogenous heme synthesis on heterologous hemoglobin production in *S. cerevisiae*, a series of recombinant strains were constructed by transforming different expression plasmids in the yeast strain CEN.PK 113-11C (Fig. 1, Table 1).

3.1. Increased endogenous heme level by overexpression of heme genes

By transforming the heme overexpression plasmids H2, H3, H12, H2H3, H2H12, and H3H2H12 into CEN.PK 113-11C, strains with the engineered heme biosynthesis pathway were constructed (Table 1). The strains were designated with the same names as the corresponding plasmids H2, H3, H12, H2H3, H2H12, and H3H2H12 (Table 1). The cytosolic intermediate of the heme biosynthetic pathway, coproporphyrin was measured in all the engineered strains and used as an indicator for heme and porphyrin levels in the cells (Fig. 2A). All strains were compared with the strain transformed with pLYC04 (empty plasmid) and the fold values were calculated relative to it. Strain H3 showed a four times increase on coproporphyrin level compared to the control strain, pLYC04, followed by H12 and H3H2H12 whose coproporphyrin levels were 3.2 and 2.2 times to pLYC04 respectively (Fig. 2A). In terms of bound heme level, H3 still stood out with almost 5 fold of pLYC04, followed by H3H2H12, 2.8 fold, and H12, 2.4 fold (Fig. 2B).

HEM13 transcription is known to be negatively regulated by oxygen and heme (Zagorec and Labbe-Bois, 1986) through *HAP1* and *ROX1* (Keng, 1992). Overexpression of the genes in the heme biosynthesis pathway (strains H2, H3, H12, H2H3, H2H12 and H3H2H12) resulted in certain improvements on heme levels (Fig. 2B) which might induce feedback inhibition on the synthesis of Hem13p which converts coproporphyrinogen III into protoporphyrinogen IX, causing subsequent accumulation of coproporphyrins (Michener et al., 2012). The coproporphyrin levels (Fig. 2A) may therefore represent reliable indicators of the heme levels in the strains, and Hem13p might be another potential bottleneck in addition to the reported rate-limiting steps catalyzed by Hem2p

and Hem3p. Both driven by *TEF1* promoter, *HEM3* overexpression (H3) showed a better effect than *HEM12* overexpression (H12) on coproporphyrin accumulation suggesting a greater rate-limiting effect of Hem3p than Hem12p during heme biosynthesis. Overexpression of *HEM3* released the bottleneck in this step providing more intermediate for further (catalyzed by Hem4p and Hem12p) conversion to coproporphyrin. On the other hand, also driven by the *TEF1* promoter, *HEM2* overexpression (H2) exhibited a very limited improvement on the coproporphyrin level in the cells although the Hem2p has been confirmed as a rate-controlling enzyme under aerobic conditions (Hoffman et al., 2003; Zhang and Hach, 1999). This is very likely due to the enzymes blocking the subsequent steps, especially Hem3p and Hem12p, were not overexpressed in H2. The point could be supported by the higher coproporphyrin levels presented in H2H3 and H2H12 than in H2. Since *HEM3* in H2H3 and *HEM12* in H2H12 are both under the *PGK1* promoter which has a reported weaker strength than *TEF1* promoter (Nacken et al., 1996) driving *HEM2* in these strains, the steps catalyzed by Hem3p and Hem12p may still being blocked to certain extends which could partly explain the not so obvious improvements on the coproporphyrin levels in these strains. Additional reason could be that with more gene cassettes compulsory for expression, the resource for expressing one gene has to be distributed to more genes and therefore each gene gets lesser/weaker expression. The expressions of *HEM2*, *HEM3* and *HEM12* in H2H3 and H2H12 might have been reduced to certain levels resulting in the minor improvement of coproporphyrin accumulation than H2. Similarly, the expression burden was even heavier for the cells in the H3H2H12 strain and consequently the proteins especially Hem3p which has been indicated to be the most effective enzyme compared to Hem2p and Hem12p to block heme synthesis, were less expressed, which could be the reason that




Fig. 2. Porphyrin/heme levels in heme over-expression strains. (A) Level of coproporphyrin. (B) Level of free porphyrins and bound heme. The relative fold change is calculated based on the fluorescence reads (excitation at $\lambda = 400$ nm and emission at $\lambda = 600$ nm) of the sample strains against the control strain, pLYC04.




Fig. 3. Transcription level of (A) *HEM3* and (B) *HEM12* in the engineered strains transformed with Heme plasmids. In both cases, the relative fold change is calculated based on the transcription levels of the genes (relative to *ACT1*) in the sample strains against the control strain, pLYC04.

H3H2H12 did not show higher or even comparable level of coproporphyrin than H3 and H12.

The transcriptional levels of *HEM3* and *HEM12* were examined and as shown in Fig. 3, the mRNA levels of the heme genes *HEM3* and *HEM12* were much higher in the strains transformed with the corresponding vectors than with the empty vector. The transcriptional level of *HEM3* in strain H3 is extremely high (131.6 folds) correlating to the highest coproporphyrin accumulation in the H3 strain.

The acidic-heat method can discriminate between heme bound to proteins and the free porphyrins in cells. As shown in Fig. 2B, the heme level in H3 was the most abundant followed by H3H2H12 and H12, showing the same trend as that for coproporphyrin. The higher heme level in H3 than in H12 might indicate a bigger flux of coproporphyrins converted to protoporphyrin IX than in H12, even though the coproporphyrins accumulated in H3 were also more than in H12. The free porphyrins include more than coproporphyrins which might explain the higher free porphyrin levels in H12 compared to H3. Both assays indicate that *HEM3* overexpression alone generates the highest heme levels in cells compared to the other strains.

3.2. Co-overexpression of heme and globin constructs increased heterologous hemoglobin production

The top three heme overexpression plasmids H3, H3H2H12 and H12 were selected to be co-transformed together with a range of globin overexpression plasmids B/A, B/AA, B/AA/B, B/A/A and the empty vector pSP-GM1 (Fig. 1B). The combination of α and β globin genes on the globin vectors were designed based on (i) α globin is more unstable than β globin (Vasseur-Godbillon et al., 2006) and (ii) $\alpha\beta$ dissociation can be prevented by linking two α globin genes with a glycine linker (Looker et al., 1992). The expression levels were examined by SDS-PAGE (Fig. 4). By co-overexpression with H3, the globin construct B/A/A showed the highest protein expression, however, there was nearly no band of the right size (around 16 kDa) when H3 was expressed in combination with other globin constructs (Fig. 4A). Very weak signal of α and β globins were detected by ELISA (data not shown) and based on this result, co-overexpressions of B/A/A with the other two heme overexpression plasmids H12 and H3H2H12 were also checked by SDS-PAGE: with heme overexpressed, the globin synthesis was also improved (Fig. 4B). Expression of B/A/A with the empty plasmid pLYC04 did not show clear expression, thus further supporting the hypothesis that endogenous heme levels are insufficient to achieve high heterologous globin production (Smith et al., 2011).

3.3. Effect of B/A/A co-overexpression on heme/porphyrin level

In order to understand if an improved hemoglobin expression is correlated with an increased heme level, the heme/porphyrin level in the strains co-overexpressed with B/A/A was examined.




Fig. 4. SDS-PAGE analysis of the engineered strains with heme and globin co-overexpressed. (A) co-overexpression of H3 with the globin constructs. (B) co-expression of B/A/A with the heme constructs. M stands for the protein ladder, Hb stands for the standard hemoglobin (16 kDa) and the numbers indicate the strains listed in the table on the side. 50 μ g of total protein was loaded per lane.

The relative fold change was calculated relative to the heme level in pLYC04+pSP-GM1. As shown in Fig. 5, both the bound heme level and the free porphyrins level in H3+B/A/A are higher than in H3+pSP-GM1, and in the H3H2H12+B/A/A strain compared with H3H2H12+pSP-GM1. With B/A/A co-overexpressed, we found an increase of 87% with H3 co-overexpression. In erythroid cells, heme export from the mitochondria requires a corresponding synthesis of heme-binding proteins (Lynch et al., 2009). One hypothesis could be that with higher levels of globin expressed, an initial decrease in the pool of free cytosolic heme occurs which induces an increase in the production rate of porphyrins resulting in a global increase of both free porphyrins and consequently the hemoproteins. The bound heme levels that we measured would be thus higher.

3.4. Physiological characterization and active hemoglobin production in batch fermentations

The physiological properties of the recombinant hemoglobin strains were characterized in batch fermentations, as shown in Table 2. The specific growth rates did not vary significantly which were 0.25, 0.28, 0.28 and 0.27 respectively for H3+B/A/A, H3H2H12+B/A/A, pLYC04+B/A/A and pLYC04+pSP-GM1. At the end of cultivation, the percentages of the active hemoglobin in the whole extracted soluble proteins were assayed and calculated. The result showed that the strain H3+B/A/A had the most active hemoglobin accounting for 4.09% of the whole cell soluble protein followed by H3H2H12+B/A/A with 3.82%. There was activity accounted for 2.8% of the whole cell protein in the pLYC04+pSP-GM1 strain, most probably contributed by the endogenous flavo-hemoglobin (Zhao et al., 1996).

The specific growth rate of strain H3+B/A/A was slightly lower when compared with other strains. It also had the most soluble hemoglobin production and less biomass yield. Declines on




Fig. 5. Effects of B/A/A plasmid co-overexpression on heme/porphyrin level. The relative fold change is calculated based on the fluorescence reads (excitation at $\lambda=400$ nm and emission at $\lambda=600$ nm) of the sample strains to that of the control strain, pLYC04.

Table 2

The physiological parameters and soluble hemoglobin concentration.

Strain	$\mu_{\max}^a$	Y_{sx}^b	Y_{se}^b	Y_{sc}^b	[Hb]% ^c
H3+B/A/A	0.25 ± 0.001	0.10 ± 0.005	0.29 ± 0.01	0.04 ± 0.006	4.09 ± 0.04
H3H2H12+B/A/A	0.28 ± 0.01	0.11 ± 0.006	0.30 ± 0.004	0.04 ± 0.001	3.82 ± 0.05
pIYC04+B/A/A	0.28 ± 0.006	0.12 ± 0.003	0.34 ± 0.004	0.04 ± 0.001	3.14 ± 0.06
pIYC04+pSP-GM1	0.27 ± 0.002	0.11 ± 0.004	0.32 ± 0.003	0.04 ± 0.001	2.8 ± 0.05


^a The maximum specific growth rate (h^{-1}) on glucose.^b Yield of biomass(X), ethanol(E), and glycerol(G) from glucose (S)(g/g).^c Percentage of active hemoglobin on whole soluble cell protein (g/g).

Fig. 6. Heme and hemoglobin production during batch fermentations. (A) Offgas profile: showing that glucose phase finished at around 20 h and ethanol phase started at around 36 h. (B) Heme and porphyrin levels in strains after inoculation for 20 and 36 h respectively. (C) SDS-PAGE analysis of the globin patterns produced at 20 and 36 h respectively. 4, 14, 20 represent strains H3+B/A/A, H3H2H12+B/A/A and pIYC04+pSP-GM1 respectively. a and b represent 20 and 36 h after inoculation. 25 μ g protein loaded.

specific growth rate due to plasmid-encoded recombinant gene expression have been extensively noticed in bacteria, yeast and filamentous fungi (Hoffmann and Rinas, 2004; Pakula et al., 2005; Tyo et al., 2012) all pointing to a higher stress and impairment to cellular process. In the strain H3+B/A/A, a coordinated over production of heme and globin units have been achieved which further assembled to the correct tetramer. This process might result in a flux switching from normal cellular processes to recombinant hemoglobin production, and consequently affect cell performance. The heme and globins production is less effective in the other two recombinant strains thus the normal growth have not been impaired significantly.

The cell growth reached the end of the glucose phase and the beginning of the ethanol phase at around 20–36 h after inoculation respectively (Fig. 6A). Interestingly, we found that the heme level in H3+B/A/A significantly decreased before the ethanol phase started (Fig. 6B) indicating a negative correlation between heme/porphyrin synthesis and respiration. In contrast, in H3H2H12+B/A/A and pIYC04+pSP-GM1, the heme level kept improving after the glucose phase was finished with a decrease in the free porphyrin level suggesting that porphyrin synthesized in the glucose phase were gradually converted into heme. The correlation between the heme levels and the hemoglobin production was also measured by SDS-PAGE (Fig. 6C). Clearly, the level of hemoglobin produced in the glucose phase is higher than in the beginning of the ethanol phase (about 1.3 fold) for H3+B/A/A, confirming that efficient heterologous hemoglobin production requires a low rate of respiration, as it occurs during the glucose consumption phase in the fermentation process.

4. Conclusion

Previous studies have reported successful expression of recombinant human hemoglobin in the yeast *S. cerevisiae* (Adachi et al., 1992; Coghlan et al., 1992; Martin de Llano et al., 1993; Ogden et al., 1994; Wagenbach et al., 1991). In this study, yeast heme

synthesis pathway was engineered for the first time to facilitate the synthesis of recombinant hemoglobin, resulting in an increase of 1.3 times (H3+B/A/A against pIYC04+B/A/A) in hemoglobin production. This work has generated a plasmid platform for further improving heterologous hemoglobin production in yeast. Further improvement might be achieved by engineering the heme membrane transport and/or the ion trans-membrane pathway (e.g. ferric reductase and permease-oxidase complex required for iron uptake in *S. cerevisiae*) (Dancis et al., 1990; Ginzburg et al., 2009; Graves et al., 2008; Stearman et al., 1996; Villarreal et al., 2008). In addition to hemoglobin, a similar strategy can be applied to enhance the production of other hemoproteins, e.g. cytochromes, catalases and P450 enzymes which are powerful catalysts for synthesis of drug metabolites (Jung et al., 2011). This work is further demonstrating the possibility for efficient production of complex proteins (e.g. multimer with a prosthetic group), as we have demonstrated that the production of a prosthetic group can be increased through metabolic engineering and hence eliminate the need for exogenous supplementation with expensive precursors.

Acknowledgment

The authors would like to thank J. K. Michener for assistance with heme measurement experiment. We thank the financial support from the European Research Council ERC project INSYSBIO (Grant No. 247013), the Novo Nordisk Foundation, the Chalmers Foundation, and the Knut and Alice Wallenberg Foundation

Appendix A. Supporting information

Supplementary data associated with this article can be found in the online version at <http://dx.doi.org/10.1016/j.ymben.2013.10.010>.

References

- Adachi, K., Konitzer, P., Lai, C.H., Kim, J., Surrey, S., 1992. Oxygen binding and other physical properties of human hemoglobin made in yeast. *Protein Eng.* 5, 807–810.
- Behringer, R.R., Ryan, T.M., Reilly, M.P., Asakura, T., Palmiter, R.D., Brinster, R.L., Townes, T.M., 1989. Synthesis of functional human hemoglobin in transgenic mice. *Science* 245, 971.
- Belcher, J.D., Beckman, J.D., Balla, G., Balla, J., Vercellotti, G., 2010. Heme degradation and vascular injury. *Antioxid. Redox Signal.* 12, 233–248.
- Benesch, R., Benesch, R.E., 1963. Some relations between structure and function in hemoglobin. *J. Mol. Biol.* 6, 498–505.
- Biro, G.P., 1993. Perfluorocarbon-based red blood cell substitutes. *Transfus. Med. Rev.* 7, 84–95.
- Chemler, J.A., Yan, Y., Koffas, M.A., 2006. Biosynthesis of isoprenoids, polyunsaturated fatty acids and flavonoids in *Saccharomyces cerevisiae*. *Microbiol. Cell Fact.* 5, 20.
- Chen, Y., Daviet, L., Schalk, M., Siewers, V., Nielsen, J., 2013. Establishing a platform cell factory through engineering of yeast acetyl-CoA metabolism. *Metab. Engineering* 15, 48–54.
- Chen, Y., Partow, S., Scalcinati, G., Siewers, V., Nielsen, J., 2012. Enhancing the copy number of episomal plasmids in *Saccharomyces cerevisiae* for improved protein production. *FEMS Yeast Res.* 12, 598–607.
- Coghlan, D., Jones, G., Denton, K.A., Wilson, M.T., Chan, B., Harris, R., Woodrow, J.R., Ogden, J.E., 1992. Structural and functional characterisation of recombinant human haemoglobin A expressed in *Saccharomyces cerevisiae*. *Eur. J. Biochem.* 207, 931–936.
- Dancis, A., Klausner, R., Hinnebusch, A., Barriocanal, J., 1990. Genetic evidence that ferric reductase is required for iron uptake in *Saccharomyces cerevisiae*. *Mol. Cell. Biol.* 10, 2294–2301.
- Daniel, Gietz, R., Woods, R.A., 2002. Transformation of yeast by lithium acetate/single-stranded carrier DNA/polyethylene glycol method. *Methods Enzymol.* 350, 87–96.
- Dieryck, W., Pagnier, J., Poyart, C., Marden, M.C., Gruber, V., Bournat, P., Baudino, S., Mérot, B., 1997. Human haemoglobin from transgenic tobacco. *Nature* 386, 29.
- Entian, K.D., Kötter, P., 1998. 23 yeast mutant and plasmid collections. *Method Microbiol.* 26, 431–449.
- Feng, L., Gell, D.A., Zhou, S., Gu, L., Kong, Y., Li, J., Hu, M., Yan, N., Lee, C., Rich, A.M., 2004. Molecular mechanism of AHSP-mediated stabilization of α -hemoglobin. *Cell* 119, 629–640.
- Feng, L., Zhou, S., Gu, L., Gell, D.A., Mackay, J.P., Weiss, M.J., Gow, A.J., Shi, Y., 2005. Structure of oxidized α -hemoglobin bound to AHSP reveals a protective mechanism for haem. *Nature* 435, 697–701.
- Fibach, E., Kolli, P., Schechter, A., Noguchi, C., Rodgers, G., 1995. Hemin-induced acceleration of hemoglobin production in immature cultured erythroid cells: preferential enhancement of fetal hemoglobin. *Blood* 85, 2967.
- Ginzburg, Y.Z., Rybicki, A.C., Suzuka, S.M., Hall, C.B., Breuer, W., Cabantchik, Z.I., Bouhassira, E.E., Fabry, M.E., Nagel, R.L., 2009. Exogenous iron increases hemoglobin in beta-thalassemic mice. *Exp. Hematol.* 37, 172–183.
- Graves, P.E., Henderson, D.P., Horstman, M.J., Solomon, B.J., Olson, J.S., 2008. Enhancing stability and expression of recombinant human hemoglobin in *Escherichia coli*: progress in the development of a recombinant HBCO source. *Biochim. Biophys. Acta* 1784, 1471–1479.
- Harris, D.R., Palmer, A.F., 2008. Modern cross-linking strategies for synthesizing acellular hemoglobin-based oxygen carriers. *Biotechnol. Prog.* 24, 1215–1225.
- Hernan, R., Hui, H., Andracki, M., Noble, R., Sligar, S., Walder, J., Walder, R., 1992. Human hemoglobin expression in *Escherichia coli*: importance of optimal codon usage. *Biochemistry* 31, 8619–8628.
- Hoffman, M., Gora, M., Rytka, J., 2003. Identification of rate-limiting steps in yeast heme biosynthesis. *Biochem. Biophys. Res. Commun.* 310, 1247–1253.
- Hoffman, S.J., Looker, D.L., Roehrich, J.M., Cozart, P.E., Durfee, S.L., Tedesco, J.L., Stetler, G.L., 1990. Expression of fully functional tetrameric human hemoglobin in *Escherichia coli*. *Proc. Natl. Acad. Sci. USA* 87, 8521–8525.
- Hoffmann, F., Rinas, U., 2004. Stress induced by recombinant protein production in *Escherichia coli*. *Physiol. Stress Responses Bioprocess.*, 73–92.
- Hou, J., Tyo, K., Liu, Z., Petranovic, D., Nielsen, J., 2012. Engineering of vesicle trafficking improves heterologous protein secretion in *Saccharomyces cerevisiae*. *Metab. Eng.* 14, 120–127.
- Jung, S.T., Lauchli, R., Arnold, F.H., 2011. Cytochrome P450: taming a wild type enzyme. *Curr. Opin. Biotechnol.* 22, 809–817.
- Keng, T., 1992. HAP1 and ROX1 form a regulatory pathway in the repression of HEM13 transcription in *Saccharomyces cerevisiae*. *Mol. Cell. Biol.* 12, 2616.
- Komar, A.A., Kommer, A., Krashennikov, I.A., Spirin, A.S., 1997. Cotranslational folding of globin. *J. Biol. Chem.* 272, 10646–10651.
- Kroeger, K.S., Kundrot, C.E., 1997. Structures of a hemoglobin-based blood substitute: insights into the function of allosteric proteins. *Structure* 5, 227–237.
- Kumar, R., 1995. Recombinant hemoglobins as blood substitutes—a biotechnology perspective. *Proc. Soc. Exp. Biol. Med.* 208, 150–158.
- Lee, S.K., Chou, H., Ham, T.S., Lee, T.S., Keasling, J.D., 2008. Metabolic engineering of microorganisms for biofuels production: from bugs to synthetic biology to fuels. *Curr. Opin. Biotechnol.* 19, 556–563.
- Looker, D., Abbott-Brown, D., Cozart, P., Durfee, S., Hoffman, S., Mathews, A.J., Miller-Roehrkh, J., Shoemaker, S., Trimble, S., Fermi, G., 1992. A human recombinant haemoglobin designed for use as a blood substitute. *Nature* 356, 258–260.
- Lu, X., Ma, G., Su, Z., 2011. Hemoglobin based blood substitutes-preparation technologies and challenges. In: Murray, M.-Y., (Ed.), *Comprehensive Biotechnology*, (second ed.). Academic Press, Burlington, pp. 713–727.
- Lynch, J., Fukuda, Y., Krishnamurthy, P., Du, G.Q., Schuetz, J.D., 2009. Cell survival under stress is enhanced by a mitochondrial ATP-binding cassette transporter that regulates hemoproteins. *Cancer Res.* 69, 5560–5567.
- Martin de Llano, J., Schneewind, O., Stetler, G., Manning, J., 1993. Recombinant human sickle hemoglobin expressed in yeast. *Proc. Nebraska Acad. Sci.* 90, 918.
- Martinez, J.L., Liu, L., Petranovic, D., Nielsen, J., 2012. Pharmaceutical protein production by yeast: towards production of human blood proteins by microbial fermentation. *Curr. Opin. Biotechnol.*
- Michener, J. K., Nielsen, J., Smolke, C.D., 2012. Identification and treatment of heme depletion attributed to overexpression of a lineage of evolved P450 monooxygenases. In: *Proceedings of the National Academy of Sciences*, vol. no. 109, pp. 19504–19509.
- Nacken, V., Achstetter, T., Degryse, E., 1996. Probing the limits of expression levels by varying promoter strength and plasmid copy number in *Saccharomyces cerevisiae*. *Gene* 175, 253–260.
- Nevoigt, E., 2008. Progress in metabolic engineering of *Saccharomyces cerevisiae*. *Microbiol. Mol. Biol. Rev.* 72, 379–412.
- Ogden, J.E., Harris, R., Wilson, M.T., 1994. Production of recombinant human hemoglobin A in *Saccharomyces cerevisiae*. *Methods Enzymol.* 231, 374–390.
- Pakula, T.M., Salonen, K., Uusitalo, J., Penttilä, M., 2005. The effect of specific growth rate on protein synthesis and secretion in the filamentous fungus *Trichoderma reesei*. *Microbiology* 151, 135–143.
- Petranovic, D., Nielsen, J., 2008. Can yeast systems biology contribute to the understanding of human disease? *Trends Biotechnol.* 26, 584–590.
- Porro, D., Sauer, M., Branduardi, P., Mattanovich, D., 2005. Recombinant protein production in yeasts. *Mol. Biotechnol.* 31, 245–259.
- Sassa, S., 1976. Sequential induction of heme pathway enzymes during erythroid differentiation of mouse Friend leukemia virus-infected cells. *J. Exp. Med.* 143, 305–315.
- Shen, T.J., Ho, N.T., Simplaceanu, V., Zou, M., Green, B.N., Tam, M.F., Ho, C., 1993. Production of unmodified human adult hemoglobin in *Escherichia coli*. *Proc. Natl. Acad. Sci. USA* 90, 8108–8112.
- Smith, B.J., Gutierrez, P., Guerrero, E., Brewer, C.J., Henderson, D.P., 2011. Development of a method to produce hemoglobin in a bioreactor culture of *Escherichia coli* BL21(DE3) transformed with a plasmid containing *Plesiomonas shigelloides* heme transport genes and modified human hemoglobin genes. *Appl. Environ. Microbiol.* 77, 6703–6705.
- Stearman, R., Yuan, D.S., Yamaguchi-Iwai, Y., Klausner, R.D., Dancis, A., 1996. A permease-oxidase complex involved in high-affinity iron uptake in yeast. *Science* 271, 1552–1557.
- Tyo, K.E.J., Liu, Z., Petranovic, D., Nielsen, J., 2012. Imbalance of heterologous protein folding and disulfide bond formation rates yields runaway oxidative stress. *BMC Biol.* 10, 16.
- Vasseur-Godbillon, C., Hamdane, D., Marden, M.C., Baudin-Creux, V., 2006. High-yield expression in *Escherichia coli* of soluble human alpha-hemoglobin complexed with its molecular chaperone. *Protein Eng. Des. Sel.* 19, 91–97.
- Villarreal, D.M., Phillips, C.L., Kelley, A.M., Villarreal, S., Villalobos, A., Hernandez, P., Olson, J.S., Henderson, D.P., 2008. Enhancement of recombinant hemoglobin production in *Escherichia coli* BL21(DE3) containing the *Plesiomonas shigelloides* heme transport system. *Appl. Environ. Microbiol.* 74, 5854–5856.
- Wagenbach, M., O'Rourke, K., Vitez, L., Wiczorek, A., Hoffman, S., Durfee, S., Tedesco, J., Stetler, G., 1991. Synthesis of wild type and mutant human hemoglobins in *Saccharomyces cerevisiae*. *Nat. Biotechnol.* 9, 57–61.
- Walsh, G., 2010. Biopharmaceutical benchmarks 2010. *Nat. Biotechnol.* 28, 917–924.
- Zagorec, M., Labbe-Bois, R., 1986. Negative control of yeast coproporphyrinogen oxidase synthesis by heme and oxygen. *J. Biol. Chem.* 261, 2506–2509.
- Zhang, L., Hach, A., 1999. Molecular mechanism of heme signaling in yeast: the transcriptional activator Hap1 serves as the key mediator. *Cell. Mol. Life Sci.* 56, 415–426.
- Zhao, X.J., Raitt, D., Clewell, P.V.B., Kwast, A.S., Poyton, K.E., R. O., 1996. Function and expression of flavohemoglobin in *Saccharomyces cerevisiae*. Evidence for a role in the oxidative stress response. *J. Biol. Chem.* 271, 25131–25138.

Table S1. Primers used in this study.

Primer name	Sequence (5'→3')	Description	Restriction site	Template
HEM2_TEF_fw	GCGGCCGCGCcaaaaccatgcatcacgtg	HEM2 fragment for insertion between <i>TEF1</i> and <i>ADHI</i> on pLYC04 to make plasmid H2	<i>NorI</i> , <i>SacI</i>	Genomic DNA of CENPK113-11C
HEM2_TEF_re	GAGCTCCTATTACAAAGGTAAGCACGT			
HEM3_TEF_fw	GCGGCCGCGCACTATGGGCCCTGAAACT	HEM3 fragment for insertion between <i>TEF1</i> and <i>ADHI</i> on pLYC04 to make plasmid H3	<i>NorI</i> , <i>SacI</i>	Genomic DNA of CENPK113-11C
HEM3_TEF_re	GAGCTCCGGGAATTCTGATCGCGTT			
HEM12_IF_fw	TAATTACAAGCGGCCGCAAAACAATGGGTAACT TTCCAGCTC	HEM12 fragment for insertion between <i>TEF1</i> and <i>ADHI</i> on pLYC04 to make plasmid H12	<i>NorI</i> , <i>SacI</i>	Genomic DNA of CENPK113-11C
HEM12_IF_re	GAATTGTTAATTAAGAGCTCCCTTAGAAATCAG CGCAT			
H3_Sall	ACGCgtcgacACTATGGGCCCTGAAACT	HEM3 fragment for insertion between <i>PGK1</i> and <i>CYC1</i> on plasmid H2 to make H2H3	<i>Sall</i> , <i>SacII</i>	Plasmid H3
H3_SacII	AATGcgcggCGGAATTCTGATCGCGTT			
H12_Sall	ACGCgtcgacACCATGGGTAACTTTCCAGCTC	HEM12 fragment for insertion between <i>PGK1</i> and <i>CYC1</i> on plasmid H2 to make H2H12	<i>Sall</i> , <i>SacII</i>	Plasmid H12
H12_SacII	AATGcgcggCCTTAGAAATCAGCGCAT			
H3_TEF_AscI_re	TTggcgcgcceTAGGAAATTGGAGCGACCTC	HEM3 expression cassette under <i>TEF1</i> promoter for insertion into plasmid H2H12 to make H3H2H12	<i>AscI</i>	Plasmid H3
H3_TEF_AscI	TTggcgcgcceTTGAAGGTACTTCCAGGC			
HBA_MfeI	AGCCcaattgCGCACACACCATAGCTTC	HBA expression cassette under <i>TEF1</i> promoter for insertion into plasmid B/A to make B/A/A	<i>MfeI</i> , <i>PstI</i>	Plasmid B/A
HBA_PstI	TGCActgcagggaattggagcgacctcat			
HBB_MfeI	AGCCcaattgaagctatgtgtgtgcgg	HBB expression cassette under <i>PGK1</i> promoter for insertion into plasmid B/AA to make B/AA/B	<i>MfeI</i>	Plasmid B/A
HBB_MfeI_re	AGCCcaattgAGGATCTTTCGAGCGTCC			

Table S2. Synthetic gene sequences.

HBA

GCGGCCGCACTAGTAAACAATGTCGTCTTGTCTCTGCGGATAAACTAATGTAAAAGCTGCCTGGGGTAAAGTAGGTGCCCCA
CGCCGGTGAATATCACGGTTCAGCACAAAGTCAAGGGTCATGGTAAAAAGGTAGCCGACGCTTTGACCAATGCAGTTGCCAC
GTCGATGACATGCCAAACGCTTTGTCCGCATTAAGTGATTGTCATGCTCACAAGTTGAGAGTAGACCCTGTAACTTCAAGTT
GTTGTCCCATTTGTTGTTAGTTACCTTAGCTGCACACTTGCCAGCCGAATTACTCCTGCTGTCCATGCTCTTTAGACAAGTTT
TTAGCATCCGTTTCCACCGTTTGTACCTCCAAGTATAGAATAAGAGCTCTTAATTAAGTCGACCTCGAG

HBB

CCCGGGGTCGACAAAACAATGTCGTACATTTGACTCCAGAAGAAAAGTCCGCCGTAACAGCATTATGGGGTAAAGTAAACGTAG
ACGAAGTTGGTGGTGAAGCCTTAGGTAGATTGTTAGTTGTCTATCCATGGACCCAAAGATTTTCGAATCTTTGGTGACTTGT
CTACTCCAGACGCCGTAATGGGTAATCCTAAAGTTAAGGCTCATGGTAAAAAGGTTTGGGTGCTTTTCCGATGGTTTGGCA
CACTTAGACAACCTTGAAGGTACTTTGCAACATTGTCTGAATTGCATTGTGATAAGTTACACGTTGATCCTGAAAACCTCAG
ATTGTTGGGTAACGTATTAGTTTGGCTCTTGGCTCATCACTTTGGTAAAGAATTAAGTCCACCTGTTCAAGCAGCCTATCAAAA
GGTTGTCGCAGGTGTTGCTAACGCATTGGCACATAAGTATCACATACTGAGAGCTCTTAATTAAGTCGACCTCGAG

HBAA

GCGGCCGCACTAGTAAACAATGTCGTATTATCTCCAGCAGACAAGACAAATGTAAAAGCAGCATGGGGTAAAGTTGGTGCCCCA
CGCCGGTGAATATGGTGCAGAAAGCTTTGGAAAGAATGTTCTTGTTCATTCCAACTACAAAGACATACTTCCCTCATTTGATT
TGTCACCGGTAGTGCTCAAGTCAAGGGTCATGGTAAAAAGGTAGCTGACGCATTAACCAATGCCGTTGCTCACGTCGATGA
CATGCCAAACGCTTTGTCTGCATTATCAGATTTGCATGCACACAAATTGAGAGTAGACCCTGTAACTTCAAGTTGTTGTCTCA
TTGTTTGTAGTCACTTTGGCTGCACACTTACCAGCCGAATTAAGTCCAGCTGATAAACTAACGTTAAGGCCGCTTGGGGTAA
AGTCGGTGCACATGCCGGTGAATATGGTGTGAAGCATTGGAAAGAATGTTTTATCCTTCCCAACCACTAAACCTTACTTTC
CACACTTTGATTTGTCTCACGGTAGTGCTCAAGTAAAAAGGTCACGGTAAAAAGGTTGCCGATGCTTTAACAATGCAGTTGCC
CATGTCGATGATATGCCTAACGCATTATCTGCCTTGTGAGATTGTCATGCTCACAAGTTAAGAGTAGACCCAGTCAATTTAA
ATTGTTGTCTATTGCTTGTAGTTACCTTGGCAGCCCACTTACCAGCTGAATTAAGTCCCTGCAGTTTCATGCATCATTGGACAA
GTTTTTGGCTTCAGTCTCTACCGTTTTGACCTCTAAATACAGATAAGAGCTCTTAATTAAGTCGACCTCGAG
