



Inactivated virus vaccines from chemistry to prophylaxis: merits, risks and challenges

Iris Delrue, Dieter Verzele, Annemieke Madder & Hans J Nauwynck

To cite this article: Iris Delrue, Dieter Verzele, Annemieke Madder & Hans J Nauwynck (2012) Inactivated virus vaccines from chemistry to prophylaxis: merits, risks and challenges, Expert Review of Vaccines, 11:6, 695-719, DOI: [10.1586/erv.12.38](https://doi.org/10.1586/erv.12.38)

To link to this article: <https://doi.org/10.1586/erv.12.38>



Published online: 09 Jan 2014.



Submit your article to this journal [↗](#)



Article views: 13010



View related articles [↗](#)



Citing articles: 71 View citing articles [↗](#)

Inactivated virus vaccines from chemistry to prophylaxis: merits, risks and challenges

Expert Rev. Vaccines 11(6), 695–719 (2012)

Iris Delrue^{†1},
Dieter Verzele^{†2},
Annemieke Madder²
and Hans J Nauwynck^{*1}

¹Laboratory of Virology, Department of Virology, Parasitology and Immunology, Ghent University, Merelbeke, Belgium

²Laboratory for Organic and Biomimetic Chemistry, Department of Organic Chemistry, Ghent University, Ghent, Belgium

*Author for correspondence:

Tel.: +32 9 264 73 66

Fax: +32 9 264 74 95

hans.nauwynck@UGent.be

[†]These authors contributed equally

The aim of this review is to make researchers aware of the benefits of an efficient quality control system for prediction of a developed vaccine's efficacy. Two major goals should be addressed when inactivating a virus for vaccine purposes: first, the infectious virus should be inactivated completely in order to be safe, and second, the viral epitopes important for the induction of protective immunity should be conserved after inactivation in order to have an antigen of high quality. Therefore, some problems associated with the virus inactivation process, such as virus aggregate formation, protein crosslinking, protein denaturation and degradation should be addressed before testing an inactivated vaccine *in vivo*.

KEYWORDS: 2,2'-dithiodipyridine • β -propiolactone • binary ethylene imine • formaldehyde • gamma irradiation • glutaraldehyde • pH • temperature • UV • vaccine • virus inactivation

Classically, two main types of vaccines are used to protect man and animals clinically and virologically on challenge: modified live virus (attenuated virus) vaccines and killed virus (KV; inactivated virus) vaccines. KV vaccines are often preferred because of safety reasons, but some problems may be encountered using them. First, virus inactivation may turn out to be incomplete with outbreaks after vaccination as a consequence [1–3]. Second, the viral-neutralizing epitopes may be destroyed during inactivation, resulting in a poor neutralizing antibody response and a poor protection on challenge [4–6]. Therefore, efficient quality control of the inactivated antigen is necessary to evaluate virus inactivation and the effect of the inactivation procedure on neutralizing epitopes.

In the following paragraphs, different and most commonly used inactivation procedures for viruses will be discussed and their usefulness for the development of a KV vaccine will be evaluated. An overview of the discussed inactivation methods and the relevant chemical conditions involved is given in TABLE 1.

Cross-linkers

Formaldehyde




KV vaccine development based on inactivation with formaldehyde has been investigated for many viruses. A few of them are

discussed below and presented in TABLE 2. Some formaldehyde-inactivated vaccines work well and protect against disease on challenge. This accounts for inactivated Ross River virus (RRV) [7], West Nile virus [8], dengue-2 virus [9], equine herpes virus type 1 [10], papilloma virus [11], Hantaan virus [12], influenza virus [13], hepatitis A virus [14], tick-borne encephalitis virus [15,16] and Japanese encephalitis virus [17]. However, with some vaccines inactivated in this way, problems have been encountered. In order to gain insight into the reasons for failure, a more detailed understanding of the chemistry is needed.

Formaldehyde (FIGURE 1A), also known as formalin when diluted in water, has an electron-deficient central carbon atom and is therefore electrophilic, as illustrated through one of the resonance forms. As a consequence, a nucleophile, such as a nonprotonated amino group, can attack the central carbonyl carbon. This chemical reaction is called a nucleophilic addition.

Formaldehyde has an effect on both genome and proteins. First of all, it monohydroxymethylates adenine [18]: the nonprotonated exocyclic amine on adenine (N6) acts as a nucleophile and reacts with the carbocation of formaldehyde [19]. Because of this nucleophilic addition, adenine is monohydroxymethylated, which blocks genome reading [20]. This

Table 1. Overview of inactivation methods for the development of killed virus vaccines.

Method		Type	Mechanism	Ref.
Formaldehyde		Alkylating agent	Monohydroxymethylation of adenine	[18]
		Crosslinker	Crosslinking of RNA to capsid proteins, causing a block of genome reading Crosslinking of proteins by formation of inter- and intra-molecular methylene bridges between hydroxymethylated amines	[25–27,107] [20]
Glutaraldehyde		Crosslinker	Crosslinking of proteins by the same mechanism as formaldehyde (described above)	[47]
2,2'-dithiodipyridine		Crosslinker	Crosslinking of proteins by oxidation of S-H groups causing formation of S-S bridges, which results in a covalent modification and functional inactivation of S-H-containing internal viral proteins	[54]
β-propiolactone		Alkylating agent Crosslinker	Alkylation of RNA and DNA as a consensus mechanism Crosslinking of proteins	[58] [59]
Binary ethylene imine		Alkylating agent	Alkylation of RNA and DNA at low concentrations. Most likely genome reading is blocked by alkylation of guanine or adenine by binary ethylene imine Alkylation of proteins (nucleocapsid) at high concentrations	[72,73] [72]
pH		Denaturation agent RNA degradation	Denaturation of viral functionally active proteins The close proximity of the hydroxyl group to the phosphor center of each internucleotide linkage facilitates transesterification under strongly acidic or strongly basic conditions, with a breakage of the phosphodiester bond as a consequence	[80] [81]
Temperature		Denaturation agent RNA degradation	A high temperature denatures viral functionally active proteins Virus inactivation at 'low' temperature (below 41°C) is considered to be caused by degradation of the nucleic acid	[82,83] [84–86]
Gamma irradiation		Radiation	Viruses are inactivated primarily by direct damage, via disruption of the genome Formation of free radicals that damage proteins	[95] [95]
Ultraviolet light		Radiation	Induction of dimer formation between adjacent uracils in RNA. Dimer formation leads to pressure and breakage of the sugar backbone causing a block of genome reading Works more slowly, ultraviolet light also causes structural modifications of the capsid proteins resulting in the formation of large and small photoproducts	[89,90] [90,94]

monohydroxymethylation of adenine (FIGURE 1A), also known as alkylation, can occur on either DNA or RNA [21]. Monohydroxymethyladenine (NH–CH₂OH) is stable for days at room temperature and is more likely to exist than the Schiff base (N=CH₂) [22]. However, if adenine and formaldehyde are stored for days at room temperature and at a pH of 4.5 then methylene bis-adenine can be formed [19].

Formaldehyde can also react with nonprotonated amino groups of the N-terminal amino acid residue and the amino acids featuring N-containing functionality in their side chain, such as lysine, arginine, glutamine, tryptophan and histidine, or a sulfhydryl group in their side chain, such as cysteine. More specifically, as exemplary illustrated for the lysine side chain (FIGURE 1B),

a nonprotonated amine acts as a nucleophile and reacts with the carbocation of formaldehyde. As a result, the amino group is monohydroxymethylated, as described above for adenine. The loss of a water molecule from the unstable hemiaminal results in Schiff base formation [23], also called an imine intermediate. The resulting Schiff base or imine intermediate can crosslink with arginine and tyrosine and to a lesser extent with glutamine, asparagine, tryptophan and histidine residues by a nucleophilic addition reaction [23]. In this way, inter- and intra-molecular methylene bridges can be formed (FIGURE 1B) [24]. Because of these bridges, proteins become inter- and intra-molecularly crosslinked.

In addition, these reactions can also cause crosslinking between genome and proteins through combining the above described

Table 2. Effect of formaldehyde on different viruses: inactivation, viral protein modification, induction of virus-specific and virus-neutralization antibodies and protection on challenge (reduction of viremia and fewer clinical signs).

Study (year)	Genome	Envelope	Capsid	Virus family	Example	Inactivation (concentration)	Viral protein modification	Induction		Protection on challenge		Ref.
								VS Ab	VN Ab	Reduced viremia	Fewer clinical signs	
Pollock et al. (1982)	DNA (ss)	No	I	Parvo Circo	CPV No	Yes	Yes	NT	NT	NT	NT	[43]
Skinner et al. (2000)	DNA (ds)	Yes	I	Hepadna	No							
				Herpes	EHV-1	Yes (0.04%)	NT	Yes	Yes	NT	Yes	[10]
				Asfor	No							
				Pox	No							
Bell et al. (1994)	RNA (segmented, ss, -)	No	I	Adeno	No							
				Papova	Papilloma	Yes	NT	NT	NT	NT	Yes	[11]
				Arena	No							
Choi et al. (2003)	RNA (nonsegmented, ss, -)	Yes	H	Bunya	Hantaan	Yes (0.0185%)	NT	Yes	Yes	NT	[12]	
Orthomyxo				Influenza	Yes	NT	Yes	NT	Yes	Yes	[13]	
Openshaw et al. (2001)	RNA (nonsegmented, ss, -)	Yes	H	Filo	No							
				Paramyxo	RSV	Yes	NT	Yes	Poor	NT	No	[34]
				Rhabdo	No							
Kistner et al. (2007)	RNA (nonsegmented, ss, +)	Yes	I	Borna	No							
Toga				RRV	Yes (0.1%)	No	Yes	Yes	Yes	NT	[7]	
Brown et al. (1993)	RNA (nonsegmented, ss, +)	Yes			VEEV	Incomplete	NT	NT	NT	NT	No	[28]
Jahrling et al. (1984)					Yes	NT	Yes	Yes	Yes	No	[6]	
Delrue et al. (2009)					PRRSV	Yes (1 × 10 ⁻⁶ %)	Yes	NT	NT	NT	NT	[41]
Ishizaka et al. (1999)					SIV	Yes	NT	Yes	No	No	NT	[35]
Rossio et al. (1998)					HIV	Yes (1:80)	Yes	NT	NT	NT	NT	[42]

+: Positive stranded; -: Negative stranded; C: Complex; CPV: Canine parvovirus; DEN: Dengue virus; ds: Double stranded; EHV: Equine herpes virus; FMDV: Foot-and-mouth disease virus; H: Helical; HAV: Hepatitis A virus; I: Icosahedral; JEV: Japanese encephalitis virus; NT: Not tested; PRRSV: Porcine reproductive and respiratory virus; PV: Poliovirus; RRV: Ross river virus; RSV: Respiratory syncytial virus; SARS-CoV: Severe acute respiratory syndrome coronavirus; ss: Single stranded; TBEV: Tick-borne encephalitis virus; VEEV: Venezuelan equine encephalitis virus; VN Ab: Virus neutralizing antibodies; VS Ab: Virus-specific antibodies; WNV: West Nile virus.

Table 2. Effect of formaldehyde on different viruses: inactivation, viral protein modification, induction of virus-specific and virus-neutralization antibodies and protection on challenge (reduction of viremia and fewer clinical signs) (cont.).

Study (year)	Genome	Envelope	Capsid	Virus family	Example	Inactivation (concentration)	Viral protein modification	Induction		Protection on challenge		Ref.
								VS Ab	VN Ab	Reduced viremia	Fewer clinical signs	
Niedrig <i>et al.</i> (1993)						Yes	NT	Yes	Yes	Yes	No	[5]
Putnak <i>et al.</i> (1996)				Flavi	DEN-2	Yes (0.05%)	NT	Yes	Yes	Yes	Yes	[9]
Appaiahgari <i>et al.</i> (2004)					JEV	Yes (1:40)	NT	Yes	Yes	NT	Yes	[17]
Samina <i>et al.</i> (2005)					WNV	Yes (0.1%)	NT	NT	NT	Yes	Yes	[8]
Goncharova <i>et al.</i> (2006)					TBEV	Yes	NT	Yes	Yes	NT	Yes	[15]
Qu <i>et al.</i> (2005)			H	Corona	SARS-CoV	Yes (1:2000)	NT	NT	Yes	NT	NT	[32]
Zhang <i>et al.</i> (2004)						Yes (0.4%)	NT	Yes	Yes	NT	NT	[33]
Darnell <i>et al.</i> (2004)						Incomplete	NT	NT	NT	NT	NT	[30]
Patil <i>et al.</i> (2002)		No	I	Calici	No							
King <i>et al.</i> (1981)				Picornia	FMDV	Yes (0.06%)	Yes	Poor	NT	NT	NT	[3]
Tano <i>et al.</i> (2007)					PV	Incomplete	NT	NT	NT	NT	No	[1]
Furesz <i>et al.</i> (1995)					HAV	Yes (0.025%)	Yes	NT	NT	NT	NT	[39]
						Yes	NT	Yes	Yes	NT	Yes	[14]
				Astro	No							
	RNA (segmented, ds)	No	I	Reo	No							
				Birna	No							

+: Positive stranded; -: Negative stranded; C: Complex; CPV: Canine parvovirus; DEN: Dengue virus; ds: Double stranded; EHV: Equine herpes virus; FMDV: Foot-and-mouth disease virus; H: Helical; HAV: Hepatitis A virus; I: Icosahedral; JEV: Japanese encephalitis virus; NT: Not tested; PRRSV: Porcine reproductive and respiratory virus; PV: Poliovirus; RRV: Ross river virus; RSV: Respiratory syncytial virus; SARS-CoV: Severe acute respiratory syndrome coronavirus; ss: Single stranded; TBEV: Tick-borne encephalitis virus; VEEV: Venezuelan equine encephalitis virus; VN Ab: Virus neutralizing antibodies; VS Ab: Virus-specific antibodies; WNV: West Nile virus.

functionalities [25,26], which prevents the transcriptional machinery from reaching the genome [27].

One of the problems observed with formaldehyde is that some vaccines based on formaldehyde inactivation can contain incompletely inactivated virus particles, which can cause outbreaks of virus infections on vaccination [28]. This is reported for several viruses such as foot-and-mouth disease virus (FMDV) [1] and Venezuelan equine encephalitis virus (VEEV) [28]. Molecular analysis proved clearly that outbreaks of FMDV in western Europe in the 1980s and VEEV in central America in 1970s are a consequence of incompletely inactivated virus vaccines [28]. This can be due to crosslinking of nucleocapsid proteins with RNA. As a consequence, RNA cannot be degraded, and some virus particles escape from inactivation with formaldehyde [29]. It is also possible that the safety testing was not sufficient and that a higher concentration of formaldehyde is necessary to inactivate these viruses properly. Darnell *et al.* examined formaldehyde inactivation of the severe acute respiratory syndrome coronavirus (SARS-CoV) [30]. They showed that formaldehyde can inactivate the virus in a temperature-dependent manner. The virus could not be inactivated at a low temperature of 4°C even after 3 days. However, when using a higher temperature of 25 or 37°C, formaldehyde could inactivate most of the virus after 1 day [30]. Although not investigated in this particular study, in addition to temperature, the pH during inactivation can play a major role, as is further illustrated in this review (see 'pH' section). When SARS-CoV is treated with 0.05% formaldehyde for 48 h, it could be inactivated properly [31]. Vaccines based on formaldehyde-inactivated SARS-CoV are shown to induce virus-specific and virus-neutralizing antibodies [32,33]; however, some virus still remained infectious on day 3 of inactivation [30].

Some KV vaccines based on formaldehyde inactivation do not induce a virus-neutralizing antibody response and do not protect on challenge, for example respiratory syncytial virus (RSV) [34], inactivated HIV [5], SIV [35] and VEEV vaccines [6]. A poor immune response on vaccination can even lead to atypical or enhanced disease after infection of vaccinated recipients. This is most likely due to the poor induction of neutralizing antibody responses together with the formation of immune complexes between virus and antibodies, and further facilitated uptake and infection of these complexes by monocytic cells, termed antibody-dependent enhanced infection [36]. This is observed with inactivated measles [37] and RSV [34,38] vaccines.




Figure 1. Reaction mechanism of formaldehyde with either DNA/RNA or amino acids. Reaction with (A) DNA or RNA (adenine) and, in a similar fashion, (B) amino acids (e.g., lysine) of proteins with a primary amine group: monohydroxy methylation and methylene bridge formation. A: Adenine; Lys: Lysine.

Formaldehyde can destroy the viral structure, which can result in a poor immune response. When poliovirus (PV) is treated with formaldehyde, the extent of the modification of the antigenic sites is dependent on the strain used and correlates with the immune response that can be induced [39]. Treatment of porcine reproductive and respiratory syndrome virus (PRRSV) with formaldehyde inactivates the virus but does not preserve the viral entry-associated domains. These domains may be important for the induction of protective immunity, as neutralizing antibodies are directed to the viral epitopes that are involved in the attachment to the PRRSV receptors and internalization into the host cell [40]. Since crosslinking of viral entry-associated domains might interfere with the subsequent presentation of viral-neutralizing epitopes to cells of the adaptive immune system, this procedure is not useful for inactivated PRRSV vaccine development [41]. Modification of the virion structure on inactivation with formaldehyde is also reported for FMDV [3], HIV [42] and canine parvovirus [43].

In conclusion, although formaldehyde has been shown to correctly inactivate some viruses, given the problems discussed

Table 3. Effect of glutaraldehyde on different viruses: inactivation, viral protein modification, induction of virus-specific and virus-neutralization antibodies and protection on challenge (reduction of viremia and fewer clinical signs).

Study (year)	Genome	Envelope	Capsid	Virus family	Example	Inactivation (concentration)	Viral protein modification	Induction		Protection on challenge		Ref.
								VS Ab	VN Ab	Reduced viremia	Fewer clinical signs	
Cham <i>et al.</i> (2006)	DNA (ss)	No	I	Parvo	No							[4]
				Circo	No							
	DNA (ds)	Yes	I	Hepadna	DHBV	Yes	NT	Yes	Yes	No	NT	
				Herpes	No							
				Asfor	No							
				Pox	No							
				Adeno	No							
				Papova	No							
				Arena	No							
Keles <i>et al.</i> (1998)	RNA (segmented, ss, -)	Yes	H	Bunya	No							[51]
				Orthomyxo	No							
	RNA (nonsegmented, ss, -)	Yes	H	Filo	No							
				Paramyxo	BRSV	Yes (0.075%)	NT	Yes	Yes	Yes	NT	
				Rhabdo	No							
				Borna	No							
				Toga	No							
		Yes	I									
Delrue <i>et al.</i> (2009)				Arteri	PRRSV	Yes (1 × 10 ⁻⁶ %)	Yes	NT	NT	NT	NT	[41]
Darnell <i>et al.</i> (2004)		Yes	H	Corona	SARS-CoV	Yes (1:4000)	NT	NT	NT	NT	NT	[30]
				Retro	No							
				Flavi	No							

+: Positive stranded; -: Negative stranded; BRSV: Bovine respiratory syncytial virus; C: Complex; DHBV: Duck HBV; ds: Double stranded; H: Helical; I: Icosahedral; IBDV: Infectious bursal disease virus; NT: Not tested; PRRSV: Porcine reproductive and respiratory syndrome virus; SARS-CoV: Severe acute respiratory syndrome coronavirus; ss: Single stranded; VN Ab: Virus-neutralizing antibodies; VS Ab: Virus-specific antibodies.

Table 3. Effect of glutaraldehyde on different viruses: inactivation, viral protein modification, induction of virus-specific and virus-neutralization antibodies and protection on challenge (reduction of viremia and fewer clinical signs) (cont.).

Study (year)	Genome	Envelope	Capsid	Virus family	Example	Inactivation (concentration)	Viral protein modification	Induction		Protection on challenge		Ref.
								VS Ab	VN Ab	Reduced viremia	Fewer clinical signs	
Cepica <i>et al.</i> (1998)	RNA (segmented, ds)	No	I	Calici	No	Yes (0.01%)	Yes	NT	NT	NT	NT	[52]
				Picornia	No							
				Astro	No							
				Reo	No							
		No	I	Birna	IBDV							

+: Positive stranded; -: Negative stranded; BRSV: Bovine respiratory syncytial virus; C: Complex; DHBV: Duck HBV; ds: Double stranded; H: Helical; I: Icosahedral; IBDV: Infectious bursal disease virus; NT: Not tested; PRRSV: Porcine reproductive and respiratory syndrome virus; SARS-CoV: Severe acute respiratory syndrome coronavirus; ss: Single stranded; VN Ab: Virus-neutralizing antibodies; VS Ab: Virus-specific antibodies.

above, care must be taken regarding the exact protocol used. In part, some failures could be due to an insufficient inactivation protocol. Problems, such as incomplete virus inactivation, that lead to outbreaks have to be kept in mind when using this inactivation method for KV vaccine development. Formaldehyde can be used for KV development for some viruses, but several KV vaccines based on formaldehyde inactivation do not protect the host on challenge with virulent virus. This could be due to RNA–protein crosslinking [26,27] or modification of viral proteins by formaldehyde, as reported for PV, FMDV and HIV [3,39,42].

Glutaraldehyde

Examples of virus inactivation with glutaraldehyde are summarized in TABLE 3. Although there are few examples, complete virus inactivation with glutaraldehyde seems to be less problematic than with formaldehyde.

Glutaraldehyde (FIGURE 2) is a saturated, five-carbon dialdehyde. The carbonyl carbons are electrophilic and therefore a nucleophile, such as a nonprotonated amino group, might attack these carbons in a nucleophilic addition reaction, just like formaldehyde.

Genomic DNA or RNA is the target of glutaraldehyde. Although the exact mechanism of action has not clearly been described, it is likely that glutaraldehyde acts in the same way as formaldehyde, since similar chemical groups are present on formaldehyde and glutaraldehyde [44]. The exocyclic amino group on adenine (N6) is the reactive group, which attacks glutaraldehyde [45]. It is certain that glutaraldehyde can have an effect on the genome, since RNA and protein synthesis can be blocked by this aldehyde [46].

Glutaraldehyde can have an effect on proteins as well, because nonprotonated amines of amino acids, such as lysine, can be cross-linked with each other [47]. However, the reaction mechanism of glutaraldehyde with proteins is different from that of formaldehyde (FIGURE 2). First, glutaraldehyde forms unsaturated aldehydes by aldol condensation and elimination of water [48]. Then, several amino acids can be linked to glutaraldehyde via a Michael-type addition, the nonprotonated amines acting as nucleophiles and reacting with the double bond of the glutaraldehyde oligomer [49]. In this way, the amino acids are coupled by formation of a bridge [50]. Unlike the reaction with formaldehyde, intermediate Schiff bases or imines are not formed [44,50], but both pathways result in covalently linked amino acid side chains.

In addition, the reactions described above can also cause crosslinking of the genome and proteins [25], which blocks genome reading [27].

To investigate virus inactivation with glutaraldehyde, SARS-CoV was incubated with glutaraldehyde at two different dilutions and at different temperatures. The glutaraldehyde exhibited a temperature dependence in its ability to inactivate SARS-CoV. The virus could not be inactivated by glutaraldehyde at 4°C even after 3 days. However, when a higher temperature of 25 or 37°C was used, the virus could be inactivated in 2 and 1 days, respectively [30].

As glutaraldehyde is a bi-functional or, after aldol condensation, multifunctional crosslinking agent, virus damage might be more




Figure 2. Reaction mechanism of glutaraldehyde with amino acids of proteins (e.g., with the primary amine group of lysine).

Lys: Lysine.

extensive. Therefore, the induction of a protective immune response can be more of a problem with glutaraldehyde-inactivated virus vaccines. Duck HBV (DHBV) and bovine RSV (BRVS) are viruses for which the use of glutaraldehyde has been tested for the production of a KV vaccine [451]. Glutaraldehyde-inactivated DHBV and BRVS developed virus-specific antibodies, but ducks vaccinated with glutaraldehyde-inactivated DHBV and lambs vaccinated with glutaraldehyde-inactivated BRVS were not fully protected against viremia following challenge. This indicates that there was no induction of protective immunity in these animals [451].

The destruction of the viral structure by glutaraldehyde may be a problem. Similar to formaldehyde, glutaraldehyde could inactivate PRRSV but did not preserve the viral entry-associated domains and is therefore not useful for inactivated PRRSV vaccine development [41]. Glutaraldehyde was also shown to change the conformational structure of infectious bursal disease virus [52].

In conclusion, glutaraldehyde can inactivate viruses properly, but there is not always protection against infection on challenge for the KV vaccines discussed here. This could be due to destruction of viral proteins by glutaraldehyde.

Aldrithiol or 2,2'-dithiodipyridine

Examples of virus inactivation with aldrithiol, dithiodipyridine (AT-2), are summarized in TABLE 4. This procedure for virus inactivation for KV vaccines has been investigated for HIV [42], PRRSV [41] and RSV [53].

AT-2 oxidizes S-H groups, which results in formation of S-S bridges that crosslink proteins (FIGURE 3). As the aromatic thiolate is the better leaving group, the reaction equilibrium is shifted toward the aliphatic disulfides. The internal viral proteins are subjected to the intracellular (reducing) environment, and as a consequence the cysteine residues are present in thiol form (S-H). By contrast, the surface proteins of viruses are subjected to the extracellular (oxidizing) environment, and as a consequence, the cysteines are present as disulfides (S-S). Treatment with AT-2 results in covalent modification and functional inactivation of S-H-containing internal viral proteins, such as the nucleocapsid protein, which is required for HIV infectivity, whereas the envelope glycoproteins with disulfide bonded cysteines remain unaffected [54].

HIV preparations inactivated with AT-2 did not retain detectable infectivity, but showed maintenance of conformational and functional integrity. This suggests that such virions may be useful as vaccine antigens [42]. An RSV vaccine based on AT-2 inactivation was found to be effective in the protection of cotton rats against virus replication in the lungs. This suggests that compounds that inactivate retroviruses by targeting the zinc finger motif in their nucleocapsid proteins are also effective against RSV [53].

However, AT-2 cannot inactivate all viruses. PRRSV is not sensitive for AT-2 and could not be inactivated by it [41]. For PRRSV, disulfide bridges between nucleocapsid proteins are important for virus infection, but as stated above, these disulfide bridges remain unaffected by AT-2 [55]. It seems there are no free thiol groups in

Table 4. Effect of 2,2'-dithiodipyridine on different viruses: inactivation, viral protein modification, induction of virus-specific and virus-neutralization antibodies and protection on challenge (reduction of viremia and fewer clinical signs).

Study (year)	Genome	Envelope	Capsid	Virus family	Example	Inactivation (concentration)	Viral protein modification	Induction		Protection on challenge		Ref.
								VS Ab	VN Ab	Reduced viremia	Fewer clinical signs	
Boukhvalova et al. (2009)	DNA (ss)	No	I	Parvo	No							
				Circo	No							
	DNA (ds)	Yes	I	Hepadna	No							
				Herpes	No							
				Asfor	No							
			C	Pox	No							
		No	I	Adeno	No							
				Papova	No							
	RNA (segmented, ss, -)	Yes	H	Arena	No							
				Bunya	No							
Delrue et al. (2009)				Orthomyxo	No							
				Filo	No							
				Paramyxo	RSV	Yes (10 mM)	No	NT	NT	Yes	NT	[53]
				Rhabdo	No							
				Borna	No							
Rossio et al. (1998)				Toga	No							
	RNA (nonsegmented, ss, +)	Yes	I									
				Arteri	PRRSV	No (2 mM)	No	NT	NT	NT	NT	[41]
Lifson et al. (2004)				Retro	HIV	Yes (300 µM)	No	NT	NT	NT	NT	[42]
				SIV	SIV	Yes (1 mM)	NT	Yes	Yes	Yes	NT	[56]

+: Positive stranded; -: Negative stranded; C: Complex; ds: Double stranded; H: Helical; I: Icosahedral; NT: Not tested; PRRSV: Porcine reproductive and respiratory virus; RSV: Respiratory syncytial virus; SIV: Simian immunodeficiency virus; ss: Single stranded; VN Ab: Virus-neutralizing antibodies; VS Ab: Virus-specific antibodies.

Table 4. Effect of 2,2'-dithiodipyridine on different viruses: inactivation, viral protein modification, induction of virus-specific and virus-neutralization antibodies and protection on challenge (reduction of viremia and fewer clinical signs) (cont.).

Study (year)	Genome	Envelope	Capsid	Virus family	Example	Inactivation (concentration)	Viral protein modification	Induction		Protection on challenge		Ref.
								VS Ab	VN Ab	Reduced viremia	Fewer clinical signs	
Vanhee et al. (2009)						Yes (300 µM)	NT	No	No	Yes	NT	[57]
				Flavi	No							
			H	Corona	No							
			I	Calici	No							
		No		Picornia	No							
					No							
				Astro	No							
				Reo	No							
	RNA (segmented, ds)	No	I									
				Birna	No							

+: Positive stranded; -: Negative stranded; C: Complex; ds: Double stranded; H: Helical; I: Icosahedral; NT: Not tested; PRRSV: Porcine reproductive and respiratory virus; RSV: Respiratory syncytial virus; SIV: Simian immunodeficiency virus; ss: Single stranded; VN Ab: Virus-neutralizing antibodies; VS Ab: Virus-specific antibodies.

PRRSV that are important for infection. Probably, only viruses where free thiol groups are important for virus infection can be inactivated by AT-2.

There are also some vaccination studies where no full protection after vaccination can be assured. A vaccination study with AT-2 inactivated SIV showed that an immune response could be induced, and there was virological protection on challenge with homologous, but not heterologous, virus [56]. Another study could not detect an immune response before challenge. There was an antibody response on challenge, but vaccination only partly protected against viremia and infection on challenge [57].

In conclusion, AT-2 seems to inactivate HIV and SIV properly with preservation of the virion structure, but vaccination studies cannot prove that AT-2 inactivated virus can protect the host on challenge. AT-2 was not effective with other viruses.

Alkylating agents

β-propiolactone

β-propiolactone (BPL) is widely used to inactivate viruses for vaccine purposes (TABLE 5), mainly acting as an alkylating agent on guanine of viral DNA or RNA [58,59]. As illustrated in FIGURE 4, the nucleophilic guanine of nucleic acids reacts with the electrophilic BPL through a nucleophilic substitution mechanism causing the ring opening of the strained BPL and the *N*-alkylation of guanine. As BPL is much more toxic as a carcinogen [60] and displays the same mode of action as binary ethylene imine (BEI) discussed in the section 'Aminoethyl ethylene imine' the interested reader is referred to the more specialized literature. Interestingly, the mechanistic details and the chemical modifications of BPL treatment have recently been elucidated through systematic studies on nucleobase, nucleoside and peptide model systems, revising the above G-alkylation consensus pathway and providing a more nuanced view on the chemistry involved [59]. In fact, acylation and cross-linking reactions also occur, to a significant extent, with proteins (cysteine, methionine and histidine as most reactive residues) being modified more extensively than originally agreed from earlier literature, which can have serious implications for vaccine production. Indeed, modifications on amino acids by BPL treatment can, for example, lead to the disability of influenza viruses to enter cells by virus-membrane fusion and modification of hemagglutinin [61,62].

BPL-inactivated virus vaccines against yellow fever [63], SARS-CoV [64], influenza [65] and rabies [66] were able to induce the production of neutralizing antibodies. However, BPL-inactivated influenza showed reduced hemagglutinin and neuraminidase activity [62]. Vaccines based on BPL-inactivated virus against SARS-CoV [64], hematopoietic necrosis virus (HNV) [67] and influenza [65], were able to protect on challenge.

In conclusion, BPL seems to inactivate viruses properly and is able to induce a protective immune response. This can be expected, as BPL can alkylate the viral genome. However, protein modification should be kept in mind when using BPL.

Aminoethyl ethylene imine

Virus inactivation by ethylene imines was first introduced over 30 years ago [68,69]. Originally, two chemically similar compounds,

ethylene imine (aziridine) and acetylene imine, were proposed for use in the development of KV vaccines [70,71]. However, the selectivity of aziridine toward nucleic acids is not very high. To improve this selectivity, a product with more protonizable amino groups was necessary, and aminoethyl ethylene imine, also called BEI, which has two protonizable amino groups while aziridine only has one, was selected.

If a low concentration of BEI is used, the virus capsid is not alkylated, but BEI passes through the capsid and alkylates the genome [72]. N7-guanine of the genome acts as a nucleophile and reacts with the electrophile BEI. The nucleophilic substitution reaction performed by N7-guanine causes an opening of the BEI ring and guanine becomes alkylated (FIGURE 5) [73]. In addition, adenine is alkylated via a ring-opening reaction with BEI [72]. This ring-opening reaction in RNA nucleosides is approximately two- to three-times faster than that in DNA nucleosides [74]. So far, there are no reports on any interaction of BEI with proteins, at least if low concentrations are used, which suggests that the neutralizing viral epitopes are preserved after treatment of a virus with BEI.

BEI is used for the inactivation of many viruses, such as FMDV [70], RRV [75], sheep pox [76], HIV [77], Nipah virus [78], HNV [67] and PRRSV (TABLE 6) [79].

As mentioned above, the preservation of the viral structure should not be a problem if a low concentration of BEI is used. This has also been shown for PRRSV, where BEI could inactivate PRRSV with the preservation of the viral entry-associated domains [41]. Furthermore, the inactivation of FMDV with BEI did not affect the antigenicity of FMDV [70].

Most vaccines based on BEI-inactivated virus are able to induce a protective immune response. Mice vaccinated with a purified BEI-inactivated RRV vaccine were protected against challenge with live virus. The vaccine induced significant levels of neutralizing antibody in all strains of mice tested [75]. A local Egyptian sheep pox virus strain inactivated with BEI could induce virus-specific antibodies starting from 1-week post-vaccination until 4-weeks post-challenge. Vaccinated lambs were protected on challenge up to 6-months post-vaccination [76]. Race *et al.* have shown that the experimental BEI-inactivated HIV vaccine induces virus-neutralizing antibodies against both the homologous vaccine strain and a heterologous virus strain [77]. Also, virus-specific antibodies could be induced against BEI-inactivated Nipah virus; however, these did not neutralize the virus *in vitro* [78]. After vaccination of pigs with BEI-inactivated PRRSV, they induced a virus-specific and virus-neutralize antibody




Figure 3. Reaction mechanism of 2,2'-dithiodipyridine with cysteine.
Cys: Cysteine.

response, which was able to reduce viremia after challenge [79]. However, rainbow trout vaccinated with BEI-inactivated HNV and challenged with live HNV showed no statistically significant differences between the cumulative percentage mortality in the group of fish immunized with the BEI vaccines and the mortality of fish in the mock-vaccinated control groups when tested 28 or 56 days after vaccination [67].

In conclusion, BEI seems to inactivate viruses properly and most KV vaccines based on BEI inactivation protect the host after challenge with virulent virus, which is expected since BEI does not interfere with the virus structure if a low concentration is used. Only one KV vaccine did not protect the host on challenge, but the possible reason for this remains to be elucidated.

Denaturing procedures

pH

Increasing or decreasing the pH has an effect on proteins by protein denaturation. This means that the proteins adopt a different 3D structure. A low (acidic) or a high (alkaline) pH can inactivate viruses via denaturation of the secondary structures of proteins, thereby altering the conformation of viral proteins that are involved in attachment to and replication in a host cell. The conformation




Figure 4. Reaction mechanism of β -propiolactone with DNA or RNA (guanine).
G: Guanine.

Table 5. Effect of β -propiolactone on different viruses: inactivation, viral protein modification, induction of virus-specific and virus-neutralization antibodies and protection on challenge (reduction of viremia and fewer clinical signs).

Study (year)	Genome	Envelope	Capsid	Virus family	Example	Inactivation (concentration)	Viral protein modification	Induction		Protection on challenge		Ref.
								VS Ab	VN Ab	Reduced viremia	Fewer clinical signs	
Bodewes <i>et al.</i> (2010)	DNA (ss)	No	I	Parvo	No							
				Circo	No							
	DNA (ds)	Yes	I	Hepadna	No							
				Herpes	No							
				Asfor	No							
Jonges <i>et al.</i> (2010)			C	Pox	No							
		No	I	Adeno	No							
				Papova	No							
	RNA (segmented, ss, -)	Yes	H	Arena	No							
				Bunya	No							
Anderson <i>et al.</i> (2008)				Orthomyxo	Influenza	Yes	NT	Yes	Yes	Yes	Yes	[65]
	RNA (nonsegmented, ss, -)	Yes	H	Filo	No							
				Paramyxo	No							
				Rhabdo	HNV	Yes (2.7 mM)	NT	NT	NT	NT	Yes	[67]
				Rabies	Rabies	Yes	NT	Yes	Yes	NT	NT	[66]
Sabchareon <i>et al.</i> (1999)				Borna	No							
	RNA (nonsegmented, ss, +)	Yes	I	Toga	No							
				Arteri	No							
				Retro	No							

+: Positive stranded; -: Negative stranded; C: Complex; SARS-CoV: Severe acute respiratory syndrome coronavirus; ds: Double stranded; H: Helical; HNV: Hematopoietic necrosis virus; I: Icosahedral; NT: Not tested; RRV: Ross river virus; ss: Single stranded; VN Ab: Virus-neutralizing antibodies; VS Ab: Virus-specific antibodies.

Table 5. Effect of β -propiolactone on different viruses: inactivation, viral protein modification, induction of virus-specific and virus-neutralization antibodies and protection on challenge (reduction of viremia and fewer clinical signs) (cont.).

Study (year)	Genome	Envelope	Capsid	Virus family	Example	Inactivation (concentration)	Viral protein modification	Induction		Protection on challenge		Ref.
								VS Ab	VN Ab	Reduced viremia	Fewer clinical signs	
Monath <i>et al.</i> (2011)				Flavi	Yellow fever	Yes	NT	Yes	Yes	NT	NT	[63]
Roberts <i>et al.</i> (2010)			H	Corona	SARS-CoV	Yes	NT	Yes	Yes	Yes	NT	[64]
		No	I	Calici	No							
				Picornia	No							
					No							
				Astro	No							
			I	Reo	No							
	RNA (segmented, ds)	No		Birna	No							

+: Positive stranded; -: Negative stranded; C: Complex; SARS-CoV: Severe acute respiratory syndrome coronavirus; ds: Double stranded; H: Helical; HNV: Hematopoietic necrosis virus; I: Icosahedral; NT: Not tested; RRV: Ross river virus; ss: Single stranded; VN Ab: Virus-neutralizing antibodies; VS Ab: Virus-specific antibodies.

of spike proteins of coronaviruses for instance changes at pH 8 and, as a result, entry of the virus into the host cell is initiated, but a lower or higher pH is associated with a loss of reactivity [80].

The pH also has an effect on the genomic level (FIGURE 6). The close proximity of the hydroxyl group of RNA to the phosphorous center of each internucleotide linkage facilitates transesterification under strongly acidic or strongly basic conditions [81]. Base-catalyzed reactions proceed via a nucleophilic addition mechanism where the oxygen attacks the adjacent phosphorus center, with a breakage of the phosphodiester bond as a consequence (FIGURE 6).

To date, there are no reports about the use of extreme pH conditions for the inactivation of viruses for KV vaccines. However, inactivation kinetics were performed for SARS-CoV [30]. The infectivity of SARS-CoV is sensitive to pH extremes, as moderate pH variations from 5 to 9 could not inactivate the virus, whereas extremely alkaline (pH 12 and 14) or acidic (pH 1 and 3) conditions could inactivate the virus (TABLE 7).

Preserving the viral structure is a problem if viruses are inactivated under extreme pH conditions. PRRSV inactivation by changing the pH can inactivate the virus, but does not preserve the viral entry-associated domains on PRRSV (TABLE 7). This suggests that, at least in the case of PRRSV, inactivation by changing the pH is not suitable to generate a proper KV vaccine [41].

In conclusion, extreme pH can inactivate viruses, but the effect of a KV vaccine based on this inactivation method in terms of protection against challenge is unknown. It is possible that the outcome of a vaccine based on pH inactivation will give a poor immune response, as the virus structure can be destroyed.

Temperature

By heating or pasteurizing viruses, they can become inactivated by denaturation of the secondary and/or tertiary structures of the viral proteins. It is possible that the conformation of the viral proteins involved in attachment to and replication in a host cell is changed in this process [82,83]. Thermal inactivation of viruses by RNA degradation via breakage of the phosphodiester bond has also been described (FIGURE 6) [84]. Virus inactivation at 'low' temperatures (below 41°C) is considered to be caused by degradation of the nucleic acid, whereas virus inactivation at 'high' temperature is related to protein denaturation [84–86].

Heat inactivation has been studied as a method to develop a KV vaccine for SARS-CoV [30], HIV [87] and HNV (TABLE 7) [67]. To test the ability of heat to inactivate the SARS-CoV, Darnell *et al.* incubated the virus at three temperatures (56, 65 and 75°C) for increasing periods of time [30]. They found that at 56 and 65°C most of the virus was inactivated. However, some virus remained infectious at a level close to the detection limit for the assay, suggesting that some virus particles were stable at 56 and 65°C. One possible explanation for this result may be the presence of aggregates that slowly dissociate. Although the virus was incompletely inactivated at 56 and 65°C, even at 60 min of incubation, it was completely inactivated at 75°C after 45 min of incubation [30]. Taken together, these results suggest that viral inactivation by pasteurization might be very effective if the treatment is long enough.

Table 6. Effect of binary ethylene imine on different viruses: inactivation, viral protein modification, induction of virus-specific and virus-neutralization antibodies and protection on challenge (reduction of viremia and fewer clinical signs).

Study (year)	Genome	Envelope	Capsid	Virus family	Example	Inactivation (concentration)	Viral protein modification	Induction		Protection on challenge		Ref.
								VS Ab	VN Ab	Reduced viremia	Fewer clinical signs	
Awad <i>et al.</i> (2003)	DNA (ss)	No	I	Parvo	No							
				Circo	No							
	DNA (ds)	Yes	I	Hepadna	No							
				Herpes	No							
				Asfor	No							
			C	Pox	Sheep pox	Yes (2%)	NT	Yes	Yes	NT	Yes	[76]
Berhane <i>et al.</i> (2006)			I	Adeno	No							
		No		Papova	No							
	RNA (segmented, ss, -)	Yes	H	Arena	No							
				Bunya	No							
				Orthomyxo	No							
	RNA (nonsegmented, ss, -)	Yes	H	Filo	No							
Berhane <i>et al.</i> (2006)				Paramyxo	Nipah virus	Yes (3 mM)	NT	Yes	No	NT	NT	[78]
Anderson <i>et al.</i> (2008)				Rhabdo	HNW	Yes (1.5 mM)	NT	NT	NT	NT	No	[67]
Aaskov <i>et al.</i> (1997)	RNA (nonsegmented, ss, +)	Yes	I	Toga	RRV	Yes	NT	Yes	Yes	Yes	NT	[75]
Delrue <i>et al.</i> (2009); Vanhee <i>et al.</i> (2009)				Arteri	PRRSV	Yes (1 mM)	No	Yes	Yes	Yes	Yes	[41,79]

+: Positive stranded; -: Negative stranded; C: Complex; ds: Double stranded; FMDV: Foot-and-mouth disease virus; H: Helical; HNW: Hematopoietic necrosis virus; I: Icosahedral; NT: Not tested; PRRSV: Porcine reproductive and respiratory syndrome virus; RRV: Ross river virus; ss: Single stranded; VN Ab: Virus-neutralizing antibodies; VS Ab: Virus-specific antibodies.

Table 6. Effect of binary ethylene imine on different viruses: inactivation, viral protein modification, induction of virus-specific and virus-neutralization antibodies and protection on challenge (reduction of viremia and fewer clinical signs) (cont.).

Study (year)	Genome	Envelope	Capsid	Virus family	Example	Inactivation (concentration)	Viral protein modification	Induction		Protection on challenge		Ref.
								VS Ab	VN Ab	Reduced viremia	Fewer clinical signs	
Race et al. (1995)				Retro	HIV	Yes (10 mM)	No	Yes	Yes	NT	NT	[77]
				Flavi	No							
				Corona	No							
				Calici	No							
Bahnmann et al. (1975)		No		Picornia	FMDV	Yes (1.5 mM)	No	NT	NT	NT	NT	[70]
					No							
				Astro	No							
	RNA (segmented, ds)	No		Reo	No							
				Birna	No							

+: Positive stranded; -: Negative stranded; C: Complex; ds: Double stranded; FMDV: Foot-and-mouth disease virus; H: Helical; HN: Hematopoietic necrosis virus; I: Icosahedral; NT: Not tested; PRRSV: Porcine reproductive and respiratory syndrome virus; RRV: Ross river virus; ss: Single stranded; VN Ab: Virus-neutralizing antibodies; VS Ab: Virus-specific antibodies.

For the preservation of the viral structure, a high temperature is suggested to have an effect on the viral proteins, whereas an inactivation temperature below 41°C is suggested to have no effect on protein level [84–86]. *In vitro* infection assays and western blot analysis showed that in the case of PRRSV the entry-associated domains were indeed preserved after treatment at 37°C and that the virus did not replicate anymore [41]. However, heat-inactivated influenza virus (>55°C) lost hemagglutinin and neuraminidase activity [62].

When it comes to induction of neutralizing antibodies and protection after vaccination, the outcome is vaccine dependent. When BALB/c mice and rhesus macaques (*Macaca mulatta*) were vaccinated with HIV samples inactivated at 62°C, virus-specific antibodies were observed after the first vaccination. In addition, the sera contained antibodies capable of neutralizing the vaccine strain [87]. In another study, HNV inactivated at 50°C did not protect fish against challenge. The percentage mortality of fish in the vaccinated groups was not statistically different compared with the mortality of fish in either the mock-vaccinated or unhandled control groups [67].

In conclusion, it has been shown that temperature can inactivate viruses, but some heat-resistant particles can remain in the inactivated samples. Some heat-inactivated vaccines could induce a neutralizing antibody response, whereas others could not prevent disease. This is probably due to protein modification, because a high inactivation temperature was used. The inactivation temperature should be below 41°C if protein modifications are not desirable.

Irradiation

Ultraviolet light

Based on wavelength, Ultraviolet (UV) light is subdivided into three classifications: UVA (320–400 nm), UVB (280–320 nm) and UVC (200–280 nm) [30]. UVC is absorbed by RNA and DNA bases and can lead to formation of dimers between two adjacent pyrimidines (uracil and thymine) (FIGURE 7). UVB can also induce formation of pyrimidine dimers, but is 20- to 100-fold less efficiently than UVC [88]. UVA is weakly absorbed by RNA and DNA and forms pyrimidine dimers much less efficiently than UVB and UVC [88].

Thus, in general, UV causes formation of pyrimidine dimers between two adjacent or opposing pyrimidines [89,90]. The two pyrimidines react through a pericyclic $2\pi-2\pi$ cycloaddition reaction resulting in the formation of a cyclobutane ring (FIGURE 7). The dimers cause a certain strain in the sugar backbone of the genome, which possibly leads to breaks in the genome. In addition, the uracil dimers formed by UV irradiation inactivate the RNA molecule as a transcription template [91].

This procedure for virus inactivation has been tested to develop a KV vaccine against viruses such as rabbit hemorrhagic disease virus (RHDV) [92], murine leukemia virus strain Cas-Br-M [93] and PRRSV [79] and are summarized in TABLE 8.

Some vaccines based on UV-inactivated viruses are able to induce an immune response. UV-inactivated murine leukemia virus strain Cas-Br-M (UV-Cas), induced a strong, Cas-specific cytotoxic




Figure 5. Reaction mechanism of binary ethylene imine with DNA or RNA (guanine).

G: Guanine.

T-lymphocyte response in newborn NFS/N mice. Vaccination with UV-Cas protected mice against disease and reduced virus replication [93]. In addition, vaccination of pigs with UV-inactivated PRRSV induced virus-specific and virus-neutralizing antibody responses, which were able to reduce viremia after challenge [79].

However, there are also UV-inactivated vaccines that do not give protection against virus infection. Rabbits that were vaccinated with UV-inactivated RHDV and challenged with virulent virus, all died within 82 h after challenge. No antibodies were detected at the time of death. These findings show that vaccination with UV-inactivated RHDV does not protect rabbits against challenge with virulent virus in this study [92].

In terms of the preservation of the viral structure after UV inactivation, caution is necessary. UV could inactivate PRRSV with preservation of the viral entry-associated domains [41], but this is not the case for all viruses. The infectivity of mengovirus is lost very rapidly on exposure to UV irradiation, probably due to dimer formation in the viral RNA [90]. Besides this rapid effect on RNA and DNA, UV irradiation also causes structural modifications of the capsid proteins, resulting in the formation of large and small photoproducts, which is a much slower process [90]. The formation of large photoproduct proteins has also been reported for UV-irradiated reovirus [94].

In conclusion, it has been shown that UV irradiation inactivates viruses properly. If the virus structure is preserved after

inactivation, the KV vaccine based on this UV-inactivated virus can protect the host on challenge. Some KV vaccines based on UV inactivation, however, do not protect the host after challenge with virulent virus. This might be due to the formation of photoproducts that interfere with the viral proteins.

Gamma irradiation

There are two mechanisms by which gamma irradiation can inactivate biological material. The first mechanism is a direct result of a photon depositing energy into

the target. The transfer of this energy results in the dislocation of electrons and breakage of covalent bonds. The second pathway involves indirect damage via free radicals formed after breakage of covalent bonds. Viruses appear to be inactivated primarily by direct damage, via disruption of the genome [95].

This inactivation procedure has been investigated for FMDV, Rauscher leukemia virus, HSV [96] and Lassa virus (TABLE 8) [97]. Inactivation kinetics showed that Rauscher leukemia virus and HSV were inactivated at a dose of 25 kGy. FMDV was inactivated with a dose of 40 kGy, and this was used successfully as antigen in the preparation of a vaccine against FMDV. The vaccine effectively protected the inoculated cattle against the disease [96]. In another study, rhesus monkeys were vaccinated with gamma irradiation-inactivated Lassa virus. The vaccinated animals had antibodies against the three major viral proteins. However, after challenge all the monkeys showed viremia and died. The inactivated Lassa virus vaccine based on gamma irradiation could not protect the animals after challenge [97].

Precaution is necessary when it comes to the preservation of the viral structure on inactivation by gamma irradiation. Gamma irradiation could inactivate PRRSV with the preservation of the viral entry-associated domains [41], but for other viruses the formation of free radicals should be investigated, as they might destroy the viral proteins [95].

In conclusion, gamma irradiation seems to inactivate viruses properly. Some KV vaccines based on gamma irradiation inactivation protect the host against disease after challenge with virulent virus, whereas others cannot prevent viremia and disease. This can be due to the formation of free radicals, which destroy viral proteins.

Expert commentary

Some issues that are important for virus inactivation for vaccine purposes have been defined. A crucial step in the development of a KV vaccine is a balanced inactivation of the virus. All virus particles need to be completely inactivated, but the inactivation




Figure 6. RNA degradation in an alkaline environment.

B: Nucleobase.

Table 7. Effect of denaturing agents on different viruses: inactivation, viral protein modification, induction of virus-specific and virus-neutralization antibodies and protection on challenge (reduction of viremia and fewer clinical signs).


Study (year)	Genome	Envelope	Capsid	Virus Family	Example	Inactivation (concentration)	Viral protein modification	Induction		Protection on challenge		Ref.
								VS Ab	VN Ab	Reduced viremia	Fewer clinical signs	
Jonges <i>et al.</i> (2010)	DNA (ss)	No	I	Parvo	No							
				Circo	No							
	DNA (ds)	Yes	I	Hepadna	No							
				Herpes	No							
				Asfor	No							
				Pox	No							
	RNA (segmented, ss, -)	Yes	H	Adeno	No							
				Papova	No							
				Arena	No							
				Bunya	No							
Anderson <i>et al.</i> (2008)	RNA (nonsegmented, ss, -)	Yes	H	Orthomyxo	Influenza	Yes (>55°C)	Yes	NT	NT	NT	NT	[62]
				Filo	No							
				Paramyxo	No							
				Rhabdo	HNV	Yes (50°C)	NT	NT	NT	NT	No	[67]
Delrue <i>et al.</i> (2009)	RNA (nonsegmented, ss, +)	Yes	I	Borna	No							
				Toga	No							
				No	No							
				Arteri	PRRSV	Yes (pH2 or 12)	Yes	NT	NT	NT	NT	[41]
Delrue <i>et al.</i> (2009)						Yes (37°C)	No	NT	NT	NT	NT	[41]
						Yes (62°C)	NT	Yes	Yes	NT	NT	[87]
Poon <i>et al.</i> (2005)				Retro	HIV							
				Flavi	No							

+: Positive stranded; -: Negative stranded; C: Complex; ds: Double stranded; H: Helical; HNV: Hematopoietic necrosis virus; I: Icosahedral; NT: Not tested; PRRSV: Porcine reproductive and respiratory syndrome virus; SARS-CoV: Severe acute respiratory syndrome virus; ss: Single stranded; VN Ab: Virus-neutralizing antibodies; VS Ab: Virus-specific antibodies.

Table 7. Effect of denaturing agents on different viruses: inactivation, viral protein modification, induction of virus-specific and virus-neutralization antibodies and protection on challenge (reduction of viremia and fewer clinical signs) (cont.).

Study (year)	Genome	Envelope	Capsid	Virus Family	Example	Inactivation (concentration)	Viral protein modification	Induction		Protection on challenge		Ref.
								VS Ab	VN Ab	Reduced viremia	Fewer clinical signs	
Damell <i>et al.</i> (2004)		H		Corona	SARS-CoV	Yes (pH1–3 or 12–14)	NT	NT	NT	NT	NT	[30]
Damell <i>et al.</i> (2004)		No	I	Calici	No	Incomplete (56, 65 and 75°C)	NT	NT	NT	NT	NT	[30]
				Picornia	No							
					No							
				Astro	No							
				Reo	No							
				Birna	No							
	RNA (segmented, ds)	No	I									

+: Positive stranded; -: Negative stranded; C: Complex; ds: Double stranded; H: Helical; HNV: Hematopoietic necrosis virus; I: Icosahedral; NT: Not tested; PRRSV: Porcine reproductive and respiratory syndrome virus; SARS-CoV: Severe acute respiratory syndrome virus; ss: Single stranded; VN Ab: Virus-neutralizing antibodies; VS Ab: Virus-specific antibodies.

**Figure 7. Ultraviolet irradiation results in pyrimidine dimer formation.**

U: Uracil.

method should only have a minor effect on the viral antigenic properties, as the immune system of the host has to recognize the neutralizing epitopes to produce neutralizing antibodies against the antigen. As the antibody response is mostly investigated in vaccination studies, this review focused on the induction of antibodies on vaccination. However, the effect on cell-mediated immune response is also of crucial importance and should be investigated in more depth when evaluating candidate vaccines. The virus inactivation as well as the virus structure preservation can be a problem that should be kept in mind when developing a KV vaccine.

First, during the inactivation process one should be aware of possible virus aggregate formation, which can lead to viruses escaping inactivation. A single virus titration does not always detect escaping virus. A quality control system to investigate the safety of an inactivated virus vaccine can be obtained by passaging the treated virus *in vitro*, which can give more information about the completeness of the virus inactivation. Only if the vaccine passes such a test, can further animal experiments be considered to confirm vaccine safety.

Second, in the development of a KV vaccine, the preservation of the virus structure is very important. Inactivation procedures acting on viral proteins, such as cross-linkers or denaturing procedures [24,39,47,80,82,83], may modify the viral-neutralizing epitopes that are essential for induction of neutralizing antibodies. Thus, it is better to inactivate the virus for KV vaccines with methods that do not affect the viral proteins. Inactivation procedures mainly acting on the genome, such as radiation and alkylating agents [72,90,95], will most likely preserve viral-neutralizing epitopes and should be preferred. Nevertheless, protein degradation after inactivation should always be checked.

Different quality control systems to investigate the antigen quality are available. For influenza, the effect of inactivation is investigated by measuring the hemagglutinating activity before and after inactivation [98]. ELISA may be used for the determination of the attachment of neutralizing antibodies to viral epitopes after inactivation. This is used as quality control of the antigens of HIV [87,99], rabies virus [100,101] and PV vaccines [102]. However, for the previous methods, knowledge of the neutralizing epitopes is necessary. Sometimes, the neutralizing epitopes are not known. As

Table 8. Effect of irradiation on different viruses: inactivation, viral protein modification, induction of virus-specific and virus-neutralization antibodies and protection on challenge (reduction of viremia and fewer clinical signs).

Study (year)	Genome	Envelope	Capsid	Virus Family	Example	Inactivation (concentration)	Viral protein modification	Induction		Protection on challenge		Ref.
								VS Ab	VN Ab	Reduced viremia	Fewer clinical signs	
Smolko <i>et al.</i> (2005)	DNA (ss)	No	I	Parvo	No							
				Circo	No							
	DNA (ds)	Yes	I	Hepadna	No							
				Herpes	HSV	Yes (25 kGy)	NT	NT	NT	NT	NT	[96]
				Asfor	No							
			C	Pox	No							
McCormick <i>et al.</i> (1992)		No	I	Adeno	No							
				Papova	No							
	RNA (segmented, ss, -)	Yes	H	Arena	Lassa	Yes (gamma)	NT	Yes	NT	No	No	[97]
				Bunya	No							
				Orthomyxo	No							
	RNA (nonsegmented, ss, -)	Yes	H	Filo	No							
				Paramyxo	No							
				Rhabdo	No							
				Borna	No							
	RNA (nonsegmented, ss, +)	Yes	I	Toga	No							
					No							

+: Positive stranded; -: Negative stranded; C: Complex; ds: Double stranded; FMDV: Foot-and-mouth disease virus; H: Helical; I: Icosahedral; MLV: Murine leukemia virus; NT: Not tested; PRRSV: Porcine reproductive and respiratory syndrome virus; RHDV: Rabbit hemorrhagic disease virus; RLV: Rauscher leukemia virus; ss: Single stranded; VN Ab: Virus-neutralizing antibodies; VS Ab: Virus-specific antibodies.

Table 8. Effect of irradiation on different viruses: inactivation, viral protein modification, induction of virus-specific and virus-neutralization antibodies and protection on challenge (reduction of viremia and fewer clinical signs) (cont.).

Study (year)	Genome	Envelope	Capsid	Virus Family	Example	Inactivation (concentration)	Viral protein modification	Induction		Protection on challenge		Ref.
								VS Ab	VN Ab	Reduced viremia	Fewer clinical signs	
Delrue <i>et al.</i> (2009)				Arteri	PRRSV	Yes (0.25 kGy)	No	NT	NT	NT	NT	[41]
Delrue <i>et al.</i> (2009); Vanhee <i>et al.</i> (2009)						Yes (100 mJ/cm ²)	No	Yes	Yes	Yes	NT	[41,79]
Sarzotti <i>et al.</i> (1994)				Retro	MLV	Yes (120 ergs/s/mm ²)	NT	NT	NT	Yes	Yes	[93]
Smolko <i>et al.</i> (2005)					RLV	Yes (25 kGy)	NT	NT	NT	NT	NT	[96]
				Flavi	No							
			H	Corona	No							
Henning <i>et al.</i> (2005)		No	I	Calici	RHDV	Yes (0.0078 W/cm ²)	NT	No	No	No	No	[92]
Smolko <i>et al.</i> (2005)				Picorna	FMDV	Yes (40 kGy)	NT	NT	NT	NT	Yes	[96]
Miller <i>et al.</i> (1974)					Mengovirus	Yes (7 kergs/min/mm ²)	Yes	NT	NT	NT	NT	[90]
					No							
				Astro	No							
Subasinghe <i>et al.</i> (1972)	RNA segmented, ds)	No	I	Reo	Reovirus	Yes (ultraviolet)	Yes	NT	NT	NT	NT	[94]
				Birna	No							

+: Positive stranded; -: Negative stranded; C: Complex; ds: Double stranded; FMDV: Foot-and-mouth disease virus; H: Helical; I: Icosahedral; MLV: Murine leukemia virus; NT: Not tested; PRRSV: Porcine reproductive and respiratory syndrome virus; RHDV: Rabbit hemorrhagic disease virus; RLV: Rauscher leukemia virus; ss: Single stranded; VN Ab: Virus-neutralizing antibodies; VS Ab: Virus-specific antibodies.

virus-neutralizing antibodies are often directed to viral proteins involved in virus entry [41,103–105], it can be advised to preserve virus entry-associated domains to induce a virus-neutralizing antibody response. In this context, a quality control system for investigating the antigen quality of the inactivated virus can be developed by confirming the preservation of entry-associated domains of a virus [41]. This can be achieved by investigating attachment and internalization of virus into the host cells before and after virus inactivation by immunofluorescence staining and confocal microscopy [41].

In conclusion, many methods have been developed that are potentially suitable to inactivate a certain vaccine virus, but the success of these methods may vary depending on the particular virus and the conditions used (i.e., temperature, pH and concentration). Nonetheless, successful methods to inactivate a virus for vaccine development are best chosen from methods that mainly have an effect on the genome. Since formaldehyde, glutaraldehyde, pH and heat have an influence on protein level, it is possible that the neutralizing epitopes are modified by cross-linking (formaldehyde or glutaraldehyde) or denaturation (pH or temperature). In our opinion, UV irradiation, gamma irradiation, BPL and BEI are the most successful methods to inactivate a virus to create a KV vaccine, as these inactivation procedures mainly target the genome and have no or minor effects on viral proteins. However, it has to be considered that viral proteins may be cross-linked by BPL, or damaged by UV photoproducts and free radicals formed by gamma irradiation. The radiation target theory, however, predicts that the radiation sensitivity of biomolecules depends on their mass [106]. Therefore, viral genomes, with a significantly lower mass, would be significantly more sensitive to damage than proteins.

Five-year view

Until recently, the search for an effective KV vaccine was often a process of trial and error, in which a virus was inactivated by a certain inactivation method and then immediately tested in animals

to investigate complete virus inactivation, antibody response on vaccination and protection on challenge. Currently, a series of *in vitro* tools is available, allowing the creation of a quality control system, in which virus inactivation (safety) and antigen quality (immune response) can be investigated. In this way, vaccine efficacy can be predicted prior to animal experiments. As a consequence, outbreaks due to incomplete virus inactivation and poor immune responses due to viral protein destruction can be avoided as much as possible.

It appears that virus inactivation methods exerting their main effects on the viral genome have most potential in generating an effective immune response and protecting the host against disease. On *in vitro* testing of several inactivation methods following a quality control system for an inactivated PRRSV vaccine, we have observed that only a few of the available methods allow complete virus inactivation with preservation of the virion structure. Considering the molecular and mechanistic basis of these evaluated methods, it is clear that they all mainly affect the viral genome, leaving the viral proteins unaffected. In our opinion, it is therefore of crucial importance for the induction of a proper immune response and protection against disease, to evolve the development methods for KV vaccines based on viral genome inactivation, which can guarantee an intact virion structure.

Financial & competing interests disclosure

I Delrue was funded by the Agricultural Research Program from the Institute for the Promotion of Innovation by Science and Technology in Flanders (IWT-LO 040721). D Verzele and A Madder acknowledge financial support from Ghent University (BOF-01J06111). H Nauwynck acknowledges the EU (Seventh Framework Program; Project No. 245141) for financial support. The authors have no other relevant affiliations or financial involvement with any organization or entity with a financial interest in or financial conflict with the subject matter or materials discussed in the manuscript apart from those disclosed.

No writing assistance was utilized in the production of this manuscript

Key issues

- For safety reasons, killed virus (KV) vaccines are preferred over modified live virus vaccines. However, KV vaccines can suffer from problems related to efficacy.
- A good *in vitro* quality control system is required for prediction of the effectiveness of a KV vaccine. This quality control system should allow verification of the completeness of virus inactivation and the quality of the resulting antigen.
- In order to create a safe KV vaccine, the virus inactivation method should inactivate the virus mixture completely without risk of resistant virus particles. Due to aggregate formation, incompletely inactivated virus particles can still be present in the vaccine with outbreaks as a consequence.
- For the generation of a KV vaccine able to induce a proper immune response, antigen quality is of high importance. Due to protein crosslinking, protein denaturation and protein degradation, the viral antigen can be completely destroyed, with a poor immune response as a consequence.
- Crosslinkers, such as formaldehyde and glutaraldehyde, or denaturing processes, such as pH and heat, can have an influence on viral proteins and destroy the virion structure. As a consequence, KV vaccines based on these inactivation methods might not induce a proper immune response.
- Next to alkylating agents, such as β -propiolactone and binary ethylene imine, ultraviolet and gamma irradiation mainly influence the viral genome, preserving the virion structure. These methods are thus highly recommended for generation of an efficient and safe KV vaccine.

References

Papers of special note have been highlighted as:
• of interest

- King AM, Underwood BO, McCahon D, Newman JW, Brown F. Biochemical identification of viruses causing the 1981 outbreaks of foot and mouth disease in the UK. *Nature* 293(5832), 479–480 (1981).
- Beck E, Strohmaier K. Subtyping of European foot-and-mouth disease virus strains by nucleotide sequence determination. *J. Virol.* 61(5), 1621–1629 (1987).
- Patil PK, Suryanarayana V, Bist P, Bayry J, Natarajan C. Integrity of GH-loop of foot-and-mouth disease virus during virus inactivation: detection by epitope specific antibodies. *Vaccine* 20(7–8), 1163–1168 (2002).
- Cham BE, Vickery K, Tohidi-Esfahani R, Cossart Y. Delipidation of a hepadnavirus: Viral inactivation and vaccine development. *J. Virol. Methods* 137(1), 160–163 (2006).
- Niedrig M, Gregersen JP, Fultz PN, Bröker M, Mehdi S, Hilfenhaus J. Immune response of chimpanzees after immunization with the inactivated whole immunodeficiency virus (HIV-1), three different adjuvants and challenge. *Vaccine* 11(1), 67–74 (1993).
- Jahrling PB, Stephenson EH. Protective efficacies of live attenuated and formaldehyde-inactivated Venezuelan equine encephalitis virus vaccines against aerosol challenge in hamsters. *J. Clin. Microbiol.* 19(3), 429–431 (1984).
- Kistner O, Barrett N, Brühmann A *et al.* The preclinical testing of a formaldehyde inactivated Ross River virus vaccine designed for use in humans. *Vaccine* 25(25), 4845–4852 (2007).
- Samina I, Khinich Y, Simanov M, Malkinson M. An inactivated West Nile virus vaccine for domestic geese—efficacy study and a summary of 4 years of field application. *Vaccine* 23(41), 4955–4958 (2005).
- Putnak R, Barvir DA, Burrous JM *et al.* Development of a purified, inactivated, dengue-2 virus vaccine prototype in Vero cells: immunogenicity and protection in mice and rhesus monkeys. *J. Infect. Dis.* 174(6), 1176–1184 (1996).
- Skinner GR, Davies J. Efficacy of an inactivated vaccine for equine herpesvirus type 1 in a novel hamster model. *Intervirology* 43(1), 27–35 (2000).
- Bell JA, Sundberg JP, Ghim SJ, Newsome J, Jensen AB, Schlegel R. A formalin-inactivated vaccine protects against mucosal papillomavirus infection: a canine model. *Pathobiology* 62(4), 194–198 (1994).
- Choi Y, Ahn CJ, Seong KM, Jung MY, Ahn BY. Inactivated Hantaan virus vaccine derived from suspension culture of Vero cells. *Vaccine* 21(17–18), 1867–1873 (2003).
- Bahgat MM, Kutkat MA, Nasraa MH *et al.* Characterization of an avian influenza virus H5N1 Egyptian isolate. *J. Virol. Methods* 159(2), 244–250 (2009).
- Furesz J, Scheifele DW, Palkonyay L. Safety and effectiveness of the new inactivated hepatitis A virus vaccine. *CMAJ* 152(3), 343–348 (1995).
- Goncharova E, Ryzhikov E, Poryvaev V *et al.* Intranasal immunization with inactivated tick-borne encephalitis virus and the antigenic peptide 89–119 protects mice against intraperitoneal challenge. *Int. J. Med. Microbiol.* 296(Suppl. 40), 195–201 (2006).
- Heinz F, Holtzmann H, Essl A, Kundt M. [Analysis of the efficiency of tick-borne encephalitis vaccination in the population in the natural foci of Austria]. *Vopr. Virusol.* 53(2), 19–27 (2008).
- Appaiahgari MB, Vratsi S. Immunogenicity and protective efficacy in mice of a formaldehyde-inactivated Indian strain of Japanese encephalitis virus grown in Vero cells. *Vaccine* 22(27–28), 3669–3675 (2004).
- Alderson T. The mechanism of formaldehyde-induced mutagenesis. The monohydroxymethylation reaction of formaldehyde with adenylic acid as the necessary and sufficient condition for the mediation of the mutagenic activity of formaldehyde. *Mutat. Res.* 106, 77–85 (1964).
- Progress in the nucleic acid research and molecular biology.* Davidson JN (Ed.). Academic Press, New York, NY (1973).
- Fraenkel-Conrat H. Reaction of nucleic acid with formaldehyde. *Biochim. Biophys. Acta* 15(2), 307–309 (1954).
- Fraenkel-Conrat H. Chemical modification of viral ribonucleic acid. I. Alkylating agents. *Biochim. Biophys. Acta* 49, 169–180 (1961).
- Michelson AM, Letters R. Synthesis of nucleotide-aminoacyl anhydrides. *Biochim. Biophys. Acta* 80, 242–246 (1964).
- Metz B, Kersten GF, Hoogerhout P *et al.* Identification of formaldehyde-induced modifications in proteins: reactions with model peptides. *J. Biol. Chem.* 279(8), 6235–6243 (2004).
- In-depth study of the modifications in proteins caused by formaldehyde treatment, revising the chemical details through model peptides.
- Fraenkel-Conrat H, Mecham DK. The reaction of formaldehyde with proteins; demonstration of intermolecular cross-linking by means of osmotic pressure measurements. *J. Biol. Chem.* 177(1), 477–486 (1949).
- Kuykendall JR, Bogdanffy MS. Efficiency of DNA-histone crosslinking induced by saturated and unsaturated aldehydes *in vitro*. *Mutat. Res.* 283(2), 131–136 (1992).
- Ma TH, Harris MM. Review of the genotoxicity of formaldehyde. *Mutat. Res.* 196(1), 37–59 (1988).
- Permana PA, Snapka RM. Aldehyde-induced protein-DNA crosslinks disrupt specific stages of SV40 DNA replication. *Carcinogenesis* 15(5), 1031–1036 (1994).
- Brown F. Review of accidents caused by incomplete inactivation of viruses. *Dev. Biol. Stand.* 81, 103–107 (1993).
- Twomey T, Newman J, Burrage T, Piatti P, Lubroth J, Brown F. Structure and immunogenicity of experimental foot-and-mouth disease and poliomyelitis vaccines. *Vaccine* 13(16), 1603–1610 (1995).
- Darnell ME, Subbarao K, Feinstone SM, Taylor DR. Inactivation of the coronavirus that induces severe acute respiratory syndrome, SARS-CoV. *J. Virol. Methods* 121(1), 85–91 (2004).
- Spruth M, Kistner O, Savidis-Dacho H *et al.* A double-inactivated whole virus candidate SARS coronavirus vaccine stimulates neutralising and protective antibody responses. *Vaccine* 24(5), 652–661 (2006).
- Qu D, Zheng B, Yao X *et al.* Intranasal immunization with inactivated SARS-CoV (SARS-associated coronavirus) induced local and serum antibodies in mice. *Vaccine* 23(7), 924–931 (2005).
- Zhang CH, Guo ZM, Zheng HY *et al.* Humoral immune responses in rabbits induced by an experimental inactivated severe acute respiratory syndrome coronavirus vaccine prepared from F69 strain. *Chin. Med. J.* 117(11), 1625–1629 (2004).
- Openshaw PJ, Culley FJ, Olszewska W. Immunopathogenesis of vaccine-enhanced RSV disease. *Vaccine* 20(Suppl. 1), S27–S31 (2001).
- Ishizaka ST, Israel ZR, Gettie A *et al.* Induction of mucosal antibody responses by microsphere-encapsulated

- formalin-inactivated simian immunodeficiency virus in a male urethral challenge model. *Vaccine* 17(22), 2817–2825 (1999).
- 36 Huisman W, Martina BE, Rimmelzwaan GF, Gruters RA, Osterhaus AD. Vaccine-induced enhancement of viral infections. *Vaccine* 27(4), 505–512 (2009).
 - 37 Griffin DE, Pan CH, Moss WJ. Measles vaccines. *Front. Biosci.* 13, 1352–1370 (2008).
 - 38 Kalina WV, Woolums AR, Berghaus RD, Gershwin LJ. Formalin-inactivated bovine RSV vaccine enhances a Th2 mediated immune response in infected cattle. *Vaccine* 22(11–12), 1465–1474 (2004).
 - 39 Tano Y, Shimizu H, Martin J, Nishimura Y, Simizu B, Miyamura T. Antigenic characterization of a formalin-inactivated poliovirus vaccine derived from live-attenuated Sabin strains. *Vaccine* 25(41), 7041–7046 (2007).
 - 40 Delpitte PL, Meerts P, Costers S, Nauwynck HJ. Effect of virus-specific antibodies on attachment, internalization and infection of porcine reproductive and respiratory syndrome virus in primary macrophages. *Vet. Immunol. Immunopathol.* 102(3), 179–188 (2004).
 - 41 Delrue I, Delpitte PL, Nauwynck HJ. Assessing the functionality of viral entry-associated domains of porcine reproductive and respiratory syndrome virus during inactivation procedures, a potential tool to optimize inactivated vaccines. *Vet. Res.* 40(6), 62 (2009).
 - **Describes a quality control system for virus inactivation and viral protein preservation to evaluate different inactivation methods for the development of a porcine reproductive and respiratory syndrome virus vaccine.**
 - 42 Rossio JL, Esser MT, Suryanarayana K *et al.* Inactivation of human immunodeficiency virus type 1 infectivity with preservation of conformational and functional integrity of virion surface proteins. *J. Virol.* 72(10), 7992–8001 (1998).
 - 43 Pollock RV, Carmichael LE. Dog response to inactivated canine parvovirus and feline panleukopenia virus vaccines. *Cornell Vet.* 72(1), 16–35 (1982).
 - 44 *Embalming Chemistry: Glutaraldehyde Versus Formaldehyde.* Bedino JH (Ed.). The Champion Company, Springfield, OH, USA (2003).
 - 45 Hemminki K, Suni R. Sites of reaction of glutaraldehyde and acetaldehyde with nucleosides. *Arch. Toxicol.* 55(3), 186–190 (1984).
 - 46 McGucken PV, Woodside W. Studies on the mode of action of glutaraldehyde on *Escherichia coli*. *J. Appl. Bacteriol.* 36(3), 419–426 (1973).
 - 47 Cheung DT, Nimni ME. Mechanism of crosslinking of proteins by glutaraldehyde II. Reaction with monomeric and polymeric collagen. *Connect. Tissue Res.* 10(2), 201–216 (1982).
 - 48 Richards FM, Knowles JR. Glutaraldehyde as a protein cross-linkage reagent. *J. Mol. Biol.* 37(1), 231–233 (1968).
 - 49 *Advances in Protein Chemistry.* Anfinsen CB (Ed.). Academic Press, New York, NY, USA (1970).
 - 50 *Bioconjugate Techniques.* Hermanson GT (Ed.). Academic Press, San Diego, CA, USA (2008).
 - 51 Keles I, Woldehiwet Z, Murray RD. Vaccination with glutaraldehyde-fixed bovine respiratory syncytial virus (BRSV)-infected cells stimulates a better immune response in lambs than vaccination with heat-inactivated cell-free BRSV. *Vaccine* 16(11–12), 1172–1178 (1998).
 - 52 Cepica A, Beauregard M, Qian B. Fluorescence spectroscopy monitoring of the conformational restraint of formaldehyde- and glutaraldehyde-treated infectious bursal disease virus proteins. *Vaccine* 16(20), 1957–1961 (1998).
 - 53 Boukhvalova MS, Prince GA, Blanco JC. The cotton rat model of respiratory viral infections. *Biologicals* 37(3), 152–159 (2009).
 - 54 Chertova E, Crise BJ, Morcock DR, Bess JW Jr, Henderson LE, Lifson JD. Sites, mechanism of action and lack of reversibility of primate lentivirus inactivation by preferential covalent modification of virion internal proteins. *Curr. Mol. Med.* 3(3), 265–272 (2003).
 - 55 Wootton SK, Yoo D. Homo-oligomerization of the porcine reproductive and respiratory syndrome virus nucleocapsid protein and the role of disulfide linkages. *J. Virol.* 77(8), 4546–4557 (2003).
 - 56 Lifson JD, Rossio JL, Piatak M Jr *et al.* Evaluation of the safety, immunogenicity, and protective efficacy of whole inactivated simian immunodeficiency virus (SIV) vaccines with conformationally and functionally intact envelope glycoproteins. *AIDS Res. Hum. Retroviruses* 20(7), 772–787 (2004).
 - 57 Vagenas P, Williams VG, Piatak M Jr *et al.* Tonsillar application of AT-2 SIV affords partial protection against rectal challenge with SIVmac239. *J. Acquir. Immune Defic. Syndr.* 52(4), 433–442 (2009).
 - 58 Perrin P, Morgeaux S. Inactivation of DNA by beta-propiolactone. *Biologicals* 23(3), 207–211 (1995).
 - 59 Uittenbogaard JP, Zomer B, Hoogerhout P, Metz B. Reactions of beta-propiolactone with nucleobase analogues, nucleosides, and peptides: implications for the inactivation of viruses. *J. Biol. Chem.* 286(42), 36198–36214 (2011).
 - **Recent elucidation and revision of the chemical implications on DNA, RNA and proteins by β -propiolactone virus inactivation, via a systematic study on nucleobase, nucleoside and amino acid test systems.**
 - 60 Larghi OP, Nebel AE. Rabies virus inactivation by binary ethylenimine: new method for inactivated vaccine production. *J. Clin. Microbiol.* 11(2), 120–122 (1980).
 - 61 Desbat B, Lancelot E, Krell T *et al.* Effect of the β -propiolactone treatment on the adsorption and fusion of influenza A/Brisbane/59/2007 and A/New Caledonia/20/1999 virus H1N1 on a dimyristoylphosphatidylcholine/ganglioside GM3 mixed phospholipids monolayer at the air-water interface. *Langmuir* 27(22), 13675–13683 (2011).
 - 62 Jonges M, Liu WM, van der Vries E *et al.* Influenza virus inactivation for studies of antigenicity and phenotypic neuraminidase inhibitor resistance profiling. *J. Clin. Microbiol.* 48(3), 928–940 (2010).
 - 63 Monath TP, Fowler E, Johnson CT *et al.* An inactivated cell-culture vaccine against yellow fever. *N. Engl. J. Med.* 364(14), 1326–1333 (2011).
 - 64 Roberts A, Lamirande EW, Vogel L *et al.* Immunogenicity and protective efficacy in mice and hamsters of a β -propiolactone inactivated whole virus SARS-CoV vaccine. *Viral Immunol.* 23(5), 509–519 (2010).
 - 65 Bodewes R, Kreijtz JH, van Amerongen G *et al.* A single immunization with CoVaccine HT-adjuvanted H5N1 influenza virus vaccine induces protective cellular and humoral immune responses in ferrets. *J. Virol.* 84(16), 7943–7952 (2010).
 - 66 Sabchareon A, Lang J, Attanath P *et al.* A new Vero cell rabies vaccine: results of a comparative trial with human diploid cell rabies vaccine in children. *Clin. Infect. Dis.* 29(1), 141–149 (1999).
 - 67 Anderson E, Clouthier S, Shewmaker W, Weighall A, LaPatra S. Inactivated infectious haematopoietic necrosis virus (IHNV) vaccines. *J. Fish Dis.* 31(10), 729–745 (2008).

- 68 Warrington RE, Cunliffe HR, Bachrach HL. Derivatives of aziridine as inactivants for foot-and-mouth disease virus vaccines. *Am. J. Vet. Res.* 34(8), 1087–1091 (1973).
- 69 Bahnemann HG. Inactivation of viruses in serum with binary ethyleneimine. *J. Clin. Microbiol.* 3(2), 209–210 (1976).
- 70 Bahnemann HG. Binary ethyleneimine as an inactivant for foot-and-mouth disease virus and its application for vaccine production. *Arch. Virol.* 47(1), 47–56 (1975).
- 71 Bahnemann HG. Inactivation of viral antigens for vaccine preparation with particular reference to the application of binary ethyleneimine. *Vaccine* 8(4), 299–303 (1990).
- 72 Broo K, Wei J, Marshall D *et al.* Viral capsid mobility: a dynamic conduit for inactivation. *Proc. Natl Acad. Sci. USA* 98(5), 2274–2277 (2001).
- 73 Gates KS, Noonan T, Dutta S. Biologically relevant chemical reactions of N7-alkylguanine residues in DNA. *Chem. Res. Toxicol.* 17(7), 839–856 (2004).
- 74 Hendler S, Furer E, Srinivasan PR. Synthesis and chemical properties of monomers and polymers containing 7-methylguanine and an investigation of their substrate or template properties for bacterial deoxyribonucleic acid or ribonucleic acid polymerases. *Biochemistry* 9(21), 4141–4153 (1970).
- 75 Aaskov J, Williams L, Yu S. A candidate Ross River virus vaccine: preclinical evaluation. *Vaccine* 15(12–13), 1396–1404 (1997).
- 76 Awad M, Michael A, Soliman SM, Samir SS, Daoud AM. Trials for preparation of inactivated sheep pox vaccine using binary ethyleneimine. *Egypt. J. Immunol.* 10(2), 67–72 (2003).
- 77 Race E, Frezza P, Stephens DM *et al.* An experimental chemically inactivated HIV-1 vaccine induces antibodies that neutralize homologous and heterologous viruses. *Vaccine* 13(1), 54–60 (1995).
- 78 Berhane Y, Berry JD, Ranadheera C *et al.* Production and characterization of monoclonal antibodies against binary ethyleneimine inactivated Nipah virus. *J. Virol. Methods* 132(1–2), 59–68 (2006).
- 79 Vanhee M, Delpitte PL, Delrue I, Geldhof MF, Nauwynck HJ. Development of an experimental inactivated PRRSV vaccine that induces virus-neutralizing antibodies. *Vet. Res.* 40(6), 63 (2009).
- **Vaccination study with an inactivated porcine reproductive and respiratory syndrome virus vaccine developed based on the results of an *in vitro* quality control system to evaluate virus inactivation and viral protein preservation.**
- 80 Weismiller DG, Sturman LS, Buchmeier MJ, Fleming JO, Holmes KV. Monoclonal antibodies to the peplomer glycoprotein of coronavirus mouse hepatitis virus identify two subunits and detect a conformational change in the subunit released under mild alkaline conditions. *J. Virol.* 64(6), 3051–3055 (1990).
- 81 Li Y, Breaker RR. Kinetics of RNA degradation by specific base catalysis of transesterification involving the 2'-hydroxyl group. *J. Am. Chem. Soc.* 121, 5364–5372 (1999).
- 82 Lelie PN, Reesink HW, Lucas CJ. Inactivation of 12 viruses by heating steps applied during manufacture of a hepatitis B vaccine. *J. Med. Virol.* 23(3), 297–301 (1987).
- 83 Schlegel A, Immelmann A, Kempf C. Virus inactivation of plasma-derived proteins by pasteurization in the presence of guanidine hydrochloride. *Transfusion* 41(3), 382–389 (2001).
- 84 Fleming P. Thermal inactivation of Semliki Forest virus. *J. Gen. Virol.* 13(3), 385–391 (1971).
- 85 Dimmock NJ. Differences between the thermal inactivation of picornaviruses at “high” and “low” temperatures. *Virology* 31(2), 338–353 (1967).
- 86 Laude H. Thermal inactivation studies of a coronavirus, transmissible gastroenteritis virus. *J. Gen. Virol.* 56(Pt 2), 235–240 (1981).
- 87 Poon B, Safrit JT, McClure H *et al.* Induction of humoral immune responses following vaccination with envelope-containing, formaldehyde-treated, thermally inactivated human immunodeficiency virus type 1. *J. Virol.* 79(8), 4927–4935 (2005).
- 88 Perdiz D, Grof P, Mezzina M, Nikaido O, Moustacchi E, Sage E. Distribution and repair of bipyrimidine photoproducts in solar UV-irradiated mammalian cells. Possible role of Dewar photoproducts in solar mutagenesis. *J. Biol. Chem.* 275(35), 26732–26742 (2000).
- 89 Sinha RP, Häder DP. UV-induced DNA damage and repair: a review. *Photochem. Photobiol. Sci.* 1(4), 225–236 (2002).
- 90 Miller RL, Plagemann PG. Effect of ultraviolet light on mengovirus: formation of uracil dimers, instability and degradation of capsid, and covalent linkage of protein to viral RNA. *J. Virol.* 13(3), 729–739 (1974).
- 91 Sauerbier W, Hercules K. Gene and transcription unit mapping by radiation effects. *Annu. Rev. Genet.* 12, 329–363 (1978).
- 92 Henning J, Meers J, Davies PR. Exposure of rabbits to ultraviolet light-inactivated rabbit haemorrhagic disease virus (RHDV) and subsequent challenge with virulent virus. *Epidemiol. Infect.* 133(4), 731–735 (2005).
- 93 Sarzotti M, Dean TA, Remington M, Hoffman PM. Ultraviolet-light-inactivated Cas-Br-M murine leukemia virus induces a protective CD8⁺ cytotoxic T lymphocyte response in newborn mice. *AIDS Res. Hum. Retroviruses* 10(12), 1695–1702 (1994).
- 94 Subasinghe HA, Loh PC. Reovirus cytotoxicity: some properties of the UV-irradiated reovirus and its capsid proteins. *Arch. Gesamte Virusforsch.* 39(1), 172–189 (1972).
- 95 Grieb T, Forng RY, Brown R *et al.* Effective use of gamma irradiation for pathogen inactivation of monoclonal antibody preparations. *Biologicals* 30(3), 207–216 (2002).
- 96 Smolko EE, Lombardo JH. Virus inactivation studies using ion beams, electron and gamma irradiation. *Nuclear Instrum. Methods Physics Res. Sec. B-Beam Interact. Mater. Atoms* 236, 249–253 (2005).
- 97 McCormick JB, Mitchell SW, Kiley MP, Ruos S, Fisher-Hoch SP. Inactivated Lassa virus elicits a non protective immune response in rhesus monkeys. *J. Med. Virol.* 37(1), 1–7 (1992).
- 98 Di Trani L, Cordioli P, Falcone E *et al.* Standardization of an inactivated H1N1 avian influenza vaccine and efficacy against A/Chicken/Italy/13474/99 high-pathogenicity virus infection. *Avian Dis.* 47(Suppl. 3), 1042–1046 (2003).
- 99 Grovit-Ferbas K, Hsu JF, Ferbas J, Gudeman V, Chen IS. Enhanced binding of antibodies to neutralization epitopes following thermal and chemical inactivation of human immunodeficiency virus type 1. *J. Virol.* 74(13), 5802–5809 (2000).
- 100 Rooijakkers EJ, Uittenbogaard JP, Groen J, Osterhaus AD. Rabies vaccine potency control: comparison of ELISA systems for antigenicity testing. *J. Virol. Methods* 58(1–2), 111–119 (1996).
- 101 Fournier-Caruaña J, Poirier B, Haond G *et al.* Inactivated rabies vaccine control and release: use of an ELISA method. *Biologicals* 31(1), 9–16 (2003).

- 102 Morgeaux S, Milne C, Daas A. Feasibility study to develop a common *in vitro* D antigen assay for inactivated poliomyelitis vaccines. *Pharmeuropa Bio* 2005(1), 19–26 (2005).
- 103 Owsianka AM, Tarr AW, Keck ZY *et al.* Broadly neutralizing human monoclonal antibodies to the hepatitis C virus E2 glycoprotein. *J. Gen. Virol.* 89(Pt 3), 653–659 (2008).
- 104 Lee JE, Saphire EO. Neutralizing ebolavirus: structural insights into the envelope glycoprotein and antibodies targeted against it. *Curr. Opin. Struct. Biol.* 19(4), 408–417 (2009).
- 105 Coughlin MM, Babcook J, Prabhakar BS. Human monoclonal antibodies to SARS-coronavirus inhibit infection by different mechanisms. *Virology* 394(1), 39–46 (2009).
- 106 Osborne JC Jr, Miller JH, Kempner ES. Molecular mass and volume in radiation target theory. *Biophys. J.* 78(4), 1698–1702 (2000).
- 107 Feron VJ, Til HP, de Vrijer F, Woutersen RA, Cassee FR, van Bladeren PJ. Aldehydes: occurrence, carcinogenic potential, mechanism of action and risk assessment. *Mutat. Res.* 259(3–4), 363–385 (1991).