


FRANCIS A.
CAREY



Organic Chemistry

SIXTH EDITION

Organic Chemistry

Sixth Edition

Francis A. Carey
University of Virginia



Higher Education

Boston Burr Ridge, IL Dubuque, IA Madison, WI New York San Francisco St. Louis
Bangkok Bogotá Caracas Kuala Lumpur Lisbon London Madrid Mexico City
Milan Montreal New Delhi Santiago Seoul Singapore Sydney Taipei Toronto

The McGraw-Hill Companies



Higher Education

ORGANIC CHEMISTRY, SIXTH EDITION

Published by McGraw-Hill, a business unit of The McGraw-Hill Companies, Inc., 1221 Avenue of the Americas, New York, NY 10020. Copyright © 2006, 2003, 2000, 1996, 1992, 1987 by The McGraw-Hill Companies, Inc. All rights reserved. No part of this publication may be reproduced or distributed in any form or by any means, or stored in a database or retrieval system, without the prior written consent of The McGraw-Hill Companies, Inc., including, but not limited to, in any network or other electronic storage or transmission, or broadcast for distance learning.

Some ancillaries, including electronic and print components, may not be available to customers outside the United States.

This book is printed on acid-free paper.

2 3 4 5 6 7 8 9 0 DOW/DOW 0 9 8 7 6 5

ISBN 0-07-282837-4

Editorial Director: *Kent A. Peterson*

Sponsoring Editor: *Thomas D. Timp*

Managing Developmental Editor: *Shirley R. Oberbroeckling*

Senior Developmental Editor: *Mary E. Hurley*

Senior Marketing Manager: *Tamara L. Good-Hodge*

Project Manager: *Jodi Rhomberg*

Production Supervisor: *Kara Kudronowicz*

Lead Media Project Manager: *Judi David*

Senior Media Technology Producer: *Jeffrey Schmitt*

Senior Designer: *David W. Hash*

(USE) Cover Image: *Baldrige, K.K.; Siegel, J.S. Corannulene-Based Fullerene Fragments C₂₀H₁₀-C₅₀H₁₀:*

When Does a Buckybowl become a Buckytube? Theor. Chem. Acta., 1997, 97, 67-71.

Senior Photo Research Coordinator: *John C. Leland*

Photo Research: *Mary Reeg*

Supplement Producer: *Brenda A. Ernzen*

Compositor: *The GTS Companies/Los Angeles, CA Campus*

Typeface: *10.5/12 Times Roman*

Printer: *R. R. Donnelley Willard, OH*

The credits section for this book begins on page C1 and is considered an extension of the copyright page.

Library of Congress Cataloging-in-Publication Data

Carey, Francis A., 1937-

Organic chemistry / Francis A. Carey. — 6th ed.

p. cm.

Includes index.

ISBN 0-07-282837-4 (acid-free paper)

1. Chemistry, Organic. I. Title.

QD251.3.C37 2006

547—dc22

2004015260

CIP


ABOUT THE AUTHOR

Francis A. Carey is a native of Pennsylvania, educated in the public schools of Philadelphia, at Drexel University (B.S. in chemistry, 1959), and at Penn State (Ph.D. 1963). Following postdoctoral work at Harvard and military service, he was appointed to the chemistry faculty of the University of Virginia in 1966. Prior to retiring in 2000, he regularly taught the two-semester lecture courses in general chemistry and organic chemistry.

With his students, Professor Carey published over 40 research papers in synthetic and mechanistic organic chemistry. In addition to this text, he is coauthor (with Robert C. Atkins) of *Organic Chemistry: A Brief Course* and (with Richard J. Sundberg) of *Advanced Organic Chemistry*, a two-volume treatment designed for graduate students and advanced undergraduates. He was a member of the Committee of Examiners of the Graduate Record Examination in Chemistry from 1993–2000.

Frank and his wife Jill, who is a teacher/director of a preschool and a church organist, are the parents of Andy, Bob, and Bill and the grandparents of Riyad and Ava.

This edition is dedicated to all those teachers, students, colleagues, and friends who helped me in ways that none of us realized at the time.



Stereochemistry

Chapter Outline

7.1	MOLECULAR CHIRALITY: ENANTIOMERS	298
7.2	THE CHIRALITY CENTER	300
7.3	SYMMETRY IN ACHIRAL STRUCTURES	302
7.4	OPTICAL ACTIVITY	303
7.5	ABSOLUTE AND RELATIVE CONFIGURATION	305
7.6	THE CAHN-INGOLD-PRELOG <i>R-S</i> NOTATIONAL SYSTEM	306
7.7	FISCHER PROJECTIONS	308
7.8	PROPERTIES OF ENANTIOMERS	310
■	<i>Chiral Drugs</i>	311
7.9	REACTIONS THAT CREATE A CHIRALITY CENTER	313
7.10	CHIRAL MOLECULES WITH TWO CHIRALITY CENTERS	314
7.11	ACHIRAL MOLECULES WITH TWO CHIRALITY CENTERS	316
7.12	MOLECULES WITH MULTIPLE CHIRALITY CENTERS	320
■	<i>Chirality of Disubstituted Cyclohexanes</i>	320
7.13	REACTIONS THAT PRODUCE DIASTEREOMERS	322
7.14	RESOLUTION OF ENANTIOMERS	324

CHAPTER 7

7.15	STEREOREGULAR POLYMERS	327
7.16	CHIRALITY CENTERS OTHER THAN CARBON	328
7.17	SUMMARY	329
	PROBLEMS	332
	LEARNING BY MODELING	339

Stereochemistry is chemistry in three dimensions. Its foundations were laid by Jacobus van't Hoff* and Joseph Achille Le Bel in 1874. Van't Hoff and Le Bel independently proposed that the four bonds to carbon were directed toward the corners of a tetrahedron. One consequence of a tetrahedral arrangement of bonds to carbon is that two compounds may be different because the arrangement of their atoms in space is different. Isomers that have the same constitution but differ in the spatial arrangement of their atoms are called **stereoisomers**. We have already had considerable experience with certain types of stereoisomers—those involving cis and trans substitution patterns in alkenes and in cycloalkanes.

Our major objectives in this chapter are to develop a feeling for molecules as three-dimensional objects and to become familiar with stereochemical principles, terms, and notation. A full understanding of organic and biological chemistry requires an awareness of the spatial requirements for interactions between molecules; this chapter provides the basis for that understanding.

*Van't Hoff was the recipient of the first Nobel Prize in chemistry in 1901 for his work in chemical dynamics and osmotic pressure—two topics far removed from stereochemistry.

7.1 MOLECULAR CHIRALITY: ENANTIOMERS

Everything has a mirror image, but not all things are superimposable on their mirror images. Mirror-image superimposability characterizes many objects we use every day. Cups and saucers, forks and spoons, chairs and beds are all identical with their mirror images. Many other objects though—and this is the more interesting case—are not. Your left hand and your right hand, for example, are mirror images of each other but can't be made to coincide point for point, palm to palm, knuckle to knuckle, in three dimensions. In 1894, William Thomson (Lord Kelvin) coined a word for this property. He defined an object as **chiral** if it is not superimposable on its mirror image. Applying Thomson's term to chemistry, we say that a *molecule is chiral if its two mirror-image forms are not superimposable in three dimensions*. The word *chiral* is derived from the Greek word *cheir*, meaning "hand," and it is entirely appropriate to speak of the "handedness" of molecules. The opposite of chiral is **achiral**. A molecule that is superimposable on its mirror image is achiral.

In organic chemistry, chirality most often occurs in molecules that contain a carbon atom that is attached to four different groups. An example is bromochlorofluoromethane (BrClFCH).



Bromochlorofluoromethane

As shown in Figure 7.1, the two mirror images of bromochlorofluoromethane cannot be superimposed on each other. *Because the two mirror images of bromochlorofluoromethane are not superimposable, BrClFCH is chiral.*

The mirror images of bromochlorofluoromethane have the same constitution. That is, the atoms are connected in the same order. But they differ in the arrangement of their atoms in space; they are **stereoisomers**. Stereoisomers that are related as an object and its non-superimposable mirror image are classified as **enantiomers**. The word *enantiomer* describes a particular relationship between two objects. One cannot look at a single molecule in isolation and ask if it is an enantiomer any more than one can look at an individual human being and ask, "Is that person a cousin?" Furthermore, just as an object has one, and only one, mirror image, a chiral molecule can have one, and only one, enantiomer.


Notice in Figure 7.1c, where the two enantiomers of bromochlorofluoromethane are similarly oriented, that the difference between them corresponds to an interchange of the positions of bromine and chlorine. It will generally be true for species of the type $\text{C}(w, x, y, z)$, where w, x, y , and z are different atoms or groups, that an exchange of two of them converts a structure to its enantiomer, but an exchange of three returns the original structure, albeit in a different orientation.

Consider next a molecule such as chlorodifluoromethane (ClF_2CH), in which two of the atoms attached to carbon are the same. Figure 7.2 shows two molecular models of ClF_2CH drawn so as to be mirror images. As is evident from these drawings, it is a simple matter to merge the two models so that all the atoms match. *Because mirror image representations of chlorodifluoromethane are superimposable on each other, ClF_2CH is achiral.*


The surest test for chirality is a careful examination of mirror-image forms for superimposability. Working with models provides the best practice in dealing with molecules as three-dimensional objects and is strongly recommended.

Bromochlorofluoromethane is a known compound, and samples selectively enriched in each enantiomer have been described in the chemical literature. In 1989 two chemists at Polytechnic University (Brooklyn, New York) described a method for the preparation of BrClFCH that is predominantly one enantiomer.

(a) Structures A and B are mirror-image representations of bromochlorofluoromethane (BrClFCH).



(b) To test for superimposability, reorient B by turning it 180° .



(c) Compare A and B'. The two do not match. A and B' cannot be superimposed on each other. Bromochlorofluoromethane is therefore a chiral molecule. The two mirror-image forms are enantiomers of each other.






FIGURE 7.1 A molecule with four different groups attached to a single carbon is chiral. Its two mirror-image forms are not superimposable.



FIGURE 7.2 Mirror-image forms of chlorodifluoromethane are superimposable on each other. Chlorodifluoromethane is achiral.

7.2 THE CHIRALITY CENTER



As we've just seen, molecules of the general type



The 1996 IUPAC recommendations for stereochemical terms can be viewed at <http://www.chem.qmw.ac.uk/iupac/stereo>

are chiral when w , x , y , and z are different. In 1996, the IUPAC recommended that a tetrahedral carbon atom that bears four different atoms or groups be called a **chirality center**, which is the term that we will use. Several earlier terms, including *asymmetric center*, *asymmetric carbon*, *chiral center*, *stereogenic center*, and *stereocenter*, are still widely used.

Noting the presence of one (but not more than one) chirality center is a simple, rapid way to determine if a molecule is chiral. For example, C-2 is a chirality center in 2-butanol; it bears H, OH, CH₃, and CH₃CH₂ as its four different groups. By way of contrast, none of the carbon atoms bear four different groups in the achiral alcohol 2-propanol.





PROBLEM 7.1



Examine the following for chirality centers:

- (a) 2-Bromopentane (c) 1-Bromo-2-methylbutane
(b) 3-Bromopentane (d) 2-Bromo-2-methylbutane


SAMPLE SOLUTION A carbon with four different groups attached to it is a chirality center. (a) In 2-bromopentane, C-2 satisfies this requirement. (b) None of the carbons in 3-bromopentane has four different substituents, and so none of its atoms is a chirality center.




Molecules with chirality centers are very common, both as naturally occurring substances and as the products of chemical synthesis. (Carbons that are part of a double bond or a triple bond can't be chirality centers.)



A carbon atom in a ring can be a chirality center if it bears two different groups and the path traced around the ring from that carbon in one direction is different from that traced in the other. The carbon atom that bears the methyl group in 1,2-epoxypropane, for example, is a chirality center. The sequence of groups is $\text{O}-\text{CH}_2$ as one proceeds clockwise around the ring from that atom, but is $\text{H}_2\text{C}-\text{O}$ in the counterclockwise direction. Similarly, C-4 is a chirality center in limonene.



1,2-Epoxypropane
(product of epoxidation of propene)



Limonene
(a constituent of lemon oil)




Examine the molecular models of the two enantiomers of 1,2-epoxypropane on *Learning By Modeling* and test them for superimposability.

PROBLEM 7.2


Identify the chirality centers, if any, in

- (a) 2-Cyclopentenol and 3-cyclopentenol
(b) 1,1,2-Trimethylcyclobutane and 1,1,3-trimethylcyclobutane

SAMPLE SOLUTION (a) The hydroxyl-bearing carbon in 2-cyclopentenol is a chirality center. There is no chirality center in 3-cyclopentenol, because the sequence of atoms $1 \rightarrow 2 \rightarrow 3 \rightarrow 4 \rightarrow 5$ is equivalent regardless of whether one proceeds clockwise or counterclockwise.




2-Cyclopentenol



3-Cyclopentenol
(does not have a chirality center)

Even isotopes qualify as different substituents at a chirality center. The stereochemistry of biological oxidation of a derivative of ethane that is chiral because of deuterium ($\text{D} = {}^2\text{H}$) and tritium ($\text{T} = {}^3\text{H}$) atoms at carbon, has been studied and shown to proceed as follows:



The stereochemical relationship between the reactant and the product, revealed by the isotopic labeling, shows that oxygen becomes bonded to carbon on the same side from which H is lost. As you will see in this and the chapters to come, determining the three-dimensional aspects of a chemical or biochemical transformation can be a subtle, yet powerful, tool for increasing our understanding of how these reactions occur.

One final, very important point: *Everything we have said in this section concerns molecules that have one and only one chirality center; molecules with more than one chirality center may or may not be chiral.* Molecules that have more than one chirality center will be discussed in Sections 7.10 through 7.13.

7.3 SYMMETRY IN ACHIRAL STRUCTURES

Certain structural features can sometimes help us determine by inspection whether a molecule is chiral or achiral. For example, a molecule that has a *plane of symmetry* or a *center of symmetry* is superimposable on its mirror image and is achiral.


A **plane of symmetry** bisects a molecule so that one half of the molecule is the mirror image of the other half. The achiral molecule chlorodifluoromethane, for example, has the plane of symmetry shown in Figure 7.3.

PROBLEM 7.3


Locate any planes of symmetry in each of the following compounds. Which of the compounds are chiral? Which are achiral?

- (a) (*E*)-1,2-Dichloroethene (c) *cis*-1,2-Dichlorocyclopropane
(b) (*Z*)-1,2-Dichloroethene (d) *trans*-1,2-Dichlorocyclopropane

SAMPLE SOLUTION (a) (*E*)-1,2-Dichloroethene is planar. The molecular plane is a plane of symmetry. Identifying a plane of symmetry tells us the molecule is achiral.




A point in the center of a molecule is a **center of symmetry** if any line drawn from it to some element of the structure will, when extended an equal distance in the opposite direction, encounter an identical element. (*E*)-1,2-Dichloroethene, for example, has a center of symmetry in addition to the plane of symmetry described in Problem 7.3. Both carbons are equidistant from the center of the molecule and lie on a line (red) that passes through the center. Likewise, both hydrogens are equally distant from the center and lie on the blue line, and both chlorines are equally distant and lie on the green line.



Planes of symmetry are easier to identify and more common than centers of symmetry. Because either one is sufficient to make a molecule achiral, it makes sense to look first for a plane of symmetry. A molecule lacking a plane or center of symmetry is *likely* to be chiral, but the superimposability test should be applied to be certain.

Among the compounds given in Problem 7.3, only (*E*)-1,2-dichloroethene has a center of symmetry.

FIGURE 7.3 A plane of symmetry defined by the atoms H—C—Cl divides chlorodifluoromethane into two mirror-image halves.



7.4 OPTICAL ACTIVITY

The experimental facts that led van't Hoff and Le Bel to propose that molecules having the same constitution could differ in the arrangement of their atoms in space concerned the physical property of optical activity. **Optical activity** is the ability of a chiral substance to rotate the plane of **plane-polarized light** and is measured using an instrument called a **polarimeter**. (Figure 7.4).

The light used to measure optical activity has two properties: it consists of a single wavelength and it is plane-polarized. The wavelength used most often is 589 nm (called the *D line*), which corresponds to the yellow light produced by a sodium lamp. Except for giving off light of a single wavelength, a sodium lamp is like any other lamp in that its light is unpolarized, meaning that the plane of its electric field vector can have any orientation along the line of travel. A beam of unpolarized light is transformed to plane-polarized light by passing it through a polarizing filter, which removes all the waves except those that have their electric field vector in the same plane. This plane-polarized light now passes through the sample tube containing the substance to be examined, either in the liquid phase or as a solution in a suitable solvent (usually water, ethanol, or chloroform). The sample is "optically active" if it rotates the plane of polarized light. The direction and magnitude of rotation are measured using a second polarizing filter (the "analyzer") and cited as α , the observed rotation.

To be optically active, the sample must contain a chiral substance and one enantiomer must be present in excess of the other. A substance that does not rotate the plane of polarized light is said to be optically inactive. *All achiral substances are optically inactive.*

What causes optical rotation? The plane of polarization of a light wave undergoes a minute rotation when it encounters a chiral molecule. Enantiomeric forms of a chiral molecule cause a rotation of the plane of polarization in exactly equal amounts but in opposite directions. A solution containing equal quantities of enantiomers therefore exhibits no net rotation because all the tiny increments of clockwise rotation produced by molecules of one "handedness" are canceled by an equal number of increments of counterclockwise rotation produced by molecules of the opposite handedness.

Mixtures containing equal quantities of enantiomers are called **racemic mixtures**. Racemic mixtures are optically inactive. Conversely, when one enantiomer is present in excess, a net rotation of the plane of polarization is observed. At the limit, where all the

The phenomenon of optical activity was discovered by the French physicist Jean-Baptiste Biot in 1815.




FIGURE 7.4 The sodium lamp emits light moving in all planes. When the light passes through the first polarizing filter, only one plane emerges. The plane-polarized beam enters the sample compartment, which contains a solution enriched in one of the enantiomers of a chiral substance. The plane rotates as it passes through the solution. A second polarizing filter (called the analyzer) is attached to a movable ring calibrated in degrees that is used to measure the angle of rotation α . (Adapted from M. Silberberg, *Chemistry*, 3rd ed., McGraw-Hill Higher Education, New York, 2003, p. 618.)

molecules are of the same handedness, we say the substance is **optically pure**. Optical purity, or *percent enantiomeric excess*, is defined as:

$$\begin{aligned}\text{Optical purity} &= \text{percent enantiomeric excess} \\ &= \text{percent of one enantiomer} - \text{percent of other enantiomer}\end{aligned}$$

Thus, a material that is 50% optically pure contains 75% of one enantiomer and 25% of the other.

Rotation of the plane of polarized light in the clockwise sense is taken as positive (+), and rotation in the counterclockwise sense is taken as a negative (−) rotation. Older terms for positive and negative rotations were *dextrorotatory* and *levorotatory*, from the Latin prefixes *dextro*- (“to the right”) and *levo*- (“to the left”), respectively. At one time, the symbols *d* and *l* were used to distinguish between enantiomeric forms of a substance. Thus the dextrorotatory enantiomer of 2-butanol was called *d*-2-butanol, and the levorotatory form *l*-2-butanol; a racemic mixture of the two was referred to as *dl*-2-butanol. Current custom favors using algebraic signs instead, as in (+)-2-butanol, (−)-2-butanol, and (±)-2-butanol, respectively.

The observed rotation α of an optically pure substance depends on how many molecules the light beam encounters. A filled polarimeter tube twice the length of another produces twice the observed rotation, as does a solution twice as concentrated. To account for the effects of path length and concentration, chemists have defined the term **specific rotation**, given the symbol $[\alpha]$. Specific rotation is calculated from the observed rotation according to the expression

$$[\alpha] = \frac{100\alpha}{cl}$$

If concentration is expressed as grams per milliliter of solution instead of grams per 100 mL, an equivalent expression is

$$[\alpha] = \frac{\alpha}{cl}$$

where *c* is the concentration of the sample in grams per 100 mL of solution, and *l* is the length of the polarimeter tube in decimeters. (One decimeter is 10 cm.)

Specific rotation is a physical property of a substance, just as melting point, boiling point, density, and solubility are. For example, the lactic acid obtained from milk is exclusively a single enantiomer. We cite its specific rotation in the form $[\alpha]_D^{25} = +3.8^\circ$. The temperature in degrees Celsius and the wavelength of light at which the measurement was made are indicated as superscripts and subscripts, respectively.

PROBLEM 7.4

Cholesterol, when isolated from natural sources, is obtained as a single enantiomer. The observed rotation α of a 0.3-g sample of cholesterol in 15 mL of chloroform solution contained in a 10-cm polarimeter tube is -0.78° . Calculate the specific rotation of cholesterol.


PROBLEM 7.5

A sample of synthetic cholesterol was prepared consisting entirely of (+)-cholesterol. This synthetic (+)-cholesterol was mixed with some natural (−)-cholesterol. The mixture had a specific rotation $[\alpha]_D^{20}$ of -13° . What fraction of the mixture was (+)-cholesterol? (Note: You need to use the solution to Problem 7.4 for the specific rotation of (−)-cholesterol.)


It is convenient to distinguish between enantiomers by prefixing the sign of rotation to the name of the substance. For example, we refer to one of the enantiomers of 2-butanol as (+)-2-butanol and the other as (−)-2-butanol. Optically pure (+)-2-butanol has a specific rotation $[\alpha]_D^{27}$ of $+13.5^\circ$; optically pure (−)-2-butanol has an exactly opposite specific rotation $[\alpha]_D^{27}$ of -13.5° .

7.5 ABSOLUTE AND RELATIVE CONFIGURATION

The exact three-dimensional spatial arrangement of substituents at a chirality center is its **absolute configuration**. Neither the sign nor the magnitude of rotation by itself can tell us the absolute configuration of a substance. Thus, one of the following structures is (+)-2-butanol and the other is (–)-2-butanol, but without additional information we can't tell which is which.




Although no absolute configuration was known for any substance until the mid-twentieth century, organic chemists had experimentally determined the configurations of thousands of compounds relative to one another (their **relative configurations**) through chemical interconversion. To illustrate, consider (+)-3-buten-2-ol. Hydrogenation of this compound yields (+)-2-butanol.




Because hydrogenation of the double bond does not involve any of the bonds to the chirality center, the spatial arrangement of substituents in (+)-3-buten-2-ol must be the same as that of the substituents in (+)-2-butanol. The fact that these two compounds have the same sign of rotation when they have the same relative configuration is established by the hydrogenation experiment; it could not have been predicted in advance of the experiment.

Compounds that have the same relative configuration often have optical rotations of opposite sign. For example, treatment of (–)-2-methyl-1-butanol with hydrogen bromide converts it to (+)-1-bromo-2-methylbutane.



This reaction does not involve any of the bonds to the chirality center, and so both the starting alcohol (–) and the product bromide (+) have the same relative configuration.

An elaborate network connecting signs of rotation and relative configurations was developed that included the most important compounds of organic and biological chemistry. When, in 1951, the absolute configuration of a salt of (+)-tartaric acid was determined, the absolute configurations of all the compounds whose configurations had been related to (+)-tartaric acid stood revealed as well. Thus, returning to the pair of 2-butanol enantiomers that began this section, their absolute configurations are now known to be as shown.



In several places throughout the chapter we will use red and blue frames to call attention to structures that are enantiomeric.



Make a molecular model of one of the enantiomers of 3-buten-2-ol and the 2-butanol formed from it.




Make a molecular model of one of the enantiomers of 2-methyl-1-butanol and the 1-bromo-2-methylbutane formed from it.



PROBLEM 7.6

Does the molecular model shown represent (+)-2-butanol or (–)-2-butanol?




7.6 THE CAHN-INGOLD-PRELOG *R-S* NOTATIONAL SYSTEM

Just as it makes sense to have a nomenclature system by which we can specify the constitution of a molecule in words rather than pictures, so too is it helpful to have one that lets us describe stereochemistry. We have already had some experience with this idea when we distinguished between *E* and *Z* stereoisomers of alkenes.


In the *E-Z* system, substituents are ranked by atomic number according to a set of rules devised by R. S. Cahn, Sir Christopher Ingold, and Vladimir Prelog (Section 5.4). Actually, Cahn, Ingold, and Prelog first developed their ranking system to deal with the problem of the absolute configuration at a chirality center, and this is the system's major application. Table 7.1 shows how the Cahn-Ingold-Prelog system, called the **sequence rules**, is used to specify the absolute configuration at the chirality center in (+)-2-butanol.

As outlined in Table 7.1, (+)-2-butanol has the *S* configuration. Its mirror image is (–)-2-butanol, which has the *R* configuration.



(*S*)-2-Butanol

and



(*R*)-2-Butanol


Often, the *R* or *S* configuration and the sign of rotation are incorporated into the name of the compound, as in (*R*)-(–)-2-butanol and (*S*)-(+)-2-butanol.

The January 1994 issue of the *Journal of Chemical Education* contains an article that describes how to use your hands to assign *R* and *S* configurations.


Learning By Modeling includes an *R,S*-labeling feature. Make models of the structures shown in Problem 7.7, select **Configure** under the **Model** menu and click the **R,S** button.

PROBLEM 7.7


Assign absolute configurations as *R* or *S* to each of the following compounds:




(+)-2-Methyl-1-butanol



(+)-1-Fluoro-2-methylbutane



(+)-1-Bromo-2-methylbutane




(+)-3-Buten-2-ol

SAMPLE SOLUTION (a) The highest ranking substituent at the chirality center of 2-methyl-1-butanol is CH_2OH ; the lowest is H. Of the remaining two, ethyl outranks methyl.

Order of precedence: $\text{CH}_2\text{OH} > \text{CH}_3\text{CH}_2 > \text{CH}_3 > \text{H}$

The lowest ranking group (hydrogen) points away from us in the drawing. The three highest ranking groups trace a clockwise path from $\text{CH}_2\text{OH} \rightarrow \text{CH}_3\text{CH}_2 \rightarrow \text{CH}_3$.




This compound therefore has the *R* configuration. It is (*R*)-(+)-2-methyl-1-butanol.

TABLE 7.1 Absolute Configuration According to the Cahn-Ingold-Prelog Notational System

Step number

Example


Given that the absolute configuration of (+)-2-butanol is



(+)-2-Butanol


1. Identify the substituents at the chirality center, and rank them in order of decreasing precedence according to the system described in Section 5.4. Precedence is determined by atomic number, working outward from the point of attachment at the chirality center.

In order of decreasing precedence, the four substituents attached to the chirality center of 2-butanol are




2. Orient the molecule so that the lowest ranked atom or group points away from you.

As represented in the wedge-and-dash drawing at the top of this table, the molecule is already appropriately oriented. Hydrogen is the lowest ranked atom attached to the chirality center and points away from us.




3. Draw the three highest ranked substituents as they appear to you when the molecule is oriented so that the lowest ranked group points away from you.

The order of decreasing precedence is *counterclockwise*. The configuration at the chirality center is *S*.




Compounds in which a chirality center is part of a ring are handled in an analogous fashion. To determine, for example, whether the configuration of (+)-4-methylcyclohexene is *R* or *S*, treat the right- and left-hand paths around the ring as if they were independent groups.

Verify that (+)-4-methylcyclohexene is *R* using the **Configuration** feature on *Learning By Modeling*.



(+)-4-Methylcyclohexene




With the lowest ranked group (hydrogen) directed away from us, we see that the order of decreasing sequence rule precedence is *clockwise*. The absolute configuration is *R*.




PROBLEM 7.8

Draw three-dimensional representations or make molecular models of

(a) The *R* enantiomer of




(b) The *S* enantiomer of




SAMPLE SOLUTION


(a) The chirality center is the one that bears the bromine. In order of decreasing precedence, the substituents attached to the chirality center are



When the lowest ranked substituent (the methyl group) is away from us, the order of decreasing precedence of the remaining groups must appear in a clockwise sense in the *R* enantiomer.



which leads to the structure



(*R*)-2-Bromo-2-methylcyclohexanone

The Cahn–Ingold–Prelog system is the standard method of stereochemical notation. It replaced an older system based on analogies to specified reference compounds that used the prefixes *D* and *L*. As long as the substance in question is structurally similar to the reference, the *D,L* system was adequate. It is ambiguous, however, when applied to structures that are much different from the references, and as a *general* method for specifying configuration, the *D,L* system is largely obsolete. When *limited* to carbohydrates and amino acids though, *D,L* notation is firmly entrenched and works very well. We will use it when we get to Chapters 25–28, but won't need it until then.

7.7 FISCHER PROJECTIONS

Stereochemistry deals with the three-dimensional arrangement of a molecule's atoms, and we have attempted to show stereochemistry with wedge-and-dash drawings and computer-generated models. It is possible, however, to convey stereochemical





FIGURE 7.5 Ball-and-spoke models (*left*), wedge-and-dash drawings (*center*), and Fischer projections (*right*) of the *R* and *S* enantiomers of bromochlorofluoromethane.


information in an abbreviated form using a method devised by the German chemist Emil Fischer.

Let's return to bromochlorofluoromethane as a simple example of a chiral molecule. The two enantiomers of BrClFCH_3 are shown as ball-and-spoke models, as wedge-and-dash drawings, and as **Fischer projections** in Figure 7.5. Fischer projections are always generated the same way: the molecule is oriented so that the vertical bonds at the chirality center are directed away from you and the horizontal bonds point toward you. A projection of the bonds onto the page is a cross. The chirality center lies at the center of the cross but is not explicitly shown.


It is customary to orient the molecule so that the carbon chain is vertical with the lowest numbered carbon at the top as shown for the Fischer projection of (*R*)-2-butanol.



To verify that the Fischer projection has the *R* configuration at its chirality center, rotate the three-dimensional representation so that the lowest-ranked atom (H) points away from you. Be careful to maintain the proper stereochemical relationships during the operation.




With H pointing away from us, we can see that the order of decreasing precedence $\text{OH} > \text{CH}_2\text{CH}_3 > \text{CH}_3$ traces a clockwise path, verifying the configuration as *R*.



You may find it helpful to use a molecular model of (*R*)-2-butanol to guide you through this procedure.

PROBLEM 7.9

What is the absolute configuration (*R* or *S*) of the compound represented by the Fischer projection shown here?




As you work with Fischer projections, you may notice that some routine structural changes lead to predictable outcomes—outcomes that may reduce the number of manipulations you need to do to solve stereochemistry problems. Instead of listing these shortcuts, Problem 7.10 invites you to discover some of them for yourself.

PROBLEM 7.10

Using the Fischer projection of (*R*)-2-butanol given in the margin, explain how each of the following affects the configuration of the chirality center.

- Switching the positions of H and OH.
- Switching the positions of CH₃ and CH₂CH₃.
- Switching the positions of three groups.
- Switching H with OH, and CH₃ with CH₂CH₃.
- Rotating the Fischer projection 180° around an axis perpendicular to the page.

SAMPLE SOLUTION (a) Exchanging the positions of H and OH in the Fischer projection of (*R*)-2-butanol converts it to the mirror-image Fischer projection. The configuration of the chirality center goes from *R* to *S*.




Switching the positions of two groups in a Fischer projection reverses the configuration of the chirality center.

We mentioned in Section 7.6 that the D,L system of stereochemical notation, while outdated for most purposes, is still widely used for carbohydrates and amino acids. Likewise, Fischer projections find their major application in these same two families of compounds.


7.8 PROPERTIES OF ENANTIOMERS

The usual physical properties such as density, melting point, and boiling point are identical for both enantiomers of a chiral compound.

Enantiomers can have striking differences, however, in properties that depend on the arrangement of atoms in space. Take, for example, the enantiomeric forms of carvone. (*R*)-(-)-Carvone is the principal component of spearmint oil. Its enantiomer, (*S*)-(+)-carvone, is the principal component of caraway seed oil. The two enantiomers do not smell the same; each has its own characteristic odor.



(*R*)-(-)-Carvone
(from spearmint oil)




(*S*)-(+)-Carvone
(from caraway seed oil)


The difference in odor between (*R*)- and (*S*)-carvone results from their different behavior toward receptor sites in the nose. It is believed that volatile molecules occupy only those odor receptors that have the proper shape to accommodate them. Because the receptor sites are themselves chiral, one enantiomer may fit one kind of receptor while the other enantiomer fits a different kind. An analogy that can be drawn is to hands and gloves. Your left hand and your right hand are enantiomers. You can place your left hand into a left glove but not into a right one. The receptor (the glove) can accommodate one enantiomer of a chiral object (your hand) but not the other.

The term *chiral recognition* refers to a process in which some chiral receptor or reagent interacts selectively with one of the enantiomers of a chiral molecule. Very high levels of chiral recognition are common in biological processes. (-)-Nicotine, for example, is much more toxic than (+)-nicotine, and (+)-adrenaline is more active than (-)-adrenaline in constricting blood vessels. (-)-Thyroxine, an amino acid of the thyroid gland that speeds up metabolism, is one of the most widely used of all prescription drugs—about 10 million people in the United States take (-)-thyroxine on a daily basis. Its enantiomer, (+)-thyroxine has none of the metabolism-regulating effects, but was formerly given to heart patients to lower their cholesterol levels.


An article entitled "When Drug Molecules Look in the Mirror" in the June 1996 issue of the *Journal of Chemical Education* (pp. 481–484) describes numerous examples of common drugs in which the two enantiomers have different biological properties.



Nicotine



Adrenaline



Thyroxine

(Can you find the chirality center in each of these?)

Chiral Drugs



A recent estimate places the number of prescription and over-the-counter drugs marketed throughout the world at about 2000. Approximately one third of these are either naturally occurring substances themselves or are prepared by chemical modification of natural products. Most of the drugs derived from natural sources are chiral and are almost always obtained as a single enantiomer rather than as a racemic mixture. Not so with the over 500 chiral substances represented among the more than 1300 drugs that are

the products of synthetic organic chemistry. Until recently, such substances were, with few exceptions, prepared, sold, and administered as racemic mixtures even though the desired therapeutic activity resided in only one of the enantiomers. Spurred by a number of factors ranging from safety and efficacy to synthetic methodology and economics, this practice is undergoing rapid change as more and more chiral synthetic drugs become available in enantiomerically pure form.

—Cont.

(Continued)


Because of the high degree of chiral recognition inherent in most biological processes (Section 7.8), it is unlikely that both enantiomers of a chiral drug will exhibit the same level, or even the same kind, of effect. At one extreme, one enantiomer has the desired effect, and the other exhibits no biological activity at all. In this case, which is relatively rare, the racemic form is simply a drug that is 50% pure and contains 50% "inert ingredients." Real cases are more complicated. For example, the *S* enantiomer is responsible for the pain-relieving properties of ibuprofen, normally sold as a racemic mixture. The 50% of racemic ibuprofen that is the *R* enantiomer is not completely wasted, however, because enzyme-catalyzed reactions in our body convert much of it to active (*S*)-ibuprofen.



Ibuprofen

A much more serious drawback to using chiral drugs as racemic mixtures is illustrated by thalidomide, briefly employed as a sedative and anti-nausea drug in Europe during the period 1959–1962. The desired properties are those of (*R*)-thalidomide. (*S*)-Thalidomide, however, has a very different spectrum of biological activity and was shown to be

responsible for over 2000 cases of serious birth defects in children born to women who took it while pregnant.




Thalidomide


Basic research aimed at controlling the stereochemistry of chemical reactions has led to novel methods for the synthesis of chiral molecules in enantiomerically pure form. Aspects of this work were recognized with the award of the 2001 Nobel Prize in chemistry to William S. Knowles (Monsanto), Ryoji Noyori (Nagoya University), and K. Barry Sharpless (Scripps Research Institute). Most major pharmaceutical companies are examining their existing drugs to see which are the best candidates for synthesis as single enantiomers and, when preparing a new drug, design its synthesis so as to provide only the desired enantiomer. One incentive to developing enantiomerically pure versions of existing drugs is that the novel production methods they require may make them eligible for patent protection separate from that of the original drugs. Thus the temporary monopoly position that patent law views as essential to fostering innovation can be extended by transforming a successful chiral, but racemic, drug into an enantiomerically pure version.

**PROBLEM 7.11**

Both naproxen (antiinflammatory) and captopril (used to treat high-blood pressure) are synthetic drugs prepared and sold as single enantiomers. All chirality centers (one in naproxen and two in captopril) have the *S* configuration. Use *Spartan* to make molecular models of both and display their *R,S* labels to verify their configurations.




Naproxen



Captopril

7.9 REACTIONS THAT CREATE A CHIRALITY CENTER

Many of the reactions we've already encountered can yield a chiral product from an achiral starting material. Epoxidation of propene, for example, creates a chirality center by adding oxygen to the double bond.



In this, as in other reactions in which achiral reactants yield chiral products, the product is formed as a *racemic mixture* and is *optically inactive*. Remember, for a substance to be optically active, not only must it be chiral but one enantiomer must be present in excess of the other.

It is a general principle that *optically active products cannot be formed when optically inactive substrates react with optically inactive reagents*. This principle holds irrespective of whether the addition is syn or anti, concerted or stepwise. No matter how many steps are involved in a reaction, if the reactants are achiral, formation of one enantiomer is just as likely as the other, and a racemic mixture results.

Figure 7.6 shows why equal amounts of (*R*)- and (*S*)-1,2-epoxypropane are formed in the epoxidation of propene. There is no difference between the top face of the double bond and the bottom face. Peroxyacetic acid can transfer oxygen to either face with equal facility, the rates of formation of the *R* and *S* enantiomers of the product are the same, and the product is racemic.

In this example, addition to the double bond of an alkene converted an achiral molecule to a chiral one. The general term for a structural feature, the alteration of which





FIGURE 7.6 Epoxidation of propene produces equal amounts of (*R*)- and (*S*)-1,2-epoxypropane.

The *pro-* in *prochiral* means "before" or "in advance of" in roughly the same way as in "proactive."

introduces a chirality center in a molecule, is **prochiral**. A chirality center is introduced when the double bond of propene reacts with a peroxy acid. The double bond is a prochiral structural unit, and we speak of the top and bottom faces of the double bond as *prochiral faces*. Because attack at one prochiral face gives the enantiomer of the compound formed by attack at the other face, we classify the relationship between the two faces as **enantiotopic**.

In a second example, addition of hydrogen bromide converts 2-butene, which is achiral, to 2-bromobutane, which is chiral. But, as before, the product is racemic because both enantiomers are formed at equal rates. This is true regardless of whether the starting alkene is *cis*- or *trans*-2-butene or whether the mechanism is electrophilic addition or free-radical addition of HBr.




Whatever happens at one enantiotopic face of the double bond of *cis*- or *trans*-2-butene happens at the same rate at the other, resulting in a 1:1 mixture of (*R*)- and (*S*)-2-bromobutane.

PROBLEM 7.12


What two stereoisomeric alkanes are formed in the catalytic hydrogenation of (*E*)-3-methyl-2-hexene? What are the relative amounts of each?

Addition to double bonds is not the only kind of reaction that converts an achiral molecule to a chiral one. Other possibilities include substitution reactions such as the formation of 2-chlorobutane by free radical chlorination of butane. Here again, the product is chiral, but racemic.

1-Chlorobutane is also formed in this reaction.




We can view this reaction as the replacement of one or the other of the two methylene hydrogens at C-2 of butane. These hydrogens are *prochiral atoms* and, as the red and blue hydrogens in the Newman projection indicate, occupy mirror-image environments.



Replacing one of them by some different atom or group gives the enantiomer of the structure obtained by replacing the other; therefore, the methylene hydrogens at C-2 of butane are enantiotopic. The same is true for the hydrogens at C-3.


PROBLEM 7.13

Citric acid has three CO_2H groups. Which, if any, of them are enantiotopic?




Citric acid

When a reactant is chiral but optically inactive because it is *racemic*, any products derived from its reactions with optically inactive reagents will be *optically inactive*. For example, 2-butanol is chiral and may be converted with hydrogen bromide to 2-bromobutane, which is also chiral. If racemic 2-butanol is used, each enantiomer will react at the same rate with the achiral reagent. Whatever happens to (*R*)-(-)-2-butanol is mirrored in a corresponding reaction of (*S*)-(+)-2-butanol, and a racemic, optically inactive product results.




Optically inactive starting materials can give optically active products only if they are treated with an optically active reagent or if the reaction is catalyzed by an optically active substance. The best examples are found in biochemical processes. Most biochemical reactions are catalyzed by enzymes. Enzymes are chiral and enantiomerically homogeneous; they provide an asymmetric environment in which chemical reaction can take place. Ordinarily, enzyme-catalyzed reactions occur with such a high level of stereoselectivity that one enantiomer of a substance is formed exclusively even when the substrate is achiral. The enzyme *fumarase*, for example, catalyzes hydration of the double bond of fumaric acid to malic acid in apples and other fruits. Only the *S* enantiomer of malic acid is formed in this reaction.



The reaction is reversible, and its stereochemical requirements are so pronounced that neither the *cis* isomer of fumaric acid (maleic acid) nor the *R* enantiomer of malic acid can serve as a substrate for the fumarase-catalyzed hydration–dehydration equilibrium.

PROBLEM 7.14


Biological reduction of pyruvic acid, catalyzed by the enzyme lactate dehydrogenase, gives (+)-lactic acid, represented by the Fischer projection shown. What is the configuration of (+)-lactic acid according to the Cahn–Ingold–Prelog *R*–*S* notational system? Making a molecular model of the Fischer projection will help.



We'll continue with the three-dimensional details of chemical reactions later in this chapter. First though, we need to develop some additional stereochemical principles concerning structures with more than one chirality center.

7.10 CHIRAL MOLECULES WITH TWO CHIRALITY CENTERS

When a molecule contains two chirality centers, as does 2,3-dihydroxybutanoic acid, how many stereoisomers are possible?




2,3-Dihydroxybutanoic acid

We can use straightforward reasoning to come up with the answer. The absolute configuration at C-2 may be *R* or *S*. Likewise, C-3 may have either the *R* or the *S* configuration. The four possible combinations of these two chirality centers are

(*2R,3R*) (stereoisomer I) (*2S,3S*) (stereoisomer II)
 (*2R,3S*) (stereoisomer III) (*2S,3R*) (stereoisomer IV)

Figure 7.7 presents structural formulas for these four stereoisomers. Stereoisomers I and II are enantiomers of each other; the enantiomer of (*R,R*) is (*S,S*). Likewise stereoisomers III and IV are enantiomers of each other, the enantiomer of (*R,S*) being (*S,R*).

FIGURE 7.7 Stereoisomeric 2,3-dihydroxybutanoic acids. Stereoisomers I and II are enantiomers. Stereoisomers III and IV are enantiomers. All other relationships are diastereomeric (see text).






FIGURE 7.8 Representations of (2*R*,3*R*)-dihydroxybutanoic acid. (a) The staggered conformation is the most stable, but is not properly arranged to show stereochemistry as a Fischer projection. (b) Rotation about the C-2–C-3 bond gives the eclipsed conformation, and projection of the eclipsed conformation onto the page gives (c) a correct Fischer projection.


Stereoisomer I is not a mirror image of III or IV, so it is not an enantiomer of either one. Stereoisomers that are not related as an object and its mirror image are called diastereomers; **diastereomers** are stereoisomers that are not mirror images. Thus, stereoisomer I is a diastereomer of III and a diastereomer of IV. Similarly, II is a diastereomer of III and IV.

To convert a molecule with two chirality centers to its enantiomer, the configuration at *both* centers must be changed. Reversing the configuration at only one chirality center converts it to a diastereomeric structure.

Enantiomers must have equal and opposite specific rotations. Diastereomers can have different rotations, with respect to both sign and magnitude. Thus, as Figure 7.7 shows, the (2*R*,3*R*) and (2*S*,3*S*) enantiomers (I and II) have specific rotations that are equal in magnitude but opposite in sign. The (2*R*,3*S*) and (2*S*,3*R*) enantiomers (III and IV) likewise have specific rotations that are equal to each other but opposite in sign. The magnitudes of rotation of I and II are different, however, from those of their diastereomers III and IV.



In writing Fischer projections of molecules with two chirality centers, the molecule is arranged in an *eclipsed* conformation for projection onto the page, as shown in Figure 7.8. Again, horizontal lines in the projection represent bonds coming toward you; vertical bonds point away.

Organic chemists use an informal nomenclature system based on Fischer projections to distinguish between diastereomers. When the carbon chain is vertical and like substituents are on the same side of the Fischer projection, the molecule is described as the **erythro** diastereomer. When like substituents are on opposite sides of the Fischer projection, the molecule is described as the **threo** diastereomer. Thus, as seen in the Fischer projections of the stereoisomeric 2,3-dihydroxybutanoic acids, compounds I and II are erythro stereoisomers and III and IV are threo.



Erythro and threo describe the *relative configuration* (Section 7.5) of two chirality centers within a single molecule.

Because diastereomers are not mirror images of each other, they can have quite different physical and chemical properties. For example, the (2*R*,3*R*) stereoisomer of 3-amino-2-butanol is a liquid, but the (2*R*,3*S*) diastereomer is a crystalline solid.






PROBLEM 7.15

Draw Fischer projections or make molecular models of the four stereoisomeric 3-amino-2-butanols, and label each erythro or threo as appropriate.

PROBLEM 7.16


One other stereoisomer of 3-amino-2-butanol is a crystalline solid. Which one?

The situation is the same when the two chirality centers are present in a ring. There are four stereoisomeric 1-bromo-2-chlorocyclopropanes: a pair of enantiomers in which the halogens are trans and a pair in which they are cis. The cis compounds are diastereomers of the trans.




(1*R*,2*R*)-1-Bromo-2-chlorocyclopropane

Enantiomers




(1*S*,2*S*)-1-Bromo-2-chlorocyclopropane



(1*R*,2*S*)-1-Bromo-2-chlorocyclopropane

Enantiomers




(1*S*,2*R*)-1-Bromo-2-chlorocyclopropane

A good thing to remember is that the cis and trans isomers of a particular compound are diastereomers of each other.




7.11 ACHIRAL MOLECULES WITH TWO CHIRALITY CENTERS

Now think about a molecule, such as 2,3-butanediol, which has two chirality centers that are equivalently substituted.



Only *three*, not four, stereoisomeric 2,3-butanediols are possible. These three are shown in Figure 7.9. The (2*R*,3*R*) and (2*S*,3*S*) forms are enantiomers of each other and have equal and opposite optical rotations. A third combination of chirality centers, (2*R*,3*S*), however, gives an *achiral* structure that is superimposable on its (2*S*,3*R*) mirror image. Because it is achiral, this third stereoisomer is *optically inactive*. We call achiral molecules that have chirality centers **meso forms**. The meso form in Figure 7.9 is known as *meso*-2,3-butanediol.

One way to demonstrate that *meso*-2,3-butanediol is achiral is to recognize that its eclipsed conformation has a plane of symmetry that passes through and is perpendicular to the C-2-C-3 bond, as illustrated in Figure 7.10*a*. The anti conformation is achiral as well. As Figure 7.10*b* shows, this conformation is characterized by a center of symmetry at the midpoint of the C-2-C-3 bond.

(1*R*,2*R*)-1,2-Dibromocyclopropane(1*S*,2*S*)-1,2-Dibromocyclopropane*cis*-1,2-Dibromocyclopropane**PROBLEM 7.18**

One of the stereoisomers of 1,3-dimethylcyclohexane is a meso form. Which one?

7.12 MOLECULES WITH MULTIPLE CHIRALITY CENTERS


Many naturally occurring compounds contain several chirality centers. By an analysis similar to that described for the case of two chirality centers, it can be shown that the maximum number of stereoisomers for a particular constitution is 2^n , where n is equal to the number of chirality centers.

PROBLEM 7.19

Using *R* and *S* descriptors, write all the possible combinations for a molecule with three chirality centers.

When two or more of a molecule's chirality centers are equivalently substituted, meso forms are possible, and the number of stereoisomers is then less than 2^n . Thus, 2^n represents the *maximum* number of stereoisomers for a molecule containing n chirality centers.

The best examples of substances with multiple chirality centers are the *carbohydrates*. One class of carbohydrates, called *aldohexoses*, has the constitution




An aldohexose

Because there are four chirality centers and no possibility of meso forms, there are 2^4 or 16, stereoisomeric aldohexoses. All 16 are known, having been isolated either as natural products or as the products of chemical synthesis.

PROBLEM 7.20

A second category of six-carbon carbohydrates, called *keto*hexoses, has the constitution shown. How many stereoisomeric 2-ketohexoses are possible?



A 2-ketohexose


Steroids are another class of natural products with multiple chirality centers. One such compound is *cholic acid*, which can be obtained from bile. Its structural formula

Chirality of Disubstituted Cyclohexanes

Disubstituted cyclohexanes present us with a challenging exercise in stereochemistry. Consider the seven possible dichlorocyclohexanes: 1,1-; *cis*- and *trans*-1,2-; *cis*- and *trans*-1,3-; and *cis*- and *trans*-1,4-. Which are chiral? Which are achiral?


Four isomers—the ones that are achiral because they have a plane of symmetry—are relatively easy to identify:

ACHIRAL DICHLOROCYCLOHEXANES




The remaining three isomers are chiral:

CHIRAL DICHLOROCYCLOHEXANES



Among all the isomers, *cis*-1,2-dichlorocyclohexane is unique in that the ring-flipping process typical of cyclohexane derivatives converts it to its enantiomer.



Structures A and A' are nonsuperimposable mirror images of each other. Thus although *cis*-1,2-dichlorocyclohexane is chiral, it is optically inactive when chair–chair interconversion occurs. Such interconversion is rapid at room temperature and converts optically active A to a racemic mixture of A and A'. Because A and A' are enantiomers interconvertible by a conformational change, they are sometimes referred to as **conformational enantiomers**.

The same kind of spontaneous racemization occurs for any *cis*-1,2 disubstituted cyclohexane in which both substituents are the same. Because such compounds are chiral, it is incorrect to speak of them as meso compounds, which are achiral molecules that have chirality centers. Rapid chair–chair interconversion, however, converts them to a 1:1 mixture of enantiomers, and this mixture is optically inactive.

is given in Figure 7.11. Cholic acid has 11 chirality centers, and so a total (including cholic acid) of 2^{11} , or 2048, stereoisomers have this constitution. Of these 2048 stereoisomers, how many are diastereomers of cholic acid? Remember! Diastereomers are stereoisomers that are not enantiomers, and any object can have only one mirror image. Therefore, of the 2048 stereoisomers, one is cholic acid, one is its enantiomer, and the other 2046 are diastereomers of cholic acid. Only a small fraction of these compounds are known, and (+)-cholic acid is the only one ever isolated from natural sources.

Eleven chirality centers may seem like a lot, but it is nowhere close to a world record. It is a modest number when compared with the more than 100 chirality centers typical for most small proteins and the thousands of chirality centers present in nucleic acids.




FIGURE 7.12 Anti addition of Br_2 to (E) -2-butene gives *meso*-2,3-dibromobutane.

Two factors combine to determine which stereoisomers are actually formed in the reaction.

1. The (E) - or (Z) -configuration of the starting alkene
2. The anti stereochemistry of addition (Section 6.16)

Figures 7.12 and 7.13 depict the stereochemical relationships associated with anti addition of bromine to (E) - and (Z) -2-butene, respectively. The trans alkene (E) -2-butene yields only *meso*-2,3-dibromobutane, but the cis alkene (Z) -2-butene gives a racemic mixture of $(2R,3R)$ - and $(2S,3S)$ -2,3-dibromobutane.

Bromine addition to alkenes is a stereospecific reaction. A **stereospecific reaction** is one in which stereoisomeric starting materials yield products that are stereoisomers of each other. In this case the starting materials, in separate reactions, are the E and Z stereoisomers of 2-butene. The chiral dibromides from (Z) -2-butene are stereoisomers (diastereomers) of the *meso* dibromide formed from (E) -2-butene.

Notice further that, consistent with the principle developed in Section 7.9, optically inactive starting materials (achiral alkenes and bromine) yield optically inactive products (a racemic mixture or a *meso* structure) in these reactions.

PROBLEM 7.21

Epoxidation of alkenes is a stereospecific syn addition. Which stereoisomer of 2-butene reacts with peroxyacetic acid to give *meso*-2,3-epoxybutane? Which one gives a racemic mixture of $(2R,3R)$ - and $(2S,3S)$ -2,3-epoxybutane?





FIGURE 7.13 Anti addition of Br_2 to (Z) -2-butene gives a racemic mixture of $(2R,3R)$ - and $(2S,3S)$ -2,3-dibromobutane.

A reaction that introduces a second chirality center into a starting material that already has one need not produce equal quantities of two possible diastereomers. Consider catalytic hydrogenation of 2-methyl(methylene)cyclohexane. As you might expect, both *cis*- and *trans*-1,2-dimethylcyclohexane are formed.

Make molecular models of the reactant and both products shown in the equation.



The relative amounts of the two products, however, are not equal; more *cis*-1,2-dimethylcyclohexane is formed than *trans*-. The reason for this is that it is the less hindered face of the double bond that approaches the catalyst surface and is the face to which hydrogen is transferred. Hydrogenation of 2-methyl(methylene)cyclohexane occurs preferentially at the side of the double bond opposite that of the methyl group and leads to a faster rate of formation of the *cis* stereoisomer of the product.

PROBLEM 7.22

Could the fact that hydrogenation of 2-methyl(methylene)cyclohexane gives more *cis*-1,2-dimethylcyclohexane than *trans*- be explained on the basis of the relative stabilities of the two stereoisomeric products?

The double bond in 2-methyl(methylene)cyclohexane is prochiral. The two faces, however, are not enantiotopic as they were for the alkenes we discussed in Section 7.9. In those earlier examples, when addition to the double bond created a new chirality center, attack at one face gave one enantiomer; attack at the other gave the other enantiomer. In the case of 2-methyl(methylene)cyclohexane, which already has one chirality center, attack at opposite faces of the double bond gives two products that are diastereomers of each other. Prochiral faces of this type are called **diastereotopic**.

The hydrogenation of 2-methyl(methylene)cyclohexane is an example of a *stereoselective reaction*, meaning one in which stereoisomeric products are formed in unequal amounts from a single starting material (Section 5.11).


A common misconception is that a stereospecific reaction is simply one that is 100% stereoselective. The two terms are not synonymous, however. A stereospecific reaction is one which, when carried out with stereoisomeric starting materials, gives a product from one reactant that is a stereoisomer of the product from the other. A stereoselective reaction is one in which a single starting material gives a predominance of a single stereoisomer when two or more are possible. *Stereospecific* is more closely connected with features of the reaction than with the reactant. Thus terms such as *syn addition* and *anti elimination* describe the stereospecificity of reactions. *Stereoselective* is more closely connected with structural effects in the reactant as expressed in terms such as *addition to the less hindered side*. A stereospecific reaction can also be stereoselective. For example, *syn addition* describes stereospecificity in the catalytic hydrogenation of alkenes, whereas the preference for addition to the less hindered face of the double bond describes stereoselectivity.

Note that the terms *regioselective* and *regiospecific*, however, are defined in terms of each other. A *regiospecific* reaction is one that is 100% *regioselective*.

7.14 RESOLUTION OF ENANTIOMERS

The separation of a racemic mixture into its enantiomeric components is termed **resolution**. The first resolution, that of tartaric acid, was carried out by Louis Pasteur in

1848. Tartaric acid is a byproduct of wine making and is almost always found as its dextrorotatory 2*R*,3*R* stereoisomer, shown here in a perspective drawing and in a Fischer projection.



(2*R*,3*R*)-Tartaric acid (mp 170°C, $[\alpha]_D +12^\circ$)

PROBLEM 7.23

There are two other stereoisomeric tartaric acids. Write their Fischer projections, and specify the configuration at their chirality centers.

Occasionally, an optically inactive sample of tartaric acid was obtained. Pasteur noticed that the sodium ammonium salt of optically inactive tartaric acid was a mixture of two mirror-image crystal forms. With microscope and tweezers, Pasteur carefully separated the two. He found that one kind of crystal (in aqueous solution) was dextrorotatory, whereas the mirror-image crystals rotated the plane of polarized light an equal amount but were levorotatory.

Although Pasteur was unable to provide a structural explanation—that had to wait for van't Hoff and Le Bel a quarter of a century later—he correctly deduced that the enantiomeric quality of the crystals was the result of enantiomeric molecules. The rare form of tartaric acid was optically inactive because it contained equal amounts of (+)-tartaric acid and (−)-tartaric acid. It had earlier been called *racemic acid* (from Latin *racemus*, meaning “a bunch of grapes”), a name that subsequently gave rise to our present term for an equal mixture of enantiomers.

PROBLEM 7.24

Could the unusual, optically inactive form of tartaric acid studied by Pasteur have been meso-tartaric acid?

Pasteur's technique of separating enantiomers not only is laborious but requires that the crystals of the enantiomers be distinguishable. This happens very rarely. Consequently, alternative and more general approaches for resolving enantiomers have been developed. Most are based on a strategy of temporarily converting the enantiomers of a racemic mixture to diastereomeric derivatives, separating these diastereomers, then regenerating the enantiomeric starting materials.

Figure 7.14 illustrates this strategy. Say we have a mixture of enantiomers, which, for simplicity, we label as C(+) and C(−). Assume that C(+) and C(−) bear some functional group that can combine with a reagent P to yield adducts C(+)-P and C(−)-P. Now, if reagent P is chiral, and if only a single enantiomer of P, say, P(+), is added to a racemic mixture of C(+) and C(−), as shown in the first step of Figure 7.14, then the products of the reaction are C(+)-P(+) and C(−)-P(+). These products are not mirror images; they are diastereomers. Diastereomers can have different physical properties, which can serve as a means of separating them. The mixture of diastereomers is separated, usually by recrystallization from a suitable solvent. In the last step, an appropriate chemical transformation liberates the enantiomers and restores the resolving agent.

Whenever possible, the chemical reactions involved in the formation of diastereomers and their conversion to separate enantiomers are simple acid–base reactions.

A description of Pasteur's work, as part of a broader discussion concerning crystal structure, can be found in the article “Molecules, Crystals, and Chirality” in the July 1997 issue of the *Journal of Chemical Education*, pp. 800–806.






FIGURE 7.14 The general procedure for resolving a chiral substance into its enantiomers. Reaction with a single enantiomer of a chiral resolving agent $P(+)$ converts the racemic mixture of enantiomers $C(+)$ and $C(-)$ to a mixture of diastereomers $C(+)-P(+)$ and $C(-)-P(+)$. The mixture of diastereomers is separated—by fractional crystallization, for example. A chemical reaction is then carried out to convert diastereomer $C(+)-P(+)$ to $C(+)$ and the resolving agent $P(+)$. Likewise, diastereomer $C(-)-P(+)$ is converted to $C(-)$ and $P(+)$. $C(+)$ has been separated from $C(-)$, and the resolving agent $P(+)$ can be recovered for further use.

For example, naturally occurring (S) - $(-)$ -malic acid is often used to resolve amines. One such amine that has been resolved in this way is 1-phenylethylamine. Amines are bases, and malic acid is an acid. Proton transfer from (S) - $(-)$ -malic acid to a racemic mixture of (R) - and (S) -1-phenylethylamine gives a mixture of diastereomeric salts.



The diastereomeric salts are separated and the individual enantiomers of the amine liberated by treatment with a base:




PROBLEM 7.25

In the resolution of 1-phenylethylamine using (–)-malic acid, the compound obtained by recrystallization of the mixture of diastereomeric salts is (*R*)-1-phenylethylammonium (*S*)-malate. The other component of the mixture is more soluble and remains in solution. What is the configuration of the more soluble salt?

This method is widely used for the resolution of chiral amines and carboxylic acids. Analogous methods based on the formation and separation of diastereomers have been developed for other functional groups; the precise approach depends on the kind of chemical reactivity associated with the functional groups present in the molecule.

As the experimental tools for biochemical transformations have become more powerful and procedures for carrying out these transformations in the laboratory more routine, the application of biochemical processes to mainstream organic chemical tasks including the production of enantiomerically pure chiral molecules has grown.

One approach, called **enzymatic resolution**, involves treating a racemic mixture with an enzyme that catalyzes the reaction of only one of the enantiomers. Some of the most commonly used ones are *lipases* and *esterases*, enzymes that catalyze the hydrolysis of esters. In a typical procedure, one enantiomer of the acetate ester of a racemic alcohol undergoes hydrolysis and the other is left unchanged when hydrolyzed in the presence of an esterase from hog liver.




The November 2002 issue of the *Journal of Chemical Education* (pp. 1351–1352) describes the resolution of an alcohol by lipase-catalyzed ester hydrolysis in an experiment suitable for introductory laboratories.

High yields of the enantiomerically pure alcohol and enantiomerically pure ester are regularly achieved. The growing interest in chiral drugs (see the boxed essay on this topic, p. 311) has stimulated the development of large-scale enzymatic resolution as a commercial process.

7.15 STEREOREGULAR POLYMERS

Before the development of the Ziegler–Natta catalyst systems (Section 6.22), polymerization of propene was not a reaction of much value. The reason for this has a stereochemical basis. Consider a section of *polypropylene*:



Representing the polymer chain in an extended zigzag conformation, as shown in Figure 7.15, reveals several distinct structural possibilities differing with respect to the relative configurations of the carbons that bear the methyl groups.

One structure, represented in Figure 7.15*a*, has all the methyl groups oriented in the same direction with respect to the polymer chain. This stereochemical arrangement is called **isotactic**. Another form called **syndiotactic**, shown in Figure 7.15*b*, has its methyl groups alternating front and back along the chain. Both isotactic and syndiotactic polypropylene are known as **stereoregular polymers** because each is characterized by a precise stereochemistry at the carbon atom that bears the methyl group. A third

TABLE 7.2 Classification of Isomers



Definition	Example
Isomers are different compounds that have the same molecular formula. They may be either constitutional isomers or stereoisomers.	
1. <i>Constitutional isomers</i> are isomers that differ in the order in which their atoms are connected.	Three constitutionally isomeric compounds have the molecular formula C_3H_8O : $CH_3CH_2CH_2OH$ 1-Propanol $CH_3CH(OH)CH_3$ 2-Propanol $CH_3CH_2OCH_3$ Ethyl methyl ether
2. <i>Stereoisomers</i> are isomers that have the same constitution but differ in the arrangement of their atoms in space.	
(a) <i>Enantiomers</i> are stereoisomers that are related as an object and its nonsuperimposable mirror image.	The two enantiomeric forms of 2-chlorobutane are  (R)-(-)-2-Chlorobutane and  (S)-(+)-2-Chlorobutane
(b) <i>Diastereomers</i> are stereoisomers that are not mirror images.	The <i>cis</i> and <i>trans</i> isomers of 4-methylcyclohexanol are stereoisomers, but they are not related as an object and its mirror image; they are diastereomers.  cis-4-Methylcyclohexanol  trans-4-Methylcyclohexanol

Table 7.2 shows the enantiomers of 2-chlorobutane. C-2 is a chirality center in 2-chlorobutane.

Section 7.3 A molecule that has a plane of symmetry or a center of symmetry is achiral. *cis*-4-Methylcyclohexanol (Table 7.2) has a plane of symmetry that bisects the molecule into two mirror-image halves and is achiral. The same can be said for *trans*-4-methylcyclohexanol.

Section 7.4 **Optical activity**, or the degree to which a substance rotates the plane of polarized light, is a physical property used to characterize chiral substances. Enantiomers have equal and opposite **optical rotations**. To be optically active a substance must be chiral, and one enantiomer must be present in excess of the other. A **racemic mixture** is optically inactive and contains equal quantities of enantiomers.


Section 7.5 **Relative configuration** compares the arrangement of atoms in space to some reference. The prefix *cis* in *cis*-4-methylcyclohexanol, for example, describes relative configuration by referencing the orientation of the CH_3 group to the OH. **Absolute configuration** is an exact description of the arrangement of atoms in space.

Section 7.6

Absolute configuration in chiral molecules is best specified using the prefixes *R* and *S* of the Cahn–Ingold–Prelog notational system. Substituents at a chirality center are ranked in order of decreasing precedence. If the three highest ranked substituents trace a clockwise path (highest→second highest→third highest) when the lowest ranked substituent is held away from you, the configuration is *R*. If the path is counterclockwise, the configuration is *S*. Table 7.2 shows the *R* and *S* enantiomers of 2-chlorobutane.

Section 7.7

A **Fischer projection** shows how a molecule would look if its bonds were projected onto a flat surface. Horizontal lines represent bonds coming toward you; vertical bonds point away from you. The projection is normally drawn so that the carbon chain is vertical, with the lowest numbered carbon at the top.




Section 7.8

Both enantiomers of the same substance are identical in most of their physical properties. The most prominent differences are biological ones, such as taste and odor, in which the substance interacts with a chiral receptor site in a living system. Enantiomers also have important consequences in medicine, in which the two enantiomeric forms of a drug can have much different effects on a patient.


Section 7.9

A chemical reaction can convert an achiral substance to a chiral one. If the product contains a single chirality center, it is formed as a racemic mixture. Optically active products can be formed from optically inactive starting materials only if some optically active agent is present. The best examples are biological processes in which enzymes catalyze the formation of only a single enantiomer.






Section 7.10

When a molecule has two chirality centers and these two chirality centers are not equivalent, four stereoisomers are possible.



Stereoisomers that are not mirror images are classified as **diastereomers**. Each enantiomer of *erythro*-3-bromo-2-butanol is a diastereomer of each enantiomer of *threo*-3-bromo-2-butanol.

Section 7.11 Achiral molecules that contain chirality centers are called **meso forms**. Meso forms typically contain (but are not limited to) two equivalently substituted chirality centers. They are optically inactive.

*meso*-2,3-Dibromobutane(2*R*,3*R*)-2,3-Dibromobutane(2*S*,3*S*)-2,3-Dibromobutane

Section 7.12 For a particular constitution, the maximum number of stereoisomers is 2^n , where n is the number of structural units capable of stereochemical variation—usually this is the number of chirality centers, but can include *E* and *Z* double bonds as well. The number of stereoisomers is reduced to less than 2^n when there are meso forms.

Section 7.13 Addition reactions of alkenes may generate one (Section 7.9) or two (Section 7.13) chirality centers. When two chirality centers are produced, their relative stereochemistry depends on the configuration (*E* or *Z*) of the alkene and whether the addition is syn or anti.

Section 7.14 **Resolution** is the separation of a racemic mixture into its enantiomers. It is normally carried out by converting the mixture of enantiomers to a mixture of diastereomers, separating the diastereomers, then regenerating the enantiomers.

Section 7.15 Certain polymers such as polypropylene contain chirality centers, and the relative configurations of these centers affect the physical properties of the polymers. Like substituents appear on the same side of a zigzag carbon chain in an **isotactic** polymer, alternate along the chain in a **syndiotactic** polymer, and appear in a random manner in an **atactic** polymer. Isotactic and syndiotactic polymers are referred to as **stereoregular** polymers.

Section 7.16 Atoms other than carbon can be chirality centers. Examples include those based on tetracoordinate silicon and tricoordinate sulfur as the chirality center. In principle, tricoordinate nitrogen can be a chirality center in compounds of the type $\text{N}(x, y, z)$, where x , y , and z are different, but inversion of the nitrogen pyramid is so fast that racemization occurs virtually instantly at room temperature.



PROBLEMS


7.26 Which of the isomeric alcohols having the molecular formula $\text{C}_5\text{H}_{12}\text{O}$ are chiral? Which are achiral?



7.27 Write structural formulas or make molecular models for all the compounds that are trichloro derivatives of cyclopropane. (Don't forget to include stereoisomers.) Which are chiral? Which are achiral?


7.28 In each of the following pairs of compounds one is chiral and the other is achiral. Identify each compound as chiral or achiral, as appropriate.





7.29 Compare 2,3-pentanediol and 2,4-pentanediol with respect to the number of stereoisomers possible for each constitution. Which stereoisomers are chiral? Which are achiral?

7.30 In 1996, it was determined that the absolute configuration of (–)-bromochlorofluoromethane is *R*. Which of the following is (are) (–)-BrClFCH?




7.31 Specify the configuration of the chirality center as *R* or *S* in each of the following.

(a) (–)-2-Octanol




(b) Monosodium L-glutamate (only this stereoisomer is of any value as a flavor-enhancing agent)



7.32 A subrule of the Cahn–Ingold–Prelog system specifies that higher mass number takes precedence over lower when distinguishing between isotopes.












(a) Determine the absolute configurations of the reactant and product in the biological oxidation of isotopically labeled ethane described in Section 7.2.














(b) Because OH becomes bonded to carbon at the same side from which H is lost, the oxidation proceeds with retention of configuration (Section 6.14). Compare this fact with the *R* and *S* configurations you determined in part (a) and reconcile any *apparent* conflicts.




7.33 Identify the relationship in each of the following pairs. Do the drawings represent constitutional isomers or stereoisomers, or are they just different ways of drawing the same compound? If they are stereoisomers, are they enantiomers or diastereomers? (Molecular models may prove useful in this problem.)








7.34 Chemical degradation of chlorophyll gives a number of substances including *phytol*. The constitution of phytol is shown. How many stereoisomers have this constitution?




7.35 *Muscarine* is a poisonous substance present in the mushroom *Amanita muscaria*. Its structure is represented by the constitution shown here.




- Including muscarine, how many stereoisomers have this constitution?
- One of the substituents on the ring of muscarine is *trans* to the other two. How many of the stereoisomers satisfy this requirement?
- Muscarine has the configuration *2S,3R,5S*. Write a structural formula or build a molecular model of muscarine showing its correct stereochemistry.

7.36 *Ectocarpene* is a volatile, sperm cell-attracting material released by the eggs of the seaweed *Ectocarpus siliculosus*. Its constitution is




All the double bonds are *cis*, and the absolute configuration of the chirality center is *S*. Write a stereochemically accurate representation of ectocarpene.

7.37 *Multifidene* is a sperm cell-attracting substance released by the female of a species of brown algae (*Cutleria multifida*). The constitution of multifidene is



- How many stereoisomers are represented by this constitution?
- Multifidene has a *cis* relationship between its alkenyl substituents. Given this information, how many stereoisomers are possible?
- The butenyl side chain has the *Z* configuration of its double bond. On the basis of all the data, how many stereoisomers are possible?
- Draw stereochemically accurate representations of all the stereoisomers that satisfy the structural requirements of multifidene.
- How are these stereoisomeric multifidenes related (enantiomers or diastereomers)?

7.38 *Streptimidone* is an antibiotic and has the structure shown. How many diastereomers of streptimidone are possible? How many enantiomers? Using the *E,Z* and *R,S* descriptors, specify all essential elements of stereochemistry of streptimidone.




7.39 In Problem 4.26 you were asked to draw the preferred conformation of menthol on the basis of the information that menthol is the most stable stereoisomer of 2-isopropyl-5-methylcyclohexanol. We can now completely describe (–)-menthol structurally by noting that it has the *R* configuration at the hydroxyl-substituted carbon.




- (a) Draw or construct a molecular model of the preferred conformation of (–)-menthol.
 (b) (+)-Isomenthol has the same constitution as (–)-menthol. The configurations at C-1 and C-2 of (+)-isomenthol are the opposite of the corresponding chirality centers of (–)-menthol. Write the preferred conformation of (+)-isomenthol.



7.40 A certain natural product having $[\alpha]_D + 40.3^\circ$ was isolated. Two structures have been independently proposed for this compound. Which one do you think is more likely to be correct? Why?



7.41 One of the principal substances obtained from archaea (one of the oldest forms of life on Earth) is derived from a 40-carbon diol. Given the fact that this diol is optically active, is it compound A or is it compound B?




- 7.42 (a) An aqueous solution containing 10 g of optically pure fructose was diluted to 500 mL with water and placed in a polarimeter tube 20 cm long. The measured rotation was -5.20° . Calculate the specific rotation of fructose.
 (b) If this solution were mixed with 500 mL of a solution containing 5 g of racemic fructose, what would be the specific rotation of the resulting fructose mixture? What would be its optical purity?
- 7.43 Each of the following reactions gives a mixture of two stereoisomers. Write their structures. Are they enantiomers or diastereomers? Are they chiral or achiral? Are they formed in equal amounts?




7.44 Write the organic products of each of the following reactions. If two stereoisomers are formed, show both. Label all chirality centers *R* or *S* as appropriate.

- 1-Butene and hydrogen iodide
- (*E*)-2-Pentene and bromine in carbon tetrachloride
- (*Z*)-2-Pentene and bromine in carbon tetrachloride
- 1-Butene and peroxyacetic acid in dichloromethane
- (*Z*)-2-Pentene and peroxyacetic acid in dichloromethane
- 1,5,5-Trimethylcyclopentene and hydrogen in the presence of platinum
- 1,5,5-Trimethylcyclopentene and diborane in tetrahydrofuran followed by oxidation with hydrogen peroxide


7.45 The enzyme *aconitase* catalyzes the hydration of aconitic acid to two products: citric acid and isocitric acid. Isocitric acid is optically active; citric acid is not. What are the respective constitutions of citric acid and isocitric acid?




7.46 Consider the ozonolysis of *trans*-4,5-dimethylcyclohexene having the configuration shown.



Structures A, B, and C are three stereoisomeric forms of the reaction product.



- Which, if any, of the compounds A, B, and C are chiral?
 - What product is formed in the reaction?
 - What product would be formed if the methyl groups were *cis* to each other in the starting alkene?
- 7.47 (a) On being heated with potassium ethoxide in ethanol (70°C), the deuterium-labeled alkyl bromide shown gave a mixture of 1-butene, *cis*-2-butene, and *trans*-2-butene. On the basis of your knowledge of the E2 mechanism, predict which alkene(s), if any, contained deuterium.



- (b) The bromide shown in part (a) is the erythro diastereomer. How would the deuterium content of the alkenes formed by dehydrohalogenation of the threo diastereomer differ from those produced in part (a)?

7.48 A compound (C_6H_{10}) contains a five-membered ring. When Br_2 adds to it, two diastereomeric dibromides are formed. Suggest reasonable structures for the compound and the two dibromides.

7.49 When optically pure 2,3-dimethyl-2-pentanol was subjected to dehydration, a mixture of two alkenes was obtained. Hydrogenation of this alkene mixture gave 2,3-dimethylpentane, which was 50% optically pure. What were the two alkenes formed in the elimination reaction, and what were the relative amounts of each?

7.50 When (*R*)-3-buten-2-ol is treated with a peroxy acid, two stereoisomeric epoxides are formed in a 60:40 ratio. The minor stereoisomer has the structure shown.



- Write the structure of the major stereoisomer.
- What is the relationship between the two epoxides? Are they enantiomers or diastereomers?
- What four stereoisomeric products are formed when racemic 3-buten-2-ol is epoxidized under the same conditions? How much of each stereoisomer is formed?



LEARNING BY MODELING

Spartan 7.1 *Spartan* can help you not only visualize molecules as three-dimensional objects, but also with *R,S* notation. The configuration of all the chirality centers in a molecule can be “turned on” at once by clicking **R,S** in the **Configure** menu under **Model**.

- Make a model of 2-bromobutane and select **R,S** in **Configure**. Which enantiomer is it?
- Select —Br in the atom library and double click on one of the C-2 hydrogens to convert 2-bromobutane to 2,2-dibromobutane. What happens to the *R,S* label?
- Delete the original Br. What label appears?
- Replace one of the hydrogens at C-3 by Br. What are the chirality labels?


Spartan 7.2 Make molecular models of each of the following and check your answer by selecting **R,S** from the **Configure** menu under **Model**.

- (*R*)-1-Fluoro-2-butanol
- (*S*)-3-Chlorocyclopentene

Spartan 7.3 Make molecular models of all of the stereoisomeric 1,2,3-trichlorocyclopropanes. How many are there? Identify each structure as chiral or achiral. What are the configurations at the chirality centers? Check your assignment using the **R,S** labels feature of the **Configure** menu.


Spartan 7.4 Make models of the three stereoisomers of 2,3-dichlorobutane and use the **Configure** feature of *Spartan* to ensure that one is *2R,3R*, another is *2S,3S*, and the third is *2R,3S*. **Minimize** each. Which two are of the same energy? Which one is different?

Spartan 7.5 Is the 2,3-dibromobutane stereoisomer shown the product of anti addition of Br_2 to *cis*-2-butene or to *trans*-2-butene? You can determine the answer by carrying out the following operations.




- (1) Make a model of the compound as shown.
- (2) Adjust the $\text{Br}-\text{C}-\text{C}-\text{Br}$ dihedral angle so that the Br atoms are anti.
- (3) **Delete** both bromines.
- (4) Select **Make bond** and connect the bonds from which the bromine atoms were deleted.
- (5) **Minimize** and the correct stereoisomer of 2-butene appears.

Spartan 7.6 Modify the procedure in the preceding problem to figure out if the starting alkene in the reaction shown has the *E* or *Z* configuration. Assume syn addition occurs.



Spartan 7.7 Molecular models of L-serine and L-cysteine, represented by the Fischer projections shown, are given in the **Database**. What are their configurations according to the Cahn-Ingold-Prelog convention? Is there a correlation between D,L and R,S notation?



Spartan 7.8 *Fischer projections*: Making molecular models of Fischer projections requires adjusting dihedral angles so as to produce eclipsed conformations. We can illustrate one way to do this by making a model of the Fischer projection of *meso*-2,3-dibromobutane.

Step 1: Starting with cyclobutane, make *cis*-1,2-dibromocyclobutane.


Step 2: Break the $\text{H}_2\text{C}-\text{CH}_2$ bond and **Minimize**.

Step 3: Set the $\text{Br}-\text{C}-\text{C}-\text{Br}$ dihedral angle to 0° .

Step 4: Rotate the molecule so that the carbon chain is vertical and the horizontal bonds at C-2 and C-3 point toward you. Both bromines should point toward you and be on the same side of the carbon chain as in the Fischer projection of the *meso* form.

- (a) What is the Cahn-Ingold-Prelog designation of each chirality center?
- (b) Change the $\text{Br}-\text{C}-\text{C}-\text{Br}$ dihedral angle from 0° to 180° . What conformation results? Are the CH_3 groups *gauche* or *anti*? Does changing the conformation change the *R* or *S* designation of either chirality center?
- (c) How could you modify this procedure so as to produce one of the chiral diastereomers of 2,3-dibromobutane (*R,R* or *S,S*)?

Spartan 7.9 We are accustomed to seeing the carbohydrate D-ribose written as a Fischer projection. Assign *R,S* configurations to the chirality centers in D-ribose. Check your answers against the model of D-ribose given in the **Database**. (Although the structure in the **Database** shows D-ribose in a staggered conformation, *conformation* does not affect the *configuration* of chirality centers.)



D-Ribose

Spartan 7.10 Locate planes of symmetry or a center of symmetry in each of the following. Which ones have only planes of symmetry? Which have both a plane and a center of symmetry? Which have only a center of symmetry? What are their calculated dipole moments? What can you say about symmetry elements and the presence or absence of a dipole moment?

- (a) *cis*-1,2-Dichloroethene
- (b) *trans*-1,2-Dichloroethene
- (c) 1,1,2-Trichloroethene
- (d) Tetrachloroethene
- (e) 1,1,1-Trichloroethane
- (f) Hexachloroethane