

Declaration of Rachel J. Watters on Authentication of Publication

I, Rachel J. Watters, am a librarian, and the Head of Resource Sharing for the University of Wisconsin-Madison General Library System, Memorial Library, located at 728 State Street, Madison, Wisconsin, 53706. I have worked as a librarian at the University of Wisconsin-Madison library system since 1998, starting as a graduate student employee in Wisconsin TechSearch (“WTS”), an interlibrary loan department at the Kurt F. Wendt Engineering Library. In 1999, I began work as a professional librarian in the Interlibrary Loan department at UW-Madison’s Memorial Library. In 2002, I returned to WTS as a professional librarian and became WTS Director in 2011. In 2019, I became of Head of Resource Sharing for UW-Madison’s General Library System at Memorial Library and part of my current job responsibilities include oversight of WTS, now housed in Memorial Library. I have a master’s degree in Library and Information Studies from the University of Wisconsin-Madison. Through the course of my studies and employment, I have become well informed about the operations of the University of Wisconsin library system, which follows standard library practices.

This Declaration relates to the dates of receipt and availability of the following:

van Balken, M.R., Vandoninck, V., Gisolf, K.W.H., Vergunst, H., Kiemeney, L.A.L.M., Debruyne, F.M.J., and Bemelmans, B.L.H. (2001). Posterior tibial nerve stimulation as neuromodulative treatment of lower urinary tract dysfunction. *Journal of Urology*, 166(3), 914-918.

Declaration of Rachel J. Watters on Authentication of Publication

Standard operating procedures for materials at the University of Wisconsin-Madison Libraries. When an issue was received by the Library, it would be checked in, stamped with the date of receipt, added to library holdings records, and made available to readers as soon after its arrival as possible. The procedure normally took a few days or at most 2 to 3 weeks.

Exhibit A to this Declaration is a true and accurate copy of the journal issue cover with library date stamp of *Journal of Urology* (2001), from the University of Wisconsin-Madison Library collection. Exhibit A also includes an excerpt of pages 914 to 918 of that issue, showing the article entitled *Posterior tibial nerve stimulation as neuromodulative treatment of lower urinary tract dysfunction* (2001). Based on this information, the date stamp on the journal issue cover indicates *Posterior tibial nerve stimulation as neuromodulative treatment of lower urinary tract dysfunction* (2001) was received by Weston Library, University of Wisconsin-Madison Libraries on August 16, 2001.

Based on the information in Exhibit A, it is clear that the volume was received by the library on or before August 16, 2001, catalogued and available to library patrons within a few days or at most 2 to 3 weeks after August 16, 2001.

Members of the interested public could locate the *Journal of Urology* (2001) publication after it was cataloged by searching the public library catalog or requesting a search through WTS. The search could be done by title and/or subject key words.

Declaration of Rachel J. Watters on Authentication of Publication

Members of the interested public could access the publication by locating it on the library's shelves or requesting it from WTS.

I declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code.

Date: September 10, 2021

Memorial Library
728 State Street
Madison, Wisconsin 53706



Rachel J. Watters
Head of Resource Sharing

LTL


Volume 166

September 2001

Number 3

The Journal of UROLOGY®

Official Journal of the American Urological Association, Inc.®
Founded In 1917 By Hugh Hampton Young



WESTON LIBRARY

AUG 16 2001

J5/120 CLINICAL SCIENCE CENTER
600 HIGHLAND AVE MADISON WI 53792



Annual Meeting, American Urological Association, Inc.®,
Orlando, Florida, May 25-30, 2002


LIPPINCOTT
WILLIAMS & WILKINS
A Wolters Kluwer Company

The Journal of UROLOGY®

Editor

Jay Y. Gillenwater
1120 North Charles Street
Baltimore, Maryland 21201

Associate Editors

Jean B. deKernion
Los Angeles, California

Edward J. McGuire
Ann Arbor, Michigan

Martin I. Resnick
Cleveland, Ohio

Section Editors

Jack S. Elder
Cleveland, Ohio

Anthony Atala
Boston, Massachusetts

Ralph V. Clayman
St. Louis, Missouri

Editorial Board

Mid-Atlantic

Burkhardt H. Zorn
Washington, D. C.

New England

Kevin R. Loughlin
Boston, Massachusetts

New York

Carl A. Olsson
New York, New York

North Central

Robert C. Flanigan
Maywood, Illinois

Drogo K. Montague
Cleveland, Ohio

Northeastern

William C. Hulbert, Jr.
Rochester, New York

South Central

H. Barton Grossman
Houston, Texas

Southeastern

Joseph A. Smith, Jr.
Nashville, Tennessee

Jerry W. Sullivan
New Orleans, Louisiana

Donald E. Novicki
Scottsdale, Arizona

Western

Brent W. Snow
Salt Lake City, Utah

Board of Consultants

David G. Bostwick
Rochester, Minnesota

Lance K. Heilbrun
Detroit, Michigan

Charles Y. C. Pak
Dallas, Texas

Michael B. Cohen
Iowa City, Iowa

Michael W. Kattan
New York, New York

Derek Raghavan
Los Angeles, California

Fred Dorey
Los Angeles, California

Steven A. Leibel
New York, New York

Arthur T. Rosenfield
New Haven, Connecticut

Robert Elashoff
Los Angeles, California

Seth P. Lerner
Houston, Texas

Deborah S. Smith
New Haven, Connecticut

Jonathan Epstein
Baltimore, Maryland

Ronald R. Townsend
Denver, Colorado

Former Editors

Hugh H. Young
1917-1945

J. A. Campbell Colston
1946-1965

Hugh J. Jewett
1966-1977

William W. Scott
1977-1983

Herbert Brendler
1983-1985

John T. Grayhack
1985-1994

The Journal of Urology® (ISSN 0022-5347) is the Official Journal of the American Urological Association Education and Research, Inc.®, and is published monthly by Lippincott Williams & Wilkins, 12107 Insurance Way, Hagerstown, MD 21740. Business offices are located at 227 East Washington Square, Philadelphia, PA 19106-3780. Periodicals postage paid at Hagerstown, MD, and at additional mailing offices. Subscription rates individual \$434.00 (\$545.00 foreign); institutions \$479.00 (\$589.00 foreign); in-training \$243.00 (\$337.00 foreign); US Site License \$621.00 (\$731 Foreign Site License) add \$35.00 for air freight delivery outside of North America. All subscription rates include access to *The Journal of Urology*® Online (jurology.com); single copy \$55.00. Foreign prices exclude Japan (See Information for Subscribers). Printed in USA. Subscription prices subject to change. The GST number for Canadian subscribers is 89552 4239 RT. To order call 1-800-638-3030 from anywhere in the U.S. Outside North America, call 301-714-2300 or fax 301-824-7390. POSTMASTER: Send address changes to: Lippincott Williams & Wilkins, 12107 Insurance Way, Hagerstown, MD 21740. Indexed by *Index Medicus* (abridged), *Current Contents*, *Clinical Medicine*, *Life Sciences*, *Science Citation Index*, *Research Alert*, *ISI/BioMed*, *EMBASE/Excerpta Medica*, *BIOSIS*, and *Reference Update*. On-line text is available through Ovid Technologies, Inc. and directly from the publisher at jurology.com. Printed on acid-free paper. Copyright © 2001 by American Urological Association Education and Research, Inc.®

The Journal of UROLOGY®

Volume 166

September 2001

Number 3

CLINICAL UROLOGY

This Month in Clinical Urology. <i>J. Y. Gillenwater</i>	799
This Month in Investigative Urology. <i>A. Atala</i>	801
Review Article	
Genitourinary Complications of Sickle Cell Disease. <i>D. Bruno, D. R. Wigfall, S. A. Zimmerman, P. M. Rosoff and J. S. Wiener</i>	803
Original Articles	
Lymphatic Mapping and Detection of Sentinel Nodes in Patients With Bladder Cancer. <i>A. Sherif, M. de la Torre, P.-U. Malmström and M. Thörn</i>	812
Incisional Hernia and Its Repair With Polypropylene Mesh in Renal Transplant Recipients. <i>E. Mazzucchi, W. C. Nahas, I. Antonopoulos, L. E. Ianhez and S. Arap</i>	816
Eight-Year Experience With Transperitoneal Laparoscopic Adrenal Surgery. <i>G. Guazzoni, A. Cestari, F. Montorsi, R. Lanzi, L. Nava, A. Centemero and P. Rigatti</i>	820
Clinical Role of F18-Fluorodeoxyglucose Positron Emission Tomography for Detection and Management of Renal Cell Carcinoma. <i>S. Ramdave, G. W. Thomas, S. U. Berlangieri, D. M. Bolton, I. Davis, H. Tochon-Danguy, D. MacGregor and A. M. Scott</i>	825
Complex Perioperative Immuno-Dysfunction in Patients With Renal Cell Carcinoma. <i>M. Böhm, A. Ittenson, C. Philipp, F.-W. Röhl, S. Ansorge and E. P. Allhoff</i>	831
Sexuality Preserving Cystectomy and Neobladder: Initial Results. <i>S. Horenblas, W. Meinhardt, W. Ijzerman and L. F. M. Moonen</i>	837
Telomerase Activity and Cytokeratin 20 as Markers for Detection and Followup of Transitional Cell Carcinoma: Unfulfilled Promise. <i>A. Cassel, M. A. Rahat, N. Lahat, N. Lindendorf, Y. Mecz and A. Stein</i> (Editorial Comment by M. J. Droller)	841
Value of Single Biopsy With 12 Transperineal Cores for Detecting Prostate Cancer in Patients With Elevated Prostate Specific Antigen. <i>P. Emiliozzi, S. Longhi, P. Scarpone, A. Pansadoro, F. DePaula and V. Pansadoro</i> (Editorial Comment by M. O. Koch)	845
Prospective Detection of Clinically Relevant Prostate Cancer in Prostate Specific Antigen Range 1 to 3 Ng./Ml. Combined With Free-to-Total Ratio 20% or Less: Aarau Experience. <i>F. Recker, M. K. Kwiatkowski, A. Huber, B. Stamm, K. Lehmann and R. Tscholl</i> (Editorial Comment by A. R. Zlotta)	851
Safety and Morbidity of First and Repeat Transrectal Ultrasound Guided Prostate Needle Biopsies: Results of Prospective European Prostate Cancer Detection Study. <i>B. Djavan, M. Waldert, A. Zlotta, P. Dobronski, C. Seitz, M. Remzi, A. Borkowski, C. Schulman and M. Marberger</i>	856
Interval After Prostate Specific Antigen Testing and Subsequent Risk of Incurable Prostate Cancer. <i>S.-L. Yao and G. Lu-Yao</i>	861
Repeat Biopsy Strategy in Patients With Atypical Small Acinar Proliferation or High Grade Prostatic Intraepithelial Neoplasia on Initial Prostate Needle Biopsy. <i>P. G. Borboroglu, R. L. Sur, J. L. Roberts and C. L. Amling</i>	866
Expression of Neuropeptides in Hyperplastic and Malignant Prostate Tissue and Its Possible Clinical Implications. <i>D.-S. Yu, D.-S. Hsieh, H.-I. Chen and S.-Y. Chang</i>	871
High Dose Radiation Delivered by Intensity Modulated Conformal Radiotherapy Improves Outcome of Localized Prostate Cancer. <i>M. J. Zelefsky, Z. Fuks, M. Hunt, H. J. Lee, D. Lombardi, C. C. Ling, V. E. Reuter, E. S. Venkatraman and S. A. Leibel</i> (Editorial Comment by A. L. Zeitman)	876
Safety and Efficacy of Exisulind for Treatment of Recurrent Prostate Cancer After Radical Prostatectomy. <i>E. T. Goluboff, D. Prager, D. Rukstalis, B. Giantonio, M. Madorsky, I. Barken, I. B. Weinstein, A. W. Partin, C. A. Olsson and UCLA Oncology Research Network</i>	882
Glanuloplasty With Scrotal Flap for Partial Penectomy. <i>O. N. Mazza and G. M. J. Cheliz</i>	887

Front cover: Location of metastatic sentinel nodes in patients with bladder cancer (page 812).

**PLEASE NOTE THE CME ENROLLMENT FORM AND QUESTIONNAIRE
PRINTED ON PAGES 1066 AND 1067 IN THIS ISSUE**

Contents continued on page A10

Optimal Prevention and Management of Proximal Ureteral Stent Migration and Remigration. <i>R. H. Breau and R. W. Norman</i> (Editorial Comment by J. W. Segura)	890
Periprosthetic Nerve Blockade for Transrectal Ultrasound Guided Biopsy of Prostate: Randomized, Double-Blind, Placebo Controlled Study. <i>G. Pareek, N. A. Armenakas and J. A. Fracchia</i>	894
Nonrefluxing, Serous Lined Extramural Tunnel for Ureteroileal Anastomosis in Ileal Conduit Urinary Diversion: First Clinical Experience in 10 Patients. <i>K. Türkölmez, S. Baltaci, Y. Bedük, C. Göğüş and O. Göğüş</i>	898
Influence of Stent Size on Success of Antegrade Endopyelotomy for Primary Ureteropelvic Junction Obstruction: Results of 2 Consecutive Series. <i>H. Danuser, W. W. Hochreiter, D. K. Ackermann and U. E. Studer</i> (Editorial Comment by M. S. Pearle)	902
Videourodynamic Studies in Men With Lower Urinary Tract Symptoms: Comparison of Community Based Versus Referral Urological Practices. <i>F. Fusco, A. Groutz, J. G. Blaivas, D. C. Chaikin and J. P. Weiss</i> (Editorial Comment by S. Madersbacher)	910
Posterior Tibial Nerve Stimulation as Neuromodulative Treatment of Lower Urinary Tract Dysfunction. <i>M. R. van Balken, V. Vandoninck, K. W. H. Gisolf, H. Vergunst, L. A. L. M. Kiemeney, F. M. J. Debruyne and B. L. H. Bemelmans</i>	914
Expansion and Bioabsorption of Self-Reinforced Lactic and Glycolic Acid Copolymer Prostatic Spiral Stent. <i>S. Laaksovirta, M. Talja, T. Välimaa, T. Isotalo, P. Törmälä and T. L. J. Tammela</i>	919
Improving Accuracy of Vascular Testing in Impotent Men: Correction Hemodynamic Alterations Using Vasoactive Medication Re-Dosing Schedule. <i>J. P. Mulhall, A. Abdel-Moneim, R. Abobakr and I. Goldstein</i>	923
Long-Term Efficacy of Sildenafil and Tachyphylaxis Effect. <i>R. El-Galley, H. Rutland, R. Talic, T. Keane and H. Clark</i>	927
Mechanical Reliability and Safety of, and Patient Satisfaction With Ambicor Inflatable Penile Prosthesis: Results of 2 Center Study. <i>L. A. Levine, C. R. Estrada and A. Morgentaler</i>	932
Radical Cystectomy is Safe in Elderly Patients at High Risk. <i>S. S. Chang, G. Alberts, M. S. Cookson and J. A. Smith, Jr.</i> (Editorial Comment by D. P. Wood)	938
Predictors of Utilities for Health States in Early Stage Prostate Cancer. <i>C. S. Saigal, J. Gornbein, R. Nease and M. S. Litwin</i>	942
Quality of Life After Treatment for Localized Prostate Cancer: Differences Based on Treatment Modality. <i>J. W. Davis, D. A. Kuban, D. F. Lynch and P. F. Schellhammer</i>	947
Patient Reported Complications After Prostate Brachytherapy. <i>B. H. Han, K. C. Demel, K. Wallner, W. Ellis, L. Young and K. Russell</i> (Editorial Comment by M. J. Barry)	953
Assessment of Early Continence Recovery After Radical Prostatectomy: Patient Reported Symptoms and Impairment. <i>S. Kielb, R. L. Dunn, M. G. Rashid, S. Murray, M. G. Sanda, J. E. Montie and J. T. Wei</i>	958
Editorial: Quality of Life. <i>J. Adolfsson</i>	962
Inguinal Hernia After Radical Retropubic Prostatectomy for Prostate Cancer: Study of Incidence and Risk Factors in Comparison to No Operation and Lymphadenectomy. <i>P. Lodding, C. Bergdahl, M. Nyberg, E. Pileblad, J. Stranne and J. Hugosson</i> (Editorial Comment by P. N. Schlegel)	964
Risk Adjustment for Evaluating Outcome of Urological Operative Procedures. <i>F. Sommer, A. Ehsan, H.-P. Caspers, T. Klotz and U. Engelmann</i>	968
Urologists at Work	
Reconstruction of Ureter With Transverse Retubularized Colon Segments. <i>B. Ubrig, M. Waldner and S. Roth</i>	973
Glutaraldehyde Cross-Linked Collagen Occlusion of Ureteral Orifices With Percutaneous Nephrostomy: Minimally Invasive Option for Treatment Refractory Hemorrhagic Cystitis. <i>C. M. Gonzalez, J. R. Case and R. B. Nadler</i>	977
Case Reports	
Ureteral Quadruplication With Contralateral Triplicate Ureter. <i>R. I. Lopes, R. N. Lopes and C. M. Barbosa Filho</i>	979
Vesical Calculus Caused by Vascular Prosthetic Graft. <i>J. Bashir, G. Gilling-Smith and A. D. Desmond</i>	981
Iatrogenic Pneumocephalus After Laparoscopic Renal Biopsy. <i>V. S. Ricchiuti and A. D. Seftel</i>	982
Rupture of Adrenal Carcinoma After Biopsy. <i>A. H. Kardar</i>	984
Local Recurrence of Oncocytic Adrenocortical Carcinoma With Ovary Metastasis. <i>R. Kurek, R. Von Knobloch, U. Feek, A. Heidenreich and R. Hofmann</i>	985
Complete Genitourinary Exenteration for Multifocal Transitional Cell Carcinoma in Renal Transplant Recipient. <i>D. T. Glassman and G. N. Sklar</i>	986
Subtotal Radical Cystectomy After Radical Prostatectomy: Source of Pelvic Recurrence of Bladder Cancer With Life Threatening Bleeding. <i>T. H. S. Hsu and E. A. Klein</i>	988
Squamous Cell Carcinoma of Phallus in Patient With Congenital Adrenal Hyperplasia. <i>R. J. Dimitriou and C. L. Coogan</i>	990

Hemangiopericytoma of Seminal Vesicle Presenting With Hypoglycemia. <i>M. Arya, D. Hayne, R. S. D. Brown, P. J. O'Donnell and A. R. Mundy</i>	992
* Undetectable Prostate Specific Antigen in Disseminated Prostate Cancer. <i>P. Beardo, P. L. Fernández, J. M. Corral, X. Filella and J. Alcover</i>	993
✓ Prostatic Sarcoma With Rapid Tumor Progression After Nerve Sparing Radical Cystoprostatectomy. <i>M. D. Young, P. Dahm and C. N. Robertson</i>	994
✓ Local Tract Metastasis of Prostatic Adenocarcinoma 8 Years After ¹²⁵ Iodine Brachytherapy. <i>W. J. Wiersinga, T. M. de Reijke and L. E. C. M. Blank</i>	995
Arteriovesical Fistula: Complication of Bladder Trauma. <i>O. Shah, M. Ficazzola, P. Torre and S. Al-Askari</i>	996
Intrarenal Varix Presenting as Enhancing Renal Mass With Calcifications. <i>A. R. Deibler, S. N. Nadig, N. Curry, N. K. Bissada and G. W. Hull</i>	997
Transrectal Ultrasound Findings of Pelvic Arteriovenous Malformation. <i>T. Ishii, T. Higashionna and Y. Hiratsuka</i>	999
Plasma Cell Granuloma of Testis: Unusual Localization. <i>P. K. Aksoy, B. H. Özdemir, C. Aygün and M. Agildere</i>	1000
Prostatosymphysal Fistula After Transurethral Resection of Prostate. <i>R. Gillitzer, S. W. Melchior, J. Jones, J. Fichtner and J. W. Thüroff</i>	1001
Tension-Free Vaginal Tape: Is it Truly Tension-Free? <i>J. M. Choe</i>	1003
Reoperation After Complicated Tension-Free Vaginal Tape Procedures. <i>M. Wyczolkowski, W. Klima and Z. Piasecki</i>	1004
Recurrent Bulbar Urethral Stricture in Region of Artificial Urinary Sphincter. <i>M. DeBell and H. Wessells</i>	1006
Female Urethral Hemangioma. <i>K. Uchida, F. Fukuta, M. Ando and M. Miyake</i>	1008

Letters to the Editor

Re: Randomized Prospective Trial of Intrarectal Lidocaine for Pain Control During Transrectal Prostate Biopsy: Emory University Experience. <i>E. Stewart, U. Otite and V. Nargund</i>	1009
Re: Percutaneous Sacral Nerve Root Neuromodulation for Intractable Interstitial Cystitis. <i>T. C. Chai</i>	1009
Re: Protracted Urinary Retention Necessitating Urethrolisis Following Tension-Free Vaginal Tape Surgery. <i>Ö. Öge</i>	1009
Re: Development of Erectile Dysfunction in Men Treated for Prostate Cancer. <i>M. S. Anscher</i>	1010
Re: Is Human Chorionic Gonadotropin Useful for Identifying and Treating Nonpalpable Testis? <i>S. G. Docimo</i>	1010
Re: Third Year Medical Student Attitudes Toward Learning Urology. <i>W. J. Huang</i>	1011

PEDIATRIC UROLOGY

Histological Analysis of Urethral Healing After Tubularized Incised Plate Urethroplasty. <i>J. F. Lopes, A. Schned, P. I. Ellsworth and M. Cendron</i> (Editorial Comment by W. T. Snodgrass)	1015
Prepubertal High Flow Priapism: Incidence, Diagnosis and Treatment. <i>B. G. Volkmer, T. Nessler, S. C. Kraemer, J. Goerich, S. Basche and H.-W. Gottfried</i>	1019
Magnetic Resonance Imaging for Evaluation of Hydronephrosis, Reflux and Renal Scarring in Children. <i>L. V. Rodríguez, D. Spielman, R. J. Herfkens and L. D. Shortliffe</i>	1024
Concealed Umbilical Stoma: Long-Term Evaluation of Stomal Stenosis. <i>D. T. Glassman and S. G. Docimo</i> (Editorial Comment by M. Cain)	1029
Outcome of Ureteral Reimplantation in Children With Urge Syndrome. <i>U. Barroso, Jr., R. Jednak, J. S. Barthold and R. González</i>	1032
Outcome Analysis of Vesicoamniotic Shunting in Comprehensive Population. <i>G. McLorie, W. Farhat, A. Khoury, D. Geary and G. Ryan</i> (Editorial Comment by A. L. Freedman)	1037
Duodenal Atresia Associated With Midgut Deletion in Cloacal Exstrophy: New Association? <i>M. McHoney, P. Ransley, P. Cuckow, E. Kiely and L. Spitz</i>	1042

Papers Presented at Fall 2000 Meeting of Section on Urology, American Academy of Pediatrics

Extraperitoneal Placement of Renal Allografts in Children Weighing Less Than 15 Kg. <i>P. D. Furness, III, J. B. Houston, S. A. Grampsas, F. M. Karrer, C. F. Firlit and M. A. Koyle</i> (Editorial Comment by J. M. Barry)	1043
Urological Complications After Renal Transplantation Using Ureteroureteral Anastomosis in Children. <i>S. P. Lapointe, M. Charbit, D. Jan, S. Lortat-Jacob, J.-L. Michel, D. Beurton, M.-F. Gagnadoux, P. Niaudet, M. Broyer and Y. Révillon</i>	1047
Management of Grade IV Renal Injury in Children. <i>R. S. Russell, A. Gomelsky, D. R. McMahon, D. Andrews and P. F. Nasrallah</i>	1050
Severe Hypospadias Repair With Meatal Based Paracoronal Skin Flap: Modified Koyanagi Repair. <i>Y. Sugita, S. Tanikaze, K. Yoshino and F. Yamamichi</i> (Editorial Comment by J. Wacksman)	1052
Sonographic Characteristics of Urinary Tract in Healthy Neonates. <i>Y. Yamazaki, R. Yago and H. Toma</i>	1055

Measurement of Urinary Flow Rate Using Ultrasound in Young Boys and Infants. <i>K. P. Wolffenbuttel, D. J. Kok, R. van Mastrigt, E. van den Berg and R. J. M. Nijman</i>	1059
Can Standard Open Pediatric Urological Procedures be Performed on Outpatient Basis? <i>J. K. Sprunger, C. T. Reese and R. M. Decter</i>	1063
CME Enrollment Form and Questionnaire	1066

INVESTIGATIVE UROLOGY

Novel Expression Patterns of Myc/Max/Mad Transcription Factor Network in Developing Murine Prostate Gland. <i>Q. Luo, E. Harmon, B. G. Timms and L. Kretzner</i>	1071
Influence of Sex Hormones on Renal Osteopontin Expression and Urinary Constituents in Experimental Urolithiasis. <i>T. Yagisawa, F. Ito, Y. Osaka, H. Amano, C. Kobayashi and H. Toma</i> ..	1078
Effects of Isoproterenol and Butylscopolamine on Friction Between Artificial Stone and Intraureteral Wall in Anesthetized Rabbits. <i>R. Miyatake, Y. Tomiyama, M. Murakami, Y.-C. Park and T. Kurita</i>	1083
Frequently Deleted Loci on Chromosome 9 May Harbor Several Tumor Suppressor Genes in Human Renal Cell Carcinoma. <i>B. Grady, R. Goharderakhshan, J. Chang, L. A. Ribeiro-Filho, G. Perinchery, J. Franks, J. Presti, P. Carroll and R. Dahiya</i>	1088
Potent Antitumor Effects of CD154 Transduced Tumor Cells in Experimental Bladder Cancer. <i>A. Loskog, A. Björkland, M. P. Brown, O. Korsgren, P.-U. Malmström and T. H. Tötterman</i>	1093
Effect of Antisense Bcl-2 Oligonucleotides on Bcl-2 Protein Expression and Apoptosis in Human Bladder Transitional Cell Carcinoma. <i>B. J. Duggan, P. Maxwell, J. D. Kelly, P. Canning, N. H. Anderson, P. F. Keane, S. R. Johnston and K. E. Williamson</i>	1098
Pyridinoline Cross-Linked Carboxyterminal Telopeptide of Type I Collagen as Useful Marker for Monitoring Metastatic Bone Activity in Men With Prostate Cancer. <i>M. Noguchi and S. Noda</i>	1106
Histological and Neurotrophic Changes Triggered by Varying Models of Bladder Inflammation. <i>M. C. Dupont, J. M. Spitsbergen, K. B. Kim, J. B. Tuttle and W. D. Steers</i>	1111
Prevention of Further Cyclophosphamide Induced Hemorrhagic Cystitis by Hyperbaric Oxygen and Mesna in Guinea Pigs. <i>A. Korkmaz, S. Oter, S. Deveci, C. Goksoy and H. Bilgic</i>	1119
Does Cyclooxygenase-2 Inhibitor Prevent Renal Tissue Damage in Unilateral Ureteral Obstruction? <i>A. Miyajima, K. Ito, T. Asano, K. Seta, A. Ueda and M. Hayakawa</i>	1124
Combined Evaluation of Detrusor Pressure and Bladder Wall Thickness as Parameter for Assessment of Detrusor Function: Experimental In Vivo Study. <i>S. Bross, S. Schumacher, J. R. Scheepe, P. M. Bruan, P. Alken and K.-P. Juenemann</i>	1130
Role of Ovarian Hormones in Pathogenesis of Impaired Detrusor Contractility: Evidence in Ovariectomized Rodents. <i>Q. Zhu, J. Ritchie, N. Marouf, S. B. Dion, N. M. Resnick, A. Elbadawi and G. A. Kuchel</i>	1136
Efficacy of $\beta 3$ -Adrenergic Receptor Agonist CL-316243 on Experimental Bladder Hyperreflexia and Detrusor Instability in Rat. <i>M. Woods, N. Carson, N. W. Norton, J. H. Sheldon and T. M. Argentieri</i>	1142
Expression of Neural Plasticity Related Gene in Pontine Tegmental Area of Rats With Overactive Bladder After Cerebral Infarction. <i>S. Yotsuyanagi, O. Yokoyama, K. Komatsu, K. Kodama, S. Niikura and M. Namiki</i>	1148
Response of Fetal Sheep Bladder Tissue to Partial Outlet Obstruction. <i>R. M. Levin, E. Macarak, P. Howard, P. Horan and B. A. Kogan</i>	1156
Spontaneous Ca^{2+} Activated Cl^{-} Currents in Isolated Urethral Smooth Muscle Cells. <i>G. P. Sergeant, M. A. Hollywood, N. G. McHale and K. D. Thornbury</i>	1161
Potassium Outward Currents in Freshly Dissociated Rabbit Corpus Cavernosum Myocytes. <i>J. Malysz, S. J. Gibbons, S. M. Miller, M. Gettman, A. Nehra, J. H. Szurszewski and G. Farrugia</i>	1167
Differences in Antigen Pattern Recognized by Antisperm Antibodies in Patients With Infertility and Vasectomy. <i>C. Bohring and W. Krause</i>	1178
Canine Model of Infertility After Spinal Cord Injury: Time Course of Acute Changes in Semen Quality and Spermatogenesis. <i>D. A. Ohl, J. Sønksen, G. Wedemeyer, M. C. Zaborniak, T. N. Dam, A. C. Menge, M. J. Putzi and S. M. Papadopoulos</i>	1181
Evidence of Inhibitory Effect of Diet and Exercise on Prostate Cancer Cell Growth. <i>C. N. Tymchuk, R. J. Barnard, D. Heber and W. J. Aronson</i>	1185

UROLOGICAL SURVEY

Urological Oncology: Testis Cancer and Advances in Oncological Therapy. <i>J. P. Richie</i>	1192
Adrenal and Renal Physiology, and Medical Renal Disease. <i>W. S. McDougal</i>	1195
Renal Transplantation and Renovascular Hypertension. <i>D. A. Goldfarb</i>	1197

POSTERIOR TIBIAL NERVE STIMULATION AS NEUROMODULATIVE TREATMENT OF LOWER URINARY TRACT DYSFUNCTION

MICHAEL R. VAN BALKEN, VERA VANDONINCK, KAREL W. H. GISOLF, HENK VERGUNST, LAMBERTUS A. L. M. KIEMENEY, FRANS M. J. DEBRUYNE AND BART L. H. BEMELMANS

From the Department of Urology, Canisius-Wilhemina Hospital and Departments of Urology and Epidemiology, University Medical Centre, Nijmegen and Department of Urology, University Hospital, Utrecht, The Netherlands

ABSTRACT

Purpose: Recently, intermittent percutaneous posterior tibial nerve stimulation was introduced as a treatment modality filling the gap between conservative and surgical therapies in patients with certain types of lower urinary tract dysfunction.

Materials and Methods: In a prospective multicenter trial posterior tibial nerve stimulation was evaluated in 37 patients who presented with symptoms of bladder overactivity, that is the urgency and frequency syndrome and/or urge incontinence, and 12 with nonobstructive urinary retention. Results were recorded in voiding diaries and on quality of life questionnaires before and after treatment. Patients were classified as responders, including those in whom therapy was successful and chose to continue treatment after the initial 12 weeks, and nonresponders, those who chose to stop treatment.

Results: Overall, a positive response was seen in 60% of all patients. In patients with bladder overactivity a statistically significant decrease was observed in leakage episodes, number of pads used, voiding frequency and nocturia, and an equal increase in mean and smallest volume voided. Improvements were also seen in nonobstructive urinary retention, including number of catheterizations, total and mean volume catheterized, and total and mean volume voided. Disease specific quality of life and some domains of general quality of life improved, especially of bladder overactivity. Only mild side effects were observed.

Conclusions: Posterior tibial nerve stimulation is a minimally invasive and successful treatment option for patients with certain types of lower urinary tract dysfunction.

KEY WORDS: bladder, neurotransmitters, transcutaneous electric nerve stimulation, tibial nerve

Non-neurogenic lower urinary tract dysfunction is a common urological problem that strongly affects quality of life. Patients can complain of urgency and frequency, urge incontinence or present with urinary retention. In most patients the etiology of these symptoms remains unclear.¹ Conservative treatment options for bladder overactivity, that is the urgency and frequency syndrome and/or urge incontinence, consist of behavioral techniques with and without biofeedback, bladder reeducation, pelvic muscle exercise or pharmacotherapy involving anticholinergics, antispasmodics and tricyclic antidepressants. Patients with nonobstructive urinary retention can be treated with clean intermittent or permanent catheterization. For refractory cases more aggressive surgical procedures, including bladder distension, ileocystoplasty or urinary diversion have been advocated. However, a high recurrence and complication rate limits the widespread application of these treatments.

Recently, continuous sacral root stimulation (Medtronic, Inc., Minneapolis, Minnesota) has been proposed as an alternative, less invasive therapeutic option for patients with non-neurogenic lower urinary tract dysfunction not responding to conservative treatment.² Although highly effective in select patients, this technique is expensive and requires explicit surgical skill. Posterior tibial nerve stimulation is technically less demanding and probably more cost-effective for management of lower urinary tract dysfunction. We present our initial experience with this new neuromodulation technique in a prospective clinical trial.

PATIENTS AND METHODS

Between November 1999 and March 2000, 15 male and 34 female patients were enrolled in a prospective clinical multicenter trial that received approval by the institutional review board. All patients were evaluated for urgency and frequency, urge incontinence and nonobstructive retention by history, voiding diaries consisting of frequency volume charts, and physical as well as urological examination, including urodynamics. Urgency and frequency was defined as greater than 8 voids per 24 hours and the sudden urge to void could hardly be suppressed. Urge incontinence was characterized by urgency leading to urinary leakage occurring at least 3 times weekly and/or phasic involuntary bladder contractions with concomitant incontinence on cystometry. Non-obstructive retention was distinguished by urinary retention necessitating intermittent catheterization at least 4 times daily without urodynamic signs of outflow obstruction, as defined on the Abrams-Griffiths nomogram.³ The use of parasympatholytic medication or other pharmaceuticals influencing bladder function, including antidepressant agents, should have been stopped 2 weeks or longer before posterior tibial nerve stimulation or continued without dose changes during the entire study. Specific exclusion criteria for our study are given in the Appendix.

Patients. A total of 10 men and 27 women with a mean age of 52.5 years (range 23 to 74) were treated for bladder overactivity. These patients lived with symptoms for a median of 4 years (range 1 to 30). The symptoms were said to be induced by childbirth in 3, pelvic surgery in 9, including 7 hysterectomies, 1 anterior prolapse correction and 1 Burch colposuspension, and other events in 3 patients, including

fracture of the coccygeal bone, urinary tract infection and bowel infection. The remaining patients had no history of any pelvic event before the onset of symptoms. Of the 37 patients 32 (91.4%) received unsuccessful prior medical therapy, including anticholinergics, with a mean number of 1.9 (range 1 to 5) drugs used. Almost half of all (48.6%) patients underwent up to 3 surgical procedures, most frequently colposuspension, for symptoms. Other unsuccessful therapies included physiotherapy and/or biofeedback in 22 (59.4%), electrical stimulation, including transcutaneous electrical nerve stimulation or peripheral nerve evaluation, in 5 (13.5%) and alternative therapies, such as Chinese herbs in 5 (13.5%) patients. Physical examination did not show overt or neuro-urological abnormalities in any patients.

There were 5 men and 7 women with a mean age of 50.8 years (range 36 to 64) treated for nonobstructive retention. These patients had symptoms for a median of 3.5 years (range 1 to 36). Initiation of symptoms was related to childbirth in 1, pelvic surgery, including Wertheim's-Meigs operation in 1 and Burch colposuspension in 1, other surgery, including hip replacement in 1 and inguinal hernia repair in 1, and benign prostatic hyperplasia without any effect of transurethral prostatic resection in 1 patient. The remaining 6 patients had no history of any pelvic event before the onset of symptoms. Of the 12 patients 11 (92%) received unsuccessful medical therapy previously, including α -blockers or sympathomimetics. Almost half the patients (5 of 12) had undergone lower urinary tract surgery, mostly urethral dilatation or urethrotomy, for symptoms previously. Other unsuccessful therapies included physiotherapy and/or biofeedback in 3 (25%) and electrical stimulation in 1 patient. All patients had clean intermittent catheterization. In addition, physical examination did not show overt or neuro-urological abnormalities in these patients. Urodynamic investigations revealed hypo-contractile detrusors in all patients.

Treatment method. Posterior tibial nerve stimulation was applied according to Stoller.⁴ Patients are in the supine position with the soles of the feet together, and knees abducted and flexed ("frog-position"). A 34 gauge stainless steel needle is inserted approximately 3 to 4 cm., about 3 fingerbreadths cephalad to the medial malleolus, between the posterior margin of the tibia and soleus muscle. A stick on electrode is placed on the same leg near the arch of the foot. The needle and electrode are connected to a low voltage (9 volts) Uro-surge stimulator (Uro-surge, Coralville, Iowa) with an adjustable pulse intensity of 0 to 10 mA., a fixed pulse width of 200 microseconds and a frequency of 20 Hz. The amplitude is slowly increased until the large toe starts to curl or toes start to fan. If the large toe does not curl or pain occurs near the insertion site the stimulation device is switched off and the procedure is repeated. If the large toe curls or toes start to fan stimulation is applied at an intensity well tolerated by the patient. If necessary the amplitude can be increased during the session. Patients underwent 12 weekly outpatient treatment sessions, each lasting for 30 minutes. If a good response occurred the patient was offered chronic treatment.

Evaluation of results. After providing informed consent all patients completed a voiding diary, as well as general and disease specific quality of life questionnaires at study entry and completion of the 12-week treatment. In the 24-hour voiding diary frequency and voided volumes, number and volume of catheterizations, number of leakage periods, leakage severity as well as the number of pads or diapers used were recorded. The leakage severity was scored on a scale of 0 to 3 and included 0—no leakage, 1—leakage of some drops, 2—loss of a small amount and 3—loss of a massive amount, that is change of clothes. General and disease specific quality of life questionnaires were evaluated with the 36-item short-form health survey (SF-36) and the adjusted quality of life questionnaire, respectively.^{5,6} The SF-36 consists of 36 items regarding 8 distinct health status concepts and 1 item measuring self-reported health transition, including physical function, role physical, pain, general health, emotional well-being, role emotional, social functioning, energy and fatigue and change in general health. The maximum score for each domain is 100, and a higher score relates to better quality of life. The quality of life questionnaire consists of 22 items, each with a 5 point response scale. As with the SF-36, a high score means good quality of life.

Success was defined as the patient request for continued chronic treatment for maintenance. Analysis was done on an intention-to-treat basis. Within group comparisons of parametric results at baseline and 12-week treatment were conducted with the paired sample t-test. Statistical analysis was performed with commercial software.

RESULTS

Bladder overactivity. Of the 444 treatment sessions stimulation was performed on the right posterior tibial nerve with a mean pulse intensity of 3.7 mA. (range 1.4 to 6.8) per session in 265 (60%). Complications, including minor bleeding or a temporary painful feeling at the insertion site, were seen only rarely. Of the 37 patients posterior tibial nerve stimulation in 7 men and 15 women (59.4%) was considered successful because they requested continuation of therapy after completion of the 12-week treatment (group 1, responders). In 3 male and 12 female patients treatment was unsuccessful since they did not choose maintenance therapy (group 2, nonresponders).

At completion of the 12-week treatment all 33 patients with urgency and frequency had significant improvement of voiding frequency during the day and night (table 1 and fig. 1, A). When discriminating group 1 (21 responders) from group 2 (12 nonresponders), these improvements appeared to be significant only in group 1. The smallest mean values and mean volumes voided also improved, whereas the largest volumes voided did not change in this group.

A statistically significant overall decrease in incontinence was seen in the 30 patients who were urge incontinent (table 2 and fig. 1, B). When comparing group 1 (18 responders) with group 2 (12 nonresponders), improvements found in

TABLE 1. Voiding parameters in 33 patients with urgency and frequency

	Frequency	Nocturia	Vol. Voided		
			Smallest	Largest	Mean
All pts:					
Mean $\pm$ SD (0 wks.)	16.5 $\pm$ 6.8	2.6 $\pm$ 2.3	65 $\pm$ 49.3	289.8 $\pm$ 161	140 $\pm$ 82.2
Mean change (95% CI)	-2.8 (-0.8, -4.9)	-1 (-0.2, -1.8)	14.2 (32.8, -4.5)	13.9 (48.7, -20.9)	19.3 (40.3, -1.7)
p Value	<0.05	<0.05			
Group 1:					
Mean $\pm$ SD (0 wks.)	16.1 $\pm$ 6.1	2.5 $\pm$ 2.7	62.6 $\pm$ 52.1	268.8 $\pm$ 146.4	133 $\pm$ 82.8
Mean change (95% CI)	-4.8 (-2.7, 6.8)	-1.4 (-0.4, 2.3)	29.8 (49.2, 10.3)	9.5 (50.6, -31.6)	35.1 (62.6, 7.7)
p Value	<0.0005	<0.05	<0.005		<0.05
Group 2:					
Mean $\pm$ SD (0 wks.)	17.1 $\pm$ 8.3	2.7 $\pm$ 1.6	71 $\pm$ 44.8	334 $\pm$ 188.5	153.6 $\pm$ 83.7
Mean change (95% CI)	0.8 (5, -3.3)	-0.2 (1.3, -1.6)	-18.5 (19.4, -56.4)	23 (100.6, -54.6)	-14 (8.5, -36.5)




FIG. 1. Before and after posterior tibial nerve stimulation for responders, nonresponders and both combined. A, voiding frequency. B, number of leakage episodes. Error bar represents SD. ns, not significant.

TABLE 2. Voiding parameters in 30 patients with urge incontinence

	Leakage Episodes	Pads Used	Leakage Severity
All pts:			
Mean ± SD (0 wks.)	9.8 ± 5.6	6.3 ± 3.2	1.7 ± 0.5
Mean change (95% CI)	-4.8 (-2.2, -7.5)	-2.5 (-1.2, -3.9)	-0.7 (-0.4, -1.0)
p Value	<0.005	<0.005	<0.005
Group 1:			
Mean ± SD (0 wks.)	9.8 ± 6.4	5.8 ± 2.6	1.6 ± 0.5
Mean change (95% CI)	-6.2 (-2.9, -9.5)	-3.3 (-1.5, -5.1)	-0.8 (-0.3, -1.3)
p Value	<0.005	<0.005	<0.005
Group 2:			
Mean ± SD (0 wks.)	9.8 ± 4.4	7.1 ± 3.9	1.9 ± 0.5
Mean change (95% CI)	-2.4 (2.4, -7.3)	-1.3 (0.9, -3.4)	-0.5 (-0.2, -0.8)
p Value			<0.05

leakage episodes and number of pads used appeared to be greater in group 1. Leakage severity scores showed improvement in all patients who were urge incontinent, and groups 1 and 2. General quality of life improved in regard to physical and social functioning in all patients with bladder overactivity. Mean physical functioning plus or minus standard deviation (SD) was 53.7 ± 27.9 (mean change +9.1, 95% confidence interval [CI] +14.1, +4, $p < 0.005$) and social functioning 64.7 ± 31.2 (mean change +10.3, +16.5, +4.1, respectively, $p < 0.005$) at 0 weeks. In group 1 both scores also increased. Mean physical functioning plus or minus SD was 61.6 ± 26.3 (mean change +11.2, 95% CI +18.8, +3.5, $p < 0.05$) and social functioning 70.6 ± 26.7 (mean change +11.6, +20.9, +2.3, respectively, $p < 0.05$) at 0 weeks. Only social functioning improved (55.9 ± 35.9 , mean change +8.4, 95% CI +16.6, +0.2, respectively, $p < 0.05$) at 0 weeks in group 2. Disease specific quality of life showed improvement in all patients (60.3 ± 19.3 , mean change +12.3, 95% CI +17.3, +6.8, $p = 0.00004$), as well as in group 1 (60.6 ± 18.2 , mean change +14.4, +22.3, +6.6, $p < 0.005$) and group 2 (59.9 ± 21.5 , mean change +8.5, +15.1, +2, respectively, $p < 0.05$) at 0 weeks (fig. 2, A).

Nonobstructive retention. Of the 144 treatment sessions stimulation was performed on the right posterior tibial nerve with a mean pulse intensity of 4 mA. (range 2.5 to 7) per session in 87 (60%). As seen in patients treated for urgency and frequency and/or urge incontinence, complications were rarely noted. Of the 12 patients posterior tibial nerve stimulation was considered successful in 1 man and 6 women (58.3%) because they requested continuation of therapy after completion of the 12-week treatment (group 3, responders). Treatment was unsuccessful in 4 men and 1 woman since they did not choose maintenance therapy (group 4, nonresponders).

At completion of the 12-week treatment all patients clearly had a decrease in the number of catheterizations, and mean and total volumes catheterized, and an increase in mean and total volumes voided. However, statistical significance could not be obtained (table 3). When discriminating group 3 (7 responders) from group 4 (5 nonresponders) these changes could also be observed in group 3 but not in group 4. In regard to quality of life hardly any significant improvement could be found in the adjusted disease specific (fig. 2, B) or general quality of life questionnaire. The only exceptions appeared to




FIG. 2. Before and after posterior tibial nerve stimulation for responders, nonresponders and both combined. A, disease specific quality of life scores for patients with bladder overactivity. B, nonobstructive retention. Maximum disease specific quality of life score was 110. Error bar represents SD. ns, not significant.

TABLE 3. *Voiding parameters in 12 patients with nonobstructive retention*

	All Pts.		Group 3		Group 4	
	Mean $\pm$ SD (0 wks.)	Mean Change (95% CI)	Mean $\pm$ SD (0 wks.)	Mean Change (95% CI)	Mean $\pm$ SD (0 wks.)	Mean Change (95% CI)
No. catheterizations	5.3 $\pm$ 2.4	-0.8 (+0.6, -2.3)	5.9 $\pm$ 2.6	-1.4 (1.2, -4.1)	4.4 $\pm$ 2	0 (0.9, -0.9)
Mean vol. catheterized	336 $\pm$ 171	-83.3 (27.4, -194)	325 $\pm$ 181	-131.9 (63.2, -326.9)	351 $\pm$ 176	-15.4 (87.5, -118.4)
Total vol. catheterized	1,552 $\pm$ 776	-537 (175, -1249)	1,707 $\pm$ 915	-921.4 (284.5, -2127.4)	1,335 $\pm$ 547	1 (659.7, -657.7)
Mean vol. voided	206 $\pm$ 192	5.9 (113.2, -101.4)	141 $\pm$ 107	94.3 (233.3, -44.7)	297 $\pm$ 258	-117.8 (16.5, 252.1)
Total vol. voided	919 $\pm$ 658	173.8 (639.2, -291.7)	758 $\pm$ 567	482.1 (1164.7, -200.4)	1,145 $\pm$ 774	-258 (374.6, -890.6)

be the SF-36 score for emotional well-being, which increased (58.3 ± 12.2 , mean change $+14.3$, 95% CI $+20.6$, $+7.9$, $p < 0.005$) in patients who thought treatment was successful (group 3), and the score for change of health, which significantly increased in all (45.8 ± 27.9 , mean change $+22.9$, $+44.9$, $+1$, $p < 0.05$) but not in group 3 or 4 at 0 week, respectively.

DISCUSSION

For treatment of refractory urinary tract dysfunction various methods of intermittent neuromodulation have been advocated, including intravesical, anal, vaginal, penile and perineal stimulation, and transcutaneous electrical stimulation of the suprapubic or sacral region and posterior tibial nerve.⁷⁻¹⁵ Because of poor results or uncomfortable stimulation sites, most of the aforementioned treatment modalities did not gain widespread acceptance. In the 1980s implantable neurostimulation electrodes became available for modulation of the sacral spinal nerves, especially S3.¹⁶ Stimulation could be performed continuously, leading to better results but drawbacks included invasiveness of the procedure, the high costs involved and limited service life of the stimulation device, which was 7 to 10 years. In addition, symptoms appear to recur almost immediately after discontinuation of stimulation.² Therefore, the development of an easily applicable, noninvasive or minimally invasive, cost-effective neuromodulation device with good treatment outcome was anticipated. The recent advent of posterior tibial nerve stimulation may offer urologists a treatment modality that meets the aforementioned criteria.^{4,17}

Stimulation of the tibial nerve was first described by McGuire et al in 1983.¹⁵ In studies of nonhuman primates with spinal cord injury bipolar anal sphincter stimulation resulted in the inhibition of detrusor activity. Similar results could be obtained by applying current through a transcutaneous electrode over the common peroneal or posterior tibial nerve, with a ground electrode being contralaterally placed over the same nerves. The idea of stimulating these nerves was based on the traditional Chinese practice of using acupuncture points over the common peroneal or posterior tibial nerves to affect bladder activity.¹⁵ Electrical stimulation of the tibial nerve was subsequently performed in 22 patients with detrusor instability, interstitial or radiation cystitis or neurological disease with promising results, including continence in 12 and urodynamics improvement in 7. However, in a prospective study by Geirsson et al in patients with interstitial cystitis, transcutaneous tibial nerve stimulation or traditional Chinese acupuncture revealed no difference in voiding frequency, mean and maximal voided volumes and visual analog symptom scores before or after either treatment modality.¹⁸

In our study percutaneous stimulation resulted in a success rate of about 60% of 49 patients with bladder overactivity or nonobstructive urinary retention. Although our definition of success was a subjective one, responders differed significantly from nonresponders in almost every semi-objective parameter. The same result was observed in disease specific and several domains of general quality of life scores.

Because our study was not placebo controlled, the observed

improvements might theoretically be the result of regression to the mean, synchronous other treatment or placebo effect rather than a positive response to posterior tibial nerve stimulation. Regression to the mean is not likely as all patients included had to present with symptoms for at least 6 months (median 4 years). To prevent interference of other treatment modalities, concomitant physiotherapy during posterior tibial nerve stimulation was not permitted, as was the case with starting parasympatholytic medication or other pharmaceuticals influencing bladder function, including antidepressant agents, within 2 weeks before treatment. In patients who were already using any of the aforementioned drugs medication had to be stopped 2 weeks before beginning posterior tibial nerve stimulation or continued without dose changes during the entire 12 weeks. In placebo controlled studies success rates in the placebo group are usually much lower than the response rate of about 60% in our study. Because most patients had symptoms for a long period and undergone various treatments before trying neuromodulation, it is not likely that they were highly susceptible to the placebo effect of this treatment modality. Nevertheless, to validate these data a properly designed placebo controlled study is warranted. However, it can be anticipated that the feasibility of such studies will be difficult.

Notwithstanding the promising results and only mild complications of this neuromodulation technique found in our study, this therapy also has some disadvantages, including percutaneous insertion of the stimulation electrode, necessity of regular visits to the outpatient clinic and, thus far, unknown results of chronic treatment of initially successful cases. In the near future most of the aforementioned problems may be solved with an implantable subcutaneous electrode, which can be radiographically stimulated by the patients at home and on a regular basis. Studies regarding long-term effects are ongoing but results are not yet available. In conclusion, although thorough research is anticipated to improve patient selection criteria, optimize the technique and help learn more about the precise mode of action, posterior tibial nerve stimulation is a promising, cost-effective and easily applicable treatment option for patients with lower urinary tract dysfunction.

APPENDIX: EXCLUSION CRITERIA

- Younger than 18 years
- Symptoms existing for less than 6 months
- Pregnancy or intention to become pregnant during the course of study
- Active urinary tract or recurrent urinary tract infection (5 or greater recurrent infections during the last 12 months), carcinoma in situ, bladder malignancy, interstitial cystitis
- Bladder or kidney stone
- Severe cardiopulmonary disease
- Use of pentosan polysulfate sodium or bladder installations, including dimethyl sulfoxide, bacillus Calmette-Guerin, Chloropectin or heparin
- Uncontrolled diabetes
- Diabetes with peripheral nerve involvement

Neurological disease like multiple sclerosis, Parkinson's disease, cerebrovascular accident, bifid spine or spinal cord lesion

Change in parasympatholytic medication or other pharmaceuticals influencing bladder function, including antidepressant agents, within 2 weeks before or during the study

Physiotherapy during the study

Bladder outlet obstruction (Abrams-Griffiths nomogram)

Transurethral instrumentation 4 weeks or less before or during the study

REFERENCES

- Hasan, S. T., Robson, W. A., Pridie, A. K. et al: Transcutaneous electrical nerve stimulation and temporary S3 neuromodulation in idiopathic detrusor instability. *J Urol*, **155**: 2005, 1996
- Bemelmans, B. L., Mundy, A. R. and Craggs, M. D.: Neuromodulation by implant for treating lower urinary tract symptoms and dysfunction. *Eur Urol*, **36**: 81, 1999
- Abrams, P. H. and Griffiths, D. J.: The assessment of prostatic obstruction from urodynamic measurements and from residual urine. *Br J Urol*, **51**: 129, 1979
- Stoller, M. L.: Afferent nerve stimulation for pelvic floor dysfunction. *Eur Urol*, suppl., **35**: 16, 1999
- Ware, J. E. and Sherbourne, C. D.: The MOS 36-item short-form health survey (SF-36). I. Conceptual framework and item selection. *Med Care*, **30**: 473, 1992
- Wagner, T. H., Patrick, D. L., Bavendam, T. G. et al: Quality of life of persons with urinary incontinence: development of a new measure. *Urology*, **47**: 67, 1996
- Madersbacher, H.: Intravesical electrical stimulation for the rehabilitation of the neuropathic bladder. *Paraplegia*, **28**: 349, 1990
- Merrill, D. C.: The treatment of detrusor incontinence by electrical stimulation. *J Urol*, **122**: 515, 1979
- Janez, J., Plevnik, S. and Suhel, P.: Urethral and bladder responses to anal electrical stimulation. *J Urol*, **122**: 192, 1979
- Lindström, S., Fall, M., Carlsson, C. A. et al: The neurophysiological basis of bladder inhibition in response to intravaginal electrical stimulation. *J Urol*, **129**: 405, 1983
- Nakamura, M. and Sakurai, T.: Bladder inhibition by penile electrical stimulation. *Br J Urol*, **56**: 413, 1984
- Vodušek, D. B., Light, J. K. and Libby, J. M.: Detrusor inhibition induced by stimulation of pudendal nerve afferents. *Neurourol Urodyn*, **5**: 381, 1986
- Fall, M. and Lindström, S.: Transcutaneous electrical nerve stimulation in classic and nonulcer interstitial cystitis. *Urol Clin North Am*, **21**: 131, 1994
- Walsh, I. K., Johnston, R. S. and Keane, P. F.: Transcutaneous sacral neurostimulation for irritative voiding dysfunction. *Eur Urol*, **35**: 192, 1999
- McGuire, E. J., Zhang, S. C., Horwinski, E. R. et al: Treatment of motor and sensory detrusor instability by electrical stimulation. *J Urol*, **129**: 78, 1983
- Schmidt, R. A.: Applications of neurostimulation in urology. *Neurourol Urodyn*, **7**: 585, 1988
- Klingler, H. C., Pycha, A., Schmidbauer, J. et al: Use of peripheral neuromodulation of the S3 region for treatment of detrusor overactivity: a urodynamic-based study. *Urology*, **56**: 766, 2000
- Geirsson, G., Wang, Y. H., Lindström, S. et al: Traditional acupuncture and electrical stimulation of the posterior tibial nerve. A trial in chronic interstitial cystitis. *Scand J Urol Nephrol*, **27**: 67, 1993