



US009254338B2

(12) **United States Patent**
Yancopoulos

(10) **Patent No.:** **US 9,254,338 B2**
(45) **Date of Patent:** **Feb. 9, 2016**

(54) **USE OF A VEGF ANTAGONIST TO TREAT
ANGIOGENIC EYE DISORDERS**

- (71) Applicant: **REGENERON
PHARMACEUTICALS, INC.,**
Tarrytown, NY (US)
- (72) Inventor: **George D. Yancopoulos**, Yorktown
Heights, NY (US)
- (73) Assignee: **Regeneron Pharmaceuticals, Inc.,**
Tarrytown, NY (US)
- (*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 132 days.

(21) Appl. No.: **13/940,370**

(22) Filed: **Jul. 12, 2013**

(65) **Prior Publication Data**
US 2013/0295094 A1 Nov. 7, 2013

- Related U.S. Application Data**
- (63) Continuation-in-part of application No.
PCT/US2012/020855, filed on Jan. 11, 2012.
- (60) Provisional application No. 61/432,245, filed on Jan.
13, 2011, provisional application No. 61/434,836,
filed on Jan. 21, 2011, provisional application No.
61/561,957, filed on Nov. 21, 2011.

- (51) **Int. Cl.**
A61K 38/18 (2006.01)
C07K 14/71 (2006.01)
A61K 47/48 (2006.01)
A61K 38/17 (2006.01)
C07K 16/22 (2006.01)
A61K 39/00 (2006.01)
- (52) **U.S. Cl.**
CPC **A61K 47/48415** (2013.01); **A61K 38/179**
(2013.01); **C07K 14/71** (2013.01); **C07K 16/22**
(2013.01); **A61K 2039/505** (2013.01)

- (58) **Field of Classification Search**
None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

7,396,664 B2	7/2008	Daly et al.
2003/0171320 A1	9/2003	Guyer
2005/0163798 A1	7/2005	Papadopoulos et al.
2005/0260203 A1	11/2005	Wiegand et al.
2006/0058234 A1	3/2006	Daly et al.
2006/0172944 A1	8/2006	Wiegand et al.
2007/0190058 A1	8/2007	Shams

FOREIGN PATENT DOCUMENTS

JP	2010-509369	3/2010
WO	00/75319	12/2000
WO	2008/063932	5/2008

OTHER PUBLICATIONS

Anonymous "Lucentis (ranibizumab injection) Intravitreal Injection" pp. 103 (Jun. 2006).

Do et al., "An exploratory study of the safety, tolerability and bioactivity of a single intravitreal injection of vascular endothelial growth factor Trap-Eye in patients with diabetic macular oedema" *Br J Ophthalmol.* 93(2):144-1449 (Feb. 2009).

Do et al., "The DA VINCI Study: phase 2 primary results of VEGF Trap-Eye in patients with diabetic macular edema" *Ophthalmology* 118(9):1819-1826 (Sep. 2011).

The Eyetech Study Group, "Anti-Vascular Endothelial Growth Factor Therapy for Subfoveal Choroidal Neovascularization Secondary to Age-related Macular Degeneration" *American Academy of Ophthalmology*, 110 (5):979-986 (May 2003).

Heier et al., "rhuFab V2 (anti-VEGF Antibody) for Treatment of Exudative AMD" *Symposium 8:Experimental and Emerging Treatments for Choroidal Neovascularization*, 10 pp (2002).

Heier et al., "RhuFab V2 in Wet AMD—6 Month Continued Improvement Following Multiple Intravitreal Injections" *Invest Ophthalmol Vis Sci*, 44:E-Abstract 972 (2003).

Krzystolik et al., "Prevention of Experimental Choroidal Neovascularization With Intravitreal Anti-Vascular Endothelial Growth Factor Antibody Fragment" *Arch Ophthalmol.*, 120:338-346 (Mar. 2002).

Nguyen et al., "A Phase I Study of Intravitreal Vascular Endothelial Growth Factor Trap-Eye in Patients with Neovascular Age-Related Macular Degeneration" *Ophthalmology*, J.B. Lippincott Co., Philadelphia, PA, US, 116(11):2141-2148 (Nov. 1, 2009).

Nichols, Earl R., "AAO: Ranibizumab (rhuRab) May Improve Vision in Age-Related Macular Degeneration" *Doctor's Guide Global Edition*, www.pslgroup.com/dg/23f2aa.htm, pp. 1-2 (Nov. 24, 20013).

Pai et al., "Current concepts in intravitreal drug therapy for diabetic retinopathy" *Saudi Journal of Ophthalmology* 24(4):143-149 (Jun. 30, 2010).

Stewart, "The expanding role of vascular endothelial growth factor inhibitors in ophthalmology" *Mayo Clin Proc.* 87 (1):77-88 (Jan. 2012).

Thomas Reuters Integrity "VEGF Trap-Eye final phase II results in age-related macular degeneration presented at 2008 Retina Society Meeting" (Sep. 28, 2008).

Nguyen et al., "A phase I trial of an IV-administered vascular endothelial growth factor trap for treatment in patients with choroidal neovascularization due to age-related macular degeneration" *Ophthalmology* (Sep. 2006) 113 (9):1522e1-1522e14 (epub Jul. 28, 2006).

Primary Examiner — Christine J Saoud


Assistant Examiner — Jon M Lockard

(74) *Attorney, Agent, or Firm* — Frank Cottingham; Karl Bozicevic

(57) **ABSTRACT**

The present invention provides methods for treating angiogenic eye disorders by sequentially administering multiple doses of a VEGF antagonist to a patient. The methods of the present invention include the administration of multiple doses of a VEGF antagonist to a patient at a frequency of once every 8 or more weeks. The methods of the present invention are useful for the treatment of angiogenic eye disorders such as age related macular degeneration, diabetic retinopathy, diabetic macular edema, central retinal vein occlusion, branch retinal vein occlusion, and corneal neovascularization.

26 Claims, 1 Drawing Sheet



1

USE OF A VEGF ANTAGONIST TO TREAT ANGIOGENIC EYE DISORDERS

CROSS-REFERENCE TO RELATED APPLICATIONS

This application is a continuation-in-part of International Patent Application No. PCT/US2012/020855, filed on Jan. 11, 2012, which claims the benefit of U.S. Provisional Application Nos. 61/432,245, filed on Jan. 13, 2011, 61/434,836, filed on Jan. 21, 2011, and 61/561,957, filed on Nov. 21, 2011, the contents of which are hereby incorporated by reference in their entireties.

FIELD OF THE INVENTION

The present invention relates to the field of therapeutic treatments of eye disorders. More specifically, the invention relates to the administration of VEGF antagonists to treat eye disorders caused by or associated with angiogenesis.

BACKGROUND

Several eye disorders are associated with pathological angiogenesis. For example, the development of age-related macular degeneration (AMD) is associated with a process called choroidal neovascularization (CNV). Leakage from the CNV causes macular edema and collection of fluid beneath the macula resulting in vision loss. Diabetic macular edema (DME) is another eye disorder with an angiogenic component. DME is the most prevalent cause of moderate vision loss in patients with diabetes and is a common complication of diabetic retinopathy, a disease affecting the blood vessels of the retina. Clinically significant DME occurs when fluid leaks into the center of the macula, the light-sensitive part of the retina responsible for sharp, direct vision. Fluid in the macula can cause severe vision loss or blindness. Yet another eye disorder associated with abnormal angiogenesis is central retinal vein occlusion (CRVO). CRVO is caused by obstruction of the central retinal vein that leads to a back-up of blood and fluid in the retina. The retina can also become ischemic, resulting in the growth of new, inappropriate blood vessels that can cause further vision loss and more serious complications. Release of vascular endothelial growth factor (VEGF) contributes to increased vascular permeability in the eye and inappropriate new vessel growth. Thus, inhibiting the angiogenic-promoting properties of VEGF appears to be an effective strategy for treating angiogenic eye disorders.

FDA-approved treatments of angiogenic eye disorders such as AMD and CRVO include the administration of an anti-VEGF antibody called ranibizumab (Lucentis®, Genentech, Inc.) on a monthly basis by intravitreal injection.

Methods for treating eye disorders using VEGF antagonists are mentioned in, e.g., U.S. Pat. Nos. 7,303,746; 7,306,799; 7,300,563; 7,303,748; and US 2007/0190058. Nonetheless, there remains a need in the art for new administration regimens for angiogenic eye disorders, especially those which allow for less frequent dosing while maintaining a high level of efficacy.

BRIEF SUMMARY OF THE INVENTION

The present invention provides methods for treating angiogenic eye disorders. The methods of the invention comprise sequentially administering multiple doses of a VEGF antagonist to a patient over time. In particular, the methods of the invention comprise sequentially administering to the patient a

2

single initial dose of a VEGF antagonist, followed by one or more secondary doses of the VEGF antagonist, followed by one or more tertiary doses of the VEGF antagonists. The present inventors have surprisingly discovered that beneficial therapeutic effects can be achieved in patients suffering from angiogenic eye disorders by administering a VEGF antagonist to a patient at a frequency of once every 8 or more weeks, especially when such doses are preceded by about three doses administered to the patient at a frequency of about 2 to 4 weeks. Thus, according to the methods of the present invention, each secondary dose of VEGF antagonist is administered 2 to 4 weeks after the immediately preceding dose, and each tertiary dose is administered at least 8 weeks after the immediately preceding dose. An example of a dosing regimen of the present invention is shown in FIG. 1. One advantage of such a dosing regimen is that, for most of the course of treatment (i.e., the tertiary doses), it allows for less frequent dosing (e.g., once every 8 weeks) compared to prior administration regimens for angiogenic eye disorders which require monthly administrations throughout the entire course of treatment. (See, e.g., prescribing information for Lucentis® [ranibizumab], Genentech, Inc.).

The methods of the present invention can be used to treat any angiogenic eye disorder, including, e.g., age related macular degeneration, diabetic retinopathy, diabetic macular edema, central retinal vein occlusion, corneal neovascularization, etc.

The methods of the present invention comprise administering any VEGF antagonist to the patient. In one embodiment, the VEGF antagonist comprises one or more VEGF receptor-based chimeric molecule(s), (also referred to herein as a “VEGF-Trap” or “VEGFT”). An exemplary VEGF antagonist that can be used in the context of the present invention is a multimeric VEGF-binding protein comprising two or more VEGF receptor-based chimeric molecules referred to herein as “VEGFR1R2-FcAC1(a)” or “aflibercept.”

Various administration routes are contemplated for use in the methods of the present invention, including, e.g., topical administration or intraocular administration (e.g., intravitreal administration).

Aflibercept (EYLEA™, Regeneron Pharmaceuticals, Inc) was approved by the FDA in November 2011, for the treatment of patients with neovascular (wet) age-related macular degeneration, with a recommended dose of 2 mg administered by intravitreal injection every 4 weeks for the first three months, followed by 2 mg administered by intravitreal injection once every 8 weeks.

Other embodiments of the present invention will become apparent from a review of the ensuing detailed description.

BRIEF DESCRIPTION OF THE FIGURE

FIG. 1 shows an exemplary dosing regimen of the present invention. In this regimen, a single “initial dose” of VEGF antagonist (“VEGFT”) is administered at the beginning of the treatment regimen (i.e. at “week 0”), two “secondary doses” are administered at weeks 4 and 8, respectively, and at least six “tertiary doses” are administered once every 8 weeks thereafter, i.e., at weeks 16, 24, 32, 40, 48, 56, etc.).

DETAILED DESCRIPTION

Before the present invention is described, it is to be understood that this invention is not limited to particular methods and experimental conditions described, as such methods and conditions may vary. It is also to be understood that the

3

terminology used herein is for the purpose of describing particular embodiments only, and is not intended to be limiting, since the scope of the present invention will be limited only by the appended claims.

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. As used herein, the term "about," when used in reference to a particular recited numerical value, means that the value may vary from the recited value by no more than 1%. For example, as used herein, the expression "about 100" includes 99 and 101 and all values in between (e.g., 99.1, 99.2, 99.3, 99.4, etc.).

Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, the preferred methods and materials are now described.

Dosing Regimens

The present invention provides methods for treating angiogenic eye disorders. The methods of the invention comprise sequentially administering to a patient multiple doses of a VEGF antagonist. As used herein, "sequentially administering" means that each dose of VEGF antagonist is administered to the patient at a different point in time, e.g., on different days separated by a predetermined interval (e.g., hours, days, weeks or months). The present invention includes methods which comprise sequentially administering to the patient a single initial dose of a VEGF antagonist, followed by one or more secondary doses of the VEGF antagonist, followed by one or more tertiary doses of the VEGF antagonist.

The terms "initial dose," "secondary doses," and "tertiary doses," refer to the temporal sequence of administration of the VEGF antagonist. Thus, the "initial dose" is the dose which is administered at the beginning of the treatment regimen (also referred to as the "baseline dose"); the "secondary doses" are the doses which are administered after the initial dose; and the "tertiary doses" are the doses which are administered after the secondary doses. The initial, secondary, and tertiary doses may all contain the same amount of VEGF antagonist, but will generally differ from one another in terms of frequency of administration. In certain embodiments, however, the amount of VEGF antagonist contained in the initial, secondary and/or tertiary doses will vary from one another (e.g., adjusted up or down as appropriate) during the course of treatment.

In one exemplary embodiment of the present invention, each secondary dose is administered 2 to 4 (e.g., 2, 2½, 3, 3½, or 4) weeks after the immediately preceding dose, and each tertiary dose is administered at least 8 (e.g., 8, 8½, 9, 9½, 10, 10½, 11, 11½, 12, 12½, 13, 13½, 14, 14½, or more) weeks after the immediately preceding dose. The phrase "the immediately preceding dose," as used herein, means, in a sequence of multiple administrations, the dose of VEGF antagonist which is administered to a patient prior to the administration of the very next dose in the sequence with no intervening doses.

In one exemplary embodiment of the present invention, a single initial dose of a VEGF antagonist is administered to a patient on the first day of the treatment regimen (i.e., at week 0), followed by two secondary doses, each administered four weeks after the immediately preceding dose (i.e., at week 4 and at week 8), followed by at least 5 tertiary doses, each administered eight weeks after the immediately preceding dose (i.e., at weeks 16, 24, 32, 40 and 48). The tertiary doses may continue (at intervals of 8 or more weeks) indefinitely during the course of the treatment regimen. This exemplary administration regimen is depicted graphically in FIG. 1.

4

The methods of the invention may comprise administering to a patient any number of secondary and/or tertiary doses of a VEGF antagonist. For example, in certain embodiments, only a single secondary dose is administered to the patient. In other embodiments, two or more (e.g., 2, 3, 4, 5, 6, 7, 8, or more) secondary doses are administered to the patient. Likewise, in certain embodiments, only a single tertiary dose is administered to the patient. In other embodiments, two or more (e.g., 2, 3, 4, 5, 6, 7, 8, or more) tertiary doses are administered to the patient.

In embodiments involving multiple secondary doses, each secondary dose may be administered at the same frequency as the other secondary doses. For example, each secondary dose may be administered to the patient 4 weeks after the immediately preceding dose. Similarly, in embodiments involving multiple tertiary doses, each tertiary dose may be administered at the same frequency as the other tertiary doses. For example, each tertiary dose may be administered to the patient 8 weeks after the immediately preceding dose. Alternatively, the frequency at which the secondary and/or tertiary doses are administered to a patient can vary over the course of the treatment regimen. For example, the present invention includes methods which comprise administering to the patient a single initial dose of a VEGF antagonist, followed by one or more secondary doses of the VEGF antagonist, followed by at least 5 tertiary doses of the VEGF antagonist, wherein the first four tertiary doses are administered 8 weeks after the immediately preceding dose, and wherein each subsequent tertiary dose is administered from 8 to 12 (e.g., 8, 8½, 9, 9½, 10, 10½, 11, 11½, 12) weeks after the immediately preceding dose. The frequency of administration may also be adjusted during the course of treatment by a physician depending on the needs of the individual patient following clinical examination.

VEGF Antagonists

The methods of the present invention comprise administering to a patient a VEGF antagonist according to specified dosing regimens. As used herein, the expression "VEGF antagonist" means any molecule that blocks, reduces or interferes with the normal biological activity of VEGF.

VEGF antagonists include molecules which interfere with the interaction between VEGF and a natural VEGF receptor, e.g., molecules which bind to VEGF or a VEGF receptor and prevent or otherwise hinder the interaction between VEGF and a VEGF receptor. Specific exemplary VEGF antagonists include anti-VEGF antibodies, anti-VEGF receptor antibodies, and VEGF receptor-based chimeric molecules (also referred to herein as "VEGF-Traps").

VEGF receptor-based chimeric molecules include chimeric polypeptides which comprise two or more immunoglobulin (Ig)-like domains of a VEGF receptor such as VEGFR1 (also referred to as Flt1) and/or VEGFR2 (also referred to as Flk1 or KDR), and may also contain a multimerizing domain (e.g., an Fc domain which facilitates the multimerization [e.g., dimerization] of two or more chimeric polypeptides). An exemplary VEGF receptor-based chimeric molecule is a molecule referred to as VEGFR1R2-FcΔC1(a) which is encoded by the nucleic acid sequence of SEQ ID NO:1. VEGFR1R2-FcΔC1(a) comprises three components: (1) a VEGFR1 component comprising amino acids 27 to 129 of SEQ ID NO:2; (2) a VEGFR2 component comprising amino acids 130 to 231 of SEQ ID NO:2; and (3) a multimerization component ("FcΔC1(a)") comprising amino acids 232 to 457 of SEQ ID NO:2 (the C-terminal amino acid of SEQ ID NO:2 [i.e., K458] may or may not be included in the VEGF antago-

5

nist used in the methods of the invention; see e.g., U.S. Pat. No. 7,396,664). Amino acids 1-26 of SEQ ID NO:2 are the signal sequence.

The VEGF antagonist used in the Examples set forth herein below is a dimeric molecule comprising two VEGFR1R2-FcΔC1(a) molecules and is referred to herein as "VEGFT." Additional VEGF receptor-based chimeric molecules which can be used in the context of the present invention are disclosed in U.S. Pat. Nos. 7,396,664, 7,303,746 and WO 00/75319.

Angiogenic Eye Disorders

The methods of the present invention can be used to treat any angiogenic eye disorder. The expression "angiogenic eye disorder," as used herein, means any disease of the eye which is caused by or associated with the growth or proliferation of blood vessels or by blood vessel leakage. Non-limiting examples of angiogenic eye disorders that are treatable using the methods of the present invention include age-related macular degeneration (e.g., wet AMD, exudative AMD, etc.), retinal vein occlusion (RVO), central retinal vein occlusion (CRVO; e.g., macular edema following CRVO), branch retinal vein occlusion (BRVO), diabetic macular edema (DME), choroidal neovascularization (CNV; e.g., myopic CNV), iris neovascularization, neovascular glaucoma, post-surgical fibrosis in glaucoma, proliferative vitreoretinopathy (PVR), optic disc neovascularization, corneal neovascularization, retinal neovascularization, vitreal neovascularization, panus, pterygium, vascular retinopathy, and diabetic retinopathies.

Pharmaceutical Formulations

The present invention includes methods in which the VEGF antagonist that is administered to the patient is contained within a pharmaceutical formulation. The pharmaceutical formulation may comprise the VEGF antagonist along with at least one inactive ingredient such as, e.g., a pharmaceutically acceptable carrier. Other agents may be incorporated into the pharmaceutical composition to provide improved transfer, delivery, tolerance, and the like. The term "pharmaceutically acceptable" means approved by a regulatory agency of the Federal or a state government or listed in the U.S. Pharmacopeia or other generally recognized pharmacopeia for use in animals, and more particularly, in humans. The term "carrier" refers to a diluent, adjuvant, excipient, or vehicle with which the antibody is administered. A multitude of appropriate formulations can be found in the formulary known to all pharmaceutical chemists: Remington's Pharmaceutical Sciences (15th ed, Mack Publishing Company, Easton, Pa., 1975), particularly Chapter 87 by Blaug, Seymour, therein. These formulations include, for example, powders, pastes, ointments, jellies, waxes, oils, lipids, lipid (cationic or anionic) containing vesicles (such as LIPOFECTIN™), DNA conjugates, anhydrous absorption pastes, oil-in-water and water-in-oil emulsions, emulsions carbowax (polyethylene glycols of various molecular weights), semi-solid gels, and semi-solid mixtures containing carbowax. Any of the foregoing mixtures may be appropriate in the context of the methods of the present invention, provided that the VEGF antagonist is not inactivated by the formulation and the formulation is physiologically compatible and tolerable with the route of administration. See also Powell et al. PDA (1998) J Pharm Sci Technol. 52:238-311 and the citations therein for additional information related to excipients and carriers well known to pharmaceutical chemists.

Pharmaceutical formulations useful for administration by injection in the context of the present invention may be prepared by dissolving, suspending or emulsifying a VEGF

6

antagonist in a sterile aqueous medium or an oily medium conventionally used for injections. As the aqueous medium for injections, there are, for example, physiological saline, an isotonic solution containing glucose and other auxiliary agents, etc., which may be used in combination with an appropriate solubilizing agent such as an alcohol (e.g., ethanol), a polyalcohol (e.g., propylene glycol, polyethylene glycol), a nonionic surfactant [e.g., polysorbate 80, HCO-50 (polyoxyethylene (50 mol) adduct of hydrogenated castor oil)], etc. As the oily medium, there may be employed, e.g., sesame oil, soybean oil, etc., which may be used in combination with a solubilizing agent such as benzyl benzoate, benzyl alcohol, etc. The injection thus prepared can be filled in an appropriate ampoule if desired.

Modes of Administration

The VEGF antagonist (or pharmaceutical formulation comprising the VEGF antagonist) may be administered to the patient by any known delivery system and/or administration method. In certain embodiments, the VEGF antagonist is administered to the patient by ocular, intraocular, intravitreal or subconjunctival injection. In other embodiments, the VEGF antagonist can be administered to the patient by topical administration, e.g., via eye drops or other liquid, gel, ointment or fluid which contains the VEGF antagonist and can be applied directly to the eye. Other possible routes of administration include, e.g., intradermal, intramuscular, intraperitoneal, intravenous, subcutaneous, intranasal, epidural, and oral.

Amount of VEGF Antagonist Administered

Each dose of VEGF antagonist administered to the patient over the course of the treatment regimen may contain the same, or substantially the same, amount of VEGF antagonist. Alternatively, the quantity of VEGF antagonist contained within the individual doses may vary over the course of the treatment regimen. For example, in certain embodiments, a first quantity of VEGF antagonist is administered in the initial dose, a second quantity of VEGF antagonist is administered in the secondary doses, and a third quantity of VEGF antagonist is administered in the tertiary doses. The present invention contemplates dosing schemes in which the quantity of VEGF antagonist contained within the individual doses increases over time (e.g., each subsequent dose contains more VEGF antagonist than the last), decreases over time (e.g., each subsequent dose contains less VEGF antagonist than the last), initially increases then decreases, initially decreases then increases, or remains the same throughout the course of the administration regimen.

The amount of VEGF antagonist administered to the patient in each dose is, in most cases, a therapeutically effective amount. As used herein, the phrase "therapeutically effective amount" means a dose of VEGF antagonist that results in a detectable improvement in one or more symptoms or indicia of an angiogenic eye disorder, or a dose of VEGF antagonist that inhibits, prevents, lessens, or delays the progression of an angiogenic eye disorder. In the case of an anti-VEGF antibody or a VEGF receptor-based chimeric molecule such as VEGFR1R2-FcΔC1(a), a therapeutically effective amount can be from about 0.05 mg to about 5 mg, e.g., about 0.05 mg, about 0.1 mg, about 0.15 mg, about 0.2 mg, about 0.25 mg, about 0.3 mg, about 0.35 mg, about 0.4 mg, about 0.45 mg, about 0.5 mg, about 0.55 mg, about 0.6 mg, about 0.65 mg, about 0.7 mg, about 0.75 mg, about 0.8 mg, about 0.85 mg, about 0.9 mg, about 1.0 mg, about 1.05 mg, about 1.1 mg, about 1.15 mg, about 1.2 mg, about 1.25 mg, about 1.3 mg, about 1.35 mg, about 1.4 mg, about 1.45 mg, about 1.5 mg, about 1.55 mg, about 1.6 mg, about 1.65 mg, about 1.7 mg, about 1.75 mg, about 1.8 mg, about 1.85

7

mg, about 1.9 mg, about 2.0 mg, about 2.05 mg, about 2.1 mg, about 2.15 mg, about 2.2 mg, about 2.25 mg, about 2.3 mg, about 2.35 mg, about 2.4 mg, about 2.45 mg, about 2.5 mg, about 2.55 mg, about 2.6 mg, about 2.65 mg, about 2.7 mg, about 2.75 mg, about 2.8 mg, about 2.85 mg, about 2.9 mg, about 3.0 mg, about 3.5 mg, about 4.0 mg, about 4.5 mg, or about 5.0 mg of the antibody or receptor-based chimeric molecule.

The amount of VEGF antagonist contained within the individual doses may be expressed in terms of milligrams of antibody per kilogram of patient body weight (i.e., mg/kg). For example, the VEGF antagonist may be administered to a patient at a dose of about 0.0001 to about 10 mg/kg of patient body weight.

Treatment Population and Efficacy

The methods of the present invention are useful for treating angiogenic eye disorders in patients that have been diagnosed with or are at risk of being afflicted with an angiogenic eye disorder. Generally, the methods of the present invention demonstrate efficacy within 104 weeks of the initiation of the treatment regimen (with the initial dose administered at “week 0”), e.g., by the end of week 16, by the end of week 24, by the end of week 32, by the end of week 40, by the end of week 48, by the end of week 56, etc. In the context of methods for treating angiogenic eye disorders such as AMD, CRVO, and DME, “efficacy” means that, from the initiation of treatment, the patient exhibits a loss of 15 or fewer letters on the Early Treatment Diabetic Retinopathy Study (ETDRS) visual acuity chart. In certain embodiments, “efficacy” means a gain of one or more (e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or more) letters on the ETDRS chart from the time of initiation of treatment.

EXAMPLES

The following examples are put forth so as to provide those of ordinary skill in the art with a complete disclosure and description of how to make and use the methods and compositions of the invention, and are not intended to limit the scope of what the inventors regard as their invention. Efforts have been made to ensure accuracy with respect to numbers used (e.g., amounts, temperature, etc.) but some experimental errors and deviations should be accounted for. Unless indicated otherwise, parts are parts by weight, molecular weight is average molecular weight, temperature is in degrees Centigrade, and pressure is at or near atmospheric. The exemplary VEGF antagonist used in all Examples set forth below is a dimeric molecule having two functional VEGF binding units. Each functional binding unit is comprised of Ig domain 2 from VEGFR1 fused to Ig domain 3 from VEGFR2, which in turn is fused to the hinge region of a human IgG1 Fc domain (VEGFR1R2-FcΔC1(a); encoded by SEQ ID NO:1). This VEGF antagonist is referred to in the examples below as “VEGFT”. For purposes of the following Examples, “monthly” dosing is equivalent to dosing once every four weeks.

Example 1

Phase I Clinical Trial of Intravitreally Administered VEGF Receptor-Based Chimeric Molecule (VEGFT) in Subjects with Neovascular AMD

In this Phase I study, 21 subjects with neovascular AMD received a single intravitreal (IVT) dose of VEGFT. Five groups of three subjects each received either 0.05, 0.15, 0.5, 2 or 4 mg of VEGFT, and a sixth group of six subjects received

8

1 mg. No serious adverse events related to the study drug, and no identifiable intraocular inflammation was reported. Preliminary results showed that, following injection of VEGFT, a rapid decrease in foveal thickness and macular volume was observed that was maintained through 6 weeks. At Day 43 across all dose groups, mean excess retinal thickness [excess retinal thickness=(retinal thickness–179μ)] on optical coherence tomography (OCT) was reduced from 119μ to 27μ as assessed by Fast Macular Scan and from 194μ to 60μ as assessed using a single Posterior Pole scan. The mean increase in best corrected visual acuity (BCVA) was 4.75 letters, and BCVA was stable or improved in 95% of subjects. In the 2 highest dose groups (2 and 4 mg), the mean increase in BCVA was 13.5 letters, with 3 of 6 subjects demonstrating improvement of >3 lines.

Example 2

Phase II Clinical Trial of Repeated Doses of Intravitreally Administered VEGF Receptor-Based Chimeric Molecule (VEGFT) in Subjects with Neovascular AMD

This study was a double-masked, randomized study of 3 doses (0.5, 2, and 4 mg) of VEGFT tested at 4-week and/or 12-week dosing intervals. There were 5 treatment arms in this study, as follows: 1) 0.5 mg every 4 weeks, 2) 0.5 mg every 12 weeks, 3) 2 mg every 4 weeks, 4) 2 mg every 12 weeks and 5) 4 mg every 12 weeks. Subjects were dosed at a fixed interval for the first 12 weeks, after which they were evaluated every 4 weeks for 9 months, during which additional doses were administered based on pre-specified criteria. All subjects were then followed for one year after their last dose of VEGFT. Preliminary data from a pre-planned interim analysis indicated that VEGFT met its primary endpoint of a statistically significant reduction in retinal thickness after 12 weeks compared with baseline (all groups combined, decrease of 135μ, p<0.0001). Mean change from baseline in visual acuity, a key secondary endpoint of the study, also demonstrated statistically significant improvement (all groups combined, increase of 5.9 letters, p<0.0001). Moreover, patients in the dose groups that received only a single dose, on average, demonstrated a decrease in excess retinal thickness (p<0.0001) and an increase in visual acuity (p=0.012) at 12 weeks. There were no drug-related serious adverse events, and treatment with the VEGF antagonists was generally well-tolerated. The most common adverse events were those typically associated with intravitreal injections.

Example 3

Phase I Clinical Trial of Systemically Administered VEGF Receptor-Based Chimeric Molecule (VEGFT) in Subjects with Neovascular AMD

This study was a placebo-controlled, sequential-group, dose-escalating safety, tolerability and bioeffect study of VEGFT by IV infusion in subjects with neovascular AMD. Groups of 8 subjects meeting eligibility criteria for subfoveal choroidal neovascularization (CNV) related to AMD were assigned to receive 4 IV injections of VEGFT or placebo at dose levels of 0.3, 1, or 3 mg/kg over an 8-week period.

Most adverse events that were attributed to VEGFT were mild to moderate in severity, but 2 of 5 subjects treated with 3 mg/kg experienced dose-limiting toxicity (DLT) (one with Grade 4 hypertension and one with Grade 2 proteinuria); therefore, all subjects in the 3 mg/kg dose group did not enter

the study. The mean percent changes in excess retinal thickness were: -12%, -10%, -66%, and -60% for the placebo, 0.3, 1, and 3 mg/kg dose groups at day 15 (ANOVA $p < 0.02$), and -5.6%, +47.1%, and -63.3% for the placebo, 0.3, and 1 mg/kg dose groups at day 71 (ANOVA $p < 0.02$). There was a numerical improvement in BCVA in the subjects treated with VEGFT. As would be expected in such a small study, the results were not statistically significant.

Example 4

Phase III Clinical Trials of the Efficacy, Safety, and Tolerability of Repeated Doses of Intravitreal VEGFT in Subjects with Neovascular Age-Related Macular Degeneration

A. Objectives, Hypotheses and Endpoints

Two parallel Phase III clinical trials were carried out to investigate the use of VEGFT to treat patients with the neovascular form of age-related macular degeneration (Study 1 and Study 2). The primary objective of these studies was to assess the efficacy of IVT administered VEGFT compared to ranibizumab (Lucentis®, Genentech, Inc.), in a non-inferiority paradigm, in preventing moderate vision loss in subjects with all subtypes of neovascular AMD.

The secondary objectives were (a) to assess the safety and tolerability of repeated IVT administration of VEGFT in subjects with all sub-types of neovascular AMD for periods up to 2 years; and (b) to assess the effect of repeated IVT administration of VEGFT on Vision-Related Quality of Life (QOL) in subjects with all sub-types of neovascular AMD.

The primary hypothesis of these studies was that the proportion of subjects treated with VEGFT with stable or improved BCVA (<15 letters lost) is similar to the proportion treated with ranibizumab who have stable or improved BCVA, thereby demonstrating non-inferiority.

The primary endpoint for these studies was the prevention of vision loss of greater than or equal to 15 letters on the ETDRS chart, compared to baseline, at 52 weeks. Secondary endpoints were as follows: (a) change from baseline to Week 52 in letter score on the ETDRS chart; (b) gain from baseline to Week 52 of 15 letters or more on the ETDRS chart; (c) change from baseline to Week 52 in total NEI VFQ-25 score; and (d) change from baseline to Week 52 in CNV area.

B. Study Design

For each study, subjects were randomly assigned in a 1:1:1 ratio to 1 of 4 dosing regimens: (1) 2 mg VEGFT administered every 4 weeks (2Q4); (2) 0.5 mg VEGFT administered every 4 weeks (0.5Q4); (3) 2 mg VEGFT administered every 4 weeks to week 8 and then every 8 weeks (with sham injection at the interim 4-week visits when study drug was not administered (2Q8); and (4) 0.5 mg ranibizumab administered every 4 weeks (RQ4). Subjects assigned to (2Q8) received the 2 mg injection every 4 weeks to week 8 and then a sham injection at interim 4-week visits (when study drug is not to be administered) during the first 52 weeks of the studies. (No sham injection were given at Week 52).

The study duration for each subject was scheduled to be 96 weeks plus the recruitment period. For the first 52 weeks (Year 1), subjects received an IVT or sham injection in the study eye every 4 weeks. (No sham injections were given at Week 52). During the second year of the study, subjects will be evaluated every 4 weeks and will receive IVT injection of study drug at intervals determined by specific dosing criteria, but at least every 12 weeks. (During the second year of the study, sham injections will not be given.) During this period, injections may be given as frequently as every 4 weeks, but no

less frequently than every 12 weeks, according to the following criteria: (i) increase in central retinal thickness of $>100 \mu\text{m}$ compared to the lowest previous value as measured by optical coherence tomography (OCT); or (ii) a loss from the best previous letter score of at least 5 ETDRS letters in conjunction with recurrent fluid as indicated by OCT; or (iii) new or persistent fluid as indicated by OCT; or (iv) new onset classic neovascularization, or new or persistent leak on fluorescein angiography (FA); or (v) new macular hemorrhage; or (vi) 12 weeks have elapsed since the previous injection. According to the present protocol, subjects must receive an injection at least every 12 weeks.

Subjects were evaluated at 4 weeks intervals for safety and best corrected visual acuity (BCVA) using the 4 meter ETDRS protocol. Quality of Life (QOL) was evaluated using the NEI VFQ-25 questionnaire. OCT and FA examinations were conducted periodically.

Approximately 1200 subjects were enrolled, with a target enrollment of 300 subjects per treatment arm.

To be eligible for this study, subjects were required to have subfoveal choroidal neovascularization (CNV) secondary to AMD. "Subfoveal" CNV was defined as the presence of subfoveal neovascularization, documented by FA, or presence of a lesion that is juxtafoveal in location angiographically but affects the fovea. Subject eligibility was confirmed based on angiographic criteria prior to randomization.

Only one eye was designated as the study eye. For subjects who met eligibility criteria in both eyes, the eye with the worse VA was selected as the study eye. If both eyes had equal VA, the eye with the clearest lens and ocular media and least amount of subfoveal scar or geographic atrophy was selected. If there was no objective basis for selecting the study eye, factors such as ocular dominance, other ocular pathology and subject preference were considered in making the selection.

Inclusion criteria for both studies were as follows: (i) signed Informed consent; (ii) at least 50 years of age; (iii) active primary subfoveal CNV lesions secondary to AMD, including juxtafoveal lesions that affect the fovea as evidenced by FA in the study eye; (iv) CNV at least 50% of total lesion size; (v) early treatment diabetic retinopathy study (ETDRS) best-corrected visual acuity of: 20/40 to 20/320 (letter score of 73 to 25) in the study eye; (vi) willing, committed, and able to return for all clinic visits and complete all study-related procedures; and (vii) able to read, understand and willing to sign the informed consent form (or, if unable to read due to visual impairment, be read to verbatim by the person administering the informed consent or a family member).

Exclusion criteria for both studies were as follows: 1. Any prior ocular (in the study eye) or systemic treatment or surgery for neovascular AMD except dietary supplements or vitamins. 2. Any prior or concomitant therapy with another investigational agent to treat neovascular AMD in the study eye, except dietary supplements or vitamins. 3. Prior treatment with anti-VEGF agents as follows: (a) Prior treatment with anti-VEGF therapy in the study eye was not allowed; (b) Prior treatment with anti-VEGF therapy in the fellow eye with an investigational agent (not FDA approved, e.g. bevacizumab) was allowed up to 3 months prior to first dose in the study, and such treatments were not allowed during the study. Prior treatment with an approved anti-VEGF therapy in the fellow eye was allowed; (c) Prior systemic anti-VEGF therapy, investigational or FDA/Health Canada approved, was only allowed up to 3 months prior to first dose, and was not allowed during the study. 4. Total lesion size >12 disc areas (30.5 mm², including blood, scars and neovascularization) as assessed by FA in the study eye. 5. Subretinal hem-

11

orrhage that is either 50% or more of the total lesion area, or if the blood is under the fovea and is 1 or more disc areas in size in the study eye. (If the blood is under the fovea, then the fovea must be surrounded 270 degrees by visible CNV.) 6. Scar or fibrosis, making up >50% of total lesion in the study eye. 7. Scar, fibrosis, or atrophy involving the center of the fovea. 8. Presence of retinal pigment epithelial tears or rips involving the macula in the study eye. 9. History of any vitreous hemorrhage within 4 weeks prior to Visit 1 in the study eye. 10. Presence of other causes of CNV, including pathologic myopia (spherical equivalent of -8 diopters or more negative, or axial length of 25 mm or more), ocular histoplasmosis syndrome, angioid streaks, choroidal rupture, or multifocal choroiditis in the study eye. 11. History or clinical evidence of diabetic retinopathy, diabetic macular edema or any other vascular disease affecting the retina, other than AMD, in either eye. 12. Prior vitrectomy in the study eye. 13. History of retinal detachment or treatment or surgery for retinal detachment in the study eye. 14. Any history of macular hole of stage 2 and above in the study eye. 15. Any intraocular or periocular surgery within 3 months of Day 1 on the study eye, except lid surgery, which may not have taken place within 1 month of day 1, as long as it was unlikely to interfere with the injection. 16. Prior trabeculectomy or other filtration surgery in the study eye. 17. Uncontrolled glaucoma (defined as intraocular pressure greater than or equal to 25 mm Hg despite treatment with anti-glaucoma medication) in the study eye. 18. Active intraocular inflammation in either eye. 19. Active ocular or periocular infection in either eye. 20. Any ocular or periocular infection within the last 2 weeks prior to Screening in either eye. 21. Any history of uveitis in either eye. 22. Active scleritis or episcleritis in either eye. 23. Presence or history of scleromalacia in either eye. 24. Aphakia or pseudophakia with absence of posterior capsule (unless it occurred as a result of a yttrium aluminum garnet [YAG] posterior capsulotomy) in the study eye. 25. Previous therapeutic radiation in the region of the study eye. 26. History of corneal transplant or corneal dystrophy in the study eye. 27. Significant media opacities, including cataract, in the study eye which might interfere with visual acuity, assessment of safety, or fundus photography. 28. Any concurrent intraocular condition in the study eye (e.g. cataract) that, in the opinion of the investigator, could require either medical or surgical intervention during the 96 week study period. 29. Any concurrent ocular condition in the study eye which, in the opinion of the investigator, could either increase the risk to the subject beyond what is to be expected from standard procedures of intraocular injection, or which otherwise may interfere with the injection procedure or with evaluation of efficacy or safety. 30. History of other disease, metabolic dysfunction, physical examination finding, or clinical laboratory finding giving reasonable suspicion of a disease or condition that contraindicates the use of an investigational drug or that might affect interpretation of the results of the study or render the subject at high risk for treatment complications. 31. Participation as a subject in any clinical study within the 12 weeks prior to Day 1. 32. Any systemic or ocular treatment with an investigational agent in the past 3 months prior to Day 1. 33. The use of long acting steroids, either systemically or intraocularly, in the 6 months prior to day 1. 34. Any history of allergy to povidone iodine. 35. Known serious allergy to the fluorescein sodium for injection in angiography. 36. Presence of any contraindications indicated in the FDA Approved label for ranibizumab (Lucentis®). 37. Females who were pregnant, breastfeeding, or of childbearing potential, unwilling to practice adequate contraception throughout the study. Adequate contraceptive measures

12

include oral contraceptives (stable use for 2 or more cycles prior to screening); IUD; Depo-Provera®; Norplant® System implants; bilateral tubal ligation; vasectomy; condom or diaphragm plus either contraceptive sponge, foam or jelly.

Subjects were not allowed to receive any standard or investigational agents for treatment of their AMD in the study eye other than their assigned study treatment with VEGFT or ranibizumab as specified in the protocol until they completed the Completion/Early Termination visit assessments. This includes medications administered locally (e.g., IVT, topical, juxtasceral or periorbital routes), as well as those administered systemically with the intent of treating the study and/or fellow eye.

The study procedures are summarized as follows:

Best Corrected Visual Acuity: Visual function of the study eye and the fellow eye were assessed using the ETDRS protocol (The Early Treatment Diabetic Retinopathy Study Group) at 4 meters. Visual Acuity examiners were certified to ensure consistent measurement of BCVA. The VA examiners were required to remain masked to treatment assignment.

Optical Coherence Tomography: Retinal and lesion characteristics were evaluated using OCT on the study eye. At the Screen Visit (Visit 1) images were captured and transmitted for both eyes. All OCT images were captured using the Zeiss Stratus OCT™ with software Version 3 or greater. OCT images were sent to an independent reading center where images were read by masked readers at visits where OCTs were required. All OCTs were electronically archived at the site as part of the source documentation. A subset of OCT images were read. OCT technicians were required to be certified by the reading center to ensure consistency and quality in image acquisition. Adequate efforts were made to ensure that OCT technicians at the site remained masked to treatment assignment.

Fundus Photography and Fluorescein Angiography (FA): The anatomical state of the retinal vasculature of the study eye was evaluated by funduscopic examination, fundus photography and FA. At the Screen Visit (Visit 1) funduscopic examination, fundus photography and FA were captured and transmitted for both eyes. Fundus and angiographic images were sent to an independent reading center where images were read by masked readers. The reading center confirmed subject eligibility based on angiographic criteria prior to randomization. All FAs and fundus photographs were archived at the site as part of the source documentation. Photographers were required to be certified by the reading center to ensure consistency and quality in image acquisition. Adequate efforts were made to ensure that all photographers at the site remain masked to treatment assignment.

Vision-Related Quality of Life: Vision-related QOL was assessed using the National Eye Institute 25-Item Visual Function Questionnaire (NEI VFQ-25) in the interviewer-administered format. NEI VFQ-25 was administered by certified personnel at a contracted call center. At the screening visit, the sites assisted the subject and initiated the first call to the call center to collect all of the subject's contact information and to complete the first NEI VFQ-25 on the phone prior to randomization and IVT injection. For all subsequent visits, the call center called the subject on the phone, prior to IVT injection, to complete the questionnaire.

Intraocular Pressure: Intraocular pressure (IOP) of the study eye was measured using applanation tonometry or Tonopen. The same method of IOP measurement was used in each subject throughout the study.

C. Results Summary (52 Week Data)

The primary endpoint (prevention of moderate or severe vision loss as defined above) was met for all three VEGFT

13

groups (2Q4, 0.5Q4 and 2Q8) in this study. The results from both studies are summarized in Table 1.

TABLE 1

	Ranibizumab 0.5 mg monthly (RQ4)	VEGFT 0.5 mg monthly (0.5Q4)	VEGFT 2 mg monthly (2Q4)	VEGFT 2 mg every 8 weeks ^[a] (2Q8)
Maintenance of vision* (% patients losing <15 letters) at week 52 versus baseline				
Study 1	94.4%	95.9%**	95.1%**	95.1%**
Study 2	94.4%	96.3%**	95.6%**	95.6%**
Mean improvement in vision* (letters) at 52 weeks versus baseline (p-value vs RQ4)***				
Study 1	8.1	6.9 (NS)	10.9 (p < 0.01)	7.9 (NS)
Study 2	9.4	9.7 (NS)	7.6 (NS)	8.9 (NS)

^[a]Following three initial monthly doses

*Visual acuity was measured as the total number of letters read correctly on the Early Treatment Diabetic Retinopathy Study (ETDRS) eye chart.

**Statistically non-inferior based on a non-inferiority margin of 10%, using confidence interval approach (95.1% and 95% for Study 1 and Study 2, respectively)

***Test for superiority

NS = non-significant

In Study 1, patients receiving VEGFT 2 mg monthly (2Q4) achieved a statistically significant greater mean improvement in visual acuity at week 52 versus baseline (secondary endpoint), compared to ranibizumab 0.5 mg monthly (RQ4); patients receiving VEGFT 2 mg monthly on average gained 10.9 letters, compared to a mean 8.1 letter gain with ranibizumab 0.5 mg dosed every month (p<0.01). All other dose groups of VEGFT in Study 1 and all dose groups in Study 2 were not statistically different from ranibizumab in this secondary endpoint.

A generally favorable safety profile was observed for both VEGFT and ranibizumab. The incidence of ocular treatment emergent adverse events was balanced across all four treatment groups in both studies, with the most frequent events associated with the injection procedure, the underlying disease, and/or the aging process. The most frequent ocular adverse events were conjunctival hemorrhage, macular degeneration, eye pain, retinal hemorrhage, and vitreous floaters. The most frequent serious non-ocular adverse events were typical of those reported in this elderly population who receive intravitreal treatment for wet AMD; the most frequently reported events were falls, pneumonia, myocardial infarction, atrial fibrillation, breast cancer, and acute coronary syndrome. There were no notable differences among the study arms.

Example 5

Phase II Clinical Trial of VEGFT in Subjects with Diabetic Macular Edema (DME)

In this study, 221 patients with clinically significant DME with central macular involvement were randomized, and 219 patients were treated with balanced distribution over five groups. The control group received macular laser therapy at baseline, and patients were eligible for repeat laser treatments, but no more frequently than at 16 week intervals. The remaining four groups received VEGFT by intravitreal injection as follows: Two groups received 0.5 or 2 mg of VEGFT once every four weeks throughout the 12-month dosing period (0.5Q4 and 2Q4, respectively). Two groups received three initial doses of 2 mg VEGFT once every four weeks (i.e., at baseline, and weeks 4 and 8), followed through week 52 by either once every 8 weeks dosing (2Q8) or as needed

14

dosing with very strict repeat dosing criteria (PRN). Mean gains in visual acuity versus baseline were as shown in Table 2:

TABLE 2

	n	Mean change in visual acuity at week 24 versus baseline (letters)	Mean change in visual acuity at week 52 versus baseline (letters)
Laser	44	2.5	-1.3
VEGFT 0.5 mg monthly (0.5Q4)	44	8.6**	11.0**
VEGFT 2 mg monthly (2Q4)	44	11.4**	13.1**
VEGFT 2 mg every 8 weeks ^[a] (2Q8)	42	8.5**	9.7**
VEGFT 2 mg as needed ^[a] (PRN)	45	10.3**	12.0**

^[a]Following three initial monthly doses

**p < 0.01 versus laser

In this study, the visual acuity gains achieved with VEGFT administration at week 24 were maintained or numerically improved up to completion of the study at week 52 in all VEGFT study groups, including 2 mg dosed every other month

As demonstrated in the foregoing Examples, the administration of VEGFT to patients suffering from angiogenic eye disorders (e.g., AMD and DME) at a frequency of once every 8 weeks, following a single initial dose and two secondary doses administered four weeks apart, resulted in significant prevention of moderate or severe vision loss or improvements in visual acuity.

Example 6

A Randomized, Multicenter, Double-Masked Trial in Treatment Naïve Patients with Macular Edema Secondary to CRVO

In this randomized, double-masked, Phase 3 study, patients received 6 monthly injections of either 2 mg intravitreal VEGFT (114 patients) or sham injections (73 patients). From Week 24 to Week 52, all patients received 2 mg VEGFT as-needed (PRN) according to retreatment criteria. Thus, “sham-treated patients” means patients who received sham injections once every four weeks from Week 0 through Week 20, followed by intravitreal VEGFT as needed from Week 24 through Week 52. “VEGFT-treated patients” means patients who received VEGFT intravitreal injections once every four weeks from Week 0 through Week 20, followed by intravitreal VEGFT as needed from Week 24 through Week 52. The primary endpoint was the proportion of patients who gained ≥15 ETDRS letters from baseline at Week 24. Secondary visual, anatomic, and Quality of Life NEI VFQ-25 outcomes at Weeks 24 and 52 were also evaluated.

At Week 24, 56.1% of VEGFT-treated patients gained >15 ETDRS letters from baseline vs 12.3% of sham-treated patients (P<0.0001). Similarly, at Week 52, 55.3% of VEGFT-treated patients gained ≥15 letters vs 30.1% of sham-treated patients (P<0.01). At Week 52, VEGFT-treated patients gained a mean of 16.2 letters vs 3.8 letters for sham-treated patients (P<0.001). Mean number of injections was 2.7 for VEGFT-treated patients vs 3.9 for sham-treated patients. Mean change in central retinal thickness was -413.0 μm for VEGFT-treated patients vs -381.8 μm for sham-treated patients. The proportion of patients with ocular

15

neovascularization at Week 24 were 0% for VEGFT-treated patients and 6.8% for sham-treated patients, respectively; at Week 52 after receiving VEGFT PRN, proportions were 0% and 6.8% for VEGFT-treated and sham-treated. At Week 24, the mean change from baseline in the VFQ-25 total score was 7.2 vs 0.7 for the VEGFT-treated and sham-treated groups; at Week 52, the scores were 7.5 vs 5.1 for the VEGFT-treated and sham-treated groups.

This Example confirms that dosing monthly with 2 mg intravitreal VEGFT injection resulted in a statistically significant improvement in visual acuity at Week 24 that was maintained through Week 52 with PRN dosing compared with sham PRN treatment. VEGFT was generally well tolerated and had a generally favorable safety profile.

Example 7

Dosing Regimens

Specific, non-limiting examples of dosing regimens within the scope of the present invention are as follows:

VEGFT 2 mg (0.05 mL) administered by intravitreal injection once every 4 weeks (monthly).

VEGFT 2 mg (0.5 mL) administered by intravitreal injection once every 4 weeks for the first 8 weeks, followed by 2 mg (0.05 mL) via intravitreal injection once every 8 weeks.

VEGFT 2 mg (0.5 mL) administered by intravitreal injection once every 4 weeks for the first 8 weeks, followed by 2 mg (0.05 mL) via intravitreal injection on a less frequent basis based on visual and/or anatomical outcomes (as assessed by a physician or other qualified medical professional).

VEGFT 2 mg (0.5 mL) administered by intravitreal injection once every 4 weeks for the first 8 weeks, followed by 2 mg (0.05 mL) via intravitreal injection administered pro re nata (PRN) based on visual and/or anatomical outcomes (as assessed by a physician or other qualified medical professional).

VEGFT 2 mg (0.5 mL) administered by intravitreal injection once every 4 weeks for the first 12 weeks, followed by 2 mg (0.05 mL) via intravitreal injection once every 8 weeks.

VEGFT 2 mg (0.5 mL) administered by intravitreal injection once every 4 weeks for the first 12 weeks, followed by 2 mg (0.05 mL) via intravitreal injection on a less frequent basis based on visual and/or anatomical outcomes (as assessed by a physician or other qualified medical professional).

VEGFT 2 mg (0.5 mL) administered by intravitreal injection once every 4 weeks for the first 12 weeks, followed by 2 mg (0.05 mL) via intravitreal injection administered pro re nata (PRN) based on visual and/or anatomical outcomes (as assessed by a physician or other qualified medical professional).

VEGFT 2 mg (0.5 mL) administered by intravitreal injection once every 4 weeks for the first 16 weeks, followed by 2 mg (0.05 mL) via intravitreal injection once every 8 weeks.

VEGFT 2 mg (0.5 mL) administered by intravitreal injection once every 4 weeks for the first 16 weeks, followed by 2 mg (0.05 mL) via intravitreal injection on a less frequent basis based on visual and/or anatomical outcomes (as assessed by a physician or other qualified medical professional).

VEGFT 2 mg (0.5 mL) administered by intravitreal injection once every 4 weeks for the first 16 weeks, followed by 2 mg (0.05 mL) via intravitreal injection administered pro re nata (PRN) based on visual and/or anatomical outcomes (as assessed by a physician or other qualified medical professional).

16

VEGFT 2 mg (0.5 mL) administered by intravitreal injection once every 4 weeks for the first 20 weeks, followed by 2 mg (0.05 mL) via intravitreal injection once every 8 weeks.

VEGFT 2 mg (0.5 mL) administered by intravitreal injection once every 4 weeks for the first 20 weeks, followed by 2 mg (0.05 mL) via intravitreal injection on a less frequent basis based on visual and/or anatomical outcomes (as assessed by a physician or other qualified medical professional).

VEGFT 2 mg (0.5 mL) administered by intravitreal injection once every 4 weeks for the first 20 weeks, followed by 2 mg (0.05 mL) via intravitreal injection administered pro re nata (PRN) based on visual and/or anatomical outcomes (as assessed by a physician or other qualified medical professional).

VEGFT 2 mg (0.5 mL) administered by intravitreal injection once every 4 weeks for the first 24 weeks, followed by 2 mg (0.05 mL) via intravitreal injection once every 8 weeks.

VEGFT 2 mg (0.5 mL) administered by intravitreal injection once every 4 weeks for the first 24 weeks, followed by 2 mg (0.05 mL) via intravitreal injection on a less frequent basis based on visual and/or anatomical outcomes (as assessed by a physician or other qualified medical professional).

VEGFT 2 mg (0.5 mL) administered by intravitreal injection once every 4 weeks for the first 24 weeks, followed by 2 mg (0.05 mL) via intravitreal injection administered pro re nata (PRN) based on visual and/or anatomical outcomes (as assessed by a physician or other qualified medical professional).

VEGFT 2 mg (0.5 mL) administered by intravitreal injection once every 4 weeks for the first 28 weeks, followed by 2 mg (0.05 mL) via intravitreal injection once every 8 weeks.

VEGFT 2 mg (0.5 mL) administered by intravitreal injection once every 4 weeks for the first 28 weeks, followed by 2 mg (0.05 mL) via intravitreal injection on a less frequent basis based on visual and/or anatomical outcomes (as assessed by a physician or other qualified medical professional).

VEGFT 2 mg (0.5 mL) administered by intravitreal injection once every 4 weeks for the first 28 weeks, followed by 2 mg (0.05 mL) via intravitreal injection administered pro re nata (PRN) based on visual and/or anatomical outcomes (as assessed by a physician or other qualified medical professional).

VEGFT 2 mg (0.05 mL) administered by intravitreal injection as a single initial dose, followed by additional doses administered pro re nata (PRN) based on visual and/or anatomical outcomes (as assessed by a physician or other qualified medical professional).

Variations on the above-described dosing regimens would be appreciated by persons of ordinary skill in the art and are also within the scope of the present invention. For example, the amount of VEGFT and/or volume of formulation administered to a patient may be varied based on patient characteristics, severity of disease, and other diagnostic assessments by a physician or other qualified medical professional.

Any of the foregoing administration regimens may be used for the treatment of, e.g., age-related macular degeneration (e.g., wet AMD, exudative AMD, etc.), retinal vein occlusion (RVO), central retinal vein occlusion (CRVO; e.g., macular edema following CRVO), branch retinal vein occlusion (BRVO), diabetic macular edema (DME), choroidal neovascularization (CNV; e.g., myopic CNV), iris neovascularization, neovascular glaucoma, post-surgical fibrosis in glaucoma, proliferative vitreoretinopathy (PVR), optic disc neovascularization, corneal neovascularization, retinal neovascularization, vitreal neovascularization, pannus, pterygium, vascular retinopathy, etc.

Sequences

SEQ ID NO: 1

(DNA sequence having 1377 nucleotides):
 ATGGTCAGCTACTGGACACCGGGTCTGTGTGCGCTGCTCAGCTGTCTGCTTCTCAC
 AGGATCTAGTTCCGGAAGTGATACCGGTAGACCTTTCGTAGAGATGTACAGTGAAATCCCCGA
 AATTATACACATGACTGAAGGAAGGAGCTCGTCATTCCCTGCCGGGTACGTACCTAACAT
 CACTGTTACTTTAAAAAGTTTCCACTTGACACTTTGATCCCTGATGGAACGCATAATCTGG
 GACAGTAGAAGGGCTTCATCATATCAATGCAACGTACAAAGAAATAGGGCTTCTGACCTGT
 GAAGCAACAGTCAATGGGCATTTGTATAAGACAACTATCTCACATCGACAAACCAATACAA
 TCATAGATGTGTTCTGAGTCCGTCTCATGGAATTGAACTATCTGTTGGAGAAAAGCTTGTCTT
 AAATTGTACAGCAAGAACTGAACTAAATGTGGGGATTGACTTCAACTGGGAATACCCTTCTTCG
 AAGCATCAGCATAAGAACTTGTAAACCGAGACCTAAAAACCCAGTCTGGGAGTGAGATGAAG
 AAATTTTGTAGACCTTAAGTATAGATGGTGAACCCGAGTGACCAAGGATTGTACACCTGTG
 CAGCATCCAGTGGGCTGATGACCAAGAAGAACAGCACATTGTGAGGTCATGAAAAGGACA
 AAATCACACATGCCACCGTGCCAGCACCTGAACCTCTGGGGGACCGTCAGTCTTCTCT
 TCCCCCAAAACCCAGGACACCTCATGATCTCCCGACCCCTGAGGTACATGCGTGGTG
 GTGGACGTGAGCCACGAAGACCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGGAGGT
 GCATAATGCCAAGACAAAGCCGCGGGAGGAGCAGTACAAAGCAGTACCGTGTGGTCAGCG
 TCCTCACCGTCTGCACAGGACTGGCTGAATGGCAAGGAGTACAAGTGCAAGGTCTCCAAC
 AAGCCCTCCAGCCCCATCGAGAAAACCATCTCCAAAGCCAAAGGGCAGCCCCGAGAACC
 ACAGGTGTACACCTGCCCCATCCCGGATGAGCTGACCAAGAACCAGGTGAGCTGACCT
 GCCTGGTCAAGGCTTCTATCCAGCGACATCGCGTGGAGTGGGAGCAATGGGACGCG
 GAGAACAACCTACAAGACCACGCTCCCGTGTGACTCCGACGGCTCTTCTCTCTACAGC
 AAGTTCACCGTGGACAAGAGCAGGTGGCAGCAGGGGAACGTCTTCTCATGTCCGTGATGCA
 TGAGGCTCTGCACAACCACTACACGAGAAGAGCCTCTCCCTGTCTCCGGTAAATGA

SEQ ID NO: 2

(polypeptide sequence having 458 amino acids):
 MVSYWDTGVLSCALLSCLLLTGSSSGSDTGRPFVEMYSEIPEIIHMTGRELVIPCRVTSFNITVTLK
 KPPLDTLIPDGKRIIWDNRKFIISNATYKEIGLLTCEATVNGHLYKTNLYTHRQNTIIDVVLSPSHGI
 ELSVGEKLVNCTARTELVNVDGIDFNWEYPSKQHKLVNRDLKTQSGSEMKKFLSTLTIDGVTRS
 DQGLYTCASSGLMTKKNSTFVRVHEKDKHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEV
 TCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKV
 SNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPEN
 NYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFSQVMHEALHNHYTQKSLSLSPGK

The present invention is not to be limited in scope by the specific embodiments described herein. Indeed, various modifications of the invention in addition to those described herein will become apparent to those skilled in the art from

the foregoing description and the accompanying figures. Such modifications are intended to fall within the scope of the appended claims.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 2

<210> SEQ ID NO 1

<211> LENGTH: 1377

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 1

```

atggtcagct actgggacac cggggtcctg ctgtgcgcgc tgcacagctg tctgtttctc   60
acaggatcta gttccggaag tgataccggt agacctttcg tagagatgta cagtgaatac   120
cccgaaatta tacacatgac tgaaggaagg gagctcgtea ttccctgccg ggttacgtea   180
cctaacaatca ctgttacttt aaaaaagttt ccacttgaca ctttgatccc tgatggaaaa   240
cgcataatct gggacagtag aaagggcttc atcatatcaa atgcaacgta caaagaaata   300
gggcttctga cctgtgaagc aacagtcaat gggcatttgt ataagacaaa ctatctcaca   360
catcgacaaa ccaatacaat catagatgtg gttctgagtc cgtctcatgg aattgaacta   420
tctgttgagg aaaagcttgt cttaaatgtt acagcaagaa ctgaactaaa tgtggggatt   480
gacttcaact gggaataccc ttcttcgaag catcagcata agaaacttgt aaaccgagac   540
ctaaaaaccc agtctgggag tgagatgaag aaatttttga gcaccttaac tatagatggt   600
gtaacccgga gtgaccaagg attgtacacc tgtgcagcat ccagtgggct gatgaccaag   660
aagaacagca catttgtcag ggtccatgaa aaggacaaaa ctcacacatg cccaccgtgc   720
ccagcacctg aactcctggg gggaccgtca gtcttcctct tcccccaaa acccaaggac   780
accctcatga tctcccgga cctgaggtc acatgcgtgg tggaggacgt gagccacgaa   840
gaccctgagg tcaagttaa ctggtacgtg gacggcgtgg aggtgcataa tgccaagaca   900
aagccgcggg aggagcagta caacagcacg taccgtgtgg tcagcgtcct caccgtcctg   960
caccaggact ggctgaatgg caaggagtac aagtgcaggg tctccaacaa agccctccca  1020
gcccccatcg agaaaacccat ctccaaagcc aaagggcagc cccgagaacc acaggtgtac  1080
accctgcccc catcccgga tgagctgacc aagaaccagg tcagcctgac ctgcctggtc  1140
aaaggcttct atcccagca catcgccgtg gagtgggaga gcaatgggca gccggagaa  1200
aactacaaga ccagcctcc cgtgctggac tccgacggct ccttcttct ctacagcaag  1260
ctcaccgtgg acaagagcag gtggcagcag gggaacgtct tctcatgctc cgtgatgcac  1320
gaggctctgc acaaccacta cagcagaag agcctctccc tgtctccggg taaatga  1377

```

<210> SEQ ID NO 2

<211> LENGTH: 458

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 2

```

Met Val Ser Tyr Trp Asp Thr Gly Val Leu Leu Cys Ala Leu Leu Ser
 1             5             10             15
Cys Leu Leu Leu Thr Gly Ser Ser Ser Gly Ser Asp Thr Gly Arg Pro
 20             25             30
Phe Val Glu Met Tyr Ser Glu Ile Pro Glu Ile Ile His Met Thr Glu
 35             40             45
Gly Arg Glu Leu Val Ile Pro Cys Arg Val Thr Ser Pro Asn Ile Thr
 50             55             60
Val Thr Leu Lys Lys Phe Pro Leu Asp Thr Leu Ile Pro Asp Gly Lys
 65             70             75             80
Arg Ile Ile Trp Asp Ser Arg Lys Gly Phe Ile Ile Ser Asn Ala Thr
 85             90             95

```

Tyr	Lys	Glu	Ile	Gly	Leu	Leu	Thr	Cys	Glu	Ala	Thr	Val	Asn	Gly	His
			100					105					110		
Leu	Tyr	Lys	Thr	Asn	Tyr	Leu	Thr	His	Arg	Gln	Thr	Asn	Thr	Ile	Ile
			115					120					125		
Asp	Val	Val	Leu	Ser	Pro	Ser	His	Gly	Ile	Glu	Leu	Ser	Val	Gly	Glu
			130					135					140		
Lys	Leu	Val	Leu	Asn	Cys	Thr	Ala	Arg	Thr	Glu	Leu	Asn	Val	Gly	Ile
			145					150					155		
Asp	Phe	Asn	Trp	Glu	Tyr	Pro	Ser	Ser	Lys	His	Gln	His	Lys	Lys	Leu
			165					170					175		
Val	Asn	Arg	Asp	Leu	Lys	Thr	Gln	Ser	Gly	Ser	Glu	Met	Lys	Lys	Phe
			180					185					190		
Leu	Ser	Thr	Leu	Thr	Ile	Asp	Gly	Val	Thr	Arg	Ser	Asp	Gln	Gly	Leu
			195					200					205		
Tyr	Thr	Cys	Ala	Ala	Ser	Ser	Gly	Leu	Met	Thr	Lys	Lys	Asn	Ser	Thr
			210					215					220		
Phe	Val	Arg	Val	His	Glu	Lys	Asp	Lys	Thr	His	Thr	Cys	Pro	Pro	Cys
			225					230					235		
Pro	Ala	Pro	Glu	Leu	Leu	Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro	Pro
			245					250					255		
Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys
			260					265					270		
Val	Val	Val	Asp	Val	Ser	His	Glu	Asp	Pro	Glu	Val	Lys	Phe	Asn	Trp
			275					280					285		
Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu
			290					295					300		
Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val	Leu
			305					310					315		
His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn
			325					330					335		
Lys	Ala	Leu	Pro	Ala	Pro	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly
			340					345					350		
Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser	Arg	Asp	Glu
			355					360					365		
Leu	Thr	Lys	Asn	Gln	Val	Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr
			370					375					380		
Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn
			385					390					395		
Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe	Phe
			405					410					415		
Leu	Tyr	Ser	Lys	Leu	Thr	Val	Asp	Lys	Ser	Arg	Trp	Gln	Gln	Gly	Asn
			420					425					430		
Val	Phe	Ser	Cys	Ser	Val	Met	His	Glu	Ala	Leu	His	Asn	His	Tyr	Thr
			435					440					445		
Gln	Lys	Ser	Leu	Ser	Leu	Ser	Pro	Gly	Lys						
			450					455							

23

What is claimed is:

1. A method for treating an angiogenic eye disorder in a patient, said method comprising sequentially administering to the patient a single initial dose of a VEGF antagonist, followed by one or more secondary doses of the VEGF antagonist, followed by one or more tertiary doses of the VEGF antagonist;

wherein each secondary dose is administered 2 to 4 weeks after the immediately preceding dose; and

wherein each tertiary dose is administered at least 8 weeks after the immediately preceding dose;

wherein the VEGF antagonist is a VEGF receptor-based chimeric molecule comprising (1) a VEGFR1 component comprising amino acids 27 to 129 of SEQ ID NO:2; (2) a VEGFR2 component comprising amino acids 130-231 of SEQ ID NO:2; and (3) a multimerization component comprising amino acids 232-457 of SEQ ID NO:2.

2. The method of claim 1, wherein only a single secondary dose is administered to the patient, and wherein the single secondary dose is administered 4 weeks after the initial dose of the VEGF antagonist.

3. The method of claim 1, wherein only two secondary doses are administered to the patient, and wherein each secondary dose is administered 4 weeks after the immediately preceding dose.

4. The method of claim 3, wherein each tertiary dose is administered 8 weeks after the immediately preceding dose.

5. The method of claim 1, wherein at least 5 tertiary doses of the VEGF antagonist are administered to the patient, and wherein the first four tertiary doses are administered 8 weeks after the immediately preceding dose, and wherein each subsequent tertiary dose is administered 8 or 12 weeks after the immediately preceding dose.

6. The method of claim 1, wherein the angiogenic eye disorder is selected from the group consisting of: age related macular degeneration, diabetic retinopathy, diabetic macular edema, central retinal vein occlusion, branch retinal vein occlusion, and corneal neovascularization.

7. The method of claim 6, wherein the angiogenic eye disorder is age related macular degeneration.

8. The method of claim 1, wherein all doses of the VEGF antagonist are administered to the patient by topical administration or by intraocular administration.

9. The method of claim 8, wherein all doses of the VEGF antagonist are administered to the patient by intraocular administration.

10. The method of claim 9, wherein the intraocular administration is intravitreal administration.

11. The method of claim 10, wherein all doses of the VEGF antagonist comprise from about 0.5 mg to about 2 mg of the VEGF antagonist.

12. The method of claim 11, wherein all doses of the VEGF antagonist comprise 0.5 mg of the VEGF antagonist.

24

13. The method of claim 11, wherein all doses of the VEGF antagonist comprise 2 mg of the VEGF antagonist.

14. A method for treating an angiogenic eye disorder in a patient, said method comprising sequentially administering to the patient a single initial dose of a VEGF antagonist, followed by one or more secondary doses of the VEGF antagonist, followed by one or more tertiary doses of the VEGF antagonist;

wherein each secondary dose is administered 2 to 4 weeks after the immediately preceding dose; and

wherein each tertiary dose is administered at least 8 weeks after the immediately preceding dose;

wherein the VEGF antagonist is a VEGF receptor-based chimeric molecule comprising VEGFR1R2-FcΔC1(a) encoded by the nucleic acid sequence of SEQ ID NO:1.

15. The method of claim 14, wherein only a single secondary dose is administered to the patient, and wherein the single secondary dose is administered 4 weeks after the initial dose of the VEGF antagonist.

16. The method of claim 14, wherein only two secondary doses are administered to the patient, and wherein each secondary dose is administered 4 weeks after the immediately preceding dose.

17. The method of claim 16, wherein each tertiary dose is administered 8 weeks after the immediately preceding dose.

18. The method of claim 17, wherein the angiogenic eye disorder is age related macular degeneration.

19. The method of claim 14, wherein at least 5 tertiary doses of the VEGF antagonist are administered to the patient, and wherein the first four tertiary doses are administered 8 weeks after the immediately preceding dose, and wherein each subsequent tertiary dose is administered 8 or 12 weeks after the immediately preceding dose.

20. The method of claim 14, wherein the angiogenic eye disorder is selected from the group consisting of: age related macular degeneration, diabetic retinopathy, diabetic macular edema, central retinal vein occlusion, branch retinal vein occlusion, and corneal neovascularization.

21. The method of claim 14, wherein all doses of the VEGF antagonist are administered to the patient by topical administration or by intraocular administration.

22. The method of claim 21, wherein all doses of the VEGF antagonist are administered to the patient by intraocular administration.

23. The method of claim 22, wherein the intraocular administration is intravitreal administration.

24. The method of claim 23, wherein all doses of the VEGF antagonist comprise from about 0.5 mg to about 2 mg of the VEGF antagonist.

25. The method of claim 24, wherein all doses of the VEGF antagonist comprise 0.5 mg of the VEGF antagonist.

26. The method of claim 24, wherein all doses of the VEGF antagonist comprise 2 mg of the VEGF antagonist.

* * * * *

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 9,254,338 B2
APPLICATION NO. : 13/940370
DATED : February 9, 2016
INVENTOR(S) : Yancopoulos

Page 1 of 1

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

In the Specification

Column 8, Line 15, delete “of >3 lines” and insert --of ≥ 3 lines--.

Column 14, Line 57, delete “gained> 15 ETDRS” and insert --gained ≥ 15 ETDRS--.

Signed and Sealed this
Twenty-ninth Day of March, 2022



Drew Hirshfeld
*Performing the Functions and Duties of the
Under Secretary of Commerce for Intellectual Property and
Director of the United States Patent and Trademark Office*