

Antivascular endothelial growth factor therapy for neovascular age-related macular degeneration

Thomas A. Ciulla^a and Philip J. Rosenfeld^b

^aVitreoretinal Service, Midwest Eye Institute, Indianapolis, Indiana and ^bDepartment of Ophthalmology, Bascom Palmer Eye Institute, University of Miami Miller School of Medicine, Miami, Florida, USA

Correspondence to Thomas A. Ciulla, MD, Vitreoretinal Service, Midwest Eye Institute, 201 Pennsylvania Parkway, Indianapolis, IN 46280, USA
Tel: +1 317 506 0334; fax: +1 317 817 1898;
e-mail: thomasciulla@yahoo.com

Current Opinion in Ophthalmology 2009, 20:158–165

Purpose of review

The most important recent advance in the treatment of neovascular age-related macular degeneration (AMD) is the development of antivascular endothelial growth factor (anti-VEGF) therapeutic agents that preserve and improve visual acuity by arresting choroidal neovascular growth and reducing vascular permeability. This review describes the current literature on the use of this therapeutic approach in the management of neovascular AMD.

Recent findings

Two anti-VEGF agents, pegaptanib sodium and ranibizumab, are currently approved by the United States Food and Drug Administration for the treatment of neovascular AMD. In addition, off-label use of a third anti-VEGF agent, bevacizumab, as a treatment option for neovascular AMD has become common worldwide. Other anti-VEGF agent strategies that have shown efficacy include small interfering RNA agents to silence the VEGF gene and receptor and the fusion protein VEGF trap.

Summary

The accumulation of preclinical and clinical evidence implicating VEGF-A in the pathogenesis of neovascular AMD has provided a strong rationale for the development of anti-VEGF agents for this disease. Anti-VEGF therapies have been used successfully in the clinic, encouraging their use in the treatment of other neovascular eye diseases.

Keywords

bevacizumab, macular degeneration, neovascularization, pegaptanib sodium, ranibizumab

Curr Opin Ophthalmol 20:158–165
© 2009 Wolters Kluwer Health | Lippincott Williams & Wilkins
1040-8738

Introduction

Ocular diseases involving neovascularization, including neovascular age-related macular degeneration (AMD), diabetic retinopathy, diabetic macular edema (DME), and retinal vein occlusion (RVO), are the primary causes of clinically significant vision loss in the developed world [1–3], particularly among the working and elderly populations. In the United States, neovascular AMD affects 1.75 million individuals aged more than 40 years and is the leading cause of blindness in those over 65 years [4]. It is characterized by the growth of choroidal blood vessels through Bruch's membrane into the subretinal pigment epithelial (RPE) space, usually leading to accumulation of fluid in the sub-RPE space and detachment of the RPE [5].

Inhibition of angiogenesis

Normal and pathologic angiogenesis is a complex balance of positive and negative regulators, and vascular endothelial growth factor-A (VEGF-A; also referred to as VEGF) is one of the most important positive regulators of angiogenesis [6] and vascular permeability [7,8].

VEGF-A has been implicated in the pathogenesis of a variety of disorders including neovascular AMD, proliferative diabetic retinopathy (PDR), and other neovascular eye diseases, as well as in tumorigenesis. Therefore, VEGF inhibition may be expected to lead not only to inhibition of further angiogenesis but also to regression of newly formed blood vessels.

Therapeutic agents that inhibit VEGF-A – pegaptanib sodium (Macugen; OSI/Eyetech, Inc., Melville, New York, USA) and ranibizumab (Lucentis; Genentech, Inc., South San Francisco, California, USA) – are Food and Drug Administration (FDA) approved for the treatment of choroidal neovascularization (CNV) secondary to AMD. Of the two agents, ranibizumab offers substantial clinical benefit in neovascular AMD. A third anti-VEGF agent, bevacizumab, has been used off label for neovascular AMD and other exudative retinal diseases. These anti-VEGF agents, as well as others in clinical development, have great potential to treat eye diseases characterized by neovascularization. Here, we present an overview of the current knowledge on anti-VEGF therapies in neovascular AMD.

Vascular endothelial growth factor-A in vascular permeability, inflammation, and ocular disease

In the various pathologic conditions in which it is implicated, overexpression of VEGF-A has been found to promote angiogenesis by inducing endothelial cell proliferation, migration, and survival [9].

Vascular endothelial growth factor-A isoforms

VEGF-A is a member of the VEGF family of growth factors that also includes VEGF-B, VEGF-C, VEGF-D, and placental growth factor (PlGF), which have different binding affinities for the three VEGF receptors, VEGFR1, VEGFR2, and VEGFR3 [10,11]. VEGF-A also binds to neuropilin-1, a membrane protein on developing neurons that plays a role in embryonic neural blood vessel formation as well as neural tip guidance [12,13]. Alternative RNA splicing of the human *VEGF-A* gene results in the formation of four major isoforms (VEGF₁₂₁, VEGF₁₆₅, VEGF₁₈₉, and VEGF₂₀₆) and at least five minor isoforms (VEGF₁₄₅, VEGF₁₄₈, VEGF₁₆₂, VEGF_{165b}, and VEGF₁₈₃) [14–16]. VEGF₁₆₅ and VEGF₁₂₁ have been suggested to have the strongest mitogenic and vascular permeability-promoting potential [17–19].

Vascular endothelial growth factor-A and vascular permeability

In addition to promoting angiogenesis, VEGF-A also affects vascular permeability by inducing formation of pores in the vascular endothelial cells [20] and by disrupting the intercellular junction between these cells [21]. The angiogenic and vascular permeability effects of VEGF-A on the endothelium are mediated by the transmembrane receptor VEGFR2 [flk-1/kinase insert domain receptor (KDR)] and involve diverse downstream signaling partners, such as Src family kinases and/or protein tyrosine phosphatases, disrupting and uncoupling the endothelial cell–cell junctions [22]. This, in turn, leads to extravasation of fluid, proteins, and circulating cells [22]. In neovascular ocular diseases, the edema from new, permeable blood vessels can disrupt the retinal anatomy and separate the retina from underlying structures, potentially causing severe vision loss.

Vascular endothelial growth factor-A and inflammation

Chronic inflammation occurs in response to infection, autoimmune disease, injury, tumors, and other diseases and involves the release of various cytokines at specific sites in the body by inflammatory cells such as T cells, B cells, macrophages, natural killer cells, neutrophils, and granulocytes. The proinflammatory cytokines include tumor necrosis factor (TNF)- α , interleukin (IL)-6, IL-8, and IL-1 α , IL-1 β , and oncostatin M, which participate in a cascade of events leading to increased levels of

VEGF-A, promotion of local angiogenesis, and increasing the severity of inflammation.

Vascular endothelial growth factor-A and neovascular ocular diseases

Evidence from preclinical and clinical studies implicates VEGF-A in the pathogenesis of neovascular eye diseases. In streptozotocin-induced diabetic rats, *VEGF-A* gene expression was significantly increased in the ganglion and inner nuclear retinal cell layers compared with control rats [23]. Laser-induced RVO in rabbits [24] and monkeys [25] also led to increased VEGF-A mRNA expression; VEGF-A protein expression was localized to ischemic regions of the retinal layers affected by laser treatment [24]. Furthermore, VEGF-A inhibition prevented retinal neovascularization in an ischemia-induced mouse model [26] and iris neovascularization in a monkey model [27]. VEGF-A inhibition also prevented laser-induced CNV in monkeys or shortened its duration [28].

In clinical studies, increased VEGF-A expression was found in the RPE [29], subfoveal fibroblasts [30], and surgically excised CNV membranes [31] from eyes of neovascular AMD patients. VEGF-A is also overexpressed in the aqueous and vitreous fluid of patients with subretinal neovascularization, diabetic retinopathy, central RVO (CRVO), branch RVO (BRVO), iris neovascularization, retinal detachment, and retinopathy of prematurity (ROP) [32–35] and in all retinal nuclear layers of ischemic CRVO eyes [36]. The consistent association of pathologic ocular neovascularization with increased VEGF-A expression provides a strong rationale for exploring the therapeutic potential of anti-VEGF agents in neovascular AMD.

Genetic case–control studies [37] have shown that the *VEGF* gene may influence an individual's tendency to develop AMD. Analyses of single nucleotide polymorphisms (SNPs) in the VEGF-A promoter and gene have associated specific VEGF-A haplotypes with neovascular AMD [38]. In particular, the VEGF SNP 936C/T [when present with the complement factor H (CFH) Y402H] has been associated with an increased risk of developing wet AMD [39••].

Antivascular endothelial growth factor therapies in neovascular age-related macular degeneration

Intravitreal pegaptanib sodium, an RNA aptamer that targets VEGF₁₆₅ and possibly its larger isoforms, and intravitreal ranibizumab, a mAb antigen-binding fragment that targets all VEGF isoforms and their bioactive cleavage products, received FDA approval for the treatment of neovascular AMD in 2004 and 2006, respectively (Table 1). Bevacizumab (Avastin; Genentech, Inc.), a

Table 1 Antivascular endothelial growth factor agents currently available or in development for the treatment of neovascular age-related macular degeneration

Agent	Class	Molecular target	Drug development stage	Manufacturer
Pegaptanib (Macugen)	Aptamer	VEGF ₁₆₅ isoform	FDA approved	OSI/Eyetech
Ranibizumab (Lucentis)	mAb fragment	All VEGF-A isoforms	FDA approved	Genentech
Bevacizumab (Avastin)	mAb	All VEGF-A isoforms	Various international phase III trials	Genentech ^a
VEGF trap	Decoy receptor	All VEGF-A isoforms, VEGF-B, and PlGF	Phase III	Regeneron
Triamcinolone acetonide (Kenalog-40)	Corticosteroid	Antiangiogenic; specific targets unknown, possibly VEGF and others	Phase III	Bristol-Myers Squibb
Bevasiranib (Cand5)	siRNA	VEGF-A mRNA	Phase III	OPKO Health
AGN211745 (Sirna-027)	siRNA	VEGF-R1 mRNA	Phase II	Allergan
ATG003	Topical mecamlamine	Antiangiogenic; $\alpha 3\beta 4$ nicotinic Ach receptor	Phase II	CoMentis, South San Francisco, California, USA
Quark	siRNA		Phase I, II	Quark Inc., Denver, Colorado, USA; Pfizer, New York, New York, USA
JSM6427	Small molecule integrin $\alpha 5\beta 1$ -antagonist	Antiangiogenic; integrin $\alpha 5\beta 1$	Phase I	Jerini AG, Berlin, Germany

Ach, acetylcholine; FDA, Food and Drug Administration; PlGF, placental growth factor; siRNA, small interfering RNA; VEGF, vascular endothelial growth factor; VEGF-R1, vascular endothelial growth factor receptor 1.

^aInvestigations of the clinical use of bevacizumab in ocular diseases are independent of Genentech.

full-length humanized mAb, was derived from the same murine anti-VEGF mAb as ranibizumab. Similar to ranibizumab, bevacizumab targets all isoforms and bioactive cleavage products of VEGF-A. Bevacizumab is FDA approved for use in combination with chemotherapy for the systemic treatment of metastatic colorectal, lung, and breast cancer. Systemic and intravitreal bevacizumab have also recently been used off label in neovascular AMD and other exudative eye diseases. The successful treatment of neovascular AMD with these agents as presented in the overview below serves as a proof-of-concept for the clinical use of anti-VEGF in neovascular eye diseases.

Pegaptanib sodium

The landmark phase III VEGF Inhibition Study In Ocular Neovascularization (VISION) clinical trials showed that intravitreal pegaptanib sodium slows visual loss in neovascular AMD, with 70% of treated patients losing more than 15 letters of visual acuity compared with 55% of controls [40,41]. Moreover, 6% of pegaptanib sodium-treated patients gained at least 15 letters compared with 2% of the patients in the control group [40]. Fluorescein angiography at 30 and 54 weeks showed that the pegaptanib-treated group had a significant reduction ($P < 0.01$) in the rate of growth of the total area of their CNV lesions and the severity of leakage compared with the control group [40].

Ranibizumab

The pivotal phase III Minimally Classic/Occult Trial of the Anti-VEGF Antibody Ranibizumab in the treatment of Neovascular AMD (MARINA) [42] and Anti-VEGF

Antibody for the Treatment of Predominantly Classic CNV in AMD (ANCHOR) [43] trials established ranibizumab as the first FDA-approved agent that prevents vision loss and improves vision in a substantial proportion of patients with all subtypes of neovascular AMD. At 12 and 24 months in the MARINA trial, 90–95% of patients treated with 0.3 or 0.5 mg ranibizumab lost less than 15 letters of visual acuity compared with 53–64% of control patients; also at 12 and 24 months, 25–34% of ranibizumab-treated patients had gained at least 15 letters of visual acuity compared with 4–5% of control patients [42]. The ANCHOR trial, which compared ranibizumab with verteporfin photodynamic therapy (PDT), had similar findings at 12 and 24 months: 90–96% of the ranibizumab-treated versus 64–66% of the PDT-treated patients lost less than 15 letters of visual acuity, whereas 34–41% of the ranibizumab group versus 6% of the PDT group gained more than 15 letters [43] (Brown DM, personal communication).

Analyses of fluorescein angiography in both the MARINA and ANCHOR studies also revealed statistically significant decreases in mean area and total CNV, leakage from CNV, serous sensory retinal detachment (SSRD), and disciform scar/subretinal fibrosis at both 12 and 24 months after ranibizumab treatment [43,44]. A retrospective analysis of optical coherence tomography (OCT)/fluorescein angiography prospectively collected in a subset of 46 patients from the MARINA study showed a statistically significant decrease at 12 months (final OCT) in mean foveal center point thickness of the ranibizumab-treated group compared with the sham-treated group [44].

Clinical studies such as Prospective OCT Imaging of Patients With Neovascular AMD Treated With Intraocular Lucentis (PrONTO) and Phase IIIb, Multicenter, Randomized, Double-Masked, Sham Injection–Controlled Study of the Efficacy and Safety of Ranibizumab in Patients with Subfoveal CNV with or without Classic CNV Secondary to AMD (PIER) have investigated alternative, less-frequent ranibizumab dosing strategies in an attempt to lower rates of potential treatment-related adverse events. The 2-year phase I/II open-label PrONTO trial evaluated an OCT-guided, variable-dosing regimen of monthly intravitreal ranibizumab (0.5 mg) for 3 months followed by intravitreal ranibizumab as needed based on OCT-defined retreatment criteria in 40 patients with all subtypes of neovascular AMD [45•]. At 1 year, 95% (38/40) of treated patients had lost less than 15 letters of visual acuity; 35% (14/40) of treated patients had gained at least 15 letters of visual acuity, and the mean change in visual acuity was +9.3 letters for treated patients. The mean number of injections for the first year was 5.6 (range 3–13); the most common reason for reinjection was a loss of at least five letters of visual acuity in association with presence of macular fluid. The earliest signs of recurrent fluid in the macula following cessation of treatment were OCT detectable. At 12 months after treatment, the mean central retinal thickness (CRT) as measured by OCT decreased by 178 μm ($P < 0.001$). At 24 months, the results were virtually identical [a mean visual acuity change of +11.1 letters, a mean CRT/OCT decrease of 215 μm , and a mean number of injections of 10 (range 3–25) over 2 years] (Rosenfeld PJ, personal communication). The PrONTO trial showed that in future clinical trials it may be possible to use qualitative OCT to determine the basis for retreatment.

The 2-year PIER trial [46•] examined the efficacy and safety of 0.3 or 0.5 mg ranibizumab monthly for 3 months followed by quarterly dosing. The first-year data showed that a significantly greater proportion of patients receiving ranibizumab lost less than 15 letters of visual acuity (83.3% of patients in the 0.3 mg group and 90.2% of patients in the 0.5 mg group) compared with 49.2% of patients in the sham treatment group ($P < 0.0001$ for each dose level versus sham). However, there was no significant difference in the proportion of patients who gained at least 15 letters: 11.7 and 13.1% of treated patients (0.3 and 0.5 mg, respectively) compared with 9.5% in the sham group. Although the overall safety profile of ranibizumab in the PIER trial was similar to the first year of the MARINA and ANCHOR trials, the efficacy outcomes of the PIER trial were less beneficial than the MARINA and ANCHOR trials because some patients required more frequent dosing than quarterly dosing to achieve maximal benefit. On the basis of the PrONTO and PIER trials, OCT-guided administration

of less-frequent ranibizumab retreatment appears to be beneficial.

The recently completed Safety Assessment of Intravitreal Lucentis for AMD (SAILOR) trial was a single-masked, multicenter phase IIIb study to evaluate the safety and tolerability of two doses of intravitreal ranibizumab in patients with neovascular AMD. Safety assessments showed higher stroke rates with increased dosage (0.7 and 1.2% for the 0.3 and 0.5 mg groups, respectively), but the difference was not statistically significant [47••]. Prior stroke was the most significant risk factor for stroke. Frequencies of cardiovascular events and ocular serious AEs were similar for the two dose groups [47••,48]. A subgroup analysis of this study showed an association between greater improvements in visual acuity and central foveal thickness in patients presenting with a higher baseline visual acuity [49].

Intravitreal bevacizumab

Although systemic bevacizumab (5 mg/kg) has been shown to reduce leakage from CNV, decrease CRT, and significantly improve vision in neovascular AMD [50–52], intravitreal administration is perceived to be safer and requires less frequent retreatment [50,51]. In the first-reported case study [53] of intravitreal bevacizumab, a patient with recurrent CNV secondary to AMD, who had previously been treated with pegaptanib and with PDT in combination with triamcinolone acetonide, experienced resolution of visual distortion within 1 week in parallel with resolution of subretinal fluid and a reduction in CRT following a single injection of 1.0 mg bevacizumab. Subsequently, several retrospective [54–67] and prospective [68–79] studies of intravitreal bevacizumab (dose range 1.0–2.5 mg) in neovascular AMD have been published, invariably demonstrating clinically significant improvement in mean visual acuity, reduction in fluorescein angiography leakage and CRT, and resolution of edema in up to 90% of bevacizumab-treated patients. Most studies were small (up to 100 patients) uncontrolled studies with different retreatment criteria and outcome measures. Systemic and ocular AEs were rare; the most common ocular side effects were endophthalmitis, uveitis, submacular hemorrhage, and RPE tears. In a recent retrospective safety assessment of intravitreal bevacizumab involving 1173 patients, 18 (1.5%) reported systemic AEs, including five (0.4%) deaths, whereas ocular AEs included subconjunctival hemorrhage [838 cases (19% of 4303 injections)] and increased intraocular pressure (IOP), endophthalmitis, and tractional retinal detachment [seven cases (0.16%) each] [67]. The low rates of systemic complications in these studies were consistent with those reported in an earlier survey of 5228 patients [80].

A randomized, prospective clinical trial compared verteporfin PDT with bevacizumab (2.5 mg) for the treatment of predominantly classic CNV secondary to AMD and found that at month 6, all 32 eyes (100%) receiving bevacizumab lost 15 letters or less in visual acuity compared with 73.3% of the PDT-receiving eyes ($P=0.002$) [76]. Mean CRT was significantly better at 3 and 6 months in patients treated with bevacizumab versus the PDT group ($P=0.04$ and $P=0.002$, respectively). The study showed overall benefit of treatment with bevacizumab compared with PDT.

Combination therapy

Small short-term studies evaluating the combination of intravitreal bevacizumab with PDT versus PDT alone [81–83] and bevacizumab as well as pegaptanib sodium [84] for neovascular AMD have demonstrated clinical benefits (visual acuity improvement and resolution of edema), suggesting possible synergistic effects. However, these findings require confirmation in large randomized controlled trials. Currently, combinations of two or three therapies, compared with anti-VEGF monotherapy, are being tested for their ability to reduce the intervention rate with equivalent efficacy and safety outcomes. These include the combination of ranibizumab as well as PDT, bevacizumab monotherapy versus bevacizumab as well as PDT, or bevacizumab as well as PDT and triamcinolone, and bevacizumab as well as triamcinolone.

An ongoing phase I trial (<http://www.clinicaltrials.gov>, NCT00569140) is evaluating the safety, tolerability, and pharmacokinetic profile of intravitreal E10030 (Ophthotech Corporation, Princeton, New Jersey, USA) in patients with neovascular AMD receiving ranibizumab or bevacizumab. E10030 is a pegylated aptamer that targets platelet-derived growth factor (PDGF). The rationale for the study is that PDGF and VEGF have independent angiogenic activities, and preclinical studies have demonstrated that the combination of E10030 and anti-VEGF agents has more potent antiangiogenic results than anti-VEGF treatment alone (Ophthotech Corporation press release, 12 February 2008).

Sorafenib (Nexavar; Bayer HealthCare Pharmaceuticals, Wayne, New Jersey, USA) is a tyrosine kinase inhibitor that diminishes VEGF signaling by inhibiting VEGFR and has been approved for the treatment of certain cancers. Recently, a report described two neovascular AMD patients who received off-label oral sorafenib (200 mg three times per week) in combination with ongoing intravitreal ranibizumab treatment in an attempt to decrease the number of ranibizumab injections [85]. Both patients experienced stable or improved visual acuity and improvement in CRT after initiation of sorafenib therapy. Therefore, it will be of interest to evalu-

ate the effects of sorafenib in combination with ranibizumab or as monotherapy in larger patient cohorts, especially in patients with refractory or recurrent neovascular AMD.

The safety and efficacy of a unique combination therapy consisting of focal strontium-90 beta radiation delivered via the Epi-Rad90 Ophthalmic System (NeoVista, Fremont, California, USA) as well as ranibizumab, versus ranibizumab alone, is being evaluated in neovascular AMD in the ongoing randomized, open-label, active control phase III CNV secondary to AMD treated with Beta Radition Epiretinal Therapy (CABERNET) trial (<http://www.clinicaltrials.gov>; NCT00454389).

Other strategies to inhibit vascular endothelial growth factor signal transduction

Several different anti-VEGF and antiangiogenic agents have demonstrated efficacy in neovascular AMD. A phase II randomized, double-masked trial of VEGF trap (Regeneron Pharmaceuticals, Tarrytown, New Jersey, USA), a fusion protein of VEGF receptor ligand-binding domains and the immunoglobulin G type 1 (IgG1) Fc region, demonstrated increases in visual acuity from baseline of 9.0 letters ($P<0.0001$) and 5.4 letters ($P<0.085$) at 52 weeks after fixed monthly or quarterly dosing (0.5, 2, or 4 mg) for 12 weeks followed by another 40 weeks on an as-needed basis in neovascular AMD (AZ-Regeneron press release, 28 September 2008). Patients receiving monthly VEGF trap (2.0 or 0.5 mg for the first 12 weeks) also experienced mean reductions in CRT of 143 and 125 μm , respectively, at week 52 ($P<0.0001$ for both from baseline). Currently, two randomized, international phase III studies (VIEW-1 and VIEW-2) (<http://www.clinicaltrials.gov>; NCT00509795, NCT00637377) are comparing intravitreal VEGF trap with ranibizumab.

Small interfering RNA (siRNA) agents designed to silence the *VEGF* gene and the VEGFR have been shown in preclinical studies to inhibit ocular neovascularization and vascular permeability in animal models [86,87]. Bevasiranib (previously called Cand5; OPKO Health, Inc./Acuity Pharmaceuticals, Miami, Florida, USA) and AGN211745 (previously called Sirna-027; Allergan, Inc., Irvine, California, USA) are currently being studied in phase II and III clinical trials, respectively, in neovascular AMD. In a phase II randomized trial, bevasiranib (single intravitreal injections at baseline and week 6) was shown to be well tolerated after 12 weeks of follow up, with mild AEs related primarily to the injection procedure and with no systemic exposure. Bevasiranib stabilized visual acuity in most patients and improved visual acuity in more than one-third of patients (Acuity Pharmaceuticals press release, 11 September 2006). The Combining Bevasiranib and Lucentis (ranibizumab)

Therapy in Wet AMD (COBALT) phase III trial (<http://www.clinicaltrials.gov>; NCT00499590) is currently evaluating the safety and efficacy of bevasiranib given every 8 or 12 weeks after ranibizumab pretreatment compared with ranibizumab alone every 4 weeks. Another phase III study (CARBON) (<http://www.clinicaltrials.gov>; NCT00557791) will assess the safety and efficacy of three doses of bevasiranib as maintenance therapy for AMD following three doses of ranibizumab. Data from a phase I study of AGN211745 in neovascular AMD indicated that after 3 months of follow up, single injections of AGN211745 stabilized visual acuity in 24 of 26 (92%) patients and improved vision in five of 26 (19%) patients (Sirna Therapeutics, Inc., press release, 10 August 2006). A recently completed phase I/II trial (<http://www.clinicaltrials.gov>; NCT00363714) evaluated the safety and effect of AGN211745 on retinal anatomy, CNV, and visual acuity; another phase II trial assessing the efficacy of multiple doses of this agent is ongoing (<http://www.clinicaltrials.gov>; NCT0035057). However, the specificity of the aforementioned siRNA drugs was recently called into question by a recent study [88**] showing that the antiangiogenic activity of siRNA may be due to a class immune effect associated with activation of the cell surface toll-like receptor 3 (TLR3) rather than to a target sequence-specific interaction. Moreover, these siRNA drugs could theoretically cause vision loss via activation of the TLR3, which may promote the formation of geographic atrophy in genetically susceptible individuals [89].

In early clinical studies [90–93], triamcinolone acetonide (Kenalog; Bristol-Myers Squibb, New York, New York, USA), an antiangiogenic corticosteroid, helped stabilize and, to a lesser extent, improve visual acuity when used in combination with PDT in neovascular AMD. Intravitreal triamcinolone acetonide (IVTA) combined with PDT can stabilize vision in the majority of treated eyes, whereas only a small proportion (9%) of patients had improved visual acuity [92]. IVTA did not prevent visual loss in a substantial number of patients, although most studies assessed only single-dose regimens, unlike the periodic-dosing regimens used for the anti-VEGF agents discussed above. Additionally, IVTA introduces AEs not seen with PDT alone, including increases in IOP and cataract progression [92]. Currently, triamcinolone acetonide appears to be most useful in reducing the need for PDT retreatment [90–92]. Other agents being investigated for neovascular ocular diseases are listed in Table 1.

Conclusion

The accumulation of preclinical and clinical evidence implicating VEGF-A in the pathogenesis of neovascular eye diseases has provided a strong rationale for exploring the therapeutic potential of anti-VEGF agents in these

diseases, including AMD. Over the past decade, the management of neovascular AMD has progressed considerably from being limited to laser photocoagulation to contain the spread of CNV to the use of anti-VEGF-A agents to inhibit neovascularization and prevent vision loss. Three such agents – pegaptanib sodium, ranibizumab, and bevacizumab – have been used successfully for this purpose. Their clinical success in neovascular AMD has provided the proof-of-concept needed to pursue similar treatment strategies in other neovascular eye diseases.

Acknowledgements

The support for third-party medical writing assistance was provided by Genentech USA, Inc.

Dr Ciulla has received grant support from Alcon, Alimera, Allergan, Genentech, GlaxoSmithKline, Jerini, OPKO, Othera, and Regeneron, is a consultant for Neovista, and serves on the advisory boards for Pfizer and Regeneron. Dr Rosenfeld has received grant support from Alcon, CoMentis, Genentech, Othera, Potentia, and Quark/Pfizer, is on the speakers bureau of Carl Zeiss Meditec, and is an advisory board member of CoMentis, Othera, and Neovista.

References and recommended reading

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
- of outstanding interest

- 1 Bird A. Age-related macular disease. *Br J Ophthalmol* 1996; 80:2–3.
- 2 Ciulla TA, Amador AG, Zinman B. Diabetic retinopathy and diabetic macular edema: pathophysiology, screening, and novel therapies. *Diabetes Care* 2003; 26:2653–2664.
- 3 Cugati S, Wang JJ, Rochtchina E, Mitchell P. Ten-year incidence of retinal vein occlusion in an older population: the Blue Mountains Eye Study. *Arch Ophthalmol* 2006; 124:726–732.
- 4 Chakravarthy U, Augood C, Bentham GC, *et al.* Cigarette smoking and age-related macular degeneration in the EUREYE Study. *Ophthalmology* 2007; 114:1157–1163.
- 5 Campochiaro PA, Soloway P, Ryan SJ, Miller JW. The pathogenesis of choroidal neovascularization in patients with age-related macular degeneration. *Mol Vis* 1999; 5:34.
- 6 Leung DW, Cachianes G, Kuang WJ, *et al.* Vascular endothelial growth factor is a secreted angiogenic mitogen. *Science* 1989; 246:1306–1309.
- 7 Senger DR, Galli SJ, Dvorak AM, *et al.* Tumor cells secrete a vascular permeability factor that promotes accumulation of ascites fluid. *Science* 1983; 219:983–985.
- 8 Keck PJ, Hauser SD, Krivi G, *et al.* Vascular permeability factor, an endothelial cell mitogen related to PDGF. *Science* 1989; 246:1309–1312.
- 9 Neufeld G, Cohen T, Gengrinovitch S, Poltorak Z. Vascular endothelial growth factor (VEGF) and its receptors. *FASEB J* 1999; 13:9–22.
- 10 Eriksson U, Alitalo K. Structure, expression and receptor-binding properties of novel vascular endothelial growth factors. *Curr Top Microbiol Immunol* 1999; 237:41–57.
- 11 Ferrara N, Gerber HP, LeCouter J. The biology of VEGF and its receptors. *Nat Med* 2003; 9:669–676.
- 12 Gerhardt H, Ruhrberg C, Abramsson A, *et al.* Neuropilin-1 is required for endothelial tip cell guidance in the developing central nervous system. *Dev Dyn* 2004; 231:503–509.
- 13 Kawasaki T, Kitsukawa T, Bekku Y, *et al.* A requirement for neuropilin-1 in embryonic vessel formation. *Development* 1999; 126:4895–4902.
- 14 Tischer E, Mitchell R, Hartman T, *et al.* The human gene for vascular endothelial growth factor. Multiple protein forms are encoded through alternative exon splicing. *J Biol Chem* 1991; 266:11947–11954.

- 15 Ferrara N. Vascular endothelial growth factor: basic science and clinical progress. *Endocr Rev* 2004; 25:581–611.
 - 16 Takahashi H, Shibuya M. The vascular endothelial growth factor (VEGF)/VEGF receptor system and its role under physiological and pathological conditions. *Clin Sci (Lond)* 2005; 109:227–241.
 - 17 Houck KA, Ferrara N, Winer J, *et al.* The vascular endothelial growth factor family: identification of a fourth molecular species and characterization of alternative splicing of RNA. *Mol Endocrinol* 1991; 5:1806–1814.
 - 18 Zhang HT, Scott PA, Morbidelli L, *et al.* The 121 amino acid isoform of vascular endothelial growth factor is more strongly tumorigenic than other splice variants in vivo. *Br J Cancer* 2000; 83:63–68.
 - 19 Park JE, Keller GA, Ferrara N. The vascular endothelial growth factor (VEGF) isoforms: differential deposition into the subepithelial extracellular matrix and bioactivity of extracellular matrix-bound VEGF. *Mol Biol Cell* 1993; 4:1317–1326.
 - 20 Roberts WG, Palade GE. Neovasculation induced by vascular endothelial growth factor is fenestrated. *Cancer Res* 1997; 57:765–772.
 - 21 Monsky WL, Fukumura D, Gohongi T, *et al.* Augmentation of transvascular transport of macromolecules and nanoparticles in tumors using vascular endothelial growth factor. *Cancer Res* 1999; 59:4129–4135.
 - 22 Weis S, Shintani S, Weber A, *et al.* Src blockade stabilizes a flk/cadherin complex, reducing edema and tissue injury following myocardial infarction. *J Clin Invest* 2004; 113:885–894.
 - 23 Gilbert RE, Vranes D, Berka JL, *et al.* Vascular endothelial growth factor and its receptors in control and diabetic rat eyes. *Lab Invest* 1998; 78:1017–1027.
 - 24 Pe'er J, Shweiki D, Itin A, *et al.* Hypoxia-induced expression of vascular endothelial growth factor by retinal cells is a common factor in neovascularizing ocular diseases. *Lab Invest* 1995; 72:638–645.
 - 25 Shima DT, Gougos A, Miller JW, *et al.* Cloning and mRNA expression of vascular endothelial growth factor in ischemic retinas of *Macaca fascicularis*. *Invest Ophthalmol Vis Sci* 1996; 37:1334–1340.
 - 26 Ozaki H, Seo MS, Ozaki K, *et al.* Blockade of vascular endothelial cell growth factor receptor signaling is sufficient to completely prevent retinal neovascularization. *Am J Pathol* 2000; 156:697–707.
 - 27 Adamis AP, Shima DT, Tolentino MJ, *et al.* Inhibition of vascular endothelial growth factor prevents retinal ischemia-associated iris neovascularization in a nonhuman primate. *Arch Ophthalmol* 1996; 114:66–71.
 - 28 Krzystolik MG, Afshari MA, Adamis AP, *et al.* Prevention of experimental choroidal neovascularization with intravitreal anti-vascular endothelial growth factor antibody fragment. *Arch Ophthalmol* 2002; 120:338–346.
 - 29 Kliffen M, Sharma HS, Mooy CM, *et al.* Increased expression of angiogenic growth factors in age-related maculopathy. *Br J Ophthalmol* 1997; 81:154–162.
 - 30 Kvant A, Alverre PV, Berglin L, Seregard S. Subfoveal fibrovascular membranes in age-related macular degeneration express vascular endothelial growth factor. *Invest Ophthalmol Vis Sci* 1996; 37:1929–1934.
 - 31 Rakic JM, Lambert V, Devy L, *et al.* Placental growth factor, a member of the VEGF family, contributes to the development of choroidal neovascularization. *Invest Ophthalmol Vis Sci* 2003; 44:3186–3193.
 - 32 Wells JA, Murthy R, Chibber R, *et al.* Levels of vascular endothelial growth factor are elevated in the vitreous of patients with subretinal neovascularisation. *Br J Ophthalmol* 1996; 80:363–366.
 - 33 Adamis AP, Miller JW, Bernal MT, *et al.* Increased vascular endothelial growth factor levels in the vitreous of eyes with proliferative diabetic retinopathy. *Am J Ophthalmol* 1994; 118:445–450.
 - 34 Aiello LP, Avery RL, Arrigg PG, *et al.* Vascular endothelial growth factor in ocular fluid of patients with diabetic retinopathy and other retinal disorders. *N Engl J Med* 1994; 331:1480–1487.
 - 35 Noma H, Minamoto A, Funatsu H, *et al.* Intravitreal levels of vascular endothelial growth factor and interleukin-6 are correlated with macular edema in branch retinal vein occlusion. *Graefes Arch Clin Exp Ophthalmol* 2006; 244:309–315.
 - 36 Pe'er J, Folberg R, Itin A, *et al.* Vascular endothelial growth factor upregulation in human central retinal vein occlusion. *Ophthalmology* 1998; 105:412–416.
 - 37 Haines JL, Schnetz-Boutaud N, Schmidt S, *et al.* Functional candidate genes in age-related macular degeneration: significant association with VEGF, VLDLR, and LRP6. *Invest Ophthalmol Vis Sci* 2006; 47:329–335.
 - 38 Churchill AJ, Carter JG, Lovell HC, *et al.* VEGF polymorphisms are associated with neovascular age-related macular degeneration. *Hum Mol Genet* 2006; 15:2955–2961.
 - 39 Lin JM, Wan L, Tsai YY, *et al.* Vascular endothelial growth factor gene polymorphisms in age-related macular degeneration. *Am J Ophthalmol* 2008; 145:1045–1051.
- This study revealed a VEGF SNP (+936T) that, together with the complement factor polymorphism Y402H, could be a potential predictor of increased risk for developing wet AMD in elderly populations. This study adds to the growing list of risk factors for AMD and strengthens the basis for a potentially more fruitful pharmacogenomic approach to the management of ocular diseases.
- 40 Gragoudas ES, Adamis AP, Cunningham ET Jr, *et al.* Pegaptanib for neovascular age-related macular degeneration. *N Engl J Med* 2004; 351:2805–2816.
 - 41 Chakravarthy U, Adamis AP, Cunningham ET Jr, *et al.* Year 2 efficacy results of 2 randomized controlled clinical trials of pegaptanib for neovascular age-related macular degeneration. *Ophthalmology* 2006; 113:1508–1525.
 - 42 Rosenfeld PJ, Brown DM, Heier JS, *et al.* Ranibizumab for neovascular age-related macular degeneration. *N Engl J Med* 2006; 355:1419–1431.
 - 43 Brown DM, Kaiser PK, Michels M, *et al.* Ranibizumab versus verteporfin for neovascular age-related macular degeneration. *N Engl J Med* 2006; 355:1432–1444.
 - 44 Kaiser PK, Blodi BA, Shapiro H, Acharya NR. Angiographic and optical coherence tomographic results of the MARINA study of ranibizumab in neovascular age-related macular degeneration. *Ophthalmology* 2007; 114:1868–1875.
 - 45 Fung AE, Lalwani GA, Rosenfeld PJ, *et al.* An optical coherence tomography-guided, variable dosing regimen with intravitreal ranibizumab (Lucentis) for neovascular age-related macular degeneration. *Am J Ophthalmol* 2007; 143:566–583.
- OCT can be used to accurately detect the earliest sign of recurrent macular fluid accumulation after cessation of ranibizumab injections. OCT-guided, variable dosing retreatment resulted in visual acuity outcomes similar to the phase III trials using fewer intravitreal injections.
- 46 Regillo CD, Brown DM, Abraham P, *et al.* Randomized, double-masked, sham-controlled trial of ranibizumab for neovascular age-related macular degeneration: PIER Study year 1. *Am J Ophthalmol* 2008; 145:239–248.
- Ranibizumab (0.3 or 0.5 mg) administered monthly and then quarterly thereafter was well tolerated and provided significant and clinically meaningful visual acuity improvement to patients with all angiographic subtypes of AMD-related CNV. However, the quarterly treatments were associated with a decline from the benefit achieved with monthly therapy.
- 47 Michels M, Francom S, Wilson L. Systemic safety and risk factors associated with intravitreal ranibizumab in patients with choroidal neovascularization (CNV) secondary to age-related macular degeneration (AMD) [abstract]. In: 26th Annual Meeting of the American Society of Retina Specialists; 11–15 October 2008; Maui, Hawaii.
- The SAILOR study showed no significant differences in the rates of cardiovascular and nonocular adverse events between patients receiving high-dose (0.5 mg) and low-dose (0.3 mg) ranibizumab. A history of stroke, heart failure, and arterial thromboembolic events was predictive of stroke, regardless of the treatment dose. This may alleviate concerns regarding the low rates of arterial thromboembolic events earlier reported in the MARINA and ANCHOR studies.
- 48 Kokame G, Francom S, Yee W. Ocular safety of ranibizumab (LUCENTIS) in patients with wet age-related macular degeneration: SAILOR and 3 additional randomized, multicenter trials [abstract]. In: 26th Annual Meeting of the American Society of Retina Specialists; 11–15 October 2008; Maui, Hawaii.
 - 49 Ferrone PJ, Francom S, Tuomi L. Efficacy of ranibizumab (LUCENTIS) in patients with wet age-related macular degeneration: subgroup analysis of SAILOR study cohort 1 [abstract]. In: 26th Annual Meeting of the American Society of Retina Specialists; 11–15 October 2008; Maui, Hawaii.
 - 50 Michels S, Rosenfeld PJ, Puliafito CA, *et al.* Systemic bevacizumab (Avastin) therapy for neovascular age-related macular degeneration: twelve-week results of an uncontrolled open-label clinical study. *Ophthalmology* 2005; 112:1035–1047.
 - 51 Moshfeghi AA, Rosenfeld PJ, Puliafito CA, *et al.* Systemic bevacizumab (Avastin) therapy for neovascular age-related macular degeneration: twenty-four-week results of an uncontrolled open-label clinical study. *Ophthalmology* 2006; 113:e1–e12.
 - 52 Geitzenauer W, Michels S, Prager F, *et al.* Early effects of systemic and intravitreal bevacizumab (Avastin) therapy for neovascular age-related macular degeneration [in German]. *Klin Monatsbl Augenheilkd* 2006; 223:822–827.
 - 53 Rosenfeld PJ, Moshfeghi AA, Puliafito CA. Optical coherence tomography findings after an intravitreal injection of bevacizumab (Avastin) for neovascular age-related macular degeneration. *Ophthalmic Surg Lasers Imaging* 2005; 36:331–335.
 - 54 Rich RM, Rosenfeld PJ, Puliafito CA, *et al.* Short-term safety and efficacy of intravitreal bevacizumab (Avastin) for neovascular age-related macular degeneration. *Retina* 2006; 26:495–511.

- 55 Spaide RF, Laud K, Fine HF, *et al.* Intravitreal bevacizumab treatment of choroidal neovascularization secondary to age-related macular degeneration. *Retina* 2006; 26:383–390.
- 56 Yoganathan P, Deramo VA, Lai JC, *et al.* Visual improvement following intravitreal bevacizumab (Avastin) in exudative age-related macular degeneration. *Retina* 2006; 26:994–998.
- 57 Aggio FB, Farah ME, Silva WC, Melo GB. Intravitreal bevacizumab for exudative age-related macular degeneration after multiple treatments. *Graefes Arch Clin Exp Ophthalmol* 2007; 245:215–220.
- 58 Chen E, Kaiser RS, Vander JF. Intravitreal bevacizumab for refractory pigment epithelial detachment with occult choroidal neovascularization in age-related macular degeneration. *Retina* 2007; 27:445–450.
- 59 Emerson MV, Lauer AK, Flaxel CJ, *et al.* Intravitreal bevacizumab (Avastin) treatment of neovascular age-related macular degeneration. *Retina* 2007; 27:439–444.
- 60 Goff MJ, Johnson RN, McDonald HR, *et al.* Intravitreal bevacizumab for previously treated choroidal neovascular from age-related macular degeneration. *Retina* 2007; 27:432–438.
- 61 Madhusudhana KC, Hannan SR, Williams CP, *et al.* Intravitreal bevacizumab (Avastin) for the treatment of choroidal neovascularization in age-related macular degeneration: results from 118 cases. *Br J Ophthalmol* 2007; 91:1716–1717.
- 62 Pedersen R, Soliman W, Lund-Andersen H, Larsen M. Treatment of choroidal neovascularization using intravitreal bevacizumab. *Acta Ophthalmol Scand* 2007; 85:526–533.
- 63 Cleary CA, Jungkim S, Ravikumar K, *et al.* Intravitreal bevacizumab in the treatment of neovascular age-related macular degeneration, 6- and 9-month results. *Eye* 2007; 22:82–86.
- 64 Ghazi NG, Knappe RM, Kirk TQ, *et al.* Intravitreal bevacizumab (Avastin) treatment of retinal angiomatous proliferation. *Retina* 2008; 28:689–695.
- 65 Jonas JB, Libondi T, Ihloff AK, *et al.* Visual acuity change after intravitreal bevacizumab for exudative age-related macular degeneration in relation to subfoveal membrane type. *Acta Ophthalmol Scand* 2007; 85:563–565.
- 66 Melamud A, Stinnett S, Fekrat S. Treatment of neovascular age-related macular degeneration with intravitreal bevacizumab: efficacy of three consecutive monthly injections. *Am J Ophthalmol* 2008; 146:91–95.
- 67 Wu L, Martinez-Castellanos MA, Quiroz-Mercado H, *et al.* Twelve-month safety of intravitreal injections of bevacizumab (Avastin): results of the Pan-American Collaborative Retina Study Group (PACORES). *Graefes Arch Clin Exp Ophthalmol* 2008; 246:81–87.
- 68 Avery RL, Pieramici DJ, Rabena MD, *et al.* Intravitreal bevacizumab (Avastin) for neovascular age-related macular degeneration. *Ophthalmology* 2006; 113:363–372e5.
- 69 Bashshur ZF, Bazarbachi A, Schakal A, *et al.* Intravitreal bevacizumab for the management of choroidal neovascularization in age-related macular degeneration. *Am J Ophthalmol* 2006; 142:1–9.
- 70 Costa RA, Jorge R, Calucci D, *et al.* Intravitreal bevacizumab for choroidal neovascularization caused by AMD (IBeNA Study): results of a phase 1 dose-escalation study. *Invest Ophthalmol Vis Sci* 2006; 47:4569–4578.
- 71 Aisenbrey S, Ziemssen F, Volker M, *et al.* Intravitreal bevacizumab (Avastin) for occult choroidal neovascularization in age-related macular degeneration. *Graefes Arch Clin Exp Ophthalmol* 2007; 245:941–948.
- 72 Algrever PV, Steen B, Seregard S, Kvanta A. A prospective study on intravitreal bevacizumab (Avastin) for neovascular age-related macular degeneration of different durations. *Acta Ophthalmol* 2008; 86:482–489.
- 73 Chen CY, Wong TY, Heriot WJ. Intravitreal bevacizumab (Avastin) for neovascular age-related macular degeneration: a short-term study. *Am J Ophthalmol* 2007; 143:510–512.
- 74 Giansanti F, Virgili G, Bini A, *et al.* Intravitreal bevacizumab therapy for choroidal neovascularization secondary to age-related macular degeneration: 6-month results of an open-label uncontrolled clinical study. *Eur J Ophthalmol* 2007; 17:230–237.
- 75 Lazic R, Gabric N. Intravitreally administered bevacizumab (Avastin) in minimally classic and occult choroidal neovascularization secondary to age-related macular degeneration. *Graefes Arch Clin Exp Ophthalmol* 2006; 245:68–73.
- 76 Bashshur ZF, Schakal A, Hamam RN, *et al.* Intravitreal bevacizumab vs verteporfin photodynamic therapy for neovascular age-related macular degeneration. *Arch Ophthalmol* 2007; 125:1357–1361.
- 77 Azad RV, Khan MA, Chanana B, Azad S. Intravitreal bevacizumab for subfoveal choroidal neovascularization secondary to age-related macular degeneration in an Indian population. *Jpn J Ophthalmol* 2008; 52:52–56.
- 78 Bashshur ZF, Haddad ZA, Schakal A, *et al.* Intravitreal bevacizumab for treatment of neovascular age-related macular degeneration: a one-year prospective study. *Am J Ophthalmol* 2008; 145:249–256.
- 79 Weigert G, Michels S, Sacu S, *et al.* Intravitreal bevacizumab (Avastin) therapy versus photodynamic therapy plus intravitreal triamcinolone for neovascular age-related macular degeneration: 6-month results of a prospective, randomised, controlled clinical study. *Br J Ophthalmol* 2008; 92:356–360.
- 80 Fung AE, Rosenfeld PJ, Reichel E. The international intravitreal bevacizumab safety survey: using the Internet to assess drug safety worldwide. *Br J Ophthalmol* 2007; 90:1344–1349.
- 81 Costa RA, Jorge R, Calucci D, *et al.* Intravitreal bevacizumab (Avastin) in combination with verteporfin photodynamic therapy for choroidal neovascularization associated with age-related macular degeneration (IBeVe Study). *Graefes Arch Clin Exp Ophthalmol* 2007; 245:1273–1280.
- 82 Dhalla MS, Shah GK, Blinder KJ, *et al.* Combined photodynamic therapy with verteporfin and intravitreal bevacizumab for choroidal neovascularization in age-related macular degeneration. *Retina* 2006; 26:988–993.
- 83 Ladewig MS, Karl SE, Hamelmann V, *et al.* Combined intravitreal bevacizumab and photodynamic therapy for neovascular age-related macular degeneration. *Graefes Arch Clin Exp Ophthalmol* 2008; 246:17–25.
- 84 Hughes MS, Sang DN. Safety and efficacy of intravitreal bevacizumab followed by pegaptanib maintenance as a treatment regimen for age-related macular degeneration. *Ophthalmic Surg Lasers Imaging* 2006; 37:446–454.
- 85 Diago T, Pulido JS, Molina JR, *et al.* Ranibizumab combined with low-dose sorafenib for exudative age-related macular degeneration. *Mayo Clin Proc* 2008; 83:231–234.
- 86 Reich SJ, Fosnot J, Kuroki A, *et al.* Small interfering RNA (siRNA) targeting VEGF effectively inhibits ocular neovascularization in a mouse model. *Mol Vis* 2003; 9:210–216.
- 87 Tolentino MJ, Brucker AJ, Fosnot J, *et al.* Intravitreal injection of vascular endothelial growth factor small interfering RNA inhibits growth and leakage in a nonhuman primate, laser-induced model of choroidal neovascularization. *Retina* 2004; 24:132–138.
- 88 Kleinman ME, Yamada K, Takeda A, *et al.* Sequence- and target-independent angiogenesis suppression by siRNA via TLR3. *Nature* 2008; 452:591–597.
- siRNAs of at least 21 nucleotides long are effective for treating angiogenic diseases regardless of target specificity via a class effect that may also lead to unanticipated and undesirable vascular or immune effects.
- 89 Yang Z, Stratton C, Francis PJ, *et al.* Toll-like receptor 3 and geographic atrophy in age-related macular degeneration. *N Engl J Med* 2008; 359:1456–1463.
- 90 Ruiz-Moreno JM, Montero JA, Barile S, Zarbin MA. Photodynamic therapy and high-dose intravitreal triamcinolone to treat exudative age-related macular degeneration: 1-year outcome. *Retina* 2006; 26:602–612.
- 91 Ergun E, Maar N, Ansari-Shahrezaei S, *et al.* Photodynamic therapy with verteporfin and intravitreal triamcinolone acetate in the treatment of neovascular age-related macular degeneration. *Am J Ophthalmol* 2006; 142:10–16.
- 92 Schmidt-Erfurth U, Michels S, Augustin A. Perspectives on verteporfin therapy combined with intravitreal corticosteroids. *Arch Ophthalmol* 2006; 124:561–563.
- 93 Spaide RF, Sorenson J, Maranan L. Combined photodynamic therapy and intravitreal triamcinolone for nonsubfoveal choroidal neovascularization. *Retina* 2005; 25:685–690.