

The causes and consequences of genetic heterogeneity in cancer evolution

Rebecca A. Burrell^{1*}, Nicholas McGranahan^{1,2*}, Jiri Bartek^{3,4} & Charles Swanton^{1,5}

Recent studies have revealed extensive genetic diversity both between and within tumours. This heterogeneity affects key cancer pathways, driving phenotypic variation, and poses a significant challenge to personalized cancer medicine. A major cause of genetic heterogeneity in cancer is genomic instability. This instability leads to an increased mutation rate and can shape the evolution of the cancer genome through a plethora of mechanisms. By understanding these mechanisms we can gain insight into the common pathways of tumour evolution that could support the development of future therapeutic strategies.

Extensive genetic and phenotypic variation exist not only between tumours (intertumour heterogeneity) but also within individual tumours (intratumour heterogeneity; Fig. 1). In tumours, populations of genetically distinct subclones can intermingle or be spatially separated, and this subclonal architecture varies dynamically throughout the disease course^{1,2}. Inter- and intratumour heterogeneity have significant implications for the choice of biomarkers to guide clinical decision-making in cancer medicine (see Siu and colleagues on page 355).

Genomic instability is a prominent source of genetic diversity within tumours, generating a diverse cell population that can be subject to selection in a given micro-environmental (see Review by Junttila and de Sauvage on page 346) or therapeutic context^{3–5}. Genomic instability can arise through various routes, leaving distinct genomic footprints and differentially affecting tumour evolution and patient outcome. Different instability mechanisms can operate over the course of tumour development, and may be influenced by exposure to cancer drugs^{6,7}. However, genomic instability might also be an attractive therapeutic target⁸.

In this Review we summarize the evidence for genetic and phenotypic variation both within and between tumours, and discuss how this might affect tumour biology, drug response and patient outcome. We focus on the role of genomic instability in generating inter- and intratumour heterogeneity, and review the array of instability mechanisms observed in cancer and their contribution to tumour evolution.

Heterogeneity between different tumour types

It has long been appreciated that there are both genetic and functional differences between tumours. Tumours that originate from different tissues and cell types vary in terms of their genomic landscapes, prognosis and their response to cytotoxic therapies, probably resulting from the fact that the genetic events of transformation interact with cell-intrinsic biological properties. Reflecting this, the site of origin of the primary tumour frequently determines treatment decisions. However, considerable variation in terms of genomic aberrations, aggressiveness and drug sensitivity is also observed between tumours that originate from the same tissue and cell type^{9–11}.

Mutational frequencies of oncogenes and tumour suppressors vary between tumours of different tissues, probably reflecting the importance of distinct signalling pathways within specific tissues or cellular contexts. Mutations in the tumour suppressor gene *APC*, for instance,

are common in gastrointestinal malignancies such as colorectal cancer, probably owing to the importance of WNT-signalling pathways in intestinal epithelial homeostasis¹². The specific cell type of origin can also profoundly influence tumour genetics and biology; for example, endocrine and exocrine pancreatic tumours exhibit considerable divergence in their mutational spectra and clinical behaviour¹³. Intrinsic differences between cell types may affect therapeutic response; the ability of cancer cells to undergo apoptosis in response to DNA-damaging agents seems to be substantially influenced by the cell type of origin¹⁴. Furthermore, drugs that target the mutant BRAF(V600E) oncoprotein seem to be less efficient in colorectal cancer cells than in melanoma cells¹⁵. This could be attributable to feedback upregulation of epidermal growth factor receptor (EGFR) after BRAF(V600E) inhibition in epithelial colorectal cancer cells but not in melanoma cells, which are derived from the neural crest and have lower basal EGFR expression¹⁵.

The clinical challenge that intertumour variation presents can be met to some extent by classifying tumours into subgroups. These may predict, for example, patient outcome or drug sensitivity on the basis of mutations, copy number changes, protein or RNA expression profiles, or patterns of genomic instability^{10,16,17}. For example, breast cancers are routinely classified according to expression of hormone receptors as well as amplification of the gene *HER2* (also known as *ERBB2*) and expression of its protein¹⁰. Similarly, the discovery that around 10% of lung cancers harbour *EGFR*-activating mutations has led to the use of small molecule EGFR inhibitors in the treatment of this disease¹⁸.

However, recent advances in next-generation sequencing have revealed greater than expected inter- and intratumour genetic heterogeneity — very few mutations were observed in more than 5–10% of tumours of a particular tissue type⁹. In addition, within a cohort of tumours, the same gene may be affected by point mutation, DNA methylation, copy number alteration, or a combination of the three, underscoring the need for integrative approaches to analysing somatic aberrations in the cancer genome. Efforts to define tumour subgroups based on specific mutations may be confounded by epistasis, which involves the action of one gene on another. For example, in acute myeloid leukaemia (AML), *NPM1* mutations confer a favourable prognosis only in the presence of a co-occurring *IDH1* or *IDH2* mutation¹⁹.

Although genetic heterogeneity between tumours is known to be widespread, it has long been appreciated that phenotypic heterogeneity

¹Translational Cancer Therapeutics Laboratory, Cancer Research UK London Research Institute, 44 Lincoln's Inn Fields, London WC2A 3LY, UK. ²Centre for Mathematics and Physics in the Life Sciences and Experimental Biology (CoMPLEX), University College London, Physics Building, Gower Street, London WC1E 6BT, UK. ³Danish Cancer Society Research Center, Strandboulevarden 49, DK-2100 Copenhagen, Denmark. ⁴Institute of Molecular and Translational Medicine, Palacky University Olomouc CZ-775 15, Czech Republic. ⁵UCL Cancer Institute, Paul O'Gorman Building, Huntley Street, London WC1E 6BT, UK. *These authors contributed equally to this work.




Figure 1 | Intertumour and intratumour heterogeneity. Genetic and phenotypic variation are observed between tumours of different tissue and cell types, as well as between individuals with the same tumour type (intertumour heterogeneity). Within a tumour, subclonal diversity may be observed (intratumour heterogeneity). Subclones may intermingle (as shown by subclones 1 and 2) or be spatially separated (as shown by subclone 3). Separation between subclones could reflect physical barriers such as blood vessels or micro-environmental changes. Tumour subclones may show differential gene expression due to both genetic and epigenetic heterogeneity. Within a subclonal

population of tumour cells — shown here as a tumour section, hybridized to two fluorescent probes for the centromeres of two chromosomes (chromosome 2, red; chromosome 18, green) with DNA (blue) — there is intercellular genetic and non-genetic variation of, for example, chromosome copy number, somatic point mutations or epigenetic modifications that results in phenotypic diversity. Intercellular genetic heterogeneity is exacerbated by genomic instability, and may foster the emergence of tumour subclones. Genomic instability and tumour subclonal architecture may vary further over time if influenced by, for example, cancer treatment.

may not be as extensive, with the same cellular pathways often repeatedly affected^{9,20–24}. Nevertheless, the impact of a given mutation will depend on the position and relative importance of the affected protein within the signalling pathway. An emerging theme from recent studies is that histone-modifying genes are recurrently mutated^{1,9,22,25}, suggesting that altered chromatin regulation could be a common phenotype in a range of tumours. Another unifying feature of many genetically diverse malignancies is genomic instability — the same pattern of instability can occur through various distinct routes. For instance, deregulation of the G1–S transition, which is observed in almost all cancers, can be achieved through multiple means, simultaneously overriding cellular senescence, promoting proliferation and destabilizing the genome through DNA replication stress^{26–28}.

Despite the evidence for functional convergence, it will nevertheless be challenging to identify biomarkers to define phenotypically similar, yet genetically diverse, lesions to guide treatment. In addition, the task of identifying and validating biomarkers may be complicated further by intratumour heterogeneity (See Review by Siu and colleagues on page 355).

Intratumour heterogeneity and tumour evolution

Within tumours, diversity occurs in the expression of protein biomarkers¹⁰, and at multiple and different genetic^{29,30} and epigenetic^{30,31} levels. This diversity might complicate our ability to resolve the full spectrum of cancer pathway aberrations and confound biomarker validation through tumour sampling bias.

Genetically distinct subclonal populations of cells arise through intercellular genetic variation, followed by selective outgrowth of clones that have a phenotypic advantage within a given tumour micro-environmental context^{32–34}. Fluctuations in subclonal architecture can occur, for example, in the context of disease progression (such as metastasis, see Review by Klein on page 365) or drug treatment^{1,6,21,35–37}. A clonal sweep, whereby a new clone takes over the entire population, replacing ancestral clones, will result in a homogeneous cell population. During linear evolution, if a new clone fails to outcompete its predecessor, or

predecessors, a degree of heterogeneity will be observed³⁸. Branched tumour evolution, in which distinct subclones evolve in parallel, however, will result in extensive subclonal diversity³.

A tumour's subclonal architecture can be reconstructed from sequencing approaches^{22,36,39–41} or multiplex fluorescence *in situ* hybridization (FISH)², providing insight into cancer evolution. Branched evolution has now been documented in a range of tumour types, including adenoma-to-carcinoma transition of the colon⁴², childhood acute lymphoblastic leukaemia (ALL)², chronic lymphoblastic leukaemia (CLL)¹, pancreatic cancer³⁵ and breast cancer^{39,41}. In cases of ALL that exhibit branched evolutionary trajectories, heterogeneity has been observed between leukaemia-propagating cells². This has important implications for cancer stem-cell models because it suggests that each subclonal branch was probably sustained by genetically distinct leukaemia-propagating cells². It is conceivable that genetically heterogeneous subclonal lineages in solid tumours could also be maintained by distinct stem cells (See Review by Meacham and Morrison on page 328).

Branched evolutionary trajectories with spatial separation of tumour subclones (Fig. 1) have also been described. In clear-cell renal cell carcinoma, sequencing multiple biopsies from the same primary tumour revealed spatially separated subclones, harbouring heterogeneous somatic mutations and copy-number events⁴⁰. Likewise, multiregion sampling in glioblastoma documented heterogeneous copy number events between different regions of the same tumour⁴³. Evidence for clonal diversity between primary and metastatic sites has also been demonstrated in breast cancer³⁶, pancreatic cancer^{21,35} and medulloblastoma⁴⁴. Intratumour heterogeneity of spatially separated subclones, at the DNA copy-number level, may be so profound that the copy number profiles of single tumour biopsies may more closely resemble those of tumours from different patients than those of adjacent biopsies of the same tumour⁴⁵.

At metastatic sites, the new micro-environment, together with the selection pressures imposed by the process of metastasis itself, offer a plausible explanation for genetic differences between metastatic and primary sites⁴⁶ (see Perspective by Klein on page 365). However, it is

BOX 1

Main pathways implicated in genomic instability

- **Base and nucleotide excision repair.** These pathways function to excise and repair abnormal bases or nucleotides, such as deaminated cytosines or ultraviolet radiation induced pyrimidine dimers. Germline mutations in components of these pathways predispose people to colonic polyposis⁶⁴ or skin cancers⁶⁶.
- **Mismatch repair.** This process acts during DNA replication to correct base mismatches, as well as insertions and deletions at repetitive sequences (microsatellites)⁶⁸. Loss of function of *MSH2* and *MLH1*, which are required for mismatch repair, results in hypermutation and microsatellite instability. Mutations abrogating the proofreading ability of the DNA polymerases δ and ϵ increase base mismatches without affecting microsatellite stability¹²⁶⁷.
- **Telomere maintenance.** Telomere erosion or uncapping results in catastrophic chromosomal instability through end-on chromosome fusions, followed by bridging during mitosis and further breakage, in 'breakage-fusion-bridge' cycles. Reactivation of telomerase expression in cancer cells can alleviate this instability⁹⁵.
- **Double-strand break repair.** Homologous recombination repair of double-strand breaks (DSBs) uses the sister DNA molecule as a template to repair the break, and is important for repair of stalled or collapsed replication forks. Non-homologous end joining directly ligates

the DSB in an error-prone manner. Defects in DSB detection or repair result in chromosomal instability^{8,81}, with an elevated rate of structural chromosomal rearrangements¹²⁴ and point mutations⁷⁰.

- **DNA replication.** Deregulated DNA replication can result in replication fork stalling, reversal and collapse. Deregulation can occur through oncogene activation, loss of certain tumour suppressors, DNA polymerase inhibition, nucleoside imbalances, replication-blocking DNA lesions and clashes of replication forks with ongoing transcription. Such replication stress can trigger DNA DSB formation (particularly at genomic fragile sites), unscheduled recombination events and chromosomal rearrangements^{26-28,87,97,125}.

- **Chromosome segregation.** Defects in chromosome segregation can arise directly through defects in the mitotic checkpoint^{73,74}, sister chromatid cohesion¹¹⁸, spindle geometry^{75,78,79} and spindle dynamics^{76,77}. These defects result in aberrant chromosome-spindle attachments, and missegregation of chromatids during anaphase, generating aneuploid daughter cells. Indirect segregation defects can be caused by structural chromosome rearrangements generated before mitosis, via replication stress^{87,89}, defective repair⁹⁰ or telomere fusion⁹⁰. Further damage to missegregating chromosomes can occur during cytokinesis⁸⁵ or aberrant replication and fragmentation of micronuclei⁸⁶.

unclear precisely how spatial separation of genetically distinct clones arises in primary tumours⁴⁰. It is possible that the separation reflects the presence of distinct micro-environmental niches in the primary tumour (see Junttila and de Sauvage on page 346), such that the sub-clones occupying each niche evolve relatively independently of one another. The similarity between this mode of tumour evolution and allopatric speciation (whereby geographically isolated populations from the same ancestral species evolve independently) raises the possibility that physical barriers in the tumour — such as blood vessels or tissue planes — prevent the mixing of tumour subclones (Fig. 1).

Nevertheless, intermingled heterogeneous clones are observed within single biopsies^{22,39,41,47} to the extent that neighbouring cells can display amplification of different receptor tyrosine kinases — proteins that regulate key signalling processes and whose dysregulation has a crucial role in tumorigenesis^{48,49}. At this intercellular level, genomic instability is thought to play a crucial part in generating enhanced genetic diversity. This intercellular heterogeneity increases phenotypic variation, broadening the pool of cells that are subject to selection, and therefore the likelihood of selective expansion of multiple different subclones and the emergence of a complex subclonal tumour architecture⁴.

Genomic instability and tumour evolution

The genome is replicated and divided with high fidelity, such that the endogenous point-mutation rate in somatic cells is relatively low (0.77×10^{-9} per site per cell division⁵⁰) and errors in chromosome segregation are both rare (roughly 1 per 100 cell divisions) and poorly tolerated in non-transformed proliferating cells⁵¹⁻⁵³. Disruption of mechanisms that maintain genome integrity or exposure to exogenous mutagens will elevate the mutation rate (mutation here is used in the broadest sense to include any genome aberration). Most solid tumours and haematopoietic malignancies display at least one form of genomic instability⁵⁴.

Despite the clear association between genomic instability and cancer, there has been considerable debate as to whether elevated mutation rates are necessary for tumorigenesis^{55,56}. Even with a normal somatic point mutation rate, the diversity and mutational burden within a large tumour cell population is likely to be extensive⁵⁵, and sequencing

studies in AML and colorectal cancer have found that there is no need to invoke elevated mutation rates to explain the number of mutations observed^{20,38}. Nevertheless, although not strictly necessary for carcinogenesis, the observation that most tumours display some form of instability suggests that it provides some benefit to tumour evolution. This hypothesis is supported by mouse models, which have revealed that genomic instability increases the risk of carcinogenesis^{57,58}, leads to tumour regrowth after oncogene withdrawal⁵⁹ and promotes loss of heterozygosity of tumour suppressor genes⁶⁰.

Different genomic instabilities result in different genetic lesions, ranging from increased point mutation frequency to small insertions and deletions, large-scale chromosomal rearrangements and alterations in ploidy (see Box 1 and Table 1). However, the phenotypic penetrance of these genetic changes may vary; a chromosome gain or loss is more likely to have functional consequences than a point mutation, most of which are neutral. As different patterns of genomic instability have distinct genomic footprints, it is possible to interrogate sequencing and copy-number data to examine how genomic instability has shaped tumour growth and evolution⁶¹⁻⁶³. Given that most studies rely on examining a pooled population of cells, the true level of intratumour genetic heterogeneity driven by genomic instability is likely to be much greater than estimated, owing to both sampling bias and the inability to resolve heterogeneity between single cells. A recent study analysed an extensive catalogue of mutations in 30 of the most common cancer types and uncovered more than 20 signatures of processes that mutate DNA, driven by both exogenous and endogenous factors⁶³.

In addition to global effects on genome architecture and mutational patterns, specific mutations may occur at an increased rate in the context of a particular instability mechanism. This is illustrated in colorectal tumours, in which those with defective mismatch repair exhibit a strikingly different mutational profile from those tumours with intact repair¹². Similarly, colorectal tumours with defective base excision repair, due to *MYH* (also known as *MUTYH*) mutations (Box 1) show a preponderance of C:G→A:T transversions in *APC*⁶⁴. In lung cancer, smoking increases the mutation rate and the frequency of C:G→A:T transversions²² such that mutations in *TP53* can often be directly attributed to DNA damage from cigarette smoke carcinogens⁶⁵. This effect

is likely to be compounded by epistasis, whereby a given mutation that occurs more frequently in the context of one pattern of instability may then lead to selection for specific combinations of mutations.

Mechanisms of genomic instability

Increased point mutation frequencies can be caused by both exogenous factors^{22,63} and defects in DNA repair pathways, such as mismatch repair and base and nucleotide excision repair^{12,64,66–68} (see Box 1 and Table 1). In addition to hypermutation, mismatch repair deficiency is associated with expansion and contraction of repetitive tracts of DNA (microsatellite instability), leading to an increased frequency of frameshift mutations⁶⁸. Elevated point mutation frequencies may also arise through mechanisms that affect gross chromosome structure, such as DNA replication stress⁶⁹ and defects in homologous recombination⁷⁰.

Different mechanisms of instability can lead to particular distributions of point mutations — replication stress is associated with an enrichment of mutations in large genes⁶⁹, whereas microsatellite instability may affect specific genes more frequently than others owing to the nucleotide composition of those genes⁶⁸. Highly localized clusters of mutations that are associated with chromosomal rearrangement, known as kataegis, may be caused by the action of the APOBEC family of cytidine deaminases in breast cancer⁶¹. A recent study has highlighted the need to consider variation in mutation rate across the genome when identifying putative driver mutations; mutation frequency across the genome was found to strongly correlate with DNA replication timing and transcriptional activity¹¹.

Chromosomal instability occurs at high frequencies across many cancer types and is associated with aggressive disease, drug resistance and poor prognosis^{54,71}. This form of instability refers to an increased rate of change in chromosome number or structure, and results in a range of karyotypic abnormalities, including whole-chromosome and segmental aneuploidies, as well as translocations, inversions and deletions (Table 1). A substantial proportion of chromosomally unstable tumours also show increased ploidy, leading to suggestions that genome doubling could act as a precursor to chromosomal instability⁷².

Chromosomal instability can arise through a range of both mitotic and pre-mitotic defects. Mitotic defects result in inaccurate partitioning of

whole chromosomes between daughter cells — mediated through incorrect attachment of chromosomes to the mitotic spindle. This may occur through inadequate detection and correction^{73,74}, or through increased formation, of incorrect attachments as a result of altered microtubule dynamics or transiently abnormal spindle geometry^{75–79}. Pre-mitotic defects that converge on defective repair or increased generation of DNA damage can affect chromosome structural integrity. DNA damage may be acquired from exogenous sources (such as radiation), but in general chromosomal rearrangements in cancer cells are thought to arise through endogenous DNA damage. This may be, for example, through telomere dysfunction⁸⁰ or defects in DNA repair pathways (such as *BRCA1* and *BRCA2* mutations, which result in defective homologous recombination^{8,81}; Box 1). A likely source of much of the DNA damage in tumours is DNA replication stress^{26,27} (Box 1), which is reflected by the enrichment of chromosome breakpoints and loss of heterozygosity in cancer genomes at fragile sites (regions of the genome that are particularly susceptible to replication stress)^{82–84}.

In recent years, there has been a growing appreciation that, from a mechanistic perspective, numerical and structural instability cannot be considered in isolation^{80,85–87}. Mitotic defects in chromosome attachment, typically thought to lead to whole-chromosome aneuploidies, have been shown to result in DNA damage and structural rearrangements, through entrapment of chromosomes in the cytokinesis furrow⁸⁵ or through aberrant replication and fragmentation of micronuclei⁸⁶. Conversely, structural chromosome rearrangements can also compromise the accuracy of chromosome segregation during mitosis^{80,87–90}. Therefore, the relative contributions of mitotic defects and pre-mitotically generated structural chromosome abnormalities to chromosomal instability in tumours need to be explored.

Variation in genomic instability over time

Tumour evolution is a dynamic process that contributes to both spatial and temporal tumour diversity. The type of genomic instability operating in a tumour at any point in time is likely to have a significant impact on tumour evolution, not only in terms of which specific pathways are dysfunctional⁹¹, but also in terms of the likelihood of a tumour

Table 1 | Characteristic features of cancer genomes

Genomic feature	Description	Possible mechanisms
Whole-genome duplication	Duplication of the entire DNA complement	Cell fusions, cytokinesis failure, metaphase or anaphase defects, or endoreduplication ⁷²
Chromosomal loss and gain	Loss and gain of whole chromosomes and segments of chromosomes	Mitotic defects: abnormal spindle geometry ^{75,78,79} , centriole amplification ^{75,78} , mitotic checkpoint defects ^{73,74} or hyperstable microtubules ^{76,77} Pre-mitotic defects: cohesin mutations ¹¹⁸ , DNA replication stress ^{27,28,82,87} , DNA damage repair defects ⁸¹ or telomere dysfunction ^{80,95}
Chromothripsis	Extensive genomic rearrangements involving one or more chromosomes	Ionizing radiation, micronucleus fragmentation or replication defects ^{86,94,119}
Translocations	Chromosomal rearrangements resulting in fusion of two chromosome arms	Non-homologous end joining, aberrant lymphocyte DNA recombination mechanisms and cytokinesis-induced DNA damage ^{84,85,120}
Tandem duplications	Sequential copies of a genomic segment that often spans several genes	Unknown ^{121,122}
Loss of heterozygosity at fragile sites	Copy number imbalance or gene conversion at genomic loci that are particularly sensitive to replication stress	DNA replication stress ^{27,82}
Focal deletions	Deletion of less than 5 megabases	Chromosome breakage or DNA replication stress ⁸²
Focal amplification	Amplification less than 5 megabases	Double minute chromosomes generated through chromosome breakage ¹²³
Hypermutation	Elevated mutation rate, leading to abundance of somatic mutations	DNA mismatch repair defects ⁶⁸ , <i>BRCA2</i> mutations ⁷⁰ and proofreading defects ⁶⁷
Kataegis	Localized hypermutation	AID/APOBEC activity ⁶¹
C-G>T-A transitions	Base changes	Ultraviolet radiation-damage ⁶³ , alkylating agents ⁷ and spontaneous deamination of methylated cytosines ⁵⁰
C-G>A-T transversions	Base changes	Base excision repair defects ⁶⁴ , tobacco-related DNA damage ²² and cytotoxic chemotherapy ⁶

metastasizing⁹² and its response to treatment^{7,93}.

Recent studies have revealed evidence for changing patterns of instability during tumour progression^{31,39,40}. For example, in a study of clear-cell renal carcinoma, we found that chromosomally unstable metastases were probably derived from a tetraploid subclone in the primary tumour, whereas the remainder of the sampled regions in the primary tumour were diploid⁴⁰. In a study of breast cancer, researchers made use of variant allele frequencies and copy number events that were derived from whole-genome sequencing reads to dissect the temporal sequence of mutations³⁹. The mutations that occurred early in cancer development — on the trunk of the tumour's evolutionary tree — were found to exhibit a mutation pattern that was largely similar to germline mutations⁵⁰. However, the mutations that occurred later during tumour progression had significantly altered mutational patterns, involving decreased frequencies of C-G→T-A base transitions³⁹. These studies suggest that temporally separated biopsies may give strikingly different pictures of a tumour's genomic landscape.

Sequencing studies have also allowed researchers to analyse how tumour evolution and genomic instability could be affected by therapy. The analysis of pairs of primary and relapsed tumour cells from eight patients with AML found relapse-specific mutations and an increase in base transversions after chemotherapy, highlighting how treatment itself can act as a transient initiator of exogenous instability⁶. A study of nine glioblastomas from patients who had undergone chemotherapy with the alkylating agent temozolomide found that two cases exhibited an elevated frequency of mutations with a pattern that was consistent with induction by alkylating agents⁷. Both of these hypermutator gliomas relapsed after chemotherapy, and mutations in the mismatch repair gene *MSH6* (Box 1) were detected in the relapsed but not in pretreatment samples. This suggests that treatment with alkylating agents can impose selection for loss of mismatch repair function, allowing propagation of therapy-induced mutations. Because chromosomal instability is associated with multidrug resistance⁹³ and poor prognosis⁷¹, this raises the possibility that this form of genome instability may also be selected for during treatment.

It is important to note that observations of high numbers of mutations or chromosomal rearrangements do not necessarily imply ongoing genomic instability, but may instead reflect a historical prolonged period of mutational acquisition, or alternatively the occurrence of transient mutational bursts^{38,94}. The evolution of the cancer genome may proceed not only in a gradual manner through clonal selection of point mutations or chromosomal alterations, but may also be subject to punctuated phases characterized by drastic changes. It has been postulated that chromothripsis, in which tens to hundreds of chromosomal rearrangements occur across one or a few chromosomes during a single catastrophic event, represents a major evolutionary leap in terms of tumour development⁹⁴. Similarly, telomere dysfunction catalysed by eroded chromosome ends may be transient, leading to catastrophic instability until telomerase expression is activated to stabilize chromosome ends⁹⁵ (Box 1). Nevertheless, although these punctuated events are likely to contribute to cancer-genome evolution, observations of an active DNA damage response, and variation in chromosome copy number in both early stage and advanced malignancies, suggest that ongoing instability occurs in a substantial proportion of tumours^{27,96,97}.

Drastic genomic change may also result from genome doubling, which has been estimated to occur at frequencies as high as 60% in some cancer types⁹⁸. Some studies have proposed that the process is an intermediate stage on the path to aneuploidy, precipitating genetic and functional intratumour heterogeneity through chromosomal instability (see ref. 72). Although genome doubling has been observed as an early event in cancer development³⁹, recent sequencing studies have found evidence for it at later stages in disease progression³⁹. Genome doubling has been observed after, rather than before, the onset of chromosomal instability, indicating that there may be additional evolutionary benefits to genome doubling beyond catalysing chromosome reassortment^{39,98}. Furthermore, genome doublings are proposed to play a major part during the evolution of eukaryotes, creating evolutionary opportunities

by providing new genetic material for mutation and selection to act upon¹⁰⁰. In tumour evolution, it is possible that both genome doublings and tandem duplications (Table 1) might facilitate subfunctionalization of duplicated genes.

Intertumour variation in genomic instability

There is considerable heterogeneity in both the patterns and extent of genomic instability that are prevalent in different tumour types. Microsatellite instability is observed most frequently in colorectal tumours; however, it occurs only in a minority of these tumours¹⁶. This instability seems to preferentially occur in tumours of the proximal colon, suggesting an association with epigenetic or environmental factors⁶⁸. By contrast, defective homologous recombination is observed more frequently in breast and ovarian cancers⁸. A given pattern of instability can be caused by multiple mechanisms; for example, chromosomal instability may be driven by different defects in different tumours^{52,75,87}. Furthermore, based on cell-line studies, multiple defects can operate simultaneously in a given tumour cell^{52,75,78,87}.

Different instability mechanisms are also associated with variations in clinical outcome, for example, colorectal tumours with microsatellite instability are associated with a better patient prognosis compared with chromosomally unstable tumours⁷. Differences in disease outcome could relate to the profound infiltration of immune cells observed in microsatellite unstable tumours compared with colorectal cancers that display chromosomal instability. This immune infiltrate may be driven by an increased repertoire of neoantigens in these tumours as a result of mismatch repair deficiency¹⁰¹. Alternatively, the hypermutator phenotype of microsatellite-instability tumours could place them at the limit of tolerable instability and treatment with mutagenic chemotherapy may push them over the threshold⁴. Consistent with this idea, *in vitro*, microsatellite-instability colorectal cancer cell lines are more sensitive than chromosomally unstable cell lines to a range of different targeted and cytotoxic agents⁹³. Whatever the explanation, these observations suggest that not all forms of genetic heterogeneity are equal in terms of their impact on patient outcome.

In addition to variation in the pattern of instability between tumours, variation also occurs in the degree of instability between tumours. For example, in breast cancer, oestrogen receptor positive (ER⁺) tumours display significantly lower karyotypic complexity (a genomic footprint of chromosomal instability), compared to ER⁻ tumours⁹⁶. This variation in the degree of chromosomal instability is associated with different outcomes; we have found evidence in ER⁻ breast cancer, squamous non-small-cell lung cancer, and serous ovarian and gastric cancer that tumours within the highest quartile of chromosomal instability have a better prognosis than those in intermediate quartiles of chromosomal instability^{96,102}. These data suggest that the benefits of genomic instability must be balanced against its costs, although it is unclear whether the link with prognosis is independent of treatment or whether therapy itself can push the level of instability beyond the tolerable threshold.

An elevated mutation rate not only increases the supply of potentially beneficial mutations but also deleterious ones, and there may be an optimal level of instability, which if exceeded impedes tumour growth, a scenario analogous to mutational meltdown in bacteria or error catastrophe in viruses⁴. Examples of excessive instability resulting in cell lethality can be observed in the context of mitotic checkpoint dysfunction^{57,103} or multipolar anaphases in cells with extra centrosomes⁷⁵. Cells may adapt to limit catastrophic instability by clustering extra centrosomes to form a bipolar spindle and avoid lethal multipolar division^{75,78}, and so efforts to prevent centrosome clustering for therapeutic purposes to exacerbate instability may be effective⁵⁴. Similarly, excessive instability in the context of *BRCA1* and *BRCA2* mutations may be averted by downregulation of the p53-binding protein 53BP1, to rescue, in part, homologous recombination repair¹⁰⁴. By understanding cellular processes and pathways such as these — which are required for tolerance or averting excessive levels of genomic instability

— therapeutic opportunities may be revealed. This is exemplified by the efficacy of poly(ADP-ribose) polymerase inhibitors in breast and ovarian cancers that bear *BRCA1* and *BRCA2* mutations^{8,105,106}. Other possible means of targeting genomic instability include inhibiting replication checkpoints in cells undergoing replication stress¹⁰⁷ and perturbing pathways that are essential for the survival of aneuploid cells^{51,53,93}. Further research is required to identify new strategies to target genomic instability in cancer, which may represent an attractive approach to treating heterogeneous tumours.

Consequences of intratumour heterogeneity

Similar to that observed between different tumours, some degree of phenotypic convergence may also occur between genetically distinct tumour subclones⁴⁰. However, intratumour genetic heterogeneity also results in phenotypic diversity, and this has been documented in a range of tumour types, affecting clinically relevant parameters such as gene-expression signatures that reflect prognosis, and response to therapeutic agents^{6,40,48,108}. In renal cancer, distinct transcriptomic profiles that reflect both good and poor prognosis have been described within one primary tumour⁴⁰, and sampling of 11 glioblastomas revealed that many patients displayed multiple transcriptomic subtypes within different biopsies from the same tumour⁴³.

Increasing evidence suggests that efforts to forecast outcome require the identification of low-frequency genetically and functionally distinct subclones at diagnosis^{1,37}. This is illustrated by a recent study in chronic lymphocytic leukaemia, which revealed the selection of subclones that harbour distinct driver events during therapy¹. The eight patients that harboured a detectable subclonal driver in their pretreatment samples showed significantly shorter time to re-treatment or death compared with the four patients with no evidence of subclonal driver events. Similarly, in a case of multiple myeloma, a subclone that was barely detectable at diagnosis was shown to be dominant towards the end of the disease³⁷. These data underscore the need for longitudinal tumour sampling over the disease course and throughout treatment because the subclone that influences outcome may not be detectable in a single biopsy, or indeed even in multiple biopsies, taken before treatment.

Subclonal driver mutations result directly in phenotypic variation, for example in alterations to signal transduction pathways. In clear-cell renal cell cancer, a heterogeneous *MTOR* mutation (resulting in constitutive mTOR kinase activity *in vitro*) was associated with differential mTOR signalling pathway activation after 6 weeks of mTOR inhibitor therapy⁴⁰. In a heterogeneous glioblastoma multiforme cell population harbouring mosaic amplification of *EGFR* and *PDGFRA*, combined inhibition of both kinases was necessary to attenuate PI(3)K pathway activity *in vitro*⁴⁸. Intermingling of tumour cells harbouring amplification of distinct receptor tyrosine kinases (cells harbouring amplification of both genes were rare) hints at a possible functional codependency between the two genetically distinct populations^{48,49}, as has been demonstrated in models using the fruitfly *Drosophila melanogaster*¹⁰⁹. Intriguingly, *EGFR* mutant tissue, cell lines and stem cells release interleukin 6 (IL-6) and leukaemia inhibitory factor, which contribute to enhanced growth of adjacent subclones through activation of wild-type *EGFR*¹⁰⁸, providing further evidence of possible codependencies between minor and major subclones in tumours.

However, it is important to note that phenotypic heterogeneity is not only mediated through genetic diversity. Genetically homogeneous subclones can behave in functionally distinct ways after exposure to chemotherapy, mediated by varied patterns of proliferation, with dormant resting cells surviving cytotoxic exposure¹¹⁰. Phenotypic heterogeneity is not determined solely through genetic distinctions between subclones, but also through stochastic events in gene expression and protein stability, epigenetic divergence and micro-environmental fluctuations^{30,31,110,111}. Indeed, a study examining the fate of daughter cells derived from the same mother cell after antimetabolic drug exposure found that the fates of the sister cells seemed to be independent of each other¹¹².

Intratumour heterogeneity and treatment strategies

Observations of intratumour heterogeneity pose a challenge to targeted therapies and raise important questions regarding future drug-development strategies, which may benefit from considering whether clonally dominant events may be more effective targets than subclonal lesions²⁹. Founder events might include *EGFR* and *KRAS* mutations in lung cancer, VHL loss-of-function through mutation, methylation or homozygous copy number loss in clear-cell renal cell carcinoma⁴⁰, and *KRAS* and *BRAF* mutations in colorectal carcinoma¹¹³. Of these examples, only *EGFR* and *BRAF* mutations are currently directly targetable.

In addition, targeting clonally dominant genetic events is complicated by observations that a founder event in one tumour can be a later event in others. For example, in lung cancer, activating *EGFR* mutations were found to display intratumour heterogeneity and spatial discordance in 6.3–28% of cases¹¹⁴, suggesting that in these tumours *EGFR* mutation was not a founder event. Even in tumours in which *EGFR* mutations are founder events, the presence of low-frequency subclones harbouring the T790M gatekeeper mutation, which confers resistance to *EGFR* tyrosine kinase inhibitors, have been shown to be associated with early disease progression in patients treated with *EGFR* inhibitors¹¹⁴. In patients with colorectal cancer who have undergone anti-*EGFR* antibody therapy, mutations in *KRAS* (which confer drug resistance) were detectable in circulating tumour DNA 5–6 months after therapy in 9 out of 24 tumours, 3 of which developed multiple distinct *KRAS* mutations. Mathematical modelling suggested that low-frequency subclones harbouring these *KRAS* mutations were present in the tumour before treatment¹¹⁵. The same challenges are likely to affect the efficacy of other targeted therapies.

Future directions

As genomics technologies advance further and become more affordable, our understanding of how genomic instability shapes tumour evolution will be greatly enhanced. Both mathematical and experimental modelling approaches are also likely to inform our understanding of how heterogeneity affects tumour progression^{5,110,115,116}. Longitudinal tumour sampling approaches will be essential to decipher the impact of tumour heterogeneity on cancer evolution, and developing minimally invasive methods to profile heterogeneous tumour genomes will play a major part in deciphering changes in the genomic architecture of tumours in real time, exemplified by circulating free tumour DNA analysis¹¹⁵. In addition, there is a need to develop clinically useful measures of heterogeneity¹¹⁷. The prevalence of different patterns of instability will be more robustly assessed in large patient cohorts, and new patterns will probably be identified as the wealth of genomic data is analysed⁶³. It will be important to examine in greater depth the relationship between the rate of instability and the complexity of subclonal architectures, as well as whether certain forms of genomic instability are associated with a greater degree of intratumour heterogeneity. There is a crucial need to understand mechanisms driving genomic instability so that therapeutic approaches to limit cancer diversity, adaptation and drug resistance can be developed. ■

Received 19 April; accepted 13 July 2013.

- Landau, D. A. *et al.* Evolution and impact of subclonal mutations in chronic lymphocytic leukemia. *Cell* **152**, 714–726 (2013).
This article examines how subclonal driver mutations influence outcome.
- Anderson, K. *et al.* Genetic variegation of clonal architecture and propagating cells in leukaemia. *Nature* **469**, 356–361 (2011).
An article demonstrates branched cancer evolution and genetic heterogeneity between leukaemia-propagating cells in ALL.
- Greaves, M. & Maley, C. C. Clonal evolution in cancer. *Nature* **481**, 306–313 (2012).
- Cahill, D. P., Kinzler, K. W., Vogelstein, B. & Lengauer, C. Genetic instability and Darwinian selection in tumours. *Trends Biochem. Sci.* **24**, M57–M60 (1999).
An early reference placing genomic instability within the framework of Darwinian evolution, and recognizing that there may be distinct genomic footprints for different instability mechanisms.
- Merlo, L. M., Pepper, J. W., Reid, B. J. & Maley, C. C. Cancer as an evolutionary and ecological process. *Nature Rev. Cancer* **6**, 924–935 (2006).

6. Ding, L. *et al.* Clonal evolution in relapsed acute myeloid leukaemia revealed by whole-genome sequencing. *Nature* **481**, 506–510 (2012).
This article demonstrates the impact of treatment upon both tumour evolution and mutational spectra.
7. Hunter, C. *et al.* A hypermutation phenotype and somatic *MSH6* mutations in recurrent human malignant gliomas after alkylator chemotherapy. *Cancer Res.* **66**, 3987–3991 (2006).
8. Lord, C. J. & Ashworth, A. The DNA damage response and cancer therapy. *Nature* **481**, 287–294 (2012).
9. Vogelstein, B. *et al.* Cancer genome landscapes. *Science* **339**, 1546–1558 (2013).
10. Weigelt, B. & Reis-Filho, J. S. Histological and molecular types of breast cancer: is there a unifying taxonomy? *Nature Rev. Clin. Oncol.* **6**, 718–730 (2009).
11. Lawrence, M. S. *et al.* Mutational heterogeneity in cancer and the search for new cancer-associated genes. *Nature* **499**, 214–218 (2013).
12. The Cancer Genome Atlas Research Network. Comprehensive molecular characterization of human colon and rectal cancer. *Nature* **487**, 330–337 (2012).
13. Jiao, Y. *et al.* *DAXX/ATRX*, *MEN1*, and mTOR pathway genes are frequently altered in pancreatic neuroendocrine tumors. *Science* **331**, 1199–1203 (2011).
14. Brown, J. M. & Attardi, L. D. The role of apoptosis in cancer development and treatment response. *Nature Rev. Cancer* **5**, 231–237 (2005).
15. Prahallad, A. *et al.* Unresponsiveness of colon cancer to BRAF(V600E) inhibition through feedback activation of EGFR. *Nature* **483**, 100–103 (2012).
16. Lengauer, C., Kinzler, K. W. & Vogelstein, B. Genetic instability in colorectal cancers. *Nature* **386**, 623–627 (1997).
17. Popat, S., Hubner, R. & Houlston, R. S. Systematic review of microsatellite instability and colorectal cancer prognosis. *J. Clin. Oncol.* **23**, 609–618 (2005).
18. Pao, W. & Chmielecki, J. Rational, biologically based treatment of EGFR-mutant non-small-cell lung cancer. *Nature Rev. Cancer* **10**, 760–774 (2010).
19. Patel, J. P. *et al.* Prognostic relevance of integrated genetic profiling in acute myeloid leukemia. *N. Engl. J. Med.* **366**, 1079–1089 (2012).
20. Jones, S. *et al.* Comparative lesion sequencing provides insights into tumor evolution. *Proc. Natl Acad. Sci. USA* **105**, 4283–4288 (2008).
21. Yachida, S. *et al.* Distant metastasis occurs late during the genetic evolution of pancreatic cancer. *Nature* **467**, 1114–1117 (2010).
22. Govindan, R. *et al.* Genomic landscape of non-small cell lung cancer in smokers and never-smokers. *Cell* **150**, 1121–1134 (2012).
23. Hanahan, D. & Weinberg, R. A. The hallmarks of cancer. *Cell* **100**, 57–70 (2000).
24. Strauss, M., Lukas, J. & Bartek, J. Unrestricted cell cycling and cancer. *Nature Med.* **1**, 1245–1246 (1995).
25. Dalglish, G. L. *et al.* Systematic sequencing of renal carcinoma reveals inactivation of histone modifying genes. *Nature* **463**, 360–363 (2010).
26. Halazonetis, T. D., Gorgoulis, V. G. & Bartek, J. An oncogene-induced DNA damage model for cancer development. *Science* **319**, 1352–1355 (2008).
27. Bartkova, J. *et al.* DNA damage response as a candidate anti-cancer barrier in early human tumorigenesis. *Nature* **434**, 864–870 (2005).
28. Bartkova, J. *et al.* Oncogene-induced senescence is part of the tumorigenesis barrier imposed by DNA damage checkpoints. *Nature* **444**, 633–637 (2006).
29. Swanton, C. Intratumor heterogeneity: evolution through space and time. *Cancer Res.* **72**, 4875–4882 (2012).
30. Marusyk, A., Almendro, V. & Polyak, K. Intra-tumour heterogeneity: a looking glass for cancer? *Nature Rev. Cancer* **12**, 323–334 (2012).
31. Varley, K. E., Mutch, D. G., Edmonston, T. B., Goodfellow, P. J. & Mitra, R. D. Intra-tumour heterogeneity of *MLH1* promoter methylation revealed by deep single molecule bisulfite sequencing. *Nucleic Acids Res.* **37**, 4603–4612 (2009).
32. Gatenby, R. A., Gillies, R. J. & Brown, J. S. Of cancer and cave fish. *Nature Rev. Cancer* **11**, 237–238 (2011).
33. Nowell, P. C. The clonal evolution of tumor cell populations. *Science* **194**, 23–28 (1976).
34. Cairns, J. Mutation selection and the natural history of cancer. *Nature* **255**, 197–200 (1975).
35. Campbell, P. J. *et al.* The patterns and dynamics of genomic instability in metastatic pancreatic cancer. *Nature* **467**, 1109–1113 (2010).
36. Shah, S. P. *et al.* Mutational evolution in a lobular breast tumour profiled at single nucleotide resolution. *Nature* **461**, 809–813 (2009).
This article demonstrates clonal diversity between primary and metastatic sites using next-generation sequencing.
37. Keats, J. J. *et al.* Clonal competition with alternating dominance in multiple myeloma. *Blood* **120**, 1067–1076 (2012).
38. Welch, J. S. *et al.* The origin and evolution of mutations in acute myeloid leukemia. *Cell* **150**, 264–278 (2012).
39. Nik-Zainal, S. *et al.* The life history of 21 breast cancers. *Cell* **149**, 994–1007 (2012).
40. Gerlinger, M. *et al.* Intratumor heterogeneity and branched evolution revealed by multiregion sequencing. *N. Engl. J. Med.* **366**, 883–892 (2012).
41. Shah, S. P. *et al.* The clonal and mutational evolution spectrum of primary triple-negative breast cancers. *Nature* **486**, 395–399 (2012).
42. Thirlwell, C. *et al.* Clonality assessment and clonal ordering of individual neoplastic crypts shows polyclonality of colorectal adenomas. *Gastroenterology* **138**, 1441–1454 (2010).
43. Sottoriva, A. *et al.* Intratumor heterogeneity in human glioblastoma reflects cancer evolutionary dynamics. *Proc. Natl Acad. Sci. USA* **110**, 4009–4014 (2013).
44. Wu, X. *et al.* Clonal selection drives genetic divergence of metastatic medulloblastoma. *Nature* **482**, 529–533 (2012).
45. Martinez, P. *et al.* Parallel evolution of tumor subclones mimics diversity between tumors. *J. Pathol.* **230**, 356–364 (2013).
46. Yates, L. R. & Campbell, P. J. Evolution of the cancer genome. *Nature Rev. Genet.* **13**, 795–806 (2012).
47. Navin, N. *et al.* Tumour evolution inferred by single-cell sequencing. *Nature* **472**, 90–94 (2011).
48. Szerlip, N. J. *et al.* Intratumoral heterogeneity of receptor tyrosine kinases EGFR and PDGFRA amplification in glioblastoma defines subpopulations with distinct growth factor response. *Proc. Natl Acad. Sci. USA* **109**, 3041–3046 (2012).
49. Snuderl, M. *et al.* Mosaic amplification of multiple receptor tyrosine kinase genes in glioblastoma. *Cancer Cell* **20**, 810–817 (2011).
This article reports intermingled heterogeneous subclones with mutually exclusive amplification of targetable receptor tyrosine kinases.
50. Lynch, M. Rate, molecular spectrum, and consequences of human mutation. *Proc. Natl Acad. Sci. USA* **107**, 961–968 (2010).
51. Thompson, S. L. & Compton, D. A. Proliferation of aneuploid human cells is limited by a p53-dependent mechanism. *J. Cell Biol.* **188**, 369–381 (2010).
52. Thompson, S. L. & Compton, D. A. Examining the link between chromosomal instability and aneuploidy in human cells. *J. Cell Biol.* **180**, 665–672 (2008).
53. Torres, E. M., Williams, B. R. & Amon, A. Aneuploidy: cells losing their balance. *Genetics* **179**, 737–746 (2008).
54. Gordon, D. J., Resio, B. & Pellman, D. Causes and consequences of aneuploidy in cancer. *Nature Rev. Genet.* **13**, 189–203 (2012).
55. Tomlinson, I. P., Novelli, M. R. & Bodmer, W. F. The mutation rate and cancer. *Proc. Natl Acad. Sci. USA* **93**, 14800–14803 (1996).
56. Loeb, L. A. Mutator phenotype in cancer: origin and consequences. *Semin. Cancer Biol.* **20**, 279–280 (2010).
57. Weaver, B. A., Silk, A. D., Montagna, C., Verdier-Pinard, P. & Cleveland, D. W. Aneuploidy acts both oncogenically and as a tumor suppressor. *Cancer Cell* **11**, 25–36 (2007).
This article describes a mouse model that illustrates the complex relationship between the level of chromosomal instability and tumorigenesis.
58. Liu, X. *et al.* Somatic loss of BRCA1 and p53 in mice induces mammary tumors with features of human BRCA1-mutated basal-like breast cancer. *Proc. Natl Acad. Sci. USA* **104**, 12111–12116 (2007).
59. Sotillo, R., Schwartzman, J. M., Socci, N. D. & Benezra, R. Mad2-induced chromosome instability leads to lung tumour relapse after oncogene withdrawal. *Nature* **464**, 436–440 (2010).
60. Baker, D. J., Jin, F., Jegannathan, K. B. & van Deursen, J. M. Whole chromosome instability caused by Bub1 insufficiency drives tumorigenesis through tumor suppressor gene loss of heterozygosity. *Cancer Cell* **16**, 475–486 (2009).
61. Nik-Zainal, S. *et al.* Mutational processes molding the genomes of 21 breast cancers. *Cell* **149**, 979–993 (2012).
References 39 and 61 demonstrate how deep sequencing can be used to elucidate the evolutionary history of tumours, and develop tools to identify mutational signatures using whole-genome sequencing data; they also describe a novel highly localized mutational process, kataegis
62. Stephens, P. J. *et al.* The landscape of cancer genes and mutational processes in breast cancer. *Nature* **486**, 400–404 (2012).
63. Alexandrov, L. B. *et al.* Signatures of mutational processes in human cancer. *Nature* **500**, 415–421 (2013).
64. Sieber, O. M. *et al.* Multiple colorectal adenomas, classic adenomatous polyposis, and germ-line mutations in *MYH*. *N. Engl. J. Med.* **348**, 791–799 (2003).
65. Pfeifer, G. P. *et al.* Tobacco smoke carcinogens, DNA damage and p53 mutations in smoking-associated cancers. *Oncogene* **21**, 7435–7451 (2002).
66. Dumaz, N., Drougard, C., Sarasin, A. & Daya-Grosjean, L. Specific UV-induced mutation spectrum in the p53 gene of skin tumors from DNA-repair-deficient xeroderma pigmentosum patients. *Proc. Natl Acad. Sci. USA* **90**, 10529–10533 (1993).
67. Palles, C. *et al.* Germline mutations affecting the proofreading domains of POLE and POLD1 predispose to colorectal adenomas and carcinomas. *Nature Genet.* **45**, 136–144 (2013).
68. Shah, S. N., Hile, S. E. & Eckert, K. A. Defective mismatch repair, microsatellite mutation bias, and variability in clinical cancer phenotypes. *Cancer Res.* **70**, 431–435 (2010).
69. Nikolaev, S. I. *et al.* A single-nucleotide substitution mutator phenotype revealed by exome sequencing of human colon adenomas. *Cancer Res.* **72**, 6279–6289 (2012).
70. Yang, D. *et al.* Association of *BRCA1* and *BRCA2* mutations with survival, chemotherapy sensitivity, and gene mutator phenotype in patients with ovarian cancer. *J. Am. Med. Assoc.* **306**, 1557–1565 (2011).
71. McGranahan, N., Burrell, R. A., Endesfelder, D., Novelli, M. R. & Swanton, C. Cancer chromosomal instability: therapeutic and diagnostic challenges. *EMBO Rep.* **13**, 528–538 (2012).
72. Ganem, N. J., Storchova, Z. & Pellman, D. Tetraploidy, aneuploidy and cancer. *Curr. Opin. Genet. Dev.* **17**, 157–162 (2007).
73. Cahill, D. P. *et al.* Mutations of mitotic checkpoint genes in human cancers. *Nature* **392**, 300–303 (1998).
74. Sotillo, R. *et al.* Mad2 overexpression promotes aneuploidy and tumorigenesis in mice. *Cancer Cell* **11**, 9–23 (2007).
75. Ganem, N. J., Godinho, S. A. & Pellman, D. A mechanism linking extra centrosomes to chromosomal instability. *Nature* **460**, 278–282 (2009).
76. Bakhom, S. F., Genovese, G. & Compton, D. A. Deviant kinetochore microtubule dynamics underlie chromosomal instability. *Curr. Biol.* **19**, 1937–1942 (2009).
77. Bakhom, S. F., Thompson, S. L., Manning, A. L. & Compton, D. A. Genome stability is ensured by temporal control of kinetochore–microtubule dynamics. *Nature Cell Biol.* **11**, 27–35 (2008).

This article demonstrates direct attenuation of chromosomal instability in cancer cells, through overexpression of microtubule depolymerases.

78. Silkworth, W. T., Nardi, I. K., Scholl, L. M. & Cimini, D. Multipolar spindle pole coalescence is a major source of kinetochore mis-attachment and chromosome mis-segregation in cancer cells. *PLoS ONE* **4**, e6564 (2009).
 79. Kaseda, K., McAlinsh, A. D. & Cross, R. A. Dual pathway spindle assembly increases both the speed and the fidelity of mitosis. *Biol. Open* **1**, 12–18 (2012).
 80. Pampalona, J., Soler, D., Genescà, A. & Tusell, L. Whole chromosome loss is promoted by telomere dysfunction in primary cells. *Genes Chromosomes Cancer* **49**, 368–378 (2010).
 81. Hoelijmakers, J. H. Genome maintenance mechanisms for preventing cancer. *Nature* **411**, 366–374 (2001).
 82. Dereli-Öz, A., Versini, G. & Halazonetis, T. D. Studies of genomic copy number changes in human cancers reveal signatures of DNA replication stress. *Mol. Oncol.* **5**, 308–314 (2011).
 83. Bignell, G. R. et al. Signatures of mutation and selection in the cancer genome. *Nature* **463**, 893–898 (2010).
 84. Barlow, J. H. et al. Identification of early replicating fragile sites that contribute to genome instability. *Cell* **152**, 620–632 (2013).
 85. Janssen, A., van der Burg, M., Szuha, K., Kops, G. J. & Medema, R. H. Chromosome segregation errors as a cause of DNA damage and structural chromosome aberrations. *Science* **333**, 1895–1898 (2011).
 86. Crasta, K. et al. DNA breaks and chromosome pulverization from errors in mitosis. *Nature* **482**, 53–58 (2012).
- References 85 and 86 highlight different mechanisms that contribute to the complex interplay between structural and numerical chromosomal instability.**
87. Burrell, R. A. et al. Replication stress links structural and numerical cancer chromosomal instability. *Nature* **494**, 492–496 (2013).
 88. Gisselsson, D. Classification of chromosome segregation errors in cancer. *Chromosoma* **117**, 511–519 (2008).
 89. Kawabata, T. et al. Stalled fork rescue via dormant replication origins in unchallenged S phase promotes proper chromosome segregation and tumor suppression. *Mol. Cell* **41**, 543–553 (2011).
 90. Chan, K. L., North, P. S. & Hickson, I. D. BLM is required for faithful chromosome segregation and its localization defines a class of ultrafine anaphase bridges. *EMBO J.* **26**, 3397–3409 (2007).
- This study provides evidence that pre-mitotic defects can result in chromosome segregation errors.**
91. Markowitz, S. et al. Inactivation of the type II TGF- β receptor in colon cancer cells with microsatellite instability. *Science* **268**, 1336–1338 (1995).
 92. Guttenberg, A. et al. High chromosomal instability in brain metastases of colorectal carcinoma. *Cancer Genet. Cytogenet.* **198**, 47–51 (2010).
 93. Lee, A. J. et al. Chromosomal instability confers intrinsic multidrug resistance. *Cancer Res.* **71**, 1858–1870 (2011).
 94. Stephens, P. J. et al. Massive genomic rearrangement acquired in a single catastrophic event during cancer development. *Cell* **144**, 27–40 (2011).
 95. Artandi, S. E. & DePinho, R. A. Telomeres and telomerase in cancer. *Carcinogenesis* **31**, 9–18 (2010).
 96. Roylance, R. et al. Relationship of extreme chromosomal instability with long-term survival in a retrospective analysis of primary breast cancer. *Cancer Epidemiol. Biomarkers Prev.* **20**, 2183–2194 (2011).
 97. Gorgoulis, V. G. et al. Activation of the DNA damage checkpoint and genomic instability in human precancerous lesions. *Nature* **434**, 907–913 (2005).
 98. Carter, S. L. et al. Absolute quantification of somatic DNA alterations in human cancer. *Nature Biotechnol.* **30**, 413–421 (2012).
 99. Galipeau, P. C. et al. 17p (p53) allelic losses, 4N (G2/tetraploid) populations, and progression to aneuploidy in Barrett's esophagus. *Proc. Natl Acad. Sci. USA* **93**, 7081–7084 (1996).
 100. Taylor, J. S. & Raes, J. Duplication and divergence: the evolution of new genes and old ideas. *Annu. Rev. Genet.* **38**, 615–643 (2004).
 101. Ishikawa, T. et al. Tumor-specific immunological recognition of frameshift-mutated peptides in colon cancer with microsatellite instability. *Cancer Res.* **63**, 5564–5572 (2003).
 102. Birkbak, N. J. et al. Paradoxical relationship between chromosomal instability and survival outcome in cancer. *Cancer Res.* **71**, 3447–3452 (2011).
 103. Kops, G. J., Foltz, D. R. & Cleveland, D. W. Lethality to human cancer cells through massive chromosome loss by inhibition of the mitotic checkpoint. *Proc. Natl Acad. Sci. USA* **101**, 8699–8704 (2004).
 104. Bouwman, P. et al. 53BP1 loss rescues BRCA1 deficiency and is associated with triple-negative and BRCA-mutated breast cancers. *Nature Struct. Mol. Biol.* **17**, 688–695 (2010).
 105. Fong, P. C. et al. Inhibition of poly(ADP-ribose) polymerase in tumors from BRCA mutation carriers. *N. Engl. J. Med.* **361**, 123–134 (2009).
 106. Bryant, H. E. et al. Specific killing of BRCA2-deficient tumours with inhibitors of poly(ADP-ribose) polymerase. *Nature* **434**, 913–917 (2005).
 107. Ma, C. X., Janetka, J. W. & Piwnicka-Worms, H. Death by releasing the breaks: CHK1 inhibitors as cancer therapeutics. *Trends Mol. Med.* **17**, 88–96 (2011).
 108. Inda, M. M. et al. Tumor heterogeneity is an active process maintained by a mutant EGFR-induced cytokine circuit in glioblastoma. *Genes Dev.* **24**, 1731–1745 (2010).
 109. Wu, M., Pastor-Pareja, J. C. & Xu, T. Interaction between Ras(V12) and scribbled clones induces tumour growth and invasion. *Nature* **463**, 545–548 (2010).
 110. Kreso, A. et al. Variable clonal repopulation dynamics influence chemotherapy response in colorectal cancer. *Science* **339**, 543–548 (2013).
 111. Meads, M. B., Gatenby, R. A. & Dalton, W. S. Environment-mediated drug resistance: a major contributor to minimal residual disease. *Nature Rev. Cancer* **9**, 665–674 (2009).
 112. Gascoigne, K. E. & Taylor, S. S. Cancer cells display profound intra- and interline variation following prolonged exposure to antimetabolic drugs. *Cancer Cell* **14**, 111–122 (2008).
 113. Vakiani, E. et al. Comparative genomic analysis of primary versus metastatic colorectal carcinomas. *J. Clin. Oncol.* **30**, 2956–2962 (2012).
 114. Chen, Z. Y. et al. EGFR mutation heterogeneity and the mixed response to EGFR tyrosine kinase inhibitors of lung adenocarcinomas. *Oncologist* **17**, 978–985 (2012).
 115. Diaz, L. A. et al. The molecular evolution of acquired resistance to targeted EGFR blockade in colorectal cancers. *Nature* **486**, 537–540 (2012).
 116. Ding, L. et al. Genome remodelling in a basal-like breast cancer metastasis and xenograft. *Nature* **464**, 999–1005 (2010).
 117. Merlo, L. M. et al. A comprehensive survey of clonal diversity measures in Barrett's esophagus as biomarkers of progression to esophageal adenocarcinoma. *Cancer Prev. Res. (Phila.)* **3**, 1388–1397 (2010).
- This study develops tools to quantify intratumour heterogeneity, and relates these indices to patient outcome.**
118. Solomon, D. A. et al. Mutational inactivation of STAG2 causes aneuploidy in human cancer. *Science* **333**, 1039–1043 (2011).
 119. Liu, P. et al. Chromosome catastrophes involve replication mechanisms generating complex genomic rearrangements. *Cell* **146**, 889–903 (2011).
 120. Wang, J. H. Mechanisms and impacts of chromosomal translocations in cancers. *Front. Med.* **6**, 263–274 (2012).
 121. McBride, D. J. et al. Tandem duplication of chromosomal segments is common in ovarian and breast cancer genomes. *J. Pathol.* **227**, 446–455 (2012).
 122. Ng, C. K. et al. The role of tandem duplicator phenotype in tumour evolution in high-grade serous ovarian cancer. *J. Pathol.* **226**, 703–712 (2012).
 123. Hahn, P. J. Molecular biology of double-minute chromosomes. *Bioessays* **15**, 477–484 (1993).
 124. Xia, F. et al. Deficiency of human BRCA2 leads to impaired homologous recombination but maintains normal nonhomologous end joining. *Proc. Natl Acad. Sci. USA* **98**, 8644–8649 (2001).
 125. Bermejo, R., Lai, M. S. & Foiani, M. Preventing replication stress to maintain genome stability: resolving conflicts between replication and transcription. *Mol. Cell* **45**, 710–718 (2012).

Acknowledgements We would like to apologize to those whose work we have not been able to cite owing to space limitations. C.S. is funded by Cancer Research UK, The Medical Research Council, EU FP7 (projects PREDICT and RESPONSIFY), Prostate Cancer Foundation and the Breast Cancer Research Foundation (BCRF). J.B. is funded by the Danish Cancer Society and the European Commission (projects DDResponse and Biomedreg). We thank E. Grönroos and P. Gorman for the use of the tumour section image in Fig. 1.

Author Information Reprints and permissions information is available at www.nature.com/reprints. The authors declare no competing financial interests. Readers are welcome to comment on the online version of this article at go.nature.com/8sjatd. Correspondence should be addressed to C.S. (charles.swanton@cancer.org.uk).