



US010471160B2

(12) **United States Patent**
Eder et al.

(10) **Patent No.:** **US 10,471,160 B2**
(45) **Date of Patent:** **Nov. 12, 2019**

(54) **LABELED INHIBITORS OF PROSTATE SPECIFIC MEMBRANE ANTIGEN (PSMA), THEIR USE AS IMAGING AGENTS AND PHARMACEUTICAL AGENTS FOR THE TREATMENT OF PROSTATE CANCER**

(71) Applicants: **DEUTSCHES KREBSFORSCHUNGSZENTRUM**, Heidelberg (DE); **RUPRECHT-KARLS-UNIVERSITÄT HEIDELBERG**, Heidelberg (DE)

(72) Inventors: **Matthias Eder**, Mannheim (DE); **Klaus Kopka**, Dossenheim (DE); **Martin Schäfer**, Neckarsteinach (DE); **Ulrike Bauder-Wüst**, Schriesheim (DE); **Uwe Haberkorn**, Schwetzingen (DE); **Michael Eisenhut**, Heidelberg (DE); **Walter Mier**, Bensheim (DE); **Martina Benesova**, Heidelberg (DE); **Hans-Christian Kliem**, Heppenheim (DE); **Clemens Kratochwil**, Hirschberg a.d.B (DE)

(73) Assignees: **DEUTSCHES KREBSFORSCHUNGSZENTRUM**, Heidelberg (DE); **RUPRECHT-KARLS-UNIVERSITÄT HEIDELBERG**, Heidelberg (DE)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **16/038,729**

(22) Filed: **Jul. 18, 2018**

(65) **Prior Publication Data**
US 2019/0008988 A1 Jan. 10, 2019

Related U.S. Application Data

(63) Continuation of application No. 15/131,118, filed on Apr. 18, 2016, now abandoned, which is a continuation-in-part of application No. PCT/EP2014/002808, filed on Oct. 17, 2014.

Foreign Application Priority Data

Oct. 18, 2013 (EP) 13004991
Jul. 3, 2014 (EP) 14175612

(51) **Int. Cl.**
A61K 51/04 (2006.01)
C07D 257/02 (2006.01)
C07B 59/00 (2006.01)
C07D 295/145 (2006.01)

(52) **U.S. Cl.**
CPC **A61K 51/0482** (2013.01); **C07B 59/002** (2013.01); **C07D 257/02** (2013.01); **C07D 295/145** (2013.01)

(58) **Field of Classification Search**
CPC **A61K 51/04**; **A61K 49/00**
See application file for complete search history.

References Cited

U.S. PATENT DOCUMENTS

4,691,024 A	9/1987	Kunikatsu et al.
4,713,249 A	12/1987	Schröder
5,103,018 A	4/1992	Motomichi et al.
5,266,333 A	11/1993	Cady et al.
5,418,982 A	5/1995	Kishi
5,627,165 A	5/1997	Glazier
5,795,877 A	8/1998	Jackson et al.
5,863,536 A	1/1999	Jackson et al.
5,866,679 A	2/1999	DeFeo-Jones et al.
5,902,817 A	5/1999	Jackson et al.
5,948,750 A	9/1999	Garsky et al.
5,962,237 A	10/1999	Ts'o et al.
5,962,521 A	10/1999	Jackson et al.
5,968,915 A	10/1999	Jackson et al.
5,998,362 A	12/1999	Feng et al.
6,054,444 A	4/2000	Jackson et al.
6,068,830 A	5/2000	Diamandis et al.
6,127,333 A	10/2000	Brady et al.
6,174,858 B1	1/2001	Brady et al.
6,177,404 B1	1/2001	DeFeo-Jones et al.
6,232,287 B1	5/2001	Ruoslahti et al.
6,265,540 B1	7/2001	Isaacs et al.
6,342,491 B1	1/2002	Dey et al.
6,355,611 B1	3/2002	Karki et al.
6,368,598 B1	4/2002	D'Amico et al.

(Continued)

FOREIGN PATENT DOCUMENTS

CA	2606138	10/2005
CN	101863924	10/2010

(Continued)

OTHER PUBLICATIONS

Aggarwal, S., Singh, P., Topaloglu, O., Isaacs, J. T., and Denmeade, S. R. (2006) A dimeric peptide that binds selectively to prostate-specific membrane antigen and inhibits its enzymatic activity. *Cancer Res* 66, 9171-7.

Zhou, J., Neale, J. H., Pomper, M. G., and Kozikowski, A. P. (2005) NAAG peptidase inhibitors and their potential for diagnosis and therapy. *Nat Rev Drug Discov* 4, 1015-26.

Yamaguchi, T., Lee, J., Uemura, H., Sasaki, T., Takahashi, N., Oka, T., Shizukuishi, K., Endou, H., Kubota, Y., and Inoue, T. (2005) Prostate cancer: a comparative study of 11C-choline PET and MR imaging combined with proton MR spectroscopy. *Eur J Nucl Med Mol Imaging* 32, 742-8.

(Continued)

Primary Examiner — Michael G. Hartley

Assistant Examiner — Jagadishwar R Samala

(74) *Attorney, Agent, or Firm* — The Marbury Law Group, PLLC

(57) ABSTRACT

The present invention generally relates to the field of radio-pharmaceuticals and their use in nuclear medicine as tracers, imaging agents and for the treatment of various disease states of prostate cancer. Thus, the present invention concerns compounds that are represented by the general Formulae (Ia) or (Ib).

28 Claims, 20 Drawing Sheets

Fusion Pharmaceuticals, Inc.

Exhibit 1025.001

(56)

References Cited

U.S. PATENT DOCUMENTS

6,391,305 B1	5/2002	Feng et al.	8,273,363 B2	9/2012	Zale et al.
6,428,785 B1	8/2002	Gocken	8,313,728 B2	11/2012	Leamon et al.
6,479,470 B1	11/2002	Kozikowski et al.	8,388,977 B2	3/2013	Low et al.
6,504,014 B1	1/2003	Isaacs et al.	8,394,922 B2	3/2013	Cheng et al.
6,511,676 B1	1/2003	Boulikas	8,404,817 B2	3/2013	Sherman et al.
6,518,033 B1	2/2003	Gromeier et al.	8,414,864 B2	4/2013	Cappelletti et al.
6,528,499 B1	3/2003	Kozikowski et al.	8,414,898 B2	4/2013	Afar et al.
6,548,260 B1	4/2003	Tewari	8,445,851 B2	5/2013	Rouso et al.
6,596,755 B2	7/2003	Burman et al.	8,450,290 B2	5/2013	Worm et al.
6,602,274 B1	8/2003	Chen	8,465,725 B2	6/2013	Babich et al.
6,613,793 B2	9/2003	Burman et al.	8,487,128 B2	7/2013	Babich et al.
6,692,724 B1	2/2004	Yang et al.	8,487,129 B2	7/2013	Babich et al.
6,875,886 B2	4/2005	Frangioni	8,491,896 B2	7/2013	Goldenberg et al.
6,946,133 B1	9/2005	Schlom et al.	8,507,434 B2	8/2013	Popel et al.
7,005,429 B2	2/2006	Dey et al.	8,557,772 B2	10/2013	Popel et al.
7,008,765 B1	3/2006	Bussemakers et al.	8,565,945 B2	10/2013	Babich et al.
7,041,786 B2	5/2006	Shailubhai et al.	8,603,499 B2	12/2013	Zale et al.
7,045,605 B2	5/2006	Bander et al.	8,603,500 B2	12/2013	Zale et al.
7,052,703 B1	5/2006	Pastan et al.	8,603,501 B2	12/2013	Zale et al.
7,060,284 B1	6/2006	Kaumaya	8,606,349 B2	12/2013	Rouso et al.
7,128,893 B2	10/2006	Leamon et al.	8,644,910 B2	2/2014	Rouso et al.
7,129,254 B2	10/2006	Berger et al.	8,685,416 B2	4/2014	Klinman et al.
7,147,837 B2	12/2006	Lauffer et al.	8,685,891 B2	4/2014	Muraca
7,153,841 B2	12/2006	Roncucci et al.	8,703,918 B2	4/2014	Colombatti et al.
7,160,885 B2	1/2007	Currie et al.	8,709,483 B2	4/2014	Farokhzad et al.
7,166,691 B2	1/2007	Koochekpour et al.	8,772,226 B2	6/2014	Denmeade et al.
7,192,586 B2	3/2007	Bander	8,772,459 B2	6/2014	Ho et al.
7,232,805 B2	6/2007	Weinschenker et al.	8,778,305 B2	7/2014	Pomper et al.
7,238,785 B2	7/2007	Govindan et al.	8,802,153 B2	8/2014	Cheng et al.
7,282,567 B2	10/2007	Goldenberg et al.	8,816,095 B2	8/2014	Brown et al.
7,361,338 B2	4/2008	Jakobovits et al.	8,834,842 B2	9/2014	Leamon et al.
7,381,745 B2	6/2008	Kozikowski et al.	8,840,865 B2	9/2014	Babich et al.
7,399,460 B2	7/2008	Wedeking et al.	8,852,630 B2	10/2014	Spiegel et al.
7,408,079 B2	8/2008	Pomper et al.	8,858,509 B2	10/2014	Spiegel et al.
7,468,354 B2	12/2008	Isaacs	8,865,126 B2	10/2014	Leamon et al.
7,485,299 B2	2/2009	Afar et al.	8,877,970 B2	11/2014	Zimmerman et al.
7,514,078 B2	4/2009	Bander et al.	8,901,294 B2	12/2014	Kim et al.
7,534,580 B2	5/2009	Reeves et al.	8,907,058 B2	12/2014	Low et al.
7,547,773 B2	6/2009	Schlom et al.	8,916,167 B2	12/2014	Low et al.
7,585,491 B2	9/2009	Govindan	8,921,378 B2	12/2014	Toermakaengas et al.
7,601,332 B2	10/2009	Vlahov et al.	8,926,944 B2	1/2015	Babich et al.
7,635,682 B2	12/2009	Denmeade et al.	8,926,945 B2	1/2015	Port et al.
7,638,122 B2	12/2009	Yu et al.	8,940,871 B2	1/2015	Wu et al.
7,638,525 B2	12/2009	Jiang et al.	8,946,388 B2	2/2015	Sahin et al.
7,659,395 B2	2/2010	Pajouheshs et al.	8,962,799 B2	2/2015	Babich et al.
7,662,795 B2	2/2010	Rodriguez et al.	8,987,319 B2	3/2015	Miller
7,691,962 B2	4/2010	Boyd et al.	9,006,415 B2	4/2015	Ren et al.
7,696,185 B2	4/2010	Berkman	9,029,340 B2	5/2015	Lupold et al.
7,713,944 B2	5/2010	Kinberger et al.	9,044,468 B2	6/2015	Pomper et al.
7,740,847 B2	6/2010	Allan et al.	9,056,841 B2	6/2015	Pomper et al.
7,767,202 B2	8/2010	Pardoll et al.	9,074,000 B2	7/2015	Scheinberg et al.
7,767,803 B2	8/2010	Diener et al.	9,120,837 B2	9/2015	Babich et al.
7,794,929 B2	9/2010	Baylin et al.	9,123,725 B2	9/2015	Cho et al.
7,862,798 B2	1/2011	Leamon et al.	9,180,203 B2	11/2015	Cui et al.
7,872,235 B2	1/2011	Rouso et al.	9,180,214 B1	11/2015	Miao
7,875,586 B2	1/2011	Kovbasnjuk et al.	9,193,763 B2	11/2015	Low et al.
7,879,981 B2	2/2011	Obata	9,216,218 B2	12/2015	Sahin et al.
7,910,594 B2	3/2011	Vlahov et al.	9,226,981 B2	1/2016	Pomper et al.
7,990,533 B2	8/2011	Maier et al.	9,242,012 B2	1/2016	Ma et al.
8,000,773 B2	8/2011	Rouso et al.	9,255,262 B2	2/2016	Wong et al.
8,013,991 B2	9/2011	Maier et al.	9,278,067 B2	3/2016	Boulikas
8,088,387 B2	1/2012	Steeves et al.	9,295,727 B2	3/2016	Zale et al.
8,101,369 B2	1/2012	Nam et al.	9,309,193 B2	4/2016	Babich et al.
8,101,713 B2	1/2012	Cuello et al.	9,346,846 B1	5/2016	Herzon et al.
8,105,568 B2	1/2012	Vlahov et al.	9,371,360 B2	6/2016	Pomper et al.
8,211,402 B2	3/2012	Babich et al.	9,387,344 B2	7/2016	Sgouros et al.
8,153,595 B2	4/2012	Chen	9,422,234 B2	8/2016	Chandran et al.
8,194,660 B2	6/2012	Birze et al.	9,429,575 B2	8/2016	Ban et al.
8,211,401 B2	7/2012	Babich et al.	9,433,594 B2	9/2016	Babich et al.
8,211,473 B2	7/2012	Troiano et al.	9,447,121 B2	9/2016	Babich et al.
8,211,635 B2	7/2012	Barton	9,498,546 B2	11/2016	Pomper et al.
8,227,634 B2	7/2012	Pomper et al.	9,556,167 B2	1/2017	Spiegel et al.
8,236,330 B2	8/2012	Zale et al.	9,567,402 B2	2/2017	Liu
8,246,968 B2	8/2012	Zale et al.	9,580,467 B2	2/2017	Cheng et al.
8,258,111 B2	9/2012	Shen et al.	9,580,474 B2	2/2017	Viscidi et al.
			9,585,957 B2	3/2017	Powell et al.
			9,617,602 B2	4/2017	Joseph et al.
			9,636,413 B2	5/2017	Vlahov et al.
			9,687,572 B2	6/2017	Babich et al.

(56)

References Cited

U.S. PATENT DOCUMENTS

9,717,484 B2	8/2017	Kaloo et al.	2007/0225213 A1	9/2007	Kosak
9,745,380 B2	8/2017	Govindan et al.	2007/0244055 A1	10/2007	Brady et al.
9,757,084 B2	9/2017	Sgouros et al.	2007/0254316 A1	11/2007	Rodriguez et al.
9,764,039 B2	9/2017	Thanos et al.	2007/0254317 A1	11/2007	Busseret-Michel et al.
9,770,467 B2	9/2017	Dubensky, Jr. et al.	2008/0008649 A1	1/2008	Cappelletti et al.
9,770,517 B2	9/2017	Govindan et al.	2008/0008719 A1	1/2008	Bowditch et al.
9,808,516 B2	11/2017	Brockstedt et al.	2008/0089869 A1	4/2008	Denmeade et al.
9,814,759 B2	11/2017	Wong et al.	2008/0089892 A1	4/2008	Allan et al.
9,861,444 B2	1/2018	Kaloo et al.	2008/0114153 A1	5/2008	Steeves et al.
9,889,199 B2	2/2018	Basilion	2008/0175789 A1	7/2008	Frangioni
9,932,411 B2	4/2018	Terrell et al.	2008/0176821 A1	7/2008	Kozikowski et al.
9,951,049 B2	4/2018	Kamal et al.	2008/0193381 A1	8/2008	Babich et al.
9,951,324 B2	4/2018	Low et al.	2008/0214436 A1	9/2008	Yu et al.
9,956,305 B2	5/2018	Babich et al.	2008/0248052 A1	10/2008	Vlahov et al.
9,988,407 B2	6/2018	Slusher et al.	2008/0269105 A1	10/2008	Taft et al.
10,010,624 B2	7/2018	Howard et al.	2008/0311037 A1	12/2008	Heston et al.
10,011,632 B2	7/2018	Wang et al.	2009/0117042 A1	5/2009	Pomper et al.
10,016,519 B2	7/2018	Kopka et al.	2009/0123467 A1	5/2009	Bedi et al.
10,029,023 B2	7/2018	Pomper et al.	2009/0180951 A1	7/2009	Zimmerman et al.
10,064,957 B2	9/2018	Govindan et al.	2009/0214636 A1	8/2009	Low et al.
2003/0049203 A1	3/2003	Elmaleh et al.	2009/0247614 A1	10/2009	Manoharan et al.
2003/0220241 A1	11/2003	DeFeo-Jones et al.	2009/0258002 A1	10/2009	Barrett et al.
2003/0232760 A1	12/2003	Garsky et al.	2009/0274625 A1	11/2009	Denmeade et al.
2004/0001846 A1	1/2004	Israeli et al.	2010/0047170 A1	2/2010	Denmeade et al.
2004/0002478 A1	1/2004	Kozikowski et al.	2010/0048490 A1	2/2010	Vlahov et al.
2004/0002587 A1	1/2004	Watkins et al.	2010/0092496 A1	4/2010	Boyd et al.
2004/0018203 A1	1/2004	Pastan et al.	2010/0178246 A1	7/2010	Babich et al.
2004/0029778 A1	2/2004	Isaacs	2010/0183509 A1	7/2010	Babich et al.
2004/0033195 A1	2/2004	Leamon et al.	2010/0183517 A1	7/2010	Berkman
2004/0052727 A1	3/2004	Dalton et al.	2010/0209343 A1	8/2010	Bander et al.
2004/0054190 A1	3/2004	Pomper et al.	2010/0240701 A1	9/2010	Vlahov et al.
2004/0058857 A1	3/2004	Yao	2010/0260677 A1	10/2010	Bhatia et al.
2004/0110723 A1	6/2004	Frangioni	2010/0324008 A1	12/2010	Low et al.
2004/0146516 A1	7/2004	Roben et al.	2011/0008253 A1	1/2011	Babich et al.
2004/0213791 A1	10/2004	Bander et al.	2011/0027180 A1	2/2011	Magnani
2004/0229845 A1	11/2004	Frangioni	2011/0027274 A1	2/2011	Cheng et al.
2004/0242582 A1	12/2004	Green et al.	2011/0064657 A1	3/2011	Pomper et al.
2005/0002942 A1	1/2005	Vlahov et al.	2011/0124948 A1	5/2011	Yokell
2005/0069889 A1	3/2005	Nihei et al.	2011/0142760 A1	6/2011	Pomper et al.
2005/0107325 A1	5/2005	Manoharan et al.	2011/0172254 A1	7/2011	Leamon et al.
2005/0119166 A1	6/2005	Brady et al.	2011/0176998 A1	7/2011	Pomper et al.
2005/0158780 A1	7/2005	Lupold et al.	2011/0200677 A1	8/2011	Chandran et al.
2005/0234247 A1	10/2005	Klar et al.	2011/0288152 A1	11/2011	Low et al.
2005/0239138 A1	10/2005	Hess et al.	2011/0305768 A1	12/2011	Mao et al.
2005/0239739 A1	10/2005	Matulic-Adamic	2012/0009121 A1	1/2012	Pomper et al.
2005/0245486 A1	11/2005	Frangioni	2012/0276162 A1	11/2012	Zale et al.
2005/0255042 A1	11/2005	Lam et al.	2012/0322741 A1	12/2012	Low et al.
2006/0024317 A1	2/2006	Boyd et al.	2013/0034494 A1	2/2013	Babich et al.
2006/0045883 A1	3/2006	Moldrem et al.	2013/0302409 A1	11/2013	Fuchs et al.
2006/0051380 A1	3/2006	Schulick et al.	2013/0336888 A1	12/2013	Babich et al.
2006/0052312 A1	3/2006	Erhardt et al.	2014/0107316 A1	4/2014	Vlahov et al.
2006/0062793 A1	3/2006	Webb et al.	2014/0113322 A1	4/2014	Cui et al.
2006/0105975 A1	5/2006	Pendergrast et al.	2014/0140925 A1	5/2014	Leamon et al.
2006/0106047 A1	5/2006	Jiang et al.	2014/0154702 A1	6/2014	Parker et al.
2006/0140871 A1	6/2006	Sillerud	2014/0187501 A1	7/2014	Bilodeau et al.
2006/0148718 A1	7/2006	Brady et al.	2014/0314864 A1	10/2014	Cheng et al.
2006/0148741 A1	7/2006	Barrett et al.	2015/0023875 A1	1/2015	Farokhzad et al.
2006/0155021 A1	7/2006	Lenges et al.	2015/0079001 A1	3/2015	Pomper et al.
2006/0155146 A1	7/2006	Lenges et al.	2015/0104387 A1	4/2015	Pomper et al.
2007/0010014 A1	1/2007	Wood et al.	2015/0110715 A1	4/2015	Eder et al.
2007/0020327 A1	1/2007	Fikes et al.	2015/0110716 A1	4/2015	Armor
2007/0031326 A1	2/2007	Shirvan et al.	2015/0246144 A1	9/2015	Pomper et al.
2007/0031438 A1	2/2007	Junghans	2015/0297735 A1	10/2015	Vlahov et al.
2007/0117153 A1	5/2007	Bieniarz et al.	2015/0315196 A1	11/2015	Howard
2007/0128670 A1	6/2007	Klatzmann et al.	2015/0366968 A1	12/2015	Basilion et al.
2007/0134332 A1	6/2007	Turnell et al.	2016/0067341 A1	3/2016	Low et al.
2007/0141052 A1	6/2007	Watkins et al.	2016/0074526 A1	3/2016	Bilodeau et al.
2007/0142296 A1	6/2007	McBride et al.	2016/0114060 A1	4/2016	Pomper et al.
2007/0148662 A1	6/2007	Israeli et al.	2016/0151508 A1	6/2016	Low et al.
2007/0160617 A1	7/2007	Ma et al.	2016/0220694 A1	8/2016	Vlahov et al.
2007/0172422 A1	7/2007	Glazier	2016/0235865 A1	8/2016	Pomper et al.
2007/0179100 A1	8/2007	Manoharan	2016/0287731 A1	10/2016	Vlahov et al.
2007/0190029 A1	8/2007	Pardoll et al.	2016/0361376 A1	12/2016	Vlahov et al.
2007/0212337 A1	9/2007	Bedi et al.	2016/0361432 A1	12/2016	Vlahov et al.
2007/0219165 A1	9/2007	Berkman et al.	2016/0361433 A1	12/2016	Vlahov et al.
			2017/0081298 A1	3/2017	Ray et al.
			2017/0151356 A1	6/2017	Govindan et al.
			2017/0218464 A1	8/2017	Pomper et al.
			2017/0224709 A1	8/2017	Slusher et al.

(56)

References Cited

U.S. PATENT DOCUMENTS

2017/0266316 A1 9/2017 Govindan et al.
 2017/0275673 A1 9/2017 Luo et al.
 2017/0281789 A1 10/2017 Basilion et al.
 2017/0281791 A1 10/2017 Govindan et al.
 2017/0333576 A1 11/2017 Pomper et al.
 2018/0008668 A1 1/2018 Isaacs et al.
 2018/0051039 A1 2/2018 Pomper et al.
 2018/0064831 A1 3/2018 Basilion et al.
 2018/0111895 A1 4/2018 Babich et al.
 2018/0118847 A1 5/2018 Bander
 2018/0133296 A1 5/2018 Barrett et al.
 2018/0148480 A1 5/2018 Issaacs et al.
 2018/0185511 A1 7/2018 Woodworth et al.
 2018/0207299 A1 7/2018 Babich et al.
 2018/0258134 A1 9/2018 Wang et al.

FOREIGN PATENT DOCUMENTS

CN 101863924 A 10/2010
 DE 202014008232 3/2015
 EP 0116208 8/1984
 EP 1177200 6/2005
 EP 1472541 9/2009
 EP 2187965 5/2010
 EP 2318366 5/2011
 EP 2136788 10/2011
 EP 2373621 A1 10/2011
 EP 2373622 10/2011
 EP 2389361 11/2011
 EP 2408755 1/2012
 EP 2644192 10/2013
 EP 2644594 10/2013
 EP 2648766 A1 10/2013
 EP 2436376 7/2014
 EP 2759535 7/2014
 EP 2240171 8/2014
 EP 2170075 12/2014
 EP 2823826 1/2015
 EP 2097111 7/2015
 EP 2921482 9/2015
 EP 2938364 A1 11/2015
 EP 2942065 11/2015
 EP 2958596 A1 12/2015
 EP 2993171 A1 3/2016
 EP 2706057 4/2016
 EP 3038996 A1 7/2016
 JP 2002506204 2/2002
 JP 2004536034 12/2004
 JP 2005274569 10/2005
 JP 2006501149 1/2006
 JP 2006514961 A 5/2006
 JP 2006518712 8/2006
 JP 2007521803 8/2007
 JP 2009519209 A 5/2009
 JP 2010515732 A 5/2010
 JP 2010518112 A 5/2010
 JP 2010532754 10/2010
 JP 2012511023 5/2012
 WO 8801622 3/1988
 WO 9107418 4/1991
 WO 95/33766 12/1995
 WO 9945374 9/1999
 WO 2000/066091 A1 11/2000
 WO 2000064911 11/2000
 WO 2002043773 2/2002
 WO 2002062398 8/2002
 WO 2002098885 12/2002
 WO 2003060523 7/2003
 WO 2003092742 11/2003
 WO 2003097647 11/2003
 WO 2004010957 2/2004
 WO 2004069159 8/2004
 WO 2004069285 A1 8/2004
 WO 2005082023 9/2005
 WO 2006012527 2/2006

WO 2006093991 9/2006
 WO 2006096754 9/2006
 WO 2006136564 12/2006
 WO 2007006041 A2 1/2007
 WO 2007022494 2/2007
 WO 2007106869 9/2007
 WO 2008057437 5/2008
 WO 2008058192 5/2008
 WO 2008088648 A2 7/2008
 WO 2008098112 A2 8/2008
 WO 2008101231 A2 8/2008
 WO 2008121949 10/2008
 WO 2009002529 12/2008
 WO 2009026177 2/2009
 WO 2009082606 2/2009
 WO 2009070302 6/2009
 WO 2009089383 A2 7/2009
 WO 2009002529 12/2009
 WO 2009002993 12/2009
 WO 2010014933 2/2010
 WO 2010/065899 A2 6/2010
 WO 2010/065906 A2 6/2010
 WO 2010065899 6/2010
 WO 2010065902 6/2010
 WO 2010108125 9/2010
 WO 2011/014821 A1 2/2011
 WO 2010108125 3/2011
 WO 2011106639 9/2011
 WO 2012078534 6/2012
 WO 2012166923 12/2012
 WO 2013022797 2/2013
 WO 2013028664 2/2013
 WO 2013/130776 A1 9/2013
 WO 2014062697 4/2014
 WO 2014078484 5/2014
 WO 2014106208 7/2014
 WO 2014127365 8/2014
 WO 2014134543 9/2014
 WO 2015053318 4/2015
 WO 2015057250 4/2015
 WO 2015171792 11/2015
 WO 2016030329 3/2016
 WO 2016040179 3/2016

OTHER PUBLICATIONS

Jadvar, H., Xiankui, L., Shahinian, A., Park, R., Tohme, M., Pinski, J., and Conti, P. S. (2005) Glucose metabolism of human prostate cancer mouse xenografts. *Mol Imaging* 4, 91-7.
 Humblet, V., Lapidus, R., Williams, L. R., Tsukamoto, T., Rojas, C., Majer, P., Hin, B., Ohnishi, S., De Grand, A. M., Zaheer, A., Renze, J. T., Nakayama, A., Slusher, B. S., and Frangioni, J. V. (2005) High-affinity near-infrared fluorescent small-molecule contrast agents for in vivo imaging of prostate-specific membrane antigen. *Mol Imaging* 4, 448-62.
 Gregor, P. D., Wolchok, J. D., Turaga, V., Latouche, J. B., Sadelain, M., Bacich, D., Heston, W. D., Houghton, A. N., and Scher, H. I. (2005) Induction of autoantibodies to syngeneic prostate-specific membrane antigen by xenogeneic vaccination. *Int J Cancer* 116, 415-21.
 Foss, C. A., Mease, R. C., Fan, H., Wang, Y., Ravert, H. T., Dannals, R. F., Olszewski, R. T., Heston, W. D., Kozikowski, A. P., and Pomper, M. G. (2005) Radiolabeled small-molecule ligands for prostate-specific membrane antigen: in vivo imaging in experimental models of prostate cancer. *Clin Cancer Res* 11, 4022-8.
 Vallabhajosula, S., Smith-Jones, P. M., Navarro, V., Goldsmith, S. J., and Bander, N. H. (2004) Radioimmunotherapy of prostate cancer in human xenografts using monoclonal antibodies specific to prostate specific membrane antigen (PSMA): studies in nude mice. *Prostate* 58, 145-55.
 Scheffel, U., and Pomper, M. G. (2004) PET imaging of GRP receptor expression in prostate cancer. *J Nucl Med* 45, 1277-8.
 O'Keefe, D. S., Bacich, D. J., and Heston, W. D. (2004) Comparative analysis of prostate-specific membrane antigen (PSMA) versus a prostate-specific membrane antigen-like gene. *Prostate* 58, 200-10.

(56)

References Cited

OTHER PUBLICATIONS

- Milowsky, M. I., Nanus, D. M., Kostakoglu, L., Vallabhajosula, S., Goldsmith, S. J., and Bander, N. H. (2004) Phase I trial of yttrium-90-labeled anti-prostate-specific membrane antigen monoclonal antibody J591 for androgen-independent prostate cancer. *J Clin Oncol* 22, 2522-31.
- Matthies, A., Ezziddin, S., Ulrich, E. M., Palmedo, H., Biersack, H. J., Bender, H., and Gohlke, S. (2004) Imaging of prostate cancer metastases with 18F-fluoroacetate using PET/CT. *Eur J Nucl Med Mol Imaging* 31, 797.
- Kozikowski, A. P., Zhang, J., Nan, F., Petukhov, P. A., Grajkowska, E., Wroblewski, J. T., Yamamoto, T., Bzdega, T., Wroblewska, B., and Neale, J. H. (2004) Synthesis of urea-based inhibitors as active site probes of glutamate carboxypeptidase II: efficacy as analgesic agents. *J Med Chem* 47, 1729-38.
- Henry, M. D., Wen, S., Silva, M. D., Chandra, S., Milton, M., and Worland, P. J. (2004) A prostate-specific membrane antigen-targeted monoclonal antibody-chemotherapeutic conjugate designed for the treatment of prostate cancer. *Cancer Res* 64, 7995-8001.
- Ghosh, A., and Heston, W. D. (2004) Tumor target prostate specific membrane antigen (PSMA) and its regulation in prostate cancer. *J Cell Biochem* 91, 528-39.
- Chopra, A. (2010) 68Ga-Labeled 2-[3-[5-(7-[1-benzoyloxycarbonyl-5-[2-(4,7,10-tris-carboxymethyl-1,4,7,10-tetraazacyclododec-1-yl)acetylaminopentyl]carbamoyl]-heptanoylamino)-1-carboxypentyl]ureido]pentanedioic acid. Molecular Imaging and Contrast Agent Database (MICAD) [database online]. Bethesda (MD): National Center for Biotechnology Information (US); 2004-2013.
- Chopra, A. (2010) 68Ga-Labeled 2-[3-(1-carboxy-5-[7-[5-carboxy-5-(3-phenyl-2-[3-phenyl-2-[2-(4,7,10-tris-carboxymethyl-1,4,7,10-tetraazacyclododec-1-yl)acetylaminopropionylamino]propionylamino)pentyl]carbamoyl]heptanoylamino)pentyl]ureido]pentanedioic acid. Molecular Imaging and Contrast Agent Database (MICAD) [database online]. Bethesda (MD): National Center for Biotechnology Information (US); 2004-2013.
- Bzdega, T., Crowe, S. L., Ramadan, E. R., Sciarretta, K. H., Olszewski, R. T., Ojefo, O. A., Rafalski, V. A., Wroblewska, B., and Neale, J. H. (2004) The cloning and characterization of a second brain enzyme with NAAG peptidase activity. *Journal of neurochemistry* 89, 627-35.
- Tang, H., Brown, M., Ye, Y., Huang, G., Zhang, Y., Wang, Y., Zhai, H., Chen, X., Shen, T. Y., and Tenniswood, M. (2003) Prostate targeting ligands based on N-acetylated alpha-linked acidic dipeptidase. *Biochem Biophys Res Commun* 307, 8-14.
- Schulke, N., Varlamova, O. A., Donovan, G. P., Ma, D., Gardner, J. P., Morrissey, D. M., Arrigale, R. R., Zhan, C., Chodera, A. J., Surowitz, K. G., Maddon, P. J., Heston, W. D., and Olson, W. C. (2003) The homodimer of prostate-specific membrane antigen is a functional target for cancer therapy. *Proc Natl Acad Sci U S A* 100, 12590-5.
- Rajasekaran, S. A., Anilkumar, G., Oshima, E., Bowie, J. U., Liu, H., Heston, W., Bander, N. H., and Rajasekaran, A. K. (2003) A novel cytoplasmic tail MXXXL motif mediates the internalization of prostate-specific membrane antigen. *Mol Biol Cell* 14, 4835-45.
- Oyama, N., Miller, T. R., Dehdashti, F., Siegel, B. A., Fischer, K. C., Michalski, J. M., Kibel, A. S., Andriole, G. L., Picus, J., and Welch, M. J. (2003) 11C-acetate PET imaging of prostate cancer: detection of recurrent disease at PSA relapse. *J Nucl Med* 44, 549-55.
- Meighan, M. A., Dickerson, M. T., Glinskii, O., Glinsky, V. V., Wright, G. L., Jr., and Deutscher, S. L. (2003) Recombinant glutamate carboxypeptidase II (prostate specific membrane antigen-PSMA)—cellular localization and bioactivity analyses. *J Protein Chem* 22, 317-26.
- Hain, S. F., and Maisey, M. N. (2003) Positron emission tomography for urological tumours. *BJU Int* 92, 159-64.
- Dimitrakopoulou-Strauss, A., and Strauss, L. G. (2003) PET imaging of prostate cancer with 11C-acetate. *J Nucl Med* 44, 556-8.
- Shvarts, O., Han, K. R., Seltzer, M., Pantuck, A. J., and Belldgrun, A. S. (2002) Positron emission tomography in urologic oncology. *Cancer Control* 9, 335-42.
- Rong, S. B., Zhang, J., Neale, J. H., Wroblewski, J. T., Wang, S., and Kozikowski, A. P. (2002) Molecular modeling of the interactions of glutamate carboxypeptidase II with its potent NAAG-based inhibitors. *J Med Chem* 45, 4140-52.
- Pomper, M. G., Musachio, J. L., Zhang, J., Scheffel, U., Zhou, Y., Hilton, J., Maini, A., Dannals, R. F., Wong, D. F., and Kozikowski, A. P. (2002) 11C-MCG: synthesis, uptake selectivity, and primate PET of a probe for glutamate carboxypeptidase II (NAALADase). *Mol Imaging* 1, 96-101.
- Oyama, N., Akino, H., Kanamaru, H., Suzuki, Y., Muramoto, S., Yonekura, Y., Sadato, N., Yamamoto, K., and Okada, K. (2002) 11C-acetate PET imaging of prostate cancer. *J Nucl Med* 43, 181-6.
- Kotzerke, J., Gschwend, J. E., and Neumaier, B. (2002) PET for prostate cancer imaging: still a quandary or the ultimate solution? *J Nucl Med* 43, 200-2.
- Hara, T., Kosaka, N., and Kishi, H. (2002) Development of (18F)-fluoroethylcholine for cancer imaging with PET: synthesis, biochemistry, and prostate cancer imaging. *J Nucl Med* 43, 187-99.
- Hara, T. (2002) 11C-choline and 2-deoxy-2-[18F]fluoro-D-glucose in tumor imaging with positron emission tomography. *Mol Imaging Biol* 4, 267-73.
- Chang, S. S., and Heston, W. D. (2002) the clinical role of prostate-specific membrane antigen (PSMA). *Urol Oncol* 7, 7-12.
- DeGrado, T. R., Coleman, R. E., Wang, S., Baldwin, S. W., Orr, M. D., Robertson, C. N., Polascik, T. J., and Price, D. T. (2001) Synthesis and evaluation of 18F-labeled choline as an oncologic tracer for positron emission tomography: initial findings in prostate cancer. *Cancer Res* 61, 110-7.
- DeGrado, T. R., Baldwin, S. W., Wang, S., Orr, M. D., Liao, R. P., Friedman, H. S., Reiman, R., Price, D. T., and Coleman, R. E. (2001) Synthesis and evaluation of (18F)F-labeled choline analogs as oncologic PET tracers. *J Nucl Med* 42, 1805-14.
- Bacich, D. J., Pinto, J. T., Tong, W. P., and Heston, W. D. W. (2001) Cloning, expression, genomic localization, and enzymatic activities of the mouse homolog of prostate-specific membrane antigen/NAALADase/folate hydrolase. *Mammalian Genome* 12, 117-123.
- Slusher, B. S., Vornov, J. J., Thomas, A. G., Hum, P. D., Harukuni, I., Bhardwaj, A., Traystman, R. J., Robinson, M. B., Britton, P., Lu, X. C., Tortella, F. C., Wozniak, K. M., Yudkoff, M., Potter, B. M., and Jackson, P. F. (1999) Selective inhibition of NAALADase, which converts NAAG to glutamate, reduces ischemic brain injury. *Nat Med* 5, 1396-402.
- Dumas, F., Gala, J. L., Berteau, P., Brasseur, F., Eschwege, P., Paradis, V., Lacour, B., Philippe, M., and Loric, S. (1999) Molecular expression of PSMA mRNA and protein in primary renal tumors. *Int J Cancer* 80, 799-803.
- Chang, S. S., Reuter, V. E., Heston, W. D., Bander, N. H., Grauer, L. S., and Gaudin, P. B. (1999) Five different anti-prostate-specific membrane antigen (PSMA) antibodies confirm PSMA expression in tumor-associated neovasculature. *Cancer Res* 59, 3192-8.
- Sweat, S. D., Pacelli, A., Murphy, G. P., and Bostwick, D. G. (1998) Prostate-specific membrane antigen expression is greatest in prostate adenocarcinoma and lymph node metastases. *Urology* 52, 637-40.
- Murphy, G. P., Elgamal, A. A., Su, S. L., Bostwick, D. G., and Holmes, E. H. (1998) Current evaluation of the tissue localization and diagnostic utility of prostate specific membrane antigen. *Cancer* 83, 2259-69.
- Liu, H., Rajasekaran, A. K., Moy, P., Xia, Y., Kim, S., Navarro, V., Rahmati, R., and Bander, N. H. (1998) Constitutive and antibody-induced internalization of prostate-specific membrane antigen. *Cancer Res* 58, 4055-60.
- Kahn, D., Williams, R. D., Manyak, M. J., Haseman, M. K., Seldin, D. W., Libertino, J. A., and Maguire, R. T. (1998) 111Indium-capromab pendetide in the evaluation of patients with residual or recurrent prostate cancer after radical prostatectomy. The ProstaScint Study Group. *J Urol* 159, 2041-6; discussion 2046-7.
- Hara, T., Kosaka, N., and Kishi, H. (1998) PET imaging of prostate cancer using carbon-11-choline. *J Nucl Med* 39, 990-5.

(56)

References Cited

OTHER PUBLICATIONS

- Bostwick, D. G., Pacelli, A., Blute, M., Roche, P., and Murphy, G. P. (1998) Prostate specific membrane antigen expression in prostatic intraepithelial neoplasia and adenocarcinoma: a study of 184 cases. *Cancer* 82, 2256-61.
- Silver, D. A., Pellicer, I., Fair, W. R., Heston, W. D., and Cordon-Cardo, C. (1997) Prostate-specific membrane antigen expression in normal and malignant human tissues. *Clin Cancer Res* 3, 81-5.
- Fair, W. R., Israeli, R. S., and Heston, W. D. (1997) Prostate-specific membrane antigen. *Prostate* 32, 140-8.
- Jackson P.F., Cole D.C., Slusher B.S., Stetz S.L., Ross L.E., Donzanti B.A., and D.A., T. (1996) Design, synthesis, and biological activity of a potent inhibitor of the neuropeptidase N-acetylated alpha-linked acidic dipeptidase.
- Wright, G. L., Jr., Haley, C., Beckett, M. L., and Schellhammer, P. F. (1995) Expression of prostate-specific membrane antigen in normal, benign, and malignant prostate tissues. *Urol Oncol* 1, 18-28.
- Leek, J., Lench, N., Maraj, B., Bailey, A., Carr, I. M., Andersen, S., Cross, J., Whelan, P., MacLennan, K. A., Meredith, D. M., and et al. (1995) Prostate-specific membrane antigen: evidence for the existence of a second related human gene. *Br J Cancer* 72, 583-8.
- Vees, H., et al., "18F-choline and/or 11 C-acetate positron emission tomography: detection of residual or progressive subclinical disease at very low prostate-specific antigen values (<1 ng/mL) after radical prostatectomy," *BJU Int*, 2007, 99, 1415-1420.
- Larson, S. M., et al., "Tumor Localization of 16 β -18F-Fluoro-5 α -Dihydrotestosterone Versus 18F-FDG in Patients with Progressive, Metastatic Prostate Cancer," *J Nucl Med*, 2004, 45, 366-373.
- Schuster, D.M., et al., "Initial Experience with the Radiotracer Anti-1-Am ino-3-18F-Fluorocyclobutane-1-Carboxylic Acid with PET/CT in Prostate Carcinoma," *J. Nucl. Med.*, 2007, 48, 56-63.
- Tehrani, O.S., et al., "Tumor Imaging Using 1-(2'-deoxy-2'-18F-Fluoro- β -D-Arabinofuranosyl) Thymine and PET," *J. Nucl. Med.*, 2007, 48, 1436-1441.
- Mease RC., et al., "N-[N-[(S)-1, 3-Dicarboxypropyl]Carbam oyl]-4-[18F]F luorobenzyl-L-Cysteine, [18F]DCFBC: A New Imaging Probe for Prostate Cancer," *Clin. Cancer Res.*, 2008, 14, 3036-3043.
- Zhou, J., et al., "NAAG Peptidase Inhibitors and Their Potential for Diagnosis and Therapy," *Nat Rev Drug Discovery*, 2005, 4, 1015-1026.
- Schulke, N., et al., "The homodimer of prostate-specific membrane antigen is a functional target for cancer therapy," *Proc. Natl. Acad. Sci., USA*, 2003, 100, 12590-12595.
- Nan, F., et al., "Dual Function Glutamate-Related Ligands: Discovery of a Novel, Potent Inhibitor of Glutamate Carboxypeptidase II Possessing mGluR3 Agonist Activity," *J. Med. Chem.*, 2000, 43, 772-774.
- Lange, P.H., "ProstaScint scan for staging prostate cancer," *Urology*, 2001, 57, 402-406.
- Haseman, M.K., et al., "Capromab Pendetide Imaging of Prostate Cancer," *Cancer Bi other Radiopharm*, 2000, 15, 131-140.
- Mier W., et al., "Conjugation of DOTA Using Isolated Phenolic Active Esters: The Labeling and Biodistribution of Albumin as Blood Pool Marker," *Bioconjugate Chem.*, 2005, 16: 237-240.
- Schafer et al., "A dimerized urea-based inhibitor of the prostate-specific membrane antigen for 68Ga-PET imaging of prostate cancer," *EJNMMI Research*, 2012, 2, 23, 11 pages.
- Humblet, V. et al., "An HPLC/mass spectrometry platform for the development of multimodality contrast agents and targeted therapeutics: prostate-specific membrane antigen small molecule derivatives," *Contrast Med. Mol. Imaging*, 2006, 1, 196-211.
- Pomper, M.G., et al., "11C-MCG: synthesis, uptake selectivity, and primate PET of a probe for glutamate carboxypeptidase II (NAALADase)," *Mal Imaging*, 2002, 1, 96-101.
- Japanese Office Action dated Mar. 23, 2017, which issued during prosecution of Japanese Application No. P016-524427.
- Humblet, et al., "An HPLC/mass spectrometry platform for the development of multimodality contrast agents and targeted therapeutics: prostate-specific membrane antigen small molecule derivatives" *Contrast Media & Molecular Imaging*, Jul. 2006 1:196-211.
- Eurasian Office Action dated Feb. 10, 2017, which issued during prosecution of Eurasian Application No. 201690495/28.
- Office Action for copending Eurasian Patent Application No. 201690495/28, dated Jul. 14, 2017.
- Scher, B. et al., "Value of 11C-choline PET and PET/CT in patients with suspected prostate cancer," *Eur. J. Nucl. Med. Mol. Imaging.*, 2007, 34, 45-53.
- Tasch, J. et al., "A Unique Folate Hydrolase, Prostage-Specific Membrane Antigen (PSMA): A Target for Immunotherapy?" *Crit. Rev. Immunol.*, 2001, 21, 249-261.
- Rosenthal, S.A. et al., "Utility of Capromab Pendetide (ProstaScint) Imaging in the Management of Prostate Cancer," *Tech Urol*, 2001, 7, 27-37.
- Wiberg et al. "A comparison of some properties of C=O and C=S bonds," *ARKIVOC*, 2011, (v) 45-56.
- Zechmann, C. M., Afshar-Oromieh, A., Armor, T., Stubbs, J. B., Mier, W., Hadaschik, B., Joyal, J., Kopka, K., Debus, J., Babich, J. W., and Haberkorn, U. (2014) Radiation dosimetry and first therapy results with a I/ I-labeled small molecule (MIP-1095) targeting PSMA for prostate cancer therapy. *Eur J Nucl Med Mol Imaging* 41, 1280-92.
- Huang, S. S., Wang, X., Zhang, Y., Doke, A., DiFilippo, F. P., and Heston, W. D. (2014) Improving the biodistribution of PSMA-targeting tracers with a highly negatively charged linker. *Prostate* 74, 702-13.
- Wiehr, S., Buhler, P., Gierschner, D., Wolf, P., Rolle, A. M., Kesenheimer, C., Pichler, B. J., and Elsasser-Beile, U. (2014) Pharmacokinetics and PET imaging properties of two recombinant anti-PSMA antibody fragments in comparison to their parental antibody. *Prostate* 74, 743-55.
- Weinisen, M., Simecek, J., Schottelius, M., Schwaiger, M., and Wester, H. J. (2014) Synthesis and preclinical evaluation of DOTAGA-conjugated PSMA ligands for functional imaging and endoradiotherapy of prostate cancer. *EJNMMI Res* 4, 63.
- Wang, X., Huang, S. S., Heston, W. D., Guo, H., Wang, B. C., and Basilion, J. P. (2014) Development of targeted near-infrared imaging agents for prostate cancer. *Mol Cancer Ther* 13(11), 2595-606.
- Uprimny, C., Kroiss, A., Nilica, B., Buxbaum, S., Decristoforo, C., Horninger, W., and Virgolini, I. J. (2014) Ga-PSMA ligand PET versus F-NaF PET: evaluation of response to Ra therapy in a prostate cancer patient. *Eur J Nucl Med Mol Imaging* 42, 362-63.
- Hillier, S. M., Maresca, K. P., Lu, G., Merkin, R. D., Marquis, J. C., Zimmerman, C. N., Eckelman, W. C., Joyal, J. L., and Babich, J. W. (2013) 99mTc-labeled small-molecule inhibitors of prostate-specific membrane antigen for molecular imaging of prostate cancer. *J Nucl Med* 54, 1369-76.
- Jadvar, H. (2013) Imaging evaluation of prostate cancer with (18)F-fluorodeoxyglucose PET/CT: utility and limitations. *Eur J Nucl Med Mol Imaging* 40 (Suppl 1), S5-S10.
- Jemaa, A. B., Bouraoui, Y., Sallami, S., Banasr, A., Nouria, Y., Horchani, A., and Oueslati, R. (2013) Cellular distribution and heterogeneity of Psa and Psma expression in normal, hyperplasia and human prostate cancer. *La tunisie Medicate* 91, 458-463.
- Kasperzyk, J. L., Finn, S. P., Flavin, R., Fiorentino, M., Lis, R., Hendrickson, W. K., Clinton, S. K., Sesso, H. D., Giovannucci, E. L., Stampfer, M. J., Loda, M., and Mucci, L. A. (2013) Prostate-specific membrane antigen protein expression in tumor tissue and risk of lethal prostate cancer. *Cancer Epidemiol Biomarkers Prev* 22, 2354-63.
- Kasten, B. B., Liu, T., Nedrow-Byers, J. R., Benny, P. D., and Berkman, C. E. (2013) Targeting prostate cancer cells with PSMA inhibitor-guided gold nanoparticles. *Bioorg Med Chem Lett* 23, 565-8.
- Tykvar, J., Schimer, J., Barinkova, J., Pahl, P., Postova-Slavetinska, L., Majer, P., Konvalinka, J., and Sacha, P. (2014) Rational design of urea-based glutamate carboxypeptidase II (GCPII) inhibitors as versatile tools for specific drug targeting and delivery. *Bioorg Med Chem* 22, 4099-108.
- Ristau, B. T., O'Keefe, D. S., and Bacich, D. J. (2014) The prostate-specific membrane antigen: Lessons and current clinical implications from 20 years of research. *Urol Oncol* 32, 272-279.
- Ray Banerjee, S., Pullambhatla, M., Foss, C. A., Nimmagadda, S., Ferdani, R., Anderson, C. J., Mease, R. C., and Pomper, M. G.

(56)

References Cited

OTHER PUBLICATIONS

(2014) ⁶⁴Cu-Labeled Inhibitors of Prostate-Specific Membrane Antigen for PET Imaging of Prostate Cancer. *J Med Chem* 57, 2657-69.

Rais, R., Rojas, C., Wozniak, K., Wu, Y., Zhao, M., Tsukamoto, T., Rudek, M. A., and Slusher, B. S. (2014) Bioanalytical method for evaluating the pharmacokinetics of the GCP-II inhibitor 2-phosphonomethyl pentanedioic acid (2-PMPA). *Journal of pharmaceutical and biomedical analysis* 88, 162-9.

Pavlicek, J., Ptacek, J., Cerny, J., Byun, Y., Skultetyova, L., Pomper, M. G., Lubkowski, J., and Barinka, C. (2014) Structural characterization of P1'-diversified urea-based inhibitors of glutamate carboxypeptidase II. *Bioorg Med Chem Lett* 24, 2340-5.

Lutje, S., Rijpkema, M., Franssen, G. M., Fracasso, G., Helfrich, W., Eek, A., Oyen, W. J., Colombatti, M., and Boerman, O. C. (2014) Dual-Modality Image-Guided Surgery of Prostate Cancer with a Radiolabeled Fluorescent Anti-PSMA Monoclonal Antibody. *J Nucl Med* 55, 995-1001.

Liu, T., Mendes, D. E., and Berkman, C. E. (2014) Functional prostate-specific membrane antigen is enriched in exosomes from prostate cancer cells. *Int J Oncol* 44, 918-22.

Krohn, T., Verburg, F. A., Pufe, T., Neuhofer, W., Vogg, A., Heinzel, A., Mottaghy, F. M., and Behrendt, F. F. (2014) [Ga]PSMA-HBED uptake mimicking lymph node metastasis in coeliac ganglia: an important pitfall in clinical practice. *Eur J Nucl Med Mol Imaging* 42, 210-14.

Kovar, J. L., Cheung, L. L., Simpson, M. A., and Olive, D. M. (2014) Pharmacokinetic and Biodistribution Assessment of a Near Infrared-Labeled PSMA-Specific Small Molecule in Tumor-Bearing Mice. *Prostate cancer* 2014, 104248.

Huang, B., Otis, J., Joice, M., Kotlyar, A., and Thomas, T. P. (2014) PSMA-Targeted Stably Linked "Dendrimer-Glutamate Urea-Methotrexate" as a Prostate Cancer Therapeutic. *Biomacromolecules* 15, 915-23.

Haberkorn, U., Mier, W., Dimitrakopoulou-Strauss, A., Eder, M., Kopka, K., and Altmann, A. (2014) Mechanistic and high-throughput approaches for the design of molecular imaging probes and targeted therapeutics. *Clinical and Translational Imaging* 2, 33-41.

El-Zaria, M. E., Genady, A. R., Janzen, N., Petlura, C. I., Beckford Vera, D. R., and Valliant, J. F. (2014) Preparation and evaluation of carborane-derived inhibitors of prostate specific membrane antigen (PSMA). *Dalton Trans* 43, 4950-61.

Eiber, M., Nekolla, S. G., Maurer, T., Weirich, G., Wester, H. J., and Schwaiger, M. (2014) Ga-PSMA PET/MR with multimodality image analysis for primary prostate cancer. *Abdom Imaging* 40, 1769-71.

Eder, M., Schäfer, M., Bauder-Wüst, U., Haberkorn, U., Eisenhut, M., and Kopka, K. (2014) Preclinical evaluation of a bispecific low-molecular heterodimer targeting both PSMA and GRPR for improved PET imaging and therapy of prostate cancer. *The Prostate* 74, 659-68.

Eder, M., Neels, O., Muller, M., Bauder-Wüst, U., Remde, Y., Schäfer, M., Hennrich, U., Eisenhut, M., Afshar-Oromieh, A., Haberkorn, U., and Kopka, K. (2014) Novel Preclinical and Radiopharmaceutical Aspects of [⁶⁸Ga]GA-PSMA-HBED-CC: A New PET Tracer for Imaging of Prostate Cancer. *Pharmaceuticals* 7, 779-96.

Afshar-Oromieh, A., Zechmann, C. M., Macher, A., Eder, M., Eisenhut, M., Linhart, H. G., Holland-Letz, T., Hadaschik, B. A., Giesel, F. L., Debus, J., and Haberkorn, U. (2014) Comparison of PET imaging with a Ga-labelled PSMA ligand and F-choline-based PET/CT for the diagnosis of recurrent prostate cancer. *Eur J Nucl Med Mol Imaging* 41, 11-20.

Afshar-Oromieh, A., Haberkorn, U., Schlemmer, H. P., Fenchel, M., Eder, M., Eisenhut, M., Hadaschik, B. A., Kopp-Schneider, A., and Rothke, M. (2014) Comparison of PET/CT and PET/MRI hybrid systems using a ⁶⁸Ga-labelled PSMA ligand for the diagnosis of recurrent prostate cancer: initial experience. *Eur J Nucl Med Mol Imaging* 41, 887-97.

Afshar-Oromieh, A., Avtzi, E., Giesel, F. L., Holland-Letz, T., Linhart, H. G., Eder, M., Eisenhut, M., Boxler, S., Hadaschik, B. A., Kratochwil, C., Weichert, W., Kopka, K., Debus, J., and Haberkorn, U. (2014) The diagnostic value of PET/CT imaging with the Ga-labelled PSMA ligand HBED-CC in the diagnosis of recurrent prostate cancer. *Eur J Nucl Med Mol Imaging* 42, 197-209.

Whitaker, H. C., Shiong, L. L., Kay, J. D., Gronberg, H., Warren, A. Y., Seipel, A., Wiklund, F., Thomas, B., Wiklund, P., Miller, J. L., Menon, S., Ramos-Montoya, A., Vowler, S. L., Massie, C., Egevad, L., and Neal, D. E. (2013) N-acetyl-L-aspartyl-L-glutamate peptidase-like 2 is overexpressed in cancer and promotes a pro-migratory and pro-metastatic phenotype. *Oncogene* 33, 5274-87.

Simone, C. B., 2nd, and Hahn, S. M. (2013) What's in a Label? Radioimmunotherapy for metastatic prostate cancer. *Clin Cancer Res* 19, 4908-10.

Rybalov, M., Breeuwsma, A. J., Leliveld, A. M., Pruim, J., Dierckx, R. A., and de Jong, I. J. (2013) Impact of total PSA, PSA doubling time and PSA velocity on detection rates of (11)C-Choline positron emission tomography in recurrent prostate cancer. *World J Urol* 31, 319-23.

Rothke, M. C., Afshar-Oromieh, A., and Schlemmer, H. P. (2013) [Potential of PET/MRI for diagnosis of prostate cancer]. *Radiologe* 53, 676-81.

Roethke, M. C., Kuru, T. H., Afshar-Oromieh, A., Schlemmer, H. P., Hadaschik, B. A., and Fenchel, M. (2013) Hybrid positron emission tomography-magnetic resonance imaging with gallium 68 prostate-specific membrane antigen tracer: a next step for imaging of recurrent prostate cancer-preliminary results. *Eur Urol* 64, 862-4.

Kim, D. Kim, S. K., Valencia, C. A., and Liu, R. (2013) Tribody: robust self-assembled trimeric targeting ligands with high stability and significantly improved target-binding strength. *Biochemistry* 52, 7283-94.

Ray Banerjee, S., Pullambhatla, M., Foss, C. A., Falk, A., Byun, Y., Nimmagadda, S., Mease, R. C., and Pomper, M. G. (2013) Effect of Chelators on the Pharmacokinetics of Tc-Labeled Imaging Agents for the Prostate-Specific Membrane Antigen (PSMA). *J Med Chem* 56, 6108-21.

Parker, S. A., Diaz, I. L., Anderson, K. A., and Batt, C. A. (2013) Design, production, and characterization of a single-chain variable fragment (ScFv) derived from the prostate specific membrane antigen (PSMA) monoclonal antibody U591. *Protein Expr Purif* 89, 136-145.

Osborne, J. R., Green, D. A., Spratt, D. E., Lyashchenko, S., Fareedy, S. B., Robinson, B. D., Beattie, B. J., Jain, M., Lewis J. S., Christos, P., Larson, S. M., Bander, N. H., and Scherr, D. S. (2014) A Prospective Pilot Study of Zr-J591/ Prostate Specific Membrane Antigen Positron Emission Tomography in Men with Localized Prostate Cancer Undergoing Radical Prostatectomy. *J Urol* 191, 1439-46.

Osborne, J. R., Akhtar, N. H., Vallabhajosula, S., Anand, A., Deh, K., and Tagawa, S. T. (2013) Prostate-specific membrane antigen-based imaging. *Urol Oncol* 31, 144-54.

Nedrow-Byers, J. R., Moore, A. L., Ganguly, T., Hopkins, M. R., Fulton, M. D., Benny, P. D., and Berkman, C. E. (2013) PSMA-targeted SPECT agents: mode of binding effect on in vitro performance. *Prostate* 73, 355-62.

Mease RC, Foss CA, and Mg, P. (2013) PET imaging in prostate cancer: focus on prostate-specific membrane antigen. *Curr Top Med Chem* 13, 951-962.

Lu, G., Maresca, K. P., Hillier, S. M., Zimmerman, C. N., Eckelman, W. C., Joyal, J. L., and Babich, J. W. (2013) Synthesis and SAR of (9)(9)mTc/Re-labeled small molecule prostate specific membrane antigen inhibitors with novel Solar chelates. *Bioorg Med Chem Lett* 23, 1557-63.

Lesche, R., Kettischau, G., Gromov, A. V., Bohnke, N., Borkowski, S., Monning, U., Hegele-Hartung, C., Dohr, O., Dinkelborg, L. M., and Graham, K. (2014) Preclinical evaluation of Bay 1075553, a novel F-labelled inhibitor of prostate-specific membrane antigen for PET imaging of prostate cancer. *Eur J Nucl Med Mol Imaging* 41, 89-101.

First Office Action in counterpart Canadian Application No. 2924360, dated Jan. 11, 2018 (4 pages).

(56)

References Cited

OTHER PUBLICATIONS

Second Office Action in counterpart Chinese Application No. 201480056250.5, dated Aug. 15, 2017 (4 pages).

Second Office Action in counterpart Eurasian Application No. 201690495/28, dated Feb. 10, 2017 (5 pages).

First Office Action in counterpart Korean Application No. 1020167012314, dated Mar. 6, 2017 (3 pages).

Second Office Action in counterpart Korean Application No. 1020167012314, dated Oct. 14, 2017 (3 pages).

Second Office Action for counterpart Georgian Application No. 14132/01, dated Mar. 2018 (2 pages).

First Office Action for counterpart Georgian Application No. 14132/01, dated Feb. 2017 (2 pages).

First Examination Report for counterpart Chilean Application No. 00883-2016, dated Apr. 12, 2018.

Third Office Action for counterpart Korean Application No. 10-2016-7012314, dated May 2, 2018 (2 pages).

Silvola J. M.U. et al., "A118F-NOTA-Folate Accumulates in Atherosclerotic Plaques and Can be Detected by PET/CT", Poster presented Nov. 7, 2015 in Orlando, FL at the 2015 American Heart Association ReSuscitation Science Symposium (<http://newsroom.heart.org/events/scientific-sessions-2015-newsroom-2942760>). (3 pages).

Silvola J. M.U. et al., "A118F-NOTA-Folate Accumulates in Atherosclerotic Plaques and Can be Detected by PET/CT", Published reference of poster, Nov. 10, 2015, at http://circ.ahajournals.org/content/132/Suppl_3/A18873?cited-by=yes&legid=circulationaha;132/Suppl_3/A18873; Circulation, 2015; 132:A18873. (2 pages).

Chinese Patent Application No. 201480071256.X, by Endocyte, Inc. et al.: Office Action dated Apr. 20, 2017; English translation (9 pages).

Eurasian Patent Application No. 201690862/28, by Endocyte, Inc. et al.: Office Action, dated May 22, 2017; English translation (2 pages).

European Patent Application No. EP 14861854, by Endocyte, Inc. et al: Partial Supplementary Search Report with Opinion; dated May 19, 2017 (15 pages).

Slusher, B. S., Tsai, G., Yoo, G., and Coyle, J. T. (1992) Immunocytochemical localization of the N-acetyl-aspartyl-glutamate (NAAG) hydrolyzing enzyme N-acetylated alpha-linked acidic dipeptidase (NAALADase). *J Comp Neurol* 315, 217-29.

Hwang D. R. Mathias, C. J., Welch, M. J., McGuire, A. H., and Kadmon, D. (1990) Imaging prostate derived tumors with PET and N-(3-[18F]fluoropropyl)putrescine. *Int J Rad Appl Instrum B* 17, 525-32.

Hwang, D. R., Lang, L. X., Mathias, C. J., Kadmon, D., and Welch, M. J. (1989) N-3-[18F]fluoropropylputrescine as potential PET imaging agent for prostate and prostate derived tumors. *J Nucl Med* 30, 1205-10.

Office Action for counterpart Chinese Application No. 201610184873.1, dated Jul. 24, 2018.

Shelley James, "Urea Based Rhenium Tricarbonyl Dipeptide Compounds as Potential Radiopharmaceuticals for PSMA Imaging"; Poster: INOR 258; The 229th ACS National Meeting, San Diego, CA, Mar. 13-17, 2005.

PCT International Search Report and Written Opinion for PCT/US2011/026238, dated Apr. 27, 2011.

PCT International Search Report and Written Opinion for PCT/US2013/070007, dated Mar. 5, 2014.

PCT International Search Report for PCT/US2008/073375 dated Oct. 26, 2008.

PCT International Search Report for PCT/US2016/012653 dated Mar. 11, 2016.

PCT International Search Report/Written Opinion for PCT/US2009/061067, completed May 28, 2010.

PCT International Search Report/Written Opinion for PCT/US2009/061049, completed Mar. 15, 2010.

PCT Search Report & Written Opinion issued in App. No. PCT/US2014/065467.

Silverman, "The Organic Chemistry of Drug Design and Drug Action," Elsevier Academic Press (2nd Ed. 2003).

Beheshti, M., Langsteger, W., and Fogelman, I. (2009) Prostate cancer: role of SPECT and PET in imaging bone metastases. *Semin Nucl Med* 39, 396-407.

Ananias, H. J., van den Heuvel, M. C., Helfrich, W., and de Jong, I. J. (2009) Expression of the gastrin-releasing peptide receptor, the prostate stem cell antigen and the prostate-specific membrane antigen in lymph node and bone metastases of prostate cancer. *Prostate* 69, 1101-8.

Vavere, A. L., Kridel, S. J., Wheeler, F. B., and Lewis, J. S. (2008) 1-11C-acetate as a PET radiopharmaceutical for imaging fatty acid synthase expression in prostate cancer. *J Nucl Med* 49, 327-34.

Tang, Q. L., and Yao, M. Y. (2008) [Updated application of prostate-specific membrane antigen to the diagnosis and treatment of prostate cancer]. *Zhonghua Nan Ke Xue* 14, 79-82.

Soloviev, D., Fini, A., Chierichetti, F., Al-Nahhas, A., and Rubello, D. (2008) PET imaging with 11C-acetate in prostate cancer: a biochemical, radiochemical and clinical perspective. *Eur J Nucl Med Mol Imaging* 35, 942-9.

Scher, B., and Seitz, M. (2008) PET/CT imaging of recurrent prostate cancer. *Eur J Nucl Med Mol Imaging* 35, 5-8.

Rinnab, L., Mottaghy, F. M., Simon, J., Volkmer, B. G., de Petriconi, R., Hautmann, R. E., Wittbrodt, M., Eggert, G., Moeller, P., Blumstein, N., Reske, S., and Kuefer, R. (2008) [11C]Choline PET/CT for targeted salvage lymph node dissection in patients with biochemical recurrence after primary curative therapy for prostate cancer. Preliminary results of a prospective study. *Urol Int* 81, 191-7.

Reske, S. N., Blumstein, N. M., and Glatting, G. (2008) [11C]choline PET/CT imaging in occult local relapse of prostate cancer after radical prostatectomy. *Eur J Nucl Med Mol Imaging* 35, 9-17.

Reske S.N. (2008) [11C]choline uptake with PET/CT for the initial diagnosis of prostate cancer: relation to PSA levels, tumour stage and anti-androgenic therapy. *Eur J Nucl Med Mol Imaging* 35, 1740-1.

Mease, R. C., Dusich, C. L., Foss, C. A., Ravert, H. T., Dannals, R. F., Seidel, J., Prideaux, A., Fox, J. J., Sgouros, G., Kozikowski, A. P., and Pomper, M. G. (2008) N-[N- [(S)-1,3-Dicarboxypropyl]carbonyl]-4-[18F]fluorobenzyl-L-cysteine, [18F]DCFBC: a new imaging probe for prostate cancer. *Clin Cancer Res* 14, 3036-43.

Luboldt, W., Kufer, R., Blumstein, N., Toussaint, T. L., Kluge, A., Seemann, M. D., and Luboldt, H. J. (2008) Prostate carcinoma: diffusion-weighted imaging as potential alternative to conventional MR and 11C-choline PET/CT for detection of bone metastases. *Radiology* 249, 1017-25.

Liu T. Toriyabe, Y., Kazak, M., and Berkman, C. E. (2008) Pseudoirreversible inhibition of prostate-specific membrane antigen by phosphoramidate peptidomimetics. *Biochemistry* 47, 12658-60.

Li, X., Liu, Q., Wang, M., Jin, X., Yao, S., Liu, S., and Li, J. (2008) C-11 choline PET/CT imaging for differentiating malignant from benign prostate lesions. *Clin Nucl Med* 33, 671-6.

Igerc, I., Kohlfurst, S., Gallowitsch, H. J., Matschnig, S., Kresnik, E., Gomez-Segovia, I., and Lind, P. (2008) The value of 18F-choline PET/CT in patients with elevated PSA-level and negative prostate needle biopsy for localisation of prostate cancer. *Eur J Nucl Med Mol Imaging* 35, 976-83.

Husarik, D. B., Miralbell, R., Dubs, M., John, H., Giger, O. T., Gelet, A., Cserevenyak, T., and Hany, T. F. (2008) Evaluation of [(18F)]-choline PET/CT for staging and restaging of prostate cancer. *Eur J Nucl Med Mol Imaging* 35, 253-63.

Hospers, G. A., Helmond, F. A., de Vries, E. G., Dierckx, R. A., and de Vries, E. F. (2008) PET imaging of steroid receptor expression in breast and prostate cancer *Curr Pharm Des* 14, 3020-32.

Colabufo, N. A., Abate, C., Contino, M., Inglese, C., Niso, M., Berardi, F., and Perrone, R. (2008) PB183, a sigma receptor ligand, as a potential PET probe for the imaging of prostate adenocarcinoma. *Bioorg Med Chem Lett* 18, 1990-3.

Chen, Y., Foss, C. A., Byun, Y., Nimmagadda, S., Pullambhatla, M., Fox, J. J., Castanares, M., Lupold, S. E., Babich, J. W., Mease, R. C., and Pomper, M. G. (2008) Radiohalogenated prostate-specific membrane antigen (PSMA)-based areas as imaging agents for prostate cancer. *J Med Chem* 51, 7933-43.

(56)

References Cited

OTHER PUBLICATIONS

- Chandran, S. S., Banerjee, S. R., Mease, R. C., Pomper, M. G., and Denmeade, S. R. (2008) Characterization of a targeted nanoparticle functionalized with a urea-based inhibitor of prostate-specific membrane antigen (PSMA). *Cancer Biol Ther* 7, 974-82.
- Barinka, C., Byun, Y., Dusich, C. L., Banerjee, S. R., Chen, Y., Castanares, M., Kozikowski, A. P., Mease, R. C., Pomper, M. G., and Lubkowski, J. (2008) Interactions between human glutamate carboxypeptidase II and urea-based inhibitors: structural characterization. *J Med Chem* 51, 7737-43.
- Banerjee, S. R., Foss, C. A., Castanares, M., Mease, R. C., Byun, Y., Fox, J. J., Hilton, J., Lupold, S. E., Kozikowski, A. P., and Pomper, M. G. (2008) Synthesis and evaluation of technetium-99m- and rhenium-labeled inhibitors of the prostate-specific membrane antigen (PSMA). *J Med Chem* 51, 4504-17.
- Wu, L. Y., Anderson, M. O., Toriyabe, Y., Maung, J., Campbell, T. Y., Tajon, C., Kazak, M., Moser, J., and Berkman, C. E. (2007) The molecular pruning of a phosphoramidate peptidomimetic inhibitor of prostate-specific membrane antigen. *Bioorg Med Chem* 15, 7434-43.
- Vees, H., Buchegger, F., Albrecht, S., Khan, H., Husarik, D., Zaidi, H., Soloviev, D., Hany, T. F., and Miralbell, R. (2007) 18F-choline and/or 11C-acetate positron emission tomography: detection of residual or progressive subclinical disease at very low prostate-specific antigen values (<1 ng/mL) after radical prostatectomy. *BJU Int* 99, 1415-20.
- Testa, C., Schiavina, R., Lodi, R., Salizzoni, E., Corti, B., Farsad, M., Kurhanewicz, J., Manferrari, F., Brunocilla, E., Tonon, C., Monetti, N., Castellucci, P., Fanti, S., Coe, M., Grigioni, E. F., Martorana, G., Canini, R., and Barbiroli, B. (2007) Prostate cancer: sextant localization with MR imaging, MR spectroscopy, and 11C-choline PET/CT. *Radiology* 244, 797-806.
- Scattoni, V., Picchio, M., Suardi, N., Messa, C., Freschi, M., Roscigno, M., Da Pozzo, L., Bocciardi, A., Rigatti, P., and Fazio, F. (2007) Detection of lymph-node metastases with integrated [11C]choline PET/CT in patients with PSA failure after radical retropubic prostatectomy: results confirmed by open pelvic-retroperitoneal lymphadenectomy. *Eur Urol* 52, 423-9.
- Sacha, P., Zamecnik, J., Barinka, C., Hlouchova, K., Vicha, A., Mlcochova, P., Hilgert, I., Eckschlager, T., and Konvalinka, J. (2007) Expression of glutamate carboxypeptidase II in human brain. *Neuroscience* 144, 1361-72.
- Reske, S. N. (2007) [Nuclear imaging of prostate cancer: current status]. *Der Urologe* 46, 1485-99.
- Ponde, D. E., Dence, C. S., Oyama, N., Kim, J., Tai, Y. C., Laforest, R., Siegel, B. A., and Welch, M. J. (2007) 18F-fluoroacetate: a potential acetate analog for prostate tumor imaging—in vivo evaluation of 18F-fluoroacetate versus 11C-acetate. *J Nucl Med* 48, 420-8.
- Pemer, S., Hofer, M. D., Kim, R., Shah, R. B., Li, H., Moller, P., Hautmann, R. E., Gschwend, J. E., Kuefer, R., and Rubin, M. A. (2007) Prostate-specific membrane antigen expression as a predictor of prostate cancer progression. *Hum Pathol* 38, 696-701.
- Oehr, P., and Bouchelouche, K. (2007) Imaging of prostate cancer. *Curr Opin Oncol* 19, 259-64.
- Morris, M. J., and Scher, H. I. (2007) (11)C-acetate PET imaging in prostate cancer. *Eur J Nucl Med Mol Imaging* 34, 181-4.
- Mlcochova, P., Plechanovova, A., Barinka, C., Mahadevan, D., Saldanha, J. W., Rulisek, L., and Konvalinka, J. (2007) Mapping of the active site of glutamate carboxypeptidase II by site-directed mutagenesis. *FEBS J* 274, 4731-41.
- Mhawech-Fauceglia, P., Zhang, S., Terracciano, L., Sauter, G., Chadhuri, A., Herrmann, F. R., and Penetrante, R. (2007) Prostate-specific membrane antigen (PSMA) protein expression in normal and neoplastic tissues and its sensitivity and specificity in prostate adenocarcinoma: an immunohistochemical study using multiple tumour tissue microarray technique. *Histopathology* 50, 472-83.
- Lambert, L. A., and Mitchell, S. L. (2007) Molecular evolution of the transferrin receptor/glutamate carboxypeptidase II family. *Journal of molecular evolution* 64, 113-28.
- Hlouchova, K., Barinka, C., Klusak, V., Sacha, P., Mlcochova, P., Majer, P., Rulisek, L., and Konvalinka, J. (2007) Biochemical characterization of human glutamate carboxypeptidase III. *Journal of neurochemistry* 101, 682-96.
- Goodman, O. B., Jr., Barwe, S. P., Ritter, B., McPherson, P. S., Vasko, A. J., Keen, J. H., Nanus, D. M., Bander, N. H., and Rajasekaran, A. K. (2007) Interaction of prostate specific membrane antigen with clathrin and the adaptor protein complex-2. *Int J Oncol* 31, 1199-203.
- Fall, K., Garmo, H., Andren, O., Bill-Axelsson, A., Adolfsson, J., Adami, H. O., Johansson, J. E., and Holmberg, L. (2007) Prostate-specific antigen levels as a predictor of lethal prostate cancer. *J Natl Cancer Inst* 99, 526-32.
- Barinka, C., Starkova, J., Konvalinka, J., and Lubkowski, J. (2007) A high-resolution structure of ligand-free human glutamate carboxypeptidase II. *Acta Crystallogr Sect F Struct Biol Cryst Commun* 63, 150-3.
- Barinka, C., Rovenska, M., Mlcochova, P., Hlouchova, K., Plechanovova, A., Majer, P., Tsukamoto, T., Slusher, B. S., Konvalinka, J., and Lubkowski, J. (2007) Structural insight into the pharmacophore pocket of human glutamate carboxypeptidase II. *J Med Chem* 50, 3267-73.
- Anderson, M. O., Wu, L. Y., Santiago, N. M., Moser, J. M., Rowley, J. A., Bolstad, E. S., and Berkman, C. E. (2007) Substrate specificity of prostate-specific membrane antigen. *Bioorg Med Chem* 15, 6678-86.
- Reske, S. N., Blumstein, N. M., Neumaier, B., Gottfried, H. W., Finsterbusch, F., Kocot, D., Moller, P., Glatting, G., and Perner, S. (2006) Imaging prostate cancer with 11C-choline PET/CT. *J Nucl Med* 47, 1249-54.
- Reske, S. N., Blumstein, N. M., and Glatting, G. (2006) [Advancement of PET and PET/CT in prostate carcinoma]. *Der Urologe. Aug. A* 45, 707-10, 712-4.
- Reske, S. N., Blumstein, N. M., and Glatting, G. (2006) [PET and PET/CT in relapsing prostate carcinoma]. *Der Urologe. Aug. A* 45, 1240, 1242-4, 1246-8, 1250.
- Oyama, N., Kaneda, T., Nakai, M., Shioyama, R., Matsuta, Y., Tanase, K., Aoki, Y., Miwa, Y., Akino, H., Yokoyama, O., Okazawa, H., Fujibayashi, Y., and Yonekura, Y. (2006) [Pet imaging in prostate cancer]. *Hinyokika Kyo* 52, 503-5.
- Mesters, J. R., Barinka, C., Li, W., Tsukamoto, T., Majer, P., Slusher, B. S., Konvalinka, J., and Hilgenfeld, R. (2006) Structure of glutamate carboxypeptidase II, a drug target in neuronal damage and prostate cancer. *EMBO J* 25, 1375-84.
- Kinoshita, Y., Kuratsukuri, K., Landas, S., Imaida, K., Rovito, P. M., Jr., Wang, C. Y., and Haas, G. P. (2006) Expression of prostate-specific membrane antigen in normal and malignant human tissues. *World J Surg* 30, 628-36.
- Elsasser-Beile, U., Wolf, P., Gierschner, D., Buhler, P., Schultze-Seemann, W., and Wetterauer, U. (2006) A new generation of monoclonal and recombinant antibodies against cell-adherent prostate specific membrane antigen for diagnostic and therapeutic targeting of prostate cancer. *Prostate* 66, 1359-70.
- Cunha, A. C., Weigle, B., Kiessling, A., Bachmann, M., and Rieber, E. P. (2006) Tissue-specificity of prostate specific antigens: comparative analysis of transcript levels in prostate and non-prostatic tissues. *Cancer Lett* 236, 229-38.
- Cimitan, M., Bortolus, R., Morassut, S., Canzonieri, V., Garbeglio, A., Baresic, T., Borsatti, E., Drigo, A., and Trovo, M. G. (2006) [18F]fluorocholine PET/CT imaging for the detection of recurrent prostate cancer at PSA relapse: experience in 100 consecutive patients. *Eur J Nucl Med Mol Imaging* 33, 1387-98.
- Extended European Search Report for European Application No. 18175078, dated Sep. 14, 2018.
- Office Action for counterpart Indonesia Patent Application No. P00201603202, dated Jan. 28, 2019 (with English Translation).
- Office Action for co-pending Canadian Application No. 2924360, dated Sep. 11, 2018.
- C. Dusich et al., General Approach for the Preparation of Fluorescent PSMA/GCPII Inhibitors, Abstract ID: 470 Poster board space: 29, *Molecular Imaging*, vol. 5, No. 3, pp. 322-323, Jul. 2006.
- Liu M., et al., "Synthesis and Biological Evaluation of Diethylenetriamine Pentaacetic acid-Polyethylene Glycol Folate: A

(56)

References Cited

OTHER PUBLICATIONS

new Folate-Derived, 99mTc-Based Radiopharmaceutical," *Bioconjugate Chem.*, 2005 vol. 16, p. 1126-1132.

Muller C., et al. "Synthesis and in Vitro/in Vivo Evaluation of Novel 99mTc(CO)₃-Folates," *Bioconjugate Chem.*, 2006 vol. 17, p. 797-806.

Viola-Villegas N., et al. "Targeting Gallium to Cancer Cells through the Folate Receptor," *Drug Target Insights*, 2008 vol. 3, p. 13-25.
Viola-Villegas N., et al. "Targeting the Folate Receptor (FR): Imaging and Cytotoxicity of Rel Conjugates in FR-Overexpressing Cancer Cells," *ChemMedChem*, 2008 vol. 3, p. 1387-1394.

Zhou J., "In vivo evaluation of medical device-associated inflammation using macrophage-specific position emission tomography (PET) imaging," *Bioorganic and Medicinal Chemistry Letters*, 2013 vol. 23, p. 2044-2047.

Kularatne S.A., et al. "Comparative Analysis of Folate Derived PET Imaging Agents with [18F]-2-Fluoro-2-deoxy-D-glucose Using Rodent Inflammatory Paw Model," *Molecular Pharmaceutics*, 2013 vol. 10, p. 3103-3111.

Office Action for counterpart Mexican Patent Application No. MX/a/2016/005013, dated Apr. 24, 2019 (with English Translation).
C. Foss et al., Synthesis and Validation of a Novel Small-Molecule Fluorescent Probe for PSMA Expression in Human Tumor Neovasculature, Abstract ID: 362, *Molecular Imaging*, vol. 4, No. 3, p. 321, Jul. 2005.

Office Action for counterpart Eurasian Patent Application No. 201690495, dated Dec. 20, 2018 (with English translation).

Harada, N., Kimura, H., Ono, M., and Saji, H. (2013) Preparation of asymmetric urea derivatives that target prostate-specific membrane antigen for SPECT imaging. *J Med Chem* 56, 7890-901.

Frigerio, B., Fracasso, G., Luisson, E., Cingarlini, S., Mortarino, M., Coliva, A., Seregini, E., Bombardieri, E., Zuccolotto, G., Rosato, A., Colombatti, M., Canevari, S., and Figini, M. (2013) A single-chain fragment against prostate specific membrane antigen as a tool to build theranostic reagents for prostate cancer. *Eur J Cancer* 49, 2223-32.

Franc, B. L., Cho, S. Y., Rosenthal, S. A., Cui, Y., Tsui, B., Vandewalker, K. M., Holz, A. L., Poonamallee, U., Pomper, M. G., and James, R. B. (2013) Detection and localization of carcinoma within the prostate using high resolution transrectal gamma imaging (TRGI) of monoclonal antibody directed at prostate specific membrane antigen (PSMA)-Proof of concept and initial imaging results. *European Journal of Radiology* 82, 1877-84.

Eder, M., Eisenhut, M., Babich, J., and Haberkorn, U. (2013) PSMA as a target for radiolabelled small molecules. *Eur J Nucl Med Mol Imaging* 40, 819-23.

Divyaa, S., Naushad, S. M., Murthy, P. V., Reddy Ch, R., and Kutala, V. K. (2013) GCPII modulates oxidative stress and prostate cancer susceptibility through changes in methylation of RASSF1, BNIP3, GSTP1 and Ec-SOD. *Molecular Biology Reports* 40, 5541-50.

Ceci, F., Castellucci, P., Mamede, M., Schiavina, R., Rubello, D., Fuccio, C., Ambrosini, V., Boschi, S., Martorana, G., and Fanti, S. (2013) (11)C-Choline PET/CT in patients with hormone-resistant prostate cancer showing biochemical relapse after radical prostatectomy. *Eur J Nucl Med Mol Imaging* 40, 149-55.

Benesova, M., Schafer, M., Bauder-Wüst, U., Mier, W., Haberkorn, U., Eisenhut, M., Kopka, K., and Eder, M. (2013) Linker Modifications of DOTA-conjugated Inhibitors of the Prostate-Specific Membrane Antigen (PSMA). *Eur J Nucl Med Mol Imaging* 40 (Suppl 2), S193.

Ben Jemaa, A., Sallami, S., Ceraline, J., and Oueslati, R. (2013) A novel regulation of PSMA and PSA expression by Q640X AR in 22Rv1 and LNCaP prostate cancer cells. *Cell biology international* 37, 464-70.

Ben Jemaa, A., Bouraoui, Y., Sallami, S., Noura, Y., and Oueslati, R. (2013) A comparison of the biological features of prostate cancer with (PSA+, PSMA+) profile according to RKIP. *BioMed Research International* 2013, 409179.

Barrett, J. A., Coleman, R. E., Goldsmith, S. J., Vallabhajosula, S., Petry, N. A., Cho, S., Armor, T., Stubbs, J. B., Maresca, K. P., Stabin, M. G., Joyal, J. L., Eckelman, W. C., and Babich, J. W. (2013) First-in-Man Evaluation of 2 High-Affinity PSMA-Avid Small Molecules for Imaging Prostate Cancer. *J Nucl Med* 54, 380-7.

Antunes, A. A., Reis, S. T., Leite, K. R., Real, D. M., Sousa-Canavez, J. M., Camara-Lopes, L. H., Dall'Oglio, M. F., and Srougi, M. (2013) PGC and PSMA in prostate cancer diagnosis: tissue analysis from biopsy samples. *International Braz J Urol : official journal of the Brazilian Society of Urology* 39, 649-56.

Afshar-Oromieh, A., Matchar, A., Eder, M., Eisenhut, M., Linhart, H. G., Hadaschik, B. A., Holland-Letz, T., Giesel, F. L., Kratochwil, C., Haufe, S., Haberkorn, U., and Zechmann, C. M. (2013) PET imaging with a [(68)Ga]gallium-labelled PSMA ligand for the diagnosis of prostate cancer biodistribution in humans and first evaluation of tumour lesions. *Eur J Nucl Med Mol Imaging* 40, 486-95.

Afshar-Oromieh, A., Haberkorn, U., Hadaschik, B., Habl, G., Eder, M., Eisenhut, M., Schlemmer, H. P., and Roethke, M. C. (2013) PET/MRI with a 68Ga-PSMA ligand for the detection of prostate cancer. *Eur J Nucl Med Mol Imaging* 40, 1629-30.

Abstracts of the Annual Congress of the European Association of Nuclear Medicine. Oct. 19-23, 2013. Lyon, France. *Eur J Nucl Med Mol Imaging* 40 Suppl 2, S1-S477.

Zhang, Y., Guo, Z., Du, T., Chen, J., Wang, W., Xu, K., Lin, T., and Huang, H. (2012) Prostate specific membrane antigen (PSMA): A novel modulator of p38 for proliferation, migration, and survival in prostate cancer cells. *Prostate* 73, 835-41.

Weissbach, L. (2012) [Which components should living guidelines contain?]. *Der Urologe*. 51, 57-9.

Thalmann, G. (2012) [Advanced prostate cancer: where are we going?]. *Der Urologe* 51, 7.

Taylor, R. M., Sevems, V., Brown, D. C., Bisoffi, M., and Sillerud, L. O. (2012) Prostate cancer targeting motifs: expression of alphanu beta3, neurotensin receptor 1, prostate specific membrane antigen, and prostate stem cell antigen in human prostate cancer cell lines and xenografts. *Prostate* 72, 523-32.

Spahn, M., and Krebs, M. (2012) [How should hormone therapy for castration-resistant prostate cancer be continued?]. *Der Urologe*. 51, 15-9.

Schafer, M., Bauder-Wüst, U., Leotta, K., Zoller, F., Mier, W., Haberkorn, U., Eisenhut, M., and Eder, M. (2012) A dimerized urea-based inhibitor of the prostate-specific membrane antigen for 68Ga-PET imaging of prostate cancer. *EJNMMI Res* 2, 23.

Cressler, S., Putora, P. M., Plasswilm, L., and Schmid, H. P. (2012) [Castration-resistant prostate cancer: surgical and radio-oncological therapeutic options]. *Der Urologe*. 51, 27-31.

Poulsen, M. H., Bouchelouche, K., Hoiland-Carlsen, P. F., Petersen, H., Gerke, O., Steffansen, S. I., Marcussen, N., Svolgaard, N., Vach, W., Geertsen, U., and Walter, S. (2012) [18F]fluoromethylcholine (FCH) positron emission tomography/computed tomography (PET/CT) for lymph node staging of prostate cancer: a prospective study of 210 patients. *BJU Intl* 110, 1666-71.

Pinto, F., Totaro, A., Palermo, G., Calarco, A., Sacco, E., D'Addessi, A., Racioppi, M., Valentini, A., Gui, B., and Bassi, P. (2012) Imaging in prostate cancer staging: present role and future perspectives. *Urol Int* 88, 125-36.

Pavlicek, J., Ptacek, J., and Barinka, C. (2012) Glutamate carboxypeptidase II: an overview of structural studies and their importance for structure-based drug design and deciphering the reaction mechanism of the enzyme. *Curr Med Chem* 19, 1300-9.

Omlin, A., and Gillesen, S. (2012) [Inhibitors of androgen and estrogen biosynthesis in castration-resistant prostate cancer]. *Der Urologe*. 51, 8-14.

Nedrow-Byers, J. R., Jabbes, M., Jewett, C., Ganguly, T., He, H., Liu, T., Benny, P., Bryan, J. N., and Berkman, C. E. (2012) A phosphoramidate-based prostate-specific membrane antigen-targeted SPECT agent. *Prostate* 72, 904-12.

Moltzahn, F., and Thalmann, G. N. (2012) [Bone metastasis in prostate cancer]. *Der Urologe*. 51, 20-6.

Meinhardt, W., van der Poel, H. G., Valdes Olmos, R. A., Bex, A., Brouwer, O. R., and Horenblas, S. (2012) Laparoscopic sentinel

(56)

References Cited

OTHER PUBLICATIONS

- lymph node biopsy for prostate cancer: the relevance of locations outside the extended dissection area. *Prostate Cancer* 2012, 751753.
- Malik, N., Zlatopolskiy, B., Machulla, H.-J., Reske, S. N., and Solbach, C. (2012) One pot radiofluorination of a new potential PSMA ligand [Al18F]NOTA-DUPA-Pep. *Journal of Labelled Compounds and Radiopharmaceuticals* 55, 320-325.
- Lutje, S., Boerman, O. C., van Rij, C. M., Sedelaar, M., Helfrich, W., Oyen, W. J., and Mulders, P. F. (2012) Prospects in radionuclide imaging of prostate cancer *Prostate* 72, 1262-72.
- Liu, T., Wu, L. Y., Fulton, M. D., Johnson, J. M., and Berkman, C. E. (2012) Prolonged androgen deprivation leads to downregulation of androgen receptor and prostate-specific membrane antigen in prostate cancer cells. *Int J Oncol* 41, 2087-92.
- Liu, T., Nedrow-Byers, J. R., Hopkins, M. R., Wu, L. Y., Lee, J., Reilly, P. T., and Berkman, C. E. (2012) Targeting prostate cancer cells with a multivalent PSMA inhibitor-guided streptavidin conjugate. *Bioorg Med Chem Lett* 22, 3931-4.
- Lees, K., Durve, M., and Parker, C. (2012) Active surveillance in prostate cancer: patient selection and triggers for intervention. *Curr Opin Urol* 22, 210-5.
- Kuru, T. H., Tulea, C., Simpfordorfer, T., Popeneciu, V., Roethke, M., Hadaschik, B. A., and Hohenfellner, M. (2012) [MRI navigated stereotactic prostate biopsy: fusion of MRI and real-time transrectal ultrasound images for perineal prostate biopsies]. *Der Urologe*. 51, 50-6.
- Kosuri, S., Akhtar, N. H., Smith, M., Osborne, J. R., and Tagawa, S. T. (2012) Review of salvage therapy for biochemically recurrent prostate cancer: the role of imaging and rationale for systemic salvage targeted anti-prostate-specific membrane antigen radioimmunotherapy. *Advances in Urology* 2012, 921674.
- Klotz, L. (2012) Cancer overdiagnosis and overtreatment. *Curr Opin Urol* 22, 203-9.
- Jadvar, H. (2012) Molecular imaging of prostate cancer: PET radiotracers. *AJR Am J Roentgenol* 199, 278-91.
- Hlouchova, K., Navratil, V., Tykvart, J., Sacha, P., and Konvalinka, J. (2012) GCPII variants, paralogs and orthologs. *Curr Med Chem* 19, 1316-22.
- Heidenreich, A. (2012) [Immunotherapy for metastatic prostate cancer: do we really need this?]. *Der Urologe*. 51, 32-8.
- Grant, C. L., Caromile, L. A., Ho, V., Durrani, K., Rahman, M. M., Claffey, K. P., Fong, G. H., and Shapiro, L. H. (2012) Prostate specific membrane antigen (PSMA) regulates angiogenesis independently of VEGF during ocular neovascularization. *PLoS One* 7, e41285.
- Graham, K., Lesche, R., Gromov, A. V., Bohnke, N., Schafer, M., Hassfeld, J., Dinkelborg, L., and Ketschau, G. (2012) Radiofluorinated derivatives of 2-(phosphonomethyl)pentanedioic acid as inhibitors of prostate specific membrane antigen (PSMA) for the imaging of prostate cancer. *J Med Chem* 55, 9510-20.
- Foss, C. A., Mease, R. C., Cho, S. Y., Kim, H. J., and Pomper, M. G. (2012) GCPII imaging and cancer. *Curr Med Chem* 19, 1346-59.
- Fortuin, A. S., Deserno, W. M., Meijer, H. J., Jager, G. J., Takahashi, S., Debats, O. A., Reske, S. N., Schick, C., Krause, B. J., van Oort, I., Wijtes, A. J., Hoogeveen, Y. L., van Lin, E. N., and Barentsz, J. O. (2012) Value of PET/CT and MR lymphography in treatment of prostate cancer patients with lymph node metastases. *Int J Radiat Oncol Biol Phys* 84(3), 712-8.
- Eder, M., Schafer, M., Bauder-Wust, U., Hull, W. E., Wangler, C., Mier, W., Haberkorn, U., and Eisenhut, M. (2012) (68)Ga-Complex Lipophilicity and the Targeting Property of a Urea-Based PSMA Inhibitor for PET Imaging. *Bioconjugate Chem* 23, 688-97.
- De Santis, M., and Bachner, Y. (2012) [Role of chemotherapy in castration-resistant prostate cancer: are there new approaches?]. *Der Urologe*. 51, 39-43.
- Chen, Z., Penet, M. F., Nimmagadda, S., Li, C., Banerjee, S. R., Winnard, P. T., Jr., Artemov, D., Glunde, K., Pomper, M. G., and Bhujwalla, Z. M. (2012) PSMA-targeted theranostic nanoplex for prostate cancer therapy. *ACS Nano* 6, 7752-62.
- Chen, Y., Pullambhatla, M., Banerjee, S. R., Byun, Y., Stathis, M., Rojas, C., Slusher, B. S., Mease, R. C., and Pomper, M. G. (2012) Synthesis and biological evaluation of low molecular weight fluorescent imaging agents for the prostate-specific membrane antigen. *Bioconjugate Chem* 23, 2377-85.
- Chen, J., Zhao, Y., Li, X., Sun, P., Wang, M., Wang, R., and Jin, X. (2012) Imaging primary prostate cancer with 11C-Choline PET/CT: relation to tumour stage, Gleason score and biomarkers of biologic aggressiveness. *Radiology and Oncology* 46(3), 179-88.
- Baiz, D., Pinder, T. A., Hassan, S., Karpova, Y., Salisbury, F., Welker, M. E., and Kulik, G. (2012) Synthesis and characterization of a novel prostate cancer-targeted phosphatidylinositol-3-kinase inhibitor prodrug. *J Med Chem* 55, 8038-46.
- Afshar-Oromieh, A., Haberkorn, U., Eder, M., Eisenhut, M., and Zechmann, C. M. (2012) [68Ga]Gallium-labelled PSMA ligand as superior PET tracer for the diagnosis of prostate cancer: comparison with 18F-FECH. *Eur J Nucl Med Mol Imaging* 39, 1085-6.
- Search Report for Counterpart European Application No. 18184296, dated Jan. 23, 2019.
- Minner, S., Wittmer, C., Graefen, M., Salomon, G., Steuber, T., Haese, A., Huland, H., Bokemeyer, C., Yekebas, E., Dierlamm, J., Balabanov, S., Kilic, E., Wilczak, W., Simon, R., Sauter, G., and Schlomm, T. (2011) High level PSMA expression is associated with early PSA recurrence in surgically treated prostate cancer. *Prostate* 71, 281-8.
- Malik, N., Machulla, H. J., Solbach, C., Winter, G., Reske, S. N., and Zlatopolskiy, B. (2011) Radiosynthesis of a new PSMA targeting ligand ([18F]FPy-DUPA-Pep). *Appl Radiat Isot* 69, 1014-8.
- Lord, M., Ratib, O., and Vallee, J. P. (2011) (1)(8)F-Fluorocholine integrated PET/MRI for the initial staging of prostate cancer. *Eur J Nucl Med Mol Imaging* 38, 2288.
- Ho, C. L., Liu, I. H., Wu, Y. H., Chen, L. C., Chen, C. L., Lee, W. C., Chuang, C. H., Lee, T. W., Lin, W. J., Shen, L. H., and Chang, C. H. (2011) Molecular imaging, pharmacokinetics, and dosimetry of In-AMBA in human prostate tumor-bearing mice. *J Biomed Biotechnol* 2011, 101497.
- Hillier, S. M., Kern, A. M., Maresca, K. P., Marquis, J. C., Eckelman, W. C., Joyal, J. L., and Babich, J. W. (2011) 123I-MIP-1072, a small-molecule inhibitor of prostate-specific membrane antigen, is effective at monitoring tumor response totaxane therapy. *J Nucl Med* 52, 1087-93.
- Fortmuller, K., Alt, K., Gierschner, D., Wolf, P., Baum, V., Freudenberg, N., Wetterauer, U., Elsasser-Beile, U., and Buhler, P. (2011) Effective targeting of prostate cancer by lymphocytes redirected by a PSMA x CD3 bispecific single-chain diabody. *Prostate* 71, 588-96.
- Evans, M. J., Smith-Jones, P. M., Wongvipat, J., Navarro, V., Kim, S., Bander, N. H., Larson, S. M., and Sawyers, C. L. (2011) Noninvasive measurement of androgen receptor signaling with a positron-emitting radiopharmaceutical that targets prostate-specific membrane antigen. *Proc Natl Acad Sci U S A* 108, 9578-82.
- Emonds, K. M., Swinnen, J. V., van Weerden, W. M., Vanderhoydonc, F., Nuyts, J., Mortelmans, L., and Mottaghy, F. M. (2011) Do androgens control the uptake of 18F-FDG, 11C-choline and 11C-acetate in human prostate cancer cell lines? *Eur J Nucl Med Mol Imaging* 38, 1842-53.
- Dahl, M., Bouchelouche, P., Kramer-Marek, G., Capala, J., Nordling, J., and Bouchelouche, K. (2011) Sarcosine induces increase in HER2/neu expression in androgen-dependent prostate cancer cells. *Molecular Biology Reports* 38, 4237-43.
- Chuu, C. P., Kokontis, J. M., Hiipakka, R. A., Fukuchi, J., Lin, H. P., Lin, C. Y., Huo, C., Su, L. C., and Liao, S. (2011) Androgen suppresses proliferation of castration-resistant LNCaP 104-R2 prostate cancer cells through androgen receptor, Skp2, and c-Myc. *Cancer science* 102, 2022-8.
- Chen, Y., Pullambhatla, M., Byun, Y., Foss, C. A., Nimmagadda, S., Senthamizchelvan, S., Sgouros, G., Mease, R. C., and Pomper, M. G. (2011) 2-(3-(1-Carboxy-5-[(6-[18F]fluoro-pyridine-3-carbonyl)-amino]-pentyl)-urei do)-pentanedioic acid, [18F]DCFPyL, a PSMA-based PET Imaging Agent for Prostate Cancer. *Clin Cancer Res* 17(24), 7645-53.
- Bouchelouche, K., Tagawa, S. T., Goldsmith, S. J., Turkbey, B., Capala, J., and Choyke, P. (2011) PET/CT Imaging and Radioimmunotherapy of Prostate Cancer. *Semin Nucl Med* 41, 29-44.

(56)

References Cited

OTHER PUBLICATIONS

- Banerjee, S. R., Pullambhatla, M., Shallal, H., Lisok, A., Mease, R. C., and Pomper, M. G. (2011) A Modular Strategy to Prepare Multivalent Inhibitors of Prostate-Specific Membrane Antigen (PSMA). *Oncotarget* 2(12), 1244-53.
- Zhang, A. X., Murelli, R. P., Barinka, C., Michel, J., Cocleaza, A., Jorgensen, W. L., Lubkowski, J., and Spiegel, D. A. (2010) A remote arene-binding site on prostate specific membrane antigen revealed by antibody-recruiting small molecules. *J Am Chem Soc* 132, 12711-6.
- Wang, H., Byun, Y., Barinka, C., Pullambhatla, M., Bhang, H. E., Fox, J. J., Lubkowski, J., Mease, R. C., and Pomper, M. G. (2010) Bioisosterism of urea-based GCPII inhibitors: Synthesis and structure-activity relationship studies. *Bioorg Med Chem Lett* 20, 392-7.
- Rioja, J., Rodriguez-Fraile, M., Lima-Favaretto, R., Rincon-Mayans, A., Penuelas-Sanchez, I., Zudaire-Bergera, J. J., and Parra, R. O. (2010) Role of positron emission tomography in urological oncology. *BJU Int* 106, 1578-93.
- Poulsen, M. H., Bouchelouche, K., Gerke, O., Petersen, H., Svolgaard, B., Marcussen, N., Svolgaard, N., Ogren, M., Vach, W., Hoiland-Carlson, P. F., Geertsen, U., and Walter, S. (2010) [18F]-fluorocholine positron-emission/computed tomography for lymph node staging of patients with prostate cancer: preliminary results of a prospective study. *BJU Int* 106, 639-44.
- Mertens, K., Slaets, D., Lambert, B., Acou, M., De Vos, F., and Goethals, I. (2010) PET with (18)F-labelled choline-based tracers for tumour imaging: a review of the literature. *Eur J Nucl Med Mol Imaging* 37, 2188-93.
- Maresca, K., Hillier, S., Lu, G., Marquis, J., Zimmerman, G., Eckelman, W., Joyal, J., and Babich, J. (2010) Influence of functionalized chelators on affinity and pharmacokinetics of 99mTc(CO)3-labeled small molecules targeting prostate specific membrane antigen (PSMA). *J Nucl Med* 51, 250.
- Liu, T., Wu, L. Y., Hopkins, M. R., Choi, J. K., and Berkman, C. E. (2010) A targeted low molecular weight near-infrared fluorescent probe for prostate cancer. *Bioorg Med Chem Lett* 20, 7124-6.
- Kularatne, S. A., Venkatesh, C., Santhapuram, H. K., Wang, K., Vaitilingam, B., Henne, W. A., and Low, P. S. (2010) Synthesis and biological analysis of prostate-specific membrane antigen-targeted anticancer prodrugs. *J Med Chem* 53, 7767-77.
- Jambor, I., Borra, R., Kemppainen, J., Lepomaki, V., Parkkola, R., Dean, K., Alanen, K., Arponen, E., Nurmi, M., Aronen, H. J., and Minn, H. (2010) Functional imaging of localized prostate cancer aggressiveness using 11C-acetate PET/CT and 1H-MR spectroscopy. *J Nucl Med* 51, 1676-83.
- Hong, H., Zhang, Y., Sun, J., and Cai, W. (2010) Positron emission tomography imaging of prostate cancer. *Amino Acids* 39, 11-27.
- Holland, J. P., Divilov, V., Bander, N. H., Smith-Jones, P. M., Larson, S. M., and Lewis, J. S. (2010) 89Zr-DFO-J591 for immunoPET of prostate-specific membrane antigen expression in vivo. *J Nucl Med* 51, 1293-300.
- Giovacchini, G., Picchio, M., Coradeschi, E., Bettinardi, V., Gianolli, L., Scattoni, V., Cozzarini, C., Di Muzio, N., Rigatti, P., Fazio, F., and Messa, C. (2010) Predictive factors of [(11)C]choline PET/CT in patients with biochemical failure after radical prostatectomy. *Eur J Nucl Med Mol Imaging* 37, 301-9.
- Bouchelouche, K., Turkbey, B., Choyke, P., and Capala, J. (2010) Imaging prostate cancer: an update on positron emission tomography and magnetic resonance imaging. *Current urology reports* 11, 180-90.
- Bouchelouche, K., Choyke, P. L., and Capala, J. (2010) Prostate specific membrane antigen—a target for imaging and therapy with radionuclides. *Discov Med* 9, 55-61.
- Bouchelouche, K., and Capala, J. (2010) 'Image and treat': an individualized approach to urological tumors. *Curr Opin Oncol* 22, 274-80.
- Banerjee, S. R., Pullambhatla, M., Byun, Y., Nimmagadda, S., Green, G., Fox, J. J., Horti, A., Mease, R. C., and Pomper, M. G. (2010) 68Ga-labeled inhibitors of prostate-specific membrane antigen (PSMA) for imaging prostate cancer. *J Med Chem* 53, 5333-41.
- Alt, K., Wiehr, S., Ehrlichmann, W., Reischl, G., Wolf, P., Pichler, B. J., Elsasser-Beile, U., and Buhler, P. (2010) High-resolution animal PET imaging of prostate cancer xenografts with three different 64Cu-labeled antibodies against native cell-adherent PSMA. *Prostate* 70, 1413-21.
- Zaheer, A., Cho, S. Y., and Pomper, M. G. (2009) New agents and techniques for imaging prostate cancer. *J Nucl Med* 50, 1387-90.
- Rinnab, L., Simon, J., Hautmann, R. E., Cronauer, M. V., Hohl, K., Buck, A. K., Reske, S. N., and Mottaghy, F. M. (2009) [(11)C]choline PET/CT in prostate cancer patients with biochemical recurrence after radical prostatectomy. *World J Urol* 27, 619-25.
- Pillarsetty, N., Punzalan, B., and Larson, S. M. (2009) 2-18F-Fluoropropionic acid as a PET imaging agent for prostate cancer. *J Nucl Med* 50, 1709-14.
- Maresca, K. P., Hillier, S. M., Femia, F. J., Keith, D., Barone, C., Joyal, J. L., Zimmerman, C. N., Kozikowski, A. P., Barrett, J. A., Eckelman, W. C., and Babich, J. W. (2009) A series of halogenated heterodimeric inhibitors of prostate specific membrane antigen (PSMA) as radiolabeled probes for targeting prostate cancer. *J Med Chem* 52, 347-57.
- Mannweiler, S., Amersdorfer, P., Trajanoski, S., Terrell, J. A., King, D., and Mehes, G. (2009) Heterogeneity of prostate-specific membrane antigen (PSMA) expression in prostate carcinoma with distant metastasis. *Pathology oncology research : POR* 15, 167-72.
- Lapi, S. E., Wahnische, H., Pham, D., Wu, L. Y., Nedrow-Byers, J. R., Liu, T., Vejdani, K., VanBrocklin, H. F., Berkman, C. E., and Jones, E. F. (2009) Assessment of an 18F-labeled phosphoramidate peptidomimetic as a new prostate-specific membrane antigen-targeted imaging agent for prostate cancer. *J Nucl Med* 50, 2042-8.
- Kwee, S. A., Coel, M. N., Ly, B. H., and Lim, J. (2009) (18)F-Choline PET/CT imaging of RECIST measurable lesions in hormone refractory prostate cancer. *Ann Nucl Med* 23, 541-8.
- Kularatne, S. A., Zhou, Z., Yang, J., Post, C. B., and Low, P. S. (2009) Design, synthesis, and preclinical evaluation of prostate-specific membrane antigen targeted (99m)Tc-radioimaging agents. *Mol Pharm* 6, 790-800.
- Kularatne, S. A., Wang, K., Santhapuram, H. K., and Low, P. S. (2009) Prostate-specific membrane antigen targeted imaging and therapy of prostate cancer using a PSMA inhibitor as a homing ligand. *Mol Pharm* 6, 780-9.
- Klusak, V., Barinka, C., Plechanovova, A., Mlcochova, P., Konvalinka, J., Rulisek, L., and Lubkowski, J. (2009) Reaction mechanism of glutamate carboxypeptidase II revealed by mutagenesis, X-ray crystallography, and computational methods. *Biochemistry* 48, 4126-38.
- Jadvar, H. (2009) Molecular imaging of prostate cancer with 18F-fluorodeoxyglucose PET. *Nat Rev Urol* 6, 317-23.
- Humblet, V., Misra, P., Bhushan, K. R., Nasr, K., Ko, Y. S., Tsukamoto, T., Pannier, N., Frangioni, J. V., and Maison, W. (2009) Multivalent scaffolds for affinity maturation of small molecule cell surface binders and their application to prostate tumor targeting. *J Med Chem* 52, 544-50.
- Hlouchova, K., Barinka, C., Konvalinka, J., and Lubkowski, J. (2009) Structural insight into the evolutionary and pharmacologic homology of glutamate carboxypeptidases II and III. *FEBS J* 276, 4448-62.
- Hillier, S. M., Maresca, K. P., Femia, F. J., Marquis, J. C., Foss, C. A., Nguyen, N., Zimmerman, C. N., Barrett, J. A., Eckelman, W. C., Pomper, M. G., Joyal, J. L., and Babich, J. W. (2009) Preclinical evaluation of novel glutamate-urea-lysine analogues that target prostate-specific membrane antigen as molecular imaging pharmaceuticals for prostate cancer. *Cancer Res* 69, 6932-40.
- Haffner, M. C., Kronberger, I. E., Ross, J. S., Sheehan, C. E., Zitt, M., Muhlmann, G., Ofner, D., Zelger, B., Ensinger, C., Yang, X. J., Geley, S., Margreiter, R., and Bander, N. H. (2009) Prostate-specific membrane antigen expression in the neovasculature of gastric and colorectal cancers. *Hum Pathol* 40, 1754-61.
- Elsasser-Beile, U., Reischl, G., Wiehr, S., Buhler, P., Wolf, P., Alt, K., Shively, J., Judenhofer, M. S., Machulla, H. J., and Pichler, B. J. (2009) PET imaging of prostate cancer xenografts with a highly specific antibody against the prostate-specific membrane antigen. *J Nucl Med* 50, 606-11.

(56)

References Cited

OTHER PUBLICATIONS

- Elsasser-Beile, U., Buhler, P., and Wolf, P. (2009) Targeted therapies for prostate cancer against the prostate specific membrane antigen. *Curr Drug Targets* 10, 118-25.
- Chen, Y., Dhara, S., Banerjee, S. R., Byun, Y., Pullambhatla, M., Mease, R. C., and Pomper, M. G. (2009) A low molecular weight PSMA-based fluorescent imaging agent for cancer. *Biochem Biophys Res Commun* 390, 624-9.
- Bouchelouche, K., Capala, J., and Oehr, P. (2009) Positron emission tomography/computed tomography and radioimmunotherapy of prostate cancer. *Curr Opin Oncol* 21, 469-74.
- Belloli, S., Jachetti, E., Moresco, R. M., Picchio, M., Lecchi, M., Valtorta, S., Freschi, M., Michelini, R. H., Bellone, M., and Fazio, F. (2009) Characterization of preclinical models of prostate cancer using PET-based molecular imaging. *Eur J Nucl Med Mol Imaging* 36, 1245-55.
- Office Action for counterpart Israel Patent Application No. 245113, dated Jan. 10, 2019 (with English Translation).
- Office Action for counterpart Georgian Patent Application No. AP 2014 014132, dated Jan. 11, 2019 (with English Translation).
- Second Examination Report for counterpart Chilean Patent Application No. 00883-2016, dated Jan. 16, 2019 (with English Translation).
- M. De Santis et al., "Role of Chemotherapy in Castration-Resistant Prostate Cancer." *Der Urologe* 1:39-43, 2012.
- A. Heidenreich, "Immunotherapy for Metastatic Prostate Cancer—Do we really need this?" *Der Urologe* 1(1):32-38, 2012.
- T.H. Kuru et al., "MRI Navigated Stereotactic Prostate Biopsy." *Der Urologe* 1(1):50-56, 2012.
- F. Moltzahn et al., "Bone Metastasis in Prostate Cancer." *Der Urologe* 1(1):20-26, 2012.
- A. Omlin et al., "Inhibitors of Androgen and Estrogen Biosynthesis in Castration-Resistant Prostate Cancer." *Der Urologe* 1(1):8-14, 2012.
- S.N. Reske et al., "Advancement of PET and PET/CT in Prostate Carcinoma." *Der Urologe* 45(6):707-714, 2006.
- S.N. Reske, "Nuclear Imaging of Prostate Cancer." *Urologe* 46(11):1485-1499, 2007.
- S.N. Reske et al., "PET and PET/CT in Relapsing Prostate Carcinoma." *Der Urologe* 45(10):1240-1250-1499, 2006.
- M. Spahn et al., "How Should Hormone Therapy for Castration-Resistant Cancer be Continued?" *Der Urologe* 51 (1):15-19, 2012.
- G. Thalmann, "Advanced Prostate Cancer Where are we going?" *Der Urologe* 51(1):7, 2012.
- L. Weissbach, "Which Components Should 'Living Guidelines' Contain?" *Der Urologe* 51(1):57-59, 2012.
- S. Preusser et al., "Castration-Resistant Prostate Cancer." *Der Urologe* 51(1):27-31, 2012.
- Y-q. Wu et al., "A Mild Deprotection Procedure for Tert-butyl Esters and Tert-butyl Ethers Using ZnBr₂ in Methylene Chloride." *Tetrahedron Letters* 41, 2847-2849, 2000.
- J. Meienhofer et al., "Solid-Phase Synthesis with Attachment of Peptide to Resin Through an Amino Acid Side Chain: [8-Lysine]-Vasopressin." *Proc. Nat. Acad. Sci., USA*, vol. 68, No. 5, pp. 1006-1009, May 1971.
- Banerjee, S. et al., "Sequential SPECT and Optical Imaging of Experimental Models of Prostate Cancer with a Dual Modality Inhibitor of the Prostate-Specific Membrane Antigen." *Angewandte Chemie International Edition*, 2011, 50, 9167-9170.
- Banerjee, S.R. et al. "Synthesis and Evaluation of Technetium-99m- and Rhenium-Labeled Inhibitors of the Prostate-Specific Membrane Antigen (PSMA)." *J. Med. Chem.* Aug. 14, 2008; 51(15): 4504-4517.
- Bennett, V.J., "Analysis of fluorescently labeled substance P analogs: binding, imaging and receptor activation." *BMC Chemical Biology*, 2001, 1:1. doi: 10.1186/1472-6769-1-1.
- Chen, Ying, et al., "Radiohalogenated Prostate-Specific Membrane Antigen (PSMA)-Based Ureas as Imaging Agents for Prostate Cancer." *J. Med. Chem.*, 2008, 51 (24), pp. 7933-7943.
- Cole et al., "Cancer theranostics: the rise of targeted magnetic nanoparticles," *Trends in Biotechnology*, 2011, 29, 323-332.
- Davis, Mindy I., et al., "Crystal Structure of Prostate-Specific Membrane Antigen, A Tumor Marker and Peptidase", *Apr. 26, 2005*, PNAS, vol. 102, No. 17, pp. 5981-5986.
- Definition of ligand, Random House Kernerman Webster's College IMMM Dictionary, downloaded on Jan. 25, 2014 from <http://www.thefreedictionary.com/ligand>, 1 page.
- Eder et al., "68Ga-complex lipophilicity and the targeting property of a urea-based PSMA inhibitor for PET imaging," *Bioconjugate Chemistry*, 2012; 23:688-697.
- Farokhzad, et al., "Nanoparticle-Aptamer Bioconjugates: A New Approach for Targeting Prostate Cancer Cells," *Cancer Research* 64, pp. 7668-7672 (2004).
- Foss, Catherine A., et al. "Radiolabeled small-molecule ligands for prostate-specific membrane antigen: in vivo maging in experimental models of prostate cancer." *Clinical cancer research* 11.11 (2005): 4022-4028.
- Gomez-Hens et al., "Long wavelength fluorophores: new trends in their analytical use," *Trends in Analytical Chemistry*, 2004; 23: 127-136.
- Henne, et al., "Synthesis and activity of a folate peptide camptothecin prodrug," *ScienceDirect, Bioorganic & Medical Chemistry Letters* 16 (2006) 5350-5355.
- Hillier, Shawn M., et al., "Preclinical Evaluation of Novel Glutamate-Urea-Lysine Analogues That Target Prostate-Specific Membrane Antigen as Molecular Imaging Pharmaceuticals for Prostate Cancer," *Cancer Res. Sep. 1, 2009; 69 (17):6932-40*.
- Jackson, Paul F., et al., "Design of NAALADase Inhibitors: A Novel Neuroprotective Strategy", 2001, *Current Medicinal Chemistry*, vol. 8, No. 8, pp. 949-957.
- Jayaprakash, Sarva, et al. "Design and synthesis of a PSMA inhibitor-doxorubicin conjugate for targeted prostate cancer therapy." *ChemMedChem* 1.3 (2006): 299-302.
- Kaur, G. et al., "Biological evaluation of tubulysin A: a potential anticancer and antiangiogenic natural product," *Biochem. J.*, 2006, 396, 235-242.
- Kozikowski, Alan P., et al., "Design of Remarkably Simple, Yet Potent Urea-Based Inhibitors of Glutamate Carboxypeptidase II (NAALADase)" 2, 001, *Journal of Medicinal Chemistry*, vol. 44, No. 3, pp. 298-301.
- Kozikowski, Alan P., et al., "Synthesis of Urea-Based Inhibitors as Active Site Probes of Glutamate Carboxypeptidase II: Efficacy as Analgesic Agents", 2004, *Journal of Medicinal Chemistry*, vol. 47, No. 7, pp. 1729-1738.
- Kularatne et al., "Prostate-specific membrane antigen targeted imaging and therapy of prostate cancer using a PSMA inhibitor as a homing ligand," *Molecular Pharmaceutics*, 6(3): 780-789 (2009).
- Kularatne, S., "Synthesis and Biological Analysis of Prostate-Specific Membrane Antigen-Targeted Anticancer Prodrugs," *J. Med. Chem.* 2010, 53(21), 7767-7777.
- Larock, "Comprehensive Organic Transformations, a guide to functional group preparations," VCH Publishers, Inc. New York (1989).
- Lu, G. et al., "Synthesis and SAR of 99mTc/Re-labeled small molecule prostate specific membrane antigen inhibitors with novel polar chelates," *Bioorganic and Medicinal Chemistry Letters*, 2013, 23, 1557-1563.
- Lupold, et al., "Identification and characterization of nuclease-stabilized RNA molecules that bind human prostate cancer cells via the prostate-specific membrane antigen." *Cancer Res.* 2002; 62:4029-4033.
- Majer, Pavel., et al., "Synthesis and Biological Evaluation of Thiol-Based Inhibitors of Glutamate Carboxypeptidase II: Discovery of an Orally Active GCP II Inhibitor", 2003, *Journal of Medicinal Chemistry*, vol. 46, No. 10, pp. 1989-1996.
- Maresca, K. P., "A Series of Halogenated Heterodimeric Inhibitors of Prostate Specific Membrane Antigen (PSMA) as Radiolabeled Probes for Targeting Prostate Cancer," *J. Med. Chem.*, 2009, 52 (2), pp. 347-357.
- Maresca, K., et al., "Molecular targeting of prostate cancer with small molecule inhibitors of prostate specific membrane antigen (PSMA)," *J. Nucl. Med.* 2007, 48 (Supplement 2):25P.
- Martin, et al., *Helv. Chim. Acta*, 78, 486-504 (1995) and Abstract.

(56)

References Cited**OTHER PUBLICATIONS**

McNamara et al, Cell type specific delivery of siRNAs with aptamer-siRNA chimeras, *Nature Biotechnology*, 2006; 24: 1005-1015.

Melby, et al., *Cancer Research* 53(8), pp. 1755-1760 (1993).

Mesters, et al., et al., "Structure of Glutamate Carboxypeptidase II, a Drug Target in Neuronal Damage and Prostate Cancer" 2006, *The EMBO Journal*, vol. 25, No. 6, pp. 1375-1384.

Olsnes, S., et al., *Immunology Today*, 10, pp. 291-295 (1989).

Paranjpe, et al., "Tumor-targeted bioconjugate based delivery of camptothecin: design, synthesis and in vitro evaluation," *ScienceDirect Journal of Controlled Release* 100 (2004) 275-292.

Pathalk et al., Enzymic protecting group techniques in organic synthesis, *Stereosel. Biocatal.* 775-797 (2000).

Peltier et al., "The Total Synthesis of Tubulysin D," *J. Am. Chem. Soc.* 128:16018-19 (2006).

Pubchem, Compound summary for: CID 58099954, Aug. 19, 2012.

Radioisotopes in Medicine, from http://www.word-nuclear.org/information-library/non-power-nuclear_applications/radioisotopes-research/radioisotopes-in-medicine.aspx, Dec. 28, 2016, pp. 1-20.

Ranasinghe, M. G., et al., "Facile Synthesis of Unsymmetrical Thiol-sulfonates via Sulfenylation of Mercaptans", 1988, *Synthetic Communications*, vol. 18, No. 3, pp. 227-232.

Reddy et al., "PS MA-specific anti-tumor activity of the targeted-tubulysin conjugate, EC 1169," American Association for Cancer Research Annual Meeting (Apr. 8, 2013) Poster.

Reddy et al., "PS MA-specific anti-tumor activity of the targeted-tubulysin conjugate, EC 1169," American Association for Cancer Research Annual Meeting (Apr. 8, 2013) Presentation Abstract.

Roy, et al., "DUPA Conjugation of a Cytotoxic Indenoisoquinoline Topoisomerase I Inhibitor for Selective Prostate Cancer Cell Targeting," *J. Med. Chem.* 58 (2015) 3094-3103.

Theodora E. Greene & Peter G.M. Wuts, "Protective Groups in Organic Synthesis," 2d edition, John Wiley & Sons, Inc. New York (1991).

Truffert, et al., *Tetrahedron*, 52:3005 (1996).

Vlahov, et al., "Design and regioselective synthesis of a new generation of targeted chemotherapeutics. Part 1: EC145, a folic acid conjugate of desacetylvinblastine monohydrazide," *ScienceDirect, Bioorganic & Medical Chemistry Letters* 16 (2006) 5093-5096.

Wang et al., "Prostate-Specific Membrane Antigen Targeted Tubulysin Conjugates for Cancer Therapy," 246th ACS National Meeting and Exposition (Sep. 8, 2013) Poster.

Yang, et al., "Characterization of the pH of Folate Receptor-Containing Endosomes and the Rate of Hydrolysis of Internalized Acid-Labile Folate-Drug Conjugates," *JPET* 321: 462-468, 2007.

Benesova, M. et al., "Linker Modifications of DOTA-conjugated Inhibitors of the Prostate-Specific Membrane Antigen (PSMA)," poster, presented at the European Association of Nuclear Medicine Conference on Oct. 21, 2013.

Benesova, M. et al., "Linker Modifications of DOTA-conjugated Inhibitors of the Prostate-Specific Membrane Antigen (PSMA)," abstract, *Eur. J. Nucl. Med. Mol. Imaging*, available Oct. 16, 2013, 40, Suppl. 2, S193.

Rinnab, L. et al., "Evaluation of [11C]-choline positron-emission/computed tomography in patients with increasing prostate-specific antigen levels after primary treatment for prostate cancer," *BJU Int*, 2007, 100, 786,793.

Reske, S.N. et al., "Imaging Prostate Cancer with 11C-Choline PET/CT," *J. Nucl. Med.*, 2006, 47, 1249-1254.

Zophel, K. et al., "Is 11C-choline the most appropriate tracer for KP prostate cancer?" *Eur J Nucl Med Mo/ Imaging*, 2004, 31, 756-759. Examination Report for counterpart Australian Patent Application No. 2018200419, dated Oct. 24, 2018.

Office Action for counterpart Japanese Patent Application No. 2017-210775, dated Nov. 6, 2018, with English Translation.

Supplementary Partial European Search Report for counterpart European Application No. 14861854.9, dated May 19, 2017.

Zophel et al., "Is 11C-choline the most appropriate tracer for prostate cancer? Against." *Eur J Nucl Med Mol Imaging*, 2004, 31: 756-759.

First Examination Report for counterpart Saudi Arabian Application No. 516370842, with English translation.

MB17

PET-imaging

Time-activity-curves




FIG. 1

Organ distribution at 1 h post injection

	MB17			MB17 blocked		
	%ID/g	SD	n	%ID/g	SD	n
Blood	1.08	0.25	3	1.24	0.92	3
Heart	0.53	0.15	3	0.49	0.27	3
Lung	1.41	0.41	3	1.10	0.53	3
Spleen	2.13	0.16	3	0.53	0.36	3
Liver	1.17	0.10	3	0.73	0.07	3
Kidney	113.34	24.45	3	2.38	1.40	3
Muscle	0.50	0.07	3	0.24	0.09	3
Intestine	0.46	0.13	3	0.35	0.23	3
Brain	0.12	0.01	3	0.07	0.01	3
LNCaP Tumor	8.47	4.09	3	0.98	0.32	3




FIG. 2






FIG. 3



24 h	%ID/g	SD	n
Blood	0,01	0,00	5
Heart	0,02	0,01	5
Lung	0,11	0,13	5
Spleen	0,13	0,05	5
Liver	0,08	0,03	5
Kidney	2,13	1,36	5
Muscle	0,02	0,01	5
Intestine	0,02	0,01	5
Brain	0,05	0,03	5
LNCaP Tumor	10,58	4,50	5

FIG. 4

**FIG. 5**




FIG. 6




FIG. 7




FIG. 8




FIG. 9




FIG. 10




FIG. 11




FIG. 12




FIG. 13




FIG. 14




FIG. 15

MB-17 vs MB-17.D

PET-imaging after 2 hours post injection, comparison of whole-body scan




FIG. 16





FIG. 17A



PSA level 0.7

FIG. 17B




FIG. 18A




FIG. 18B

1

**LABELLED INHIBITORS OF PROSTATE
SPECIFIC MEMBRANE ANTIGEN (PSMA),
THEIR USE AS IMAGING AGENTS AND
PHARMACEUTICAL AGENTS FOR THE
TREATMENT OF PROSTATE CANCER**

**RELATED APPLICATIONS AND
INCORPORATION BY REFERENCE**

This application is a continuation of U.S. application Ser. No. 15/131,118 filed Apr. 18, 2016, which is a continuation-in-part application of international patent application Serial No. PCT/EP2014/002808 filed Oct. 17, 2014, which published as PCT Publication No. WO 2015/055318 on Apr. 23, 2015, which claims benefit of European patent application Serial Nos. 13004991.9 filed Oct. 18, 2013 and 14175612.2 filed Jul. 3, 2014.

The foregoing applications, and all documents cited therein are incorporated herein by reference in their entireties.

FIELD OF THE INVENTION

The present invention generally relates to the field of radiopharmaceuticals and their use in nuclear medicine as tracers, imaging agents and for the treatment of various disease states of prostate cancer.

BACKGROUND OF THE INVENTION

Prostate cancer (PCa) is the leading cancer in the US and European population. At least 1-2 million men in the western hemisphere suffer from prostate cancer and it is estimated that the disease will strike one in six men between the ages of 55 and 85. There are more than 300,000 new cases of prostate cancer diagnosed each year in USA. The mortality from the disease is second only to lung cancer. Currently anatomic methods, such as computed tomography (CT), magnetic resonance (MR) imaging and ultrasound, predominate for clinical imaging of prostate cancer. An estimated \$2 billion is currently spent worldwide on surgical, radiation, drug therapy and minimally invasive treatments. However, there is presently no effective therapy for relapsing, metastatic, androgen-independent prostate cancer.

A variety of experimental low molecular weight PCa imaging agents are currently being pursued clinically, including radiolabeled choline analogs [^{18}F]fluorodihydrotestosterone ([^{18}F]FDHT), anti-1-amino-3-[^{18}F]fluorocyclobutyl-1-carboxylic acid (anti [^{18}F]F-FACBC, [^{11}C]acetate and 1-(2-deoxy-2-[^{18}F]fluoro-L-arabino-furanosyl)-5-methyluracil ([^{18}F]FMAU) (Scher, B.; et al. *Eur J Nucl Med Mol Imaging* 2007, 34, 45-53; Rinnab, L.; et al. *BJU Int* 2007, 100, 786, 793; Reske, S. N.; et al. *J Nucl Med* 2006, 47, 1249-1254; Zophel, K.; Kotzerke, J. *Eur J Nucl Med Mol Imaging* 2004, 31, 756-759; Veas, H.; et al. *BJU Int* 2007, 99, 1415-1420; Larson, S. M.; et al. *J Nucl Med* 2004, 45, 366-373; Schuster, D. M.; et al. *J Nucl Med* 2007, 48, 56-63; Tehrani, O. S.; et al. *J Nucl Med* 2007, 48, 1436-1441). Each operates by a different mechanism and has certain advantages, e.g., low urinary excretion for [^{11}C]choline, and disadvantages, such as the short physical half-life of positron-emitting radionuclides.

It is well known that tumors may express unique proteins associated with their malignant phenotype or may over-express normal constituent proteins in greater number than normal cells. The expression of distinct proteins on the surface of tumor cells offers the opportunity to diagnose and

2

characterize disease by probing the phenotypic identity and biochemical composition and activity of the tumor. Radioactive molecules that selectively bind to specific tumor cell surface proteins provide an attractive route for imaging and treating tumors under non-invasive conditions. A promising new series of low molecular weight imaging agents targets the prostate-specific membrane antigen (PSMA) (Mease R. C. et al. *Clin Cancer Res* 2008, 14, 3036-3043; Foss, C. A.; et al. *Clin Cancer Res* 2005, 11, 4022-4028; Pomper, M. G.; et al. *Mol Imaging* 2002, 1, 96-101; Zhou, J.; et al. *Nat Rev Drug Discov* 2005, 4, 1015-1026; WO 2013/022797).

PSMA is a trans-membrane, 750 amino acid type II glycoprotein that has abundant and restricted expression on the surface of PCa, particularly in androgen-independent, advanced and metastatic disease (Schulke, N.; et al. *Proc Natl Acad Sci USA* 2003, 100, 12590-12595). The latter is important since almost all PCa become androgen independent over the time. PSMA possesses the criteria of a promising target for therapy, i.e., abundant and restricted (to prostate) expression at all stages of the disease, presentation at the cell surface but not shed into the circulation and association with enzymatic or signaling activity (Schulke, N.; et al. *Proc. Natl. Acad. Sci. USA* 2003, 100, 12590-12595). The PSMA gene is located on the short arm of chromosome 11 and functions both as a folate hydrolase and neuropeptidase. It has neuropeptidase function that is equivalent to glutamate carboxypeptidase II (GCPII), which is referred to as the "brain PSMA", and may modulate glutamatergic transmission by cleaving N-acetylaspartylglutamate (NAAG) to N-acetylaspartate (NAA) and glutamate (Nan, F.; et al. *J Med Chem* 2000, 43, 772-774). There are up to 10^6 PSMA molecules per cancer cell, further suggesting it as an ideal target for imaging and therapy with radionuclide-based techniques (Tasch, J.; et al. *Crit Rev Immunol* 2001, 21, 249-261).

The radio-immunoconjugate of the anti-PSMA monoclonal antibody (mAb) 7E11, known as the PROSTASCINT® scan, is currently being used to diagnose prostate cancer metastasis and recurrence. However, this agent tends to produce images that are challenging to interpret (Lange, P. H. PROSTASCINT scan for staging prostate cancer. *Urology* 2001, 57, 402-406; Haseman, M. K.; et al. *Cancer Biother Radiopharm* 2000, 15, 131-140; Rosenthal, S. A.; et al. *Tech Urol* 2001, 7, 27-37). More recently, monoclonal antibodies have been developed that bind to the extracellular domain of PSMA and have been radiolabeled and shown to accumulate in PSMA-positive prostate tumor models in animals. However, diagnosis and tumor detection using monoclonal antibodies has been limited by the low permeability of the monoclonal antibody in solid tumors.

The selective targeting of cancer cells with radiopharmaceuticals, either for imaging or therapeutic purposes is challenging. A variety of radionuclides are known to be useful for radio-imaging or cancer radiotherapy, including ^{111}In , ^{90}Y , ^{68}Ga , ^{177}Lu , $^{99\text{m}}\text{Tc}$, ^{123}I and ^{131}I . Recently it has been shown that some compounds containing a glutamate-urea-glutamate (GUG) or a glutamate-urea-lysine (GUL) recognition element linked to a radionuclide-ligand conjugate exhibit high affinity for PSMA.

Citation or identification of any document in this application is not an admission that such document is available as prior art to the present invention.

SUMMARY OF THE INVENTION

New agents that will enable rapid visualization of prostate cancer and specific targeting to allow radiotherapy present are needed.

Thus, the object of the present invention is to develop ligands that interact with PSMA and carry appropriate radionuclides which provide a promising and novel targeting option for the detection, treatment and management of prostate cancer.

The solution of said object is achieved by providing the embodiments characterized in the claims.

The inventors found new compounds which are useful radiopharmaceuticals and their use in nuclear medicine as tracers, imaging agents and for the treatment of various disease states of prostate cancer.

The novel imaging agents with structural modifications in the linker region have improved tumor targeting properties and pharmacokinetics. The pharmacophore presents three carboxylic groups able to interact with the respective side chains of PSMA and an oxygen as part of zinc complexation in the active center. Besides these obligatory interactions, the inventors were able to optimize the lipophilic interactions in the linker region.

It is noted that in this disclosure and particularly in the claims and/or paragraphs, terms such as "comprises", "comprising", "comprising" and the like can have the meaning attributed to it in U.S. Patent law; e.g., they can mean "includes", "included", "including", and the like; and that terms such as "consisting essentially of" and "consists essentially of" have the meaning ascribed to them in U.S. Patent law, e.g., they allow for elements not explicitly recited, but exclude elements that are found in the prior art or that affect a basic or novel characteristic of the invention.

These and other embodiments are disclosed or are obvious from and encompassed by, the following Detailed Description.

BRIEF DESCRIPTION OF THE DRAWINGS

The following detailed description, given by way of example, but not intended to limit the invention solely to the specific embodiments described, may best be understood in conjunction with the accompanying drawings.

FIG. 1: PET—Imaging of MB17. Whole-body coronal microPET images of an athymic male nude mice bearing LNCaP tumor xenografts. The tumor-targeting efficacy and pharmacokinetic properties of [^{68}Ga]MB17 were evaluated by dynamic microPET scans. Approximately 15 MBq/mouse were injected. Graph A shows the respective time-activity-curves of kidney and bladder and graph B the respective time-activity-curves of heart, muscle and tumor. The values are expressed as mean SUV (standardized uptake values).

FIG. 2: Organ Distribution at 1 h post injection. Organ distribution at one hour post injection of 0.06 nmol of the ^{68}Ga labeled PSMA inhibitor MB17. PSMA-blocking by co-administration of 2 mg/kg body weight 2-PMPA indicates PSMA-specific uptake in the tumor and the kidneys. Data are expressed as mean % ID/g tissue $\pm$ SD (n=3).

FIG. 3: PET—Imaging of MB4. Whole-body coronal microPET images of an athymic male nude mice bearing LNCaP tumor xenografts. The tumor-targeting efficacy and pharmacokinetic properties of [^{68}Ga]MB4 were evaluated by dynamic microPET scans. Approximately 15 MBq/mouse were injected. Graph A shows the respective time-activity-curves of kidney and bladder and graph B the respective time-activity-curves of heart, muscle and tumor. The values are expressed as mean SUV (standardized uptake values)

FIG. 4: Organ distribution expressed as % ID/g tissue $\pm$ SD (n=5) 24 h post injection of 0.06 nmol of the ^{177}Lu -labeled

MB17. Organ distribution with ^{177}Lu shows that the high initial kidney uptake is nearly completely washed out ($2.13\pm 1.36\%$ ID/g) after 24 hours while the tumor uptake remained high and even increased ($10.58\pm 4.50\%$ ID/g).

5 Other organs as liver ($0.08\pm 0.03\%$ ID/g), lung ($0.11\pm 0.13\%$ ID/g) and spleen ($0.13\pm 0.05\%$ ID/g) showed very low uptake. The favourable pharmacokinetics led to extremely high tumor-to-background ratios (Tumor/Blood: 1058; Tumor/Muscle: 529) after 24 hours

10 FIG. 5: PET—Imaging of MB 2. Whole-body coronal microPET images of an athymic male nude mouse bearing LNCaP tumor xenografts. The tumor-targeting efficacy and pharmacokinetic properties of [^{68}Ga]MB2 were evaluated by dynamic microPET scans. Approximately 15 MBq/mouse were injected.

15 FIG. 6: PET—Imaging of MB 3. Whole-body coronal microPET images of an athymic male nude mouse bearing LNCaP tumor xenografts. The tumor-targeting efficacy and pharmacokinetic properties of [^{68}Ga]MB 3 were evaluated by dynamic microPET scans. Approximately 15 MBq/mouse were injected.

20 FIG. 7: PET—Imaging of MB10. Whole-body coronal microPET images of an athymic male nude mouse bearing LNCaP tumor xenografts. The tumor-targeting efficacy and pharmacokinetic properties of [^{68}Ga]MB10 were evaluated by dynamic microPET scans. Approximately 15 MBq/mouse were injected.

25 FIG. 8: PET—Imaging of MB17.D. Whole-body coronal microPET images of an athymic male nude mouse bearing LNCaP tumor xenografts. The tumor-targeting efficacy and pharmacokinetic properties of [^{68}Ga]MB17.D were evaluated by dynamic microPET scans. Approximately 15 MBq/mouse were injected. MB17D: stereoisomer of MB17(L); synthesis based on Fmoc-3(2-naphthyl)-D-alanine

30 FIG. 9: PET—Imaging of MB22. Whole-body coronal microPET images of an athymic male nude mouse bearing LNCaP tumor xenografts. The tumor-targeting efficacy and pharmacokinetic properties of [^{68}Ga]MB22 were evaluated by dynamic microPET scans. Approximately 15 MBq/mouse were injected.

35 FIG. 10: PET—Imaging of MB 24. Whole-body coronal microPET images of an athymic male nude mouse bearing LNCaP tumor xenografts. The tumor-targeting efficacy and pharmacokinetic properties of [^{68}Ga]MB 24 were evaluated by dynamic microPET scans. Approximately 15 MBq/mouse were injected.

40 FIG. 11: PET—Imaging of MB25. Whole-body coronal microPET images of an athymic male nude mouse bearing LNCaP tumor xenografts. The tumor-targeting efficacy and pharmacokinetic properties of [^{68}Ga]MB25 were evaluated by dynamic microPET scans. Approximately 15 MBq/mouse were injected.

45 FIG. 12: PET—Imaging of MB31. Whole-body coronal microPET images of an athymic male nude mouse bearing LNCaP tumor xenografts. The tumor-targeting efficacy and pharmacokinetic properties of [^{68}Ga]MB31 were evaluated by dynamic microPET scans. Approximately 15 MBq/mouse were injected.

50 FIG. 13: PET—Imaging of MB33. Whole-body coronal microPET images of an athymic male nude mouse bearing LNCaP tumor xenografts. The tumor-targeting efficacy and pharmacokinetic properties of [^{68}Ga]MB33 were evaluated by dynamic microPET scans. Approximately 15 MBq/mouse were injected.

55 FIG. 14: PET—Imaging of MB35. Whole-body coronal microPET images of an athymic male nude mouse bearing LNCaP tumor xenografts. The tumor-targeting efficacy and

5

pharmacokinetic properties of [^{68}Ga]MB35 were evaluated by dynamic microPET scans. Approximately 15 MBq/mouse were injected.

FIG. 15: PET scan of a mouse injected with ^{68}Ga -CHX-DTPA. On the left the caudal, in the centre the dorsal and on the right the lateral view. The pictures cover the time spans of 20-40 min (top), 40-60 min (centre) and 120-140 min (bottom).

FIG. 16: MB-17 vs MB-17.D. Whole-body coronal microPET images of athymic male nude mice bearing LNCaP tumor xenografts. The tumor-targeting efficacy and pharmacokinetic properties of the stereoisomers MB-17 and MB-17 D were directly compared at 2 hours post injection.

FIGS. 17A-B: Human PET/CT imaging ^{68}Ga -labeled MB17. (a) First clinical experience with ^{68}Ga -labeled MB17 PET/CT demonstrates the detection of small lymph node metastases 1 hour post injection, primarily due to a high radiotracer uptake. Red arrows point to a representative lesion with a SUVmax of 36.5 and a tumor-to-background ratio of 52.1 one hour post injection. MIP=maximum intensity projection of the PET 1 h post injection. (b) The significant advantage of ^{68}Ga -labeled MB17 PET/CT is the sensitive detection of lesions even at low PSA level.

FIGS. 18A-B: PET imaging of patient with multiple prostate cancer metastasis. (a) First scan demonstrate initial PET imaging of the patient with multiple prostate cancer metastases with blood PSA value of 14. Two months later


6

DETAILED DESCRIPTION OF THE INVENTION


The present invention relates to radiopharmaceuticals and their use in nuclear medicine as tracers, imaging agents and for the treatment of various disease states of prostate cancer.

Thus, the present invention concerns compounds that are represented by the general Formulae (Ia) or (Ib):

Formula (Ia)



Formula (Ib)



with:

n:	0, 1
m:	1, 2, 3, 4
Z:	—CO ₂ H, —SO ₂ H, —SO ₃ H, —SO ₄ H, —PO ₂ H, —PO ₃ H, —PO ₄ H ₂
X:	Naphthyl, Phenyl, Biphenyl, Indolyl (=2,3-benzopyrrolyl), Benzothiazolyl
Y:	Aryl, Alkylaryl, Cyclopentyl, Cyclohexyl, Cycloheptyl
Chelator radical of:	1,4,7,10-tetraazacyclododecane-N,N',N'',N'''-tetraacetic acid (=DOTA), N,N'-bis[2-hydroxy-5-(carboxyethyl)benzyl]ethylenediamine-N,N'-diacetic acid (=HBED-CC), 1,4,7-triazacyclononane-1,4,7-triacetic acid (=NOTA), 2-(4,7-bis(carboxymethyl)-1,4,7-triazonan-1-yl)pentanedioic acid (NODAGA), 2-(4,7,10-tris(carboxymethyl)-1,4,7,10-tetraazacyclododecan-1-yl)pentanedioic acid (DOTAGA), 1,4,7-triazacyclononane phosphinic acid (TRAP), 1,4,7-triazacyclononane-1-[methyl(2-carboxyethyl)phosphinic acid]-4,7-bis[methyl(2-hydroxymethyl)phosphinic acid] (NOPO), 3,6,9,15-tetraazabicyclo[9.3.1][.]]pentadeca-1(15),11,13-triene-3,6,9-triacetic acid (=PCTA), N'-[5-[Acetyl(hydroxyamino)pentyl]-N-[5-[(5-aminopentyl)(hydroxyamino)-4-oxobutanoyl]amino]pentyl]-N-hydroxysuccinamide (DFO), Diethylenetriaminepentaacetic acid (DTPA) Trans-cyclohexyl-diethylenetriaminepentaacetic acid (CHX-DTPA) 1-oxa-4,7,10-triazacyclododecane-4,7,10-triacetic acid (oxo-Do3A) p-isothiocyanatobenzyl-DTPA (SCN-Bz-DTPA) 1-(p-isothiocyanatobenzyl)-3-methyl-DTPA (1B3M) 2-(p-isothiocyanatobenzyl)-4-methyl-DTPA (1M3B) 1-(2-methyl-4-isocyanatobenzyl)-DTPA (MX-DTPA)

3.3 GBq of ^{177}Lu -labeled MB17 was applied. At this time point, the amount of PSA in blood reached a value of 38. After the first cycle, the PSA level decreased to 8. Three months after the first cycle another 4 GBq of ^{177}Lu -labeled MB17 was applied. The control PET scan was performed one month after the second cycle. The treatment has shown a significant impact on the tumor lesions and PSA value and resulted in a reduction of bone pain. (b) The graph demonstrates the significant impact on the PSA value which decreased after the first application of the therapeutic dose of ^{177}Lu -labeled MB17.

If not stated otherwise, in the present invention the "alkyl" residue (preferably: C₁ to C₁₀) can be linear or branched, unsubstituted or substituted. Preferred alkyl residues are methyl, ethyl, n-propyl, iso-propyl, n-butyl, tert-butyl, n-pentanyl, n-hexanyl. The same also applies to the corresponding cycloalkyl compounds having preferably 3 to 10 carbon atoms.

"Aryl" refers to an aromatic monocyclic or polycyclic ring system having 6 to 14 carbon atoms, preferably 6 to 10 carbon atoms. The aryl group can be substituted, where


Fusion Pharmaceuticals, Inc.

Exhibit 1025.037


Although it is preferred that the Z-Group is $\text{—CO}_2\text{H}$ it may be easily replaced with biosteric replacements such as $\text{—SO}_2\text{H}$, $\text{—SO}_3\text{H}$, $\text{—SO}_4\text{H}$, $\text{—PO}_2\text{H}$, $\text{—PO}_3\text{H}$, $\text{—PO}_4\text{H}_2$, see e.g. “The Practice of Medicinal Chemistry” (Academic Press New York, 1996), page 203.

In a preferred embodiment, the motif specifically binding to cell membranes of neoplastic cells is a motif specifically binding to cell membranes of cancerous cells, preferably wherein said motif may comprise a prostate-specific membrane antigen (PSMA), in particular wherein said PSMA may comprise a glutamate-urea-lysine motif according to the following formula in Scheme 1.

Scheme 1: Structure of Preferred Compounds of the Present Invention




The different preferred linkers are shown below, wherein R=radical of Glu-urea-Lys and R'=radical of DOTA (as a preferred example for the chelator), as shown above.




Molecular Weight: 972.03 g/mol

MB3



Molecular Weight: 1105.18 g/mol

MB4





Molecular Weight: 1238.33 g/mol

MB10


9

-continued




Linker

Chemical Formula: $C_{49}H_{71}N_9O_{16}$
 Molecular Weight: 1042.16 g/mol



Linker

Chemical Formula: $C_{36}H_{60}N_8O_{15}$
 Molecular Weight: 844.92 g/mol

**10**

-continued

MB17

MB31

5

10

15

20

25

MB22

30

35

MB24

40

45




50

MB25

55

60

65




11


Preferred compounds of the present invention are e.g.

12


MB17




or




MB 4




MB 10




-continued




MB 22



MB24



MB25



Fusion Pharmaceuticals, Inc.
Exhibit 1025.043

saccharides, formic acids or glucuronic acid, etc.). Such a prodrug of a compound according to the invention can be applied to a patient, and this prodrug can be transformed into a substance of general formula (Ia) and/or (Ib) so as to obtain the desired pharmacological effect.

Some compounds of Formula (Ia) and/or (Ib) are encompassed in form of the racemates, their enantiomers and optionally in form of their diastereomers and all possible mixtures thereof.

According to the invention all chiral C-atoms shall have D- and/or L-configuration; also combinations within one compound shall be possible, i.e. some of the chiral C-atoms may be D- and others may be L-configuration.

The obtained compounds can be optionally separated by known methods (e.g. Allinger, N. L. and Elliel E. L. in, "Topics in Stereochemistry" Vol. 6, Wiley Interscience, 1971) in their enantiomers and/or diastereomers. One possible method of enantiomeric separation is the use of chromatography.

The invention also relates to pharmaceutical preparations which contain a therapeutically effective amount of the active ingredients (compound according to the invention of formula (Ia) or (Ib) together with organic or inorganic solid or liquid, pharmaceutically acceptable carriers which are suited for the intended administration and which interact with the active ingredients without drawbacks.

The phrase "pharmaceutically acceptable" is employed herein to refer to those compounds, material, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of a patient without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio.

A "patient" includes an animal, such as a human, monkey, cow, horse, cat or dog. The animal can be a mammal such as a non-primate and a primate (e.g., monkey and human). In one embodiment, a patient is a human being.

In general, the Formula (Ia) or (Ib) compound or pharmaceutical compositions thereof, may be administered orally or via a parenteral route, usually injection or infusion.

A "parenteral administration route" means modes of administration other than enteral and topical administration, usually by injection, and includes, without limitation, intravenous, intramuscular, intraarterial, intrathecal, intracapsular, intraorbital, intracardiac, intradermal, intraperitoneal, transtracheal, subcutaneous, subcuticular, intraarticular, subcapsular, subarachnoid, intraspinal and intrasternal injection and infusion.

The dosage of the compounds according to the invention is determined by the physician on the basis of the patient-specific parameters, such as age, weight, sex, severity of the disease, etc. The dosage is preferably from 0.00001 mg/kg to 100 mg/kg body weight, preferably from 0.001 to 50 mg/kg body weight and most preferably from 0.01 to 10 mg/kg body weight.

Corresponding to the kind of administration, the medication is suitably formulated, e.g. in the form of solutions or suspensions, simple tablets or dragees, hard or soft gelatine capsules, suppositories, ovules, preparations for injection, which are prepared according to common galenic methods.

The compounds according to the invention can be formulated, where appropriate, together with further active substances and with excipients and carriers common in pharmaceutical compositions, e.g.—depending on the preparation to be produced—talcum, gum arabic, lactose, starch, magnesium stearate, cocoa butter, aqueous and non-aqueous carriers, fatty bodies of animal or vegetable origin,

paraffin derivatives, glycols (in particular polyethylene glycol), various plasticizers, dispersants or emulsifiers, pharmaceutically compatible gases (e.g. air, oxygen, carbon dioxide, etc.), preservatives.

In order to produce liquid preparations, additives, such as sodium chloride solution, ethanol, sorbitol, glycerine, olive oil, almond oil, propylene glycol or ethylene glycol, can be used.

When solutions for infusion or injection are used, they are preferably aqueous solutions or suspensions, it being possible to produce them prior to use, e.g. from lyophilized preparations which contain the active substance as such or together with a carrier, such as mannitol, lactose, glucose, albumin and the like. The ready made solutions are sterilized and, where appropriate, mixed with excipients, e.g. with preservatives, stabilizers, emulsifiers, solubilizers, buffers and/or salts for regulating the osmotic pressure. The sterilization can be obtained by sterile filtration using filters having a small pore size according to which the composition can be lyophilized, where appropriate. Small amounts of antibiotics can also be added to ensure the maintenance of sterility.

The phrases "effective amount" or "therapeutically-effective amount" as used herein means that amount of a compound, material, or composition which may comprise a compound of the invention, or other active ingredient which is effective for producing some desired therapeutic effect in at least a sub-population of cells in an animal at a reasonable benefit/risk ratio applicable to any medical treatment. A therapeutically effective amount with respect to a compound of the invention means that amount of therapeutic agent alone, or in combination with other therapies, that provides a therapeutic benefit in the treatment of prevention of a disease. Used in connection with a compound of the invention, the term can encompass an amount that improves overall therapy, reduces or avoids symptoms or causes of disease, or enhances the therapeutic efficacy of or synergies with another therapeutic agent.

As used herein, the terms "treating" or "treatment" is intended to encompass also diagnosis, prophylaxis, prevention, therapy and cure.

The terms "prevent", "preventing," and "prevention" refer to the prevention of the onset, recurrence, or spread of the disease in a patient resulting from the administration of a prophylactic or therapeutic agent.

Depending on whether the inventive Formula (Ia) and/or (Ib) compounds are to be used as radio-imaging agents or radio-pharmaceuticals different radionuclides are complexed to the chelator. Illustrative radionuclides include, for example, ^{89}Zr , ^{44}Sc , ^{111}In , ^{90}Y , ^{66}Ga , ^{67}Ga , ^{177}Lu , $^{99\text{m}}\text{Tc}$, ^{61}Cu , ^{62}Cu , ^{64}Cu , ^{67}Cu , ^{149}Tb , ^{152}Tb , ^{155}Tb , ^{161}Tb , ^{153}Gd , ^{155}Gd , ^{157}Gd , ^{213}Bi , ^{225}Ac , ^{230}U , ^{223}Ra , ^{165}Er and Fe. According to one aspect of this invention, the radionuclide is ^{111}In , ^{90}Y , ^{68}Ga , ^{64}Cu , ^{153}Gd , ^{155}Gd , ^{213}Bi , ^{225}Ac , Fe, or ^{177}Lu .

As noted above, complexes of the compounds according to Formula (Ia) or (Ib) may contain one or more radionuclides which are suitable for use as radio-imaging agents or as therapeutics for the treatment of rapidly proliferating cells, for example, PSMA expressing prostate cancer cells. According to the present invention they are called "metal complexes" or "radiopharmaceuticals".

Preferred imaging methods are positron emission tomography (PET) or single photon emission computed tomography (SPECT).

Accordingly, in one embodiment, a pharmaceutical composition is provided including a complex that includes a

21

radionuclide and a compound of Formula (Ia) or Formula (Ib), a salt, solvate, stereoisomer, or tautomer thereof, and a pharmaceutically acceptable carrier.

According to another aspect, a pharmaceutical composition is provided, which is suitable for in vivo imaging and radiotherapy. Suitable pharmaceutical compositions may contain a radio imaging agent, or a radiotherapeutic agent that has a radionuclide either as an element, i.e. radioactive iodine, or a radioactive metal chelate complex of the compound of Formula (Ia) and/or (Ib) in an amount sufficient for imaging, together with a pharmaceutically acceptable radiological vehicle. The radiological vehicle should be suitable for injection or aspiration, such as human serum albumin; aqueous buffer solutions, e.g., tris(hydromethyl) aminomethane (and its salts), phosphate, citrate, bicarbonate, etc; sterile water physiological saline; and balanced ionic solutions containing chloride and or dicarbonate salts or normal blood plasma cautions such as calcium potassium, sodium and magnesium.

The concentration of the imaging agent or the therapeutic agent in the radiological vehicle should be sufficient to provide satisfactory imaging. For example, when using an aqueous solution, the dosage is about 1.0 to 100 millicuries. The actual dose administered to a patient for imaging or therapeutic purposes, however, is determined by the physician administering treatment. The imaging agent or therapeutic agent should be administered so as to remain in the patient for about 1 hour to 10 days, although both longer and shorter time periods are acceptable. Therefore, convenient ampoules containing 1 to 10 mL of aqueous solution may be prepared.

Imaging may be carried out in the normal manner, for example by injecting a sufficient amount of the imaging composition to provide adequate imaging and then scanning with a suitable imaging or scanning machine, such as a tomograph or gamma camera. In certain embodiments, a method of imaging a region in a patient includes the steps of: (i) administering to a patient a diagnostically effective amount of a compound complexed with a radionuclide;

22

exposing a region of the patient to the scanning device; and (ii) obtaining an image of the region of the patient. In certain embodiments of the region imaged is the head or thorax. In other embodiments, the compounds and complexes of Formula I(a) and/or (Ib) target the PSMA protein.


Thus, in some embodiments, a method of imaging tissue such as spleen tissue, kidney tissue, or PSMA-expressing tumor tissue is provided including contacting the tissue with a complex synthesized by contacting a radionuclide and a Formula (Ia) and/or Formula (Ib) compound.

The amount of the compound of the present invention, or a formulation which may comprise a complex of a metal and a compound according to Formula (Ia) and/or (Ib), or its salt, solvate, stereoisomer, or tautomer that is administered to a patient depends on several physiological factors that are routinely used by the physician, including the nature of imaging to be carried out, tissue to be targeted for imaging or therapy and the body weight and medical history of the patient to be imaged or treated using a radiopharmaceutical.

Accordingly in another aspect, the invention provides a method for treating a patient by administering to a patient a therapeutically effective amount of a Formula (Ia) and/or (Ib) compound complexed to a radionuclide, or a pharmaceutically acceptable salt or solvate of the complex to treat a patient suffering from a cell proliferative disease or disorder. Specifically, the cell proliferative disease or disorder to be treated or imaged using a compound, pharmaceutical composition or radiopharmaceutical in accordance with this invention is a cancer, for example, prostate cancer and/or prostate cancer metastasis in e.g. lung, liver, kidney, bones, brain, spinal cord, bladder, etc.

The synthesis of the compounds of the present invention is described in detail in the example section. An overview of the synthesis is exemplified in Scheme 2 concerning DOTA conjugated-PSMA inhibitors. However, a person skilled in the art would be able to modify the reactions e.g. by using another chelator. Thus, this scheme shall not be understood to limit the compounds of the present invention to the DOTA chelator only.

Scheme 2



25

of tris(t-bu)-DOTA (Chematech) relative to the resin loading were reacted in a final volume of 3 mL DMF. The product was cleaved from the resin in a 2 mL mixture consisting of trifluoroacetic acid, triisopropylsilane, and water (95:2.5:2.5).

The chelator was also conjugated by using HBTU activated DOTA-NHS ester (CheMatech) or DOTA-TFP ester (Mier W., Hoffend J., Krämer S., Schuhmacher J., Hull W. E., Eisenhut M., Haberkorn U., *Bioconjugate Chem.* 2005, 16: 237-240).

Analysis of the synthesized molecules was performed using reversed-phase high performance liquid chromatography (RP-HPLC; Chromolith RP-18e, 100×4.6 mm; Merck, Darmstadt, Germany) with a linear A-B gradient (0% B to 100% B in 6 min) at a flow rate of 4 mL/min (analysis) or 6 mL/min (purification). Solvent A consisted of 0.1% aqueous TFA and solvent B was 0.1% TFA in CH₃CN. The HPLC system (L6200 A; Merck-Hitachi, Darmstadt, Germany) was equipped with a UV and a gamma detector (Bioscan; Washington, USA). UV absorbance was measured at 214 nm. Mass spectrometry was performed with a MALDI-MS Daltonics Microflex system (Bruker Daltonics, Bremen, Germany).

Example 2

Radiolabeling

Typically, 1.5 nmol of a synthesized compound of Example 1 (dissolved in 0.1 M HEPES buffer pH 7.5) was added in a volume of 100 μ L to a mixture of 10 μ L 2.1 M HEPES solution and 40 μ L [⁶⁸Ga]Ga³⁺ eluate (40 MBq). The pH of the labeling solution was adjusted to 4.5.

The radiolabeling of the compounds resulted in a radiochemical yield of >97% after 15 minutes at 95° C. and was determined by RP-HPLC and TLC. Subsequent purification was done using Sep-Pak C18 cartridges.

Example 3

Synthesis of Compounds MB4 and MB17

The isocyanate of the glutamyl moiety was generated *in situ* by adding a mixture of 3 mmol of bis(tert-butyl) L-glutamate hydrochloride and 1.5 mL of N-ethyldiisopropylamine (DIPEA) in 200 mL of dry CH₂Cl₂ to a solution of 1 mmol triphosgene in 10 mL of dry CH₂Cl₂ at 0° C. over 4 h. After agitation of the reaction mixture for 1 h at 25° C., 0.5 mmol of the resin-immobilized (2-chloro-tritylresin) ϵ -allyloxycarbonyl protected lysine in 4 mL DCM was added and reacted for 16 h with gentle agitation. The resin was filtered off and the allyloxy-protecting group was removed using 30 mg tetrakis(triphenyl)palladium(0) and 400 μ L morpholine in 4 mL CH₂Cl₂ for 3 hours. The following coupling of 3 times 4-(Fmoc-aminomethyl)benzoic acid (in case of MB4) or Fmoc-3-(2-naphthyl)-L-alanine and trans-4-(Fmoc-aminomethyl)cyclohexanecarboxylic acid (in case of MB17), respectively, was performed stepwise using 2 mmol of the Fmoc-protected acid, 1.96 mmol of HBTU and 2 mmol of N-ethyldiisopropylamine in a final volume of 4 mL DMF. After activation with 3.95 eq of HBTU and DIPEA for 2 h, 4 eq of tris(t-bu)-DOTA (Chematech) relative to the resin loading were reacted for 3 h in a final volume of 3 mL DMF. The product was cleaved from the resin in a 2 mL mixture consisting of trifluoroacetic acid, triisopropylsilane, and water (95:2.5:2.5). Purification


26

was performed using RP-HPLC and the purified product was analysed by analytical RP-HPLC and MALDI-MS.

For preparing MB-17D which is the stereoisomer of MB17(L), the synthesis was based on Fmoc-3(2-naphthyl)-D-alanine. If not stated otherwise, in the present description MB17 means the L-stereoisomer.

Example 4

Coupling to Various Chelators



The chelators (DOTA, NOTA, NODAGA, DTPA, CHX-DTPA, PCTA, Do3A) were coupled to the MB17 linker by solid phase synthesis. In general, 13 μ mol of resin which was coupled with the PSMA binding motif was swollen with DCM in a syringe with a filter. After washing the resin 5× with DMF, it was incubated 2× for 5 min with 20% of piperidine in DMF to deprotect the N-terminus. Another 5× washing with DMF followed.

Between 1.5 and 4 equivalents of the chelator (depending of the chelator), 0.98×n_{chelator} HATU (if needed) and 10 equivalents of DIPEA were dissolved in 500 μ L of DMF, the solution was drawn up into the syringe containing the resin and incubated overnight. Next, the resin was washed 5× each with DMF, methanol, DCM and diethyl ether and dried over vacuum.

To check the state of the reaction, test separations were used. This was achieved by washing a small amount of resin with DCM into a filter tip and adding 100 μ L of separation solution containing 95% TFA, 2.5% water and 2.5% TIPS. After 30 min of incubation, the solution was pipetted into ice cold diethyl ether and centrifuged. The diethyl ether was decanted and the remaining pellet was dissolved in 35 μ L of ACN:H₂O (1:1) and analysed by HPLC (0-100% ACN in water within 5 min) and LC/MS.

If the desired product was obtained, the complete peptide was separated from the resin. The dried resin was incubated with 500 μ L of the separation solution (95% TFA, 2.5% H₂O, 2.5% TIPS) for 2 hours. The resulting solution was mixed with ice cold diethyl ether and centrifuged (4000 min⁻¹, 5 min). The supernatant was discarded, new diethyl ether was added and the receptacle was shaken vigorously to resuspend the pellet. Again, the solution was centrifuged (4000 min⁻¹, 5 min) and the resulting supernatant discarded. The pellet was then vacuum dried and finally resuspended in 1 mL of ACN:H₂O (1:1).

Purification was achieved by preparative HPLC, the peaks were analysed by analytic HPLC (0-100% ACN in water within 5 min) and LC/MS and those containing the product were pooled and lyophilized.

Example 5

Radiolabelling

¹⁷⁷Lu-labelling

¹⁷⁷Lu (approx. 100 MBq) was mixed with 200 µl of 0.4 M sodium acetate buffer containing Chelex (pH=5). 10 µl of a 1 mM solution of the compound in 10% DMSO in water, 2 µl of a saturated solution of ascorbic acid and 40 µl of the solution containing ¹⁷⁷Lu were mixed and heated to 95° C. for 10 min. The labelling was checked by radio-HPLC (0-100% ACN in water within 5 min, Monolith column).

⁶⁸Ga-labelling

For the PET scan CHX-DTPA was labelled with ⁶⁸Ga. 1 ml of ⁶⁸Ga was eluted from a ⁶⁸Ge/⁶⁸Ga generator with 0.6 M HCl. 298 µl NaOAc buffer and 1 µl of a 10 mM solution of CHX-DTPA in DMSO was added and incubated for 5 min. Afterwards the product was purified using a SOLA cartridge. Washing was done with a 0.9% NaCl solution and for elution ethanol was used. The ethanol then was vaporized and the remaining product was dissolved in 100 µl of a 0.9% NaCl solution and 10 µl of phosphate buffer.

Example 6

Determination of the IC₅₀ Value

A filter plate MultiScreen_{HTS}-DV was incubated at room temperature with 100 µl PBS with 1% BSA per well for 30 min. After removing the PBS/BSA solution 10⁵ LNCaP cells in 50 µl of Opti-MEM were applied to each well. Different concentrations of the compounds (leading to concentrations of 0, 0.5, 1, 2.5, 5, 10, 25, 50, 100, 500, 1000 and 5000 nM in each well) in 300 µl of Opti-MEM were mixed with 3 µl of a 150 nM solution of ¹²⁵I-labeled MIP—1466 in Opti-MEM. 50 µl of the resulting solution were added to each well, each concentration was pipetted in quadruples. Each well now contained the radioactively labelled ligand in a concentration of 0.75 nM and the competitive, not labelled ligand in the concentration mentioned above. The plate was then incubated for 45 min at room temperature on a shaker.

After the incubation, the cells were washed 2× with 100 µl of ice cold PBS and 1× with 200 µl of ice cold PBS. Finally, the filters were collected and the remaining radioactivity was measured with a gamma counter. Each tube was measured for 5 min.

The data measured by the gamma counter were evaluated with Graphpad Prism to achieve an inhibition concentration 50 (IC₅₀) against the radioactively labelled MIP-1095.

Conjugate	IC ₅₀ [nM]
MB17-DOTA	0.13 ± 0.08
MB17-NOTA	0.14 ± 0.08
MB17-DTPA	0.12 ± 0.05
MB17-CHX-DTPA	0.06 ± 0.04
MB17-PCTA	0.10 ± 0.06
MB17-DO3A	0.10 ± 0.05
MB17-NODAGA	0.09 ± 0.08

Example 7

µPET—Imaging Using CHX-DTPA-MB17

Before injection into the mouse, the solution containing the purified ⁶⁸Ga-CHX-DTPA-coupled PSMA inhibitor was sterile-filtered. 100 µl of this solution was taken up into a syringe and then injected into a BALB/c nude mouse LNCaP xenograft, intravenously into the tail vein. The PET scan was recorded for 140 min with a Siemens Inveon PET (FIG. 15)

Example 8

Determination of the Competitive Binding Affinity

In order to compare the series of novel compounds the competitive binding affinity and the specific internalization was analyzed using the PSMA expressing cell line LNCaP. To determine specific cellular uptake, cells were blocked with 2-(phosphonomethyl)-pentanedioic acid (PMPA). The inhibition potency was also investigated by the enzyme-based NAALADase assay.

Cell Culture

For binding studies and in vivo experiments LNCaP cells (metastatic lesion of human prostatic adenocarcinoma, ATCC CRL-1740) were cultured in RPMI medium supplemented with 10% fetal calf serum and Glutamax (PAA, Austria). During cell culture, cells were grown at 37° C. in an incubator with humidified air, equilibrated with 5% CO₂. The cells were harvested using trypsin-ethylenediaminetetraacetic acid (trypsin-EDTA; 0.25% trypsin, 0.02% EDTA, all from PAA, Austria) and washed with PBS.

Cell Binding and Internalization

The competitive cell binding assay and internalization experiments were performed as described previously (Eder et al. 2012). Briefly, the respective cells (10⁵ per well) were incubated with the radioligand (⁶⁸Ga-labeled [Glu-urea-Lys (Ahx)]2-HBED-CC (Schafer et al., 2012) in the presence of 12 different concentrations of analyte (0-5000 nM, 100 µL/well). After incubation, washing was carried out using a multiscreen vacuum manifold (Millipore, Billerica, Mass.). Cell-bound radioactivity was measured using a gamma counter (Packard Cobra II, GMI, Minnesota, USA). The 50% inhibitory concentration (IC₅₀) was calculated by fitting the data using a nonlinear regression algorithm (GraphPad Software). Experiments were performed three times.

To determine the specific cell uptake and internalization, 10⁵ cells were seeded in poly-L-lysine coated 24-well cell culture plates 24 h before incubation. After washing, the cells were incubated with 25 nM of the radiolabeled compounds for 45 min at 37° C. and at 4° C., respectively. Specific cellular uptake was determined by competitive blocking with 2-(phosphonomethyl)pentanedioic acid (500 µM final concentration, PMPA, Axxora, Loerrach, Germany). Cellular uptake was terminated by washing 4 times with 1 mL of ice-cold PBS. Cells were subsequently incubated twice with 0.5 mL glycine-HCl in PBS (50 mM, pH=2.8) for 5 min to remove the surface-bound fraction. The cells were washed with 1 mL of ice-cold PBS and lysed using 0.3 N NaOH (0.5 mL). The surface-bound and the internalized fractions were measured in a gamma counter. The cell uptake was calculated as percent of the initially added radioactivity bound to 10⁶ cells [% ID/10⁶ cells].

Naaladase Assay

Recombinant human PSMA (rhPSMA, R&D systems, Wiesbaden, Germany) was diluted in assay buffer (50 mM HEPES, 0.1 M NaCl, pH 7.5) to 0.4 µg/mL. The substrate Ac-Asp-Glu (Sigma, Taufkirchen, Germany, 40 µM final concentration) was mixed with natGa labeled analyte at concentrations ranging from 0.05 nM to 1000 nM in a final volume of 125 µL assay buffer. The mixtures were combined with 125 µL of the rhPSMA solution (0.4 µg/mL) and incubated for one hour at 37° C. The reaction was stopped by heating at 95° C. for 5 minutes. 250 µL of a 15 mM solution of ortho-phthaldialdehyde (Sigma, Taufkirchen, Germany) was added to all vials and incubated for 10 minutes at ambient temperature. Finally, 200 µL of the reaction solutions were loaded onto a F16 Black Maxisorp Plate (Nunc, Langenselbold, Germany) and read at excitation and emission wavelengths of 330 nm and 450 nm, respectively, using a microplate reader (DTX-880, Beckman Coulter, Krefeld, Germany). The data were analyzed by a one site-total binding regression algorithm of GraphPad (GraphPad Software, California, USA).

Biodistribution

7- to 8-week-old male BALB/c nu/nu mice (Charles River Laboratories) were subcutaneously inoculated into the right trunk with 5×10⁶ cells of LNCaP (in 50% Matrigel; Becton Dickinson, Heidelberg, Germany). The tumors were allowed to grow until approximately 1 cm³ in size. The radiolabeled compounds were injected into the tail vein (approx. 1 MBq per mouse; 0.06 nmol). At 1 h after injection the animals were sacrificed. Organs of interest were dissected, blotted dry, and weighed. The radioactivity was measured using a gamma counter and calculated as % ID/g.

MicroPET

For the microPET studies, 10-25 MBq of the radiolabeled compounds in a volume of 0.15 ml (~0.5 nmol) were injected via a lateral tail vein into mice bearing LNCaP tumor xenografts. The anesthetized animals (2% sevoflurane, Abbott, Wiesbaden, Germany) were placed in prone position into the Inveon small animal PET scanner (Siemens, Knoxville, Tenn, USA) to perform dynamic micro-PET scans and 20 min-static scans; c.f. FIG. 1, 3, 5-14

TABLE A

Substance	IC ₅₀ [nM]	Internalization [% IA/10 ⁶ cells]
MB2	2.75 ± 0.82	8.78 ± 3.96 for Ga-68 5.22 ± 0.67 for Lu-177
MB3	10.51 ± 6.06	3.65 ± 1.32 for Lu-177
MB4	0.74 ± 0.50	14.18 ± 0.98 for Ga-68 14.25 ± 4.61 for Lu-177
MB10	8.67 ± 1.58	6.96 ± 3.90 for Lu-177
MB17	0.13 ± 0.08	17.02 ± 4.36 for Ga-68 17.51 ± 3.99 for Lu-177
MB17.D	12.41 ± 5.10	2.60 ± 0.14 for Lu-177
MB22	52.80	1.15 ± 0.19 for Lu-177
MB24	3.33	7.26 ± 2.76 for Lu-177
MB25	6.64	3.91 ± 0.54 for Lu-177
MB31	91.80	0.53 ± 0.48 for Lu-177
MB33	59.33	1.96 ± 0.20 for Lu-177
MB35	26.18	0.97 ± 0.17 for Lu-177

The present example shows that the binding affinity of PSMA inhibitors can be influenced by linker modifications. Two cyclic moieties and at least one aromatic moiety in the

linker region of the substance seem to be preferable and resulted in the high affinity compounds MB4 and MB17. These novel variants show low nanomolar affinity to LNCap cell line and were specifically internalized at 37° C. up to 48% ID/10⁶ cells. Former studies showed that besides binding affinity the internalization properties of PSMA-targeting probes are highly important and high internalization rates are essential for high in vivo tumor uptake and retention. Thus, MB17 represents a novel PSMA-targeting probe with optimal characteristics which was also confirmed by organ distribution and small animal PET imaging. MB 17 shows a high PSMA-specific tumor uptake (FIG. 2). In addition, dynamic PET imaging of MB17 (FIG. 2) shows an early enrichment in the bladder and also the maximum kidney uptake (highest point in the time-activity-curve) is as early as 15 min after injection of the radiotracer and diminishes substantially already after 20 minutes. With regard to therapeutic use, this gives clear clinical advantages for MB17 compared to other PSMA-inhibitors. In the PET diagrams (FIG. 1) MB17 shows a rapid background clearance as well as a substantial reduction of the enrichment in the kidney after 2 hours while it is further accumulated and retained in the PSMA-expressing tumor.

In addition, organ distribution with ¹⁷⁷Lu (FIG. 4) showed that the high initial kidney uptake is nearly completely washed out (2.13±1.36% ID/g) after 24 hours while the tumor uptake remained high and even increased (10.58±4.50% ID/g). Other organs as liver (0.08±0.03% ID/g), lung (0.11±0.13% ID/g) and spleen (0.13±0.05% ID/g) showed very low uptake. The favourable pharmacokinetics led to extremely high tumor-to-background ratios (Tumor/Blood: 1058; Tumor/Muscle: 529) after 24 hours

Table A clearly confirms that the chemical modifications in the linker region of the molecule affect the biological properties, e.g. affinity and internalization efficacy. MB17 and MB4 show the most promising binding properties on cells.

Example 9

Clinical Data Concerning MB17

PET/CT imaging was performed using the radiotracer MB17 labeled with Ga-68 (c.f. FIG. 17)

The ⁶⁸Ge/⁶⁸Ga-generator used for radiopharmaceutical production was purchased from IDB-Holland BV (Baarle-Nassau, The Netherlands). Disposable cassette kits and chemicals including the precursor in GMP-compliant grade used for the radiosynthesis were obtained from ABX advanced biochemical compounds (Radeberg, Germany). An Ultimate 3000 HPLC system (Dionex) (acetonitrile (A), water+0.1% TFA (B); gradient: 0.5 min 95% B, 10.0 min 80% A, flowrate: 2 mL/min) equipped with a Chromolith Performance RP-18e column (100×4.6 mm, Merck) and a NaI radiodetector (Raytest) was used to determine the radiochemical purity. Residual solvents were determined using a 6850 Series gas chromatograph (Agilent Technologies). Endotoxin testing was performed with an Endosafe®-PTS device (Charles River).

2 µg of MB17 were dissolved in 1.5 M acetate buffer pH 4.5 (1 mL) and 1 M ascorbic acid (10 µL) and transferred into the reaction vessel. The ⁶⁸Ge/⁶⁸Ga-generator was eluted with 10 mL of 0.6 M HCl and the eluate diluted with 9 mL

31

of ultrapure water. The mixture was then transferred to a cation exchange cartridge (Macherey-Nagel PS-H+, Size M) and eluted with 5 M NaCl solution (1.2 mL) into the preheated reaction vessel (100° C.). The reaction mixture was heated for 10 minutes. The crude reaction mixture was then removed from the reaction vessel and transferred to a pre-conditioned (10 mL EtOH/10 mL ultrapure water) C18 cartridge (Waters Sep-Pak light). 9 mL ultrapure water was used to rinse the reaction vessel and passed over the C18 cartridge. The C18 cartridge was washed with another 5 mL of ultrapure water. The final product was eluted from the C18 cartridge with 2 mL of EtOH/H₂O (v:v 1:1), sterile filtered (Millipore Cathivex-GV, 0.22 µm) and diluted with 10 mL of phosphate buffered saline (PBS) solution pH 7.4 (according to Eur. Ph. 8.0 (4005000)). The ⁶⁸Ga-MB17 complex solution was applied to patients via an intravenous bolus.

Example 10

Human Therapy with ¹⁷⁷Lu-labeled MB17

For therapy, the PSMA ligand MB17 was radiolabeled with Lu-177. ¹⁷⁷LuCl₃ was obtained from Perkin Elmer (4 GBq, NEZ307D, 0.04 M HCl). 80 nmoles of MB17 were dissolved in 400 µL sodium acetate buffer (0.4 M, pH 5) supplemented with 5 µL of 20% ascorbic acid. The solution was transferred to the ¹⁷⁷LuCl₃ and incubated for 10 minutes



32

at 95° C. Finally, 2 mL 0.9% NaCl was added. For quality control, ITLC and radio-HPLC was performed.

The ¹⁷⁷Lu-labeled MB17 was applied to patients via an intravenous bolus (5 mL, slowly within 30 seconds). The intravenous application was accompanied by an infusion of 0.9% NaCl for 4.5 h starting at 0.5 h before injection. Reference is made to FIG. 18.

The invention is further described by the following numbered paragraphs:


1. A compound of Formula (Ia) or (Ib):




n:	0, 1
m:	1, 2, 3, 4
Z:	—CO ₂ H, —SO ₂ H, —SO ₃ H, —SO ₄ H, —PO ₂ H, —PO ₃ H, —PO ₄ H ₂
X:	Naphthyl, Phenyl, Biphenyl, Indolyl (=2,3-benzopyrrolyl), Benzothiazolyl
Y:	Aryl, Alkylaryl, Cyclopentyl, Cyclohexyl, Cycloheptyl
Chelator radical of:	1,4,7,10-tetraazacyclododecane-N,N',N'',N'''-tetraacetic acid (=DOTA), N,N'-bis[2-hydroxy-5-(carboxyethyl)benzyl]ethylenediamine-N,N'-diacetic acid (=HBED-CC), 1,4,7-triazacyclononane-1,4,7-triacetic acid (=NOTA), 2-(4,7-bis(carboxymethyl)-1,4,7-triazonan-1-yl)pentanedioic acid (NODAGA), 2-(4,7,10-tris(carboxymethyl)-1,4,7,10-tetraazacyclododecan-1-yl)pentanedioic acid (DOTAGA), 1,4,7-triazacyclononane phosphinic acid (TRAP), 1,4,7-triazacyclononane-1-[methyl(2-carboxyethyl)phosphinic acid]-4,7-bis[methyl(2-hydroxymethyl)phosphinic acid] (NOPO), 3,6,9,15-tetraazabicyclo[9.3.1]pentadeca-1(15),11,13-triene-3,6,9-triacetic acid (=PCTA), N'-{5-[Acetyl(hydroxy)amino]pentyl}-N-[5-({4-[(5-aminopentyl)(hydroxy)amino]-4-oxobutanoyl}amino)pentyl]-N-hydroxysuccinamide (DFO), Diethylenetriaminepentaacetic acid (DTPA) Trans-cyclohexyl-diethylenetriaminepentaacetic acid (CHX-DTPA) 1-oxa-4,7,10-triazacyclododecane-4,7,10-triacetic acid (oxo-Do3A) p-isothiocyanatobenzyl-DTPA (SCN-Bz-DTPA) 1-(p-isothiocyanatobenzyl)-3-methyl-DTPA (1B3M) 2-(p-isothiocyanatobenzyl)-4-methyl-DTPA (1M3B) 1-(2-methyl-4-isocyanatobenzyl)-DTPA (MX-DTPA).

33


2. The compound of paragraph 1 having the structure R'-LINKER-R, wherein R'=radical of DOTA and R=radical of Glu-Urea-Lys:



wherein the compound R'-LINKER-R is selected from:


**34**

-continued



35

-continued

**36**

-continued

5


10

15



20



25

30






3. The compound of paragraph 1 or 2, selected from the following:









-continued




41

-continued

42





4. Use of the compound of any of paragraphs 1 to 3 for the preparation of radiolabeled compounds.

5. A metal complex comprising a radionuclide and a compound of any of paragraphs 1 to 3.

6. The metal complex of paragraph 5, wherein the radionuclide is ^{111}In , ^{90}Y , ^{68}Ga , ^{171}Lu , $^{99\text{m}}\text{Tc}$, ^{64}Cu , ^{153}Gd , ^{155}Gd , ^{157}Gd , ^{213}Bi , ^{225}Ac or Fe .

7. A pharmaceutical composition comprising a compound of any of paragraphs 1 to 3 or metal complex of paragraph 5 or 6, or a pharmaceutically acceptable salt, or ester thereof, and a pharmaceutically acceptable carrier.

8. Compound of any of paragraphs 1 to 3 or metal complex of paragraph 5 or 6 for use in a method of imaging in a patient.

9. Compound of any of paragraphs 1 to 3 or metal complex of paragraph 5 or 6 for use in a method of diagnosing prostate cancer and/or metastasis thereof.



10. Compound of any of paragraphs 1 to 3 or metal complex of paragraph 5 or 6 for use in a method of treating prostate cancer and/or metastasis thereof.

* * *

Having thus described in detail preferred embodiments of the present invention, it is to be understood that the invention defined by the above paragraphs is not to be limited to particular details set forth in the above description as many apparent variations thereof are possible without departing from the spirit or scope of the present invention.

What is claimed is:

1. A compound of Formula (Ia) or (Ib):



wherein:

n is chosen from 0 or 1;


m is chosen from 1, 2, 3, or 4;

X is chosen from Naphthyl, Phenyl, Biphenyl, Indolyl (=2,3-benzopyrrolyl), or Benzothiazolyl;

Y is chosen from Aryl, Alkylaryl, Cyclopentyl, Cyclohexyl, or Cycloheptyl; and

Z is independently chosen from $-\text{CO}_2\text{H}$, $-\text{SO}_2\text{H}$, $-\text{SO}_3\text{H}$, $-\text{SO}_4\text{H}$, $-\text{PO}_2\text{H}$, $-\text{PO}_3\text{H}$, or $-\text{PO}_4\text{H}_2$, with the proviso that the compound is not



45



2. The compound of claim 1, wherein Chelator is a radical of:

46



- 1,4,7,10-tetraazacyclododecane-N,N',N'',N'''-tetraacetic acid (DOTA);
 - N,N''-bis[2-hydroxy-5-(carboxyethyl)benzyl]ethylenediamine-N,N''-diacetic acid (HBED-CC);
 - 1,4,7-triazacyclononane-1,4,7-triacetic acid (NOTA);
 - 2-(4,7-bis(carboxymethyl)-1,4,7-triazonan-1-yl)pentanedioic acid (NODAGA);
 - 2-(4,7,10-tris(carboxymethyl)-1,4,7,10-tetraazacyclodecan-1-yl)pentanedioic acid (DOTAGA);
 - 1,4,7-triazacyclononane phosphinic acid (TRAP);
 - 1,4,7-triazacyclononane-1-[methyl(2-carboxyethyl)phosphinic acid]-4,7-bis[methyl(2-hydroxymethyl)phosphinic acid] (NOPO);
 - 3,6,9,15-tetraazabicyclo[9.3.1]pentadeca-1(15),11,13-triene-3,6,9-triacetic acid (PCTA);
 - N'-{5-[Acetyl(hydroxy)amino]pentyl}-N-[5-({4-[(5-aminopentyl)(hydroxy)amino]-4-oxobutanoyl}amino)pentyl]-N-hydroxysuccinamide (DFO);
 - diethylenetriaminepentaacetic acid (DTPA);
 - trans-cyclohexyl-diethylenetriaminepentaacetic acid (CHX-DTPA);
 - 1-oxa-4,7,10-triazacyclododecane-4,7,10-triacetic acid (oxo-Do3A);
 - p-isothiocyanatobenzyl-DTPA (SCN-Bz-DTPA);
 - 1-(p-isothiocyanatobenzyl)-3-methyl-DTPA (1B3M);
 - 2-(p-isothiocyanatobenzyl)-4-methyl-DTPA (1M3B); or
 - 1-(2-methyl-4-isocyanatobenzyl)-DTPA (MX-DTPA).
3. A compound selected from the following:




47

-continued





48



49



50



51

4. A radiolabeled compound of claim 1.
5. A radiolabeled compound of claim 2.
6. A metal complex comprising a radionuclide and a compound of claim 1.
7. The metal complex of claim 6, wherein the radionuclide is selected from ^{89}Zr , ^{44}Sc , ^{111}In , ^{90}Y , ^{68}Ga , ^{177}Lu , $^{99\text{m}}\text{Tc}$, ^{64}Cu , ^{67}Cu , ^{153}Gd , ^{155}Gd , ^{157}Gd , ^{213}Bi , or ^{225}Ac .
8. A pharmaceutical composition comprising the compound of claim 1 or a pharmaceutically acceptable salt, or ester thereof, and a pharmaceutically acceptable carrier.
9. A pharmaceutical composition comprising the metal complex of claim 6, or a pharmaceutically acceptable salt, or ester thereof, and a pharmaceutically acceptable carrier.
10. A method of imaging a region in a patient comprising: administering to the patient a diagnostically effective amount of the metal complex of claim 6, exposing the region of the patient to a scanning device, and obtaining an image of the region in the patient.
11. A method of diagnosing prostate cancer and/or metastasis thereof comprising: administering to the patient a diagnostically effective amount of the metal complex of claim 6, exposing the region of the patient to a scanning device, and obtaining an image of the region in the patient, wherein the image of a PSMA-expressing tumor indicates a diagnosis of prostate cancer and/or metastasis thereof.
12. A method of treating prostate cancer and/or metastasis thereof comprising administering to a patient in need thereof a therapeutically effective amount of the metal complex of claim 6, or a pharmaceutically acceptable salt or solvate of the complex.
13. A method of treating prostate cancer and/or metastasis thereof comprising administering to a patient in need thereof a therapeutically effective amount of the pharmaceutical composition of claim 9.
14. The method of claim 10 wherein the region of the patient is a head, thorax, spleen, kidney, or PSMA-expressing tumor tissue.
15. The method of claim 11 wherein the region of the patient is a head, thorax, spleen, kidney, or PSMA-expressing tumor tissue.
16. The method of claim 12 wherein the metastasis is in lung, liver, kidney, bone, brain, spinal cord, or bladder.
17. A metal complex comprising a radionuclide and a compound of claim 1.
18. The metal complex of claim 17, wherein the radionuclide is selected from ^{89}Zr , ^{44}Sc , ^{111}In , ^{90}Y , ^{68}Ga , ^{177}Lu , $^{99\text{m}}\text{Tc}$, ^{64}Cu , ^{67}Cu , ^{153}Gd , ^{155}Gd , ^{157}Gd , ^{213}Bi , or ^{225}Ac .
19. A pharmaceutical composition comprising the compound of claim 1 or a pharmaceutically acceptable salt, or ester thereof, and a pharmaceutically acceptable carrier.
20. A pharmaceutical composition comprising the metal complex of claim 17, or a pharmaceutically acceptable salt, or ester thereof, and a pharmaceutically acceptable carrier.
21. A method of imaging a region in a patient comprising: administering to the patient a diagnostically effective amount of the metal complex of claim 17, exposing the region of the patient to a scanning device, and obtaining an image of the region in the patient.
22. A method of diagnosing prostate cancer and/or metastasis thereof comprising: administering to the patient a diagnostically effective amount of the metal complex of claim 17, exposing the region of the patient to a scanning device, and obtaining an image of the region in the patient,

52

wherein the image of a PSMA-expressing tumor indicates a diagnosis of prostate cancer and/or metastasis thereof.

23. A method of treating prostate cancer and/or metastasis thereof comprising administering to a patient in need thereof a therapeutically effective amount of the metal complex of claim 17, or a pharmaceutically acceptable salt or solvate of the complex.


24. A method of treating prostate cancer and/or metastasis thereof comprising administering to a patient in need thereof a therapeutically effective amount of the pharmaceutical composition of claim 20.

25. The method of claim 21, wherein the region of the patient is a head, thorax, spleen, kidney, or PSMA-expressing tumor tissue.


26. The method of claim 22, wherein the region of the patient is a head, thorax, spleen, kidney, or PSMA-expressing tumor tissue.

27. The method of claim 23, wherein the metastasis is in lung, liver, kidney, bone, brain, spinal cord, or bladder.

28. An $\text{R}'\text{-LINKER-R}$ compound with $\text{R}'=\text{radical of DOTA}$ and $\text{R}=\text{radical of Glu-Urea-Lys}$, and having a structure of:




wherein the $\text{R}'\text{-LINKER-R}$ compound is selected from:




53

-continued

**54**

-continued



* * * * *