

Angiogenesis

An Integrative Approach From Science to Medicine

Edited by

William D. Figg

Medical Oncology Branch, Center for Cancer Research,
National Cancer Institute, Bethesda, Maryland, USA

Judah Folkman

Department of Surgery, Harvard Medical School and
Vascular Biology Program, Children's Hospital Boston,
Boston, Massachusetts, USA



Editors

William D. Figg
Medical Oncology Branch
Center for Cancer Research
National Cancer Institute
Bethesda, Maryland
USA

Judah Folkman
Department of Surgery
Harvard Medical School and
Vascular Biology Program
Children's Hospital Boston
Boston, Massachusetts
USA

ISBN: 978-0-387-71517-9 e-ISBN: 978-0-387-71518-6
DOI: 10.1007/978-0-387-71518-6

Library of Congress Control Number: 2007934647

© 2008 Springer Science+Business Media, LLC

All rights reserved. This work may not be translated or copied in whole or in part without the written permission of the publisher (Springer Science+Business Media, LLC, 233 Spring Street, New York, NY 10013, USA), except for brief excerpts in connection with reviews or scholarly analysis. Use in connection with any form of information storage and retrieval, electronic adaptation, computer software, or by similar or dissimilar methodology now known or hereafter developed is forbidden.

The use in this publication of trade names, trademarks, service marks, and similar terms, even if they are not identified as such, is not to be taken as an expression of opinion as to whether or not they are subject to proprietary rights.

While the advice and information in this book are believed to be true and accurate at the date of going to press, neither the authors nor the editors nor the publisher can accept any legal responsibility for any errors or omissions that may be made. The publisher makes no warranty, express or implied, with respect to the material contained herein.

Printed on acid-free paper

9 8 7 6 5 4 3 2 1

springer.com

Chapter 36

Clinical Development of VEGF Trap

John S. Rudge, Ella Ioffe, Jingtai Cao, Nick Papadopoulos, Gavin Thurston, Stanley J. Wiegand, and George D. Yancopoulos

Keywords: VEGF Trap, vascular eye disease, kinase inhibitors

Abstract: The inhibition of angiogenesis is proving to be an effective strategy in treating diseases involving pathological angiogenesis such as cancer and ocular vascular diseases. Since its discovery in the 1980s, vascular endothelial cell growth factor (VEGF) has been shown to play a vital role in both physiological and pathological angiogenesis, resulting in the development of numerous approaches to block VEGF and VEGF signaling, ranging from small molecule tyrosine kinase inhibitors to protein-based and RNA-based therapeutic candidates. VEGF Trap is one such protein-based agent that has been engineered to bind and sequester VEGF, as well as placental growth factor (PlGF), with high affinity. VEGF Trap has been shown to effectively inhibit pathological angiogenesis in numerous preclinical models of cancer and eye disease, and is now being evaluated in clinical trials in several types of cancer, as well as the ‘wet’ or neovascular form of age-related macular degeneration (AMD). This chapter will summarize the basic biology of VEGF and the progress of the VEGF Trap from the bench to the clinic.

Introduction

Angiogenesis is a vital process not only during development, but also in the adult in settings of wound repair and reproduction [1, 2]. However, in diseases characterized by uncontrolled, pathological angiogenesis, such as solid tumors or vascular diseases of the eye, inhibition of angiogenesis would be of therapeutic benefit. While diverse factors might be expected to regulate angiogenesis in different settings, a clear consensus is emerging that a single growth factor, VEGF is the

critical requisite driver of both physiological and pathological angiogenesis in most settings [3]. Thus, while blocking VEGF signaling during early development can lead to severe growth retardation, it can also produce highly beneficial effects when applied to disease states characterized by pathological neovascularization [3]. Recent clinical studies have validated VEGF as a bona fide target for therapeutic intervention in cancer as well as in vascular diseases affecting the eye, such as wet AMD. These studies have led to approval by the U.S. Food and Drug Administration (FDA) of drugs that target the VEGF pathway, specifically antibodies directed against VEGF, or kinase inhibitors which block activation of the VEGF receptors [4–7]. Emerging therapeutic candidates that otherwise target VEGF signaling have also reported encouraging results [8].

Biology of VEGF and Its Receptors

VEGF is widely known to promote angiogenesis, by stimulating vascular endothelial cell proliferation, migration and tube formation, and it can also markedly increase vascular permeability [9]. VEGF-A is the prototypical member of a family of factors that also consists of VEGF-B, VEGF-C, VEGF-D and PlGF, which bind differentially to VEGF receptors 1, 2 and 3 and neuropilin with different specificities [10,11]. In addition, alternative exon splicing results in the production of four major isoforms of human VEGF-A – VEGF₁₂₁, VEGF₁₆₅, VEGF₁₈₉, VEGF₂₀₆ differentiated by their heparin binding affinity [12]. VEGF₁₂₁ differs from the other isoforms in that it does not bind heparin and thus is freely diffusible. As the isoforms increase in molecular weight, they show increasing heparin binding affinity and consequently bind extracellular matrix such that they tend to be sequestered near their site of production [13–15]. Matrix metalloproteases (MMPs) appear to play an important role in regulating the release of these isoforms from matrix [16]. Although endothelial cells are the primary targets of VEGF actions, more recently it has been shown that other cell types, including monocytes, hematopoietic stem cells and neurons, can also respond to VEGF [17–19].

Regeneron Pharmaceuticals, 777 Old Saw Mill River Road
Tarrytown, NY 10591, USA
E-mail: John.Rudge@regeneron.com

The VEGF receptor 2 (VEGFR2) appears to be the major mediator of the mitogenic, angiogenic and pro-permeability actions of VEGF-A. Indeed, genetic deletion of VEGFR2 results in profound defects in embryonic vasculogenesis, including a failure to develop blood islands and organized blood vessels [20]. In contrast, the role of VEGFR1, which was the first VEGF receptor identified, is more difficult to discern [21]. To date, VEGFR1 has been implicated in MMP9 induction, hematopoiesis, and monocyte chemotaxis, and also appears to act as a 'biological sink' which sequesters VEGF from binding to the lower affinity receptor, VEGFR2 [11]. A further level of complexity was revealed when VEGF was shown to bind neuropilin, a molecule previously identified as a receptor for the collapsin/semaphorin family of ligands involved in axon guidance. The binding of VEGF to neuropilin is thought to result in presentation of VEGF to VEGFR2, augmenting VEGFR2 signal transduction [22].

The Evidence for VEGF as a Key Player in Tumor Angiogenesis

In situ hybridization studies have revealed that VEGF is highly expressed in a number of human tumors [23–27]. This is most apparent in renal cell carcinoma where mutations in the von Hippel-Lindau (VHL) tumor suppressor gene result in increased transcription of hypoxia-inducible factor (HIF1) genes [28,29].

Numerous preclinical studies have shown that the growth of many different tumor types can be inhibited using agents that variously inhibit VEGF signaling (small molecule inhibitors, VEGFR2 antibodies, soluble VEGF receptors including the VEGF Trap) [30–35]. These studies have also shown that VEGF derived from the stromal compartment, as well as the tumor itself, plays an important role in mediating the angiogenesis which supports tumor growth [32,36].

Not surprisingly, antiangiogenic therapy results in a cytostatic effect in many tumor types rather than frank regression. Thus, in many models, combination of VEGF blockade with chemotherapy or radiation therapy results in greater efficacy than either approach alone [37,38].

VEGF Pathway Inhibitors Approved for the Treatment of Cancer

Bevacizumab (Avastin®)

The positive results from preclinical studies have led to the testing of several VEGF inhibitors in clinical trials. These include the humanized anti-VEGF-A monoclonal antibody (bevacizumab, Avastin®), an anti-VEGFR antibody [39], small molecules that inhibit VEGFR signaling and VEGF Trap [4,40–42]. The first clinical validation of the anti-VEGF approach came with FDA approval of bevacizumab in 2004, based on the results of a randomized double-blind phase III trial in which

bevacizumab was combined with bolus IFL (irinotecan, 5FU, leucovorin) chemotherapy as first line therapy for metastatic colorectal cancer. Median survival increased from 15.6 months in the IFL alone arm to 20.3 months in the IFL + bevacizumab arm. Severe hypertension was observed in about 10% of bevacizumab treated patients, and gastrointestinal perforation was noted in ~2% of patients. In addition, the incidence of arterial thromboembolic complications (stroke, myocardial infarction, transient ischemic attacks, unstable angina) was double the incidence seen with chemotherapy alone [43].

Kinase Inhibitors

An alternative approach to using antibodies that bind and neutralize VEGF is the use of tyrosine kinase inhibitors that target the VEGF and other receptors. To date, two such kinase inhibitors have been approved by the FDA: SU11248 (sunitinib; Sutent®) and Bay 43–9006 (sorafenib; Nexavar®). Sunitinib inhibits tyrosine phosphorylation of VEGFR1, VEGFR2, platelet derived growth factor receptor (PDGFR), c-kit and Flt3, and has been approved for the treatment of Gleevec-resistant gastrointestinal stromal tumors (GIST) and metastatic renal cell carcinoma [7]. Sorafenib, which inhibits tyrosine phosphorylation of raf, VEGFR2/3, PDGFR, kit and Flt3, was approved in 2006 for the treatment of metastatic renal cell carcinoma [5,6]. It should be pointed out that these kinase inhibitors do not block VEGF signaling selectively or completely, and much of their clinical benefit could derive from the inhibition of other kinase pathways.

Development and Application of VEGF Trap in Preclinical Animal Models

Despite the important benefits in patient care provided by the currently approved VEGF pathway blockers, current evidence suggests that optimal VEGF blockade may not have yet been achieved (e.g., dose response studies do not indicate that saturation of benefit has been reached) [44,45], raising the possibility that more potent VEGF blockade could provide even more benefit for patients. The VEGF Trap may provide the opportunity to test this hypothesis, as it was designed to bind VEGF with exceedingly high-affinity. The VEGF Trap is a soluble chimeric receptor in which key domains of VEGFR1 (domain 2) and VEGFR2 (domain 3) are fused to the constant region (Fc portion) of human IgG1 [34]. This fully human protein is capable of binding all isoforms of VEGF-A with very high affinity (KD <0.5 pM for hVEGF₁₆₅). Moreover, and in contrast to antibodies directed against VEGF-A, the VEGF Trap also binds PlGF, a VEGF family member also implicated in pathological angiogenesis [46] (KD ~25 pM for hPlGF2). In addition, the VEGF Trap was engineered to exhibit excellent pharmacokinetic properties, and has a circulating half-life in humans of approximately 2–3 weeks, allowing for bi-weekly or even less frequent dosing.

Once VEGF Trap had been optimized, it was tested and shown to have significant antiangiogenic and anti-tumor efficacy in

numerous preclinical tumor models [34,37,38,47–50]. When administered in combination with chemotherapy or radiation, VEGF Trap also produced a significant additive impact on tumor growth. Moreover, in an ovarian cancer model, VEGF Trap potently inhibited ascites formation [37,38,50]. The impressive efficacy profile of VEGF Trap in preclinical animal models justified its progression into clinical trials.

Interestingly, VEGF Trap also has an additional property that distinguishes it from antibodies: while antibodies form multimeric complexes with their antigens that are rapidly cleared from the systemic circulation, VEGF Trap forms an inert 1:1 complex with VEGF that remains in the circulation and can thus be readily measured. Thus, when given in doses that saturate VEGF binding, this unique property of the VEGF:VEGF Trap complex allows for the accurate determination of tumor and whole body VEGF production rates [51]. Somewhat surprisingly, when non-tumor bearing animals and humans are given the VEGF Trap, we find that total body production rates of VEGF are quite high, challenging previous claims that systemic VEGF levels can be used as a sensitive measure of tumor burden [51]. This finding has the important corollary that agents designed to bind and inactivate tumor-derived VEGF must be provided in amounts sufficient to ensure that they are not effectively consumed by the large amounts of VEGF normally produced by the body. Building on this finding, we have determined that measurement of the levels of free and bound VEGF Trap provides a clear index of the doses required to capture all available VEGF, offering a useful guide for dosing of this antiangiogenic agent. Based on these assays, the doses currently being used in the clinic for cancer appear to be in the therapeutic range.

VEGF Trap in Clinical Trials for Cancer

Multiple Phase 1 trials have been carried out using the VEGF Trap as a single agent, or in combination with various chemotherapeutic regimens, in patients with advanced cancers, with more than 300 patients treated to date. Numerous objective responses as well as cases of prolonged stable disease have been noted in these trials. Interestingly, in addition to anti-tumor responses, improvements in tumor-associated co-morbidities have been observed: in particular, substantial or complete resolution of tumor-associated ascites. The major adverse events noted in these trials are consistent with those seen with other anti-VEGF agents, and include hypertension and proteinuria. Importantly, no antibodies to VEGF Trap were observed in any patients.

VEGF Trap is currently in Phase II clinical trials and is about to initiate multiple large Phase 3 trials. Preliminary results from two of these studies were recently reported at the American Society of Clinical Oncology 2007 annual meeting [41,42]. Results were reported from an interim analysis of a randomized, double-blind, multi-center Phase 2 trial comparing two doses of VEGF Trap in patients with recurrent, platinum-resistant epithelial ovarian cancer. The patients selected for this study

were heavily pre-treated and had failed several other treatment regimens. While the study remains blinded with respect to dose, the pooled preliminary results from both the high and low dose groups demonstrated anti-tumor activity, as evidenced by an 8% objective tumor response rate, with a 13% response rate as measured by a 50% reduction in circulating levels of the tumor marker CA-125. Disease was judged to be stable in 77% of patients at 4 weeks, and in 41% of patients at 14 weeks. In addition, of 24 patients in the study who had tumor-associated ascites, 29% experienced complete resolution of their ascites while 54% experienced no increase in ascites during treatment. Tolerability was similar to other molecules in this class with hypertension being the most common grade 3/4 event (16%). Two patients (1.2%) experienced bowel perforation, both of whom recovered. In a collaboration involving Sanofi-Aventis and Regeneron Pharmaceuticals, the ongoing single-agent studies of VEGF Trap will be complemented by a large Phase 3 program combining VEGF Trap with standard chemotherapy regimens in at least 5 different advanced solid tumors: colorectal, non-small cell lung, prostate, pancreas and gastric cancer, with the first of these studies initiating in 2007.

Preclinical Studies with VEGF Trap in Vascular Eye Diseases

The finding that VEGF-A was up-regulated in the aqueous and vitreous humor of patients with proliferative diabetic retinopathy extended the focus of antiangiogenic agents from cancer to vascular eye disease [52,53]. Wet age-related macular degeneration (AMD) is the most common cause of vision loss in the elderly [53,54]. In this disease, VEGF is believed to mediate the abnormal growth of choroidal vessels that, together with the associated vascular leak and edema, disrupts the normal retinal architecture. VEGF Trap has also been tested in a number of rodent models of ocular vascular disease, where it has been shown to inhibit choroidal [55] and corneal [56] neovascularization in addition to suppressing vascular leak in the retina [57]. VEGF Trap has also been shown to improve the survival of corneal transplants [58]. In a primate model of AMD, in which a laser was used to induce choroidal vascular lesions, intravitreal administration of VEGF Trap was found not only to prevent the development of vascular leak and neovascularization when administered before the time of injury but also completely resolved vascular leak when administration was delayed until after the lesions had fully developed [59].

VEGF Trap in Clinical Trials for Vascular Eye Disease

The promising results in preclinical models supported the introduction of VEGF Trap into the clinic for treatment of both wet AMD and diabetic macular edema, using a version of VEGF Trap specifically formulated for intra-ocular administration,

termed VEGF Trap-Eye. Following successful completion of Phase I studies in wet AMD, VEGF Trap-Eye has progressed to a Phase 2 trial, for which interim results were recently presented at the Association for Research in Vision and Ophthalmology (ARVO) 2007 annual meeting [60,61]. In this trial, VEGF Trap-Eye was given at monthly intervals at doses of 0.5 or 2.0 mg, or as a single injection of 0.5, 2.0 or 4.0 mg. In the interim analysis, VEGF Trap-Eye met the pre-specified primary endpoint of reducing retinal thickness as measured by ocular coherence tomography (OCT) at 12 weeks compared with baseline (all groups combined, decrease of 135 μm , $P < 0.0001$). Mean change in visual acuity, a key secondary endpoint of the study, also demonstrated a statistically significant improvement (all groups combined, increase of 5.9 letters, $P < 0.0001$). Interestingly, patients in the highest monthly dose group achieved an average vision gain of more than 10 letters at 12 weeks, and even a single injection of VEGF Trap-Eye resulted in improved mean visual acuity at 12 weeks (all dose levels combined). Thus, the increased potency of VEGF Trap-Eye may offer the potential for improved efficacy and/or longer dosing intervals compared to the current standard of care [62–64]. In the VEGF Trap-Eye Phase 2 trial, there were no drug-related serious adverse events, and treatment with the VEGF Trap-Eye was generally well-tolerated. The most common adverse events were those typically associated with intravitreal injections. A Phase III program for VEGF Trap-Eye in wet AMD will be initiated in 2007.

In addition, as reported at ARVO 2007 [60,65], a Phase 1 study of VEGF Trap-Eye in diabetic macular edema (DME) showed that a single intravitreal injection in patients with longstanding diabetes resulted in a marked decrease in mean central retinal thickness and mean macular volume throughout the 6 week observation period. Additional trials with VEGF Trap-Eye in DME are anticipated in the future.

Conclusion

The VEGF Trap is an all human, chimeric receptor-based protein engineered to potently bind all forms of VEGF-A and PlGF, and to exhibit superior pharmacokinetic properties. The promise of the VEGF Trap in preclinical models of cancer and vascular eye diseases is being maintained during its initial evaluation in Phase 1 and 2 human trials, offering the hope that this therapeutic candidate may offer additional benefit to patients suffering from cancer or blinding vascular eye diseases.

Bibliography

1. Folkman J, Klagsbrun M. Angiogenic factors. *Science* 1987;235(4787):442–7.
2. Carmeliet P. Angiogenesis in life, disease and medicine. *Nature* 2005;438(7070):932–6.
3. Ferrara N. Role of vascular endothelial growth factor in physiologic and pathologic angiogenesis: therapeutic implications. *Semin Oncol* 2002;29(6 Suppl 16):10–
4. Ferrara N, Hillan KJ, Novotny W. Bevacizumab (Avastin), a humanized anti-VEGF monoclonal antibody for cancer therapy. *Biochem Biophys Res Commun* 2005;333(2):328–35.
5. Ratain MJ, Eisen T, Stadler WM, et al. Phase II placebo-controlled randomized discontinuation trial of sorafenib in patients with metastatic renal cell carcinoma. *J Clin Oncol* 2006;24(16):2505–12.
6. Escudier B, Eisen T, Stadler WM, et al. Sorafenib in advanced clear-cell renal-cell carcinoma. *N Engl J Med* 2007;356(2):125–34.
7. Motzer RJ, Hoosen S, Bello CL, Christensen JG. Sunitinib malate for the treatment of solid tumours: a review of current clinical data. *Expert Opin Investig Drugs* 2006;15(5):553–61.
8. Ho QT, Kuo CJ. Vascular endothelial growth factor: Biology and therapeutic applications. *Int J Biochem Cell Biol* 2007.
9. Dvorak HF. Vascular permeability factor/vascular endothelial growth factor: a critical cytokine in tumor angiogenesis and a potential target for diagnosis and therapy. *J Clin Oncol* 2002;20(21):4368–80.
10. Ferrara N, Gerber HP, LeCouter J. The biology of VEGF and its receptors. *Nat Med* 2003;9(6):669–76.
11. Yancopoulos GD, Davis S, Gale NW, Rudge JS, Wiegand SJ, Holash J. Vascular-specific growth factors and blood vessel formation. *Nature* 2000;407(6801):242–8.
12. Tischer E, Mitchell R, Hartman T, et al. The human gene for vascular endothelial growth factor. Multiple protein forms are encoded through alternative exon splicing. *J Biol Chem* 1991;266(18):11947–54.
13. Park JE, Keller GA, Ferrara N. The vascular endothelial growth factor (VEGF) isoforms: differential deposition into the sub-epithelial extracellular matrix and bioactivity of extracellular matrix-bound VEGF. *Mol Biol Cell* 1993;4(12):1317–26.
14. Houck KA, Leung DW, Rowland AM, Winer J, Ferrara N. Dual regulation of vascular endothelial growth factor bioavailability by genetic and proteolytic mechanisms. *J Biol Chem* 1992;267(36):26031–7.
15. Keyt BA, Berleau LT, Nguyen HV, et al. The carboxyl-terminal domain (111–165) of vascular endothelial growth factor is critical for its mitogenic potency. *J Biol Chem* 1996;271(13):7788–95.
16. Lee S, Jilani SM, Nikolova GV, Carpizo D, Iruela-Arispe ML. Processing of VEGF-A by matrix metalloproteinases regulates bioavailability and vascular patterning in tumors. *J Cell Biol* 2005;169(4):681–91.
17. Barleon B, Sozzani S, Zhou D, Weich HA, Mantovani A, Marme D. Migration of human monocytes in response to vascular endothelial growth factor (VEGF) is mediated via the VEGF receptor flt-1. *Blood* 1996;87(8):3336–43.
18. Clauss M, Gerlach M, Gerlach H, et al. Vascular permeability factor: a tumor-derived polypeptide that induces endothelial cell and monocyte procoagulant activity, and promotes monocyte migration. *J Exp Med* 1990;172(6):1535–45.
19. Gerber HP, Malik AK, Solar GP, et al. VEGF regulates haematopoietic stem cell survival by an internal autocrine loop mechanism. *Nature* 2002;417(6892):954–8.
20. Shalaby F, Rossant J, Yamaguchi TP, et al. Failure of blood-island formation and vasculogenesis in Flk-1-deficient mice. *Nature* 1995;376(6535):62–6.
21. de Vries C, Escobedo JA, Ueno H, Houck K, Ferrara N, Williams LT. The fms-like tyrosine kinase, a receptor for vascular endothelial growth factor. *Science* 1992;255(5047):989–91.
22. Soker S, Takashima S, Miao HQ, Neufeld G, Klagsbrun M. Neuropilin-1 is expressed by endothelial and tumor cells as an

- isoform-specific receptor for vascular endothelial growth factor. *Cell* 1998;92(6):735–45.
23. Yoshiji H, Gomez DE, Shibuya M, Thorgeirsson UP. Expression of vascular endothelial growth factor, its receptor, and other angiogenic factors in human breast cancer. *Cancer Res* 1996;56(9):2013–6.
 24. Sowter HM, Corps AN, Evans AL, Clark DE, Charnock-Jones DS, Smith SK. Expression and localization of the vascular endothelial growth factor family in ovarian epithelial tumors. *Lab Invest* 1997;77(6):607–14.
 25. Volm M, Koomagi R, Mattern J. Prognostic value of vascular endothelial growth factor and its receptor Flt-1 in squamous cell lung cancer. *Int J Cancer* 1997;74(1):64–8.
 26. Ellis LM, Takahashi Y, Fenoglio CJ, Cleary KR, Bucana CD, Evans DB. Vessel counts and vascular endothelial growth factor expression in pancreatic adenocarcinoma. *Eur J Cancer* 1998;34(3):337–40.
 27. Tomisawa M, Tokunaga T, Oshika Y, et al. Expression pattern of vascular endothelial growth factor isoform is closely correlated with tumour stage and vascularisation in renal cell carcinoma. *Eur J Cancer* 1999;35(1):133–7.
 28. Iliopoulos O, Levy AP, Jiang C, Kaelin WG, Jr., Goldberg MA. Negative regulation of hypoxia-inducible genes by the von Hippel-Lindau protein. *Proc Natl Acad Sci USA* 1996;93(20):10595–9.
 29. Lonser RR, Glenn GM, Walther M, et al. von Hippel-Lindau disease. *Lancet* 2003;361(9374):2059–67.
 30. Kim KJ, Li B, Winer J, et al. Inhibition of vascular endothelial growth factor-induced angiogenesis suppresses tumour growth in vivo. *Nature* 1993;362(6423):841–4.
 31. Prewett M, Huber J, Li Y, et al. Antivascular endothelial growth factor receptor (fetal liver kinase 1) monoclonal antibody inhibits tumor angiogenesis and growth of several mouse and human tumors. *Cancer Res* 1999;59(20):5209–18.
 32. Gerber HP, Kowalski J, Sherman D, Eberhard DA, Ferrara N. Complete inhibition of rhabdomyosarcoma xenograft growth and neovascularization requires blockade of both tumor and host vascular endothelial growth factor. *Cancer Res* 2000;60(22):6253–8.
 33. Smith JK, Mamoon NM, Duhe RJ. Emerging roles of targeted small molecule protein-tyrosine kinase inhibitors in cancer therapy. *Oncol Res* 2004;14(4–5):175–225.
 34. Holash J, Davis S, Papadopoulos N, et al. VEGF-Trap: a VEGF blocker with potent antitumor effects. *Proc Natl Acad Sci U S A* 2002;99(17):11393–8.
 35. Rudge JS, Thurston G, Davis S, et al. VEGF trap as a novel anti-angiogenic treatment currently in clinical trials for cancer and eye diseases, and VelociGene- based discovery of the next generation of angiogenesis targets. *Cold Spring Harb Symp Quant Biol* 2005;70:411–8.
 36. Liang WC, Wu X, Peale FV, et al. Cross-species vascular endothelial growth factor (VEGF)-blocking antibodies completely inhibit the growth of human tumor xenografts and measure the contribution of stromal VEGF. *J Biol Chem* 2006;281(2):951–61.
 37. Byrne AT, Ross L, Holash J, et al. Vascular endothelial growth factor-trap decreases tumor burden, inhibits ascites, and causes dramatic vascular remodeling in an ovarian cancer model. *Clin Cancer Res* 2003;9(15):5721–8.
 38. Wachsberger PR, Burd R, Cardi C, et al. VEGF Trap in combination with radiotherapy improves tumor control in U87 glioblastoma. *Int J Radiat Oncol Biol Phys* 2007.
 39. Shaheen RM, Ahmad SA, Liu W, et al. Inhibited growth of colon cancer carcinomatosis by antibodies to vascular endothelial and epidermal growth factor receptors. *Br J Cancer* 2001;85(4):584–9.
 40. Jain RK, Duda DG, Clark JW, Loeffler JS. Lessons from phase III clinical trials on anti-VEGF therapy for cancer. *Nat Clin Pract Oncol* 2006;3(1):24–40.
 41. Massarelli E, Miller VA, Leighl N, et al. Phase II Study of the Efficacy and Safety of Intravenous (IV) AVE0005 (VEGF Trap) Given Every 2 Weeks in Patients (Pts) with Platinum- and Erlotinib-Resistant Adenocarcinoma of the Lung (NSCLC). *ASCO Abs* 2007.
 42. Tew WP, Colombo N, Ray-Coquard I, et al. VEGF-Trap for patients (pts) with recurrent platinum-resistant epithelial ovarian cancer (EOC): Preliminary results of a randomized, multicenter phase III study. *ASCO Abs* 2007;In Press.
 43. Hurwitz H, Fehrenbacher L, Novotny W, et al. Bevacizumab plus irinotecan, fluorouracil, and leucovorin for metastatic colorectal cancer. *N Engl J Med* 2004;350(23):2335–42.
 44. Yang JC. Bevacizumab for patients with metastatic renal cancer: an update. *Clin Cancer Res* 2004;10(18 Pt 2):6367S–70S.
 45. Yang JC, Haworth L, Sherry RM, et al. A randomized trial of bevacizumab, an anti-vascular endothelial growth factor antibody, for metastatic renal cancer. *N Engl J Med* 2003;349(5):427–34.
 46. Luttun A, Tiwa M, Carmeliet P. Placental growth factor (PlGF) and its receptor Flt-1 (VEGFR1): novel therapeutic targets for angiogenic disorders. *Ann NY Acad Sci* 2002;979:80–93.
 47. Fukasawa M, Korc M. Vascular endothelial growth factor-trap suppresses tumorigenicity of multiple pancreatic cancer cell lines. *Clin Cancer Res* 2004;10(10):3327–32.
 48. Huang J, Frischer JS, Serur A, et al. Regression of established tumors and metastases by potent vascular endothelial growth factor blockade. *Proc Natl Acad Sci USA* 2003;100(13):7785–90.
 49. Dalal S, Berry AM, Cullinane CJ, et al. Vascular endothelial growth factor: a therapeutic target for tumors of the Ewing's sarcoma family. *Clin Cancer Res* 2005;11(6):2364–78.
 50. Hu L, Hofmann J, Holash J, Yancopoulos GD, Sood AK, Jaffe RB. Vascular endothelial growth factor trap combined with paclitaxel strikingly inhibits tumor and ascites, prolonging survival in a human ovarian cancer model. *Clin Cancer Res* 2005;11(19 Pt 1):6966–71.
 51. Rudge JS, Holash J, Hylton D, et al. VEGF Trap Complex Formation measures production rates of VEGF, providing a new biomarker for predicting efficacious angiogenic blockade. *PNAS* 2007; *Proc Natl Acad Sci USA* 2007;104(47):18363–70.
 52. Aiello LM, Cavallerano J. Diabetic retinopathy. *Curr Ther Endocrinol Metab* 1994;5:436–46.
 53. Malecize F, Clamens S, Simorre-Pinatel V, et al. Detection of vascular endothelial growth factor messenger RNA and vascular endothelial growth factor-like activity in proliferative diabetic retinopathy. *Arch Ophthalmol* 1994;112(11):1476–82.
 54. Congdon N, O'Colmain B, Klaver CC, et al. Causes and prevalence of visual impairment among adults in the United States. *Arch Ophthalmol* 2004;122(4):477–85.
 55. Saishin Y, Saishin Y, Takahashi K, et al. VEGF-TRAP(R1R2) suppresses choroidal neovascularization and VEGF-induced breakdown of the blood-retinal barrier. *J Cell Physiol* 2003;195(2):241–8.
 56. Wiegand S, Cao J, Renard R, Rudge J, Yancopoulos G. Long-lasting inhibition of corneal neovascularization following systemic administration of the VEGF Trap. *Invest Ophthalmol Vis Sci* 2003;44(Abtract):829.

57. Qaum T, Xu Q, Jousseaume AM, et al. VEGF-initiated blood-retinal barrier breakdown in early diabetes. *Invest Ophthalmol Vis Sci* 2001;42(10):2408–13.
58. Cursiefen C, Cao J, Chen L, et al. Inhibition of hemangiogenesis and lymphangiogenesis after normal-risk corneal transplantation by neutralizing VEGF promotes graft survival. *Invest Ophthalmol Vis Sci* 2004;45(8):2666–73.
59. Wiegand SJ, Zimmer E, Nork TM, et al. VEGF Trap Both Prevents Experimental Choroidal Neovascularization and Causes Regression of Established Lesions in Non-Human Primates. *Invest Ophthalmol Vis Sci* 2005;46:E-Abstract 1411.
60. Ciulla TA. VEGF Trap, Potential Therapeutic Role for Neovascular AMD and DME. ARVO 2007.
61. Benzl MS, Nguyen QD, Chu K, et al. CLEAR-IT-2: Interim Results Of The Phase II, Randomized, Controlled Dose-and Interval-ranging Study Of Repeated Intravitreal VEGF Trap Administration In Patients With Neovascular Age-related Macular Degeneration (AMD). 2007.
62. Brown DM, Kaiser PK, Michels M, et al. Ranibizumab versus verteporfin for neovascular age-related macular degeneration. *N Engl J Med* 2006;355(14):1432–44.
63. Rosenfeld PJ, Brown DM, Heier JS, et al. Ranibizumab for neovascular age-related macular degeneration. *N Engl J Med* 2006;355(14):1419–31.
64. Rosenfeld PJ, Rich RM, Lalwani GA. Ranibizumab: Phase III clinical trial results. *Ophthalmol Clin North Am* 2006;19(3):361–72.
65. Dol DV, Nguyen QD, Browning DJ, et al. An Exploratory Study of the Safety, Tolerability and Biological Effect of a Single Intravitreal Administration of VEGF Trap in Patients with Diabetic Macular Edema ARVO 2007.