



(12) **United States Patent**
Evans et al.

(10) **Patent No.: US 6,355,245 B1**
(45) **Date of Patent: Mar. 12, 2002**

(54) **C5-SPECIFIC ANTIBODIES FOR THE
TREATMENT OF INFLAMMATORY
DISEASES**

(75) Inventors: **Mark J. Evans**, Cheshire; **Louis A. Matis**, Southport; **Eileen Elliott Mueller**, East Haven, all of CT (US); **Steven H. Nye**, Mequon, WI (US); **Scott Rollins**, Monroe, CT (US); **Russell P. Rother**; **Jeremy P. Springhorn**, both of Cheshire, CT (US); **Stephen P. Squinto**, Bethany, CT (US); **Thomas C. Thomas**, Madison, CT (US); **James A. Wilkins**, Woodbridge, CT (US)

(73) Assignee: **Alexion Pharmaceuticals, Inc.**, Cheshire, CT (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **08/487,283**

(22) Filed: **Jun. 7, 1995**

Related U.S. Application Data

(63) Continuation of application No. PCT/US95/05688, filed on May 1, 1995, which is a continuation-in-part of application No. 08/236,208, filed on May 2, 1994, now Pat. No. 6,074,642.

(51) **Int. Cl.**⁷ **A61K 39/395**; C07K 16/36; C12N 5/12

(52) **U.S. Cl.** **424/145.1**; 424/130.1; 424/133.1; 424/135.1; 424/141.1; 424/145.1; 424/158.1; 424/139.1; 424/130; 530/387.1; 530/387.3; 530/387.9; 530/388.1; 530/388.23; 530/388.25; 530/388.7; 435/326; 435/328; 435/331; 435/33 L; 435/337; 435/343; 435/346

(58) **Field of Search** 530/387.1, 387.3, 530/388.1, 388.7, 387.9, 388.23; 435/69.1, 172.3, 328, 343, 70.21, 325, 331, 332, 346; 424/130.1, 141.1, 139, 133.1, 145.1

(56) **References Cited**

U.S. PATENT DOCUMENTS

4,578,335 A	3/1986	Urdal et al.	435/68
4,686,100 A	8/1987	Raffin et al.	424/85
4,816,397 A	3/1989	Boss et al.	435/68
4,816,565 A	3/1989	Honjo et al.	530/351
4,816,567 A	3/1989	Cabilly et al.	530/387
4,845,198 A	7/1989	Urdal et al.	530/387
4,867,973 A	9/1989	Goers et al.	424/85.91
5,135,916 A	8/1992	Sims et al.	514/21
5,173,490 A	12/1992	Sindelar et al.	514/462
5,198,359 A	3/1993	Taniguchi et al.	435/252.3
5,225,539 A	7/1993	Winter	530/387.3
5,395,760 A	3/1995	Smith et al.	435/240.1
5,506,247 A	4/1996	Sindelar et al.	514/374
5,530,101 A *	6/1996	Queen et al.	
5,585,089 A	12/1996	Queen et al.	424/133.1
5,610,279 A	3/1997	Brockaus et al.	530/387.3

5,635,178 A	6/1997	Sims et al.	424/145.1
5,693,761 A	12/1997	Queen et al.	536/23.53
5,693,762 A	12/1997	Queen et al.	530/387.3
5,808,029 A	9/1998	Brockaus et al.	536/23.5
5,859,205 A	1/1999	Adair et al.	530/387.3

OTHER PUBLICATIONS

Fitch et al. *Circulation* 100: 2499–2506 (1999).*

Rollins et al. *Immunopharmacology* 49: 69 (2000).*

Jain et al. *Arthritis and Rheumatism* 42: 577 (1999).*

Rollins et al. *Molecular Immunology* 35: 397 (1998).*

Wurzner et al. *Complement. Inflamm.* 8:328–340 (1991).*

Ames et al., “Isolation of neutralizing anti-C5a monoclonal antibodies from a filamentous phage monovalent fab display library” *J. Immunol.* 152:4572–4581, 1994.

Auda et al., “Measurement of complement activation products in patients with chronic rheumatic diseases” *Rheumatol Int* 10:185–189, 1990.

Baker et al., “Depletion of C6 prevents development of proteinuria in experimental membranous nephropathy in rats” *Am J Path* 135:185–194, 1989.

Bhakdi et al., “Electroimmunoassay-immunoblotting (EIA-IB) for utilization of monoclonal antibodies in quantitative immunoelectrophoresis: the method and its applications” *J Immunol Methods*, 80:25–32, 1985.

Biesecker et al., “The release of C5a in complement-activated serum does not require C6” *J Immunol*, 143:1228–1232, 1989.

Cochrane et al., “A role of polymorphonuclear leukocytes and complement in nephrotoxic nephritis” *J Exp Med*, 122:99–116, 1965.

Couser et al., “C6 depletion reduces proteinuria in experimental nephropathy induced by a nonglomerular antigen” *J Am Soc Nephrol*, 2:894–901, 1991.

Couser et al., “Complement and the direct mediation of immune glomerular injury: A new perspective” *Kidney Internat*, 28:879–890, 1985.

Couser et al., “Role of C_{5b-9} in experimental membranous nephropathy” *Nephrol Dial Transplant*, Suppl 1:25–31, 1992.

Dalmaso et al., “Complement channels in membranes: inhibition with a monoclonal antibody to a neoantigen of polymerized C9” *Biochem Biophys Res Commun*, 125:1013–1019, 1984.

(List continued on next page.)

Primary Examiner—Phillip Gambel

(57) **ABSTRACT**

The use of anti-C5 antibodies, e.g., monoclonal antibodies, to treat glomerulonephritis (GN) is disclosed. The administration of such antibodies at low dosage levels has been found to significantly reduce glomerular inflammation/enlargement and other pathologic conditions associated with GN. Also disclosed are anti-C5 antibodies and anti-C5 antibody-encoding nucleic acid molecules. These antibodies are useful in the treatment of GN and other inflammatory conditions involving pathologic activation of the complement system.

23 Claims, 19 Drawing Sheets

OTHER PUBLICATIONS

- Discipio et al., "The activation of human complement component C5 by a fluid phase C5 convertase." 258(17), 10629-10636, 1983.
- Discipio et al., "The conversion of human complement component C5 into fragment C5b by the alternative-pathway C5 convertase" *Biochem J*, 199:497-504, 1981.
- Discipio, "Formation and structure of the C5b-7 complex of the lytic pathway of complement" *J Biol Chem* 267:17087-17094, 1992.
- Falk and Jennette, "Immune complex induced glomerular lesions in C5 sufficient and deficient mice" *Kidney Internat*, 30:678-686, 1986.
- Floege et al., "Markers of complement-dependent and complement-independent glomerular visceral epithelial cell injury in vivo" *Lab Invest*, 67:486-497, 1992.
- Frei et al., "Generation of a monoclonal antibody to mouse C5 application in an ELISA assay for detection of anti-C5 antibodies" *Mol Cell Probes*, 1:141-149, 1987.
- Garrad et al., "Synthesis of C3, C5, C6, C7, C8, and C9 by human fibroblast." *Scand J. Immunol* 32(5): 555-560, 1990.
- Giclas et al., "Preparation and characterization of monoclonal antibodies against the fifth component of rabbit complement (C5)" *J Immunol Meth* 105:201-209, 1987.
- Goldman and Goldman, "Antibody-induced suppression of the fifth component of complement in mice" *J Immunol*, 120:400-407, 1978.
- Groggel et al., "Role of the terminal complement pathway in experimental membranous nephropathy in the rabbit" *J Clin Invest*, 72:1948-1957, 1983.
- Hong et al., "An anticomplement agent, K-76 monocarboxylic acid: Its site and mechanism of inhibition of the complement activation cascade." *J. Immunol* 122:2418-2433, 1979.
- Hugo et al., "Monoclonal antibodies against neoantigens of the terminal C5b-9 complex of human complement" *Bio-Science Rep*, 5:649-658, 1985.
- Hugo et al., "Sensitive ELISA for quantitating the terminal membrane C5b-9 and fluid-phase SC5b-9 complex of human complement" *J Immunol Methods* 99:243-251 1987.
- Inoue, "C5 neoepitopes appearing during activation." *Complement Inflamm* 6(3):219-222, 1989.
- Jennette et al., "Amelioration of immune complex-mediated glomerulonephritis by synthetic protease inhibitors" *Am J Path*, 127:499-506, 1987.
- Jones et al., "Replacing the complementarity-determining regions in a human antibody with those from a mouse." *Nature*, 321:522-525, 1986.
- Kitamura et al., "The activation of C5 in the fluid phase and in the absence of C3 through the classical pathway of the complement system." *Immunology* 58(3): 459-465, 1986.
- Klos et al., "Detection of native human complement components C3 and C5 and their primary activation peptides C3a and C5a (anaphylatoxic peptides) by ELISAs with monoclonal antibodies" *J Immuno Methods* 111:241-252, 1988.
- Knicker and Cochrane, "Pathogenic factors in vascular lesions of experimental serum sickness" *J Exp Med*, 122:83-97, 1965.
- Kolb and Muller-Eberhard, "The Membrane Attack Mechanism of Complement" *J Exper Med*, 141:724-735, 1975.
- Minta and Man, "Cleavage of Human C5 By Trypsin: Characterization of the Digestion Products by Gel Electrophoresis" *J Immunol*, 119:1597-1602, 1977.
- Mollnes et al., "Identification of a human C5 β -chain epitope exposed in the native complement component but concealed in the SC5b-9 complex" *Scand J. Immunol*, 28:307-312, 1988.
- Mollnes et al., "Quantification of the terminal complement complex in human plasma by an enzyme-linked immunosorbent assay based on monoclonal antibodies against a neoantigen of the complex." *Scand J Immunol*, 22(2): 197-202, 1985.
- Montz et al., "Regulation of the human autologous T cell proliferation by endogeneously generated C5a" *Cell Immunol*, 127:337-351, 1990.
- Moongkarndi et al., "Monoclonal antibodies against the fifth component of human complement" *Immunobiol* 162:397, 1982.
- Moongkarndi et al., "Immunological and functional properties of two monoclonal antibodies against human C5." *Immunobiol* 165:323, 1983.
- Morrison, "In vitro antibodies strategies for production and application" *Ann Rev Immunol* 10:239-265, 1992.
- Morgan, "Clinical complementology recent progress and future trends." *EPO J. Clinical Invesg*. 24: 219-228, 1994.
- Morgan et al., "Inhibition of complement-induced [14C] sucrose release by intracellular and extracellular monoclonal antibodies to C9: evidence that C9 is a transmembrane protein." *Biochem Biophys Res Commun*, 118(2):616-622, 1984.
- Passwell et al., "Local extrahepatic expression of complement genes C3, factor B, C2, and C4 is increased in murine lupus nephritis" *J Clin Invest*, 82:1676-1684, 1988.
- Peake et al., "Differences in the metabolism of C4 in patients with complement activation." *Clin Exp Immunol* 78:49-53, 1989.
- Perez et al., "Complement (C5)-derived chemotactic activity in serum from patients with pancreatitis" *J Lab Clin Med*, 101:123-129, 1983.
- Perez et al., "Radioimmuno-electrophoresis, a sensitive method of detecting cleavage of the fifth component of human complement (C5)" *J Immunol Methods*, 56:55-62, 1983.
- Reed et al., "Synthesis of complement component C5 by human B and T lymphoblastoid cell lines" *Immunogenetics*, 31:145-151, 1990.
- Riechmann et al., "Reshaping human antibodies for therapy." *Nature*, 332: 323-327, 1988.
- Rinder et al., "Blockade of C5 and C5b-9 generation inhibits leukocyte and platelet activation during extracorporeal circulation." *J. Clinical Invesg*. 96(3): 1564-1572.
- Rodrigues et al., "Engineering Fab' fragments for efficient F(ab)₂ formation in *Escherichia coli* and for improved in vivo stability." *J. Immunology*, 151(12): 6954-6961, 1993.
- Rottini et al., "Monoclonal antibodies as probes to investigate the molecular changes of C5 associated with the different stability of the molecule on sheep erythrocytes and *Escherichia coli* 0111:B4" *J. Immunol* 146:643-647, 1991.
- Salant et al., "A new role for complement in experimental membranous nephropathy in rats" *J Clin Invest*, 66:1339-1350, 1980.
- Schrijver et al., "Anti-GBM nephritis in the mouse: role of granulocytes in the heterologous phase" *Kidney Internat*, 38:86-95, 1990.

- Schrijver et al., "Antiglomerular basement membrane nephritis in the mouse" *Lab Invest*, 59:484-491, 1988.
- Seeger et al., "Noncytolytic terminal complement complexes may serve as calcium gates to elicit leukotriene B4 generation in human polymorphonuclear leukocytes." *J. Immunol*, 137(4): 1286-1293, 1986.
- Stahl et al., "Role of granulocytes and C5 in myocardial response to zymosan-activated serum." *Am J. Physiol*, 261(1 Pt2): H29-H37, 1991.
- Sundsmo, "Leukocyte complement: a possible role for C5 in lymphocyte stimulation" *J Immunol*, 131:886-891, 1983.
- Takeda et al., "Rapid and simple measurement of human C5a-des-Arg level in plasma or serum using monoclonal antibodies" *J Immunol Methods*, 101:265-270, 1987.
- Unanue and Dixon, "Participation of complement in nephrotoxic nephritis" *J Exp Med*, 119:965-982, 1964.
- Vogt et al., "'Inactivated' third component of complement (C3b-like C3; C3i) acquires C5 binding capacity and supports C5 activation upon covalent fixation to a solid surface" *Complement*, 1:87-96, 1984.
- Vogt et al., "Non-enzymic activation of the fifth component of human complement. by oxygen radicals. some properties of the activation product. C5b-like C5" *Mol Immunol* 26(12): 1133-1142, 1989.
- Wetsel and Kolb, "Complement-independent activation of the fifth component (C5) of human complement: limited trypsin digestion resulting in the expression of biologic activity" *J Immunol* 128:2209-2216, 1982.
- Wurzner et al., "Inhibition of terminal complement complex formation and cell lysis by monoclonal antibodies" *Complement Inflamm*, 8:328-340, 1991.
- * cited by examiner




FIG. 1A




FIG. 1B




FIG. 1C




FIG. 2A




FIG. 2B




FIG. 2C




FIG.3




FIG.4




FIG. 5A



FIG. 5B



FIG. 5C




FIG.6




FIG.7A




FIG.7B




FIG.8




FIG.9




FIG.10




FIG. 11




FIG. 12




FIG.13




FIG.14




FIG. 15




FIG. 16




FIG. 17

1	10																		20		
GAC	ATC	CAG	ATG	ACT	CAG	TCT	CCA	GCT	TCA	CTG	TCT	GCA	TCT	GTG	GGA	GAA	ACT	GTC	ACC		
D	I	Q	M	T	Q	S	P	A	S	L	S	A	S	V	G	E	T	V	T		
24				30						34				40							
ATC	ACA	TGT	GGA	GCA	AGT	GAG	AAT	ATT	TAC	GGT	GCT	TTA	AAT	TGG	TAT	CAG	CGG	AAA	CAG		
I	T	C	G	A	S	E	N	I	Y	G	A	L	N	W	Y	Q	R	K	Q		
CDR-L1																					
50										56										60	
GGA	AAA	TCT	CCT	CAG	CTC	CTG	ATC	TAT	GGT	GCA	ACC	AAC	TTG	GCA	GAT	GGC	ATG	TCA	TCG		
G	K	S	P	Q	L	L	I	Y	G	A	T	N	L	A	D	G	M	S	S		
CDR-L2																					
70														80							
AGG	TTC	AGT	GGC	AGT	GGA	TCT	GGT	AGA	CAG	TAT	TAT	CTC	AAG	ATC	AGT	AGC	CTG	CAT	CCT		
R	F	S	G	S	G	S	G	R	Q	Y	Y	L	K	I	S	S	L	H	P		
89										97										100	
GAC	GAT	GTT	GCA	ACG	TAT	TAC	TGT	CAA	AAT	GTG	TTA	AAT	ACT	CCT	CTC	ACG	TTC	GGT	GCT		
D	D	V	A	T	Y	Y	C	Q	N	V	L	N	T	P	L	T	F	G	A		
CDR-L3																					
GGG ACC AAG TTG GAG CTG AAA																					
G T K L E L K																					

FIG. 18

-19
 atg aaa tgg agc tgg gtt att ctc ttC CTC CTG TCA GTA ACT GCA GGT GTC CAC TCC CAG
 M K W S W V I L F L L S V T A G V H S Q

10
 GTT CAG CTG CAG CAG TCT GGA GCT GAG CTG ATG AAG CCT GGG GCC TCA GTG AAG ATG TCC
 V Q L Q Q S G A E L M K P G A S V K M S

26 30 35 40
 TGC AAG GCT ACT GGC TAC ATA TTC AGT AAC TAC TGG ATA CAG TGG ATA AAG CAG AGG CCT
 C K A T G Y I F S N Y W I Q W I K Q R P

CDR-H1
 50 52a 60
 GGA CAT GGC CTT GAG TGG ATT GGT GAG ATT TTA CCT GGA AGT GGT TCT ACT GAG TAC ACT
 G H G L E W I G E I L P G S G S T E Y T

CDR-H2
 65 70 80
 GAG AAC TTC AAG GAC AAG GCC GCA TTC ACT GCA GAT ACA TCC TCC AAC ACA GCC TAC ATG
 E N F K D K A A F T A D T S S N T A Y M

82a 82b 82c 90 95
 CAA CTC AGC AGC CTG ACA TCA GAG GAC TCT GCC GTC TAT TAC TGT GCA AGA TAT TTC TTC
 Q L S S L T S E D S A V Y Y C A R Y F F

100 a b c d e 102 110
 GGT AGT AGC CCC AAC TGG TAC TTC GAT GTC TGG GGC GCA GGG ACC ACG GTC ACC GTC TCC
 G S S P N W Y F D V W G A G T T V T V S

CDR-H3
 TCA
 S

FIG.19

1

C5-SPECIFIC ANTIBODIES FOR THE TREATMENT OF INFLAMMATORY DISEASES

This is a continuation in part of International application Ser. No. PCT/US95/05688, filed May 1, 1995, which is a continuation in part of U.S. application Ser. No. 08/236,208, filed May 2, 1994, which issued as U.S. Pat. No. 6,074,642 on Jun. 13, 2000. International application Serial No. PCT/US95/05688 was published in English under PCT Article 21(2) on Nov. 9, 1995 as WO 95/29697.

FIELD OF THE INVENTION

The present invention relates to the treatment of glomerulonephritis (GN) and other inflammatory diseases, and more generally to therapeutic treatments involving the pharmacologic inhibition of a patient's complement system. In particular, the invention relates to the use of antibodies specific to human complement component C5 to accomplish such therapeutic treatment. The invention also relates to compositions comprising native monoclonal antibodies (mAbs) specific to human complement component C5 that block complement hemolytic activity and C5a generation at concentrations that substantially reach the theoretical one to two stoichiometric limit of antibody to antigen that can be achieved by a bivalent antibody. The invention further provides recombinant mAbs that are derivatives (including monovalent derivatives) of these native mAbs that provide substantially the same blocking activities as the native mAbs.

BACKGROUND OF THE INVENTION

I. Immune Complex Mediated Disease

The formation of immune complexes is the typical consequence of the interaction of antigens with specific antibodies. The inflammatory response that ensues when such complexes accumulate in a limited area is an important element of normal host defenses, leading to immune complex clearance and antigen destruction by phagocytic cells. In contrast, immune complex diseases are reflections of excess complex formation or retarded clearance, usually under conditions of exceptional antigen challenge or immunologic dysregulation. Under such circumstances, immune complexes are deposited or formed at specific tissue sites and resulting inflammatory responses lead to disease states due to localized or systemic tissue damage. The kidney, and more specifically the kidney structure known as the glomerulus, is a particularly important site of immune complex deposition resulting in the development of serious disease conditions.

Human studies, and studies using animal models of human diseases, have implicated the complement system in the pathologies associated with a number of immune complex associated disorders. The activation of complement that mediates the pathology associated with these disorders may be a consequence of an autoimmune mechanism, or can be non-immunologic in origin.

The hypersensitivity response that occurs when antibodies bind to antigens either in tissues or in the circulation results from the activation of complement and the release of molecules that mediate inflammation. This process is classified as either being mediated by the binding of antibody to fixed tissue or cell bound antigens (Type II hypersensitivity) or to circulating antigens, resulting in the formation of circulating immune complexes and their subsequent pathogenic deposition in tissues (Type III hypersensitivity).

Type II hypersensitivity is mediated through the activation of complement following the binding of antibodies to

2

fixed tissue antigens. The inflammatory response that ensues results from the activation of the proinflammatory and lytic components of the complement system and the subsequent recruitment of stimulated leukocytes to the sites of immune complex formation. The increased vascular permeability that results from the anaphylatoxic activities of C3a and C5a further enhances immune complex deposition and leukocyte recruitment.

The cross-linking of antibody bound cells or tissues to effector cells such as neutrophils, platelets, NK cells, and monocytes via their Fc receptors also plays a proinflammatory role. Such cross-linking activates effector cells, stimulating the release of oxygen radicals, prostaglandins, and leukotrienes, which release is further potentiated by the actions of activated complement components.

Examples of Type II hypersensitivity-mediated conditions include hyperacute rejection of transplanted organs, autoimmune hemolytic and thrombocytopenic states, Goodpasture's syndrome (and associated glomerulonephritis and pulmonary hemorrhage), myasthenia gravis, pathologic sequelae associated with insulin-dependent diabetes mellitus, and pemphigus vulgaris.

Type III hypersensitivity reactions involving circulating antigens can also result in the development of numerous pathologic conditions. These include glomerulonephritis (discussed in detail below), vasculitis (a potentially life-threatening inflammatory condition of large and/or small blood vessels), rheumatoid arthritis, dermatitis, and other disorders.

Other diseases associated with type III hypersensitivity reactions include autoimmune diseases such as systemic lupus erythematosus (SLE), many infectious diseases, neoplastic diseases, and a wide variety of other conditions (Dixon, et al. *Immune Complex Injury*, in Samter, (ed.) *Immunological Diseases*, 4th ed. Little Brown & Co. Boston, 1987).

II. Glomerulonephritis

The glomerulus is a key structural and functional element of the kidney. Each glomerulus is found as part of a larger structure that serves as the main functional unit of the kidney and is called a nephron. About a million nephrons are found in each kidney. Each glomerulus is a network of up to fifty parallel capillaries encased in a structure known as Bowman's capsule. The area inside Bowman's capsule that is not taken up by the glomerular capillaries is known as Bowman's space. The glomerulus functions as a filter, separating water and certain solutes from the proteins and cells of the blood into Bowman's space for further processing in the convoluted tubules, loop of Henle, and collecting duct of the nephron.

Glomerulonephritis (GN) is a disease of the kidney characterized by inflammation and resulting enlargement of the glomeruli that is typically due to immune complex formation. The accumulation of immune complexes in the glomeruli results in inflammatory responses, involving inter alia hypercellularity, that can cause total or partial blockage of the glomerulus through, among other factors, narrowing of capillary lumens. One result of this process is the inhibition of the normal filtration function of the glomerulus. Blockage may occur in large numbers of glomeruli, directly compromising kidney function and often causing the abnormal deposition of proteins in the walls of the capillaries making up the glomerulus. Such deposition can, in turn, cause damage to glomerular basement membranes. Those glomeruli that are not blocked develop increased permeability, allowing large amounts of protein to pass into the urine, a condition referred to as proteinuria.

3

In many cases of severe GN, pathological structures called crescents are formed within the Bowman's space, further impeding glomerular filtration. These structures can only be seen by microscopic examination of tissue samples obtained by biopsy or necropsy, and are thus not always observed in those patients in which they occur. Crescents are a manifestation of hypercellularity and are thought to arise from the extensive abnormal proliferation of parietal epithelial cells, the cells that form the inner lining of the Bowman's capsule. Clinical research has shown that there is a rough correlation between the percentage of glomeruli with crescents and the clinical severity of the disease, and thus the patient's prognosis. When present in large numbers, crescents are a poor prognostic sign.

Symptoms of GN include: proteinuria; reduced glomerular filtration rate (GFR); serum electrolyte changes including azotemia (uremia, excessive blood urea nitrogen—BUN) and salt retention, leading to water retention resulting in hypertension and edema; hematuria and abnormal urinary sediments including red cell casts; hypoalbuminemia; hyperlipidemia; and lipiduria.

In 1990, over 210,000 patients in the United States required hemodialysis or transplantation for chronic renal failure at an annual cost in excess of 7 billion dollars, according to the United States Renal Data System (USRDS). The USRDS compiles data on kidney disease in the United States in conjunction with the National Institute of Diabetes and Digestive and Kidney Diseases, Division of Kidney, Urologic, and Hematologic Diseases, of the National Institutes of Health (NIDDKD). The USRDS estimates that the costs of treatment for renal failure are now increasing by 20 percent annually.

GN is the third leading cause of death in end-stage renal disease patients, exceeded only by diabetes and hypertension. As a result, there is a clear and long felt need in the medical community for effective treatments for this condition. Research aimed at the development of new treatments for GN is ongoing worldwide. In the United States, the NIDDKD, the National Kidney Foundation, and several other public and private organizations sponsor research in this area. The National Kidney Foundation alone supplies over two million dollars annually to fund the efforts of kidney researchers.

III. Current Treatments for GN

Corticosteroid administration, typically as high doses of "pulse" intravenous methylprednisolone or oral prednisone therapy, is currently considered the most effective pharmacologic agent available for the treatment of GN. Such steroid therapy is often administered in combination with cytotoxic general immunosuppressive agents such as azathioprine or cyclophosphamide. The overall immune suppression and resulting increased susceptibility to infection, along with other debilitating side effects associated with both steroid and cytotoxic drug administration, limit the effective use of these drugs.

Aspirin-like non-steroidal anti-inflammatory drugs (NSAIDs) have also been used to reduce the glomerular inflammation and enlargement of GN. These drugs are not routinely used for this purpose, however, probably because of their relatively weak anti-inflammatory effects and propensity to cause gastrointestinal and other side effects in many patients.

The administration of anticoagulants such as heparin or warfarin sodium, and antithrombotic agents such as cyproheptadine, dipyridamole, or sulfapyrazone, has been used on the basis of evidence suggesting the involvement of the coagulation process in the genesis of glomerular cres-

4

cents. However, objective evidence of benefit from such therapies in animals afflicted with experimentally induced crescentic GN has been inconsistent. Also, anticoagulants are dangerous drugs, as they can potentiate life-threatening bleeding episodes. They are especially hazardous in this regard in patients with advanced renal failure.

In addition to pharmacologic approaches, intensive plasma exchange (plasmapheresis) of 2 to 4 liters of plasma daily (or in some cases three times a week) can dramatically reduce high levels of circulating immune complexes when acute intervention in the inflammatory process is needed. Such treatment is expensive and requires that the patient be connected to the plasmapheresis machine for many hours each week. In addition, all procedures in which blood is removed from and returned to a patient are associated with an increased risk of infection. Nonetheless, plasma exchange is currently considered the most effective non-pharmacological treatment for removal of circulating immune complexes which can cause GN.

Circulating immune complex levels can also be decreased by eliminating or reducing the source of the antigen or antigens contained in the complexes by, for example, effective therapy of an underlying infection or change in an antibiotic. However, while such therapy is almost always a treatment of choice, great care must be taken since reduction of the antigen load alters the molar ratio of antigen to antibody involved in forming immune complexes and thus a dangerous temporary exacerbation of the inflammatory process may occur (see discussion below in Background Physiology & Pathology).

IV. Antibody Engineering

Native antibodies are multi-subunit animal protein molecules with highly specific antigen-binding properties. Animals make multiple classes of antibodies. There are five major classes (IgA, IgD, IgE, IgG and IgM) and a variety of subclasses. Native antibodies are made up of two or more heterodimeric subunits each containing one heavy (H) and one light (L) chain. The differences between antibody classes derive from their different H chains. H chains have a molecular weight of about 53 kDa, while L chains are about 23 kDa in mass.

Every individual native antibody has one type of L chain and one type of H chain, which are held together by disulfide bonds to form a heterodimeric subunit. Typically a native antibody (e.g., an IgG) has two such subunits, which are also held together by disulfide bonds. Within each chain, units of about 110 amino acid residues fold so as to form compact domains. Each domain is held together by a single intrachain disulfide bond. L chains have two domains, while H chains have four or five. Most H chains have a hinge region after the first (i.e., most amino-terminally located) two domains. The disulfide bonds linking together the heterodimeric subunits are located at the hinge regions. The hinge region is particularly sensitive to proteolytic cleavage, such proteolysis yielding two or three fragments (depending on the precise site of cleavage), a non-antigen binding fragment containing only H chain C regions (Fc) and one bivalent (Fab') or two monovalent (Fab) antigen binding fragments. The hinge region allows the antigen binding regions (each made up of a light chain and the first two domains of a heavy chain) to move freely relative to the rest of the native antibody, which includes the remaining heavy chain domains.

The first domain of each chain is highly variable in amino acid sequence, providing the vast spectrum of antibody binding specificities found in each individual. These are known as variable heavy (VH) and variable light (VL)

5

domains. The second and subsequent (if any) domains of each chain are relatively invariant in amino acid sequence. These are known as constant heavy (CH) and constant light (CL) domains.

Each variable region contains three loops of hypervariable sequence that provide a complementary structure to that of the antigen and are critical in determining the antigen binding specificity of the antibody, as they are the contact sites for binding to the antigen. These loops are known as complementarity determining regions, or CDRs. Each variable domain is made up of three CDRs embedded in four much less variable framework segments (FRs). Together, the sets of collinear CDRs and FRs are in large part responsible for determining the three dimensional conformation of the variable regions of antibody molecules.

CDRs and FRs are features that have been deduced from structural properties of antibody variable regions. Both amino acid sequence (primary structure) and three dimensional modeling (deduced secondary and tertiary structure) of antibody variable regions have been used by various researchers to define CDRs and, by default, FRs. While the positions of the CDRs are beyond question, not all workers in the art agree upon the precise locations of the boundaries of each CDR in VH or VL regions; there is no clear cut structural marker delineating CDR/FR boundaries.

Two definitions of CDR location are currently in general use in the art. These are the "sequence variability" definition of Kabat et al. ("Sequences of Proteins of Immunological Interest," 4th ed. Washington, D.C.: Public Health Service, N.I.H.) and the "structural variability" definition of Chothia and Lesk (J. Mol. Biol. 1987, 196:901). As used herein, the terms VL CDR1, VL CDR2, VL CDR3, VH CDR1, VH CDR2, and VH CDR3 refer minimally to the region of overlap between the regions designated for each CDR by each of these two definitions, and maximally to the total region spanned by the combination of the regions designated for each CDR by each of these two definitions.

One problem that antibody engineering attempts to address is the immune activity of a human patient that occurs in response to a native murine (or other non-human animal) antibody, typically a mAb, that is being administered to the patient for therapeutic purposes. This activity against murine antibodies is characterized by a human anti-mouse antibody (HAMA) response that can have deleterious effects on treatment efficacy and patient health. It has been found that almost all such human anti-non-human antibody ("HAMA type") activity is directed at the constant domains and at the FR regions of the variable domains of native non-human antibodies.

By manipulating the nucleic acid molecules encoding antibody H and L chains it is possible to incorporate non-human variable regions into antibodies otherwise made up of human constant regions. The resulting antibodies are referred to as "chimeric antibodies," and are typically less prone to eliciting HAMA type responses than are the non-human antibodies from which the variable regions are derived.

An even more effective approach to eliminating the potential of a non-human antibody to elicit a HAMA type response is to "humanize" it, i.e., to replace its non-human framework regions with human ones. One way of achieving such humanization involves the insertion of polynucleotide fragments encoding the non-human CDRs of the antibody to be humanized into a nucleic acid molecule encoding an otherwise human antibody (with human constant regions if desired) so as to replace the human CDRs and to use the resulting nucleic acid molecule to express the encoded "humanized" antibody.

6

Unfortunately, however, humanization of non-human antibodies has unpredictable effects on antibody antigen interactions, e.g., antigen binding properties. Some of this unpredictability stems from the properties of the CDRs. Certain CDRs may be more amenable to the construction of humanized antibodies that retain the properties of the non-human CDR donor antibody than others. While the CDRs are key to the antigen binding properties of an antibody, CDRs and FRs must interact appropriately if the antigen specificity of an antibody is to be retained following humanization. The effects of combination with particular human FRs on uncharacterized non-human CDRs cannot be reliably predicted by any known method. However, the successful humanization of an antibody provides information that, in general, facilitates the successful humanization of the CDRs of that antibody using other human or altered human FRs. In addition, approaches are available that facilitate tailoring human FRs to enhance the likelihood of successful humanization.

Other problems addressed by antibody engineering include efficient antibody production and alteration of antibody pharmacokinetics. Recombinant protein production is generally most efficiently carried out in bacterial hosts. The large size and multimeric nature of native antibodies makes their production in bacteria difficult. One approach to dealing with production problems is to use recombinant DNA methods to construct antibodies that have their H and L chains joined by a linker peptide to form a single chain (sc) antibody. As described below, there are several types of sc antibodies that can be constructed.

As is the case for humanization, the effects on antigen binding properties of constructing a particular type of sc antibody using H and L chains that have not been characterized with regard to their ability to function as part of an sc antibody cannot be reliably predicted by any known method. However, the successful construction of any one type of sc antibody from a particular native antibody provides information that, in general, facilitates the successful construction of other types of sc antibodies from that native antibody.

Single chain antibodies may include one each of only VH and VL domains, in which case they are referred to as scFv antibodies; they may include only one each of VH, VL, CH, and CL domains, in which case they are referred to as scFab antibodies; or they may contain all of the variable and constant regions of a native antibody, in which case they are referred to as full length sc antibodies.

The differing sizes of these antibodies imparts each with differing pharmacokinetic properties. In general, smaller proteins are cleared from the circulation more rapidly than larger proteins of the same general composition. Thus, full length sc antibodies and native antibodies generally have the longest duration of action, scFab antibodies have shorter durations of action, and scFv antibodies have even shorter durations of action. Of course, depending upon the illness being treated, longer or shorter acting therapeutic agents may be desired. For example, therapeutic agents for use in the prevention of immune and hemostatic disorders associated with extracorporeal circulation procedures (which are typically of brief duration) are preferably relatively short acting, while antibodies for the treatment of long term chronic conditions (such as inflammatory joint disease or GN) are preferably relatively long acting.

Detailed discussions of antibody engineering may be found in numerous recent publications including: Borrebaek, "Antibody Engineering, A Practical Guide," 1992, W.H. Freeman and Co. NY; and Borrebaek, "Antibody Engineering," 2nd ed. 1995, Oxford University Press, NY, Oxford.

SUMMARY OF THE INVENTION

In view of the foregoing, it is an object of the present invention to provide a new approach for reducing the glomerular inflammation and kidney dysfunction associated with GN.

The method of the invention involves the use of preparations containing antibodies to human complement component C5 as pharmaceutical agents. More particularly, the invention provides for the use of anti-C5 antibodies that bind to complement component C5 or active fragments thereof. Preferably, the antibodies block the generation and/or activity of complement components C5a and C5b. For most applications, the antibody is a monoclonal antibody.

In the preferred embodiments of the invention, the administration of the anti-C5 antibody preparation is started after the appearance of GN symptoms, e.g., after the appearance of proteinuria. Alternatively, the invention can be used prophylactically to treat patients who are at risk for an acute exacerbation of existing GN, e.g., patients experiencing a flare-up of symptoms of systemic lupus erythematosus or similar autoimmune diseases that have resulted in GN.

As shown in the examples presented below, anti-C5 antibodies administered subsequent to the onset of GN essentially eliminate glomerular inflammation/enlargement and reduce kidney dysfunction (see Examples 1 and 2).

Although not wishing to be bound by any particular theory of operation, it is believed that the anti-C5 antibodies have these and other therapeutic effects through their activity in blocking the generation or activity of the C5a and/or C5b active fragments of complement component C5. Through this blocking effect, the antibodies inhibit the proinflammatory (anaphylatoxic) effects of C5a and the generation of the C5b-9 membrane attack complex (MAC). Significantly, the blockage effected by the anti-C5 antibodies, since it occurs at the level of complement component C5, has the advantage of maintaining important opsonic, anti-infective, and immune complex clearance functions of the complement system mediated by, inter alia, complement component C3.

The invention additionally provides compositions comprising anti-C5 antibodies that block complement hemolytic activity and C5a generation. These antibodies are useful for the treatment of GN as well as a number of other conditions. These include treatment of immune and hemostatic dysfunctions associated with extracorporeal circulation (see copending U.S. patent application Ser. No. 08/217,391, now U.S. Pat. No. 5,853,722 which is incorporated herein by reference), treatment of inflammatory joint diseases (see copending U.S. patent application Ser. No. 08/311,489, which is incorporated herein by reference), and other complement associated conditions, particularly inflammatory diseases.

Although other antibodies can be used to treat GN in accordance with the present invention, the novel antibodies of the invention are preferred. Preferably, these novel antibodies bind to the alpha chain of C5, but do not exhibit substantial binding to the alpha chain cleavage product C5a (referred to hereinafter and in the claims as "free C5a"). Other preferred targets for antibody binding include fragments of the alpha chain of human C5 that are immunoreactive with the most preferred antibody of the invention, the 5G1.1 antibody discussed below. Such preferred targets include the 46 kDa acid hydrolysis fragment of C5 (the "5G46k" fragment), the 27 kDa tryptic digestion fragment of C5 (the "5G27k" fragment), the 325aa peptide spanning amino acid residues 725-1049 of SEQ ID NO:2 (the

"5G325aa" peptide), the 200 amino acid peptide spanning amino acids residues 850 to 1049 of SEQ ID NO:2 (the "5G200aa" peptide)—as discussed below in Example 13.

The novel antibodies of the invention include antibodies that bind to an epitope within the amino acid sequence Val Ile Asp His Gln Gly Thr Lys Ser Ser Lys Cys Val Arg Gln Lys Val Glu Gly Ser Ser, (SEQ ID NO:1) hereinafter referred to as the KSSKC epitope. These novel antibodies that bind to the KSSKC epitope (SEQ ID NO:1) are hereinafter referred to as anti-KSSKC antibodies, and monoclonal antibodies binding to the KSSKC epitope are hereinafter referred to as anti-KSSKC mAbs.

The novel antibodies of the invention have many advantages over other anti-C5 antibodies, particularly with regard for their use as anti-inflammatory therapeutic agents. These include the ability to substantially block both complement hemolytic activity and the generation of the proinflammatory complement cleavage product C5a to substantially the same extent at the same concentration of antibody. Some of the preferred antibodies of the invention have the additional advantageous property of blocking the binding of C5 to C3 or C4.

Particularly preferred antibodies of the invention are monospecific native anti-KSSKC antibodies. The 5G1.1 native anti-KSSKC mAb has the distinct advantage of substantially blocking both complement hemolytic activity and the generation of C5a at a stoichiometric ratio of antibody to C5 that approaches the theoretical one to two (antibody to antigen) limit of binding that can be achieved by a bivalent antibody. This is a desirable property because it allows smaller doses of antibody to achieve therapeutic effects than would be required of otherwise similar antibodies that cannot function at such a ratio.

The invention further provides recombinant mAbs that are derivatives (including monovalent derivatives) of these native mAbs. These include anti-KSSKC recombinant mAbs. Preferably the antibodies of the invention provide a level of blockade of both complement hemolytic activity and C5a generation (on a per mole of binding site basis) that is obtained when the antibody concentration is within an order of magnitude of that of the native mAbs. Particularly preferred anti-KSSKC recombinant mAbs provide a level of such blockade when the antibody concentration is no more than three fold that of the native mAbs of the invention.

The invention further provides nucleic acid sequences of polynucleotides encoding such recombinant anti-KSSKC mAbs, as well as amino acid sequences of the polypeptides encoded by these nucleic acid molecules of the invention.

The invention further provides CDR sequences that are useful in the construction of the humanized antibodies of the invention, as well as peptides and oligopeptides that are useful in the preparation and characterization of the antibodies of the invention.

Anti-C5 antibodies of the invention have activity in blocking the generation or activity of the C5a and/or C5b active fragments of complement component C5. Through this blocking effect, the antibodies inhibit the proinflammatory (anaphylatoxic) effects of C5a and the generation of the C5b-9 membrane attack complex (MAC). Significantly, the blockage effected by the anti-C5 antibodies, since it occurs at the level of complement component C5, has the advantage of maintaining important opsonic, anti-infective, and immune complex clearance functions of the complement system mediated by, inter alia, complement component C3.

The accompanying figures, which are incorporated in and constitute part of the specification, illustrate certain aspects

of the invention, and together with the description, serve to explain the principles of the invention. It is to be understood, of course, that both the figures and the description are explanatory only and are not restrictive of the invention.

BRIEF DESCRIPTION OF THE DRAWINGS

FIGS. 1A, 1B, and 1C—Photomicrographs of PAS stained sections of mouse kidneys. FIG. 1A—uninduced untreated mouse. FIG. 1B—GN-induced PBS-(control)-treated mouse. FIG. 1C—GN-induced anti-C5 treated mouse. Magnification for each is the same, approximately 400x.

FIGS. 2A, 2B, and 2C—Photomicrographs of immunofluorescence stained sections of mouse kidneys. FIG. 2A—uninduced untreated mouse. FIG. 2B—GN-induced PBS-(control)-treated mouse. FIG. 2C—GN-induced anti-C5 treated mouse. Magnification for each is the same, approximately 200x.

FIG. 3—Results of hemolytic (cell lysis) assays of serum from GN-induced animals treated with either anti-C5 antibodies in PBS (“Anti-C5”) or PBS alone (“PBS control”). Also shown are the results of assays performed with normal serum.

FIG. 4—Results of soluble C5b-9 (“sC5b-9”) assays. “ND” indicates not determined.

FIGS. 5A, 5B, and 5C—Immunofluorescence photomicrographs of kidney sections stained for mouse C3. FIG. 5A—uninduced untreated mouse. FIG. 5B—GN-induced PBS-(control)-treated mouse. FIG. 5C—GN-induced anti-C5 treated mouse. Magnification for each is the same, approximately 400x.

FIG. 6—Results of C3a assays of samples of circulating human blood. “ND” indicates not determined.

FIGS. 7A and 7B—Pharmacokinetic analyses of the reduction of the cell lysis ability of mouse (FIG. 7A) or human (FIG. 7B) blood after treatment with anti-C5 antibodies.

The immunofluorescent staining of FIGS. 2 and 5 is confined to the glomerular capillary network (tuft) and thus the enlargement of the glomerulus seen in FIG. 1B is not visible in FIGS. 2B and 5B.

FIG. 8—Scatchard analysis of native 5G1.1 binding to C5.

FIG. 9—Scatchard analysis of native N19/8 binding to C5.

FIG. 10—C3a generation in samples of circulating human blood in the presence of native 5G1.1.

FIG. 11—sC5b-9 generation in samples of circulating human blood in the presence of native 5G1.1.

FIG. 12—Serum hemolytic activity of samples of circulating human blood in the presence of native 5G1.1.

FIG. 13—Serum hemolytic activity in the presence of m5G1.1 scFv.

FIG. 14—C5a generation in the presence of m5G1.1 scFv.

FIG. 15—C3a generation in samples of circulating human blood in the presence of m5G1.1 scFv.

FIG. 16—Serum hemolytic activity of samples of circulating human blood in the presence of 5G1.1 scFv.

FIG. 17—sC5b-9 generation in samples of circulating human blood in the presence of m5G1.1 scFv.

FIG. 18—The light chain variable region of the antibody 5G1.1. Sequence derived from the 5' oligonucleotide primer used for PCR amplification of the variable region is shown

in lower case. Amino acids are numbered according to Kabat et al., supra. Boxed amino acids correspond to peptide sequences obtained from the mature 5G1.1 light chain or from an endoproteinase Lys C peptide of 5G1.1. The complementarity determining region (CDR) residues according to the sequence variability definition and the structural variability definition are underlined and overlined, respectively.

FIG. 19—The heavy chain variable region of the antibody 5G1.1. Sequence derived from the 5' oligonucleotide primer used for PCR amplification of the variable region is shown in lower case. Amino acids are numbered using the scheme of Kabat et al. supra with +1 denoting the first amino acid of the processed mature variable region. Boxed amino acids correspond to peptide sequence obtained from the 5G1.1 heavy chain after treatment with pyroglutamate aminopeptidase. The complementarity determining region (CDR) residues according to the sequence variability definition or according to the structural variability definition are underlined and overlined, respectively.

BACKGROUND PHYSIOLOGY & PATHOLOGY

The discussion in this section is not limited to subject matter that qualifies as “prior art” against the present invention. Therefore, no admission of such prior art status shall be implied or inferred by reason of inclusion of particular subject matter in this discussion, and no declaration against the present inventors’ interests shall be implied by reason of such inclusion.

I. Introduction

As described above, the present invention relates to therapeutic treatments for GN and other immune complex mediated diseases, as well as to the treatment of other complement mediated diseases and to the inhibition of complement component C5. To provide background for the description of the preferred embodiments and the examples presented below, we turn first to general discussions of the complement arm of the immune system, the pathophysiologic features of GN, and previous studies of the role of complement in GN pathogenesis.

General discussions of the complement system and GN can be found in, for example, Glasscock and Brenner, 1994; Couser, 1993; Couser, 1992; Couser, et al, 1992; Rich, 1992; Glasscock and Brenner, 1987; Robbins and Cotran, 1979; and Guyton, 1971.

II. The Complement System

The complement system acts in conjunction with other immunological systems of the body to defend against intrusion of cellular and viral pathogens. There are at least 25 complement proteins, which are found as a complex collection of plasma proteins and membrane cofactors. The plasma proteins make up about 10% of the globulins in vertebrate serum. Complement components achieve their immune defensive functions by interacting in a series of intricate but precise enzymatic cleavage and membrane binding events. The resulting complement cascade leads to the production of products with opsonic, immunoregulatory, and lytic functions.

The complement cascade progresses via the classical pathway or the alternative pathway. These pathways share many components, and while they differ in their initial steps, they converge and share the same “terminal complement” components (C5 through C9) responsible for the activation and destruction of target cells.

The classical complement pathway is typically initiated by antibody recognition of and binding to an antigenic site on a target cell. The alternative pathway is usually antibody

independent, and can be initiated by certain molecules on pathogen surfaces. Both pathways converge at the point where complement component C3 is cleaved by an active protease (which is different in each pathway) to yield C3a and C3b. Other pathways activating complement attack can act later in the sequence of events leading to various aspects of complement function.

C3a is an anaphylatoxin (see discussion below). C3b binds to bacterial and other cells, as well as to certain viruses and immune complexes, and tags them for removal from the circulation. (C3b in this role is known as opsonin.) The opsonic function of C3b is considered to be the most important anti-infective action of the complement system. Patients with genetic lesions that block C3b function are prone to infection by a broad variety of pathogenic organisms, while patients with lesions later in the complement cascade sequence, i.e., patients with lesions that block C5 functions, are found to be more prone only to *Neisseria* infection, and then only somewhat more prone (Fearon, in *Intensive Review of Internal Medicine*, 2nd Ed. Fanta and Minaker, eds. Bringham and Women's and Beth Israel Hospitals, 1983).

C3b also forms a complex with other components unique to each pathway to form classical or alternative C5 convertase, which cleaves C5 into C5a and C5b. C3 is thus regarded as the central protein in the complement reaction sequence since it is essential to both the alternative and classical pathways (Wurzner, et al., *Complement Inflamm.* 8:328-340, 1991). This property of C3b is regulated by the serum protease Factor I, which acts on C3b to produce iC3b. While still functional as opsonin, iC3b cannot form an active C5 convertase.

C5 is a 190 kDa beta globulin found in normal serum at approximately 75 $\mu\text{g/ml}$ (0.4 μM). C5 is glycosylated, with about 1.5-3 percent of its mass attributed to carbohydrate. Mature C5 is a heterodimer of a 999 amino acid 115 kDa alpha chain that is disulfide linked to a 656 amino acid 75 kDa beta chain. C5 is synthesized as a single chain precursor protein product of a single copy gene (Haviland et al. *J. Immunol.* 1991, 146:362-368). The cDNA sequence of the transcript of this gene predicts a secreted pro-C5 precursor of 1659 amino acids along with an 18 amino acid leader sequence (SEQ ID NO:2).

The pro-C5 precursor is cleaved after amino acid 655 and 659, to yield the beta chain as an amino terminal fragment (amino acid residues +1 to 655 of SEQ ID NO:2) and the alpha chain as a carboxyl terminal fragment (amino acid residues 660 to 1658 of SEQ ID NO:2), with four amino acids (amino acid residues 656-659 of SEQ ID NO:2) deleted between the two.

C5a is cleaved from the alpha chain of C5 by either alternative or classical C5 convertase as an amino terminal fragment comprising the first 74 amino acids of the alpha chain (i.e., amino acid residues 660-733 of SEQ ID NO:2). Approximately 20 percent of the 11 kDa mass of C5a is attributed to carbohydrate. The cleavage site for convertase action is at or immediately adjacent to amino acid residue 733 of SEQ ID NO:2. A compound that would bind at or adjacent to this cleavage site would have the potential to block access of the C5 convertase enzymes to the cleavage site and thereby act as a complement inhibitor.

C5 can also be activated by means other than C5 convertase activity. Limited trypsin digestion (Minta and Man, *J. Immunol.* 1977, 119:1597-1602; Wetsel and Kolb, *J. Immunol.* 1982, 128:2209-2216) and acid treatment (Yammamoto and Gewurz, *J. Immunol.* 1978, 120:2008; Damerau et al., *Molec. Immunol.* 1989, 26:1133-1142) can also cleave C5 and produce active C5b.

C5a is another anaphylatoxin (see discussion below). C5b combines with C6, C7, and C8 to form the C5b-8 complex at the surface of the target cell. Upon binding of several C9 molecules, the membrane attack complex (MAC, C5b-9, terminal complement complex—TCC) is formed. When sufficient numbers of MACs insert into target cell membranes the openings they create (MAC pores) mediate rapid osmotic lysis of the target cells. Lower, non-lytic concentrations of MACs can produce other effects. In particular, membrane insertion of small numbers of the C5b-9 complexes into endothelial cells and platelets can cause deleterious cell activation. In some cases activation may precede cell lysis.

As mentioned above, C3a and C5a are anaphylatoxins. These activated complement components can trigger mast cell degranulation, which releases histamine and other mediators of inflammation, resulting in smooth muscle contraction, increased vascular permeability, leukocyte activation, and other inflammatory phenomena including cellular proliferation resulting in hypercellularity. C5a also functions as a chemotactic peptide that serves to attract pro-inflammatory granulocytes to the site of complement activation.

III. Pathophysiology of GN

Although GN may accompany an extraordinary range of pathologic processes, in general it is encountered most commonly in the course of infectious diseases, in autoimmunity, and as a consequence of therapy for some other disease process. The causative mechanism for GN is typically the deposit of circulating immune complexes in the kidney. Factors involved in the pathogenesis of GN include the specific antigen and antibody involved and the inflammatory processes that occur as a consequence of immune complex deposition.

Antigens Involved in the Formation of Immune Complexes that Cause GN

Antigens involved in the development of GN can be broadly classified as endogenous, infectious, and iatrogenic (those encountered as a consequence of medical practice). In many cases the specific antigen is unknown, although the general class can usually be identified.

The best known example of the formation of endogenous immune complexes is the DNA anti-DNA complexes produced in connection with systemic lupus erythematosus (lupus, SLE). Other important sources of endogenous antigens include malignancies in which immune complex formation may contribute to the development of paraneoplastic syndromes.

Infections with organisms of many types, particularly chronic infections, are also associated with the development of immune complexes that can cause GN. Bacterial and fungal infections that can produce such complexes include infection with certain strains of streptococci, *Pseudomonas*, disseminated gonococcal infection, lepromatous leprosy, subacute bacterial endocarditis, bronchopulmonary aspergillosis, secondary syphilis, and chronic infections in patients with cystic fibrosis.

Viral diseases in which immune complex deposition may be a prominent feature include hepatitis B infection, dengue, infectious mononucleosis, and subacute sclerosing panencephalitis. GN is also a prominent feature of many parasitic infestations such as the GN seen in children with quartan malaria, as well as toxoplasmosis, trypanosomiasis, and schistosomiasis.

Iatrogenic antigens constitute a special class of exogenous antigens. These include those responsible for the prototype immune complex disease, serum sickness, which follows

formation of immune complexes between heterologous serum constituents and autologous antibodies. Serum sickness was regularly seen earlier in this century when infectious diseases were frequently treated with heterologous antisera.

An iatrogenic disease essentially indistinguishable from classic serum sickness can occur as a consequence of high-dose antibiotic therapy. The serum sickness-like manifestations of immune responses to these drugs include GN and reflect the fact that certain drugs, particularly the β -lactam and sulfonamide antibiotics, are effective haptens that are capable of inducing antibody responses upon spontaneous conjugation to autologous proteins.

Factors Affecting Immune Complex Formation and Deposition

Features of both antigen and antibody determine the likelihood of pathologic immune complex formation and subsequent deposition in the kidney. Chief among these are the absolute concentrations of the reactants and their relative molar ratios.

Most antigens display multiple epitopes and typically stimulate a polyclonal antibody response. All naturally occurring antibody molecules are at least bivalent. These properties allow for the formation of an extensive antigen-antibody lattice, the size of which is determined largely by the affinity of the antibodies and the molar ratio of antigen to antibody.

In general, antibody responses begin under conditions in which antigen is present in excess to antibody, and this relative ratio changes as the antibody response increases in magnitude. Complexes formed initially are usually small and exhibit little or no pathogenic activity. In contrast, very large complexes are often formed as the amount of antigen becomes limiting, late in the course of an antibody response under conditions of antibody excess. Because these very large complexes are readily cleared by the reticuloendothelial system in the liver, they are also relatively nonpathogenic.

The formation of immune complexes that can cause GN is believed to occur during conditions of slight antigen excess or near the point of antibody-antigen equivalence, where lattice formation is maximal and lattice size is large, but not very large.

Several factors influence the speed and location of immune complex precipitation. Interactions between Fc regions of antibody molecules promote rapid precipitation of immune complexes. The role of Fc-Fc interactions in immune complex precipitation is illustrated by studies of the properties of F(ab')₂ antibody fragments, which do not contain Fc regions. Although the valence of F(ab')₂ fragments does not differ from that of most whole immunoglobulins, F(ab')₂ antibody fragments form lattices more slowly.

Antigen charge plays a role in determining the tissue localization of sites of deposition of immune complex precipitates. Complexes with a substantial positive charge are preferentially attracted to the strong negative charge of basement membranes, particularly in the renal glomerulus.

Localized presence of antigen may largely account for certain cases of organ specific immune complex deposition. Diseases such as Goodpasture's syndrome (a rare form of GN) are typically not classified as immune complex diseases because the complexes are formed in situ in the kidney rather than being preformed in the circulation and then deposited. Once the immune complexes are formed, the subsequent inflammatory process is believed to be essentially the same as that seen following deposition of pre-

formed complexes. However, the different mode of deposition distinguishes this syndrome from typical GN caused by circulating immune complexes.

Features of blood flow and vascular structure are also important in determining the localization of immune complex deposits. Chief among these is capillary permeability. Because their capillary endothelium is fenestrated, renal glomeruli are preferential sites for the deposition of immune complexes. Hemodynamic variables enhancing immune complex localization include turbulence of flow and increased blood pressure, both of which are present in the renal glomeruli.

Complement and Complement Receptors as Regulators of Immune Complex Deposition

In addition to their proinflammatory functions, complement components can also inhibit immune complex deposition and resolubilize immune complex precipitates from sites of deposition. In addition, it is known that erythrocyte receptors for C3b, e.g., CR1, are important for reticuloendothelial clearance of opsonized circulating immune complexes.

Analysis of the clinical pattern of immune complex disease in patients with deficiencies of particular complement components provides information regarding the normal role of these components in the prevention of complex deposition. The incidence of immune complex disease in patients with deficiencies of Clq, C1r, C1s, C4, C2, or C3 varies from 60 to 90 percent, with the majority of these patients exhibiting a lupus-like syndrome. Immune complex disease is rarely associated with deficiencies of late-acting or alternative pathway components.

The binding of complement components to immune complexes prevents the formation of large antigen-antibody lattices and inhibits immune precipitation. This process requires activation via the classical pathway; serum that is deficient for Clq, C4, or C2 does not effectively inhibit lattice formation and complex precipitation. Classical pathway dependence may reflect the initial binding of C1 components, impeding the Fc-Fc interactions between IgG molecules that contribute to immune precipitation. This is followed by covalent binding of C3b to the complexes, which further inhibits immune precipitation and leads to solubilization of previously deposited complexes.

The solubilization process also depends upon activation of components of the alternative pathway. Consequently, by promoting clearance of immune complexes and inhibiting their deposition at sites of inflammation, complement components and their receptors serve as negative regulators of immune complex diseases that may retard disease development.

It should be noted that the present invention involves blocking the activities of complement component C5. The targeting of this component does not alter the functions of the early complement components, and thus does not compromise the negative regulatory effects on immune complex deposition of those early components.

Immune Complex-Mediated Inflammation

Basophils are important in the initiation of immune complex-mediated inflammatory responses, as capillary permeability is markedly increased by the action of vasoactive amines such as histamine and platelet-activating factor, which are released by these cells. Vascular permeability is also promoted by aggregation of platelets at sites of an inflammatory lesion, with the release of platelet-activating factor and the formation of microthrombi.

Basophil degranulation may reflect the effects of IgE antibodies, as well as the elaboration of the anaphylatoxin components of complement, C3a and C5a.

In addition to basophils and platelets, the primary cellular effectors of immune complex-mediated inflammation are polymorphonuclear leukocytes, monocytes, and macrophages.

IV. Previous Studies of the Role of Complement in GN Pathogenesis

Extensive work has been performed in an attempt to understand the possible role of complement in the development of GN. This work has included studies of GN using a number of animal models by, among others, Unanue, et al., (1964); Cochrane, et al., (1965); Kniker, et al., (1965); Salant, et al., (1980); Groggel, et al., (1983); Falk and Jennette (1986); Jennette, et al., (1987); Passwell, et al., (1988); Schrijver, et al., (1988); Baker, et al., (1989); Schrijver, et al., (1990); Couser, et al., (1991); and Couser, et al., (1992).

These studies have shown that complement plays a role in GN pathogenesis. However, they have not established specific unequivocal roles for the various complement components. In particular, the relative roles of C3 and other anaphylatoxins compared to the roles of the terminal complement components in GN pathogenesis have not been unequivocally established. Also, some researchers have reported that complement depletion does not diminish glomerular injury. See Kniker, et al., (1965).

The foregoing work includes that of Falk and Jennette (1986), who reported results of experiments in which attempts were made to induce GN in mice having a genetic defect that resulted in a deficiency of complement component C5. The report concludes that C5 or some terminal complement component dependent on C5 plays a role in the pathogenesis of GN.

Significantly, with regard to the present invention, Falk and Jennette in no way disclose or suggest that an antibody to C5 can be used to treat GN. Indeed, it would be counterintuitive to use an antibody to treat disease which typically involves the formation and deposition of circulating antibody-antigen immune complexes. Plainly, the creation of more circulating immune complexes would seem to be the last way to go to solve a problem that can be caused by circulating immune complexes. Yet, as demonstrated by the surprising results presented below, anti-C5 antibodies have been found to effectively block GN, even though the creation of additional circulating immune complexes is inherent in their mode of action.

Baker et al. (1989), Couser et al. (1991), and Couser et al. (1992) (hereinafter referred to collectively as the "C6" work) discuss experiments in which high levels of an anti-C6 polyclonal antibody preparation were administered to rats, following which immune complexes were formed in situ in the rats' kidneys. Significantly, with regard to the present invention, the anti-C6 antibody preparation was not administered to animals with pre-existing kidney disease, i.e., it was not used as a therapeutic treatment. Moreover, the experimental protocol used in the C6 experiments did not involve circulating immune complexes, but rather involved complexes formed in situ. Accordingly, the experiments did not disclose or suggest the counterintuitive approach of the present invention wherein more circulating immune complexes are formed in the process of treating a disease state caused by circulating immune complexes.

Further, the anti-C6 antibody dosages used in the C6 work were too high for practical medical use. Specifically, these antibodies were used at a dosage of 1 gm/kg, a dosage which would correspond to 70 gm of antibody for a 70 kg (155 lb) individual. In contrast, the anti-C5 antibodies used in the practice of the present invention are used at concentrations

at or below 0.1 gm/kg, i.e., a factor of at least ten times less than used in the C6 work. Indeed, as shown by the examples presented below, anti-C5 antibody dosages as low as 0.03 gm/kg, i.e., 33 times less than those used in the C6 work, have been found to achieve the therapeutic effects of the invention in treating GN. For a 70 kg individual, this antibody level corresponds to a dose of just 2.1 gms.

The novel anti-KSSKC antibodies of the invention allow the use of even lower dosage levels to treat GN and other inflammatory conditions. Based upon their level of activity in human blood, they are expected to provide complete complement inhibition at dosages below 0.005 g/kg, and to provide therapeutically effective complement inhibition at dosages below 0.003 g/kg. This 3 mg/kg dosage is one tenth the dosage discussed below in Examples 4 and 5 for the anti-C5 (beta chain specific) mAb N19/8. Some of the full length anti-KSSKC mAbs of the invention will provide therapeutic benefits even at dosages below 0.0022 g/kg. This is the minimum dose providing complete complement inhibition as calculated from the data obtained using the anti-KSSKC 5G1.1 mAb in human blood in a CPB circuit, as discussed below in Example 9.

Accordingly, dosages of less than 0.005 g/kg are preferred, with dosages of below 0.003 g/kg being more preferred, and dosages below 0.0022 g/kg being particularly preferred. For a 70 kg individual, these antibody dosage levels correspond to a dose of less than 0.35 gms for the highest dosage of the preferred dosages, less than 0.21 gms for the more preferred dosage, and less than or equal to 0.15 gms for the most preferred dosage.

Of course, dosage levels of single chain and other recombinant mAbs of the invention must be adjusted according to their level of activity (e.g., their binding affinity, their ability to block C5 activation, and/or their ability to block complement hemolytic activity), their valency, and their molecular weight. For example, the humanized scFv anti-KSSKC mAbs of Example 11 are approximately 27 kDa, about one sixth the approximately 155 kDa mass of a native, full length IgG antibody. These antibodies completely block complement hemolytic activity and C5a generation at a ratio of 3:1, six fold greater than for native 5G1.1 (but only three fold greater when viewed in terms of numbers of antibody-antigen binding sites).

Thus, the number of molecules of each of these scFvs required to equal the effect of a single molecule of native 5G1.1 must be increased by a factor of six to adjust for the ratio at which blocking is complete. Since the mass of these molecules is approximately one sixth of the mass of native 5G1.1, dosages of the scFvs are in the same range as those for the native 5G1.1 mAb.

In addition to lowering dosage levels, the anti-C5 antibodies used in the practice of the present invention (i.e., in treating GN) achieve important therapeutic effects not achieved with the anti-C6 antibodies. Specifically, the control and test animals in the C6 work exhibited both hypercellularity and narrowing of capillary lumens. In direct contrast, no such hypercellularity or narrowing of capillary lumens was seen when diseased individuals were treated with anti-C5 antibodies (see FIG. 1).

Moreover, the anti-C5 antibodies used in the present invention achieve a reduction in glomerular enlargement, thus providing a clear demonstration of the unexpectedly powerful anti-inflammatory effects of the anti-C5 antibodies used in the practice of the invention. Nowhere in the C6 work is there any disclosure or suggestion of such a powerful anti-inflammatory effect.

V. Anti-C5 Monoclonal Antibodies that Block Complement Hemolytic Activity and Block the Generation of C5a

Anti-C5 mabs that have the desirable ability to block complement hemolytic activity and to block the generation of C5a (and are thus preferred for use in the treatment of GN and other inflammatory conditions in accordance with the present invention) have been known in the art since at least 1982 (Moongkarndi et al. *Immunobiol.* 1982, 162:397; Moongkarndi et al. *Immunobiol.* 1983, 165:323). Antibodies known in the art that are immunoreactive against C5 or C5 fragments include antibodies against the C5 beta chain (Moongkarndi et al. *Immunobiol.* 1982, 162:397; Moongkarndi et al. *Immunobiol.* 1983, 165:323; Wurzner et al. 1991, *supra*; Mollnes et al. *Scand. J. Immunol.* 1988, 28:307-312); C5a (see for example, Ames et al. *J. Immunol.* 1994, 152:4572-4581, U.S. Pat. No. 4,686,100, and European patent publication No. 0 411 306); and antibodies against non-human C5 (see for example, Giclas et al. *J. Immunol. Meth.* 1987, 105:201-209). Significantly, none of these anti-C5 mAbs has the properties of the novel anti-C5 mabs of the invention, i.e., none of them binds to the C5 alpha chain but not to the C5 cleavage product C5a, none of them has the ability to substantially block both complement hemolytic activity and the generation of C5a to substantially the same extent at the same concentration of antibody. It is noteworthy that an scFv derivative of the N19/8 antibody of Wurzner et al. 1991, *supra*, has been prepared, and that the N19/8 scFv has 50% less inhibitory activity towards C5a generation than the native N19/8 antibody (see Example 15). This is in contrast to the 5G1.1 scFv, which retained substantially all of its inhibitory activity towards C5a generation (see Example 12).

While not wishing to be bound by any particular theory of operation, it is believed that these distinctions are due to the specific binding characteristics of the antibodies of the invention. Accordingly, antibodies that do not bind to sites within the alpha chain of C5, and antibodies that bind to the C5 cleavage product C5a (free C5a), are believed to lack the ability to substantially block both complement hemolytic activity and the generation of C5a to substantially the same extent at the same concentration of antibody.

DESCRIPTION OF THE PREFERRED EMBODIMENTS

As discussed above, the present invention relates to the use of anti-C5 antibodies in treating patients suffering from GN and other diseases, and to specific C5 antibodies and antibody preparations. Preferably, and when used to treat GN, the anti-C5 antibodies are used in an amount effective to substantially reduce (e.g., reduce by at least about 50%) the cell-lysing ability of complement present in the patient's blood (the "cell-lysing ability of complement present in the patient's blood" is also referred to herein as the "serum complement activity of the patient's blood"). Reduction of the cell-lysing ability of complement present in the patient's blood can be measured by methods well known in the art such as, for example, by the chicken erythrocyte hemolysis method described below under the heading "Cell Lysis Assays."

To achieve the desired reductions, the anti-C5 antibodies can be administered in a variety of unit dosage forms. The dose will vary according to the particular antibody. For example, different antibodies may have different masses and/or affinities, and thus require different dosage levels. Antibodies prepared as Fab' fragments will also require differing dosages than the equivalent intact immunoglobulins, as they are of considerably smaller mass

than intact immunoglobulins, and thus require lower dosages to reach the same molar levels in the patient's blood.

The dose will also vary depending on the manner of administration, the particular symptoms of the patient being treated, the overall health, condition, size, and age of the patient, and the judgment of the prescribing physician. Dosage levels of the antibodies for human subjects are generally between about 1 mg per kg and about 100 mg per kg per patient per treatment, and preferably between about 5 mg per kg and about 50 mg per kg per patient per treatment. In terms of plasma concentrations, the antibody concentrations are preferably in the range from about 25 μ g/ml to about 500 μ g/ml.

Subject to the judgement of the physician, a typical therapeutic treatment includes a series of doses, which will usually be administered concurrently with the monitoring of clinical endpoints such as BUN levels, proteinuria levels, etc., with the dosage levels adjusted as needed to achieve the desired clinical outcome. Alternatively, levels of serum complement activity available in the patient's blood are monitored using the techniques set forth below under the heading "Cell Lysis Assays" to determine if additional doses or higher or lower dosage levels of antibodies are needed, with such doses being administered as required to maintain at least about a 50% reduction, and preferably about a 95% or greater reduction of serum complement activity. Other protocols can, of course, be used if desired as determined by the physician.

Administration of the anti-C5 antibodies will generally be performed by an intravascular route, e.g., via intravenous infusion by injection. Other routes of administration may be used if desired. Formulations suitable for injection are found in *Remington's Pharmaceutical Sciences*, Mack Publishing Company, Philadelphia, Pa., 17th ed. (1985). Such formulations must be sterile and non-pyrogenic, and generally will include a pharmaceutically effective carrier, such as saline, buffered (e.g., phosphate buffered) saline, Hank's solution, Ringer's solution, dextrose/saline, glucose solutions, and the like. The formulations may contain pharmaceutically acceptable auxiliary substances as required, such as, tonicity adjusting agents, wetting agents, bactericidal agents, preservatives, stabilizers, and the like.

The formulations of the invention can be distributed as articles of manufacture comprising packaging material and the anti-C5 antibodies. When prepared for use in the treatment of GN, the packaging material will include a label which indicates that the formulation is for use in the treatment of kidney disease and may specifically refer to nephritis or glomerulonephritis.

The anti-C5 antibody is preferably a monoclonal antibody, although polyclonal antibodies produced and screened by conventional techniques can also be used if desired. As discussed above, the anti-C5 antibodies must be effective in reducing the cell-lysing ability of complement present in human blood. As also discussed above, this property of the antibodies can be determined by methods well known in the art such as, for example, by the chicken erythrocyte hemolysis method described below under the heading "Cell Lysis Assays".

The anti-C5 antibodies used in the practice of the invention bind to C5 or fragments thereof, e.g., C5a or C5b. Preferably, the anti-C5 antibodies are immunoreactive against epitopes on the beta chain of purified human complement component C5 and are capable of blocking the conversion of C5 into C5a and C5b by C5 convertase. This capability can be measured using the techniques described in

Wurzner, et al., *Complement Inflamm* 8:328–340, 1991. Preferably, the anti-C5 antibodies are used to treat GN in an amount effective to reduce the C5 convertase activity available in the patient's blood by at least about 50%.

In a particularly preferred embodiment of the invention, the anti-C5 antibodies are not immunoreactive against epitopes on the beta chain, but rather are immunoreactive against epitopes within the alpha chain of purified human complement component C5. In this embodiment the antibodies are also capable of blocking the conversion of C5 into C5a and C5b by C5 convertase. In an especially preferred example of this embodiment they can provide this blockade at substantially the same concentrations needed to block hemolytic activity.

Within the alpha chain, the most preferred antibodies bind to an amino-terminal region, however, they do not bind to free C5a. Particularly preferred targets for these antibodies within the alpha chain include the 5G46k fragment, the 5G27k fragment, the 5G325aa peptide, the 5G200aa peptide, or the KSSKC epitope. The scope of the invention also includes the 5G46k fragment, the 5G27k fragment, the 5G325aa peptide, the 5G200aa peptide, or the KSSKC epitope (SEQ ID NO:1) that are useful as immunogens and screening ligands for producing the antibodies of the invention.

Hybridomas producing monoclonal antibodies reactive with complement component C5 can be obtained according to the teachings of Sims, et al., U.S. Pat. No. 5,135,916. As discussed therein, antibodies are prepared using purified components of the complement membrane attack complex as immunogens. In accordance with the present invention, complement component C5 or C5b is preferably used as the immunogen. In accordance with a particularly preferred aspect of the present invention, the immunogen is the alpha chain of C5. Within the alpha chain, the most preferred immunogens include the 5G46k fragment, the 5G27k fragment, the 5G325aa peptide, or the 5G200aa peptide. A less preferred immunogen is the KSSKC epitope.

In accordance with the invention, the antibodies of the invention all share certain required functional properties. These are the ability to substantially inhibit complement hemolytic activity and to substantially inhibit the conversion of C5 to produce C5a. Preferably, but not requisitely, they provide these functions when used at a molar ratio of antibody to antigen (C5) of 3:1 or less.

A particularly preferred antibody of the invention is the 5G1.1 antibody (5G1.1, produced by the 5G1.1 hybridoma, ATCC designation HB-11625). Other particularly preferred antibodies of the present invention share the required functional properties discussed in the preceding paragraph and have any of the following characteristics:

- (1) they compete with 5G1.1 for binding to portions of C5—the C5 alpha chain, the 5G46k fragment, the 5G27k fragment, the 5G325aa peptide (SEQ ID NO:1), the 5G200aa peptide, or the KSSKC peptide—that are specifically immunoreactive with 5G1.1; and
- (2) they specifically bind to the C5 alpha chain, the 5G46k fragment, the 5G27k fragment, the 5G325aa peptide, the 5G200aa peptide, and/or the KSSKC peptide (SEQ ID NO:1). Such specific binding, and competition for binding can be determined by various methods well known in the art, including the plasmon surface resonance method (Johne et al., *J. Immunol. Meth.* 1993, 160:191–198).
- (3) they block the binding of C5 to either C3 or C4 (which are components of C5 convertase).

Also in accordance with the invention, the antibodies preferably should prevent the cleavage of C5 to form C5a and C5b, thus preventing the generation of the anaphylatoxic activity associated with C5a and preventing the assembly of the membrane attack complex associated with C5b. In a particularly preferred embodiment, these anti-C5 antibodies will not impair the opsonization function associated with the activation of complement component C3 by a C3 convertase. Plasma C3 convertase activity can be measured by assaying plasma for the presence of C3a as described below under the heading “Histology.” Preferably, the anti-C5 antibody produces essentially no reduction in plasma C3a levels.

General methods for the immunization of animals (in this case with C5 or C5b or another preferred immunogen), isolation of polyclonal antibodies or antibody producing cells, fusion of such cells with immortal cells (e.g., myeloma cells) to generate Hybridomas secreting monoclonal antibodies, screening of hybridoma supernatants for reactivity of secreted monoclonal antibodies with a desired antigen (in this case C5 or C5b or another preferred immunogen), the preparation of quantities of such antibodies in hybridoma supernatants or ascites fluids, and for the purification and storage of such monoclonal antibodies, can be found in numerous publications. These include: Coligan, et al., eds. *Current Protocols In Immunology*, John Wiley & Sons, New York, 1992; Harlow and Lane, *Antibodies, A Laboratory Manual*, Cold Spring Harbor Laboratory, New York, 1988; Liddell and Cryer, *A Practical Guide To Monoclonal Antibodies*, John Wiley & Sons, Chichester, West Sussex, England, 1991; Montz, et al., *Cellular Immunol.* 127:337–351, 1990; Wurzner, et al., *Complement Inflamm.* 8:328–340, 1991; and Mollnes, et al., *Scand. J. Immunol.* 28:307–312, 1988.

As used herein, the term “antibodies” refers to immunoglobulins produced in vivo, as well as those produced in vitro by a hybridoma, and antigen binding fragments (e.g., Fab' preparations) of such immunoglobulins, as well as to recombinantly expressed antigen binding proteins, including immunoglobulins, chimeric immunoglobulins, “humanized” immunoglobulins, antigen binding fragments of such immunoglobulins, single chain antibodies, and other recombinant proteins containing antigen binding domains derived from immunoglobulins. As used herein, “antibodies” also refers to antigen binding synthetic peptides comprising sequences derived from the sequences of immunoglobulin antigen binding domains. As used herein, the term “recombinant mAbs” refers to recombinantly expressed antigen binding proteins. As used herein, the term “antibody-antigen binding site” refers to an antigen binding site of an antibody comprising at least one CDR sequence.

Antibodies whose amino acid sequences are full length immunoglobulin sequences that have not been truncated (e.g., to produce an scFv or an Fab) or mutated (e.g., spliced to form a chimeric antibody or humanized) are referred to herein as “native” antibodies. Publications describing methods for the preparation of such antibodies, in addition to those listed immediately above, include: Reichmann, et al., *Nature*, 332:323–327, 1988; Winter and Milstein, *Nature*, 349:293–299, 1991; Clackson, et al., *Nature*, 352:624–628, 1991; Morrison, *Annu Rev Immunol*, 10:239–265, 1992; Haber, *Immunol Rev*, 130:189–212, 1992; and Rodrigues, et al., *J. Immunol*, 151:6954–6961, 1993.

While treatment of GN in accordance with the process of the present invention may be carried out using polyclonal or monoclonal antibodies, monospecific antibodies are preferred. As used herein “monospecific antibodies” refer to

antibodies that bind to a specific region of a particular antigen. All monoclonal antibodies are monospecific, but polyclonal antibodies are typically not monospecific.

As is known in the art, however, monospecific polyclonal antibodies may be prepared by various methods. For example, a peptide (e.g., an oligopeptide—as used herein—after and in the claims, a polymer of 5 to 200 amino acids) may be used as an immunogen. Another procedure allowing the preparation of monospecific polyclonal antibodies is the use of antigen affinity purification techniques to isolate a monospecific antibody population from a polyclonal antibody mixture. In accordance with the present invention, peptides are preferred as immunogens for the production and as affinity ligands for the purification of monospecific polyclonal anti-KSSKC antibodies.

The native (i.e., non-engineered) monoclonal antibodies of the invention are preferably prepared by conventional means, with the 5G46k fragment, the 5G27k fragment, the 5G200aa peptide, the 5G325aa peptide, and/or the KSSKC peptide (SEQ ID NO:1) (e.g., immobilized on a polypropylene membrane as described below in Example 13) being used as screening ligand(s). This involves testing hybridoma supernatants for binding to each screening ligand.

In one preferred embodiment, the native mAbs of the invention are prepared using the alpha chain of human C5, or fragments thereof, as immunogen. Preferred fragments of the alpha chain of human C5 for this purpose include the 5G46k fragment, the 5G27k fragment, and/or the 5G200aa fragment. Although less preferred, the KSSKC peptide (SEQ ID NO:1) may also be used as an immunogen.

Another (albeit less preferred) immunogen and screening ligand for the preparation of antibodies within the scope of the novel antibodies of the present invention is the “cleavage site peptide,” i.e., the peptide spanning amino acids 725 through 754 of SEQ ID NO:2 (the C5a cleavage site), as discussed below in Example 13.

In another preferred embodiment of the invention, the native mAbs of the invention are prepared in transgenic mice expressing human immunoglobulins (see, for example, Green et al., *Nature Genet.* 1994, 7:13–21). In this case, the same preferred immunogens and screening ligands are used as described for the preparation of other native mAbs.

In another preferred embodiment of the invention, the recombinant mAbs of the invention are prepared by screening phage display libraries expressing recombinant mAb-encoding polynucleotides (preferably encoding human recombinant mabs). See, for example, Ames et al., 1994, *supra*; Smith and Scott, *Meth. Enzymol.* 1993, 217:228; Kay et al., *Gene*, 1993, 128:59–65. This screening is carried out with the screening ligands described above for the preparation of native mAbs. The recombinant mAbs of the invention are prepared by subcloning the recombinant mAb-encoding polynucleotides into a suitable expression vector, expressing them in a suitable host (as described below), and isolating the recombinant mAbs.

The present invention provides recombinant expression vectors which include the synthetic, genomic, or cDNA-derived nucleic acid fragments of the invention, i.e. polynucleotides encoding the mabs of the invention. The nucleotide sequence coding for any of the mAbs of the invention can be inserted into an appropriate expression vector, i.e., a vector that contains the necessary elements for the transcription and translation of the inserted protein-coding sequence. The necessary transcriptional and translational signals can also be supplied by the native or source gene and/or its flanking regions.

A variety of host vector systems may be utilized to express the recombinant expression vectors of the invention.

These include, but are not limited to, mammalian cell systems infected with recombinant virus (e.g., vaccinia virus, adenovirus, retroviruses, etc.); mammalian cell systems transfected with recombinant plasmids; insect cell systems infected with recombinant virus (e.g., baculovirus); microorganisms such as yeast containing yeast expression vectors, or bacteria transformed with recombinant bacteriophage DNA, recombinant plasmid DNA, or cosmid DNA (see, for example, Goeddel, 1990).

Useful expression vectors for bacterial use can comprise a selectable marker and bacterial origin of replication derived from commercially available plasmids comprising genetic elements of the well-known cloning vector pBR322 (American Type Culture Collection—“ATCC”—, 10801 University Boulevard, Manassas, Va. 20110-2209, United States of America; ATCC Accession No. 37017). These pBR322 “backbone sections,” or functionally equivalent sequences, are combined with an appropriate promoter and the structural gene to be expressed. Promoters commonly used in recombinant microbial expression vectors include, but are not limited to, the lactose promoter system (Chang, et al., *Nature* 275:615), the tryptophan (trp) promoter (Goeddel, et al., 1980, *Gene Expression Technology*, Volume 185. Academic Press, Inc., San Diego, Calif.) and the tac promoter, or a fusion between the tac and trp promoters referred to as the trc promoter (Maniatis, 1982, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.). Particularly preferred promoters include the T7 promoter, which is used in conjunction with host cell expression of a T7 RNA polymerase (see Studier et al. 1990, *Meth. Enzymol.* 185:60–89), and the trc promoter, which is found in several commercially available vectors, as described below.

Preferred bacterial expression vectors include, but are not limited to, the pET vectors (see Studier et al. 1990, *supra*) and the Trc vectors. Many of the pET vectors are commercially available from Stratagene Cloning Systems (La Jolla, Calif.). A particularly preferred vector is the pET Trc SO5/N1 vector described below (SEQ ID NO:18). A Trc vector, pTrc 99A, is available from Pharmacia. Other Trc vectors include the pSE vectors (Invitrogen, San Diego, Calif.).

Preferred bacteria for expression of recombinant mAbs include *Bacillus subtilis* and, most preferably, *Escherichia coli*. A particularly preferred strain of *E. coli* is strain W3110 (ATCC designation 27325). Under certain unusual conditions it may be necessary to use standard bacterial genetics methods to prepare derivative strains of W3110, for example, when a contaminating bacteriophage (“phage”) is present in the laboratory where the bacterial manipulations are being carried out. Generally, and particularly for large scale preparation of the recombinant anti-KSSKC mAbs of the invention, it is preferred to use unmodified W3110, or another fully characterized strain.

In cases where phage contamination is a problem and disinfection is not practicable or desirable, it is preferred to identify the phage contaminant and to then use a fully characterized bacterial strain having a known mutation rendering the bacterium resistant to the phage. Preferably the mutation is a null mutant for the receptor for the phage. In some instances, however, the generation use of a relatively uncharacterized phage-resistant derivative strain may be acceptable, particularly in small scale experimental work. When such derivative strains are desired, they may be prepared using the methods described below in Example 11.

For most purposes the use of unmodified W3110 or another fully characterized bacterial strain is generally preferred. This is particularly true for the preparation of phar-

maceutical agents comprising the recombinant anti-KSSKC mAbs of the invention. This is because of the problems, well known in the art, of using bacterial strains containing uncharacterized or partially characterized mutations for the production of ingredients of pharmaceutical agents.

The recombinant mAbs of the invention may also be expressed in fungal hosts, preferably yeast of the *Saccharomyces* genus such as *S. cerevisiae*. Fungi of other genera such as *Aspergillus*, *Pichia* or *Kluyveromyces* may also be employed. Fungal vectors will generally contain an origin of replication from the 2 μ m yeast plasmid or another autonomously replicating sequence (ARS), a promoter, DNA encoding a mAb of the invention, sequences directing polyadenylation and transcription termination, and a selectable marker gene. Preferably, fungal vectors will include an origin of replication and selectable markers permitting transformation of both *E. coli* and fungi.

Suitable promoter systems in fungi include the promoters for metallothionein, 3-phosphoglycerate kinase, or other glycolytic enzymes such as enolase, hexokinase, pyruvate kinase, glucokinase, the glucose-repressible alcohol dehydrogenase promoter (ADH2), the constitutive promoter from the alcohol dehydrogenase gene, ADH1, and others. See, for example, Schena, et al. 1991 *Meth. Enzymol.* 194:389-398. Secretion signals, such as those directing the secretion of yeast alpha-factor or yeast invertase, can be incorporated into the fungal vector to promote secretion of a soluble recombinant mAb into the fungal growth medium. See Moir, et al., 1991, *Meth. Enzymol.* 194:491-507.

Preferred fungal expression vectors can be assembled using DNA sequences from pBR322 for selection and replication in bacteria, and fungal DNA sequences, including the ADH1 promoter and the alcohol dehydrogenase ADH1 termination sequence, as found in vector pAAH5 (Ammerer, 1983, *Meth. Enzymol.* 101:192). The ADH1 promoter is effective in yeast in that ADH1 mRNA is estimated to be 1-2% of total poly(A) RNA.

Various mammalian or insect cell culture systems can be employed to express recombinant mAbs. Suitable baculovirus systems for production of heterologous proteins in insect cells are reviewed by Luckow, et al., 1988. Examples of suitable mammalian host cell lines include the COS cell of monkey kidney origin, mouse L cells, murine C127 mammary epithelial cells, mouse Balb/3T3 cells, Chinese hamster ovary cells (CHO), human 293 EBNA and HeLa cells, myeloma, and baby hamster kidney (BHK) cells, with myeloma cells, CHO cells, and human 293 EBNA cells being particularly preferred.

Mammalian expression vectors may comprise non-transcribed elements such as origin of replication, a suitable promoter and enhancer linked to the recombinant mAb gene to be expressed, and other 5' or 3' flanking sequences such as ribosome binding sites, a polyadenylation sequence, splice donor and acceptor sites, and transcriptional termination sequences.

The transcriptional and translational control sequences in mammalian expression vector systems to be used in transforming vertebrate cells may be provided by viral sources. For example, commonly used promoters and enhancers are derived from Polyoma virus, Adenovirus, Simian Virus 40 (SV40), and human cytomegalovirus, including the cytomegalovirus immediate-early gene 1 promoter and enhancer (CMV).

Particularly preferred eukaryotic vectors for the expression of recombinant anti-KSSKC mAbs are pAPEX-1 (SEQ ID NO:3 and, more preferably, pAPEX-3p, SEQ ID NO:4. The vector pAPEX-1 is a derivative of the vector pcDNAI/

Amp (Invitrogen) which was modified to increase protein expression levels. First, the 3'-untranslated SV40 small-t antigen intron was removed by deletion of a 601 base pair *XbaI/HpaI* fragment since this intron is susceptible to aberrant splicing into upstream coding regions (Evans and Scarpulla, 1989 *Gene* 84:135; Huang and Gorman, 1990, *Molec. Cell Biol.* 10:1805). Second, a chimeric adenovirus-immunoglobulin hybrid intron was introduced into the 5'-untranslated region by replacing a 484 base pair *NdeI-NotI* fragment with a corresponding 845 base pair *NdeI-NotI* fragment from the vector pRc/CMV7SB (Sato et al., 1994, *J. Biol. Chem.* 269:17267). Finally, to increase plasmid DNA yields from *E. coli*, the resulting CMV promoter expression cassette was shuttled into the vector pGEM-4Z (Promega Corp. Madison, Wis.).

The vector pAPEX-3 is a derivative of the vector pDR2 (Clontech Laboratories, Inc. Palo Alto, Calif.) in which the EBNA gene was first removed by deletion of a 2.4 kb *ClaI/AccI* fragment. The RSV promoter was then replaced with the CMV promoter and the adenovirus/immunoglobulin chimeric intron by exchanging a 450 bp *MluI/BamHI* fragment from pDR2 with a 1.0 kb *MluI/BamHI* fragment from the vector pAPEX-1. For construction of pAPEX-3P, a 1.7 kb *BstBI/SwaI* fragment containing the HSV tk promoter and hygromycin phosphotransferase (hyg) gene was removed from pAPEX-3 and replaced with a 1.1 kb *SnaBI/NheI* fragment containing the SV40 early promoter and puromycin acetyltransferase (pac) gene (Morgenstern and Land, 1990, *Nucleic Acids Res.* 18:3587-3596) plus a 137 bp *XbaI/ClaI* fragment containing an SV40 polyadenylation signal from the vector pAPEX-1.

A particularly preferred host cell for the expression of recombinant mAb-encoding inserts in the pAPEX vectors is the human 293 EBNA cell line (Invitrogen, San Diego, Calif.).

Another preferred eukaryotic vector for the expression of recombinant mAbs is pcDNAI/Amp (Invitrogen Corporation, San Diego, Calif.). The pcDNAI/Amp expression vector contains the human cytomegalovirus immediate-early gene I promoter and enhancer elements, the Simian Virus 40 (SV40) consensus intron donor and acceptor splice sequences, and the SV40 consensus polyadenylation signal. This vector also contains an SV40 origin of replication that allows for episomal amplification in cells (e.g., COS cells, MOP8 cells, etc.) transformed with SV40 large T antigen, and an ampicillin resistance gene for propagation and selection in bacterial hosts.

Purified recombinant mAbs are prepared by culturing suitable host/vector systems to express the recombinant mAb translation products of the nucleic acid molecules of the present invention, which are then purified from the culture media or cell extracts of the host system, e.g., the bacteria, insect cells, fungal, or mammalian cells. Fermentation of fungi or mammalian cells that express recombinant mAb proteins containing a histidine tag sequence (a sequence comprising a stretch of at least 5 histidine residues) as a secreted product greatly simplifies purification. Such a histidine tag sequence enables binding under specific conditions to metals such as nickel, and thereby to nickel (or other metal) columns for purification. Recombinant mAbs may also be purified by protein G affinity chromatography (Proudfoot et al., 1992, *Protein Express. Purif.* 3:368).

Additional preferred embodiments are numbered and set forth below as "favored embodiments."

FAVORED EMBODIMENTS

1. A method for the treatment of glomerulonephritis in a patient in need of such treatment comprising introducing an

antibody that binds to complement component C5 into the patient's bloodstream in an amount effective to substantially reduce the cell-lysing ability of complement present in the patient's blood.

2. The method of favored embodiment 1 wherein the antibody reduces the conversion of complement component C5 into complement components C5a and C5b.

3. The method of favored embodiment 1 wherein the antibody binds to C5b.

4. The method of favored embodiment 1 wherein the antibody does not substantially inhibit formation of complement component C3b.

5. The method of favored embodiment 1 wherein the antibody is introduced into the patient's bloodstream in a dose that is not greater than 0.1 grams per kilogram.

6. An article of manufacture comprising packaging material and a pharmaceutical agent contained within said packaging material, wherein:

(a) said pharmaceutical agent comprises an antibody to complement component C5, said antibody being effective in substantially reducing the cell-lysing ability of complement present in the patient's blood; and

(b) said packaging material comprises a label which indicates that said pharmaceutical agent is for use in the treatment of kidney disease.

7. The article of manufacture of favored embodiment 6 wherein the label indicates that said pharmaceutical agent is for use in the treatment of nephritis.

8. The article of manufacture of favored embodiment 7 wherein the label indicates that said pharmaceutical agent is for use in the treatment of glomerulonephritis.

9. The article of manufacture of favored embodiment 6 wherein the pharmaceutical agent is to be used at a dosage level not greater than 0.1 grams per kilogram.

10. An antibody comprising at least one antibody-antigen binding site, said antibody exhibiting specific binding to human complement component C5, said specific binding being targeted to the alpha chain of human complement component C5, wherein the antibody inhibits complement activation in a human body fluid and does not specifically bind to the human complement activation product free C5a.

11. The antibody of favored embodiment 10 wherein the inhibition of complement activation in the human body fluid is measurable as a substantial increment of blockade of C5a generation and a substantial increment of blockade of complement hemolytic activity in the body fluid, said increment of blockade of C5a generation being substantially equal to said increment of blockade of complement hemolytic activity.

12. The antibody of favored embodiment 10 wherein, upon binding to human C5, the antibody substantially inhibits the ability of C5 to bind to human complement component C3.

13. The antibody of favored embodiment 10 wherein, upon binding to human C5, the antibody substantially inhibits the ability of C5 to bind to human complement component C4.

14. The antibody of favored embodiment 10 wherein the antibody binds specifically with a 5G46k fragment.

15. The antibody of favored embodiment 10 wherein the antibody binds specifically to a 5G27k fragment.

16. The antibody of favored embodiment 10 wherein the antibody binds specifically to a 5G325aa peptide.

17. The antibody of favored embodiment 10 wherein the antibody binds specifically to a 5G200aa peptide.

18. The antibody of favored embodiment 10 wherein the antibody binds specifically to a KSSKC peptide.

19. The antibody of favored embodiment 10 wherein the inhibition of complement activation in the human body fluid is measurable as a substantially complete blockade of C5a generation in the body fluid and a substantially complete blockade of complement hemolytic activity in the body fluid when the antibody is added to the body fluid at a concentration yielding a ratio equal to or less than 10 moles of antibody-antigen binding sites of the antibody to 1 mole of human C5 in the body fluid.

20. The antibody of favored embodiment 19 wherein the concentration yields a ratio equal to or less than 3 moles of antibody-antigen binding sites of the antibody to 1 mole of human C5 in the body fluid.

21. Hybridoma 5G1.1 having ATCC designation HB-11625.

22. An antibody produced by the hybridoma of favored embodiment 21.

23. An antibody that can compete with the antibody of favored embodiment 22 for binding to the alpha chain of human C5.

24. A nucleic acid molecule comprising a nucleotide sequence encoding an scFv polypeptide comprising an amino acid sequence corresponding to amino acid 1 through amino acid 248 of SEQ ID NO:7.

25. A nucleic acid molecule comprising a nucleotide sequence encoding a polypeptide comprising a variable light chain region amino acid sequence corresponding to amino acid 3 through amino acid 110 of SEQ ID NO:9.

26. A nucleic acid molecule comprising a nucleotide sequence encoding a polypeptide comprising a variable heavy chain region amino acid sequence corresponding to amino acid 1 through amino acid 122 of SEQ ID NO:10.

27. An isolated protein comprising:

(a) a first polypeptide region comprising a variable light chain region amino acid sequence corresponding to amino acid 3 through amino acid 110 of SEQ ID NO:9; and

(b) a second polypeptide region comprising a variable heavy chain region amino acid sequence corresponding to amino acid 1 through amino acid 122 of SEQ ID NO:10.

28. An isolated polypeptide comprising an amino acid sequence encoded by the nucleic acid molecule of favored embodiment 24, favored embodiment 25, or favored embodiment 26, wherein the polypeptide is an antibody.

29. A nucleic acid vector comprising a first nucleic acid molecule covalently and operatively linked to a second nucleic acid molecule so that a host containing the vector expresses the polypeptide coded for by the first nucleic acid molecule, wherein the first nucleic acid molecule is the nucleic acid molecule of favored embodiment 24, favored embodiment 25, or favored embodiment 26.

30. A recombinant host cell containing the nucleic acid vector of favored embodiment 29.

31. A method for producing an isolated C5 antibody polypeptide comprising growing the recombinant host cell of favored embodiment 30 such that the polypeptide encoded by the first nucleic acid molecule of the vector is expressed by the host cell, and isolating the expressed polypeptide, wherein the expressed polypeptide is an anti-C5 antibody.

32. The isolated anti-C5 antibody of favored embodiment 31.

33. A nucleic acid molecule comprising a nucleotide sequence encoding an scFv comprising an amino acid sequence corresponding to amino acid 1 through amino acid 248 of SEQ ID NO:8.

34. A nucleic acid molecule comprising a nucleotide sequence encoding an scFv comprising an amino acid sequence corresponding to amino acid 1 through amino acid 248 of SEQ ID NO:17.

35. A nucleic acid molecule comprising a nucleotide sequence encoding a polypeptide comprising a variable light chain region amino acid sequence corresponding to amino acid 1 through amino acid 108 of SEQ ID NO:15.

36. A nucleic acid molecule comprising a nucleotide sequence encoding a polypeptide comprising a variable light chain region amino acid sequence corresponding to amino acid 3 through amino acid 110 of SEQ ID NO:14.

37. A nucleic acid molecule comprising a nucleotide sequence encoding a polypeptide comprising a variable heavy chain region amino acid sequence corresponding to amino acid 1 through amino acid 122 of SEQ ID NO:16.

38. A nucleic acid molecule comprising a nucleotide sequence encoding a polypeptide comprising a variable heavy chain region amino acid sequence corresponding to amino acid 1 through amino acid 122 of SEQ ID NO:12.

39. A nucleic acid molecule comprising a nucleotide sequence encoding a polypeptide comprising a variable heavy chain region amino acid sequence corresponding to amino acid 1 through amino acid 122 of SEQ ID NO:11.

40. An isolated protein comprising:

(a) a first polypeptide region comprising a variable light chain region amino acid sequence corresponding to amino acid 1 through amino acid 108 of SEQ ID NO:15; and

(b) a second polypeptide region comprising a variable heavy chain region amino acid sequence corresponding to amino acid 1 through amino acid 122 of SEQ ID NO:16.

41. An isolated protein comprising:

(a) a first polypeptide region comprising a variable light chain region amino acid sequence corresponding to amino acid 1 through amino acid 1 through amino acid 108 of SEQ ID NO:15; and

(b) a second polypeptide region comprising a variable heavy chain region amino acid sequence corresponding to amino acid 1 through amino acid 122 of SEQ ID NO:12.

42. An isolated protein comprising:

(a) a first polypeptide region comprising a variable light chain region amino acid sequence corresponding to amino acid 1 through amino acid 108 of SEQ ID NO:15; and

(b) a second polypeptide region comprising a variable heavy chain region amino acid sequence corresponding to amino acid 1 through amino acid 122 of SEQ ID NO:11.

43. An isolated protein comprising:

(a) a first polypeptide region comprising a variable light chain region amino acid sequence corresponding to amino acid 3 through amino acid 110 of SEQ ID NO:14; and

(b) a second polypeptide region comprising a variable heavy chain region amino acid sequence corresponding to amino acid 1 through amino acid 122 of SEQ ID NO:16.

44. An isolated protein comprising:

(a) a first polypeptide region comprising a variable light chain region amino acid sequence corresponding to amino acid 3 through amino acid 110 of SEQ ID NO:14; and

(b) a second polypeptide region comprising a variable heavy chain region amino acid sequence corresponding to amino acid 1 through amino acid 122 of SEQ ID NO:12.

45. An isolated protein comprising:

(a) a first polypeptide region comprising a variable light chain region amino acid sequence corresponding to amino acid 3 through amino acid 110 of SEQ ID NO:14; and

(b) a second polypeptide region comprising a variable heavy chain region amino acid sequence corresponding to amino acid 1 through amino acid 122 of SEQ ID NO:11.

46. An isolated protein comprising the amino acid sequence encoded by the nucleic acid molecule of favored embodiment 33, favored embodiment 34, favored embodiment 35, favored embodiment 36, favored embodiment 37, favored embodiment 38, or favored embodiment 39, wherein the isolated protein is an anti-C5 antibody.

47. A nucleic acid vector comprising a first nucleic acid molecule covalently and operatively linked to a second nucleic acid molecule so that a host containing the vector expresses the polypeptide coded for by the first nucleic acid molecule, wherein the first nucleic acid molecule is the nucleic acid molecule of favored embodiment 33, favored embodiment 34, favored embodiment 35, favored embodiment 36, favored embodiment 37, favored embodiment 38, or favored embodiment 39.

48. A recombinant host cell containing the nucleic acid vector of favored embodiment 47.

49. A method for producing an isolated anti-C5 antibody protein comprising growing the recombinant host cell of favored embodiment 48 such that a protein encoded by the nucleic acid molecule is expressed by the host cell, and isolating the expressed protein, wherein the expressed protein is an anti-C5 antibody.

50. The isolated anti-C5 antibody of favored embodiment 47.

51. An isolated nucleic acid molecule comprising:

(a) a nucleotide sequence encoding a variable light region CDR3 comprising an amino acid sequence corresponding to amino acid 93 through amino acid 98 of SEQ ID NO:7;

(b) a sequence complementary to (a); or

(c) both (a) and (b).

52. An isolated nucleic acid molecule comprising:

(a) a nucleotide sequence encoding a variable light region CDR3 comprising an amino acid sequence corresponding to amino acid 91 through amino acid 99 of SEQ ID NO:8;

(b) a sequence complementary to (a); or

(c) both (a) and (b).

53. An isolated nucleic acid molecule comprising:

(a) a nucleotide sequence encoding a variable heavy region CDR1 comprising an amino acid sequence corresponding to amino acid 156 through amino acid 159 of SEQ ID NO:7;

(b) a sequence complementary to (a); or

(c) both (a) and (b).

54. An isolated nucleic acid molecule comprising:

(a) a nucleotide sequence encoding a variable heavy region CDR1 comprising an amino acid sequence corresponding to amino acid 152 through amino acid 161 of SEQ ID NO:8;

(b) a sequence complementary to (a); or

(c) both (a) and (b).

55. An isolated nucleic acid molecule comprising:

(a) a nucleotide sequence encoding a variable heavy region CDR2 comprising an amino acid sequence cor-

- responding to amino acid 179 through amino acid 182 of SEQ ID NO:7;
- (b) a sequence complementary to (a); or
- (c) both (a) and (b).
56. An isolated nucleic acid molecule comprising:
- (a) a nucleotide sequence encoding a variable heavy region CDR2 comprising an amino acid sequence corresponding to amino acid 176 through amino acid 186 of SEQ ID NO:8;
- (b) a sequence complementary to (a); or
- (c) both (a) and (b).
57. An isolated nucleic acid molecule comprising:
- (a) a nucleotide sequence encoding a variable heavy region CDR3 comprising an amino acid sequence corresponding to amino acid 226 through amino acid 236 of SEQ ID NO:7;
- (b) a sequence complementary to (a); or
- (c) both (a) and (b).
58. An isolated nucleic acid molecule comprising:
- (a) a nucleotide sequence encoding a variable heavy region CDR3 comprising an amino acid sequence corresponding to amino acid 225 through amino acid 237 of SEQ ID NO:8;
- (b) a sequence complementary to (a); or
- (c) both (a) and (b).
59. An isolated nucleic acid molecule comprising:
- (a) a nucleotide sequence encoding a variable light region CDR3 comprising an amino acid sequence corresponding to amino acid 91 through amino acid 99 of SEQ ID NO:8;
- (b) a nucleotide sequence encoding a variable heavy region CDR1 comprising an amino acid sequence corresponding to amino acid 152 through amino acid 161 of SEQ ID NO:8;
- (c) a nucleotide sequence encoding a variable heavy region CDR2 comprising an amino acid sequence corresponding to amino acid 176 through amino acid 186 of SEQ ID NO:8; and
- (d) a nucleotide sequence encoding a variable heavy region CDR3 comprising an amino acid sequence corresponding to amino acid 225 through amino acid 237 of SEQ ID NO:8.
60. An isolated nucleic acid molecule comprising:
- (a) a nucleotide sequence encoding a variable light region CDR3 comprising an amino acid sequence corresponding to amino acid 91 through amino acid 99 of SEQ ID NO:8;
- (b) a nucleotide sequence encoding a variable heavy region CDR1 comprising an amino acid sequence corresponding to amino acid 152 through amino acid 161 of SEQ ID NO:8;
- (c) a nucleotide sequence encoding a variable heavy region CDR2 comprising an amino acid sequence corresponding to amino acid 176 through amino acid 192 of SEQ ID NO:8; and
- (d) a nucleotide sequence encoding a variable heavy region CDR3 comprising an amino acid sequence corresponding to amino acid 225 through amino acid 237 of SEQ ID NO:8.
61. An isolated nucleic acid molecule comprising:
- (a) a nucleotide sequence encoding a variable light region CDR3 comprising an amino acid sequence corresponding to amino acid 91 through amino acid 99 of SEQ ID NO:8;

- (b) a nucleotide sequence encoding a variable heavy region CDR1 comprising an amino acid sequence corresponding to amino acid 152 through amino acid 161 of SEQ ID NO:8;
- (c) a nucleotide sequence encoding a variable heavy region CDR2 comprising an amino acid sequence corresponding to amino acid 179 through amino acid 182 of SEQ ID NO:7; and
- (d) a nucleotide sequence encoding a variable heavy region CDR3 comprising an amino acid sequence corresponding to amino acid 225 through amino acid 237 of SEQ ID NO:8.
62. An isolated protein comprising the amino acid sequence encoded by the nucleic acid molecule of favored embodiment 51, favored embodiment 52, favored embodiment 53, favored embodiment 54, favored embodiment 55, favored embodiment 56, favored embodiment 57, favored embodiment 58, favored embodiment 59, favored embodiment 60 or favored embodiment 87.
63. The isolated protein of favored embodiment 62 wherein the protein is an anti-C5 antibody.
64. A nucleic acid vector comprising a first nucleic acid molecule, said first nucleic acid molecule corresponding to the nucleic acid molecule of favored embodiment 51, favored embodiment 52, favored embodiment 53, favored embodiment 54, favored embodiment 55, favored embodiment 56, favored embodiment 57, favored embodiment 58, or favored embodiment 87 covalently and operatively linked to a second nucleic acid molecule so that a host containing the vector expresses the protein encoded by the first nucleic acid molecule.
65. A recombinant host cell containing the nucleic acid vector of favored embodiment 64.
66. A method for producing an anti-C5 antibody comprising growing the recombinant host cell of favored embodiment 65 so that the protein encoded by the nucleic acid molecule is expressed by the host cell, and isolating the expressed protein, wherein the expressed protein is an anti-C5 antibody.
67. The anti-C5 antibody of favored embodiment 66.
68. An isolated 5G46k fragment of human complement component C5.
69. An isolated 5G27k fragment of human complement component C5.
70. An isolated 5G325aa peptide.
71. An isolated 5G200aa peptide.
72. An isolated oligopeptide comprising an amino acid sequence corresponding to amino acid 8 through amino acid 12 of SEQ ID NO:1, i.e., Lys Ser Ser Lys Cys, or in single letter notation, KSSKC.
73. A method of inducing an animal to produce an anti-C5 antibody comprising repeatedly immunizing an animal with the isolated alpha chain of human C5.
74. A method of inducing an animal to produce an anti-C5 antibody comprising immunizing an animal with the isolated 5G46k fragment of favored embodiment 68.
75. A method of inducing an animal to produce an anti-C5 antibody comprising immunizing an animal with the isolated 5G27k fragment of favored embodiment 69.
76. A method of inducing an animal to produce an anti-C5 antibody comprising immunizing an animal with the isolated 5G325aa peptide of favored embodiment 70.
77. A method of inducing an animal to produce an anti-C5 antibody comprising immunizing an animal with the isolated 5G200aa peptide of favored embodiment 71.
78. A method of inducing an animal to produce an anti-C5 antibody comprising immunizing an animal with the isolated oligopeptide of favored embodiment 72.

31

79. A method of identifying an anti-C5 antibody comprising screening candidate antibodies with the isolated alpha chain of human C5.

80. A method of identifying an anti-C5 antibody comprising screening candidate antibodies with the isolated 5G46k fragment of favored embodiment 68.

81. A method of identifying an anti-C5 antibody comprising screening candidate antibodies with the isolated 5G27k fragment of favored embodiment 69.

82. A method of identifying an anti-C5 antibody comprising screening candidate antibodies with the isolated 5G325aa peptide of favored embodiment 70.

83. A method of identifying an anti-C5 antibody comprising screening candidate antibodies with the isolated 5G200aa peptide of favored embodiment 71.

84. A method of identifying an anti-C5 antibody comprising screening candidate antibodies with the isolated oligopeptide of favored embodiment 72.

85. A method of treating a patient in need of complement inhibition comprising administering the antibody of favored embodiment 10, favored embodiment 22, favored embodiment 23, favored embodiment 28, favored embodiment 32, favored embodiment 46, favored embodiment 50, favored embodiment 63, or favored embodiment 67 to the patient in an amount effective to substantially reduce hemolytic activity in a body fluid of the patient.

86. The antibody of favored embodiment 10 wherein the antibody is a recombinant antibody that comprises a human constant domain.

87. An isolated nucleic acid molecule comprising:

(a) a nucleotide sequence encoding a variable heavy region CDR2 comprising an amino acid sequence corresponding to amino acid 176 through amino acid 192 of SEQ ID NO:8;

(b) a sequence complementary to (a); or

(c) both (a) and (b).

88. An isolated nucleic acid molecule comprising:

(a) a nucleotide sequence encoding a variable heavy region CDR2 comprising an amino acid sequence corresponding to amino acid 176 through amino acid 192 of SEQ ID NO:8;

(b) a sequence complementary to (a); or

(c) both (a) and (b).

89. An isolated antibody comprising any one of the CDR regions of CO12, CO13, CO14, Co15, DO12b, DO12C, DO12D.

90. An isolated nucleic acid molecule encoding the antibody of favored embodiment 89.

EXAMPLES

Without intending to limit it in any manner, the present invention will be more fully described by the following examples. The methods and materials which are common to various of the examples are as follows.

Induction of GN in Mice

Four month old female B10.D2/nSnJ mice averaging approximately 25 gms each were obtained from the Jackson Laboratory, Bar Harbor, Me. Mice were injected with 0.1 mL daily (six days per week) of a 40 mg/mL solution of horse apoferritin (HAF), which was prepared by dilution of a saline solution of HAF (Sigma Chemical Company Catalog No. A-3641) with PBS.

Anti-C5 Monoclonal Antibodies

Monoclonal antibodies that bind to complement component C5 of the mouse were prepared by standard methods as an IgG fraction from supernatants of cultures of hybridoma

32

BB5.1 (Frei, et al., 1987), which was obtained from Dr. Brigitta Stockinger of the National Institute for Medical Research, Mill Hill, London, England.

Histology

Kidneys were subjected to microscopic analysis using standard histochemical staining and immunofluorescence techniques. Periodic Acid Schiff (PAS) staining of 5 paraffin sections was by standard methods using a HARLECO PAS histochemical reaction set (EM Diagnostic Systems, Gibbstown, N.J., number 64945/93) according to the manufacturer's directions.

Immunofluorescence staining of 5μ cryostat sections was carried out by standard methods using FITC conjugated sheep anti-mouse C3 (Bioscience International, Kennebunk, Me, Catalog No. W90280F) to detect murine complement component C3, or FITC conjugated goat anti-mouse IgG, IgA, and IgM (Zymed Laboratories, South San Francisco, Calif., Catalog No. 65-6411) to detect immune complexes.

Urine Assays

Protein and glucose levels were determined by spotting urine samples on CHEMSTRIP 2GP dipsticks (Boehringer Mannheim Diagnostics, Indianapolis, Ind., Catalog No. 200743). The detection areas of these strips change color when exposed to urine containing protein or glucose; a lack of color change indicates no detectable protein or glucose is present. The level of analyte in the urine being tested is read out by matching changed colors with color charts supplied by the manufacturer. The urine protein chart shows colors corresponding to trace, 30, 100, and 500 mg/dL.

Cell Lysis Assays

The cell-lysing ability of complement in blood can be determined using hemolytic assays that are performed as follows: Chicken erythrocytes are washed well in GVBS (Rollins, et al., *J Immunol* 144:3478-3483, 1990, Sigma Chemical Co. St. Louis, Mo., catalog No. G-6514) and resuspended to 2×10^8 /mL in GVBS. Anti-chicken erythrocyte antibody (IgG fraction of anti-chicken-RBC antiserum, Intercell Technologies, Hopewell, N.J.) is added to the cells at a final concentration of 25 μg/mL and the cells are incubated for 15 min. at 23° C. The cells are washed 2x with GVBS and 5×10^6 cells are resuspended to 30 μL in GVBS. A 100 μL volume of serum test solution is then added to yield a final reaction mixture volume of 130 μL. As used herein, reference to the serum percentage and/or serum input in these assays indicates the percent serum in the 100 μL volume of serum test solution.

For assays of mouse serum activity, the 100 μL volume of serum test solution contained 50 μL of diluted (in GVBS) mouse serum and 50 μL of human C5 deficient serum (Quidel Corporation, San Diego, Calif.). For assays of human serum activity, the serum test solution may contain up to 100% human plasma or serum, with hybridoma supernatants and/or GVBS being added to yield the 100 μL volume. For the assays used to screen hybridoma supernatants discussed below in Example 7, each 100 μL volume of serum test solution contained 50 μL of hybridoma supernatant and 50 μL of a 10% solution of human serum in GVBS, yielding a 5% human serum input.

After incubation for 30 min. at 37° C., percent hemolysis was calculated relative to a fully lysed control sample. Hemolysis was determined by spinning the cells down and measuring released hemoglobin in the supernatant as the optical density at 415 nm.

A 50% reduction in hemolysis after treatment with the anti-C5 antibodies used in the practice of the invention means that the percent hemolysis after treatment is one half of the percent hemolysis before treatment.

33

Example 1

Anti-C5 Antibodies Inhibit Glomerular
Inflammation and Enlargement

This example illustrates that anti-C5 antibodies will inhibit glomerular inflammation and enlargement.

The protocol for these experiments was as follows. GN-induced mice were treated with anti-C5 antibodies or with PBS as a control after 2 weeks of GN induction. Each mouse received 750 μ g of anti-C5 monoclonal antibodies in PBS (30 mg/kg in a 25 gm mouse) or an equal volume of PBS alone. The amount injected was from 0.25 to 0.365 mL (the concentration of antibodies in PBS varied), which was administered by intraperitoneal injection once a day, six days a week. After an additional 2 weeks of induction and treatment, the animals were sacrificed and kidneys were harvested and prepared for histological examination as described above. Kidneys were also obtained from age-matched uninduced and untreated control mice.

FIG. 1 shows sections of mouse kidneys with a single glomerulus located centrally amidst surrounding interstitium and cross sections of convoluted tubules in each section. As can be seen therein, the kidneys of the GN-induced, PBS-treated mice (FIG. 1B) developed severe crescentic glomerular pathology, including inflammatory glomerular hypercellularity, apparent basement membrane thickening, and glomerular enlargement, while the glomeruli of the GN-induced, anti-C5-treated animals (FIG. 1C) were essentially indistinguishable from the glomeruli of the normal healthy kidneys of the uninduced untreated mice (FIG. 1A).

Note that in the glomeruli with severe crescentic pathology, the size of the glomerular capillary network (glomerular tuft) is not enlarged, but shows signs of compression by a crescent-shaped proliferation of epithelial cells and PAS-positive material, and the Bowman's capsule is dramatically enlarged. Also note that in the section of diseased glomerulus shown in FIG. 1B, the capillary network is split in half by a projection of the hypercellular crescentic mass.

The non-inflamed glomerulus of the uninduced untreated mouse shown in FIG. 1A is approximately 100 μ in diameter; the inflamed glomerulus of the GN-induced, PBS treated mouse shown in FIG. 1B is approximately 175 μ in diameter; the non-inflamed glomerulus of the GN-induced, anti-C5-treated mouse shown in FIG. 1C is approximately 90 μ in diameter.

Example 2

Anti-C5 Antibodies Prevent/Reduce Proteinuria
Associated with GN

This example demonstrates that treatment with anti-C5 antibodies results in the prevention/reduction of kidney damage as evidenced by the lack of significant amounts of protein in the urine (i.e. the presence of less than 100 mg/dL of protein in the urine).

The protocol for the experiments of this example was the same as that used in the experiments of Example 1. Five PBS-treated, GN-induced mice, 6 anti-C5-treated, GN-induced mice, and 4 age-matched untreated uninduced mice were used in this study. A first set of urine samples was analyzed prior to treatment after the initial 2 week induction period. A second set of urine samples was analyzed after the 2 week treatment period. None of the untreated uninduced control animals had detectable protein in their urine at either of these timepoints.

34

The results obtained with the GN-induced mice are set forth in Table 1. As shown therein, at the end of the 2 week PBS treatment period, 4 out of the 5 PBS treated (control) animals developed significant proteinuria, i.e., at least 100 mg/dL of protein in the urine. The fifth animal (mouse D in Table 1) did not have detectable protein in the urine at either timepoint but, unlike the other mice in the study, was found to have very high levels of glucose in the urine after the 2 week PBS treatment period, suggesting that this animal was physiologically compromised.

In the anti-C5-treated, GN-induced group, the one mouse that developed significant proteinuria at the end of the initial 2 week induction period (mouse 6 in Table 1) improved by the end of the 2 week antibody treatment period. In addition, in contrast to the development of significant proteinuria in 4 out of 5 PBS-treated, GN-induced mice, none of the anti-C5-treated, GN-induced mice exhibited significant proteinuria at the end of the 2 week antibody treatment period.

Example 3

Anti-C5 Antibodies do not Inhibit Glomerular
Immune Complex Deposition

This example demonstrates that anti-C5 antibodies used in the practice of the invention achieve their therapeutic effects even though immune complexes are deposited in the glomeruli of treated animals at equivalent levels to those seen in the glomeruli of PBS-treated animals. The example further illustrates that the mechanism of operation of the anti-C5 antibodies is not through the inhibition of immune complex deposition in the glomerulus.

The protocol used in the experiments of this example was the same as that used in the experiments of Example 1. Immunofluorescence staining as described above was performed on sections from the same kidneys harvested in Example 1.

The results are shown in FIG. 2. As can be seen in this figure, equivalent amounts of immune complexes were deposited in the glomeruli of the kidneys of both the PBS-treated, GN-induced mice (FIG. 2B) and the anti-C5-treated, GN-induced mice (FIG. 2C), but not in the untreated uninduced controls (FIG. 2A). Kidneys of GN-induced mice harvested after the 2 week induction period, but before treatment, showed immune complex deposits in the glomeruli, but at lower levels (as indicated by lower fluorescence intensity) than in the kidney sections shown in FIG. 2B and FIG. 2C.

Example 4

Anti-C5 Antibodies Inhibit C5b-9 Generation

This example demonstrates that the anti-C5 antibodies used in the practice of the invention inhibit C5b-9 generation. C5b-9 generation was assayed in 2 ways: (1) by testing the cell-lysing (hemolytic) ability of blood samples, and (2) by measuring levels of soluble C5b-9 in blood samples.

FIG. 3 shows the results of cell lysis assays performed as described above, with mouse serum added to the percentage indicated on the X axis ("serum input %"). In these assays, serum from GN-induced animals treated with either anti-C5 antibodies in PBS or PBS alone (see above) was assayed at the end of the two week treatment period. Serum from normal, uninduced, uninjected mice ("normal mouse serum") obtained from Sigma Chemical Company (St. Louis, Mo., Catalog No. S-3269) was also assayed as an additional control. These results indicate that the anti-C5

35

monoclonal antibody administered to mice at a dosage of 30 mg/Kg completely blocked the cell lysing ability of mouse blood at serum input levels 4-fold higher than the levels of normal serum that produce maximum hemolysis in the assay.

The effects of an anti-C5 monoclonal antibody raised to human C5 was evaluated in circulating human blood. Hybridoma N19/8 (Wurzner, et al., 1991) was obtained from Dr. Otto Götze, Department of Immunology, University of Göttingen, FRG. The C5 monoclonal antibody was prepared following immunization of mice with purified human C5 protein as described in Wurzner, et al., (1991). The hybridoma was propagated in mice, and the monoclonal antibody recovered and purified as an IgG fraction from mouse ascites fluid (*Antibodies, A Laboratory Manual*, Cold Spring Harbor Laboratory, New York, 1988; *Current Protocols In Immunology*, John Wiley & Sons, New York, 1992).

To carry out these experiments, as well as others described below in Examples 5 and 6, 300 mL of whole human blood was drawn from a healthy human donor and additionally a 1 mL sample was removed as a control sample for later analysis. The blood was diluted to 600 mL by the addition of Ringer's lactate solution containing 10 U/mL heparin. The anti-C5 mAb (30 mg in sterile PBS) was added to the diluted blood to a final concentration of 50 µg/mL (results using test samples obtained in this way are labeled "+anti-C5 sample" in FIG. 4 and FIG. 6). In a control experiment, an equal volume of sterile PBS was added to diluted blood (results using control samples obtained in this way are labeled "-anti-C5 sample" in FIG. 4 and FIG. 6).

The blood was then used to prime the extracorporeal circuit of a COBE CML EXCEL membrane oxygenator cardiopulmonary bypass (CPB) machine (Cobe BCT, Inc., Lakewood, Colo.) and circulation through the circuit was started. The circuit was cooled to 28° C. and circulated for 60 minutes. The circuit was then warmed to 37° C. and circulated for an additional 30 minutes, after which time the experiment was terminated. Mechanical circulation of blood in this fashion activates the complement cascade. Samples were taken at several time points.

At each time point an aliquot of blood was taken, and subaliquots were centrifuged to remove all cells and the remaining plasma diluted 1:1 in QUIDEL sample preservation solution (Quidel Corporation, San Diego, Calif.) and stored at -80° C. for subsequent evaluation of soluble C5b-9 (sC5b-9) generation. Diluted subaliquots of plasma were also frozen for evaluation of C3a generation (see Example 5, below). Undiluted subaliquots of plasma were frozen at -80° C. for analysis in hemolytic assays to evaluate the pharmacokinetics of the effects of the anti-C5 antibodies on the cell lysing ability of complement present in the blood (see Example 6, below). These experiments are also discussed in copending U.S. patent application Ser. No. 08/217,391, filed Mar. 23, 1994, now U.S. Pat. No. 5,853,722.

sC5b-9 assays were performed before the addition of the antibody or the commencement of the CPB circuit (labeled "Pre Tx" in FIG. 4 and FIG. 6) using undiluted blood (i.e. blood from the 1 mL sample taken before the blood was diluted with Ringer's lactate solution—labeled "undil" in FIG. 4 and FIG. 6) and Ringer's lactate solution diluted blood (labeled "dil" in FIG. 4 and FIG. 6). Samples of Ringer's lactate solution diluted blood to which the antibody had been added (labeled "Post Tx" in FIG. 4 and FIG. 6) were assayed at the times indicated after starting the CPB circuit.

As can be seen in FIG. 4, while sC5b-9 levels were more than 4-fold higher in untreated samples after 90 minutes of

36

circulation than before circulation, the anti-C5 antibody completely inhibited C5b-9 generation throughout the 90 minute time course of circulation so that sC5b-9 levels during circulation were essentially equivalent to control, uncirculated samples, at all timepoints.

Example 5

Anti-C5 Antibodies do not Inhibit C3 Deposition or Activation

This example demonstrates that treatment with anti-C5 antibodies does not result in the inhibition of the activation of complement component C3 or in the deposition of C3 or its activated fragments in glomeruli.

The deposition of C3, or the fragments generated by its activation (e.g., C3a and C3b), in the glomeruli of GN-induced and GN-uninduced mice was visualized by immunofluorescence staining with a FITC-conjugated sheep anti-mouse C3 antibody preparation using standard methods, as described above. As can be seen in FIG. 5, kidneys of the PBS-treated (FIG. 5B) and the anti-C5 antibody-treated (FIG. 5C) GN-induced mice had roughly equivalent levels of C3 immunoreactive material in the glomeruli, while the uninduced untreated control mice had only traces of C3 immunoreactive material in their kidneys (FIG. 5A).

Note that the print shown in FIG. 5A was overexposed compared to those of FIG. 5B and FIG. 5C to show the very slight levels of reactivity present in normal uninduced kidneys. Kidneys of GN-induced mice harvested after the 2 week induction period, but before treatment, showed C3 immunoreactive materials in the glomeruli, but at lower levels (as indicated by lower fluorescence intensity) than in the kidney sections shown in FIG. 5B and FIG. 5C.

Anti-human C5 antibodies were also tested for possible inhibition of C3 activation in human blood prepared and circulated as described above in Example 4. Activation of complement component C3 was indicated by the presence in the blood of the C3 activation product C3a. C3a assays were performed as follows.

The plasma samples that had previously been diluted in QUIDEL sample preservation solution and frozen (see Example 4) were assayed for the presence of C3a by using the QUIDEL C3A EIA kit (Quidel Corporation, San Diego, Calif.) according to the manufacturer's specifications. Concentrations of C3a in the samples is expressed as ng/well as determined by comparison to a standard curve generated from samples containing known amounts of human C3a.

As seen in FIG. 6, the addition of the anti-C5 mAb had no inhibitory effect on the production of C3a during the circulation of human blood in this experiment.

Example 6

Pharmacokinetics of Anti-C5 Antibodies

The in vivo duration of action of mAb BB5.1, and a Fab' fragment of mAb BB5.1 (prepared by standard methods) was determined in normal female BALB/cByJ mice (averaging approximately 20 gms each) which were obtained from the Jackson Laboratory, Bar Harbor, Me. The mice were given a single intravenous injection (at 35 mg/kg body weight) of the mAb or the Fab' fragment of the mAb (or an equal volume of PBS as a control). Blood samples were collected from the retroorbital plexus at 1, 4, 24, 96, and 144 hours after administration of PBS; 4, 16, and 24 hours after administration of the Fab' fragment of mAb

BB5.1; and 4, 24, 48, 72, 96, and 144 hours after administration of intact mAb BB5.1.

FIG. 7A shows the time course of inhibition of the cell-lysing ability of complement in mouse blood (determined, by testing serum obtained from the blood and diluted to 2.5%, as described above) after the in vivo administration of the mAb, the Fab' fragment, or the PBS. As shown in the figure, the mAb almost completely inhibited the hemolytic activity of the blood throughout the 6 day test period. The Fab', however, had a half-life of approximately 24 hours.

In addition to the above experiments, at the end of the 6 day testing period all of the mice were sacrificed. Kidneys, lungs, and livers were harvested and examined by gross inspection, as well as by microscopic examination of stained sections. All of the organs of the anti-C5 antibody treated animals appeared the same as those taken from PBS control treated animals. The overall appearance of the test and control mice was also indistinguishable prior to necropsy.

Anti-human C5 antibodies were also tested for pharmacokinetic properties in circulating human blood as described above in Example 4. As described therein, the hemolysis inhibiting effects of an anti-human C5 monoclonal antibody were assayed over a 90 minute period of circulation. The results of these assays are charted in FIG. 7B, and show that the N19/8 anti-C5 mAb essentially completely inhibited the cell lysing ability of the human blood during the entire 90 minute period of circulation.

The results of these experiments demonstrate that the anti-C5 antibodies will survive in the bloodstream for a substantial period of time, thus making periodic administration practical.

Example 7

Preparation of Anti-C5 Monoclonal Antibodies

A monoclonal antibody suitable for use in the practice of the present invention was prepared in accordance with the teachings of Sims, et al., U.S. Pat. No. 5,135,916, as follows.

Balb/c mice were immunized three times by intraperitoneal injection with human C5 protein (Quidel Corporation, San Diego, Calif., Cat # A403). The first injection contained 100 μ g of C5 protein in a complete Freund's adjuvant emulsion, the second immunization contained 100 μ g of C5 protein in an incomplete Freund's adjuvant emulsion, and the third immunization was 100 μ g of protein in PBS. The mice were injected at roughly 2 month intervals.

Fusions of splenocytes to myeloma cells to generate hybridomas were performed essentially as described in Current Protocols in Immunology (John Wiley & Sons, New York, 1992, pages 2.5.1 to 2.5.17). One day prior to fusion the mice were boosted IV with 100 μ g of C5 protein. On the day of fusion, the immunized mice were sacrificed and spleens was harvested. SP2/0-AG14 myeloma cells (ATCC CRL#1581) were used as the fusion partner. SP2/0-AG14 cultures were split on the day before the fusion to induce active cell division. A ratio of 1:10 (myeloma cells:splenocytes) was used in the fusions.

The cells were fused using PEG 1450 in PBS without calcium (Sigma Chemical Company, St. Louis, Mo., Catalog No. P-7181) and plated at $1-2.5 \times 10^5$ cells per well. Selection in EX-CELL 300 medium (JRH Biosciences, Lexena, Kans., Catalog No. 14337-78P) supplemented with 10% heat inactivated fetal bovine serum (FBS); glutamine, penicillin and streptomycin (GPS); and HAT (Sigma Chemical Company, St. Louis, Mo., Catalog No. H-0262) was started

the following day. The fusions were then fed every other day with fresh FBS, GPS, and HAT supplemented medium. Cell death could be seen as early as 2 days and viable cell clusters could be seen as early as 5 days after initiating selection. After two weeks of selection in HAT, surviving hybridomas chosen for further study were transferred to EX-CELL 300 medium supplemented with FBS, GPS, and HT (Sigma Chemical Company, St. Louis, Mo., Catalog No. H-0137) for 1 week and then cultured in EX-CELL 300 medium supplemented with FBS and GPS.

Hybridomas were screened for reactivity to C5 and inhibition of complement-mediated hemolysis 10-14 days after fusion, and were carried at least until the screening results were analyzed. The screen for inhibition of hemolysis was the chicken erythrocyte lysis assay described above. The screen for C5 reactivity was an ELISA, which was carried out using the following protocol:

A 50 μ L aliquot of a 2 μ g/mL solution of C5 (Quidel Corporation, San Diego, Calif.) in sodium carbonate /bicarbonate buffer, pH 9.5, was incubated overnight at 4° C. in each test well of a 96 well plate (NUNC-IMMUNO F96 POLYSORP, A/S Nunc, Roskilde, Denmark). The wells were then subjected to a wash step. (Each wash step consisted of three washes with TBST.) Next, test wells were blocked with 200 μ L of blocking solution, 1% BSA in TBS (BSA/TBS) for 1 hour at 37° C. After an additional wash step, a 50 μ L aliquot of hybridoma supernatant was incubated in each test well for 1 hour at 37° C. with a subsequent wash step. As a secondary (detection) antibody, 50 μ L of a 1:2000 dilution of horseradish peroxidase (HRP) conjugated goat anti-mouse IgG in BSA/TBS, was incubated in each test well for 1 hour at 37° C., followed by a wash step. Following the manufacturer's procedures, 10 mg of O-phenylenediamine (Sigma Chemical Company, St. Louis, Mo., Catalog No. P-8287) was dissolved in 25 mLs of phosphate-citrate buffer (Sigma Chemical Company, St. Louis, Mo., Catalog No. P-4922), and 50 μ L of this substrate solution was added to each well to allow detection of peroxidase activity. Finally, to stop the peroxidase detection reaction, a 50 μ L aliquot of 3N hydrochloric acid was added to each well. The presence of antibodies reactive with C5 in the hybridoma supernatants was read out by a spectrophotometric OD determination at 490 nm.

The supernatant from a hybridoma designated as 5G1.1 tested positive by ELISA and substantially reduced the cell-lysing ability of complement present in normal human blood in the chicken erythrocyte hemolysis assay. Further analyses revealed that the 5G1.1 antibody reduces the cell-lysing ability of complement present in normal human blood so efficiently that, even when present at roughly one-half the molar concentration of human C5 in the hemolytic assay, it can almost completely neutralize serum hemolytic activity.

Immunoblot analysis was undertaken to further characterize the 5G1.1 mAb. Human C5 (Quidel Corporation, San Diego, Calif., Catalog No. A403) was subjected to polyacrylamide gel electrophoresis under reducing conditions, transferred to a nitrocellulose membrane, and probed with the 5G1.1 mAb as a purified IgG preparation. Two bands were immunoreactive with the 5G1.1 mAb at apparent molecular weights corresponding to those of the alpha and beta chains of the human C5 protein. The two 5G1.1 immunoreactive bands seen on this Western blot were subsequently found to result from the binding of the 5G1.1 antibody to the 115 kDa C5 alpha chain and to a large fragment of the alpha chain that had the same apparent molecular weight (approximately 75 kDa) as the beta chain of C5 and was present in the C5 preparations used for the experiment.

Assays were performed to determine the relative activity of the N19/8 mAb discussed in Examples 4 and 5 with the 5G1.1 mAb in functional hemolytic assays and to assess whether these mAbs blocked the cleavage of C5 to yield C5a. To this end, the N19/8 and 5G1.1 mAbs were directly compared in human complement hemolytic and C5a release assays.

Hemolytic assays performed in the presence of 20% v/v human serum revealed that the 5G1.1 mAb effectively blocked serum hemolytic activity at a final concentration of 6.25 $\mu\text{g/ml}$ (0.5/1 molar ratio of 5G1.1/C5) whereas the N19/8 mAb blocked at a higher concentration of 25.0 $\mu\text{g/ml}$ (2.0/1 molar ratio of N19/8/C5). When the supernatants from these assays were tested for the presence of C5a, the 5G1.1 mAb was found to have effectively inhibited C5a generation at doses identical to those required for the blockade of C5b-9 mediated hemolytic activity.

In contrast, the N19/8 mAb was 10 fold less effective in blocking the release of C5a in these assays when compared to the 5G1.1 mAb. Furthermore, the ability of the N19/8 mAb to block complement mediated hemolysis was not equivalent to its capacity to block C5a generation in that a dose of 25 $\mu\text{g/ml}$ of N19/8 completely blocked hemolysis while only reducing C5a generation by 37%.

Hybridoma 5G1.1 was deposited with the American Type Culture Collection, 10801 University Boulevard, Manassas, Va. 20110-2209, United States of America, on Apr. 27, 1994, and has been assigned the designation HB-11625. This deposit was made under the Budapest Treaty on the International Recognition of the Deposit of Micro-organisms for the Purposes of Patent Procedure (1977).

Example 8

Determination of the Affinity Constants (K_D) for the Anti-human C5 Monoclonal Antibodies 5G1.1 and N19/8

The procedure utilized to determine the dissociation constant (K_D) of antibody-antigen equilibria in solution was that described by Friguet et al., J. Immunol. Meth. 1985, 77:305-319. This method was used to determine the K_D for the anti-human C5 monoclonal antibodies N19/8 and 5G1.1. The monoclonal antibodies were incubated with the antigen (C5) in solution until the equilibrium was reached. The proportion of antibody that remains unbound (free) at equilibrium was measured using a conventional Enzyme Linked Immunosorbant Assay (ELISA). The experimental values of K_D obtained by this method have been shown to be equivalent to those obtained by other methods (immunoprecipitation of the radiolabeled antigen and fluorescence transfer). This method offers the advantage of dealing with unmodified antigen.

FIGS. 8 and 9 show the Scatchard plots of the binding of the anti-human C5 monoclonal antibodies 5G1.1 and N19/8 to human C5 as measured by ELISA. In each graph (v) represents the fraction of bound antibody and (a) represents the concentration of free antigen at equilibrium. The calculated K_D for the 5G1.1 mAb was 30 pM while the calculated K_D for the N19/8 mAb was 43 pM. These results indicate that the K_D for the 5G1.1 and N19/8 mAb's are similar, and therefore the functional disparity between the two antibodies cannot be explained simply by the differences in affinity for the C5 antigen.

Example 9

Effect of 5G1.1 mAb on Complement Activation During CPB

Experiments involving recirculation of human blood in an CPB circuit, as described above in Examples 4 and 5, were

carried out using three doses of the 5G1.1 mAb (15 mg, 7.5 mg, 3.75 mg) as well as controls in the absence of the 5G1.1 mAb. In five such control experiments performed in this series, C3a FIG. 10) and sC5b-9 (FIG. 11) levels increased during the first 30 min and continued to rise throughout the entire experiment. Addition of the 5G1.1 mAb to the CPB circuit had no effect on the generation of C3a in these experiments.

Conversely, addition of the two highest doses (15 mg and 7.5 mg) of the 5G1.1 mAb completely blocked the generation of sC5b-9 in these experiments while the lowest dose (3.75 mg) only partially blocked sC5b-9 generation. Hemolytic assays performed on serum samples drawn throughout the time course of these experiments revealed that total serum complement activity was not affected in control experiments (FIG. 12). In contrast, the highest dose of the 5G1.1 mAb (15 mg) completely blocked complement hemolytic activity, while the two lower doses (7.5 mg and 3.75 mg), failed to block hemolytic activity.

These results show that the 7.5 mg dose effectively blocked C5b-9 generation in the CPB circuit but failed to block C5b-9-mediated hemolytic activity, suggesting that hemolytic assays alone may not accurately reflect the complement activation that occurs during CPB. These results further indicate that the 5G1.1 mAb can completely block complement activation in human blood, as measured by either criterion, at a dosage of 15 mg/500 ml, a dose that is approximately equivalent to a dose of 150 mg for a 70 kg patient.

Example 10

Cloning of Anti-C5 Recombinant Anti-KSSKC Variable Region Genes Amino Acid Sequencing

To determine the N-terminal amino acid sequence of the 5G1.1 mAb, a 12% acrylamide gel (37.5:1 acrylamide/N, N'-methylene-bisacrylamide) was prepared and pre-electrophoresed for 45 minutes at 10 mA using 1 $\times$ pre-electrophoresis buffer (123 mM bis-Tris, pH 6.6, with the cathode buffer reservoir supplemented with 1 mM reduced glutathione). The following day, the pre-electrophoresis buffer in the cathode reservoir was replaced with cathode reservoir buffer (44 mM N-tris-(hydroxymethyl)methyl-2-aminoethanesulfonic acid, 113 mM bis-Tris, 0.1% (w/v) sodium dodecyl sulfate (SDS), 0.067% (w/v) thioglycolic acid) and the pre-electrophoresis buffer in the anode reservoir was replaced with anode reservoir buffer (63 mM bis-Tris, pH 5.9).

75 μg 5G1.1 monoclonal antibody was added to Laemmli sample buffer (30 mM Tris-HCl pH 6.8, 3% (w/v) SDS, 10 mM EDTA, 0.02% (w/v) bromophenol blue, 5% (v/v) glycerol, 2.5% (v/v) beta-mercaptoethanol) and electrophoresed at 10 mA until the bromophenol blue tracking dye reached the bottom of the gel. The protein was transferred to a PROBLOTT membrane (Applied Biosystems, Foster City, Calif.) using 1 $\times$ transfer buffer (10 mM cyclohexylamino-propane sulfonic acid, 0.05% (w/v) dithiothreitol, 15% (v/v) methanol) at 50 V for one hour.

Protein bands were localized by staining with 0.2% Ponceau S (in 3% trichloroacetic acid, 3% sulfosalicylic acid) followed by destaining with water. Bands were excised and subjected to amino acid sequence analysis using Edman chemistry performed on a pulsed liquid protein sequencer (ABI model 477A), with the PTH amino acids thereby obtained being analyzed with an on-line microbore HPLC system (ABI model 120A).

41

To deblock the amino terminus of the 5G1.1 heavy chain, 10 mg 5G1.1 monoclonal antibody was exchanged into reducing buffer (5 M guanidine-HCl, 50 mM Tris-HCl, 10 mM dithiothreitol, pH 8.5) using a PD-10 column (Pharmacia, Piscataway, N.J.). After a one hour incubation at room temperature, 50 mM iodoacetamide was added and the incubation allowed to continue for 30 minutes. The carbamidomethylated light and heavy chains thus obtained were separated by size exclusion chromatography on a SUPEROSE 12 (Pharmacia) column equilibrated with 5 M guanidine-HCl, 50 mM Tris-HCl pH 8.5. The carbamidomethylated heavy chain was exchanged into 50 mM sodium phosphate, pH 7.0 using a PD-10 column, subjected to digestion with pyroglutamate aminopeptidase (PanVera, Madison, Wis.; 0.5 mU per nmol of heavy chain protein), and sequenced as described above.

For determination of internal amino acid sequence, the carbamidomethylated 5G1.1 light chain was exchanged into 2 M urea, 25 mM Tris-HCl, 1 mM EDTA, pH 8.0 and incubated with endoproteinase Lys-C (Promega, Madison, Wis.; protease:protein ratio of 1:40) at 37° C. overnight. The digested material was run on a C18 reversed phase HPLC column (Beckman Instruments, Fullerton, Calif.) and eluted using a linear 0–50% acetonitrile gradient in 0.1% trifluoroacetic acid. Peaks were subjected to amino acid sequence analysis as described above.

PCR Cloning

Cloning of the 5G1.1 variable heavy region was performed using a set of commercially available primers (Mouse Ig-PRIMER SET, catalogue number 69831-1, Novagen, Madison, Wis.). Total RNA was isolated from 5G1.1 hybridoma cells using the acid/guanidinium thiocyanate technique (Chomczynski and Sacchi, Anal. Biochem. 1987, 162:156–159). For first strand cDNA synthesis, ten micrograms total RNA were denatured at 65° C. for 5 min., chilled on ice, and added to a 100 µl reaction containing 10 mM Tris pH 8.3, 50 mM KCl, 1.5 mM MgCl₂, 10 mM dithiothreitol, 250 µM each dNTP, 20 units AMV reverse transcriptase (Seikagaku America, Rockville, Md.), and 10 pmole of the appropriate 3' primer (as described in the Ig-PRIMER SET kit protocol). After incubation at 37° C. for one hour, five microliters of the cDNA synthesis reaction were added to a 100 microliter PCR reaction containing: 10 mM Tris-HCl pH 9.0 at 25° C., 50 mM KCl, 1.5 mM MgCl₂, 0.1% (w/v) gelatin, 1.0% (v/v) Triton X-100, 200 µM each dNTP, 2.5 U AMPLITAQ DNA polymerase (Perkin-Elmer-Cetus, Norwalk, Conn.) and 25 pmoles of the appropriate 5' and 3' primers (as described in the Ig-PRIMER SET kit protocol). The reaction conditions were 1 minute at 95° C., 1 minute at 42° C., and 1 minute at 72° C. for 30 cycles, followed by a final extension at 72° C. for 10 minutes.

PCR products having the expected size (approximately 450 bp) were cloned into the vector PCR II (Invitrogen, San Diego, Calif.) using a T/A cloning kit (Invitrogen). DNA sequence analysis of cloned DNA fragments was performed by the dideoxy chain-termination method using double-stranded plasmid DNA as a template. A unique heavy chain variable region was isolated by this procedure, with the resulting plasmid designated p5G1.1 VH 2-1-3. Several clones obtained from independent replicate PCR reactions were sequenced to detect any mutations introduced during the PCR amplification of this variable region.

To clone the 5G1.1 light chain variable region, PCR primers were designed by using the UWGCG program TFASTA (University of Wisconsin, Madison, Wis.) to search the GenBank rodent subdirectory with the 19mer query amino acid sequence Ile Gln Met Thr Gln Ser Pro Ala Ser

42

Leu Ser Ala Ser Val Gly Glu Thr Val Thr, that was obtained by amino acid sequencing as described above. An exact match to this sequence was located in the murine germline gene encoding the v-kappa k2 variable region (Seidman et al. Proc. Natl. Acad. Sci. USA 1978 75:3881–3885). The DNA sequence of this germline gene was used to design the oligonucleotide UDEC690 (SEQ ID NO:5) for use as a variable region 5'-primer. A murine kappa gene constant region primer, UDEC395 (SEQ ID NO:6) was also synthesized and used in this reaction. Cloning of the 5G1.1 variable light region was performed using the UDEC690 variable region 5'-primer and the UDEC395 murine kappa gene constant region primer.

PolyA mRNA was isolated from hybridoma 5G1.1. The acid/guanidinium thiocyanate procedure (Chomczynski and Sacchi, supra) was used to isolate total RNA, and was followed by oligo(dT)-cellulose chromatography of 1 mg of total RNA. For first strand cDNA synthesis, one microliter of the 25 microliters of oligo(dT)-cellulose eluate (containing approximately 2 micrograms of purified 5G1.1 mRNA) was denatured at 65° C. for 5 min., chilled on ice, and incubated in extension buffer (10 mM Tris pH 8.3, 50 mM KCl, 1 mM dithiothreitol, 240 µM each dNTP) containing 100 nM UDEC395 (SEQ ID NO:6) and 25 units AMV reverse transcriptase (Seikagaku America, Rockville, Md.) at 42° C. for one hour. Five microliters of the completed first strand reaction was subjected to PCR amplification using amplification buffer supplemented with 2.5 units AMPLITAQ DNA polymerase (Perkin Elmer, Foster City, Calif.) and 500 nM each of primer UDEC690 (SEQ ID NO:5) and UDEC395 (SEQ ID NO:6). Amplification was performed using 30 cycles each consisting of 1 minute at 95° C., 1 minute at 52° C., and 1 minute at 72° C., followed by a single ten minute incubation at 72° C.

The resulting PCR product was purified using GENECLEAN according to the manufacturer's directions (Bio 101, La Jolla, Calif.), digested with Sse8387 I and Hind III, gel purified, and ligated into the vector Bluescript II SK⁺ (Stratagene, La Jolla, Calif.). Ligated plasmids were transformed into the bacterial strain DH10B by electroporation.

Plasmid DNA was purified from cultures of transformed bacteria by conventional methods including column chromatography using a QIAGEN-TIP-500 column according to the manufacturer's directions (QIAGEN, Chatsworth, Calif.) and sequenced by the Sanger dideoxy chain termination method using SEQUENASE enzyme (U.S. Biochemical, Cleveland, Ohio). Clones obtained from a second independent PCR reaction verified that no mutations were introduced during the amplification process. The resulting plasmid containing the cloned variable region was designated SK (+) 690/395. This light chain encoding insert in this plasmid coded for both the N-terminal and internal light chain sequences determined by amino acid sequencing of 5G1.1, as described above.

Example 11

Construction and Expression of Recombinant mAbs

Recombinant DNA constructions encoding the recombinant mAbs comprising the 5G1.1 CDRs are prepared by conventional recombinant DNA methods including restriction fragment subcloning and overlapping PCR procedures. The resulting recombinant mAb-encoding DNAs include:

- (1) one encoding a non-humanized (murine) scFv designated 5G1.1M1scFv (SEQ ID NO:7), wherein CDR L1 is amino acid residues 28–34 of SEQ ID NO:7, CDR L2 is amino acid residues 52–54 of SEQ ID NO:7,

43

- CDR L3 is amino acid residues 93–98 of SEQ ID NO:7, CDR H1 is amino acid residues 156–159 of SEQ ID NO:7, CDR H2 is amino acid residues 179–183 of SEQ ID NO:7, and CDR H3 is amino acid residues 226–236 of SEQ ID NO:7;
- (2) one encoding a humanized (CDR grafted) scFv designated 5G1.1 scFv CB (SEQ ID NO:8), wherein CDR L1 is amino acid residues 26–36 of SEQ ID NO:8, CDR L2 is amino acid residues 52–58 of SEQ ID NO:8, CDR L3 is amino acid residues 91–99 of SEQ ID NO:8, CDR H1 is amino acid residues 152–161 of SEQ ID NO:8, CDR H2 is amino acid residues 176–192 of SEQ ID NO:8, H3 is amino acid residues 225–237 of SEQ ID NO:8;
- (3) one encoding a chimeric light chain (which can form the light chain portion of an Fab) designated 5G1.1M1 VL HuK (SEQ ID NO:9);
- (4) one encoding a chimeric Fd (the heavy chain portion of an Fab) designated 5G1.1M1 VH HuG1 (SEQ ID NO:10);
- (5) one encoding a humanized (CDR grafted and framework sequence altered) Fd designated 5G1.1 VH+IGHRL (SEQ ID NO:11), wherein CDR H1 is amino acid residues 26–35 of SEQ ID NO:11, CDR H2 is amino acid residues 50–60 of SEQ ID NO:11, and CDR H3 is amino acid residues 99–111 of SEQ ID NO:11;
- (6) one encoding a humanized (CDR grafted, not framework altered) Fd designated 5G1.1 VH+IGHRLC (SEQ ID NO:12), CDR H1 is amino acid residues 26–35 of SEQ ID NO:12, CDR H2 is amino acid residues 50–66 of SEQ ID NO:12, and CDR H3 is amino acid residues 99–111 of SEQ ID NO:12;
- (7) one encoding a humanized (CDR grafted and framework sequence altered) light chain designated 5G1.1 VL+KLV56 (SEQ ID NO:13), wherein CDR L1 is amino acid residues 26–36 of SEQ ID NO:13, CDR L2 is amino acid residues 52–58 of SEQ ID NO:13, and CDR L3 is amino acid residues 91–99 of SEQ ID NO:13;
- (8) one encoding a humanized (CDR grafted, not framework altered) light chain designated 5G1.1 VL+KLV56B (SEQ ID NO:14), wherein CDR L1 is amino acid residues 26–36 of SEQ ID NO:14, CDR L2 is amino acid residues 52–58 of SEQ ID NO:14, and CDR L3 is amino acid residues 91–99 of SEQ ID NO:14;
- (9) one encoding a humanized (CDR grafted, not framework altered) light chain designated 5G1.1 VL+012 (SEQ ID NO:15), wherein CDR L1 is amino acid residues 24–34 of SEQ ID NO:15, CDR L2 is amino acid residues 50–56 of SEQ ID NO:15, and CDR L3 is amino acid residues 89–97 of SEQ ID NO:15; and
- (10) one encoding a humanized (CDR grafted, not framework altered) Fd designated 5G1.1 VH+IGHRLD (SEQ ID NO:16), wherein CDR H1 is amino acid residues 26–35 of SEQ ID NO:16, CDR H2 is amino acid residues 50–60 of SEQ ID NO:16, and CDR H3 is amino acid residues 99–111 of SEQ ID NO:16.
- (11) one encoding a humanized (CDR grafted) scFv designated 5G1.1 scFv DO12 (SEQ ID NO:17), wherein CDR L1 is amino acid residues 26–36 of SEQ ID NO:17, CDR L2 is amino acid residues 52–58 of SEQ ID NO:17, CDR L3 is amino acid residues 91–99 of SEQ ID NO:17, CDR H1 is amino acid residues

44

- 152–161 of SEQ ID NO:17, CDR H2 is amino acid residues 176–186 of SEQ ID NO:17, and CDR H3 is amino acid residues 225–237 of SEQ ID NO:17;
- (12) one encoding a humanized (CDR grafted and framework sequence altered) scFv designated 5G1.1 scFv CO12 (SEQ ID NO:20), wherein CDR L1 is amino acid residues 26–36 of SEQ ID NO:20, CDR L2 is amino acid residues 52–58 of SEQ ID NO:20, CDR L3 is amino acid residues 91–99 of SEQ ID NO:20, CDR H1 is amino acid residues 152–161 of SEQ ID NO:20, CDR H2 is amino acid residues 176–192 of SEQ ID NO:20, H3 is amino acid residues 225–237 of SEQ ID NO:20;
- (13) one encoding a humanized (CDR grafted) scFv designated 5G1.1 scFv DO12B (SEQ ID NO:21), wherein CDR L1 is amino acid residues 26–36 of SEQ ID NO:21, CDR L2 is amino acid residues 52–58 of SEQ ID NO:21, CDR L3 is amino acid residues 91–99 of SEQ ID NO:21, CDR H1 is amino acid residues 152–161 of SEQ ID NO:21, CDR H2 is amino acid residues 176–192 of SEQ ID NO:21, H3 is amino acid residues 225–237 of SEQ ID NO:21;
- (14) one encoding a humanized (CDR grafted) scFv designated 5G1.1 scFv DO12C (SEQ ID NO:22), wherein CDR L1 is amino acid residues 26–36 of SEQ ID NO:22, CDR L2 is amino acid residues 52–58 of SEQ ID NO:22, CDR L3 is amino acid residues 91–99 of SEQ ID NO:22, CDR H1 is amino acid residues 152–161 of SEQ ID NO:22, CDR H2 is amino acid residues 176–192 of SEQ ID NO:22, H3 is amino acid residues 225–237 of SEQ ID NO:22;
- (15) one encoding a humanized (CDR grafted) scFv designated 5G1.1 scFv DO12D (SEQ ID NO:23), wherein CDR L1 is amino acid residues 26–36 of SEQ ID NO:23, CDR L2 is amino acid residues 52–58 of SEQ ID NO:23, CDR L3 is amino acid residues 91–99 of SEQ ID NO:23, CDR H1 is amino acid residues 152–161 of SEQ ID NO:23, CDR H2 is amino acid residues 176–192 of SEQ ID NO:23, H3 is amino acid residues 225–237 of SEQ ID NO:23;
- (16) one encoding a humanized (CDR grafted and framework sequence altered) scFv designated 5G1.1 scFv CO13 (SEQ ID NO:24), wherein CDR L1 is amino acid residues 26–36 of SEQ ID NO:24, CDR L2 is amino acid residues 52–58 of SEQ ID NO:24, CDR L3 is amino acid residues 91–99 of SEQ ID NO:24, CDR H1 is amino acid residues 152–161 of SEQ ID NO:24, CDR H2 is amino acid residues 176–192 of SEQ ID NO:24, H3 is amino acid residues 225–237 of SEQ ID NO:24;
- (17) one encoding a humanized (CDR grafted and framework sequence altered) scFv designated 5G1.1 scFv CO14 (SEQ ID NO:25), wherein CDR L1 is amino acid residues 26–36 of SEQ ID NO:25, CDR L2 is amino acid residues 52–58 of SEQ ID NO:25, CDR L3 is amino acid residues 91–99 of SEQ ID NO:25, CDR H1 is amino acid residues 152–161 of SEQ ID NO:25, CDR H2 is amino acid residues 176–192 of SEQ ID NO:25, H3 is amino acid residues 225–237 of SEQ ID NO:25;
- (18) one encoding a humanized (CDR grafted and framework sequence altered) scFv designated 5G1.1 scFv CO15 (SEQ ID NO:26), wherein CDR L1 is amino acid residues 26–36 of SEQ ID NO:26, CDR L2 is amino acid residues 52–58 of SEQ ID NO:26, CDR L3 is amino acid residues 91–99 of SEQ ID NO:26, CDR H1

is amino acid residues 152–161 of SEQ ID NO:26, CDR H2 is amino acid residues 176–192 of SEQ ID NO:26, H3 is amino acid residues 225–237 of SEQ ID NO:26;

In accordance with the invention, one each of the various L1, L2 and L3 CDRs discussed in (1) to (18) above may be combined with any of the other light chain CDRs so as to make a set of 3 light chain CDRs comprising one L1, one L2, and one L3 CDR, as part of a recombinant antibody or synthetic peptide antibody (i.e., a synthetic peptide with the sequence of a recombinant peptide of the invention). Furthermore, the framework regions (i.e., the regions not included in the CDRs as described for each) of each of (1) to (18) above may be interchanged with homologous framework regions of the other recombinant antibody molecules of (1) to (18) to produce other antibodies of the invention.

In accordance with the invention, one each of the various H1, H2 and H3 CDRs discussed in (1) to (18) above may be combined with any of the other light chain CDRs so as to make a set of 3 light chain CDRs comprising one H1, one H2, and one H3 CDR, as part of a recombinant antibody or synthetic peptide antibody (i.e., a synthetic peptide with the sequence of a recombinant peptide of the invention).

In accordance with the invention, matched pairs of the variable regions (e.g., a VL and a VH region) of the various antibody molecules, Fds, and light chains described above may be combined with constant region domains by recombinant DNA or other methods known in the art to form full length antibodies of the invention. Particularly preferred constant regions for this purpose are IgG constant regions, which may be unaltered, or constructed of a mixture of constant domains from IgGs of various subtypes, e.g., IgG1 and IgG 4.

Matched pairs of the Fd and light chain encoding DNAs described immediately above—i.e. (3) and (4), (5) and (7), (6) and (8), and (6) and (9)—were subcloned together into the APEX-3P vector, essentially as described below in Example 15 for N19/8. The scFv constructs of (1) and (2) were subcloned into pET Trc SO5/NI using conventional techniques.

Plasmids so obtained were introduced by into the bacterial strain ME2 (pET plasmids) by conventional electroporation, or into human 293 EBNA cells (APEX plasmids) by lipofection using 2–3 microliters of TRANSFECTAM reagent (Promega, Madison, Wis.) per microgram of DNA according to the manufacturer's directions. Bacterial strains ME1 and ME2 are derivatives of *Escherichia coli* strain W3110 (ATCC designation 27325) prepared as follows.

Preparation of W3110 Derivatives ME1 and ME2

The non-humanized, non-chimeric murine 5G1.1-scFv “m5G1.1-scFv”—made up of light chain (3) and Fd (4)—was expressed in a derivative of *E. coli* K12 strain W3110. This derivative was prepared by inactivating an uncharacterized gene to provide protection against infections by a lytic bacteriophage. *E. coli* strain W3110 is a particularly preferred strain because it is fully characterized and is commonly used for recombinant DNA product fermentations.

A single colony of *E. coli* strain W3110 was grown overnight in L medium at 30° C. The cells were collected by centrifugation and resuspended in 10 mM MgSO₄. A total of 0.1 ml of the culture was added to 2.5 ml 0.7% L soft agar at 45° C. and quickly poured on an L plate. Fifty microliter aliquots of a plaque purified phage lysate, undiluted, diluted 10⁻² and diluted 10⁻⁴, were spotted onto the agar surface. Phage lysates had previously been filtered through 0.45 µm membranes and stored in sterile tubes with a drop of

chloroform at 4° C. The spots were allowed to dry on the soft agar surface and incubated overnight at 37° C.

The next day L plates were spread with 10⁹ phage PFU and allowed to dry. Using a sterile, flat toothpick, cells from isolated colonies growing in the zones of phage lysis on the spot plates were streaked for single colonies on the plates spread with 10⁹ phage PFU and incubated overnight at 37° C. Single colonies were rechecked for phage resistance by cross-streaking after single colony purification. The cross streak test for phage sensitivity was performed as follows. Fifty µl of phage (10⁸ pfu/ml) was spread in a vertical line in the left hand portion of the plate using a Pasteur pipette. Additional phage were tested parallel to the first and to the right. The plate was allowed to dry, and strains to be checked for sensitivity or resistance were spread perpendicular to and across the lines of all phages in a single swath from the left to the right. Resistant strains grow in the area of the phage streaks while sensitive strains lyse.

The phage resistant mutant strain ME1 was tested for phage production after overnight growth in L medium and treatment with the DNA damaging agent, mitomycin C. The strain failed to produce viable phage utilizing a standard plaque assay and *E. coli* W3110 as the phage sensitive indicator strain. These results suggest that strain ME1 does not harbor a resident prophage.

Strain ME2 was constructed by site specific integration of the lambdaDE3 prophage (Studier et al. 1990, Meth. Enzymol. 185:60–89) into the ME1 chromosome. Expression of the T7 RNA polymerase, directed by the prophage, allows expression of target genes cloned into pET vectors (Studier et al., supra) under the control of the T7 promoter in the lysogenized host. Lysogenization was accomplished in a three way infection with lambdaDE3, the lambda helper phage, lambdaB10 and the selection phage, lambdaB482 (Studier et al., supra).

lambdaDE3 (imm21) was constructed by Studier and colleagues (1990, Meth. Enzymol. 185:60–89) by inserting the T7 RNA polymerase gene behind the *E. coli* lacUV5 promoter into the BamHI cloning site of lambdaD69 (imm21). Since cloning into the BamHI site of lambdaD69 interrupts the integrase gene, lambdaDE3 cannot integrate or excise from the chromosome by itself. The helper phage lambdaB10 provides the integrase function that lambdaDE3 lacks but cannot form a lysogen by itself. The selection phage, lambdaB482, lyses any lambdaDE3 host range mutants that otherwise would be among the surviving cells, but it can neither integrate into susceptible cells nor lyse lambdaDE3 lysogens since it has the same immunity region as lambdaDE3 (imm21).

Lysoaenization Protocol

Strain ME1 was grown in L medium supplemented with 0.2% maltose and 10 mM MgSO₄ at 37° C. to a density of approximately 10⁸ cells/ml. One µl of ME1 cells were incubated with 2×10⁸ plaque forming units (pfu) of lambdaDE3 and 10⁸ pfu of lambdaB10 and lambdaB482. The host/phage mixture was incubated at 37° C. for 20 min to allow phage adsorption to ME1 cells. Several dilutions of the cell/phage suspension were spread on L plates to produce plates containing approximately 30–200 candidate lysogens as isolated colonies. The plates were inverted and incubated at 37° C. overnight. Several isolated colonies were screened for the acquisition of the lambdaDE3 prophage as described below.

Verification of lambdaDE3 Lysogens

lambdaDE3 lysogen candidates were tested for their ability to support the growth of the T7 phage 4107, a T7 phage deletion mutant that is completely defective unless active T7

RNA polymerase is provided in trans. Only lambdaDE3 lysogens will support the normal growth of the phage in the presence of IPTG (isopropyl-beta-thiogalactopyranoside). The T7 phage produces very large plaques on lambdaDE3 lysogens in the presence of IPTG, while very small plaques are observed in the absence of inducer. The size of the plaque in the absence of IPTG is an indication of the basal level of T7 RNA polymerase expression in the lysogen. Putative lambdaDE3 lysogens were grown in L broth supplemented with 0.2 % maltose and 10 mM MgSO₄ at 37° C. to a cell density of approximately 10⁸ cells/ml. A total of 0.5 ml of cells was centrifuged and the pellet was resuspended in 0.2 ml of a T7 phage lysate containing 2x10⁴ pfu. The phage was allowed to adsorb for 30 min at 37° C. One-half of suspension (0.1 ml) was added to 3.0 ml of molten top agarose at 47° C. and poured onto L plates. The remaining aliquot of cell/phage suspension was poured onto an L plate supplemented with 0.4 mM IPTG to check for induction of T7 RNA polymerase. The plates were inverted and incubated at 37° C. overnight.

Strains were also tested for the presence of the lambdaDE3 lysogen by demonstrating that each strain was resistant to infection by the phage lambdaB482, which is in the same immunity group (imm21), by the cross streak method described above. A lysogen was chosen with a low basal expression level for protein production from pET vectors. The resulting strain, designated ME2, is phage resistant and overexpresses T7 RNA polymerase in the presence of IPTG. Purification of Humanized 5G1.1-scFv from *E. coli*

The humanized 5G1.1-scFv (h5G1.1-scFv) cDNA construct was cloned into the bacterial expression plasmid pET Trc SO5/NI (SEQ ID NO:18) and transformed into *E. coli* strain ME1. The resulting strain expressing h5G1.1 scFv was grown at 37° C. in 2 liter Applikon glass vessel fermentors containing Terrific Broth (1.2 % (w/v) bacto-tryptone, 2.4% (w/v) bacto-yeast extract, 0.4% (v/v) glycerol, 90 mM KPO₄, pH 7.0) supplemented with 100 µg/ml ampicillin. The production of recombinant scFv was induced by the addition of 1 mM IPTG when the O.D.₅₅₀ of the culture reached 10. After an additional 3 h incubation at 37° C., the cells were harvested by centrifugation and the cell pellets stored at -80° C.

Cells were resuspended in 1 mM EDTA, pH 5.0 at 10 ml per gram weight and lysed by a single pass through a microfluidizer (Model M110T, Microfluidics Corp., Newton, Mass.). After centrifugation at 17,500xg for 15 min, the resulting inclusion body pellet was washed by resuspension in 20 mM Tris-HCl pH 8.0, 1 mM EDTA, 100 mM NaCl, 0.15% (w/v) deoxycholate at 10 ml per gram inclusion body using a Tekmar POLYTRON. The inclusion bodies were again pelleted by centrifugation at 17,500xg for 15 min and resuspended in 20 mM Tris-HCl pH 9.0, 8 M urea at 10 ml per g. After stirring for 1 h, the sample was centrifuged at 14,000xg for 30 min to pellet remaining insoluble material.

The extract supernatant was diluted 10-fold with 20 mM Tris-HCl pH 9.0, 7 M urea, 50 µM cupric sulfate and allowed to stir for at least 16 hours at 4° C. to refold the scFv. After addition of BIOCRYL BPA-1000 (TosoHaas, Montgomeryville, Pa.) as a flocculating agent at 3 µl per ml, the sample was centrifuged at 15,000xg for 10 minutes to remove insoluble material. The refolding mixture was exchanged into 20 mM Tris, pH 9.0, 1 mM EDTA by diafiltration and concentrated by ultrafiltration using a stirred cell fitted with a YM10 membrane (Amicon, Beverly, Mass.).

In subsequent experiments, other refolding conditions were tested. Thawed bacterial cells were resuspended with a

POLYTRON homogenizer in 1 mM EDTA at 2.5 mL per gram of cells, passed through the MICROFLUIDIZER at 18,000 psi, and centrifuged at 10,000 RPM for 15 min in a Beckman JA-10 rotor, the resulting pellet was washed by resuspension in 20 mM Tris-HCl pH 8.0, 1 mM EDTA, 100 mM NaCl, 0.15% (w/v) deoxycholate at 10 ml per gram inclusion body using a Tekmar POLYTRON. The inclusion bodies were again pelleted by centrifugation. The pellet from this centrifugation was resuspended with a POLYTRON homogenizer in 8M urea, 20 mM Tris pH9 at 10 mL per gram of pellet. After stirring for 1 hour at 4 degrees C., the resuspended pellet was diluted with 9 volumes of 7M urea, 20 mM Tris pH9. Cupric sulfate was then added to various final concentrations (0, 5, 10, 20, 25, 30, 40, 50, 100, 150, and 200 µM) before incubation overnight at 4 degrees C. with stirring. The use of 5 µM copper was found to give the highest levels of refolding of the humanized 5G1.1-scFv as assessed by analytical HPLC.

In the initial experiments, the properly refolded scFv was then separated from aggregated material and contaminating proteins by anion exchange chromatography using Q SEPHAROSE FAST FLOW (Pharmacia, Piscataway, N.J.). Bound scFv was eluted with 20 mM Tris-HCl pH 9.0, 1 mM EDTA containing a linear NaCl gradient (0 to 0.5 M). The fractions containing the scFv were combined, concentrated by ultrafiltration using a stirred cell fitted with a YM10 membrane, and applied to a SEPHACRYL S200 HR 26/100 gel filtration column (Pharmacia) equilibrated in 20 mM Tris-HCl pH 9.0, 1 mM EDTA, 150 mM NaCl. Fractions containing the scFv were combined, exchanged into phosphate-buffered saline by diafiltration, concentrated by ultrafiltration, filtered through a 0.22 µm MILLEX-GV filter (Millipore, Bedford, Mass.), and stored at 4° C.

Subsequent experiments have indicated that cation exchange chromatography (e.g., using POROS HS resin—PerSeptive Biosystems, Cambridge, Mass.) should give better yields than the Q Sepharose Fast Flow anion exchange chromatography step described in the preceding paragraph. In addition, it would be preferable to carry out the final gel filtration chromatography in a buffer that is more pharmaceutically acceptable than the Tris buffer described. A buffer such as PBS would be preferred if it does not interfere with the efficacy of the gel filtration chromatographic separation. This would reduce any trace amounts of Tris remaining in the preparation after diafiltration, and might eliminate the need for the diafiltration step.

Purification of m5G1.1-scFv from *E. coli*

Frozen bacterial cell paste was thawed and resuspended in 2.5 ml of 1 MM EDTA (pH 5) per gram of cell paste. This suspension of cells was lysed by passage through a MICROFLUIDIZER (Microfluidics) with the interaction chamber in line and a backpressure of approximately 18000 psi. The cell lysate was then centrifuged at 10,000 rpm in a JA-10 centrifuge rotor at 4° C. for 15 min. The supernatant was decanted and discarded.

The pellet was resuspended in 10 ml of 20 mM Tris, pH 8.0, 100 mM NaCl, 0.15% sodium deoxycholate per gram of pellet. This suspension was centrifuged as above for 10 min. Again the supernatant was decanted and discarded. This detergent washed pellet was then resuspended in 10 ml of 8 M urea, 20 mM Tris-HCl, pH 9 (1 mM EDTA may also be added to this buffer, but has the effect of increasing the time required to achieve a particular level of refolding). The suspension was stirred at 4° C. for 1 hr. and was then diluted 10 fold with 7 M urea, 20 mM Tris-HCl, pH 9 and stirred at 4° C. CUSO₄ was then added to a final concentration of 50 µM and stirring was continued overnight at 4° C.

The majority of contaminating proteins (including incorrectly folded versions of m5G1.1 scFv) were then removed by precipitation by diluting (with stirring) the refolded sample five fold with buffer such that the final concentrations after dilution were 1.4 M urea, 25 mM NaCl, 1 mM EDTA, and 20 mM sodium acetate at 4° C. The pH of the dilution buffer when prepared at room temperature was pH 5.0. Prior to dilution the pH of the dilution buffer is determined at 4° C. After the dilution the pH of the sample was greater than pH 5.5. The pH of the sample was then adjusted with 6. N HCl to the initial pH 5.0 of the buffer at 4° C. The solution immediately became cloudy and it was left stirring at 4–8° C. for 0.5 to 24 hours.

The precipitate was removed by filtering the sample through a 300 kDa cut-off ultrafiltration membrane (Millipore Corporation, Bedford, Mass.). The permeate was collected and concentrated 5 fold using a 10 kDa cutoff ultrafiltration membrane (Millipore). This concentrated retentate was then diluted 2 fold with 20 mM sodium acetate, 1 mM EDTA, pH 5.0 in order to lower the NaCl concentration to 12.5 mM.

The diluted retentate was then loaded at 4° C. onto a SP SEPHAROSE FF column (Pharmacia) equilibrated in 0.7 M urea, 1 mM EDTA, 10 mM NaCl, 20 mM sodium acetate, pH 5.0, at a linear flowrate of 5 cm/min. Bed height was equal to or greater than 3.5 cm. Following loading the column was washed with 40 column volumes (CV) of equilibration buffer. The column was then washed with 20 CV of 20 mM sodium acetate, pH 5.0, 1 mM EDTA. The bound scFv was then eluted using 20 mM sodium citrate, pH 5.8, 1 mM EDTA. A single peak was collected in approximately 4 column volumes.

The SP SEPHAROSE eluate was then adjusted to 20 mM Tris-HCL by addition of 1 M Tris-HCL, pH 8. The pH of the sample was adjusted to 8.0 by addition of 1 N NaOH. This sample was loaded onto a Q SEPHAROSE FF column (Pharmacia) equilibrated in 20 mM Tris-HCL, pH 8.0, 1 mM EDTA at room temperature at a flowrate of 5 cm/min. The flow through fraction containing the scFv was collected.

The Q SEPHAROSE flow through fraction was then adjusted to 150 mM NaCl and concentrated to 10 mg of scFv per ml using a 10 kDa cutoff ultrafiltration membrane. This concentrated sample was then loaded onto a SEPHACRYL S200 column equilibrated in phosphate buffered saline, pH 7.4 and eluted at 0.4 cm/min. The fractions were analyzed by SDS-PAGE and silver staining. Peak fractions were combined after discarding the front and back shoulder fractions that contained the majority of contaminants.

Example 12

Functional Analysis of the m5G1.1 scFv

Titration of the m5G1.1 scFv in hemolytic assays revealed that the m5G1.1 scFv inhibited human complement-mediated lysis in a dose dependent fashion (FIG. 13). Direct comparison of the efficacy of the m5G1.1 scFv to the 5G1.1 mAb and Fab demonstrated that the m5G1.1 scFv completely blocked C5b-9-mediated hemolysis in 20% human serum at 0.15 μ M while the 5G1.1 mAb and Fab blocked at 0.06–0.08 μ M. Analysis of C5a generation in these assays revealed similar results in that the 5G1.1 scFv completely blocked C5a generation at 0.15 μ M while the 5G1.1 mAb and Fab blocked at 0.06–0.08 μ M (FIG. 14). Taken together these experiments indicated that unlike N19/8, which lost half of its effectiveness at blocking C5a generation upon being engineered as an scFv (SEQ ID NO:19), the 5G1.1 murine scFv retained the capacity to block the generation of both C5a and C5b-9.

Additionally, these data demonstrate that the m5G1.1 scFv retained similar activity to that of the parent molecule (the native murine 5G1.1 mAb) in that the molar concentration of 5G1.1 murine scFv required to completely block C5a and C5b-9 (0.15 μ M) was within two-fold of that required for the 5G1.1 mAb and Fab (0.06–0.08 μ M).

In order to determine whether the m5G1.1 scFv retained the capacity to block the activation of complement in the ex vivo model of cardiopulmonary bypass, 4.5 mg of the purified bacterially produced 5G1.1 murine scFv was added to the CPB circuit and complement activation was monitored. In control experiments, both C3a and C5b-9 levels increased throughout the time course of the experiment. In a single experiment, addition of 4.5 mg of the m5G1.1 scFv to the CPB circuit had no effect on the generation of C3a (FIG. 15). Conversely, complement hemolytic activity as well as the generation of sC5b-9 was completely blocked in this experiment (FIG. 16 and FIG. 17).

Example 13

Characterization of the Epitope Recognized by 5G1.1

Tryptic digestion: Twenty micrograms of purified human C5 (Advanced Technologies, San Diego, Calif.) was subjected to enzymatic digestion with 1 μ g of TPCK-treated trypsin (Worthington Biochemical Corp., Freehold, N.J.). The digestion was allowed to continue for 3 minutes, after which time it was stopped by the addition of 20 μ g soy bean trypsin inhibitor (Worthington). The reaction was then denatured and reduced by the addition of protein sample buffer and immediately boiled for 5 min. The digested fragments were size fractionated through a SDS-PAGE on a 12 % gel. The gel was then electroblotted in transfer buffer (20% (v/v) methanol, 25 mM Tris-base pH 8.0, and 192 mM glycine) to nitrocellulose (Bio-Rad Laboratories, Hercules, Calif.) and subjected to ECL western blot analysis using either 5G1.1 or a C5a specific monoclonal antibody (G25/2, obtained from Dr. Otto Götze, Department of Immunology, University of Göttingen, Germany).

The filters were incubated twice for 30 minutes each in blocking solution (500 mM NaCl, 5 mM Tris p-H 7.4, 10% (v/v) nonfat dry milk, and 0.2% (v/v) TWEEN-20). The filters were then changed to fresh blocking solution (20 ml) containing the primary antibody and incubated for 40 minutes on a rocking platform. The filters were rinsed briefly with washing solution (500 mM NaCl, 35 mM Tris pH7.4, 0.1% SDS, 1% NP40, and 0.5% deoxycholic acid) to remove any milk, and then fresh wash solution was added and incubated for two 20 minute intervals on an orbiting shaker. The filters were rinsed briefly with 10 to 20 mls of secondary antibody solution (500 mM NaCl, 5 mM Tris pH 7.4, 10% (v/v) Nonfat dry milk, 0.2% (v/v) TWEEN-20, and 1% NP-40) and then incubated with fresh secondary antibody solution containing a 1:2000 dilution of HRP conjugated goat anti-mouse for 20 minutes on a rocking platform. The filters were then washed as described above, incubated in ECL reagent (Amersham Corp., Arlington Heights, Ill.) for 1 minute and then exposed to ECL HYPERFILM (Amersham).

Acid Hydrolysis: Twenty micrograms of purified human C5 (Advanced Technologies) was subjected to hydrolysis in 1N acetic acid. The 20 μ g of human C5 (1 μ g/ μ l) was added to 20 μ l of 2N acetic acid and incubated for 10 min at 100° C. The sample was denatured and reduced with protein sample buffer, also at 100° C., for 5 minutes. The acid was

51

neutralized by dropwise addition of a saturated tris base solution until the sample turned blue. The cleavage products were then size fractionated by SDS-PAGE and western blotted as described above. For N-terminal sequencing, the gel fractionated acid hydrolysate was transferred to PVDF membrane. N-terminal sequence was obtained by excising the 46 kDa acid hydrolysis fragment band from a PVDF membrane and subjecting it to amino acid sequence analysis as discussed above in Example 10.

Deglycosylation: Reduced and denatured acid hydrolyzed or tryptic fragments of human C5 were subjected to deglycosylation with N-Glycosidase F (Peptide-N-Glycosidase F, Boehringer Mannheim Corp., Indianapolis, Ind.) according to the manufacture's directions.

Results: Acid hydrolysis of human C5 yielded a fragment with an apparent molecular weight by SDS-PAGE of 46 kDa that was immunoreactive for both the anti-C5a mAb G25/2 and the anti-C5 alpha chain mAb 5G1.1. Western blots probed with both antibodies simultaneously, as well as silver stain SDS-PAGE analysis, confirmed the presence of a single 46 kDa fragment that was immunoreactive with both antibodies. The presence of a single immunoreactive fragment containing binding sites for both 5G1.1 and G25/2 strongly suggested that the 5G1.1 epitope was contained within approximately the first 46 kDa of the N-terminus of the alpha chain of C5.

As discussed above in the description of the complement system under the heading "Background Physiology & Pathology," a compound (e.g., an antibody) that binds to a site at or immediately adjacent to the C5a cleavage site would have the potential to act as a terminal complement inhibitor. The potential inhibitory activity of antibodies binding to this site led to the expectation that the C5 alpha chain-binding 5G1.1 antibody would bind to an epitope at or near the C5a cleavage site. The finding that 5G1.1 bound to the 46 kDa acid hydrolysis fragment of C5 lent support to this expectation.

Western blot analysis of the tryptic digestion products identified one proteolytic fragment migrating at approximately 27 kDa that was immunoreactive with 5G1.1. Likewise, one immunoreactive proteolytic fragment migrating at approximately 29 kDa was observed following western blot analysis with the anti-C5a mAb G25/2. Experiments in which a blot was simultaneously probed with both 5G1.1 and G25/2 demonstrated that each band was distinct and that their apparent differential mobility was not a gel anomaly. This was surprising, because the 5G1.1 mAb was thought likely to bind to the C5 convertase cleavage site. 5G1.1 was thus expected to be immunoreactive with any fragment of C5 of over 12 kDa that exhibited immunoreactivity with G25/2. Such a fragment would contain enough of the extreme amino terminus of the C5 alpha chain to bind specifically to the anti-C5a mAb, and enough beyond that to encompass a region including and extending beyond the C5 convertase cleavage site.

The immunoreactivity of G25/2 with the 29 kDa fragment indicated that that fragment contains the N-terminal region of the alpha chain of C5 that is cleaved off to yield C5a. Furthermore, because 5G1.1 was not immunoreactive with this band, the 5G1.1 epitope was not likely to be contained within approximately the first 29 kDa of the N-terminus of the alpha chain of C5, and therefore could not be located near the C5 convertase cleavage site.

These tryptic digestion and acid hydrolysis mapping data suggested that the 5G1.1 epitope was contained within a region starting about 29 kDa (including post-translational

52

modifications) from the N-terminus of the alpha chain of C5 and continuing 17 kDa in a C-terminal direction, i.e., ending 46 kDa from the N-terminus, a surprising finding in view of the expectation, discussed above, that the antibody would bind at or immediately adjacent to the point at which C5a is cleaved off of the C5 alpha chain, i.e., at or immediately adjacent to amino acid residue 733 of SEQ ID NO:2.

Post-translational modifications can alter the mobility of proteins during SDS-PAGE electrophoresis. One such modification is the addition of carbohydrate via N-linked glycosylation. As discussed above under the heading "Background Physiology & Pathology", C5 is glycosylated, as is C5a. C5a is glycosylated at an asparagine residue corresponding to amino acid number 723 of the full length pro-C5 precursor of human C5 (SEQ ID NO:2).

Computer analysis of the human C5 alpha chain suggests potential N-linked glycosylation sites at positions corresponding to amino acid numbers 893, 1097, and 1612 of SEQ ID NO:2. In order to determine the contribution of carbohydrate to the electrophoretic mobility of both the tryptic and acid fragments, enzymatic deglycosylation of the fragments was performed and followed by western blot analysis. It was determined that each tryptic fragment lost approximately 3 kDa in apparent molecular weight while the acid fragment lost approximately 6 kDa.

This result was interpreted as indicating that the tryptic fragments were each glycosylated at a single site and that the 46 kDa acid fragment was glycosylated at two sites (one of which was the known glycosylation site in C5a referred to above). The diminished mobility observed following deglycosylation agrees with the computed prediction of a second N-linked glycosylation site within the first 233 amino acids of the C5 alpha chain.

N-terminal sequence analysis determined that the first four amino acids of the 46 kDa fragment generated by 1N acetic acid treatment was Thr Leu Gln Lys. This sequence is found only once in the full length human pro-C5 precursor molecule—at a position corresponding to amino acids 660 through 663 of SEQ ID NO:2. This four amino acid sequence also corresponds to the sequence of the amino-terminus of the alpha chain of human C5 and, thus to the amino-terminus of human C5a.

In order to more precisely map the binding site of 5G1.1, overlapping peptide analysis was performed. The sequence predicted to be contained within the 17 kDa section of human C5 described above (SEQ ID NO:2; amino acids 893 through 1019) together with an extension of 43 amino acids towards the N-terminus and 30 amino acids towards the C-terminus (a total of 200 amino acids) was synthesized as a series of 88 overlapping peptides by solid phase synthesis on polypropylene filters (Research Genetics Inc., Huntsville, Ala.).

The 43 and 30 amino acid extensions were added to allow for possible inaccuracies in the prediction of the span of this 17 kDa region. Such inaccuracies are likely due to the uncertainty of the specific extent of glycosylation of each of the various regions of C5a, as well as to the aberrant gel mobility that is commonly seen when highly charged polypeptides (such as the 5G46k fragment and the 5G27k fragment) are analyzed by SDS-PAGE. As discussed above in the Summary of the Invention, a 200 amino acid peptide corresponding to the region covered by these overlapping peptides is referred to herein as the "5G200aa" peptide.

Because of the expectation that the 5G1.1 antibody would bind at the C5a cleavage site, an additional set of 8 overlapping peptides was synthesized that spanned a 30 amino

53

acid section spanning the C5a cleavage site (amino acids 725 through 754 of SEQ ID NO:2). A peptide having the sequence of this 30 amino acid section is referred to herein as the "cleavage site peptide". A 325aa peptide spanning amino acid residues 725–1049 of SEQ ID NO:2 (this peptide spans the region covered by the cleavage site peptide and the 5G200aa peptide) is referred to herein as the "5G325aa" peptide.

These filters were probed with 5G1.1 as described above for ECL western blot analysis, and a set of 4 overlapping peptides spanning the region corresponding to amino acid residues 3–19 of the KSSKC peptide (SEQ ID NO:1) each gave a positive signal indicative of monoclonal antibody binding, while peptides corresponding to the C5a cleavage site did not bind to the 5G1.1 antibody.

Example 14

C3/C4 Binding Assay

C3 and C4 are both key components of classical C5 convertase, and C3 is also a key component of alternative C5 convertase. These C5 convertases are required for the conversion of C5 to active C5a and C5b. The ability to block C5 binding to C3 and C4 is thus a desirable property for an antibody to be used in treatment of complement mediated diseases in accordance with the present invention.

96 well microtiter plates were coated with 501 μ /well, 10 μ g/ml of either purified human C3 or C4 (Quidel) for 1 hour at 37° C. The plates were then blocked with 200 μ l/well of TBS containing 1% BSA for 1 hour at room temperature. After three washes in TBS 0.1% BSA, purified human C5 (Quidel, 20 μ g/ml in TBS 1% BSA) was added to the plates in the presence (20 μ g/ml) or absence of a 5G1.1 Fab (derived from 5G1.1 by conventional papain digestion) and allowed to incubate for 1 hour at 37° C. After three washes in TBS/0.1% BSA, a monoclonal antibody directed against the C5 beta chain (N19/8, 5 μ g/ml) was added to the wells to detect C5 bound to either C3 or C4. After three final washes in TBS/0.1% BSA, the plate was developed using a horseradish peroxidase-conjugated secondary antibody and the appropriate substrate.

The results of these assays showed that the 5G1.1 mAb inhibited the binding of purified human C5 to either C3 or C4 by at least 60% to 90%. As used herein and in the claims, such a 60% to 90% reduction in C3 or C4 binding is a "substantial reduction" in C3 or C4 binding.

Example 15

Construction and Functional Analysis of N19/8 Chimeric Fab

The heavy chain and light chain variable regions from the hybridoma N19-8 were cloned by PCR using the Ig-Prime System (Novagen) as described by the manufacturer. Clones from multiple independent PCR reactions were sequenced to detect mutations introduced during the PCR amplification. An N19-8 VL/human kappa constant region chimeric cDNA was created by using a plasmid containing the N19-8 light chain variable region and the plasmid pHuCK (Hieter et al., 1980 Cell, 22:197–207) as templates in an overlapping PCR reaction.

Similarly, an N19-8 VH/human IgG1 Fd chimeric cDNA was created using a plasmid containing the N19-8 heavy chain variable region and a plasmid containing the human IgG1 gene (obtained from Ilan R. Kirsch, National Cancer Institute, Bethesda, Md.) as templates. This Fd construct

54

contained the first nine amino acids of the IgG1 hinge region, including the cysteine residue which normally forms a disulfide bond with the terminal cysteine residue of the kappa light chain.

The resulting chimeric cDNAs were separately cloned into the APEX-1 vector using appropriate flanking restriction enzyme sites introduced during the PCR amplification procedure and sequenced. A fragment containing the promoter, intron, and cDNA insert from one of these APEX vectors was subsequently subcloned into the polylinker of the other to produce a single vector directing the expression of both the light chain and Fd. The tandem expression cassette from this APEX-1 vector was subsequently subcloned into APEX-3P, which was transfected into 293 EBNA cells for expression of the chimeric Fab.

When tested for the ability to block complement hemolytic activity and C5a generation, the chimeric N19/8 Fab retained the ability to block hemolytic activity, but lost 50% of its C5a generation blocking capacity.

Throughout this application various publications and patent disclosures are referred to. The teachings and disclosures thereof, in their entirety, are hereby incorporated by reference into this application to more fully describe the state of the art to which the present invention pertains.

Although preferred and other embodiments of the invention have been described herein, further embodiments may be perceived by those skilled in the art without departing from the scope of the invention as defined by the following claims.

REFERENCES

- Baker, et al., 1989, *American Journal of Pathology*. 135, pp. 185–194.
- Clackson, et al., 1991, *Nature*. 352, pp. 624–628.
- Cochrane, et al., 1965, *Journal of Experimental Medicine*. 122, pp. 99–116.
- Coligan, et al., eds. 1992, *Current Protocols in Immunology*. John Wiley & Sons, New York.
- Couser, et al., 1985, *Kidney International*. 28, pp. 897–890.
- Couser, et al., 1991, *Journal of the American Society of Nephrology*. 2, pp. 894–901.
- Couser, 1992, in *Cecil Textbook of Medicine*. 19th Ed. (Wyngaarden, Smith, and Bennett, eds.) W.B. Saunders Co., Philadelphia, Pa., Ch. 79, pp. 551–568.
- Couser, et al., 1992, *Nephrology Dialysis Transplantation*. Suppl. 1, pp. 25–30.
- Couser, 1993, *Kidney International*. 44, Suppl. 42, pp. S19–S26.
- Falk and Jennette, 1986, *Kidney International*. 30, pp. 678–686.
- Fearon, 1983, in *Intensive Review of Internal Medicine*, 2nd Ed.; Fanta and Minaker, eds. Brigham and Women's and Beth Israel Hospitals, pp. 204–210.
- Floege, et al., 1992, *Laboratory Investigation*. 67, pp. 486–497.
- Frei, et al., 1987, *Molecular and Cellular Probes*. 1, pp. 141–149.
- Glasscock and Brenner, 1987, in *Harrison's Principles of Internal Medicine*, 11th Ed. (Braunwald, Isselbacher, Petersdorf, Wilson, Martin, and Fauci, eds.) McGraw-Hill Book Co., New York, N.Y., Ch. 222 & 223, pp. 1170–1189.
- Glasscock and Brenner, 1994, in *Harrison's Principles of Internal Medicine*, 13th Ed. (Isselbacher, Braunwald, Wilson, Martin, Fauci, and Kasper, eds.) McGraw-Hill, Inc., New York, N.Y., pp. 1292–1313.

55

- Grogel, et al., 1983, *Journal of Clinical Investigations*. 72, pp. 1948–1957.
- Guyton, 1971, *Textbook of Medical Physiology*, 4th Ed. W.B. Saunders Co., Ch. 34 & 38, pp. 393–405 & pp. 442–454.
- Haber, 1992, *Immunology Review*. 130, pp. 189–212.
- Harlow and Lane, 1988, *Antibodies, A Laboratory Manual*, Cold Spring Harbor Laboratory, New York.
- Jennette, et al., 1987, *American Journal of Pathology*. 127, pp. 499–506.
- Knicker and Cochrane, 1965, *Journal of Experimental Medicine*. 122, pp. 83–98.
- Liddell and Cryer, 1991, *A Practical Guide to Monoclonal Antibodies*, John Wiley & Sons, Chichester, West Sussex, England.
- Mollnes, et al., 1988, *Scandinavian Journal of Immunology*. 28, pp. 307–312.
- Montz, et al., 1990, *Cellular Immunology*. 127, pp. 337–351.
- Morrison, et al., 1992, *Annual Review of Immunology*. 10, pp. 239–265.
- Passwell, et al., 1988, *The American Society for Clinical Investigation, Inc.* 82, pp. 1676–1684.
- Reichmann, et al., 1988, *Nature*. 332, pp. 323–327. *Remington's Pharmaceutical Sciences*. 17th Ed. 1985, Mack Publishing Company, Philadelphia, Pa.
- Rich, 1992, in *Cecil Textbook of Medicine*, 19th Ed. (Wyngaarden, Smith, and Bennett, eds.) W.B. Saunders Co., Philadelphia, Pa., Ch. 249, pp. 1467–1470.
- Robbins and Cotran, 1979, *Pathologic Basis of Disease*. 2nd ed. W.B. Saunders Co., Philadelphia, Pa., pp. 1128–1129.
- Rodrigues, et al., 1993, *Journal of Immunology*. 151, pp. 6954–6961.
- Salant, et al., 1980, *Journal of Clinical Investigations*. 66, pp. 1339–1350.

56

- Schrijver, et al., 1988, *Laboratory Investigation*. 59, pp. 484–491.
- Schrijver, et al., 1990, *Kidney International*. 38, pp. 86–95.
- Unanue and Dixon, 1964, *Journal of Experimental Medicine*. 119, pp. 965–982.
- Winter and Milstein, 1991, *Nature*. 349, 293–299.
- Wurzner, et al., 1991, *Complement Inflammation*. 8, 328–340.

TABLE 1

Prevention/Reduction of Proteinuria by Treatment With Anti-C5 Antibodies		
	Before Treatment Urine Protein (mg/dL)	After Treatment Urine Protein (mg/dL)
<hr/>		
	PBS Control	
mouse A	none	100
mouse B	none	500
mouse C	none	500
mouse D*	trace	trace
mouse E	100	100
	Anti-C5 Treated	
mouse 1	none	none
mouse 2	none	30
mouse 3	30	trace
mouse 4	30	30
mouse 5	30	30
mouse 6	100	30

*Mouse D had more than 500 mg/dL urine glucose after treatment

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(iii) NUMBER OF SEQUENCES: 26

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21 amino acids
- (B) TYPE: Amino Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear

(ii) MOLECULE TYPE:

- (A) DESCRIPTION: KSSKC peptide

(iii) HYPOTHETICAL: No

(iv) ANTI-SENSE: No

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

Val Ile Asp His Gln Gly Thr Lys Ser Ser
5 10

Lys Cys Val Arg Gln Lys Val Glu Gly Ser Ser
15 20

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1676 Amino Acids
- (B) TYPE: Amino Acid

-continued

(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE:
(A) DESCRIPTION: Pro-C5 Polypeptide

(iii) HYPOTHETICAL: No

(iv) ANTI-SENSE: No

(vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

(x) PUBLICATION INFORMATION:
(A) AUTHORS: Haviland, D.L.
Haviland, J.C.
Fleischer, D.T.
Hunt, A.
Wetsel, R.A.
(B) TITLE: Complete cDNA Sequence of Human
Complement Pro-C5
(C) JOURNAL: Journal of Immunology
(D) VOLUME: 146
(F) PAGES: 362-368
(G) DATE: 1991

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met Gly Leu Leu Gly Ile Leu Cys Phe Leu
-15 -10

Ile Phe Leu Gly Lys Thr Trp Gly Gln Glu Gln Thr Tyr Val
-5 -1 5

Ile Ser Ala Pro Lys Ile Phe Arg Val Gly Ala Ser Glu Asn
10 15 20

Ile Val Ile Gln Val Tyr Gly Tyr Thr Glu Ala Phe Asp Ala
25 30

Thr Ile Ser Ile Lys Ser Tyr Pro Asp Lys Lys Phe Ser Tyr
35 40 45

Ser Ser Gly His Val His Leu Ser Ser Glu Asn Lys Phe Gln
50 55 60

Asn Ser Ala Ile Leu Thr Ile Gln Pro Lys Gln Leu Pro Gly
65 70 75

Gly Gln Asn Pro Val Ser Tyr Val Tyr Leu Glu Val Val Ser
80 85 90

Lys His Phe Ser Lys Ser Lys Arg Met Pro Ile Thr Tyr Asp
95 100

Asn Gly Phe Leu Phe Ile His Thr Asp Lys Pro Val Tyr Thr
105 110 115

Pro Asp Gln Ser Val Lys Val Arg Val Tyr Ser Leu Asn Asp
120 125 130

Asp Leu Lys Pro Ala Lys Arg Glu Thr Val Leu Thr Phe Ile
135 140 145

Asp Pro Glu Gly Ser Glu Val Asp Met Val Glu Glu Ile Asp
150 155 160

His Ile Gly Ile Ile Ser Phe Pro Asp Phe Lys Ile Pro Ser
165 170

Asn Pro Arg Tyr Gly Met Trp Thr Ile Lys Ala Lys Tyr Lys
175 180 185

Glu Asp Phe Ser Thr Thr Gly Thr Ala Tyr Phe Glu Val Lys
190 195 200

Glu Tyr Val Leu Pro His Phe Ser Val Ser Ile Glu Pro Glu
205 210 215

-continued

Tyr Asn Phe Ile Gly	Tyr Lys Asn Phe Lys	Asn Phe Glu Ile
220	225	230
Thr Ile Lys Ala Arg	Tyr Phe Tyr Asn Lys	Val Val Thr Glu
235	240	
Ala Asp Val Tyr Ile	Thr Phe Gly Ile Arg	Glu Asp Leu Lys
245	250	255
Asp Asp Gln Lys Glu	Met Met Gln Thr Ala	Met Gln Asn Thr
260	265	270
Met Leu Ile Asn Gly	Ile Ala Gln Val Thr	Phe Asp Ser Glu
275	280	285
Thr Ala Val Lys Glu	Leu Ser Tyr Tyr Ser	Leu Glu Asp Leu
290	295	300
Asn Asn Lys Tyr Leu	Tyr Ile Ala Val Thr	Val Ile Glu Ser
305	310	
Thr Gly Gly Phe Ser	Glu Glu Ala Glu Ile	Pro Gly Ile Lys
315	320	325
Tyr Val Leu Ser Pro	Tyr Lys Leu Asn Leu	Val Ala Thr Pro
330	335	340
Leu Phe Leu Lys Pro	Gly Ile Pro Tyr Pro	Ile Lys Val Gln
345	350	355
Val Lys Asp Ser Leu	Asp Gln Leu Val Gly	Gly Val Pro Val
360	365	370
Ile Leu Asn Ala Gln	Thr Ile Asp Val Asn	Gln Glu Thr Ser
375	380	
Asp Leu Asp Pro Ser	Lys Ser Val Thr Arg	Val Asp Asp Gly
385	390	395
Val Ala Ser Phe Val	Leu Asn Leu Pro Ser	Gly Val Thr Val
400	405	410
Leu Glu Phe Asn Val	Lys Thr Asp Ala Pro	Asp Leu Pro Glu
415	420	425
Glu Asn Gln Ala Arg	Glu Gly Tyr Arg Ala	Ile Ala Tyr Ser
430	435	440
Ser Leu Ser Gln Ser	Tyr Leu Tyr Ile Asp	Trp Thr Asp Asn
445	450	
His Lys Ala Leu Leu	Val Gly Glu His Leu	Asn Ile Ile Val
455	460	465
Thr Pro Lys Ser Pro	Tyr Ile Asp Lys Ile	Thr His Tyr Asn
470	475	480
Tyr Leu Ile Leu Ser	Lys Gly Lys Ile Ile	His Phe Gly Thr
485	490	495
Arg Glu Lys Phe Ser	Asp Ala Ser Tyr Gln	Ser Ile Asn Ile
500	505	510
Pro Val Thr Gln Asn	Met Val Pro Ser Ser	Arg Leu Leu Val
515	520	
Tyr Tyr Ile Val Thr	Gly Glu Gln Thr Ala	Glu Leu Val Ser
525	530	535
Asp Ser Val Trp Leu	Asn Ile Glu Glu Lys	Cys Gly Asn Gln
540	545	550
Leu Gln Val His Leu	Ser Pro Asp Ala Asp	Ala Tyr Ser Pro
555	560	565
Gly Gln Thr Val Ser	Leu Asn Met Ala Thr	Gly Met Asp Ser
570	575	580

-continued

Trp Val Ala Leu Ala	Ala Val Asp Ser Ala Val Tyr Gly Val
585	590
Gln Arg Gly Ala Lys	Lys Pro Leu Glu Arg Val Phe Gln Phe
595	600 605
Leu Glu Lys Ser Asp	Leu Gly Cys Gly Ala Gly Gly Gly Leu
610	615 620
Asn Asn Ala Asn Val	Phe His Leu Ala Gly Leu Thr Phe Leu
625	630 635
Thr Asn Ala Asn Ala	Asp Asp Ser Gln Glu Asn Asp Glu Pro
640	645 650
Cys Lys Glu Ile Leu	Arg Pro Arg Arg Thr Leu Gln Lys Lys
655	660
Ile Glu Glu Ile Ala	Ala Lys Tyr Lys His Ser Val Val Lys
665	670 675
Lys Cys Cys Tyr Asp	Gly Ala Cys Val Asn Asn Asp Glu Thr
680	685 690
Cys Glu Gln Arg Ala	Ala Arg Ile Ser Leu Gly Pro Arg Cys
695	700 705
Ile Lys Ala Phe Thr	Glu Cys Cys Val Val Ala Ser Gln Leu
710	715 720
Arg Ala Asn Ile Ser	His Lys Asp Met Gln Leu Gly Arg Leu
725	730
His Met Lys Thr Leu	Leu Pro Val Ser Lys Pro Glu Ile Arg
735	740 745
Ser Tyr Phe Pro Glu	Ser Trp Leu Trp Glu Val His Leu Val
750	755 760
Pro Arg Arg Lys Gln	Leu Gln Phe Ala Leu Pro Asp Ser Leu
765	770 775
Thr Thr Trp Glu Ile	Gln Gly Ile Gly Ile Ser Asn Thr Gly
780	785 790
Ile Cys Val Ala Asp	Thr Val Lys Ala Lys Val Phe Lys Asp
795	800
Val Phe Leu Glu Met	Asn Ile Pro Tyr Ser Val Val Arg Gly
805	810 815
Glu Gln Ile Gln Leu	Lys Gly Thr Val Tyr Asn Tyr Arg Thr
820	825 830
Ser Gly Met Gln Phe	Cys Val Lys Met Ser Ala Val Glu Gly
835	840 845
Ile Cys Thr Ser Glu	Ser Pro Val Ile Asp His Gln Gly Thr
850	855 860
Lys Ser Ser Lys Cys	Val Arg Gln Lys Val Glu Gly Ser Ser
865	870
Ser His Leu Val Thr	Phe Thr Val Leu Pro Leu Glu Ile Gly
875	880 885
Leu His Asn Ile Asn	Phe Ser Leu Glu Thr Trp Phe Gly Lys
890	895 900
Glu Ile Leu Val Lys	Thr Leu Arg Val Val Pro Glu Gly Val
905	910 915
Lys Arg Glu Ser Tyr	Ser Gly Val Thr Leu Asp Pro Arg Gly
920	925 930
Ile Tyr Gly Thr Ile	Ser Arg Arg Lys Glu Phe Pro Tyr Arg
935	940

-continued

Ile Pro Leu Asp Leu Val Pro Lys Thr Glu Ile Lys Arg Ile		
945	950	955
Leu Ser Val Lys Gly Leu Leu Val Gly Glu Ile Leu Ser Ala		
960	965	970
Val Leu Ser Gln Glu Gly Ile Asn Ile Leu Thr His Leu Pro		
975	980	985
Lys Gly Ser Ala Glu Ala Glu Leu Met Ser Val Val Pro Val		
990	995	1000
Phe Tyr Val Phe His Tyr Leu Glu Thr Gly Asn His Trp Asn		
1005	1010	
Ile Phe His Ser Asp Pro Leu Ile Glu Lys Gln Lys Leu Lys		
1015	1020	1025
Lys Lys Leu Lys Glu Gly Met Leu Ser Ile Met Ser Tyr Arg		
1030	1035	1040
Asn Ala Asp Tyr Ser Tyr Ser Val Trp Lys Gly Gly Ser Ala		
1045	1050	1055
Ser Thr Trp Leu Thr Ala Phe Ala Leu Arg Val Leu Gly Gln		
1060	1065	1070
Val Asn Lys Tyr Val Glu Gln Asn Gln Asn Ser Ile Cys Asn		
1075	1080	
Ser Leu Leu Trp Leu Val Glu Asn Tyr Gln Leu Asp Asn Gly		
1085	1090	1095
Ser Phe Lys Glu Asn Ser Gln Tyr Gln Pro Ile Lys Leu Gln		
1100	1105	1110
Gly Thr Leu Pro Val Glu Ala Arg Glu Asn Ser Leu Tyr Leu		
1115	1120	1125
Thr Ala Phe Thr Val Ile Gly Ile Arg Lys Ala Phe Asp Ile		
1130	1135	1140
Cys Pro Leu Val Lys Ile Asp Thr Ala Leu Ile Lys Ala Asp		
1145	1150	
Asn Phe Leu Leu Glu Asn Thr Leu Pro Ala Gln Ser Thr Phe		
1155	1160	1165
Thr Leu Ala Ile Ser Ala Tyr Ala Leu Ser Leu Gly Asp Lys		
1170	1175	1180
Thr His Pro Gln Phe Arg Ser Ile Val Ser Ala Leu Lys Arg		
1185	1190	1195
Glu Ala Leu Val Lys Gly Asn Pro Pro Ile Tyr Arg Phe Trp		
1200	1205	1210
Lys Asp Asn Leu Gln His Lys Asp Ser Ser Val Pro Asn Thr		
1215	1220	
Gly Thr Ala Arg Met Val Glu Thr Thr Ala Tyr Ala Leu Leu		
1225	1230	1235
Thr Ser Leu Asn Leu Lys Asp Ile Asn Tyr Val Asn Pro Val		
1240	1245	1250
Ile Lys Trp Leu Ser Glu Glu Gln Arg Tyr Gly Gly Gly Phe		
1255	1260	1265
Tyr Ser Thr Gln Asp Thr Ile Asn Ala Ile Glu Gly Leu Thr		
1270	1275	1280
Glu Tyr Ser Leu Leu Val Lys Gln Leu Arg Leu Ser Met Asp		
1285	1290	
Ile Asp Val Ser Tyr Lys His Lys Gly Ala Leu His Asn Tyr		
1295	1300	1305

-continued

Lys Met Thr Asp Lys	Asn Phe Leu Gly Arg	Pro Val Glu Val
1310	1315	1320
Leu Leu Asn Asp Asp	Leu Ile Val Ser Thr	Gly Phe Gly Ser
1325	1330	1335
Gly Leu Ala Thr Val	His Val Thr Thr Val	Val His Lys Thr
1340	1345	1350
Ser Thr Ser Glu Glu	Val Cys Ser Phe Tyr	Leu Lys Ile Asp
1355	1360	
Thr Gln Asp Ile Glu	Ala Ser His Tyr Arg	Gly Tyr Gly Asn
1365	1370	1375
Ser Asp Tyr Lys Arg	Ile Val Ala Cys Ala	Ser Tyr Lys Pro
1380	1385	1390
Ser Arg Glu Glu Ser	Ser Ser Gly Ser Ser	His Ala Val Met
1395	1400	1405
Asp Ile Ser Leu Pro	Thr Gly Ile Ser Ala	Asn Glu Glu Asp
1410	1415	1420
Leu Lys Ala Leu Val	Glu Gly Val Asp Gln	Leu Phe Thr Asp
1425	1430	
Tyr Gln Ile Lys Asp	Gly His Val Ile Leu	Gln Leu Asn Ser
1435	1440	1445
Ile Pro Ser Ser Asp	Phe Leu Cys Val Arg	Phe Arg Ile Phe
1450	1455	1460
Glu Leu Phe Glu Val	Gly Phe Leu Ser Pro	Ala Thr Phe Thr
1465	1470	1475
Val Tyr Glu Tyr His	Arg Pro Asp Lys Gln	Cys Thr Met Phe
1480	1485	1490
Tyr Ser Thr Ser Asn	Ile Lys Ile Gln Lys	Val Cys Glu Gly
1495	1500	
Ala Ala Cys Lys Cys	Val Glu Ala Asp Cys	Gly Gln Met Gln
1505	1510	1515
Glu Glu Leu Asp Leu	Thr Ile Ser Ala Glu	Thr Arg Lys Gln
1520	1525	1530
Thr Ala Cys Lys Pro	Glu Ile Ala Tyr Ala	Tyr Lys Val Ser
1535	1540	1545
Ile Thr Ser Ile Thr	Val Glu Asn Val Phe	Val Lys Tyr Lys
1550	1555	1560
Ala Thr Leu Leu Asp	Ile Tyr Lys Thr Gly	Glu Ala Val Ala
1565	1570	
Glu Lys Asp Ser Glu	Ile Thr Phe Ile Lys	Lys Val Thr Cys
1575	1580	1585
Thr Asn Ala Glu Leu	Val Lys Gly Arg Gln	Tyr Leu Ile Met
1590	1595	1600
Gly Lys Glu Ala Leu	Gln Ile Lys Tyr Asn	Phe Ser Phe Arg
1605	1610	1615
Tyr Ile Tyr Pro Leu	Asp Ser Leu Thr Trp	Ile Glu Tyr Trp
1620	1625	1630
Pro Arg Asp Thr Thr	Cys Ser Ser Cys Gln	Ala Phe Leu Ala
1635	1640	
Asn Leu Asp Glu Phe	Ala Glu Asp Ile Phe	Leu Asn Gly Cys
1645	1650	1655

-continued

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 4059 base pairs
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Double
- (D) TOPOLOGY: Circular

(ii) MOLECULE TYPE: Other nucleic acid

- (A) DESCRIPTION: Apex-1 Eukaryotic Expression Vector

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

ACGCGTTGAC ATTGATTATT GACTAGTTAT TAATAGTAAT CAATTACGGG	50
GTCATTAGTT CATAGCCCAT ATATGGAGTT CCGCGTTACA TAACTTACGG	100
TAAATGGCCC CGCCTGGCTG ACCGCCCAAC GACCCCGCC CATTGACGTC	150
AATAATGACG TATGTTCCCA TAGTAACGCC AATAGGGACT TTCCATTGAC	200
GTCAATGGGT GGAATATTTA CGGTAACTG CCCACTTGGC AGTACATCAA	250
GTGTATCATA TGCCAAGTAC GCCCCTATT GACGTCAATG ACGGTAAATG	300
GCCCGCCTGG CATTATGCCC AGTACATGAC CTTATGGGAC TTTCTACTT	350
GGCAGTACAT CTACGTATTA GTCATCGCTA TTACCATGGT GATGCGGTTT	400
TGGCAGTACA TCAATGGGCG TGGATAGCGG TTTGACTCAC GGGGATTTC	450
AAGTCTCCAC CCCATTGACG TCAATGGGAG TTTGTTTGG CACCAAAATC	500
AACGGGACTT TCCAAAATGT CGTAACAACT CCGCCCCATT GACGCAAATG	550
GGCGGTAGGC GTGTACGGTG GGAGGTCTAT ATAAGCAGAG CTCGTTTAGT	600
GAACCGTCAG AATTCTGTTG GGCTCGCGGT TGATTACAAA CTCTTCGCGG	650
TCTTTCCAGT ACTCTTGGAT CGGAAACCCG TCGGCCTCCG AACGGTACTC	700
CGCCACCGAG GGACCTGAGC GAGTCCGCAT CGACCGGATC GAAAACCTC	750
TCGACTGTTG GGGTGAGTAC TCCCTCTCAA AAGCGGCAT GACTTCTGCG	800
CTAAGATTGT CAGTTTCCAA AAACGAGGAG GATTTGATAT TCACCTGGCC	850
CGCGGTGATG CCTTTGAGGG TGGCCGCGTC CATCTGGTCA GAAAAGACAA	900
TCTTTTGTG GTCAAGCTTG AGGTGTGGCA GGCTTGAGAT CTGGCCATAC	950
ACTTGAGTGA CAATGACATC CACTTTGCCT TTCTCTCCAC AGGTGTCCAC	1000
TCCCAGGTCC AACTGCAGGT CGACCGGCTT GGTACCGAGC TCGGATCCAC	1050
TAGTAACGGC CGCCAGTGTG CTGGAATTCT GCAGATATCC ATCACACTGG	1100
CGGCCGCTCG AGCATGCATC TAGAACTTGT TTATTGCAGC TTATAATGGT	1150
TACAAATAAA GCAATAGCAT CACAAATTTC ACAAATAAAG CATTTTTTTC	1200
ACTGCATTCT AGTTGTGGTT TGTCCAAACT CATCAATGTA TCTTATCATG	1250
TCTGGATCGA TCCCGCCATG GTATCAACGC CATATTTCTA TTTACAGTAG	1300
GGACCTCTTC GTTGTGTAGG TACCGCTGTA TTCCTAGGGA AATAGTAGAG	1350
GCACCTTGAA CTGTCTGCAT CAGCCATATA GCCCCGCTG TTCGACTTAC	1400
AAACACAGGC ACAGTACTGA CAAACCCATA CACCTCCTCT GAAATACCCA	1450
TAGTTGCTAG GGCTGTCTCC GAACTCATTA CACCCTCCAA AGTCAGAGCT	1500
GTAATTTTCG CATCAAGGGC AGCGAGGGCT TCTCCAGATA AAATAGCTTC	1550
TGCCGAGAGT CCCGTAAGGG TAGACACTTC AGCTAATCCC TCGATGAGGT	1600
CTACTAGAAT AGTCAGTGC GCTCCCATTT TGAAAATCA CTTACTTGAT	1650

-continued

CAGCTTCAGA AGATGGCGGA GGGCCTCCAA CACAGTAATT TTCCTCCCGA	1700
CTCTTAAAT AGAAAAATGTC AAGTCAGTTA AGCAGGAAGT GGAATACTG	1750
ACGCAGCTGG CCGTGCGACA TCCTCTTTTA ATTAGTTGCT AGGCAACGCC	1800
CTCCAGAGGG CGTGTGGTTT TGCAAGAGGA AGCAAAAGCC TCTCCACCCA	1850
GGCCTAGAAT GTTTCACCC AATCATTACT ATGACAACAG CTGTTTTTTT	1900
TAGTATTAAG CAGAGGCCGG GGACCCCTGG GCCCGCTTAC TCTGGAGAAA	1950
AAGAAGAGAG GCATTGTAGA GGCTTCAGA GGCAACTTGT CAAAACAGGA	2000
CTGCTTCTAT TTCTGTCACA CTGTCTGGCC CTGTACACAAG GTCCAGCACC	2050
TCCATACCCC CTTTAATAAG CAGTTTGGGA ACGGGTGC GG GTCTTACTCC	2100
GCCCATCCCG CCCCTAACTC CGCCCACTTC CGCCCATCT CCGCCCATG	2150
GCTGACTAAT TTTTTTTAT TATGCAGAG CCGAGGCCGC CTCGGCTCT	2200
GAGCTATTCC AGAAGTAGTG AGGAGGCTTT TTTGGAGGCC TAGGCTTTTG	2250
CAAAAAGGAG CTCCCAGCAA AAGGCCAGGA ACCGTAAAAA GGCCGCGTTG	2300
CTGGCGTTTT TCCATAGGCT CGCCCCCCT GACGAGCATC AAAAAATCG	2350
ACGCTCAAGT CAGAGGTGGC GAAACCCGAC AGGACTATAA AGATACCAGG	2400
CGTTTCCCC TGGAAGCTCC CTCGTGCGCT CTCCTGTTC GACCTGCCG	2450
CTTACCGGAT ACCTGTCCGC CTTTCTCCCT TCGGGAAGCG TGGCGCTTC	2500
TCAATGCTCA CGCTGTAGGT ATCTCAGTTC GGTGTAGGTC GTTCGCTCCA	2550
AGCTGGGCTG TGTGCACGAA CCCCCCGTTC AGCCCGACCG CTGCGCCTTA	2600
TCCGGTAACT ATCGTCTTGA GTCCAACCCG GTAAGACACG ACTTATCGCC	2650
ACTGGCAGCA GCCACTGGTA ACAGATTAG CAGAGCGAGG TATGTAGGCG	2700
GTGCTACAGA GTTCTTGAAG TGGTGGCCTA ACTACGGCTA CACTAGAAGG	2750
ACAGTATTG GTATCTGCGC TCTGTGAAG CCAGTTACCT TCGGAAAAAG	2800
AGTTGGTAGC TCTTGATCCG GCAACAAAC CACCGCTGGT AGCGGTGGTT	2850
TTTTTGTTTG CAAGCAGCAG ATTACGCGCA GAAAAAAG ATCTCAAGAA	2900
GATCCTTTGA TCTTTCTAC GGGGTCTGAC GCTCAGTGA ACGAAACTC	2950
ACGTTAAGG ATTTTGGTCA TGAGATTATC AAAAAGGATC TTCACCTAGA	3000
TCCTTTTAAA TTAATAATGA AGTTTAAAT CAATCTAAAG TATATATGAG	3050
TAAACTTGGT CTGACAGTTA CCAATGCTTA ATCAGTGAGG CACCTATCTC	3100
AGCGATCTGT CTATTCGTT CATCCATAGT TGCCTGACTC CCCGTCGTGT	3150
AGATAACTAC GATACGGGAG GGCTTACCAT CTGGCCCCAG TGCTGCAATG	3200
ATACCGCGAG ACCCACGCTC ACCGCTCCA GATTATCAG CAATAAACCA	3250
GCCAGCCGGA AGGGCCGAGC GCAGAAGTGG TCCTGCAACT TTATCCGCCT	3300
CCATCCAGTC TATTAATTGT TGCCGGAAG CTAGAGTAAG TAGTTCGCCA	3350
GTTAATAGTT TGCACAACGT TGTGCCATT GCTACAGGCA TCGTGGTGT	3400
ACGCTCGTCG TTTGGTATGG CTTCAATCAG CTCCGGTTCC CAACGATCAA	3450
GGCGAGTTAC ATGATCCCC ATGTTGTGCA AAAAAGCGGT TAGCTCCTTC	3500
GGTCTCCGA TCGTTGTCAG AAGTAAGTTG GCCGAGTGT TATCACTCAT	3550
GGTTATGGCA GCACTGCATA ATTCTCTTAC TGTCATGCCA TCCGTAAGAT	3600
GCTTTTCTGT GACTGGTGAG TACTCAACCA AGTCATTCTG AGAATAGTGT	3650

-continued

ATGCGGCGAC CGAGTTGCTC TTGCCCCGCG TCAATACGGG ATAATACCGC	3700
GCCACATAGC AGAACTTTAA AAGTGCTCAT CATTGGAAAA CGTTCCTCGG	3750
GGCGAAAACT CTCAAGGATC TTACCGCTGT TGAGATCCAG TTCGATGTAA	3800
CCCACTCGTG CACCCAACTG ATCTTCAGCA TCTTTTACTT TCACCAGCGT	3850
TTCTGGGTGA GAAAAACAG GAAGGCAAAA TGCCGCAAAA AAGGGAATAA	3900
GGGCGACACG GAAATGTTGA ATACTCATAC TCTTCCTTTT TCAATATTAT	3950
TGAAGCATTT ATCAGGGTTA TTGTCTCATG AGCGGATACA TATTTGAATG	4000
TATTTAGAAA AATAAACAAA TAGGGGTCC GCGCACATTT CCCCAGAAAAG	4050
TGCCACCTG	4059

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 8540 base pairs
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Double
- (D) TOPOLOGY: Circular

(ii) MOLECULE TYPE: Other nucleic acid

- (A) DESCRIPTION: Apex-3P Eukaryotic Expression Vector

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

GTGACCAATA CAAAAACAAA GCGCCCTCG TACCAGCGAA GAAGGGGCGAG	50
AGATGCCGTA GTCAGGTTA GTTCGTCCGG CGGCGGGGA TCTGTATGGT	100
GCACTCTCAG TACAATCTGC TCTGATGCCG CATAGTTAAG CCAGTATCTG	150
CTCCCTGCTT GTGTGTGGA GGTGCTGAG TAGTGCGCGA GCAAAATTTA	200
AGCTACAACA AGGCAAGGCT TGACCGACAA TTGCATGAAG AATCTGCTTA	250
GGGTAGGCG TTTTGCCTG CTTCGCGATG TACGGGCCAG ATATACGCGT	300
TGACATTGAT TATTGACTAG TTATTAATAG TAATCAATTA CGGGGTCATT	350
AGTTCATAGC CCATATATGG AGTTCCGCGT TACATAACTT ACGGTAATG	400
GCCCCTGCG CTGACCGCCC AACGACCCC GCCCATTGAC GTCAATAATG	450
ACGTATGTTT CCATAGTAAC GCCAATAGGG ACTTTCATT GACGTCAATG	500
GGTGGACTAT TTACGGTAAA CTGCCCACTT GGCAGTACAT CAAGTGTATC	550
ATATGCCAAG TACGCCCCCT ATTGACGTCA ATGACGGTAA ATGGCCCGCC	600
TGGCATTATG CCCAGTACAT GACCTTATGG GACTTTCCTA CTTGGCAGTA	650
CATCTACGTA TTAGTCATCG CTATTACCAT GGTGATGCGG TTTTGGCAGT	700
ACATCAATGG GCGTGGATAG CGGTTTGACT CACGGGGATT TCCAAGTCTC	750
CACCCCATTG ACGTCAATGG GAGTTTGTTT TGGCACCAAA ATCAACGGGA	800
CTTTCCAAAA TGTCGTAACA ACTCCGCCCC ATTGACGCAA ATGGGCGGTA	850
GGCGTGTACG GTGGGAGGTC TATATAAGCA GAGCTCGTTT AGTGAACCGT	900
CAGAAATCTG TTGGGCTCGC GGTGATTAC AAACCTCTCG CGGTCTTTCC	950
AGTACTCTTG GATCGGAAAC CCGTCGGCCT CCGAACGGTA CTCCGCCACC	1000
GAGGGACCTG AGCGAGTCCG CATCGACCGG ATCGGAAAAC CTCTCGACTG	1050
TTGGGGTGAG TACTCCCTCT CAAAAGCGGG CATGACTTCT GCGCTAAGAT	1100
TGTCAGTTTC CAAAAACGAG GAGGATTGTA TATTCACCTG GCCCGCGGTG	1150
ATGCCTTTGA GGGTGGCCGC GTCCATCTGG TCAGAAAAGA CAATCTTTTT	1200

-continued

GTTGTCAAGC	TTGAGGTGTG	GCAGGCTTGA	GATCTGGCCA	TACACTTGAG	1250
TGACAATGAC	ATCCACTTTG	CCTTTCTCTC	CACAGGTGTC	CACTCCCAGG	1300
TCCAAC TGCA	GGTCGACCGG	CTTGGTACCG	AGCTCGGATC	CTCTAGAGTC	1350
GACCTGCAGG	CATGCAAGCT	TGGCACTGGC	CGTCGTTTTA	CAACGTCGTG	1400
ACTGGGAAAA	CCCTGGCGTT	ACCCAAC TTA	ATCGCCTTGC	AGCACATCCC	1450
CCTTTCGCCA	GCTGGCGTAA	TAGCGAAGAG	GCCCGCACCG	ATCCAGACAT	1500
GATAAGATAC	ATTGATGAGT	TTGGACAAAC	CACAACTAGA	ATGCAGTGAA	1550
AAAAATGCTT	TATTTGTGAA	ATTTGTGATG	CTATTGCTTT	ATTGTAAACC	1600
ATTATAAGCT	GCAATAAACA	AGTTAACAAC	AACAATTGCA	TTCATTTTAT	1650
GTTTCAGGTT	CAGGGGGAGG	TGTGGGAGGT	TTTTTAAAGC	AAGTAAAACC	1700
TCTACAAATG	TGGTATGGCT	GATTATGATC	CCCAGGAAGC	TCCTCTGTGT	1750
CCTCATAAAC	CCTAACCTCC	TCTACTTGAG	AGGACATTCC	AATCATAGGC	1800
TGCCCCATCA	CCCTCTGTGT	CCTCCTGTTA	ATTAGGTCAC	TTAACAAAAA	1850
GGAAATTGGG	TAGGGGTTTT	TCACAGACCG	CTTCTAAGG	GTAATTTTAA	1900
AATATCTGGG	AAGTCCCTTC	CACTGCTGTG	TTCCAGAAGT	GTTGGTAAAC	1950
AGCCCACAAA	TGTCACACGC	AGAAACATAC	AAGCTGTCAG	CTTGCACAA	2000
GGGCCCCAACA	CCCTGCTCAT	CAAGAAGCAC	TGTGGTTGCT	GTGTTAGTAA	2050
TGTGCAAAAC	AGGAGGCACA	TTTTCCCCAC	CTGTGTAGGT	TCCAAAATAT	2100
CTAGTGTTTT	CATTTT TACT	TGGATCAGGA	ACCCAGCACT	CCACTGGATA	2150
AGCATTATCC	TTATCCAAAA	CAGCCTTGTG	GTCAGTGTTT	ATCTGCTGAC	2200
TGTCAACTGT	AGCATTTTTT	GGGGTTACAG	TTTGAGCAGG	ATATTGGTC	2250
CTGTAGTTTG	CTAACACACC	CTGCAGCTCC	AAAGGTTCCC	CACCAACAGC	2300
AAAAAAATGA	AAATTTGACC	CTTGAATGGG	TTTTCCAGCA	CCATTTTCAT	2350
GAGTTTTTTG	TGTCCCTGAA	TGCAAGTTTA	ACATAGCAGT	TACCCCAATA	2400
ACCTCAGTTT	TAACAGTAAC	AGCTTCCCAC	ATCAAAATAT	TTCCACAGGT	2450
TAAGTCCTCA	TTTG TAGAAT	TCGCCAGCAC	AGTGGTCGAC	CCTGTGGATG	2500
TGTGTCAC TT	AGGGTGTGGA	AAGTCCCCAG	GCTCCCCAGC	AGGCAGAA GT	2550
ATGCAAAGCA	TGCATCTCAA	TTAGTCAGCA	ACCAGGTGTG	GAAAGTCCCC	2600
AGGCTCCCCA	GCAGGCAGAA	GTATGCAAAG	CATGCATCTC	AATTAGTCAG	2650
CAACCATAGT	CCCGCCCCTA	ACTCCGCCCA	TCCCGCCCCCT	AACTCCGCCC	2700
AGTTCCGCCC	ATTCTCCGCC	CCATGGCTGA	CTAATTTTTT	TTATTTATGC	2750
AGAGGCCGAG	GCCGCCTCGG	CCTCTGAGCT	ATTCCAGAAG	TAGTGAGGAG	2800
GCTTTTTTGG	AGGCCTAGGC	TTTTGCAAAA	GCTTACCATG	ACCGAGTACA	2850
AGCCCACGGT	GCGCCTCGCC	ACCCGCGACG	ACGTCCCCCG	GGCCGTACGC	2900
ACCTCGCCG	CCGCGTTCGC	CGACTACCCC	GCCACGCGCC	ACACCGTCGA	2950
CCCGGACCGC	CACATCGAGC	GGGTCAACGA	GCTGCAAGAA	CTCTTCCTCA	3000
CGCGCGTCGG	GCTCGACATC	GGCAAGGTGT	GGGTCGCGGA	CGACGGCGCC	3050
GCGGTGGCGG	TCTGGACCAC	GCCGAGAGC	GTCGAAGCGG	GGGCGGTGTT	3100
CGCCGAGATC	GGCCCGCGCA	TGGCCGAGTT	GAGCGGTTCC	CGGCTGGCCG	3150
CGCAGCAACA	GATGGAAGGC	CTCCTGGCGC	CGCACCGGCC	CAAGGAGCCC	3200

-continued

GCGTGGTTCC TGGCCACCGT CGGCGTCTCG CCCGACCACC AGGGCAAGGG	3250
TCTGGGCAGC GCCGTCTGTC TCCCGGAGT GGAGGCGGCC GAGCGCGCCG	3300
GGGTGCCCCG CTTCTGGAG ACCTCCGCGC CCCGCAACCT CCCCTTCTAC	3350
GAGCGGCTCG GCTTCACCGT CACCGCCGAC GTCGAGTGCC CGAAGGACCG	3400
CGCGACCTGG TGCATGACCC GCAAGCCCGG TGCCTGACGC CCGCCCCACG	3450
ACCCGCAGCG CCCGACCGAA AGGAGCGCAC GACCCCATGC ATCGATAAAA	3500
TAAAAGATTT TATTTAGTCT CCAGAAAAAG GGGGAATGA AAGACCCAC	3550
CTGTAGGTTT GGCAAGCTAG AACTTGTTTA TTGCAGCTTA TAATGGTTAC	3600
AAATAAGCA ATAGCATCAC AAATTTACA AATAAAGCAT TTTTTCCTACT	3650
GCATTCTAGT TGTGGTTTGT CCAAACTCAT CAATGTATCT TATCATGTCT	3700
GGATCGATCC CGCCATGGTA TCAACGCCAT ATTTCTATTT ACAGTAGGGA	3750
CCTCTTCGTT GTGTAGGTAC CCCGGGTTTCG AAATCGAATT CGCCAATGAC	3800
AAGACGCTGG GCGGGGTTTG TGTATCATA GAACTAAAGA CATGCAAATA	3850
TATTTCTTCC GGGGACACCG CCAGCAAACG CGAGCAACGG GCCACGGGGA	3900
TGAAGCAGCC CGGCGGCACC TCGCTAACGG ATTCACCACT CCAAGAATTG	3950
GAGCCAATCA ATTCTTGCGG AGAACTGTGA ATGCGCAAAC CAACCCTTGG	4000
CAGAACATAT CCATCGCGTC CGCCATCTCC AGCAGCCGCA CGCGGCGCAT	4050
CTCGGGGCCG ACGCGCTGGG CTACGTCTTG CTGGCGTTTCG CGACGCGAGG	4100
CTGGATGGCC TTCCCCATTA TGATTCTTCT CGCTTCCGGC GGCATCGGGA	4150
TGCCCCGCTT GCAGGCCATG CTGTCCAGGC AGGTAGATGA CGACCATCAG	4200
GGACAGCTTC AAGGATCGCT CGCGGCTCTT ACCAGCGCCA GCAAAAGGCC	4250
AGGAACCGTA AAAAGGCCGC GTTGCTGGCG TTTTTCATA GGCTCCGCC	4300
CCCTGACGAG CATCACAAAA ATCGACGCTC AAGTCAGAGG TGGCGAAACC	4350
CGACAGGACT ATAAAGATAC CAGGCGTTTC CCCCTGGAAG CTCCCTCGTG	4400
CGCTCTCCTG TTCCGACCCCT GCCGCTTACC GGATACCTGT CCGCCTTCT	4450
CCCTTCGGGA AGCGTGGCGC TTTCTCATAG CTCACGCTGT AGGTATCTCA	4500
GTTTCGGTGA GGTCTTTCGC TCCAAGCTGG GCTGTGTGCA CGAACCCCC	4550
GTTCAGCCCG ACCGCTGCGC CTTATCCGGT AACTATCGTC TTGAGTCCAA	4600
CCCGGTAAGA CACGACTTAT CGCCTGCGC AGCAGCCACT GGTAACAGGA	4650
TTAGCAGAGC GAGGTATGTA GCGGTGCTA CAGAGTTCTT GAAGTGGTGG	4700
CCTAACTACG GCTACACTAG AAGGACAGTA TTTGGTATCT GCGCTCTGCT	4750
GAAGCCAGTT ACCTTCGGAA AAAGAGTTGG TAGCTCTTGA TCCGGCAAAC	4800
AAACCACCGC TGGTAGCGGT GGTTTTTTTT TTTGCAAGCA GCAGATTACG	4850
CGCAGAAAAA AAGGATCTCA AGAAGATCCT TTGATCTTTT CTACGGGGTC	4900
TGACGCTCAG TGAACGAAA ACTCACGTTA AGGGATTTTG GTCATGAGAT	4950
TATCAAAAAG GATCTTCACC TAGATCCTTT TAAATTAAAA ATGAAGTTTT	5000
AAATCAATCT AAAGTATATA TGAGTAAACT TGGTCTGACA GTTACCAATG	5050
CTTAATCAGT GAGGCACCTA TCTCAGCGAT CTGTCTATTT CGTTCATCCA	5100
TAGTTGCCTG ACTCCCCGTC GTGTAGATAA CTACGATACG GGAGGGCTTA	5150
CCATCTGGCC CCAGTGCTGC AATGATACCG CGAGACCCAC GCTCACCGGC	5200

-continued

TCCAGATTTA	TCAGCAATAA	ACCAGCCAGC	CGGAAGGGCC	GAGCGCAGAA	5250
GTGGTCCTGC	AACTTTATCC	GCCTCCATCC	AGTCTATTAA	TTGTTGCCGG	5300
GAAGCTAGAG	TAAGTAGTTC	GCCAGTTAAT	AGTTTGCGCA	ACGTTGTTC	5350
CATTGCTGCA	GGCATCGTGG	TGTCACGCTC	GTCGTTTGGT	ATGGCTTCAT	5400
TCAGCTCCGG	TTCCCAACGA	TCAAGGCGAG	TTACATGATC	CCCCATGTTG	5450
TGCAAAAAAG	CGGTTAGCTC	CTTCGGTCCT	CCGATCGTTG	TCAGAAGTAA	5500
GTTGGCCGCA	GTGTTATCAC	TCATGTTTAT	GGCAGCACTG	CATAATTCTC	5550
TTACTGTCAT	GCCATCCGTA	AGATGCTTTT	CTGTGACTGG	TGAGTACTCA	5600
ACCAAGTCAT	TCTGAGAATA	GTGTATGCGG	CGACCGAGTT	GCTCTTGCCC	5650
GGCGTCAACA	CGGGATAATA	CCGCGCCACA	TAGCAGAACT	TTAAAAGTGC	5700
TCATCATTGG	AAAACGTTCT	TCGGGGCGAA	AACTCTCAAG	GATCTTACCG	5750
CTGTTGAGAT	CCAGTTCGAT	GTAACCCACT	CGTGCACCCA	ACTGATCTTC	5800
AGCATCTTTT	ACTTTCACCA	CGGTTTCTGG	GTGAGCAAAA	ACAGGAAGGC	5850
AAAATGCCGC	AAAAAAGGGA	ATAAGGGCGA	CACGGAAATG	TTGAATACTC	5900
ATACTCTTCC	TTTTTCAATA	TTATTGAAGC	ATTTATCAGG	GTTATTGTCT	5950
CATGAGCGGA	TACATATTTG	AATGTATTTA	GAAAAATAAA	CAATAGGGG	6000
TTCCGCGCAC	ATTTCCCCGA	AAAGTGCCAC	CTGACGTCTA	AGAAACCATT	6050
ATTATCATGA	CATTAAACCTA	TAAAAATAGG	CGTATCACGA	GGCCCTTTCG	6100
TCTTCAAGAA	TTCTCATGTT	TGACAGCTTA	TCGTAGACAT	CATGCGTGCT	6150
GTTGGTGTAT	TTCTGGCCAT	CTGTCTTGTC	ACCATTTTCG	TCCTCCCAAC	6200
ATGGGGCAAT	TGGGCATACC	CATGTTGTCA	CGTCACTCAG	CTCCGCGCTC	6250
AACACCTTCT	CGCGTTGGAA	AACATTAGCG	ACATTTACCT	GGTGAGCAAT	6300
CAGACATGCG	ACGGCTTTAG	CCTGGCCTCC	TTAAATTACAC	CTAAGAATGG	6350
GAGCAACCAG	CAGGAAAAGG	ACAAGCAGCG	AAAATTCACG	CCCCCTTGGG	6400
AGGTGGCGGC	ATATGCAAAG	GATAGCACTC	CCACTCTACT	ACTGGGTATC	6450
ATATGCTGAC	TGTATATGCA	TGAGGATAGC	ATATGCTACC	CGGATACAGA	6500
TTAGGATAGC	ATATACTACC	CAGATATAGA	TTAGGATAGC	ATATGCTACC	6550
CAGATATAGA	TTAGGATAGC	CTATGCTACC	CAGATATAAA	TTAGGATAGC	6600
ATATACTACC	CAGATATAGA	TTAGGATAGC	ATATGCTACC	CAGATATAGA	6650
TTAGGATAGC	CTATGCTACC	CAGATATAGA	TTAGGATAGC	ATATGCTACC	6700
CAGATATAGA	TTAGGATAGC	ATATGCTATC	CAGATATTTG	GGTAGTATAT	6750
GCTACCCAGA	TATAAATTAG	GATAGCATAT	ACTACCCTAA	TCTCTATTAG	6800
GATAGCATAT	GCTACCCGGA	TACAGATTAG	GATAGCATAT	ACTACCCAGA	6850
TATAGATTAG	GATAGCATAT	GCTACCCAGA	TATAGATTAG	GATAGCCTAT	6900
GCTACCCAGA	TATAAATTAG	GATAGCATAT	ACTACCCAGA	TATAGATTAG	6950
GATAGCATAT	GCTACCCAGA	TATAGATTAG	GATAGCCTAT	GCTACCCAGA	7000
TATAGATTAG	GATAGCATAT	GCTATCCAGA	TATTTGGGTA	GTATATGCTA	7050
CCCATGGCAA	CATTAGCCCA	CCGTGCTCTC	AGCGACCTCG	TGAATATGAG	7100
GACCAACAAC	CCTGTGCTTG	GCGCTCAGGC	GCAAGTGTGT	GTAATTTGTC	7150
CTCCAGATCG	CAGCAATCGC	GCCCCTATCT	TGGCCCGCCC	ACCTACTTAT	7200

-continued

GCAGGTATTC CCCGGGGTGC CATTAGTGGT TTTGTGGGCA AGTGGTTTGA	7250
CCGCAGTGGT TAGCGGGGTT ACAATCAGCC AAGTTATTAC ACCCTTATTT	7300
TACAGTCCAA AACCGCAGGG CGGCGTGTGG GGGCTGACGC GTGCCCCAC	7350
TCCACAATTT CAAAAAAG AGTGGCCACT TGTCTTTGTT TATGGGCCCC	7400
ATTGGCGTGG AGCCCCGTTT AATTTTCGGG GGTGTTAGAG ACAACCAAGT	7450
GAGTCCGCTG CTGTCGGCGT CCACTCTCTT TCCCCTTGTT ACAAATAGAG	7500
TGTAACAACA TGGTTCACCT GTCTTGGTCC CTGCCTGGGA CACATCTTAA	7550
TAACCCCAAGT ATCATATTGC ACTAGGATTA TGTGTGCCCC ATAGCCATAA	7600
ATTTCGTGTGA GATGGACATC CAGTCTTTAC GGCTTGTCCC CACCCCATGG	7650
ATTTCATATTG TTAAAGATAT TCAGAATGTT TCATTCCTAC ACTAGTATTT	7700
ATTGCCCAAG GGGTTTGTGA GGGTTATATT GGTGTCATAG CACAATGCCA	7750
CCACTGAACC CCCCGTCCAA ATTTTATTCT GGGGGCGTCA CCTGAAACCT	7800
TGTTTTCGAG CACCTCACAT ACACCTTACT GTTCACAACT CAGCAGTTAT	7850
TCTATTAGCT AAACGAAGGA GAATGAAGAA GCAGGCGAAG ATTCAGGAGA	7900
GTTCACTGCC CGCTCCTTGA TCTTCAGCCA CTGCCCTTGT GACTAAAATG	7950
GTTCACTACC CTCGTGGAAT CCTGACCCCA TGTAATAAAA ACCGTGACAG	8000
CTCATGGGGT GGGAGATATC GCTGTTCTTT AGGACCCTTT TACTAACCCCT	8050
AATTCGATAG CATATGCTTC CCGTTGGGTA ACATATGCTA TTGAATTAGG	8100
GTTAGTCTGG ATAGTATATA CTACTACCCG GGAAGCATAT GCTACCCGTT	8150
TAGGGTTAAC AAGGGGGCCT TATAAACACT ATTGCTAATG CCCTCTTGAG	8200
GGTCCGCTTA TCGGTAGCTA CACAGGCCCC TCTGATTGAC GTTGGTGTAG	8250
CCTCCCGTAG TCTTCCTGGG CCCCTGGGAG GTACATGTCC CCCAGCATTG	8300
GTGTAAGAGC TTCAGCCAAG AGTTACACAT AAAGGCAATG TTGTGTTGCA	8350
GTCCACAGAC TGCAAAGTCT GCTCCAGGAT GAAAGCCACT CAGTGTGGC	8400
AAATGTGCAC ATCCATTTAT AAGGATGTCA ACTACAGTCA GAGAACCCT	8450
TTGTGTTTGG TCCCCCCCCG TGTACATGT GGAACAGGGC CCAGTTGGCA	8500
AGTTGTACCA ACCAACTGAA GGGATTACAT GCACTGCCCC	8540

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 30 bases
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Other nucleic acid

(A) DESCRIPTION: Oligonucleotide primer UDEC690

(iii) HYPOTHETICAL: No

(iv) ANTI-SENSE: No

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

CGCCTGCAGG ACATCCAGAT GACTCAGTCT	30
----------------------------------	----

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 30 bases
- (B) TYPE: Nucleic Acid

-continued

(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Other nucleic acid
(A) DESCRIPTION: Oligonucleotide primer UDEC395

(iii) HYPOTHETICAL: No

(iv) ANTI-SENSE: Yes

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

CCCAAGCTTA CTGGATGGTG GGAAGATGGA 30

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 747 base pairs
(B) TYPE: Nucleic Acid
(C) STRANDEDNESS: Double
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other nucleic acid
(A) DESCRIPTION: 5G1.1M1 scFv (murine)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

ATG GCC GAC ATC CAG ATG ACT CAG TCT CCA 30
Met Ala Asp Ile Gln Met Thr Gln Ser Pro
1 5 10

GCT TCA CTG TCT GCA TCT GTG GGA GAA ACT 60
Ala Ser Leu Ser Ala Ser Val Gly Glu Thr
15 20

GTC ACC ATC ACA TGT GGA GCA AGT GAG AAT 90
Val Thr Ile Thr Cys Gly Ala Ser Glu Asn
25 30

ATT TAC GGT GCT TTA AAT TGG TAT CAG CGG 120
Ile Tyr Gly Ala Leu Asn Trp Tyr Gln Arg
35 40

AAA CAG GGA AAA TCT CCT CAG CTC CTG ATC 150
Lys Gln Gly Lys Ser Pro Gln Leu Leu Ile
45 50

TAT GGT GCA ACC AAC TTG GCA GAT GGC ATG 180
Tyr Gly Ala Thr Asn Leu Ala Asp Gly Met
55 60

TCA TCG AGG TTC AGT GGC AGT GGA TCT GGT 210
Ser Ser Arg Phe Ser Gly Ser Gly Ser Gly
65 70

AGA CAG TAT TAT CTC AAG ATC AGT AGC CTG 240
Arg Gln Tyr Tyr Leu Lys Ile Ser Ser Leu
75 80

CAT CCT GAC GAT GTT GCA ACG TAT TAC TGT 270
His Pro Asp Asp Val Ala Thr Tyr Tyr Cys
85 90

CAA AAT GTG TTA AAT ACT CCT CTC ACG TTC 300
Gln Asn Val Leu Asn Thr Pro Leu Thr Phe
95 100

GGT GCT GGG ACC AAG TTG GAG CTG AAA CGG 330
Gly Ala Gly Thr Lys Leu Glu Leu Lys Arg
105 110

ACC GGA GGT GGC GGG TCG GGT GGC GGG GGA 360
Thr Gly Gly Gly Gly Ser Gly Gly Gly Gly
115 120

TCG GGT GGC GGA GGG TCG CAG GTT CAG CTG 390
Ser Gly Gly Gly Gly Ser Gln Val Gln Leu
125 130

-continued

CAG CAG TCT GGA GCC GAG CTG ATG AAG CCT Gln Gln Ser Gly Ala Glu Leu Met Lys Pro 135 140	420
GGG GCC TCA GTG AAG ATG TCC TGC AAG GCT Gly Ala Ser Val Lys Met Ser Cys Lys Ala 145 150	450
ACT GGC TAC ATA TTC AGT AAC TAC TGG ATA Thr Gly Tyr Ile Phe Ser Asn Tyr Trp Ile 155 160	480
CAG TGG ATA AAG CAG AGG CCT GGA CAT GGC Gln Trp Ile Lys Gln Arg Pro Gly His Gly 165 170	510
CTT GAG TGG ATT GGT GAG ATT TTA CCT GGA Leu Glu Trp Ile Gly Glu Ile Leu Pro Gly 175 180	540
AGT GGT TCT ACT GAG TAC ACT GAG AAC TTC Ser Gly Ser Thr Glu Tyr Thr Glu Asn Phe 185 190	570
AAG GAC AAG GCC GCA TTC ACT GCA GAT ACA Lys Asp Lys Ala Ala Phe Thr Ala Asp Thr 195 200	600
TCC TCC AAC ACA GCC TAC ATG CAA CTC AGC Ser Ser Asn Thr Ala Tyr Met Gln Leu Ser 205 210	630
AGC CTG ACA TCA GAG GAC TCT GCC GTC TAT Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr 215 220	660
TAC TGT GCA AGA TAT TTC TTC GGT AGT AGC Tyr Cys Ala Arg Tyr Phe Phe Gly Ser Ser 225 230	690
CCC AAC TGG TAC TTC GAT GTC TGG GGC GCA Pro Asn Trp Tyr Phe Asp Val Trp Gly Ala 235 240	720
GGG ACC ACG GTC ACC GTC TCC TCA TGA Gly Thr Thr Val Thr Val Ser Ser 245	747

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 747 base pairs
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other nucleic acid

- (A) DESCRIPTION: 5G1.1 scFv CB (humanized)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

ATG GCC GAT ATC CAG ATG ACC CAG TCC CCG Met Ala Asp Ile Gln Met Thr Gln Ser Pro 1 5 10	30
TCC TCC CTG TCC GCC TCT GTG GGC GAT AGG Ser Ser Leu Ser Ala Ser Val Gly Asp Arg 15 20	60
GTC ACC ATC ACC TGC GGC GCC AGC GAA AAC Val Thr Ile Thr Cys Gly Ala Ser Glu Asn 25 30	90
ATC TAT GGC GCG CTG AAC TGG TAT CAA CGT Ile Tyr Gly Ala Leu Asn Trp Tyr Gln Arg 35 40	120

-continued

AAA CCT GGG AAA GCT CCG AAG CTT CTG ATT Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile 45 50	150
TAC GGT GCG ACG AAC CTG GCA GAT GGA GTC Tyr Gly Ala Thr Asn Leu Ala Asp Gly Val 55 60	180
CCT TCT CGC TTC TCT GGA TCC GGC TCC GGA Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly 65 70	210
ACG GAT TTC ACT CTG ACC ATC AGC AGT CTG Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu 75 80	240
CAG CCT GAA GAC TTC GCT ACG TAT TAC TGT Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys 85 90	270
CAG AAC GTT TTA AAT ACT CCG TTG ACT TTC Gln Asn Val Leu Asn Thr Pro Leu Thr Phe 95 100	300
GGA CAG GGT ACC AAG GTG GAA ATA AAA CGT Gly Gln Gly Thr Lys Val Glu Ile Lys Arg 105 110	330
ACT GGC GGT GGT GGT TCT GGT GGC GGT GGA Thr Gly Gly Gly Gly Ser Gly Gly Gly Gly 115 120	360
TCT GGT GGT GGC GGT TCT CAA GTC CAA CTG Ser Gly Gly Gly Gly Ser Gln Val Gln Leu 125 130	390
GTG CAA TCC GGC GCC GAG GTC AAG AAG CCA Val Gln Ser Gly Ala Glu Val Lys Lys Pro 135 140	420
GGG GCC TCA GTC AAA GTG TCC TGT AAA GCT Gly Ala Ser Val Lys Val Ser Cys Lys Ala 145 150	450
AGC GGC TAT ATT TTT TCT AAT TAT TGG ATT Ser Gly Tyr Ile Phe Ser Asn Tyr Trp Ile 155 160	480
CAA TGG GTG CGT CAG GCC CCC GGG CAG GGC Gln Trp Val Arg Gln Ala Pro Gly Gln Gly 165 170	510
CTG GAA TGG ATG GGT GAG ATC TTA CCG GGC Leu Glu Trp Met Gly Glu Ile Leu Pro Gly 175 180	540
TCT GGT AGC ACC GAA TAT ACC GAA AAT TTT Ser Gly Ser Thr Glu Tyr Thr Glu Asn Phe 185 190	570
AAA GAC CGT GTT ACT ATG ACG CGT GAC ACT Lys Asp Arg Val Thr Met Thr Arg Asp Thr 195 200	600
TCG ACT AGT ACA GTA TAC ATG GAG CTC TCC Ser Thr Ser Thr Val Tyr Met Glu Leu Ser 205 210	630
AGC CTG CGA TCG GAG GAC ACG GCC GTC TAT Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr 215 220	660
TAT TGC GCG CGT TAT TTT TTT GGT TCT AGC Tyr Cys Ala Arg Tyr Phe Phe Gly Ser Ser 225 230	690
CCG AAT TGG TAT TTT GAT GTT TGG GGT CAA Pro Asn Trp Tyr Phe Asp Val Trp Gly Gln 235 240	720

-continued

GGA ACC CTG GTC ACT GTC TCG AGC TGA 747
 Gly Thr Leu Val Thr Val Ser Ser
 245

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 726 base pairs
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other nucleic acid

(A) DESCRIPTION:5G1.1M1 VL HuK (chimeric light chain)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

ATG GGA ATC CAA GGA GGG TCT GTC CTG TTC Met Gly Ile Gln Gly Gly Ser Val Leu Phe -25 -20	30
GGG CTG CTG CTC GTC CTG GCT GTC TTC TGC Gly Leu Leu Leu Val Leu Ala Val Phe Cys -15 -10	60
CAT TCA GGT CAT AGC CTG CAG GAC ATC CAG His Ser Gly His Ser Leu Gln Asp Ile Gln -5 1 5	90
ATG ACT CAG TCT CCA GCT TCA CTG TCT GCA Met Thr Gln Ser Pro Ala Ser Leu Ser Ala 10 15	120
TCT GTG GGA GAA ACT GTC ACC ATC ACA TGT Ser Val Gly Glu Thr Val Thr Ile Thr Cys 20 25	150
GGA GCA AGT GAG AAT ATT TAC GGT GCT TTA Gly Ala Ser Glu Asn Ile Tyr Gly Ala Leu 30 35	180
AAT TGG TAT CAG CGG AAA CAG GGA AAA TCT Asn Trp Tyr Gln Arg Lys Gln Gly Lys Ser 40 45	210
CCT CAG CTC CTG ATC TAT GGT GCA ACC AAC Pro Gln Leu Leu Ile Tyr Gly Ala Thr Asn 50 55	240
TTG GCA GAT GGC ATG TCA TCG AGG TTC AGT Leu Ala Asp Gly Met Ser Ser Arg Phe Ser 60 65	270
GGC AGT GGA TCT GGT AGA CAG TAT TAT CTC Gly Ser Gly Ser Gly Arg Gln Tyr Tyr Leu 70 75	300
AAG ATC AGT AGC CTG CAT CCT GAC GAT GTT Lys Ile Ser Ser Leu His Pro Asp Asp Val 80 85	330
GCA ACG TAT TAC TGT CAA AAT GTG TTA AAT Ala Thr Tyr Tyr Cys Gln Asn Val Leu Asn 90 95	360
ACT CCT CTC ACG TTC GGT GCT GGG ACC AAG Thr Pro Leu Thr Phe Gly Ala Gly Thr Lys 100 105	390
TTG GAG CTG AAA CGA ACT GTG GCT GCA CCA Leu Glu Leu Lys Arg Thr Val Ala Ala Pro 110 115	420
TCT GTC TTC ATC TTC CCG CCA TCT GAT GAG Ser Val Phe Ile Phe Pro Pro Ser Asp Glu 120 125	450

-continued

CAG TTG AAA TCT GGA ACT GCC TCT GTT GTG Gln Leu Lys Ser Gly Thr Ala Ser Val Val 130 135	480
TGC CTG CTG AAT AAC TTC TAT CCC AGA GAG Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu 140 145	510
GCC AAA GTA CAG TGG AAG GTG GAT AAC GCC Ala Lys Val Gln Trp Lys Val Asp Asn Ala 150 155	540
CTC CAA TCG GGT AAC TCC CAG GAG AGT GTC Leu Gln Ser Gly Asn Ser Gln Glu Ser Val 160 165	570
ACA GAG CAG GAC AGC AAG GAC AGC ACC TAC Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr 170 175	600
AGC CTC AGC AGC ACC CTG ACG CTG AGC AAA Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys 180 185	630
GCA GAC TAC GAG AAA CAC AAA GTC TAC GCC Ala Asp Tyr Glu Lys His Lys Val Tyr Ala 190 195	660
TGC GAA GTC ACC CAT CAG GGC CTG AGC TCG Cys Glu Val Thr His Gln Gly Leu Ser Ser 200 205	690
CCC GTC ACA AAG AGC TTC AAC AGG GGA GAG Pro Val Thr Lys Ser Phe Asn Arg Gly Glu 210 215	720
TGT TAG Cys	726

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 750 base pairs
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other nucleic acid

(A) DESCRIPTION: 5G1.1M1 VH +HuG1 (chimeric Fd)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

ATG AAA TGG AGC TGG GTT ATT CTC TTC CTC Met Lys Trp Ser Trp Val Ile Leu Phe Leu -15 -10	30
CTG TCA GTA ACT GCA GGT GTC CAC TCC CAG Leu Ser Val Thr Ala Gly Val His Ser Gln -5 1	60
GTT CAG CTG CAG CAG TCT GGA GCT GAG CTG Val Gln Leu Gln Gln Ser Gly Ala Glu Leu 5 10	90
ATG AAG CCT GGG GCC TCA GTG AAG ATG TCC Met Lys Pro Gly Ala Ser Val Lys Met Ser 15 20	120
TGC AAG GCT ACT GGC TAC ATA TTC AGT AAC Cys Lys Ala Thr Gly Tyr Ile Phe Ser Asn 25 30	150
TAC TGG ATA CAG TGG ATA AAG CAG AGG CCT Tyr Trp Ile Gln Trp Ile Lys Gln Arg Pro 35 40	180
GGA CAT GGC CTT GAG TGG ATT GGT GAG ATT Gly His Gly Leu Glu Trp Ile Gly Glu Ile 45 50	210

-continued

TTA CCT GGA AGT GGT TCT ACT GAG TAC ACT Leu Pro Gly Ser Gly Ser Thr Glu Tyr Thr 55 60	240
GAG AAC TTC AAG GAC AAG GCC GCA TTC ACT Glu Asn Phe Lys Asp Lys Ala Ala Phe Thr 65 70	270
GCA GAT ACA TCC TCC AAC ACA GCC TAC ATG Ala Asp Thr Ser Ser Asn Thr Ala Tyr Met 75 80	300
CAA CTC AGC AGC CTG ACA TCA GAG GAC TCT Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser 85 90	330
GCC GTC TAT TAC TGT GCA AGA TAT TTC TTC Ala Val Tyr Tyr Cys Ala Arg Tyr Phe Phe 95 100	360
GGT AGT AGC CCC AAC TGG TAC TTC GAT GTC Gly Ser Ser Pro Asn Trp Tyr Phe Asp Val 105 110	390
TGG GGC GCA GGG ACC ACG GTC ACC GTC TCC Trp Gly Ala Gly Thr Thr Val Thr Val Ser 115 120	420
TCA GCC TCC ACC AAG GGC CCA TCG GTC TTC Ser Ala Ser Thr Lys Gly Pro Ser Val Phe 125 130	450
CCC CTG GCG CCC TCC TCC AAG AGC ACC TCT Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser 135 140	480
GGG GGC ACA GCG GCC CTG GGC TGC CTG GTC Gly Gly Thr Ala Ala Leu Gly Cys Leu Val 145 150	510
AAG GAC TAC TTC CCC GAA CCG GTG ACG GTG Lys Asp Tyr Phe Pro Glu Pro Val Thr Val 155 160	540
TCG TGG AAC TCA GGC GCC CTG ACC AGC GGC Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly 165 170	570
GTG CAC ACC TTC CCG GCT GTC CTA CAG TCC Val His Thr Phe Pro Ala Val Leu Gln Ser 175 180	600
TCA GGA CTC TAC TCC CTC AGC AGC GTG GTG Ser Gly Leu Tyr Ser Leu Ser Ser Val Val 185 190	630
ACC GTG CCC TCC AGC AGC TTG GGC ACC CAG Thr Val Pro Ser Ser Ser Leu Gly Thr Gln 195 200	660
ACC TAC ATC TGC AAC GTG AAT CAC AAG CCC Thr Tyr Ile Cys Asn Val Asn His Lys Pro 205 210	690
AGC AAC ACC AAG GTG GAC AAG AAA GTT GAG Ser Asn Thr Lys Val Asp Lys Lys Val Glu 215 220	720
CCC AAA TCT TGT GAC AAA ACT CAC ACA TAA Pro Lys Ser Cys Asp Lys Thr His Thr 225	750

(2) INFORMATION FOR SEQ ID NO:11:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 750 base pairs
 - (B) TYPE: Nucleic Acid
 - (C) STRANDEDNESS: Double
 - (D) TOPOLOGY: linear

-continued

(ii) MOLECULE TYPE: Other nucleic acid
 (A) DESCRIPTION:5G1.1 VH + IGHL (Humanized Fd)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

ATG AAG TGG AGC TGG GTT ATT CTC TTC CTC Met Lys Trp Ser Trp Val Ile Leu Phe Leu -15 -10	30
CTG TCA GTA ACT GCC GGC GTC CAC TCC CAA Leu Ser Val Thr Ala Gly Val His Ser Gln -5 1	60
GTC CAA CTG GTG CAA TCC GGC GCC GAG GTC Val Gln Leu Val Gln Ser Gly Ala Glu Val 5 10	90
AAG AAG CCA GGG GCC TCA GTC AAA GTG TCC Lys Lys Pro Gly Ala Ser Val Lys Val Ser 15 20	120
TGT AAA GCT AGC GGC TAT ATT TTT TCT AAT Cys Lys Ala Ser Gly Tyr Ile Phe Ser Asn 25 30	150
TAT TGG ATT CAA TGG GTG CGT CAG GCC CCC Tyr Trp Ile Gln Trp Val Arg Gln Ala Pro 35 40	180
GGG CAG GGC CTG GAA TGG ATG GGT GAG ATC Gly Gln Gly Leu Glu Trp Met Gly Glu Ile 45 50	210
TTA CCG GGC TCT GGT AGC ACC GAA TAT GCC Leu Pro Gly Ser Gly Ser Thr Glu Tyr Ala 55 60	240
CAA AAA TTC CAG GGC CGT GTT ACT ATG ACT Gln Lys Phe Gln Gly Arg Val Thr Met Thr 65 70	270
GCG GAC ACT TCG ACT AGT ACA GCC TAC ATG Ala Asp Thr Ser Thr Ser Thr Ala Tyr Met 75 80	300
GAG CTC TCC AGC CTG CGA TCG GAG GAC ACG Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr 85 90	330
GCC GTC TAT TAT TGC GCG CGT TAT TTT TTT Ala Val Tyr Tyr Cys Ala Arg Tyr Phe Phe 95 100	360
GGT TCT AGC CCG AAT TGG TAT TTT GAT GTT Gly Ser Ser Pro Asn Trp Tyr Phe Asp Val 105 110	390
TGG GGT CAA GGA ACC CTG GTC ACT GTC TCG Trp Gly Gln Gly Thr Leu Val Thr Val Ser 115 120	420
AGC GCC TCC ACC AAG GGC CCA TCG GTC TTC Ser Ala Ser Thr Lys Gly Pro Ser Val Phe 125 130	450
CCC CTG GCG CCC TCC TCC AAG AGC ACC TCT Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser 135 140	480
GGG GGC ACA GCG GCC CTG GGC TGC CTG GTC Gly Gly Thr Ala Ala Leu Gly Cys Leu Val 145 150	510
AAG GAC TAC TTC CCC GAA CCG GTG ACG GTG Lys Asp Tyr Phe Pro Glu Pro Val Thr Val 155 160	540

-continued

TCG TGG AAC TCA GGC GCC CTG ACC AGC GGC Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly 165 170	570
GTG CAC ACC TTC CCG GCT GTC CTA CAG TCC Val His Thr Phe Pro Ala Val Leu Gln Ser 175 180	600
TCA GGA CTC TAC TCC CTC AGC AGC GTG GTG Ser Gly Leu Tyr Ser Leu Ser Ser Val Val 185 190	630
ACC GTG CCC TCC AGC AGC TTG GGC ACC CAG Thr Val Pro Ser Ser Ser Leu Gly Thr Gln 195 200	660
ACC TAC ATC TGC AAC GTG AAT CAC AAG CCC Thr Tyr Ile Cys Asn Val Asn His Lys Pro 205 210	690
AGC AAC ACC AAG GTG GAC AAG AAA GTT GAG Ser Asn Thr Lys Val Asp Lys Lys Val Glu 215 220	720
CCC AAA TCT TGT GAC AAA ACT CAC ACA TAA Pro Lys Ser Cys Asp Lys Thr His Thr 225 230	750
(2) INFORMATION FOR SEQ ID NO:12:	
(i) SEQUENCE CHARACTERISTICS:	
(A) LENGTH: 750 base pairs	
(B) TYPE: Nucleic Acid	
(C) STRANDEDNESS: Double	
(D) TOPOLOGY: linear	
(ii) MOLECULE TYPE: Other nucleic acid	
(A) DESCRIPTION:5G1.1 VH + IGHRCL (Humanized Fd)	
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:	
ATG AAG TGG AGC TGG GTT ATT CTC TTC CTC Met Lys Trp Ser Trp Val Ile Leu Phe Leu -15 -10	30
CTG TCA GTA ACT GCC GGC GTC CAC TCC CAA Leu Ser Val Thr Ala Gly Val His Ser Gln -5 1	60
GTC CAA CTG GTG CAA TCC GGC GCC GAG GTC Val Gln Leu Val Gln Ser Gly Ala Glu Val 5 10	90
AAG AAG CCA GGG GCC TCA GTC AAA GTG TCC Lys Lys Pro Gly Ala Ser Val Lys Val Ser 15 20	120
TGT AAA GCT AGC GGC TAT ATT TTT TCT AAT Cys Lys Ala Ser Gly Tyr Ile Phe Ser Asn 25 30	150
TAT TGG ATT CAA TGG GTG CGT CAG GCC CCC Tyr Trp Ile Gln Trp Val Arg Gln Ala Pro 35 40	180
GGG CAG GGC CTG GAA TGG ATG GGT GAG ATC Gly Gln Gly Leu Glu Trp Met Gly Glu Ile 45 50	210
TTA CCG GGC TCT GGT AGC ACC GAA TAT ACC Leu Pro Gly Ser Gly Ser Thr Glu Tyr Thr 55 60	240
GAA AAT TTT AAA GAC CGT GTT ACT ATG ACG Glu Asn Phe Lys Asp Arg Val Thr Met Thr 65 70	270

-continued

CGT GAC ACT TCG ACT AGT ACA GTA TAC ATG Arg Asp Thr Ser Thr Ser Thr Val Tyr Met 75 80	300
GAG CTC TCC AGC CTG CGA TCG GAG GAC ACG Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr 85 90	330
GCC GTC TAT TAT TGC GCG CGT TAT TTT TTT Ala Val Tyr Tyr Cys Ala Arg Tyr Phe Phe 95 100	360
GGT TCT AGC CCG AAT TGG TAT TTT GAT GTT Gly Ser Ser Pro Asn Trp Tyr Phe Asp Val 105 110	390
TGG GGT CAA GGA ACC CTG GTC ACT GTC TCG Trp Gly Gln Gly Thr Leu Val Thr Val Ser 115 120	420
AGC GCC TCC ACC AAG GGC CCA TCG GTC TTC Ser Ala Ser Thr Lys Gly Pro Ser Val Phe 125 130	450
CCC CTG GCG CCC TCC TCC AAG AGC ACC TCT Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser 135 140	480
GGG GGC ACA GCG GCC CTG GGC TGC CTG GTC Gly Gly Thr Ala Ala Leu Gly Cys Leu Val 145 150	510
AAG GAC TAC TTC CCC GAA CCG GTG ACG GTG Lys Asp Tyr Phe Pro Glu Pro Val Thr Val 155 160	540
TCG TGG AAC TCA GGC GCC CTG ACC AGC GGC Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly 165 170	570
GTG CAC ACC TTC CCG GCT GTC CTA CAG TCC Val His Thr Phe Pro Ala Val Leu Gln Ser 175 180	600
TCA GGA CTC TAC TCC CTC AGC AGC GTG GTG Ser Gly Leu Tyr Ser Leu Ser Ser Val Val 185 190	630
ACC GTG CCC TCC AGC AGC TTG GGC ACC CAG Thr Val Pro Ser Ser Ser Leu Gly Thr Gln 195 200	660
ACC TAC ATC TGC AAC GTG AAT CAC AAG CCC Thr Tyr Ile Cys Asn Val Asn His Lys Pro 205 210	690
AGC AAC ACC AAG GTG GAC AAG AAA GTT GAG Ser Asn Thr Lys Val Asp Lys Lys Val Glu 215 220	720
CCC AAA TCT TGT GAC AAA ACT CAC ACA TAA Pro Lys Ser Cys Asp Lys Thr His Thr 225 230	750

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 726 base pairs
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other nucleic acid

- (A) DESCRIPTION: 5G1.1 VL +KLV56
(Humanized light chain)

-continued

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:	
ATG GGA ATC CAA GGA GGG TCT GTC CTG TTC Met Gly Ile Gln Gly Gly Ser Val Leu Phe -25 -20	30
GGG CTG CTG CTC GTC CTG GCT GTC TTC TGC Gly Leu Leu Leu Val Leu Ala Val Phe Cys -15 -10	60
CAT TCA GGT CAT AGC CTG CAG GAT ATC CAG His Ser Gly His Ser Leu Gln Asp Ile Gln -5 1 5	90
ATG ACC CAG TCC CCG TCC TCC CTG TCC GCC Met Thr Gln Ser Pro Ser Ser Leu Ser Ala 10 15	120
TCT GTG GGC GAT AGG GTC ACC ATC ACC TGC Ser Val Gly Asp Arg Val Thr Ile Thr Cys 20 25	150
GGC GCC AGC GAA AAC ATC TAT GGC GCG CTG Gly Ala Ser Glu Asn Ile Tyr Gly Ala Leu 30 35	180
AAC TGG TAT CAA CGT AAA CCT GGG AAA GCT Asn Trp Tyr Gln Arg Lys Pro Gly Lys Ala 40 45	210
CCG AAG CTT CTG ATT TAC GGT GCG ACG AAC Pro Lys Leu Leu Ile Tyr Gly Ala Thr Asn 50 55	240
CTG GCA GAT GGA GTC CCT TCT CGC TTC TCT Leu Ala Asp Gly Val Pro Ser Arg Phe Ser 60 65	270
GGA TCC GGC TCC GGA ACG GAT TAC ACT CTG Gly Ser Gly Ser Gly Thr Asp Tyr Thr Leu 70 75	300
ACC ATC AGC AGT CTG CAA CCT GAG GAC TTC Thr Ile Ser Ser Leu Gln Pro Glu Asp Phe 80 85	330
GCT ACG TAT TAC TGT CAG AAC GTT TTA AAT Ala Thr Tyr Tyr Cys Gln Asn Val Leu Asn 90 95	360
ACT CCG TTG ACT TTC GGA CAG GGT ACC AAG Thr Pro Leu Thr Phe Gly Gln Gly Thr Lys 100 105	390
GTG GAA ATA AAA CGA ACT GTG GCT GCA CCA Val Glu Ile Lys Arg Thr Val Ala Ala Pro 110 115	420
TCT GTC TTC ATC TTC CCG CCA TCT GAT GAG Ser Val Phe Ile Phe Pro Pro Ser Asp Glu 120 125	450
CAG TTG AAA TCT GGA ACT GCC TCT GTT GTG Gln Leu Lys Ser Gly Thr Ala Ser Val Val 130 135	480
TGC CTG CTG AAT AAC TTC TAT CCC AGA GAG Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu 140 145	510
GCC AAA GTA CAG TGG AAG GTG GAT AAC GCC Ala Lys Val Gln Trp Lys Val Asp Asn Ala 150 155	540
CTC CAA TCG GGT AAC TCC CAG GAG AGT GTC Leu Gln Ser Gly Asn Ser Gln Glu Ser Val 160 165	570

-continued

ACA GAG CAG GAC AGC AAG GAC AGC ACC TAC	600
Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr	
170 175	
AGC CTC AGC AGC ACC CTG ACG CTG AGC AAA	630
Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys	
180 185	
GCA GAC TAC GAG AAA CAC AAA GTC TAC GCC	660
Ala Asp Tyr Glu Lys His Lys Val Tyr Ala	
190 195	
TGC GAA GTC ACC CAT CAG GGC CTG AGC TCG	690
Cys Glu Val Thr His Gln Gly Leu Ser Ser	
200 205	
CCC GTC ACA AAG AGC TTC AAC AGG GGA GAG	720
Pro Val Thr Lys Ser Phe Asn Arg Gly Glu	
210 215	
TGT TAG	726
Cys	

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 726 base pairs
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other nucleic acid

- (A) DESCRIPTION: 5G1.1 VL +KL56B
- (Humanized light chain)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

ATG GGA ATC CAA GGA GGG TCT GTC CTG TTC	30
Met Gly Ile Gln Gly Gly Ser Val Leu Phe	
-25 -20	
GGG CTG CTG CTC GTC CTG GCT GTC TTC TGC	60
Gly Leu Leu Leu Val Leu Ala Val Phe Cys	
-15 -10	
CAT TCA GGT CAT AGC CTG CAG GAT ATC CAG	90
His Ser Gly His Ser Leu Gln Asp Ile Gln	
-5 1 5	
ATG ACC CAG TCC CCG TCC TCC CTG TCC GCC	120
Met Thr Gln Ser Pro Ser Ser Leu Ser Ala	
10 15	
TCT GTG GGC GAT AGG GTC ACC ATC ACC TGC	150
Ser Val Gly Asp Arg Val Thr Ile Thr Cys	
20 25	
GGC GCC AGC GAA AAC ATC TAT GGC GCG CTG	180
Gly Ala Ser Glu Asn Ile Tyr Gly Ala Leu	
30 35	
AAC TGG TAT CAA CGT AAA CCT GGG AAA GCT	210
Asn Trp Tyr Gln Arg Lys Pro Gly Lys Ala	
40 45	
CCG AAG CTT CTG ATT TAC GGT GCG ACG AAC	240
Pro Lys Leu Leu Ile Tyr Gly Ala Thr Asn	
50 55	
CTG GCA GAT GGA GTC CCT TCT CGC TTC TCT	270
Leu Ala Asp Gly Val Pro Ser Arg Phe Ser	
60 65	
GGA TCC GGC TCC GGA ACG GAT TTC ACT CTG	300
Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu	
70 75	

-continued

ACC ATC AGC AGT CTG CAG CCT GAA GAC TTC Thr Ile Ser Ser Leu Gln Pro Glu Asp Phe 80 85	330
GCT ACG TAT TAC TGT CAG AAC GTT TTA AAT Ala Thr Tyr Tyr Cys Gln Asn Val Leu Asn 90 95	360
ACT CCG TTG ACT TTC GGA CAG GGT ACC AAG Thr Pro Leu Thr Phe Gly Gln Gly Thr Lys 100 105	390
GTG GAA ATA AAA CGA ACT GTG GCT GCA CCA Val Glu Ile Lys Arg Pro Val Ala Ala Pro 110 115	420
TCT GTC TTC ATC TTC CCG CCA TCT GAT GAG Ser Val Phe Ile Phe Pro Pro Ser Asp Glu 120 125	450
CAG TTG AAA TCT GGA ACT GCC TCT GTT GTG Gln Leu Lys Ser Gly Thr Ala Ser Val Val 130 135	480
TGC CTG CTG AAT AAC TTC TAT CCC AGA GAG Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu 140 145	510
GCC AAA GTA CAG TGG AAG GTG GAT AAC GCC Ala Lys Val Gln Trp Lys Val Asp Asn Ala 150 155	540
CTC CAA TCG GGT AAC TCC CAG GAG AGT GTC Leu Gln Ser Gly Asn Ser Gln Glu Ser Val 160 165	570
ACA GAG CAG GAC AGC AAG GAC AGC ACC TAC Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr 170 175	600
AGC CTC AGC AGC ACC CTG ACG CTG AGC AAA Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys 180 185	630
GCA GAC TAC GAG AAA CAC AAA GTC TAC GCC Ala Asp Tyr Glu Lys His Lys Val Tyr Ala 190 195	660
TGC GAA GTC ACC CAT CAG GGC CTG AGC TCG Cys Glu Val Thr His Gln Gly Leu Ser Ser 200 205	690
CCC GTC ACA AAG AGC TTC AAC AGG GGA GAG Pro Val Thr Lys Ser Phe Asn Arg Gly Glu 210 215	720
TGT TAG Cys	726
(2) INFORMATION FOR SEQ ID NO:15:	
(i) SEQUENCE CHARACTERISTICS:	
(A) LENGTH: 711 base pairs	
(B) TYPE: Nucleic Acid	
(C) STRANDEDNESS: Double	
(D) TOPOLOGY: linear	
(ii) MOLECULE TYPE: Other nucleic acid	
(A) DESCRIPTION:5G1.1 VL + O12	
(Humanized light chain)	
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:	
ATG GAC ATG AGG GTC CCC GCT CAG CTC CTG Met Asp Met Arg Val Pro Ala Gln Leu Leu -20 -15	30

-continued

GGG CTC CTG CTA CTC TGG CTC CGA GGT GCC Gly Leu Leu Leu Leu Trp Leu Arg Gly Ala -10 -5	60
AGA TGT GAT ATC CAG ATG ACC CAG TCC CCG Arg Cys Asp Ile Gln Met Thr Gln Ser Pro 1 5	90
TCC TCC CTG TCC GCC TCT GTG GGC GAT AGG Ser Ser Leu Ser Ala Ser Val Gly Asp Arg 10 15	120
GTC ACC ATC ACC TGC GGC GCC AGC GAA AAC Val Thr Ile Thr Cys Gly Ala Ser Glu Asn 20 25	150
ATC TAT GGC GCG CTG AAC TGG TAT CAA CAG Ile Tyr Gly Ala Leu Asn Trp Tyr Gln Gln 30 35	180
AAA CCC GGG AAA GCT CCG AAG CTT CTG ATT Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile 40 45	210
TAC GGT GCG ACG AAC CTG GCA GAT GGA GTC Tyr Gly Ala Thr Asn Leu Ala Asp Gly Val 50 55	240
CCT TCT CGC TTC TCT GGA TCC GGC TCC GGA Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly 60 65	270
ACG GAT TTC ACT CTG ACC ATC AGC AGT CTG Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu 70 75	300
CAG CCT GAA GAC TTC GCT ACG TAT TAC TGT Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys 80 85	330
CAG AAC GTT TTA AAT ACT CCG TTG ACT TTC Gln Asn Val Leu Asn Thr Pro Leu Thr Phe 90 95	360
GGA CAG GGT ACC AAG GTG GAA ATA AAA CGA Gly Gln Gly Thr Lys Val Glu Ile Lys Arg 100 105	390
ACT GTG GCT GCA CCA TCT GTC TTC ATC TTC Thr Val Ala Ala Pro Ser Val Phe Ile Phe 110 115	420
CCG CCA TCT GAT GAG CAG TTG AAA TCT GGA Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly 120 125	450
ACT GCC TCT GTT GTG TGC CTG CTG AAT AAC Thr Ala Ser Val Val Cys Leu Leu Asn Asn 130 135	480
TTC TAT CCC AGA GAG GCC AAA GTA CAG TGG Phe Tyr Pro Arg Glu Ala Lys Val Gln Trp 140 145	510
AAG GTG GAT AAC GCC CTC CAA TCG GGT AAC Lys Val Asp Asn Ala Leu Gln Ser Gly Asn 150 155	540
TCC CAG GAG AGT GTC ACA GAG CAG GAC AGC Ser Gln Glu Ser Val Thr Glu Gln Asp Ser 160 165	570
AAG GAC AGC ACC TAC AGC CTC AGC AGC ACC Lys Asp Ser Thr Tyr Ser Leu Ser Ser Thr 170 175	600
CTG ACG CTG AGC AAA GCA GAC TAC GAG AAA Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys 180 185	630

-continued

CAC AAA GTC TAC GCC TGC GAA GTC ACC CAT His Lys Val Tyr Ala Cys Glu Val Thr His 190 195	660
CAG GGC CTG AGC TCG CCC GTC ACA AAG AGC Gln Gly Leu Ser Ser Pro Val Thr Lys Ser 200 205	690
TTC AAC AGG GGA GAG TGT TAG Phe Asn Arg Gly Glu Cys 210	711
(2) INFORMATION FOR SEQ ID NO:16:	
(i) SEQUENCE CHARACTERISTICS:	
(A) LENGTH: 750 base pairs	
(B) TYPE: Nucleic Acid	
(C) STRANDEDNESS: Double	
(D) TOPOLOGY: linear	
(ii) MOLECULE TYPE: Other nucleic acid	
(A) DESCRIPTION:5G1.1 VH + IGHRD (Humanized Fd)	
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:	
ATG AAG TGG AGC TGG GTT ATT CTC TTC CTC Met Lys Trp Ser Trp Val Ile Leu Phe Leu -15 -10	30
CTG TCA GTA ACT GCC GGC GTC CAC TCC CAA Leu Ser Val Thr Ala Gly Val His Ser Gln -5 1	60
GTC CAA CTG GTG CAA TCC GGC GCC GAG GTC Val Gln Leu Val Gln Ser Gly Ala Glu Val 5 10	90
AAG AAG CCA GGG GCC TCA GTC AAA GTG TCC Lys Lys Pro Gly Ala Ser Val Lys Val Ser 15 20	120
TGT AAA GCT AGC GGC TAT ATT TTT TCT AAT Cys Lys Ala Ser Gly Tyr Ile Phe Ser Asn 25 30	150
TAT TGG ATT CAA TGG GTG CGT CAG GCC CCC Tyr Trp Ile Gln Trp Val Arg Gln Ala Pro 35 40	180
GGG CAG GGC CTG GAA TGG ATG GGT GAG ATC Gly Gln Gly Leu Glu Trp Met Gly Glu Ile 45 50	210
TTA CCG GGC TCT GGT AGC ACC GAA TAT GCC Leu Pro Gly Ser Gly Ser Thr Glu Tyr Ala 55 60	240
CAA AAA TTC CAG GGC CGT GTT ACT ATG ACT Gln Lys Phe Gln Gly Arg Val Thr Met Thr 65 70	270
CGT GAC ACT TCG ACT AGT ACA GTA TAC ATG Arg Asp Thr Ser Thr Ser Thr Val Tyr Met 75 80	300
GAG CTC TCC AGC CTG CGA TCG GAG GAC ACG Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr 85 90	330
GCC GTC TAT TAT TGC GCG CGT TAT TTT TTT Ala Val Tyr Tyr Cys Ala Arg Tyr Phe Phe 95 100	360
GGT TCT AGC CCG AAT TGG TAT TTT GAT GTT Gly Ser Ser Pro Asn Trp Tyr Phe Asp Val 105 110	390

-continued

TGG GGT CAA GGA ACC CTG GTC ACT GTC TCG	420
Trp Gly Gln Gly Thr Leu Val Thr Val Ser	
115 120	
AGC GCC TCC ACC AAG GGC CCA TCG GTC TTC	450
Ser Ala Ser Thr Lys Gly Pro Ser Val Phe	
125 130	
CCC CTG GCG CCC TCC TCC AAG AGC ACC TCT	480
Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser	
135 140	
GGG GGC ACA GCG GCC CTG GGC TGC CTG GTC	510
Gly Gly Thr Ala Ala Leu Gly Cys Leu Val	
145 150	
AAG GAC TAC TTC CCC GAA CCG GTG ACG GTG	540
Lys Asp Tyr Phe Pro Glu Pro Val Thr Val	
155 160	
TCG TGG AAC TCA GGC GCC CTG ACC AGC GGC	570
Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly	
165 170	
GTG CAC ACC TTC CCG GCT GTC CTA CAG TCC	600
Val His Thr Phe Pro Ala Val Leu Gln Ser	
175 180	
TCA GGA CTC TAC TCC CTC AGC AGC GTG GTG	630
Ser Gly Leu Tyr Ser Leu Ser Ser Val Val	
185 190	
ACC GTG CCC TCC AGC AGC TTG GGC ACC CAG	660
Thr Val Pro Ser Ser Ser Leu Gly Thr Gln	
195 200	
ACC TAC ATC TGC AAC GTG AAT CAC AAG CCC	690
Thr Tyr Ile Cys Asn Val Asn His Lys Pro	
205 210	
AGC AAC ACC AAG GTG GAC AAG AAA GTT GAG	720
Ser Asn Thr Lys Val Asp Lys Lys Val Glu	
215 220	
CCC AAA TCT TGT GAC AAA ACT CAC ACA TAA	750
Pro Lys Ser Cys Asp Lys Thr His Thr	
225 230	

(2) INFORMATION FOR SEQ ID NO:17:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 747 base pairs
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other nucleic acid

- (A) DESCRIPTION: 5G1.1 scFv D012
(Humanized scFv)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

ATG GCC GAT ATC CAG ATG ACC CAG TCC CCG	30
Met Ala Asp Ile Gln Met Thr Gln Ser Pro	
1 5 10	
TCC TCC CTG TCC GCC TCT GTG GGC GAT AGG	60
Ser Ser Leu Ser Ala Ser Val Gly Asp Arg	
15 20	
GTC ACC ATC ACC TGC GGC GCC AGC GAA AAC	90
Val Thr Ile Thr Cys Gly Ala Ser Glu Asn	
25 30	
ATC TAT GGC GCG CTG AAC TGG TAT CAA CAG	120
Ile Tyr Gly Ala Leu Asn Trp Tyr Gln Gln	
35 40	

-continued

AAA CCT GGG AAA GCT CCG AAG CTT CTG ATT Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile 45 50	150
TAC GGT GCG ACG AAC CTG GCA GAT GGA GTC Tyr Gly Ala Thr Asn Leu Ala Asp Gly Val 55 60	180
CCT TCT CGC TTC TCT GGA TCC GGC TCC GGA Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly 65 70	210
ACG GAT TTC ACT CTG ACC ATC AGC AGT CTG Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu 75 80	240
CAG CCT GAA GAC TTC GCT ACG TAT TAC TGT Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys 85 90	270
CAG AAC GTT TTA AAT ACT CCG TTG ACT TTC Gln Asn Val Leu Asn Thr Pro Leu Thr Phe 95 100	300
GGA CAG GGT ACC AAG GTG GAA ATA AAA CGT Gly Gln Gly Thr Lys Val Glu Ile Lys Arg 105 110	330
ACT GGC GGT GGT GGT TCT GGT GGC GGT GGA Thr Gly Gly Gly Gly Ser Gly Gly Gly Gly 115 120	360
TCT GGT GGT GGC GGT TCT CAA GTC CAA CTG Ser Gly Gly Gly Gly Ser Gln Val Gln Leu 125 130	390
GTG CAA TCC GGC GCC GAG GTC AAG AAG CCA Val Gln Ser Gly Ala Glu Val Lys Lys Pro 135 140	420
GGG GCC TCA GTC AAA GTG TCC TGT AAA GCT Gly Ala Ser Val Lys Val Ser Cys Lys Ala 145 150	450
AGC GGC TAT ATT TTT TCT AAT TAT TGG ATT Ser Gly Tyr Ile Phe Ser Asn Tyr Trp Ile 155 160	480
CAA TGG GTG CGT CAG GCC CCC GGG CAG GGC Gln Trp Val Arg Gln Ala Pro Gly Gln Gly 165 170	510
CTG GAA TGG ATG GGT GAG ATC TTA CCG GGC Leu Glu Trp Met Gly Glu Ile Leu Pro Gly 175 180	540
TCT GGT AGC ACC GAA TAT GCC CAA AAA TTC Ser Gly Ser Thr Glu Tyr Ala Gln Lys Phe 185 190	570
CAG GGC CGT GTT ACT ATG ACG CGT GAC ACT Gln Gly Arg Val Thr Met Thr Arg Asp Thr 195 200	600
TCG ACT AGT ACA GTA TAC ATG GAG CTC TCC Ser Thr Ser Thr Val Tyr Met Glu Leu Ser 205 210	630
AGC CTG CGA TCG GAG GAC ACG GCC GTC TAT Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr 215 220	660
TAT TGC GCG CGT TAT TTT TTT GGT TCT AGC Tyr Cys Ala Arg Tyr Phe Phe Gly Ser Ser 225 230	690
CCG AAT TGG TAT TTT GAT GTT TGG GGT CAA Pro Asn Trp Tyr Phe Asp Val Trp Gly Gln 235 240	720

-continued

GGA ACC CTG GTC ACT GTC TCG AGC TGA
 Gly Thr Leu Val Thr Val Ser Ser
 245

747

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 5248 base pairs
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Double
- (D) TOPOLOGY: Circular

(ii) MOLECULE TYPE: Other nucleic acid

- (A) DESCRIPTION: pET Trc SO5/NI
 prokaryotic expression vector

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

TGGCGAATGG GACGCGCCCT GTAGCGGCGC ATTAAGCGCG GCGGGTGTGG	50
TGGTTACGCG CAGCGTGACC GCTACACTTG CCAGCGCCCT AGCGCCCGCT	100
CCTTTCGCTT TCTTCCCTTC CTTTCTCGCC ACGTTCGCCG GCTTCCCCCG	150
TCAAGCTCTA AATCGGGGGC TCCCTTTAGG GTTCCGATTT AGTGCTTTAC	200
GGCACCTCGA CCCCAAAAAA CTTGATTAGG GTGATGGTTC ACGTAGTGGG	250
CCATCGCCCT GATAGACGGT TTTTCGCCCT TTGACGTTGG AGTCCACGTT	300
CTTTAATAGT GGACTCTTGT TCCAACTGG AACAACTC AACCTATCT	350
CGGTCTATTC TTTTGATTTA TAAGGATTT TGCCGATTTC GGCCTATTGG	400
TTAAAAATG AGCTGATTTA ACAAAAATTT AACGCGAATT TTAACAAAAT	450
ATTAACGTTT ACAATTTCAG GTGGCACTTT TCGGGGAAAT GTGCGCGGAA	500
CCCCTATTTC TTTATTTTTC TAAATACATT CAAATATGTA TCCGCTCATG	550
AGACAATAAC CCTGATAAAT GCTTCAATAA TATTGAAAAA GGAAGAGTAT	600
GAGTATTCAA CATTTCCGTG TCGCCCTTAT TCCCTTTTTT GCGGCATTTT	650
GCCTTCCTGT TTTTGCTCAC CCAGAAACGC TGGTGAAAGT AAAAGATGCT	700
GAAGATCAGT TGGGTGCACG AGTGGGTAC ATCGAACTGG ATCTCAACAG	750
CGGTAAGATC CTTGAGAGTT TTCGCCCGA AGAACGTTTT CCAATGATGA	800
GCACTTTTAA AGTTCTGCTA TGTGGCGCGG TATTATCCCG TATTGACGCC	850
GGGCAAGAGC AACTCGGTCG CCGCATACAC TATTCTCAGA ATGACTTGGT	900
TGAGTACTCA CCAGTCACAG AAAAGCATCT TACGGATGGC ATGACAGTAA	950
GAGAATTATG CAGTGCTGCC ATAACCATGA GTGATAACAC TCGGCCAAC	1000
TTACTTCTGA CAACGATCGG AGGACCGAAG GAGCTAACCG CTTTTTTGCA	1050
CAACATGGGG GATCATGTAA CTCGCCTTGA TCGTTGGGAA CCGGAGCTGA	1100
ATGAAGCCAT ACCAAACGAC GAGCGTGACA CCACGATGCC TGCAGCAATG	1150
GCAACAACGT TGCGCAAACT ATTAAGTGGC GAACTACTTA CTCTAGCTTC	1200
CCGGCAACAA TTAATAGACT GGATGGAGGC GGATAAAGTT GCAGGACCAC	1250
TTCTGCGCTC GGCCCTTCCG GCTGGCTGGT TTATTGCTGA TAAATCTGGA	1300
GCCGGTGAGC GTGGGTCTCG CGGTATCATT GCAGCACTGG GGCCAGATGG	1350
TAAGCCCTCC CGTATCGTAG TTATCTACAC GACGGGGAGT CAGGCAACTA	1400
TGGATGAACG AAATAGACAG ATCGCTGAGA TAGGTGCCTC ACTGATTAAG	1450
CATTGGTAAC TGTGAGACCA AGTTTACTCA TATATACTTT AGATTGATTT	1500
AAAACCTCAT TTTTAATTTA AAAGGATCTA GGTGAAGATC CTTTTTGATA	1550

-continued

ATCTCATGAC CAAATCCCT TAACGTGAGT TTTCGTTCCA CTGAGCGTCA	1600
GACCCCGTAG AAAAGATCAA AGGATCTTCT TGAGATCCTT TTTTCTGCG	1650
CGTAATCTGC TGCTTGCAAA CAAAAAACC ACCGCTACCA GCGGTGGTTT	1700
GTTTGCCGGA TCAAGAGCTA CCAACTCTTT TTCCGAAGGT AACTGGCTTC	1750
AGCAGAGCGC AGATACCAAA TACTGTCCTT CTAGTGTAGC CGTAGTTAGG	1800
CCACCACTTC AAGAACTCTG TAGCACCGCC TACATACCTC GCTCTGCTAA	1850
TCCTGTTACC AGTGGCTGCT GCCAGTGGCG ATAAGTCGTG TCTTACCGG	1900
TTGGACTIONA GACGATAGTT ACCGGATAAG GCGCAGCGGT CGGGCTGAAC	1950
GGGGGGTTCG TGCACACAGC CCAGCTTGGA GCGAACGACC TACACCGAAC	2000
TGAGATACCT ACAGCGTGAG CTATGAGAAA GCGCCACGCT TCCCGAAGG	2050
AGAAAGGCGG ACAGGTATCC GGTAAAGCGC AGGGTCGGA CAGGAGAGCG	2100
CACGAGGGAG CTTCCAGGGG GAAACGCCTG GTATCTTTAT AGTCCTGTCTG	2150
GGTTTCGCCA CCTCTGACTT GAGCGTCGAT TTTGTGATG CTCGTCAGG	2200
GGGCGGAGCC TATGGAAAAA CGCCAGCAAC GCGGCCTTTT TACGGTTCCT	2250
GGCCTTTTGC TGGCCTTTTG CTCACATGTT CTTTCTGCG TTATCCCCTG	2300
ATTCTGTGGA TAACCGTATT ACCGCCTTTG AGTGAGCTGA TACCGCTCGC	2350
CGCAGCCGAA CGACCGAGCG CAGCGAGTCA GTGAGCGAGG AAGCGGAAGA	2400
GCGCCTGATG CGGTATTTTC TCCTTACGCA TCTGTGCGGT ATTTACACC	2450
GCATATATGG TGCACTCTCA GTACAATCTG CTCTGATGCC GCATAGTTAA	2500
GCCAGTATAC ACTCCGCTAT CGCTACGTGA CTGGTCATG GCTGCGCCCC	2550
GACACCCGCC AACACCCGCT GACGCGCCCT GACGGGCTTG TCTGCTCCCG	2600
GCATCCGCTT ACAGACAAGC TGTGACCGTC TCCGGGAGCT GCATGTGTCA	2650
GAGGTTTTCA CCGTCATCAC CGAAACGCGC GAGGAGCTG CGGTAAAGCT	2700
CATCAGCGTG GTCGTGAAGC GATTACAGA TGTCTGCCTG TTCATCCGCG	2750
TCCAGCTCGT TGAGTTTCTC CAGAAGCGTT AATGTCTGGC TTCTGATAAA	2800
GCGGGCCATG TTAAGGGCGG TTTTCTCTG TTTGGTCACT GATGCCTCCG	2850
TGTAAGGGGG ATTTCTGTTC ATGGGGGTAA TGATACCGAT GAAACGAGAG	2900
AGGATGTCTA CGATACGGGT TACTGATGAT GAACATGCCC GGTACTGGA	2950
ACGTTGTGAG GGTAAACAAC TGGCGGTATG GATGCGGCGG GACCAGAGAA	3000
AAATCACTCA GGTCAATGC CAGCGCTTCG TTAATACAGA TGTAGGTGTT	3050
CCACAGGTA GCCAGCAGCA TCCTGCGATG CAGATCCGGA ACATAATGGT	3100
GCAGGGCGCT GACTTCCGCG TTTCCAGACT TTACGAAACA CGGAAACCGA	3150
AGACCATTCA TGTGTTGCT CAGGTCGAG ACGTTTTGCA GCAGCAGTCG	3200
CTTCACGTTT GCTCGCGTAT CGGTGATTCA TTCTGCTAAC CAGTAAGGCA	3250
ACCCCGCCAG CCTAGCCGGG TCCTCAACGA CAGGAGCAG ATCATGCGCA	3300
CCCGTGGGGC CGCCATGCCG GCGATAATGG CCTGCTTCTC GCCGAAACGT	3350
TTGGTGGCGG GACCAGTGAC GAAGGCTTGA GCGAGGCGT GCAAGATTCC	3400
GAATACCGCA AGCGACAGGC CGATCATCGT CGCGCTCCAG CGAAAGCGGT	3450
CCTCGCCGAA AATGACCCAG AGCGCTGCCG GCACCTGTCC TACGAGTTGC	3500
ATGATAAAGA AGACAGTCAT AAGTGCGGCG ACGATAGTCA TGCCCCGCG	3550

-continued

CCACCGAAG GAGCTGACTG GGTGAAGGC TCTCAAGGC ATCGGTCGAG	3600
ATCCCGGTGC CTAATGAGTG AGCTAACTTA CATTAATTGC GTTGCGCTCA	3650
CTGCCCCTT TCCAGTCGGG AAACCTGTCG TGCCAGCTGC ATTAATGAAT	3700
CGGCCAACGC GCGGGGAGAG GCGGTTTTCG TATTGGGCGC CAGGGTGGTT	3750
TTTCTTTTCA CCAGTGAGAC GGGCAACAGC TGATTGCCCT TCACCGCCTG	3800
GCCCTGAGAG AGTTGCAGCA AGCGGTCCAC GCTGGTTTGC CCCAGCAGGC	3850
GAAATCCTG TTTGATGGTG GTTAACGGCG GGATATAACA TGAGCTGTCT	3900
TCGGTATCGT CGTATCCAC TACCGAGATA TCCGCACCAA CGCGCAGCCC	3950
GGACTCGGTA ATGGCGCGCA TTGCGCCAG CGCCATCTGA TCGTTGGCAA	4000
CCAGCATCGC AGTGGGAACG ATGCCCTCAT TCAGCATTTG CATGGTTTGT	4050
TGAAAACCGG ACATGGCACT CCAGTCGCCT TCCCGTTCCG CTATCGGCTG	4100
AATTGTATTG CGAGTGAGAT ATTTATGCCA GCCAGCCAGA CGCAGACGCG	4150
CCGAGACAGA ACTTAATGGG CCCGCTAACA GCGCGATTTG CTGGTGACCC	4200
AATGCGACCA GATGCTCCAC GCCCAGTCGC GTACCGTCTT CATGGGAGAA	4250
AATAATACTG TTGATGGGTG TCTGGTCAGA GACATCAAGA AATAACGCCG	4300
GAACATTAGT GCAGGCAGCT TCCACAGCAA TGGCATCCTG GTCATCCAGC	4350
GGATAGTTAA TGATCAGCCC ACTGACGCGT TGCGCGAGAA GATTGTGCAC	4400
CGCCGCTTTA CAGGCTTCGA CGCCGCTTCG TTCTACCATC GACACCACCA	4450
CGCTGGCACC CAGTTGATCG GCGCGAGATT TAATCGCCGC GACAATTTGC	4500
GACGGCGCGT GCAGGGCCAG ACTGGAGGTG GCAACGCCAA TCAGCAACGA	4550
CTGTTTGCCC GCCAGTTGTT GTGCCACGCG GTTGGGAATG TAATTCAGCT	4600
CCGCCATCGC CGCTTCCACT TTTTCCCGCG TTTTCGAGA AACGTGGCTG	4650
GCCTGGTTCA CCACGCGGGA AACGGTCTGA TAAGAGACAC CGGCATACTC	4700
TGCGACATCG TATAACGTTA CTGGTTTCAC ATTCACCACC CTGAATTGAC	4750
TCTCTCCGG GCGCTATCAT GCCATACCGC GAAAGGTTTT GCGCCATTCTG	4800
ATGGTGTCG GATCTCGAC GCTCTCCCTT ATGCGACTCC TGCATTAGGA	4850
AGCAGCCCAG TAGTAGGTTG AGGCCGTTGA GCACCGCCGC CGCAAGGAAT	4900
GGTGCATGCG GTACCAGCTG TTGACAATTA ATCATCCGGC TCGTATAATA	4950
GTACTGTGTG GAATTGTGAG CGCTCACAAAT TCCACACATC TAGAAATAAT	5000
TTTGTTTAAC TTTAAGAAGG AGATATACCA TGGAGATCTG GATCCATCGA	5050
TGAATTCGAG CTCCGTCGAC AAGCTTGCGG CCGCACTCGA GCACCACCAC	5100
CACCACCACT GAGATCCGGC TGCTAACAAA GCCCGAAAGG AAGCTGAGTT	5150
GGCTGCTGCC ACCGCTGAGC AATAACTAGC ATAACCCCTT GGGGCCTCTA	5200
AACGGGTCTT GAGGGGTTTT TTGCTGAAAG GAGGAATAT ATCCGGAT	5248

(2) INFORMATION FOR SEQ ID NO:19:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 783 base pairs
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other nucleic acid

- (A) DESCRIPTION: N19/8 scFv (His Tagged)

-continued

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:	
ATG GCC AAT ATT GTG CTG ACC CAA TCT CCA Met Ala Asn Ile Val Leu Thr Gln Ser Pro 1 5 10	30
GCT TCT TTG GCT GTG TCT CTA GGG CAG AGG Ala Ser Leu Ala Val Ser Leu Gly Gln Arg 15 20	60
GCC ACC ATA TCC TGC AGA GCC AGT GAA AGT Ala Thr Ile Ser Cys Arg Ala Ser Glu Ser 25 30	90
GTT GAT AGT TAT GAC AAT AGT TTT ATG CAC Val Asp Ser Tyr Asp Asn Ser Phe Met His 35 40	120
TGG TAC CAG CAG AAA CCA GGA CAG CCA CCC Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro 45 50	150
AAA CTC CTC ATC TTT CTT GCA TCC AAC CTA Lys Leu Leu Ile Phe Leu Ala Ser Asn Leu 55 60	180
GAA TCT GGG GTC CCT GCC AGG TTC AGT GGC Glu Ser Gly Val Pro Ala Arg Phe Ser Gly 65 70	210
AGT GGG TCT AGG ACA GAC TTC ACC CTC ACC Ser Gly Ser Arg Thr Asp Phe Thr Leu Thr 75 80	240
ATT GAT CCT GTG GAG GCT GAT GAT GCT GCA Ile Asp Pro Val Glu Ala Asp Asp Ala Ala 85 90	270
ACC TAT TAC TGT CAG CAA AAT AAT GAG GTT Thr Tyr Tyr Cys Gln Gln Asn Asn Glu Val 95 100	300
CCG AAC ACG TTC GGA GGG GGG ACC AAG CTG Pro Asn Thr Phe Gly Gly Gly Thr Lys Leu 105 110	330
GAA ATA AAA CGG ACC GGA GGT GGC GGG TCG Glu Ile Lys Arg Thr Gly Gly Gly Gly Ser 115 120	360
GGT GGC GGG GGA TCG GGT GGC GGA GGG TCG Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser 125 130	390
GAC GTC AAG CTC GTG GAG TCT GGG GGA GAC Asp Val Lys Leu Val Glu Ser Gly Gly Asp 135 140	420
TTA GTG AAG CTT GGA GGG TCC CTG AAA CTC Leu Val Lys Leu Gly Gly Ser Leu Lys Leu 145 150	450
TCC TGT GCA GCC TCT GGA TTC ACC TTC AGT Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser 155 160	480
AGC TAT TAT ATG TCT TGG GTT CGC CAG ATT Ser Tyr Tyr Met Ser Trp Val Arg Gln Ile 165 170	510
TCA GAG AAG AGG CTG GAG TTG GTC GCA GCC Ser Glu Lys Arg Leu Glu Leu Val Ala Ala 175 180	540
ATT AAT AGT AAT GGT GAT AGC ACC TAC TAT Ile Asn Ser Asn Gly Asp Ser Thr Tyr Tyr 185 190	570

-continued

CCA GAC ACT GTG AAG GGC CGA TTC ACC ATC Pro Asp Thr Val Lys Gly Arg Phe Thr Ile 195 200	600
TCC AGA GAC AAT GCC AAG AGC ACC CTG GAT Ser Arg Asp Asn Ala Lys Ser Thr Leu Asp 205 210	630
CTG CAA ATG AGC AGT CTG AAG TCT GAG GAC Leu Gln Met Ser Ser Leu Lys Ser Glu Asp 215 220	660
ACA GCC TTG TAT TTC TGT GTA AGA GAG ACT Thr Ala Leu Tyr Phe Cys Val Arg Glu Thr 225 230	690
TAT TAC TAC GGG ATT AGT CCC GTC TTC GAT Tyr Tyr Tyr Gly Ile Ser Pro Val Phe Asp 235 240	720
GTC TGG GGC ACA GGG ACC ACG GTC ACC GTC Val Trp Gly Thr Gly Thr Thr Val Thr Val 245 250	750
TCC TCA CTC GAG CAC CAC CAC CAC CAC CAC Ser Ser Leu Glu His His His His His His 255 260	780
TGA	783

(2) INFORMATION FOR SEQ ID NO:20:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 747 base pairs

(B) TYPE: Nucleic Acid

(C) STRANDEDNESS: Double

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other nucleic acid

(A) DESCRIPTION: 5G1.1 scFv C012 (humanized)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

ATG GCC GAT ATC CAG ATG ACC CAG TCC CCG Met Ala Asp Ile Gln Met Thr Gln Ser Pro 1 5 10	30
TCC TCC CTG TCC GCC TCT GTG GGC GAT AGG Ser Ser Leu Ser Ala Ser Val Gly Asp Arg 15 20	60
GTC ACC ATC ACC TGC GGC GCC AGC GAA AAC Val Thr Ile Thr Cys Gly Ala Ser Glu Asn 25 30	90
ATC TAT GGC GCG CTG AAC TGG TAT CAA CAG Ile Tyr Gly Ala Leu Asn Trp Tyr Gln Gln 35 40	120
AAA CCC GGG AAA GCT CCG AAG CTT CTG ATT Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile 45 50	150
TAC GGT GCG ACG AAC CTG GCA GAT GGA GTC Tyr Gly Ala Thr Asn Leu Ala Asp Gly Val 55 60	180
CCT TCT CGC TTC TCT GGA TCC GGC TCC GGA Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly 65 70	210
ACG GAT TTC ACT CTG ACC ATC AGC AGT CTG Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu 75 80	240
CAG CCT GAA GAC TTC GCT ACG TAT TAC TGT Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys 85 90	270

-continued

CAG AAC GTT TTA AAT ACT CCG TTG ACT TTC Gln Asn Val Leu Asn Thr Pro Leu Thr Phe 95 100	300
GGA CAG GGT ACC AAG GTG GAA ATA AAA CGT Gly Gln Gly Thr Lys Val Glu Ile Lys Arg 105 110	330
ACT GGC GGT GGT GGT TCT GGT GGC GGT GGA Thr Gly Gly Gly Gly Ser Gly Gly Gly Gly 115 120	360
TCT GGT GGT GGC GGT TCT CAA GTC CAA CTG Ser Gly Gly Gly Gly Ser Gln Val Gln Leu 125 130	390
GTG CAA TCC GGC GCC GAG GTC AAG AAG CCA Val Gln Ser Gly Ala Glu Val Lys Lys Pro 135 140	420
GGG GCC TCA GTC AAA GTG TCC TGT AAA GCT Gly Ala Ser Val Lys Val Ser Cys Lys Ala 145 150	450
AGC GGC TAT ATT TTT TCT AAT TAT TGG ATT Ser Gly Tyr Ile Phe Ser Asn Tyr Trp Ile 155 160	480
CAA TGG GTG CGT CAG GCC CCC GGG CAG GGC Gln Trp Val Arg Gln Ala Pro Gly Gln Gly 165 170	510
CTG GAA TGG ATG GGT GAG ATC TTA CCG GGC Leu Glu Trp Met Gly Glu Ile Leu Pro Gly 175 180	540
TCT GGT AGC ACC GAA TAT ACC GAA AAT TTT Ser Gly Ser Thr Glu Tyr Thr Glu Asn Phe 185 190	570
AAA GAC CGT GTT ACT ATG ACG CGT GAC ACT Lys Asp Arg Val Thr Met Thr Arg Asp Thr 195 200	600
TCG ACT AGT ACA GTA TAC ATG GAG CTC TCC Ser Thr Ser Thr Val Tyr Met Glu Leu Ser 205 210	630
AGC CTG CGA TCG GAG GAC ACG GCC GTC TAT Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr 215 220	660
TAT TGC GCG CGT TAT TTT TTT GGT TCT AGC Tyr Cys Ala Arg Tyr Phe Phe Gly Ser Ser 225 230	690
CCG AAT TGG TAT TTT GAT GTT TGG GGT CAA Pro Asn Trp Tyr Phe Asp Val Trp Gly Gln 235 240	720
GGA ACC CTG GTC ACT GTC TCG AGC TGA Gly Thr Leu Val Thr Val Ser Ser 245	747

(2) INFORMATION FOR SEQ ID NO:21:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 747 base pairs
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other nucleic acid

- (A) DESCRIPTION: 5G1.1 scFv DO12B
(Humanized scFv)

-continued

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:									
ATG GCC GAT ATC CAG ATG ACC CAG TCC CCG									30
Met Ala Asp Ile Gln Met Thr Gln Ser Pro									
1 5 10									
TCC TCC CTG TCC GCC TCT GTG GGC GAT AGG									60
Ser Ser Leu Ser Ala Ser Val Gly Asp Arg									
15 20									
GTC ACC ATC ACC TGC CGT GCT AGC GAA AAC									90
Val Thr Ile Thr Cys Arg Ala Ser Glu Asn									
25 30									
ATC TAT GGC GCG CTG AAC TGG TAT CAA CAG									120
Ile Tyr Gly Ala Leu Asn Trp Tyr Gln Gln									
35 40									
AAA CCT GGG AAA GCT CCG AAG CTT CTG ATT									150
Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile									
45 50									
TAC GGT GCG ACG AAC CTG GCA GAT GGA GTC									180
Tyr Gly Ala Thr Asn Leu Ala Asp Gly Val									
55 60									
CCT TCT CGC TTC TCT GGA TCC GGC TCC GGA									210
Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly									
65 70									
ACG GAT TTC ACT CTG ACC ATC AGC AGT CTG									240
Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu									
75 80									
CAG CCT GAA GAC TTC GCT ACG TAT TAC TGT									270
Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys									
85 90									
CAG AAC GTT TTA AAT ACT CCG TTG ACT TTC									300
Gln Asn Val Leu Asn Thr Pro Leu Thr Phe									
95 100									
GGA CAG GGT ACC AAG GTG GAA ATA AAA CGT									330
Gly Gln Gly Thr Lys Val Glu Ile Lys Arg									
105 110									
ACT GGC GGT GGT GGT TCT GGT GGC GGT GGA									360
Thr Gly Gly Gly Gly Ser Gly Gly Gly Gly									
115 120									
TCT GGT GGT GGC GGT TCT CAA GTC CAA CTG									390
Ser Gly Gly Gly Gly Ser Gln Val Gln Leu									
125 130									
GTG CAA TCC GGC GCC GAG GTC AAG AAG CCA									420
Val Gln Ser Gly Ala Glu Val Lys Lys Pro									
135 140									
GGG GCC TCA GTC AAA GTG TCC TGT AAA GCT									450
Gly Ala Ser Val Lys Val Ser Cys Lys Ala									
145 150									
AGC GGC TAT ATT TTT TCT AAT TAT TGG ATT									480
Ser Gly Tyr Ile Phe Ser Asn Tyr Trp Ile									
155 160									
CAA TGG GTG CGT CAG GCC CCC GGG CAG GGC									510
Gln Trp Val Arg Gln Ala Pro Gly Gln Gly									
165 170									
CTG GAA TGG ATG GGT GAG ATC TTA CCG GGC									540
Leu Glu Trp Met Gly Glu Ile Leu Pro Gly									
175 180									
TCT GGT AGC ACC GAA TAT GCC CAA AAA TTC									570
Ser Gly Ser Thr Glu Tyr Ala Gln Lys Phe									
185 190									

-continued

CAG GGC CGT GTT ACT ATG ACG CGT GAC ACT Gln Gly Arg Val Thr Met Thr Arg Asp Thr 195 200	600
TCG ACT AGT ACA GTA TAC ATG GAG CTC TCC Ser Thr Ser Thr Val Tyr Met Glu Leu Ser 205 210	630
AGC CTG CGA TCG GAG GAC ACG GCC GTC TAT Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr 215 220	660
TAT TGC GCG CGT TAT TTT TTT GGT TCT AGC Tyr Cys Ala Arg Tyr Phe Phe Gly Ser Ser 225 230	690
CCG AAT TGG TAT TTT GAT GTT TGG GGT CAA Pro Asn Trp Tyr Phe Asp Val Trp Gly Gln 235 240	720
GGA ACC CTG GTC ACT GTC TCG AGC TGA Gly Thr Leu Val Thr Val Ser Ser 245	747

(2) INFORMATION FOR SEQ ID NO:22:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 747 base pairs
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other nucleic acid

- (A) DESCRIPTION: 5G1.1 scFv D012C
(Humanized scFv)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

ATG GCC GAT ATC CAG ATG ACC CAG TCC CCG Met Ala Asp Ile Gln Met Thr Gln Ser Pro 1 5 10	30
TCC TCC CTG TCC GCC TCT GTG GGC GAT AGG Ser Ser Leu Ser Ala Ser Val Gly Asp Arg 15 20	60
GTC ACC ATC ACC TGC GGC GCC AGC GAA AAC Val Thr Ile Thr Cys Gly Ala Ser Glu Asn 25 30	90
ATC TAT GGC GCG CTG AAC TGG TAT CAA CAG Ile Tyr Gly Ala Leu Asn Trp Tyr Gln Gln 35 40	120
AAA CCT GGG AAA GCT CCG AAG CTT CTG ATT Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile 45 50	150
TAC GGT GCG ACG AGC CTG CAG TCT GGA GTC Tyr Gly Ala Thr Ser Leu Gln Ser Gly Val 55 60	180
CCT TCT CGC TTC TCT GGA TCC GGC TCC GGA Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly 65 70	210
ACG GAT TTC ACT CTG ACC ATC AGC AGT CTG Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu 75 80	240
CAG CCT GAA GAC TTC GCT ACG TAT TAC TGT Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys 85 90	270
CAG AAC GTT TTA AAT ACT CCG TTG ACT TTC Gln Asn Val Leu Asn Thr Pro Leu Thr Phe 95 100	300

-continued

GGA CAG GGT ACC AAG GTG GAA ATA AAA CGT Gly Gln Gly Thr Lys Val Glu Ile Lys Arg 105 110	330
ACT GGC GGT GGT GGT TCT GGT GGC GGT GGA Thr Gly Gly Gly Gly Ser Gly Gly Gly Gly 115 120	360
TCT GGT GGT GGC GGT TCT CAA GTC CAA CTG Ser Gly Gly Gly Gly Ser Gln Val Gln Leu 125 130	390
GTG CAA TCC GGC GCC GAG GTC AAG AAG CCA Val Gln Ser Gly Ala Glu Val Lys Lys Pro 135 140	420
GGG GCC TCA GTC AAA GTG TCC TGT AAA GCT Gly Ala Ser Val Lys Val Ser Cys Lys Ala 145 150	450
AGC GGC TAT ATT TTT TCT AAT TAT TGG ATT Ser Gly Tyr Ile Phe Ser Asn Tyr Trp Ile 155 160	480
CAA TGG GTG CGT CAG GCC CCC GGG CAG GGC Gln Trp Val Arg Gln Ala Pro Gly Gln Gly 165 170	510
CTG GAA TGG ATG GGT GAG ATC TTA CCG GGC Leu Glu Trp Met Gly Glu Ile Leu Pro Gly 175 180	540
TCT GGT AGC ACC GAA TAT GCC CAA AAA TTC Ser Gly Ser Thr Glu Tyr Ala Gln Lys Phe 185 190	570
CAG GGC CGT GTT ACT ATG ACG CGT GAC ACT Gln Gly Arg Val Thr Met Thr Arg Asp Thr 195 200	600
TCG ACT AGT ACA GTA TAC ATG GAG CTC TCC Ser Thr Ser Thr Val Tyr Met Glu Leu Ser 205 210	630
AGC CTG CGA TCG GAG GAC ACG GCC GTC TAT Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr 215 220	660
TAT TGC GCG CGT TAT TTT TTT GGT TCT AGC Tyr Cys Ala Arg Tyr Phe Phe Gly Ser Ser 225 230	690
CCG AAT TGG TAT TTT GAT GTT TGG GGT CAA Pro Asn Trp Tyr Phe Asp Val Trp Gly Gln 235 240	720
GGA ACC CTG GTC ACT GTC TCG AGC TGA Gly Thr Leu Val Thr Val Ser Ser 245	747
(2) INFORMATION FOR SEQ ID NO:23:	
(i) SEQUENCE CHARACTERISTICS:	
(A) LENGTH: 747 base pairs	
(B) TYPE: Nucleic Acid	
(C) STRANDEDNESS: Double	
(D) TOPOLOGY: linear	
(ii) MOLECULE TYPE: Other nucleic acid	
(A) DESCRIPTION: 5G1.1 scFv DO12D	
(Humanized scFv)	
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:	
ATG GCC GAT ATC CAG ATG ACC CAG TCC CCG Met Ala Asp Ile Gln Met Thr Gln Ser Pro 1 5 10	30

-continued

TCC TCC CTG TCC GCC TCT GTG GGC GAT AGG Ser Ser Leu Ser Ala Ser Val Gly Asp Arg 15 20	60
GTC ACC ATC ACC TGC CGT GCT AGC GAA AAC Val Thr Ile Thr Cys Arg Ala Ser Glu Asn 25 30	90
ATC TAT GGC GCG CTG AAC TGG TAT CAA CAG Ile Tyr Gly Ala Leu Asn Trp Tyr Gln Gln 35 40	120
AAA CCT GGG AAA GCT CCG AAG CTT CTG ATT Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile 45 50	150
TAC GGT GCG ACG AGC CTG CAG TCT GGA GTC Tyr Gly Ala Thr Ser Leu Gln Ser Gly Val 55 60	180
CCT TCT CGC TTC TCT GGA TCC GGC TCC GGA Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly 65 70	210
ACG GAT TTC ACT CTG ACC ATC AGC AGT CTG Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu 75 80	240
CAG CCT GAA GAC TTC GCT ACG TAT TAC TGT Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys 85 90	270
CAG AAC GTT TTA AAT ACT CCG TTG ACT TTC Gln Asn Val Leu Asn Thr Pro Leu Thr Phe 95 100	300
GGA CAG GGT ACC AAG GTG GAA ATA AAA CGT Gly Gln Gly Thr Lys Val Glu Ile Lys Arg 105 110	330
ACT GGC GGT GGT GGT TCT GGT GGC GGT GGA Thr Gly Gly Gly Gly Ser Gly Gly Gly Gly 115 120	360
TCT GGT GGT GGC GGT TCT CAA GTC CAA CTG Ser Gly Gly Gly Gly Ser Gln Val Gln Leu 125 130	390
GTG CAA TCC GGC GCC GAG GTC AAG AAG CCA Val Gln Ser Gly Ala Glu Val Lys Lys Pro 135 140	420
GGG GCC TCA GTC AAA GTG TCC TGT AAA GCT Gly Ala Ser Val Lys Val Ser Cys Lys Ala 145 150	450
AGC GGC TAT ATT TTT TCT AAT TAT TGG ATT Ser Gly Tyr Ile Phe Ser Asn Tyr Trp Ile 155 160	480
CAA TGG GTG CGT CAG GCC CCC GGG CAG GGC Gln Trp Val Arg Gln Ala Pro Gly Gln Gly 165 170	510
CTG GAA TGG ATG GGT GAG ATC TTA CCG GGC Leu Glu Trp Met Gly Glu Ile Leu Pro Gly 175 180	540
TCT GGT AGC ACC GAA TAT GCC CAA AAA TTC Ser Gly Ser Thr Glu Tyr Ala Gln Lys Phe 185 190	570
CAG GGC CGT GTT ACT ATG ACG CGT GAC ACT Gln Gly Arg Val Thr Met Thr Arg Asp Thr 195 200	600
TCG ACT AGT ACA GTA TAC ATG GAG CTC TCC Ser Thr Ser Thr Val Tyr Met Glu Leu Ser 205 210	630

-continued

AGC CTG CGA TCG GAG GAC ACG GCC GTC TAT Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr 215 220	660
TAT TGC GCG CGT TAT TTT TTT GGT TCT AGC Tyr Cys Ala Arg Tyr Phe Phe Gly Ser Ser 225 230	690
CCG AAT TGG TAT TTT GAT GTT TGG GGT CAA Pro Asn Trp Tyr Phe Asp Val Trp Gly Gln 235 240	720
GGA ACC CTG GTC ACT GTC TCG AGC TGA Gly Thr Leu Val Thr Val Ser Ser 245	747
(2) INFORMATION FOR SEQ ID NO:24:	
(i) SEQUENCE CHARACTERISTICS:	
(A) LENGTH: 747 base pairs	
(B) TYPE: Nucleic Acid	
(C) STRANDEDNESS: Double	
(D) TOPOLOGY: linear	
(ii) MOLECULE TYPE: Other nucleic acid	
(A) DESCRIPTION:5G1.1 scFv C013 (humanized)	
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:	
ATG GCC GAT ATC CAG ATG ACC CAG TCC CCG Met Ala Asp Ile Gln Met Thr Gln Ser Pro 1 5 10	30
TCC TCC CTG TCC GCC TCT GTG GGC GAT AGG Ser Ser Leu Ser Ala Ser Val Gly Asp Arg 15 20	60
GTC ACC ATC ACC TGC CGT GCT AGC GAA AAC Val Thr Ile Thr Cys Arg Ala Ser Glu Asn 25 30	90
ATC TAT GGC GCG CTG AAC TGG TAT CAA CAG Ile Tyr Gly Ala Leu Asn Trp Tyr Gln Gln 35 40	120
AAA CCC GGG AAA GCT CCG AAG CTT CTG ATT Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile 45 50	150
TAC GGT GCG ACG AAC CTG GCA GAT GGA GTC Tyr Gly Ala Thr Asn Leu Ala Asp Gly Val 55 60	180
CCT TCT CGC TTC TCT GGA TCC GGC TCC GGA Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly 65 70	210
ACG GAT TTC ACT CTG ACC ATC AGC AGT CTG Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu 75 80	240
CAG CCT GAA GAC TTC GCT ACG TAT TAC TGT Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys 85 90	270
CAG AAC GTT TTA AAT ACT CCG TTG ACT TTC Gln Asn Val Leu Asn Thr Pro Leu Thr Phe 95 100	300
GGA CAG GGT ACC AAG GTG GAA ATA AAA CGT Gly Gln Gly Thr Lys Val Glu Ile Lys Arg 105 110	330
ACT GGC GGT GGT GGT TCT GGT GGC GGT GGA Thr Gly Gly Gly Gly Ser Gly Gly Gly Gly 115 120	360

-continued

TCT GGT GGT GGC GGT TCT CAA GTC CAA CTG Ser Gly Gly Gly Gly Ser Gln Val Gln Leu 125 130	390
GTG CAA TCC GGC GCC GAG GTC AAG AAG CCA Val Gln Ser Gly Ala Glu Val Lys Lys Pro 135 140	420
GGG GCC TCA GTC AAA GTG TCC TGT AAA GCT Gly Ala Ser Val Lys Val Ser Cys Lys Ala 145 150	450
AGC GGC TAT ATT TTT TCT AAT TAT TGG ATT Ser Gly Tyr Ile Phe Ser Asn Tyr Trp Ile 155 160	480
CAA TGG GTG CGT CAG GCC CCC GGG CAG GGC Gln Trp Val Arg Gln Ala Pro Gly Gln Gly 165 170	510
CTG GAA TGG ATG GGT GAG ATC TTA CCG GGC Leu Glu Trp Met Gly Glu Ile Leu Pro Gly 175 180	540
TCT GGT AGC ACC GAA TAT ACC GAA AAT TTT Ser Gly Ser Thr Glu Tyr Thr Glu Asn Phe 185 190	570
AAA GAC CGT GTT ACT ATG ACG CGT GAC ACT Lys Asp Arg Val Thr Met Thr Arg Asp Thr 195 200	600
TCG ACT AGT ACA GTA TAC ATG GAG CTC TCC Ser Thr Ser Thr Val Tyr Met Glu Leu Ser 205 210	630
AGC CTG CGA TCG GAG GAC ACG GCC GTC TAT Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr 215 220	660
TAT TGC GCG CGT TAT TTT TTT GGT TCT AGC Tyr Cys Ala Arg Tyr Phe Phe Gly Ser Ser 225 230	690
CCG AAT TGG TAT TTT GAT GTT TGG GGT CAA Pro Asn Trp Tyr Phe Asp Val Trp Gly Gln 235 240	720
GGA ACC CTG GTC ACT GTC TCG AGC TGA Gly Thr Leu Val Thr Val Ser Ser 245	747

(2) INFORMATION FOR SEQ ID NO:25:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 747 base pairs
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other nucleic acid

- (A) DESCRIPTION: 5G1.1 scFv C014 (humanized)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

ATG GCC GAT ATC CAG ATG ACC CAG TCC CCG Met Ala Asp Ile Gln Met Thr Gln Ser Pro 1 5 10	30
TCC TCC CTG TCC GCC TCT GTG GGC GAT AGG Ser Ser Leu Ser Ala Ser Val Gly Asp Arg 15 20	60
GTC ACC ATC ACC TGC GGC GCC AGC GAA AAC Val Thr Ile Thr Cys Gly Ala Ser Glu Asn 25 30	90

-continued

ATC TAT GGC GCG CTG AAC TGG TAT CAA CAG Ile Tyr Gly Ala Leu Asn Trp Tyr Gln Gln 35 40	120
AAA CCC GGG AAA GCT CCG AAG CTT CTG ATT Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile 45 50	150
TAC GGT GCG ACG AGC CTG CAG TCT GGA GTC Tyr Gly Ala Thr Ser Leu Gln Ser Gly Val 55 60	180
CCT TCT CGC TTC TCT GGA TCC GGC TCC GGA Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly 65 70	210
ACG GAT TTC ACT CTG ACC ATC AGC AGT CTG Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu 75 80	240
CAG CCT GAA GAC TTC GCT ACG TAT TAC TGT Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys 85 90	270
CAG AAC GTT TTA AAT ACT CCG TTG ACT TTC Gln Asn Val Leu Asn Thr Pro Leu Thr Phe 95 100	300
GGA CAG GGT ACC AAG GTG GAA ATA AAA CGT Gly Gln Gly Thr Lys Val Glu Ile Lys Arg 105 110	330
ACT GGC GGT GGT GGT TCT GGT GGC GGT GGA Thr Gly Gly Gly Gly Ser Gly Gly Gly Gly 115 120	360
TCT GGT GGT GGC GGT TCT CAA GTC CAA CTG Ser Gly Gly Gly Gly Ser Gln Val Gln Leu 125 130	390
GTG CAA TCC GGC GCC GAG GTC AAG AAG CCA Val Gln Ser Gly Ala Glu Val Lys Lys Pro 135 140	420
GGG GCC TCA GTC AAA GTG TCC TGT AAA GCT Gly Ala Ser Val Lys Val Ser Cys Lys Ala 145 150	450
AGC GGC TAT ATT TTT TCT AAT TAT TGG ATT Ser Gly Tyr Ile Phe Ser Asn Tyr Trp Ile 155 160	480
CAA TGG GTG CGT CAG GCC CCC GGG CAG GGC Gln Trp Val Arg Gln Ala Pro Gly Gln Gly 165 170	510
CTG GAA TGG ATG GGT GAG ATC TTA CCG GGC Leu Glu Trp Met Gly Glu Ile Leu Pro Gly 175 180	540
TCT GGT AGC ACC GAA TAT ACC GAA AAT TTT Ser Gly Ser Thr Glu Tyr Thr Glu Asn Phe 185 190	570
AAA GAC CGT GTT ACT ATG ACG CGT GAC ACT Lys Asp Arg Val Thr Met Thr Arg Asp Thr 195 200	600
TCG ACT AGT ACA GTA TAC ATG GAG CTC TCC Ser Thr Ser Thr Val Tyr Met Glu Leu Ser 205 210	630
AGC CTG CGA TCG GAG GAC ACG GCC GTC TAT Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr 215 220	660
TAT TGC GCG CGT TAT TTT TTT GGT TCT AGC Tyr Cys Ala Arg Tyr Phe Phe Gly Ser Ser 225 230	690

-continued

CCG AAT TGG TAT TTT GAT GTT TGG GGT CAA Pro Asn Trp Tyr Phe Asp Val Trp Gly Gln 235 240	720
GGA ACC CTG GTC ACT GTC TCG AGC TGA Gly Thr Leu Val Thr Val Ser Ser 245	747
(2) INFORMATION FOR SEQ ID NO:26:	
(i) SEQUENCE CHARACTERISTICS:	
(A) LENGTH: 747 base pairs	
(B) TYPE: Nucleic Acid	
(C) STRANDEDNESS: Double	
(D) TOPOLOGY: linear	
(ii) MOLECULE TYPE: Other nucleic acid	
(A) DESCRIPTION:5G1.1 scFv C015 (humanized)	
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:	
ATG GCC GAT ATC CAG ATG ACC CAG TCC CCG Met Ala Asp Ile Gln Met Thr Gln Ser Pro 1 5 10	30
TCC TCC CTG TCC GCC TCT GTG GGC GAT AGG Ser Ser Leu Ser Ala Ser Val Gly Asp Arg 15 20	60
GTC ACC ATC ACC TGC CGT GCT AGC GAA AAC Val Thr Ile Thr Cys Arg Ala Ser Glu Asn 25 30	90
ATC TAT GGC GCG CTG AAC TGG TAT CAA CAG Ile Tyr Gly Ala Leu Asn Trp Tyr Gln Gln 35 40	120
AAA CCC GGG AAA GCT CCG AAG CTT CTG ATT Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile 45 50	150
TAC GGT GCG ACG AGC CTG CAG TCT GGA GTC Tyr Gly Ala Thr Ser Leu Gln Ser Gly Val 55 60	180
CCT TCT CGC TTC TCT GGA TCC GGC TCC GGA Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly 65 70	210
ACG GAT TTC ACT CTG ACC ATC AGC AGT CTG Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu 75 80	240
CAG CCT GAA GAC TTC GCT ACG TAT TAC TGT Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys 85 90	270
CAG AAC GTT TTA AAT ACT CCG TTG ACT TTC Gln Asn Val Leu Asn Thr Pro Leu Thr Phe 95 100	300
GGA CAG GGT ACC AAG GTG GAA ATA AAA CGT Gly Gln Gly Thr Lys Val Glu Ile Lys Arg 105 110	330
ACT GGC GGT GGT GGT TCT GGT GGC GGT GGA Thr Gly Gly Gly Gly Ser Gly Gly Gly Gly 115 120	360
TCT GGT GGT GGC GGT TCT CAA GTC CAA CTG Ser Gly Gly Gly Gly Ser Gln Val Gln Leu 125 130	390
GTG CAA TCC GGC GCC GAG GTC AAG AAG CCA Val Gln Ser Gly Ala Glu Val Lys Lys Pro 135 140	420

-continued

GGG GCC TCA GTC AAA GTG TCC TGT AAA GCT Gly Ala Ser Val Lys Val Ser Cys Lys Ala 145 150	450
AGC GGC TAT ATT TTT TCT AAT TAT TGG ATT Ser Gly Tyr Ile Phe Ser Asn Tyr Trp Ile 155 160	480
CAA TGG GTG CGT CAG GCC CCC GGG CAG GGC Gln Trp Val Arg Gln Ala Pro Gly Gln Gly 165 170	510
CTG GAA TGG ATG GGT GAG ATC TTA CCG GGC Leu Glu Trp Met Gly Glu Ile Leu Pro Gly 175 180	540
TCT GGT AGC ACC GAA TAT ACC GAA AAT TTT Ser Gly Ser Thr Glu Tyr Thr Glu Asn Phe 185 190	570
AAA GAC CGT GTT ACT ATG ACG CGT GAC ACT Lys Asp Arg Val Thr Met Thr Arg Asp Thr 195 200	600
TCG ACT AGT ACA GTA TAC ATG GAG CTC TCC Ser Thr Ser Thr Val Tyr Met Glu Leu Ser 205 210	630
AGC CTG CGA TCG GAG GAC ACG GCC GTC TAT Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr 215 220	660
TAT TGC GCG CGT TAT TTT TTT GGT TCT AGC Tyr Cys Ala Arg Tyr Phe Phe Gly Ser Ser 225 230	690
CCG AAT TGG TAT TTT GAT GTT TGG GGT CAA Pro Asn Trp Tyr Phe Asp Val Trp Gly Gln 235 240	720
GGA ACC CTG GTC ACT GTC TCG AGC TGA Gly Thr Leu Val Thr Val Ser Ser 245	747

What is claimed is:

1. An antibody comprising at least one antibody-antigen binding site, said antibody exhibiting specific binding to human complement component C5, said specific binding being targeted to the alpha chain of human complement component C5, wherein the antibody 1) inhibits complement activation in a human body fluid, 2) inhibits the binding of purified human complement component C5 to either human complement component C3 or human complement component C4, and 3) does not specifically bind to the human complement activation product free C5a.

2. The antibody of claim 1 wherein the inhibition of complement activation in the human body fluid is measurable as an increment of blockade of C5a generation and an increment of blockade of complement hemolytic activity in the body fluid, said increment of blockade of C5a generation being substantially equal to said increment of blockade of complement hemolytic activity.

3. The antibody of claim 1 wherein, upon binding to human C5, there is a 60% to 90% reduction in the ability of C5 to bind to human complement component C3.

4. The antibody of claim 1 wherein, upon binding to human C5, there is a 60% to 90% reduction in the ability of C5 to bind to human complement component C4.

5. The antibody of claim 1 wherein the antibody binds specifically to an isolated oligopeptide comprising an amino acid sequence corresponding to amino acid 8 through amino acid 12 of SEQ ID NO:1.

6. The antibody of claim 1 wherein the inhibition of complement activation in the human body fluid is measurable as a substantially complete blockade of C5a generation in the body fluid and a substantially complete blockade of complement hemolytic activity in the body fluid when the antibody is added to the body fluid at a concentration yielding a ratio equal to or less than 10 moles of antibody-antigen binding sites of the antibody to 1 mole of human C5 in the body fluid.

7. The antibody of claim 1 wherein the antibody is a humanized antibody.

8. The antibody of claim 1 wherein the antibody is an scFv.

9. The antibody of claim 1, wherein, when administered to a human patient via intravenous infusion, the antibody provides complete complement inhibition at dosages below 0.005 g/kg.

10. The antibody of claim 1, wherein, when administered to a human patient via intravenous infusion, the antibody provides therapeutic benefits at dosages below 0.0022 g/kg.

11. The antibody of claim 10, wherein the antibody is administered in association with an extracorporeal circulation procedure.

12. The antibody of claim 1 wherein the inhibition of complement activation in the human body fluid is measurable as a substantially complete blockade of C5a generation in the body fluid and a substantially complete blockade of complement hemolytic activity in the body fluid when the

143

antibody is added to the body fluid at a concentration yielding a ratio equal to or less than 3 moles of antibody-antigen binding sites of the antibody to 1 mole of human C5 in the body fluid.

13. The antibody of claim 1, wherein, when administered to a human patient via intravenous infusion, the antibody provides therapeutically effective complement inhibition at dosages below 0.003 g/kg.

14. A sterile non-pyrogenic therapeutic agent comprising the antibody of claim 1 in a formulation suitable for administration to a human.

15. The therapeutic agent of claim 14 wherein the antibody is a humanized immunoglobulin.

16. The therapeutic agent of claim 14 wherein the antibody is an scFv.

17. The therapeutic agent of claim 14 wherein the antibody is made up of two or more heterodimeric subunits each containing one heavy and one light chain.

18. Antibody 5G1.1 scFv CB (humanized) having the amino acid sequence encoded by the nucleic acid sequence of SEQ ID NO:8.

19. An isolated antigen binding protein comprising:

- 1) a variable light region CDR1 comprising an amino acid sequence corresponding to amino acid residues 26–36 of SEQ ID NO:8,
- 2) a variable light region CDR2 comprising an amino acid sequence corresponding to amino acid residues 52–58 of SEQ ID NO:8,
- 3) a variable light region CDR3 comprising an amino acid sequence corresponding to amino acid residues 91 through amino acid 99 of SEQ ID NO:8,
- 4) a variable heavy region CDR1 comprising an amino acid sequence corresponding to amino acid residues 152 through amino acid 161 of SEQ ID NO:8,

144

5) a variable heavy region CDR2 comprising an amino acid sequence corresponding to amino acid residues 176 through amino acid 192 of SEQ ID NO:8,

6) a variable heavy region CDR3 comprising an amino acid sequence corresponding to amino acid residues 225 through amino acid 237 of SEQ ID NO:8,

said protein exhibiting specific binding to human complement component C5, said specific binding being targeted to the alpha chain of human complement component C5, wherein the protein inhibits complement activation in a human body fluid and does not specifically bind to the human complement activation product free C5a.

20. A nucleic acid molecule comprising a nucleotide sequence encoding a polypeptide comprising a variable heavy chain region amino acid sequence corresponding to amino acid 1 through amino acid 122 of SEQ ID NO: 12.

21. Hybridoma 5G1.1 having ATCC designation HB-11625.

22. An antibody produced by the hybridoma of claim 21.

23. An antibody comprising at least one antibody-antigen binding site, said antibody exhibiting specific binding to human complement component C5, said specific binding being targeted to the alpha chain of human complement component C5, wherein:

(A) the antibody inhibits (i) C5b-9-mediated hemolysis and (ii) C5a generation in a fluid comprising human serum; and

(B) the antibody does not specifically bind to the human complement activation product free C5a.

* * * * *

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 1 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Title page, and Column 1, lines 1-3.

Delete the title "**C5-SPECIFIC ANTIBODIES FOR THE TREATMENT OF INFLAMMATORY DISEASES**" and insert therefor -- **ANTIBODIES TO HUMAN COMPLEMENT COMPONENT C5** --.

Column 19.

Line 54, delete "(SEQ ID NO:1)".

Line 55, following "KSSKC peptide", insert -- (SEQ ID NO:1) --.

Column 55.

Please delete lines 50-53, and insert therefor:

Val Ile Asp His Gln Gly Thr Lys Ser Ser
5 10

Lys Cys Val Arg Gln Lys Val Glu Gly Ser Ser
15 20

Column 57.

Please delete lines 22-37, and insert therefor:

Met Gly Leu Leu Gly Ile Leu Cys Phe Leu
-15 -10
Ile Phe Leu Gly Lys Thr Trp Gly Gln Glu Gln Thr Tyr Val
-5 -1 5
Ile Ser Ala Pro Lys Ile Phe Arg Val Gly Ala Ser Glu Asn
10 15 20
Ile Val Ile Gln Val Tyr Gly Tyr Thr Glu Ala Phe Asp Ala
25 30
Thr Ile Ser Ile Lys Ser Tyr Pro Asp Lys Lys Phe Ser Tyr
35 40 45
Ser Ser Gly His Val His Leu Ser Ser Glu Asn Lys Phe Gln
50 55 60
Asn Ser Ala Ile Leu Thr Ile Gln Pro Lys Gln Leu Pro Gly
65 70 75
Gly Gln Asn Pro Val Ser Tyr Val Tyr Leu Glu Val Val Ser
80 85 90

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 2 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 57,

Please delete lines 38-55, and insert therefor:

```
Lys His Phe Ser Lys Ser Lys Arg Met Pro Ile Thr Tyr Asp
                      95                      100

Asn Gly Phe Leu Phe Ile His Thr Asp Lys Pro Val Tyr Thr
105                      110                      115

Pro Asp Gln Ser Val Lys Val Arg Val Tyr Ser Leu Asn Asp
120                      125                      130

Asp Leu Lys Pro Ala Lys Arg Glu Thr Val Leu Thr Phe Ile
135                      140                      145

Asp Pro Glu Gly Ser Glu Val Asp Met Val Glu Glu Ile Asp
150                      155                      160

His Ile Gly Ile Ile Ser Phe Pro Asp Phe Lys Ile Pro Ser
165                      170

Asn Pro Arg Tyr Gly Met Trp Thr Ile Lys Ala Lys Tyr Lys
175                      180                      185

Glu Asp Phe Ser Thr Thr Gly Thr Ala Tyr Phe Glu Val Lys
190                      195                      200

Glu Tyr Val Leu Pro His Phe Ser Val Ser Ile Glu Pro Glu
205                      210                      215
```

Column 59,

Please delete lines 1-4, and insert therefor:

```
Tyr Asn Phe Ile Gly Tyr Lys Asn Phe Lys Asn Phe Glu Ile
220                      225                      230

Thr Ile Lys Ala Arg Tyr Phe Tyr Asn Lys Val Val Thr Glu
235                      240
```

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 3 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 59.

Please delete lines 5-28, and insert therefor:

Ala	Asp	Val	Tyr	Ile	Thr	Phe	Gly	Ile	Arg	Glu	Asp	Leu	Lys
245					250					255			
Asp	Asp	Gln	Lys	Glu	Met	Met	Gln	Thr	Ala	Met	Gln	Asn	Thr
260						265					270		
Met	Leu	Ile	Asn	Gly	Ile	Ala	Gln	Val	Thr	Phe	Asp	Ser	Glu
	275						280					285	
Thr	Ala	Val	Lys	Glu	Leu	Ser	Tyr	Tyr	Ser	Leu	Glu	Asp	Leu
		290						295					300
Asn	Asn	Lys	Tyr	Leu	Tyr	Ile	Ala	Val	Thr	Val	Ile	Glu	Ser
			305						310				
Thr	Gly	Gly	Phe	Ser	Glu	Glu	Ala	Glu	Ile	Pro	Gly	Ile	Lys
315					320					325			
Tyr	Val	Leu	Ser	Pro	Tyr	Lys	Leu	Asn	Leu	Val	Ala	Thr	Pro
	330					335					340		
Leu	Phe	Leu	Lys	Pro	Gly	Ile	Pro	Tyr	Pro	Ile	Lys	Val	Gln
	345						350					355	
Val	Lys	Asp	Ser	Leu	Asp	Gln	Leu	Val	Gly	Gly	Val	Pro	Val
		360						365					370
Ile	Leu	Asn	Ala	Gln	Thr	Ile	Asp	Val	Asn	Gln	Glu	Thr	Ser
			375						380				
Asp	Leu	Asp	Pro	Ser	Lys	Ser	Val	Thr	Arg	Val	Asp	Asp	Gly
385					390					395			
Val	Ala	Ser	Phe	Val	Leu	Asn	Leu	Pro	Ser	Gly	Val	Thr	Val
	400					405					410		

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 4 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 59.

Please delete lines 29-52, and insert therefor:

Leu Glu Phe Asn Val Lys Thr Asp Ala Pro Asp Leu Pro Glu
415 420 425

Glu Asn Gln Ala Arg Glu Gly Tyr Arg Ala Ile Ala Tyr Ser
430 435 440

Ser Leu Ser Gln Ser Tyr Leu Tyr Ile Asp Trp Thr Asp Asn
445 450

His Lys Ala Leu Leu Val Gly Glu His Leu Asn Ile Ile Val
455 460 465

Thr Pro Lys Ser Pro Tyr Ile Asp Lys Ile Thr His Tyr Asn
470 475 480

Tyr Leu Ile Leu Ser Lys Gly Lys Ile Ile His Phe Gly Thr
485 490 495

Arg Glu Lys Phe Ser Asp Ala Ser Tyr Gln Ser Ile Asn Ile
500 505 510

Pro Val Thr Gln Asn Met Val Pro Ser Ser Arg Leu Leu Val
515 520

Tyr Tyr Ile Val Thr Gly Glu Gln Thr Ala Glu Leu Val Ser
525 530 535

Asp Ser Val Trp Leu Asn Ile Glu Glu Lys Cys Gly Asn Gln
540 545 550

Leu Gln Val His Leu Ser Pro Asp Ala Asp Ala Tyr Ser Pro
555 560 565

Gly Gln Thr Val Ser Leu Asn Met Ala Thr Gly Met Asp Ser
570 575 580

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 5 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 61,

Please delete lines 1-26, and insert therefor:

Trp Val Ala Leu Ala Ala Val Asp Ser Ala Val Tyr Gly Val
585 590

Gln Arg Gly Ala Lys Lys Pro Leu Glu Arg Val Phe Gln Phe
595 600 605

Leu Glu Lys Ser Asp Leu Gly Cys Gly Ala Gly Gly Gly Leu
610 615 620

Asn Asn Ala Asn Val Phe His Leu Ala Gly Leu Thr Phe Leu
625 630 635

Thr Asn Ala Asn Ala Asp Asp Ser Gln Glu Asn Asp Glu Pro
640 645 650

Cys Lys Glu Ile Leu Arg Pro Arg Arg Thr Leu Gln Lys Lys
655 660

Ile Glu Glu Ile Ala Ala Lys Tyr Lys His Ser Val Val Lys
665 670 675

Lys Cys Cys Tyr Asp Gly Ala Cys Val Asn Asn Asp Glu Thr
680 685 690

Cys Glu Gln Arg Ala Ala Arg Ile Ser Leu Gly Pro Arg Cys
695 700 705

Ile Lys Ala Phe Thr Glu Cys Cys Val Val Ala Ser Gln Leu
710 715 720

Arg Ala Asn Ile Ser His Lys Asp Met Gln Leu Gly Arg Leu
725 730

His Met Lys Thr Leu Leu Pro Val Ser Lys Pro Glu Ile Arg
735 740 745

Ser Tyr Phe Pro Glu Ser Trp Leu Trp Glu Val His Leu Val
750 755 760

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 6 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 61,

Please delete lines 27-52, and insert therefor:

```
Pro Arg Arg Lys Gln Leu Gln Phe Ala Leu Pro Asp Ser Leu
      765                      770          775

Thr Thr Trp Glu Ile Gln Gly Ile Gly Ile Ser Asn Thr Gly
      780                      785          790

Ile Cys Val Ala Asp Thr Val Lys Ala Lys Val Phe Lys Asp
      795                      800

Val Phe Leu Glu Met Asn Ile Pro Tyr Ser Val Val Arg Gly
805                      810          815

Glu Gln Ile Gln Leu Lys Gly Thr Val Tyr Asn Tyr Arg Thr
820                      825          830

Ser Gly Met Gln Phe Cys Val Lys Met Ser Ala Val Glu Gly
835                      840          845

Ile Cys Thr Ser Glu Ser Pro Val Ile Asp His Gln Gly Thr
850                      855          860

Lys Ser Ser Lys Cys Val Arg Gln Lys Val Glu Gly Ser Ser
865                      870

Ser His Leu Val Thr Phe Thr Val Leu Pro Leu Glu Ile Gly
875                      880          885

Leu His Asn Ile Asn Phe Ser Leu Glu Thr Trp Phe Gly Lys
890                      895          900

Glu Ile Leu Val Lys Thr Leu Arg Val Val Pro Glu Gly Val
905                      910          915

Lys Arg Glu Ser Tyr Ser Gly Val Thr Leu Asp Pro Arg Gly
920                      925          930

Ile Tyr Gly Thr Ile Ser Arg Arg Lys Glu Phe Pro Tyr Arg
935                      940
```

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 7 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 63,

Please delete lines 1-26, and insert therefor:

```
Ile Pro Leu Asp Leu Val Pro Lys Thr Glu Ile Lys Arg Ile
945                      950                      955

Leu Ser Val Lys Gly Leu Leu Val Gly Glu Ile Leu Ser Ala
960                      965                      970

Val Leu Ser Gln Glu Gly Ile Asn Ile Leu Thr His Leu Pro
975                      980                      985

Lys Gly Ser Ala Glu Ala Glu Leu Met Ser Val Val Pro Val
990                      995                      1000

Phe Tyr Val Phe His Tyr Leu Glu Thr Gly Asn His Trp Asn
1005                      1010

Ile Phe His Ser Asp Pro Leu Ile Glu Lys Gln Lys Leu Lys
1015                      1020                      1025

Lys Lys Leu Lys Glu Gly Met Leu Ser Ile Met Ser Tyr Arg
1030                      1035                      1040

Asn Ala Asp Tyr Ser Tyr Ser Val Trp Lys Gly Gly Ser Ala
1045                      1050                      1055

Ser Thr Trp Leu Thr Ala Phe Ala Leu Arg Val Leu Gly Gln
1060                      1065                      1070

Val Asn Lys Tyr Val Glu Gln Asn Gln Asn Ser Ile Cys Asn
1075                      1080

Ser Leu Leu Trp Leu Val Glu Asn Tyr Gln Leu Asp Asn Gly
1085                      1090                      1095

Ser Phe Lys Glu Asn Ser Gln Tyr Gln Pro Ile Lys Leu Gln
1100                      1105                      1110

Gly Thr Leu Pro Val Glu Ala Arg Glu Asn Ser Leu Tyr Leu
1115                      1120                      1125
```

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 8 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 63,

Please delete lines 27-52, and insert therefor:

Thr Ala Phe Thr Val Ile Gly Ile Arg Lys Ala Phe Asp Ile
1130 1135 1140

Cys Pro Leu Val Lys Ile Asp Thr Ala Leu Ile Lys Ala Asp
1145 1150

Asn Phe Leu Leu Glu Asn Thr Leu Pro Ala Gln Ser Thr Phe
1155 1160 1165

Thr Leu Ala Ile Ser Ala Tyr Ala Leu Ser Leu Gly Asp Lys
1170 1175 1180

Thr His Pro Gln Phe Arg Ser Ile Val Ser Ala Leu Lys Arg
1185 1190 1195

Glu Ala Leu Val Lys Gly Asn Pro Pro Ile Tyr Arg Phe Trp
1200 1205 1210

Lys Asp Asn Leu Gln His Lys Asp Ser Ser Val Pro Asn Thr
1215 1220

Gly Thr Ala Arg Met Val Glu Thr Thr Ala Tyr Ala Leu Leu
1225 1230 1235

Thr Ser Leu Asn Leu Lys Asp Ile Asn Tyr Val Asn Pro Val
1240 1245 1250

Ile Lys Trp Leu Ser Glu Glu Gln Arg Tyr Gly Gly Gly Phe
1255 1260 1265

Tyr Ser Thr Gln Asp Thr Ile Asn Ala Ile Glu Gly Leu Thr
1270 1275 1280

Glu Tyr Ser Leu Leu Val Lys Gln Leu Arg Leu Ser Met Asp
1285 1290

Ile Asp Val Ser Tyr Lys His Lys Gly Ala Leu His Asn Tyr
1295 1300 1305

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 9 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 65,

Please delete lines 1-26, and insert therefor:

Lys Met Thr Asp Lys Asn Phe Leu Gly Arg Pro Val Glu Val
1310 1315 1320

Leu Leu Asn Asp Asp Leu Ile Val Ser Thr Gly Phe Gly Ser
1325 1330 1335

Gly Leu Ala Thr Val His Val Thr Thr Val Val His Lys Thr
1340 1345 1350

Ser Thr Ser Glu Glu Val Cys Ser Phe Tyr Leu Lys Ile Asp
1355 1360

Thr Gln Asp Ile Glu Ala Ser His Tyr Arg Gly Tyr Gly Asn
1365 1370 1375

Ser Asp Tyr Lys Arg Ile Val Ala Cys Ala Ser Tyr Lys Pro
1380 1385 1390

Ser Arg Glu Glu Ser Ser Ser Gly Ser Ser His Ala Val Met
1395 1400 1405

Asp Ile Ser Leu Pro Thr Gly Ile Ser Ala Asn Glu Glu Asp
1410 1415 1420

Leu Lys Ala Leu Val Glu Gly Val Asp Gln Leu Phe Thr Asp
1425 1430

Tyr Gln Ile Lys Asp Gly His Val Ile Leu Gln Leu Asn Ser
1435 1440 1445

Ile Pro Ser Ser Asp Phe Leu Cys Val Arg Phe Arg Ile Phe
1450 1455 1460

Glu Leu Phe Glu Val Gly Phe Leu Ser Pro Ala Thr Phe Thr
1465 1470 1475

Val Tyr Glu Tyr His Arg Pro Asp Lys Gln Cys Thr Met Phe
1480 1485 1490

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 10 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 65,

Please delete lines 27-50, and insert therefor:

Tyr Ser Thr Ser Asn Ile Lys Ile Gln Lys Val Cys Glu Gly
1495 1500

Ala Ala Cys Lys Cys Val Glu Ala Asp Cys Gly Gln Met Gln
1505 1510 1515

Glu Glu Leu Asp Leu Thr Ile Ser Ala Glu Thr Arg Lys Gln
1520 1525 1530

Thr Ala Cys Lys Pro Glu Ile Ala Tyr Ala Tyr Lys Val Ser
1535 1540 1545

Ile Thr Ser Ile Thr Val Glu Asn Val Phe Val Lys Tyr Lys
1550 1555 1560

Ala Thr Leu Leu Asp Ile Tyr Lys Thr Gly Glu Ala Val Ala
1565 1570

Glu Lys Asp Ser Glu Ile Thr Phe Ile Lys Lys Val Thr Cys
1575 1580 1585

Thr Asn Ala Glu Leu Val Lys Gly Arg Gln Tyr Leu Ile Met
1590 1595 1600

Gly Lys Glu Ala Leu Gln Ile Lys Tyr Asn Phe Ser Phe Arg
1605 1610 1615

Tyr Ile Tyr Pro Leu Asp Ser Leu Thr Trp Ile Glu Tyr Trp
1620 1625 1630

Pro Arg Asp Thr Thr Cys Ser Ser Cys Gln Ala Phe Leu Ala
1635 1640

Asn Leu Asp Glu Phe Ala Glu Asp Ile Phe Leu Asn Gly Cys
1645 1650 1655

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 11 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 81,

Please delete lines 18-47, and insert therefor:

ATG GCC GAC ATC CAG ATG ACT CAG TCT CCA	30
Met Ala Asp Ile Gln Met Thr Gln Ser Pro	
1 5 10	
GCT TCA CTG TCT GCA TCT GTG GGA GAA ACT	60
Ala Ser Leu Ser Ala Ser Val Gly Glu Thr	
15 20	
GTC ACC ATC ACA TGT GGA GCA AGT GAG AAT	90
Val Thr Ile Thr Cys Gly Ala Ser Glu Asn	
25 30	
ATT TAC GGT GCT TTA AAT TGG TAT CAG CGG	120
Ile Tyr Gly Ala Leu Asn Trp Tyr Gln Arg	
35 40	
AAA CAG GGA AAA TCT CCT CAG CTC CTG ATC	150
Lys Gln Gly Lys Ser Pro Gln Leu Leu Ile	
45 50	
TAT GGT GCA ACC AAC TTG GCA GAT GGC ATG	180
Tyr Gly Ala Thr Asn Leu Ala Asp Gly Met	
55 60	
TCA TCG AGG TTC AGT GGC AGT GGA TCT GGT	210
Ser Ser Arg Phe Ser Gly Ser Gly Ser Gly	
65 70	
AGA CAG TAT TAT CTC AAG ATC AGT AGC CTG	240
Arg Gln Tyr Tyr Leu Lys Ile Ser Ser Leu	
75 80	
CAT CCT GAC GAT GTT GCA ACG TAT TAC TGT	270
His Pro Asp Asp Val Ala Thr Tyr Tyr Cys	
85 90	
CAA AAT GTG TTA AAT ACT CCT CTC ACG TTC	300
Gln Asn Val Leu Asn Thr Pro Leu Thr Phe	
95 100	

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 12 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 81,

Please delete line 48, through column 83, line 21, and insert therefor:

GGT GCT GGG ACC AAG TTG GAG CTG AAA CGG	330
Gly Ala Gly Thr Lys Leu Glu Leu Lys Arg	
105 110	
ACC GGA GGT GGC GGG TCG GGT GGC GGG GGA	360
Thr Gly Gly Gly Gly Ser Gly Gly Gly Gly	
115 120	
TCG GGT GGC GGA GGG TCG CAG GTT CAG CTG	390
Ser Gly Gly Gly Gly Ser Gln Val Gln Leu	
125 130	
CAG CAG TCT GGA GCC GAG CTG ATG AAG CCT	420
Gln Gln Ser Gly Ala Glu Leu Met Lys Pro	
135 140	
GGG GCC TCA GTG AAG ATG TCC TGC AAG GCT	450
Gly Ala Ser Val Lys Met Ser Cys Lys Ala	
145 150	
ACT GGC TAC ATA TTC AGT AAC TAC TGG ATA	480
Thr Gly Tyr Ile Phe Ser Asn Tyr Trp Ile	
155 160	
CAG TGG ATA AAG CAG AGG CCT GGA CAT GGC	510
Gln Trp Ile Lys Gln Arg Pro Gly His Gly	
165 170	
CTT GAG TGG ATT GGT GAG ATT TTA CCT GGA	540
Leu Glu Trp Ile Gly Glu Ile Leu Pro Gly	
175 180	
AGT GGT TCT ACT GAG TAC ACT GAG AAC TTC	570
Ser Gly Ser Thr Glu Tyr Thr Glu Asn Phe	
185 190	
AAG GAC AAG GCC GCA TTC ACT GCA GAT ACA	600
Lys Asp Lys Ala Ala Phe Thr Ala Asp Thr	
195 200	

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 13 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 83,

Please delete lines 22-36, and insert therefor:

TCC TCC AAC ACA GCC TAC ATG CAA CTC AGC	630
Ser Ser Asn Thr Ala Tyr Met Gln Leu Ser	
205 210	
AGC CTG ACA TCA GAG GAC TCT GCC GTC TAT	660
Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr	
215 220	
TAC TGT GCA AGA TAT TTC TTC GGT AGT AGC	690
Tyr Cys Ala Arg Tyr Phe Phe Gly Ser Ser	
225 230	
CCC AAC TGG TAC TTC GAT GTC TGG GGC GCA	720
Pro Asn Trp Tyr Phe Asp Val Trp Gly Ala	
235 240	
GGG ACC ACG GTC ACC GTC TCC TCA TGA	747
Gly Thr Thr Val Thr Val Ser Ser	
245	

Please delete lines 46-57, and insert therefor:

ATG GCC GAT ATC CAG ATG ACC CAG TCC CCG	30
Met Ala Asp Ile Gln Met Thr Gln Ser Pro	
1 5 10	
TCC TCC CTG TCC GCC TCT GTG GGC GAT AGG	60
Ser Ser Leu Ser Ala Ser Val Gly Asp Arg	
15 20	
GTC ACC ATC ACC TGC GGC GCC AGC GAA AAC	90
Val Thr Ile Thr Cys Gly Ala Ser Glu Asn	
25 30	
ATC TAT GGC GCG CTG AAC TGG TAT CAA CGT	120
Ile Tyr Gly Ala Leu Asn Trp Tyr Gln Arg	
35 40	

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 14 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 85,

Please delete lines 1-30, and insert therefor:

AAA CCT GGG AAA GCT CCG AAG CTT CTG ATT	150
Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile	
45 50	
TAC GGT GCG ACG AAC CTG GCA GAT GGA GTC	180
Tyr Gly Ala Thr Asn Leu Ala Asp Gly Val	
55 60	
CCT TCT CGC TTC TCT GGA TCC GGC TCC GGA	210
Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly	
65 70	
ACG GAT TTC ACT CTG ACC ATC AGC AGT CTG	240
Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu	
75 80	
CAG CCT GAA GAC TTC GCT ACG TAT TAC TGT	270
Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys	
85 90	
CAG AAC GTT TTA AAT ACT CCG TTG ACT TTC	300
Gln Asn Val Leu Asn Thr Pro Leu Thr Phe	
95 100	
GGA CAG GGT ACC AAG GTG GAA ATA AAA CGT	330
Gly Gln Gly Thr Lys Val Glu Ile Lys Arg	
105 110	
ACT GGC GGT GGT GGT TCT GGT GGC GGT GGA	360
Thr Gly Gly Gly Gly Ser Gly Gly Gly Gly	
115 120	
TCT GGT GGT GGC GGT TCT CAA GTC CAA CTG	390
Ser Gly Gly Gly Gly Ser Gln Val Gln Leu	
125 130	
GTG CAA TCC GGC GCC GAG GTC AAG AAG CCA	420
Val Gln Ser Gly Ala Glu Val Lys Lys Pro	
135 140	

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 15 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 85,

Please delete lines 31-60, and insert therefor:

GGG GCC TCA GTC AAA GTG TCC TGT AAA GCT	450
Gly Ala Ser Val Lys Val Ser Cys Lys Ala	
145 150	
AGC GGC TAT ATT TTT TCT AAT TAT TGG ATT	480
Ser Gly Tyr Ile Phe Ser Asn Tyr Trp Ile	
155 160	
CAA TGG GTG CGT CAG GCC CCC GGG CAG GGC	510
Gln Trp Val Arg Gln Ala Pro Gly Gln Gly	
165 170	
CTG GAA TGG ATG GGT GAG ATC TTA CCG GGC	540
Leu Glu Trp Met Gly Glu Ile Leu Pro Gly	
175 180	
TCT GGT AGC ACC GAA TAT ACC GAA AAT TTT	570
Ser Gly Ser Thr Glu Tyr Thr Glu Asn Phe	
185 190	
AAA GAC CGT GTT ACT ATG ACG CGT GAC ACT	600
Lys Asp Arg Val Thr Met Thr Arg Asp Thr	
195 200	
TCG ACT AGT ACA GTA TAC ATG GAG CTC TCC	630
Ser Thr Ser Thr Val Tyr Met Glu Leu Ser	
205 210	
AGC CTG CGA TCG GAG GAC ACG GCC GTC TAT	660
Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr	
215 220	
TAT TGC GCG CGT TAT TTT TTT GGT TCT AGC	690
Tyr Cys Ala Arg Tyr Phe Phe Gly Ser Ser	
225 230	
CCG AAT TGG TAT TTT GAT GTT TGG GGT CAA	720
Pro Asn Trp Tyr Phe Asp Val Trp Gly Gln	
235 240	

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 16 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 87,

Please delete lines 1-3, and insert therefor:

GGA ACC CTG GTC ACT GTC TCG AGC TGA	747
Gly Thr Leu Val Thr Val Ser Ser	
245	

Please delete lines 13-36, and insert therefor:

ATG GGA ATC CAA GGA GGG TCT GTC CTG TTC	30
Met Gly Ile Gln Gly Gly Ser Val Leu Phe	
-25 -20	

GGG CTG CTG CTC GTC CTG GCT GTC TTC TGC	60
Gly Leu Leu Leu Val Leu Ala Val Phe Cys	
-15 -10	

CAT TCA GGT CAT AGC CTG CAG GAC ATC CAG	90
His Ser Gly His Ser Leu Gln Asp Ile Gln	
-5 1 5	

ATG ACT CAG TCT CCA GCT TCA CTG TCT GCA	120
Met Thr Gln Ser Pro Ala Ser Leu Ser Ala	
10 15	

TCT GTG GGA GAA ACT GTC ACC ATC ACA TGT	150
Ser Val Gly Glu Thr Val Thr Ile Thr Cys	
20 25	

GGA GCA AGT GAG AAT ATT TAC GGT GCT TTA	180
Gly Ala Ser Glu Asn Ile Tyr Gly Ala Leu	
30 35	

AAT TGG TAT CAG CGG AAA CAG GGA AAA TCT	210
Asn Trp Tyr Gln Arg Lys Gln Gly Lys Ser	
40 45	

CCT CAG CTC CTG ATC TAT GGT GCA ACC AAC	240
Pro Gln Leu Leu Ile Tyr Gly Ala Thr Asn	
50 55	

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 17 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 87,

Please delete line 37, through column 89, line 9, and insert therefor:

TTG GCA GAT GGC ATG TCA TCG AGG TTC AGT	270
Leu Ala Asp Gly Met Ser Ser Arg Phe Ser	
60 65	
GGC AGT GGA TCT GGT AGA CAG TAT TAT CTC	300
Gly Ser Gly Ser Gly Arg Gln Tyr Tyr Leu	
70 75	
AAG ATC AGT AGC CTG CAT CCT GAC GAT GTT	330
Lys Ile Ser Ser Leu His Pro Asp Asp Val	
80 85	
GCA ACG TAT TAC TGT CAA AAT GTG TTA AAT	360
Ala Thr Tyr Tyr Cys Gln Asn Val Leu Asn	
90 95	
ACT CCT CTC ACG TTC GGT GCT GGG ACC AAG	390
Thr Pro Leu Thr Phe Gly Ala Gly Thr Lys	
100 105	
TTG GAG CTG AAA CGA ACT GTG GCT GCA CCA	420
Leu Glu Leu Lys Arg Thr Val Ala Ala Pro	
110 115	
TCT GTC TTC ATC TTC CCG CCA TCT GAT GAG	450
Ser Val Phe Ile Phe Pro Pro Ser Asp Glu	
120 125	
CAG TTG AAA TCT GGA ACT GCC TCT GTT GTG	480
Gln Leu Lys Ser Gly Thr Ala Ser Val Val	
130 135	
TGC CTG CTG AAT AAC TTC TAT CCC AGA GAG	510
Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu	
140 145	
GCC AAA GTA CAG TGG AAG GTG GAT AAC GCC	540
Ala Lys Val Gln Trp Lys Val Asp Asn Ala	
150 155	

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 18 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 89.

Please delete lines 10-29, and insert therefor:

CTC CAA TCG GGT AAC TCC CAG GAG AGT GTC	570
Leu Gln Ser Gly Asn Ser Gln Glu Ser Val	
160 165	
ACA GAG CAG GAC AGC AAG GAC AGC ACC TAC	600
Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr	
170 175	
AGC CTC AGC AGC ACC CTG ACG CTG AGC AAA	630
Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys	
180 185	
GCA GAC TAC GAG AAA CAC AAA GTC TAC GCC	660
Ala Asp Tyr Glu Lys His Lys Val Tyr Ala	
190 195	
TGC GAA GTC ACC CAT CAG GGC CTG AGC TCG	690
Cys Glu Val Thr His Gln Gly Leu Ser Ser	
200 205	
CCC GTC ACA AAG AGC TTC AAC AGG GGA GAG	720
Pro Val Thr Lys Ser Phe Asn Arg Gly Glu	
210 215	
TGT TAG	726
Cys	

Please delete lines 39-44, and insert therefor:

ATG AAA TGG AGC TGG GTT ATT CTC TTC CTC	30
Met Lys Trp Ser Trp Val Ile Leu Phe Leu	
-15 -10	
CTG TCA GTA ACT GCA GGT GTC CAC TCC CAG	60
Leu Ser Val Thr Ala Gly Val His Ser Gln	
-5 1	

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 19 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 89.

Please delete line 45, through column 91, line 15, and insert therefor:

GTT CAG CTG CAG CAG TCT GGA GCT GAG CTG	90
Val Gln Leu Gln Gln Ser Gly Ala Glu Leu	
5 10	
ATG AAG CCT GGG GCC TCA GTG AAG ATG TCC	120
Met Lys Pro Gly Ala Ser Val Lys Met Ser	
15 20	
TGC AAG GCT ACT GGC TAC ATA TTC AGT AAC	150
Cys Lys Ala Thr Gly Tyr Ile Phe Ser Asn	
25 30	
TAC TGG ATA CAG TGG ATA AAG CAG AGG CCT	180
Tyr Trp Ile Gln Trp Ile Lys Gln Arg Pro	
35 40	
GGA CAT GGC CTT GAG TGG ATT GGT GAG ATT	210
Gly His Gly Leu Glu Trp Ile Gly Glu Ile	
45 50	
TTA CCT GGA AGT GGT TCT ACT GAG TAC ACT	240
Leu Pro Gly Ser Gly Ser Thr Glu Tyr Thr	
55 60	
GAG AAC TTC AAG GAC AAG GCC GCA TTC ACT	270
Glu Asn Phe Lys Asp Lys Ala Ala Phe Thr	
65 70	
GCA GAT ACA TCC TCC AAC ACA GCC TAC ATG	300
Ala Asp Thr Ser Ser Asn Thr Ala Tyr Met	
75 80	
CAA CTC AGC AGC CTG ACA TCA GAG GAC TCT	330
Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser	
85 90	
GCC GTC TAT TAC TGT GCA AGA TAT TTC TTC	360
Ala Val Tyr Tyr Cys Ala Arg Tyr Phe Phe	
95 100	

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 20 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 91,

Please delete lines 16-45, and insert therefor:

GGT AGT AGC CCC AAC TGG TAC TTC GAT GTC	390
Gly Ser Ser Pro Asn Trp Tyr Phe Asp Val	
105 110	
TGG GGC GCA GGG ACC ACG GTC ACC GTC TCC	420
Trp Gly Ala Gly Thr Thr Val Thr Val Ser	
115 120	
TCA GCC TCC ACC AAG GGC CCA TCG GTC TTC	450
Ser Ala Ser Thr Lys Gly Pro Ser Val Phe	
125 130	
CCC CTG GCG CCC TCC TCC AAG AGC ACC TCT	480
Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser	
135 140	
GGG GGC ACA GCG GCC CTG GGC TGC CTG GTC	510
Gly Gly Thr Ala Ala Leu Gly Cys Leu Val	
145 150	
AAG GAC TAC TTC CCC GAA CCG GTG ACG GTG	540
Lys Asp Tyr Phe Pro Glu Pro Val Thr Val	
155 160	
TCG TGG AAC TCA GGC GCC CTG ACC AGC GGC	570
Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly	
165 170	
GTG CAC ACC TTC CCG GCT GTC CTA CAG TCC	600
Val His Thr Phe Pro Ala Val Leu Gln Ser	
175 180	
TCA GGA CTC TAC TCC CTC AGC AGC GTG GTG	630
Ser Gly Leu Tyr Ser Leu Ser Ser Val Val	
185 190	
ACC GTG CCC TCC AGC AGC TTG GGC ACC CAG	660
Thr Val Pro Ser Ser Ser Leu Gly Thr Gln	
195 200	

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 21 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 91,

Please delete lines 46-54, and insert therefor:

ACC TAC ATC TGC AAC GTG AAT CAC AAG CCC	690
Thr Tyr Ile Cys Asn Val Asn His Lys Pro	
205 210	
AGC AAC ACC AAG GTG GAC AAG AAA GTT GAG	720
Ser Asn Thr Lys Val Asp Lys Lys Val Glu	
215 220	
CCC AAA TCT TGT GAC AAA ACT CAC ACA TAA	750
Pro Lys Ser Cys Asp Lys Thr His Thr	
225	

Column 93

Please delete lines 4-21, and insert therefor:

ATG AAG TGG AGC TGG GTT ATT CTC TTC CTC	30
Met Lys Trp Ser Trp Val Ile Leu Phe Leu	
-15 -10	
CTG TCA GTA ACT GCC GGC GTC CAC TCC CAA	60
Leu Ser Val Thr Ala Gly Val His Ser Gln	
-5 1	
GTC CAA CTG GTG CAA TCC GGC GCC GAG GTC	90
Val Gln Leu Val Gln Ser Gly Ala Glu Val	
5 10	
AAG AAG CCA GGG GCC TCA GTC AAA GTG TCC	120
Lys Lys Pro Gly Ala Ser Val Lys Val Ser	
15 20	
TGT AAA GCT AGC GGC TAT ATT TTT TCT AAT	150
Cys Lys Ala Ser Gly Tyr Ile Phe Ser Asn	
25 30	
TAT TGG ATT CAA TGG GTG CGT CAG GCC CCC	180
Tyr Trp Ile Gln Trp Val Arg Gln Ala Pro	
35 40	

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 22 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 93,

Please delete lines 22-51, and insert therefor:

GGG CAG GGC CTG GAA TGG ATG GGT GAG ATC Gly Gln Gly Leu Glu Trp Met Gly Glu Ile 45 50	210
TTA CCG GGC TCT GGT AGC ACC GAA TAT GCC Leu Pro Gly Ser Gly Ser Thr Glu Tyr Ala 55 60	240
CAA AAA TTC CAG GGC CGT GTT ACT ATG ACT Gln Lys Phe Gln Gly Arg Val Thr Met Thr 65 70	270
GCG GAC ACT TCG ACT AGT ACA GCC TAC ATG Ala Asp Thr Ser Thr Ser Thr Ala Tyr Met 75 80	300
GAG CTC TCC AGC CTG CGA TCG GAG GAC ACG Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr 85 90	330
GCC GTC TAT TAT TGC GCG CGT TAT TTT TTT Ala Val Tyr Tyr Cys Ala Arg Tyr Phe Phe 95 100	360
GGT TCT AGC CCG AAT TGG TAT TTT GAT GTT Gly Ser Ser Pro Asn Trp Tyr Phe Asp Val 105 110	390
TGG GGT CAA GGA ACC CTG GTC ACT GTC TCG Trp Gly Gln Gly Thr Leu Val Thr Val Ser 115 120	420
AGC GCC TCC ACC AAG GGC CCA TCG GTC TTC Ser Ala Ser Thr Lys Gly Pro Ser Val Phe 125 130	450
CCC CTG GCG CCC TCC TCC AAG AGC ACC TCT Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser 135 140	480

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 23 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 93,

Please delete line 52, through column 95, line 27, and insert therefor:

GGG GGC ACA GCG GCC CTG GGC TGC CTG GTC	510
Gly Gly Thr Ala Ala Leu Gly Cys Leu Val	
145 150	
AAG GAC TAC TTC CCC GAA CCG GTG ACG GTG	540
Lys Asp Tyr Phe Pro Glu Pro Val Thr Val	
155 160	
TCG TGG AAC TCA GGC GCC CTG ACC AGC GGC	570
Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly	
165 170	
GTG CAC ACC TTC CCG GCT GTC CTA CAG TCC	600
Val His Thr Phe Pro Ala Val Leu Gln Ser	
175 180	
TCA GGA CTC TAC TCC CTC AGC AGC GTG GTG	630
Ser Gly Leu Tyr Ser Leu Ser Ser Val Val	
185 190	
ACC GTG CCC TCC AGC AGC TTG GGC ACC CAG	660
Thr Val Pro Ser Ser Ser Leu Gly Thr Gln	
195 200	
ACC TAC ATC TGC AAC GTG AAT CAC AAG CCC	690
Thr Tyr Ile Cys Asn Val Asn His Lys Pro	
205 210	
AGC AAC ACC AAG GTG GAC AAG AAA GTT GAG	720
Ser Asn Thr Lys Val Asp Lys Lys Val Glu	
215 220	
CCC AAA TCT TGT GAC AAA ACT CAC ACA TAA	750
Pro Lys Ser Cys Asp Lys Thr His Thr	
225 230	

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 24 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 95,

Please delete line 31, through column 97, line 3, and insert therefor:

ATG AAG TGG AGC TGG GTT ATT CTC TTC CTC	30
Met Lys Trp Ser Trp Val Ile Leu Phe Leu	
-15 -10	
CTG TCA GTA ACT GCC GGC GTC CAC TCC CAA	60
Leu Ser Val Thr Ala Gly Val His Ser Gln	
-5 1	
GTC CAA CTG GTG CAA TCC GGC GCC GAG GTC	90
Val Gln Leu Val Gln Ser Gly Ala Glu Val	
5 10	
AAG AAG CCA GGG GCC TCA GTC AAA GTG TCC	120
Lys Lys Pro Gly Ala Ser Val Lys Val Ser	
15 20	
TGT AAA GCT AGC GGC TAT ATT TTT TCT AAT	150
Cys Lys Ala Ser Gly Tyr Ile Phe Ser Asn	
25 30	
TAT TGG ATT CAA TGG GTG CGT CAG GCC CCC	180
Tyr Trp Ile Gln Trp Val Arg Gln Ala Pro	
35 40	
GGG CAG GGC CTG GAA TGG ATG GGT GAG ATC	210
Gly Gln Gly Leu Glu Trp Met Gly Glu Ile	
45 50	
TTA CCG GGC TCT GGT AGC ACC GAA TAT ACC	240
Leu Pro Gly Ser Gly Ser Thr Glu Tyr Thr	
55 60	
GAA AAT TTT AAA GAC CGT GTT ACT ATG ACG	270
Glu Asn Phe Lys Asp Arg Val Thr Met Thr	
65 70	
CGT GAC ACT TCG ACT AGT ACA GTA TAC ATG	300
Arg Asp Thr Ser Thr Ser Thr Val Tyr Met	
75 80	

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 25 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 97,

Please delete lines 4-33, and insert therefor:

GAG CTC TCC AGC CTG CGA TCG GAG GAC ACG Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr 85 90	330
GCC GTC TAT TAT TGC GCG CGT TAT TTT TTT Ala Val Tyr Tyr Cys Ala Arg Tyr Phe Phe 95 100	360
GGT TCT AGC CCG AAT TGG TAT TTT GAT GTT Gly Ser Ser Pro Asn Trp Tyr Phe Asp Val 105 110	390
TGG GGT CAA GGA ACC CTG GTC ACT GTC TCG Trp Gly Gln Gly Thr Leu Val Thr Val Ser 115 120	420
AGC GCC TCC ACC AAG GGC CCA TCG GTC TTC Ser Ala Ser Thr Lys Gly Pro Ser Val Phe 125 130	450
CCC CTG GCG CCC TCC TCC AAG AGC ACC TCT Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser 135 140	480
GGG GGC ACA GCG GCC CTG GGC TGC CTG GTC Gly Gly Thr Ala Ala Leu Gly Cys Leu Val 145 150	510
AAG GAC TAC TTC CCC GAA CCG GTG ACG GTG Lys Asp Tyr Phe Pro Glu Pro Val Thr Val 155 160	540
TCG TGG AAC TCA GGC GCC CTG ACC AGC GGC Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly 165 170	570
GTG CAC ACC TTC CCG GCT GTC CTA CAG TCC Val His Thr Phe Pro Ala Val Leu Gln Ser 175 180	600

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 26 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 97,

Please delete lines 34-48, and insert therefor:

TCA GGA CTC TAC TCC CTC AGC AGC GTG GTG	630
Ser Gly Leu Tyr Ser Leu Ser Ser Val Val	
185 190	
ACC GTG CCC TCC AGC AGC TTG GGC ACC CAG	660
Thr Val Pro Ser Ser Ser Leu Gly Thr Gln	
195 200	
ACC TAC ATC TGC AAC GTG AAT CAC AAG CCC	690
Thr Tyr Ile Cys Asn Val Asn His Lys Pro	
205 210	
AGC AAC ACC AAG GTG GAC AAG AAA GTT GAG	720
Ser Asn Thr Lys Val Asp Lys Lys Val Glu	
215 220	
CCC AAA TCT TGT GAC AAA ACT CAC ACA TAA	750
Pro Lys Ser Cys Asp Lys Thr His Thr	
225 230	

Column 99,

Please delete lines 2-13, and insert therefor:

ATG GGA ATC CAA GGA GGG TCT GTC CTG TTC	30
Met Gly Ile Gln Gly Gly Ser Val Leu Phe	
-25 -20	
GGG CTG CTG CTC GTC CTG GCT GTC TTC TGC	60
Gly Leu Leu Leu Val Leu Ala Val Phe Cys	
-15 -10	
CAT TCA GGT CAT AGC CTG CAG GAT ATC CAG	90
His Ser Gly His Ser Leu Gln Asp Ile Gln	
-5 1 5	
ATG ACC CAG TCC CCG TCC TCC CTG TCC GCC	120
Met Thr Gln Ser Pro Ser Ser Leu Ser Ala	
10 15	

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 27 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 99,

Please delete lines 14-43, and insert therefor:

TCT GTG GGC GAT AGG GTC ACC ATC ACC TGC	150
Ser Val Gly Asp Arg Val Thr Ile Thr Cys	
20 25	
GGC GCC AGC GAA AAC ATC TAT GGC GCG CTG	180
Gly Ala Ser Glu Asn Ile Tyr Gly Ala Leu	
30 35	
AAC TGG TAT CAA CGT AAA CCT GGC AAA GCT	210
Asn Trp Tyr Gln Arg Lys Pro Gly Lys Ala	
40 45	
CCG AAG CTT CTG ATT TAC GGT GCG ACG AAC	240
Pro Lys Leu Leu Ile Tyr Gly Ala Thr Asn	
50 55	
CTG GCA GAT GGA GTC CCT TCT CGC TTC TCT	270
Leu Ala Asp Gly Val Pro Ser Arg Phe Ser	
60 65	
GGA TCC GGC TCC GGA ACG GAT TAC ACT CTG	300
Gly Ser Gly Ser Gly Thr Asp Tyr Thr Leu	
70 75	
ACC ATC AGC AGT CTG CAA CCT GAG GAC TTC	330
Thr Ile Ser Ser Leu Gln Pro Glu Asp Phe	
80 85	
GCT ACG TAT TAC TGT CAG AAC GTT TTA AAT	360
Ala Thr Tyr Tyr Cys Gln Asn Val Leu Asn	
90 95	
ACT CCG TTG ACT TTC GGA CAG GGT ACC AAG	390
Thr Pro Leu Thr Phe Gly Gln Gly Thr Lys	
100 105	
GTG GAA ATA AAA CGA ACT GTG GCT GCA CCA	420
Val Glu Ile Lys Arg Thr Val Ala Ala Pro	
110 115	

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 28 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 99,

Please delete line 44, through column 101, line 17, and insert therefor:

TCT GTC TTC ATC TTC CCG CCA TCT GAT GAG	450
Ser Val Phe Ile Phe Pro Pro Ser Asp Glu	
120 125	
CAG TTG AAA TCT GGA ACT GCC TCT GTT GTG	480
Gln Leu Lys Ser Gly Thr Ala Ser Val Val	
130 135	
TGC CTG CTG AAT AAC TTC TAT CCC AGA GAG	510
Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu	
140 145	
GCC AAA GTA CAG TGG AAG GTG GAT AAC GCC	540
Ala Lys Val Gln Trp Lys Val Asp Asn Ala	
150 155	
CTC CAA TCG GGT AAC TCC CAG GAG AGT GTC	570
Leu Gln Ser Gly Asn Ser Gln Glu Ser Val	
160 165	
ACA GAG CAG GAC AGC AAG GAC AGC ACC TAC	600
Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr	
170 175	
AGC CTC AGC AGC ACC CTG ACG CTG AGC AAA	630
Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys	
180 185	
GCA GAC TAC GAG AAA CAC AAA GTC TAC GCC	660
Ala Asp Tyr Glu Lys His Lys Val Tyr Ala	
190 195	
TGC GAA GTC ACC CAT CAG GGC CTG AGC TCG	690
Cys Glu Val Thr His Gln Gly Leu Ser Ser	
200 205	
CCC GTC ACA AAG AGC TTC AAC AGG GGA GAG	720
Pro Val Thr Lys Ser Phe Asn Arg Gly Glu	
210 215	
TGT TAG	726
Cys	

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 29 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 101,

Please delete lines 28-57, and insert therefor:

ATG GGA ATC CAA GGA GGG TCT GTC CTG TTC Met Gly Ile Gln Gly Gly Ser Val Leu Phe -25 -20	30
GGG CTG CTG CTC GTC CTG GCT GTC TTC TGC Gly Leu Leu Leu Val Leu Ala Val Phe Cys -15 -10	60
CAT TCA GGT CAT AGC CTG CAG GAT ATC CAG His Ser Gly His Ser Leu Gln Asp Ile Gln -5 1 5	90
ATG ACC CAG TCC CCG TCC TCC CTG TCC GCC Met Thr Gln Ser Pro Ser Ser Leu Ser Ala 10 15	120
TCT GTG GGC GAT AGG GTC ACC ATC ACC TGC Ser Val Gly Asp Arg Val Thr Ile Thr Cys 20 25	150
GGC GCC AGC GAA AAC ATC TAT GGC GCG CTG Gly Ala Ser Glu Asn Ile Tyr Gly Ala Leu 30 35	180
AAC TGG TAT CAA CGT AAA CCT GGG AAA GCT Asn Trp Tyr Gln Arg Lys Pro Gly Lys Ala 40 45	210
CCG AAG CTT CTG ATT TAC GGT GCG ACG AAC Pro Lys Leu Leu Ile Tyr Gly Ala Thr Asn 50 55	240
CTG GCA GAT GGA GTC CCT TCT CGC TTC TCT Leu Ala Asp Gly Val Pro Ser Arg Phe Ser 60 65	270
GGA TCC GGC TCC GGA ACG GAT TTC ACT CTG Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu 70 75	300

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 30 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 103,

Please delete lines 1-30, and insert therefor:

ACC ATC AGC AGT CTG CAG CCT GAA GAC TTC	330
Thr Ile Ser Ser Leu Gln Pro Glu Asp Phe	
80 85	
GCT ACG TAT TAC TGT CAG AAC GTT TTA AAT	360
Ala Thr Tyr Tyr Cys Gln Asn Val Leu Asn	
90 95	
ACT CCG TTG ACT TTC GGA CAG GGT ACC AAG	390
Thr Pro Leu Thr Phe Gly Gln Gly Thr Lys	
100 105	
GTG GAA ATA AAA CGA ACT GTG GCT GCA CCA	420
Val Glu Ile Lys Arg Thr Val Ala Ala Pro	
110 115	
TCT GTC TTC ATC TTC CCG CCA TCT GAT GAG	450
Ser Val Phe Ile Phe Pro Pro Ser Asp Glu	
120 125	
CAG TTG AAA TCT GGA ACT GCC TCT GTT GTG	480
Gln Leu Lys Ser Gly Thr Ala Ser Val Val	
130 135	
TGC CTG CTG AAT AAC TTC TAT CCC AGA GAG	510
Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu	
140 145	
GCC AAA GTA CAG TGG AAG GTG GAT AAC GCC	540
Ala Lys Val Gln Trp Lys Val Asp Asn Ala	
150 155	
CTC CAA TCG GGT AAC TCC CAG GAG AGT GTC	570
Leu Gln Ser Gly Asn Ser Gln Glu Ser Val	
160 165	
ACA GAG CAG GAC AGC AAG GAC AGC ACC TAC	600
Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr	
170 175	

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 31 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 103,

Please delete lines 31-44, and insert therefor:

AGC CTC AGC AGC ACC CTG ACG CTG AGC AAA	630
Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys	
180 185	
GCA GAC TAC GAG AAA CAC AAA GTC TAC GCC	660
Ala Asp Tyr Glu Lys His Lys Val Tyr Ala	
190 195	
TGC GAA GTC ACC CAT CAG GGC CTG AGC TCG	690
Cys Glu Val Thr His Gln Gly Leu Ser Ser	
200 205	
CCC GTC ACA AAG AGC TTC AAC AGG GGA GAG	720
Pro Val Thr Lys Ser Phe Asn Arg Gly Glu	
210 215	
TGT TAG	726
Cys	

Please delete line 55, through column 105, line 9, and insert therefor:

ATG GAC ATG AGG GTC CCC GCT CAG CTC CTG	30
Met Asp Met Arg Val Pro Ala Gln Leu Leu	
-20 -15	
GGG CTC CTG CTA CTC TGG CTC CGA GGT GCC	60
Gly Leu Leu Leu Leu Trp Leu Arg Gly Ala	
-10 -5	
AGA TGT GAT ATC CAG ATG ACC CAG TCC CCG	90
Arg Cys Asp Ile Gln Met Thr Gln Ser Pro	
1 5	
TCC TCC CTG TCC GCC TCT GTG GGC GAT AGG	120
Ser Ser Leu Ser Ala Ser Val Gly Asp Arg	
10 15	

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 32 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 105.

Please delete lines 10-39, and insert therefor:

GTC ACC ATC ACC TGC GGC GCC AGC GAA AAC	150
Val Thr Ile Thr Cys Gly Ala Ser Glu Asn	
20 25	
ATC TAT GGC GCG CTG AAC TGG TAT CAA CAG	180
Ile Tyr Gly Ala Leu Asn Trp Tyr Gln Gln	
30 35	
AAA CCC GGG AAA GCT CCG AAG CTT CTG ATT	210
Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile	
40 45	
TAC GGT GCG ACG AAC CTG GCA GAT GGA GTC	240
Tyr Gly Ala Thr Asn Leu Ala Asp Gly Val	
50 55	
CCT TCT CGC TTC TCT GGA TCC GGC TCC GGA	270
Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly	
60 65	
ACG GAT TTC ACT CTG ACC ATC AGC AGT CTG	300
Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu	
70 75	
CAG CCT GAA GAC TTC GCT ACG TAT TAC TGT	330
Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys	
80 85	
CAG AAC GTT TTA AAT ACT CCG TTG ACT TTC	360
Gln Asn Val Leu Asn Thr Pro Leu Thr Phe	
90 95	
GGA CAG GGT ACC AAG GTG GAA ATA AAA CGA	390
Gly Gln Gly Thr Lys Val Glu Ile Lys Arg	
100 105	
ACT GTG GCT GCA CCA TCT GTC TTC ATC TTC	420
Thr Val Ala Ala Pro Ser Val Phe Ile Phe	
110 115	

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 33 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 105,

Please delete line 40, through column 107, line 9, and insert therefor:

CCG CCA TCT GAT GAG CAG TTG AAA TCT GGA	450
Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly	
120 125	
ACT GCC TCT GTT GTG TGC CTG CTG AAT AAC	480
Thr Ala Ser Val Val Cys Leu Leu Asn Asn	
130 135	
TTC TAT CCC AGA GAG GCC AAA GTA CAG TGG	510
Phe Tyr Pro Arg Glu Ala Lys Val Gln Trp	
140 145	
AAG GTG GAT AAC GCC CTC CAA TCG GGT AAC	540
Lys Val Asp Asn Ala Leu Gln Ser Gly Asn	
150 155	
TCC CAG GAG AGT GTC ACA GAG CAG GAC AGC	570
Ser Gln Glu Ser Val Thr Glu Gln Asp Ser	
160 165	
AAG GAC AGC ACC TAC AGC CTC AGC AGC ACC	600
Lys Asp Ser Thr Tyr Ser Leu Ser Ser Thr	
170 175	
CTG ACG CTG AGC AAA GCA GAC TAC GAG AAA	630
Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys	
180 185	
CAC AAA GTC TAC GCC TGC GAA GTC ACC CAT	660
His Lys Val Tyr Ala Cys Glu Val Thr His	
190 195	
CAG GGC CTG AGC TCG CCC GTC ACA AAG AGC	690
Gln Gly Leu Ser Ser Pro Val Thr Lys Ser	
200 205	
TTC AAC AGG GGA GAG TGT TAG	711
Phe Asn Arg Gly Glu Cys	
210	

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 34 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 107.

Please delete lines 20-49, and insert therefor:

ATG AAG TGG AGC TGG GTT ATT CTC TTC CTC	30
Met Lys Trp Ser Trp Val Ile Leu Phe Leu	
-15 -10	
CTG TCA GTA ACT GCC GGC GTC CAC TCC CAA	60
Leu Ser Val Thr Ala Gly Val His Ser Gln	
-5 1	
GTC CAA CTG GTG CAA TCC GGC GCC GAG GTC	90
Val Gln Leu Val Gln Ser Gly Ala Glu Val	
5 10	
AAG AAG CCA GGG GCC TCA GTC AAA GTG TCC	120
Lys Lys Pro Gly Ala Ser Val Lys Val Ser	
15 20	
TGT AAA GCT AGC GGC TAT ATT TTT TCT AAT	150
Cys Lys Ala Ser Gly Tyr Ile Phe Ser Asn	
25 30	
TAT TGG ATT CAA TGG GTG CGT CAG GCC CCC	180
Tyr Trp Ile Gln Trp Val Arg Gln Ala Pro	
35 40	
GGG CAG GGC CTG GAA TGG ATG GGT GAG ATC	210
Gly Gln Gly Leu Glu Trp Met Gly Glu Ile	
45 50	
TTA CCG GGC TCT GGT AGC ACC GAA TAT GCC	240
Leu Pro Gly Ser Gly Ser Thr Glu Tyr Ala	
55 60	
CAA AAA TTC CAG GGC CGT GTT ACT ATG ACT	270
Gln Lys Phe Gln Gly Arg Val Thr Met Thr	
65 70	
CGT GAC ACT TCG ACT AGT ACA GTA TAC ATG	300
Arg Asp Thr Ser Thr Ser Thr Val Tyr Met	
75 80	

CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 35 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 107,

Please delete line 50, through column 109, line 21, and insert therefor:

GAG CTC TCC AGC CTG CGA TCG GAG GAC ACG 330
Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr
85 90

GCC GTC TAT TAT TGC GCG CGT TAT TTT TTT 360
Ala Val Tyr Tyr Cys Ala Arg Tyr Phe Phe
95 100

GGT TCT AGC CCG AAT TGG TAT TTT GAT GTT 390
Gly Ser Ser Pro Asn Trp Tyr Phe Asp Val
105 110

TGG GGT CAA GGA ACC CTG GTC ACT GTC TCG 420
 Trp Gly Gln Gly Thr Leu Val Thr Val Ser
 115 120

AGC GCC TCC ACC AAG GGC CCA TCG GTC TTC 450
Ser Ala Ser Thr Lys Gly Pro Ser Val Phe
125 130

CCC CTG GCG CCC TCC TCC AAG AGC ACC TCT 480
Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser
135 140

GGG GGC ACA GCG GCC CTG GGC TGC CTG GTC 510
Gly Gly Thr Ala Ala Leu Gly Cys Leu Val
145 150

AAG GAC TAC TTC CCC GAA CCG GTG ACG GTG 540
Lys Asp Tyr Phe Pro Glu Pro Val Thr Val
155 160

TCG TGG AAC TCA GGC GCC CTG ACC AGC GGC 570
Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly
165 170

GTG CAC ACC TTC CCG GCT GTC CTA CAG TCC 600
Val His Thr Phe Pro Ala Val Leu Gln Ser
175 180

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 36 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 109.

Please delete lines 22-36, and insert therefor:

TCA GGA CTC TAC TCC CTC AGC AGC GTG GTG	630
Ser Gly Leu Tyr Ser Leu Ser Ser Val Val	
185 190	
ACC GTG CCC TCC AGC AGC TTG GGC ACC CAG	660
Thr Val Pro Ser Ser Ser Leu Gly Thr Gln	
195 200	
ACC TAC ATC TGC AAC GTG AAT CAC AAG CCC	690
Thr Tyr Ile Cys Asn Val Asn His Lys Pro	
205 210	
AGC AAC ACC AAG GTG GAC AAG AAA GTT GAG	720
Ser Asn Thr Lys Val Asp Lys Lys Val Glu	
215 220	
CCC AAA TCT TGT GAC AAA ACT CAC ACA TAA	750
Pro Lys Ser Cys Asp Lys Thr His Thr	
225 230	

Please delete lines 47-58, and insert therefor:

ATG GCC GAT ATC CAG ATG ACC CAG TCC CCG	30
Met Ala Asp Ile Gln Met Thr Gln Ser Pro	
1 5 10	
TCC TCC CTG TCC GCC TCT GTG GGC GAT AGG	60
Ser Ser Leu Ser Ala Ser Val Gly Asp Arg	
15 20	
GTC ACC ATC ACC TGC GGC GCC AGC GAA AAC	90
Val Thr Ile Thr Cys Gly Ala Ser Glu Asn	
25 30	
ATC TAT GGC GCG CTG AAC TGG TAT CAA CAG	120
Ile Tyr Gly Ala Leu Asn Trp Tyr Gln Gln	
35 40	

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 37 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 111,

Please delete lines 1-30, and insert therefor:

AAA CCT GGG AAA GCT CCG AAG CTT CTG ATT	150
Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile	
45 50	
TAC GGT GCG ACG AAC CTG GCA GAT GGA GTC	180
Tyr Gly Ala Thr Asn Leu Ala Asp Gly Val	
55 60	
CCT TCT CGC TTC TCT GGA TCC GGC TCC GGA	210
Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly	
65 70	
ACG GAT TTC ACT CTG ACC ATC AGC AGT CTG	240
Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu	
75 80	
CAG CCT GAA GAC TTC GCT ACG TAT TAC TGT	270
Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys	
85 90	
CAG AAC GTT TTA AAT ACT CCG TTG ACT TTC	300
Gln Asn Val Leu Asn Thr Pro Leu Thr Phe	
95 100	
GGA CAG GGT ACC AAG GTG GAA ATA AAA CGT	330
Gly Gln Gly Thr Lys Val Glu Ile Lys Arg	
105 110	
ACT GGC GGT GGT GGT TCT GGT GGC GGT GGA	360
Thr Gly Gly Gly Gly Ser Gly Gly Gly Gly	
115 120	
TCT GGT GGT GGC GGT TCT CAA GTC CAA CTG	390
Ser Gly Gly Gly Gly Ser Gln Val Gln Leu	
125 130	
GTG CAA TCC GGC GCC GAG GTC AAG AAG CCA	420
Val Gln Ser Gly Ala Glu Val Lys Lys Pro	
135 140	

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 38 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 111,

Please delete lines 31-60, and insert therefor:

GGG GCC TCA GTC AAA GTG TCC TGT AAA GCT	450
Gly Ala Ser Val Lys Val Ser Cys Lys Ala	
145 150	
AGC GGC TAT ATT TTT TCT AAT TAT TGG ATT	480
Ser Gly Tyr Ile Phe Ser Asn Tyr Trp Ile	
155 160	
CAA TGG GTG CGT CAG GCC CCC GGG CAG GGC	510
Gln Trp Val Arg Gln Ala Pro Gly Gln Gly	
165 170	
CTG GAA TGG ATG GGT GAG ATC TTA CCG GGC	540
Leu Glu Trp Met Gly Glu Ile Leu Pro Gly	
175 180	
TCT GGT AGC ACC GAA TAT GCC CAA AAA TTC	570
Ser Gly Ser Thr Glu Tyr Ala Gln Lys Phe	
185 190	
CAG GGC CGT GTT ACT ATG ACG CGT GAC ACT	600
Gln Gly Arg Val Thr Met Thr Arg Asp Thr	
195 200	
TCG ACT AGT ACA GTA TAC ATG GAG CTC TCC	630
Ser Thr Ser Thr Val Tyr Met Glu Leu Ser	
205 210	
AGC CTG CGA TCG GAG GAC ACG GCC GTC TAT	660
Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr	
215 220	
TAT TGC GCG CGT TAT TTT TTT GGT TCT AGC	690
Tyr Cys Ala Arg Tyr Phe Phe Gly Ser Ser	
225 230	
CCG AAT TGG TAT TTT GAT GTT TGG GGT CAA	720
Pro Asn Trp Tyr Phe Asp Val Trp Gly Gln	
235 240	

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 39 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 113,

Please delete lines 1-3, and insert therefor:

GGA ACC CTG GTC ACT GTC TCG AGC TGA	747
Gly Thr Leu Val Thr Val Ser Ser	
245	

Column 119,

Please delete lines 2-25, and insert therefor:

ATG GCC AAT ATT GTG CTG ACC CAA TCT CCA	30
Met Ala Asn Ile Val Leu Thr Gln Ser Pro	
1 5 10	
GCT TCT TTG GCT GTG TCT CTA GGG CAG AGG	60
Ala Ser Leu Ala Val Ser Leu Gly Gln Arg	
15 20	
GCC ACC ATA TCC TGC AGA GCC AGT GAA AGT	90
Ala Thr Ile Ser Cys Arg Ala Ser Glu Ser	
25 30	
GTT GAT AGT TAT GAC AAT AGT TTT ATG CAC	120
Val Asp Ser Tyr Asp Asn Ser Phe Met His	
35 40	
TGG TAC CAG CAG AAA CCA GGA CAG CCA CCC	150
Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro	
45 50	
AAA CTC CTC ATC TTT CTT GCA TCC AAC CTA	180
Lys Leu Leu Ile Phe Leu Ala Ser Asn Leu	
55 60	
GAA TCT GGG GTC CCT GCC AGG TTC AGT GGC	210
Glu Ser Gly Val Pro Ala Arg Phe Ser Gly	
65 70	
AGT GGG TCT AGG ACA GAC TTC ACC CTC ACC	240
Ser Gly Ser Arg Thr Asp Phe Thr Leu Thr	
75 80	

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 40 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 119,

Please delete lines 26-55, and insert therefor:

ATT GAT CCT GTG GAG GCT GAT GAT GCT GCA	270
Ile Asp Pro Val Glu Ala Asp Asp Ala Ala	
85 90	
ACC TAT TAC TGT CAG CAA AAT AAT GAG GTT	300
Thr Tyr Tyr Cys Gln Gln Asn Asn Glu Val	
95 100	
CCG AAC ACG TTC GGA GGG GGG ACC AAG CTG	330
Pro Asn Thr Phe Gly Gly Gly Thr Lys Leu	
105 110	
GAA ATA AAA CGG ACC GGA GGT GGC GGG TCG	360
Glu Ile Lys Arg Thr Gly Gly Gly Gly Ser	
115 120	
GGT GGC GGG GGA TCG GGT GGC GGA GGG TCG	390
Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser	
125 130	
GAC GTC AAG CTC GTG GAG TCT GGG GGA GAC	420
Asp Val Lys Leu Val Glu Ser Gly Gly Asp	
135 140	
TTA GTG AAG CTT GGA GGG TCC CTG AAA CTC	450
Leu Val Lys Leu Gly Gly Ser Leu Lys Leu	
145 150	
TCC TGT GCA GCC TCT GGA TTC ACC TTC AGT	480
Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser	
155 160	
AGC TAT TAT ATG TCT TGG GTT CGC CAG ATT	510
Ser Tyr Tyr Met Ser Trp Val Arg Gln Ile	
165 170	
TCA GAG AAG AGG CTG GAG TTG GTC GCA GCC	540
Ser Glu Lys Arg Leu Glu Leu Val Ala Ala	
175 180	

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 41 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 119,

Please delete lines 56, through column 121, line 22, and insert therefor:

ATT AAT AGT AAT GGT GAT AGC ACC TAC TAT	570
Ile Asn Ser Asn Gly Asp Ser Thr Tyr Tyr	
185 190	
CCA GAC ACT GTG AAG GGC CGA TTC ACC ATC	600
Pro Asp Thr Val Lys Gly Arg Phe Thr Ile	
195 200	
TCC AGA GAC AAT GCC AAG AGC ACC CTG GAT	630
Ser Arg Asp Asn Ala Lys Ser Thr Leu Asp	
205 210	
CTG CAA ATG AGC AGT CTG AAG TCT GAG GAC	660
Leu Gln Met Ser Ser Leu Lys Ser Glu Asp	
215 220	
ACA GCC TTG TAT TTC TGT GTA AGA GAG ACT	690
Thr Ala Leu Tyr Phe Cys Val Arg Glu Thr	
225 230	
TAT TAC TAC GGG ATT AGT CCC GTC TTC GAT	720
Tyr Tyr Tyr Gly Ile Ser Pro Val Phe Asp	
235 240	
GTC TGG GGC ACA GGG ACC ACG GTC ACC GTC	750
Val Trp Gly Thr Gly Thr Thr Val Thr Val	
245 250	
TCC TCA CTC GAG CAC CAC CAC CAC CAC	780
Ser Ser Leu Glu His His His His His	
255 260	
TGA	783

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 42 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 121,

Please delete lines 32, through column 123, line 3, and insert therefor:

ATG GCC GAT ATC CAG ATG ACC CAG TCC CCG	30
Met Ala Asp Ile Gln Met Thr Gln Ser Pro	
1 5 10	
TCC TCC CTG TCC GCC TCT GTG GGC GAT AGG	60
Ser Ser Leu Ser Ala Ser Val Gly Asp Arg	
15 20	
GTC ACC ATC ACC TGC GGC GCC AGC GAA AAC	90
Val Thr Ile Thr Cys Gly Ala Ser Glu Asn	
25 30	
ATC TAT GGC GCG CTG AAC TGG TAT CAA CAG	120
Ile Tyr Gly Ala Leu Asn Trp Tyr Gln Gln	
35 40	
AAA CCC GGG AAA GCT CCG AAG CTT CTG ATT	150
Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile	
45 50	
TAC GGT GCG ACG AAC CTG GCA GAT GGA GTC	180
Tyr Gly Ala Thr Asn Leu Ala Asp Gly Val	
55 60	
CCT TCT CGC TTC TCT GGA TCC GGC TCC GGA	210
Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly	
65 70	
ACG GAT TTC ACT CTG ACC ATC AGC AGT CTG	240
Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu	
75 80	
CAG CCT GAA GAC TTC GCT ACG TAT TAC TGT	270
Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys	
85 90	
CAG AAC GTT TTA AAT ACT CCG TTG ACT TTC	300
Gln Asn Val Leu Asn Thr Pro Leu Thr Phe	
95 100	

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 43 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 123,

Please delete lines 4-33, and insert therefor:

GGA CAG GGT ACC AAG GTG GAA ATA AAA CGT	330
Gly Gln Gly Thr Lys Val Glu Ile Lys Arg	
105 110	
ACT GGC GGT GGT GGT TCT GGT GGC GGT GGA	360
Thr Gly Gly Gly Gly Ser Gly Gly Gly Gly	
115 120	
TCT GGT GGT GGC GGT TCT CAA GTC CAA CTG	390
Ser Gly Gly Gly Gly Ser Gln Val Gln Leu	
125 130	
GTG CAA TCC GGC GCC GAG GTC AAG AAG CCA	420
Val Gln Ser Gly Ala Glu Val Lys Lys Pro	
135 140	
GGG GCC TCA GTC AAA GTG TCC TGT AAA GCT	450
Gly Ala Ser Val Lys Val Ser Cys Lys Ala	
145 150	
AGC GGC TAT ATT TTT TCT AAT TAT TGG ATT	480
Ser Gly Tyr Ile Phe Ser Asn Tyr Trp Ile	
155 160	
CAA TGG GTG CGT CAG GCC CCC GGG CAG GGC	510
Gln Trp Val Arg Gln Ala Pro Gly Gln Gly	
165 170	
CTG GAA TGG ATG GGT GAG ATC TTA CCG GGC	540
Leu Glu Trp Met Gly Glu Ile Leu Pro Gly	
175 180	
TCT GGT AGC ACC GAA TAT ACC GAA AAT TTT	570
Ser Gly Ser Thr Glu Tyr Thr Glu Asn Phe	
185 190	
AAA GAC CGT GTT ACT ATG ACG CGT GAC ACT	600
Lys Asp Arg Val Thr Met Thr Arg Asp Thr	
195 200	

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 44 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 123,

Please delete lines 34-48, and insert therefor:

TCG ACT AGT ACA GTA TAC ATG GAG CTC TCC	630
Ser Thr Ser Thr Val Tyr Met Glu Leu Ser	
205 210	
AGC CTG CGA TCG GAG GAC ACG GCC GTC TAT	660
Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr	
215 220	
TAT TGC GCG CGT TAT TTT TTT GGT TCT AGC	690
Tyr Cys Ala Arg Tyr Phe Phe Gly Ser Ser	
225 230	
CCG AAT TGG TAT TTT GAT GTT TGG GGT CAA	720
Pro Asn Trp Tyr Phe Asp Val Trp Gly Gln	
235 240	
GGA ACC CTG GTC ACT GTC TCG AGC TGA	747
Gly Thr Leu Val Thr Val Ser Ser	
245	

Column 125,

Please delete lines 2-13, and insert therefor:

ATG GCC GAT ATC CAG ATG ACC CAG TCC CCG	30
Met Ala Asp Ile Gln Met Thr Gln Ser Pro	
1 5 10	
TCC TCC CTG TCC GCC TCT GTG GGC GAT AGG	60
Ser Ser Leu Ser Ala Ser Val Gly Asp Arg	
15 20	
GTC ACC ATC ACC TGC CGT GCT AGC GAA AAC	90
Val Thr Ile Thr Cys Arg Ala Ser Glu Asn	
25 30	
ATC TAT GGC GCG CTG AAC TGG TAT CAA CAG	120
Ile Tyr Gly Ala Leu Asn Trp Tyr Gln Gln	
35 40	

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 45 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 125,

Please delete lines 14-43, and insert therefor:

AAA CCT GGG AAA GCT CCG AAG CTT CTG ATT	150
Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile	
45 50	
TAC GGT GCG ACG AAC CTG GCA GAT GGA GTC	180
Tyr Gly Ala Thr Asn Leu Ala Asp Gly Val	
55 60	
CCT TCT CGC TTC TCT GGA TCC GGC TCC GGA	210
Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly	
65 70	
ACG GAT TTC ACT CTG ACC ATC AGC AGT CTG	240
Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu	
75 80	
CAG CCT GAA GAC TTC GCT ACG TAT TAC TGT	270
Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys	
85 90	
CAG AAC GTT TTA AAT ACT CCG TTG ACT TTC	300
Gln Asn Val Leu Asn Thr Pro Leu Thr Phe	
95 100	
GGA CAG GGT ACC AAG GTG GAA ATA AAA CGT	330
Gly Gln Gly Thr Lys Val Glu Ile Lys Arg	
105 110	
ACT GGC GGT GGT GGT TCT GGT GGC GGT GGA	360
Thr Gly Gly Gly Gly Ser Gly Gly Gly Gly	
115 120	
TCT GGT GGT GGC GGT TCT CAA GTC CAA CTG	390
Ser Gly Gly Gly Gly Ser Gln Val Gln Leu	
125 130	
GTG CAA TCC GGC GCC GAG GTC AAG AAG CCA	420
Val Gln Ser Gly Ala Glu Val Lys Lys Pro	
135 140	

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 46 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 125,

Please delete line 44, through column 127, line 15, and insert therefor:

GGG GCC TCA GTC AAA GTG TCC TGT AAA GCT	450
Gly Ala Ser Val Lys Val Ser Cys Lys Ala	
145 150	
AGC GGC TAT ATT TTT TCT AAT TAT TGG ATT	480
Ser Gly Tyr Ile Phe Ser Asn Tyr Trp Ile	
155 160	
CAA TGG GTG CGT CAG GCC CCC GGG CAG GGC	510
Gln Trp Val Arg Gln Ala Pro Gly Gln Gly	
165 170	
CTG GAA TGG ATG GGT GAG ATC TTA CCG GGC	540
Leu Glu Trp Met Gly Glu Ile Leu Pro Gly	
175 180	
TCT GGT AGC ACC GAA TAT GCC CAA AAA TTC	570
Ser Gly Ser Thr Glu Tyr Ala Gln Lys Phe	
185 190	
CAG GGC CGT GTT ACT ATG ACG CGT GAC ACT	600
Gln Gly Arg Val Thr Met Thr Arg Asp Thr	
195 200	
TCG ACT AGT ACA GTA TAC ATG GAG CTC TCC	630
Ser Thr Ser Thr Val Tyr Met Glu Leu Ser	
205 210	
AGC CTG CGA TCG GAG GAC ACG GCC GTC TAT	660
Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr	
215 220	
TAT TGC GCG CGT TAT TTT TTT GGT TCT AGC	690
Tyr Cys Ala Arg Tyr Phe Phe Gly Ser Ser	
225 230	
CCG AAT TGG TAT TTT GAT GTT TGG GGT CAA	720
Pro Asn Trp Tyr Phe Asp Val Trp Gly Gln	
235 240	

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 47 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 127,

Please delete lines 16-18, and insert therefor:

GGA ACC CTG GTC ACT GTC TCG AGC TGA	747
Gly Thr Leu Val Thr Val Ser Ser	
245	

Please delete lines 29-52, and insert therefor:

ATG GCC GAT ATC CAG ATG ACC CAG TCC CCG	30
Met Ala Asp Ile Gln Met Thr Gln Ser Pro	
1 5 10	

TCC TCC CTG TCC GCC TCT GTG GGC GAT AGG	60
Ser Ser Leu Ser Ala Ser Val Gly Asp Arg	
15 20	

GTC ACC ATC ACC TGC GGC GCC AGC GAA AAC	90
Val Thr Ile Thr Cys Gly Ala Ser Glu Asn	
25 30	

ATC TAT GGC GCG CTG AAC TGG TAT CAA CAG	120
Ile Tyr Gly Ala Leu Asn Trp Tyr Gln Gln	
35 40	

AAA CCT GGG AAA GCT CCG AAG CTT CTG ATT	150
Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile	
45 50	

TAC GGT GCG ACG AGC CTG CAG TCT GGA GTC	180
Tyr Gly Ala Thr Ser Leu Gln Ser Gly Val	
55 60	

CCT TCT CGC TTC TCT GGA TCC GGC TCC GGA	210
Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly	
65 70	

ACG GAT TTC ACT CTG ACC ATC AGC AGT CTG	240
Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu	
75 80	

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 48 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 127,

Please delete line 53, through column 129, line 24, and insert therefor:

CAG CCT GAA GAC TTC GCT ACG TAT TAC TGT Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys 85 90	270
CAG AAC GTT TTA AAT ACT CCG TTG ACT TTC Gln Asn Val Leu Asn Thr Pro Leu Thr Phe 95 100	300
GGA CAG GGT ACC AAG GTG GAA ATA AAA CGT Gly Gln Gly Thr Lys Val Glu Ile Lys Arg 105 110	330
ACT GGC GGT GGT GGT TCT GGT GGC GGT GGA Thr Gly Gly Gly Gly Ser Gly Gly Gly Gly 115 120	360
TCT GGT GGT GGC GGT TCT CAA GTC CAA CTG Ser Gly Gly Gly Gly Ser Gln Val Gln Leu 125 130	390
GTG CAA TCC GGC GCC GAG GTC AAG AAG CCA Val Gln Ser Gly Ala Glu Val Lys Lys Pro 135 140	420
GGG GCC TCA GTC AAA GTG TCC TGT AAA GCT Gly Ala Ser Val Lys Val Ser Cys Lys Ala 145 150	450
AGC GGC TAT ATT TTT TCT AAT TAT TGG ATT Ser Gly Tyr Ile Phe Ser Asn Tyr Trp Ile 155 160	480
CAA TGG GTG CGT CAG GCC CCC GGG CAG GGC Gln Trp Val Arg Gln Ala Pro Gly Gln Gly 165 170	510
CTG GAA TGG ATG GGT GAG ATC TTA CCG GGC Leu Glu Trp Met Gly Glu Ile Leu Pro Gly 175 180	540

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 49 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 129,

Please delete lines 25-45, and insert therefor:

TCT GGT AGC ACC GAA TAT GCC CAA AAA TTC	570
Ser Gly Ser Thr Glu Tyr Ala Gln Lys Phe	
185 190	
CAG GGC CGT GTT ACT ATG ACG CGT GAC ACT	600
Gln Gly Arg Val Thr Met Thr Arg Asp Thr	
195 200	
TCG ACT AGT ACA GTA TAC ATG GAG CTC TCC	630
Ser Thr Ser Thr Val Tyr Met Glu Leu Ser	
205 210	
AGC CTG CGA TCG GAG GAC ACG GCC GTC TAT	660
Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr	
215 220	
TAT TGC GCG CGT TAT TTT TTT GGT TCT AGC	690
Tyr Cys Ala Arg Tyr Phe Phe Gly Ser Ser	
225 230	
CCG AAT TGG TAT TTT GAT GTT TGG GGT CAA	720
Pro Asn Trp Tyr Phe Asp Val Trp Gly Gln	
235 240	
GGA ACC CTG GTC ACT GTC TCG AGC TGA	747
Gly Thr Leu Val Thr Val Ser Ser	
245	

Please delete line 56, through column 131, line 3, and insert therefor:

ATG GCC GAT ATC CAG ATG ACC CAG TCC CCG	30
Met Ala Asp Ile Gln Met Thr Gln Ser Pro	
1 5 10	
TCC TCC CTG TCC GCC TCT GTG GGC GAT AGG	60
Ser Ser Leu Ser Ala Ser Val Gly Asp Arg	
15 20	

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 50 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 131,

Please delete lines 4-33, and insert therefor:

GTC ACC ATC ACC TGC CGT GCT AGC GAA AAC	90
Val Thr Ile Thr Cys Arg Ala Ser Glu Asn	
25 30	
ATC TAT GGC GCG CTG AAC TGG TAT CAA CAG	120
Ile Tyr Gly Ala Leu Asn Trp Tyr Gln Gln	
35 40	
AAA CCT GGG AAA GCT CCG AAG CTT CTG ATT	150
Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile	
45 50	
TAC GGT GCG ACG AGC CTG CAG TCT GGA GTC	180
Tyr Gly Ala Thr Ser Leu Gln Ser Gly Val	
55 60	
CCT TCT CGC TTC TCT GGA TCC GGC TCC GGA	210
Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly	
65 70	
ACG GAT TTC ACT CTG ACC ATC AGC AGT CTG	240
Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu	
75 80	
CAG CCT GAA GAC TTC GCT ACG TAT TAC TGT	270
Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys	
85 90	
CAG AAC GTT TTA AAT ACT CCG TTG ACT TTC	300
Gln Asn Val Leu Asn Thr Pro Leu Thr Phe	
95 100	
GGA CAG GGT ACC AAG GTG GAA ATA AAA CGT	330
Gly Gln Gly Thr Lys Val Glu Ile Lys Arg	
105 110	
ACT GGC GGT GGT GGT TCT GGT GGC GGT GGA	360
Thr Gly Gly Gly Gly Ser Gly Gly Gly Gly	
115 120	

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 51 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 131,

Please delete line 34, through column 133, line 3, and insert therefor:

TCT GGT GGT GGC GGT TCT CAA GTC CAA CTG	390
Ser Gly Gly Gly Gly Ser Gln Val Gln Leu	
125 130	
GTG CAA TCC GGC GCC GAG GTC AAG AAG CCA	420
Val Gln Ser Gly Ala Glu Val Lys Lys Pro	
135 140	
GGG GCC TCA GTC AAA GTG TCC TGT AAA GCT	450
Gly Ala Ser Val Lys Val Ser Cys Lys Ala	
145 150	
AGC GGC TAT ATT TTT TCT AAT TAT TGG ATT	480
Ser Gly Tyr Ile Phe Ser Asn Tyr Trp Ile	
155 160	
CAA TGG GTG CGT CAG GCC CCC GGG CAG GGC	510
Gln Trp Val Arg Gln Ala Pro Gly Gln Gly	
165 170	
CTG GAA TGG ATG GGT GAG ATC TTA CCG GGC	540
Leu Glu Trp Met Gly Glu Ile Leu Pro Gly	
175 180	
TCT GGT AGC ACC GAA TAT GCC CAA AAA TTC	570
Ser Gly Ser Thr Glu Tyr Ala Gln Lys Phe	
185 190	
CAG GGC CGT GTT ACT ATG ACG CGT GAC ACT	600
Gln Gly Arg Val Thr Met Thr Arg Asp Thr	
195 200	
TCG ACT AGT ACA GTA TAC ATG GAG CTC TCC	630
Ser Thr Ser Thr Val Tyr Met Glu Leu Ser	
205 210	
AGC CTG CGA TCG GAG GAC ACG GCC GTC TAT	660
Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr	
215 220	

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 52 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 133.

Please delete lines 4-12, and insert therefor:

TAT TGC GCG CGT TAT TTT TTT GGT TCT AGC	690
Tyr Cys Ala Arg Tyr Phe Phe Gly Ser Ser	
225 230	
CCG AAT TGG TAT TTT GAT GTT TGG GGT CAA	720
Pro Asn Trp Tyr Phe Asp Val Trp Gly Gln	
235 240	
GGA ACC CTG GTC ACT GTC TCG AGC TGA	747
Gly Thr Leu Val Thr Val Ser Ser	
245	

Please delete lines 22-45, and insert therefor:

ATG GCC GAT ATC CAG ATG ACC CAG TCC CCG	30
Met Ala Asp Ile Gln Met Thr Gln Ser Pro	
1 5 10	
TCC TCC CTG TCC GCC TCT GTG GGC GAT AGG	60
Ser Ser Leu Ser Ala Ser Val Gly Asp Arg	
15 20	
GTC ACC ATC ACC TGC CGT GCT AGC GAA AAC	90
Val Thr Ile Thr Cys Arg Ala Ser Glu Asn	
25 30	
ATC TAT GGC GCG CTG AAC TGG TAT CAA CAG	120
Ile Tyr Gly Ala Leu Asn Trp Tyr Gln Gln	
35 40	
AAA CCC GGG AAA GCT CCG AAG CTT CTG ATT	150
Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile	
45 50	
TAC GGT GCG ACG AAC CTG GCA GAT GGA GTC	180
Tyr Gly Ala Thr Asn Leu Ala Asp Gly Val	
55 60	

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 53 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 133,

Please delete line 46, through column 135, line 12, and insert therefor:

CCT TCT CGC TTC TCT GGA TCC GGC TCC GGA	210
Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly	
65 70	
ACG GAT TTC ACT CTG ACC ATC AGC AGT CTG	240
Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu	
75 80	
CAG CCT GAA GAC TTC GCT ACG TAT TAC TGT	270
Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys	
85 90	
CAG AAC GTT TTA AAT ACT CCG TTG ACT TTC	300
Gln Asn Val Leu Asn Thr Pro Leu Thr Phe	
95 100	
GGA CAG GGT ACC AAG GTG GAA ATA AAA CGT	330
Gly Gln Gly Thr Lys Val Glu Ile Lys Arg	
105 110	
ACT GGC GGT GGT GGT TCT GGT GGC GGT GGA	360
Thr Gly Gly Gly Gly Ser Gly Gly Gly Gly	
115 120	
TCT GGT GGT GGC GGT TCT CAA GTC CAA CTG	390
Ser Gly Gly Gly Gly Ser Gln Val Gln Leu	
125 130	
GTG CAA TCC GGC GCC GAG GTC AAG AAG CCA	420
Val Gln Ser Gly Ala Glu Val Lys Lys Pro	
135 140	
GGG GCC TCA GTC AAA GTG TCC TGT AAA GCT	450
Gly Ala Ser Val Lys Val Ser Cys Lys Ala	
145 150	
AGC GGC TAT ATT TTT TCT AAT TAT TGG ATT	480
Ser Gly Tyr Ile Phe Ser Asn Tyr Trp Ile	
155 160	

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 54 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 135.

Please delete lines 13-39, and insert therefor:

CAA TGG GTG CGT CAG GCC CCC GGG CAG GGC	510
Gln Trp Val Arg Gln Ala Pro Gly Gln Gly	
165 170	
CTG GAA TGG ATG GGT GAG ATC TTA CCG GGC	540
Leu Glu Trp Met Gly Glu Ile Leu Pro Gly	
175 180	
.TCT GGT AGC ACC GAA TAT ACC GAA AAT TTT	570
Ser Gly Ser Thr Glu Tyr Thr Glu Asn Phe	
185 190	
AAA GAC CGT GTT ACT ATG ACG CGT GAC ACT	600
Lys Asp Arg Val Thr Met Thr Arg Asp Thr	
195 200	
TCG ACT AGT ACA GTA TAC ATG GAG CTC TCC	630
Ser Thr Ser Thr Val Tyr Met Glu Leu Ser	
205 210	
AGC CTG CGA TCG GAG GAC ACG GCC GTC TAT	660
Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr	
215 220	
TAT TGC GCG CGT TAT TTT TTT GGT TCT AGC	690
Tyr Cys Ala Arg Tyr Phe Phe Gly Ser Ser	
225 230	
CCG AAT TGG TAT TTT GAT GTT TGG GGT CAA	720
Pro Asn Trp Tyr Phe Asp Val Trp Gly Gln	
235 240	
GGA ACC CTG GTC ACT GTC TCG AGC TGA	747
Gly Thr Leu Val Thr Val Ser Ser	
245	

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 55 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 135,

Please delete line 49, through column 137, line 21, and insert therefor:

ATG GCC GAT ATC CAG ATG ACC CAG TCC CCG	30
Met Ala Asp Ile Gln Met Thr Gln Ser Pro	
1 5 10	
TCC TCC CTG TCC GCC TCT GTG GGC GAT AGG	60
Ser Ser Leu Ser Ala Ser Val Gly Asp Arg	
15 20	
GTC ACC ATC ACC TGC GGC GCC AGC GAA AAC	90
Val Thr Ile Thr Cys Gly Ala Ser Glu Asn	
25 30	
ATC TAT GGC GCG CTG AAC TGG TAT CAA CAG	120
Ile Tyr Gly Ala Leu Asn Trp Tyr Gln Gln	
35 40	
AAA CCC GGG AAA GCT CCG AAG CTT CTG ATT	150
Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile	
45 50	
TAC GGT GCG ACG AGC CTG CAG TCT GGA GTC	180
Tyr Gly Ala Thr Ser Leu Gln Ser Gly Val	
55 60	
CCT TCT CGC TTC TCT GGA TCC GGC TCC GGA	210
Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly	
65 70	
ACG GAT TTC ACT CTG ACC ATC AGC AGT CTG	240
Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu	
75 80	
CAG CCT GAA GAC TTC GCT ACG TAT TAC TGT	270
Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys	
85 90	
CAG AAC GTT TTA AAT ACT CCG TTG ACT TTC	300
Gln Asn Val Leu Asn Thr Pro Leu Thr Phe	
95 100	

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 56 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 137.

Please delete lines 22-51, and insert therefor:

GGA CAG GGT ACC AAG GTG GAA ATA AAA CGT	330
Gly Gln Gly Thr Lys Val Glu Ile Lys Arg	
105 110	
ACT GGC GGT GGT GGT TCT GGT GGC GGT GGA	360
Thr Gly Gly Gly Gly Ser Gly Gly Gly Gly	
115 120	
TCT GGT GGT GGC GGT TCT CAA GTC CAA CTG	390
Ser Gly Gly Gly Gly Ser Gln Val Gln Leu	
125 130	
GTG CAA TCC GGC GCC GAG GTC AAG AAG CCA	420
Val Gln Ser Gly Ala Glu Val Lys Lys Pro	
135 140	
GGG GCC TCA GTC AAA GTG TCC TGT AAA GCT	450
Gly Ala Ser Val Lys Val Ser Cys Lys Ala	
145 150	
AGC GGC TAT ATT TTT TCT AAT TAT TGG ATT	480
Ser Gly Tyr Ile Phe Ser Asn Tyr Trp Ile	
155 160	
CAA TGG GTG CGT CAG GCC CCC GGC CAG GGC	510
Gln Trp Val Arg Gln Ala Pro Gly Gln Gly	
165 170	
CTG GAA TGG ATG GGT GAG ATC TTA CCG GGC	540
Leu Glu Trp Met Gly Glu Ile Leu Pro Gly	
175 180	
TCT GGT AGC ACC GAA TAT ACC GAA AAT TTT	570
Ser Gly Ser Thr Glu Tyr Thr Glu Asn Phe	
185 190	
AAA GAC CGT GTT ACT ATG ACG CGT GAC ACT	600
Lys Asp Arg Val Thr Met Thr Arg Asp Thr	
195 200	

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 57 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 137,

Please delete line 52, through column 139, line 6, and insert therefor:

TCG ACT AGT ACA GTA TAC ATG GAG CTC TCC	630
Ser Thr Ser Thr Val Tyr Met Glu Leu Ser	
205 210	
AGC CTG CGA TCG GAG GAC ACG GCC GTC TAT	660
Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr	
215 220	
TAT TGC GCG CGT TAT TTT TTT GGT TCT AGC	690
Tyr Cys Ala Arg Tyr Phe Phe Gly Ser Ser	
225 230	
CCG AAT TGG TAT TTT GAT GTT TGG GGT CAA	720
Pro Asn Trp Tyr Phe Asp Val Trp Gly Gln	
235 240	
GGA ACC CTG GTC ACT GTC TCG AGC TGA	747
Gly Thr Leu Val Thr Val Ser Ser	
245	

Column 139,

Please delete lines 16-27, and insert therefor:

ATG GCC GAT ATC CAG ATG ACC CAG TCC CCG	30
Met Ala Asp Ile Gln Met Thr Gln Ser Pro	
1 5 10	
TCC TCC CTG TCC GCC TCT GTG GGC GAT AGG	60
Ser Ser Leu Ser Ala Ser Val Gly Asp Arg	
15 20	
GTC ACC ATC ACC TGC CGT GCT AGC GAA AAC	90
Val Thr Ile Thr Cys Arg Ala Ser Glu Asn	
25 30	
ATC TAT GGC GCG CTG AAC TGG TAT CAA CAG	120
Ile Tyr Gly Ala Leu Asn Trp Tyr Gln Gln	
35 40	

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 58 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 139,

Please delete lines 28-57, and insert therefor:

AAA CCC GGG AAA GCT CCG AAG CTT CTG ATT	150
Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile	
45 50	
TAC GGT GCG ACG AGC CTG CAG TCT GGA GTC	180
Tyr Gly Ala Thr Ser Leu Gln Ser Gly Val	
55 60	
CCT TCT CGC TTC TCT GGA TCC GGC TCC GGA	210
Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly	
65 70	
ACG GAT TTC ACT CTG ACC ATC AGC AGT CTG	240
Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu	
75 80	
CAG CCT GAA GAC TTC GCT ACG TAT TAC TGT	270
Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys	
85 90	
CAG AAC GTT TTA AAT ACT CCG TTG ACT TTC	300
Gln Asn Val Leu Asn Thr Pro Leu Thr Phe	
95 100	
GGA CAG GGT ACC AAG GTG GAA ATA AAA CGT	330
Gly Gln Gly Thr Lys Val Glu Ile Lys Arg	
105 110	
ACT GGC GGT GGT GGT TCT GGT GGC GGT GGA	360
Thr Gly Gly Gly Gly Ser Gly Gly Gly Gly	
115 120	
TCT GGT GGT GGC GGT TCT CAA GTC CAA CTG	390
Ser Gly Gly Gly Gly Ser Gln Val Gln Leu	
125 130	
GTG CAA TCC GGC GCC GAG GTC AAG AAG CCA	420
Val Gln Ser Gly Ala Glu Val Lys Lys Pro	
135 140	

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 59 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 141,

Please delete lines 1-30, and insert therefor:

GGG GCC TCA GTC AAA GTG TCC TGT AAA GCT	450
Gly Ala Ser Val Lys Val Ser Cys Lys Ala	
145 150	
AGC GGC TAT ATT TTT TCT AAT TAT TGG ATT	480
Ser Gly Tyr Ile Phe Ser Asn Tyr Trp Ile	
155 160	
CAA TGG GTG CGT CAG GCC CCC GGG CAG GGC	510
Gln Trp Val Arg Gln Ala Pro Gly Gln Gly	
165 170	
CTG GAA TGG ATG GGT GAG ATC TTA CCG GGC	540
Leu Glu Trp Met Gly Glu Ile Leu Pro Gly	
175 180	
TCT GGT AGC ACC GAA TAT ACC GAA AAT TTT	570
Ser Gly Ser Thr Glu Tyr Thr Glu Asn Phe	
185 190	
AAA GAC CGT GTT ACT ATG ACG CGT GAC ACT	600
Lys Asp Arg Val Thr Met Thr Arg Asp Thr	
195 200	
TCG ACT AGT ACA GTA TAC ATG GAG CTC TCC	630
Ser Thr Ser Thr Val Tyr Met Glu Leu Ser	
205 210	
AGC CTG CGA TCG GAG GAC ACG GCC GTC TAT	660
Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr	
215 220	
TAT TGC GCG CGT TAT TTT TTT GGT TCT AGC	690
Tyr Cys Ala Arg Tyr Phe Phe Gly Ser Ser	
225 230	
CCG AAT TGG TAT TTT GAT GTT TGG GGT CAA	720
Pro Asn Trp Tyr Phe Asp Val Trp Gly Gln	
235 240	

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 60 of 60

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 141,

Please delete lines 31-33, and insert therefor:

GGA ACC CTG GTC ACT GTC TCG AGC TGA
Gly Thr Leu Val Thr Val Ser Ser
245

747

Signed and Sealed this

Nineteenth Day of November, 2002

Attest:

A handwritten signature in black ink, appearing to read "James E. Rogan", with a horizontal line drawn underneath it.

Attesting Officer

JAMES E. ROGAN
Director of the United States Patent and Trademark Office

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,355,245 B1
APPLICATION NO. : 08/487283
DATED : March 12, 2002
INVENTOR(S) : Mark J. Evans et al.

Page 1 of 1

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

On the title page, item (63) should read:

Related U.S. Application Data

- (63) Continuation-in-part of application No. 08/311,489, filed on September 23, 1994, and continuation-in-part of application No. PCT/US95/05688, filed on May 1, 1995, which is a continuation-in-part of application No. 08/236,208, filed on May 2, 1994, now Pat. No. 6,074,642.


In the Specification,

At column 1, the paragraph at lines 5-11 should read:

This is a continuation-in-part of International application Serial No. PCT/US95/05688, filed May 1, 1995, which is a continuation-in-part of U.S. application Serial No. 08/236,208, filed May 2, 1994, which issued as U.S. Pat. No. 6,074,642 on June 13, 2000. International application Serial No. PCT/US95/05688 was published in English under PCT Article 21(2) on November 9, 1995 as WO 95/29697.

This is also a continuation-in-part of U.S. application Serial No. 08/311,489, filed on September 23, 1994.

Signed and Sealed this
Twentieth Day of December, 2016



Michelle K. Lee
Director of the United States Patent and Trademark Office