IMMUNO BIOLOGY

the immune system in health and disease

6th EDITION



JANEWAY • TRAVERS • WALPORT • SHLOMCHIK

ALXN2031
IPR2023-00999

ALXN2031
IPR2019-00739

IMMUNOBIOLOGY

the immune system in health and disease

6th EDITION

IMMUNOBIOLOGY

the immune system in health and disease

6th EDITION

Charles A. Janeway, Jr.
Yale University School of Medicine

Paul Travers
Anthony Nolan Research Institute, London

Mark Walport
Wellcome Trust, London

Mark J. Shlomchik
Yale University School of Medicine

GS Garland Science
Taylor & Francis Group
NEW YORK AND LONDON

Vice President:	Denise Schanck
Text Editor:	Eleanor Lawrence
Senior Publisher:	Jackie Harbor
Editorial Assistants:	Emma Catherall & Sbita Reandi
Production Editor:	Georgia Bushell
New Media Editor:	Michael Morales
Copyeditor:	Bruce Goatly
Indexer:	Merrall-Ross International Ltd
Illustration and Layout:	Matthew McClements, Blink Studio Ltd
Manufacturing:	Andrea Bosher & Georgia Bushell

***Immunobiology Interactive*:**

Storyboards by:	Paul Travers
Narrated by:	Julie Theriot
Animations, Interface Design & Programming:	Matthew McClements & Tom McElderry, Blink Studio Ltd

© 2005 by Garland Science Publishing.
All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means—electronic, mechanical, photocopying, recording, or otherwise—without the prior written permission of the copyright holder.

Distributors:
Inside North America: Garland Science Publishing, 29 West 35th Street, New York, NY 10001-2299, USA.
Inside Japan: Nankodo Co. Ltd., 42-6, Hongo 3-Chrome, Bunkyo-ku, Tokyo, 113-8410, Japan.
Outside North America and Japan: Elsevier Ltd, Books Customer Services, Linacre House, Jordan Hill, Oxford, OX2 8DP, UK.

ISBN 0 8153 4101 6 (paperback) Garland Science Publishing
ISBN 0 4430 7310 4 (paperback) Churchill Livingstone
ISBN 0 4430 7309 0 (paperback) Churchill Livingstone International Student Edition

Library of Congress Cataloging-in-Publication Data
Immunobiology : the immune system in health and disease / Charles A. Janeway, Jr. .. [et al.].-- 6th ed.
p. cm.
Includes bibliographical references and index.
ISBN 0-8153-4101-6 (pbk.) -- ISBN 0-443-07310-4 (pbk.) -- ISBN 0-443-07309-0 (pbk.)
1. Immunology. 2. Immunity. I. Janeway, Charles.
QR181.I397 2005
616.07'9--dc22
2004007144

Published by Garland Science Publishing, a member of the Taylor & Francis Group, 29 West 35th Street, New York, NY 10001-2299, USA and 4 Park Square, Milton Park, Abingdon, Oxon, OX14 4RN, UK.

Printed in the United States of America.
15 14 13 12 11 10 9 8 7 6 5 4 3 2 1



Fig. A.13 The Coombs direct and indirect anti-globulin tests for antibody to red blood cell antigens. A Rh⁻ mother of a Rh⁺ fetus can become immunized to fetal red blood cells that enter the maternal circulation at the time of delivery. In a subsequent pregnancy with a Rh⁺ fetus, IgG anti-Rh antibodies can cross the placenta and damage the fetal red blood cells. In contrast to anti-Rh antibodies, maternal anti-ABO antibodies are of the IgM isotype and cannot cross the placenta, and so do not cause harm. Anti-Rh antibodies do not agglutinate red blood cells but their presence on the fetal red blood cell surface can be shown by washing away unbound immunoglobulin and then adding antibody to human immunoglobulin, which agglutinates the antibody-coated cells. Anti-Rh antibodies can be detected in the mother's serum in an indirect Coombs test; the serum is incubated with Rh⁺ red blood cells, and once the antibody binds, the red blood cells are treated as in the direct Coombs test.

Since the Rh antigens are widely spaced on the red blood cell surface, the IgG anti-Rh antibodies cannot fix complement and cause lysis of red blood cells *in vitro*. Furthermore, for reasons that are not fully understood, antibodies to Rh blood group antigens do not agglutinate red blood cells as do antibodies to the ABO blood group antigens. Thus, detecting these antibodies was difficult until anti-human immunoglobulin antibodies were developed. With these, maternal IgG antibodies bound to the fetal red blood cells can be detected after washing the cells to remove unbound immunoglobulin that is present in the fetal serum. Adding anti-human immunoglobulin antibodies to the washed fetal red blood cells agglutinates any cells to which maternal antibodies are bound. This is the **direct Coombs test** (Fig. A.13), so called because it directly detects antibody bound to the surface of the fetal red blood cells. An **indirect Coombs test** is used to detect nonagglutinating anti-Rh antibody in maternal serum; the serum is first incubated with Rh-positive red blood cells, which bind the anti-Rh antibody, after which the antibody-coated cells are washed to remove unbound immunoglobulin and are then agglutinated with anti-immunoglobulin antibody (see Fig. A.13). The indirect Coombs test allows Rh incompatibilities that might lead to hemolytic disease of the newborn to be detected, and this knowledge allows the disease to be prevented (see Section 10-25). The Coombs test is also commonly used to detect antibodies to drugs that bind to red blood cells and cause hemolytic anemia.

A-12 Monoclonal antibodies.

The antibodies generated in a natural immune response or after immunization in the laboratory are a mixture of molecules of different specificities and affinities. Some of this heterogeneity results from the production of antibodies that bind to different epitopes on the immunizing antigen, but even antibodies directed at a single antigenic determinant such as a hapten can be markedly heterogeneous, as shown by **isoelectric focusing**. In this technique, proteins are separated on the basis of their isoelectric point, the pH at which their net charge is zero. By electrophoresing proteins in a pH gradient for long enough, each molecule migrates along the pH gradient until it reaches the pH at which it is neutral and is thus concentrated (focused) at that point. When antiserum containing anti-hapten antibodies is treated in this way and then transferred to a solid support such as nitrocellulose paper, the anti-hapten antibodies can be detected by their ability to bind labeled hapten. The binding of antibodies of various isoelectric points to the hapten shows that even antibodies that bind the same antigenic determinant can be heterogeneous.

Antisera are valuable for many biological purposes but they have certain inherent disadvantages that relate to the heterogeneity of the antibodies they contain. First, each antiserum is different from all other antisera, even if

raised in a genetically identical animal by using the identical preparation of antigen and the same immunization protocol. Second, antisera can be produced in only limited volumes, and thus it is impossible to use the identical serological reagent in a long or complex series of experiments or clinical tests. Finally, even antibodies purified by affinity chromatography (see Section A-5) may include minor populations of antibodies that give unexpected cross-reactions, which confound the analysis of experiments. To avoid these problems, and to harness the full potential of antibodies, it was necessary to develop a way of making an unlimited supply of antibody molecules of homogeneous structure and known specificity. This has been achieved through the production of monoclonal antibodies from cultures of hybrid antibody-forming cells or, more recently, by genetic engineering.

Biochemists in search of a homogeneous preparation of antibody that they could subject to detailed chemical analysis turned early to proteins produced by patients with multiple myeloma, a common tumor of plasma cells. It was known that antibodies are normally produced by plasma cells and since this disease is associated with the presence of large amounts of a homogeneous gamma globulin called a **myeloma protein** in the patient's serum, it seemed likely that myeloma proteins would serve as models for normal antibody molecules. Thus, much of the early knowledge of antibody structure came from studies on myeloma proteins. These studies showed that monoclonal antibodies could be obtained from immortalized plasma cells. However, the antigen specificity of most myeloma proteins was unknown, which limited their usefulness as objects of study or as immunological tools.

This problem was solved by Georges Köhler and Cesar Milstein, who devised a technique for producing a homogeneous population of antibodies of known antigenic specificity. They did this by fusing spleen cells from an immunized mouse to cells of a mouse myeloma to produce hybrid cells that both proliferated indefinitely and secreted antibody specific for the antigen used to immunize the spleen cell donor. The spleen cell provides the ability to make specific antibody, while the myeloma cell provides the ability to grow indefinitely in culture and secrete immunoglobulin continuously. By using a myeloma cell partner that produces no antibody proteins itself, the antibody produced by the hybrid cells comes only from the immune spleen cell partner. After fusion, the hybrid cells are selected using drugs that kill the myeloma parental cell, while the unfused parental spleen cells have a limited life-span and soon die, so that only hybrid myeloma cell lines or **hybridomas** survive. Those hybridomas producing antibody of the desired specificity are then identified and cloned by regrowing the cultures from single cells (Fig. A.14). Since each hybridoma is a **clone** derived from fusion with a single B cell, all the antibody molecules it produces are identical in structure, including their antigen-binding site and isotype. Such antibodies are called **monoclonal antibodies**. This technology has revolutionized the use of antibodies by providing a limitless supply of antibody of a single and known specificity. Monoclonal antibodies are now used in most serological assays, as diagnostic probes, and as therapeutic agents. So far, however, only mouse monoclonals are routinely produced and efforts to use this same approach to make human monoclonal antibodies have met with very limited success.

A-13 Phage display libraries for antibody V-region production.

This is a technique for producing antibody-like molecules. Gene segments encoding the antigen-binding variable or V domains of antibodies are fused to genes encoding the coat protein of a bacteriophage. Bacteriophage containing such gene fusions are used to infect bacteria, and the resulting phage particles have coats that express the antibody-like fusion protein, with