



MOLECULAR BIOLOGY OF  
THE CELL

ALBERTS JOHNSON LEWIS RAFF ROBERTS WALTER

F O U R T H E D I T I O N

## The Molecular Biology of the Cell

### [Acknowledgments](#)

### [Preface](#)

### [A Note to the Reader](#)

## I. [Introduction to the Cell](#)

### 1. [Cells and Genomes](#)

[The Universal Features of Cells on Earth](#)

[The Diversity of Genomes and the Tree of Life](#)

[Genetic Information in Eucaryotes](#)

[References](#)

### 2. [Cell Chemistry and Biosynthesis](#)

[The Chemical Components of a Cell](#)

[Catalysis and the Use of Energy by Cells](#)

[How Cells Obtain Energy from Food](#)

[References](#)

### 3. [Proteins](#)

[The Shape and Structure of Proteins](#)

[Protein Function](#)

[References](#)

## II. [Basic Genetic Mechanisms](#)

### 4. [DNA and Chromosomes](#)

[The Structure and Function of DNA](#)

[Chromosomal DNA and Its Packaging in the Chromatin Fiber](#)

[The Global Structure of Chromosomes](#)

[References](#)

5. [DNA Replication, Repair, and Recombination](#)

[The Maintenance of DNA Sequences](#)

[DNA Replication Mechanisms](#)

[The Initiation and Completion of DNA Replication in Chromosomes](#)

[DNA Repair](#)

[General Recombination](#)

[Site-Specific Recombination](#)

[References](#)

6. [How Cells Read the Genome: From DNA to Protein](#)

[From DNA to RNA](#)

[From RNA to Protein](#)

[The RNA World and the Origins of Life](#)

[References](#)

7. [Control of Gene Expression](#)

[An Overview of Gene Control](#)

[DNA-Binding Motifs in Gene Regulatory Proteins](#)

[How Genetic Switches Work](#)

[The Molecular Genetic Mechanisms That Create Specialized Cell Types](#)

[Posttranscriptional Controls](#)

[How Genomes Evolve](#)

[References](#)

### III. [Methods](#)

#### 8. [Manipulating Proteins, DNA, and RNA](#)

[Isolating Cells and Growing Them in Culture](#)

[Fractionation of Cells](#)

[Isolating, Cloning, and Sequencing DNA](#)

[Analyzing Protein Structure and Function](#)

[Studying Gene Expression and Function](#)

[References](#)

#### 9. [Visualizing Cells](#)

[Looking at the Structure of Cells in the Microscope](#)

[Visualizing Molecules in Living Cells](#)

[References](#)

### IV. [Internal Organization of the Cell](#)

#### 10. [Membrane Structure](#)

[The Lipid Bilayer](#)

[Membrane Proteins](#)

[References](#)

#### 11. [Membrane Transport of Small Molecules and the Electrical Properties of Membranes](#)

[Principles of Membrane Transport](#)

[Carrier Proteins and Active Membrane Transport](#)

[Ion Channels and the Electrical Properties of Membranes](#)

[References](#)

12. [Intracellular Compartments and Protein Sorting](#)

[The Compartmentalization of Cells](#)

[The Transport of Molecules between the Nucleus and the Cytosol](#)

[The Transport of Proteins into Mitochondria and Chloroplasts](#)

[Peroxisomes](#)

[The Endoplasmic Reticulum](#)

[References](#)

13. [Intracellular Vesicular Traffic](#)

[The Molecular Mechanisms of Membrane Transport and the Maintenance of Compartmental Diversity](#)

[Transport from the ER through the Golgi Apparatus](#)

[Transport from the \*Trans\* Golgi Network to Lysosomes](#)

[Transport into the Cell from the Plasma Membrane: Endocytosis](#)

[Transport from the \*Trans\* Golgi Network to the Cell Exterior: Exocytosis](#)

[References](#)

14. [Energy Conversion: Mitochondria and Chloroplasts](#)

[The Mitochondrion](#)

[Electron-Transport Chains and Their Proton Pumps](#)

[Chloroplasts and Photosynthesis](#)

[The Genetic Systems of Mitochondria and Plastids](#)

[The Evolution of Electron-Transport Chains](#)

[References](#)

15. [Cell Communication](#)

[General Principles of Cell Communication](#)

[Signaling through G-Protein-Linked Cell-Surface Receptors](#)

[Signaling through Enzyme-Linked Cell-Surface Receptors](#)

[Signaling Pathways That Depend on Regulated Proteolysis](#)

[Signaling in Plants](#)

[References](#)

16. [The Cytoskeleton](#)

[The Self-Assembly and Dynamic Structure of Cytoskeletal Filaments](#)

[How Cells Regulate Their Cytoskeletal Filaments](#)

[Molecular Motors](#)

[The Cytoskeleton and Cell Behavior](#)

[References](#)

17. [The Cell Cycle and Programmed Cell Death](#)

[An Overview of the Cell Cycle](#)

[Components of the Cell-Cycle Control System](#)

[Intracellular Control of Cell-Cycle Events](#)

[Programmed Cell Death \(Apoptosis\)](#)

[Extracellular Control of Cell Division, Cell Growth, and Apoptosis](#)

[References](#)

18. [The Mechanics of Cell Division](#)

[An Overview of M Phase](#)

[Mitosis](#)

[Cytokinesis](#)

[References](#)

V. [Cells in Their Social Context](#)

19. [Cell Junctions, Cell Adhesion, and the Extracellular Matrix](#)

[Cell Junctions](#)

[Cell-Cell Adhesion](#)

[The Extracellular Matrix of Animals](#)

[Integrins](#)

[The Plant Cell Wall](#)

[References](#)

20. [Germ Cells and Fertilization](#)

[The Benefits of Sex](#)

[Meiosis](#)

[Primordial Germ Cells and Sex Determination in Mammals](#)

[Eggs](#)

[Sperm](#)

[Fertilization](#)

[References](#)

## 21. [Development of Multicellular Organisms](#)

[Universal Mechanisms of Animal Development](#)

[Caenorhabditis Elegans: Development from the Perspective of the Individual Cell](#)

[Drosophila and the Molecular Genetics of Pattern Formation: Genesis of the Body Plan](#)

[Homeotic Selector Genes and the Patterning of the Anteroposterior Axis](#)

[Organogenesis and the Patterning of Appendages](#)

[Cell Movements and the Shaping of the Vertebrate Body](#)

[The Mouse](#)

[Neural Development](#)

[Plant Development](#)

[References](#)

## 22. [Histology: The Lives and Deaths of Cells in Tissues](#)

[Epidermis and Its Renewal by Stem Cells](#)

[Sensory Epithelia](#)

[The Airways and the Gut](#)

[Blood Vessels and Endothelial Cells](#)

[Renewal by Multipotent Stem Cells: Blood Cell Formation](#)

[Genesis, Modulation, and Regeneration of Skeletal Muscle](#)

[Fibroblasts and Their Transformations: The Connective-Tissue Cell Family](#)

[Stem-Cell Engineering](#)

[References](#)

## 23. [Cancer](#)

[Cancer as a Microevolutionary Process](#)

[The Preventable Causes of Cancer](#)

[Finding the Cancer-Critical Genes](#)

[The Molecular Basis of Cancer-Cell Behavior](#)

[Cancer Treatment: Present and Future](#)

[References](#)

## 24. [The Adaptive Immune System](#)

[Lymphocytes and the Cellular Basis of Adaptive Immunity](#)

[B Cells and Antibodies](#)

[The Generation of Antibody Diversity](#)

[T Cells and MHC Proteins](#)

[Helper T Cells and Lymphocyte Activation](#)

[References](#)

## 25. [Pathogens, Infection, and Innate Immunity](#)

[Introduction to Pathogens](#)

[Cell Biology of Infection](#)

[Innate Immunity](#)

[References](#)

[Glossary](#)

© 2002 by Bruce Alberts, Alexander Johnson, Julian Lewis, Martin Raff, Keith Roberts, and Peter Walter.

■

|                                                                                   |                                         |                                                          |                                                                                                                                                           |
|-----------------------------------------------------------------------------------|-----------------------------------------|----------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------|
|  | MOLECULAR BIOLOGY OF<br><b>THE CELL</b> | ALBERTS<br>JOHNSON<br>LEWIS<br>RAFF<br>ROBERTS<br>WALTER | <br><a href="http://www.garlandscience.com">www.garlandscience.com</a> |
|                                                                                   | FOURTH EDITION                          |                                                          |                                                                                                                                                           |

[Short Contents](#) | [Full Contents](#)

[Other books @ NCBI](#)

| Navigation                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                               |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <a href="#">About this book</a><br><b>I. Introduction to the Cell</b><br>1. Cells and Genomes<br>The Universal Features of Cells on Earth<br>The Diversity of Genomes and the Tree of Life<br><b>➔ Genetic Information in Eucaryotes</b><br><a href="#">References</a>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
| Figures                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
|  <a href="#">Figure 1-31. The major features of eucaryotic...</a><br> <a href="#">Figure 1-32. Phagocytosis.</a><br> <a href="#">Figure 1-33. A single-celled eucaryote that eats...</a><br> <a href="#">Figure 1-34. A mitochondrion.</a><br> <a href="#">Figure 1-35. The origin of mitochondria.</a><br> <a href="#">Figure 1-36. Chloroplasts.</a><br> <a href="#">Figure 1-37. The origin of chloroplasts.</a> |

***Molecular Biology of the Cell* ➔ I. Introduction to the Cell ➔ 1. Cells and Genomes**

## Genetic Information in Eucaryotes

Eucaryotic cells, in general, are bigger and more elaborate than procaryotic cells, and their genomes are bigger and more elaborate, too. The greater size is accompanied by radical differences in cell structure and function. Moreover, many classes of eucaryotic cells form multicellular organisms that attain a level of complexity unmatched by any procaryote.

Because they are so complex, eucaryotes confront molecular biologists with a special set of challenges, which will concern us in the rest of this book. Increasingly, biologists meet these challenges through the analysis and manipulation of the genetic information within cells and organisms. It is therefore important at the outset to know something of the special features of the eucaryotic genome. We begin by briefly reviewing how eucaryotic cells are organized, how this reflects their way of life, and how their genomes differ from those of procaryotes. This leads us to an outline of the strategy by which molecular biologists, by exploiting genetic information, are attempting to discover how eucaryotic organisms work.

### Eucaryotic Cells May Have Originated as Predators

By definition, eucaryotic cells keep their DNA in a separate internal compartment, the nucleus. The DNA is separated from the cytoplasm by the *nuclear envelope*, which consists of a double layer of membrane. Eucaryotes also have other features that set them apart from procaryotes ([Figure 1-31](#)). Their cells are, typically, 10 times bigger in linear dimension, and 1000 times larger in volume. They have a *cytoskeleton*—a system of protein filaments crisscrossing the cytoplasm and forming, together with the many proteins that attach to them, a system of girders, ropes, and motors that gives the cell mechanical strength, controls its shape, and drives and guides its movements. The nuclear envelope is only one part of an elaborate set of *internal membranes*, each structurally similar to the plasma membrane and enclosing different types of spaces inside the cell, many of them involved in processes related to digestion and secretion. Lacking the tough cell wall of most bacteria, animal cells and the free-living eucaryotic cells called *protozoa* can change their shape rapidly and engulf other cells and small objects by *phagocytosis*

[Figure 1-38.](#) [Genome sizes compared.](#)[Figure 1-39. The puffer fish \(\*Fugu rubripes\*\)....](#)[Figure 1-40.](#) [Controlling gene readout by environmental...](#)[Figure 1-41. Genetic control of the program...](#)[Figure 1-42. An assortment of protists: a...](#)[Figure 1-43. The yeast \*Saccharomyces cerevisiae\*....](#)[Figure 1-44. The reproductive cycles of the...](#)[Figure 1-45.](#) [Monitoring changes in yeast gene...](#)[Figure 1-46.](#) [Arabidopsis thaliana, the plant chosen...](#)[Figure 1-47.](#) [Caenorhabditis elegans, the first multicellular...](#)[Figure 1-48.](#) [Drosophila melanogaster....](#)[Figure 1-49. Giant chromosomes from salivary gland...](#) [Figure 1-50. Two species of the frog...](#)[\(Figure 1-32\).](#)

It is still a mystery how all these properties evolved, and in what sequence. One plausible view, however, is that they are all reflections of the way of life of a primordial eucaryotic cell that was a predator, living by capturing other cells and eating them (Figure 1-33). Such a way of life requires a large cell with a flexible plasma membrane, as well as an elaborate cytoskeleton to support and move this membrane. It may also require that the cell's long, fragile DNA molecules be sequestered in a separate nuclear compartment, to protect the genome from damage by the movements of the cytoskeleton. [↑ TOP](#)

## Eucaryotic Cells Evolved from a Symbiosis

A predatory way of life helps to explain another feature of eucaryotic cells. Almost all such cells also contain *mitochondria* (Figure 1-34). These are small bodies in the cytoplasm, enclosed by a double layer of membrane, that take up oxygen and harness energy from the oxidation of food molecules—such as sugars—to produce most of the ATP that powers the cell's activities. Mitochondria are similar in size to small bacteria, and, like bacteria, they have their own genome in the form of a circular DNA molecule, their own ribosomes that are different from those elsewhere in the eucaryotic cell, and their own transfer RNAs. It is virtually certain that mitochondria originated from free-living oxygen-metabolizing (*aerobic*) eubacteria that were engulfed by an ancestral eucaryotic cell that could otherwise make no such use of oxygen (that is, was *anaerobic*). Escaping digestion, these bacteria evolved in symbiosis with the engulfing cell and its progeny, receiving shelter and nourishment in return for the power generation they performed for their hosts (Figure 1-35). This partnership between a primitive anaerobic eucaryotic predator cell and an aerobic bacterial cell is thought to have been established about 1.5 billion years ago, when the Earth's atmosphere first became rich in oxygen.

Many eucaryotic cells—specifically, those of plants and algae—also contain another class of small membrane-bounded organelles somewhat similar to mitochondria—the *chloroplasts* (Figure 1-36). Chloroplasts perform photosynthesis, using the energy of sunlight to synthesize carbohydrates from atmospheric carbon dioxide and water, and deliver the products to the host cell as food. Like mitochondria, chloroplasts have their own genome and almost certainly originated as symbiotic photosynthetic bacteria, acquired by cells that already possessed mitochondria (Figure 1-37).

A eucaryotic cell equipped with chloroplasts has no need to chase after other cells as prey; it is nourished by the captive chloroplasts it has inherited from its ancestors. Correspondingly, plant cells, although they possess the cytoskeletal equipment for movement, have lost the ability to change shape rapidly and to engulf other cells by phagocytosis. Instead, they create around themselves a tough, protective cell wall. If the ancestral eucaryote was indeed a predator on

[Figure 1-51. The consequences of gene duplication...](#)

[Figure 1-52. Times of divergence of different...](#)

[Figure 1-53. Human and mouse: similar genes...](#)

other organisms, we can view plant cells as eucaryotes that have made the transition from hunting to farming.

Fungi represent yet another eucaryotic way of life. Fungal cells, like animal cells, possess mitochondria but not chloroplasts; but in contrast with animal cells and protozoa, they have a tough outer wall that limits their ability to move rapidly or to swallow up other cells. Fungi, it seems, have turned from hunters into scavengers: other cells secrete nutrient molecules or release them upon death, and fungi feed on these leavings—performing whatever digestion is necessary extracellularly, by secreting digestive enzymes to the exterior. [↑ TOP](#)

Search

Go

☒ This book
 ☐ All books

☐ PubMed

## Eucaryotes Have Hybrid Genomes

The genetic information of eucaryotic cells has a hybrid origin—from the ancestral anaerobic eucaryote, and from the bacteria that it adopted as symbionts. Most of this information is stored in the nucleus, but a small amount remains inside the mitochondria and, for plant and algal cells, in the chloroplasts. The mitochondrial DNA and the chloroplast DNA can be separated from the nuclear DNA and individually analyzed and sequenced. The mitochondrial and chloroplast genomes are found to be degenerate, cut-down versions of the corresponding bacterial genomes, lacking genes for many essential functions. In a human cell, for example, the mitochondrial genome consists of only 16,569 nucleotide pairs, and codes for only 13 proteins, two ribosomal RNA components, and 22 transfer RNAs.

The genes that are missing from the mitochondria and chloroplasts have not all been lost; instead, many of them have been somehow moved from the symbiont genome into the DNA of the host cell nucleus. The nuclear DNA of humans contains many genes coding for proteins that serve essential functions inside the mitochondria; in plants, the nuclear DNA also contains many genes specifying proteins required in chloroplasts. [↑ TOP](#)

## Eucaryotic Genomes Are Big

Natural selection has evidently favored mitochondria with small genomes, just as it has favored bacteria with small genomes. By contrast, the nuclear genomes of most eucaryotes seem to have been free to enlarge. Perhaps the eucaryotic way of life has made large size an advantage: predators need to be bigger than their prey, and cell size generally increases in proportion to genome size. Whatever the explanation, the fact is that the genomes of most eucaryotes are orders of magnitude larger than those of bacteria and archaea (Figure 1-38). And the freedom to be extravagant with DNA has had profound implications.

Eucaryotes not only have more genes than procaryotes; they also have vastly more DNA that does not code for protein or for any other functional product

molecule. The human genome contains a 1000 times as many nucleotide pairs as the genome of a typical bacterium, 20 times as many genes, and about 10,000 times as much noncoding DNA (~98.5% of the genome for a human is noncoding, as opposed to 11% of the genome for the bacterium *E. coli*). [↑ TOP](#)

## Eucaryotic Genomes Are Rich in Regulatory DNA

Much of our noncoding DNA is almost certainly dispensable junk, retained like a mass of old papers because, when there is little pressure to keep an archive small, it is easier to retain everything than to sort out the valuable information and discard the rest. Certain exceptional eucaryotic species, such as the puffer fish ([Figure 1-39](#)), bear witness to the profligacy of their relatives; they have somehow managed either to rid themselves of large quantities of noncoding DNA, or to have avoided acquiring it in the first place. Yet they appear similar in structure, behavior, and fitness to related species that have vastly more such DNA.

Even in compact eucaryotic genomes such as that of puffer fish, there is more noncoding DNA than coding DNA, and at least some of the noncoding DNA certainly has important functions. In particular, it serves to regulate the expression of adjacent genes. With this regulatory DNA, eucaryotes have evolved distinctive ways of controlling when and where a gene is brought into play. This sophisticated gene regulation is crucial for the formation of complex multicellular organisms. [↑ TOP](#)

## The Genome Defines the Program of Multicellular Development

The cells in an individual animal or plant are extraordinarily varied. Fat cells, skin cells, bone cells, nerve cells—they seem as dissimilar as any cells could be. Yet all these cell types are generated during embryonic development from a single fertilized egg cell, and all (with minor exceptions) contain identical copies of the genome of the species.

The explanation lies in the way in which these cells make selective use of their genetic instructions according to the cues they get from their surroundings. The DNA is not just a shopping list specifying the molecules that every cell must have, and the cell is not an assembly of all the items on the list. Rather, the cell behaves as a multipurpose machine, with sensors to receive environmental signals and highly developed abilities to call different sets of genes into action according to the sequences of signals to which the cell has been exposed. A large fraction of the genes in the eucaryotic genome code for proteins that serve to regulate the activities of other genes. Most of these *gene regulatory proteins* act by binding, directly or indirectly, to the regulatory DNA adjacent to the genes that are to be controlled ([Figure 1-40](#)), or by interfering with the abilities of other proteins to do so. The expanded genome of eucaryotes therefore serves not only to specify the hardware of the cell, but

also to store the software that controls how that hardware is used.

In a developing multicellular organism, each cell is governed by the same control system, but with different consequences depending on the signals the cell exchanges with its neighbors. The outcome, astonishingly, is a precisely patterned array of cells in different states, each displaying a character appropriate to its position in the multicellular structure. The genome in each cell is big enough to accommodate the information that specifies an entire multicellular organism, but in any individual cell only part of that information is used (Figure 1-41). [↑ TOP](#)

## Many Eucaryotes Live as Solitary Cells: the Protists

Many species of eucaryotic cells lead a solitary life—some as hunters (the *protozoa*), some as photosynthesizers (the unicellular *algae*), some as scavengers (the unicellular fungi, or *yeasts*). Even though they do not form multicellular organisms, these single-celled eucaryotes, or *protists*, can be very complex. Figure 1-42 conveys something of the variety of forms they can take. The anatomy of protozoa, especially, is often elaborate and includes such structures as sensory bristles, photoreceptors, sinuously beating cilia, leglike appendages, mouth parts, stinging darts, and musclelike contractile bundles. Although they are single cells, protozoa can be as intricate, as versatile, and as complex in their behavior as many multicellular organisms (see Figure 1-33).

In terms of their ancestry and DNA sequences, protists are far more diverse than the multicellular animals, plants, and fungi, which arose as three comparatively late branches of the eucaryotic pedigree (see Figure 1-21). As with procaryotes, humans have tended to neglect the protists because they are microscopic. Only now, with the help of genome analysis, are we beginning to understand their positions in the tree of life, and to put into context the glimpses these strange creatures offer us of our distant evolutionary past. [↑ TOP](#)

## A Yeast Serves as a Minimal Model Eucaryote

The molecular and genetic complexity of eucaryotes is daunting. Even more than for procaryotes, biologists need to concentrate their limited resources on a few selected model organisms to fathom this complexity.

To analyze the internal workings of the eucaryotic cell, without being distracted by the additional problems of multicellular development, it makes sense to use a species that is unicellular and as simple as possible. The popular choice for this role of minimal model eucaryote has been the yeast *Saccharomyces cerevisiae* (Figure 1-43)—the same species that is used by brewers of beer and bakers of bread.

*S. cerevisiae* is a small, single-celled member of the kingdom of fungi and thus, according to modern views, at least as closely related to animals as it is to plants. It is robust and easy to grow in a simple nutrient medium. Like other fungi, it has a tough cell wall, is relatively immobile, and possesses mitochondria but not chloroplasts. When nutrients are plentiful, it grows and divides almost as rapidly as a bacterium. It can reproduce either vegetatively (that is, by simple cell division), or sexually: two yeast cells that are *haploid* (possessing a single copy of the genome) can fuse to create a cell that is *diploid* (containing a double genome); and the diploid cell can undergo *meiosis* (a reduction division) to produce cells that are once again haploid (Figure 1-44). In contrast with higher plants and animals, the yeast can divide indefinitely in either the haploid or the diploid state, and the process leading from the one state to the other can be induced at will by changing the growth conditions.

All these features make this yeast an extremely convenient organism for genetic studies, as does one further property: its genome, by eucaryotic standards, is exceptionally small. Nevertheless, it suffices for all the basic tasks that every eucaryotic cell must perform. As we shall see later in this book, studies on yeasts (using both *S. cerevisiae* and other species) have provided a key to the understanding of many crucial processes. These include the eucaryotic cell-division cycle—the critical chain of events by which the nucleus and all the other components of a cell are duplicated and parceled out to create two daughter cells from one. The control system that governs this process has been so well conserved over the course of evolution that many of its components can function interchangeably in yeast and human cells: if a mutant yeast lacking an essential yeast cell-division-cycle gene is supplied with a copy of the homologous cell-division-cycle gene from a human, the yeast is cured of its defect and becomes able to divide normally.  [TOP](#)

## The Expression Levels of All The Genes of An Organism Can Be Monitored Simultaneously

The complete genome sequence of *S. cerevisiae* was determined in 1997. It consists of approximately 13,117,000 nucleotide pairs, including the small contribution (78,520 nucleotide pairs) of the mitochondrial DNA. This total is only about 2.5 times as much DNA as there is in *E. coli*, and it codes for only 1.5 times as many distinct proteins (about 6300 in all). The way of life of *S. cerevisiae* is similar in many ways to that of a bacterium, and it seems that this yeast has likewise been subject to selection pressures that have kept its genome compact.

Knowledge of the complete genome sequence of any organism—be it a yeast or a human—opens up new perspectives on the workings of the cell: things that once seemed impossibly complex now seem within our grasp. One example will suffice. It is possible to array on a glass microscope slide a set of samples of DNA sequence corresponding to each of the ~6300 proteins the

yeast has encoded in its genome. A DNA molecule will bind selectively, by base-pairing, with DNA or RNA that has a complementary sequence. Therefore, if a mixture of nucleic acid molecules is passed over such a *DNA array*, each type of molecule in the mixture will stick to its corresponding DNA spot. If the molecules in the mixture have a fluorescent chemical tag attached to them, the result will be a set of spots of fluorescence, each localized at a specific point in the array (Figure 1-45). In this way, it is possible to monitor, simultaneously, the amount of mRNA transcript that is produced from every gene in the genome by a yeast under any chosen conditions, and to see how this whole pattern of gene activity changes when conditions change. The analysis can be repeated with mRNA prepared from mutants lacking a chosen gene—any gene that we care to test. In principle, this approach provides a way to reveal the entire system of control relationships that govern gene expression—not only in these yeast cells, but in any organism whose genome sequence is known. [↑ TOP](#)

## *Arabidopsis* Has Been Chosen Out of 300,000 Species As a Model Plant

The large multicellular organisms that we see around us—the flowers and trees and animals—seem fantastically varied, but they are much closer to one another in their evolutionary origins, and more similar in their basic cell biology, than the great host of microscopic single-celled organisms. Thus, while bacteria and eucaryotes are separated by more than 3000 million years of divergent evolution, vertebrates and insects are separated by about 700 million years, fish and mammals by about 450 million years, and the different species of flowering plants by only about 150 million years.

Because of the close evolutionary relationship between all flowering plants, we can, once again, get insight into the cell and molecular biology of this whole class of organisms by focusing on just one or a few species for detailed analysis. Out of the several hundred thousand species of flowering plants on Earth today, molecular biologists have chosen to concentrate their efforts on a small weed, the common wall cress *Arabidopsis thaliana* (Figure 1-46), which can be grown indoors in large numbers, and produces thousands of offspring per plant after 8–10 weeks. *Arabidopsis* has a genome of approximately 140 million nucleotide pairs, about 11 times as much as yeast, and its complete sequence is known. [↑ TOP](#)

## The World of Animal Cells Is Represented By a Worm, a Fly, a Mouse, and a Human

Multicellular animals account for the majority of all named species of living organisms, and for the largest part of the biological research effort. Four species have emerged as the foremost model organisms for molecular genetic studies and have had their genomes sequenced. In order of increasing size, they are the nematode worm *Caenorhabditis elegans*, the fly *Drosophila*

*melanogaster*, the mouse *Mus musculus*, and the human, *Homo sapiens*.

*Caenorhabditis elegans* (Figure 1-47) is a small, harmless relative of the eelworm that attacks crops. With a life cycle of only a few days, an ability to survive in a freezer indefinitely in a state of suspended animation, a simple body plan, and an unusual life cycle that is well suited for genetic studies (described in Chapter 21), it is an ideal model organism. *C. elegans* develops with clockwork precision from a fertilized egg cell into an adult worm with exactly 959 body cells (plus a variable number of egg and sperm cells)—an unusual degree of regularity for an animal. We now have a minutely detailed description of the sequence of events by which this occurs, as the cells divide, move, and change their characters according to strict and predictable rules. The genome of 97 million nucleotide pairs code for about 19,000 proteins and a wealth of mutants are available for the testing of gene functions. Although the worm has a body plan very different from our own, the conservation of biological mechanisms has been sufficient for the worm to provide a valuable model for many of the processes that occur in the human body. Studies of the worm help us to understand, for example, the programs of cell division and cell death that determine the numbers of cells in the body—a topic of great importance in developmental biology and cancer research. [↑ TOP](#)

## Studies in *Drosophila* Provide a Key to Vertebrate Development

The fruitfly *Drosophila melanogaster* (Figure 1-48) has been in use as a model genetic organism for longer than any other; in fact, the foundations of classical genetics were built to a large extent on studies of this insect. Over 80 years ago, it provided, for example, definitive proof that genes—the abstract units of hereditary information—are carried on chromosomes, concrete physical objects whose behavior had been closely followed in the eucaryotic cell with the light microscope, but whose function was at first unknown. The proof depended on one of the many features that make *Drosophila* peculiarly convenient for genetics—the giant chromosomes, with characteristic banded appearance, that are visible in some of its cells (Figure 1-49). Specific changes in the hereditary information, manifest in families of mutant flies, were found to correlate exactly with the loss or alteration of specific giant-chromosome bands.

In more recent times, *Drosophila*, more than any other organism, has shown us how to trace the chain of cause and effect from the genetic instructions encoded in the chromosomal DNA to the structure of the adult multicellular body. *Drosophila* mutants with body parts strangely misplaced or mispatterned provided the key to the identification and characterization of the genes required to make a properly structured body, with gut, limbs, eyes, and all the other parts in their correct places. Once these *Drosophila* genes were sequenced, the genomes of vertebrates could be scanned for homologs. These were found, and their functions in vertebrates were then tested by analyzing

mice in which the genes had been mutated. The results, as we see later in the book, reveal an astonishing degree of similarity in the molecular mechanisms of insect and vertebrate development.

The majority of all named species of living organisms are insects. Even if *Drosophila* had nothing in common with vertebrates, but only with insects, it would still be an important model organism. But if understanding the molecular genetics of vertebrates is the goal, why not simply tackle the problem head-on? Why sidle up to it obliquely, through studies in *Drosophila*?

*Drosophila* requires only 9 days to progress from a fertilized egg to an adult; it is vastly easier and cheaper to breed than any vertebrate, and its genome is much smaller—about 170 million nucleotide pairs, compared with 3200 million for a human. This codes for about 14,000 proteins, and mutants can now be obtained for essentially any gene. But there is also another, deeper reason why genetic mechanisms that are hard to discover in a vertebrate are often readily revealed in the fly. This relates, as we now explain, to the frequency of gene duplication, which is substantially greater in vertebrate genomes than in the fly genome and has probably been crucial in making vertebrates the complex and subtle creatures that they are. [↑ TOP](#)

## The Vertebrate Genome Is a Product of Repeated Duplication

Almost every gene in the vertebrate genome is found to have paralogs—other genes in the same genome that are unmistakably related and must have arisen by gene duplication. In many cases, a whole cluster of genes is closely related to similar clusters present elsewhere in the genome, suggesting that genes have been duplicated in linked groups rather than as isolated individuals. According to one hypothesis, at an early stage in the evolution of the vertebrates, the entire genome underwent duplication twice in succession, giving rise to four copies of every gene. In some groups of vertebrates, such as fish of the salmon and carp families (including the zebrafish, a popular research animal), it has been suggested that there was yet another duplication, creating an eightfold multiplicity of genes.

The precise course of vertebrate genome evolution remains uncertain, because many further evolutionary changes have occurred since these ancient events. Genes that were once identical have diverged; many of the gene copies have been lost through disruptive mutations; some have undergone further rounds of local duplication; and the genome, in each branch of the vertebrate family tree, has suffered repeated rearrangements, breaking up most of the original gene orderings. A careful comparison of the gene order in two related organisms, such as the human and the mouse, reveals that—on the time scale of vertebrate evolution—chromosomes frequently fuse and fragment to move large blocks of DNA sequence around. Indeed, it is entirely possible, as we shall discuss in [Chapter 7](#), that the present state of affairs is the result of many separate

duplications of fragments of the genome, rather than duplications of the genome as a whole.

There is, however, no doubt that such whole-genome duplications do occur from time to time in evolution, for we can see recent instances in which duplicated chromosome sets are still clearly identifiable as such. The frog genus *Xenopus*, for example, comprises a set of closely similar species related to one another by repeated duplications or triplications of the whole genome. Among these frogs are *X. tropicalis*, with an ordinary diploid genome; the common laboratory species *X. laevis*, with a duplicated genome and twice as much DNA per cell; and *X. ruwenzoriensis*, with a sixfold reduplicated genome and six times as much DNA per cell (108 chromosomes, compared with 36 in *X. laevis*, for example). These species are estimated to have diverged from one another within the past 120 million years ([Figure 1-50](#)). [↑ TOP](#)

## Genetic Redundancy Is a Problem for Geneticists, But It Creates Opportunities for Evolving Organisms

Whatever the details of the evolutionary history, it is clear that most genes in the vertebrate genome exist in several versions that were once identical. The related genes often remain functionally interchangeable for many purposes. This phenomenon is called **genetic redundancy**. For the scientist struggling to discover all the genes involved in some particular process, it has a dire consequence. If gene A is mutated and no effect is seen, one can no longer conclude that gene A is functionally irrelevant—it may simply be that this gene normally works in parallel with its relatives, and these suffice for near-normal function even when gene A is defective (just as an airliner continues to fly when one of its engines fails). In the less repetitive genome of *Drosophila*, where gene duplication is less common, the analysis is more straightforward: single gene functions are revealed directly by the consequences of single-gene mutations (the single-engined plane stops flying when the engine fails).

Genome duplication has clearly allowed the development of more complex life forms; it provides an organism with a cornucopia of spare gene copies, which are free to mutate to serve divergent purposes. While one copy becomes optimized for use in the liver, say, another can become optimized for use in the brain or adapted for a novel purpose. In this way, the additional genes allow for increased complexity and sophistication. As the genes take on divergent functions, they cease to be redundant. Often, however, while the genes acquire individually specialized roles, they also continue to perform some aspects of their original core function in parallel, redundantly. Mutation of a single gene then causes a relatively minor abnormality that reveals only a part of the gene's function ([Figure 1-51](#)). Families of genes with divergent but partly overlapping functions are a pervasive feature of vertebrate molecular biology, and they are encountered repeatedly in this book. [↑ TOP](#)

## The Mouse Serves as a Model for Mammals

Mammals have typically three or four times as many genes as *Drosophila*, a genome that is 20 times larger, and millions or billions of times as many cells in their adult bodies. In terms of genome size and function, cell biology, and molecular mechanisms, mammals are nevertheless a highly uniform group of organisms. Even anatomically, the differences among mammals are chiefly a matter of size and proportions; it is hard to think of a human body part that does not have a counterpart in elephants and mice, and vice versa. Evolution plays freely with quantitative features, but it does not readily change the logic of the structure.

To get a more exact measure of how closely mammalian species resemble one another genetically, we can compare the nucleotide sequences of corresponding (orthologous) genes, or the amino acid sequences of the proteins that these genes encode. The results for individual genes and proteins vary widely. But typically, if we line up the amino acid sequence of a human protein with that of the orthologous protein from, say, an elephant, about 85% of the amino acids are identical. A similar comparison between human and bird shows an amino acid identity of about 70%—twice as many differences, because the bird and the mammalian lineages have had twice as long to diverge as those of the elephant and the human ([Figure 1-52](#)).

The mouse, being small, hardy, and a rapid breeder, has become the foremost model organism for experimental studies of vertebrate molecular genetics. Many naturally occurring mutations are known, often mimicking the effects of corresponding mutations in humans ([Figure 1-53](#)). Methods have been developed, moreover, to test the function of any chosen mouse gene, or of any non-coding portion of the mouse genome, by artificially creating mutations in it, as we explain later in the book.

One made-to-order mutant mouse can provide a wealth of information for the cell biologist. It reveals the effects of the chosen mutation in a host of different contexts, simultaneously testing the action of the gene in all the different kinds of cells in the body that could in principle be affected. [↑ TOP](#)

## Humans Report on Their Own Peculiarities

As humans, we have, of course, a special interest in the human genome. We want to know the full set of parts from which we ourselves are made, and to discover how they work. But even if you were a mouse, preoccupied with the molecular biology of mice, humans would be attractive as model genetic organisms, because of one special property: through medical examinations and self-reporting, we catalog our own genetic (and other) disorders. The human population is enormous, consisting today of some 6 billion individuals, and this self-documenting property means that a huge database of information is available on human mutations. The complete human genome sequence of more

than 3 billion nucleotide pairs has recently been determined in a rough draft form, making it easier than ever before to identify at a molecular level the precise gene responsible for each human mutant characteristic.

By drawing together the insights from humans, mice, flies, worms, yeasts, plants, and bacteria—using gene sequence similarities to map out the correspondences between one model organism and another—we enrich our understanding of them all. [↑ TOP](#)

## We Are All Different in Detail

What precisely do we mean when we speak of *the* human genome? Whose genome? On average, any two people taken at random differ in about one or two in every 1000 nucleotide pairs in their DNA sequence. The Human Genome Project has arbitrarily selected DNA from a small number of anonymous individuals for sequencing. The human genome—the genome of the human species—is, properly speaking, a more complex thing, embracing the entire pool of variant genes that are found in the human population and continually exchanged and reassorted in the course of sexual reproduction. Ultimately, we can hope to document this variation too. Knowledge of it will help us understand, for example, why some people are prone to one disease, others to another; why some respond well to a drug, others badly. It will also provide new clues to our history—the population movements and minglings of our ancestors, the infections they suffered, the diets they ate. All these things leave traces in the variant forms of genes that have survived in human communities.

Knowledge and understanding bring the power to intervene—with humans, to avoid or prevent disease; with plants, to create better crops; with bacteria, to turn them to our own uses. All these biological enterprises are linked, because the genetic information of all living organisms is written in the same language. The new-found ability of molecular biologists to read and decipher this language has already begun to transform our relationship to the living world. The cell biology to be presented in the subsequent chapters will, we hope, prepare you to understand, and possibly to contribute to, the great scientific adventure of the twenty-first century. [↑ TOP](#)

## Summary

Eucaryotic cells, by definition, keep their DNA in a separate membrane-bounded compartment, the nucleus. They have, in addition, a cytoskeleton for movement, elaborate intracellular compartments for digestion and secretion, the capacity (in many species) to engulf other cells, and a metabolism that depends on the oxidation of organic molecules by mitochondria. These properties suggest that eucaryotes originated as predators on other cells. Mitochondria—and, in plants, chloroplasts—contain their own genetic material, and evidently evolved from bacteria that were taken up into the

cytoplasm of the eucaryotic cell and survived as symbionts. Eucaryotic cells have typically 3–30 times as many genes as procaryotes, and often thousands of times more noncoding DNA. The noncoding DNA allows for complex regulation of gene expression, as required for the construction of complex multicellular organisms. Many eucaryotes are, however, unicellular, among them the yeast *Saccharomyces cerevisiae*, which serves as a simple model organism for eucaryotic cell biology, revealing the molecular basis of conserved fundamental processes such as the eucaryotic cell division cycle. A small number of other organisms have been chosen as primary models for multicellular plants and animals, and the sequencing of their entire genomes has opened the way to systematic and comprehensive analysis of gene functions, gene regulation, and genetic diversity. As a result of gene duplications during vertebrate evolution, vertebrate genomes contain multiple closely related homologs of most genes. This genetic redundancy has allowed diversification and specialization of genes for new purposes, but it also makes gene functions harder to decipher. There is less genetic redundancy in the nematode *Caenorhabditis elegans* and the fly *Drosophila melanogaster*, which have thus played a key part in revealing universal genetic mechanisms of animal development. [↑ TOP](#)

© 2002 by Bruce Alberts, Alexander Johnson, Julian Lewis, Martin Raff, Keith Roberts, and Peter Walter.

MOLECULAR BIOLOGY OF  
**THE CELL**  
FOURTH EDITION

ALBERTS  
JOHNSON  
LEWIS  
RAFF  
ROBERTS  
WALTER

GS  
www.garlandscience.com

[Short Contents](#) | [Full Contents](#)
[Other books @ NCBI](#)

**Navigation**

[About this book](#)

**I. Introduction to the Cell**

1. Cells and Genomes

The Universal Features of Cells on Earth

The Diversity of Genomes and the Tree of Life

Genetic Information in Eucaryotes

[References](#)

*Molecular Biology of the Cell* → I. Introduction to the Cell → 1. Cells and Genomes → Genetic Information in Eucaryotes




**Figure 1-31. The major features of eucaryotic cells.** The drawing depicts a typical animal cell, but almost all the same components are found in plants and fungi and in single-celled eucaryotes such as yeasts and protozoa. Plant cells contain chloroplasts in addition to the components shown here, and their plasma membrane is surrounded by a tough external wall formed of cellulose.

**Search**

[Go](#)

☒ This book

☐ All books

☐ PubMed

© 2002 by Bruce Alberts, Alexander Johnson, Julian Lewis, Martin Raff, Keith Roberts, and Peter Walter.

MOLECULAR BIOLOGY OF

# THE CELL

FOURTH EDITION

ALBERTS  
JOHNSON  
LEWIS  
RAFF  
ROBERTS  
WALTER

GS  
[www.garlandscience.com](http://www.garlandscience.com)

[Short Contents](#) | [Full Contents](#)
[Other books @ NCBI](#)

Navigation

About this book

**I. Introduction to the Cell**

1. Cells and Genomes

The Universal Features of Cells on Earth

The Diversity of Genomes and the Tree of Life

Genetic Information in Eucaryotes

[References](#)

*Molecular Biology of the Cell* ➔ **I. Introduction to the Cell** ➔ **1. Cells and Genomes** ➔ Genetic Information in Eucaryotes




**Figure 1-32. Phagocytosis.** This series of stills from a movie shows a human white blood cell (a neutrophil) engulfing a red blood cell (artificially colored *red*) that has been treated with antibody. (Courtesy of Stephen E. Malawista and Anne de Boisfleury Chevance.)

Search

**Go**

☒ This book

☐ All books

☐ PubMed

© 2002 by Bruce Alberts, Alexander Johnson, Julian Lewis, Martin Raff, Keith Roberts, and Peter Walter.

MOLECULAR BIOLOGY OF  
**THE CELL**  
FOURTH EDITION

ALBERTS  
JOHNSON  
LEWIS  
RAFF  
ROBERTS  
WALTER

GS  
www.garlandscience.com

Short Contents | Full Contents | Other books @ NCBI

Navigation

About this book

I. Introduction to the Cell

1. Cells and Genomes

The Universal Features of Cells on Earth

The Diversity of Genomes and the Tree of Life

Genetic Information in Eucaryotes

References

*Molecular Biology of the Cell* → I. Introduction to the Cell → 1. Cells and Genomes → Genetic Information in Eucaryotes



**Figure 1-33. A single-celled eucaryote that eats other cells.** (A) *Didinium* is a carnivorous protozoan, belonging to the group known as *ciliates*. It has a globular body, about 150  $\mu\text{m}$  in diameter, encircled by two fringes of cilia—sinuous, whiplike appendages that beat continually; its front end is flattened except for a single protrusion, rather like a snout. (B) *Didinium* normally swims around in the water at high speed by means of the synchronous beating of its cilia. When it encounters a suitable prey, usually another type of protozoan, it releases numerous small paralyzing darts from its snout region. Then, the *Didinium* attaches to and devours the other cell by phagocytosis, inverting like a hollow ball to engulf its victim, which is almost as large as itself. (Courtesy of D. Barlow.)

Search

Go

☒ This book

☐ All books

☐ PubMed

© 2002 by Bruce Alberts, Alexander Johnson, Julian Lewis, Martin Raff, Keith Roberts, and Peter Walter.

*Molecular Biology of the Cell* → **I. Introduction to the Cell** → 1. Cells and Genomes → Genetic Information in Eucaryotes




**Figure 1-34. A mitochondrion.** (A) A cross section, as seen in the electron microscope. (B) A drawing of a mitochondrion with part of it cut away to show the three-dimensional structure. (C) A schematic eucaryotic cell, with the interior space of a mitochondrion, containing the mitochondrial DNA and ribosomes, colored. Note the smooth outer membrane and the convoluted inner membrane, which houses the proteins that generate ATP from the oxidation of food molecules. (A, courtesy of Daniel S. Friend.)

© 2002 by Bruce Alberts, Alexander Johnson, Julian Lewis, Martin Raff, Keith Roberts, and Peter Walter.

MOLECULAR BIOLOGY OF

# THE CELL

FOURTH EDITION

ALBERTS  
JOHNSON  
LEWIS  
RAFF  
ROBERTS  
WALTER

GS  
www.garlandscience.com

[Short Contents](#) | [Full Contents](#)
[Other books @ NCBI](#)

Navigation

About this book

**I. Introduction to the Cell**

1. Cells and Genomes

The Universal Features of Cells on Earth

The Diversity of Genomes and the Tree of Life

Genetic Information in Eucaryotes

References

*Molecular Biology of the Cell* ➔ **I. Introduction to the Cell** ➔ 1. Cells and Genomes ➔ Genetic Information in Eucaryotes




**Figure 1-35. The origin of mitochondria.** An ancestral eucaryotic cell is thought to have engulfed the bacterial ancestor of mitochondria, initiating a symbiotic relationship.

Search

**Go**

☒ This book    ☐ All books

☐ PubMed

© 2002 by Bruce Alberts, Alexander Johnson, Julian Lewis, Martin Raff, Keith Roberts, and Peter Walter.




**Figure 1-36. Chloroplasts.** These organelles capture the energy of sunlight in plant cells and some single-celled eucaryotes. (A) Leaf cells in a moss seen in the light microscope, showing the green chloroplasts. (B) An electron micrograph of a chloroplast in a leaf of grass, showing the highly folded system of internal membranes containing the chlorophyll molecules by which light is absorbed. (C) An interpretative drawing of (B). (B, courtesy of Eldon Newcomb.)

MOLECULAR BIOLOGY OF

# THE CELL

FOURTH EDITION

ALBERTS  
JOHNSON  
LEWIS  
RAFF  
ROBERTS  
WALTER

GS  
www.garlandscience.com

[Short Contents](#) | [Full Contents](#)
[Other books @ NCBI](#)

Navigation

About this book

**I. Introduction to the Cell**

1. Cells and Genomes


The Universal Features of Cells on Earth

The Diversity of Genomes and the Tree of Life

Genetic Information in Eucaryotes

References

*Molecular Biology of the Cell* → **I. Introduction to the Cell** → 1. Cells and Genomes → Genetic Information in Eucaryotes



Search

**Go**

☒ This book ☐ All books

☐ PubMed

**Figure 1-37. The origin of chloroplasts.** An early eucaryotic cell, already possessing mitochondria, engulfed a photosynthetic bacterium (a cyanobacterium) and retained it in symbiosis. All present-day chloroplasts are thought to trace their ancestry back to a single species of cyanobacterium that was adopted as an internal symbiont (an endosymbiont) over a billion years ago.

© 2002 by Bruce Alberts, Alexander Johnson, Julian Lewis, Martin Raff, Keith Roberts, and Peter Walter.

MOLECULAR BIOLOGY OF

# THE CELL

FOURTH EDITION

ALBERTS  
JOHNSON  
LEWIS  
RAFF  
ROBERTS  
WALTER

GS  
www.garlandscience.com

[Short Contents](#)
[Full Contents](#)
[Other books @ NCBI](#)

**Navigation**

About this book

**I. Introduction to the Cell**

1. Cells and Genomes

The Universal Features of Cells on Earth

The Diversity of Genomes and the Tree of Life

Genetic Information in Eucaryotes

References

*Molecular Biology of the Cell* → **I. Introduction to the Cell** → 1. Cells and Genomes → Genetic Information in Eucaryotes




**Figure 1-38. Genome sizes compared.** Genome size is measured in nucleotide pairs of DNA per haploid genome, that is, per single copy of the genome. (The cells of sexually reproducing organisms such as ourselves are generally diploid: they contain two copies of the genome, one inherited from the mother, the other from the father.) Closely related organisms can vary widely in the quantity of DNA in their genomes, even though they contain similar numbers of functionally distinct genes. (Data from W.-H. Li, *Molecular Evolution*, pp. 380–383. Sunderland, MA: Sinauer, 1997.)

**Search**

Go

☒ This book

☐ All books

☐ PubMed

© 2002 by Bruce Alberts, Alexander Johnson, Julian Lewis, Martin Raff, Keith Roberts, and Peter Walter.

The puffer fish (*Fugu rubripes*).

MOLECULAR BIOLOGY OF  
**THE CELL**  
FOURTH EDITION

ALBERTS  
JOHNSON  
LEWIS  
RAFF  
ROBERTS  
WALTER

GS  
[www.garlandscience.com](http://www.garlandscience.com)

[Short Contents](#) | [Full Contents](#) [Other books @ NCBI](#)

**Navigation**

[About this book](#)

**I. Introduction to the Cell**

1. Cells and Genomes

The Universal Features of Cells on Earth

The Diversity of Genomes and the Tree of Life

 Genetic Information in Eucaryotes

[References](#)

*Molecular Biology of the Cell* ➔ **I. Introduction to the Cell** ➔ 1. Cells and Genomes ➔ Genetic Information in Eucaryotes



**Figure 1-39. The puffer fish (*Fugu rubripes*).** This organism has a genome size of 400 million nucleotide pairs—about one-quarter as much as a zebrafish, for example, even though the two species of fish have similar numbers of genes. (From a woodcut by Hiroshige, courtesy of Arts and Designs of Japan.)

**Search**



☒ This book ☐ All books

☐ PubMed

© 2002 by Bruce Alberts, Alexander Johnson, Julian Lewis, Martin Raff, Keith Roberts, and Peter Walter.

MOLECULAR BIOLOGY OF

# THE CELL

FOURTH EDITION

ALBERTS  
JOHNSON  
LEWIS  
RAFF  
ROBERTS  
WALTER

GS  
[www.garlandscience.com](http://www.garlandscience.com)

[Short Contents](#) | [Full Contents](#)
[Other books @ NCBI](#)

Navigation

About this book

**I. Introduction to the Cell**

1. Cells and Genomes

The Universal Features of Cells on Earth

The Diversity of Genomes and the Tree of Life

↻ Genetic Information in Eucaryotes

References

Search

☒ This book
☐ All books

☐ PubMed

*Molecular Biology of the Cell* ➔ I. Introduction to the Cell ➔ 1. Cells and Genomes ➔ Genetic Information in Eucaryotes




**Figure 1-40. Controlling gene readout by environmental signals.**

Regulatory DNA allows gene expression to be controlled by regulatory proteins, which are in turn the products of other genes. This diagram shows how a cell's gene expression is adjusted according to a signal from the cell's environment. The initial effect of the signal is to activate a regulatory protein already present in the cell; the signal may, for example, trigger the attachment of a phosphate group to the regulatory protein, altering its chemical properties.

© 2002 by Bruce Alberts, Alexander Johnson, Julian Lewis, Martin Raff, Keith Roberts, and Peter Walter.

MOLECULAR BIOLOGY OF  
**THE CELL**  
FOURTH EDITION

ALBERTS  
JOHNSON  
LEWIS  
RAFF  
ROBERTS  
WALTER

GS  
www.garlandscience.com

Short Contents | Full Contents | Other books @ NCBI

Navigation

About this book

I. Introduction to the Cell

1. Cells and Genomes

The Universal Features of Cells on Earth

The Diversity of Genomes and the Tree of Life

Genetic Information in Eucaryotes

References

Search

Go

☒ This book ☐ All books

☐ PubMed

*Molecular Biology of the Cell* ➔ I. Introduction to the Cell ➔ 1. Cells and Genomes ➔ Genetic Information in Eucaryotes



**Figure 1-41. Genetic control of the program of multicellular development.** The role of a regulatory gene is demonstrated in the snapdragon *Antirrhinum*. In this example, a mutation in a single gene coding for a regulatory protein causes leafy shoots to develop in place of flowers: because a regulatory protein has been changed, the cells adopt characters that would be appropriate to a different location in the normal plant. The mutant is on the left, the normal plant on the right. (Courtesy of Enrico Coen and Rosemary Carpenter.)

© 2002 by Bruce Alberts, Alexander Johnson, Julian Lewis, Martin Raff, Keith Roberts, and Peter Walter.

beautifully simple process in which strand S separates from strand S', and each separated strand then serves as a template for the production of a new complementary partner strand that is identical to its former partner.

The ability of each strand of a DNA molecule to act as a template for producing a complementary strand enables a cell to copy, or *replicate*, its genes before passing them on to its descendants. In the next chapter we describe the elegant machinery the cell uses to perform this enormous task. [↑ TOP](#)

## In Eucaryotes, DNA Is Enclosed in a Cell Nucleus

Nearly all the DNA in a eucaryotic cell is sequestered in a nucleus, which occupies about 10% of the total cell volume. This compartment is delimited by a *nuclear envelope* formed by two concentric lipid bilayer membranes that are punctured at intervals by large nuclear pores, which transport molecules between the nucleus and the cytosol. The nuclear envelope is directly connected to the extensive membranes of the endoplasmic reticulum. It is mechanically supported by two networks of intermediate filaments: one, called the *nuclear lamina*, forms a thin sheetlike meshwork inside the nucleus, just beneath the inner nuclear membrane; the other surrounds the outer nuclear membrane and is less regularly organized ([Figure 4-9](#)).

The nuclear envelope allows the many proteins that act on DNA to be concentrated where they are needed in the cell, and, as we see in subsequent chapters, it also keeps nuclear and cytosolic enzymes separate, a feature that is crucial for the proper functioning of eucaryotic cells. Compartmentalization, of which the nucleus is an example, is an important principle of biology; it serves to establish an environment in which biochemical reactions are facilitated by the high concentration of both substrates and the enzymes that act on them. [↑ TOP](#)

## Summary

Genetic information is carried in the linear sequence of nucleotides in DNA. Each molecule of DNA is a double helix formed from two complementary strands of nucleotides held together by hydrogen bonds between G-C and A-T base pairs. Duplication of the genetic information occurs by the use of one DNA strand as a template for formation of a complementary strand. The genetic information stored in an organism's DNA contains the instructions for all the proteins the organism will ever synthesize. In eucaryotes, DNA is contained in the cell nucleus. [↑ TOP](#)

**NK cell** *see* **natural killer cell**

**N-linked oligosaccharide** Chain of sugars attached to a protein through the NH<sub>2</sub> group of the side chain of an asparagine residue.

**NMR (nuclear magnetic resonance)** Resonant absorption of electromagnetic radiation at a specific frequency by atomic nuclei in a magnetic field, due to flipping of the orientation of their magnetic dipole moments. The NMR spectrum provides information about the chemical environment of the nuclei. Two-dimensional NMR is used widely to determine the three-dimensional structure of small proteins.

**nm** *see* **nanometer**

**noncovalent attraction** Chemical bond in which, in contrast to a covalent bond, no electrons are shared. Noncovalent bonds are relatively weak, but they can sum together to produce strong, highly specific interactions between molecules.

**noncyclic photophosphorylation** Photosynthetic process that produces both ATP and NADPH in plants and cyanobacteria.

**nonenveloped virus** Virus consisting of a nucleic acid core and protein capsid only.

**nonpolar (apolar)** Lacking any asymmetric accumulation of positive and negative charge. Nonpolar molecules are generally insoluble in water.

**nonsense-mediated mRNA decay** Mechanism for removing aberrant mRNAs containing in-frame internal stop codons before they can be translated.

**Northern blotting** Technique in which RNA fragments separated by electrophoresis are immobilized on a paper sheet. A specific RNA is then detected by hybridization with a labeled nucleic acid probe.

**NO** *see* **nitric oxide**

**Notch** Receptor protein involved in many instances of choice of cell fate in animal development, for example in the specification of nerve cells from ectodermal epithelium. Its ligands are cell-surface proteins such as Delta and Serrate.

**notochord** Stiff rod of mesoderm that runs along the back of all chordate embryos. In vertebrates it does not persist and becomes incorporated into the vertebral column.

**NSF** Protein with ATPase activity that disassembles a complex of a v-SNARE and a t-SNARE.

**nuclear envelope** Double membrane surrounding the nucleus. Consists of an outer and inner membrane and is perforated by nuclear pores.