

Erythropoietin treatment during complement inhibition with eculizumab in a patient with paroxysmal nocturnal hemoglobinuria

Paroxysmal nocturnal hemoglobinuria (PNH) is characterized by intravascular hemolysis leading to anemia and other clinical manifestations. Transfusions are often required to support hemoglobin at tolerable levels. A PNH patient with aplastic anemia was treated with the complement inhibitor eculizumab, followed by concurrent treatment with recombinant human erythropoietin (rHuEpo). Eculizumab alone reduced hemolysis, increased PNH red blood cell (RBC) mass, and decreased transfusions. Addition of rHuEpo during eculizumab therapy, enhanced erythropoiesis, further increased PNH RBC mass and hemoglobin levels, and rendered the patient transfusion independent for more than two years. These data show that driving erythropoiesis during eculizumab treatment provided further benefit to a patient with PNH and underlying bone marrow failure.

Haematologica 2007; 92:e31-e33

Paroxysmal nocturnal hemoglobinuria (PNH) is an acquired genetic disorder resulting from clonal expansion of a hematopoietic stem cell containing a somatic mutation of the PIG-A gene, which results in a deficiency of glycosylphosphatidylinositol (GPI) anchored proteins from the surface of red blood cells (RBCs).¹ Absence of the GPI-anchored terminal complement inhibitor CD59 on RBCs results in an increase in complement-mediated RBC lysis leading to anemia and other clinical manifestations of the disease.²⁻⁵ Currently, no approved therapies effectively reduce intravascular hemolysis in PNH, and patients are often dependent on transfusions to artificially support hemoglobin levels.²

Eculizumab is a novel humanized monoclonal antibody directed against the complement protein C5.⁶ Eculizumab was previously shown to safely and effectively reduce intravascular hemolysis and transfusion requirements in patients with PNH.⁷⁻⁹ Herein, we report

a case study of a PNH patient with evidence of underlying aplastic anemia who was treated with eculizumab and thereafter received recombinant human erythropoietin (rHuEpo) in an effort to increase PNH RBC production during complement inhibition and thereby further reduce transfusion requirements.

Case report

A 48-year old transfusion-dependent male was diagnosed with aplastic anemia in May 1988 and with PNH in September 1993. He has been transfusion-dependent due to PNH starting in September 1993, requiring transfusions of packed red blood cells (PRBCs) every 4 to 6 weeks. He received eculizumab infusions starting May 22, 2002 and is currently dosed at 900 mg every other week. On November 6, 2002, after 6 months of receiving eculizumab, rHuEpo (NeoRecormon®) therapy was initiated at the following doses: 450 IU/kg/week in 3 divided doses during the first 2 months; 900 IU/kg/week in 3 divided doses during the next 15 months; and 750 IU/kg/week in 3 divided doses until May 3, 2006. At that time he was switched to Aranesp® at a dose of 300 mcg every 2 weeks. The dose was increased to 500 mcg every 2 weeks on 28th June 2006.

Intravascular hemolysis was assessed by measuring levels of the enzyme lactate dehydrogenase (LDH). Levels of erythropoiesis were determined by measuring reticulocyte counts. The PNH RBC count was calculated by multiplying the absolute number of RBCs by the proportion of PNH type III RBCs as assessed by flow cytometry.¹⁰ Hemoglobin levels and PRBC transfusion requirements were also monitored. All assessments have been collected to the present date and results are reported through August 2006.

During the year prior to eculizumab therapy, the mean LDH level was 2,075 IU/L (more than 4 times that of the upper limit of the normal range), the mean hemoglobin level was 10.5 g/dL, and the mean reticulocyte count was $77.5 \times 10^9/L$ (Table 1). The absolute number of PNH type III RBCs was $1.13 \times 10^{12}/L$, and the proportion of these cells constituted less than 50% of the total RBC mass. The patient required 1.8 units of PRBCs per month during the pre-treatment period (Table 1), receiving a

Table 1. Hematological parameters before and after eculizumab and rHuEpo therapies.

Parameter	Mean $\pm$ SD				
	Upon Diagnosis of AA (May 1988)	Upon Diagnosis of PNH (Sept 1993)	Pre-treatment (1 year)	Eculizumab alone (0.5 year)	Eculizumab + rHuEpo (3.7 years)
LDH, IU/L (normal range 150-480)	-	-	2075 $\pm$ 1590	456 $\pm$ 76	679 $\pm$ 146
Hemoglobin, g/dL (normal range 13.5-18.0)	7.5	9.3	10.5 $\pm$ 1.5	10.2 $\pm$ 0.9	11.4 $\pm$ 1.1
Reticulocytes, $\times 10^9/L$ (normal range 20-80)	-	-	77.5 $\pm$ 10.6	96.4 $\pm$ 29.5	205.3 $\pm$ 43.6
White Blood Cell Count, $\times 10^9/L$	2.9	3.9	4.47 $\pm$ 0.53	3.49 $\pm$ 0.50	3.84 $\pm$ 0.58
Neutrophil Count, $\times 10^9/L$	0.6	2.2	2.83 $\pm$ 0.52	2.02 $\pm$ 0.28	2.14 $\pm$ 0.33
Platelet Count, $\times 10^9/L$	20	140	113.60 $\pm$ 14.53	96.88 $\pm$ 8.06	114.40 $\pm$ 15.62
PNH Neutrophil Proportions (%)	-	-	99.82*	99.84 $\pm$ 0.14	99.84 $\pm$ 0.10
PNH Red Cell Proportions (%)	-	-	35.10 $\pm$ 10.30	65.95 $\pm$ 11.68	87.90 $\pm$ 9.23
PNH type III RBCs, $\times 10^{12}/L^\dagger$	-	-	1.13 $\pm$ 0.31	1.93 $\pm$ 0.13	2.52 $\pm$ 0.27
Units transfused per month	1.6	2.5	1.8	1.0	0.1

*Single reading; $\dagger$ Calculated as (proportion of PNH type III RBCs) $\times$ (total number of RBCs) $\div$ 100.




Figure 1. Effect of eculizumab and recombinant human erythropoietin on PNH type III RBC counts and transfusion requirements. The diamonds represent PNH type III RBC counts and solid bars represent the number of PRBC units transfused. The x-axis indicates time in years. Type and dose of erythropoietin was as follows: *NeoRecormon® at 450 IU/kg/week in 3 divided doses; †NeoRecormon® at 900 IU/kg/week in 3 divided doses; ‡NeoRecormon® at 750 IU/kg/week in 3 divided doses; §Aranesp® at 300 mcg every 2 weeks; ^Aranesp® at 500 mcg every 2 weeks.

total of 9 transfusions and 22 units (Figure 1).

After starting eculizumab treatment, hemolysis was rapidly and consistently reduced as indicated by a 78% decrease in the mean LDH level (Table 1). A concomitant increase (73%) in the PNH type III RBC mass was also demonstrated, supporting enhanced survival of these cells. Further, the average number of transfusions required each month was reduced by 44%. RBC hemoglobin was stable even though transfusion requirement decreased, indicating a net increase in endogenous hemoglobin levels (Table 1).

After 6 months of eculizumab treatment, the patient received concomitant rHuEpo therapy resulting in a mean reticulocyte count increase of 113% (Table 1). This increase in erythropoiesis was associated with an additional 32% increase in the PNH type III RBC mass over that achieved with eculizumab treatment alone. In addition, RBC hemoglobin levels showed an increase from 10.2 g/dL to 11.4 g/dL during the same period. This improvement in anemia resulted in a further decrease in transfusion requirements, eventually leading to transfusion-independence for more than two years (Figure 1). This transfusion coincided with a transient decrease in erythropoiesis, as evidenced by a drop in the reticulocyte count (data not shown). There was no evidence of an increase in intravascular hemolysis and LDH levels have remained within the normal range or just above the upper limit of the normal range during the entire treatment period. This patient continues to receive eculizumab and rHuEpo and has received only 1 transfusion in more than 3 years.

In patients with PNH, transfusions are often required to help maintain tolerable RBC hemoglobin. However, the level of hemoglobin in these patients is not only a function of RBC transfusions but also the underlying

bone marrow dysfunction and the reduced survival of the complement-sensitive PNH RBC. Attempts to drive erythropoiesis have not generally been successful as a treatment for anemia in PNH patients. Although some patients respond vigorously to erythropoietin administration, this does not translate into a sustainable increase in the endogenous RBC mass as the PNH type III RBC is destroyed by the unopposed activation of terminal complement. In addition, increasing the PNH RBC mass in a setting where these cells are vulnerable to destruction may lead to exacerbation of the disease as intravascular hemolysis is central to the morbidities of PNH.¹¹

Eculizumab, a recombinant humanized monoclonal antibody has been shown to reduce chronic intravascular hemolysis, increase PNH RBC survival, improve patient quality of life, and reduce the need for transfusion in patients with PNH.⁷⁻⁹ In the recent pivotal phase 3 trial, of 43 patients on eculizumab for up to 26 weeks, the reduction in transfusions was sustained in all patients, with 51% maintaining transfusion independence.⁷

We hypothesized that reduction of intravascular hemolysis during eculizumab treatment combined with enhanced erythropoiesis should result in a net increase in PNH RBC mass and higher endogenous hemoglobin levels, thereby further reducing the need for transfusion in patients who were not rendered transfusion independent by eculizumab alone. PNH patients with clear clinical evidence of underlying bone marrow failure, and reticulocyte counts that are either low, normal or only marginally elevated, are likely the best candidates to respond to erythropoietin therapy and therefore benefit in a setting where the PNH RBC is protected. Endogenous serum erythropoietin levels may also be a valuable indicator of those more likely to respond to rHuEpo

therapy although they were not measured in this patient.¹² However, in a meta-analysis by Littlewood *et al*, it was found that 55% of patients with erythropoietin levels >100 mU/mL still responded to rHuEpo by 8 weeks, and even 42% of patients with baseline erythropoietin levels >500 mU/mL ultimately responded;¹³ although this study was performed on patients with cancer-associated anemia and may not be directly comparable to patients with AA.

Results of this case study involving a PNH patient with aplastic anemia, demonstrate that control of intravascular hemolysis with eculizumab while driving erythropoiesis further improves anemia by increasing the endogenous PNH RBC mass, thereby reducing and/or eliminating the need for transfusion support.

A. Hill,¹ S.J. Richards,¹ R.P. Rother,² P. Hillmen¹

¹Department of Haematology, Leeds Teaching Hospitals NHS Trust, Leeds, United Kingdom; ²Alexion Pharmaceuticals, Inc, Cheshire, Connecticut, USA

Address correspondence to

Dr. Anita Hill, MBChB (Hons), MRCP, DipRCPath, Leeds General Infirmary, Department of Haematology, D Floor, Brotherton Wing, Great George Street, Leeds, LS1 3EX, UK
Phone: +44.113.3925153, fax: +44.113.3926349
E-mail: anitahill@nhs.net

Key words: Complement, eculizumab, erythropoietin, intravascular hemolysis, paroxysmal nocturnal hemoglobinuria

Acknowledgements: The Authors would like to acknowledge Dr. Henk-Andre Kroon for medical monitoring, Dr. Jing Jing Wang for clinical operational oversight, Dr. Kerry Quinn-Senger for assistance with technical writing, and Dr. Leonard Bell for critical review of the manuscript.

References

1. Takeda J, Miyata T, Kawagoe K, Iida Y, Endo Y, Fujita T et al. Deficiency of the GPI anchor caused by a somatic mutation of the PIG-A gene in paroxysmal nocturnal hemoglobinuria. *Cell* 1993;73:703-11.
2. Brodsky RA. Paroxysmal Nocturnal Hemoglobinuria. In: Hematology: Basic Principles and Practice 4th ed edn (Hoffman R, Benz EJ, Shattil SJ, Furie B, Cohen H, eds). Churchill Livingstone, Philadelphia, pp. 419-27.
3. Yamashina M, Ueda E, Kinoshita T, Takami T, Ojima A, Ono H et al. Inherited complete deficiency of 20-kilodalton homologous restriction factor (CD59) as a cause of paroxysmal nocturnal hemoglobinuria. *N Engl J Med* 1990;323:1184-9.
4. Okada N, Harada R, Okada H. Erythrocytes of patients with paroxysmal nocturnal haemoglobinuria acquire resistance to complement attack by purified 20-kD homologous restriction factor. *Clin Exp Immunol* 1990;80:109-13.
5. Rosenfeld SI, Jenkins DE Jr, Leddy JP. Enhanced reactive lysis of paroxysmal nocturnal hemoglobinuria erythrocytes. Studies on C9 binding and incorporation into high molecular weight complexes. *J Exp Med* 1986;164:981-97.
6. Thomas TC, Rollins SA, Rother RP, Giannoni MA, Hartman SL, Elliott EA et al. Inhibition of complement activity by humanized anti-C5 antibody and single-chain Fv Mol Immunol 1996;33:1389-401.
7. Hillmen P, Young NS, Schubert J, Brodsky RA, Socie G, Muus P et al. The complement inhibitor eculizumab in paroxysmal nocturnal hemoglobinuria. *N Engl J Med* 2006;355:1233-43.
8. Hill A, Hillmen P, Richards SJ, Elebute D, Marsh JC, Chan J et al. Sustained response and long-term safety of eculizumab in paroxysmal nocturnal hemoglobinuria. *Blood* 2005;106:2559-65.
9. Hillmen P, Hall C, Marsh JC, Elebute M, Bombara MP, Petro BE et al. Effect of eculizumab on hemolysis and transfusion requirements in patients with paroxysmal nocturnal hemoglobinuria. *N Engl J Med* 2004;350:552-9.
10. Richards SJ, Hillmen P. Advances in laboratory diagnosis of paroxysmal nocturnal hemoglobinuria. *Clin Applied Immunol Rev* 2001;1:315-30.
11. Parker C, Omine M, Richards S, Nishimura J, Bessler M, Ware R et al. Diagnosis and management of paroxysmal nocturnal hemoglobinuria. *Blood* 2005;106:3699-709.
12. Ludwig H, Fritz E, Leitgeb C, Pecherstorfer M, Samonigg H, Schuster J. Prediction of response to erythropoietin treatment in chronic anemia of cancer. *Blood* 1994;84:1056-63.
13. Littlewood TJ, Zagari M, Pallister C, Perkins A. Baseline and early treatment factors are not clinically useful for predicting individual response to erythropoietin in anemic cancer patients. *Oncologist* 2003;8:99-107.

[Advanced Search](#)

haematologica

Journal of The Ferrata Storti Foundation

[Home](#) [Current Issue](#) [Ahead Of Print](#) [Archive](#) [Submit a Manuscript](#)

[About Us](#) [More](#)

Table Of Contents

March 1, 2007 ; 92 (3)

Editorials and Perspectives

✓ [Current treatment of acute promyelocytic leukemia](#)

Francesco Lo-Coco, Emanuele Ammatuna, Miguel A. Sanz
Haematologica March 2007 92: 289-291;
Doi:10.3324/haematol.11226

✓ [Investigating people with mucocutaneous bleeding suggestive of primary hemostatic defects: a low likelihood of a definitive diagnosis?](#)

Emmanuel J. Favaloro
Haematologica March 2007 92: 292-296;
Doi:10.3324/haematol.11231

✓ [The metabolic syndrome and the risk of thrombosis](#)

Francesco Dentali, Erica Romualdi, Walter Ageno
Haematologica March 2007 92: 297-299;
Doi:10.3324/haematol.11243

Research Papers

Hematopoietic Stem Cells

✓ [The characteristics of hematopoietic stem cells from autoimmune-prone mice and the role of neural cell adhesion molecules in abnormal proliferation of these cells in MRL/lpr mice](#)

Xiaoli Wang, Hiroko Hisha, Wenhao Cui, Changye Song, Tomomi Mizokami, Satoshi Okazaki, Qing Li, Wei Feng, Junko Kato, Shiwen Jiang, Hongxue Fan, Susumu Ikehara
Haematologica March 2007 92: 300-307;
Doi:10.3324/haematol.10603

Acute Leukemia

In this issue



[Table of Contents](#)

[Table of Contents \(PDF\)](#)

[Index by author](#)



Jump to

- [Editorials and Perspectives](#)
- [Research Papers](#)
- [Hematopoietic Stem Cells](#)
- [Acute Leukemia](#)
- [Acute Myeloid Leukemia](#)
- [Acute Promyelocytic Leukemia](#)
- [Acute Lymphoblastic Leukemia](#)
- [Chronic Lymphocytic Leukemia](#)
- [Disorders of Hemostasis](#)
- [Thrombosis](#)
- [Progress in Hematology](#)
- [Immunotherapy](#)
- [Decision Making and Problem Solving](#)
- [Acute Myeloid Leukemia](#)
- [Brief Reports](#)
- [Aplastic Anemia](#)

- ✔ The potential effect of gender in combination with common genetic polymorphisms of drug-metabolizing enzymes on the risk of developing acute leukemia

Pascual Bolufer, Maria Collado, Eva Barragán, José Cervera, María-José Calasanz, Dolors Colomer, José Roman-Gómez, Miguel A. Sanz

Haematologica March 2007 92: 308-314;
Doi:10.3324/haematol.10752

Acute Myeloid Leukemia

- ✔ Using quantification of the *PML-RAR α* transcript to stratify the risk of relapse in patients with acute promyelocytic leukemia

Carlos Santamaría, Maria Carmen Chillón, Carina Fernández, Patricia Martín-Jiménez, Ana Balanzategui, Ramón García Sanz, Jesús F. San Miguel, Marcos-Gonzalez González

Haematologica March 2007 92: 315-322;
Doi:10.3324/haematol.10734

- ✔ Subclassification of patients with acute myelogenous leukemia based on chemokine responsiveness and constitutive chemokine release by their leukemic cells

Øystein Bruserud, Anita Rynningen, Astrid Marta Olsnes, Laila Stordrange, Anne Margrete Øyan, Karl Henning Kalland, Bjørn Tore Gjertsen

Haematologica March 2007 92: 332-341;
Doi:10.3324/haematol.10148

[Supplementary Data](#)

Acute Promyelocytic Leukemia

- ✔ Arsenic but not all-trans retinoic acid overcomes the aberrant stem cell capacity of PML/RAR α -positive leukemic stem cells

Xiaomin Zheng, Anita Seshire, Brigitte Rüster, Gesine Bug, Tim Beissert, Elena Puccetti, Dieter Hoelzer, Reinhard Henschler, Martin Ruthardt

Haematologica March 2007 92: 323-331;
Doi:10.3324/haematol.10541

Acute Lymphoblastic Leukemia

- ✔ Absence of prognostic impact of CD13 and/or CD33 antigen expression in adult acute lymphoblastic leukemia. Results of the GIMEMA ALL 0496 trial

Antonella Vitale, Anna Guarini, Cristina Ariola, Giovanna Meloni, Omar Perbellini, Michele Pizzuti, Cinzia De Gregoris, Vincenzo Mettivier, Alessandro Pastorini, Giovanni Pizzolo, Marco Vignetti, Franco Mandelli, Robin Foà

Haematologica March 2007 92: 342-348;
Doi:10.3324/haematol.10385

Chronic Lymphocytic Leukemia


- ✔ The prognostic role of CXCR3 expression by chronic lymphocytic leukemia B cells

Esther Ocaña, Luis Delgado-Pérez, Antonio Campos-Caro, Juan Muñoz, Rafael Franco, José A. Brieva

Haematologica March 2007 92: 349-356;
Doi:10.3324/haematol.10649

- Chronic Myeloid Leukemia
- Malignant Lymphomas
- Thrombosis
- Stem Cell Transplantation
- Letters to the Editor
- Disorders of Iron Metabolism
- Red Cell Disorders
- Chronic Myeloid Leukemia
- Myeloproliferative Disorders
- Online Only Articles

What about you?
Tell us your interests and get
all the new contents of
Haematologica in advance



European Myeloma Network
recommendations on the
evaluation and treatment of
newly diagnosed patients with
multiple myeloma

The predictive role of interim
positron emission tomography
for Hodgkin lymphoma
treatment outcome is confirmed

Disorders of Hemostasis

- ✔ [High prevalence of bleeders of unknown cause among patients with inherited mucocutaneous bleeding. A prospective study of 280 patients and 299 controls](#)

Teresa Quiroga, Manuela Goycoolea, Olga Panes, Eduardo Aranda, Carlos Martínez, Sabine Belmont, Blanca Muñoz, Pamela Zúñiga, Jaime Pereira, Diego Mezzano

Haematologica March 2007 92: 357-365;
Doi:10.3324/haematol.10816

- ✔ [Surgical bleeding after pre-operative unfractionated heparin and low molecular weight heparin for coronary bypass surgery](#)

Giulia Renda, Raffaele Di Pillo, Alberto D'Alleva, Adolfo Sciarilli, Marco Zimarino, Erica De Candia, Raffaele Landolfi, Gabriele Di Giammarco, Antonio Calafiore, Raffaele De Caterina

Haematologica March 2007 92: 366-373;
Doi:10.3324/haematol.10913

Thrombosis

- ✔ [Venous thromboembolism – a manifestation of the metabolic syndrome](#)

Cihan Ay, Theres Tengler, Rainer Vormittag, Ralph Simanek, Wolfgang Dorda, Thomas Vukovich, Ingrid Pabinger

Haematologica March 2007 92: 374-380;
Doi:10.3324/haematol.10828

Progress in Hematology

Immunotherapy

- ✔ [Chimeric T-cell receptors: new challenges for targeted immunotherapy in hematologic malignancies](#)

Ettore Biagi, Virna Marin, Greta Maria Paola Giordano Attianese, Erica Dander, Giovanna D'Amico, Andrea Biondi

Haematologica March 2007 92: 381-388;
Doi:10.3324/haematol.10873

Decision Making and Problem Solving

Acute Myeloid Leukemia

- ✔ [Autologous stem cell transplantation after complete remission and first consolidation in acute myeloid leukemia patients aged 61–70 years: results of the prospective EORTC –GIMEMA AML–13 study](#)

Xavier Thomas, Stefan Suciu, Bernard Rio, Giuseppe Leone, Giorgio Broccia, George Fillet, Ulrich Jehn, Walter Feremans, Giovanna Meloni, Marco Vignetti, Theo de Witte, Sergio Amadori

Haematologica March 2007 92: 389-396;
Doi:10.3324/haematol.10552

Brief Reports

Aplastic Anemia

- ✓ [Immunosuppressive therapy with anti-thymocyte globulin and cyclosporine A in selected children with hypoplastic refractory cytopenia](#)

Ayami Yoshimi, Irith Baumann, Monika Führer, Eva Bergsträsser, Ulrich Göbel, Karl-Walter Sykora, Thomas Klingebiel, Ute Gross-Wieltsch, Marry M van den Heuvel-Eibrink, Alexandra Fischer, Peter Nöllke, Charlotte Niemeyer

Haematologica March 2007 92: 397-400;
Doi:10.3324/haematol.10683

Chronic Myeloid Leukemia

- ✓ [Resistance to dasatinib in Philadelphia-positive leukemia patients and the presence or the selection of mutations at residues 315 and 317 in the BCR-ABL kinase domain](#)

Simona Soverini, Sabrina Colarossi, Alessandra Gnani, Fausto Castagnetti, Gianantonio Rosti, Costanza Bosi, Stefania Paolini, Michela Rondoni, Pier Paolo Piccaluga, Francesca Palandri, Panagiota Giannoulia, Giulia Marzocchi, Simona Luatti, Nicoletta Testoni, Ilaria Iacobucci, Daniela Cilloni, Giuseppe Saglio, Michele Baccarani, Giovanni Martinelli

Haematologica March 2007 92: 401-404;
Doi:10.3324/haematol.10822

Malignant Lymphomas

- ✓ [The translocations t\(6;18;11\)\(q24;q21;q21\) and t\(11;14;18\)\(q21;q32;q21\) lead to a fusion of the *API2* and *MALT1* genes and occur in MALT lymphomas](#)

Eva M. Murga Penas, Evelyne Callet-Bauchu, Hongtao Ye, Kristina Hinz, Nadine Albert, Christiane Copie-Bergman, Sophie Gazzo, Françoise Berger, Gilles Salles, Carsten Bokemeyer, Ming-Qing Du, Judith Dierlamm

Haematologica March 2007 92: 405-409;
Doi:10.3324/haematol.10268

Thrombosis

- ✓ [Deferral of assessment of pulmonary embolism](#)

Sergio Siragusa, Alessandra Malato, Francesco Falaschi, Fernando Porro, Raffaella Anastasio, Antonino Giarratano, Lucio Lo Coco, Maria Cristina Buonanno, Elena Maggi, Maria Antonietta Bressan, Guglielmo Mariani

Haematologica March 2007 92: 410-413;
Doi:10.3324/haematol.09899

Stem Cell Transplantation

- ✓ [Donor lymphocyte infusion for the treatment of leukemia relapse after HLA-mismatched/haploidentical T-cell-replete hematopoietic stem cell transplantation](#)

Xiao-Jun Huang, Dai-Hong Liu, Kai-Yan Liu, Lan-Ping Xu, Huan Chen, Wei Han

Haematologica March 2007 92: 414-417;
Doi:10.3324/haematol.10570



Reduced intensity stem cell transplantation for advanced soft tissue sarcomas in adults: a retrospective analysis of the European Group for Blood and Marrow Transplantation

Simona Secondino, Matteo G. Carrabba, Paolo Pedrazzoli, Luca Castagna, Francesco Spina, Federica Grosso, Alexia Bertuzzi, Jacques-Olivier Bay, Salvatore Siena, Paolo Corradini, Dietger Niederwieser, Taner Demirer
Haematologica March 2007 92: 418-420;
Doi:10.3324/haematol.10521

Letters to the Editor

Disorders of Iron Metabolism

- ✔ Juvenile hemochromatosis caused by a novel combination of hemojuvelin G320V/R176C mutations in a 5-year old girl

Patricia Aguilar-Martinez, Chun Yu Lok, Séverine Cunat, Estelle Cadet, Kathryn Robson, Jacques Rochette
Haematologica March 2007 92: 421-422;
Doi:10.3324/haematol.10701

Red Cell Disorders

- ✔ A 65 bp duplication/insertion in exon II of the β -globin gene causing β^0 -thalassemia

Hannes Frischknecht, Fabrizio Dutly
Haematologica March 2007 92: 423-424;
Doi:10.3324/haematol.10785

-
- ✔ Steroid treatment in children with sickle-cell disease

Sonia Couillard, Malika Benkerrou, Robert Giro, Valentine Brousse, Alicia Ferster, Brigitte Bader-Meunier
Haematologica March 2007 92: 425-426;
Doi:10.3324/haematol.10800

-
- ✔ Abnormalities of erythrocyte glycoconjugates are identical in two families with congenital dyserythropoietic anemia type II with different chromosomal localizations of the disease gene

Ewa Zdebska, Achille Iolascon, Justyna Spychalska, Silverio Perrotta, Carmen Lanzara, Gabriela Smolenska-Sym, Jerzy Koscielak
Haematologica March 2007 92: 427-428;
Doi:10.3324/haematol.10803

Chronic Myeloid Leukemia

- ✔ Monitoring *BCR-ABL* transcript levels by real-time quantitative polymerase chain reaction: a linear regression equation to convert from *BCR-ABL/B2M* ratio to estimated *BCR-ABL/ABL* ratio

Ilaria Iacobucci, Livia Galletti, Marilina Amabile, Simona Soverini, Michele Baccarani, Giovanni Martinelli
Haematologica March 2007 92: 429-430;
Doi:10.3324/haematol.10632

Myeloproliferative Disorders



JAK2V^{617F} detection and dosage of serum erythropoietin: first steps of the diagnostic work-up for patients consulting for elevated hematocrit

François Girodon, Eric Lippert, Pascal Mossuz, Irène Dobo, Nathalie Boiret-Dupré, Jean-François Lesesve, Sylvie Hermouet, Vincent Praloran
Haematologica March 2007 92: 431-432;
Doi:10.3324/haematol.10660

Online Only Articles

✓ Paraplegia due to extramedullary hematopoiesis in thalassemia treated successfully with radiation therapy

M. Malik, L.S. Pillai, N. Gogia, T. Puri, M. Mahapatra, D.N. Sharma, R. Kumar
Haematologica March 2007 92: e28-e30;
Doi:10.3324/haematol.10199

✓ Erythropoietin treatment during complement inhibition with eculizumab in a patient with paroxysmal nocturnal hemoglobinuria

A. Hill, S.J. Richards, R.P. Rother, P. Hillmen
Haematologica March 2007 92: e31-e33;
Doi:10.3324/haematol.10836

✓ Clinically and genetically atypical T-cell prolymphocytic leukemia underlines the relevance of a multidisciplinary diagnostic approach

Y. Sandberg, K.L. Wu, F. Heule, R.R. van den Bos, K.H. Lam, A..W. Langerak, V.H. van der Velden, K. van Lom, H.B. Beverloo
Haematologica March 2007 92: e34-e36;
Doi:10.3324/haematol.11057

✓ Primary T-cell lymphoma of the thyroid gland with chemokine receptors of Th1 phenotype complicating autoimmune thyroiditis

S. Koida, K. Tsukasaki, T. Tsuchiya, H. Harasawa, T. Fukushima, Y. Yamada, K. Ohshima, S. Kamihira, M. Kikuchi, M. Tomonaga
Haematologica March 2007 92: e37-e40;
Doi:10.3324/haematol.10351

✓ Cocaine abuse may influence the response to imatinib in CML patients

M. Breccia, F. Gentilini, G. Alimena
Haematologica March 2007 92: e41-e42;
Doi:10.3324/haematol.10976

✓ Large granular lymphocyte proliferation and revertant mosaicism: two rare events in a Wiskott-Aldrich syndrome patient

K. Boztug, U. Baumann, M. Ballmaier, D. Webster, I. Sandrock, R. Jacobs, T. Lion, S. Preuner, M. Germeshausen, G. Hansen, K. Welte, C. Klein
Haematologica March 2007 92: e43-e45;
Doi:10.3324/haematol.11222

Navigate

[Home](#)
[Current issue](#)
[Ahead of print](#)
[Archive](#)
[Info for](#)
 [Authors](#)
 [Reviewers](#)
 [Advertisers](#)
 [Subscribers](#)
[About us](#)
 [About](#)
 [Haematologica](#)
 [Editorial Board](#)
 [Our policies](#)

For Authors

[Author guidelines](#)
[Submit Manuscript](#)
[Track Manuscript](#)

For Reviewers

[Reviewer Guidelines](#)
[Access Your Profile](#)
[Access Your Tasks](#)
[2014 reviewers](#)

For Advertisers

[Information For
Advertising](#)

Education

[Review Articles](#)
[Guideline Articles](#)

More

[Rights & Permissions](#)
[Advertising](#)
[Alerts](#)
[Feedback](#)
[Contact](#)
[App](#)

Copyright © 2019 by the Ferrata Storti Foundation

ISSN 0390-6078 print

ISSN 1592-8721 online

Powered by