



nature

INTERNATIONAL WEEKLY JOURNAL OF SCIENCE

Volume 332 No. 6162 24 March 1988 \$4.95



RECOGNITION OF QUEEN BEES



nature

24 March 1988
Vol. 332 Issue no. 6162

A retinue of worker honey bees in the court of a glass pseudo-queen (in centre of photo) treated with a five-component synthetic queen mandibular gland blend. The blend initiates retinue formation, recognized here by the body alignment of the workers, head towards the stimulus. See page 354. And how bees see the third dimension, page 356.

THIS WEEK

Serotherapy hope

By engineering the hypervariable regions of a rat antibody raised against a human antigen into a human antibody sequence, it is possible to produce an antibody specific for human lymphocytes and with potential applications in serotherapy, page 323.

Proteins in stereo

An improvement in resolution means that three-dimensional NMR spectroscopy can now tackle the structures of macromolecules such as a 46-residue protein, pages 374 and 303.

Out of register

Is the present system for registering endangered species,



like this tube-nosed fruit bat, inadequate? See page 304.

Risk of quakes


The traditional methods of measuring the magnitudes of earthquakes are of only limited value, so some estimates of seismic risk are suspect. New more rigorous scales are now needed, page 319.

Tin in the air

Decaying algal material in naturally reducing environments like salt marshes may volatilize tin as stannane — and other metals might be lost to atmosphere via similar routes, pages 339 and 309.

Microscopy by force

This colour molecular-resolution image shows the surface of a leucine crystal, as seen by the atomic force microscope, now being used on biological



molecules. White spots are highspots sensed by the atomic force probe, page 332.

Oven-to-table

A domestic microwave oven cooks up superconducting material, page 311. More on the latest cuprate superconductors, pages 334 and 305. And thin film technology, page 295.

Tree ring dating

Oak trees from bogs in Northern Ireland provide an accurate date for the violent volcanic eruption of the Aegean island Santorini, page 344.

Author Index

The Author Index for January and February, facing page 380.

Nature® (ISSN 0028-0836) is published weekly on Thursday, except the last week in December, by Macmillan Magazines Ltd (4 Little Essex Street, London WC2R 3LF). Annual subscription for USA and Canada US\$250 (institutional/corporate), US\$125 (individual making personal payment). USA and Canadian orders to: *Nature*, Subscription Dept, PO Box 7663, Teaneck, NJ 07666-9857, USA. Other orders to *Nature*, Brunel Road, Basingstoke, Hants RG21 2XS, UK. Second class postage paid at New York, NY 10012 and additional mailing offices. Authorization to photocopy material for internal or personal use, or internal or personal use of specific clients, is granted by *Nature* to libraries and others registered with the Copyright Clearance Center (CCC) Transactional Reporting Service, provided the base fee of \$1.00 a copy plus \$0.10 a page is paid direct to CCC, 21 Congress Street, Salem, MA 01970, USA. Identification code for *Nature*: 0028-0836/88 \$1.00 + \$0.10. US Postmaster send address changes to: *Nature*, 65 Bleeker Street, New York, NY 10012. Published in Japan by Nature Japan K.K., Shin-Mitsuke Bldg, 36 Ichigaya Tamachi, Shinjuku-ku, Tokyo 162, Japan. © 1988 Macmillan Magazines Ltd.

OPINION

Where the world stands on ozone ■ Stations in space ■ Living with civil war 291–292

NEWS

Space station partners ■ Ozone decline ■ Foreigners in SSC ■ US social science research ■ AIDS in France, London and USA ■ Superconductors ■ Indian space science ■ UK cash windfall ■ SDI five years old ■ French biology ■ West German education ■ Soviet education ■ Natal court ruling ■ West German graduate colleges ■ Electricity privatisation ■ UK biotechnology ■ Correspondence 293–300

NEWS AND VIEWS

Making the geoid respectable again 301
Archaeology: Triple Czech burial Paul G Bahn 302
An extra dimension to NMR Dagmar Ringe 303
Conservation biology: Red books or green lists? Jared M Diamond 304
Cuprate superconductors: Structure and superstructure Colin Greaves & Ted Forgan 305
Immunology: Global or directed exocytosis? Michail V Sitkovsky & William E Paul 306
Superfluidity of ³He films P V E McClintock 307
Information processing: Neural populations revealed Terrence J Sejnowski 308
Natural volatilization of tin Peter Craig 309
A R J P Ubbelohde (1907–1988) J M Thomas 310
Daedalus: Abstract concrete 310

SCIENTIFIC CORRESPONDENCE

Microwave syntheses for superconducting ceramics D R Baghurst, A M Chippindale & D M P Mingos 311
Punctuation in perspective J Maynard Smith 311
Solitons and energy transfer in DNA K F Baverstock & R B Cundall 312
AIDS incubation period in haemophiliacs M Rees 312
Wood treatment used in Cremonese instruments C Y Barlow, P P Edwards, G R Millward, R A Raphael & D J Rubio 313
Ultra-high energy radiation from young supernovae T K Gaisser, T Stanev & F Halzen 314

BOOK REVIEWS

The Problems of Physics by A J Leggett Sudip Chakravarty 315
A Climate Modelling Primer by A Henderson-Sellers & K McGuffie A Slingo ■ Immobilized Cells: Principles and Applications by J Tampion & M D Tampion and Immobilised Enzymes and Cells by A Rosevear et al Peter Dunnill 316
Simple Curiosity: Letters from George Gaylord Simpson to His Family L F Laporte ed Colin Patterson 317
Cancer Cytogenetics by S Heim & F Mitelman K W Jones ■ Vectors: A Survey of Molecular Cloning Vectors and Their Uses R L Rodriguez & D T Denhardt eds Tim Harris 318

ARTICLES

Evidence of bias in estimations of earthquake size G Ekström & A M Dziewonski 319
Reshaping human antibodies for therapy L Riechmann, M Clark, H Waldmann & G Winter 323
Contents continued ►

LETTERS TO NATURE

- Probes of large-scale structure in the Universe**
Y Suto, K Górski, R Juszkievicz & J Silk 328
- Neutron star/red giant encounters in globular clusters**
CD Bailyn 330
- Molecular resolution images of amino acid crystals with the atomic force microscope**
S Gould, O Marti, B Drake, L Hellemans, CE Bracker, PK Hansma, NL Keder, MM Eddy & GD Stucky 332
- Crystallography, chemistry and structural disorder in the new high- T_c Bi-Ca-Sr-Cu-O superconductor**
DR Veblen, PJ Heaney, RJ Angel, LW Finger, RM Hazen, CT Prewitt, NL Ross, CW Chu, PH Hor & RL Meng [N&V] 334
- The mechanism of operation of tin(IV) oxide carbon monoxide sensors**
PG Harrison & MJ Willett 337
- Volatilization of tin as stannane in anoxic environments**
OFX Donard & JH Weber [N&V] 339
- The barite-opal-organic carbon association in oceanic particulate matter**
JKB Bishop 341
- Irish tree rings, Santorini and volcanic dust veils**
MGL Baillie & MAR Munro 344
- The dependence of convection planform on mode of heating**
G Houseman 346
- Measurement of reduced peridotite-C-O-H solidus and implications for redox melting of the mantle**
WR Taylor & DH Green 349
- Extreme ^{18}O depletion in calcite and chert clasts from the Elephant Moraine on the East Antarctic ice sheet**
G Faure, J Hoefs, LM Jones, JB Curtis & DE Pride 352
- Semiochemical basis of the retinue response to queen honey bees**
KN Slessor, L-A Kaminski, GGS King, JH Borden & ML Winston 354
- Motion cues provide the bee's visual world with a third dimension**
M Lehrer, MV Srinivasan, SW Zhang & GA Horridge 356
- Population coding of saccadic eye movements by neurons in the superior colliculus**
C Lee, WH Rohrer & DL Sparks [N&V] 357

- Basic fibroblast growth factor prevents death of lesioned cholinergic neurons *in vivo***
KJ Anderson, D Dam, S Lee & CW Cotman 360
- Stimulus-dependent myristoylation of a major substrate for protein kinase C**
AA Aderem, KA Albert, MM Keum, JKT Wang, P Greengard & ZA Cohn 362
- Intracellular free calcium rise triggers nuclear envelope breakdown in the sea urchin embryo**
RA Steinhardt & J Alderton 364
- Translational control of InsP_3 -induced chromatin condensation during the early cell cycles of sea urchin embryos**
J Twigg, R Patel & M Whitaker 366
- Allele-specific and *Mutator*-associated instability at the *Rp1* disease-resistance locus of maize**
JL Bennetzen, M-M Qin, S Ingels & AH Ellingboe 369
- The *GLI* gene is a member of the *Kruppel* family of zinc finger proteins**
KW Kinzler, JM Ruppert, SH Bigner & B Vogelstein 371
- Three-dimensional NMR spectroscopy of a protein in solution**
H Oschkinat, C Griesinger, PJ Kraulis, O W Sørensen, R R Ernst, A M Gronenborn & G M Clore [N&V] 374
- Heparan sulphate bound growth factors: a mechanism for stromal cell mediated haemopoiesis**
R Roberts, J Gallagher, E Spooncer, TD Allen, F Bloomfield & TM Dexter 376
- Receptor-directed focusing of lymphokine release by helper T cells**
W-J Poo, L Conrad & CA Janeway Jr [N&V] 378
- [N&V] See News and Views

CLASSIFIED


Back pages: Professional appointments ■ Research posts
■ Studentships ■ Fellowships ■ Conferences ■ Courses ■ Seminars ■ Symposia

NEXT WEEK

Oort cloud comets ■ Transient detection microscopy ■ Organic suspensions in rivers ■ Human gene expression preimplantation ■ A novel vasoconstrictor ■ Superconductor structures ■ Digging up Londinium

SUBSCRIPTION SERVICE

Your mailing label contains your unique subscriber identification number



To ensure prompt service, please quote this number whenever you speak or write to us about your subscription

HELP US TO HELP YOU

Annual Subscription Rates (including Annual Index)

USA & Canada
Institutional/corporate rate US\$250
Individual rate US\$125

The individual rate is available only to subscribers paying by personal check or personal credit card.

Orders (with remittance), subscription inquiries and change of address to:

Nature
Subscription Department
PO Box 7663
Teaneck
NJ 07666-9857
USA

Call toll-free (USA only); (800) 524-0384
Please allow 6-8 weeks for your subscription to start.

Back issues
US\$9.00

Binder orders to: Nature, Jesse Jones Ind. Dept
NAT 499 E. Erie Ave. Phila. PA 19134 Charge
orders call toll free 1-800-972-5858. Single binder:
USA \$10.95; Canada \$12.45. Set of Four:
USA \$39.95; Canada \$45.95.

Annual indexes (1971-1987)

US\$10.00 each

Nature first issue facsimile
US\$4.00

Orders (with remittance) to:
Circulation Offices
Nature

65 Bleeker Street
New York, NY 10012, USA
Tel: (212) 477-9628

Details of subscription rates in all other countries are available from this address

Nature in microform

For information contact:
University Microfilms International
300 North Zeeb Road
Ann Arbor
MI 48106, USA

Washington Editorial Office

Nature
1137 National Press Building
DC 20045, USA
Tel: (202) 737-2355

different paths, and by ensuring that an azimuthally uniform coverage of stations is used in the averaging calculation. To compensate for other factors, such as focal depth, fault geometry and corner frequency would require such a detailed knowledge of the earthquake source that the M_s measurement itself would be redundant.

The results of this analysis can be summarized in five points.

(1) A global average moment-magnitude relationship M_s has been defined which can be used to predict M_0 over a wide range of magnitudes and scalar moments.

(2) The variance of surface wave measurements for an event of a particular scalar moment is ~ 0.2 magnitude units.

(3) Large regional biases in M_s exist.

(4) Differences in source scaling may explain some of the differences. Specifically, observations show that the transition from a slope of unity to a smaller value occurs at large moments for continental events than for ridge and fracture zone events, suggesting systematic differences in stress drop.

(5) Other systematic factors affecting the calculation of M_s also appear to contribute to the observed regional bias.

We thank Professor J. H. Woodhouse for reading and correcting the manuscript and Professor H. Kanamori for constructive criticism throughout our work on this subject. This work was supported by the NSF.

Received 20 October 1987; accepted 4 February 1988.

1. Richter, C. F. *Bull. seism. Soc. Am.* **25**, 1-32 (1935).
2. Vanek, J. et al. *Izv. akad. Nauk. USSR, Ser. Geophys.* **2**, 153-158 (1962).
3. Aki, K. *Bull. Earthq. Res. Inst. Tokyo Univ.* **44**, 23-88 (1966).
4. Agnew, D., Berger, J., Buland, R., Farrell, W. & Gilbert, F. *Eos* **57**, 280-288 (1976).
5. Peterson, J., Butler, H. M., Holcomb, L. G. & Hutt, C. R. *Bull. seism. Soc. Am.* **66**, 2049-2068 (1976).
6. Kanamori, H. & Given, J. W. *Phys. Earth planet. Inter.* **27**, 8-31 (1981).
7. Dziewonski, A. M., Chou, T. A. & Woodhouse, J. H. *J. geophys. Res.* **86**, 2825-2852 (1981).
8. Woodhouse, J. H. & Dziewonski, A. M. *J. geophys. Res.* **88**, 3247-3271 (1983).
9. Woodhouse, J. H. & Dziewonski, A. M. *J. geophys. Res.* **89**, 5953-5986 (1984).
10. Dziewonski, A. M., Franzen, J. E. & Woodhouse, J. H. *Phys. Earth planet. Inter.* **34**, 209-219 (1984).
11. Dziewonski, A. M., Ekström, G., Franzen, J. E. & Woodhouse, J. H. *Phys. Earth planet. Inter.* **45**, 11-36 (1987).
12. Dziewonski, A. M., Ekström, G., Woodhouse, J. H. & Zwart, G. *Phys. Earth planet. Inter.* (in the press).
13. Kanamori, H. *J. geophys. Res.* **82**, 2981-2987 (1977).
14. Richter, C. F. *Elementary Seismology* (W. H. Freeman, San Francisco, 1958).
15. Lienkamper, J. J. *Bull. seism. Soc. Am.* **74**, 2357-2378 (1984).
16. Kanamori, H., Anderson, D. L. *Bull. seism. Soc. Am.* **65**, 1073-1095 (1975).
17. Ekström, G. & Dziewonski, A. M. *Bull. seism. Soc. Am.* **75**, 23-39 (1985).
18. Sipkin, S. A. *Bull. seism. Soc. Am.* **76**, 1515-1541 (1986).
19. Harkrider, D. G. *Bull. seism. Soc. Am.* **54**, 627-679 (1964).
20. Gutenberg, B. & Richter, C. F. *Gerlands Beitr. z. Geophysik* **47**, 73-131 (1936).
21. Gutenberg, B. *Bull. seism. Soc. Am.* **35**, 3-12 (1945).
22. Von Seggern, D. *Bull. seism. Soc. Am.* **60**, 503-516 (1970).
23. Nuttli, O. *Tectonophysics* **118**, 161-174 (1985).
24. Kanamori, H. & Allen, C. R. in *Maurice Ewing Series Vol. 6, Earthquake Source Mechanics* (American Geophysical Union, Washington, DC, 1986).
25. Zhuo, T. & Kanamori, H. *Bull. seism. Soc. Am.* **77**, 514-529 (1987).

Reshaping human antibodies for therapy

Lutz Riechmann[†], Michael Clark^{*}, Herman Waldmann^{*} & Greg Winter[‡]

MRC Laboratory of Molecular Biology, Hills Road, Cambridge CB2 2QH, UK

^{*} Department of Pathology, University of Cambridge, Tennis Court Road, Cambridge CB2 1QP, UK

A human IgG1 antibody has been reshaped for serotherapy in humans by introducing the six hypervariable regions from the heavy- and light-chain variable domains of a rat antibody directed against human lymphocytes. The reshaped human antibody is as effective as the rat antibody in complement and is more effective in cell-mediated lysis of human lymphocytes.

IN 1890 it was shown that resistance to diphtheria toxin could be transferred from one animal to another by the transfer of serum. It was concluded that the immune serum contained an anti-toxin, later called an antibody¹. For many years animal antisera were used in the treatment of microbial infections and for the neutralization of toxins in man². More recently rodent monoclonal antibodies (mAbs)³ have been used as 'magic bullets'⁴ to kill and to image tumours^{5,6}. The foreign immunoglobulin, however, can elicit an anti-globulin response which may interfere with therapy⁷ or cause allergic or immune complex hypersensitivity². Thus ideally human antibodies would be used. Human immunoglobulins are widely used as both prophylactic and microbicidal agents⁸, but it would be far better to have available human mAbs of the desired specificity. It has proven difficult, however, to make such mAbs by the conventional route of immortalization of human antibody-producing cells⁹.


There is an alternative approach. Antibody genes have been transfected into lymphoid cells, and the encoded antibodies expressed and secreted; by shuffling genomic exons, simple chimaeric antibodies with mouse variable regions and human constant regions have been made¹⁰⁻¹². Such chimaeric antibodies

have at least two advantages over mouse antibodies. First, the effector functions can be selected or tailored as desired. For example, of the human IgG isotypes, IgG1 and IgG3 appear to be the most effective for complement and cell-mediated lysis¹³⁻¹⁵, and therefore for killing tumour cells. Second, the use of human rather than mouse isotypes should minimize the anti-globulin responses during therapy^{16,17} by avoiding anti-isotypic antibodies. The extent to which anti-idiotypic responses to rodent antibodies in therapy are dictated by foreign components of the variable versus the constant region is not known, but the use of human isotypes should reduce the anti-idiotypic response. For example, when mice were made tolerant to rat immunoglobulin constant-region determinants, administration of rat anti-lymphocyte antibodies did evoke anti-idiotypic responses, but these were delayed and weaker than in animals that had not been made tolerant¹⁸. Nevertheless, it is likely that a chimaeric antibody would provoke a greater immune response than a human mAb.

We have attempted to build rodent antigen binding sites directly into human antibodies by transplanting only the antigen binding site, rather than the entire variable domain, from a rodent antibody. The antigen binding site is essentially encoded by the hypervariable loops at one end of the β -sheet framework. The hypervariable regions of the heavy chain of mouse antibodies against a hapten¹⁹ or a protein antigen⁴⁷ were previously transplanted into a human heavy chain, and, in association with the mouse light chain, the antigen binding site was retained.

[†] Address from April 1988: Department of Molecular Biology, The Research Institute of Scripps Clinic, North Torrey Pines Road, La Jolla, California 02937, USA

[‡] To whom correspondence should be addressed.






Fig. 2 Strategy for reshaping a human antibody for therapy. Sequences of rat origin are marked in black, and those of human origin in white. The recombinant heavy and light chains are also marked using a systematic nomenclature. See text for description of stages 1, 2 and 3. The genes encoding the variable domains were excised from the M13 vectors as *Hind*III-*Bam*HI fragments, and recloned into pSV2gpt²⁹ (heavy chains) or pSV2neo³⁰ (light chains), expression vectors containing the immunoglobulin enhancer¹². The human γ 1 (ref. 40), γ 2 (ref. 41), γ 3 (ref. 42), γ 4 (ref. 41) and κ (ref. 36) and the rat γ 2b (ref. 43) constant domains were introduced as *Bam*HI fragments. The following plasmids were constructed and transfected into lymphoid cell lines by electroporation⁴⁴. In stage 1, the pSVgpt plasmids HuVHCAMP-RaIgG2b, HuVHCAMP(Ser $\rightarrow$ Phe)-RaIgG2b, HuVHCAMP(Ser27 $\rightarrow$ Phe, Ser30 $\rightarrow$ Thr)-RaIgG2b were introduced into the heavy chain loss variant of YTH 34.5HL. In stage 2, the pSVgpt plasmids RaVHCAMP-RaIgG2b, RaVHCAMP-HuIgG1, RaVHCAMP-HuIgG2, RaVHCAMP-HuIgG3, RaVHCAMP-HuIgG4 were transfected as above. In stage 3, the pSV-gpt plasmid Hu(Ser27 $\rightarrow$ Phe, Ser30 $\rightarrow$ Thr)VHCAMP-HuIgG1 was co-transfected with the pSV-neo plasmid HuVLCAMP-HuIgK into the rat myeloma cell line Y0 (Y B2/3.0 Ag 20 (ref. 31)). In each of the three stages, clones resistant to mycophenolic acid were selected and screened for antibody production by ELISA assays. Clones secreting antibody were subcloned by limiting dilution (for Y0) or the soft agar method (for the loss variant) and assayed again before 1 litre growth in roller bottles.

Since, to a first approximation, the sequences of hypervariable regions do not contain characteristic rodent or human motifs, such 'reshaped' antibodies should be indistinguishable in sequence from human antibodies.

There are mAbs to many cell-type-specific differentiation antigens, but only a few have therapeutic potential. Of particular interest is a group of rat mAbs directed against an antigen, the 'CAMPATH-1' antigen, which is strongly expressed on virtually all human lymphocytes and monocytes, but is absent from other blood cells including the haemopoietic stem cells²⁰. The CAMPATH-1 series contains rat mAb of IgM, IgG2a and IgG2c isotypes²¹, and more recently IgG1 and IgG2b isotypes which were isolated as class-switch variants from the IgG2a-secreting cell line YTH 34.5HL²². All of these antibodies, except the rat IgG2c isotype, are able to lyse human lymphocytes efficiently with human complement. Also the IgG2b antibody YTH 34.5HL-G2b, but not the other isotypes, is effective in antibody-dependent cell-mediated cytotoxicity (ADCC) with human effector cells²². These rat mAbs have important applications in problems of immunosuppression: for example control of graft-versus-host disease in bone-marrow transplantation²⁰; the management of organ rejection²³; the prevention of marrow rejection; and the treatment of various lymphoid malignancies (ref. 24 and M. J. Dyer, Hale, G., Hayhoe, F. G. J. and Waldmann, H., unpublished observations). The IgG2b antibody YTH 34.5HL-G2b seems to be the most effective at depleting lymphocytes *in vivo* but the use of all of these antibodies is limited by the anti-globulin response which can occur within two weeks of the initiation of treatment²⁴. Here we describe the reshaping of human heavy and light chains towards binding the

Table 1 Reshaping the heavy-chain variable domain

Heavy chain variable domain	Concentration of antibody in $\mu\text{g ml}^{-1}$ at	
	50% antigen binding	50% complement lysis
RaVHCAMP	0.7	2.1
HuVHCAMP	27.3	*
HuVHCAMP (Ser27 $\rightarrow$ Phe)	1.8	16.3
HuVHCAMP (Ser 27 $\rightarrow$ Phe, Ser 30 $\rightarrow$ Thr)	2.0	17.6

Antibodies with the heavy-chain variable domains listed above, rat IgG2b constant domains and rat light chains were collected from supernatants of cells at stationary phase and concentrated by precipitation with ammonium sulphate, followed by ion exchange chromatography on a Pharmacia MonoQ column. The yields of antibody were measured by an enzyme-linked immunosorbent assay (ELISA) directed against the rat IgG2b isotype, and each was adjusted to the same concentration³⁵. To measuring binding to antigen, partially purified CAMPATH-1 antigen was coated onto microtitre wells and bound antibody was detected via a biotin-labelled anti-rat IgG2b mAb³⁵, developed with a streptavidin-peroxidase conjugate (Amersham). Complement lysis of human lymphocytes was with human serum as the complement source²¹. For both binding and complement assays, antibody titres were determined by fitting the data to a sigmoid curve by at least squares iterative procedure²¹.

* Complement lysis with the HuVHCAMP variable domain was too weak for the estimation of lytic titre.

CAMPATH-1 antigen and the selection of human effector functions to match the lytic potential of the rat IgG2b isotype.

Strategy

The amino-acid sequences of the heavy- and light-chain variable domains of the rat IgG2a CAMPATH-1 antibody YTH 34.5HL were determined from the cloned complementary DNA (Fig. 1), and the hypervariable regions were identified according to Kabat²⁵. In the heavy-chain variable domain there is an unusual feature in the framework region. In most known heavy-chain sequences Pro41 and Leu45 are highly conserved: Pro41 helps turn a loop distant from the antigen binding site and Leu45 is in the β bulge which forms part of the conserved packing between heavy- and light-chain variable domains²⁶. In YTH 34.5HL these residues are replaced by Ala41 and Pro45 and presumably this could have some effect on the packing of the heavy- and light-chain variable domains. Working at the level of the gene and using three large mutagenic oligonucleotides for each variable domain, the rat hypervariable regions were mounted in a single step on the human heavy- or light-chain framework regions taken from the crystallographically solved proteins NEW²⁷ and REI²⁸ respectively (Fig. 1). The REI light chain was used because there is a deletion at the beginning of the third framework region in NEW. The reshaped human heavy- and light-chain variable domains were then assembled with constant domains in three stage (Fig. 2). This permits a step-wise check on the reshaping of the heavy-chain variable domain (stage 1), the selection of the human isotype (stage 2), and the reshaping of the light-chain variable domain and the assembly of human antibody (stage 3). The plasmid constructions were genomic, with the sequences encoding variable domains cloned as *Hind*III-*Bam*HI fragments and those encoding the constant domains as *Bam*HI-*Bam*HI fragments in either pSVgpt (heavy chain)²⁹ or pSVneo (light chain)³⁰ vectors. The heavy-chain enhancer sequence was included on the 5' side of the variable domain, and expression of both light and heavy chains was driven from the heavy-chain promoter and the heavy-chain signal sequence.

Heavy-chain variable domain

In stage 1, the reshaped heavy-chain variable domain (HuVHCAMP) was attached to constant domains of the rat




Fig. 3 Loop Phe27 to Tyr35 in the heavy-chain variable domain of the human myeloma protein KOL, which has been solved crystallographically⁴⁵. The backbone of the hypervariable region according to Kabat²⁵ is highlighted, and a 200% van der Waal surface is thrown around Phe 27 to show the interactions with Tyr 32 and Met 34 of the Kabat hypervariable region. In the rat YTH 34.5HL heavy chain, these three side chains are conserved in character, but in HuVHCAMP, Phe27 is replaced by Ser.

isotype IgG2b and transfected into a heavy-chain loss variant of the YTH 34.5 hybridoma. This variant carries two light chains, one derived from the Y3 fusion partner³¹. The cloned rat heavy-chain variable domain (RaVHCAMP) was also expressed as above, and the antibodies were purified and quantified (Table 1). The HuVHCAMP and RaVHCAMP antibodies, each of the rat IgG2b isotype, were compared to the CAMPATH-1 antigen in a direct binding assay and in complement lysis of human lymphocytes (Table 1). Compared with the original rat antibody, or the engineered equivalent, the antibody with the reshaped heavy-chain domain bound poorly to the CAMPATH-1 antigen and was weakly lytic. This suggested an error in the design of the reshaped domain.

There are several assumptions underlying the transfer of hypervariable loops from one antibody to another⁴⁷, in particular the assumption that the antigen binds mainly to the hypervariable regions. These are defined as regions of sequence²⁵ or structural³² hypervariability, the locations of hypervariable regions being similar by both criteria except for the first hypervariable loop of the heavy chain. By sequence the first hypervariable loop extends from residues 31–35 (ref. 25) whereas by structure it extends from residues 26–32 (ref. 32). Residues 29 and 30 form part of the surface loop, and residue 27, which is phenylalanine or tyrosine in most sequences, including YTH 35.5HL, helps pack against residues 32 and 34 (Fig. 3). Unlike most human heavy chains, in NEW (see note in Fig. 1) the phenylalanine is replaced by serine, which would be unable to pack in the same way. To restore the packing of the loop, we made both a Ser 27→Phe mutation, and a Ser 27→Phe, Ser 30→Thr double mutation in HuVHCAMP. These two mutants showed a significant increase in binding to CAMPATH-1 antigen and lysed human lymphocytes with human complement (Table 1). Thus the affinity of the reshaped antibody could be restored by a single Ser 27→Phe mutation, possibly as a consequence of an altered packing between the hypervariable regions and the framework. This suggests that alterations in the 'Kabat' framework region can enhance the affinity of the antibody and extends previous work in which an engineered change in the hypervariable region yielded an antibody with increased affinity³³.

Heavy-chain constant domains

In stage 2 (Fig. 2), the rat heavy-chain variable domain was attached to constant domains of the human isotypes IgG1, 2, 3 and 4, and transfected into the heavy-chain loss variant of the YTH 34.5 hybridoma. In complement lysis (Fig. 4a), the human IgG1 isotype proved similar to the YTH 34.5HL-G2b, with the human IgG3 isotype being less effective. The human IgG2 isotype was only weakly lytic and the IgG4 isotype was non-lytic. In ADCC (Fig. 4b) the human IgG1 was more lytic than the YTH 34.5HL-G2b antibody. The decrease in lysis at higher concentrations of the rat IgG2b and the human IgG1 antibody is due to an excess of antibody, which causes the lysis of effector cells. The human IgG3 antibody was weakly lytic, and the IgG2 and IgG4 isotypes were non-lytic.

We therefore selected the human IgG1 isotype for the reshaped antibody. Other recent work also favours the use of IgG1 isotype for therapeutic application. When the effector functions of human isotypes were compared using a set of chimaeric antibodies with an anti-hapten variable domain, the IgG1 isotype appeared superior to the IgG3 in both complement and cell-mediated lysis¹⁵. Also, of two mouse chimaeric antibodies with human IgG1 or IgG3 isotypes directed against cell surface antigens as tumour cell markers, only the IgG1 isotype mediated complement lysis^{13,14}.




Fig. 4 *a*, Complement lysis and *b*, ADCC for antibodies with rat light-chain and rat heavy-chain variable domain attached to human IgG1 (□), IgG2 (○), IgG3 (■), or IgG4 (▽) isotypes. Lysis with the YTH 34.5HL antibody (●) is also shown.

Methods. Antibody was collected from cells in stationary phase, concentrated by precipitation with ammonium sulphate and desalted into phosphate buffered saline (PBS). Antibodies bound to the CAMPATH-1 antigen-coated on microtitre plates, were assayed in ELISA directed against the rat κ light chain³⁵, and each adjusted to the same concentration. The antibodies were assayed in complement lysis (Table 1) and ADCC with activated human peripheral blood mononuclear cells^{35,46}. Briefly, 5×10^4 human peripheral blood cells were labelled with ^{51}Cr and incubated for 30 min at room temperature with different concentrations of antibody. Excess antibody was removed and a 20-fold excess of activated cells added as effectors. After 4 h at 37 °C cell death was estimated by ^{51}Cr release.




Fig. 5 *a*, Complement lysis and *b*, ADCC of the reshaped human (□) and rat YTH 34.5HL (●) antibodies. Antibody HuVHCAMP (Ser27→Phe, Thr30→Ser)-HuIgG1, HuVLCAMP-HuIgG1 was purified from supernatants of cells in stationary phase by affinity chromatography on protein-A Sepharose. The yield (about 10 mg l⁻¹) was measured spectrophotometrically. Complement and ADCC assays were performed as in Fig. 4.

Light chain

In stage 3 (Fig. 2), the reshaped heavy chain was completed by attaching the reshaped HuVHCAMP (Ser27→Phe, Ser30→Thr) domain to the human IgG1 isotype. The reshaped light-chain domain HuVLCAMP was attached to the human C κ domain. The two clones were co-transfected into the non-secreting rat Y0 myeloma line. The resultant antibody, bound to CAMPATH-1 antigen (data not shown), and proved almost identical to the YTH 34.5HL-G2b antibody in complement lysis (Fig. 5*a*). In cell-mediated lysis the reshaped human antibody was more effective than the rat antibody (Fig. 5*b*). Similar results were

obtained with three different donors of target and effector cells (data not shown). Also, the antibody was as effective as YTH 34.5HL-G2b in killing leukaemic cells from three patients with B-cell lymphocytic leukaemia by complement-mediated lysis with human serum. Thus, by transplanting the hypervariable regions from a rodent to a human antibody of the IgG1 subtype, we have reshaped the antibody for therapeutic application.

Prospects

The availability of a reshaped human antibody with specificity for the CAMPATH-1 antigen should permit a full analysis of the *in vivo* potency and immunogenicity of an anti-lymphocyte antibody with wide therapeutic potential. Even if anti-idiotypic responses are eventually observed, considerable therapeutic benefit could be derived from an extended course of treatment. Also, it should be possible to circumvent an anti-globulin response restricted to idiotype by using a series of antibodies with different idiotypes³⁴. In principle, the idiotype of the reshaped CAMPATH-1 could be changed by altering the hypervariable regions or the framework regions—evidence from a reshaped antibody specific for the hapten nitrophenyl acetate suggests that recognition by anti-idiotypic antisera and anti-idiotypic mAbs is influenced by residues in the framework region¹⁹. Thus, recycling the hypervariable regions on different human framework regions should change the idiotype, although ultimately it might focus the response directly onto the binding site for the CAMPATH-1 antigen. Although such focusing would be undesirable for CAMPATH-1 antibodies, it could be an advantage for the development of anti-idiotypic vaccines. It is likely that the answers to some of these questions will emerge from the use of this reshaped antibody in therapy.

We thank J. Foote for the synthetic gene encoding the framework regions of a human light chain; P. Leder, T. R. Rabbitts, T. Honjo, M. P. Lefranc respectively for clones encoding the constant regions of human κ chain, human IgG2 and IgG4, human IgG1, human IgG3; G. Hale for CAMPATH-1 antigen and for advice; M. Bruggemann and M. S. Neuberger for subclones of the heavy-chain constant regions¹⁵ and for advice; M. Frewin for technical assistance and C. Milstein for encouragement. L.R. is supported by a fellowship from the German 'Sonderprogramm Gentechnologie des DAAD'. The work was supported by the Medical Research Council and by Wellcome Biotechnology Ltd. 'CAMPATH' is a trademark of The Wellcome Foundation Ltd.

Received 3 December 1987; accepted 12 February 1988.

- Behring, E. von & Kitasato, S. *Dtsch. Med. Wochenschr.* **16**, 1113–1114 (1890).
- Ratner, B. *Allergy Anaphylaxis and Immunotherapy* (Williams and Wilkins, Baltimore, 1943).
- Köhler, G. & Milstein, C. *Nature* **256**, 495–497 (1975).
- Ehrlich, P. in *Immunology and Cancer Research* vol. 2 (ed. Himmelweit, F.) (Pergamon, London, 1957).
- Deland, F. H. & Goldenberg, D. M. in *Radionuclide Imaging* (ed. Kuhl, D. E.) 289–307 (Pergamon, Paris, 1982).
- Levy, R. & Miller, R. A. *Rev. Med. Sci.* **34**, 107–116 (1983).
- Miller, R. A., Oseroff, A. R., Stratte, P. T. & Levy, R. *Blood* **62**, 988–995 (1983).
- Leung, D. Y. M., Rosen, G. & Geha, R. S. in *Primary and Secondary Immunodeficiency Disorders* (ed. Chandra, R. K.) 44–61 (Churchill Livingstone, New York, 1983).
- Carson, D. A. & Freemark, B. D. *Adv. Immun.* **38**, 275–311 (1986).
- Morrison, S. L., Johnson, M. J., Herzenberg, S. A. & Oi, V. T. *Proc. natn. Acad. Sci. U.S.A.* **81**, 6851–6855 (1984).
- Boulianne, G. L., Hozumi, N. & Shulman, M. J. *Nature* **312**, 643–646 (1984).
- Neuberger, M. S. *et al. Nature* **314**, 268–270 (1985).
- Liu, A. Y. *et al. Proc. natn. Acad. Sci. U.S.A.* **84**, 3439–3443 (1987).
- Shaw, D. R. *et al. J. Immun.* **138**, 4534–4538 (1987).
- Bruggemann, M. *et al. J. exp. Med.* **166**, 1351–1361 (1987).
- Hale, G., Swirsky, D. M., Hayhoe, F. G. J. & Waldmann, H. *Molec. biol. Med.* **1**, 321–334 (1983).
- Giorgi, J. V. *et al. Transplant Proc.* **15**, 639–642 (1983).
- Benjamin, R., Cobbold, S. P., Clark, M. R. & Waldmann, H. *J. exp. Med.* **163**, 1539–1552 (1986).
- Jones, P. T., Dear, P. H., Foote, J., Neuberger, M. S. & Winter, G. *Nature* **321**, 522–525 (1986).
- Hale, G. *et al. Blood* **62**, 873–882 (1983).
- Hale, G., Hoang, T., Prospero, T., Watt, S. M. & Waldmann, H. *Molec. biol. Med.* **1**, 305–319 (1983).

- Hale, G. *et al. J. Immun. Meth.* **103**, 59–67 (1987).
- Hale, G., Waldmann, H., Friend, P. & Calne, R. *Transplantation* **42**, 308–311 (1986).
- Hale, G., Swirsky, D. M., Hayhoe, F. G. J. & Waldmann, H. *Molec. biol. Med.* **1**, 321–334 (1983).
- Kabat, E. A., Wu, T. T., Reid-Miller, M., Perry, H. M. & Gottesman, K. S. in *Sequences of Proteins of Immunological Interest* (US Dept. of Health and Human Services, US Government Printing Office, 1987).
- Chothia, C., Novotny, J., Brucoleri, R. & Karplus, M. *J. molec. Biol.* **186**, 651–663 (1985).
- Saul, F. A., Amzel, M. & Poljak, R. J. *J. biol. Chem.* **253**, 585–597 (1978).
- Epp, O. *et al. Eur. J. Biochem.* **45**, 513–524 (1974).
- Mulligan, R. C. & Berg, P. *Proc. natn. Acad. Sci. U.S.A.* **78**, 2072–2076 (1981).
- Southern, P. J. & Berg, P. *J. molec. appl. Genetics* **1**, 327–345 (1981).
- Galfre, G. & Milstein, C. *Meth. Enzym.* **73**, 1–46 (1981).
- Chothia, C. & Lesk, A. *J. Mol. Biol.* **196**, 901–917 (1987).
- Roberts, S., Cheetham, J. C. & Rees, A. R. *Nature* **328**, 731–734 (1987).
- Jonker, M. & den Brok, J. H. A. M. *Eur. J. Immun.* **17**, 1547–1552 (1987).
- Clark, M. & Waldmann, H. *J.N.C.I.* **79**, 1393–1398 (1987).
- Hieter, P. A., Max, E. E., Seidmann, J. G., Maizel, J. V. Jr & Leder, P. *Cell* **22**, 197–207 (1980).
- Kaartinen, M. *et al. J. Immun.* **130**, 320–324 (1983).
- Gubler, U. & Hoffmann, B. *J. Gene* **25**, 263–269 (1983).
- Sanger, F., Nicklen, S. A. & Coulson, A. R. *Proc. natn. Acad. Sci. U.S.A.* **74**, 5463–5467 (1977).
- Takahashi, N., Ueda, N. S., Obata, M., Nikaido, T. & Honjo, T. *Cell* **29**, 671–679 (1982).
- Flanagan, J. G. & Rabbitts, T. H. *Nature* **300**, 709–713 (1982).
- Huck, S., Fort, P., Crawford, D. H., Lefranc, M.-P. & Lefranc, G. *Nucleic Acids Res.* **14**, 1779–1789 (1986).
- Bruggemann, M. *et al. Proc. natn. Acad. Sci. U.S.A.* **83**, 6075–6079 (1986).
- Potter, H., Weir, L. & Leder, P. *Proc. natn. Acad. Sci. U.S.A.* **81**, 7161–7163 (1984).
- Marquardt, M., Deisenhofer, J., Huber, R. & Palm, W. *J. molec. Biol.* **141**, 369–391 (1980).
- Hale, G., Clark, M. & Waldmann, H. *J. Immun.* **123**, 3056–3061 (1985).
- Verhoeyen, M., Milstein, C. & Winter, G. *Science* (in the press).