



US 20160194701A1

(19) **United States**

(12) **Patent Application Publication**
BEECHEM et al.

(10) **Pub. No.: US 2016/0194701 A1**

(43) **Pub. Date: Jul. 7, 2016**

(54) **ENZYME- AND AMPLIFICATION-FREE
SEQUENCING**

Publication Classification

(71) Applicant: **NanoString Technologies, Inc.**, Seattle,
WA (US)

(51) **Int. Cl.**
C12Q 1/68 (2006.01)

(72) Inventors: **Joseph M. BEECHEM**, Eugene, OR
(US); **Rustem KHAFIZOV**, Seattle, WA
(US)

(52) **U.S. Cl.**
CPC **C12Q 1/6874** (2013.01)

(21) Appl. No.: **14/946,386**

(57) **ABSTRACT**

(22) Filed: **Nov. 19, 2015**

Related U.S. Application Data

(60) Provisional application No. 62/082,883, filed on Nov.
21, 2014.

The present invention relates to sequencing probes, methods,
kits, and apparatuses that provide enzyme-free, amplifica-
tion-free, and library-free nucleic acid sequencing that has
long-read-lengths and with low error rate.




Figure 1




Figure 2




Figure 3





Figure 4






Figure 5

Figure 6A




Figure 6B




Figure 6C




Figure 6D





Figure 7



10 specific bases: $4^{10} \sim 1$ million probes

8 specific bases: $4^8 \sim 65$ thousand probes

6 specific bases: $4^6 \sim 4$ thousand probes

U_b = universal base (e.g., Inosine and nitro-indole)

Figure 8A




Figure 8B




Figure 8C




Figure 8D




Figure 8E




Figure 9A




Figure 9B

Reporter complex comprising a detectable label




Figure 9C




Figure 9D




Figure 9E




Figure 9F




Figure 9G





Figure 10



Flank = Flanking single-stranded polynucleotide
 AR = Attachment Region
 Detect = Complementary Nucleic Acid Comprising Detectable Label
 Lack = Hybridizing Nucleic Acid Lacking Detectable Label

Figure 11

	Target Binding Domain	(Optional) dsDNA spacer	Flank 1	AR-1	AR-1/ Flank 2	AR-2	AR-2/ Flank 3	AR-3	AR-3/ Flank 4	AR-4	AR-4/ Flank 5	...
V2 Design	TTCACGTTAG	CTGTCTCAATTTGCTGCATCCCTGTCCTGTCACGTTG	GAGCTT	GTGATC	CGTCCT	CTTTTC	ACTGCT	AGGCAT	TGGCCT	ATTGGG	CGTCCT	





Figure 12




~ 20 FOV (for 100X coverage of a gene) X 8 mins / FOV = 2.7 hour run = 3 Gb/24hrs
(~ 100% MiSeq)

~ 20 FOV X 5000 genes/FOV = 100,000 gene reads (yields 100X of 1000 genes)

~ 20 FOV (for 100X coverage of a gene) X 8 mins / FOV = 2.7 hour run = 12 Gb/24hrs*
(~ 400% MiSeq)

*(ordered-array NSTG throughput estimate)

Figure 13



*Need for fragmentation will be dependent on sequencing application (i.e. long-read apps such as HLA sequencing will not require this step)

Figure 14




Figure 15

Cycling calculator based on probe occupancy


Total # of single molecule targets on flow-cell	30000	
Average length of 'gapped' target (bp)	100	
Barcode occupancy (%) per seq cycle	0.75	
# of barcode cycle required to 'hit' >95% of targets on flow-cell at least once	3	
Dye occupancy (%) per position	0.8	
Total number of Dye positions	6	
Total # of 1 bp encoded read per seq cycle	6	35
Total # of 2 bp encoded read per seq cycle	23	346
Total # of 3 bp encoded read per seq cycle	92	1843
Total # of 4 bp encoded read per seq cycle	369	5530
Total # of 5 bp encoded read per seq cycle	1475	8847
Total # of 6 bp encoded read per seq cycle	5898	5898
Total number of 'usable' read per seq cycle		22439
# of barcode cycle required to cover given single molecule target with ~3X coverage	143	
# of barcode cycle required to cover given single molecule target with ~5X coverage	238	
# of barcode cycle required to cover all single molecule target on flow-cell with ~3X coverage	430	
# of barcode cycle required to cover all single molecule target on flow-cell with ~5X coverage	713	

Figure 16

	Gigabases	time hrs.	# Reads	Clinical utility
HiSeq 2500	900Gb	11 days	3 Billion !	Genome sequence cancer sequence
Next Seq	120GB	27hrs	400M	Genome sequence exome Transcriptome
Proton PI	10Gb	2-4	70M	exome Transcriptome
Present Invention	3 to 12 Gb	3	0.5M	Targeted (Dx, cDx) plus multi-analyte plus sample-to-answer, long-reads
MiSeq	3.5Gb	24hrs	25M	transcriptome targeted (Dx)
PGM 318	up to 2Gb	7 3/4 4	4M	Targeted
PacBio RS II	2.2Gb/day	3 hrs	50k	Niche genomes, long-reads, high error rate
PGM 314	0.1Gb	3.7/2 3	400k	Targeted

Figure 17

All 16 possible
di-nucleotide
combination:



Template: C G A T C T G G T T

All 10 base positions
interrogated by reading
di-nucleotides (n+1
sliding window)

G	T	A	G	A	C	C	A	A
G	T	A	G	A	C	C	A	A
G	C	T	A	G	A	C	C	A
G	C	T	A	G	A	C	C	A
G	C	T	A	G	A	C	C	A
G	C	T	A	G	A	C	C	A
G	C	T	A	G	A	C	C	A
G	C	T	A	G	A	C	C	A
G	C	T	A	G	A	C	C	A
G	C	T	A	G	A	C	C	A

(%)

G	T	A	G	A	C	C	A	A
1.5	2.3	1.7	3.1	1.0	3.8	2.2	2.8	3.0
0.8	2.0	2.1	2.0	2.0	1.8	0.6	3.7	

9 of 16 di-nucleotides
measured

Raw Error

1.5%	2.3%
0.8%	2.0%
1.7%	3.1%
2.1%	2.0%
1.0%	3.8%
2.0%	1.8%
2.2%	2.8%
0.6%	3.7%
3.0%	3.2%

base-calling positions in bold,
blue edge-bases, not called)

Raw error rate
(average: 2.1%, without
terminal positions)

Figure 18




Figure 19

		Position:									
		1	2	3	4	5	6	7	8	9	10
Template Sequence (3' – 5'): 'Perfect Match' → Probe Sequence (5' – 3'): Probe Sequence (5' – 3'): 'Single mis-match' → Probe Sequence (5' – 3'): Probe Sequence (5' – 3'):		C	G	A	T	C	T	G	G	T	T
		G	C	T	A	G	A	C	C	A	A
		G	C	T	G	G	A	C	C	A	A
		G	C	T	C	G	A	C	C	A	A
		G	C	T	T	G	A	C	C	A	A

Counts observed*	% of Total valid counts
2543	98%
40	1.5%
13	0.5%
1	0%
2636	100%


Total:

* Counts observed value represent counts after background count subtraction


****All hybridization done under buffer condition: 1xSSPE/0.1% Tween20**

Figure 20A


5x5
(1 primary nucleic acid molecule, 5 secondary nucleic acid molecules, 5 tertiary nucleic acid molecule with detectable label(s)).



4x3



3x4



Spacer 3x4




Figure 20B




Figure 20C

Exemplary recipe for constructing reporter complexes comprising detectable labels

Volume (uL)	Primary n. a. molecule (10uM)	Secondary n. a. molecule (10uM)	Tertiary n. a. molecule (100uM)	H2O
5x4	1	4.5	2.25	92.25
5x3	1	4.5	1.8	92.7
4x4	1.28	4.5	2.25	91.97
4x3	1.28	4.5	1.8	92.42
3x4	1.8	4.5	2.25	91.45

Colors in table correspond to colors in diagrams of complexes shown at left

Figure 21A




Figure 21B




Figure 22A




Figure 22B





Figure 23






Figure 24

Figure 25

code	sequences
YGBYGR	GAGACTGTAC - original
BGYRGB	GAGACAGTAC
YBRYGB	GAGACCGTAC
RGBYGR	GAGACGGTAC
YBGGYR	GAGAGTGTAC
GRYGRB	GAGAATGTAC
RGYBRG	GAGATTGTAC

code	counts	sequences
YGBYGR	16074	GAGACTGTAC
RGYBRG	1132	GAGATTGTAC
YBRYGB	998	GAGACCGTAC
YBGGYR	784	GAGAGTGTAC
YBRYBG	712	
YBGRBY	699	
YGRYBG	692	
YBRGBR	674	
Background Threshold	717	

Figure 26

DNA	Sequence	code	count
12 mer	CTGAGACTGTAC	RYGBRG	80845
11 mer	TGAGACTGTAC	GBRYGB	24474
10 mer	GAGACTGTAC	YGBYGR	18339
9 mer	AGACTGTAC	YRBYRG	1667
Mirror of "12 mer"	CATGTCAGAGTC	GRBGYR	1473
Mirror of "11 mer"	CATGTCAGAGT	BGYRBG	1384
8 mer	GACTGTAC	GBRYBG	900
		Background Threshold	622

Figure 27

Full length Reporter

20 pM full length (6 spot) **33** counts/fov

Single Spot Reporter

Single spot [fM]	Counts/FOV
20,000	Saturated
2,000	12000
200	308
20	40
2	11

20pM NS reporter $\approx$ 20fM Single spot

ENZYME- AND AMPLIFICATION-FREE SEQUENCING

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the benefit of U.S. Provisional Application No. 62/082,883, filed Nov. 21, 2014. The contents of the aforementioned patent application are incorporated herein by reference in their entireties.

SEQUENCE LISTING

[0002] The instant application contains a Sequence Listing which has been submitted in ASCII format via EFS-Web and is hereby incorporated by reference in its entirety. Said ASCII copy, created on Nov. 19, 2015, is named NATE-025_ST25.txt and is 20,860 bytes in size.

BACKGROUND OF THE INVENTION

[0003] There are currently a variety of methods for nucleic acid sequencing, i.e., the process of determining the precise order of nucleotides within a nucleic acid molecule. Current methods require amplifying a nucleic acid enzymatically, e.g., PCR, and/or by cloning. Further enzymatic polymerizations are required to produce a detectable signal by a light detection means. Such amplification and polymerization steps are costly and/or time-consuming. Thus, there is a need in the art for a method of nucleic acid sequencing that is amplification- and enzyme-free. The present invention addresses these needs.

SUMMARY OF THE INVENTION

[0004] The present invention provides sequencing probes, methods, kits, and apparatuses that provide enzyme-free, amplification-free, and library-free nucleic acid sequencing that has long-read-lengths and with low error rate. Moreover, the methods, kits, and apparatuses have rapid sample-to-answer capability. These features are particularly useful for sequencing in a clinical setting.

[0005] Provided herein are sequencing probes comprising a target binding domain and a barcode domain. The target binding domain and the barcode domain may be operably linked, e.g., covalently linked. A sequencing probe optionally comprises a spacer between the target binding domain and the barcode domain. The spacer can be any polymer with appropriate mechanical properties, for example, a single- or double-stranded DNA spacer (of 1 to 100 nucleotides, e.g., 2 to 50 nucleotides). Non-limiting examples of double-stranded DNA spacers include the sequences covered by SEQ ID NO: 25 to SEQ ID NO: 29.

[0006] The target binding domain comprises at least four nucleotides (e.g., 4, 5, 6, 7, 8, 9, 10, 11, 12, or more) and is capable of binding a target nucleic acid (e.g., DNA, RNA, and PNA). The barcode domain comprises a synthetic backbone, the barcode domain having at least a first position which comprises one or more attachment regions. The barcode domain may have one, two, three, four, five, six, seven, eight, nine, ten, eleven, twelve, or more positions; each position having one or more (e.g., one to fifty) attachment regions; each attachment region comprises at least one (i.e., one to fifty, e.g., ten to thirty copies of a nucleic acid sequence(s)) capable of reversibly binding to a complementary nucleic acid molecule (RNA or DNA). Certain positions in a barcode domain may have more attachment regions than other posi-

tions; alternately, each position in a barcode domain has the same number of attachment regions. The nucleic acid sequence of a first attachment region determines the position and identity of a first nucleotide in the target nucleic acid that is bound by a first nucleotide of the target binding domain, whereas the nucleic acid sequence of a second attachment region determines the position and identity of a second nucleotide in the target nucleic acid that is bound by a second nucleotide of the target binding domain. Likewise, the nucleic acid sequence of a sixth attachment region determines the position and identity of a sixth nucleotide in the target nucleic acid that is bound by a sixth nucleotide of the target binding domain. In embodiments, the synthetic backbone comprises a polysaccharide, a polynucleotide (e.g., single or double stranded DNA or RNA), a peptide, a peptide nucleic acid, or a polypeptide. The number of nucleotides in a target binding domain equals to or is greater than (e.g., 1, 2, 3, 4, or more) the number of positions in the barcode domain. Each attachment region in a specific position of the barcode domain may include one copy of the same nucleic acid sequence and/or multiple copies of the same nucleic acid sequence. However, an attachment region will include a different nucleic acid sequence than an attachment region in a different position of the barcode domain, even when both attachment regions identify the same type of nucleotide, e.g., adenine, thymine, cytosine, guanine, uracil, and analogs thereof. An attachment region may be linked to a modified monomer, e.g., a modified nucleotide, in the synthetic backbone, thereby creating a branch relative to the backbone. An attachment region may be part of a synthetic backbone's polynucleotide sequence. One or more attachment regions may be adjacent to at least one flanking single-stranded polynucleotide, that is, an attachment region may be operably linked to a 5' flanking single-stranded polynucleotide and/or to a 3' flanking single-stranded polynucleotide. An attachment region with or without one or two flanking single-stranded polynucleotides may be hybridized to a hybridizing nucleic acid molecule lacking a detectable label. A hybridizing nucleic acid molecule lacking a detectable label may be between about 4 and about 20 nucleotides in length, e.g., 12 nucleotides, or longer.

[0007] An attachment region may be bound by a complementary nucleic acid comprising a detectable label. Each complementary nucleic acid may comprise a detectable label.

[0008] Alternately, an attachment region may be bound by a complementary nucleic acid that is part of a reporter complex (comprising detectable labels). A complementary nucleic acid (either comprising a detectable label or of a reporter complex) may be between about 4 and about 20 nucleotides in length, e.g., about 8, 10, 12, and 14 nucleotides, or more. In a reporter complex, a complementary nucleic acid is linked (directly or indirectly) to a primary nucleic acid molecule. A complementary nucleic acid may be indirectly linked to a primary nucleic acid molecule via a single or double-stranded nucleic acid linker (e.g., a polynucleotide comprising 1 to 100 nucleotides). A primary nucleic acid is hybridized to one or more (e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) secondary nucleic acids. Each secondary nucleic acid is hybridized to one or more (e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) tertiary nucleic acids; the tertiary nucleic acids comprise one or more detectable labels. A or each secondary nucleic acid may comprise a region that does not hybridize to a primary nucleic acid molecule and does not hybridize to a tertiary nucleic acid molecule (an "extra-handle"); this region may be four or more (e.g., about 6 to about 40, e.g., about 8,

10, 12, and 14) nucleotides in length. The region that does not hybridize to a primary nucleic acid molecule and does not hybridize to a tertiary nucleic acid molecule may comprise the nucleotide sequence of the complementary nucleic acid molecule that is linked to the primary nucleic acid molecule. This region may be located near the end of the secondary nucleic acid distal to its end that hybridizes to the primary nucleic acid. By having "extra-handles" comprising the nucleotide sequence of the complementary nucleic acid, the likelihood and speed at which a reporter complex binds to a sequencing probe is greatly increased. In any embodiment or aspect of the present invention, when a reporter complex comprises "extra-handles", the reporter complex can hybridize to a sequencing probe either via the reporter complex's complementary nucleic acid or via the "extra-handle." Thus, for example, the phrase "binding to the first attachment region . . . a first complementary nucleic acid molecule of a first reporter complex" would be understood according to its plain meaning and also understood to mean "binding to the first attachment region . . . an 'extra handle' of a first reporter complex."

[0009] In embodiments, the terms "barcode domain" and "synthetic backbone" are synonymous.

[0010] Provided herein is a method for sequencing a nucleic acid using a sequencing probe of the present invention. The method comprises steps of: (1) hybridizing at least one sequencing probe, of the present invention, to a target nucleic acid that is immobilized (e.g., at one, two, three, four, five, six, seven, eight, nine, ten or more positions) to a substrate; (2) binding to the first attachment region a first complementary nucleic acid molecule (RNA or DNA) which has a detectable label (e.g., a fluorescent label) or a first complementary nucleic acid molecule of a first reporter complex comprising detectable labels (e.g., fluorescent labels); (3) detecting the detectable label(s), and (4) identifying the position and identity of the first nucleotide in the immobilized target nucleic acid. Optionally, the immobilized target nucleic acid is elongated prior to being bound by the probe. The method further comprises steps of: (5) contacting the first attachment region (with or without one or two flanking single-stranded polynucleotides) with a first hybridizing nucleic acid molecule lacking a detectable label, thereby unbinding the first complementary nucleic acid molecule having a detectable label or the first complementary nucleic acid molecule of a first reporter complex comprising detectable labels and binding to, at least, the first attachment region a first hybridizing nucleic acid lacking a detectable label; (6) binding to the second attachment region a second complementary nucleic acid molecule having a detectable label or a complementary nucleic acid molecule of a second reporter complex comprising detectable labels; (7) detecting the detectable label(s); and (8) identifying the position and identity of the second nucleotide in the immobilized target nucleic acid. Steps (5) to (8) are repeated until each nucleotide in the immobilized target nucleic acid and corresponding to the target binding domain has been identified. Steps (5) and (6) may occur concurrently or sequentially. Each (e.g., first, second, third, fourth, fifth, sixth, seventh, eighth, ninth, tenth, or higher) complementary nucleic acid molecule (having a detectable label or part of a reporter complex) has the same nucleic acid sequence as its corresponding (i.e., first, second, third, fourth, fifth, sixth, seventh, eighth, ninth, tenth, or higher) hybridizing nucleic acid molecule lacking a detectable label. The target nucleic acid is immobilized to a sub-

strate by binding a first position and/or second position of the target nucleic acid with a first and/or a second capture probe; each capture probe comprises an affinity tag that selectively binds to a substrate. The first and/or second positions may be at or near a terminus of a target nucleic acid. The substrate can be any solid support known in the art, e.g., a coated slide and microfluidic device (e.g., coated with streptavidin). Other positions which are located distant from a terminus of a target nucleic acid may be selectively bound to the substrate. The nucleic acid may be elongated by applying a force (e.g., gravity, hydrodynamic force, electromagnetic force, flow-stretching, a receding meniscus technique, and combinations thereof) sufficient to extend the target nucleic acid.

[0011] Provided herein is a method for sequencing a nucleic acid using one population of probes of the present invention or a plurality of populations of probes of the present invention. The method comprises steps of: (1) hybridizing a first population of sequencing probes (of the present invention) to a target nucleic acid that is immobilized to a substrate (with each sequencing probe in the first population de-hybridizing from the immobilized target nucleic acid under about the same conditions, e.g., level of chaotropic agent, temperature, salt concentration, pH, and hydrodynamic force); (2) binding a plurality of first complementary nucleic acid molecules each having a detectable label or a plurality of first complementary nucleic acid molecules of a plurality of first reporter complexes each complex comprising detectable labels to a first attachment region in each sequencing probe in the first population; (3) detecting the detectable label(s); (4) identifying the position and identity of a plurality of first nucleotides in the immobilized target nucleic acid hybridized by sequencing probes in the first population; (5) contacting each first attachment region of each sequencing probe of the first population with a plurality of first hybridizing nucleic acid molecules lacking a detectable label thereby unbinding the first complementary nucleic acid molecules having a detectable label or of a reporter complex and binding to each first attachment region a first hybridizing nucleic acid molecule lacking a detectable label (6) binding a plurality of second complementary nucleic acid molecules each having a detectable label or a plurality of second complementary nucleic acid molecules of a plurality of second reporter complexes each complex comprising detectable labels to a second attachment region in each sequencing probe in the first population; (7) detecting the detectable label(s); and (8) identifying the position and identity of a plurality of second nucleotides in the immobilized target nucleic acid hybridized by sequencing probes in the first population. In step (9), steps (5) to (8) are repeated until each nucleotide in the immobilized target nucleic acid and corresponding to the target binding domain of each sequencing probe in the first population has been identified. Steps (5) and (6) may occur concurrently or sequentially. Thereby, the linear order of nucleotides is identified for regions of the immobilized target nucleic acid that were hybridized by the target binding domain of sequencing probes in the first population of sequencing probes.

[0012] In embodiments, when a plurality of populations (i.e., more than one population) of probes are used, the method further comprises steps of: (10) de-hybridizing each sequencing probe of the first population from the nucleic acid; (11) removing each de-hybridized sequencing probe of the first population; (12) hybridizing at least a second population of sequencing probes of the present invention, where each sequencing probe in the second population de-hybrid-

izes from the immobilized target nucleic acid under about the same conditions and de-hybridizes from the immobilized target nucleic acid under different conditions from the sequencing probes in the first population; (13) binding a plurality of first complementary nucleic acid molecules each having a detectable label or a plurality of first complementary nucleic acid molecules of a plurality of first reporter complexes each complex comprising detectable labels to a first attachment region in each sequencing probe in the second population; (14) detecting the detectable label(s) (15) identifying the position and identity of a plurality of first nucleotides in the immobilized target nucleic acid hybridized by sequencing probes in the second population; (16) contacting each first attachment region of each sequencing probe of the second population with a plurality of first hybridizing nucleic acid molecules lacking a detectable label thereby unbinding the first complementary nucleic acid molecules (having a detectable label or from a reporter complex) and binding to each first attachment region a first hybridizing nucleic acid molecule lacking detectable label; (17) binding a plurality of second complementary nucleic acid molecules each having a detectable label or a plurality of second complementary nucleic acid molecules of a plurality of second reporter complexes each complex comprising detectable labels to a second attachment region in each sequencing probe in the second population; (18) detecting the detectable label(s); (19) identifying the position and identity of a plurality of second nucleotides in the immobilized target nucleic acid hybridized by sequencing probes in the second population; and (20) repeating steps (16) to (19) until the linear order of nucleotides has been identified for regions of the immobilized target nucleic acid that were hybridized by the target binding domain of sequencing probes in the second population of sequencing probes. Steps (16) and (17) may occur concurrently or sequentially.

[0013] Each sequencing probe in the second population may de-hybridize from the immobilized target nucleic acid at a different condition (e.g., a higher temperature, higher level of chaotropic agent, higher salt concentration, higher flow rate, and different pH) than the average condition for which the sequencing probes in the first population de-hybridize from the target nucleic acid.

[0014] However, when more than two populations of probes are used, then probes in two sequential populations may de-hybridize at different conditions and probes in non-sequential populations may de-hybridize at similar conditions. As an example, probes in a first population and third population may de-hybridize under similar conditions. In embodiments, sequential populations of probes de-hybridized at increasingly more stringent conditions (e.g., higher levels of chaotropic agent, salt concentration, and temperature). For a microfluidic device, using temperature as an example, a first population of probes may remain hybridized at a first temperature but de-hybridize at a second temperature, which is higher than the first. A second population of probes may remain hybridized at the second temperature but de-hybridize at a third temperature, which is higher than the second. In this example, solutions (comprising reagents required by the present method) flowing over a target nucleic acid for initial probe populations are at a lower temperature than solutions flowing over the target nucleic acid for later probe populations.

[0015] In some embodiments, after a population of probes has been used, the population of probes is de-hybridized from

the target nucleic acid and a new aliquot of the same population of probes is used. For example, after a first population of probes has been hybridized, detected, and de-hybridized, a subsequent aliquot of the first population of probes is hybridized. Alternately, as an example, a first population of probes may be de-hybridized and replaced with a second population of probes; once the second population has been detected and de-hybridized, a subsequent aliquot of the first population of probes is hybridized to the target nucleic acid. Thus, a probe in the subsequent population may hybridize to a region of the target nucleic acid that had been previously sequenced (thereby gaining duplicative and/or confirmatory sequence information) or a probe in the subsequent population may hybridize to a region of the target nucleic acid that had not previously been sequenced (thereby gaining new sequence information). Accordingly, a population of probes may be re-aliquoted when a prior read was unsatisfactory (for any reason) and/or to improve the accuracy of the alignment resulting from the sequencing reads.

[0016] The probes hybridizing and de-hybridizing under similar conditions may have similar lengths of their target binding domain, GC content, or frequency of repeated bases and combinations thereof. Relationships between T_m and length of an oligonucleotide are taught, for example, in Sugimoto et al., *Biochemistry*, 34, 11211-6.

[0017] When more than two populations of probes are used, steps, as described for the first and second populations of sequencing probes, are repeated with additional populations of probes (e.g., 10 to 100 to 1000 populations). The number of populations of probes used will depend on a variety of factors, including but not limited to the size of the target nucleic acid, the number of unique probes in each population, the degree of overlap among sequencing probes desired, and the enrichment of probes to regions of interest.

[0018] A population of probes may contain extra sequencing probes directed to a specific region of interest in a target nucleic acid, e.g., a region containing a mutation (e.g., a point mutation) or a SNP allele. A population of probes may contain fewer sequencing probes directed to a specific region of less interest in a target nucleic acid.

[0019] A population of sequencing probes may be compartmentalized into discrete smaller pools of sequencing probes. The compartmentalization may be based upon predicted melting temperature of the target binding domain in the sequencing probes and/or upon sequence motif of the target binding domain in the sequencing probes. The compartmentalization may be based on empirically-derived rules. The different pools of sequencing probes can be reacted with the target nucleic acid using different reaction conditions, e.g., based on temperature, salt concentration, and/or buffer content. The compartmentalization may be performed to cover target nucleic acid with uniform coverage. The compartmentalization may be performed to cover target nucleic acid with known coverage profile.

[0020] The lengths of target binding domains in a population of sequencing probes may be reduced to increase coverage of probes in a specific region of a target nucleic acid. The lengths of target binding domains in a population of sequencing probes may be increased to decrease coverage of probes in a specific region of a target nucleic acid, e.g., to above the resolution limit of the sequencing apparatus.

[0021] Alternately or additionally, the concentration of sequencing probes in a population may be increased to increase coverage of probes in a specific region of a target

nucleic acid. The concentration of sequencing probes may be reduced to decrease coverage of probes in a specific region of a target nucleic acid, e.g., to above the resolution limit of the sequencing apparatus.

[0022] The methods for sequencing a nucleic acid further comprises steps of assembling each identified linear order of nucleotides for each region of the immobilized target nucleic acid, thereby identifying a sequence for the immobilized target nucleic acid. Steps of assembling use a non-transitory computer-readable storage medium with an executable program stored thereon which instructs a microprocessor to arrange each identified linear order of nucleotides, thereby obtaining the sequence of the nucleic acid. Assembling can occur in “real time”, i.e., while data is being collected from sequencing probes rather than after all data has been collected.

[0023] The target nucleic acid, i.e., that is sequenced, may be between about 4 and 1,000,000 nucleotides in length. The target may include a whole, intact chromosome or a fragment thereof either of which is greater than 1,000,000 nucleotides in length.

[0024] Provided herein are apparatuses for performing a method of the present invention.

[0025] Provided herein are kits including sequencing probes of the present invention and for performing methods of the present invention. In embodiments, the kits include a substrate capable of immobilizing a nucleic acid via a capture probe, a plurality of sequencing probes of the present invention, at least one capture probe, at least one complementary nucleic acid molecule having a detectable label, at least one complementary nucleic acid molecule which lacks a detectable label, and instructions for use. In embodiments, the kit comprises about or at least 4096 unique sequencing probes. 4096 is the minimum number of unique probes necessary to include each possible hexameric combination (i.e., for probes each having six attachment regions in the barcode domains). Here, “4096” is achieved since there are four nucleotides options for six positions: 4^6 . For a set of probes having four attachment regions in the barcode domains, only 256 (i.e., 4^4) unique probes will be needed. For a set of probes having eight nucleotides in their target binding domains, 4^8 (i.e., 65,536) unique probes will be needed. For a set of probes having ten nucleotides in their target binding domains, 4^{10} (i.e., 1,048,576) unique probes will be needed.

[0026] In embodiments, the kit comprises about or at least twenty four distinct complementary nucleic acid molecule having a detectable label and about or at least twenty four distinct hybridizing nucleic acid molecule lacking a detectable label. A complementary nucleic acid may bind to an attachment region having a sequence of one of SEQ ID NO: 1 to 24, as non-limiting examples. Additional exemplary sequences that may be included in a barcode domain are listed in SEQ ID NO: 42 to SEQ ID NO: 81. Indeed, the nucleotide sequence is not limited; preferably it lacks substantial homology (e.g., 50% to 99.9%) with a known nucleotide sequence; this helps avoid undesirable hybridization of a complementary nucleic acid and a target nucleic acid.

[0027] Any of the above aspects and embodiments can be combined with any other aspect or embodiment.

[0028] Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. In the Specification, the singular forms also include the plural unless the context clearly dictates

otherwise; as examples, the terms “a,” “an,” and “the” are understood to be singular or plural and the term “or” is understood to be inclusive. By way of example, “an element” means one or more element. Throughout the specification the word “comprising,” or variations such as “comprises” or “comprising,” will be understood to imply the inclusion of a stated element, integer or step, or group of elements, integers or steps, but not the exclusion of any other element, integer or step, or group of elements, integers or steps. About can be understood as within 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1%, 0.5%, 0.1%, 0.05%, or 0.01% of the stated value. Unless otherwise clear from the context, all numerical values provided herein are modified by the term “about.”

[0029] Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, suitable methods and materials are described below. All publications, patent applications, patents, and other references mentioned herein are incorporated by reference in their entirety. The references cited herein are not admitted to be prior art to the claimed invention. In the case of conflict, the present Specification, including definitions, will control. In addition, the materials, methods, and examples are illustrative only and are not intended to be limiting. Other features and advantages of the invention will be apparent from the following detailed description and claim.

BRIEF DESCRIPTION OF THE DRAWINGS

[0030] The patent or application file contains at least one drawing executed in color. Copies of this patent or patent application publication with color drawings will be provided by the Office upon request and payment of the necessary fee.

[0031] The above and further features will be more clearly appreciated from the following detailed description when taken in conjunction with the accompanying drawings.

[0032] FIG. 1 shows a schematic of an exemplary sequencing probe of the present invention.

[0033] FIG. 2 shows a schematic of an exemplary sequencing probe of the present invention.

[0034] FIG. 3 shows a schematic of an exemplary sequencing probe of the present invention.

[0035] FIG. 4 shows a schematic of an exemplary sequencing probe of the present invention.

[0036] FIG. 5 shows a schematic of an exemplary sequencing probe of the present invention.

[0037] FIG. 6A is a schematic showing a sequencing probe variant of the present invention.

[0038] FIG. 6B is a schematic showing a sequencing probe variant of the present invention.

[0039] FIG. 6C is a schematic showing a sequencing probe variant of the present invention.

[0040] FIG. 6D is a schematic showing a sequencing probe variant of the present invention.

[0041] FIG. 7 shows schematics of target binding domains of sequencing probes of the present invention; the domains include zero, two, or four nucleotides having universal bases.

[0042] FIG. 8A illustrates a step of a sequencing method of the present invention.

[0043] FIG. 8B illustrates a step of a sequencing method of the present invention begun in FIG. 8A.

[0044] FIG. 8C illustrates a step of a sequencing method of the present invention begun in FIG. 8A.

[0045] FIG. 8D illustrates a step of a sequencing method of the present invention begun in FIG. 8A.

[0046] FIG. 8E illustrates a step of a sequencing method of the present invention begun in FIG. 8A.

[0047] FIG. 9A shows an initial step of a sequencing method of the present invention.

[0048] FIG. 9B shows a schematic of a reporter complex comprising detectable labels.

[0049] FIG. 9C shows a plurality of reporter complexes each comprising detectable labels.

[0050] FIG. 9D shows a further step of the sequencing method begun in FIG. 9A.

[0051] FIG. 9E shows a further step of the sequencing method begun in FIG. 9A.

[0052] FIG. 9F shows a further step of the sequencing method begun in FIG. 9A.

[0053] FIG. 9G shows a further step of the sequencing method begun in FIG. 9A.

[0054] FIG. 10 shows an alternate illustration of the steps shown in FIG. 9D and FIG. 9E and exemplary data obtained therefrom. The fragment of the sequencing probe shown has the sequence of SEQ ID NO: 82.

[0055] FIG. 11 illustrates a variation of the method shown in FIG. 10. The fragment of the sequencing probe shown likewise has the sequence of SEQ ID NO: 82.

[0056] FIG. 12 illustrates a method of the present invention.

[0057] FIG. 13 compares steps required in a sequencing method of the present invention with steps required with other sequencing methods.

[0058] FIG. 14 exemplifies performance measurements obtainable by the present invention.

[0059] FIG. 15 exemplifies performance measurements obtainable by the present invention.

[0060] FIG. 16 compares the sequencing rate, number of reads, and clinical utility for the present invention and various other sequencing methods/apparatuses.

[0061] FIG. 17 demonstrates the low raw error rate of sequencing methods of the present invention. The template sequence shown has the sequence of SEQ ID NO: 83.

[0062] FIG. 18 compares sequencing data obtainable from the present invention with other sequencing methods.

[0063] FIG. 19 demonstrates single-base specificity of sequencing methods of the present invention. The template and probe sequences shown (from top to bottom) have the sequences of SEQ ID NO: 84 to SEQ ID NO: 88.

[0064] FIG. 20A shows various designs of reporter complexes of the present invention.

[0065] FIG. 20B shows fluorescent counts obtained from the reporter complexes shown in FIG. 20A.

[0066] FIG. 20C shows exemplary recipes for constructing reporter complexes of the present invention.

[0067] FIG. 21A shows designs of reporter complexes comprising "extra-handles".

[0068] FIG. 21B shows fluorescent counts obtained from the reporter complexes having "extra-handles".

[0069] FIG. 22A shows hybridization kinetics of two exemplary designs of reporter complexes of the present invention.

[0070] FIG. 22B shows hybridization kinetics of two exemplary designs of reporter complexes of the present invention.

[0071] FIG. 23 shows a schematic of a sequencing probe of the present invention used in a method distinct from that shown in FIG. 8 through FIG. 12.

[0072] FIG. 24 shows a schematic of a consumable sequencing card useful in the present invention.

[0073] FIG. 25 shows the mismatch detection of a 10 mer, as described in Example 3. The nucleotides shown (top to bottom) have the sequences of SEQ ID NO: 89 to SEQ ID NO: 99.

[0074] FIG. 26 shows hybridization ability depending on the size of a target binding domain, as described in Example 3. The background is high due to very high reporter concentration and there was no prior purification. The nucleotides shown (top to bottom) have the sequences of SEQ ID NO: 100 to SEQ ID NO: 104.

[0075] FIG. 27 shows a comparison between a single spot vs a full-length reporter. Results for single spots show speed of hybridization is 1000x greater than for a full length barcode (Conditions 100 nM target, 30 minute hybridization).

DETAILED DESCRIPTION OF THE INVENTION

[0076] The present invention provides sequencing probes, methods, kits, and apparatuses that provide enzyme-free, amplification-free, and library-free nucleic acid sequencing that has long-read-lengths and with low error rate.

Sequencing Probe

[0077] The present invention relates to a sequencing probe comprising a target binding domain and a barcode domain. Non-limiting examples of sequencing probes of the present invention are shown in FIGS. 1 to 6.

[0078] FIG. 1 shows a schematic of a sequencing probe of the present invention. This exemplary sequencing probe has a target binding domain of six nucleotides, each of which corresponds to a position in the barcode domain (which comprises one or more attachment regions). A first attachment region is noted; it corresponds to the nucleotide of a target nucleic acid bound by a first nucleotide in the target binding domain. The third position on the barcode domain is noted. A fifth position comprising two attachment regions is noted. Each position on a barcode domain can have multiple attachment regions. For example, a position may have 1 to 50 attachment regions. Certain positions in a barcode domain may have more attachment regions than other positions (as shown here in position 5 relative to positions 1 to 4 and 6); alternately, each position in a barcode domain has the same number of attachment regions (see, e.g., FIGS. 2, 3, 5, and 6). Although not shown, each attachment region comprises at least one (i.e., one to fifty, e.g., ten to thirty) copies of a nucleic acid sequence(s) capable of reversibly binding to a complementary nucleic acid molecule (RNA or DNA). In FIG. 1, the attachment regions are integral to the linear polynucleotide molecule that makes up the barcode domain.

[0079] FIG. 2 shows a schematic of a sequencing probe of the present invention. This exemplary sequencing probe has a target binding domain of six nucleotides, each of which corresponds to an attachment region in the barcode domain. A first attachment region is noted; it corresponds to the nucleotide of a target nucleic acid bound by a first nucleotide in the target binding domain. The fourth position on the barcode domain, which comprises a portion of the barcode domain and two fourth attachment regions are encircled. Two sixth attachment regions are noted. Here, each position has two attachment regions; however, each position on a barcode domain can have one attachment region or multiple attachment regions, e.g., 2 to 50 attachment regions. Although not shown, each attachment region comprises at least one (i.e., one to fifty, e.g., ten to thirty) copies of a nucleic acid

sequence(s) capable of reversibly binding to a complementary nucleic acid molecule (RNA or DNA). In FIG. 2, the barcode domain is a linear polynucleotide molecule to which the attachment regions are linked; the attachment regions are not integral to the polynucleotide molecule.

[0080] FIG. 3 shows another a schematic of a sequencing probe of the present invention. This exemplary sequencing probe has a target binding domain of four nucleotides, with these four nucleotides in the corresponding to four positions in the barcode domain. Each position is shown with three linked attachment regions.

[0081] FIG. 4 shows yet another schematic of a sequencing probe of the present invention. This exemplary sequencing probe has a target binding domain of ten nucleotides. However, only the first six nucleotides correspond to six positions in the barcode domain. The seventh to tenth nucleotides (indicated by “n₁ to n₄”) are added to increase the length of the target binding domain thereby affecting the likelihood that a probe will hybridize and remain hybridized to a target nucleic acid. In embodiments, “n” nucleotides may precede the nucleotides corresponding to positions in the barcode domain. In embodiments, “n” nucleotides may follow the nucleotides corresponding to positions in the barcode domain. In FIG. 4, four “n” nucleotides are shown; however, a target binding domain may include more than four “n” nucleotides. The “n” nucleotides may have universal bases (e.g., inosine, 2'-deoxyinosine (hypoxanthine deoxynucleotide) derivatives, nitroindole, nitroazole analogues, and hydrophobic aromatic non-hydrogen-bonding bases) which can base pair with any of the four canonical bases.

[0082] Another sequencing probe of the present invention is shown in FIG. 5. Here, the “n” nucleotides precede and follow the nucleotides corresponding to positions in the barcode domain. The exemplary sequencing probe shown has a target binding domain of ten nucleotides. However, only the third to eighth nucleotides in the target binding domain correspond to six positions (first to sixth) in the barcode domain. The first, second, ninth, and tenth nucleotides (indicated by “n₁ to n₄”) are added to increase the length of the target binding domain. In FIG. 5, four “n” nucleotides are shown; however, a target binding domain may include more or less than four “n” nucleotides.

[0083] FIG. 6A to FIG. 6D show variants of a sequencing probe of FIG. 1. In FIG. 6A, the linear order of nucleotides in the target binding domain and linear order of attachment regions in the barcode domain progress from left to right (with respect to the illustration). In FIG. 6B, the linear order of nucleotides in the target binding domain and linear order of attachment regions in the barcode domain progress from right to left (with respect to the illustration). In FIG. 6C, the linear order of nucleotides in the target binding domain is reversed relative to the linear order of attachment regions in the barcode domain. In any probe of the present invention, there may be a lack of strict order of the nucleotides in the target binding domain and of attachment regions in barcode domain as long as the probe is designed such that each nucleotide in the target binding domain corresponds to an attachment domain or attachment domains in the barcode domain; lacks of strict order is shown in FIG. 6D. Any probe of the present invention (e.g., those exemplified in FIGS. 1 to 5) may have an ordering of nucleotides and attachment regions as shown in FIG. 6.

[0084] The target binding domain has at least four nucleotides, e.g., at least, 4, 5, 6, 7, 8, 9, 10, 11, 12, or more

nucleotides. The target binding domain preferable is a polynucleotide. The target binding domain is capable of binding a target nucleic acid.

[0085] A probe may include multiple copies of the target binding domain operably linked to a synthetic backbone.

[0086] Probes can be designed to control the likelihood of hybridization and/or de-hybridization and the rates at which these occur. Generally, the lower a probe's T_m, the faster and more likely that the probe will de-hybridize to/from a target nucleic acid. Thus, use of lower T_m probes will decrease the number of probes bound to a target nucleic acid.

[0087] The length of a target binding domain, in part, affects the likelihood of a probe hybridizing and remaining hybridized to a target nucleic acid. Generally, the longer (greater number of nucleotides) a target binding domain is, the less likely that a complementary sequence will be present in the target nucleotide. Conversely, the shorter a target binding domain is, the more likely that a complementary sequence will be present in the target nucleotide. For example, there is a 1/256 chance that a four-mer sequence will be located in a target nucleic acid versus a 1/4096 chance that a six-mer sequence will be located in the target nucleic acid. Consequently, a collection of shorter probes will likely bind in more locations for a given stretch of a nucleic acid when compared to a collection of longer probes.

[0088] FIG. 7 shows 10-mer target binding domains. In some embodiments, the target binding domain includes four universal bases (identified as “U_b”) which base pair with any of the four canonical nucleotides (A, G, C, and T). In embodiments, the target binding domain includes one to six (e.g., 2 and 4) universal bases. A target binding domain may include no universal nucleotides. FIG. 7 notes that a “complete” population of probes having 6 specific nucleotides in the target binding domain will require 4096 unique probes and a “complete” population of probes having 10 specific nucleotides will require ~1 million unique probes.

[0089] In circumstances, it is preferable to have probes having shorter target binding domains to increase the number of reads in the given stretch of the nucleic acid, thereby enriching coverage of a target nucleic acid or a portion of the target nucleic acid, especially a portion of particular interest, e.g., when detecting a mutation or SNP allele.

[0090] However, it may be preferable to have fewer numbers of probes bound to a target nucleic acid since there are occasions when too many probes in a region may cause overlap of their detectable label, thereby preventing resolution of two nearby probes. This is explained as follows. Given that one nucleotide is 0.34 nm in length and given that the lateral (x-y) spatial resolution of a sequencing apparatus is about 200 nm, a sequencing apparatus's resolution limit is about 588 base pair (i.e., a 1 nucleotide/0.34 nm×200 nm). That is to say, the sequencing apparatus mentioned above would be unable to resolve signals from two probes hybridized to a target nucleic acid when the two probes are within about 588 base pair of each other. Thus, two probes, depending on the resolution of the sequencing apparatus, will need be spaced approximately 600 bp's apart before their detectable label can be resolved as distinct “spots”. So, at optimal spacing, there should be a single probe per 600 bp of target nucleic-acid. A variety of software approaches (e.g., utilize fluorescence intensity values and wavelength dependent ratios) can be used to monitor, limit, and potentially deconvolve the number of probes hybridizing inside a resolvable region of a target nucleic acid and to design probe populations accordingly.

Moreover, detectable labels (e.g., fluorescent labels) can be selected that provide more discrete signals. Furthermore, methods in the literature (e.g., Small and Parthasarthy: "Superresolution localization methods." *Annu. Rev. Phys. Chem.*, 2014; 65:107-25) describe structured-illumination and a variety of super-resolution approaches which decrease the resolution limit of a sequencing microscope up to 10's-of-nanometers. Use of higher resolution sequencing apparatuses allow for use of probes with shorter target binding domains.

[0091] As mentioned above, designing the T_m of probes can affect the number of probes hybridized to a target nucleic acid. Alternately or additionally, the concentration of sequencing probes in a population may be increased to increase coverage of probes in a specific region of a target nucleic acid. The concentration of sequencing probes may be reduced to decrease coverage of probes in a specific region of a target nucleic acid, e.g., to above the resolution limit of the sequencing apparatus.

[0092] The term "target nucleic acid" shall mean a nucleic acid molecule (DNA, RNA, or PNA) whose sequence is to be determined by the probes, methods, and apparatuses of the invention. In general, the terms "target nucleic acid", "nucleic acid molecule", "nucleic acid sequence", "nucleic acid", "nucleic acid fragment", "oligonucleotide" and "polynucleotide" are used interchangeably and are intended to include, but not limited to, a polymeric form of nucleotides that may have various lengths, either deoxyribonucleotides or ribonucleotides, or analogs thereof. Non-limiting examples of nucleic acids include a gene, a gene fragment, an exon, an intron, intergenic DNA (including, without limitation, heterochromatic DNA), messenger RNA (mRNA), transfer RNA, ribosomal RNA, ribozymes, small interfering RNA (siRNA), non-coding RNA (ncRNA), cDNA, recombinant polynucleotides, branched polynucleotides, plasmids, vectors, isolated DNA of a sequence, isolated RNA of a sequence, nucleic acid probes, and primers.

[0093] The present methods directly sequence a nucleic acid molecule obtained from a sample, e.g., a sample from an organism, and, preferably, without a conversion (or amplification) step. As an example, for RNA-based sequencing, the present methods do not require conversion of an RNA molecule to a DNA molecule (i.e., via synthesis of cDNA) before a sequence can be obtained. Since no amplification or conversion is required, a nucleic acid sequenced in the present invention will retain any unique base and/or epigenetic marker present in the nucleic acid when the nucleic acid is in the sample or when it was obtained from the sample. Such unique bases and/or epigenetic markers are lost in sequencing methods known in the art.

[0094] The target nucleic acid can be obtained from any sample or source of nucleic acid, e.g., any cell, tissue, or organism, in vitro, chemical synthesizer, and so forth. The target nucleic acid can be obtained by any art-recognized method. In embodiments, the nucleic acid is obtained from a blood sample of a clinical subject. The nucleic acid can be extracted, isolated, or purified from the source or samples using methods and kits well known in the art.

[0095] A nucleic acid molecule comprising the target nucleic acid may be fragmented by any means known in the art. Preferably, the fragmenting is performed by an enzymatic or a mechanical means. The mechanical means may be sonication or physical shearing. The enzymatic means may be

performed by digestion with nucleases (e.g., Deoxyribonuclease I (DNase I)) or one or more restriction endonucleases.

[0096] When a nucleic acid molecule comprising the target nucleic acid is an intact chromosome, steps should be taken to avoid fragmenting the chromosome.

[0097] The target nucleic acid can include natural or non-natural nucleotides, comprising modified nucleotides, as well-known in the art.

[0098] Probes of the present invention may have overall lengths (including target binding domain, barcode domain, and any optional domains) of about 20 nanometers to about 50 nanometers. A probe's backbone may be a polynucleotide molecule comprising about 120 nucleotides.

[0099] The barcode domain comprises a synthetic backbone. The synthetic backbone and the target binding domain are operably linked, e.g., are covalently attached or attached via a linker. The synthetic backbone can comprise any material, e.g., polysaccharide, polynucleotide, polymer, plastic, fiber, peptide, peptide nucleic acid, or polypeptide. Preferably, the synthetic backbone is rigid. In embodiments, the backbone comprises "DNA origami" of six DNA double helices (See, e.g., Lin et al, "Submicrometre geometrically encoded fluorescent barcodes self-assembled from DNA." *Nature Chemistry*; 2012 October; 4(10): 832-9). A barcode can be made of DNA origami tiles (Jungmann et al, "Multiplexed 3D cellular super-resolution imaging with DNA-PAINT and Exchange-PAINT", *Nature Methods*, Vol. 11, No. 3, 2014).

[0100] The barcode domain comprises a plurality of positions, e.g., one, two, three, four, five, six, seven, eight, nine, ten, or more positions. The number of positions may be less than, equal to, or more than the number of nucleotides in the target binding domain. It is preferable to include additional nucleotides in a target binding domain than number of positions in the backbone domain, e.g., one, two, three, four, five, six, seven, eight, nine, ten, or more nucleotides. The length of the barcode domain is not limited as long as there is sufficient space for at least four positions, as described above.

[0101] Each position in the barcode domain corresponds to a nucleotide in the target binding domain and, thus, to a nucleotide in the target nucleic acid. As examples, the first position in the barcode domain corresponds to the first nucleotide in the target binding domain and the sixth position in the barcode domain corresponds to the sixth nucleotide in the target binding domain.

[0102] Each position in the barcode domain comprises at least one attachment region, e.g., one to 50, or more, attachment regions. Certain positions in a barcode domain may have more attachment regions than other positions (e.g., a first position may have three attachment regions whereas a second position may have two attachment positions); alternately, each position in a barcode domain has the same number of attachment regions. Each attachment region comprises at least one (i.e., one to fifty, e.g., ten to thirty) copies of a nucleic acid sequence(s) capable of being reversibly bound by a complementary nucleic acid molecule (e.g., DNA or RNA). In examples, the nucleic acid sequence in a first attachment region determines the position and identity of a first nucleotide in the target nucleic acid that is bound by a first nucleotide of the target binding domain. Each attachment region may be linked to a modified monomer (e.g., modified nucleotide) in the synthetic backbone such that the attachment region branches from the synthetic backbone. In embodiments, the attachment regions are integral to a poly-

nucleotide backbone; that is to say, the backbone is a single polynucleotide and the attachment regions are parts of the single polynucleotide's sequence. In embodiments, the terms "barcode domain" and "synthetic backbone" are synonymous.

[0103] The nucleic acid sequence in an attachment region identifies the position and identity of a nucleotide in the target nucleic acid that is bound by a nucleotide in the target binding domain of a sequencing probe. In a probe, each attachment region will have a unique overall sequence. Indeed, each position on a barcode domain can have an attachment region comprising a nucleic acid sequence that encodes one of four nucleotides, i.e., specific to one of adenine, thymine/uracil, cytosine, and guanine. Also, the attachment region of a first position (and encoding cytosine, for example) will include a nucleic acid sequence different from the attachment region of a second position (and encoding cytosine, for example). Thus, to a nucleic acid sequence in an attachment region in a first position that encodes a thymine, there will be no binding of a complementary nucleic acid molecule that identifies an adenine in a target nucleic acid corresponding to the first nucleotide of a target binding domain. Also, to an attachment region in a second position, there will be no binding of a complementary nucleic acid molecule that identifies an adenine in a target nucleic acid corresponding to the first nucleotide of a target binding domain.

[0104] Each position on a barcode domain may include one or more (up to fifty, preferably ten to thirty) attachment region; thus, each attachment region may bind one or more (up to fifty, preferably ten to thirty) complementary nucleic acid molecules. As examples, the probe in FIG. 1 has a fifth position comprising two attachment regions and the probe in FIG. 2 has a second position having six attachment regions. In embodiments, the nucleic acid sequences of attachment regions at a position are identical; thus, the complementary nucleic acid molecules that bind those attachment regions are identical. In alternate embodiments, the nucleic acid sequences of attachment regions at a position are not identical; thus, the complementary nucleic acid molecules that bind those attachment regions are not identical, e.g., each comprises a different nucleic acid sequence and/or detectable label. Therefore, in the alternate embodiment, the combination of non-identical nucleic acid molecules (e.g., their detectable labels) attached to an attachment region together provides a code for identifying a nucleotide in the target nucleic acid.

[0105] Table 1 provides exemplary sequences, for illustration purposes only, for attachments regions for sequencing probes having up to six positions in its barcode domain and detectable labels on complementary nucleic acid that bind thereto.

TABLE 1

Nucleotide in target binding domain/position in barcode domain	Nucleotide	Nucleic Acid Sequence (5' to 3')	Detectable label of complementary nucleic acid	SEQ ID NO
1	A	ATACATCTAG	GFP	1
1	G	GATCTACATA	RFP	2
1	C	TTAGGTAAAG	CFP	3
1	U/T	TCTTCATTAC	YFP	4
2	A	ATGAATCTAC	GFP	5
2	G	TCAATGTATG	RFP	6
2	C	AATTGAGTAC	CFP	7
2	U/T	ATGTTAATGG	YFP	8
3	A	AATTAGGATG	GFP	9
3	G	ATAATGGATC	RFP	10
3	C	TAATAAGGTG	CFP	11
3	U/T	TAGTTAGAGC	YFP	12
4	A	ATAGAGAAGG	GFP	13
4	G	TTGATGATAC	RFP	14
4	C	ATAGTGATTC	CFP	15
4	U/T	TATAACGATG	YFP	16
5	A	TTAAGTTTAG	GFP	17
5	G	ATACGTTATG	RFP	18
5	C	TGTACTATAG	CFP	19
5	U/T	TTACAAGTG	YFP	20

TABLE 1-continued

Nucleotide in target binding domain/position in barcode domain	Nucleotide	Nucleic Acid Sequence (5' to 3') in Attachment Region	Detectable label of complementary nucleic acid	SEQ ID NO
6	A	AACATGTAC	GFP	21
6	G	TAACATGAC	RFP	22
6	C	ACTAATGTTT	CFP	23
6	U/T	TCATTGAATG	YFP	24

[0106] As seen in Table 1, the nucleic acid sequence of a first attachment region may be one of SEQ ID NO: 1 to SEQ ID NO: 4 and the nucleic acid sequence of a second attachment region may be one of SEQ ID NO: 5 to SEQ ID NO: 8. When the first nucleotide in the target nucleic acid is adenine, the nucleic acid sequence of the first attachment region would have the sequence of SEQ ID NO: 1 and when the second nucleotide in the target nucleic acid is adenine, the nucleic acid sequence of the second attachment region would have the sequence of SEQ ID NO: 5.

[0107] In embodiments, a complementary nucleic acid molecule may be bound by a detectable label. In alternate embodiments, a complementary nucleic acid is associated with a reporter complex comprising detectable labels.

[0108] The nucleotide sequence of a complementary nucleic acid is not limited; preferably it lacks substantial homology (e.g., 50% to 99.9%) with a known nucleotide sequence; this helps avoid undesirable hybridization of a complementary nucleic acid and a target nucleic acid.

[0109] An example of the reporter complex useful in the present invention is shown in FIG. 9B. In this example, a complementary nucleic acid is linked to a primary nucleic acid molecule, which in turn is hybridized to a plurality of secondary nucleic acid molecules, each of which is in turn hybridized to a plurality of tertiary nucleic acid molecules having attached thereto one or more detectable labels.

[0110] In embodiments, a primary nucleic acid molecule may comprise about 90 nucleotides. A secondary nucleic acid molecule may comprise about 87 nucleotides. A tertiary nucleic acid molecule may comprise about 15 nucleotides.

[0111] FIG. 9C shows a population of exemplary reporter complexes. Included in the top left panel of FIG. 9C are the four complexes that hybridize to attachment region 1 of a probe. There is one type of reporter complex for each possible nucleotide that can be present in nucleotide position 1 of a probe's target binding domain. Here, while performing a sequence method of the present invention, if the position 1 of a probe's reporter domain is bound by a reporter complex having a "blue-colored" detectable label, then the first nucleotide in the target binding domain is identified as Adenine. Alternately, if the position 1 is bound by a reporter complex having a "green-colored" detectable label, then the first nucleotide in the target binding domain is identified as Thymine.

[0112] Reporter complexes can be of various designs. For example, a primary nucleic acid molecule can be hybridized to at least one (e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) secondary nucleic acid molecules. Each secondary nucleic acid molecule may be hybridized to at least one (e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) tertiary nucleic acid molecules.

Exemplary reporter complexes are shown in FIG. 20A. Here, the "4x3" reporter complex has one primary nucleic acid molecule (that is linked to a complementary nucleic acid molecule) hybridized to four secondary nucleic acid molecules, each of which is hybridized to three tertiary nucleic acid molecules (each comprising a detectable label). In this figure, each complementary nucleic acid of a complex is 12 nucleotides long ("12 bases"); however, the length of the complementary nucleic acid is non-limited and can be less than 12 or more than 12 nucleotides. The bottom-right complex includes a spacer region between its complementary nucleic acid and its primary nucleic acid molecule. The spacer is identified as 20 to 40 nucleotides long; however, the length of a spacer is non-limiting and it can be shorter than 20 nucleotides or longer than 40 nucleotides.

[0113] FIG. 20B shows variable average (fluorescent) counts obtained from the four exemplary reporter complexes shown in FIG. 20A. In FIG. 20B, 10 pM of biotinylated target template was attached onto a streptavidin-coated flow-cell surface, 10 nM of a reporter complex was flowed onto the flow-cell; after a one minute incubation, the flow-cell was washed, the flow-cell was imaged, and fluorescent features were counted.

[0114] In embodiments, the reporter complexes are "pre-constructed". That is, each polynucleotide in the complex is hybridized prior to contacting the complex with a probe. An exemplary recipe for pre-constructing five exemplary reporter complexes is shown in FIG. 20C.

[0115] FIG. 21A shows alternate reporter complexes in which the secondary nucleic acid molecules have "extra-handles" that are not hybridized to a tertiary nucleic acid molecule and are distal to the primary nucleic acid molecule. In this figure, each "extra-handle" is 12 nucleotides long ("12 mer"); however, their lengths are non-limited and can be less than 12 or more than 12 nucleotides. In embodiments, the "extra-handles" each comprise the nucleotide sequence of the complementary nucleic acid; thus, when a reporter complex comprises "extra-handles", the reporter complex can hybridize to a sequencing probe either via the reporter complex's complementary nucleic acid or via an "extra-handle." Accordingly, the likelihood that a reporter complex binds to a sequencing probe is increased. The "extra-handle" design may also improve hybridization kinetics. Without being bound to theory, the "extra-handles" essentially increase the effective concentration of the reporter complex's complementary nucleic acid.

[0116] FIG. 21B shows variable average (fluorescent) counts obtained from the five exemplary reporter complexes having "extra-handles" using the procedure described for FIG. 20B.

[0117] FIGS. 22A and 22B show hybridization kinetics and fluorescent intensities for two exemplary reporter complexes. By about 5 minutes, total counts start to plateau indicating that most reporter complex added have found an available target.

[0118] A detectable moiety, label or reporter can be bound to a complementary nucleic acid or to a tertiary nucleic acid molecule in a variety of ways, including the direct or indirect attachment of a detectable moiety such as a fluorescent moiety, colorimetric moiety and the like. One of skill in the art can consult references directed to labeling nucleic acids. Examples of fluorescent moieties include, but are not limited to, yellow fluorescent protein (YFP), green fluorescent protein (GFP), cyan fluorescent protein (CFP), red fluorescent protein (RFP), umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, cyanines, dansyl chloride, phycocyanin, phycoerythrin and the like. Fluorescent labels and their attachment to nucleotides and/or oligonucleotides are described in many reviews, including Haugland, *Handbook of Fluorescent Probes and Research Chemicals*, Ninth Edition (Molecular Probes, Inc., Eugene, 2002); Keller and Manak, *DNA Probes*, 2nd Edition (Stockton Press, New York, 1993); Eckstein, editor, *Oligonucleotides and Analogues: A Practical Approach* (IRL Press, Oxford, 1991); and Wetmur, *Critical Reviews in Biochemistry and Molecular Biology*, 26:227-259 (1991). Particular methodologies applicable to the invention are disclosed in the following sample of references: U.S. Pat. Nos. 4,757,141; 5,151,507; and 5,091,519. In one aspect, one or more fluorescent dyes are used as labels for labeled target sequences, e.g., as disclosed by U.S. Pat. No. 5,188,934 (4,7-dichlorofluorescein dyes); U.S. Pat. No. 5,366,860 (spectrally resolvable rhodamine dyes); U.S. Pat. No. 5,847,162 (4,7-dichlororhodamine dyes); U.S. Pat. No. 4,318,846 (ether-substituted fluorescein dyes); U.S. Pat. No. 5,800,996 (energy transfer dyes); Lee et al. U.S. Pat. No. 5,066,580 (xanthine dyes); U.S. Pat. No. 5,688,648 (energy transfer dyes); and the like. Labelling can also be carried out with quantum dots, as disclosed in the following patents and patent publications: U.S. Pat. Nos. 6,322,901; 6,576,291; 6,423,551; 6,251,303; 6,319,426; 6,426,513; 6,444,143; 5,990,479; 6,207,392; 2002/0045045; and 2003/0017264. As used herein, the term "fluorescent label" comprises a signaling moiety that conveys information through the fluorescent absorption and/or emission properties of one or more molecules. Such fluorescent properties include fluorescence intensity, fluorescence lifetime, emission spectrum characteristics, energy transfer, and the like.

[0119] Commercially available fluorescent nucleotide analogues readily incorporated into nucleotide and/or oligonucleotide sequences include, but are not limited to, Cy3-dCTP, Cy3-dUTP, Cy5-dCTP, Cy5-dUTP (Amersham Biosciences, Piscataway, N.J.), fluorescein-12-dUTP, tetramethylrhodamine-6-dUTP, TEXAS RED™-5-dUTP, CASCADE BLUE™-7-dUTP, BODIPY TMFL-14-dUTP, BODIPY TMR-14-dUTP, BODIPY TMTR-14-dUTP, RHODAMINE GREEN™-5-dUTP, OREGON GREEN™-488-5-dUTP, TEXAS RED™-12-dUTP, BODIPY™ 630/650-14-dUTP, BODIPY™ 650/665-14-dUTP, ALEXA FLUOR™ 488-5-dUTP, ALEXA FLUOR™ 532-5-dUTP, ALEXA FLUOR™ 568-5-dUTP, ALEXA FLUOR™ 594-5-dUTP, ALEXA FLUOR™ 546-14-dUTP, fluorescein-12-UTP, tetramethylrhodamine-6-UTP, TEXAS RED™-5-UTP, mCherry, CASCADE BLUE™-7-UTP, BODIPY™ FL-14-

UTP, BODIPY TMR-14-UTP, BODIPY™ TR-14-UTP, RHODAMINE GREEN™-5-UTP, ALEXA FLUOR™ 488-5-UTP, LEXA FLUOR™ 546-14-UTP (Molecular Probes, Inc. Eugene, Oreg.) and the like. Alternatively, the above fluorophores and those mentioned herein may be added during oligonucleotide synthesis using for example phosphoramidite or NHS chemistry. Protocols are known in the art for custom synthesis of nucleotides having other fluorophores (See, Henegariu et al. (2000) *Nature Biotechnol.* 18:345). 2-Aminopurine is a fluorescent base that can be incorporated directly in the oligonucleotide sequence during its synthesis. Nucleic acid could also be stained, a priori, with an intercalating dye such as DAPI, YOYO-1, ethidium bromide, cyanine dyes (e.g., SYBR Green) and the like.

[0120] Other fluorophores available for post-synthetic attachment include, but are not limited to, ALEXA FLUOR™ 350, ALEXA FLUOR™ 405, ALEXA FLUOR™ 430, ALEXA FLUOR™ 532, ALEXA FLUOR™ 546, ALEXA FLUOR™ 568, ALEXA FLUOR™ 594, ALEXA FLUOR™ 647, BODIPY 493/503, BODIPY FL, BODIPY R6G, BODIPY 530/550, BODIPY TMR, BODIPY 558/568, BODIPY 558/568, BODIPY 564/570, BODIPY 576/589, BODIPY 581/591, BODIPY TR, BODIPY 630/650, BODIPY 650/665, Cascade Blue, Cascade Yellow, Dansyl, lissamine rhodamine B, Marina Blue, Oregon Green 488, Oregon Green 514, Pacific Blue, Pacific Orange, rhodamine 6G, rhodamine green, rhodamine red, tetramethyl rhodamine, Texas Red (available from Molecular Probes, Inc., Eugene, Oreg.), Cy2, Cy3, Cy3.5, Cy5, Cy5.5, Cy7 (Amersham Biosciences, Piscataway, N.J.) and the like. FRET tandem fluorophores may also be used, including, but not limited to, PerCP-Cy5.5, PE-Cy5, PE-Cy5.5, PE-Cy7, PE-Texas Red, APC-Cy7, PE-Alexa dyes (610, 647, 680), APC-Alexa dyes and the like.

[0121] Metallic silver or gold particles may be used to enhance signal from fluorescently labeled nucleotide and/or oligonucleotide sequences (Lakowicz et al. (2003) *BioTechniques* 34:62).

[0122] Other suitable labels for an oligonucleotide sequence may include fluorescein (FAM, FITC), digoxigenin, dinitrophenol (DNP), dansyl, biotin, bromodeoxyuridine (BrdU), hexahistidine (6xHis), phosphor-amino acids (e.g., P-tyr, P-ser, P-thr) and the like. In one embodiment the following hapten/antibody pairs are used for detection, in which each of the antibodies is derivatized with a detectable label: biotin/a-biotin, digoxigenin/a-digoxigenin, dinitrophenol (DNP)/a-DNP, 5-Carboxyfluorescein (FAM)/a-FAM.

[0123] Detectable labels described herein are spectrally resolvable. "Spectrally resolvable" in reference to a plurality of fluorescent labels means that the fluorescent emission bands of the labels are sufficiently distinct, i.e., sufficiently non-overlapping, that molecular tags to which the respective labels are attached can be distinguished on the basis of the fluorescent signal generated by the respective labels by standard photodetection systems, e.g., employing a system of band pass filters and photomultiplier tubes, or the like, as exemplified by the systems described in U.S. Pat. Nos. 4,230,558; 4,811,218; or the like, or in Wheelless et al., pgs. 21-76, in *Flow Cytometry: Instrumentation and Data Analysis* (Academic Press, New York, 1985). In one aspect, spectrally resolvable organic dyes, such as fluorescein, rhodamine, and the like, means that wavelength emission maxima are spaced at least 20 nm apart, and in another aspect, at least 40 nm apart. In another aspect, chelated lanthanide compounds,

quantum dots, and the like, spectrally resolvable means that wavelength emission maxima are spaced at least 10 nm apart, and in a further aspect, at least 15 nm apart.

Sequencing Method

[0124] The present invention relates to methods for sequencing a nucleic acid using a sequencing probe of the present invention. Examples of the method are shown in FIGS. 8 to 12.

[0125] The method comprises reversibly hybridizing at least one sequencing probe, of the present invention, to a target nucleic acid that is immobilized (e.g., at one, two, three, four, five, six, seven, eight, nine, ten, or more positions) to a substrate.

[0126] The substrate can be any solid support known in the art, e.g., a coated slide and a microfluidic device, which is capable of immobilizing a target nucleic acid. In certain embodiments, the substrate is a surface, membrane, bead, porous material, electrode or array. The target nucleic acid can be immobilized onto any substrate apparent to those of skill in the art.

[0127] In embodiments, the target nucleic acid is bound by a capture probe which comprises a domain that is complementary to a portion of the target nucleic acid. The portion may be an end of the target nucleic acid or not towards an end.

[0128] Exemplary useful substrates include those that comprise a binding moiety selected from the group consisting of ligands, antigens, carbohydrates, nucleic acids, receptors, lectins, and antibodies. The capture probe comprises a binding moiety capable of binding with the binding moiety of the substrate. Exemplary useful substrates comprising reactive moieties include, but are not limited to, surfaces comprising epoxy, aldehyde, gold, hydrazide, sulfhydryl, NETS-ester, amine, thiol, carboxylate, maleimide, hydroxymethyl phosphine, imidoester, isocyanate, hydroxyl, pentafluorophenyl-ester, psoralen, pyridyl disulfide or vinyl sulfone, polyethylene glycol (PEG), hydrogel, or mixtures thereof. Such surfaces can be obtained from commercial sources or prepared according to standard techniques. Exemplary useful substrates comprising reactive moieties include, but are not limited to, OptArray-DNA NHS group (Accelr8), Nexterion Slide AL (Schott) and Nexterion Slide E (Schott).

[0129] In embodiments, the capture probe's binding moiety is biotin and the substrate comprises avidin (e.g., streptavidin). Useful substrates comprising avidin are commercially available including TB0200 (Accelr8), SAD6, SAD20, SAD100, SAD500, SAD2000 (Xantec), SuperAvidin (Array-It), streptavidin slide (catalog #MPC 000, Xenopore) and STREPTAVIDINslide (catalog #439003, Greiner Bio-one).

[0130] In embodiments, the capture probe's binding moiety is avidin (e.g., streptavidin) and the substrate comprises biotin. Useful substrates comprising biotin that are commercially available include, but are not limited to, Optarray-biotin (Accelr8), BD6, BD20, BD100, BD500 and BD2000 (Xantec).

[0131] In embodiments, the capture probe's binding moiety can comprise a reactive moiety that is capable of being bound to the substrate by photoactivation. The substrate could comprise the photoreactive moiety, or the first portion of the nanopore could comprise the photoreactive moiety. Some examples of photoreactive moieties include aryl azides, such as N((2-pyridyl)dithio)ethyl)-4-azidosalicylamide; fluorinated aryl azides, such as 4-azido-2,3,5,6-tetrafluorobenzoic

acid; benzophenone-based reagents, such as the succinimidyl ester of 4-benzoylbenzoic acid; and 5-Bromo-deoxyuridine.

[0132] In embodiments, the capture probe's binding moiety can be immobilized to the substrate via other binding pairs apparent to those of skill in the art.

[0133] After binding to the substrate, the target nucleic acid may be elongated by applying a force (e.g., gravity, hydrodynamic force, electromagnetic force "electrostretching", flow-stretching, a receding meniscus technique, and combinations thereof) sufficient to extend the target nucleic acid.

[0134] The target nucleic acid may be bound by a second capture probe which comprises a domain that is complementary to a second portion of the target nucleic acid. The portion may be an end of the target nucleic acid or not towards an end. Binding of a second capture probe can occur after or during elongation of the target nucleic acid or to a target nucleic acid that has not been elongated. The second capture probe can have a binding as described above.

[0135] A capture probe may comprise or be associated with a detectable label, i.e., a fiducial spot.

[0136] The capture probe is capable of isolating a target nucleic acid from a sample. Here, a capture probe is added to a sample comprising the target nucleic acid. The capture probe binds the target nucleic acid via the region of the capture probe that is complementary to a region of the target nucleic acid. When the target nucleic acid contacts a substrate comprising a moiety that binds the capture probe's binding moiety, the nucleic acid becomes immobilized onto the substrate.

[0137] To ensure that a user "captures" as many target nucleic acid molecules as possible from high fragmented samples, it is helpful to include a plurality of capture probes, each complementary to a different region of the target nucleic acid. For example, there may be three pools of capture probes, with a first pool complementary to regions of the target nucleic acid near its 5' end, a second pool complementary to regions in the middle of the target nucleic acid, and a third pool near its 3' end. This can be generalized to "n-regions-of-interest" per target nucleic acid. In this example, each individual pool of fragmented target nucleic acid bound to a capture probe comprising or bound to a biotin tag, 1/nth of input sample (where n=the number of distinct regions in target nucleic acid) is isolated for each pool chamber. The capture probe binds the target nucleic acid of interest. Then the target nucleic acid is immobilized, via the capture probe's biotin, to an avidin molecule adhered to the substrate. Optionally, the target nucleic acid is stretched, e.g., via flow or electrostatic force. All n-pools can be stretched-and-bound simultaneously, or, in order to maximize the number of fully stretched molecules, pool 1 (which captures most 5' region) can be stretched and bound first; then pool 2, (which captures the middle-of-target region) is then can be stretched and bound; finally, pool 3 is can be stretched and bound.

[0138] The number of distinct capture probes required is inversely related to the size of target nucleic acid fragment. In other word, more capture probes will be required for a highly-fragmented target nucleic acid. For sample types with highly fragmented and degraded target nucleic acids (e.g., Formalin-Fixed Paraffin Embedded Tissue) it may be useful to include multiple pools of capture probes. On the other hand, for samples with long target nucleic acid fragments, e.g., in vitro obtained isolated nucleic acids, a single capture probe at a 5' end may be sufficient.

[0139] The region of the target nucleic acid between two capture probes or after one capture probe and before a terminus of the target nucleic acid is referred herein as a “gap”. The gap is a portion of the target nucleic acid that is available to be bound by a sequencing probe of the present invention. The minimum gap is a target binding domain length (e.g., 4 to 10 nucleotides) and a maximum gap is the majority of a whole chromosome.

[0140] An immobilized target nucleic acid is shown in FIG. 12. Here, the two capture probes are identified as “5' capture probe” and “3' capture probe”.

[0141] FIG. 8A shows a schematic of a sequencing probe bound to a target nucleic acid. Here, the target nucleic acid has a thymidine (T). A first pool of complementary nucleic acids comprising a detectable label or reporter complexes is shown at the top, each member of the pool has a different detectable label (e.g., thymidine is identified by a green signal) and a different nucleotide sequence. The first nucleotide in the target binding domain binds the T in the target nucleic acid. The first attachment regions of the probe include one or more nucleotide sequence(s) that specifies that the first nucleotide in the probe's target binding domain binds a thymidine. Thus, only the complementary nucleic acid for thymidine binds the first position of the barcode domain. As shown, a thymidine-encoding first complementary nucleic acid comprising a detectable label or reported complexes comprising detectable labels are bound to attachment regions in the first position of the probe's barcode domain.

[0142] The number of pools of complementary nucleic acids or reporter complexes is identical to the number of positions in the barcode domain. Thus, for a barcode domain having six positions, six pools will be cycled over the probes.

[0143] Alternately, prior to contacting a target nucleic acid with a probe, the probe may be hybridized at its first position to a complementary nucleic acid comprising a detectable label or a reporter complex. Thus, when contacted with its target nucleic acid, the probe is capable of emitting a detectable signal from its first position and it is unnecessary to provide a first pool of complementary nucleic acids or reporter complexes that are directed to the first position on the barcode domain.

[0144] FIG. 8B continues the method shown in FIG. 8A. Here, the first complementary nucleic acids (or reporter complexes) for thymidine that were bound to attachment regions in the first position of the barcode domain have been replaced with a first hybridizing nucleic acid for thymidine and lacking a detectable label. The first hybridizing nucleic acid for thymidine and lacking a detectable label displaces the previously-bound complementary nucleic acids comprising a detectable label or the previously-bound reporter complexes. Thereby, position 1 of the barcode domain no longer emits a detectable signal.

[0145] In embodiments, the complementary nucleic acids comprising a detectable label or reporter complexes may be removed from the attachment region but not replaced with a hybridizing nucleic acid lacking a detectable label. This can occur, for example, by adding a chaotropic agent, increasing the temperature, changing salt concentration, adjusting pH, and/or applying a hydrodynamic force. In these embodiments fewer reagents (i.e., hybridizing nucleic acids lacking detectable labels) are needed.

[0146] FIG. 8C continues the method of the claimed invention. Here, the target nucleic acid has a cytidine (C) following its thymidine (T). A second pool of complementary nucleic

acids or reporter complexes is shown at the top, each member of the pool has a different detectable label and a different nucleotide sequence. Moreover, the nucleotide sequences for the complementary nucleic acids or complementary nucleic acids of the reporter complexes of the first pool are different from the nucleotide sequences for those of the second pool. However, the base specific detectable labels are common to the pools of complementary nucleic acids, e.g., thymidines are identified by green signals. Here, the second nucleotide in the target binding domain binds the C in the target nucleic acid. The second attachment regions of the probe have a nucleotide sequence that specifies that the second nucleotide in the probe's target binding domain binds a cytidine. Thus, only the complementary nucleic acids comprising a detectable label or reporter complexes from the second pool and for cytidine binds the second position of the barcode domain. As shown, the cytidine-encoding second complementary nucleic acid or reporter complex is bound at the second position of the probe's barcode domain.

[0147] In embodiments, the steps shown in FIG. 8C are subsequent to steps shown in FIG. 8B. Here, once the first pool of complementary nucleic acids or reporter complexes (of FIG. 8A) has been replaced with first hybridizing nucleic acids lacking a detectable label (in FIG. 8B), then a second pool of complementary nucleic acids or reporter complexes is provided (as shown in FIG. 8C). Alternately, the steps shown in FIG. 8C are concurrent with steps shown in FIG. 8B. Here, the first hybridizing nucleic acids lacking a detectable label (in FIG. 8B) are provided simultaneously with a second pool of complementary nucleic acids or reporter complexes (as shown in FIG. 8C).

[0148] FIG. 8D continues the method shown in FIG. 8C. Here, the first through fifth positions on the barcode domain were bound by complementary nucleic acids comprising a detectable labels or reporter complexes and have been replaced with hybridizing nucleic acids lacking detectable labels. The sixth position of the barcode domain is currently bound by a complementary nucleic acid comprising a detectable label or reporter complex, which identifies the sixth position in the target binding domain as being bound to a guanine (G).

[0149] As mentioned above, complementary nucleic acids comprising detectable labels or reporter complexes can be removed from attachment regions but not replaced with hybridizing nucleic acid lacking detectable labels.

[0150] If needed, the rate of detectable label exchange can be accelerated by incorporating small single-stranded oligonucleotides that accelerate the rate of exchange of detectable labels (e.g., “Toe-Hold” Probes; see, e.g., Seeling et al., “Catalyzed Relaxation of a Metastable DNA Fuel”; *J. Am. Chem. Soc.* 2006, 128(37), pp 12211-12220).

[0151] It is possible to replace the complementary nucleic acids or reporter complexes on a final position on a barcode domain (the sixth position in FIG. 8D); however, this may be unnecessary when a sequencing probe is to be replaced with another sequencing probe. Indeed, the sequencing probe of FIG. 8D can now be de-hybridized and removed from the target nucleic acid and replaced with a second (overlapping or non-overlapping) sequencing probe that has not yet been bound by any complementary nucleic acids, as shown in FIG. 8E. The probe in FIG. 8E may be included in a second population of probes.

[0152] Like FIGS. 8A to 8E, FIGS. 9A and 9D to 9G show method steps of the present invention; however, FIGS. 9A and

9D to 9G clearly show that reporter complexes (comprising detectable labels) are bound to attachment regions of sequencing probes. FIGS. 9D and 9E show fluorescent signals emitted from probes hybridized to reporter complexes. FIGS. 9D and 9E show that the target nucleic acid has a sequence of "T-A".

[0153] FIG. 10 summarizes the steps shown in FIGS. 9D and 9E. At the top of the figure is shown the nucleotide sequence of an exemplary probe and identifies significant domains of the probe. The probe includes an optional double-stranded DNA spacer between its target binding domain and its barcode domain. The barcode domain comprises, in order, a "Flank 1" portion, an "AR-1" portion, an "AR-1/Flank 2" portion, an "AR-2" portion, and an "AR-2/Flank 3" portion. In Step 1, the "AR-1 Detect" is hybridized to the probe's "AR-1" and "AR-1/Flank 2" portions. "AR-1 Detect" corresponds to a reporter complex or complementary nucleic acid comprising a detectable label that encodes a first position thymidine. Thus, Step 1 corresponds to FIG. 9D. In Step 2, the "Lack 1" is hybridized to the probe's "Flank 1" and "AR-1" portions. "Lack 1" corresponds to the hybridizing nucleic acid lacking a detectable label that is specific to the probe's first attachment region (as shown in FIG. 9E as a black bar covering the first attachment region). By hybridizing to the "Flank 1" position, which is 5' to the reporter complex or complementary nucleic acid, the hybridizing nucleic acid more efficiently displaces the reporter complex/complementary nucleic acid from the probe. The "Flank" portions are also known as "Toe-Holds". In Step 3, the "AR-2 Detect" is hybridized to the probe's "AR-2" and "AR-2/Flank 3" portions. "AR-2 Detect" corresponds to a reporter complex or complementary nucleic acid comprising a detectable label that encodes a second position Guanine. Thus, Step 3 corresponds to FIG. 9E. In this embodiment, hybridizing nucleic acid lacking a detectable label and complementary nucleic acids comprising detectable labels/reporter complexes are provided sequentially.

[0154] Alternately, hybridizing nucleic acid lacking a detectable label and complementary nucleic acids comprising detectable labels/reporter complexes are provided concurrently. This alternate embodiment is shown in FIG. 11. In Step 2, the "Lack 1" (hybridizing nucleic acid lacking a detectable label) is provided along with the "AR-2 Detect" (reporter complex that encodes a second position Guanine). This alternate embodiment may be more time effective than the embodiment illustrated in FIG. 10 because it combines two steps into one.

[0155] FIG. 12 illustrates the methods of the present invention. Here, a target nucleic acid is captured and immobilized at two positions, thereby producing a "gap" to which a probe is able to bind. A first population of probes is hybridized onto the target nucleic acid and detectable labels are detected. The initial steps are repeated with a second population of probes, a third population of probes, to more than 100 populations of probes. Use of about 100 populations of probes provides about 5x coverage of each nucleotide in a target nucleic acid. FIG. 12 provides estimated rates of read times based on the time required to detect signals from one Field of View (FOV).

[0156] The distribution of probes along a length of target nucleic acid is critical for resolution of detectable signal. As discussed above, the resolution limit for two detectable labels is about 600 nucleotides. Preferably, each sequencing probe in a population of probes will bind no closer than 600 nucleotides from each other. As discussed above, 600 nucleotides is

the resolution limit of a typical sequencing apparatus. In this case, a sequencing probe will provide a single read; this is shown in FIG. 12 in the left-most resolution-limited spot.

[0157] Randomly, but in part depending on the length of the target binding domain, the T_m of the probes, and concentration of probes applied, it is possible for two distinct sequencing probes in a population to bind within 600 nucleotides of each other. In this case, unordered multiple reads will emit from a single resolution-limited spot; this is shown in FIG. 12 in the second resolution-limited spot.

[0158] Alternately or additionally, the concentration of sequencing probes in a population may be reduced to decrease coverage of probes in a specific region of a target nucleic acid, e.g., to above the resolution limit of the sequencing apparatus, thereby producing a single read from a resolution-limited spot.

[0159] FIG. 23 shows a schematic of a sequencing probe distinct from that used in FIGS. 8 through 12. Here, each position on a barcode domain is bound by complementary nucleic acids comprising detectable labels or by reporter complexes. Thus, in this example, a six nucleotide sequence can be read without needing to sequentially replace complementary nucleic acids. Use of this sequencing probe would reduce the time to obtain sequence information since many steps of the described method are omitted. However, this probe would benefit from detectable labels that are non-overlapping, e.g., fluorophores are excited by non-overlapping wavelengths of light or the fluorophores emit non-overlapping wavelengths of light.

[0160] The method further comprising steps of assembling each identified linear order of nucleotides for each region of the immobilized target nucleic acid, thereby identifying a sequence for the immobilized target nucleic acid. The steps of assembling uses a non-transitory computer-readable storage medium with an executable program stored thereon. The program instructs a microprocessor to arrange each identified linear order of nucleotides for each region of the target nucleic acid, thereby obtaining the sequence of the nucleic acid. Assembling can occur in "real time", i.e., while data is being collected from sequencing probes rather than after all data has been collected.

[0161] Any of the above aspects and embodiments can be combined with any other aspect or embodiment as disclosed here in the Summary and/or Detailed Description sections.

DEFINITIONS

[0162] In certain exemplary embodiments, the terms "annealing" and "hybridization," as used herein, are used interchangeably to mean the formation of a stable duplex. In one aspect, stable duplex means that a duplex structure is not destroyed by a stringent wash under conditions such as a temperature of either about 5° C. below or about 5° C. above the T_m of a strand of the duplex and low monovalent salt concentration, e.g., less than 0.2 M, or less than 0.1 M or salt concentrations known to those of skill in the art. The term "perfectly matched," when used in reference to a duplex means that the polynucleotide and/or oligonucleotide strands making up the duplex form a double stranded structure with one another such that every nucleotide in each strand undergoes Watson-Crick base pairing with a nucleotide in the other strand. The term "duplex" comprises, but is not limited to, the pairing of nucleoside analogs, such as deoxyinosine, nucleosides with 2-aminopurine bases, PNAs, and the like, that may be employed. A "mismatch" in a duplex between two oligo-

nucleotides means that a pair of nucleotides in the duplex fails to undergo Watson-Crick bonding.

[0163] As used herein, the term “hybridization conditions,” will typically include salt concentrations of less than about 1 M, more usually less than about 500 mM and even more usually less than about 200 mM. Hybridization temperatures can be as low as 5° C., but are typically greater than 22° C., more typically greater than about 30° C., and often in excess of about 37° C. Hybridizations are usually performed under stringent conditions, e.g., conditions under which a probe will specifically hybridize to its target subsequence. Stringent conditions are sequence-dependent and are different in different circumstances. Longer fragments may require higher hybridization temperatures for specific hybridization. As other factors may affect the stringency of hybridization, including base composition and length of the complementary strands, presence of organic solvents and extent of base mismatching, the combination of parameters is more important than the absolute measure of any one alone.

[0164] Generally, stringent conditions are selected to be about 5° C. lower than the T_m for the specific sequence at a defined ionic strength and pH. Exemplary stringent conditions include salt concentration of at least 0.01 M to no more than 1 M Na ion concentration (or other salts) at a pH 7.0 to 8.3 and a temperature of at least 25° C. For example, conditions of 5×SSPE (750 mM NaCl, 50 mM Na phosphate, 5 mM EDTA, pH 7.4) and a temperature of 25–30° C. are suitable for allele-specific probe hybridizations. For stringent conditions, see for example, Sambrook, Fritsche and Maniatis, “Molecular Cloning A Laboratory Manual, 2nd Ed.” Cold Spring Harbor Press (1989) and Anderson Nucleic Acid Hybridization, 1st Ed., BIOS Scientific Publishers Limited (1999). As used herein, the terms “hybridizing specifically to” or “specifically hybridizing to” or similar terms refer to the binding, duplexing, or hybridizing of a molecule substantially to a particular nucleotide sequence or sequences under stringent conditions.

[0165] Detectable labels associated with a particular position of a probe can be “readout” (e.g., its fluorescence detected) once or multiple times; a “readout” may be synonymous with the term “basecall”. Multiple reads improve accuracy. A target nucleic acid sequence is “read” when a contiguous stretch of sequence information derived from a single original target molecule is detected; typically, this is generated via multi-pass consensus (as defined below). As used herein, the term “coverage” or “depth of coverage” refers to the number of times a region of target has been sequenced (via discrete reads) and aligned to a reference sequence. Read coverage is the total number of reads that map to a specific reference target sequence; base coverage is the total number of basecalls made at a specific genomic position.

[0166] As used herein, a “hybe and seq cycle” refers to all steps required to detect each attachment region on a particular probe or population of probes. For example, for a probe capable of detecting six positions on a target nucleic acid, one “hybe and seq cycle” will include, at least, hybridizing the probe to the target nucleic acid, hybridizing complementary nucleic acids/reporter complexes to attachment region at each of the six positions on the probe’s barcode domain, and detecting the detectable labels associated with each of the six positions.

[0167] The term “k-mer probe” is synonymous with a probe of the present invention.

[0168] When two or more sequences from discrete reads are aligned, the overlapping portions can be combined to create a single consensus sequence. In positions where overlapping portions have the same base (a single column of the alignment), those bases become the consensus. Various rules may be used to generate the consensus for positions where there are disagreements among overlapping sequences. A simple majority rule uses the most common base in the column as the consensus. A “multi-pass consensus” is an alignment of all discrete probe readouts from a single target molecule. Depending on the total number of cycles of probe populations/polls applied, each base position within a single target molecules can be queried with different levels of redundancy or overlap; generally, redundancy increases the confidence level of a basecall.

[0169] The “Raw Accuracy” is a measure of system’s inherent ability to correctly identify a base. Raw accuracy is dependent on sequencing technology. “Consensus Accuracy” is a measure of system’s ability to correctly identify a base with the use of additional reads and statistical power. “Specificity” refers to the percentage of reads that map to the intended targets out of total reads per run. “Uniformity” refers to the variability in sequence coverage across target regions; high uniformity correlates with low variability. This feature is commonly reported as the fraction of targeted regions covered by >20% of the average coverage depth across all targeted regions. Stochastic errors (i.e., intrinsic sequencing chemistry errors) can be readily corrected with ‘multi-pass’ sequencing of same target nucleic acid; given a sufficient number of passes, substantially ‘perfect consensus’ or ‘error-free’ sequencing can be achieved.

[0170] The methods described herein may be implemented and/or the results recorded using any device capable of implementing the methods and/or recording the results. Examples of devices that may be used include but are not limited to electronic computational devices, including computers of all types. When the methods described herein are implemented and/or recorded in a computer, the computer program that may be used to configure the computer to carry out the steps of the methods may be contained in any computer readable medium capable of containing the computer program. Examples of computer readable medium that may be used include but are not limited to diskettes, CD-ROMs, DVDs, ROM, RAM, non-transitory computer-readable media, and other memory and computer storage devices. The computer program that may be used to configure the computer to carry out the steps of the methods, assemble sequence information, and/or record the results may also be provided over an electronic network, for example, over the internet, an intranet, or other network.

[0171] A “Consumable Sequencing Card” (FIG. 24) can be incorporated into a fluorescence imaging device known in the art. Any fluorescence microscope with a number of varying features is capable of performing this sequencing readout. For instance: wide-field lamp, laser, LED, multi-photon, confocal or total-internal reflection illumination can be used for excitation and/or detection. Camera (single or multiple) and/or Photomultiplier tube (single or multiple) with either filter-based or grating-based spectral resolution (one or more spectrally resolved emission wavelengths) are possible on the emission-detection channel of the fluorescence microscope. Standard computers can control both the Consumable Sequencing Card, the reagents flowing through the Card, and detection by the fluorescence microscope.

[0172] The sequencing data can be analyzed by any number of standard next-generation-sequencing assemblers (see, e.g., Wajid and Serpedin, "Review of general algorithmic features for genome assemblers for next generation sequencers" *Genomics, proteomics & bioinformatics*, 10 (2), 58-73, 2012). The sequencing data obtained within a single diffraction limited region of the microscope is "locally-assembled" to generate a consensus sequence from the multiple reads within a diffraction spot. The multiple diffraction spot assembled reads are then mapped together to generate contiguous sequences representing the entire targeted gene set, or a de-novo assembly of entire genome(s).

[0173] Additional teaching relevant to the present invention are described in one or more of the following: U.S. Pat. No. 8,148,512, U.S. Pat. No. 7,473,767, U.S. Pat. No. 7,919,237, U.S. Pat. No. 7,941,279, U.S. Pat. No. 8,415,102, U.S. Pat. No. 8,492,094, U.S. Pat. No. 8,519,115, U.S. 2009/0220978, U.S. 2009/0299640, U.S. 2010/0015607, U.S. 2010/0261026, U.S. 2011/0086774, U.S. 2011/0145176, U.S. 2011/0201515, U.S. 2011/0229888, U.S. 2013/0004482, U.S. 2013/0017971, U.S. 2013/0178372, U.S. 2013/0230851, U.S. 2013/0337444, U.S. 2013/0345161, U.S. 2014/0005067, U.S. 2014/0017688, U.S. 2014/0037620, U.S. 2014/0087959, U.S. 2014/0154681, and U.S. 2014/0162251, each of which is incorporated herein by reference in their entireties.

EXAMPLES

Example 1

The Present Invention's Method of Sequencing a Target Nucleic Acid is Rapid

[0174] Below is described the timing for steps in the methods of the present invention and as shown in FIGS. 8 to 12.

[0175] The present invention requires minimal sample preparation. For example, as shown in FIG. 13, nucleic acids in a sample can begin to be read after 2 hours or less or preparation time; this is significantly less time required for Ion Torrent (AmpliSeg™) or Illumina (TruSight) sequencing, which, respectively, require about 12 or 9 hours of preparation time.

[0176] Calculations for an exemplary run are shown in FIG. 14 and calculations for cycling times are shown in FIG. 15.

[0177] Binding a population of probes to an immobilized target nucleic acid takes about sixty seconds. This reaction can be accelerated by utilizing multiple copies of the target binding domain on the synthetic backbone. With microfluidic-controlled fluid exchange device, washing away un-bound probes takes about a half a second.

[0178] Adding a first pool of complementary nucleic acids (comprising a detectable label) and binding them to attachment regions in the first position of the barcode domain takes about fifteen seconds.

[0179] Each field of view (FOV) is imaged for four different colors, each color representing a single-base. Fiducial spots placed on a 5' capture probe or 3' capture probe (or both) may be helpful for reading only those optical barcodes in-line (consistent with the presence of gapped target nucleic acid) between the two locations. Fiducial spots can also be added to each field of view in order to generate equal alignment of images upon successive steps in the sequencing process. All four images can be obtained at a single FOV and then the optical reading device may move to a new FOV, or take all

FOV in one color then reimage in a second color. A single FOV can be read in about a half a second. It takes about a half a second to move to a next FOV. Therefore, the time to read "n" FOV's equals "n" times 1 sec).

[0180] The complementary nucleic acids having detectable labels are removed from the first position of the barcode domain by addition of heat or washing with excess of complementary nucleic acids lacking detectable labels. If needed, the rate of detectable label exchange can be accelerated by incorporating small single-stranded oligonucleotides that accelerate the rate of exchange of detectable labels (e.g., "Toe-Hold" Probes; see, e.g., Seeling et al., "Catalyzed Relaxation of a Metastable DNA Fuel"; *J. Am. Chem. Soc.* 2006, 128(37), pp 12211-12220). A FOV can be reimaged to confirm that all complementary nucleic acids having detectable labels are removed before moving continuing. This takes about fifteen seconds. This step can be repeated until background signal levels are reached.

[0181] The above steps are repeated or the remaining positions in the probes' barcode domain.

[0182] The total time to read equals m (bases read) times (15 sec+n FOV's times 1 sec+15 sec). For example, when the number of positions in the barcode domain is 6 and 20 FOV's, the time to read equals 6x(30+20+15) or 390 seconds.

[0183] Probes of the first population are de-hybridized. This takes about sixty seconds.

[0184] The above steps are repeated for second and subsequent populations of probes. If populations of sequencing probes are organized by melting temperature (T_m), each population of probes will require multiple hybridizations to ensure that each base is covered to required depth (this is driven by error rate). Moreover, by analyzing the hybridization reads during a run, it is possible to recognize each individual gene that is being sequenced well before the entire sequence is actually determined. Hence cycling can be repeated until a particular desired error-frequency (or coverage) is met.

[0185] Using the timing described above, together with some gapped-nucleic acid binding density estimates, throughput of a Nanostring (NSTG)-Next Generation Sequencer of the present invention can be estimated.

[0186] Net throughput of sequencer is given by:

[0187] Fractional-Base-Occupancyx<gap-length>x number-of-gaps-per-FOVxnumber-of-bases-per-optical-barcode/[60 sec (hybridizing probes to target nucleic acid)+0.5 sec (wash)+m: positions in the barcode domainx(15 sec (binding complementary nucleic acids)+nFOV'sX1+15 sec (unbinding complementary nucleic acids))+60 sec (de-hybridizing probes to target nucleic acid)]

[0188] Therefore, in an example, a total "cycle" for a single gapped-nucleic acid (adding together from the method shown in FIG. 10):

[0189] 60 sec (hybridizing probes to target nucleic acid)+0.5 sec (wash)+m-basesx(15 sec (binding complementary nucleic acids)+nFOV's times 1+15 sec (unbinding complementary nucleic acids))+60 sec (de-hybridizing probes to target nucleic acid). Using m=6, nFOV's=20, yields time=60+0.5+390+60=510.5 sec.

[0190] Assuming: 1% occupancy of the gapped-nucleic acid region, 4000 bases per gap, and 5000 gapped nucleic-acid fragments per FOV and an m of 6 and nFOV's of 20 (as described above) yields a net throughput of:

[0191] $0.01 \times 4000 \times 5000 \times 20 = 4,000,000$ 6-base reads per 510.5 secs = 47,012.73 bases/sec.

[0192] Therefore, in this example, a net throughput per 24 hours of continuous measurement = 4.062 Gigabases (Gb) per day. Alternate estimates up to 12 Gb per day. See FIG. 12.

[0193] As shown in FIG. 14, the run-time required to sequence 100 different target nucleic acids (a "100-plex") is about 4.6 hours; the run-time required to sequence 1000 different target nucleic acids (a "1000-plex") is about 16 hours.

[0194] FIG. 16 compares the sequencing rate, number of reads, and clinical utility for the present invention and various other sequencing methods/apparatuses.

Example 2

The Present Invention's Method has a Low Error Rate

[0195] FIG. 17 shows that the present invention has a raw error rate of about 2.1%, when terminal positions are omitted.

[0196] For the claimed invention, an error rate associated with sequencing is related to the free-energy difference between a fully-matched (m+n)-mer and a single-base mismatch (m-1+n)-mer. The sum of m+n is the number of nucleotides in a target binding domain and m represents the number of positions in a barcode domain. An estimate of the selectivity of hybridization can be made using the equation (See, Owczarzy, R. (2005), *Biophys. Chem.*, 117:207-215 and Integrated DNA Technologies website: at the World Wide Web (www.idtdna.com/analyzer/Applications/Instructions/Default.aspx?AnalyzerDefinitions=true#Mismatch_hMelt-Temp);

$$\theta = 1 - \left(\frac{K_a([strand2] - [strand1]) - 1}{2K_a[strand2]} + \sqrt{\frac{K_a^2([strand1] - [strand2])^2 + 2K_a([strand1] + [strand2]) + 1}{2K_a[strand2]}} \right)$$

[0197] where K_a is the association equilibrium constant obtained from predicted thermodynamic parameters,

$$K_a = \exp\left(\frac{-(\Delta H^\circ - T\Delta S^\circ)}{RT}\right)$$

[0198] Theta represents the percent bound of the exact complement and the single base mismatch sequences, which are expected to be annealed to target at the specified hybridization temperature. The T is the hybridization temperature in Kelvins, ΔH° (enthalpy) and ΔS° (entropy) are the melting parameters calculated from the sequence and the published nearest neighbor thermodynamic parameters, R is the ideal gas constant ($1.987 \text{ cal} \cdot \text{K}^{-1} \cdot \text{mole}^{-1}$), $[strand/2]$ is the molar concentration of an oligonucleotide, and the constant of -273.15 converts temperature from Kelvin to degrees of Celsius. The most accurate, nearest-neighbor parameters were obtained from the following publications for DNA/DNA base pairs (See, Allawi, H., Santa Lucia, J. *Biochemistry*, 36, 10581), RNA/DNA base pairs (See, Sugimoto et al., *Biochemistry*, 34, 11211-6), RNA/RNA base pairs (See, Xia, T. et al., *Biochemistry*, 37, 14719),

[0199] As example of an estimate of the approximate error-rate expected from the NSTG-sequencer follows. For (m+n) equals 8-mer. Consider the following 8-mer barcode and its single-base mismatch.

5'ATCGTACG3'
(region to sequence)

3'TAGCATGC5'
(sequencing optical barcode with perfect match)

3'TAGTATGC5'
(sequencing optical barcode with single-base mismatch (G-T) pairing)

[0200] Using the IDT calculator based upon the above equations yields:

[0201] At 17.4° C. (the T_m of the perfect match case), (50%/0.3%) would be the ratio of the correct optical barcode hybridized to that sequence versus the incorrect barcode at the T_m , yielding an estimated error rate for that sequence to be 0.6%.

[0202] A very high GC content sequencing calculation yields:

5'CGCGCGCC3'
(region to sequence)

3'GCGGCGCG5'
(sequencing optical barcode with perfect match)

3'GCGGACCG5'
(sequencing optical barcode with single-base mismatch (G-A) mis-pairing)

[0203] At 41.9° C. (the T_m of the perfect match case), (50%/0.4%) would be the ratio of the correct optical barcode hybridized to that sequence versus the incorrect barcode at the T_m , yielding an estimated error rate for that sequence to be 0.8%.

[0204] Examination of a number of 8-mer pairs yields a distribution of error rates, in the range of 0.2% to 1%. While the above calculations will not be identical to the conditions used, these calculations provide an indication that the method of the present invention will have a relatively low intrinsic error rate, when compared to other single-molecule sequencing technologies, such as Pacific Biosciences and Oxford Nanopore Technologies where error rates can be significant (>>10%).

[0205] FIG. 18 demonstrates that the present invention's raw accuracy is higher than other sequencing methods. Thus, the present invention provides a consensus sequence from a single target after fewer passes than required for other sequencing methods. Additionally, the present invention may obtain "perfect consensus"/"error-free" sequencing (i.e., 99.9999%/Q60) after 30 or more passes whereas the PacBio sequencing methods (for example) cannot attain such a consensus after 70 passes.

Example 3

The Present Invention has Single Base-Pair Resolution Ability

[0206] FIG. 19 shows that the present invention has single-base resolution and with low error rates (ranging from 0% to 1.5% depending on a specific nucleotide substitution).

[0207] Additional experiments were performed using a target RNA hybridized with barcode and immobilized to the surface of cartridge using normal NanoString gene-expression binding technology (see, e.g., Geiss et al, "Direct multiplexed measurement of gene expression with color-coded probe pairs"; *Nature Biotechnology*, 26, 317-325 (2008)). The ability of a barcode with different target binding domain length and with a perfect match (YGBYGR-2 um optical bar code connected to perfect 10-mer match sequence) to hybridize to RNA-target was measured (FIG. 26). Longer length of target binding domain gives higher counts. It also shows that 10-mer target binding domain is enough to register the sequence above background. Each of the individual single-base altered matches was synthesized with alternate optical bar codes. The ratio of correct to incorrect optical barcodes was counted (FIGS. 24 and 25).

[0208] Ability of 10mer to detect a SNP the real sequence is >15000 counts over background, whilst incorrect sequences are at most >400 over background. In the presence of correct probe, error rates are expected to be <3% of real sequence. Note that this data is (in essence) a worse-case scenario. Having only a 10-base-pair hybridization sequence attached to a 6.6 Kilobase optical barcode reporter (Gen2 style). No specific condition optimizations were performed. This data, however, does reveal that the NanoString Next-Generation Sequencing approach is capable of resolving single-base pairs of sequence.

[0209] The detailed materials and methods utilized in the above study are as follows:

[0210] Hybridization Protocol Probe B plus codeset

[0211] Take 25 ul elements (194 codeset)

[0212] Add 5 ul Probe B+ complimentary sequence to target (100 uM)

[0213] Add 15 ul Hyb Buffer (14.56xSSPE 0.18% Tween 20)

[0214] SSPE (150 mM NaCl, NaH₂PO₄·xH₂O 10 mM, Na₂EDTA 10 mM)

[0215] Incubate on ice for 10 min

[0216] Add 150 ul G beads (40 ul G beads at 10 mg/ml plus 110 ul 5xSSPE 0.1% Tween 20)

[0217] Incubate for 10 min at RT

[0218] Wash three times with 0.1SSPE 0.1% Tween 20 using magnet collector

[0219] Elute in 100 ul 0.1xSSPE for 10 min at 45 C.

[0220] Target Hybridization protocol (750 mM NaCl)

[0221] Take 20 ul above eluted sample

[0222] Add 10 ul hyb buffer

[0223] Add 1 ul Target (100 nM biotinylated RNA)

[0224] Incubate on ice for 30 min

[0225] Take 15 ul and Bind to streptavidin slide for 20 min, flow stretch with G hooks, count using nCounter

[0226] Materials

[0227] Elements 194 codeset

[0228] Oligos bought from IDT

[0229] SSPE (150 mM NaCl, NaH₂PO₄·xH₂O 10 mM, Na₂EDTA 10 mM)

[0230] Hyb buffer (14.56xSSPE 0.18% Tween 20)

TABLE 2

Probe B Sequences for 12, 11, . . . , 8 mers. (SEQ ID NO: 30 to SEQ ID NO: 34)	
GCRYBG	5 GACTGTACCCACGCGATGACGTTCTGTCAGAGTCGCATAAT 3 CT
YRBYRG	5 AGACTGTACCACAAGAATCCCTGCTAGCTGAAGGAGGGTCA 3 AAC
YGBYGR	5 GAGACTGTACCCTACGTATATATCCAAGTGGTTATGTCCGA 3 CGGC
GCRYGB	5 TGAGACTGTACCACCCCTCCAAACGCATTCTTATTGGCAA 3 TGGAA
RYGBRG	5 CTGAGACTGTACCCGGGAATCGGCATTTCGCATTCTTAGGA 3 TCTAAA

TABLE 3

Target Sequence (in Bold; SEQ ID NO: 35)	
RNA	5 CAATGTGAGTCTCTTGGT TACAGTCTCAGT CTAGTCACTCCCTA 3 AG\Bio TRG\

TABLE 4

Probe B Sequences for 10 mer mismatches (in Bold; SEQ ID NO: 36 to SEQ ID NO: 41)	
10mermis2A	GAGACAGTACCCCTGGTCTAGGTATCTAATTCGTGGGTC GGTACT
10mermis2C	GAGACCGTACCGCTCATTTTGAACATACGATTGCGATT ACGGAAA
10mermis2G	GAGACGGTACCTTAAAGCTATCCACGAATGTCAAAAAT GTGGTTT
10mermis1G	GAGAGTGTACCCAATGCTTGCAATATGTATCCTGATCG TGCGTGC
10mermis1A	GAGAATGTACCCTCATACCAATGTAAAGTATAGTTAAC GCCCTGT
10mermis1T	GAGATTGTACCCTACATATATAGGAAAAGGGAAGGTAG AAGAGCT

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 104

<210> SEQ ID NO 1

<211> LENGTH: 10

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 1

atacatctag 10

<210> SEQ ID NO 2
<211> LENGTH: 10
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 2

gatctacata 10

<210> SEQ ID NO 3
<211> LENGTH: 10
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 3

ttaggtaaag 10

<210> SEQ ID NO 4
<211> LENGTH: 10
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 4

tcttcattac 10

<210> SEQ ID NO 5
<211> LENGTH: 10
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 5

atgaatctac 10

<210> SEQ ID NO 6
<211> LENGTH: 10
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 6

tcaatgtatg 10

<210> SEQ ID NO 7
<211> LENGTH: 10
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 7

-continued

aattgagtag	10
 <210> SEQ ID NO 8 <211> LENGTH: 10 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic Polynucleotide <400> SEQUENCE: 8	
atgttaatgg	10
 <210> SEQ ID NO 9 <211> LENGTH: 10 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic Polynucleotide <400> SEQUENCE: 9	
aattaggatg	10
 <210> SEQ ID NO 10 <211> LENGTH: 10 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic Polynucleotide <400> SEQUENCE: 10	
ataatggatc	10
 <210> SEQ ID NO 11 <211> LENGTH: 10 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic Polynucleotide <400> SEQUENCE: 11	
taataagggtg	10
 <210> SEQ ID NO 12 <211> LENGTH: 10 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic Polynucleotide <400> SEQUENCE: 12	
tagtttagagc	10
 <210> SEQ ID NO 13 <211> LENGTH: 10 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic Polynucleotide <400> SEQUENCE: 13	
atagagaagg	10

-continued

<210> SEQ ID NO 14
<211> LENGTH: 10
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 14

ttgatgatac 10

<210> SEQ ID NO 15
<211> LENGTH: 10
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 15

atagtgattc 10

<210> SEQ ID NO 16
<211> LENGTH: 10
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 16

tataacgatg 10

<210> SEQ ID NO 17
<211> LENGTH: 10
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 17

ttaagtttag 10

<210> SEQ ID NO 18
<211> LENGTH: 10
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 18

atacgttatg 10

<210> SEQ ID NO 19
<211> LENGTH: 10
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 19

tgtaactatag 10

<210> SEQ ID NO 20
<211> LENGTH: 10
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 20

ttaacaagtg 10

<210> SEQ ID NO 21
<211> LENGTH: 10
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 21

aactatgtac 10

<210> SEQ ID NO 22
<211> LENGTH: 10
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 22

taactatgac 10

<210> SEQ ID NO 23
<211> LENGTH: 10
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 23

actaatgttc 10

<210> SEQ ID NO 24
<211> LENGTH: 10
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 24

toattgaatg 10

<210> SEQ ID NO 25
<211> LENGTH: 14
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 25

ctgtctcacc tctt 14

<210> SEQ ID NO 26
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 26

-continued

ctgtctcacc tcttgctgca tctgtg 26

<210> SEQ ID NO 27
<211> LENGTH: 38
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 27

ctgtctcacc tcttgctgca tctgtgctgt tcaagttg 38

<210> SEQ ID NO 28
<211> LENGTH: 36
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 28

ctgtctcacc ttgtctgcat ctgtcgggtc acgttg 36

<210> SEQ ID NO 29
<211> LENGTH: 36
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 29

ctgtctcatt ttgtctgcat ctgtcgggtc acgttg 36

<210> SEQ ID NO 30
<211> LENGTH: 43
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 30

gactgtaccc acgcgatgac gttcgtcaag agtcgcataa tct 43

<210> SEQ ID NO 31
<211> LENGTH: 44
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 31

agactgtacc acaagaatcc ctgctagctg aaggagggtc aaac 44

<210> SEQ ID NO 32
<211> LENGTH: 45
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 32

gagactgtac cctacgtata tatccaagt gttatgtcgc acggc 45

-continued

<210> SEQ ID NO 33
<211> LENGTH: 46
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 33

tgagactgta ccaccctcc aaacgcattc ttattggcaa atggaa 46

<210> SEQ ID NO 34
<211> LENGTH: 47
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 34

ctgagactgt acccggaat cggcatttcg cattcttagg atctaaa 47

<210> SEQ ID NO 35
<211> LENGTH: 44
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 35

caatgtgagt ctcttggtac agtcctcagtt agtcactccc taag 44

<210> SEQ ID NO 36
<211> LENGTH: 45
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 36

gagacagtac cctgggtctag gtatctaatt cgtgggtcgg gtact 45

<210> SEQ ID NO 37
<211> LENGTH: 45
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 37

gagacgtac cgtcattttt gaacatacga ttgcgattac ggaaa 45

<210> SEQ ID NO 38
<211> LENGTH: 45
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 38

gagacggtag cttaaagcta tccacgaatg tcaaaaatgt ggttt 45

<210> SEQ ID NO 39
<211> LENGTH: 45
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence

-continued

```

<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 39

gagagtgtac ccaatgcttg cagtatgtat cctgatcgtg cgtgc          45

<210> SEQ ID NO 40
<211> LENGTH: 45
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 40

gagaatgtac cctcattacca atgtaaagta tagttaacgc cctgt          45

<210> SEQ ID NO 41
<211> LENGTH: 45
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 41

gagattgtac cctacatata taggaaaagg gaaggtagaa gagct          45

<210> SEQ ID NO 42
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 42

cacgaacgtc ag          12

<210> SEQ ID NO 43
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 43

catcgcatgc ct          12

<210> SEQ ID NO 44
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 44

gtcatctctc ac          12

<210> SEQ ID NO 45
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 45

```

-continued

gtcatccgct ac 12

<210> SEQ ID NO 46
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 46

gtcatcgact ac 12

<210> SEQ ID NO 47
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 47

gtcatcttct ac 12

<210> SEQ ID NO 48
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 48

gtcatcacct ac 12

<210> SEQ ID NO 49
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 49

gtcatcactc ac 12

<210> SEQ ID NO 50
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 50

gtcatcttcg ac 12

<210> SEQ ID NO 51
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 51

gtcatcaact ac 12

-continued

<210> SEQ ID NO 52
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 52

gtcatccgta ac 12

<210> SEQ ID NO 53
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 53

gtcatccgaa ac 12

<210> SEQ ID NO 54
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 54

gtcatcacia ac 12

<210> SEQ ID NO 55
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 55

gtcatcttgc ac 12

<210> SEQ ID NO 56
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 56

gtcatcttgc ct 12

<210> SEQ ID NO 57
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 57

gtcatccgtc ct 12

<210> SEQ ID NO 58
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 58

cttttcacct ct 12

<210> SEQ ID NO 59
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 59

cttttctctt ct 12

<210> SEQ ID NO 60
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 60

cttttgcact ct 12

<210> SEQ ID NO 61
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 61

ctttttgtct ct 12

<210> SEQ ID NO 62
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 62

ctttttgtga ct 12

<210> SEQ ID NO 63
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 63

ctttttgtg ct 12

<210> SEQ ID NO 64
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 64

-continued

cttttctgtc ct 12

<210> SEQ ID NO 65
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 65

cttttcactc ct 12

<210> SEQ ID NO 66
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 66

cttttctgtc ct 12

<210> SEQ ID NO 67
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 67

cttttctgtc ct 12

<210> SEQ ID NO 68
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 68

cttttctgtc ct 12

<210> SEQ ID NO 69
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 69

cttttctgtc ct 12

<210> SEQ ID NO 70
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 70

aggcatgcga tg 12

-continued

<210> SEQ ID NO 71
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 71

aggcattgtg ct 12

<210> SEQ ID NO 72
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 72

aggcattgct ct 12

<210> SEQ ID NO 73
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 73

aggcatttct ac 12

<210> SEQ ID NO 74
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 74

aggcatacct ac 12

<210> SEQ ID NO 75
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 75

aggcatttgc ac 12

<210> SEQ ID NO 76
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 76

aggcatcgct ct 12

<210> SEQ ID NO 77
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence

-continued

```

<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 77

tcctgtcggg tc                                     12

<210> SEQ ID NO 78
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 78

gttcaatgct ct                                     12

<210> SEQ ID NO 79
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 79

attcgggtgct ct                                    12

<210> SEQ ID NO 80
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 80

gatgcctgct ct                                    12

<210> SEQ ID NO 81
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 81

tttgcttgcct ct                                    12

<210> SEQ ID NO 82
<211> LENGTH: 100
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 82

ttcaactgtag ctgtctcatt ttgtgcata ctgtcggg acgttggagg ttgtaacgg 60
tcctcttttc actcctaggc attgcctat tcggcgctct 100

<210> SEQ ID NO 83
<211> LENGTH: 10
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

```

-continued

<400> SEQUENCE: 83
cgatctggtt 10

<210> SEQ ID NO 84
<211> LENGTH: 10
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 84
cgatctggtt 10

<210> SEQ ID NO 85
<211> LENGTH: 10
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 85
gctagaccaa 10

<210> SEQ ID NO 86
<211> LENGTH: 10
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 86
gctggaccaa 10

<210> SEQ ID NO 87
<211> LENGTH: 10
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 87
gctcgaccaa 10

<210> SEQ ID NO 88
<211> LENGTH: 10
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 88
gcttgaccaa 10

<210> SEQ ID NO 89
<211> LENGTH: 10
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 89
gagactgtac 10

-continued

<210> SEQ ID NO 90
<211> LENGTH: 10
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 90

gagacagtac 10

<210> SEQ ID NO 91
<211> LENGTH: 10
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 91

gagaccgtac 10

<210> SEQ ID NO 92
<211> LENGTH: 10
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 92

gagacggtac 10

<210> SEQ ID NO 93
<211> LENGTH: 10
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 93

gagagtgtac 10

<210> SEQ ID NO 94
<211> LENGTH: 10
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 94

gagaatgtac 10

<210> SEQ ID NO 95
<211> LENGTH: 10
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 95

gagattgtac 10

<210> SEQ ID NO 96
<211> LENGTH: 10

-continued

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 96

gagactgtac 10

<210> SEQ ID NO 97
<211> LENGTH: 10
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 97

gagattgtac 10

<210> SEQ ID NO 98
<211> LENGTH: 10
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 98

gagaccgtac 10

<210> SEQ ID NO 99
<211> LENGTH: 10
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 99

gagagtgtac 10

<210> SEQ ID NO 100
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 100

ctgagactgt ac 12

<210> SEQ ID NO 101
<211> LENGTH: 11
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 101

tgagactgta c 11

<210> SEQ ID NO 102
<211> LENGTH: 10
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Polynucleotide

-continued

<400> SEQUENCE: 102

gagactgtac

10

<210> SEQ ID NO 103

<211> LENGTH: 12

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 103

catgtcagag tc

12

<210> SEQ ID NO 104

<211> LENGTH: 11

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Polynucleotide

<400> SEQUENCE: 104

catgtcagag t

11

1-81. (canceled)

82. A method for sequencing a nucleic acid comprising steps of:

- (1) hybridizing at least one sequencing probe to a target nucleic acid that is immobilized to a substrate; wherein said sequencing probe comprises:

a target binding domain and a barcode domain;

wherein said target binding domain comprises at least four nucleotides and is capable of binding the immobilized target nucleic acid;

wherein said barcode domain comprises a synthetic backbone, said barcode domain comprising at least a first attachment region, said first attachment region comprising a nucleic acid sequence capable of being bound by a first complementary nucleic acid molecule and wherein said nucleic acid sequence of said first attachment region determines the position and identity of a first nucleotide in said immobilized target nucleic acid that is bound by a first nucleotide of said target binding domain and

- (2) binding to the first attachment region a first complementary nucleic acid molecule comprising a detectable label or a first complementary nucleic acid molecule of a first reporter complex comprising a detectable label;
- (3) detecting the detectable label of the bound first complementary nucleic acid molecule or the detectable label of the bound first complementary nucleic acid molecule of the first reporter complex;
- (4) identifying the position and identity of the first nucleotide in the immobilized target nucleic acid;
- (5) contacting the first attachment region with a first hybridizing nucleic acid molecule lacking a detectable label thereby unbinding the first complementary nucleic acid molecule and binding to the first attachment region the first hybridizing nucleic acid molecule lacking a detectable label;

- (6) binding to a second attachment region a second complementary nucleic acid molecule comprising a detectable label or a second complementary nucleic acid molecule of a second reporter complex comprising a detectable label, said second attachment region comprising a nucleic acid sequence that determines the position and identity of a second nucleotide in the immobilized target nucleic acid that is bound by a second nucleotide of the target binding domain;

- (7) detecting the detectable label of the bound second complementary nucleic acid molecule or the detectable label of the bound second complementary nucleic acid molecule of the second reporter complex; and

- (8) identifying the position and identity of the second nucleotide in the immobilized target nucleic acid.

83. (canceled)

84. The method of claim 82, wherein steps (5) and (6) occur sequentially or concurrently.

85. The method of claim 82, wherein steps (5) to (8) are repeated until each attachment region in the barcode domain has been sequentially bound by a complementary nucleic acid molecule comprising a detectable label or a complementary nucleic acid molecule of a reporter complex comprising a detectable label, and the detectable label of the sequentially bound complementary nucleic acid molecule or the detectable label of the sequentially bound complementary nucleic acid molecule of a reporter complex has been detected,

thereby identifying the linear order of nucleotides for a region of the immobilized target nucleic acid that was hybridized by the target binding domain of the sequencing probe.

86. The method of claim 82, wherein the target nucleic acid is first immobilized to a substrate by at least binding a first position of the target nucleic acid with a first capture probe that comprises a first affinity tag that selectively binds to a substrate.

87. The method of claim 86, wherein the target nucleic acid is elongated by applying a force sufficient to extend the target nucleic acid that is immobilized to a substrate at a first position.

88. The method of claim 87, wherein the force is gravity, hydrodynamic force, electromagnetic force, flow-stretching, a receding meniscus technique, or combinations thereof.

89. The method of claim 88, wherein the target nucleic acid is further immobilized to a substrate by binding an at least second position of the target nucleic acid with an at least second capture probe that comprises an affinity tag that selectively binds to the substrate.

90. (canceled)

91. The method of claim 89, wherein the force can be removed once the second position of the target nucleic acid is immobilized to the substrate.

92. (canceled)

93. The method of claim 91, wherein said immobilized target nucleic acid is elongated.

94. The method of claim 82, wherein said synthetic backbone comprises a polysaccharide, a polynucleotide, a peptide, a peptide nucleic acid, or a polypeptide.

95. The method of claim 94, wherein said synthetic backbone comprises single stranded-stranded DNA or single-stranded RNA or single-stranded PNA.

96. The method of claim 95, wherein said sequencing probe comprises a double-stranded DNA spacer between the target binding domain and the barcode domain.

97. The method of claim 82, wherein said first attachment region is adjacent to at least one flanking single-stranded polynucleotide.

98. The method of claim 82, wherein the first complementary nucleic acid is RNA, DNA or PNA or other polynucleotide analogue.

99. The method of claim 82, wherein the first nucleotide in said target binding domain is a modified nucleotide or a nucleic acid analogue.

100. The method of claim 82, wherein said barcode domain comprises at least a second attachment region, said second attachment region comprising a nucleic acid sequence capable of being bound by a second complementary nucleic acid molecule and wherein said nucleic acid sequence of said second attachment region determines the position and identity of a second nucleotide in said immobilized target nucleic acid that is bound by a second nucleotide of said target binding domain and

wherein the first complementary nucleic acid molecule is different from the second complementary nucleic acid molecule.

101-103. (canceled)

104. The method of claim 82, wherein the first complementary nucleic acid molecule and the first hybridizing nucleic acid molecule lacking a detectable label comprise the same nucleic acid sequence.

105. The method of claim 97, wherein the first hybridizing nucleic acid molecule lacking a detectable label comprises a nucleic acid sequence complementary to a flanking single-stranded polynucleotide adjacent to said first attachment region.

106-113. (canceled)

114. The method of claim 82, wherein the number of nucleotides in a target binding domain equals the number of attachment regions in the barcode domain.

115. The method of claim 82, wherein the number of nucleotides in a target binding domain is at least one more than the number of attachment regions in the barcode domain.

116. The method of claim 82, wherein at least the first attachment region branches from the synthetic backbone.

117-123. (canceled)

124. The method of claim 82, wherein at least one position in a barcode domain has a greater number of attachment regions as another position.

125-127. (canceled)

128. The method of claim 82, wherein each reporter complex comprising a detectable label comprises a complementary nucleic acid molecule directly linked to a primary nucleic acid molecule or indirectly linked to a primary nucleic acid molecule via a nucleic acid spacer.

129-134. (canceled)

135. The method of claim 128, wherein each primary nucleic acid molecule is hybridized to at least one secondary nucleic acid molecule.

136-139. (canceled)

140. The method of claim 135, wherein the at least one secondary nucleic acid molecule comprises at least one detectable label.

141. The method claim 135, wherein each secondary nucleic acid molecule is hybridized to at least one tertiary nucleic acid molecule comprising at least one detectable label.

142-147. (canceled)

148. The method of claim 135, wherein at least one secondary nucleic acid molecule comprises a region that does not hybridize to a primary nucleic acid molecule and does not hybridize to a tertiary nucleic acid molecule.

149. (canceled)

150. The method of claim 148, wherein the region that does not hybridize to a primary nucleic acid molecule and does not hybridize to a tertiary nucleic acid molecule comprises the nucleotide sequence of the complementary nucleic acid molecule that hybridizes to the primary nucleic acid molecule.

151. The method of claim 150, wherein the region that does not hybridize to a primary nucleic acid molecule and does not hybridize to a tertiary nucleic acid molecule is located at a terminus of the secondary nucleic acid molecule.

152-153. (canceled)

154. A method for sequencing a nucleic acid comprising steps of:

(1) hybridizing a first population of sequencing probes to a target nucleic acid that is immobilized to a substrate, wherein each sequencing probe in the first population comprises:

a target binding domain and a barcode domain;

wherein said target binding domain comprises at least four nucleotides and is capable of binding a target nucleic acid;

wherein said barcode domain comprises a synthetic backbone, said barcode domain comprising a first attachment region, said first attachment region comprising a nucleic acid sequence capable of being bound by a first complementary nucleic acid molecule and wherein said nucleic acid sequence of said first attachment region determines the position and identity of a first nucleotide in said target nucleic acid that is bound by a first nucleotide of said target binding domain and

said barcode domain further comprising at least a second attachment region, said second attachment region comprising a nucleic acid sequence capable of being bound by a second complementary nucleic acid molecule and wherein said nucleic acid sequence of said second attachment region determines the position and identity of a second nucleotide in said target nucleic acid that is bound by a second nucleotide of said target binding domain and

wherein the first complementary nucleic acid molecule is different from the second complementary nucleic acid molecule;

wherein each sequencing probe in the first population de-hybridizes from the immobilized target nucleic acid under about the same conditions;

- (2) binding to a first attachment region in each sequencing probe in the first population a plurality of first complementary nucleic acid molecules each comprising a detectable label or a plurality of first complementary nucleic acid molecules of a plurality of first reporter complexes each complex comprising a detectable label;
- (3) detecting the detectable label of each bound first complementary nucleic acid molecule or of each first reporter complex;
- (4) identifying the position and identity of a plurality of first nucleotides in the immobilized target nucleic acid hybridized by sequencing probes in the first population;
- (5) contacting each first attachment region of each sequencing probe of the first population with a plurality of first hybridizing nucleic acid molecules each lacking a detectable label thereby unbinding the first complementary nucleic acid molecules comprising a detectable label or the first complementary nucleic acid molecules of each first reporter complex and binding to each first attachment region a first hybridizing nucleic acid molecule lacking a detectable label;
- (6) binding to a second attachment region in each sequencing probe in the first population a plurality of second complementary nucleic acid molecules each comprising

a detectable label or a plurality of second complementary nucleic acid molecules of a plurality of second reporter complexes each complex comprising a detectable label;

- (7) detecting the detectable label of each bound second complementary nucleic acid molecule or of each second reporter complex;
- (8) identifying the position and identity of a plurality of second nucleotides in the immobilized target nucleic acid hybridized by sequencing probes in the first population; and
- (9) repeating steps (5) to (8) until each nucleotide in the immobilized target nucleic acid corresponding to the target binding domain of each sequencing probe in the first population has been identified.

155-191. (canceled)

192. An apparatus for performing the method of claim 82.

193-195. (canceled)

196. A sequencing probe comprising a target binding domain and a barcode domain;

wherein said target binding domain comprises at least four nucleotides and is capable of binding a target nucleic acid;

wherein said barcode domain comprises a synthetic backbone, said barcode domain comprising at least a first attachment region, said first attachment region comprising a nucleic acid sequence capable of being bound by a first complementary nucleic acid molecule and wherein said nucleic acid sequence of said first attachment region determines the position and identity of a first nucleotide in said target nucleic acid that is bound by a first nucleotide of said target binding domain.

197. A kit comprising a substrate, a plurality of sequencing probes of claim 196, at least one capture probe, at least one complementary nucleic acid molecule comprising a detectable label, at least one complementary nucleic acid molecule which lacks a detectable label, and instructions for use.

* * * * *

