

FOREWORD TO THE

NINTH EDITION



PDR enters its fifty-ninth year offering a wide range of pharmaceutical reference options than ever before. Available unaltered—in print, on CD-ROM, and on the Internet—PDR also provides essential prescription information in other forms as well, including the PDR mobile app.

Physicians' Desk Reference, the publisher does not warrant, guarantee, or assume any liability for the products described, or for any independent analysis in connection with any of the products described. The publisher does not assume any liability for the products described, or for any independent analysis in connection with any of the products described.

ABOUT THIS BOOK

Physicians' Desk Reference is a comprehensive, authoritative, and up-to-date source of information on the drugs and other products marketed without restriction under current FDA policies. Some dietary supplements and other products are also included.

Physicians' Desk Reference is a comprehensive, authoritative, and up-to-date source of information on the drugs and other products marketed without restriction under current FDA policies. Some dietary supplements and other products are also included.

PHYSICIANS' DESK REFERENCE®

Executive Vice President, PDR: David Duplay

Vice President, Sales and Marketing: Dikran N. Barsamian
Senior Director, Pharmaceutical Sales: Anthony Sorce
National Account Manager: Marion Reid, RPh
Senior Account Managers: Frank Karkowsky, Suzanne E. Yarrow, RN
Account Managers: Marjorie A. Jaxel, Kevin McGlynn, Elaine Musco, Lois Smith, Eileen Sullivan, Richard Zwickel
Senior Director, Brand and Product Management: Valerie E. Berger
Director, Brand and Product Management: Carmen Mazzatta
Associate Product Managers: Michael Casale, Andrea Colavecchio
Senior Director, Publishing Sales and Marketing: Michael Bennett
Director of Trade Sales: Bill Gaffney
Associate Director of Marketing: Jennifer M. Fronzaglia
Senior Marketing Manager: Kim Marich
Direct Mail Manager: Lorraine M. Loening
Manager of Marketing Analysis: Dina A. Maeder
Promotion Manager: Linda Levine
Vice President, Regulatory Affairs: Mukesh Mehta, RPh
Vice President, PDR Services: Brian Holland
Director of PDR Operations: Jeffrey D. Schaefer
Director of Operations: Robert Klein
Clinical Content Operations Manager: Thomas Fleming, PharmD

Manager, Editorial Services: Bette LaGow
Drug Information Specialists: Min Ko, PharmD; Greg Tallis, RPh
Project Editors: Neil Chesanow, Harris Fleming
Senior Editor: Lori Murray
Production Editor: Gwynned L. Kelly
Manager, Production Purchasing: Thomas Westburgh
Production Manager: Gayle Graizzaro
Production Specialist: Christina Klinger
Senior Production Coordinator: Gianna Caradonna
Production Coordinator: Yasmin Hernández
Senior Index Editors: Noel Deloughery, Shannon Reilly
Format Editor: Michelle S. Guzman
Traffic Assistant: Kim Condon
PDR Sales Coordinators: Nick W. Clark, Gary Lew
Production Design Supervisor: Adeline Rich
Senior Electronic Publishing Designer: Livio Udina
Electronic Publishing Designers: Bryan C. Dix, Rosalia Sberna
Production Associate: Joan K. Akerlind
Digital Imaging Manager: Christopher Husted
Digital Imaging Coordinator: Michael Labryere
Finance Director: Mark S. Ritchin
Director of Client Services: Stephanie Struble



Copyright © 2005 and published by Thomson PDR at Montvale, NJ 07645-1742. All rights reserved. None of the content of this publication may be reproduced, stored in a retrieval system, resold, redistributed, or transmitted in any form or by any means (electronic, mechanical, photocopying, recording, or otherwise) without the prior written permission of the publisher. Physicians' Desk Reference®, PDR®, Pocket PDR®, PDR Family Guide to Prescription Drugs®, PDR Family Guide to Women's Health and Prescription Drugs®, and PDR Family Guide to Nutrition and Health® are registered trademarks used herein under license. PDR® for Ophthalmic Medicines, PDR® for Nonprescription Drugs and Dietary Supplements, PDR® Companion Guide, PDR® Pharmacopoeia, PDR® for Herbal Medicines, PDR® for Nutritional Supplements, PDR® Medical Dictionary, PDR® Nurse's Drug Handbook, PDR® Nurse's Dictionary, PDR® Family Guide Encyclopedia of Medical Care, PDR® Family Guide to Natural Medicines and Healing Therapies, PDR® Family Guide to Common Ailments, PDR® Family Guide to Over-the-Counter Drugs, PDR® Family Guide to Nutritional Supplements, and PDR® Electronic Library are trademarks used herein under license.

Officers of Thomson Healthcare, Inc.: President and Chief Executive Officer: Robert Cullen; Chief Financial Officer: Paul Hilger; Chief Technology Officer: Fred Lauber; Executive Vice President, Medical Education: Jeff MacDonald; Executive Vice President, Micromedex: Jeff Reihl; Executive Vice President, PDR: David Duplay; Senior Vice President, Business Development: Robert Christopher; Senior Vice President, Marketing: Timothy Murray; Vice President, Human Resources: Pamela M. Bilash

ISBN: 1-56363-497-X

CELLTRION EXHIBIT - 1034

monitoring of all children treated with diuretics is indicated. See CONTRAINDICATIONS, WARNINGS, PRECAUTIONS.

Geriatric Use: Clinical studies of ZAROXOLYN did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy. This drug is known to be substantially excreted by the kidney, and the risk of toxic reactions to this drug may be greater in patients with impaired renal function. Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection, and it may be useful to monitor renal function.

ADVERSE REACTIONS

ZAROXOLYN is usually well tolerated, and most reported adverse reactions have been mild and transient. Many ZAROXOLYN related adverse reactions represent extensions of its expected pharmacologic activity and can be attributed to either its antihypertensive action or its renal/metabolic actions. The following adverse reactions have been reported. Several are single or comparably rare occurrences. Adverse reactions are listed in decreasing order of severity within body systems.

Cardiovascular: Chest pain/discomfort, orthostatic hypotension, excessive volume depletion, hemoconcentration, venous thrombosis, palpitations.

Central and Peripheral Nervous System: Syncope, neuropathy, vertigo, paresthesias, psychotic depression, impotence, dizziness/lightheadedness, drowsiness, fatigue, weakness, restlessness (sometimes resulting in insomnia), headache.

Dermatologic/Hypersensitivity: Toxic epidermal necrolysis (TEN), Stevens-Johnson Syndrome, necrotizing angitis (cutaneous vasculitis), skin necrosis, purpura, petechiae, dermatitis (photosensitivity), urticaria, pruritus, skin rashes.

Gastrointestinal: Hepatitis, intrahepatic cholestatic jaundice, pancreatitis, vomiting, nausea, epigastric distress, diarrhea, constipation, anorexia, abdominal bloating, abdominal pain.

Hematologic: Aplastic/hypoplastic anemia, agranulocytosis, leukopenia, thrombocytopenia.

Metabolic: Hypokalemia, hyponatremia, hyperuricemia, hypochloremia, hypochloremic alkalosis, hyperglycemia, glycosuria, increase in serum urea nitrogen (BUN) or creatinine, hypophosphatemia, hypomagnesemia, hypercalcemia.

Musculoskeletal: Joint pain, acute gouty attacks, muscle cramps or spasms.

Other: Transient blurred vision, chills, dry mouth.

In addition, adverse reactions reported with similar antihypertensive diuretics, but which have not been reported to date for ZAROXOLYN include: bitter taste, sialadenitis, xanthopsia, respiratory distress (including pneumonitis), and anaphylactic reactions. These reactions should be considered as possible occurrences with clinical usage of ZAROXOLYN.

Whenever adverse reactions are moderate or severe, ZAROXOLYN dosage should be reduced or therapy withdrawn.

OVERDOSAGE

Intentional overdosage has been reported rarely with metolazone and similar diuretic drugs.

Signs and Symptoms: Orthostatic hypotension, dizziness, drowsiness, syncope, electrolyte abnormalities, hemoconcentration and hemodynamic changes due to plasma volume depletion may occur. In some instances depressed respiration may be observed. At high doses, lethargy of varying degree may progress to coma within a few hours. The mechanism of CNS depression with thiazide overdosage is unknown. Also, GI irritation and hypermotility may occur. Temporary elevation of BUN has been reported, especially in patients with impairment of renal function. Serum electrolyte changes and cardiovascular and renal function should be closely monitored.

Treatment: There is no specific antidote available but immediate evacuation of stomach contents is advised. Dialysis is not likely to be effective. Care should be taken when evacuating the gastric contents to prevent aspiration, especially in the stuporous or comatose patient. Supportive measures should be initiated as required to maintain hydration, electrolyte balance, respiration, and cardiovascular and renal function.

DOSAGE AND ADMINISTRATION

Effective dosage of ZAROXOLYN should be individualized according to indication and patient response. A single daily dose is recommended. Therapy with ZAROXOLYN should be titrated to gain an initial therapeutic response and to determine the minimal dose possible to maintain the desired therapeutic response.

Usual Single Daily Dosage Schedules: Suitable initial dosages will usually fall in the ranges given.

Edema of cardiac failure:

ZAROXOLYN 5 to 20 mg once daily.

Edema of renal disease:

ZAROXOLYN 5 to 20 mg once daily.

Mild to moderate essential hypertension:

ZAROXOLYN 2½ to 5 mg once daily.

New patients—MYKROX® Tablets (metolazone tablets, USP) (see MYKROX package circular). If considered desirable to switch patients currently on ZAROXOLYN to MYKROX, the dose should be determined by titration starting at one tablet (½ mg) once daily and increasing to two tablets (1 mg) once daily if needed.

Treatment of Edematous States: The time interval required for the initial dosage to produce an effect may vary. Diuresis and saluresis usually begin within one hour and persist for 24 hours or longer. When a desired therapeutic effect has been obtained, it may be advisable to reduce the dose if possible. The daily dose depends on the severity of the patient's condition, sodium intake, and responsiveness. A decision to change the daily dose should be based on the results of thorough clinical and laboratory evaluations. If antihypertensive drugs or diuretics are given concurrently with ZAROXOLYN, more careful dosage adjustment may be necessary. For patients who tend to experience paroxysmal nocturnal dyspnea, it may be advisable to employ a larger dose to ensure prolongation of diuresis and saluresis for a full 24-hour period.

Treatment of Hypertension: The time interval required for the initial dosage regimen to show effect may vary from three or four days to three to six weeks in the treatment of elevated blood pressure. Doses should be adjusted at appropriate intervals to achieve maximum therapeutic effect.

HOW SUPPLIED

ZAROXOLYN Tablets (metolazone tablets, USP) are shallow biconvex, round tablets, and are available in three strengths:

2½ mg, pink, debossed "ZAROXOLYN" on one side, and

"2½" on reverse side.

NDC 53014-975-71 Bottle of 100's

NDC 53014-975-90 Bottle of 1000's

NDC 53014-975-72 Carton of 100's, unit dose

5 mg, blue, debossed "ZAROXOLYN" on one side, and "5" on

reverse side.

NDC 53014-850-71 Bottle of 100's

NDC 53014-850-90 Bottle of 1000's

NDC 53014-850-72 Carton of 100's, unit dose

10 mg, yellow, debossed "ZAROXOLYN" on one side, and

"10" on reverse side.

NDC 53014-835-71 Bottle of 100's

NDC 53014-835-90 Bottle of 1000's

NDC 53014-835-72 Carton of 100's, unit dose

Store at 25°C (77°F); excursions permitted to 15°-30°C (59°-86°F) [See USP Controlled Room Temperature]. Protect from light. Keep out of the reach of children.

Celltech Pharmaceuticals, Inc.

Rochester, NY 14623 USA

© Celltech Manufacturing, Inc.

© 2003, Celltech Pharmaceuticals, Inc.

All rights reserved.

Rev. 2/03

R522B

EDUCATIONAL MATERIAL

For educational information, please write to Celltech Pharmaceuticals, Inc., PO Box 31766, Rochester, NY 14603.

Centocor, Inc.

200 GREAT VALLEY PARKWAY
MALVERN, PA 19355
USA

Direct General Inquiries to:

Ph: (610) 651-6000

(888) 874-3083

Fax: (610) 651-6100

Medical Emergency Contact:

Ph: 1-(800)-457-6399

For Medical Information/Adverse Experience Reporting

Contact:

Medical Information

Ph: (800) 457-6399

REMICADE®

(rem-i-kad)

(infliximab)

for IV Injection

WARNING

RISK OF INFECTIONS

TUBERCULOSIS (FREQUENTLY DISSEMINATED OR EXTRAPULMONARY AT CLINICAL PRESENTATION), INVASIVE FUNGAL INFECTIONS, AND OTHER OPPORTUNISTIC INFECTIONS, HAVE BEEN OBSERVED IN PATIENTS RECEIVING REMICADE. SOME OF THESE INFECTIONS HAVE BEEN FATAL (SEE WARNINGS).

PATIENTS SHOULD BE EVALUATED FOR LATENT TUBERCULOSIS INFECTION WITH A TUBERCULIN SKIN TEST.¹ TREATMENT OF LATENT TUBERCULOSIS INFECTION SHOULD BE INITIATED PRIOR TO THERAPY WITH REMICADE.

DESCRIPTION

REMICADE® is a chimeric IgG1 κ monoclonal antibody with an approximate molecular weight of 149,100 daltons. It is composed of human constant and murine variable regions. Infliximab binds specifically to human tumor necrosis factor alpha (TNF α) with an association constant of 10^{10} M $^{-1}$. Infliximab is produced by a recombinant cell line cultured by continuous perfusion and is purified by a series of steps that includes measures to inactivate and remove viruses.

REMICADE is supplied as a sterile, white, lyophilized powder for intravenous infusion. Following reconstitution with 10 mL of Sterile Water for Injection, USP, the resulting pH is approximately 7.2. Each single-use vial contains 100 mg infliximab, 500 mg sucrose, 0.5 mg polysorbate 80, 2.2 mg monobasic sodium phosphate, monohydrate, and 6.1 mg dibasic sodium phosphate, dihydrate. No preservatives are present.

CLINICAL PHARMACOLOGY

General

Infliximab neutralizes the biological activity of TNF α by binding with high affinity to the soluble and transmembrane forms of TNF α and inhibits binding of TNF α with its receptors.^{2,3} Infliximab does not neutralize TNF β (lymphotxin α), a related cytokine that utilizes the same receptors as TNF α . Biological activities attributed to TNF α include: induction of pro-inflammatory cytokines such as interleukins (IL) 1 and 6, enhancement of leukocyte migration by increasing endothelial layer permeability and expression of adhesion molecules by endothelial cells and leukocytes, activation of neutrophil and eosinophil functional activity, induction of acute phase reactants and other liver proteins, as well as tissue degrading enzymes produced by synovocytes and/or chondrocytes. Cells expressing transmembrane TNF α bound by infliximab can be lysed *in vitro*⁴ or *in vivo*.⁴ Infliximab inhibits the functional activity of TNF α in a wide variety of *in vitro* bioassays utilizing human fibroblasts, endothelial cells, neutrophils, B and T lymphocytes and epithelial cells. Anti-TNF α antibodies reduce disease activity in the cotton-top tamarin colitis model, and decrease synovitis and joint erosions in a murine model of collagen-induced arthritis. Infliximab prevents disease in transgenic mice that develop polyarthritis as a result of constitutive expression of human TNF α , and when administered after disease onset, allows eroded joints to heal.

Pharmacodynamics

Elevated concentrations of TNF α have been found in the joints of rheumatoid arthritis patients and the stools of Crohn's disease patients and correlate with elevated disease activity. In rheumatoid arthritis, treatment with REMICADE reduced infiltration of inflammatory cells into inflamed areas of the joint as well as expression of molecules mediating cellular adhesion [E-selectin, intercellular adhesion molecule-1 (ICAM-1) and vascular cell adhesion molecule-1 (VCAM-1)], chemotaxis [IL-8 and monocyte chemoattractant protein (MCP-1)] and tissue degradation [matrix metalloproteinase (MMP) 1 and 3]. In Crohn's disease, treatment with REMICADE reduced infiltration of inflammatory cells and TNF α production in inflamed areas of the intestine, and reduced the proportion of mononuclear cells from the lamina propria able to express TNF α and interferon. After treatment with REMICADE, patients with rheumatoid arthritis or Crohn's disease exhibited decreased levels of serum IL-6 and C-reactive protein (CRP) compared to baseline. Peripheral blood lymphocytes from REMICADE-treated patients showed no significant decrease in number or in proliferative responses to *in vitro* mitogenic stimulation when compared to cells from untreated patients.

Pharmacokinetics

Single intravenous (IV) infusions of 3 mg/kg to 20 mg/kg showed a linear relationship between the dose administered and the maximum serum concentration. The volume of distribution at steady state was independent of dose and indicated that infliximab was distributed primarily within the vascular compartment. Median pharmacokinetic results for doses of 3 mg/kg to 10 mg/kg in rheumatoid arthritis and 5 mg/kg in Crohn's disease indicate that the terminal half-life of infliximab is 8.0 to 9.5 days.

Following an initial dose of REMICADE, repeated infusions at 2 and 6 weeks resulted in predictable concentration-time profiles following each treatment. No systemic accumulation of infliximab occurred upon continued repeated treatment with 3 mg/kg or 10 mg/kg at 4- or 8-week intervals. No major differences in clearance or volume of distribution were observed in patient subgroups defined by age, weight, or gender. It is not known if there are differences in clearance or volume of distribution in patients with marked impairment of hepatic or renal function.

A pediatric Crohn's disease pharmacokinetic study was conducted in 21 patients aged 11 to 17 years old. No notable differences in single-dose pharmacokinetic parameters were observed between pediatric and adult Crohn's disease patients (see PRECAUTIONS, Pediatric Use).

CLINICAL STUDIES

Rheumatoid Arthritis

The safety and efficacy of REMICADE when given in conjunction with methotrexate (MTX) were assessed in a multi-

Continued on next page

Remicade—Cont.

ticer, randomized, double-blind, placebo-controlled study of 428 patients with active rheumatoid arthritis despite treatment with MTX (the Anti-TNF Trial in Rheumatoid Arthritis with Concomitant Therapy or ATTRACT). Patients enrolled had a median age of 54 years, median disease duration of 8.4 years, median swollen and tender joint count of 20 and 31 respectively, and were on a median dose of 15 mg/wk of MTX. Patients received either placebo + MTX or one of 4 doses/schedules of REMICADE + MTX: 3 mg/kg or 10 mg/kg of REMICADE by IV infusion at weeks 0, 2 and 6 followed by additional infusions every 4 or 8 weeks in combination with MTX. Concurrent use of stable doses of folic acid, oral corticosteroids (≤ 10 mg/day) and/or nonsteroidal anti-inflammatory drugs was also permitted.

Data on use of REMICADE without concurrent MTX are limited (see ADVERSE REACTIONS, Immunogenicity).^{5,6}

Clinical response

All doses/schedules of REMICADE + MTX resulted in improvement in signs and symptoms as measured by the American College of Rheumatology response criteria (ACR 20) with a higher percentage of patients achieving an ACR 20, 50 and 70 compared to placebo + MTX (Table 1). This improvement was observed at week 2 and maintained through week 102. Greater effects on each component of the ACR 20 were observed in all patients treated with REMICADE + MTX compared to placebo + MTX (Table 2). Approximately 10% of patients treated with REMICADE achieved a major clinical response, defined as maintenance of an ACR 70 response over a 6-month period compared to 0% of placebo-treated patients ($p \leq 0.018$).

[See table 1 at right]

[See table 2 at right]

Radiographic response

Structural damage in both hands and feet was assessed radiographically at week 54 by the change from baseline in the van der Heijde-modified Sharp score, a composite score of structural damage that measures the number and size of joint erosions and the degree of joint space narrowing in hands/wrists and feet.⁷ Approximately 80% of patients had paired x-ray data at 54 weeks and approximately 70% at 102 weeks. The inhibition of progression of structural damage was observed at 54 weeks (Table 3) and maintained through 102 weeks.

[See table 3 at right]

Physical function response

Physical function and disability were assessed using the Health Assessment Questionnaire (HAQ) and the general health-related quality of life questionnaire SF-36. All doses/schedules of REMICADE + MTX showed significantly greater improvement from baseline in HAQ and SF-36 physical component summary score averaged over time through week 54 compared to placebo + MTX, and no worsening in the SF-36 mental component summary score.

The median (interquartile range) improvement from baseline to week 54 in HAQ was 0.1 (-0.1, 0.5) for the placebo + MTX group and 0.4 (0.1, 0.9) for REMICADE + MTX ($p < 0.001$). Both HAQ and SF-36 effects were maintained through week 102. Approximately 80% of patients in all doses/schedules of REMICADE + MTX remained in the trial through 102 weeks.

Active Crohn's Disease

The safety and efficacy of single and multiple doses of REMICADE were assessed in two randomized, double-blind, placebo-controlled clinical studies in 653 patients with moderate to severely active Crohn's disease (Crohn's Disease Activity Index (CDAI) ≥ 220 and ≤ 400) with an inadequate response to prior conventional therapies. Concomitant stable doses of aminosalicylates, corticosteroids and/or immunomodulatory agents were permitted and 92% of patients continued to receive at least one of these medications.

In the single-dose trial⁸ of 108 patients, 16% (4/25) of placebo patients achieved a clinical response (decrease in CDAI ≥ 70 points) at week 4 vs. 81% (22/27) of patients receiving 5 mg/kg REMICADE ($p < 0.001$, two-sided, Fisher's Exact test). Additionally, 4% (1/25) of placebo patients and 48% (13/27) of patients receiving 5 mg/kg REMICADE achieved clinical remission (CDAI < 150) at week 4.

In a multidose trial (ACCENT I)⁹, 545 patients received 5 mg/kg at week 0 and were then randomized to one of three treatment groups; the placebo maintenance group received placebo at weeks 2 and 6, and then every 8 weeks; the 5 mg/kg maintenance group received 5 mg/kg at weeks 2 and 6, and then every 8 weeks; and the 10 mg/kg maintenance group received 5 mg/kg at weeks 2 and 6, and then 10 mg/kg every 8 weeks. Patients in response at week 2 were randomized and analyzed separately from those not in response at week 2. Corticosteroid taper was permitted after week 6.

At week 2, 57% (311/545) of patients were in clinical response. At week 30, a significantly greater proportion of these patients in the 5 mg/kg and 10 mg/kg maintenance groups achieved clinical remission compared to patients in the placebo maintenance group (Table 4).

Additionally, a significantly greater proportion of patients in the 5 mg/kg and 10 mg/kg infliximab maintenance groups were in clinical remission and were able to discontinue corticosteroid use compared to patients in the placebo maintenance group at week 54 (Table 4).

[See table 4 at right]

Table 1
PERCENTAGE OF PATIENTS WHO ACHIEVED AN ACR RESPONSE AT WEEKS 30 AND 54
REMICADE + MTX

Response	Placebo + MTX (n=88)	3 mg/kg ^a		10 mg/kg ^a	
		q 8 wks (n=86)	q 4 wks (n=86)	q 8 wks (n=87)	q 4 wks (n=81)
ACR 20					
Week 30	20%	50%	50%	52%	58%
Week 54	17%	42%	48%	59%	59%
ACR 50					
Week 30	5%	27%	29%	31%	26%
Week 54	9%	21%	34%	40%	38%
ACR 70					
Week 30	0%	8%	11%	18%	11%
Week 54	2%	11%	18%	26%	19%

^a $p < 0.05$ for each outcome compared to placebo

Table 2
COMPONENTS OF ACR 20 AT BASELINE AND 54 WEEKS
Placebo + MTX **REMICADE + MTX^a**

Parameter (medians)	Placebo + MTX (n=88)		REMICADE + MTX ^a (n=340)	
	Baseline	Week 54	Baseline	Week 54
No. of Tender Joints	24	16	32	8
No. of Swollen Joints	19	13	20	7
Pain ^b	6.7	6.1	6.8	3.3
Physician's Global Assessment ^b	6.5	5.2	6.2	2.1
Patient's Global Assessment ^b	6.2	6.2	6.3	3.2
Disability Index (HAQ) ^c	1.8	1.5	1.8	1.3
CRP (mg/dL)	3.0	2.3	2.4	0.6

^a All doses/schedules of REMICADE + MTX

^b Visual Analog Scale (0=best, 10=worst)

^c Health Assessment Questionnaire, measurement of 8 categories: dressing and grooming, arising, eating, walking, hygiene, reach, grip, and activities (0=best, 3=worst)

Table 3
RADIOGRAPHIC CHANGE FROM BASELINE TO WEEK 54
REMICADE + MTX

Median (10, 90 percentiles)	Placebo + MTX (n=64)	3 mg/kg		10 mg/kg		p-value ^a
		q 8 wks (n=71)	q 4 wks (n=71)	q 8 wks (n=77)	q 4 wks (n=66)	
Total Score						
Baseline	55 (14, 188)	57 (15, 187)	45 (8, 162)	56 (6, 143)	43 (7, 178)	
Change from baseline	4.0 (-1.0, 19.0)	0.5 (-3.0, 5.5)	0.1 (-5.2, 9.0)	0.5 (-4.8, 5.0)	-0.5 (-5.7, 4.0)	$p < 0.001$
Erosion Score						
Baseline	25 (8, 110)	29 (9, 100)	22 (3, 91)	22 (3, 80)	26 (4, 104)	
Change from baseline	2.0 (-1.0, 9.7)	0.0 (-3.0, 4.3)	-0.3 (-3.1, 2.5)	0.5 (-3.0, 2.5)	-0.5 (-2.7, 2.5)	$p < 0.001$
JSN Score						
Baseline	26 (3, 88)	29 (4, 80)	20 (3, 83)	24 (1, 79)	25 (3, 77)	
Change from baseline	1.5 (-0.8, 8.0)	0.0 (-2.5, 4.5)	0.0 (-3.4, 5.0)	0.0 (-3.0, 2.5)	0.0 (-3.0, 3.5)	$p < 0.001$

^a For comparisons of each dose against placebo

Table 4
CLINICAL REMISSION AND STEROID WITHDRAWAL

	Single 5 mg/kg Dose ^a		Three Dose Induction ^b	
	Placebo Maintenance	Infliximab Maintenance	5 mg/kg Maintenance	10 mg/kg Maintenance
Week 30				
Clinical remission	25/102	41/104	48/105	
p-value ^c	39%	39%	46%	
		0.022	0.001	
Week 54				
Patients in remission able to discontinue corticosteroid use ^d	6/54	14/56	18/53	
p-value ^c	11%	25%	34%	
		0.059	0.005	

^a REMICADE at week 0

^b REMICADE 5 mg/kg administered at weeks 0, 2 and 6

^c p-values represent pairwise comparisons to placebo

^d Of those receiving corticosteroids at baseline

Patients in the infliximab maintenance groups (5 mg/kg and 10 mg/kg) had a longer time to loss of response than patients in the placebo maintenance group (Figure 1). At weeks 30 and 54, significant improvement from baseline was seen among the 5 mg/kg and 10 mg/kg infliximab-treated groups compared to the placebo group in the disease specific inflammatory bowel disease questionnaire (IBDQ), particularly the bowel and systemic components, and in the physical component summary score of the general health-related quality of life questionnaire SF-36.¹⁰

[See figure 1 at top of next page]

In a subset of 78 patients who had mucosal ulceration at baseline and who participated in an endoscopic substudy, 13

of 43 patients in the infliximab maintenance group had endoscopic evidence of mucosal healing compared to 1 of 28 patients in the placebo group at week 10. Of the infliximab-treated patients showing mucosal healing at week 10, 9 of 12 patients also showed mucosal healing at week 54. Patients who achieved a response and subsequently lost response were eligible to receive infliximab on an episodic basis at a dose that was 5 mg/kg higher than the dose to which they were randomized. The majority of such patients responded to the higher dose. Among patients who were not in response at week 2, 59% (92/157) of infliximab maintenance patients responded by week 14 compared to 51% (39/77) of placebo maintenance patients. Among patients who did not




Figure 1
Kaplan-Meier estimate of the proportion of patients who had not lost response through week 54

respond by week 14, additional therapy did not result in significantly more responses (see DOSAGE AND ADMINISTRATION).

Fistulizing Crohn's Disease

The safety and efficacy of REMICADE were assessed in 2 randomized, double-blind, placebo-controlled studies in patients with fistulizing Crohn's disease with fistula(s) that were of at least 3 months duration. Concurrent use of stable doses of corticosteroids, 5-aminosalicylates, antibiotics, MTX, 6-mercaptopurine (6-MP) and/or azathioprine (AZA) was permitted.

In the first trial,¹⁰ 94 patients received three doses of either placebo or REMICADE at weeks 0, 2 and 6. Fistula response ($\geq 50\%$ reduction in number of enterocutaneous fistulas draining upon gentle compression on at least two consecutive visits without an increase in medication or surgery for Crohn's disease) was seen in 68% (21/31) of patients in the 5 mg/kg REMICADE group ($p=0.002$) and 56% (18/32) of patients in the 10 mg/kg REMICADE group ($p=0.021$) vs. 26% (8/31) of patients in the placebo arm. The median time to onset of response and median duration of response in REMICADE-treated patients was 2 and 12 weeks, respectively. Closure of all fistula was achieved in 52% of REMICADE-treated patients compared with 13% of placebo-treated patients ($p<0.001$).

In the second trial (ACCENT II), patients who were enrolled had to have at least one draining enterocutaneous (perianal, abdominal) fistula. All patients received 5 mg/kg REMICADE at weeks 0, 2 and 6. Patients were randomized to placebo or 5 mg/kg REMICADE maintenance at week 14. Patients received maintenance doses at week 14 and then every eight weeks through week 46. Patients who were in fistula response (fistula response was defined the same as in the first trial) at both weeks 10 and 14 were randomized separately from those not in response. The primary endpoint was time from randomization to loss of response among those patients who were in fistula response.

Among the randomized patients (273 of the 296 initially enrolled), 87% had perianal fistulas and 14% had abdominal fistulas. Eight percent also had rectovaginal fistulas. Greater than 90% of the patients had received previous immunosuppressive and antibiotic therapy.

At week 14, 65% (177/273) of patients were in fistula response. Patients randomized to REMICADE maintenance had a longer time to loss of fistula response compared to the placebo maintenance group (Figure 2). At week 54, 38% (33/87) of REMICADE-treated patients had no draining fistulas compared with 22% (20/90) of placebo-treated patients ($p=0.02$). Compared to placebo maintenance, patients on REMICADE maintenance had a trend toward fewer hospitalizations.




Figure 2
Life table estimates of the proportion of patients who had not lost fistula response through week 54

Patients who achieved a fistula response and subsequently lost response were eligible to receive REMICADE maintenance therapy at a dose that was 5 mg/kg higher than the dose to which they were randomized. Of the placebo maintenance patients, 66% (25/38) responded to 5 mg/kg REMICADE, and 57% (12/21) of REMICADE maintenance patients responded to 10 mg/kg.

Patients who had not achieved a response by week 14 were unlikely to respond to additional doses of REMICADE.

Similar proportions of patients in either group developed new fistulas (17% overall) and similar numbers developed abscesses (15% overall).

INDICATIONS AND USAGE

Rheumatoid Arthritis

REMICADE, in combination with methotrexate, is indicated for reducing signs and symptoms, inhibiting the progression of structural damage and improving physical function in patients with moderately to severely active rheumatoid arthritis who have had an inadequate response to methotrexate.

Crohn's Disease

REMICADE is indicated for reducing signs and symptoms and inducing and maintaining clinical remission in patients with moderately to severely active Crohn's disease who have had an inadequate response to conventional therapy. REMICADE is indicated for reducing the number of draining enterocutaneous and rectovaginal fistulas and maintaining fistula closure in patients with fistulizing Crohn's disease.

CONTRAINDICATIONS

REMICADE at doses >5 mg/kg should not be administered to patients with moderate to severe heart failure. In a randomized study evaluating REMICADE in patients with moderate to severe heart failure (New York Heart Association [NYHA] Functional Class III/IV), REMICADE treatment at 10 mg/kg was associated with an increased incidence of death and hospitalization due to worsening heart failure (see WARNINGS and ADVERSE REACTIONS, Patients with Heart Failure).

REMICADE should not be administered to patients with known hypersensitivity to any murine proteins or other component of the product.

WARNINGS

RISK OF INFECTIONS

(See boxed WARNING)

SERIOUS INFECTIONS, INCLUDING SEPSIS HAVE BEEN REPORTED IN PATIENTS RECEIVING TNF-BLOCKING AGENTS. SOME OF THESE INFECTIONS HAVE BEEN FATAL. MANY OF THE SERIOUS INFECTIONS IN PATIENTS TREATED WITH REMICADE HAVE OCCURRED IN PATIENTS ON CONCOMITANT IMMUNOSUPPRESSIVE THERAPY THAT, IN ADDITION TO THEIR CROHN'S DISEASE OR RHEUMATOID ARTHRITIS, COULD PREDISPOSE THEM TO INFECTIONS.

REMICADE SHOULD NOT BE GIVEN TO PATIENTS WITH A CLINICALLY IMPORTANT, ACTIVE INFECTION. CAUTION SHOULD BE EXERCISED WHEN CONSIDERING THE USE OF REMICADE IN PATIENTS WITH A CHRONIC INFECTION OR A HISTORY OF RECURRENT INFECTION. PATIENTS SHOULD BE MONITORED FOR SIGNS AND SYMPTOMS OF INFECTION WHILE ON OR AFTER TREATMENT WITH REMICADE. NEW INFECTIONS SHOULD BE CLOSELY MONITORED. IF A PATIENT DEVELOPS A SERIOUS INFECTION, REMICADE THERAPY SHOULD BE DISCONTINUED (see ADVERSE REACTIONS, Infections).

CASES OF HISTOPLASMOIS, COCCIDIOIDOMYCOSIS, LISTERIOSIS, PNEUMOCYSTOSIS, TUBERCULOSIS, OTHER BACTERIAL, MYCOBACTERIAL AND FUNGAL INFECTIONS HAVE BEEN OBSERVED IN PATIENTS RECEIVING REMICADE. FOR PATIENTS WHO HAVE RESIDED IN REGIONS WHERE HISTOPLASMOIS OR COCCIDIOIDOMYCOSIS IS ENDEMIC, THE BENEFITS AND RISKS OF REMICADE TREATMENT SHOULD BE CAREFULLY CONSIDERED BEFORE INITIATION OF REMICADE THERAPY.

SERIOUS INFECTIONS WERE SEEN IN CLINICAL STUDIES WITH CONCURRENT USE OF ANAKINRA AND ANOTHER TNF-BLOCKING AGENT, ETANERCEPT, WITH NO ADDED CLINICAL BENEFIT COMPARED TO ETANERCEPT ALONE. BECAUSE OF THE NATURE OF THE ADVERSE EVENTS SEEN WITH COMBINATION OF ETANERCEPT AND ANAKINRA THERAPY, SIMILAR TOXICITIES MAY ALSO RESULT FROM THE COMBINATION OF REMICADE AND ANAKINRA. THEREFORE, THE COMBINATION OF REMICADE AND ANAKINRA IS NOT RECOMMENDED.

Patients with Heart Failure

REMICADE has been associated with adverse outcomes in patients with heart failure, and should be used in patients with heart failure only after consideration of other treatment options. The results of a randomized study evaluating the use of REMICADE in patients with heart failure (NYHA Functional Class III/IV) suggested higher mortality in patients who received 10 mg/kg REMICADE, and higher rates of cardiovascular adverse events at doses of 5 mg/kg and 10 mg/kg. There have been post-marketing reports of worsening heart failure, with and without identifiable precipitating factors, in patients taking REMICADE. There have also been rare post-marketing reports of new onset heart failure, including heart failure in patients without known pre-existing cardiovascular disease. Some of these patients have been under 50 years of age. If a decision is made to administer REMICADE to patients with heart failure, they should be closely monitored during therapy, and REMICADE should be discontinued if new or worsening symptoms of heart failure appear. (see CONTRAINDICATIONS and ADVERSE REACTIONS, Patients with Heart Failure.)

Hematologic Events

Cases of leukopenia, neutropenia, thrombocytopenia, and pancytopenia, some with a fatal outcome, have been reported in patients receiving REMICADE. The causal rela-

tionship to REMICADE therapy remains unclear. Although no high-risk group(s) has been identified, caution should be exercised in patients being treated with REMICADE who have ongoing or a history of significant hematologic abnormalities. All patients should be advised to seek immediate medical attention if they develop signs and symptoms suggestive of blood dyscrasias or infection (e.g., persistent fever) while on REMICADE. Discontinuation of REMICADE therapy should be considered in patients who develop significant hematologic abnormalities.

Hypersensitivity

REMICADE has been associated with hypersensitivity reactions that vary in their time of onset and required hospitalization in some cases. Most hypersensitivity reactions, which include urticaria, dyspnea, and/or hypotension, have occurred during or within 2 hours of REMICADE infusion. However, in some cases, serum sickness-like reactions have been observed in Crohn's disease patients 3 to 12 days after REMICADE therapy was reinstituted following an extended period without REMICADE treatment. Symptoms associated with these reactions include fever, rash, headache, sore throat, myalgias, polyarthralgias, hand and facial edema and/or dysphagia. These reactions were associated with marked increase in antibodies to infliximab, loss of detectable serum concentrations of infliximab, and possible loss of drug efficacy. REMICADE should be discontinued for severe reactions. Medications for the treatment of hypersensitivity reactions (e.g., acetaminophen, antihistamines, corticosteroids and/or epinephrine) should be available for immediate use in the event of a reaction (see ADVERSE REACTIONS, Infusion-related Reactions).

Neurologic Events

Infliximab and other agents that inhibit TNF have been associated in rare cases with optic neuritis, seizure and new onset or exacerbation of clinical symptoms and/or radiographic evidence of central nervous system demyelinating disorders, including multiple sclerosis, and CNS manifestation of systemic vasculitis. Prescribers should exercise caution in considering the use of REMICADE in patients with pre-existing or recent onset of central nervous system demyelinating or seizure disorders. Discontinuation of REMICADE should be considered in patients who develop significant central nervous system adverse reactions.

PRECAUTIONS

Autoimmunity

Treatment with REMICADE may result in the formation of autoantibodies and, rarely, in the development of a lupus-like syndrome. If a patient develops symptoms suggestive of a lupus-like syndrome following treatment with REMICADE, treatment should be discontinued (see ADVERSE REACTIONS, Autoantibodies/Lupus-like Syndrome).

Malignancy

Patients with long duration of Crohn's disease or rheumatoid arthritis and chronic exposure to immunosuppressant therapies are more prone to develop lymphomas (see ADVERSE REACTIONS, Malignancies/Lymphoproliferative Disease). The impact of treatment with REMICADE on these phenomena is unknown.

Vaccinations

No data are available on the response to vaccination or on the secondary transmission of infection by live vaccines in patients receiving anti-TNF therapy. It is recommended that live vaccines not be given concurrently.

Information for Patients

Patients should be provided the REMICADE Patient Information Sheet and provided an opportunity to read it prior to each treatment infusion session. Because caution should be exercised in administering REMICADE to patients with clinically important active infections, it is important that the patient's overall health be assessed at each treatment visit and any questions resulting from the patient's reading of the Patient Information Sheet be discussed.

Drug Interactions

Concurrent administration of etanercept (another TNF-blocking agent) and anakinra (an interleukin-1 antagonist) has been associated with an increased risk of serious infections, and increased risk of neutropenia and no additional benefit compared to these medicinal products alone. Other TNF-blocking agents (including REMICADE) used in combination with anakinra may also result in similar toxicities (see WARNINGS, RISK OF INFECTIONS).

Specific drug interaction studies, including interactions with MTX, have not been conducted. The majority of patients in rheumatoid arthritis or Crohn's disease clinical studies received one or more concomitant medications. In rheumatoid arthritis, concomitant medications besides MTX were nonsteroidal anti-inflammatory agents, folic acid, corticosteroids and/or narcotics. Concomitant Crohn's disease medications were antibiotics, antivirals, corticosteroids, 6-MP/AZA and aminosalicylates. Patients with Crohn's disease who received immunosuppressants tended to experience fewer infusion reactions compared to patients on no immunosuppressants (see ADVERSE REACTIONS, Immunogenicity and Infusion-related Reactions).

Serum infliximab concentrations appeared to be unaffected by baseline use of medications for the treatment of Crohn's disease including corticosteroids, antibiotics (miconazole or ciprofloxacin) and aminosalicylates.

Carcinogenesis, Mutagenesis and Impairment of Fertility
A repeat dose toxicity study was conducted with mice given cV1q anti-mouse TNF α to evaluate tumorigenicity. CV1q is

Continued on next page

Remicade—Cont.

an analogous antibody that inhibits the function of TNF α in mice. Animals were assigned to 1 of 3 dose groups: control, 10 mg/kg or 40 mg/kg cV1q given weekly for 6 months. The weekly doses of 10 mg/kg and 40 mg/kg are 2 and 8 times, respectively, the human dose of 5 mg/kg for Crohn's disease. Results indicated that cV1q did not cause tumorigenicity in mice. No clastogenic or mutagenic effects of infliximab were observed in the *in vivo* mouse micronucleus test or the *Salmonella-Escherichia coli* (Ames) assay, respectively. Chromosomal aberrations were not observed in an assay performed using human lymphocytes. The significance of these findings for human risk is unknown. It is not known whether infliximab can impair fertility in humans. No impairment of fertility was observed in a fertility and general reproduction toxicity study with the analogous mouse antibody used in the 6-month chronic toxicity study.

Pregnancy Category B

Since infliximab does not cross-react with TNF α in species other than humans and chimpanzees, animal reproduction studies have not been conducted with REMICADE. No evidence of maternal toxicity, embryotoxicity or teratogenicity was observed in a developmental toxicity study conducted in mice using an analogous antibody that selectively inhibits the functional activity of mouse TNF α . Doses of 10 to 15 mg/kg in pharmacodynamic animal models with the anti-TNF α analogous antibody produced maximal pharmacologic effectiveness. Doses up to 40 mg/kg were shown to produce no adverse effects in animal reproduction studies. It is not known whether REMICADE can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. REMICADE should be given to a pregnant woman only if clearly needed.

Nursing Mothers

It is not known whether infliximab is excreted in human milk or absorbed systemically after ingestion. Because many drugs and immunoglobulins are excreted in human milk, and because of the potential for adverse reactions in nursing infants from REMICADE, a decision should be made whether to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother.

Pediatric Use

Safety and effectiveness of REMICADE in patients with juvenile rheumatoid arthritis and in pediatric patients with Crohn's disease have not been established.

Geriatric Use

In the ATTRACT study, no overall differences were observed in effectiveness or safety in 72 patients aged 65 or older compared to younger patients. In Crohn's disease studies, there were insufficient numbers of patients aged 65 and over to determine whether they respond differently from patients aged 18 to 65. Because there is a higher incidence of infections in the elderly population in general, caution should be used in treating the elderly (see ADVERSE REACTIONS, Infections).

ADVERSE REACTIONS

The data described herein reflect exposure to REMICADE in 1678 patients, including 842 patients exposed beyond 30 weeks and 295 exposed beyond one year. The most common reason for discontinuation of treatment was infusion-related reactions (e.g. dyspnea, flushing, headache and rash). Adverse events have been reported in a higher proportion of rheumatoid arthritis patients receiving the 10 mg/kg dose than the 3 mg/kg dose, however, no differences were observed in the frequency of adverse events between the 5 mg/kg dose and 10 mg/kg dose in patients with Crohn's disease.

Infusion-related Reactions**Acute infusion reactions**

An infusion reaction was defined in clinical trials as any adverse event occurring during an infusion or within 1 to 2 hours after an infusion. Approximately 20% of REMICADE-treated patients in all clinical studies experienced an infusion reaction compared to approximately 10% of placebo-treated patients. Among all REMICADE infusions, 3% were accompanied by nonspecific symptoms such as fever or chills, 1% were accompanied by cardiopulmonary reactions (primarily chest pain, hypotension, hypertension or dyspnea), and <1% were accompanied by pruritus, urticaria, or the combined symptoms of pruritus/urticaria and cardiopulmonary reactions. Serious infusion reactions occurred in <1% of patients and included anaphylaxis, convulsions, erythematous rash and hypotension. Approximately 3% of patients discontinued REMICADE because of infusion reactions, and all patients recovered with treatment and/or discontinuation of the infusion. REMICADE infusions beyond the initial infusion were not associated with a higher incidence of reactions.

Patients who became positive for antibodies to infliximab were more likely (approximately 2- to 3-fold) to have an infusion reaction than were those who were negative. Use of concomitant immunosuppressant agents appeared to reduce the frequency of antibodies to infliximab and infusion reactions (see ADVERSE REACTIONS, Immunogenicity and PRECAUTIONS, Drug Interactions).

In post-marketing experience, cases of anaphylactic-like reactions, including laryngeal/pharyngeal edema and severe bronchospasm, and seizure have been associated with REMICADE administration.

Reactions following readministration

In a study where 37 of 41 patients with Crohn's disease were retreated with infliximab following a 2 to 4 year period without infliximab treatment, 10 patients experienced adverse events manifesting 3 to 12 days following infusion of which 6 were considered serious. Signs and symptoms included myalgia and/or arthralgia with fever and/or rash, with some patients also experiencing pruritus, facial, hand or lip edema, dysphagia, urticaria, sore throat, and headache. Patients experiencing these adverse events had not experienced infusion-related adverse events associated with their initial infliximab therapy. These adverse events occurred in 39% (9/23) of patients who had received liquid formulation which is no longer in use and 7% (1/14) of patients who received lyophilized formulation. The clinical data are not adequate to determine if occurrence of these reactions is due to differences in formulation. Patients' signs and symptoms improved substantially or resolved with treatment in all cases. There are insufficient data on the incidence of these events after drug-free intervals of 1 to 2 years. These events have been observed only infrequently in clinical studies and post-marketing surveillance with retreatment intervals up to 1 year.

Infections

In REMICADE clinical studies, treated infections were reported in 35% of REMICADE-treated patients (average of 53 weeks of follow-up) and in 26% of placebo-treated patients (average of 41 weeks of follow-up). When longer observation of patients on REMICADE was accounted for, the event rate was similar for both groups. The infections most frequently reported were respiratory tract infections (including sinusitis, pharyngitis, and bronchitis) and urinary tract infections. No increased risk of serious infections or sepsis was observed with REMICADE compared with placebo in clinical studies. Among REMICADE-treated patients, serious infections included pneumonia, cellulitis, abscess, skin ulceration, sepsis, and bacterial infection. Three opportunistic infections were reported; coccidioidomycosis (which resulted in death), nocardiosis and cytomegalovirus. Tuberculosis was reported in two patients, one of whom died due to miliary tuberculosis. Other cases of tuberculosis, including disseminated tuberculosis, also have been reported post-marketing. Most of the cases of tuberculosis occurred within the first two months after initiation of therapy with infliximab and may reflect recrudescence of latent disease (see WARNINGS, RISK OF INFECTIONS). During the 54 week ACCENT II trial, 15% of patients with fistulizing Crohn's disease developed a new fistula-related abscess.

In post-marketing experience, infections have been observed with various pathogens including viral, bacterial, fungal, and protozoal organisms. Infections have been noted in all organ systems and have been reported in patients receiving REMICADE alone or in combination with immunosuppressive agents.

Autoantibodies/Lupus-like Syndrome

Approximately 52% of 1261 infliximab-treated patients in clinical trials who were antinuclear antibody (ANA) negative at baseline developed a positive ANA during the trial compared with approximately 19% of 129 placebo-treated patients. Anti-dsDNA antibodies were newly detected in approximately 17% of 1507 infliximab-treated patients compared with 0% of 162 placebo-treated patients. Reports of lupus and lupus-like syndromes, however, remain uncommon.

Malignancies/Lymphoproliferative Disease

In completed clinical studies of REMICADE for up to 102 weeks, 18 of 1678 patients developed 19 new or recurrent malignancies of various types, such as non-Hodgkin's B-cell lymphoma, breast, melanoma, squamous, rectal and basal cell. There are insufficient data to determine whether REMICADE contributed to the development of these malignancies. The observed rates and incidences were similar to those expected for the populations studied^{11,12} (see PRECAUTIONS, Malignancy).

Patients with Heart Failure

In a randomized study evaluating REMICADE in moderate to severe heart failure (NYHA Class III/IV; left ventricular ejection fraction $\leq 35\%$), 150 patients were randomized to receive treatment with 3 infusions of REMICADE 10 mg/kg, 5 mg/kg, or placebo, at 0, 2, and 6 weeks. Higher incidences of mortality and hospitalization due to worsening heart failure were observed in patients receiving the 10 mg/kg REMICADE dose. At 1 year, 8 patients in the 10 mg/kg REMICADE group had died compared with 4 deaths each in the 5 mg/kg REMICADE and the placebo groups. There were trends towards increased dyspnea, hypotension, angina, and dizziness in both the 10 mg/kg and 5 mg/kg REMICADE treatment groups, versus placebo. REMICADE has not been studied in patients with mild heart failure (NYHA Class I/II). (See CONTRAINDICATIONS and WARNINGS, Patients with Heart Failure.)

Immunogenicity

Treatment with REMICADE can be associated with the development of antibodies to infliximab. The incidence of antibodies to infliximab in patients given a 3-dose induction regimen followed by maintenance dosing was approximately 10% as assessed through one to two years of REMICADE treatment. A higher incidence of antibodies to infliximab was observed in Crohn's disease patients receiving REMICADE after drug free intervals >16 weeks. The majority of antibody-positive patients had low titers. Patients who were antibody-positive were more likely to experience an infusion reaction (see ADVERSE REACTIONS, Infusion-related Reactions). Antibody development was

lower among rheumatoid arthritis and Crohn's disease patients receiving immunosuppressant therapies such as 6-MP/AZA or MTX.

The data reflect the percentage of patients whose test results were positive for antibodies to infliximab in an ELISA assay, and are highly dependent on the sensitivity and specificity of the assay. Additionally, the observed incidence of antibody positivity in an assay may be influenced by several factors including sample handling, timing of sample collection, concomitant medication, and underlying disease. For these reasons, comparison of the incidence of antibodies to infliximab with the incidence of antibodies to other products may be misleading.

Other Adverse Reactions

Safety data are available from 1675 REMICADE-treated patients, including 555 with rheumatoid arthritis, and 1106 with Crohn's disease and 17 with conditions other than rheumatoid arthritis or Crohn's disease. Adverse events reported in $\geq 5\%$ of all patients with rheumatoid arthritis receiving 4 or more infusions are in Table 5. The types and frequencies of adverse reactions observed were similar in REMICADE-treated rheumatoid arthritis and Crohn's disease patients except for abdominal pain which occurred in 26% of REMICADE-treated patients with Crohn's disease. In the Crohn's disease studies, there were insufficient numbers and duration of follow-up for patients who never received REMICADE to provide meaningful comparisons.

Table 5
ADVERSE EVENTS OCCURRING IN 5% OR MORE OF PATIENTS RECEIVING 4 OR MORE INFUSIONS FOR RHEUMATOID ARTHRITIS

	Placebo (n=81)	REMICADE (n=430)
Average weeks of follow-up	73	82
Gastrointestinal		
Abdominal pain	12%	17%
Nausea	23%	24%
Diarrhea	19%	19%
Dyspepsia	9%	10%
Respiratory		
Upper respiratory tract infection	35%	40%
Pharyngitis	12%	17%
Sinusitis	7%	20%
Coughing	9%	18%
Rhinitis	14%	14%
Dyspnea	2%	6%
Skin and appendages disorders		
Rash	7%	18%
Pruritis	2%	9%
Body as a whole—general disorders		
Fatigue	9%	13%
Chest pain	6%	7%
Resistance mechanism disorders		
Fever	11%	13%
Abscess	5%	6%
Moniliasis	2%	8%
Central and peripheral nervous system disorders		
Headache	21%	29%
Musculoskeletal system disorders		
Arthralgia	7%	13%
Back pain	5%	13%
Psychiatric disorders		
Insomnia	4%	6%
Depression	2%	8%
Urinary system disorders		
Urinary tract infection	12%	14%
Cardiovascular disorders, general		
Hypertension	6%	10%

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in clinical trials of a drug cannot be directly compared to rates in clinical trials of another drug and may not predict the rates observed in broader patient populations in clinical practice. The most common serious adverse events observed in clinical trials were infections (see ADVERSE REACTIONS, Infections). Other serious, medically relevant adverse events $\geq 0.2\%$ or clinically significant adverse events by body system were as follows:

Body as a whole: allergic reaction, diaphragmatic hernia, edema, surgical/procedural sequele

Blood: pancytopenia

Cardiovascular: circulatory failure, hypotension, syncope

Gastrointestinal: constipation, gastrointestinal hemorrhage, ileus, intestinal obstruction, intestinal perforation, intestinal stenosis, pancreatitis, peritonitis, proctalgia

Central & Peripheral Nervous: meningitis, neuritis, peripheral neuropathy, dizziness

Heart Rate and Rhythm: arrhythmia, bradycardia, cardiac arrest, tachycardia

Liver and Biliary: biliary pain, cholecystitis, cholelithiasis, hepatitis

Metabolic and Nutritional: dehydration

Musculoskeletal: intervertebral disk herniation, tendon disorder

Myo., Endo., Pericardial and Coronary Valve: myocardial infarction

Platelet, Bleeding and Clotting: thrombocytopenia

Neoplasms: basal cell, breast, lymphoma

Psychiatric: confusion, suicide attempt

Red Blood Cell: anemia, hemolytic anemia

Reproductive: menstrual irregularity

Resistance Mechanism: cellulitis, sepsis, serum sickness

Respiratory: adult respiratory distress syndrome, lower respiratory tract infection, pleural effusion, pleurisy, pulmonary edema, respiratory insufficiency**Skin and Appendages:** increased sweating, ulceration**Urinary:** renal calculus, renal failure**Vascular (Extracardiac):** brain infarction, pulmonary embolism, thrombophlebitis**White Cell and Reticuloendothelial:** leukopenia, lymphadenopathy

A greater proportion of patients enrolled into the ATTRACT study who received REMICADE + MTX experienced transient mild (<2 times the upper limit of normal) or moderate (≥2 but <3 times the upper limit of normal) elevations in AST or ALT (49% and 47%, respectively) compared to patients treated with placebo + MTX (27% and 35%, respectively). Six (1.8%) patients treated with REMICADE + MTX experienced more prolonged elevations in their ALT.

The following adverse events have been reported during post-approval use of REMICADE: neutropenia (see WARNINGS, Hematologic Events), interstitial pneumonitis/fibrosis, idiopathic thrombocytopenic purpura, thrombotic thrombocytopenic purpura, pericardial effusion, systemic and cutaneous vasculitis, Guillain-Barré syndrome, transverse myelitis, and neuropathies (additional neurologic events have also been observed, see WARNINGS, Neurologic Events). Because these events are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to REMICADE exposure.

OVERDOSAGE

Single doses up to 20 mg/kg have been administered without any direct toxic effect. In case of overdosage, it is recommended that the patient be monitored for any signs or symptoms of adverse reactions or effects and appropriate symptomatic treatment instituted immediately.

DOSAGE AND ADMINISTRATION**Rheumatoid Arthritis**

The recommended dose of REMICADE is 3 mg/kg given as an intravenous infusion followed with additional similar doses at 2 and 6 weeks after the first infusion then every 8 weeks thereafter. REMICADE should be given in combination with methotrexate. For patients who have an incomplete response, consideration may be given to adjusting the dose up to 10 mg/kg or treating as often as every 4 weeks.

Crohn's Disease or Fistulizing Crohn's Disease

The recommended dose of REMICADE is 5 mg/kg given as an induction regimen at 0, 2 and 6 weeks followed by a maintenance regimen of 5 mg/kg every 8 weeks thereafter for the treatment of moderately to severely active Crohn's disease or fistulizing disease. For patients who respond and then lose their response, consideration may be given to treatment with 10 mg/kg. Patients who do not respond by week 14 are unlikely to respond with continued dosing and consideration should be given to discontinue REMICADE in these patients.

Preparation and Administration Instructions**Use aseptic technique.**

REMICADE vials do not contain antibacterial preservatives. Therefore, the vials after reconstitution should be used immediately, not re-entered or stored. The diluent to be used for reconstitution is 10 mL of Sterile Water for Injection, USP. The total dose of the reconstituted product must be further diluted to 250 mL with 0.9% Sodium Chloride Injection, USP. The infusion concentration should range between 0.4 mg/mL and 4 mg/mL. The REMICADE infusion should begin within 3 hours of preparation.

1. Calculate the dose and the number of REMICADE vials needed. Each REMICADE vial contains 100 mg of infliximab. Calculate the total volume of reconstituted REMICADE solution required.

2. Reconstitute each REMICADE vial with 10 mL of Sterile Water for Injection, USP, using a syringe equipped with a 21-gauge or smaller needle. Remove the flip-top from the vial and wipe the top with an alcohol swab. Insert the syringe needle into the vial through the center of the rubber stopper and direct the stream of Sterile Water for Injection, USP, to the glass wall of the vial. Do not use the vial if the vacuum is not present. Gently swirl the solution by rotating the vial to dissolve the lyophilized powder. Avoid prolonged or vigorous agitation. DO NOT SHAKE. Foaming of the solution on reconstitution is not unusual. Allow the reconstituted solution to stand for 5 minutes. The solution should be colorless to light yellow and opalescent, and the solution may develop a few translucent particles as infliximab is a protein. Do not use if opaque particles, discoloration, or other foreign particles are present.

3. Dilute the total volume of the reconstituted REMICADE solution dose to 250 mL with 0.9% Sodium Chloride Injection, USP, by withdrawing a volume of 0.9% Sodium Chloride Injection, USP, equal to the volume of reconstituted REMICADE from the 0.9% Sodium Chloride Injection, USP, 250 mL bottle or bag. Slowly add the total volume of reconstituted REMICADE solution to the 250 mL infusion bottle or bag. Gently mix.

4. The infusion solution must be administered over a period of not less than 2 hours and must use an infusion set with an in-line, sterile, non-pyrogenic, low-protein-binding filter (pore size of 1.2 µm or less). Any unused portion of the infusion solution should not be stored for reuse.

5. No physical biochemical compatibility studies have been conducted to evaluate the co-administration of REMICADE with other agents. REMICADE should not be infused concomitantly in the same intravenous line with other agents.

6. Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit. If visibly opaque particles, discoloration or other foreign particulates are observed, the solution should not be used.

Storage

Store the lyophilized product under refrigeration at 2°C to 8°C (36°F to 46°F). Do not freeze. Do not use beyond the expiration date. This product contains no preservative.

HOW SUPPLIED

REMICADE lyophilized concentrate for IV injection is supplied in individually-boxed single-use vials in the following strength:

NDC 57894-030-01 100 mg infliximab in a 20 mL vial

REFERENCES

1. American Thoracic Society; Centers for Disease Control and Prevention. Targeted tuberculin testing and treatment of latent tuberculosis infection. *Am J Respir Crit Care Med* 2000;161:S221-S247.
2. Knight DM, Trinh H, Le J, et al. Construction and initial characterization of a mouse-human chimeric anti-TNF antibody. *Molec Immunol* 1993;30:1443-1453.
3. Scallon BJ, Moore MA, Trinh H, et al. Chimeric anti-TNFα monoclonal antibody cA2 binds recombinant transmembrane TNFα and activates immune effector functions. *Cytokine* 1995;7:251-259.
4. ten Hove T, van Montfrans C, Peppelenbosch MP, et al. Infliximab treatment induces apoptosis of lamina propria T lymphocytes in Crohn's disease. *Gut* 2002;50:206-211.
5. Maini RN, Breedveld FC, Kalden JR, et al. Therapeutic efficacy of multiple intravenous infusions of anti-tumor necrosis factor α monoclonal antibody combined with low-dose weekly methotrexate in rheumatoid arthritis. *Arthritis Rheum* 1998;41(9):1552-1563.
6. Elliott MJ, Maini RN, Feldmann M, et al. Randomised double-blind comparison of chimeric monoclonal antibody to tumour necrosis factor alpha (cA2) vs. placebo in rheumatoid arthritis. *Lancet* 1994;344(8930):1105-1110.
7. Van der Heijde DM, van Leeuwen MA, van Riel PL, et al. Biannual radiographic assessments of hands and feet in a three-year prospective follow-up of patients with early rheumatoid arthritis. *Arthritis Rheum* 1992;35(1):26-34.
8. Targan SR, Hanauer SR, van Deventer SJH, et al. A short-term study of chimeric monoclonal antibody cA2 to tumor necrosis factor α for Crohn's disease. *N Engl J Med* 1997;337(15):1029-1035.
9. Hanauer SB, Feagan BG, Lichtenstein GR, et al. Maintenance infliximab for Crohn's disease: the ACCENT I randomized trial. *Lancet* 2002;359:1541-1549.
10. Present DH, Rutgeerts P, Targan S, et al. Infliximab for the treatment of fistulas in patients with Crohn's disease. *N Engl J Med* 1999;340:1398-1405.
11. Bernstein C, Blanchard JF, Kiewer E, et al. Cancer risk in patients with inflammatory bowel disease. *Cancer* 2001;91:854-862.
12. Jones M, Symmons D, Finn J, et al. Does exposure to immunosuppressive therapy increase the 10 year malignancy and mortality risks in rheumatoid arthritis? A matched cohort study. *Br J Rheum* 1996;35:738-745.

©Centocor, Inc. 2004

Malvern, PA 19355, USA

1-800-457-6399

Rx Only

License #1242

Revised July 2004

REMICADE® (infliximab)**Patient Information Sheet**

You should read this information sheet before you start using REMICADE® (pronounced rem-eh-kaid) and before each time you are scheduled to receive REMICADE. This information sheet does not take the place of talking with your doctor. You and your doctor should talk about your health and how you are feeling before you start taking REMICADE, while you are taking it and at regular check-ups. If you do not understand any of the information in this sheet, you should ask your doctor to explain what it means.

What is REMICADE?

REMICADE is a medicine that is used to treat adults with moderate to severely active rheumatoid arthritis and Crohn's disease. Your doctor has decided to treat you with REMICADE because your disease is still active even though you have tried other treatments.

How does REMICADE work?

The medicine REMICADE is a type of protein that recognizes, attaches to and blocks the action of a substance in your body called tumor necrosis factor. Tumor necrosis factor (TNF) is made by certain blood cells in your body. REMICADE will not cure rheumatoid arthritis or Crohn's disease, but blocking TNF with REMICADE may reduce the inflammation caused by too much TNF in your body. You should also know that REMICADE may help you feel better but can also cause serious side effects and can reduce your body's ability to fight infections (see below).

What should I know about the immune system, and taking REMICADE for Rheumatoid Arthritis or Crohn's Disease?

The immune system protects the body by responding to "invaders" like bacteria, viruses and other foreign matter that

enter your body by producing antibodies and putting them into action to fight off the "invaders." In diseases like rheumatoid arthritis and Crohn's disease, your body's immune system produces too much TNF. Too much TNF can cause your immune system to attack healthy tissues in your body and cause inflammation. If this condition is left untreated, it can cause permanent damage to the body's bones, cartilage and tissue.

While taking REMICADE can block the TNF that causes inflammation, it can also lower your body's ability to fight infections. So, taking REMICADE can make you more prone to getting infections or it can make an infection that you already have worse. You should call your doctor right away if you think you have an infection.

What important information should I know about treatment with REMICADE?

REMICADE, like other medicines that affect your immune system, is a strong medicine that can cause serious side effects. Possible serious side effects include:

Serious Infections:

- Some patients have had serious infections while receiving REMICADE. Some of the patients have died from these infections. Serious infections include TB (tuberculosis), and infections caused by viruses, fungi or bacteria that have spread throughout the body. If you develop a fever, feel very tired, have a cough, or have flu-like symptoms, these could be signs that you may be getting an infection. If you have any of these symptoms while you are taking or after you have taken REMICADE, you should tell your doctor right away.

Heart Failure:

- If you have been told that you have a heart problem called congestive heart failure and you are currently being treated with REMICADE, you will need to be closely monitored by your doctor. If you develop new or worse symptoms that are related to your heart condition, such as shortness of breath or swelling of your ankles or feet, you must contact your doctor immediately.

Blood Problems:

- In some patients the body may fail to produce enough of the blood cells that help your body fight infections or help you stop bleeding. Some of the patients have died from this failure to produce blood cells. If you develop a fever that doesn't go away, bruise or bleed very easily or look very pale, call your doctor right away. Your doctor may decide to stop your treatment.

Allergic Reactions:

- Some patients have had severe allergic reactions to REMICADE. These reactions can happen while you are getting your REMICADE infusion or shortly afterwards. The symptoms of an allergic reaction may include hives (red, raised, itchy patches of skin), difficulty breathing, chest pain and high or low blood pressure. Your doctor may decide to stop REMICADE treatment and give you medicines to treat the allergic reaction.
- Some patients who have been taking REMICADE for Crohn's disease have had allergic reactions 3 to 12 days after receiving their REMICADE treatment. The symptoms of this type of delayed reaction may include fever, rash, headache and muscle or joint pain. Call your doctor right away if you develop any of these symptoms or any other unusual symptoms such as difficulty swallowing.

Nervous System Disorders:

- There have been rare cases where people taking REMICADE or other TNF blockers have developed disorders that affected their nervous system. Signs that you could be having a problem include: changes in your vision, weakness in your arms and/or legs, and numbness or tingling in any part of your body.

Other Important Information

People who have been treated for rheumatoid arthritis or Crohn's disease for a long time tend to be more prone to a type of blood cancer called lymphoma. There have been some patients that while taking REMICADE developed other types of cancer, but, the number of people taking REMICADE that developed cancer does not seem to be much different from what you would expect to see in people who are not taking REMICADE.

Some patients have developed symptoms that can resemble a disease called lupus. Lupus-like symptoms may include chest discomfort or pain that doesn't go away, shortness of breath, joint pain, or a rash on the cheeks or arms that gets worse in the sun. If you develop any of these symptoms your doctor may decide to stop your treatment with REMICADE.

What are the more common side effects of REMICADE?

The more common side effects with REMICADE are respiratory infections (that may include sinus infections and sore throat), coughing and stomach pain.

Who should not take REMICADE?

YOU SHOULD NOT take REMICADE if you have:

- Heart failure, unless your doctor has talked to you and decided that you are able to take REMICADE.
- Had an allergic reaction to REMICADE or any other product that was made with murine (mouse) proteins.

What health concerns should I talk to my doctor about?

Before receiving your first treatment with REMICADE you should tell your doctor if you:

- Have or think you may have any kind of infection. The infection could be in only one place in your body (such as an open cut or sore), or an infection that affects your whole body (such as the flu). Having an infection could put you at risk for serious side effects with REMICADE.

Continued on next page

Remicade—Cont.

- Have an infection that won't go away or a history of infection that keeps coming back.
- Have had TB (tuberculosis), or if you have recently been with anyone who might have TB. Your doctor will examine you for TB and perform a skin test. If your doctor feels that you are at risk for TB, he or she may start treating you for TB before you begin REMICADE therapy.
- Have lived in or visited an area of the country where an infection called histoplasmosis or coccidioidomycosis (an infection caused by a fungus that affects the lungs) is common. If you don't know if the area you live in is one where histoplasmosis or coccidioidomycosis is common, ask your doctor.
- Have or have previously had heart failure or other heart conditions.
- Have or have had a condition that affects your nervous system, like multiple sclerosis, or Guillain-Barré syndrome, or if you experience any numbness, or tingling, or have had a seizure.
- Are pregnant or nursing.
- Have recently received or are scheduled to receive a vaccine.

Can I take REMICADE while I am on other medicines?

Tell your doctor if you are taking any other medicines including over the counter medicines, supplements or herbal products before you are treated with REMICADE. If you start taking or plan to start taking any new medicine while you are taking REMICADE, tell your doctor. REMICADE and KINERET should not be taken together.

How will REMICADE be given to me?

REMICADE will be given to you by a healthcare professional. REMICADE will be given to you by an IV. This means that the medicine will be given to you through a needle placed in a vein in your arm. It will take about 2 hours to give you the full dose of medicine. During that time and for a period after you receive REMICADE, you will be monitored by a healthcare professional. Your doctor may ask you to take other medicines along with REMICADE. Only a health care professional should prepare the medicine and administer it to you.

How often will I receive REMICADE?

Rheumatoid Arthritis
If you are receiving REMICADE for rheumatoid arthritis you will receive your first dose followed by additional doses at 2 and 6 weeks after the first dose. You will then receive a dose every 8 weeks. Your doctor will monitor your response to REMICADE and may change your dose or dose you more frequently (as often as every 4 weeks).

Crohn's Disease or Fistulizing Crohn's Disease

If you are receiving REMICADE for active Crohn's disease or fistulizing Crohn's disease, you will receive your first dose followed by additional doses at 2 and 6 weeks after the first dose. You will then receive a dose every 8 weeks. Your doctor will monitor your response to REMICADE and may change your dose.

What if I still have questions?

If you have any questions, or problems, always talk first with your doctor. You can also visit the REMICADE internet site at www.remicade.com.

Product developed and manufactured by:

Centocor, Inc.
200 Great Valley Parkway
Malvern, PA 19355
Revised July 2004
IN04240

Shown in Product Identification Guide, page 310

Cephalon, Inc.

145 BRANDYWINE PARKWAY
WEST CHESTER, PA 19380

For Medical Information and Adverse
Drug Experience/Product Complaint Reporting Contact:
(800) 896-5855
Fax 610-738-6669

ACTIQ®

(ak' tek)

(oral transmucosal fentanyl citrate)

PHYSICIANS AND OTHER HEALTHCARE PROVIDERS MUST BECOME FAMILIAR WITH THE IMPORTANT WARNINGS IN THIS LABEL.

Actiq is indicated only for the management of breakthrough cancer pain in patients with malignancies who are already receiving and who are tolerant to opioid therapy for their underlying persistent cancer pain.

Patients considered opioid tolerant are those who are taking at least 60 mg morphine/day, 50 mcg transdermal fentanyl/hour, or an equianalgesic dose of another opioid for a week or longer.

Because life-threatening hypoventilation could occur at any dose in patients not taking chronic opiates, Actiq is contraindicated in the management of acute or postop-

erative pain. This product **must not** be used in opioid non-tolerant patients.

Actiq is intended to be used only in the care of cancer patients and only by oncologists and pain specialists who are knowledgeable of and skilled in the use of Schedule II opioids to treat cancer pain.

Patients and their caregivers must be instructed that Actiq contains a medicine in an amount which can be fatal to a child. Patients and their caregivers must be instructed to keep all units out of the reach of children and to discard opened units properly. (See Information for Patients and Their Caregivers for disposal instructions.)


WARNING: May be habit forming

DESCRIPTION

Actiq (oral transmucosal fentanyl citrate) is a solid formulation of fentanyl citrate, a potent opioid analgesic, intended for oral transmucosal administration. Actiq is formulated as a white to off-white solid drug matrix on a handle that is radiopaque and is fracture resistant (ABS plastic) under normal conditions when used as directed.

Actiq is designed to be dissolved slowly in the mouth in a manner to facilitate transmucosal absorption. The handle allows the Actiq unit to be removed from the mouth if signs of excessive opioid effects appear during administration.

Active ingredient: Fentanyl citrate, USP is N-(1-Phenethyl-4-piperidyl) propionanilide citrate (1:1). Fentanyl is a highly lipophilic compound (octanol-water partition coefficient at pH 7.4 is 816:1) that is freely soluble in organic solvents and sparingly soluble in water (1:40). The molecular weight of the free base is 336.5 (the citrate salt is 528.6). The pKa of the tertiary nitrogens are 7.3 and 8.4. The compound has the following structural formula:



Actiq is available in six strengths equivalent to 200, 400, 600, 800, 1200, or 1600 mcg fentanyl base that is identified by the text on the solid drug matrix, the dosage unit handle tag, the blister package, and the shelf carton.

Inactive ingredients: Hydrated dextrates, citric acid, dibasic sodium phosphate, artificial berry flavor, magnesium stearate, modified food starch, and confectioner's sugar.

CLINICAL PHARMACOLOGY AND PHARMACOKINETICS

Pharmacology:

Fentanyl, a pure opioid agonist, acts primarily through interaction with opioid mu-receptors located in the brain, spinal cord and smooth muscle. The primary site of therapeutic action is the central nervous system (CNS). The most clinically useful pharmacologic effects of the interaction of fentanyl with mu-receptors are analgesia and sedation.

Other opioid effects may include somnolence, hypoventilation, bradycardia, postural hypotension, pruritus, dizziness, nausea, diaphoresis, flushing, euphoria and confusion or difficulty in concentrating at clinically relevant doses.

Clinical Pharmacology

Analgesia:

The analgesic effects of fentanyl are related to the blood level of the drug, if proper allowance is made for the delay into and out of the CNS (a process with a 3-to-5-minute half-life). In opioid non-tolerant individuals, fentanyl provides effects ranging from analgesia at blood levels of 1 to 2 ng/mL, all the way to surgical anesthesia and profound respiratory depression at levels of 10-20 ng/mL.

In general, the minimum effective concentration and the concentration at which toxicity occurs rise with increasing tolerance to any and all opioids. The rate of development of tolerance varies widely among individuals. As a result, the dose of Actiq should be individually titrated to achieve the desired effect (see **DOSE AND ADMINISTRATION**).

Gastrointestinal (GI) Tract and Other Smooth Muscle:

Opioids increase the tone and decrease contractions of the smooth muscle of the gastrointestinal (GI) tract. This results in prolongation in GI transit time and may be responsible for the constipating effect of opioids. Because opioids may increase biliary tract pressure, some patients with biliary colic may experience worsening of pain.

While opioids generally increase the tone of urinary tract smooth muscle, the overall effect tends to vary, in some cases producing urinary urgency, in others, difficulty in urination.

Respiratory System:

All opioid mu-receptor agonists, including fentanyl, produce dose dependent respiratory depression. The risk of respiratory depression is less in patients receiving chronic opioid therapy who develop tolerance to respiratory depression and other opioid effects. During the titration phase of the clinical trials, somnolence, which may be a precursor to respiratory depression, did increase in patients who were treated with higher doses of Actiq. In studies of opioid non-tolerant subjects, respiratory rate and oxygen saturation typically decrease as fentanyl blood concentration increases. Typically, peak respiratory depressive effects (decrease in respiratory rate) are seen 15 to 30 minutes from the start of oral transmucosal fentanyl citrate (OTFC®) administration and may persist for several hours.

Serious or fatal respiratory depression can occur, even at recommended doses, in vulnerable individuals. As with other potent opioids, fentanyl has been associated with cases of serious and fatal respiratory depression in opioid non-tolerant individuals.

Fentanyl depresses the cough reflex as a result of its CNS activity. Although not observed with Actiq in clinical trials, fentanyl given rapidly by intravenous injection in large doses may interfere with respiration by causing rigidity in the muscles of respiration. Therefore, physicians and other healthcare providers should be aware of this potential complication. (See **BOX WARNING, CONTRAINDICATIONS, WARNINGS, PRECAUTIONS, ADVERSE REACTIONS, AND OVERDOSAGE** for additional information on hypoventilation.)

Pharmacokinetics

Absorption:


The absorption pharmacokinetics of fentanyl from the oral transmucosal dosage form is a combination of an initial rapid absorption from the buccal mucosa and a more prolonged absorption of swallowed fentanyl from the GI tract. Both the blood fentanyl profile and the bioavailability of fentanyl will vary depending on the fraction of the dose that is absorbed through the oral mucosa and the fraction swallowed.

Absolute bioavailability, as determined by area under the concentration-time curve, of 15 mcg/kg in 12 adult males was 50% compared to intravenous fentanyl.

Normally, approximately 25% of the total dose of Actiq is rapidly absorbed from the buccal mucosa and becomes systemically available. The remaining 75% of the total dose is swallowed with the saliva and then is slowly absorbed from the GI tract. About 1/3 of this amount (25% of the total dose) escapes hepatic and intestinal first-pass elimination and becomes systemically available. Thus, the generally observed 50% bioavailability of Actiq is divided equally between rapid transmucosal and slower GI absorption. Therefore, a unit dose of Actiq, if chewed and swallowed, might result in lower peak concentrations and lower bioavailability than when consumed as directed.

Dose proportionality among four of the available strengths of Actiq (200, 400, 800, and 1600 mcg) has been demonstrated in a balanced crossover design in adult subjects. Mean serum fentanyl levels following these four doses of Actiq are shown in Figure 1. The curves for each dose level are similar in shape with increasing dose levels producing increasing serum fentanyl levels. C_{max} and $AUC_{0-\infty}$ increased in a dose-dependent manner that is approximately proportional to the Actiq administered.

Figure 1
Mean Serum Fentanyl Concentration (ng/mL)
in Adult Subjects Comparing 4 Doses of Actiq



The pharmacokinetic parameters of the four strengths of Actiq tested in the dose-proportionality study are shown in Table 1. The mean C_{max} ranged from 0.39-2.51 ng/mL. The median time of maximum plasma concentration (T_{max}) across these four doses of Actiq varied from 20-40 minutes (range of 20-480 minutes) as measured after the start of administration.

(See table 1 at top of next page)

Distribution:

Fentanyl is highly lipophilic. Animal data showed that following absorption, fentanyl is rapidly distributed to the brain, heart, lungs, kidneys and spleen followed by a slower redistribution to muscles and fat. The plasma protein binding of fentanyl is 80-85%. The main binding protein is alpha-1-acid glycoprotein, but both albumin and lipoproteins contribute to some extent. The free fraction of fentanyl increases with acidosis. The mean volume of distribution at steady state (V_{ss}) was 4 L/kg.

Metabolism:

Fentanyl is metabolized in the liver and in the intestinal mucosa to norfentanyl by cytochrome P450 3A4 isoenzyme. Norfentanyl was not found to be pharmacologically active in animal studies (see **PRECAUTIONS: Drug Interactions** for additional information).

Elimination:

Fentanyl is primarily (more than 90%) eliminated by biotransformation to N-dealkylated and hydroxylated inactive metabolites. Less than 7% of the dose is excreted unchanged in the urine, and only about 1% is excreted unchanged in the feces. The metabolites are mainly excreted in the urine, while fecal excretion is less important. The total plasma clearance of fentanyl was 0.5 L/hr/kg (range 0.3-0.7 L/hr/kg). The terminal elimination half-life after OTFC administration is about 7 hours.