

540.5
JA

Coden: JACSAT

LIBRARY
FEB 14 1979
UNIVERSITY OF
WASHINGTON

101(2)

LIBRARY USE ONLY

JOURNAL

OF THE

AMERICAN CHEMICAL SOCIETY

VOL. 101

JANUARY 17, 1979

NO. 2



JOURNAL OF THE AMERICAN CHEMICAL SOCIETY

EDITOR

CHEVES WALLING

*Department of Chemistry, University of Utah, Salt Lake City, Utah 84112
(801) 581-5739*

BOARD OF EDITORS

Carl H. Brubaker, Jr.
Ernest R. Davidson
Ernest Grunwald

Jack Halpern
Emil Thomas Kaiser
Robert W. Parry
Barry M. Trost

Regitze R. Vold
Robert L. Vold
Peter J. Wagner

Peter A. S. Smith, *Book Review Editor*

EDITORIAL ADVISORY BOARD

Stephen J. Benkovic
Elkan R. Blout
John I. Brauman
F. Albert Cotton

Philip J. Elving
Christopher S. Foote
Roald Hoffmann
Robert E. Ireland

Josef Michl
Earl L. Muetterties
Koji Nakanishi
Paul v. R. Schleyer

Howard Simmons
Franz Sondheimer
Jacob Szmuszkovicz
Kenneth N. Trueblood
Robert Ullman

Published by the
AMERICAN CHEMICAL SOCIETY
BOOKS AND JOURNALS DIVISION

D. H. Michael Bowen, Director

Marjorie Laflin, Assistant to the Director

Journals Department: Charles R. Bertsch,
Head; Marianne C. Brogan, Associate Head;
Eileen B. Segal, Production Editor; Fern S.
Jackson and Deborah A. Wilson, Associate
Editors; Andrew J. D'Amelio, Editorial As-
sistant; Judith A. Bennison, Production As-
sistant

Magazine and Production Department: Basil
Guiley, Head

Research and Development Department: Seldon
W. Terrant, Head

Advertising Office: Centcom, Ltd., 25 Sylvan
Road South, Westport, Conn. 06880

© Copyright, 1979, by the American Chemical Society. Permission of the American Chemical Society is granted for libraries and other users to make reprographic copies for use beyond that permitted by Sections 107 or 108 of the U.S. Copyright Law, provided that the copying organization pay the appropriate per-copy fee through the Copyright Clearance Center, Inc. Educational institutions are generally granted permission to copy upon application to Office of the Director, Books and Journals Division, at the ACS Washington address.

Published biweekly by the American Chemical Society at 20th and Northampton Sts., Easton, Pennsylvania 18042. Second class postage paid at Washington, D.C., and at additional mailing offices.

Editorial Information

Instructions for authors are printed in the first issue of each volume. Please conform to these instructions when submitting manuscripts.

Manuscripts for publication should be submitted to the Editor, Cheves Walling, at his Salt Lake City address. Correspondence regarding accepted papers and proofs should be directed to the Journals Department at the ACS Easton address.

Page charges of \$40.00 per page may be paid for papers published in this journal. Payment does not affect acceptance or scheduling of papers. Bulk reprints or photocopies of individual articles are available. For information write to Business Operations, Books and Journals Division at the ACS Washington address.

The American Chemical Society and its Editors assume no responsibility for the statements and opinions advanced by contributors. Views expressed in the editorials are those of the author and do not necessarily represent the official position of the society.

Subscription and Business Information

1979 subscription prices, printed or microfiche, including postage. Microfiche by air mail, printed by surface mail. Printed edition air mail or air freight rates available from Membership & Subscription Services at the ACS Columbus address.

	U.S.	Foreign
Members	\$42.00	\$58.00
Nonmembers	168.00	184.00
Supplementary material (available in microfiche only)	35.00	50.00

New and renewal subscriptions should be sent with payment to the Office of the Controller at the ACS Washington address. Changes of address must include both old and new addresses with ZIP code and a recent mailing label. Send

all address changes to the ACS Columbus address. Please allow 6 weeks for change to become effective. Claims for missing numbers will not be allowed if loss was due to failure of notice of change of address to be received in the time specified; if claim is dated (a) North America: more than 90 days beyond issue date, (b) all other foreign: more than one year beyond issue date; or if the reason given is "missing from files." Hard copy claims are handled at the ACS Columbus address.

Microfilm and microfiche editions of all the ACS primary publications, by single volume or back issue collection, are available. For additional price and subscription information, contact Microforms Program at the ACS Washington address, or call (202) 872-4554.

To order single issues or back volumes of the printed edition, contact Special Issues Sales at the ACS Washington address, or call (202) 872-4365. Current year single issue \$6.50. Prior year single issue \$7.00. Back volume \$175.00. Foreign postage additional.

Supplementary material is noted in the Table of Contents with a ■. See supplementary material notice at the end of article for number of pages. Orders over 20 pages are available only on 24X microfiche. Orders must state photocopy or microfiche and give complete title of article, names of authors, journal, date, and page numbers. Prepayment is required and prices are subject to change.

	U.S.	Foreign
Microfiche	\$3.00	\$4.00
Photocopy		
1-8 pages	5.50	7.00
9-20 pages	6.50	8.00

Single microfiche or paper copies of supplementary material may be ordered from Business Operations, Books and Journals Division, at the ACS Washington address. Supplementary material also appears in the microfilm edition.

American Chemical Society
1155 16th Street, N.W.
Washington, D.C. 20036
(202) 872-4600

Membership & Subscription Services
American Chemical Society
P.O. Box 3337
Columbus, Ohio 43210
(614) 421-7230

Journals Department
American Chemical Society
20th & Northampton Sts.
Easton, Pennsylvania 18042
(215) 258-9111

Notice to Authors and sample Copyright Transfer form printed in the issue of January 3, 1979

JOURNAL

OF THE AMERICAN CHEMICAL SOCIETY

VOLUME 101, NUMBER 2
JANUARY 17, 1979

JACSAT 101(2) 279-514 (1979)
ISSN 0002-7863

Dynamic and Static Aspects of Solubilization of Neutral Arenes in Ionic Micellar Solutions Mats Almgren, Franz Grieser,* and J. Kerry Thomas	279
Molecular and Internal Reorientation in 1,1,2,2-Tetraphenylethane F. M. Llort, K. Mislow,* J. B. Wooten, A. L. Beyerlein, G. B. Savitsky, and J. Jacobus*	292
Equilibria Involving Cation Radical Ion Pairs Ignacio J. Ocasio and Paul D. Sullivan*	295
Tunnelling in Collinear Light-Heavy-Heavy Reactions Werner Jakubetz	298
Sudden Polarization: Pyramidalization of Twisted Ethylene Bernard R. Brooks and Henry F. Schaefer III*	307
Conformations, Stabilities, and Charge Distributions in 2- and 3-Monosubstituted Thiophenes. An ab Initio Molecular Orbital Study James Kao* and Leo Radom*	311
Theoretical Exploration of the Photochemical Rearrangement of Oxaziridines Esther Oliveros, Monique Riviere, J. P. Malrieu,* and Ch. Teichteil	318
An Electrochemical and Spectroscopic Study of Some Aromatic Hydrocarbons in the Room Temperature Molten Salt System Aluminum Chloride- <i>n</i> Butylpyridinium Chloride J. Robinson and R. A. Osteryoung*	323
Ion-Solvent Interaction. Effects of Added Polar Compounds on the Conductances of Several Alkali Metal Salts in 2-Butanone at 25 °C M. D. Jackson and W. R. Gilkerson*	328 ■
Lanthanide Ion Probes of Structure in Biology. Laser-Induced Luminescence Decay Constants Provide a Direct Measure of the Number of Metal-Coordinated Water Molecules William DeW. Horrocks, Jr.,* and Daniel R. Sudnick	334
Effect of Halide Complexation of Cadmium(II) on Cadmium-113 Chemical Shifts Joseph J. H. Ackerman, Thomas V. Orr, Victor J. Bartuska, and Gary E. Maciel*	341
Trifluoromethanol (CF ₃ OH) and Trifluoromethylamine (CF ₃ NH ₂) Gerhard Klöter and Konrad Seppelt*	347
Infrared Photochemistry of Halogenated Ethylenes Christopher Reiser, Frances M. Lussier, Craig C. Jensen, and Jeffrey I. Steinfeld*	350
Substituent Effects of the Trifluoromethyl Group on Electrophilic Additions to Alkenes. Solvolysis of the Trifluoromethyl Group. Protonation of Alkenes Less Basic Than Ethylene, <i>p</i> ⁺ Values of Deactivated Styrenes, and Reactivity-Selectivity Effects K. M. Koshy, Danielle Roy, and Thomas T. Tidwell*	357
Skeletal Isomerization of Platinacyclobutanes and Its Relevance to the Mechanism of Olefin Metathesis Razak J. Al-Essa, Richard J. Puddephatt,* Mohammed A. Quysar, and Charles F. H. Tipper	364
Phosphoramidate Stabilized Allylic Carbanions. New Homoenoate Anion Equivalents David A. Evans,* James M. Takacs, and Kenneth M. Hurst	371
Competing Triplet Reactions in Azidoketones Peter J. Wagner* and B. J. Scheve	378
Proton Transfers of Substituted Ammonium Salts. 14. The 1, <i>cis</i> -2,6-Trimethylpiperidinium Ion in Anhydrous Acidic Dimethyl Sulfoxide Jean-Jacques Delpuech* and Bernard Bianchin	383
Diphenylthiirene 1-Oxide. Synthesis, Characterization, and Reactivity Louis A. Carpino* and Hung-Wu Chen	390
Manifestation of Dual Aromaticity in Doubly Charged Annulated Pentalenes Itamar Willner, James Y. Becker, and Mordecai Rabinovitz*	395
Liquid Crystalline Solvents as Mechanistic Probes. 3. The Influence of Ordered Media on the Efficiency of the Photodimerization of Acenaphthylene Jeanne M. Nerbonne and Richard G. Weiss*	402
Stereospecific Acid-Catalyzed Rearrangements of 5,12-Dimethylpentacyclo[6.4.0.0 ^{2,5} .0 ^{3,12} .0 ^{4,9}]dodecane-6,11-diones with Their Strain Release to Biscordiamantane and Diprotadamantane Ring Systems Ken-ichi Hirao, Mikio Taniguchi, Osamu Yonemitsu,* Judith L. Flippen, and Bernhard Witkop	408
Mechanistic Aspects of the Thermal Equilibration of Retinylidene Imines and Immonium Salts Patrick C. Mowery* and Walther Stoeckenius	414
Relativity in Methyl Transfer Reactions. 3. Equilibria and Rates in Transfers between Substituted Thiophenoxides Edward S. Lewis* and Semyon Kukes	417
The Mechanism of Alkaline Hydrolysis of Thiazolidines Roberta Lohowy and Frank Meneghini*	420 ■
Comments on the Putative Stereoselectivity in Cysteine-Aldehyde Reactions. Selective C(2) Inversion and C(4) Epimerization in Thiazolidine-4-carboxylic Acids László Szilágyi* and Zoltán Györgydeák	427

Use of Polymers as Protecting Groups in Organic Synthesis. Application of Polystyrylboronic Acid to the One-Pot Synthesis of Acylated Carbohydrate Derivatives	Jean M. J. Fréchet,* Lucy J. Nuyens, and Elizabeth Seymour	432
Amino Acid Constituents of Ristocetin A	Constance M. Harris,* Jeffrey J. Kibby, James R. Fehlner, Austin B. Raabe, Thomas A. Barber, and Thomas M. Harris*	437
Crystal and Molecular Structure of the Cyclic Hexapeptide <i>cyclo</i> -(Gly-Pro-d-Phe) ₂	Joe N. Brown* and C. H. Yang	445 ■
Conformations of Poly(ethylene glycol) Bound Homooligo-L-alanines and -L-valines in Aqueous Solution	Claudio Toniolo,* Gian Maria Bonora, and Manfred Mutter	450
Studies of the Thermodynamics of Electron Transfer Reactions of Blue Copper Proteins	Napapon Sailasuta, F. C. Anson, and Harry B. Gray*	455
Mechanism of Alkaline Hydrolysis of <i>S</i> -Adenosyl-L-methionine and Related Sulfonium Nucleosides	Ronald T. Borchardt	458
Variation of the Linear Polarization Across the Emission Band of Nicotinamide 1, <i>N</i> ⁶ -Ethenoadenine Dinucleotide Bound to Dehydrogenases	A. Gafni,* J. Schlessinger, and I. Z. Steinberg	463
Electrochemistry of Vitamin B-12. 4. Kinetics and Mechanisms in B-12a-B-12r Oxido-Reduction	N. R. de Tacconi, D. Lexa, and J. M. Savéant*	467
A Study of the Peptide Hormone Oxytocin and of Prolylleucylglycinamide by ¹⁵ N NMR	David H. Live, Herman R. Wyssbrod, Alan J. Fischman, William C. Agosta, Craig H. Bradley, and David Cowburn*	474
Intramolecular Water Bridge between the 2'-OH and Phosphate Groups of RNA. Cyclic Nucleotides as a Model System	Philip H. Bolton and David R. Kearns*	479

COMMUNICATIONS TO THE EDITOR

Satellite Structures in the ESCA Spectra of (Diphthalocyaninato)lanthanides(III) and -actinides(IV)	K. Tatsumi, K. Kasuga, and M. Tsutsui*	484
Efficient Trapping of Silylenes through Disilene Intermediates	Yasuhiro Nakadaira, Toshiaki Kobayashi, Tatsuo Otsuka, and Hideki Sakurai*	486
<i>trans</i> - and <i>cis</i> -1,2-Dimethyl-1,2-diphenyldisilene. Is Si=Si a True Double Bond?	Hideki Sakurai,* Yasuhiro Nakadaira, and Toshiaki Kobayashi	487
Activation of Aldehyde C-H Bonds to Oxidative Addition via Formation of 3-Methyl-2-aminopyridyl Aldimines and Related Compounds: Rhodium Based Catalytic Hydroacylation	J. William Suggs	489
Reactions of Aniline with Olefins Catalyzed by Group 8 Metal Complexes: N-Alkylation and Heterocycle Formation	Steven E. Diamond,* Andrew Szalkiewicz, and Frank Mares*	490
Polyoxomolybdate-Hydrocarbon Interactions. Synthesis and Structure of the CH ₃ Mo ₄ O ₁₅ H ³⁻ Anion and Related Methylenedioxymolybdates	V. W. Day, M. F. Fredrich, W. G. Klemperer,* and R.-S. Liu	491
An Unusual Rearrangement of Ajaconine: An Example of a "Disfavored" 5-Endo-Trigonal Ring Closure	S. William Pelletier* and Naresh V. Mody	492
Cyclization Reactions via Oxo- π -allylpalladium(II) Intermediates	Yoshihiko Ito, Hirokazu Aoyama, Toshikazu Hirao, Akira Mochizuki, and Takeo Saegusa*	494
Fourier Transform Infrared Vibrational Circular Dichroism	Laurence A. Nafie,* Max Diem, and D. Warren Vidrine	496
On the Dynamic Behavior of Trans-Dihalide-Bishydrazone-Palladium(II) Complexes	Giovanni Natile,* Lucio Cattalini, Francesco Gasparrini, and Luciano Caglioti	498
Formation of Metal Chelates in Secondary Ion Mass Spectrometry. Comparisons with Solution Chemistry	R. J. Day, S. E. Unger, and R. G. Cooks*	499
Ionization of Quaternary Nitrogen Compounds by Secondary Ion Mass Spectrometry	R. J. Day, S. E. Unger, and R. G. Cooks*	501
Reactions in Dry Media. Cleavage of Carbon Carbon Single Bonds by Ozonation.	Daniel Tal, Ehud Keinan, and Yehuda Mazur*	502
Organometallic Models for Possible Fischer-Tropsch Intermediates. Synthesis, Structure, and Reactions of a Formaldehyde Complex of Osmium	Kevin L. Brown, George R. Clark, Christine E. L. Headford, Karen Marsden, and Warren R. Roper*	503
Ips0 Nitration. Preparation of 4-Methyl-4-nitrocyclohexadienols and Detection of Intramolecular Hydrogen Migration (NIH Shift) upon Solvolytic Rearomatization	K. S. Feldman, Ann McDermott, and P. C. Myhre*	505
Electrophilic Aromatic Protonation of Stable π Carbocations	Richard M. Pagni* and Robert J. Smith	506
Novel Bridging Sulfide Anion Complexes of the Hexacarbonyldiiron Unit: a New Route to Alkylthio Complexes of Iron	Dietmar Seyferth* and Richard S. Henderson	508
The Case for Stereoelectronic Axial Migration of Hydrogen in Nonrigid Cyclohexylidenes in Chair Conformation	Linda Seghers Press and Harold Shechter*	509
Stereochemical Course of Glycerol Kinase, Pyruvate Kinase, and Hexokinase: Phosphoryl Transfer from Chiral [γ (<i>S</i>)- ¹⁶ O, ¹⁷ O, ¹⁸ O]ATP	Walter A. Blättler and Jeremy P. Knowles*	510

ADDITIONS AND CORRECTIONS

Structure of Nogalamycin	Paul F. Wiley,* Ronald B. Kelly, E. Louis Caron, Veronica H. Wiley, Jian H. Johnson, Forrest A. MacKellar, and Stephen A. Mizak	512
The Microwave Spectrum, Structure, and Dipole Moment of 2-Mercaptoethanol; Evidence for an Intramolecular OH...S Hydrogen Bond	Eun-Mo Sung and Marlin D. Harmony*	512
Studies of Nitrogen Metabolism Using ^{13}C NMR Spectroscopy. 1. Streptonigrin Biosynthesis	Steven J. Gould* and Chiu Chin Chang	512
Experimental Evidence for Trapped Valences in the Mixed-Valence Complex μ -Pyrazine-bis(pentaammineruthenium) Tosylate. Electron Paramagnetic Resonance, Magnetic Susceptibility, and Nuclear Magnetic Resonance Results	Bruce C. Bunker, Russell S. Drago,* David N. Hendrickson,* Robert M. Richman, and Stephen L. Kessell	512
π,π -Biradicaloid Hydrocarbons. The Pleiadene Family. 3. A Facile Symmetry-Forbidden Thermal Conversion of a Polycyclic Butadiene Moiety to a Cyclobutene	Richard P. Steiner and Josef Michl*	512
Does a Methyl Substituent Stabilize or Destabilize Anions?	Addy Pross* and Leo Radom*	512
Characterization of the Asymmetric Nitric Oxide Dimer $\text{O}=\text{N}=\text{O}=\text{N}$ by Resonance Raman and Infrared Spectroscopy	J. R. Ohlsen and J. Laane*	512
Photochemistry of Bichromophoric Molecules. Photochemistry and Photophysics of 2-Methylenebenzborbornene and Related Nonconjugated Aryl Olefins in Nonprotic Media	Frank Scully, Tom Nylund, Fred Palensky, and Harry Morrison*	512

BOOK REVIEWS

Equilibrium in Solution and Surface and Colloid Chemistry. By G. Scatchard	Reviewed by Cecil M. Criss	513
Purine Metabolism in Man. II. Volumes 76A and 76B. By M. M. Muller, E. Kaiser, and J. E. Seegmiller	Reviewed by Irving H. Fox	513
Laser Spectroscopy III. Edited by J. L. Hall and J. L. Carlsten	Reviewed by George E. Leroi	513
Chemical Applications of Graph Theory. Edited by A. T. Balaban	Reviewed by Lawrence L. Lohr, Jr.	513
Dye Lasers. By F. P. Schafer	Reviewed by John R. Lombardi	514
Flame Emission and Atomic Absorption Spectrometry. Volume 3, Elements and Matrices. Edited by John A. Dean and Theodore C. Rains	Reviewed by Marvin Margoshes	514
Trace Analysis of Atmospheric Samples. By Kikuo Oikawa	Reviewed by David F. S. Natusch	514
Analytical Profiles of Drug Substances. Volume 6. Edited by Klaus Florey	Reviewed by Charles W. Parker	514

■ Supplementary material for this paper is available separately (consult the masthead page for ordering information); it will also appear following the paper in the microfilm edition of this journal.

* In papers with more than one author, the asterisk indicates the name of the author to whom inquiries about the paper should be addressed.

- K. Hirao, and O. Yonemitsu, *J. Am. Chem. Soc.*, **96**, 2570 (1974); (I) H. D. Becker, B. Ruge, and T. Weslöl, *Tetrahedron Lett.*, 253 (1975).
- (3) This reaction has also been interpreted in terms of the orbital interaction rule: S. Inagaki, H. Fujimoto, and K. Fukui, *J. Am. Chem. Soc.*, **98**, 4693 (1976).
- (4) K. Hirao, M. Taniguchi, T. Iwakuma, O. Yonemitsu, J. L. Flippen, I. L. Karle, and B. Witkop, *J. Am. Chem. Soc.*, **97**, 3249 (1975).
- (5) J. L. Flippen, *Acta Crystallogr., Sect. B*, **32**, 1269 (1976).
- (6) C. A. Grob and P. W. Schliess, *Angew. Chem., Int. Ed. Engl.*, **6**, 1 (1967).
- (7) E. Adler and K. Holmberg, *Acta Chem. Scand., Ser. B*, **28**, 465 (1974).
- (8) J. Karle and I. L. Karle, *Acta Crystallogr.*, **21**, 849 (1966).
- (9) K. Alder, F. H. Flock, and H. Lessenich, *Chem. Ber.*, **90**, 1709 (1957).
- (10) E. Osawa, K. Aigami, and Y. Inamoto, *J. Chem. Soc., Perkin Trans. 2*, in press.
- (11) Otherwise, the rearranged ketone **3d** may be formed directly from **3a** through a concerted 1,2 shift with deprotonation.
- (12) For a review see R. L. Cargill, T. E. Jackson, N. P. Peet, and D. M. Pond, *Acc. Chem. Res.*, **7**, 106 (1974).
- (13) B. P. Mundy and R. D. Otzenberger, *J. Org. Chem.*, **38**, 2109 (1973).
- (14) Cf. E. N. Marvell and E. Magoon, *J. Am. Chem. Soc.*, **77**, 2542 (1955); B. Miller, *ibid.*, **92**, 6252 (1970).
- (15) I. F. Halverstadt and W. D. Kumler, *J. Am. Chem. Soc.*, **64**, 2988 (1942).

This material may be protected by Copyright law (Title 17 U.S. Code)

Mechanistic Aspects of the Thermal Equilibration of Retinylidene Imines and Immonium Salts

Patrick C. Mowery* and Walther Stoeckenius

Contribution from the Cardiovascular Research Institute, University of California, San Francisco, California 94143. Received July 6, 1978

Abstract: The reactions of isomeric retinylidene *n*-butylimines and protonated imines in the dark were examined by proton nuclear magnetic resonance and ultraviolet absorption spectroscopy and by high-performance liquid chromatography. The immonium hydrochlorides were shown to decompose by a complex system of pathways. However, 13-*cis*-retinylidene-*n*-butylamine was shown to undergo thermodynamic equilibration with the trans isomer by general base catalysis, whereas the 9-*cis* isomer was thermodynamically stable. The specificity of the reaction for the 13-*cis* isomer suggests that it occurs via the 13-methylene enamine intermediate. Protonated retinylidene imines have been shown to be important as chromophores in the visual cycle and in a light-driven proton pump in certain halophilic bacteria. The mechanisms of action for these systems are briefly contrasted in the light of the result reported here.

Introduction

Retinal imines have long served an important role as models of the visual chromophore.¹ Interest in their study has been further augmented by the recent discovery of a light-driven proton pump in certain halophilic bacteria which catalyzes the photophosphorylation of ADP.² The pump has been isolated, and has been shown to consist of 25% lipid and 75% of a single protein called bacteriorhodopsin of molecular weight 25 000. Furthermore, its intense purple color is due to a retinyl moiety attached to an ϵ amino group of a lysine.³ It exists in two forms: (1) the dark-adapted form, characterizable by a maximum visible spectral transition at 558 nm, which is believed to contain a 1:1 mixture of the chromophore as 13-*cis* and all-*trans*- ϵ -retinylidene-L-lysine;⁴ and (2) a light-adapted state with a transition at 568 nm, in which it is believed to be essentially all *trans*.^{4,5}

It is unclear whether a photolytic all-*trans* to 13-*cis* isomerization is important for a proton pumping,⁶ although there is some evidence from chromophore extraction data that an intermediate exists with a 13-*cis*-retinyl chromophore.⁷ If this is so, the 13-*cis* intermediate thermally reisomerizes to *trans*, since there must be a cyclic mechanism for the bacteriorhodopsin proton pump.⁸

The purpose of this paper is to demonstrate that at least one mechanism exists for the nonphotolytic rearrangement of the 13-*cis* imine to the all-*trans* form. The immonium hydrochloride, on the other hand, decomposes by several reaction pathways. Investigators are therefore cautioned to use care in the preparation and study of these compounds. Finally, the relevance of these findings to the visual cycle and to the bacteriorhodopsin system is briefly discussed.

Materials and Methods

Materials. all-*trans*-, 13-*cis*-, and 9-*cis*-retinal (Sigma) were checked for purity on a Waters ALC/GPC 204 liquid chro-

matograph outfitted with a dual-wavelength (254 and 365 nm) detector and a μ -Porasil column, as described by Pettei et al.⁷ The eluent was 2% ether in hexane with a flow rate of 2.0 mL/min. Proton magnetic resonance spectra were taken on a Perkin-Elmer R12b spectrometer, outfitted with a temperature control, at 37 °C. Absorption spectra were recorded on a Cary 14 spectrometer. *N*-Butylamine (Mallinckrodt) was distilled prior to use and stored over molecular sieves. Retinals were used without purification for the NMR experiments, but were purified by high-performance liquid chromatography before studies involving electronic spectra were made.

Preparation of Imines. All operations were conducted in semidarkness at 0 °C. Retinal was allowed to react with a 0.05 molar excess of *n*-butylamine in anhydrous ether for 30 min. The solvent was removed on a rotary evaporator, and the excess amine was driven off by passing dry nitrogen over the residue for 30 min. The protonated imine was prepared by washing the residue with ether saturated with anhydrous hydrogen chloride and removing the solvent on the rotary evaporator. Excess acid was further removed by a nitrogen purge. Samples were dissolved in chloroform or (for NMR experiments) chloroform-*d* containing 1% tetramethylsilane.

Results

NMR Studies. Our interest in this study originated when we encountered difficulties in obtaining ¹³C NMR spectra of 13-*cis*-retinylidene-*n*-butylamine, owing to its conversion to the *trans* isomer at ambient temperatures. We were able to conveniently follow the isomerization by ¹H NMR (Figure 1), and noted that the half-life was roughly 1 h. In a separate experiment, we followed the decay of the 13-*cis* imine at 37 °C by LC, using 2% ether in hexane as the eluent, and found a decay in the 13-*cis* isomer from 85 to 45% in 80 min for a 0.1 M solution.

In NMR experiments, the immonium hydrochloride was




Figure 1. Vinyl proton NMR spectra of 0.14 M retinyl imines at 37 °C in chloroform-*d* containing 1% Me₄Si (ppm from Me₄Si): (a) 13-*cis*-retinylidene-*n*-butylimine; after (b) 37 min, (c) 3 h, (d) 6 h, (e) freshly prepared all-*trans* imine.

studied as well. The complex vinyl region of the protonated imines is unresolved at 60 MHz. However, it is qualitatively clear from Figure 2 that a 0.15 M solution of the 13-*cis* isomer decomposes by a variety of competing pathways at 35 °C and that a simple *cis*-*trans* isomerization is not immediately evident as the major route.

Electronic Spectral Studies. The molecularity of the isomerization was examined by preparing a 10 mM stock solution of the 13-*cis* imine in chloroform, and then preparing cuvettes containing a 1:25 (1-mm cuvette) and a 1:250 (1-cm cuvette) dilution. The three containers were kept at 32 °C, and their UV spectra were monitored periodically. The rearrangement was essentially complete after 3 h in the stock solution, whereas the dilute solutions were essentially unchanged. However, when the 40- μ M solution was made 100 mM in *n*-butylamine, it quickly isomerized to the *trans* imine. In identical experiments, 13-*cis*-retinylidene-*n*-butylimmonium hydrochloride did not isomerize, although the stock solution did begin to decompose, as was witnessed by line broadening of its major (460 nm) spectral transition.

The rate of *n*-butylamine-catalyzed isomerization of the 13-*cis* imine was measured at 25 °C, as shown in Figure 3.




Figure 2. Vinyl proton NMR spectra of 0.14 M protonated retinyl imines at 37 °C in chloroform-*d* (ppm from Me₄Si): (a) 13-*cis*-retinylidene-*n*-butylimmonium chloride; (b) after 5 h; (c) freshly prepared all-*trans*-immonium chloride.




Figure 3. UV spectra showing the reaction between 13-*cis*-retinal (14 μ M) and *n*-butylamine (100 mM) at 25 °C in chloroform: 13-*cis*-retinal (a); formation of 13-*cis* imine (b, c); formation of all-*trans* imine (d-i). Time after adding amine (min): (b) 5; (c) 11; (d) 22; (e) 47; (f) 65; (g) 100; (h) 130; (i) 211.

Butylamine (0.2 μ mol) was added to a solution of 13-*cis* retinal (0.028 μ mol) in chloroform (2.0 mL), and the solution was monitored spectroscopically. The *cis* imine was formed in 10 min, and then smoothly isomerized to the *trans* isomer. The spectral changes as the 13-*cis* and all-*trans* imines are formed




Figure 4. Electronic spectra showing the reaction between 9-*cis*-retinal (9.7 μ M) and *n*-butylamine (100 mM) at 25 $^{\circ}$ C in chloroform: (a) 9-*cis*-retinal; formation of 9-*cis* imine. Hours after adding amine: (b) 0.5; (c) 1.0; (d) 6.0; (e) 30.

are qualitatively consistent with the spectral data for the isomeric retinals and imines published by Schaffer et al.⁹ The pseudo-first-order rate could be conveniently monitored by adding *n*-butylamine to a solution of the imine and measuring the decrease in absorbance at 450 nm, giving a half-life of 26 min for a 0.1 mM amine and 15 μ M imine at 25 $^{\circ}$ C (correlation coefficient for a linear regression analysis: 0.998), and a second-order rate constant of $4.4 \times 10^{-3} \text{ M}^{-1} \text{ s}^{-1}$.

All of the bases used here could potentially react by a transimination mechanism, and not by simple general base catalysis. The 13-*cis* imine was therefore prepared, and triethylamine was added under conditions in which the concentrations of amine (70 μ M) and imine (20 μ M) were similar to those used for *n*-butylamine. The rearrangement was much faster ($t_{1/2} \sim 7 \text{ s}$ at 25 $^{\circ}$ C), establishing that the mechanism of action was by general base catalysis.

As a final control, solvent effects on the course of the isomerization were monitored, using ethanol, isopropyl alcohol, and acetonitrile at 25 $^{\circ}$ C. The rates relative to chloroform follow: ethanol (>100) $\gg$ isopropyl alcohol (55.3) $\gg$ acetonitrile (1.6).

The 9-*cis* isomer was investigated in the same fashion, in order to ascertain whether other *cis*-retinylidene imines isomerize. A 9.7 μ M solution of 9-*cis*-retinal in chloroform was allowed to react with *n*-butylamine (100 mM). As shown in Figure 4, the imine quickly formed, but did not noticeably isomerize over a 2-day period at room temperature. From the quantitative data of Schaffer et al.,⁹ one would expect a 9-nm blue shift of λ_{max} and a 20% increase in the extinction coefficient as the imine is formed from 9-*cis*-retinal, and followed by a 1-nm red shift in the major spectral transition accompanied by a 13% increase in ϵ_{max} as the trans imine is formed. The observed spectral changes (a 9-nm blue shift with an ϵ_{max} increase of 18%) are only in accord with the formation of the 9-*cis* imine. Note too that the characteristic *cis* peak is retained throughout.

Discussion

It is clear that 13-*cis*-retinylidene-*n*-butylamine easily isomerizes to the trans imine by a mechanism involving general base catalysis. Although the exact mechanism is unknown, the thermal stability of the 9-*cis* imine militates against mechanisms involving extensive charge delocalizations or a retro form. The simplest hypothesis would therefore consist of general-base-catalyzed thermodynamic equilibration between




Figure 5. Proposed scheme for the thermal equilibration of 13-*cis*- and all-*trans*-retinylidene-*n*-butylamine.

imines via a 13-methylene enamine intermediate (Figure 5). Such intermediates are usually not encountered in high concentrations, because C=N bonds are more stable than C=C bonds when primary amines are the reactants,¹⁰ and because enamines of aldehydes are generally not stable.¹¹ It is noteworthy in this respect that the trans imine showed signs of decomposition at 130 min (Figure 2).

The relative rates of isomerization in ethanol, 2-propanol, acetonitrile, and chloroform are in the order one might expect:¹² the reaction is faster in protic than in aprotic solvents and, within these groups, is faster in the solvent with the higher dielectric constant. We would conclude from this that nonpolar solvents such as 3-methylpentane are the most suitable for studying Schiff bases.

The manifestation of a 13-methylene enamine intermediate and not the 9-methylene or the thermodynamically more stable retroretinyl enamine may be expected under conditions of kinetic control. The 13-methyl hydrogens of the imine are relatively acidic, because they are homoallylic to the nitrogen, whereas the C4, C18, or C19 hydrogens are less so, because the electron-withdrawing nitrogen substituent effect is attenuated as it is broadcast down the chain.

In the light of the base-catalyzed isomerization mechanism discussed here, it is interesting to contrast the evolution of the mechanism of vision with that of the proton pump in halobacteria. As Lewis has pointed out,¹³ the mechanism for vision involves a signal transducer, in which the sensitivity to a single photon event must be maximized. Natural selection has taken advantage of the thermodynamically favorable path of 11-*cis* to all-*trans* isomerization of the chromophore and has discriminated against a reversible cycle and specifically against the involvement of a 13-*cis* isomer. The proton pump, on the other hand, requires light energy to maintain a proton gradient and has opted for a cycle at the expense of sensitivity to the single-photon event. Reversibility is therefore allowed, and the 13-*cis* isomer is found in dark-adapted systems and may be present in the photocycle as well.

This mechanism may play an important role in the light-dark adaptation cycle in bacteriorhodopsin, since the mixture of isomers may be a thermodynamic distribution reflecting an equilibration which is catalyzed by a weak base. Although the mechanism may be important to the photocycle as well, it is unlikely, because the cycle would then involve a kinetically unfavorable protonation of a soft carbon base (the 13-methylene) by a hard acid (an oxygen or nitrogen acid).¹² This supposition is based on the fact that carbon acids and their conjugate bases do not form hydrogen bonds with the oxygen and nitrogen acids and bases one would expect to be involved in enzyme catalysis, thereby vitiating the formation of favorable encounter complexes. Yet, both of these hypotheses can be tested.

The protonated imines react by more than one path, with isomerization competing with Michael additions and the like. However, like their conjugate bases, they are stable at low temperature or at low concentrations. Retinylidene immonium

salts do not undergo thermodynamic equilibration as readily as do retinals in the presence of acids.¹⁴ However, an enzyme-catalyzed route is possible. At any rate, care must be exercised in the preparation and handling of these compounds.

Acknowledgment. This work was supported by NHLI Program Project Grant HL06285.

References and Notes

- (1) Honig, B.; Ebrey, T. G. *Annu. Rev. Biophys. Bioeng.* **1974**, *3*, 151-177.
- (2) Henderson, R. *Annu. Rev. Biophys. Bioeng.* **1977**, *6*, 87-109.
- (3) Oesterhelt, D.; Stoekenius, W. *Nature (London), New Biol.* **1971**, *233*, 149-152.
- (4) Oesterhelt, D.; Meentzen, M.; Schuhman, L. *Eur. J. Biochem.* **1973**, *40*, 453-463.
- (5) Jan, L.-Y. *Vision Res.* **1975**, *15*, 1081-1086.
- (6) Aton, B.; Doukas, A. G.; Callender, R. H.; Becher, B.; Ebrey, T. G. *Biochemistry* **1977**, *16*, 2995-3000.
- (7) Pettei, M. J.; Yudd, A. P.; Nakanishi, K.; Henselman, R.; Stoekenius, W. *Biochemistry* **1977**, *16*, 1955-1958.
- (8) Lozier, R. H.; Bogomolni, R. A.; Stoekenius, W. *Biophys. J.* **1975**, *15*, 955-962.
- (9) Schaffer, A. M.; Waddell, W. H.; Becker, R. S. *J. Am. Chem. Soc.* **1974**, *96*, 2063-2068.
- (10) Franklin, J. L. *Ind. Eng. Chem.* **1949**, *41*, 1070-1076.
- (11) Stork, G.; Brizzolara, A.; Landesman, H.; Szmuszkowicz, J.; Terrell, R. J. *Am. Chem. Soc.* **1963**, *85*, 207-216.
- (12) Szwarc, M.; Streitwieser, A.; Mowery, P. C. "Ions and Ion Pairs in Organic Reactions", Vol. 2; Szwarc, M., Ed.; Wiley: New York, N.Y., 1974; Chapter 2.
- (13) Lewis, A. *Biophys. Struct. Mech.* **1977**, *3*, 97-100.
- (14) Patel, D. J. *Nature (London)* **1969**, *264*, 799-800.

Reactivity in Methyl Transfer Reactions. 3. Equilibria and Rates in Transfers between Substituted Thiophenoxides

Edward S. Lewis* and Semyon Kukes


Contribution from the Department of Chemistry, Rice University,
P.O. Box 1892, Houston, Texas 77001. Received June 12, 1978

Abstract: Methyl phenyl sulfide methylates substituted thiophenoxide ions reversibly at elevated temperatures. Using a GC analysis, the rates and equilibria for five substituted cases have been measured at several temperatures in the range ca. 110-210 °C, and the identity reaction with unsubstituted thiophenoxide has also been measured using a ³⁵S tracer. The Hammett ρ for the forward reaction is close to half that for the equilibrium, suggesting a transition state with a half-transferred methyl group. This represents the first measurement of substituent effect on both rate and equilibrium for a simple organic substitution reaction.

In the first two papers of this series,^{1,2} substituent effects in an attacking nucleophile were compared for leaving groups of greatly differing reactivity in some virtually quantitative reactions. Surprisingly, substituent effects were large and remarkably similar over a wide range of rate, although the reactivity-selectivity principle³ leads to an expectation of reduced substituent effects with the most reactive leaving groups, such as trifluoromethanesulfonate. The term "large" substituent effect implies, however, a standard of comparison, and the only ones available were those of one leaving group with another. A more reasonable comparison is between the substituent effect on the reaction rate and that on the equilibrium position in the same reaction, perhaps expressible as ρ_+ and ρ_{eq} , the Hammett reaction constants for the forward rate constants and the equilibrium constants, respectively (similarly ρ_- for the reverse reaction can be defined, but it is not independent, for $\rho_{eq} = \rho_+ - \rho_-$). The ratio ρ_+/ρ_{eq} can then be used to measure the extent of proton transfer with the same utility and limitations as using the Brønsted α or β (which are also comparisons of effect of structural changes on rate and equilibrium) to measure the extent of proton transfer.⁴

Equilibria have been measured for very few organic reactions, except acidities and oxidation-reduction processes. One, of complex mechanism, is listed in Hammett's tabulation,⁵ Jaffe⁶ lists six more, and most of these do not have rates measured also. More recently, both rates and equilibria have been measured for some diazonium⁷ and carbonium⁸ ion reactions with nucleophiles. There appears to be only one published report of equilibria in methyl transfers,⁹ and rates were not reported. A very relevant study of methyl iodide with pyridine is not yet published.¹⁰

We report the study of the reaction



This reaction is, of course, too slow to study near room temperature, but is fast enough to measure between about 110 and 210 °C in ethanol. Rate constants, measured as initial rates by a gas chromatographic procedure comparing PhSCH_3 and ArSCH_3 peak areas, fit the Arrhenius equation with the results expressed as $\Delta H^\ddagger$ and $\Delta S^\ddagger$, presented in Table I. The case of $\text{Ar} = \text{Ph}$ was followed using ³⁵S-labeled PhS^- . The kinetic law, assumed to be first order in each reagent, was essentially confirmed in one case illustrated in Figure 1, in which experimental points of mole fraction of ArSCH_3 as a function of time in experiments at 144 °C are shown together with the curve calculated from the equation

$$\frac{K}{2A_0} \ln \left[\frac{C\sqrt{K} - C + A_0}{C\sqrt{K} + C - A_0} \right] = k_+t \quad (2)$$

for a reversible second-order reaction with equal starting concentrations, A_0 of the two reagents. In this equation C is the product concentration, and values were $k_+ = 8.0 \times 10^{-7} \text{ M}^{-1} \text{ s}^{-1}$, $K = 2.0$, and $A_0 = 0.90 \text{ M}$. The plot also shows experimental points for the reverse reaction, starting with PhS^- and $\text{CH}_3\text{SC}_6\text{H}_4\text{CH}_3$; the calculated curve uses $k_- = 4 \times 10^{-7} \text{ M}^{-1} \text{ s}^{-1}$ and an initial concentration of 0.80 M. The time scale on Figure 1 shows why this was not done routinely. A further demonstration of the kinetic law is that no reaction was detected when thiol was used instead of the salt.

Equilibrium constants in Table I are calculated as k_+/k_- , but several were measured directly after long time periods.