



US 20100285478A1

(19) **United States**

(12) **Patent Application Publication**

Chen et al.

(10) **Pub. No.: US 2010/0285478 A1**

(43) **Pub. Date: Nov. 11, 2010**

(54) **METHODS, COMPOSITIONS, AND KITS FOR
DETECTING ALLELIC VARIANTS**

(75) Inventors: **Caifu Chen**, Palo Alto, CA (US);
Ruoying Tan, Palo Alto, CA (US)

Correspondence Address:
LIFE TECHNOLOGIES CORPORATION
C/O INTELLEVATE
P.O. BOX 52050
MINNEAPOLIS, MN 55402 (US)

(73) Assignee: **LIFE TECHNOLOGIES
CORPORATION**, Carlsbad, CA
(US)

(21) Appl. No.: **12/748,329**

(22) Filed: **Mar. 26, 2010**

Related U.S. Application Data

(63) Continuation-in-part of application No. 12/641,321,
filed on Dec. 17, 2009.

(60) Provisional application No. 61/258,582, filed on Nov.
5, 2009, provisional application No. 61/253,501, filed
on Oct. 20, 2009, provisional application No. 61/251,

623, filed on Oct. 14, 2009, provisional application No.
61/186,775, filed on Jun. 12, 2009, provisional appli-
cation No. 61/164,230, filed on Mar. 27, 2009.

(30) **Foreign Application Priority Data**

Dec. 17, 2009 (US) PCT/US2009/006652

Publication Classification

(51) **Int. Cl.**
C12Q 1/68 (2006.01)

(52) **U.S. Cl.** **435/6**

(57) **ABSTRACT**

In some embodiments, the present inventions relates gener-
ally to compositions, methods and kits for use in discriminat-
ing sequence variation between different alleles. More spe-
cifically, in some embodiments, the present invention
provides for compositions, methods and kits for quantitating
rare (e.g., mutant) allelic variants, such as SNPs, or nucle-
otide (NT) insertions or deletions, in samples comprising
abundant (e.g., wild type) allelic variants with high specificity
and selectivity. In particular, in some embodiments, the
invention relates to a highly selective method for mutation
detection referred to as competitive allele-specific TaqMan
PCR ("cast-PCR").

Schematic of cast-PCR




Fig. 1

Use of highly discriminating bases for allele detection (e.g., mutant allele)




Fig. 2

Use of MGB at the 5' end of a blocking probe




Fig. 3

Use of modified bases in MGB blockers and allele-specific PCR primers (ASP1 and ASP2)




Fig. 4A

Examples of Modified Nucleosides




Fig. 4B

TaqMan-like Sensitivity and Dynamic Range of cast-PCR




Fig. 5

KRAS mutations

```
1  gtactggtgg agtatttggat agtgtatttaa ccttatgtgt gacatgttct aatatagtcg  
61  cattttcatt atttttatta taaggcgtgg tgaaaatgac aatatataa gtttggata  
121  ttggaagcgg tgggtatgg aaagcggcc ttacataga gttacagag aaacattttt  
181  tggcctgata ttatccatga atagacata atcttgtttt aatatgcata ttactggtgc  
241  aggaccattc ttgatacag ataaaggttt cctgaccat tttcatgact  
// CDS REGION cod 12-13
```

KRAS mutations

- 12 Asp (GGT>GAT)
- 12 Val (GGT>GTT)
- 12 Cys (GGT>TGT)
- 12 Ser (GGT>AGT)
- 12 Ala (GGT>GCT)
- 12 Arg (GGT>CGT)
- 13 Asp (GGC>GAC)

Fig. 6

Specificity of KRAS-183AC cast-PCR Assays




Fig. 7

cast-PCR Assays Can Detect a Single Copy of Mutant DNA in 10⁶ Copies of Wild-type DNA





Fig. 8

Specificity of cast-PCR Assays – Tumor Cell DNA Spiking



Spiking mutated cell line DNA (KRAS-G12A) in to wild type cell line DNA

Fig. 9

Detection of Mutant DNA from Tumor Samples by cast-PCR

SNP ID	Sample ID	Ct_WT	Ct_Mutant	Normalized Δ Ct	Mutant cell DNA (%)
KRAS-38GA	Normal-1	23.2	40.0	16.8	0.00
	Tumor-01	27.8	30.3	3.6	7.62
EGFR-2573TG	Normal-1	22.9	40.0	17.1	0.00
	Tumor-02	27.7	25.4	-2	80.00
CTNNB1-134CT	Normal-1	22.8	40	17.2	0.00
	Tumor-03	27.6	31.3	-2.7	86.66
KRAS-35GA	Normal-1	23.8	40	16.2	0.00
	Tumor-04	30.9	31.7	0	50.00
BRAF-1799TA	Normal-1	24.5	40	15.5	0.00
	Tumor-05	24.5	25	1.4	27.48
NRAS-183AT	Normal-1	24.2	40	15.8	0.00
	Tumor-06	29.8	40.0	5.9	1.65
TP53-733GA	Normal-1	21.8	40	18.2	0.00
	Tumor-07	26.2	26.1	-1.2	30.33

Fig. 10

List of Primers and Probes used in cast-PCR Assays

SNP ID	Tailed Allele-specific Primer 1 (ASP1)	Sequence of ASP1	Tailed Allele-specific Primer 2 (ASP2)	Sequence of ASP2
CV11201742	ASP1	cccGCTGCTCCTGTGC	ASP2	cccGGCTGCTCCTGTGT
CV11349123	ASP1	acctAACGCACAACCTTAGCG	ASP2	gccGAACGCACAACCTTAGCA
CV1207700	ASP1	cccgCCTTGATGGATGGAGATTAC	ASP2	cccgCCTTGATGGATGGAGATTAT
CV25594064	ASP1	gcacCCCTTGGTGAAACAATTTGTTG	ASP2	tgcaCCCTTGGTGAAACAATTTGTTTC
CV25639181	ASP1	cgccGAACTGGAAGTATTTTGACAG	ASP2	cgccGAACTGGAAGTATTTTGACAT
SNP ID	Locus-specific Primer (LSP)	Sequence of LSP	Locus-specific TaqMan Probe (LST)	Sequence of LST
CV11201742	LSP	TATCCCTCACAGAGAGAAGGGAG	LST	(6-FAM)ATGGCCCTGATAAGGGAGA(MGB)
CV11349123	LSP	GCITGCAATGGCTCCAACC	LST	(6-FAM)AGGGCGGTGCTCGAG(MGB)
CV1207700	LSP	GGAGCCATTGCAAGCCAAG	LST	(6-FAM)TGTCACCTCAACCACTAAA(MGB)
CV25594064	LSP	TGGCAAGAGCGATGAGTACTC	LST	(6-FAM)AGAATCTGCCAATTACCTAGA(MGB)
CV25639181	LSP	CTCTCAAAGGTAAGTCCCCTTA	LST	(6-FAM)ACAAGTAGCTGATAATAAATG(MGB)
SNP ID	MGB Blocker 1 (MGB1)	Sequence of MGB1	MGB Blocker 2 (MGB2)	Sequence of MGB2
CV11201742	MGB1	CTCCTGTGCTGTCACC(MGB)	MGB2	GCTCCTGTGTTGTCACC(MGB)
CV11349123	MGB1	CTTAGCGGCACGCAC(MGB)	MGB2	CTTAGCAGCACGCACC(MGB)
CV1207700	MGB1	GGAGATTTACGCAATGTG(MGB)	MGB2	TGGAGATTTATGCAATGTG(MGB)
CV25594064	MGB1	ACAATTGTTGAGGGGGG(MGB)	MGB2	ACAATTGTTGAGGGGGG(MGB)
CV25639181	MGB1	AAGTATTTGACAGCTTTAC(MGB)	MGB2	AAGTATTTGACATCTTTAC(MGB)

Fig. 11A

SNP ID	Tailed Allele-specific Primer 1 (ASP1)	Sequence of ASP1 with higher T _m	Tailed Allele-specific Primer 2 (ASP2)	Sequence of ASP2 with higher T _m
BRAF-1799TA	ASP1	gccTGATTTTGGTCTAGCTACAGA	ASP2	cgcGATTTTGGTCTAGCTACAGT
CTNNB1-121AG	ASP1	cgcGAGAAGGAGCTGTGGT	ASP2	gccAGAAGGAGCTGTGGC
CTNNB1-134CT	ASP1	cgcTGCCTTTACCACTCAGAG	ASP2	cgcTGCCTTTACCACTCAGAA
EGFR-2369CT	ASP1	ggcCGTGCAGCTCATCAC	ASP2	ggcCGTGCAGCTCATCA1
EGFR-2373TG	ASP1	cggCAGCAGTTTGGCCC	ASP2	cggCAGCAGTTTGGCCA
KRAS-176CG	ASP1	gccGGATATTCTCGACACAGC	ASP2	cgcTGGATAATTCTCGACACAGG
KRAS-183AC	ASP1	ggcGACACAGCAGGTCAA	ASP2	ggcGACACAGCAGGTCAAC
KRAS-183AT	ASP1	ggcGACACAGCAGGTCAA	ASP2	ggcGACACAGCAGGTCAAT
KRAS-34GA	ASP1	cgcCTTGCCTACGCCACT	ASP2	ggcTTGCCTACGCCACC
KRAS-34GC	ASP1	cgcTGCCTACGCCACG	ASP2	ggcTTGCCTACGCCACC
KRAS-34GT	ASP1	ggcTTGCCTACGCCACC	ASP2	ggcTTGCCTACGCCACA
KRAS-35GA	ASP1	cgcCTCTTGCTACGCCAT	ASP2	ggcTCTTGCTACGCCAC
KRAS-35GC	ASP1	ggcTCTTGCTACGCCAG	ASP2	ggcTCTTGCTACGCCAC
KRAS-35GT	ASP1	ggcTCTTGCTACGCCAC	ASP2	ggcTCTTGCTACGCCAA
KRAS-38GA	ASP1	cgcGTAGTTGGAGCTGGTGA	ASP2	ggcTAGTTGGAGCTGGTGG
NRAS-181CA	ASP1	cgcATACTGGATACAGCTGGAA	ASP2	cgcATACTGGATACAGCTGGAC
NRAS-183AT	ASP1	cgcTGGATACAGCTGGACAA	ASP2	cgcCTGGATACAGCTGGACA1
NRAS-35GA	ASP1	cgcGGTGGTTGGAGCAGA	ASP2	cgcGTGGTTGGAGCAGG
NRAS-38GA	ASP1	cgcGTTGGAGCAGGTGA	ASP2	cgcGTTGGAGCAGGTGG
TP53-524GA	ASP1	cgcGGAGGTTGTGAGCCA	ASP2	ggcGAGGTTGTGAGGCG
TP53-637CT	ASP1	ggcGGATGACAGAAACACTTTTC	ASP2	ggcTGGATGACAGAAACACTTTTT
TP53-721TG	ASP1	ggcTACAACTACATGTGTAACAGTG	ASP2	ggcCTACAACTACATGTGTAACAGTT
TP53-733GA	ASP1	ggcTTCCTGCATGGGCA	ASP2	ggcTTCCTGCATGGGCG
TP53-742CT	ASP1	ggcGGCGGCATGAACC	ASP2	ggcGGCGGCATGAACCT
TP53-743GA	ASP1	ggcGCGGCATGAACCA	ASP2	ggcGCGGCATGAACCG
TP53-817CT	ASP1	ggcGCCAACAGCTTTGAGGTGC	ASP2	cgcGAACAGCTTTGAGGTGT

Fig. 11B

SNP ID	Tailed Allele-specific Primer 1 (ASP1)	Sequence of ASP1 with lower Tm	Tailed Allele-specific Primer 2 (ASP2)	Sequence of ASP2 with lower Tm
BRAF-1799TA	ASP1	cccgTGGTCTAGCTACAGA	ASP2	cccgTGGTCTAGCTACAGT
CTNNB1-121AG	ASP1	ccccAAGGAGCTGTGGT	ASP2	ccccAGGAGCTGTGGC
CTNNB1-134CT	ASP1	cggcCCTTTACCACTCAGAG	ASP2	cgccCCTTTACCACTCAGAA
EGFR-2369CT	ASP1	ggggaTGCAGCTCATCAC	ASP2	ggggaTGCAGCTCAATCAAT
EGFR-2573TG	ASP1	ggcGCAGTTTGGCCC	ASP2	ggcGCAGTTTGGCCA
KRAS-176CG	ASP1	ccccATTCTCGACACAGC	ASP2	ccccATTCTCGACACAGG
KRAS-183AC	ASP1	ggccCACAGCAGGTCAA	ASP2	ggccCACAGCAGGTCAAC
KRAS-183AT	ASP1	ggccCACAGCAGGTCAA	ASP2	ggccCACAGCAGGTCAAT
KRAS-34GA	ASP1	ggcGCCTACGCCACT	ASP2	ggcGCCTACGCCACC
KRAS-34GC	ASP1	ggccCCTACGCCACG	ASP2	ggccCCTACGCCACC
KRAS-34GT	ASP1	ggccCCTACGCCACC	ASP2	ggcGCCTACGCCACA
KRAS-35GA	ASP1	ggcTGCCTACGCCAT	ASP2	ggcTGCCTACGCCAC
KRAS-35GC	ASP1	ggcTGCCTACGCCAG	ASP2	ggcTGCCTACGCCAC
KRAS-35GT	ASP1	ggcTGCCTACGCCAC	ASP2	ggcTGCCTACGCCAA
KRAS-38GA	ASP1	ccccGTTGGAGCTGGTGA	ASP2	ccccGTTGGAGCTGGTGG
NRAS-181CA	ASP1	cggaTGGATACAGCTGGAA	ASP2	cggaGGATACAGCTGGAC
NRAS-183AT	ASP1	gggcGATACAGCTGGACAA	ASP2	gggcGATACAGCTGGACAT
NRAS-35GA	ASP1	ggccTGGTTGGAGCAGA	ASP2	ggccGGTTGGAGCAGG
NRAS-38GA	ASP1	cccGTGGAGCAGGTGA	ASP2	cccTTGGAGCAGGTGG
TP53-524GA	ASP1	ggcAGGTTGTGAGGCA	ASP2	ggcAGGTTGTGAGGGC
TP53-637CT	ASP1	ggcccTGACAGAAACACTTTTC	ASP2	ggcccTGACAGAAACACTTTTT
TP53-721TG	ASP1	ggccCTACATGTGTAAACACTG	ASP2	ggccCTACATGTGTAAACACTT
TP53-733GA	ASP1	cccCCTGCATGGGCA	ASP2	cccCTGCATGGGGC
TP53-742CT	ASP1	cccGGGCATGAACCC	ASP2	cccGGGCATGAACCT
TP53-743GA	ASP1	cccGGGCATGAACCA	ASP2	cccGGGCATGAACCG
TP53-817CT	ASP1	ggcCAGCTTTGAGGTGC	ASP2	ggcCAGCTTTGAGGTGT

Fig. 11C

SNP ID	Locus-specific Primer (LSP)	Sequence of LSP	Locus-specific TaqMan Probe (LST)	Sequence of LST
BRAF-1799TA	LSP	AGCCTCAATCTTACCATCCACAAA	LST	(6-FAM)CAGACAACTGTTCAAACTG(MGB)
CTNNB1-121AG	LSP	CATGGAAACCAGACAGAAAAGCG	LST	(6-FAM)CTGGCAGCAACAAGTCT(MGB)
CTNNB1-134CT	LSP	GTCACCTGGCAGCAACAGTCTTA	LST	(6-FAM)TAGTGGCACCAGAAATGG(MGB)
EGFR-2369CT	LSP	TGGGAGCCAATATTGTCTTTGTGTT	LST	(6-FAM)CCCTTCGGCTGCCTCC(MGB)
EGFR-2573TG	LSP	AGGAACGTACTGGTGAAAACACC	LST	(6-FAM)CAGCATGTCAAGATCAC(MGB)
KRAS-176CG	LSP	CCCTCCCCAGTCCTCATGTA	LST	(6-FAM)CTGGTCCCTCATTGCAC(MGB)
KRAS-183AC	LSP	TCCTCAAATGATTAGTATTATTTATGGCAAAATACACAAAG	LST	(6-FAM)CCCTCCCCAGTCTCA(MGB)
KRAS-183AT	LSP	TCTTCAAATGATTAGTATTATTTATGGCAAAATACACAAAG	LST	(6-FAM)CCCTCCCCAGTCTCA(MGB)
KRAS-34GA	LSP	TGTGTGACATGTTCTAATATAGTCACAT	LST	(6-FAM)CCTGCTGAAAATGAC(MGB)
KRAS-34GC	LSP	TGTGTGACATGTTCTAATATAGTCACAT	LST	(6-FAM)CCTGCTGAAAATGAC(MGB)
KRAS-34GT	LSP	TGTGTGACATGTTCTAATATAGTCACAT	LST	(6-FAM)CCTGCTGAAAATGAC(MGB)
KRAS-35GA	LSP	TGTGTGACATGTTCTAATATAGTCACAT	LST	(6-FAM)CCTGCTGAAAATGAC(MGB)
KRAS-35GC	LSP	TGTGTGACATGTTCTAATATAGTCACAT	LST	(6-FAM)CCTGCTGAAAATGAC(MGB)
KRAS-35GI	LSP	TGTGTGACATGTTCTAATATAGTCACAT	LST	(6-FAM)CCTGCTGAAAATGAC(MGB)
KRAS-38GA	LSP	GGTCCCTGCACCAATATATGCA	LST	(6-FAM)TCGTCCCAAAATGATTG(MGB)
NRAS-181CA	LSP	GCAAATGACTTGCTATTATTGATGGCAAA	LST	(6-FAM)CCTTCGCTGTCTCATG(MGB)
NRAS-183AT	LSP	GCAAATGACTTGCTATTATTGATGGCAAA	LST	(6-FAM)CCTTCGCTGTCTCATG(MGB)
NRAS-35GA	LSP	AGTGGTCTCGGATTAGCTGGATTG	LST	(6-FAM)CAGTGGCTTTTCC(MGB)
NRAS-38GA	LSP	GTGGTCTGGATTAGCTGGATTGT	LST	(6-FAM)CAGTGGCTTTTCC(MGB)
TP53-524GA	LSP	GCAACCAAGCCCTGTCGTC	LST	(6-FAM)CTGCTCACCACGCTATC(MGB)
TP53-637CT	LSP	AGACCCCAAGTCCAAACCAG	LST	(6-FAM)CCTCAGGCGGCTCATAG(MGB)
TP53-721TG	LSP	CTGGAGTCTTCAGTGATGATG	LST	(6-FAM)ATGGGCTCCGGTTTCA(MGB)
TP53-733GA	LSP	CTGGAGTCTTCAGTGATGATG	LST	(6-FAM)ACCGGAGGCCATCCT(MGB)
TP53-742CT	LSP	TGTGCAGGGTGGCAAGT	LST	(6-FAM)CACACTGGAAGACTCC(MGB)
TP53-743GA	LSP	TGTGCAGGGTGGCAAGT	LST	(6-FAM)CACACTGGAAGACTCC(MGB)
TP53-817CT	LSP	CCTTCTGCGGAGATTCTCTTCT	LST	(6-FAM)CTGTGCGCGGCTCTC(MGB)

Fig. 11D

SNP ID	MGB Blocker 1 (MGB1)	Sequence of MGB1	MGB Blocker 2 (MGB2)	Sequence of MGB2
BRAF-1799TA	MGB1	GCTACAGAGAAATCTC(MGB)	MGB2	GCTACAGTGAATCTC(MGB)
CTNNB1-121AG	MGB1	GAGCTGTGGTAATTGGCA(MGB)	MGB2	AGCTGTGGCAGTGGCA(MGB)
CTNNB1-134CT	MGB1	CACTCAGAGAAAGGAGC(MGB)	MGB2	CACTCAGAAAAGGAGC(MGB)
EGFR-2369CT	MGB1	CATCAGCAGCTCATG(MGB)	MGB2	TCATCATGCAGCTCATG(MGB)
EGFR-2573TG	MGB1	AGTTTGGCCCGCCCAA(MGB)	MGB2	AGTTTGGCCAGCCCAA(MGB)
KRAS-176CG	MGB1	CTCGACACAGCAGGTCA(MGB)	MGB2	CTCGACACAGGAGGTCA(MGB)
KRAS-183AC	MGB1	GCAGGTCAAGAGGAGTAC(MGB)	MGB2	CAGGTCAAGAGGAGTAC(MGB)
KRAS-183AT	MGB1	GCAGGTCAAGAGGAGTAC(MGB)	MGB2	GCAGGTCAAGAGGAGTAC(MGB)
KRAS-34GA	MGB1	CGCCACTAGCTCCA(MGB)	MGB2	CGCCACCAGCTCC(MGB)
KRAS-34GC	MGB1	CCACGAGCTCCAACTA(MGB)	MGB2	CCACCAAGCTCCAACTA(MGB)
KRAS-34GT	MGB1	ACGCCACCAAGCTCC(MGB)	MGB2	ACGCCACAAGCTCCA(MGB)
KRAS-35GA	MGB1	CGCCACTAGCTCC(MGB)	MGB2	CGCCACCAGCTCC(MGB)
KRAS-35GC	MGB1	CGCCAGCAGCTCC(MGB)	MGB2	CGCCACCAGCTCC(MGB)
KRAS-35GT	MGB1	CGCCACCAGCTCCA(MGB)	MGB2	CGCCAACAGCTCCA(MGB)
KRAS-38GA	MGB1	GCTGGTGACGTAGGC(MGB)	MGB2	GCTGGTGCGGTAGGC(MGB)
KRAS-38CA	MGB1	GCTGGAAAAGAGAGTAC(MGB)	MGB2	GCTGGACAAGAGAGTAC(MGB)
KRAS-38AT	MGB1	CAGCTGGACAAGAGAG(MGB)	MGB2	CAGCTGGACATGAAGAG(MGB)
KRAS-35GA	MGB1	TTGGAGCAGATGGTGT(MGB)	MGB2	TGGAGCAGGTGGTGT(MGB)
KRAS-38GA	MGB1	TGGAGCAGGTGATGT(MGB)	MGB2	TGGAGCAGGTGGTGT(MGB)
TP53-524GA	MGB1	TGAGGCACTGCC(MGB)	MGB2	TGAGGCGCTGCC(MGB)
TP53-637CJ	MGB1	GAAACACTTTTCGACATAGTGTGTG(MGB)	MGB2	AGAAACACTTTTGAACATAGTGTGTG(MGB)
TP53-721TG	MGB1	TGTGTAACAGTGCTGCATG(MGB)	MGB2	ATGTGTAACAGTTCCTGCATG(MGB)
TP53-733GA	MGB1	GCATGGGCAGCATG(MGB)	MGB2	GCATGGGCGGCATG(MGB)
TP53-742CT	MGB1	GCATGAACCGAGGCC(MGB)	MGB2	GCATGAACGGAGGCC(MGB)
TP53-743GA	MGB1	CATGAACCGAGGCC(MGB)	MGB2	CATGAACCGAGGCC(MGB)
TP53-817CT	MGB1	TTGAGGTGCTGTTTGTG(MGB)	MGB2	TTTGAGGTGTGTTTGTG(MGB)

Fig. 11E

Pre-amplification of cast-PCR Reactions

	No Preamp			Preamp			Improvement
SNP ID	Ct-A1	Ct-A2	Δ Ct	Ct-A1	Ct-A2	Δ Ct	$\Delta\Delta$ Ct
KRAS-G12A_GC	31.9	45.0	13.1	25.9	45.0	19.1	6.0
KRAS-G12R_GC	29.8	45.0	15.2	24.2	45.0	20.8	5.6
KRAS-G12D_GA	32.1	45.0	12.9	26.2	45.0	18.8	5.9
KRAS-G12S_GA	31.7	45.0	13.3	25.6	39.2	13.6	0.3
KRAS-G13D_GA	32.7	45.0	12.3	26.1	45.0	18.9	6.6
KRAS-G12C_GT	36.5	45.0	8.5	32.7	45.0	12.3	3.8
KRAS-G12V_GT	36.5	45.0	8.5	29.4	45.0	15.6	7.1
Average	33.0	45.0	12.0	27.2	44.2	17.0	5.0

Fig. 12

Effect of Tailed-ASP on cast-PCR

	Non-tailed ASP (ASP -tail)			Tailed ASP (ASP +tail)			Improvement
SNP ID	ΔCt 10 ⁶ A1/A1	ΔCt 10 ⁶ A2/A2	Ave ΔCt	ΔCt 10 ⁶ A1/A1	ΔCt 10 ⁶ A2/A2	Ave ΔCt	$\Delta \Delta Ct$
BRAF-1799TA	3.4	10.0	6.7	13.4	18.8	16.1	9.41
CTNNB1-121AG	10.6	6.8	8.7	10.1	16.2	13.2	4.46
CTNNB1-134CT	8.4	3.2	5.8	15.7	10.2	13.0	7.15
EGFR-2369CT	2.6	4.3	3.5	10.5	9.1	9.8	6.32
EGFR-2573TG	16.9	10.2	13.5	21.0	19.6	20.3	6.72
KRAS-176CG	7.2	17.9	12.5	23.0	20.8	21.9	9.37
KRAS-183AC	10.4	19.8	15.1	22.4	23.1	22.7	7.60
KRAS-34GA	11.0	9.5	10.2	11.0	16.4	13.7	3.42
KRAS-35GA	10.3	10.2	10.2	11.8	17.4	14.6	4.42
KRAS-38GA	5.6	9.1	7.3	10.8	13.8	12.3	4.92
NRAS-181CA	15.0	24.3	19.6	14.0	25.6	19.8	0.13
NRAS-183AT	9.5	11.2	10.3	16.2	20.6	18.4	8.06
Average			10.3			16.3	6.0

Fig. 13

Comparison of ASB-PCR and cast-PCR Assays

	ASB-PCR			cast-PCR			Improvement
SNP ID	ΔCt 10 ⁶ A1/A1	ΔCt 10 ⁶ A2/A2	Ave ΔCt	ΔCt 10 ⁶ A1/A1	ΔCt 10 ⁶ A2/A2	Ave ΔCt	$\Delta\Delta Ct$
BRAF-1799TA	10.6	15.7	13.2	13.4	18.8	16.1	2.93
CTNNB1-121AG	12.7	10.6	11.7	10.1	16.2	13.2	1.50
CTNNB1-134CT	13.2	7.9	10.6	15.7	10.2	13.0	2.38
EGFR-2369CT	7.6	10.0	8.8	10.5	9.1	9.8	1.00
EGFR-2573TG	17.8	17.1	17.5	21.0	19.6	20.3	2.81
KRAS-176CG	10.0	20.9	15.5	23.0	20.8	21.9	6.43
KRAS-183AC	19.7	21.5	20.6	22.4	23.1	22.7	2.15
KRAS-34GA	12.1	11.0	11.5	11.0	16.4	13.7	2.14
KRAS-35GA	12.7	13.7	13.2	11.8	17.4	14.6	1.42
KRAS-38GA	7.9	13.2	10.5	10.8	13.8	12.3	1.71
NRAS-181CA	15.1	24.6	19.9	14.0	25.6	19.8	0.09
NRAS-183AT	15.3	18.2	16.8	16.2	20.6	18.4	1.63
Average			14.1			16.3	2.2

Fig. 14

Comparison of MGB and Phosphate Blocker Probes in cast-PCR

SNP ID	Phosphate blocker probe (PHOS1 and PHOS2)			MGB blocker probe (MGB1 and MGB2)			Improvement
	ΔCt 10 ⁶ A1/A1	ΔCt 10 ⁶ A2/A2	Ave ΔCt	ΔCt 10 ⁶ A1/A1	ΔCt 10 ⁶ A2/A2	Ave ΔCt	
BRAF-1799TA	10.1	14.1	12.12	10.2	14.1	12.10	-0.02
CTNNB1-121AG	13.4	11.4	12.39	15.1	11.1	13.07	0.67
CTNNB1-134CT	13.3	8.3	10.80	14.0	12.2	13.13	2.34
EGFR-2369CT	9.7	11.3	10.50	11.3	9.9	10.59	0.09
EGFR-2573TG	18.4	23.1	20.71	18.9	21.7	20.29	-0.42
KRAS-176CG	12.0	17.7	14.86	17.9	22.0	19.94	5.08
KRAS-183AC	26.6	22.1	24.39	30.2	20.7	25.44	1.06
KRAS-34GA	15.2	10.2	12.73	14.0	13.0	13.48	0.75
KRAS-35GA	15.4	11.8	13.58	15.4	12.2	13.83	0.25
KRAS-38GA	12.4	11.7	12.05	11.7	11.9	11.82	-0.23
NRAS-181CA	17.1	22.1	19.59	16.2	24.4	20.30	0.71
NRAS-183AT	12.7	21.3	17.02	15.3	14.9	15.09	-1.93
Average			15.1			15.8	0.7

Fig. 15

Effect of LNA Modified ASPs on cast-PCR

SNP ID	cast-PCR			cast-PCR with LNA-modified ASP			Improvement
	ΔCt 10 ⁶ A1/A1	ΔCt 10 ⁶ A2/A2	Ave ΔCt	ΔCt 10 ⁶ A1/A1	ΔCt 10 ⁶ A2/A2	Ave ΔCt	$\Delta \Delta Ct$
BRAF-1799TA	13.4	18.8	16.1	21.5	25.2	23.4	7.25
CTNNB1-121AG	10.1	16.2	13.2	9.4	19.0	14.2	1.02
CTNNB1-134CT	15.7	10.2	13.0	18.2	14.3	16.3	3.31
EGFR-2369CT	10.5	9.1	9.8	14.6	10.8	12.7	2.94
EGFR-2573TG	21.0	19.6	20.3	19.3	23.0	21.1	0.89
KRAS-176CG	23.0	20.8	21.9	26.6	21.9	24.3	2.37
KRAS-183AC	22.4	23.1	22.7	26.5	21.3	23.9	1.18
KRAS-34GA	11.0	16.4	13.7	8.9	17.2	13.1	-0.59
KRAS-35GA	11.8	17.4	14.6	9.9	20.7	15.3	0.69
KRAS-38GA	10.8	13.8	12.3	15.9	12.5	14.2	1.91
NRAS-181CA	14.0	25.6	19.8	13.8	29.6	21.7	1.89
NRAS-183AT	16.2	20.6	18.4	18.0	24.7	21.3	2.96
Average			16.3			18.5	2.2

Fig. 16

Effect of Other Modified ASPs on cast-PCR

	cast-PCR no modified ASP				cast-PCR with modified ASP*			Improvement
SNP ID	ΔCt 10 ⁶ A1/A1	ΔCt 10 ⁶ A2/A2	Ave ΔCt	*Modified NT at 3' of ASP	ΔCt 10 ⁶ A1/A1	ΔCt 10 ⁶ A2/A2	Ave ΔCt	$\Delta \Delta Ct$
BRAF-1799TA	13.2	19.5	16.4	(ppA)	14.1	23.2	18.6	2.2
CTNNB1-121AG	9.6	16.5	13.0	(fdU)	11.8	8.3	10.0	-3.0
CTNNB1-134CT	15.2	10.5	12.8	(ppG)	15.6	14.9	15.3	2.4
EGFR-2369CT	8.9	9.1	9.0	(iso dC)	7.8	5.4	6.6	-2.4
EGFR-2573TG	19.3	24.1	21.7	(iso dC)	19.2	27.2	23.2	1.5
KRAS-176CG	21.2	21.1	21.1	(iso dC)	25.5	16.5	21.0	-0.1
Average			15.7				15.8	0.1

Fig. 17

List of Primers and Probes used in Pre-amplification cast-PCR Assays

SNP ID	Allele-1-specific Primer	Sequence of ASP1	Allele-2-specific Primer	Sequence of ASP2
KRAS-G12A_GC	ASP1	CCCTGGTAGTTGGAGCTGG	ASP2	CCCTGGTAGTTGGAGCTGC
KRAS-G12R_GC	ASP1	CGCTGTGGTAGTTGGAGCTG	ASP2	CGCTGTGGTAGTTGGAGCTC
KRAS-G12D_GA	ASP1	GCCCGGTAGTTGGAGCTGG	ASP2	GCCCGGTAGTTGGAGCTGA
KRAS-G12S_GA	ASP1	CGCCGTGGTAGTTGGAGCTG	ASP2	CGCCGTGGTAGTTGGAGCTA
KRAS-G13D_GA	ASP1	CCCCAGTTGGAGCTGGTGG	ASP2	CCCCAGTTGGAGCTGGTGA
KRAS-G12C_GT	ASP1	GCCTTGCCTACGCCACA	ASP2	GCCTTGCCTACGCCACC
KRAS-G12V_GT	ASP1	GCCTCTTGCCTACGCCAA	ASP2	GCCTCTTGCCTACGCCAC
SNP ID	Locus-specific Primer	Sequence of LSP	Locus-specific TaqMan Probe	Sequence of LST
KRAS-G12A_GC	LSP	AAAGAATGGTCCTGCACCAAGTAA	LST	(6-FAM)ATGCATATTAAACAAGATTAC(MGB)
KRAS-G12R_GC	LSP	GGTCCTGCACCAAGTAATATGCA	LST	(6-FAM)AACAATAGAGGTAAATCTTG(MGB)
KRAS-G12D_GA	LSP	AAAGAATGGTCCTGCACCAAGTAA	LST	(6-FAM)ATGCATATTAAACAAGATTAC(MGB)
KRAS-G12S_GA	LSP	AAAGAATGGTCCTGCACCAAGTAA	LST	(6-FAM)ATGCATATTAAACAAGATTAC(MGB)
KRAS-G13D_GA	LSP	AAAGAATGGTCCTGCACCAAGTAA	LST	(6-FAM)ATGCATATTAAACAAGATTAC(MGB)
KRAS-G12C_GT	LSP	GAGTTTGTATTAAAGGTACTGGTGGAGT	LST	(6-FAM)ACCTTATGTGTGACATGTTC(MGB)
KRAS-G12V_GT	LSP	GAGTTTGTATTAAAGGTACTGGTGGAGT	LST	(6-FAM)ACCTTATGTGTGACATGTTC(MGB)

Fig. 18A

**List of Primers and Probes used in Pre-amplification cast-PCR Assays
(Continued)**

SNP ID	Allele-1 MGB Blocker	Sequence of MGB1	Allele-2 MGB Blocker	Sequence of MGB2
KRAS-G12A_GC	MGB1	TGGAGCTGCTGGCGTA(MGB)	MGB2	TGGAGCTGGTGGCGTA(MGB)
KRAS-G12R_GC	MGB1	TGGAGCTCGTGGCGTA(MGB)	MGB2	TGGAGCTGGTGGCGTA(MGB)
KRAS-G12D_GA	MGB1	TTGGAGCTGATGGCGTA(MGB)	MGB2	TGGAGCTGGTGGCGTA(MGB)
KRAS-G12S_GA	MGB1	TAGTTGGAGCTAGTGGCGTA(MGB)	MGB2	TGGAGCTGGTGGCGTA(MGB)
KRAS-G13D_GA	MGB1	TTGGAGCTGGTGACGTA(MGB)	MGB2	TGGAGCTGGTGGCGTA(MGB)
KRAS-G12C_GT	MGB1	CGCCACCAGCTCCA(MGB)	MGB2	CGCCACAAGCTCCA(MGB)
KRAS-G12V_GT	MGB1	CTACGCCACCAGCTC(MGB)	MGB2	TACGCCAACAGCTC(MGB)

Fig. 18B

List of Primers and Probes used in cast-PCR Assays

SNP ID	Untailed Allele-1-specific Primer	Sequence of ASP1 without tails	Non-tailed Allele-2-specific Primer	Sequence of ASP2 without tails
BRAF-1799TA	ASP1 -tail	TGATTTGGTCTAGCTACAGA	ASP2 -tail	GATTTTGGTCTAGCTACAGT
CTNNB1-121AG	ASP1 -tail	GAGAAGGAGCTGTGGT	ASP2 -tail	AGAAGGAGCTGTGGC
CTNNB1-134CT	ASP1 -tail	TGCCTTTACCACTCAGAG	ASP2 -tail	TGCCTTTACCACTCAGAA
EGFR-2369CT	ASP1 -tail	CGTGCAGCTCATCAC	ASP2 -tail	CGTGCAGCTCATCAT
EGFR-2573TG	ASP1 -tail	CAGCAGTTTGGCCC	ASP2 -tail	CAGCAGTTTGGCCA
KRAS-176CG	ASP1 -tail	GGATATTCTCGACACAGC	ASP2 -tail	TGGATATTCTCGACACAGG
KRAS-183AC	ASP1 -tail	GACACAGCAGGTCAA	ASP2 -tail	GACACAGCAGGTCACT
KRAS-34GA	ASP1 -tail	CTTGCCTACGCCACT	ASP2 -tail	TTGCCTACGCCACC
KRAS-35GA	ASP1 -tail	CTCTTGCCTACGCCAT	ASP2 -tail	TCTTGCCTACGCCAC
KRAS-3GA	ASP1 -tail	GTAGTTGGAGCTGGTGA	ASP2 -tail	TAGTTGGAGCTGGTGG
NRAS-181CA	ASP1 -tail	ATACTGGATACAGCTGGAA	ASP2 -tail	ATACTGGATACAGCTGGAC
NRAS-183AT	ASP1 -tail	TGGATACAGCTGGACAA	ASP2 -tail	CTGGATACAGCTGGACAT

Fig. 19A

List of Primers and Probes used in cast-PCR Assays

SNP ID	Tailed Allele-1-specific Primer	Sequence of ASP1 with tails	Tailed Allele-2-specific Primer	Sequence of ASP2 with tails
BRAF-1799TA	ASP1 +tail	gceTGATTTTGGTCTAGCTACAGA	ASP2 +tail	cegGATTTTGGTCTAGCTACAGT
CTNNB1-121AG	ASP1 +tail	cegGAGAAGGAGCTGTGGT	ASP2 +tail	gceAGAAGGAGCTGTGGC
CTNNB1-134CT	ASP1 +tail	cegTGCCTTTACCACTCAGAG	ASP2 +tail	cegTGCCTTTACCACTCAGAA
EGFR-2369CT	ASP1 +tail	gegCGTGCAGCTCATCAC	ASP2 +tail	gegCGTGCAGCTCATCAT
EGFR-2573TG	ASP1 +tail	eggCAGCAGTTTGGCCC	ASP2 +tail	eggCAGCAGTTTGGCCA
KRAS-176CG	ASP1 +tail	gceGGATATTCTCGACACAGC	ASP2 +tail	cegTGGATATTCTCGACACAGG
KRAS-183AC	ASP1 +tail	gegGACACAGCAGGTCAA	ASP2 +tail	gegGACACAGCAGGTCAAC
KRAS-34GA	ASP1 +tail	cegCTTGCCTACGCCACT	ASP2 +tail	gegTTGCCTACGCCACC
KRAS-35GA	ASP1 +tail	cegCTCTTGCCTACGCCAT	ASP2 +tail	gegTCTTGCCTACGCCAC
KRAS-38GA	ASP1 +tail	cegGTAGTTGGAGCTGGTGA	ASP2 +tail	gceTAGTTGGAGCTGGTGG
NRAS-181CA	ASP1 +tail	cegATACTGGATACAGCTGGAA	ASP2 +tail	cegATACTGGATACAGCTGGAC
NRAS-183AT	ASP1 +tail	cegTGGATACAGCTGGACAA	ASP2 +tail	cegCTGGATACAGCTGGACAT

Fig. 19B

List of Primers and Probes used in cast-PCR Assays

SNP ID	Locus-specific Primer	Sequence of LSP	Locus-specific TaqMan Probe	Sequence of LST
BRAF-1799TA	LSP	AGCCTCAATTCTTACCATCCACAAAA	LST	(6-FAM)CAGACAACCTGTTCAAACCTG(MGB)
CTNNB1-121AG	LSP	CATGGAACCAGACAGAAAAGCG	LST	(6-FAM)CTGGCAGCAACAGTCT(MGB)
CTNNB1-134CT	LSP	GTCACTGGCAGCAACAGTCITA	LST	(6-FAM)TAGTGGCACCAGAATGG(MGB)
EGFR-2369CT	LSP	TGGGAGCCAATATTGTCTTTGTGTT	LST	(6-FAM)CCCTTCGGCTGCCTCC(MGB)
EGFR-2573TG	LSP	AGGAACGTACTGGTGAAAACACC	LST	(6-FAM)CAGCATGTCAAGATCAC(MGB)
KRAS-176CG	LSP	CCCTCCCCAGTCCTCATGTA	LST	(6-FAM)CTGGTCCCTCATTGCAC(MGB)
KRAS-183AC	LSP	TCTTCAAATGATTTAGTATTATTTATGGCAAATACACAAAG	LST	(6-FAM)CCCTCCCAGTCCTCA(MGB)
KRAS-34GA	LSP	TGTGTGACATGTTCTAATATAGTCACAT	LST	(6-FAM)CCTGCTGAAAATGAC(MGB)
KRAS-35GA	LSP	TGTGTGACATGTTCTAATATAGTCACAT	LST	(6-FAM)CCTGCTGAAAATGAC(MGB)
KRAS-38GA	LSP	GGTCCTGCACCAGTAATATGCA	LST	(6-FAM)TCGTCCACAAAATGATTC(MGB)
NRAS-181CA	LSP	GCAAATGACTTGCTATTATTGATGGCAAA	LST	(6-FAM)CCTTCGCCTGTCTCATG(MGB)
NRAS-183AT	LSP	GCAAATGACTTGCTATTATTGATGGCAAA	LST	(6-FAM)CCTTCGCCTGTCTCATG(MGB)

Fig. 19C

List of Primers and Probes used in cast-PCR Assays

SNP ID	Allele-1 MGB Blocker	Sequence of MGB1	Allele-2 MGB Blocker	Sequence of MGB2
BRAF-1799TA	MGB1	CTACAGAGAAATCT(MGB)	MGB2	CTACAGTGAAATCT(MGB)
CTNNB1-121AG	MGB1	CTGTGGTAGTGGCA(MGB)	MGB2	TGTGGCAGTGGCA(MGB)
CTNNB1-134CT	MGB1	CTCAGAGAAGGAGC(MGB)	MGB2	CTCAGAAAAGGAGC(MGB)
EGFR-2369CT	MGB1	ATCACGCAGCTCA(MGB)	MGB2	CATCATGCAGCTCA(MGB)
EGFR-2573TG	MGB1	GGCCCGCCCAA(MGB)	MGB2	TGGCCAGCCCAA(MGB)
KRAS-176CG	MGB1	GACACAGCAGGTCA(MGB)	MGB2	GACACAGGAGGTCA(MGB)
KRAS-183AC	MGB1	CAGGTCAAGAGGAGTA(MGB)	MGB2	AGGTCAAGAGGAGTA(MGB)
KRAS-34GA	MGB1	GCCACTAGCTCCA(MGB)	MGB2	CCACCAGCTCCA(MGB)
KRAS-35GA	MGB1	GCCATCAGCTCC(MGB)	MGB2	CCACCAGCTCC(MGB)
KRAS-38GA	MGB1	TGGTGACGTAGGC(MGB)	MGB2	TGGTGGCGTAGGC(MGB)
NRAS-181CA	MGB1	GCTGGAAAAGAAGAG(MGB)	MGB2	CTGGACAAGAAGAG(MGB)
NRAS-183AT	MGB1	CTGGACAAGAAGAG(MGB)	MGB2	CTGGACATGAAGAG(MGB)

Fig. 19D

List of Primers and Probes used in ASB-PCR Assays

SNP ID	Allele-1-Specific Primer (ASP1)	Sequence of ASP1	Allele-2-Specific Primer (ASP2)	Sequence of ASP2
BRAF-1799TA	ASP1	GTGATTTTGGTCTAGCTACAGA	ASP2	GTGATTTTGGTCTAGCTACAGT
CTNNB1-121AG	ASP1	TCAGAGAAGGAGCTGTGGT	ASP2	TCAGAGAAGGAGCTGTGGC
CTNNB1-134CT	ASP1	GATTGCCTTTACCACTCAGAG	ASP2	GATTGCCTTTACCACTCAGAA
EGFR-2369CT	ASP1	CACCGTGCAGCTCATCAC	ASP2	CACCGTGCAGCTCATCAT
EGFR-2573TG	ASP1	CACCCAGCAGTTTGGCCC	ASP2	CACCCAGCAGTTTGGCCA
KRAS-176CG	ASP1	TTGGATATTCTCGACACAGC	ASP2	CTTGGATATTCTCGACACAGG
KRAS-183AC	ASP1	CTCGACACAGCAGGTCAA	ASP2	TCTCGACACAGCAGGTCAC
KRAS-34GA	ASP1	ACTCTTGCCTACGCCACT	ASP2	ACTCTTGCCTACGCCACC
KRAS-35GA	ASP1	CACTCTTGCCTACGCCAT	ASP2	CACTCTTGCCTACGCCAC
KRAS-38GA	ASP1	TGGTAGTTGGAGCTGGTGA	ASP2	TGGTAGTTGGAGCTGGTGG
NRAS-181CA	ASP1	CATACTGGATACAGCTGGAA	ASP2	CATACTGGATACAGCTGGAC
NRAS-183AT	ASP1	ATACTGGATACAGCTGGACAA	ASP2	TACTGGATACAGCTGGACAT

Fig. 20A

List of Primers and Probes used in ASB-PCR Assays

SNP ID	Locus-specific Primer (LSP)	Sequence of LSP	Locus-specific TaqMan Probe (LST)	Sequence of LST
BRAF-1799TA	LSP	AGCCTCAATTCTTACCATCCACAAA	LST	(6-FAM)CAGACAACCTGTTCAAACCTG(MGB)
CTNNB1-121AG	LSP	CATGGGAACCAGACAGAAAAGCG	LST	(6-FAM)CTGGCAGCAACAGTCT(MGB)
CTNNB1-134CT	LSP	GTCACCTGGCAGCAACAGTCTTA	LST	(6-FAM)TAGTGGCACCAGAAATGG(MGB)
EGFR-2369CT	LSP	TGGGAGCCAATATTGTCCTTGTGTT	LST	(6-FAM)CCCTTCGGCTGCCTCC(MGB)
EGFR-2573TG	LSP	AGGAACGTACTGGTGAAAACACC	LST	(6-FAM)CAGCATGTCAAGATCAC(MGB)
KRAS-176CG	LSP	CCCTCCCAGTCCTCATGTA	LST	(6-FAM)CTGGTCCCTCATTCAC(MGB)
KRAS-183AC	LSP	TCTTCAAATGATTAGTATTATTATGGCAAATACACAAAG	LST	(6-FAM)CCCTCCCAGTCCTCA(MGB)
KRAS-34GA	LSP	TGTGTGACATGTTCTAATATAGTCACAT	LST	(6-FAM)CCTGCTGAAAATGAC(MGB)
KRAS-35GA	LSP	TGTGTGACATGTTCTAATATAGTCACAT	LST	(6-FAM)CCTGCTGAAAATGAC(MGB)
KRAS-38GA	LSP	GGTCCTGCACCAGTAATATGCA	LST	(6-FAM)TCGTCCACAAAATGATTC(MGB)
NRAS-181CA	LSP	GCAAATGACTTGCTATTATTGATGGCAA	LST	(6-FAM)CCCTTCGCCTGTCCTCATG(MGB)
NRAS-183AT	LSP	GCAAATGACTTGCTATTATTGATGGCAA	LST	(6-FAM)CCCTTCGCCTGTCCTCATG(MGB)

Fig. 20B

**List of Primers and Probes used in ASB-PCR Assays –
Phosphate Blocker Probes**

SNP ID	Allele-1 PO ₄ Blocker	Sequence of PHOS1	Allele-2 PO ₄ Blocker	Sequence of PHOS2
BRAF-1799TA	PHOS1	GCTACAGAGAAATCTCGATGG(Phos)	PHOS2	TAGCTACAGTGAAATCTCGATG(Phos)
CTNNB1-121AG	PHOS1	GAGCTGTGGTAGTGGCA(Phos)	PHOS2	GAGCTGTGGCAGTGG(Phos)
CTNNB1-134CT	PHOS1	TTACCACTCAGAGAAGGAGC(Phos)	PHOS2	TTACCACTCAGAAAAGGAGCT(Phos)
EGFR-2369CT	PHOS1	GCTCATCACGCAGCTCA(Phos)	PHOS2	CAGCTCA1CATGCAGCTCAT(Phos)
EGFR-2573TG	PHOS1	TTGGCCCGCCAAAAT(Phos)	PHOS2	GTTTGGCCAGCCAAAATC(Phos)
KRAS-176CG	PHOS1	GACACAGCAGGTCACGAG(Phos)	PHOS2	GACACAGGAGGTCAGGAG(Phos)
KRAS-183AC	PHOS1	CAGCAGGTCAAGAGGAGTAC(Phos)	PHOS2	CAGCAGGTCACGAGGAGTA(Phos)
KRAS-34GA	PHOS1	CTACGCCACTAGCTCCAAC(Phos)	PHOS2	TACGCCACCAGCTCCAA(Phos)
KRAS-35GA	PHOS1	CTACGCCATCAGCTCC(Phos)	PHOS2	TACGCCACCAGCTCCA(Phos)
KRAS-38GA	PHOS1	CTGGTGACGTAGGCAAG(Phos)	PHOS2	CTGGTGGCGTAGGCAAG(Phos)
NRAS-181CA	PHOS1	GGATACAGCTGGAAAAGAAGAG(Phos)	PHOS2	ATACAGCTGGACAAGAAGAGT(Phos)
NRAS-183AT	PHOS1	AGCTGGACAAGAAGAGTACA(Phos)	PHOS2	AGCTGGACATGAAGAGTACA(Phos)

Fig. 20C

List of Primers and Probes for cast-PCR Assays using LNA-modified ASPs

SNP ID	Allele-1-specific Primer	Sequence of ASP1	Allele-2-specific Primer	Sequence of ASP2
BRAF-1799TA	ASP1	CCCCTTTGGTCTAGCTACAG(+A)	ASP2	CCCCTTTGGTCTAGCTACAG(+T)
CTNNB1-121AG	ASP1	GCCCGAAGGAGCTGTGG(+T)	ASP2	GCCCGAAGGAGCTGTGG(+C)
CTNNB1-134CT	ASP1	CGCCCCTTTACACTCAGA(+G)	ASP2	CGCCCCTTTACACTCAGA(+A)
EGFR-2369CT	ASP1	GCGGGTGCAGCTCATCA(+C)	ASP2	GCGGGTGCAGCTCATCA(+T)
EGFR-2573TG	ASP1	CGGGAGCAGTTTGGCC(+C)	ASP2	CGGGAGCAGTTTGGCC(+A)
KRAS-176CG	ASP1	GCCCGATATTCTCGACACAG(+C)	ASP2	GCCCGATATTCTCGACACAG(+G)
KRAS-183AC	ASP1	GCGGACACAGCAGGTCA(+A)	ASP2	GCGGACACAGCAGGTCA(+C)
KRAS-34GA	ASP1	CGGTGCCTACGCCAC(+T)	ASP2	CGGTGCCTACGCCAC(+C)
KRAS-35GA	ASP1	CGCCTTGCCTACGCCA(+T)	ASP2	CGCCTTGCCTACGCCA(+C)
KRAS-38GA	ASP1	GCCCAGTTGGAGCTGGTG(+A)	ASP2	GCCCAGTTGGAGCTGGTG(+G)
NRAS-181CA	ASP1	CGCCACTGGATACAGCTGGA(+A)	ASP2	CGCCACTGGATACAGCTGGA(+C)
NRAS-183AT	ASP1	CCGCGGATACAGCTGGACA(+A)	ASP2	CCGCGGATACAGCTGGACA(+T)

"+" indicates an LNA modification

Fig. 21A

List of Primers and Probes for cast-PCR Assays using LNA-modified ASPs

SNP ID	Locus-specific Primer (LSP)	Sequence of LSP	Locus-specific TaqMan Probe (LST)	Sequence of LST
BRAF-1799TA	LSP	AGCCTCAATTCTTACCATCCACAAAA	LST	(6-FAM)CAGACAACTGTTCAAAC TG(MGB)
CTNNB1-121AG	LSP	CATGGAACCAGACAGAAAAGCG	LST	(6-FAM)CTGGCAGCAACAGTCT(MGB)
CTNNB1-134CT	LSP	GTCACTGGCAGCAACAGTCTTA	LST	(6-FAM)TAGTGGCACCAGAA TGG(MGB)
EGFR-2369CT	LSP	TGGGAGCCAATAITGTCTTTGTGTT	LST	(6-FAM)CCCTTCGGCTGCCTCC(MGB)
EGFR-2573TG	LSP	AGGAACGTACTGGTGAAAACACC	LST	(6-FAM)CAGCATGTCAA GATCAC(MGB)
KRAS-176CG	LSP	CCCTCCCCAGTCCTCATGTA	LST	(6-FAM)CTGGTCCCTCATTGCAC(MGB)
KRAS-183AC	LSP	TCTTCAAAATGAITTAGTATTATTTATGGCAAATACACAAAG	LST	(6-FAM)CCCTCCCCAGTCCTCA(MGB)
KRAS-34GA	LSP	TGTGTGACATGTTCTAATATAGTCACAT	LST	(6-FAM)CCTGCTGAAAAATGAC(MGB)
KRAS-35GA	LSP	TGTGTGACATGTTCTAATATAGTCACAT	LST	(6-FAM)CCTGCTGAAAAATGAC(MGB)
KRAS-38GA	LSP	GGTCCTGCACCAGTAATATGCA	LST	(6-FAM)TCGTCCACAAAATGATTG(MGB)
NRAS-181CA	LSP	GCAAATGACTTGCTATTATTGATGGCAAA	LST	(6-FAM)CCTTCGCCTGTCTCATG(MGB)
NRAS-183AT	LSP	GCAAATGACTTGCTATTATTGATGGCAAA	LST	(6-FAM)CCTTCGCCTGTCTCATG(MGB)

Fig. 21B

List of Primers and Probes for cast-PCR Assays using LNA-modified ASPs

SNP ID	Allele-1 MGB Blocker	Sequence of MGB1	Allele-2 MGB Blocker	Sequence of MGB2
BRAF-1799TA	MGB1	CTACAGAGAAATCT(MGB)	MGB2	CTACAGTGAAATCT(MGB)
CTNNB1-121AG	MGB1	CTGTGGTAGTGGCA(MGB)	MGB2	TGTGGCAGTGGCA(MGB)
CTNNB1-134CT	MGB1	CTCAGAGAAGGAGC(MGB)	MGB2	CTCAGAAAAGGAGC(MGB)
EGFR-2369CT	MGB1	ATCACGCAGCTCA(MGB)	MGB2	CATCATGCAGCTCA(MGB)
EGFR-2573TG	MGB1	GGCCCGCCCAA(MGB)	MGB2	TGGCCAGCCCAA(MGB)
KRAS-176CG	MGB1	GACACAGCAGGTCA(MGB)	MGB2	GACACAGGAGGTCA(MGB)
KRAS-183AC	MGB1	CAGGTCAAGAGGAGTA(MGB)	MGB2	AGGTCACGAGGAGTA(MGB)
KRAS-34GA	MGB1	GCCACTAGCTCCA(MGB)	MGB2	CCACCAGCTCCA(MGB)
KRAS-35GA	MGB1	GCCATCAGCTCC(MGB)	MGB2	CCACCAGCTCC(MGB)
KRAS-38GA	MGB1	TGGTGACGTAGGC(MGB)	MGB2	TGGTGGCGTAGGC(MGB)
NRAS-181CA	MGB1	GCTGGAAAAGAAGAG(MGB)	MGB2	CTGGACAAGAAGAG(MGB)
NRAS-183AT	MGB1	CTGGACAAGAAGAG(MGB)	MGB2	CTGGACATGAAGAG(MGB)

Fig. 21C

List of Primers and Probes for cast-PCR Assays using chemically modified ASPs

SNP ID	Allele-1-specific Primer	Sequence of ASP1	Allele-2-specific Primer	Sequence of ASP2
BRAF-1799TA	ASP1	CCCCTTTTGGTCTAGCTACAG(ppA)	ASP2	CCCCTTTTGGTCTAGCTACAG(fdU)
CTNNB1-121AG	ASP1	GCCCGAAGGAGCTGTGG(fdU)	ASP2	GCCCGAAGGAGCTGTGG(iso dC)
CTNNB1-134CT	ASP1	CGCCCCCTTACCACCTCAGA(ppG)	ASP2	CGCCCCCTTACCACCTCAGA(ppA)
EGFR-2369CT	ASP1	GCGGGTGCAGCTCATCA(iso dC)	ASP2	GCGGGTGCAGCTCATCA(fdU)
EGFR-2573TG	ASP1	CGGGAGCAGTTTGGCC(iso dC)	ASP2	CGGGAGCAGTTTGGCC(ppA)
KRAS-176CG	ASP1	GCCCGATATTCTCGACACAG(iso dC)	ASP2	GCCCGATATTCTCGACACAG(ppG)
SNP ID	Locus-specific Primer (LSP)	Sequence of LSP	Locus-specific TagMan Probe (LST)	Sequence of LST
BRAF-1799TA	LSP	AGCCTCAATTCTTACCATCCACAAAA	LST	(6-FAM)CAGACAACTGTTCAAACTG(MGB)
CTNNB1-121AG	LSP	CATGGAACCAAGACAGAAAAGCG	LST	(6-FAM)CTGGCAGCAACAGTCT(MGB)
CTNNB1-134CT	LSP	GTCACCTGGCAGCAACAGTCTTA	LST	(6-FAM)TAGTGGCACCAGAAATGG(MGB)
EGFR-2369CT	LSP	TGGGAGCCAATATTGTCTTGTGTT	LST	(6-FAM)CCCTTCGGCTGCCTCC(MGB)
EGFR-2573TG	LSP	AGGAACGTACTGGTGAAAACACC	LST	(6-FAM)CAGCATGTCAAGATCAC(MGB)
KRAS-176CG	LSP	CCCTCCCAGTCTCATGTA	LST	(6-FAM)CTGTCCTCATTCGCAC(MGB)
SNP ID	Allele-1 MGB Blocker	Sequence of MGB1	Allele-2 MGB Blocker	Sequence of MGB2
BRAF-1799TA	MGB1	CTACAGAGAAATCT(MGB)	MGB2	CTACAGTGAAATCT(MGB)
CTNNB1-121AG	MGB1	CTGTGGTAGTGCA(MGB)	MGB2	TGTGGCAGTGCA(MGB)
CTNNB1-134CT	MGB1	CTCAGAGAAAGGAGC(MGB)	MGB2	CTCAGAAAAGGAGC(MGB)
EGFR-2369CT	MGB1	ATCACGCAOCTCA(MGB)	MGB2	CATCATGCAGCTCA(MGB)
EGFR-2573TG	MGB1	GGCCCGCCCAA(MGB)	MGB2	TGGCCAGCCCAA(MGB)
KRAS-176CG	MGB1	GACACAGCAGGTCA(MGB)	MGB2	GACACAGGAGGTCA(MGB)

Fig. 22

METHODS, COMPOSITIONS, AND KITS FOR DETECTING ALLELIC VARIANTS

CROSS REFERENCE TO RELATED APPLICATIONS

[0001] The present application is a continuation-in-part of U.S. application Ser. No. 12/641,321, filed Dec. 17, 2009 and claims the benefit of priority under 35 U.S.C. 119 to U.S. Provisional Application Nos. 61/258,582, filed Nov. 5, 2009; 61/253,501, filed Oct. 20, 2009; 61/251,623, filed Oct. 14, 2009; 61/186,775, filed Jun. 12, 2009; and 61/164,230, filed Mar. 27, 2009, all of which are incorporated herein by reference in their entireties.

BACKGROUND

[0002] Single nucleotide polymorphisms (SNPs) are the most common type of genetic diversity in the human genome, occurring at a frequency of about one SNP in 1,000 nucleotides or less in human genomic DNA (Kwok, P-Y, *Ann Rev Genom Hum Genet* 2001, 2: 235-258). SNPs have been implicated in genetic disorders, susceptibility to different diseases, predisposition to adverse reactions to drugs, and for use in forensic investigations. Thus, SNP (or rare mutation) detection provides great potentials in diagnosing early phase diseases, such as detecting circulating tumor cells in blood, for prenatal diagnostics, as well as for detection of disease-associated mutations in a mixed cell population.

[0003] Numerous approaches for SNP genotyping have been developed based on methods involving hybridization, ligation, or DNA polymerases (Chen, X., and Sullivan, P F, *The Pharmacoeconomics Journal* 2003, 3, 77-96.). For example, allele-specific polymerase chain reaction (AS-PCR) is a widely used strategy for detecting DNA sequence variation (Wu D Y, Ugozzoli L, Pal B K, Wallace R B., *Proc Natl Acad Sci USA* 1989; 86:2757-2760). AS-PCR, as its name implies, is a PCR-based method whereby one or both primers are designed to anneal at sites of sequence variations which allows for the ability to differentiate among different alleles of the same gene. AS-PCR exploits the fidelity of DNA polymerases, which extend primers with a mismatched 3' base at much lower efficiency, from 100 to 100,000 fold less efficient, than that with a matched 3' base (Chen, X., and Sullivan, P F, *The Pharmacoeconomics Journal* 2003; 3:77-96). The difficulty in extending mismatched primers results in diminished PCR amplification that can be readily detected.

[0004] The specificity and selectivity of AS-PCR, however, is largely dependent on the nature of exponential amplification of PCR which makes the decay of allele discriminating power rapid. Even though primers are designed to match a specific variant to selectively amplify only that variant, in actuality significant mismatched amplification often occurs. Moreover, the ability of AS-PCR to differentiate between allelic variants can be influenced by the type of mutation or the sequence surrounding the mutation or SNP (Ayyadevara S, Thaden J J, Shmookler Reis R J., *Anal Biochem* 2000; 284:11-18), the amount of allelic variants present in the sample, as well as the ratio between alternative alleles. Collectively, these factors are often responsible for the frequent appearance of false-positive results, leading many researchers to attempt to increase the reliability of AS-PCR (Orou A, Fechner B, Utermann G, Menzel H J., *Hum Mutat* 1995; 6:163-169) (Imyanitov E N, Busboy K G, Suspitsin E N, Kuligina E S, Belogubova E V, Grigoriev M Y, et al., *Biotech-*

niques 2002; 33:484-490) (McKinzie P B, Parsons B L. Detection of rare K-ras codon 12 mutations using allele-specific competitive blocker PCR. *Mutat Res* 2002; 517:209-220) (Latorra D, Campbell K, Wolter A, Hurley J M., *Hum Mutat* 2003; 22:79-85).

[0005] In some cases, the selectivity of AS-PCR has been increased anywhere from detection of 1 in 10 alleles to 1 in 100,000 alleles by using SNP-based PCR primers containing locked nucleic acids (LNAs) (Latorra, D., et al., *Hum Mut* 2003, 2:79-85; Nakiandwe, J. et al., *Plant Method* 2007, 3:2) or modified bases (Koizumi, M. et al. *Anal Biochem.* 2005, 340:287-294). However, these base "mimics" or modifications increase the overall cost of analysis and often require extensive optimization.

[0006] Another technology involving probe hybridization methods used for discriminating allelic variations is Taq-Man® genotyping. However, like AS-PCR, selectivity using this method is limited and not suitable for detecting rare (1 in $\geq 1,000$) alleles or mutations in a mixed sample.

SUMMARY

[0007] In some embodiments, the present inventions relates generally to compositions, methods and kits for use in discriminating sequence variation between different alleles. More specifically, in some embodiments, the present invention provides for compositions, methods and kits for quantitating rare (e.g., mutant) allelic variants, such as SNPs, or nucleotide (NT) insertions or deletions, in samples comprising abundant (e.g., wild type) allelic variants with high specificity. In particular, in some embodiments, the invention relates to a highly selective method for mutation detection referred to as competitive allele-specific TaqMan PCR ("cast-PCR").

[0008] In one aspect, the present invention provides compositions for use in identifying and/or quantitating allelic variants in nucleic acid samples. Some of these compositions can comprise: (a) an allele-specific primer; (b) an allele-specific blocker probe; (c) a detector probe; and/or (d) a locus-specific primer.

[0009] In some embodiments of the compositions, the allele-specific primer comprises a target-specific portion and an allele-specific nucleotide portion. In some embodiments, the allele-specific primer may further comprise a tail. In some exemplary embodiments, the tail is located at the 5' end of the allele-specific primer. In other embodiments, the tail of the allele-specific primer has repeated guanine and cytosine residues ("GC-rich"). In some embodiments, the melting temperature ("T_m") of the entire allele-specific primer ranges from about 50° C. to 67° C. In some embodiments, the allele-specific primer concentration is between about 20-900 nM.

[0010] In some embodiments of the compositions, the allele-specific nucleotide portion of the allele-specific primer is located at the 3' terminus. In some embodiments, the selection of the allele-specific nucleotide portion of the allele-specific primer involves the use of a highly discriminating base (e.g., for detection of A/A, A/G, G/A, G/G, A/C, or C/A alleles). In some embodiments, for example when the allele to be detected involves A/G or C/T SNPs, A or G is used as the 3' allele-specific nucleotide portion of the allele-specific primer (e.g., if A or T is the target allele), or C or T is used as the 3' allele-specific nucleotide portion of the allele-specific primer (e.g., if C or G is the target allele). In other embodiments, A is used as the discriminating base at the 3' end of the allele-specific primer when detecting and/or quantifying A/T

SNPs. In other embodiments, G is used as the discriminating base at the 3' end of the allele-specific primer when detecting and/or quantifying C/G SNPs.

[0011] In some embodiments of the compositions, the allele-specific blocker probe comprises a non-extendable blocker moiety at the 3' terminus. In some exemplary embodiments, the non-extendable blocker moiety is a minor groove binder (MGB). In some embodiments, the target allele position is located about 6-10, such as about 6, about 7, about 8, about 9, or about 10 nucleotides away from the non-extendable blocker moiety of the allele-specific blocker probe. In some embodiments, the allele-specific blocker probe comprises an MGB moiety at the 5' terminus. In some exemplary embodiments, the allele-specific blocker probe is not cleaved during PCR amplification. In some embodiments, the T_m of the allele-specific blocker probe ranges from about 58° C. to 66° C.

[0012] In some embodiments of the compositions, the allele-specific blocker probe and/or allele-specific primer comprise at least one modified base. In some embodiments, the modified base(s) may increase the difference in the T_m between matched and mismatched target sequences and/or decrease mismatch priming efficiency, thereby improving not only assay specificity but also selectivity. Such modified base(s) may include, for example, 8-Aza-7-deaza-dA (ppA), 8-Aza-7-deaza-dG (ppG), 2'-Deoxypseudoisocytidine (iso dC), 5-fluoro-2'-deoxyuridine (FdU), locked nucleic acid (LNA), or 2'-O,4'-C-ethylene bridged nucleic acid (ENA) bases (also see, for example, FIG. 4B). In some embodiments the modified base is located at (a) the 3'-end, (b) the 5'-end, (c) at an internal position or at any combination of (a), (b) or (c) within the allele-specific blocker probe and/or the allele-specific primer.

[0013] In some embodiments of the compositions, the detector probe is a sequence-based or locus-specific detector probe. In other embodiments the detector probe is a 5' nuclease probe. In some exemplary embodiments, the detector probe can comprise an MGB moiety, a reporter moiety (e.g., FAMTM, TETTM, JOETM, VTCTM, or SYBR[®] Green), a quencher moiety (e.g., Black Hole QuencherTM or TAMRATM), and/or a passive reference (e.g., ROXTM). In some exemplary embodiments, the detector probe is designed according to the methods and principles described in U.S. Pat. No. 6,727,356 (the disclosure of which is incorporated herein by reference in its entirety). In some exemplary embodiments, the detector probe is a TaqMan[®] probe (Applied Biosystems, Foster City).

[0014] In some embodiments of the compositions, the composition can further comprise a polymerase; deoxyribonucleotide triphosphates (dNTPs); other reagents and/or buffers suitable for amplification; and/or a template sequence or nucleic acid sample. In some embodiments, the polymerase can be a DNA polymerase. In some other embodiments, the polymerase can be thermostable, such as Tag DNA polymerase. In other embodiments, the template sequence or nucleic acid sample can be DNA, such as genomic DNA (gDNA) or complementary DNA (cDNA). In other embodiments the template sequence or nucleic acid sample can be RNA, such as messenger RNA (mRNA).

[0015] In another aspect, the present disclosure provides methods for amplifying an allele-specific sequence. Some of these methods can include one or more of the following: (a) hybridizing an allele-specific primer to a first nucleic acid molecule comprising a first allele (allele-1); (b) hybridizing

an allele-specific blocker probe to a second nucleic acid molecule comprising a second allele (allele-2), wherein allele-2 corresponds to the same loci as allele-1; (c) hybridizing a detector probe to the first nucleic acid molecule; (d) hybridizing a locus-specific primer to the extension product of the allele-specific primer; and (e) PCR amplifying the first nucleic acid molecule comprising allele-1.

[0016] In another aspect, the present invention provides methods for detecting and/or quantitating an allelic variant in a pooled or mixed sample comprising other alleles. Some of these methods can include one or more of the following: (a) in a first reaction mixture hybridizing a first allele-specific primer to a first nucleic acid molecule comprising a first allele (allele-1) and in a second reaction mixture hybridizing a second allele-specific primer to a first nucleic acid molecule comprising a second allele (allele-2), wherein allele-2 corresponds to the same locus as allele-1; (b) in the first reaction mixture hybridizing a first allele-specific blocker probe to a second nucleic acid molecule comprising allele-2 and in the second reaction mixture hybridizing a second allele-specific blocker probe to a second nucleic acid molecule comprising allele-1; (c) in the first reaction mixture, hybridizing a first detector probe to the first nucleic acid molecule and in the second reaction mixture and hybridizing a second detector probe to the first nucleic acid molecule; (d) in the first reaction mixture hybridizing a first locus-specific primer to the extension product of the first allele-specific primer and in the second reaction mixture hybridizing a second locus-specific primer to the extension product of the second allele-specific primer; and (e) PCR amplifying the first nucleic acid molecule to form a first set or sample of amplicons and PCR amplifying the second nucleic acid molecule to form a second set or sample of amplicons; and (f) comparing the first set of amplicons to the second set of amplicons to quantitate allele-1 in the sample comprising allele-2 and/or allele-2 in the sample comprising allele-1.

[0017] In some embodiments of the methods, the first and/or second allele-specific primer comprises a target-specific portion and an allele-specific nucleotide portion. In some embodiments, the first and/or second allele-specific primer may further comprise a tail. In some embodiments, the T_m of the entire first and/or second allele-specific primer ranges from about 50° C. to 67° C. In some embodiments the first and/or second allele-specific primer concentration is between about 20-900 nM.

[0018] In some embodiments of the methods, the target-specific portion of the first allele-specific primer and the target-specific portion of the second allele-specific primer comprise the same sequence. In other embodiments, the target-specific portion of the first allele-specific primer and the target-specific portion of the second allele-specific primer are the same sequence.

[0019] In some embodiments of the methods, the tail is located at the 5'-end of the first and/or second allele-specific primer. In some embodiments, the 5' tail of the first allele-specific primer and the 5' tail of the second allele-specific primer comprise the same sequence. In other embodiments, the 5' tail of the first allele-specific primer and the 5' tail of the second allele-specific primer are the same sequence. In other embodiments, the tail of the first and/or second allele-specific primer is GC-rich.

[0020] In some embodiments of the methods, the allele-specific nucleotide portion of the first allele-specific primer is specific to a first allele (allele-1) of a SNP and the allele-

specific nucleotide portion of the second allele-specific primer is specific to a second allele (allele-2) of the same SNP. In some embodiments of the methods, the allele-specific nucleotide portion of the first and/or second allele-specific primer is located at the 3'-terminus. In some embodiments, the selection of the allele-specific nucleotide portion of the first and/or second allele-specific primer involves the use of a highly discriminating base (e.g., for detection of A/A, A/G, G/A, G/G, A/C, or C/A alleles). In some embodiments, for example when the allele to be detected involves A/G or C/T SNPs, A or G is used as the 3' allele-specific nucleotide portion of the first and/or second allele-specific primer (e.g., if A or T is the major allele), or C or T is used as the 3' allele-specific nucleotide portion of the first and/or second allele-specific primer (e.g., if C or G is the major allele). In other embodiments, A is used as the discriminating base at the 3' end of the first and/or second allele-specific primer when detecting and/or quantifying A/T SNPs. In other embodiments, G is used as the discriminating base at the 3' end of the first and/or second allele-specific primer when detecting and/or quantifying C/G SNPs.

[0021] In some embodiments of the methods, the first and/or second allele-specific blocker probe comprises a non-extendable blocker moiety at the 3' terminus. In some exemplary embodiments, the non-extendable blocker moiety is an MGB. In some embodiments, the target allele position is located about 6-10, such as about 6, about 7, about 8, about 9, or about 10 nucleotides away from the non-extendable blocker moiety of the first and/or second allele-specific blocker probe. In some embodiments, the first and/or second allele-specific blocker probe comprises an MGB moiety at the 5'-terminus. In other embodiments, the first and/or second allele-specific blocker probe is not cleaved during PCR amplification. In some embodiments, the T_m of the first and/or second allele-specific blocker probe ranges from about 58° C. to 66° C.

[0022] In some embodiments of the methods, the first and/or second allele-specific blocker probe and/or the first and/or second allele-specific primer comprises at least one modified base. In some embodiments, the modified base(s) may increase the difference in the T_m between matched and mismatched target sequences and/or decrease mismatch priming efficiency, thereby improving not only assay specificity, but also selectivity. Such modified base(s) may include, for example, 8-Aza-7-deaza-dA (ppA), 8-Aza-7-deaza-dG (ppG), 2'-Deoxypseudoisocytidine (iso dC), 5-fluoro-2'-deoxyuridine (fdU), locked nucleic acid (LNA), or 2'-O,4'-C-ethylene bridged nucleic acid (ENA) bases (see also, for example, FIG. 4B). In some embodiments the modified base is located at (a) the 3'-end, (b) the 5'-end, (c) at an internal position or at any combination of (a), (b) or (c) within said first and/or second allele-specific blocker probe and/or the first and/or second allele-specific primer.

[0023] In some embodiments of the methods, the specificity of allelic discrimination is improved by the inclusion of a modified base in the first and/or second allele-specific primer and/or first, and/or second allele-specific blocker probe as compared to the use of a non-modified allelic-specific primer or blocker probe. In some embodiments, the improvement in specificity is at least 2 fold.

[0024] In some embodiments of the methods, the specificity of allelic discrimination is at least 2 fold (e.g., 2 fold, 3 fold, 4, fold, 5 fold, and so on) better than the specificity of

allelic discrimination using Allele-Specific PCR with a Blocking reagent (ASB-PCR) methods.

[0025] In some embodiments, the methods further comprise a 2-stage cycling protocol. In some embodiments, the number of cycles in the first stage of the 2-stage cycling protocol comprises fewer cycles than the number of cycles used in the second stage. In other embodiments, the number of cycles in the first stage is about 90% fewer cycles than the number of cycles in the second stage. In yet other embodiments, the number of cycles in the first stage is between 3-7 cycles and the number of cycles in the second stage is between 42-48 cycles.

[0026] In some embodiments, the annealing/extension temperature used during the first cycling stage of the 2-stage cycling protocol is between 1-3° C. lower than the annealing/extension temperature used during the second stage. In preferred embodiments, the annealing/extension temperature used during the first cycling stage of the 2-stage cycling protocol is between 56-59° C. and the annealing/extension temperature used during the second stage is between 60-62° C.

[0027] In some embodiments, the methods further comprise a pre-amplification step. In preferred embodiments, the pre-amplification step comprises a multiplex amplification reaction that uses at least two complete sets of allele-specific primers and locus-specific primers, wherein each set is suitable or operative for amplifying a specific polynucleotide of interest. In other embodiments, the products of the multiplex amplification reaction are divided into secondary single-plex amplification reactions, such as a cast-PCR reaction, wherein each single-plex reaction contains at least one primer set previously used in the multiplex reaction. In other embodiments, the multiplex amplification reaction further comprises a plurality of allele-specific blocker probes. In some embodiments, the multiplex amplification reaction is carried out for a number of cycles suitable to keep the reaction within the linear phase of amplification.

[0028] In some embodiments of the methods, the first and/or second detector probes are the same. In some embodiments, the first and/or second detector probes are different. In some embodiments, the first and/or second detector probe is a sequence-based or locus-specific detector probe. In other embodiments the first and/or second detector probe is a 5' nuclease probe. In some exemplary embodiments, the first and/or second detector probes comprises an MGB moiety, a reporter moiety (e.g., FAMTM, TETTM, JOETTM, VICTM, or SYBR® Green), a quencher moiety (e.g., Black Hole QuencherTM or TAMRATM), and/or a passive reference (e.g., ROXTM). In some exemplary embodiments, the first and/or second detector probe is designed according to the methods and principles described in U.S. Pat. No. 6,727,356 (the disclosure of which is incorporated herein by reference in its entirety). In some exemplary embodiments, the detector probe is a TaqMan® probe.

[0029] In some embodiments of the methods, the first locus-specific primer and the second locus-specific primer comprise the same sequence. In some embodiments the first locus-specific primer and the second locus-specific primer are the same sequence.

[0030] In some embodiments of the methods, the first and/or second reaction mixtures can further comprises a polymerase; dNTPs; other reagents and/or buffers suitable for PCR amplification; and/or a template sequence or nucleic acid sample. In some embodiments, the polymerase can be a

DNA polymerase. In some embodiments, the polymerase can be thermostable, such as Taq DNA polymerase. In some embodiments, the template sequence or nucleic acid sample can be DNA, such as gDNA or cDNA. In other embodiments the template sequence or nucleic acid sample can be RNA, such as mRNA.

[0031] In some embodiments of the methods, the first allele-specific blocker probe binds to the same strand or sequence as the second allele-specific primer, while the second allele-specific blocker probe binds to the same strand or sequence as the first allele-specific primer. In some embodiments, the first and/or second allele-specific blocker probes are used to reduce the amount of background signal generated from either the second allele and/or the first allele, respectively. In some embodiments, first and/or second allele-specific blocker probes are non-extendable and preferentially anneal to either the second allele or the first allele, respectively, thereby blocking the annealing of, for example, the extendable first allele-specific primer to the second allele and/or the extendable second allele-specific primer to first allele.

[0032] In some exemplary embodiments, the first allele is a rare (e.g., minor) or mutant allele. In other exemplary embodiments the second allele is an abundant (e.g., major) or wild type allele.

[0033] In another aspect, the present invention provides kits for quantitating a first allelic variant in a sample comprising a second allelic variant involving: (a) a first allele-specific primer; (b) a second allele-specific primer; (c), a first locus-specific primer; (d) a second locus-specific primer; (e) a first allele-specific blocker probe; (f) a second allele-specific blocker probe; and (g) a first locus-specific detector probe and (h) a second locus-specific detector probe.

[0034] In some embodiments of the kits, the first and/or second allele-specific primer comprises a target-specific portion and an allele-specific nucleotide portion. In some embodiments, the first and/or second allele-specific primer may further comprise a tail. In some embodiments, the T_m of the entire first and/or second allele-specific primer ranges from about 50° C. to 67° C. In some embodiments the first and/or second allele-specific primer concentrations are between about 20-900 nM.

[0035] In some embodiments of the kits, the target-specific portion of the first allele-specific primer and the target-specific portion of the second allele-specific primer comprise the same sequence. In other embodiments, the target-specific portion of the first allele-specific primer and the target-specific portion of the second allele-specific primer are the same sequence.

[0036] In some embodiments of the kits, the tail is located at the 5' end of the first and/or second allele-specific primer. In some embodiments, the 5' tail of the first allele-specific primer and the 5' tail of the second allele-specific primer comprise the same sequence. In other embodiments, the 5' tail of the first allele-specific primer and the 5' tail of the second allele-specific primer are the same sequence. In other embodiments, the tail of the first and/or second allele-specific primer is GC rich.

[0037] In some embodiments of the kits, the allele-specific nucleotide portion of the first allele-specific primer is specific to a first allele (allele-1) of a SNP and the allele-specific nucleotide portion of the second allele-specific primer is specific to a second allele (allele-2) of the same SNP. In some embodiments of the disclosed methods, the allele-specific

nucleotide portion of the first and/or second allele-specific primer is located at the 3' terminus. In some embodiments, the selection of the allele-specific nucleotide portion of the first and/or second allele-specific primer involves the use of a highly discriminating base (e.g., for detection of A/A, A/G, G/A, G/G, A/C, or C/A alleles) (FIG. 2). In some embodiments, for example when the allele to be detected involves A/G or C/T SNPs, A or G is used as the 3' allele-specific nucleotide portion of the first and/or second allele-specific primer (e.g., if A or T is the major allele), or C or T is used as the 3' allele-specific nucleotide portion of the first and/or second allele-specific primer (e.g., if C or G is the major allele). In other embodiments, A is used as the discriminating base at the 3' end of the first and/or second allele-specific primer when detecting and/or quantifying A/T SNPs. In other embodiments, G is used as the discriminating base at the 3' end of the first and/or second allele-specific primer when detecting and/or quantifying C/G SNPs.

[0038] In some embodiments of the kits, the first and/or second allele-specific blocker probe comprises a non-extendable blocker moiety at the 3' terminus. In some exemplary embodiments, the non-extendable blocker moiety is an MGB. In some embodiments, the target allele position is located about 6-10, such as about 6, about 7, about 8, about 9, or about 10 nucleotides away from the non-extendable blocker moiety of the first and/or second allele-specific blocker probe. In some embodiments, the first and/or second allele-specific blocker probe comprises an MGB moiety at the 5' terminus. In other embodiments, the first and/or second allele-specific blocker probe is not cleaved during PCR amplification. In some embodiments, the T_m of the first and/or second allele-specific blocker probe ranges from about 58° C. to 66° C.

[0039] In some embodiments of the kits, the allele-specific blocker probe and/or the first and/or second allele-specific primer comprises at least one modified base. In some embodiments, the modified base(s) may increase the difference in the T_m between matched and mismatched target sequences and/or decrease mismatch priming efficiency, thereby improving not only assay specificity but also selectivity. Such modified base(s) may include, for example, 8-Aza-7-deaza-dA (ppA), 8-Aza-7-deaza-dG (ppG), 2'-Deoxypseudoisocytidine (iso dC), 5-fluoro-2'-deoxyuridine (fdU), locked nucleic acid (LNA), or 2'-O,4'-C-ethylene bridged nucleic acid (ENA) bases (see also, for example, FIG. 4B). In some embodiments the modified base is located at (a) the 3'-end, (b) the 5'-end, (c) at an internal position or at any combination of (a), (b) or (c) within said first and/or second allele-specific blocker probe and/or the first and/or second allele-specific primer.

[0040] In some embodiments of the kits, the first and/or second detector probes are the same. In some embodiments of the disclosed kits the first and/or second detector probes are different. In some embodiments of the disclosed kits, the first and/or second detector probes are sequence-based or locus-specific detector probes. In other embodiments the first and/or second detector probe are 5' nuclease probes. In some exemplary embodiments, the first and/or second detector probes comprise an MGB moiety, a reporter moiety (e.g., FAMTM, TETTM, JOETM, VICTM, or SYBR® Green), a quencher moiety (e.g., Black Hole QuencherTM or TAMRATM), and/or a passive reference (e.g., ROXTM). In some exemplary embodiments, the first and/or second detector probe are designed according to the methods and principles described in U.S. Pat. No. 6,727,356 (the disclosure of which is incorporated herein

by reference in its entirety). In some exemplary embodiments, the detector probe is a TagMan® probe.

[0041] In some embodiments of the kits, the first locus-specific primer and the second locus-specific primer comprise the same sequence. In some embodiments the first locus-specific primer and the second locus-specific primer are the same sequence.

[0042] In some embodiments of the kits, the first and/or second reaction mixture can further comprise a polymerase; dNTPs; other reagents and/or buffers suitable for PCR amplification; and/or a template sequence or nucleic acid sample. In some embodiments, the polymerase can be a DNA polymerase. In some other embodiments, the polymerase can be thermostable, such as Tag DNA polymerase.

[0043] In some embodiments, the compositions, methods and kits of the present invention provide high allelic discrimination specificity and selectivity. In some embodiments, the quantitative determination of specificity and/or selectivity comprises a comparison of Ct values between a first set of amplicons and a second set of amplicons. In some embodiments, selectivity is at a level whereby a single copy of a given allele in about 1 million copies of another allele or alleles can be detected.

[0044] The foregoing has described various embodiments of the invention that provide improved detection and discrimination of allelic variants using one or more of the following: (a) tailed allele-specific primers; (b) low allele-specific primer concentration; (c) allele-specific primers designed to have lower Tms; (d) allele-specific primers designed to target discriminating bases; (e) allele-specific blocker probes containing MGB, designed to prevent amplification from alternative, and potentially more abundant, allelic variants in a sample; and (f) allele-specific blocker probes and/or allele-specific primers designed to comprise modified bases in order to increase the delta Tm between matched and mismatched target sequences.

[0045] While particular embodiments employing several of the above improvements have been discussed herein, it will be apparent to the skilled artisan that depending on the nature of the sample to be examined, various combinations of the above improvements can be combined to arrive at a favorable result. Thus, for example, non-MGB blocker probes can be used with an embodiment that include methods employing allele-specific primers containing modified bases to increase delta Tm; such primers can also be designed to target discriminating bases; and the primers can be used at low primer concentrations. Accordingly, alternative embodiments based upon the present disclosure can be used to achieve a suitable level of allelic detection.

[0046] The present disclosure provides the advantage that any of the combinations of listed improvements could be utilized by a skilled artisan in a particular situation. For example, the current invention can include a method or reaction mixture that employs improvements a, c, d and f; improvements b, c, and e; or improvements

[0047] It is to be understood that both the foregoing general description and the following detailed description are exemplary and explanatory only and are not restrictive of the invention, as claimed.

[0048] The accompanying drawings, which are incorporated in and constitute a part of this specification, illustrate

several exemplary embodiments of the disclosure and together with the description, serve to explain certain teachings.

BRIEF DESCRIPTION OF THE DRAWINGS

[0049] The skilled artisan will understand that the drawings described below are for illustration purposes only. The drawings are not intended to limit the scope of the present teachings in any way.

[0050] FIG. 1 depicts a schematic of an illustrative embodiment of cast-PCR. In some embodiments, components of cast-PCR include the following: one locus-specific TaqMan probe (LST); two MGB blockers: one allele-1-specific MGB blocker (MGB1) and one allele-2-specific MGB blocker (MGB2); 3 PCR primers: one locus-specific PCR primer (LSP); one allele-1-specific primer (ASP1) and one allele-2-specific primer (ASP2).

[0051] FIG. 2 depicts a schematic of an illustrative embodiment of cast-PCR using allele-specific blocker probes comprising highly discriminating bases for detecting rare allelic variants. Highly discriminating bases may include, for example, A/A, A/G, G/A, G/G, A/C, C/A. The least discriminating bases may include, for example, C/C, T/C, G/T, T/G, C/T. In some embodiments, for example, for detection of A-G or C-T SNPs, A & G are used as the discriminating base if A/T is allelic variant (e.g., mutant allele); or C & T are used as the discriminating base if C/G allelic variant (e.g., mutant allele).

[0052] FIG. 3 depicts a schematic of an illustrative embodiment of cast-PCR using an allele-specific blocker probe with an MGB moiety at the 5' end. In some embodiments the blocker moiety at the 3'-end of the probe may include, for example, NH₂, biotin, MGB, PO₄, and PEG.

[0053] FIG. 4A depicts a schematic of an illustrative embodiment of cast-PCR using modified bases in an MGB blocker probe or allele-specific primer. (G* represents ppG.)

[0054] FIG. 4B depicts some examples of modified bases of an MGB blocker probe or allele-specific primer.

[0055] FIG. 5 depicts the TaqMan-like sensitivity and dynamic range of one exemplary embodiment of cast-PCR.

[0056] FIG. 6 depicts the sequence of KRAS mutations at codons 12 and 13 that are detectable using cast-PCR methods. KRAS mutations at codons 12 and 13 are associated with resistance to cetuximab or panitumumab in metastatic colorectal cancer (Di Nicolantonio F., et al., *J Clin Oncol.* 2008; 26:5705-12.)

[0057] FIG. 7 depicts the specificity of KRAS mutation detection using cast-PCR assays in one exemplary embodiment.

[0058] FIG. 8 depicts one exemplary embodiment using cast-PCR methods to detect a single copy of mutant DNA in 10⁶ copies of wild-type DNA.

[0059] FIG. 9 depicts detection of the relative copy number of mutant samples (KRAS-G12A) spiked in wild type samples using cast-PCR methods.

[0060] FIG. 10 depicts a number of different tumor markers (SNPs) detected in tumor samples using one exemplary embodiment of cast-PCR.

[0061] FIG. 11A-E shows a list of exemplary primers and probes used in cast-PCR assays. Nucleotides shown in lower case are the tailed portion of the primers. The nucleotide-portion of allele-specific primers (ASP) is at the 3'-most terminus of each primer and are indicated in bold. The allele

positions of the blocker probes (MGB) are located at various internal positions relative to the blocker moieties, in some cases, are indicated in bold.

[0062] FIG. 12 depicts the specificity of allelic discrimination for samples that were pre-amplified prior to analysis by cast-PCR.

[0063] FIG. 13 depicts the specificity of allelic discrimination for cast-PCR assays performed using tailed versus non-tailed allele-specific primers.

[0064] FIG. 14 depicts the specificity of allelic discrimination for samples analyzed by cast-PCR versus samples analyzed by ASB-PCR methods.

[0065] FIG. 15 depicts the specificity of allelic discrimination for cast-PCR assays performed using MGB blocker probes or phosphate blocker probes.

[0066] FIG. 16 depicts the specificity of allelic discrimination for cast-PCR assays performed using LNA-modified allele-specific primers.

[0067] FIG. 17 compares the specificity of allelic discrimination for cast-PCR assays performed using various chemically-modified allele-specific primers.

[0068] FIG. 18A-B shows a list of exemplary allele-specific primers and probes used in pre-amplification and cast-PCR assays.

[0069] FIG. 19A-D shows a list of exemplary primers and probes used in cast-PCR assays using either tailed (ASP +tail) or non-tailed (ASP -tail) allele-specific primers. (The tailed portion of the ASP +tail primers are indicated in lower case lettering.)

[0070] FIG. 20A-C shows a list of exemplary primers and probes used in ASB-PCR. The blocker probes used in ASB-PCR comprise a phosphate group at the 3'-end of the blocker probes (PHOS).

[0071] FIG. 21A-C shows a list of exemplary primers and probes used in cast-PCR assays performed using LNA-modified allele-specific primers. In this exemplary embodiment, the LNA modifications of the ASP are at the 3'-ends. ("+" indicates the LNA modified nucleotide and are notated in parentheses.)

[0072] FIG. 22 shows a list of exemplary primers and probes used in cast-PCR assays performed using other chemically modified allele-specific primers. In this exemplary embodiment, the chemical modifications (e.g., ppA, ppG, fdU, and iso dC) of the ASP are at the 3'-ends. (The chemically modified nucleotides are shown in parentheses.)

DETAILED DESCRIPTION

I. Introduction

[0073] The selective amplification of an allele of interest is often complicated by factors including the mispriming and extension of a mismatched allele-specific primer on an alternative allele. Such mispriming and extension can be especially problematic in the detection of rare alleles present in a sample populated by an excess of another allelic variant. When in sufficient excess, the mispriming and extension of the other allelic variant may obscure the detection of the allele of interest. When using PCR-based methods, the discrimination of a particular allele in a sample containing alternative allelic variants relies on the selective amplification of an allele of interest, while minimizing or preventing amplification of other alleles present in the sample.

[0074] A number of factors have been identified, which alone or in combination, contribute to the enhanced discrimi-

nating power of allele-specific PCR. As disclosed herein, a factor which provides a greater ΔC_t value between a mismatched and matched allele-specific primer is indicative of greater discriminating power between allelic variants. Such factors found to improve discrimination of allelic variants using the present methods include, for example, the use of one or more of the following: (a) tailed allele-specific primers; (b) low allele-specific primer concentration; (c) allele-specific primers designed to have lower Tms; (d) allele-specific primers designed to target discriminating bases; (e) allele-specific blocker probes designed to prevent amplification from alternative, and potentially more abundant, allelic variants in a sample; and (f) allele-specific blocker probes and/or allele-specific primers designed to comprise modified bases in order to increase the delta Tm between matched and mismatched target sequences.

[0075] The above-mentioned factors, especially when used in combination, can influence the ability of allele-specific PCR to discriminate between different alleles present in a sample. Thus, the present disclosure relates generally to novel amplification methods referred to as cast-PCR, which utilizes a combination of factors referred to above to improve discrimination of allelic variants during PCR by increasing ΔC_t values. In some embodiments, the present methods can involve high levels of selectivity, wherein one mutant molecule in a background of at least 1,000 to 1,000,000, such as about 1000-10,000, about 10,000 to 100,000, about 20,000 to 250,000, about 30,000 to 300,000, about 40,000 to 400,000, about 50,000 to 500,000, about 75,000 to 750,000 or about 100,000 to 1,000,000 wild type molecules, or any fractional ranges in between can be detected. In some embodiments, the comparison of a first set of amplicons and a second set of amplicons involving the disclosed methods provides improvements in specificity from 1,000 $\times$ to 100,000,000 $\times$ fold difference, such as about 1000-10,000 $\times$, about 10,000 to 100,000 $\times$, about 20,000 to 250,000 $\times$, about 30,000 to 300,000 $\times$, about 40,000 to 400,000 $\times$, about 50,000 to 500,000 $\times$, about 75,000 to 750,000 $\times$, about 100,000 to 1,000,000 $\times$, about 2,000,000 to 25,000,000 $\times$, about 3,000,000 to 30,000,000 $\times$, about 4,000,000 $\times$ to 40,000,000 $\times$, about 5,000,000 to 50,000,000 $\times$, about 7,500,000 to 75,000,000 $\times$ or about 10,000,000 to 100,000,000 $\times$ fold difference, or any fractional ranges in between.

II. Definitions

[0076] For the purposes of interpreting this specification, the following definitions will apply and whenever appropriate, terms used in the singular will also include the plural and vice versa. In the event that any definition set forth below conflicts with the usage of that word in any other document, including any document incorporated herein by reference, the definition set forth below shall always control for purposes of interpreting this specification and its associated claims unless a contrary meaning is clearly intended.

[0077] As used herein, the term "allele" refers generally to alternative DNA sequences at the same physical locus on a segment of DNA, such as, for example, on homologous chromosomes. An allele can refer to DNA sequences which differ between the same physical locus found on homologous chromosomes within a single cell or organism or which differ at the same physical locus in multiple cells or organisms ("allelic variant"). In some instances, an allele can correspond to a single nucleotide difference at a particular physical locus.

In other embodiments and allele can correspond to nucleotide (single or multiple) insertion or deletion.

[0078] As used herein, the term “allele-specific primer” refers to an oligonucleotide sequence that hybridizes to a sequence comprising an allele of interest, and which when used in PCR can be extended to effectuate first strand cDNA synthesis. Allele-specific primers are specific for a particular allele of a given target DNA or loci and can be designed to detect a difference of as little as one nucleotide in the target sequence. Allele-specific primers may comprise an allele-specific nucleotide portion, a target-specific portion, and/or a tail.

[0079] As used herein, the terms “allele-specific nucleotide portion” or “allele-specific target nucleotide” refers to a nucleotide or nucleotides in an allele-specific primer that can selectively hybridize and be extended from one allele (for example, a minor or mutant allele) at a given locus to the exclusion of the other (for example, the corresponding major or wild type allele) at the same locus.

[0080] As used herein, the term “target-specific portion” refers to the region of an allele-specific primer that hybridizes to a target polynucleotide sequence. In some embodiments, the target-specific portion of the allele-specific primer is the priming segment that is complementary to the target sequence at a priming region 5' of the allelic variant to be detected. The target-specific portion of the allele-specific primer may comprise the allele-specific nucleotide portion. In other instances, the target-specific portion of the allele-specific primer is adjacent to the 3' allele-specific nucleotide portion.

[0081] As used herein, the terms “tail” or “5'-tail” refers to the non-3' end of a primer. This region typically will, although does not have to contain a sequence that is not complementary to the target polynucleotide sequence to be analyzed. The 5' tail can be any of about 2-30, 2-5, 4-6, 5-8, 6-12, 7-15, 10-20, 15-25 or 20-30 nucleotides, or any range in between, in length.

[0082] As used herein, the term “allele-specific blocker probe” (also referred to herein as “blocker probe,” “blocker,”) refers to an oligonucleotide sequence that binds to a strand of DNA comprising a particular allelic variant which is located on the same, opposite or complementary strand as that bound by an allele-specific primer, and reduces or prevents amplification of that particular allelic variant. As discussed in greater detail herein, allele-specific blocker probes generally comprise modifications, e.g., at the 3'-OH of the ribose ring, which prevent primer extension by a polymerase. The allele-specific blocker probe can be designed to anneal to the same or opposing strand of what the allele-specific primer anneals to and can be modified with a blocking group (e.g., a “non-extendable blocker moiety”) at its 3' terminal end. Thus, a blocker probe can be designed, for example, so as to tightly bind to a wild type allele (e.g., abundant allelic variant) in order to suppress amplification of the wild type allele while amplification is allowed to occur on the same or opposing strand comprising a mutant allele (e.g., rare allelic variant) by extension of an allele-specific primer. In illustrative examples, the allele-specific blocker probes do not include a label, such as a fluorescent, radioactive, or chemiluminescent label.

[0083] As used herein, the term “non-extendable blocker moiety” refers generally to a modification on an oligonucleotide sequence such as a probe and/or primer which renders it incapable of extension by a polymerase, for example, when hybridized to its complementary sequence in a PCR reaction.

Common examples of blocker moieties include modifications of the ribose ring 3'-OH of the oligonucleotide, which prevents addition of further bases to the 3'-end of the oligonucleotide sequence a polymerase. Such 3'-OH modifications are well known in the art. (See, e.g., Josefsen, M., et al., *Molecular and Cellular Probes*, 23 (2009):201-223; McKinzie, P. et al., *Mutagenesis*, 2006, 21(6):391-7; Parsons, B. et al., *Methods Mol Biol.* 2005, 291:235-45; Parsons, B. et al., *Nucleic Acids Res.* 1992, 25:20(10):2493-6; and Morlan, J. et al., *PLoS One* 2009, 4 (2): e4584, the disclosures of which are incorporated herein by reference in their entireties.)

[0084] As used herein, the terms “MGB,” “MGB group,” “MGB compound,” or “MBG moiety” refers to a minor groove binder. When conjugated to the 3' end of an oligonucleotide, an MGB group can function as a non-extendable blocker moiety.

[0085] An MGB is a molecule that binds within the minor groove of double stranded DNA. Although a general chemical formula for all known MGB compounds cannot be provided because such compounds have widely varying chemical structures, compounds which are capable of binding in the minor groove of DNA, generally speaking, have a crescent shape three dimensional structure. Most MGB moieties have a strong preference for A-T (adenine and thymine) rich regions of the B form of double stranded DNA. Nevertheless, MGB compounds which would show preference to C-G (cytosine and guanine) rich regions are also theoretically possible. Therefore, oligonucleotides comprising a radical or moiety derived from minor groove binder molecules having preference for C-G regions are also within the scope of the present invention.

[0086] Some MGBs are capable of binding within the minor groove of double stranded DNA with an association constant of 10^3 M^{-1} or greater. This type of binding can be detected by well established spectrophotometric methods such as ultraviolet (UV) and nuclear magnetic resonance (NMR) spectroscopy and also by gel electrophoresis. Shifts in UV spectra upon binding of a minor groove binder molecule and NMR spectroscopy utilizing the “Nuclear Overhauser” (NOSEY) effect are particularly well known and useful techniques for this purpose. Gel electrophoresis detects binding of an MGB to double stranded DNA or fragment thereof, because upon such binding the mobility of the double stranded DNA changes.

[0087] A variety of suitable minor groove binders have been described in the literature. See, for example, Kutayin, et al. U.S. Pat. No. 5,801,155; Wemmer, D. E., and Dervan P. B., *Current Opinion in Structural Biology*, 7:355-361 (1997); Walker, W. L., Kopka, J. L. and Goodsell, D. S., *Biopolymers*, 44:323-334 (1997); Zimmer, C. & Wahnert, U. *Frog. Biophys. Molec. Bio.* 47:31-112 (1986) and Reddy, B. S. P., Dondhi, S. M., and Lown, J. W., *Pharmacol. Therap.*, 84:1-111 (1999) (the disclosures of which are herein incorporated by reference in their entireties). A preferred MGB in accordance with the present disclosure is DPI₃. Synthesis methods and/or sources for such MGBs are also well known in the art. (See, e.g., U.S. Pat. Nos. 5,801,155; 6,492,346; 6,084,102; and 6,727,356, the disclosures of which are incorporated herein by reference in their entireties.)

[0088] As used herein, the term “MGB blocker probe,” “MBG blocker,” or “MGB probe” is an oligonucleotide sequence and/or probe further attached to a minor groove binder moiety at its 3' and/or 5' end. Oligonucleotides conjugated to MGB moieties form extremely stable duplexes with

single-stranded and double-stranded DNA targets, thus allowing shorter probes to be used for hybridization based assays. In comparison to unmodified DNA, MGB probes have higher melting temperatures (T_m) and increased specificity, especially when a mismatch is near the MGB region of the hybridized duplex. (See, e.g., Kutayavin, I. V., et al., *Nucleic Acids Research*, 2000, Vol. 28, No. 2: 655-661).

[0089] As used herein, the term “modified base” refers generally to any modification of a base or the chemical linkage of a base in a nucleic acid that differs in structure from that found in a naturally occurring nucleic acid. Such modifications can include changes in the chemical structures of bases or in the chemical linkage of a base in a nucleic acid, or in the backbone structure of the nucleic acid. (See, e.g., Latorra, D. et al., *Hum Mut* 2003, 2:79-85. Nakiandwe, J. et al., *Plant Method* 2007, 3:2.)

[0090] As used herein, the term “detector probe” refers to any of a variety of signaling molecules indicative of amplification. For example, SYBR® Green and other DNA-binding dyes are detector probes. Some detector probes can be sequence-based (also referred to herein as “locus-specific detector probe”), for example 5' nuclease probes. Various detector probes are known in the art, for example (TaqMan® probes described herein (See also U.S. Pat. No. 5,538,848) various stem-loop molecular beacons (See, e.g., U.S. Pat. Nos. 6,103,476 and 5,925,517 and Tyagi and Kramer, *Nature Biotechnology* 1996, 14:303-308), stemless or linear beacons (See, e.g., WO 99/21881), PNA Molecular Beacons™ (See, e.g., U.S. Pat. Nos. 6,355,421 and 6,593,091), linear PNA beacons (See, e.g., Kubista et al., 2001, SPIE 4264:53-58), non-FRET probes (See, e.g., U.S. Pat. No. 6,150,097), Sunrise®/Amplifluor® probes (U.S. Pat. No. 6,548,250), stem-loop and duplex Scorpion™ probes (Salinas et al., 2001, *Nucleic Acids Research* 29:E96 and U.S. Pat. No. 6,589,743), bulge loop probes (U.S. Pat. No. 6,590,091), pseudo knot probes (U.S. Pat. No. 6,589,250), cyclocons (U.S. Pat. No. 6,383,752), MGB Eclipse™ probe (Epoch Biosciences), hairpin probes (U.S. Pat. No. 6,596,490), peptide nucleic acid (PNA) light-up probes, self-assembled nanoparticle probes, and ferrocene-modified probes described, for example, in U.S. Pat. No. 6,485,901; Mhlana et al., 2001, *Methods* 25:463-471; Whitcombe et al., 1999, *Nature Biotechnology* 17:804-807; Isacson et al., 2000, *Molecular Cell Probes* 14:321-328; Svanvik et al., 2000, *Anal Biochem* 281:26-35; Wolffs et al., 2001, *Biotechniques* 766:769-771; Tsourkas et al., 2002, *Nucleic Acids Research* 30:4208-4215; Riccelli et al., 2002, *Nucleic Acids Research* 30:4088-4093; Zhang et al., 2002 Shanghai. 34:329-332; Maxwell et al., 2002, *J. Am. Chem. Soc.* 124:9606-9612; Braude et al., 2002, *Trends Biotechnol.* 20:249-56; Huang et al., 2002, *Chem Res. Toxicol.* 15:118-126; and Yu et al., 2001, *J. Am. Chem. Soc.* 14:11155-11161. Detector probes can comprise reporter dyes such as, for example, 6-carboxyfluorescein (6-FAM) or tetrachlorofluorescein (TET). Detector probes can also comprise quencher moieties such as tetramethylrhodamine (TAMRA), Black Hole Quenchers (Biosearch), Iowa Black (IDT), QSY quencher (Molecular Probes), and Dabsyl and Dabcyl sulfonate/carboxylate Quenchers (Epoch). Detector probes can also comprise two probes, wherein for example a fluor is on one probe, and a quencher on the other, wherein hybridization of the two probes together on a target quenches the signal, or wherein hybridization on a target alters the signal signature via a change in fluorescence. Detector probes can also comprise sulfonate derivatives of fluorescein dyes with SO₃

instead of the carboxylate group, phosphoramidite forms of fluorescein, phosphoramidite forms of CY5 (available, for example, from Amersham Biosciences-GE Healthcare).

[0091] As used herein, the term “locus-specific primer” refers to an oligonucleotide sequence that hybridizes to products derived from the extension of a first primer (such as an allele-specific primer) in a PCR reaction, and which can effectuate second strand cDNA synthesis of said product. Accordingly, in some embodiments, the allele-specific primer serves as a forward PCR primer and the locus-specific primer serves as a reverse PCR primer, or vice versa. In some preferred embodiments, locus-specific primers are present at a higher concentration as compared to the allele-specific primers.

[0092] As used herein, the term “rare allelic variant” refers to a target polynucleotide present at a lower level in a sample as compared to an alternative allelic variant. The rare allelic variant may also be referred to as a “minor allelic variant” and/or a “mutant allelic variant.” For instance, the rare allelic variant may be found at a frequency less than 1/10, 1/100, 1/1,000, 1/10,000, 1/100,000, 1/1,000,000, 1/10,000,000, 1/100,000,000 or 1/1,000,000,000 compared to another allelic variant for a given SNP or gene. Alternatively, the rare allelic variant can be, for example, less than 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 50, 75, 100, 250, 500, 750, 1,000, 2,500, 5,000, 7,500, 10,000, 25,000, 50,000, 75,000, 100,000, 250,000, 500,000, 750,000, or 1,000,000 copies per 1, 10, 100, 1,000 micro liters of a sample or a reaction volume.

[0093] As used herein, the terms “abundant allelic variant” may refer to a target polynucleotide present at a higher level in a sample as compared to an alternative allelic variant. The abundant allelic variant may also be referred to as a “major allelic variant” and/or a “wild type allelic variant.” For instance, the abundant allelic variant may be found at a frequency greater than 10x, 100x, 1,000x, 10,000x, 100,000x, 1,000,000x, 10,000,000x, 100,000,000x or 1,000,000,000x compared to another allelic variant for a given SNP or gene. Alternatively, the abundant allelic variant can be, for example, greater than 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 50, 75, 100, 250, 500, 750, 1,000, 2,500, 5,000, 7,500, 10,000, 25,000, 50,000, 75,000, 100,000, 250,000, 500,000, 750,000, 1,000,000 copies per 1, 10, 100, 1,000 micro liters of a sample or a reaction volume.

[0094] As used herein, the terms “first” and “second” are used to distinguish the components of a first reaction (e.g., a “first” reaction; a “first” allele-specific primer) and a second reaction (e.g., a “second” reaction; a “second” allele-specific primer). By convention, as used herein the first reaction amplifies a first (for example, a rare) allelic variant and the second reaction amplifies a second (for example, an abundant) allelic variant or vice versa.

[0095] As used herein, both “first allelic variant” and “second allelic variant” can pertain to alleles of a given locus from the same organism. For example, as might be the case in human samples (e.g., cells) comprising wild type alleles, some of which have been mutated to form a minor or rare allele. The first and second allelic variants of the present teachings can also refer to alleles from different organisms. For example, the first allele can be an allele of a genetically modified organism, and the second allele can be the corresponding allele of a wild type organism. The first allelic variants and second allelic variants of the present teachings can be contained on gDNA, as well as mRNA and cDNA, and

generally any target nucleic acids that exhibit sequence variability due to, for example, SNP or nucleotide(s) insertion and/or deletion mutations.

[0096] As used herein, the term “thermostable” or “thermostable polymerase” refers to an enzyme that is heat stable or heat resistant and catalyzes polymerization of deoxyribonucleotides to form primer extension products that are complementary to a nucleic acid strand. Thermostable DNA polymerases useful herein are not irreversibly inactivated when subjected to elevated temperatures for the time necessary to effect destabilization of single-stranded nucleic acids or denaturation of double-stranded nucleic acids during PCR amplification. Irreversible denaturation of the enzyme refers to substantial loss of enzyme activity. Preferably a thermostable DNA polymerase will not irreversibly denature at about 90°-100° C. under conditions such as is typically required for PCR amplification.

[0097] As used herein, the term “PCR amplifying” or “PCR amplification” refers generally to cycling polymerase-mediated exponential amplification of nucleic acids employing primers that hybridize to complementary strands, as described for example in Innis et al., PCR Protocols: A Guide to Methods and Applications, Academic Press (1990). Devices have been developed that can perform thermal cycling reactions with compositions containing fluorescent indicators which are able to emit a light beam of a specified wavelength, read the intensity of the fluorescent dye, and display the intensity of fluorescence after each cycle. Devices comprising a thermal cycler, light beam emitter, and a fluorescent signal detector, have been described, e.g., in U.S. Pat. Nos. 5,928,907; 6,015,674; 6,174,670; and 6,814,934 and include, but are not limited to, the ABI Prism® 7700 Sequence Detection System (Applied Biosystems, Foster City, Calif.), the ABI GeneAmp® 5700 Sequence Detection System (Applied Biosystems, Foster City, Calif.), the ABI GeneAmp® 7300 Sequence Detection System (Applied Biosystems, Foster City, Calif.), the ABI GeneAmp® 7500 Sequence Detection System (Applied Biosystems, Foster City, Calif.), the StepOne™ Real-Time PCR System (Applied Biosystems, Foster City, Calif.) and the ABI GeneAmp® 7900 Sequence Detection System (Applied Biosystems, Foster City, Calif.).

[0098] As used herein, the term “pre-amplification” or “pre-amplify” refers to a process wherein a plurality of primer pairs are included in a multiplexed PCR amplification reaction, and the multiplexed amplification reaction undergoes a limited number of cycles so that the PCR-based pre-amplification reaction ends prior to the PCR plateau and/or reagent depletion. The term “PCR-based pre-amplification” can be considered to indicate that a secondary amplification reaction is subsequently performed, typically of lower plexy level than the PCR-based pre-amplification reaction. This secondary amplification reaction, typically a plurality of separate secondary amplification reactions, can employ primer pairs encoded by the primers used in the multiplexed PCR-based pre-amplification reaction. However, each secondary amplification reaction typically comprises a single or a few primer pairs. Further examples of PCR-based pre-amplification approaches can be found, for example, in U.S. Pat. No. 6,605,451, and U.S. patent application Ser. No. 10/723,520, the disclosures of which are herein incorporated by reference in their entireties.

[0099] As used herein, the term “Tm” or “melting temperature” of an oligonucleotide refers to the temperature (in

degrees Celsius) at which 50% of the molecules in a population of a single-stranded oligonucleotide are hybridized to their complementary sequence and 50% of the molecules in the population are not-hybridized to said complementary sequence. The Tm of a primer or probe can be determined empirically by means of a melting curve. In some cases it can also be calculated using formulas well known in the art (See, e.g., Maniatis, T., et al., Molecular cloning: a laboratory manual/Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.: 1982).

[0100] As used herein, the term “sensitivity” refers to the minimum amount (number of copies or mass) of a template that can be detected by a given assay. As used herein, the term “specificity” refers to the ability of an assay to distinguish between amplification from a matched template versus a mismatched template. Frequently, specificity is expressed as $\Delta C_t = C_{t\_mismatch} - C_{t\_match}$. An improvement in specificity or “specificity improvement” or “fold difference” is expressed herein as $2^{(\Delta C_{t\_condition1} - (\Delta C_{t\_condition2}))}$. The term “selectivity” refers to the extent to which an AS-PCR assay can be used to determine minor (often mutant) alleles in mixtures without interferences from major (often wild type) alleles. Selectivity is often expressed as a ratio or percentage. For example, an assay that can detect 1 mutant template in the presence of 100 wild type templates is said to have a selectivity of 1:100 or 1%. As used herein, assay selectivity can also be calculated as $1/2^{\Delta C_t}$ or as a percentage using $(1/2^{\Delta C_t} \times 100)$.

[0101] As used herein, the term “Ct” or “Ct value” refers to threshold cycle and signifies the cycle of a PCR amplification assay in which signal from a reporter that is indicative of amplicon generation (e.g., fluorescence) first becomes detectable above a background level. In some embodiments, the threshold cycle or “Ct” is the cycle number at which PCR amplification becomes exponential.

[0102] As used herein, the term “delta Ct” or “ ΔC_t ” refers to the difference in the numerical cycle number at which the signal passes the fixed threshold between two different samples or reactions. In some embodiments delta Ct is the difference in numerical cycle number at which exponential amplification is reached between two different samples or reactions. The delta Ct can be used to identify the specificity between a matched primer to the corresponding target nucleic acid sequence and a mismatched primer to the same corresponding target nucleic acid sequence.

[0103] In some embodiments, the calculation of the delta Ct value between a mismatched primer and a matched primer is used as one measure of the discriminating power of allele-specific PCR. In general, any factor which increases the difference between the Ct value for an amplification reaction using a primer that is matched to a target sequence (e.g., a sequence comprising an allelic variant of interest) and that of a mismatched primer will result in greater allele discrimination power.

[0104] According to various embodiments, a Ct value may be determined using a derivative of a PCR curve. For example, a first, second, or nth order derivative method may be performed on a PCR curve in order to determine a Ct value. In various embodiments, a characteristic of a derivative may be used in the determination of a Ct value. Such characteristics may include, but are not limited by, a positive inflection of a second derivative, a negative inflection of a second derivative, a zero crossing of the second derivative, or a positive inflection of a first derivative. In various embodiments, a Ct value may be determined using a thresholding and baselining

method. For example, an upper bound to an exponential phase of a PCR curve may be established using a derivative method, while a baseline for a PCR curve may be determined to establish a lower bound to an exponential phase of a PCR curve. From the upper and lower bound of a PCR curve, a threshold value may be established from which a Ct value is determined. Other methods for the determination of a Ct value known in the art, for example, but not limited by, various embodiments of a fit point method, and various embodiments of a sigmoidal method (See, e.g., U.S. Pat. Nos. 6,303,305; 6,503,720; 6,783,934, 7,228,237 and U.S. Application No. 2004/0096819; the disclosures of which are herein incorporated by reference in their entireties).

III. Compositions, Methods and Kits

[0105] In one aspect, the present invention provides compositions for use in identifying and/or quantitating an allelic variant in a nucleic acid sample. Some of these compositions can comprise: (a) an allele-specific primer; (b) an allele-specific blocker probe; (c) a detector probe; and (d) a locus-specific primer, or any combinations thereof. In some embodiments of the compositions, the compositions may further comprise a polymerase, dNTPs, reagents and/or buffers suitable for PCR amplification, and/or a template sequence or nucleic acid sample. In some embodiments, the polymerase can be thermostable.

[0106] In another aspect, the invention provides compositions comprising: (i) a first allele-specific primer, wherein an allele-specific nucleotide portion of the first allele-specific primer is complementary to the first allelic variant of a target sequence; and (ii) a first allele-specific blocker probe that is complementary to a region of the target sequence comprising the second allelic variant, wherein said region encompasses a position corresponding to the binding position of the allele-specific nucleotide portion of the first allele-specific primer, and wherein the first allele-specific blocker probe comprises a minor groove binder.

[0107] In some illustrative embodiments, the compositions can further include a locus-specific primer that is complementary to a region of the target sequence that is 3' from the first allelic variant and on the opposite strand.

[0108] In further embodiments, the compositions can further include a detector probe.

[0109] In other embodiments, compositions for pre-amplification are further provided as described in further detail below.

[0110] In another aspect, the present invention provides methods for amplifying an allele-specific sequence. Some of these methods can include: (a) hybridizing an allele-specific primer to a first nucleic acid molecule comprising a target allele; (b) hybridizing an allele-specific blocker probe to a second nucleic acid molecule comprising an alternative allele wherein the alternative allele corresponds to the same loci as the target allele; (c) hybridizing a locus-specific detector probe to the first nucleic acid molecule; (d) hybridizing a locus-specific primer to the extension product of the allele-specific primer and (e) PCR amplifying the target allele.

[0111] In another aspect, the present invention provides methods for detecting and/or quantitating an allelic variant in a mixed sample. Some of these methods can involve: (a) in a first reaction mixture hybridizing a first allele-specific primer to a first nucleic acid molecule comprising a first allele (allele-1) and in a second reaction mixture hybridizing a second allele-specific primer to a first nucleic acid molecule com-

prising a second allele (allele-2), wherein the allele-2 corresponds to the same loci as allele-1; (b) in the first reaction mixture hybridizing a first allele-specific blocker probe to a second nucleic acid molecule comprising allele-2 and in the second reaction mixture hybridizing a second allele-specific blocker probe to a second nucleic acid molecule comprising allele-1; (c) in the first reaction mixture, hybridizing a first detector probe to the first nucleic acid molecule and in the second reaction mixture, hybridizing a second detector probe to the first nucleic acid molecule; (d) in the first reaction mixture hybridizing a first locus-specific primer to the extension product of the first allele-specific primer and in the second reaction mixture hybridizing a second locus-specific primer to the extension product of the second allele-specific primer; and (e) PCR amplifying the first nucleic acid molecule to form a first set or sample of amplicons and PCR amplifying the second nucleic acid molecule to form a second set or sample of amplicons; and (f) comparing the first set of amplicons to the second set of amplicons to quantitate allele-1 in the sample comprising allele-2 and/or allele-2 in the sample comprising allele-1.

[0112] In yet another aspect, the present invention provides methods for detecting and/or quantitating allelic variants. Some of these methods can comprise: (a) PCR amplifying a first allelic variant in a first reaction comprising (i) a low-concentration first allele-specific primer, (ii) a first locus-specific primer, and (iii) a first blocker probe to form first amplicons; (b) PCR amplifying a second allelic variant in a second reaction comprising (i) a low-concentration second allele-specific primer, (ii) a second locus-specific primer, and (iii) a second blocker probe to form second amplicons; and (d) comparing the first amplicons to the second amplicons to quantitate the first allelic variant in the sample comprising second allelic variants.

[0113] In yet another aspect, the present invention provides methods for detecting a first allelic variant of a target sequence in a nucleic acid sample suspected of comprising at least a second allelic variant of the target sequence. Methods of this aspect include forming a first reaction mixture by combining the following: (i) a nucleic acid sample; (ii) a first allele-specific primer, wherein an allele-specific nucleotide portion of the first allele-specific primer is complementary to the first allelic variant of the target sequence; (iii) a first allele-specific blocker probe that is complementary to a region of the target sequence comprising the second allelic variant, wherein said region encompasses a position corresponding to the binding position of the allele-specific nucleotide portion of the first allele-specific primer, and wherein the first allele-specific blocker probe comprises a minor groove binder; (iv) a first locus-specific primer that is complementary to a region of the target sequence that is 3' from the first allelic variant and on the opposite strand; and (v) a first detector probe.

[0114] Next an amplification reaction, typically a PCR amplification reaction, is carried out on the first reaction mixture using the first locus-specific primer and the first allele-specific primer to form a first amplicon. Then, the first amplicon is detected by a change in a detectable property of the first detector probe upon binding to the amplicon, thereby detecting the first allelic variant of the target gene in the nucleic acid sample. The detector probe in some illustrative embodiments is a 5' nuclease probe. The detectable property in certain illustrative embodiments is fluorescence.

[0115] In some embodiments, the 3' nucleotide position of the 5' target region of the first allele-specific primer is an allele-specific nucleotide position. In certain other illustrative embodiments, including those embodiments where the 3' nucleotide position of the 5' target region of the first allele-specific primer is an allele-specific nucleotide position, the blocking region of the allele-specific primer encompasses the allele-specific nucleotide position. Furthermore, in illustrative embodiments, the first allele-specific blocker probe includes a minor groove binder. Furthermore, the allele-specific blocker probe in certain illustrative embodiments does not have a label, for example a fluorescent label, or a quencher.

[0116] In certain illustrative embodiments, the quantity of the first allelic variant is determined by evaluating the change in a detectable property of the first detector probe.

[0117] In certain illustrative embodiments, the method further includes forming a second reaction mixture by combining (i) the nucleic acid sample; (ii) a second allele-specific primer, wherein an allele-specific nucleotide portion of the second allele-specific primer is complementary to the second allelic variant of the target sequence; (iii) a second allele-specific blocker probe that is complementary to a region of the target sequence comprising the first allelic variant, wherein said region encompasses a position corresponding to the binding position of the allele-specific nucleotide portion of the second allele-specific primer, and wherein the second allele-specific blocker probe comprises a minor groove binder; (iv) a second locus-specific primer that is complementary to a region of the target sequence that is 3' from the second allelic variant and on the opposite strand; and (v) a second detector probe. Next, an amplification reaction is carried out on the second reaction mixture using the second allele-specific primer and the locus-specific primer, to form a second amplicon. Then the second amplicon is detected by a change in a detectable property of the detector probe.

[0118] In certain embodiments, the method further includes comparing the change in a detectable property of the first detector probe in the first reaction mixture to the change in a detectable property of the second detector probe in the second reaction mixture.

[0119] In some preferred embodiments, the methods may include a 2-stage cycling protocol. In some embodiments, the methods for detecting an allelic variant in a target sequence in a nucleic acid sample comprises forming a reaction mixture comprising:

[0120] i) a nucleic acid sample;

[0121] ii) an allele-specific primer, wherein an allele-specific nucleotide portion of the allele-specific primer is complementary to the first allelic variant of the target sequence;

[0122] iii) an allele-specific blocker probe that is complementary to a region of the target sequence comprising a second allelic variant, wherein said region encompasses a position corresponding to the binding position of the allele-specific nucleotide portion of the allele-specific primer, and wherein the allele-specific blocker probe comprises a blocking moiety;

[0123] iv) a locus-specific primer that is complementary to a region of the target sequence that is 3' from the first allelic variant and on the opposite strand; and

[0124] v) a detector probe;

[0125] b) PCR amplifying the target sequence using a 2-stage cycling protocol, comprising:

[0126] i) a first amplification step comprising a first number of cycles run at a first annealing/extension temperature; and

[0127] ii) a second amplification step comprising a second number of cycles run at a second annealing/extension temperature,

[0128] wherein the first number of cycles is fewer than the second number of cycles and the first annealing/extension temperature is lower than said second annealing/extension temperature; and

[0129] c) detecting a change in a detectable property of the detector probe in the amplified products of the target sequence produced by step (b).

[0130] In some embodiments, the cycling protocol comprises a first stage of amplification that employs an initial number of cycles at a lower annealing/extension temperature, followed by a second stage of amplification that employs a number of cycles at a higher annealing/extension temperature. Due to the lower T_m of cast-PCR allele-specific primers (e.g., 53-56° C.), PCR may not be optimal at standard annealing/extension conditions (e.g., 60-64° C.). Consequently, lower annealing/extension temperatures may be used during the initial cycling stage which improve cast-PCR efficiency significantly.

[0131] In some embodiments, the number of cycles used in the first stage of the cast-PCR cycling protocol is fewer than the number of cycles used in the second stage. In some embodiments of the cast-PCR methods, the number of cycles used in the first stage of the cycling protocol is about 2%-20%, 4%-18%, 6%-16%, 8%-14%, 10%-12%, or any percent in between, of the total number of cycles used in the second stage. In some embodiments, the first stage employs between 1 to 10 cycles, 2 to 8 cycles, 3 to 7 cycles, or 4 to 6 cycles, and all number of cycles in between, e.g., 2, 3, 4, 5, 6, or 7 cycles.

[0132] In some embodiments, the number of cycles used in the second stage of the cast-PCR cycling protocol is greater than the number of cycles used in the first stage. In some embodiments of the cast-PCR methods, the number of cycles used in the second stage of the cycling protocol is 5 times, 6 times, 8 times, 10 times, 12 times, 18 times, 25 times, or 30 times the number of cycles used in the first stage. In some embodiments, the second stage employs between 30 to 50 cycles, 35 to 48 cycles, 40 to 46 cycles, or any number of cycles in between, e.g., 42, 43, 44, 45, or 46 cycles.

[0133] In some embodiments, the lower annealing/extension temperature used during the first cycling stage is about 1° C., about 2° C., about 3° C., about 4° C., or about 5° C. lower than the annealing/extension temperature used during the second cycling stage. In some preferred embodiments, the annealing/extension temperature of the first stage is between 50° C. to 60° C., 52° C. to 58° C., or 54° C. to 56° C., e.g., 53° C., 54° C., 55° C. or 55° C. In some preferred embodiments the annealing/extension temperature of the second stage is between 56° C. to 66° C., 58° C. to 64° C., or 60° C. to 62° C., e.g., 58° C., 60° C., 62° C. or 64° C.

[0134] There are several major advantages of this 2-stage PCR cycling protocol used in cast-PCR that make it better than conventional AS-PCR methods. First, it improves the detection sensitivity by lowering the C_t value for matched targets or alleles. Next, it improves the specificity of cast-PCR by increasing the ΔC_t between C_t values of matched and

mismatched sequences. Finally, it can improve the uniformity of cast-PCR by making it similarly efficient across various assays.

[0135] In some embodiments, the 2-stage cycling protocol improves cast-PCR efficiency (e.g. specificity and sensitivity). In some embodiments, the specificity is increased by 2-8x, 3-7x, or 4-6x, or any number of times in between (e.g., 3x, 4x, 5x, 6x, or 7x), more than the specificity of cast-PCR reactions run without the 2-stage cycling protocol. In other embodiments, the sensitivity is increased by 1-3x, (e.g., 1x, 2x, or 3x) more than the sensitivity of cast-PCR reactions run without the 2-stage cycling protocol. In some embodiments, the difference in ΔC_t between assays run using cast-PCR with the 2-stage cycling protocol compared to cast-PCR assays run without the 2-stage cycling protocol is a ΔC_t of 1, 2 or 3.

[0136] In yet another aspect, the present invention provides a reaction mixture that includes the following (i) nucleic acid molecule; (ii) an allele-specific primer, wherein an allele-specific nucleotide portion of the allele-specific primer is complementary to a first allelic variant of a target sequence; (iii) an allele-specific blocker probe that is complementary to a region of the target sequence comprising a second allelic variant, wherein said region encompasses a position corresponding to the binding position of the allele-specific nucleotide portion of the allele-specific primer, and wherein the allele-specific blocker probe comprises a minor groove binder; (iv) a locus-specific primer that is complementary to a region of the target sequence that is 3' from the first allelic variant and on the opposite strand; and (v) a detector probe.

[0137] In certain embodiments, the methods of the invention are used to detect a first allelic variant that is present at a frequency less than $1/10$, $1/100$, $1/1,000$, $1/10,000$, $1/100,000$, $1/1,000,000$, $1/10,000,000$, $1/100,000,000$ or $1/1,000,000,000$, and any fractional ranges in between, of a second allelic variant for a given SNP or gene. In other embodiments, the methods are used to detect a first allelic variant that is present in less than 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 50, 75, 100, 250, 500, 750, 1,000, 2,500, 5,000, 7,500, 10,000, 25,000, 50,000, 75,000, 100,000, 250,000, 500,000, 750,000, 1,000,000 copies per 1, 10, 100, 1,000 micro liters, and any fractional ranges in between, of a sample or a reaction volume.

[0138] In some embodiments, the first allelic variant is a mutant. In some embodiments the second allelic variant is wild type. In some embodiments, the present methods can involve detecting one mutant molecule in a background of at least 1,000 to 1,000,000, such as about 1000 to 10,000, about 10,000 to 100,000, or about 100,000 to 1,000,000 wild type molecules, or any fractional ranges in between. In some embodiments, the methods can provide high sensitivity and the efficiency at least comparable to that of TaqMan®-based assays.

[0139] In some embodiments, the comparison of the first amplicons and the second amplicons involving the disclosed methods provides improvements in specificity from 1,000x to 1,000,000x fold difference, such as about 1000 to 10,000x, about 10,000 to 100,000x, or about 100,000 to 1,000,000x fold difference, or any fractional ranges in between. In some embodiments, the size of the amplicons range from about 60-120 nucleotides long.

[0140] In some embodiments, the methods can further include additional steps used for pre-amplification of cast-PCR reactions as described in further detail below.

[0141] In another aspect, the present invention provides kits for quantitating a first allelic variant in a sample comprising

an alternative second allelic variants that include: (a) a first allele-specific primer; (b) a second allele-specific primer; (c) a first locus-specific primer; (d) a second locus-specific primer; (e) a first allele-specific blocker probe; (f) a second allele-specific blocker probe; and (g) a polymerase. In some embodiments of the disclosed kits, the kit further comprises a first locus-specific detector probe and a second locus-specific detector probe.

[0142] In another aspect, the present invention provides kits that include two or more containers comprising the following components independently distributed in one of the two or more containers: (i) a first allele-specific primer, wherein an allele-specific nucleotide portion of the first allele-specific primer is complementary to the first allelic variant of a target sequence; and (ii) a first allele-specific blocker probe that is complementary to a region of the target sequence comprising the second allelic variant, wherein said region encompasses a position corresponding to the binding position of the allele-specific nucleotide portion of the first allele-specific primer, and wherein the first allele-specific blocker probe comprises a minor groove binder.

[0143] In some illustrative embodiments, the kits can further include a locus-specific primer that is complementary to a region of the target sequence that is 3' from the first allelic variant and on the opposite strand.

[0144] In other embodiments, the kits can further include a detector probe.

[0145] In some embodiments, the kits can further include additional components used for pre-amplification as described in further detail below.

[0146] In some embodiments, the compositions, methods, and/or kits can be used in detecting circulating cells in diagnosis. In one embodiment, the compositions, methods, and/or kits can be used to detect tumor cells in blood for early cancer diagnosis. In some embodiments, the compositions, methods, and/or kits can be used for cancer or disease-associated genetic variation or somatic mutation detection and validation. In some embodiments, the compositions, methods, and/or kits can be used for genotyping tri-, di- and bi-allelic SNPs. In other embodiments, the compositions, methods, and/or kits can be used for identifying single or multiple nucleotide insertion or deletion mutations. In some embodiments, the compositions, methods, and/or kits can be used for DNA typing from mixed DNA samples for QC and human identification assays, cell line QC for cell contaminations, allelic gene expression analysis, virus typing/rare pathogen detection, mutation detection from pooled samples, detection of circulating tumor cells in blood, and/or prenatal diagnostics.

[0147] In some embodiments, the compositions, methods, and/or kits are compatible with various instruments such as, for example, SDS instruments from Applied Biosystems (Foster City, Calif.).

[0148] Allele-Specific Primers

[0149] Allele-specific primers (ASPS) designed with low Tms exhibit increased discrimination of allelic variants. In some embodiments, the allele-specific primers are short oligomers ranging from about 15-30, such as about 16-28, about 17-26, about 18-24, or about 20-22, or any range in between, nucleotides in length. In some embodiments, the Tm of the allele-specific primers range from about 50° C. to 70° C., such as about 52° C. to 68° C., about 54° C. to 66° C., about 56° C. to 64° C., about 58° C. to 62° C., or any temperature in between (e.g., 53° C., 54° C., 55° C., 56° C.). In other embodi-

ments, the T_m of the allele-specific primers is about 3° C. to 6° C. higher than the anneal/extend temperature of the PCR cycling conditions employed during amplification.

[0150] Low allele-specific primer concentration can also improve selectivity. Reduction in concentration of allele-specific primers below 900 nM can increase the delta C_t between matched and mismatched sequences. In some embodiments of the disclosed compositions, the concentration of allele-specific primers ranges from about 20 nM to 900 nM, such as about 50 nM to 700 nM, about 100 nM to 500 nM, about 200 nM to 300 nM, about 400 nM to 500 nM, or any range in between. In some exemplary embodiments, the concentration of the allele-specific primers is between about 200 nM to 400 nM.

[0151] In some embodiments, allele-specific primers can comprise an allele-specific nucleotide portion that is specific to the target allele of interest. The allele-specific nucleotide portion of an allele-specific primer is complementary to one allele of a gene, but not another allele of the gene. In other words, the allele-specific nucleotide portion binds to one or more variable nucleotide positions of a gene that is nucleotide positions that are known to include different nucleotides for different allelic variants of a gene. The allele-specific nucleotide portion is at least one nucleotide in length. In exemplary embodiments, the allele-specific nucleotide portion is one nucleotide in length. In some embodiments, the allele-specific nucleotide portion of an allele-specific primer is located at the 3' terminus of the allele-specific primer. In other embodiments, the allele-specific nucleotide portion is located about 1-2, 3-4, 5-6, 7-8, 9-11, 12-15, or 16-20 nucleotides in from the 3' most-end of the allele-specific primer.

[0152] Allele-specific primers designed to target discriminating bases can also improve discrimination of allelic variants. In some embodiments, the nucleotide of the allele-specific nucleotide portion targets a highly discriminating base (e.g., for detection of A/A, A/G, G/A, G/G, A/C, or C/A alleles). Less discriminating bases, for example, may involve detection of C/C, T/C, G/T, T/G, C/T alleles. In some embodiments, for example when the allele to be detected involves A/G or C/T SNPs, A or G may be used as the 3' allele-specific nucleotide portion of the allele-specific primer (e.g., if A or T is the major allele), or C or T may be used as the 3' allele-specific nucleotide portion of the allele-specific primer (e.g., if C or G is the major allele). In other embodiments, A may be used as the nucleotide-specific portion at the 3' end of the allele specific primer (e.g., the allele-specific nucleotide portion) when detecting and/or quantifying A/T SNPs. In other embodiments, G may be used as the nucleotide-specific portion at the 3' end of the allele specific primer when detecting and/or quantifying C/G SNPs.

[0153] In some embodiments, the allele-specific primer can comprise a target-specific portion that is specific to the polynucleotide sequence (or locus) of interest. In some embodiments the target-specific portion is about 75-85%, 85-95%, 95-99% or 100% complementary to the target polynucleotide sequence of interest. In some embodiments, the target-specific portion of the allele-specific primer can comprise the allele-specific nucleotide portion. In other embodiments, the target-specific portion is located 5' to the allele-specific nucleotide portion. The target-specific portion can be about 4-30, about 5-25, about 6-20, about 7-15, or about 8-10 nucleotides in length. In some embodiments, the T_m of the target specific portion is about 5° C. below the anneal/extend temperature used for PCR cycling. In some embodiments, the T_m

of the target specific portion of the allele-specific primer ranges from about 51° C. to 60° C., about 52° C. to 59° C., about 53° C. to 58° C., about 54° C. to 57° C., about 55° C. to 56° C., or about 50° C. to about 60° C.

[0154] In some embodiments of the disclosed methods and kits, the target-specific portion of the first allele-specific primer and the target-specific portion of the second allele-specific primer comprise the same sequence. In other embodiments, the target-specific portion of the first allele-specific primer and the target-specific portion of the second allele-specific primer are the same sequence.

[0155] In some embodiments, the allele-specific primer comprises a tail. Allele-specific primers comprising tails, enable the overall length of the primer to be reduced, thereby lowering the T_m without significant impact on assay sensitivity.

[0156] In some exemplary embodiments, the tail is on the 5' terminus of the allele-specific primer. In some embodiments, the tail is located 5' of the target-specific portion and/or allele-specific nucleotide portion of the allele-specific primer. In some embodiments, the tail is about 65-75%, about 75-85%, about 85-95%, about 95-99% or about 100% non-complementary to the target polynucleotide sequence of interest. In some embodiments the tail can be about 2-40, such as about 4-30, about 5-25, about 6-20, about 7-15, or about 8-10 nucleotides in length. In some embodiments the tail is GC-rich. For example, in some embodiments the tail sequence is comprised of about 50-100%, about 60-100%, about 70-100%, about 80-100%, about 90-100% or about 95-100% G and/or C nucleotides.

[0157] The tail of the allele-specific primer may be configured in a number of different ways, including, but not limited to a configuration whereby the tail region is available after primer extension to hybridize to a complementary sequence (if present) in a primer extension product. Thus, for example, the tail of the allele-specific primer can hybridize to the complementary sequence in an extension product resulting from extension of a locus-specific primer.

[0158] In some embodiments of the disclosed methods and kits, the tail of the first allele-specific primer and the tail of the second allele-specific primer comprise the same sequence. In other embodiments, the 5' tail of the first allele-specific primer and the 5' tail of the second allele-specific primer are the same sequence.

[0159] Allele-Specific Blocker Probes

[0160] Allele-specific blocker probes (or ASBs) (herein sometimes referred to as "blocker probes") may be designed as short oligomers that are single-stranded and have a length of 100 nucleotides or less, more preferably 50 nucleotides or less, still more preferably 30 nucleotides or less and most preferably 20 nucleotides or less with a lower limit being approximately 5 nucleotides.

[0161] In some embodiments, the T_m of the blocker probes range from 58° C. to 70° C., 61° C. to 69° C., 62° C. to 68° C., 63° C. to 67° C., 64° C. to 66° C., or about 60° C. to about 63° C., or any range in between. In yet other embodiments, the T_m of the allele-specific blocker probes is about 3° C. to 6° C. higher than the anneal/extend temperature in the PCR cycling conditions employed during amplification.

[0162] In some embodiments, the blocker probes are not cleaved during PCR amplification. In some embodiments, the blocker probes comprise a non-extendable blocker moiety at their 3'-ends. In some embodiments, the blocker probes can further comprise other moieties (including, but not limited to

additional non-extendable blocker moieties, quencher moieties, fluorescent moieties, etc) at their 3'-end, 5'-end, and/or any internal position in between. In some embodiments, the allele position is located about 5-15, such as about 5-11, about 6-10, about 7-9, about 7-12, or about 9-11, such as about 6, about 7, about 8, about 9, about 10, or about 11 nucleotides away from the non-extendable blocker moiety of the allele-specific blocker probes when hybridized to their target sequences. In some embodiments, the non-extendable blocker moiety can be, but is not limited to, an amine (NH₂), biotin, PEG, DPI₃, or PO₄. In some preferred embodiments, the blocker moiety is a minor groove binder (MGB) moiety. (The oligonucleotide-MGB conjugates of the present invention are hereinafter sometimes referred to as "MGB blocker probes" or "MGB blockers.")

[0163] As disclosed herein, the use of MGB moieties in allele-specific blocker probes can increase the specificity of allele-specific PCR. One possibility for this effect is that, due to their strong affinity to hybridize and strongly bind to complementary sequences of single or double stranded nucleic acids, MGBs can lower the T_m of linked oligonucleotides (See, for example, Kuttyavin, I., et al., *Nucleic Acids Res.*, 2000, Vol. 28, No. 2: 655-661). Oligonucleotides comprising MGB moieties have strict geometric requirements since the linker between the oligonucleotide and the MGB moiety must be flexible enough to allow positioning of the MGB in the minor groove after DNA duplex formation. Thus, MGB blocker probes can provide larger T_m differences between matched versus mismatched alleles as compared to conventional DNA blocker probes.

[0164] In general, MGB moieties are molecules that bind within the minor groove of double stranded DNA. Although a generic chemical formula for all known MGB compounds cannot be provided because such compounds have widely varying chemical structures, compounds which are capable of binding in the minor groove of DNA, generally speaking, have a crescent shape three dimensional structure. Most MGB moieties have a strong preference for A-T (adenine and thymine) rich regions of the B form of double stranded DNA. Nevertheless, MGB compounds which would show preference to C-G (cytosine and guanine) rich regions are also theoretically possible. Therefore, oligonucleotides comprising a radical or moiety derived from minor groove binder molecules having preference for C-G regions are also within the scope of the present invention.

[0165] Some MGBs are capable of binding within the minor groove of double stranded DNA with an association constant of 10³M⁻¹ or greater. This type of binding can be detected by well established spectrophotometric methods such as ultraviolet (UV) and nuclear magnetic resonance (NMR) spectroscopy and also by gel electrophoresis. Shifts in UV spectra upon binding of a minor groove binder molecule and NMR spectroscopy utilizing the "Nuclear Overhauser" (NOSEY) effect are particularly well known and useful techniques for this purpose. Gel electrophoresis detects binding of an MGB to double stranded DNA or fragment thereof, because upon such binding the mobility of the double stranded DNA changes.

[0166] A variety of suitable minor groove binders have been described in the literature. See, for example, Kuttyavin, et al. U.S. Pat. No. 5,801,155; Wemmer, D. E., and Dervan P. B., *Current Opinion in Structural Biology*, 7:355-361 (1997); Walker, W. L., Kopka, J. L. and Goodsell, D. S., *Biopolymers*, 44:323-334 (1997); Zimmer, C & Wahnert, U. *Prog. Biophys.*

Molec. Bio. 47:31-112 (1986) and Reddy, B. S. P., Dondhi, S. M., and Lown, J. W., *Pharmacol. Therap.*, 84:1-111 (1999). In one group of embodiments, the MGB is selected from the group consisting of CC1065 analogs, lexitropsins, distamycin, netropsin, berenil, duocarmycin, pentamidine, 4,6-diamino-2-phenylindole and pyrrolo[2,1-c][1,4]benzodiazepines. A preferred MGB in accordance with the present disclosure is DPI₃ (see U.S. Pat. No. 6,727,356, the disclosure of which is incorporated herein by reference in its entirety).

[0167] Suitable methods for attaching MGBs through linkers to oligonucleotides or probes and have been described in, for example, U.S. Pat. Nos. 5,512,677; 5,419,966; 5,696,251; 5,585,481; 5,942,610; 5,736,626; 5,801,155 and 6,727,356. (The disclosures of each of which are incorporated herein by reference in their entireties.) For example, MGB-oligonucleotide conjugates can be synthesized using automated oligonucleotide synthesis methods from solid supports having cleavable linkers. In other examples, MGB probes can be prepared from an MGB modified solid support substantially in accordance with the procedure of Lukhtanov et al. *Bioconjugate Chem.*, 7: 564-567 (1996). (The disclosure of which is also incorporated herein by reference in its entirety.) According to these methods, one or more MGB moieties can be attached at the 5'-end, the 3'-end and/or at any internal portion of the oligonucleotide.

[0168] The location of an MGB moiety within an MGB-oligonucleotide conjugate can affect the discriminatory properties of such a conjugate. An unpaired region within a duplex will likely result in changes in the shape of the minor groove in the vicinity of the mismatched base(s). Since MGBs fit best within the minor groove of a perfectly-matched DNA duplex, mismatches resulting in shape changes in the minor groove would reduce binding strength of an MGB to a region containing a mismatch. Hence, the ability of an MGB to stabilize such a hybrid would be decreased, thereby increasing the ability of an MGB-oligonucleotide conjugate to discriminate a mismatch from a perfectly-matched duplex. On the other hand, if a mismatch lies outside of the region complementary to an MGB-oligonucleotide conjugate, discriminatory ability for un conjugated and MGB-conjugated oligonucleotides of equal length is expected to be approximately the same. Since the ability of an oligonucleotide probe to discriminate single base pair mismatches depends on its length, shorter oligonucleotides are more effective in discriminating mismatches. The first advantage of the use of MGB-oligonucleotides conjugates in this context lies in the fact that much shorter oligonucleotides compared to those used in the art (i.e., 20-mers or shorter), having greater discriminatory powers, can be used, due to the pronounced stabilizing effect of MGB conjugation. Consequently, larger delta T_ms of allele-specific blocker probes can improve AS-PCR assay specificity and selectivity.

[0169] Blocker probes having MGB at the 5' termini may have additional advantages over other blocker probes having a blocker moiety (e.g., MGB, PO₄, NH₂, PEG, or biotin) only at the 3' terminus. This is at least because blocker probes having MGB at the 5' terminus (in addition to a blocking moiety at the 3'-end that prevents extension) will not be cleaved during PCR amplification. Thus, the probe concentration can be maintained at a constant level throughout PCR, which may help maintain the effectiveness of blocking non-specific priming, thereby increasing cast-PCR assay specificity and selectivity (FIG. 3).

[0170] In some embodiments, as depicted in FIG. 4A, the allele-specific primer and/or the allele-specific blocker probe can comprise one or more modified nucleobases or nucleosidic bases different from the naturally occurring bases (i.e., adenine, cytosine, guanine, thymine and uracil). In some embodiments, the modified bases are still able to effectively hybridize to nucleic acid units that contain adenine, guanine, cytosine, uracil or thymine moieties. In some embodiments, the modified base(s) may increase the difference in the T_m between matched and mismatched target sequences and/or decrease mismatch priming efficiency, thereby improving not only assay specificity, but also selectivity.

[0171] Modified bases are considered to be those that differ from the naturally-occurring bases by addition or deletion of one or more functional groups, differences in the heterocyclic ring structure (i.e., substitution of carbon for a heteroatom, or vice versa), and/or attachment of one or more linker arm structures to the base. In some embodiments, all tautomeric forms of naturally occurring bases, modified bases and base analogues may also be included in the oligonucleotide primers and probes of the invention.

[0172] Some examples of modified base(s) may include, for example, the general class of base analogues 7-deazapurines and their derivatives and pyrazolopyrimidines and their derivatives (described in PCT WO 90/14353; and U.S. application Ser. No. 09/054,630, the disclosures of each of which are incorporated herein by reference in their entireties). Examples of base analogues of this type include, for example, the guanine analogue 6-amino-1H-pyrazolo[3,4-d]pyrimidin-4(5H)-one (ppG), the adenine analogue 4-amino-1H-pyrazolo[3,4-d]pyrimidine (ppA), and the xanthine analogue 1H-pyrazolo[4,4-d]pyrimidin-4(5H)-6(7H)-dione (ppX). These base analogues, when present in an oligonucleotide of some embodiments of this invention, strengthen hybridization and can improve mismatch discrimination.

[0173] Additionally, in some embodiments, modified sugars or sugar analogues can be present in one or more of the nucleotide subunits of an oligonucleotide conjugate in accordance with the invention. Sugar modifications include, but are not limited to, attachment of substituents to the 2', 3' and/or 4' carbon atom of the sugar, different epimeric forms of the sugar, differences in the α - or β -configuration of the glycosidic bond, and other anomeric changes. Sugar moieties include, but are not limited to, pentose, deoxypentose, hexose, deoxyhexose, ribose, deoxyribose, glucose, arabinose, pentofuranose, xylose, lyxose, and cyclopentyl.

[0174] Locked nucleic acid (LNA)-type modifications, for example, typically involve alterations to the pentose sugar of ribo- and deoxyribonucleotides that constrains, or "locks," the sugar in the N-type conformation seen in A-form DNA. In some embodiments, this lock can be achieved via a 2'-O, 4'-C methylene linkage in 1,2:5,6-di-O-isopropylene- α -D-allofuranose. In other embodiments, this alteration then serves as the foundation for synthesizing locked nucleotide phosphoramidite monomers. (See, for example, Wengel J., *Acc. Chem. Res.*, 32:301-310 (1998), U.S. Pat. No. 7,060,809; Obika, et al., *Tetrahedron Lett* 39: 5401-5405 (1998); Singh, et al., *Chem Commun* 4:455-456 (1998); Koshkin, et al., *Tetrahedron* 54: 3607-3630 (1998), the disclosures of each of which are incorporated herein by reference in their entireties.)

[0175] In some preferred embodiments, the modified bases include 8-Aza-7-deaza-dA (ppA), 8-Aza-7-deaza-dG (ppG), 2'-Deoxypseudoisocytidine (iso dC), 5-fluoro-2'-deoxyuridine (fdU), locked nucleic acid (LNA), or 2'-O,4'-C-ethylene

bridged nucleic acid (ENA) bases. Other examples of modified bases that can be used in the invention are depicted in FIG. 4B and described in U.S. Pat. No. 7,517,978 (the disclosure of which is incorporated herein by reference in its entirety).

[0176] Many modified bases, including for example, LNA, ppA, ppG, 5-Fluoro-dU (fdU), are commercially available and can be used in oligonucleotide synthesis methods well known in the art. In some embodiments, synthesis of modified primers and probes can be carried out using standard chemical means also well known in the art. For example, in certain embodiments, the modified moiety or base can be introduced by use of (a) modified nucleoside as a DNA synthesis support, (b) modified nucleoside as a phosphoramidite, (c) reagent during DNA synthesis (e.g., benzylamine treatment of a convertible amidite when incorporated into a DNA sequence), or (d) by post-synthetic modification.

[0177] In some embodiments, the primers or probes are synthesized so that the modified bases are positioned at the 3' end. In some embodiments, the modified base are located between, 1-6 nucleotides, e.g., 2, 3, 4 or 5 nucleotides away from the 3'-end of the allele-specific primer or blocker probe. In some preferred embodiments, the primers or probes are synthesized so that the modified bases are positioned at the 3'-most end of the allele-specific primer or blocker probe.

[0178] Modified internucleotide linkages can also be present in oligonucleotide conjugates of the invention. Such modified linkages include, but are not limited to, peptide, phosphate, phosphodiester, phosphodiester, alkylphosphate, alkanephosphonate, thiophosphate, phosphorothioate, phosphorodithioate, methylphosphonate, phosphoramidate, substituted phosphoramidate and the like. Several further modifications of bases, sugars and/or internucleotide linkages, that are compatible with their use in oligonucleotides serving as probes and/or primers, will be apparent to those of skill in the art.

[0179] In addition, in some embodiments, the nucleotide units which are incorporated into the oligonucleotides of the allele-specific primers and/or blocker probes of the present invention may have a cross-linking function (an alkylating agent) covalently bound to one or more of the bases, through a linking arm.

[0180] In some embodiments of the methods and kits, the first allele-specific blocker probe binds to the same strand or sequence as the first allele-specific primer, while the second allele-specific blocker probe binds to the opposite strand and/or complementary sequence as the first allele-specific primer.

[0181] Detector Probes

[0182] In some embodiments, detector probe is designed as short oligomers ranging from about 15-30 nucleotides, such as about 16, about 18, about 22, about 24, about 30, or any number in between. In some embodiments, the T_m of the detector probe ranges from about 60° C. to 70° C., about 61° C. to 69° C., about 62° C. to 68° C., about 63° C. to 67° C., or about 64° C. to 66° C., or any temperature in between.

[0183] In some embodiments, the detector probe is a locus-specific detector probes (LST). In other embodiments the detector probe is a 5' nuclease probe. In some exemplary embodiments, the detector probe can comprises an MGB moiety, a reporter moiety (e.g., FAMTM, TETTM, JOETM, VICTM, or SYBR® Green), a quencher moiety (e.g., Black Hole QuencherTM or TAMRATM), and/or a passive reference (e.g., ROXTM). In some exemplary embodiments, the detector probe is designed according to the methods and principles

described in U.S. Pat. No. 6,727,356 (the disclosure of which is incorporated herein by reference in its entirety). In some exemplary embodiments, the detector probe is a TaqMan® probe (Applied Biosystems, Foster City). In exemplary embodiments, the locus-specific detector probe can be designed according to the principles and methods described in U.S. Pat. No. 6,727,356 (the disclosure of which is incorporated herein by reference in its entirety). For example, fluorogenic probes can be prepared with a quencher at the 3' terminus of a single DNA strand and a fluorophore at the 5' terminus. In such an example, the 5'-nuclease activity of a Tag DNA polymerase can cleave the DNA strand, thereby separating the fluorophore from the quencher and releasing the fluorescent signal. In some embodiments, the detector probes are hybridized to the template strands during primer extension step of PCR amplification (e.g., at 60-65° C.). In yet other embodiments, an MGB is covalently attached to the quencher moiety of the locus-specific detector probes (e.g., through a linker).

[0184] In some embodiments of the disclosed methods and kits, the first and second detector probes are the same and/or comprise the same sequence or are the same sequence.

[0185] Locus-Specific Primers

[0186] In some embodiments, locus-specific primer (LSP) is designed as a short oligomer ranging from about 15-30 nucleotides, such as about 16, about 18, about 22, about 24, about 30, or any number in between. In some embodiments, the T_m of the locus-specific primer ranges from about 60° C. to 70° C., about 61° C. to 69° C., about 62° C. to 68° C., about 63° C. to 67° C., or about 64° C. to 66° C., or any range in between.

[0187] In some other embodiments of the disclosed methods and kits, the first locus-specific detector probe and/or second locus-specific detector probes comprise the same sequence or are the same sequence.

[0188] Additional Components

[0189] Polymerase enzymes suitable for the practice of the present invention are well known in the art and can be derived from a number of sources. Thermostable polymerases may be obtained, for example, from a variety of thermophilic bacteria that are commercially available (for example, from American Type Culture Collection, Rockville, Md.) using methods that are well-known to one of ordinary skill in the art (See, e.g., U.S. Pat. No. 6,245,533). Bacterial cells may be grown according to standard microbiological techniques, using culture media and incubation conditions suitable for growing active cultures of the particular species that are well-known to one of ordinary skill in the art (See, e.g., Brock, T. D., and Freeze, H., J. Bacterial. 98(1):289-297 (1969); Oshima, T., and Imahori, K, Int. J. Syst. Bacteriol. 24(1):102-112 (1974)). Suitable for use as sources of thermostable polymerases are the thermophilic bacteria *Thermus aquaticus*, *Thermus thermophilus*, *Thermococcus litoralis*, *Pyrococcus furiosus*, *Pyrococcus woosii* and other species of the *Pyrococcus* genus, *Bacillus stearothermophilus*, *Sulfolobus acidocaldarius*, *Thermoplasma acidophilum*, *Thermus flavus*, *Thermus ruber*, *Thermus brockianus*, *Thermotoga neapolitana*, *Thermotoga maritima* and other species of the *Thermotoga* genus, and *Methanobacterium thermoautotrophicum*, and mutants of each of these species. Preferable thermostable polymerases can include, but are not limited to, Tag DNA polymerase, The DNA polymerase, Tma DNA polymerase, or mutants, derivatives or fragments thereof.

[0190] Various Sources and/or Preparation Methods of Nucleic Acids

[0191] Sources of nucleic acid samples in the disclosed compositions, methods and/or kits include, but are not limited to, human cells such as circulating blood, buccal epithelial cells, cultured cells and tumor cells. Also other mammalian tissue, blood and cultured cells are suitable sources of template nucleic acids. In addition, viruses, bacteriophage, bacteria, fungi and other micro-organisms can be the source of nucleic acid for analysis. The DNA may be genomic or it may be cloned in plasmids, bacteriophage, bacterial artificial chromosomes (BACs), yeast artificial chromosomes (YACs) or other vectors. RNA may be isolated directly from the relevant cells or it may be produced by in vitro priming from a suitable RNA promoter or by in vitro transcription. The present invention may be used for the detection of variation in genomic DNA whether human, animal or other. It finds particular use in the analysis of inherited or acquired diseases or disorders. A particular use is in the detection of inherited diseases.

[0192] In some embodiments, template sequence or nucleic acid sample can be gDNA. In other embodiments, the template sequence or nucleic acid sample can be cDNA. In yet other embodiments, as in the case of simultaneous analysis of gene expression by RT-PCR, the template sequence or nucleic acid sample can be RNA. The DNA or RNA template sequence or nucleic acid sample can be extracted from any type of tissue including, for example, formalin-fixed paraffin-embedded tumor specimens.

[0193] Pre-Amplification

[0194] In another aspect, additional compositions, methods and kits are provided for "boosting" or pre-amplifying cast-PCR amplification reactions. In some embodiments, pre-amplification of cast-PCR reactions is employed for limited quantity specimens having very low nucleic acid copy number.

[0195] In some embodiments, said "boosting" compositions, methods and kits involve a two-step amplification process comprising a first pre-amplification reaction (see, for example, U.S. Pat. Nos. 6,605,451 and 7,087,414 and U.S. Published Application No. 2004/0175733, the disclosures of which are herein incorporated by reference in their entireties), followed by a second amplification (i.e., cast-PCR) reaction.

[0196] In some embodiments, pre-amplification composition used in cast-PCR pre-amplification reactions comprise:

[0197] i) a nucleic acid sample; and

[0198] ii) at least two sets of primers wherein each set comprises (a) a first allele-specific primer and a second allele-specific primer, wherein the allele-specific nucleotide portion of said first and second allele-specific primers is complementary to a first allele and a second allele, respectively, of a given SNP within a target sequence; and (b) a locus-specific primer that is complementary to a region of said target sequence that is 3' from the first and second alleles and on the opposite strand; wherein the at least two sets of primers is each specific for a different target sequence.

[0199] In some embodiments, the cast-PCR pre-amplification methods comprise:

[0200] a) forming in a single vessel a first reaction mixture comprising:

[0201] i) a nucleic acid sample; and

[0202] ii) at least two sets of primers wherein each set comprises (a) a first allele-specific primer and a second allele-specific primer, wherein the allele-specific

nucleotide portion of said first and second allele-specific primers is complementary to a first allele and a second allele, respectively, of a given SNP within a target sequence; and (b) a locus-specific primer that is complementary to a region of said target sequence that is 3' from the first and second alleles and on the opposite strand; wherein the at least two sets of primers is each specific for a different target sequence;

[0203] b) amplifying said different target sequences using a number of cycles suitable to keep the reaction within a linear phase;

[0204] c) dividing the amplified products of step (b) into at least two separate vessels;

[0205] d) adding to each of the divided products of step (c)

[0206] i) at least one set of primers used in step (a),

[0207] ii) an allele-specific blocker probe that is complementary to a region of said target sequence comprising a second allelic variant, wherein said region encompasses a position corresponding to the binding position of the allele-specific nucleotide portion of the first allele-specific primer, and wherein the first allele-specific blocker probe comprises a minor groove binder; and

[0208] iv) a detector probe

[0209] to form a second and a third reaction mixture in the at least two separate vessels;

[0210] e) amplifying said target sequence in said second and third reaction mixtures; and

[0211] f) detecting a change in a detectable property of the detector probe in each of the amplified products of said target sequence produced by step (e), thereby detecting the allelic variant of the target gene in the nucleic acid sample.

[0212] In some embodiments, the first reaction is a multiplex reaction and said second reaction is a single-plex reaction. In some preferred embodiments, the multiplex reaction involves the use of at least two complete sets of primers (e.g., a set comprises one forward allele-1-specific primer, one forward allele-2-specific primer and one reverse locus-specific primer), each set of which is suitable or operative for amplifying a specific polynucleotide of interest. In other embodiments, the resultant multiplex products acquired in the first step are divided into optimized secondary single-plex cast-PCR amplification reactions, each containing at least one primer set previously used in the first multiplexing step and then PCR amplified using the cast-PCR methods described herein.

[0213] In other preferred embodiments, the first multiplex reaction is a cast-PCR amplification reaction (although other well known amplification methods such as, but not limited to PCR, RT-PCR, NASBA, SDA, TMA, CRCA, Ligase Chain Reaction, etc. can be used). In certain embodiments, the first multiplex reaction comprises a plurality of allele-specific primers, and locus-specific primers, each group of which is specific for a particular allele of interest and designed according to the cast-PCR methods described herein. Unlike single-plex cast-PCR reactions that generate a single amplified sequence, multiplex cast-PCR amplification reactions, by virtue of utilizing a plurality of different primer sets, can permit the simultaneous amplification of a plurality of different sequences of interest in a single reaction. Because a plurality of different sequences is amplified simultaneously in a single reaction, the multiplex amplifications can effectively increase

the concentration or quantity of a sample available for downstream cast-PCR assays. Thus, in some preferred embodiments, significantly more analyses or assays can be performed with a pre-amplified cast-PCR sample than could have been performed with the original sample.

[0214] The number of different amplification primer pairs utilized in the multiplex amplification is not critical and can range from as few as two, to as many as tens, hundreds, thousands, or even more. Thus, depending upon the particular conditions, the multiplex amplifications permit the simultaneous amplification of from as few as two to as many as tens, hundreds, thousands, or even more polynucleotide sequences of interest.

[0215] The number of amplification cycles performed with a multiplex amplification may depend upon, among other factors, the degree of amplification desired. The degree of amplification desired, in turn, may depend upon such factors as the amount of polynucleotide sample to be amplified or the number of alleles or mutations to be detected using subsequent cast-PCR assays.

[0216] In preferred embodiments, it may be desirable to keep the multiplex amplification from progressing beyond the exponential phase or the linear phase. Indeed, in some embodiments, it may be desirable to carry out the multiplex amplification for a number of cycles suitable to keep the reaction within the exponential or linear phase. Utilization of a truncated multiplex amplification round can result in a sample having a boosted product copy number of about 100-1000 fold increase.

[0217] In many embodiments, pre-amplification permits the ability to perform cast-PCR assays or analyses that require more sample, or a higher concentration of sample, than was originally available. For example, after a 10x, 100x, 1000x, 10,000x, and so on, multiplex amplification, subsequent cast-PCR single-plex assays can then be performed using, respectively, a 10x, 100x, 1000x, 10,000x, and so on, less sample volume. In some embodiments, this allows each single-plex cast-PCR reaction to be optimized for maximum sensitivity and requires only one method of detection for each allele analyzed. This can be a significant benefit to cast-PCR analysis since, in some embodiments, it allows for the use of off-the-shelf commercially available cast-PCR reagents and kits to be pooled together and used in a multiplex amplification reaction without extensive effort toward or constraints against redesigning and/or re-optimizing cast-PCR assays for any given target sequence. Moreover, in some embodiments, the ability to carry out a multiplex amplification with reagents and kits already optimized for cast-PCR analysis permits the creation of multiplex amplification reactions that are ideally correlated or matched with subsequent single-plex cast-PCR assays.

[0218] In one aspect, kits are provided that include at least one separate container comprising the following components independently distributed in at least one of the separate containers: (i) a first allele-specific primer, wherein an allele-specific nucleotide portion of the first allele-specific primer is complementary to the first allelic variant of a target sequence; (ii) a second allele-specific primer wherein an allele-specific nucleotide portion of the second allele-specific primer is complementary to the second allelic variant of a target sequence; and (iii) a common locus-specific primer.

[0219] The following examples are intended to illustrate but not limit the invention.

EXAMPLES

I. General Cast-PCR Assay Design

[0220] The general schema for the cast-PCR assays used in the following examples is illustrated in FIG. 1. For each SNP that was analyzed, allele-specific primers (ASPs) were designed to target a first allele (i.e. allele-1) and a second allele (i.e. allele-2). The cast-PCR assay reaction mixture for allele-1 analysis included a 5'-tailed allele-1-specific primer (ASP1), one MGB allele-2 blocker probe (MGB2), one common locus-specific TaqMan probe (LST) and one common locus-specific primer (LSP). The cast-PCR assay reaction mixture for analysis of allele-2 included a 5'-tailed allele-2-specific primer (ASP2), one MGB allele-1 blocker probe (MGB1), one common locus-specific TaqMan probe (LST) and one common locus-specific primer (LSP).

II. Reaction Conditions

[0221] Each assay reaction mixture (10 μ l total) contained 1 $\times$ TaqMan Genotyping Master Mixture (Applied Biosystems, Foster City, Calif.; P/N 437135), 0.5 ng/ μ l genomic DNA or 1 million copies of plasmid DNA (or as indicated otherwise), 300 nM (unless specified otherwise) tailed-, or in some cases untail-, allele-specific primer (ASP1 for detection of allele-1 or ASP2 for detection of allele-2), 200 nM TaqMan probe (LST), 900 nM locus-specific primer (LSP), 150 nM allele-specific MGB blocker probe (MGB1 for detection of allele-2 or MGB2 for detection of allele-1). The reactions were incubated in a 384-well plate at 95° C. for 10 minutes, then for 5 cycles at 95° C. for 15 seconds and 58° C. for 1 minute, then by 45 cycles at 95° C. for 15 seconds and 60° C. for 1 minute. All reactions were run in duplicate or higher replication in an ABI PRISM 7900HT® Sequence Detection System, according to the manufacturer's instructions.

[0222] The 2-stage cycling protocol used in the following examples for cast-PCR amplification reactions is different from conventional allele-specific PCR (AS-PCR). The 2-stage cycling protocol comprises an initial 5 cycles at a lower annealing/extension temperature (e.g., 58° C.), followed by 45 standard cycles at a higher annealing/extension temperature (e.g., 60° C.). Due to the lower T_m of cast-PCR allele-specific primers (e.g., 53-56° C.), PCR is not optimal at standard annealing/extension conditions (e.g., 60° C.). Consequently, lower annealing/extension temperatures used during the initial 5 cycles increases overall cast-PCR efficiency.

III. Nucleic Acid Samples

[0223] Plasmids containing specific SNP sequences were designed and ordered from BlueHeron (Bothell, Wash.). (See Table 1 for a list of plasmids comprising SNPs used in some of the following examples.) The plasmids were quantified using TaqMan RNase P Assay (Applied Biosystems, Foster City, Calif.; P/N 4316838) according to the manufacturer's instructions and were used as templates (See Table 1, RNase P Control) to validate sensitivity, linear dynamic range, specificity, and selectivity of the given assays.

[0224] Genomic DNAs were purchased from Coriell Institute for Medical Research (Camden, N.J.; NA17203, NA17129, NA17201). The genotypes of target SNPs were validated with TaqMan SNP Genotyping Assays (Applied Biosystems, Foster City, Calif.; P/N 4332856) according to the manufacturer's instructions.

IV. Modified Oligonucleotides

[0225] Modified bases were purchased from Berry and Associates (ppA: P/N BA 0239; ppG: P/N BA 0242; fdU: P/N BA 0246; and iso dC: P/N BA 0236) or Exicon (LNA-T Amidite: P/N EQ-0064; LNA-mC Amidite: P/N EQ-0066; LNA-G Amidite: P/N EQ-0082; and LNA-A Amidite: P/N EQ-0063). Oligonucleotides comprising the modified nucleotides at their 3' ends were synthesized according to the manufacturer's instructions.

TABLE 1

Plasmid SNP Sequences (target alleles are indicated in brackets).	
SNP ID	Sequence
CV11201742	GCTCTGCTTCATTCGTCTGAAGAAGGGCAGATAGTTTGGCTGCTCCTGTG[C/T]TGTCACCTGCAATTCCTCC CTTATCAGGGCCATTGGCCTCTCCCTTCTCTCTGTGAGGGATATTTCTCTGACTTGTCAATCCACATCTTCC
CV11349123	GGCTTGCAATGGCTCCAACCGAAGGGCGGTGCTCGAGCTGTGGTGCCTGC[C/T]GCTAAGTTGTGCGTTCCA GGGTGCACTCGC
CV1207700	GCAACTATACCCCTTGATGGATGGAGATT[A/C]GCAATGTGTTTTACTGGGTAGAGTGACAGACCTT
CV25594064	CCTGAACTTATTTGGCAAGCGATGAGTACTCTTAAATTAATCTGGAATATATATTATTAGAACTGACC AATTACCTAGATCCCCCT[C/G]AACAATTGTTTACCAAGGAACCTCCTGAA
CV25639181	GAATTGGTTGTCTCCTTATGGGAAGTGAAGTATTTTACAC[G/T]CTTTACCACATTTCTTCATGGGATAGTAA GTGTTAAACAGCTCTGAGCCATTATTATCAGCTACTTGTAATATAGCAGTAGAATTTTATTTTATACCTGTAA GTGGGCAGTTACCTTTTGAGAGGAATACCTATAG
RNaseP Control	GCGGAGGGAAGCTCATCAGTGGGGCCACGAGCTGAGTGCCTCCTGCTCACTCCACTCCCATGTCCCTTGGGAAG GTCTGAGACTAGGG
BRAF-1799TA	TACTACACCTCAGATATATTTCTTCATGAAGACCTCACAGTAAAAATAGGTGATTTTGGTCTAGCTACAG [T/A]GAAATCTCGATGGAGTGGGTCCTCAGTTTGAACAGTTGTCTGGATCCATTTTGTGGATGGTAAGAATT GAGGCTATTTTCCACTGATTAAATTTTGGCCCTGAGATGCTGCTGAGTT

TABLE 1-continued

Plasmid SNP Sequences (target alleles are indicated in brackets).	
SNP ID	Sequence
CTNNB1-121AG	TGCTAATACTGTTTTCGTATTATAGCTGATTGATGGAGTTGGACATGGCCATGGAACCAGACAGAAAAGCGG CTGTTAGTCACTGGCAGCAACAGTCTTACCTGGACTCTGGAATCCATTCTGGTGCCACT[A/G]CCACAGCTCCT TCTCTGAGTGGTAAGGCAATCCTGAGGAAGAGGATGTGGATACCTCCCAAGTC
CTNNB1-134-CT	TTTGTATGGAGTTGGACATGGCCATGGAACCAGACAGAAAAGCGGCTGTTAGTCACTGGCAGCAACAGTCTTAC CTGGACTCTGGAATCCATTCTGGTGCCACTACCACAGCTCCTT[C/T]TCTGAGTGGTAAGGCAATCCTGAGG AAGAGGATGTGGATACCTCCCAAGTCTGTATGATGGGAA
EGFR-2369CT	GTGGACAACCCCCACGTGTGCGCCTGCTGGGCATCTGCCTCACCTCCACCGTGCAGCTCATCA[C/T]GCAGC TCATGCCCTTCGGCTGCCTCCTGGACTATGTCCGGGAACACAAAGACAATATTGGCTCCCAGTACCTGCTCAAC TGGTGTGTGCAGATCGCAAAGGTAATCAGGGAAGGGA
EGFR-2573TG	GCATGAACACTTGTGAGGACCGTCGCTTGGTGACCCGCGACCTGGCAGCCAGGAACGTACTGGTGAAAACACC GCAGCATGTCAAGATCACAGATTTTGGGC[T/G]GGCCAACTGCTGGGTGCGGAAGAGAAAGAATACCATGC AGAAGGAGGCCAAAGTAAGGAGGTG
KRAS-176CG	CAGGATTCTACAGGAAGCAAGTAGTAATTGATGGAGAAACCTGTCTCTTGGATATTCTCGACACAG[C/G]AG GTCAGAGGAGTACAGTGCAATGAGGACCAGTACATGAGGACTGGGGAGGGCTTCTTTGTGTATTGTCAT AAATAACTAAATCATTTGAAGATATT
KRAS-183AC	ACAGGAAGCAAGTAGTAATTGATGGAGAAACCTGTCTCTTGGATATTCTCGACACAGCAGGTCA[A/C]GAGG AGTACAGTGCAATGAGGACCAGTACATGAGGACTGGGGAGGGCTTCTTTGTGTATTGCCATAAATAATAC TAAATCATTTGAAGATATTACCATTTATAGTGGGTTAAATTGAATATAATAAGCTGACATTAA
KRAS-34GA	TATTAACCTTATGTGTGACATGTTCTAATATAGTCACATTTTCATTATTTTATTATAAGGCCTGCTGAAAATGA CTGAATATAAACTTGTGGTAGTTGGAGCT[G/A]GTGGCGTAGGCAAGAGTGCCTTGACGATACAGCTAATTCA GAATCATTTTGTGGACGAATATGA
KRAS-35GA	TATTAACCTTATGTGTGACATGTTCTAATATAGTCACATTTTCATTATTTTATTATAAGGCCTGCTGAAAATGA CTGAATATAAACTTGTGGTAGTTGGAGCTG[G/A]TGGCGTAGGCAAGAGTGCCTTGACGATACAGCTAATTCA GAATCATTTTGTGGACGAATATGATC
KRAS-38GA	CATTATTTTATTATAAGGCCTGCTGAAAATGACTGAATATAAACTTGTGGTAGTTGGAGCTGGTG[G/A]CGTA GGCAAGAGTGCCCTTGACGATACAGCTAATTGAGAAATCATTTTGTGGACGAATATGATCCAAACAATAGAGGTAA ATCTTGTTTAAATATGCATATTACTGGTGCAGGACCATTTCTTGATACAGATAAAGGTTTCTCTGACCATTTTCA TGAGTACTTAT
NRAS-181CA	ATTCTTACAGAAAACAAGTGGTTATAGATGGTGAAACCTGTTTGTGTGGACATACTGGATACAGCTGGA[C/A]A AGAAGAGTACAGTGCCATGAGAGACCAATACATGAGGACAGGCGAAGGCTTCTCTGTGTATTGTCATCAAT AATAGCAAGTCATTTGCGGATATTAACCTCTACAGGTACTAGGAGCATTATTTCTCTGAAAGGATG
NRAS-183AT	TTACAGAAAACAAGTGGTTATAGATGGTGAAACCTGTTTGTGTGGACATACTGGATACAGCTGGACA[A/T]GAA GAGTACAGTGCCATGAGAGACCAATACATGAGGACAGGCGAAGGCTTCTCTGTGTATTGTCATCAATAATA GCAAGTCATTTGCGGATATTAACCTCTACAGGTACTAGGAGCATTATTTCTCTGAAAGGATG
NRAS-35GA	TGGTTTCCAACAGGTTCTTGCTGGTGTGAAATGACTGAGTACAACTGGTGGTGGTTGGAGCAG[G/A]TGGTG TTGGGAAAAGCGCACTGACAATCCAGCTAATCCAGAACCCTTGTAGATGAATATGATCCCACCATAGAGGT GAGGCCAGTGGTAGCCCG
NRAS-38GA	TTTCCAACAGGTTCTTGCTGGTGTGAAATGACTGAGTACAACTGGTGGTGGTTGGAGCAGGTG[G/A]TGTTG GGAAAAGCGCACTGACAATCCAGCTAATCCAGAACCCTTGTAGATGAATATGATCCCACCATAGAGGTGAG GCCAGTGGTAGCCC
TP53-524GA	GGCACCCGCGTCCGCGCCATGGCCATCTACAAGCAGTCAAGCACATGACGGAGGTTGTGAGGC[G/A]CTGC CCCCACCATGAGCGCTGCTCAGATAGCGATGGTGAGCAGCTGGGGCTGGAGAGACGACAGGGCTGGTTGCCCA GGGTCCCCAGGCCCTGTGATTCCTCACTGATTGCTCTTAGTCTGGCC
TP53-637CT	CCTCTCAGCATCTTATCCGAGTGGAAAGGAAATTGCGTGTGGAGTATTGGATGACAGAAACACTTTT[C/T] GACATAGTGTGGTGGTCCCTATGAGCCGCTGAGGTCTGGTTTGCACTGGGGTCTCTGGGAGGAGGGGTTA AGGGTGGTTGTCAGTGGCCCTC
TP53-721TG	CTTGGGCTGTGTTATCTCCTAGGTTGGCTCTGACTGTACCACCATCCACTACAACATACATGTGTAACAGT[T/G] CCTGCATGGGCGGCATGAACCGGAGGCCATCCTCACCATCATCACTGGAAGACTCCAGGTGAGGAGCCA CTTGCCACCTGCACACTGGCCTGCTGTGCCCCAGCCTC
TP53-733GA	TAGGTTGGCTCTGACTGTACCACCATCCACTACAACATACATGTGTAACAGTTCTGTCATGGGC[G/A]GCATGA ACCGGAGGCCATCCTCACCATCATCACTGGAAGACTCCAGGTGAGGAGCCACTTGCCACCTGCACACTG GCTGCTGTGCCCCAGCCTC

TABLE 1-continued

Plasmid SNP Sequences (target alleles are indicated in brackets).	
SNP ID	Sequence
TP53-742CT	CTGACTGTACCACCATCCACTACAACACTACATGTGTAACAGTTCTTCATGGGCGGCATGAAC[C/T]GGAGGCC CATCCTCACCATCATCACACTGGAAGACTCCAGGTCAGGAGCCACTTGCCACCCCTGCACACTGGCCTGCTGTGC CCCAGCCTCTGCTTGCCCTC
TP53-743GA	TGACTGTACCACCATCCACTACAACACTACATGTGTAACAGTTCTTCATGGGCGGCATGAACC[G/A]GAGGCC ATCCTCACCATCATCACACTGGAAGACTCCAGGTCAGGAGCCACTTGCCACCCCTGCACACTGGCCTGCTGTGCC CCAGCCTCTGCTTGCCCTC
TP53-817CT	CCTCTTGCTTCTCTTTTCTATCCTGAGTAGTGGTAATCTACTGGGACGGAACAGCTTTGAGGTG[C/T]GTGTTT GTGCTGTCTGCGAGAGACCGGCGCACAGAGGAAGAGAATCTCCGCAAGAAAGGGGAGCCTCACCACGAGC TGCCCCAGGGAGCACTAAGCGAGGTAAGCAA

[0226] Data Analysis:

[0227] An automatic baseline and manual threshold of 0.2 were used to calculate the threshold cycle (C_t) which is defined as the fractional cycle number at which the fluorescence passes the fixed threshold. PCR reactions were run for a total of 50 cycles. For cast-PCR reactions, there was a pre-run of five cycles at a lower annealing/extension temperature followed by an additional 45 cycles at a higher annealing/extension temperature. The ΔC_t between amplification reactions for matched vs. mismatched sequences is defined as the specificity of cast-PCR ($\Delta C_t = C_{t_{mismatch}} - C_{t_{match}}$). The larger the ΔC_t between mismatched and matched targets, the better assay specificity. The $2^{\Delta C_t}$ value was used to estimate the power of discrimination (or selectivity) which is equal to $1/2^{\Delta C_t}$ or, in some cases, calculated as % ($1/2^{\Delta C_t} \times 100$).

Example 1

Tailed Primers Improve Discrimination of Allelic Variants

[0228] The following example demonstrates that the application of allele-specific primers comprising tails significantly improves the discrimination of allelic variants.

[0229] In conventional AS-PCR, the discrimination of 3' nucleotide mismatches is largely dependent on the sequence surrounding the SNP and the nature of the allele. The ΔC_t between the amplification reactions for matched and mismatched primers varies. To improve the discrimination between the amplification of matched and mismatched sequences, allele-specific primers were designed to comprise tails at their 5' termini and then tested for their suitability in AS-PCR assays.

[0230] Assays were performed using the general experimental design and reaction conditions indicated above (with the exception that no blocker probes were included and either tailed or non-tailed allele-specific primers were added), using 0.5 ng/uL genomic DNA containing the hsv 11711720 SNP comprising one of three alleles (A, C, or T) as the nucleic acid template (see Table 2A). The three genotypes are indicated in Table 2B. Primers and probes were designed according to the sequences shown in Table 3.

TABLE 2A

Genomic DNA Sequence for hsv11711720 SNP (target alleles are indicated in brackets).	
AGAAAAATAACTAAGGGAAGGAGAAAGTGGGAGGAAGGAACAGTGT	
GAAGACAATGGCCTGAAAACTGAAAAAGTCTGTTAAAGTTAATTATCAGT	
TTTTGAGTCCAAGAACTGGCTTTGCTACTTTCTGTAAGTTCTAATTAC	
TGAATAAGCATGAAAAAGATTGCTTTGAGGAATGGTTATAAACACATTCT	
TAGAGCATAGTAAGCAGTAGGGAGTAACAAAATAACACTGATTAGAATAC	
TTTACTCTACTTAATTAATCAATCATATTTAGTTTGACTCACCTTCCCAG	
[A/C/T]ACCTTCTAGTTCTTTCTTATCTTTCAGTGCTTGCCAGACAAC	
ATTTTCATTTCAACAACCTCTGCTATTGCAATGATGGGTACAATTGCTAA	
GAGTAACAGTGTTAGTTGCCAACCATAGATGAAGGATATAATTATTCCTG	
TCCAAGATTGCTATATTCTGGTAATTACAGCAAGCCTGGAACCTATA	
GCCTGCAAAACAAAACAAATTAGAGAAATTTAAAAATATTATCTTCACA	
ACTCATGCTTCTATTTTCTGAAACTCACCTTCATGAGACTATATTCATT	
ATTTTAT	

TABLE 2B

Genotypes of Genomic DNA Sequence for hsv11711720 SNP	
Genomic DNA ID	Genotype
NA17203	AA
NA17129	CC
NA17201	TT

TABLE 3

List and Sequences of Primers and Probes for genomic DNA: conventional allele-specific primers ("ASP"); tailed allele-specific primers ("tailASP"); locus-specific TaqMan probe (LST); locus-specific primer (LSP). The nucleotides shown in lower case are the tailed portion of the primers. The nucleotide-specific portion of each allele-specific primer is at the 3'-most terminus of each primer (indicated in bold).		
Primer/ Probe ID	Sequence (5' to 3')	T _m (° C.)
17129-ASP	ATATTGAGTTTGACTCACCTTCCCAGC	63.2
17129- tailASP	acc ACTCACCTTTCCCAGC	63.0
17203-ASP	ATATTGAGTTTGACTCACCTTCCCAGA	62.0
17203- tailASP	acc ACTCACCTTTCCCAGA	63.7
17201-ASP	ATATTGAGTTTGACTCACCTTCCCAGT	62.2
17201- tailASP	acc ACTCACCTTTCCCAGT	64.0
LST	(6-FAM) -TGGACAAGCACTGAAAGA- (MGB)	67.4
LSP	GCAGGAGTTGTTGAAATGAAAATGTTG	62.5

[0231] As shown in Table 4, when using non-tailed ASPs ("ASP-tail"), the discrimination of 3' nucleotide mismatch is largely dependent on the nature of the allele, as a considerable range of ΔC_t values is observed depending on the identity of the 3'-terminal base. The range of ΔC_t values between matched and mismatched nucleotides ("NT") were from -0.1 to 10. However, with tailed ASPs (ASP+tail), the discrimination of 3' nucleotide mismatch was significantly improved. In fact, as Table 4 shows, the ΔC_t value between matched and mismatched nucleotides was consistently equal to or greater than 10 when tailed ASPs were used. The C_t values for amplification of matched sequences using tailed ASPs were comparable to those using conventional or non-tailed ASPs. These results indicate that tailed ASP, can improve the specificity of AS-PCR, but may not improve the sensitivity of detection.

TABLE 4

Tailed allele-specific primers ("ASP") significantly improve discrimination of allelic variants. The specificity ("fold difference") was calculated based on the difference between C_t values using tailed vs. untail primers ($2^{(\Delta C_t(ASP+tail) - \Delta C_t(ASP-tail))}$). The mismatched nucleotides of the 3' allele-specific nucleotide portion of the ASPs (+/- tail) and the target allele are also indicated ("NT mismatch").			
NT mismatch	ΔC_t (ASP - tail)	ΔC_t (ASP + tail)	Specificity Improvement (fold difference)
C-A	0.9	11.5	1552.1
C-T	1.2	11.5	1278.3
A-C	10.0	11.9	3.7
A-G	9.8	11.9	4.3
T-G	2.3	11.5	588.1
T-C	-0.1	11.5	3104.2
Average	4.0	11.6	1088.5

Example 2

Low Primer Concentrations Improve Discrimination of Allelic Variants

[0232] Assays were performed using the general experimental design and reaction conditions indicated above, in the presence of 1 million copies of plasmid DNA containing various SNP target sequences (see Table 1) and 200 nM or 800 nM tailed ASP (as indicated). Assay primers and probes were designed according to the sequences shown in FIG. 11A.

[0233] The effect of tailed ASP concentration on discrimination of allelic variants is summarized in Table 5. The ΔC_t between the amplification reactions for matched and mismatched primers demonstrate that lower tailed ASP concentrations improve discrimination of allelic variants.

TABLE 5

Assay Results Using Different Concentrations of Tailed Allele-specific Primers			
Plasmid SNP ID	ΔC_t (ASP @ 800 nM)	ΔC_t (ASP @ 200 nM)	Specificity Improvement (fold difference)
CV11201742	14.1	15.2	2.14
CV11349123	8.2	10	3.48
CV1207700	5.2	6.6	2.64
CV25594064	20.1	19.1	0.5
CV25639181	11.9	12.9	2
Average	12.6	13.44	2.14

Example 3

Primers Designed with Reduced T_ms Improves Discrimination of Allelic Variants

[0234] Assays were performed using the general experimental design and reaction conditions indicated above, in the presence of 1 million copies of plasmid DNA containing various SNP target sequences (see Table 1) using tailed ASP with a higher T_m (~57° C.) or tailed ASP with a lower T_m (~53° C.). Assay primers and probes were designed according to the sequences shown in FIG. 11B-E (see FIG. 11B for higher T_m ASP and FIG. 11C or lower T_m ASP).

[0235] The effect of allele-specific primer T_m on discrimination of allelic variants is summarized in Table 6. The ΔC_t of allele-specific primers with a lower T_m are significantly higher than those of allele-specific primers with a higher T_m. Allele-specific primers designed with reduced T_ms improved discrimination of allelic variants by as much as 118-fold in some cases or an average of about 13-fold difference.

TABLE 6

ΔC_t Values Using Tailed ASPs with Lower T _m (~53° C.) or with Higher T _m (~57° C.)			
Plasmid SNP ID	ΔC_t (ASP w/ T _m ~57° C.)	ΔC_t (ASP w/ T _m ~53° C.)	Specificity Improvement (fold difference)
BRAF-1799TA	12.2	19.1	118.9
CTNNB1-121AG	11.6	14.9	10.0
KRAS-176CG	18.8	22.5	13.1
NRAS-35GA	13.0	14.0	1
TP53-721TG	14.7	19.1	20.6
CTNNB1-134CT	8.6	14.1	44.8
EGFR-2369CT	9.7	10.7	2

TABLE 6-continued

Δ Ct Values Using Tailed ASPs with Lower T _m (~53° C.) or with Higher T _m (~57° C.)			
Plasmid SNP ID	Δ Ct (ASP w/ T _m ~57° C.)	Δ Ct (ASP w/ T _m ~53° C.)	Specificity Improvement (fold difference)
KRAS-183AC	22.2	23.1	1.8
NRAS-38GA	14.0	14.3	1.2
TP53-733GA	13.6	13.5	1.0
EGFR-2573TG	16.7	20.2	10.9
KRAS-34GA	14	14.8	1.8
KRAS-38GA	11.2	14.4	8.9
NRAS-181CA	24.0	27.1	8.6
TP53-742CT	9.1	8.0	0.5
KRAS-35GA	11.5	15.1	12.3
NRAS-183AT	23.6	22.7	0.5
TP53-524GA	11.4	13.5	4.6
TP53-637CT	11.4	14.4	7.8
TP53-743GA	10.1	13.2	8.4
TP53-817CT	13.6	13.9	1.2
Average	14.1	16.3	13.3

Example 4

Use of Blocker Probes Improves Discrimination of Allelic Variants

[0236] The following example illustrates that the use of MGB blocker probes improves the discrimination between 3' nucleotide mismatched and matched primers to target sequences in AS-PCR reactions.

[0237] Assays were performed using the general cast-PCR schema and reaction conditions indicated above, using 1 million copies of plasmid DNA containing various SNP target sequences (see Table 1) in the presence of MGB blocker probes or in the absence of MGB blocker probes. Assay primers and probes were designed according to the sequences shown in FIG. 11C-E.

[0238] To improve the selectivity of AS-PCR, blocker probes were synthesized to comprise an MGB group at their 3' terminus. (See, for example, Kutyavin, I. V., et al., Nucleic Acids Research, 2000, Vol. 28, No. 2: 655-661, U.S. Pat. Nos. 5,512,677; 5,419,966; 5,696,251; 5,585,481; 5,942,610 and 5,736,626.)

[0239] The results of cast-PCR using MGB blocker probes are summarized in Table 7. The Δ Ct between cast-PCR with MGB blocker probes is larger than that without MGB blocker probes. As shown, MGB blocker probes improve the discrimination of allelic variants.

TABLE 7

MGB Blocker Probes Improve Discrimination of Allelic Variants			
SNP ID	Δ Ct (no MGB blocker)	Δ Ct (+MGB blocker)	Specificity Improvement (fold difference)
BRAF-1799TA	11.4	14.9	11.5
CTNNB1-121AG	11.6	14.1	5.4
KRAS-176CG	17.8	20.9	9
NRAS-35GA	13.9	14.3	1.4
TP53-721TG	12.5	14.7	4.4
CTNNB1-134CT	6.7	10.2	11.6
EGFR-2369CT	7.7	10.1	5.3
KRAS-183AC	22.4	23	1.5
NRAS-38GA	14.5	14.6	1.1
TP53-733GA	13.2	14.4	2.3
EGFR-2573TG	18.2	21.8	11.6
KRAS-34GA	14.4	15.1	1.7
KRAS-38GA	11.9	15.1	1.7
NRAS-181CA	19.3	24.2	30.2
TP53-742CT	12.7	13.6	1.9
KRAS-35GA	11.0	13.7	6.5
NRAS-183AT	20.2	21.7	2.9
TP53-524GA	13.5	13.5	1
TP53-637CT	9.3	12.1	7.0
TP53-743GA	9.9	11.5	3.1
TP53-817CT	12.6	13.2	1.5
Average	13.6	15.5	6.0

Example 5

Primers Designed to Target Discriminating Bases Improves Discrimination of Allelic Variants

[0240] Assays were performed using the general cast-PCR schema and reaction conditions indicated above, in the presence of 1 million copies of plasmid DNA containing SNP target sequences (see Table 1). Assay primers and probes were designed according to the sequences shown in FIG. 11C-E.

[0241] According to the data summarized in Table 8, the discrimination of cast-PCR was dependent on the nature of the allele being analyzed. As Table 8 indicates, the Δ Ct between mismatched and matched sequences for allele-1 were different from Δ Ct between mismatched and matched sequences for allele-2. However, both A and G bases, as compared to a T base, were highly discriminating for allele-1 and allele-2 in all four SNPs examined.

TABLE 8

Primers Designed to Target Discriminating Bases Improve Discrimination of Allelic Variants								
SNP ID	ASP design		Allele	SNP allele-1		Allele	SNP allele-2	
	3' NT of ASP1	3' NT of ASP2		ΔC_t (Ct_mismatch – Ct_match)	Specificity (fold difference)		ΔC_t (Ct_mismatch – Ct_match)	Specificity (fold difference)
KRAS-38GA	G	A	C	13.4	10809	T	8.2	294
NRAS-181CA	C	A	G	27.5	189812531	T	9.8	891
NRAS-183AT	A	T	T	17.9	244589	A	23.4	11068835
TP53-742CT	C	T	G	12.3	5043	A	8.3	315

Example 6

Determination of the Sensitivity and Dynamic Range for Cast-PCR

[0242] In this example, the sensitivity and dynamic range of cast-PCR was determined by performing cast-PCR using various copy numbers of a target plasmid.

[0243] Assays were performed using the general cast-PCR schema and reaction conditions indicated above, using 1×10^0 (1 copy) to 1×10^7 copies of plasmid DNA containing the NRAS-181CA SNP target sequence (see Table 1). Assay primers and probes were designed according to the sequences shown in FIG. 11C-E.

[0244] As shown in FIG. 5, the use of tailed primers and MGB-blocker probes does not adversely affect the sensitivity of cast-PCR, as the sensitivity of cast-PCR is comparable to TaqMan assays which do not utilize tailed primers or blocker probes. Furthermore, FIG. 5 shows that the cast-PCR assay shows a linear dynamic range over at least 7 logs.

Example 7

Determination of the Specificity of Cast-PCR

[0245] In this example, the specificity of cast-PCR was determined by comparing the amplification of particular alleles of KRAS using either matched or mismatched ASPs to a given allele in the presence of their corresponding blocker probes.

[0246] Assays were performed using the general cast-PCR schema and reaction conditions indicated above, using 1×10^6 copies of plasmid DNA containing either one of two alleles of the KRAS-183AC SNP target sequence (see Table 1). Assay primers and probes were designed according to the sequences shown in FIG. 11C-E.

[0247] The left panel of FIG. 7 shows the an amplification plot of cast-PCR on allele-1 DNA using matched (A1) primers in the presence of A2 blocker probes or mismatched (A2) primers in the presence of A1 blocker probes. The right hand panel shows a similar experiment in which cast-PCR was performed on allele-2 DNA. As indicated in the data summary in FIG. 7, a robust ΔC_t values of over 20 were observed for cast-PCR on both alleles of KRAS-183AC tested. This corresponds to a specificity as determined by a calculation of $2^{\Delta C_t}$ of 9×10^6 , and 2×10^6 , respectively, for allele-1 and allele-2. Furthermore, a calculation of selectivity ($1/2^{\Delta C_t}$) indicates

that values of $1/1.1 \times 10^7$ and $1/5.0 \times 10^7$ are observed for allele-1 and allele-2, respectively.

Example 8

Cast-PCR is Able to Detect a Single Copy Mutant DNA in One Million Copies of Wild Type DNA

[0248] In this example, the selectivity of cast-PCR, i.e., the ability of cast-PCR to detect a rare mutant DNA in an excess of wild type DNA, was determined.

[0249] Assays were performed using the general cast-PCR schema and reaction conditions indicated above, using various copy numbers of mutant KRAS-183AC plasmid DNA (1 copy to 1×10^6 copies) mixed with 1×10^6 copies of wild type KRAS-183AC plasmid DNA (see Table 1). Assay primers and probes were designed according to the sequences shown in FIG. 11C-E, and cast-PCR reactions were performed using wild type or mutant allele-specific primers and the corresponding MGB blocker probes.

[0250] FIG. 8 shows that cast-PCR is able to detect as little as one copy of a mutant DNA sequence, even when surrounded by a million-fold excess of a wild type sequence.

Example 9

Selectivity of Cast-PCR in Discriminating Tumor Cell DNA from Normal Cell DNA

[0251] In this example, the selectivity of cast-PCR was determined by performing assays on samples in which various amounts of tumor cell genomic DNA were mixed with or "spiked" into genomic DNA from normal cells. DNA samples were extracted using QIAmp DNA Mini Prep Kits (Qiagen). Wild type DNA was extracted from the SW48 cell line and mutant DNA was extracted the H1573 cell line.

[0252] The mutant DNA contained the KRAS-G12A mutation (See FIG. 6). The percentage of tumor cell DNA in the spiked samples varied from 0.5 to 100%. cast-PCR was used to detect the presence of tumor cell DNA when present in these percentages.

[0253] Assays were performed using the general cast-PCR schema and reaction conditions indicated above, using 30 ng of gDNA per reaction. Assay primers and probes were designed according to the sequences corresponding to KRAS-G12A SNP ID, as shown in FIG. 18A-B.

[0254] As shown in FIG. 9, tumor cell DNA, even when present only at a level of 0.5% as compared to normal cell DNA, is easily detected using cast-PCR.

Example 10

Use of Cast-PCR to Detect Tumor Cells in Tumor Samples

[0255] In this example, cast-PCR was used to detect and determine the percentage of tumor cells in tumor samples. Various normal and tumor samples were obtained and assayed by cast-PCR for the presence of a number of SNPs associated with cancer as shown in FIG. 10.

[0256] Assays were performed using the general cast-PCR schema and reaction conditions indicated above, using 5 ng of gDNA or 1.5 ng cDNA derived from either normal or tumor samples. Assay primers and probes corresponding to the

[0259] Prior to conducting cast-PCR amplification, multiplex reactions were performed for 7 different KRAS mutations (see Table 9).

[0260] 10 μ L multiplex reactions were prepared in a single tube by combining into one reaction 45 nM of each allele-specific primer (including the allele-1-specific primer and the allele-2-specific primer) and 45 nM of each locus-specific primer, as listed in FIG. 18A-B for each of the seven different KRAS SNPs, 0.1 ng/ μ L genomic DNA, and 1 $\times$ Preamp Master Mix (Applied Biosystems, Foster City, Calif.; P/N 437135). The 10 μ L pre-amplification reactions were then incubated in an Applied Biosystems 9700 Thermocycler in a 96- or 384-well plate for 95° C. for 10 minutes, followed by 10 cycles of 95° C. for 15 seconds, 60° C. for 4 minutes, and 99.9° C. for 10 minutes, and then held at 4° C. Next, 190 μ L of 0.1 $\times$ TE pH 8.0 was added to each 10 μ L pre-amplification reaction (20 $\times$ dilution). The diluted pre-amplification reaction products were then directly used in subsequent cast-PCR reactions or stored at -20° C. for at least one week prior to use.

TABLE 9

KRAS SNP Sequences (target alleles are indicated in brackets).	
SNP ID	SNP Sequence
KRAS-G12A_GC	TGACTGAATATAAACTTGTGGTAGTTGGAGCTG[G/C]TGGCGTAGGCAAGAGTGCCTTGACGATACAGCTAAT TCAGAATCATTTTGTGGACGAATATGATCCAACAATAGAGGTAAATCTTGTTTAAATATGCATATTACTGGTGC AGGACCATTCTTTGATACA
KRAS-G12R_GC	TGACTGAATATAAACTTGTGGTAGTTGGAGCT[G/C]GTGGCGTAGGCAAGAGTGCCTTGACGATACAGCTAAT TCAGAATCATTTTGTGGACGAATATGATCCAACAATAGAGGTAAATCTTGTTTAAATATGCATATTACTGGTGC AGGACCATTCTTTGATACA
KRAS-G12D_GA	TGACTGAATATAAACTTGTGGTAGTTGGAGCTG[G/A]TGGCGTAGGCAAGAGTGCCTTGACGATACAGCTAAT TCAGAATCATTTTGTGGACGAATATGATCCAACAATAGAGGTAAATCTTGTTTAAATATGCATATTACTGGTGC AGGACCATTCTTTGATACA
KRAS-G12S_GA	TGACTGAATATAAACTTGTGGTAGTTGGAGCT[G/A]GTGGCGTAGGCAAGAGTGCCTTGACGATACAGCTAAT TCAGAATCATTTTGTGGACGAATATGATCCAACAATAGAGGTAAATCTTGTTTAAATATGCATATTACTGGTGC AGGACCATTCTTTGATACA
KRAS-G13D_GA	TGACTGAATATAAACTTGTGGTAGTTGGAGCTGGTG[G/A]CGTAGGCAAGAGTGCCTTGACGATACAGCTAAT TCAGAATCATTTTGTGGACGAATATGATCCAACAATAGAGGTAAATCTTGTTTAAATATGCATATTACTGGTGC AGGACCATTCTTTGATACA
KRAS-G12C_GT	GTGAGTTTGTATTAAAGGTACTGGTGGAGTATNNGATAGTGTATTAACTTATGTGTGACATGTTCTAATATA GTCACATTTTCATTATTTTATTATAAGGCCTGCTGAAAATGACTGAATATAAACTTGTGGTAGTTGGAGCT[G/ T]GTGGCGTAGGCAAGAGT

SNPs shown in FIG. 10 were designed according to the sequences as shown in FIG. 11C-E.

[0257] The results shown in FIG. 10 indicate that cast-PCR has a low false positive rate as indicated by the failure of cast-PCR to detect the presence of mutant cells in normal samples. In contrast, cast-PCR was able to provide a determination of the percentage of tumor cells in various tumor samples that ranged from just under 2% for a tumor sample containing the NRAS-183AT SNP to greater than 80% for a sample containing the CTNNB1-134CT SNP.

Example 11

Use of Pre-Amplification in Cast-PCR

[0258] The following example demonstrates that pre-amplification combined with cast-PCR methods enables detection of multiple alleles from a limited amount of genomic DNA template.

[0261] Following pre-amplification, the diluted pre-amplification products were aliquotted into single-plex cast-PCR reactions. Individual assays were performed for each of the 7 different KRAS mutations using the general experimental design and reaction conditions indicated above (see section II of Examples). 10 μ L, cast-PCR reactions were run for assays containing, as the nucleic acid template, either 1 μ L 20 $\times$ diluted pre-amplification reaction product (as prepared above) or, as a comparison, 0.07 ng genomic DNA. All assay primers and probes were designed according to the sequences shown in FIG. 18A-B.

[0262] As shown in FIG. 12, for assays without pre-amplification the average Δ Ct of the 7 tested KRAS mutations was 12.0, whereas for assays using pre-amplification the average Δ Ct was 17.0. Thus, the $\Delta\Delta$ Ct between the two gave an improvement of about 5, which is about a 32 fold increase in sensitivity, for the pre-amplified reactions over those without pre-amplification.

[0263] In an ideal situation (where PCR efficiency=100%), the copy number of the target gene increases about 1000 fold

in 10 cycles. Under these conditions, if the starting copy number in a pre-amplification reaction is 0.1 ng/ μ L (or approximately 33 copies/ μ L), then after 10 cycles the copy number increases to approximately 33,000 copies/ μ L. In the example above, the copy number of a 20 fold diluted pre-amplification product is estimated to be approximately 1,650 copies/ μ L. Therefore, after adding 1 μ L of 1,650 copies/ μ L diluted pre-amplification product into a cast-PCR reaction (final volume of 10 μ L), the concentration of the cast-PCR products is approximately 165 copies/ μ L. Based on this estimation, 10 μ L of pre-amplification products from 1 ng/ μ L genomic DNA can be diluted by as much as 200 fold, and still provide up to 2000 μ L of nucleic acid template for use in subsequent cast-PCR reactions.

Example 12

Effect of Tailed ASP on Cast-PCR Specificity

[0264] Assays were performed using the general cast-PCR schema and reaction conditions indicated above (see section II of Examples), using 1 million copies of plasmid DNA containing various SNP target sequences (see Table 10). Assay primers and probes were designed according to the sequences shown in FIG. 19A-D. For each SNP analyzed, the blocker probes, locus-specific probes and locus-specific primers were the same and only the allele-specific primers varied (e.g., tailed or non-tailed).

TABLE 10

Plasmid SNP Sequences (target alleles are indicated in brackets).	
SNP ID	Sequence
BRAF-1799TA_TP	ATATTTCTTCATGAAGACCTCACAGTAAAAATAGGTGATTTTGGTCTAGCTACAG[T/A]GAAATCTCGATGGAGTGGGTCCCATCAGTTTGAACAGTTGTCTGGATCCATTTTG
CTNNB1-121AG_TP	GTATCCACATCCTCTTCTCAGGATTGCCTTTACCACTCAGAGAAGGAGCTGTGG[A/G]AGTGGCACCAG AATGGATTNCAGAGTNCAGGTAAGACTGTTGCTGCCAGTGACTA
CTNNB1-134CT_TP	CAGGACTTGGGAGGTATCCACATCCTCTTCTCAGGATTGCCTTTACCACTCAGA[C/T]AAGGAGCTGTG GTAGTGGCACCAGAATGGATTNCAGAGTNCAGGTAAGACTGTTG
EGFR-2369CT_TP	CCCACGTGTGCCGCTGCTGGGCATCTGCCTCACCTCCACCGTGCAGCTCATCA[C/T]GCAGCTCATGCC CTTCCGGCTGCCTCCTGGACTATGTCCGGGAACACAAGACAATA
EGFR-2573TG_TP	TTACTTTGCCTCCTTCTGCATGGTATTCTTTCTCTTCCGCACCCAGCAGTTTGGCC[T/G]GCCCCAAATCT GTGATCTTGACATGCTGCGGTGTTTTCACCACTACGTTCTCTGG
KRAS-176CG_TP	AGGAAGCAAGTAGTAATTGATGGAGAAACCTGTCTCTTGGATATTCTCGACACAG[C/G]AGGTCANGAG GAGTACAGTGCAATGAGGGACCAGTACATGAGGACTGGGGAGGGC
KRAS-183AC_TP	AAGTAGTAATTGATGGAGAAACCTGTCTCTTGGATATTCTCGACACAGCAGGTCA[A/C]GAGGAGTACA GTGCAATGAGGGACCAGTACATGAGGACTGGGGAGGGCTTTCTTT
KRAS-34GA_TP	CCACAAAATGATTCTGAATTAGCTGTATCGTCAAGGCACTCTTGCCTACGCCAC[G/A]AGCTCCAAC TAC CACAAGTTTATATTCAGTCATTTTCAGCAGGCCTTATAATAAAA
KRAS-35GA_TP	TCCACAAAATGATTCTGAATTAGCTGTATCGTCAAGGCACTCTTGCCTACGCCA[G/A]CAGCTCCAAC TA CCACAAGTTTATATTCAGTCATTTTCAGCAGGCCTTATAATAAA
KRAS-3GA_TP	ATTATAAGGCCTGCTGAAAATGACTGAATATAAACTTGTGGTAGTTGGAGCTGGTG[G/A]CGTAGGCAA GAGTGCCTTGACGATACAGCTAATTCAGAATCATTTTGTGGACGA
NRAS-181CA_TP	AACAAGTGGTTATAGATGGTGAAACCTGTTTGTGGACATACTGGATACAGCTGGA[C/A]AAGAAGAGT ACAGTGCCATGAGAGACCAATACATGAGGACAGGCGAAGGCTTCC
NRAS-183AT_TP	CAAGTGGTTATAGATGGTGAAACCTGTTTGTGGACATACTGGATACAGCTGGACA[A/T]GAAGAGTAC AGTGCCATGAGAGACCAATACATGAGGACAGGCGAAGGCTTCTCTC
BRAF-1799TA_TP	ATATTTCTTCATGAAGACCTCACAGTAAAAATAGGTGATTTTGGTCTAGCTACAG[T/A]GAAATCTCGAT GGAGTGGGTCCCATCAGTTTGAACAGTTGTCTGGATCCATTTTG
CTNNB1-121AG_TP	GTATCCACATCCTCTTCTCAGGATTGCCTTTACCACTCAGAGAAGGAGCTGTGG[A/G]AGTGGCACCAG AATGGATTNCAGAGTNCAGGTAAGACTGTTGCTGCCAGTGACTA
CTNNB1-134CT_TP	CAGGACTTGGGAGGTATCCACATCCTCTTCTCAGGATTGCCTTTACCACTCAGA[C/T]AAGGAGCTGTG GTAGTGGCACCAGAATGGATTNCAGAGTNCAGGTAAGACTGTTG
EGFR-2369CT_TP	CCCACGTGTGCCGCTGCTGGGCATCTGCCTCACCTCCACCGTGCAGCTCATCA[C/T]GCAGCTCATGCC CTTCCGGCTGCCTCCTGGACTATGTCCGGGAACACAAGACAATA
EGFR-2573TG_TP	TTACTTTGCCTCCTTCTGCATGGTATTCTTTCTCTTCCGCACCCAGCAGTTTGGCC[T/G]GCCCCAAATCT GTGATCTTGACATGCTGCGGTGTTTTCACCACTACGTTCTCTGG
KRAS-176CG_TP	AGGAAGCAAGTAGTAATTGATGGAGAAACCTGTCTCTTGGATATTCTCGACACAG[C/G]AGGTCANGAG GAGTACAGTGCAATGAGGGACCAGTACATGAGGACTGGGGAGGGC

TABLE 10-continued

Plasmid SNP Sequences (target alleles are indicated in brackets).	
SNP ID	Sequence
KRAS-183AC_TP	AAGTAGTAATTGATGGAGAAACCTGTCTCTTGGATATTCTCGACACAGCAGGTCA[C/G]GAGGAGTACA GTGCAATGAGGGACCAGTACATGAGGACTGGGGAGGGCTTTCTTT
KRAS-34GA_TP	CCACAAAATGATTCTGAATTAGCTGTATCGTCAAGGCACTCTTGCCCTACGCCAC[G/A]AGCTCCAACTAC CACAAAGTTTATATTAGTCATTTTCAGCAGGCCTTATAATAAAA
KRAS-35GA_TP	TCCACAAAATGATTCTGAATTAGCTGTATCGTCAAGGCACTCTTGCCCTACGCCA[G/A]CAGCTCCAACTA CCACAAAGTTTATATTAGTCATTTTCAGCAGGCCTTATAATAAA
KRAS-3GA_TP	ATTATAAGGCCTGCTGAAAATGACTGAATATAAACTTGTGGTAGTTGGAGCTGGTG[G/A]CGTAGGCAA GAGTGCCTTGACGATACAGCTAATTACAGATCATTGTTGTGGACGA
NRAS-181CA_TP	AACAAGTGGTTATAGATGGTGAAACCTGTTTGTGGACATACTGGATACAGCTGGA[C/A]AAGAAGAGT ACAGTGCCATGAGAGACCAATACATGAGGACAGGCGAAGGCTTCC
NRAS-183AT_TP	CAAGTGGTTATAGATGGTGAAACCTGTTTGTGGACATACTGGATACAGCTGGACA[A/T]GAAGAGTAC AGTGCCATGAGAGACCAATACATGAGGACAGGCGAAGGCTTCTCT
BRAF-1799TA_TP	ATATTTCTTCATGAAGACCTCACAGTAAATAGGTGATTTTGGTCTAGCTACAG[T/A]GAAATCTCGAT GGAGTGGGTCCCATCAGTTTGAACAGTTGTCTGGATCCATTTTG
CTNNB1-121AG_TP	GTATCCACATCCTCTTCTCAGGATTGCCTTTACCACTCAGAGAAGGAGCTGTGG[A/G]AGTGGCACCAG AATGGATTNCAGAGTNCAGGTAAGACTGTTGCTGCCAGTGACTA
CTNNB1-134CT_TP	CAGGACTTGGGAGGTATCCACATCCTCTTCTCAGGATTGCCTTTACCACTCAGA[C/T]AAGGAGCTGTG GTAGTGGCACCAGAATGGATTNCAGAGTNCAGGTAAGACTGTTG
EGFR-2369CT_TP	CCCACGTGTGCCGCTGCTGGGCATCTGCCTCACCTCCACCGTGCAGCTCATCA[C/T]GCAGCTCATGCC CTTCGGCTGCCTCCTGGACTATGTCCGGGAACAAAGACAATA
EGFR-2573TG_TP	TTACTTTGCCTCCTTCTGCATGGTATTCTTCTCTTCCGCACCCAGCAGTTTGGCC[C/T]GCCCCAAATCT GTGATCTTGACATGCTGCGGTGTTTCACCACTACGTTCTCTGG
KRAS-176CG_TP	AGGAAGCAAGTAGTAATTGATGGAGAAACCTGTCTCTTGGATATTCTCGACACAG[T/G]AGGTCANGAG GAGTACAGTGCAATGAGGGACCAGTACATGAGGACTGGGGAGGGC
KRAS-183AC_TP	AAGTAGTAATTGATGGAGAAACCTGTCTCTTGGATATTCTCGACACAGCAGGTCA[A/C]GAGGAGTACA GTGCAATGAGGGACCAGTACATGAGGACTGGGGAGGGCTTTCTTT
KRAS-34GA_TP	CCACAAAATGATTCTGAATTAGCTGTATCGTCAAGGCACTCTTGCCCTACGCCAC[G/A]AGCTCCAACTAC CACAAAGTTTATATTAGTCATTTTCAGCAGGCCTTATAATAAAA
KRAS-35GA_TP	TCCACAAAATGATTCTGAATTAGCTGTATCGTCAAGGCACTCTTGCCCTACGCCA[G/A]CAGCTCCAACTA CCACAAAGTTTATATTAGTCATTTTCAGCAGGCCTTATAATAAA
KRAS-3GA_TP	ATTATAAGGCCTGCTGAAAATGACTGAATATAAACTTGTGGTAGTTGGAGCTGGTG[G/A]CGTAGGCAA GAGTGCCTTGACGATACAGCTAATTACAGATCATTGTTGTGGACGA
NRAS-181CA_TP	AACAAGTGGTTATAGATGGTGAAACCTGTTTGTGGACATACTGGATACAGCTGGA[C/A]AAGAAGAGT ACAGTGCCATGAGAGACCAATACATGAGGACAGGCGAAGGCTTCC
NRAS-183AT_TP	CAAGTGGTTATAGATGGTGAAACCTGTTTGTGGACATACTGGATACAGCTGGACA[A/T]GAAGAGTAC AGTGCCATGAGAGACCAATACATGAGGACAGGCGAAGGCTTCTCT

[0265] The results of cast-PCR using non-tailed ASP and cast-PCR using tailed-ASP are summarized in FIG. 13. In the cast-PCR reactions having no tailed primers, the average ΔC_t of the 12 tested mutations was 10.3, whereas in the cast-PCR reactions with tailed primers the average ΔC_t of 12 tested assays is 16.3. Thus, the average $\Delta \Delta C_t$ between cast-PCR comprising tailed ASP versus cast-PCR comprising non-tailed ASP was about 6.0, which is about a 64 fold improvement in specificity for reactions comprising ASP +tail primers.

Example 13

Comparison of Cast-PCR and ASB-PCR

[0266] Allelic discrimination for assays using cast-PCR methods was compared to assays using other Allele-Specific PCR with a Blocking reagent (ASB-PCR) methods (see, e.g., Morlan et al., 2009).

[0267] cast-PCR assays were performed using the general schema and reaction conditions indicated above, using 1 mil-

lion copies of plasmid DNA containing various SNP target sequences (see Table 10). Assays were performed using the general experimental design and reaction conditions indicated above (see section II of Examples). Assay primers and probes were designed according to the sequences shown in FIG. 19B-D.

[0268] ASB-PCR assays were performed using 1 million copies of plasmid DNA containing various SNP target sequences (see Table 10), 900 nM non-tailed allele-specific primers (T_m 58–62° C.), 3600 nM allele-specific phosphate blocker (T_m 58–62° C.), 200 nM locus-specific TaqMan probe (T_m 70–74° C.), and 900 nM locus-specific primers (T_m 60–63° C.).

[0269] ASB-PCR assay primers and probes were designed according to the sequences shown in FIG. 20A-C. The ASB-PCR reactions were incubated in a 384-well plate at 95° C. for 10 minutes, followed by 50 cycles of at 92° C. for 20 seconds, 60° C. for 45 seconds. All reactions were run in 4 replications in an ABI PRISM 7900HT® Sequence Detection System, according to the manufacturer's instructions.

[0270] The results for this example are summarized in FIG. 14. In the ASB-PCR assays, the average ΔC_t of 12 different mutations was 14.1. In cast-PCR assays, the average ΔC_t of the same 12 mutations was 16.3. The $\Delta\Delta C_t$ between ASB-PCR and cast-PCR was 2.2, which indicates that the specificity of cast-PCR was approximately 4.6 fold higher than that of the ASB-PCR assay.

Example 14

Comparison of MGB and Phosphate Blocker Probes in Cast-PCR

[0271] The use of MGB blocker probes was compared to the use of other types of blocker probes, such as PO_4 blocker probes (e.g., Morlan et al., 2009), in cast-PCR assays.

[0272] All assays were performed using the general cast-PCR schema and reaction conditions indicated above (see Section II in Examples), using 1 million copies of plasmid DNA containing various SNP target sequences (see Table 10), except that reactions contained either 150 nM allele-specific MGB blocker probes or 150 nM allele-specific 3'-phosphate blocker probes. Assay primers and probes were designed according to the sequences shown in FIG. 19B-D (for cast-PCR using MGB blocker probes) or FIG. 19B-C and FIG. 20C (for cast-PCR using phosphate blocker probes; "PHOS1" to block allele-1 and "PHOS2" to block allele-2).

[0273] The results of assays with phosphate blocker probes or with MGB blocker probes are summarized in FIG. 15. In cast-PCR assays performed using phosphate blocker probes the average ΔC_t of 12 different mutations was 15.1. In comparison, the average ΔC_t for the same 12 mutations using cast-PCR assays performed with MGB blocker probes was slightly higher and gave a ΔC_t of 15.8.

Example 15

Improving the Specificity of Cast-PCR Using Lna Modified Asp

[0274] LNA-modified cast-PCR assays were performed using the general experimental design and reaction conditions indicated above (see Section H in Examples), using 0.5 ng/ μ L of genomic DNA. Assay primers and probes were designed according to the sequences shown in FIGS. 21A-C.

For each SNP analyzed, the blocker probes, locus-specific probes and locus-specific primers were the same and only the allele-specific primers varied (i.e., with or without an LNA-modification at the 3' end.).

[0275] The effect of LNA modification of the ASP on the specificity of cast-PCR is summarized in FIG. 16. For the 12 cast-PCR assays performed using LNA-modified allele-specific primers, the average ΔC_t was 16.3. In comparison, the average ΔC_t for the same 12 mutations using cast-PCR assays performed allele-specific primers having no modifications the ΔC_t was noticeably higher at 18.5. Based on the $\Delta\Delta C_t$ the assay specificity increased by approximately 4 fold for those assays that used LNA-modified allele-specific primers.

Example 16

Improving the Specificity of Cast-PCR Using Other Modified ASP

[0276] cast-PCR assays using other chemically-modified ASPs were performed using the general experimental design and reaction conditions indicated above conditions indicated above (see Section II in Examples), performed in the presence of 1 million copies of plasmid DNA containing various SNP target sequences (see Table 10). Assay primers and probes were designed according to the sequences shown in FIG. 22. For each SNP analyzed, the blocker probes, locus-specific probes and locus-specific primers were the same and only the allele-specific primers varied (i.e., with or without chemical modifications, i.e., ppA, ppG, iso dC or fdU, at the 3' end).

[0277] The results of cast-PCR assays using unmodified ASP and cast-PCR assays with modified ASP are summarized in FIG. 17. As shown, allele-specific primers having pyrophosphate modifications (ppA or ppG) at their 3'-ends increased ΔC_t by 2-3, which is approximately a 4-6 fold increase in assay specificity.

[0278] It is noted that, as used in this specification and the appended claims, the singular forms "a," "an," and "the," include plural referents unless expressly and unequivocally limited to one referent. The use of "or" means "and/or" unless stated otherwise. The use of "comprise," "comprises," "comprising," "include," "includes," and "including" are interchangeable and not intended to be limiting. Furthermore, where the description of one or more embodiments uses the term "comprising," those skilled in the art would understand that, in some specific instances, the embodiment or embodiments can be alternatively described using the language "consisting essentially of" and/or "consisting of."

[0279] The section headings used herein are for organizational purposes only and are not to be construed as limiting the subject matter described. All documents, or portions of documents, cited in this application, including but not limited to patents, patent applications, articles, books, and treatises are hereby expressly incorporated by reference in their entirety for any purpose. In the event that one or more of the incorporated documents defines a term that contradicts that term's definition in this application, this application controls.

[0280] All references cited herein, including patents, patent applications, papers, text books, and the like, and the references cited therein, to the extent that they are not already, are hereby incorporated by reference in their entirety. In the event that one or more of the incorporated literature and similar materials differs.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 635

<210> SEQ ID NO 1

<211> LENGTH: 144

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 1

```
gctctgcttc attcctgtct gaagaagggc agatagtttg gctgctcctg tgytgtcacc      60
tgcaattctc ccttatcagg gccattggcc tctcccttct ctctgtgagg gatattttct      120
ctgacttgtc aatccacatc ttcc                                             144
```

<210> SEQ ID NO 2

<211> LENGTH: 82

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide

<400> SEQUENCE: 2

```
ggcttgcaat ggctccaacc ggaagggcgg tgctcgagct gtggtgctg cygctaagtt      60
gtgcgttcca gggtgcactc gc                                             82
```

<210> SEQ ID NO 3

<211> LENGTH: 65

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide

<400> SEQUENCE: 3

```
gcaactatac ccttgatgga tggagattta ygcaatgtgt tttactgggt agagtgcag      60
acctt                                             65
```

<210> SEQ ID NO 4

<211> LENGTH: 124

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 4

```
cctgaactta tttggcaaga gcgatgagta ctcttaaaat tactatctgg aaattatatt      60
atttagaatc tgccaattac ctagatcccc cctsaacaat tgtttacca aggaacttcc      120
tgaa                                             124
```

<210> SEQ ID NO 5

<211> LENGTH: 180

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

-continued

<400> SEQUENCE: 5

gaattggttg tctccttatg ggaactggaa gtattttgac akctttacca cattttctca 60
tgggatagta agtggttaaag agctctgagc catttattat cagctacttg taaattagca 120
gtagaatttt attttttatac ttgtaagtgg gcagttacct tttgagagga atacctatag 180

<210> SEQ ID NO 6

<211> LENGTH: 87

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide

<400> SEQUENCE: 6

gcggaggga gctcatcagt ggggccacga gctgagtgcg tctgtcact ccactcccat 60
gtcccttggg aaggctctgag actaggg 87

<210> SEQ ID NO 7

<211> LENGTH: 192

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 7

tactacacct cagatatatt tcttcatgaa gacctcacag taaaaatagg tgattttggt 60
ctagctacag wgaatctctg atggagtggg tcccatcagt ttgaacagtt gtctggatcc 120
attttgtgga tggtaagaat tgaggctatt tttccactga ttaaatTTTT ggccctgaga 180
tgctgctgag tt 192

<210> SEQ ID NO 8

<211> LENGTH: 198

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 8

tgctaatact gtttcgtatt tatagctgat ttgatggagt tggacatggc catggaacca 60
gacagaaaag cggtctgttag tcaatggcag caacagtctt acctggactc tggaatccat 120
tctggtgcc actrccacag tcttctctg agtggtaaa gcaatcctga ggaagaggat 180
gtggatacct cccaagtc 198

<210> SEQ ID NO 9

<211> LENGTH: 184

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 9

tttgatggag ttggacatgg ccaatggaac agacagaaaa gcggctgtta gtcactggca 60
gcaacagtct tacctggact ctggaatcca ttctggtgcc actaacacag ctcttcttct 120

-continued

```

gagtggtaaa ggcaatcctg aggaagagga tgtggatacc tcccaagtcc tgtatgagtg      180
ggaa                                                                    184

<210> SEQ ID NO 10
<211> LENGTH: 181
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        polynucleotide

<400> SEQUENCE: 10
gtggacaacc cccacgtgtg cgcctgctg ggcactctgcc tcacctccac cgtgcagctc      60
atcaygcagc tcatgccctt cggtgcctc ctggactatg tccgggaaca caaagacaat      120
attggctccc agtacctgct caactggtgt gtgcagatcg caaaggtaat caggggaaggg      180
a                                                                    181

<210> SEQ ID NO 11
<211> LENGTH: 166
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        polynucleotide

<400> SEQUENCE: 11
gcatgaacta ctggaggac cgtcgcttgg tgcaccgcga cctggcagcc aggaacgtac      60
tggtgaaaac accgcagcat gtcaagatca cagattttgg gckggccaaa ctgctgggtg      120
cggaagagaa agaataccat gcagaaggag gcaaagtaag gaggtg                    166

<210> SEQ ID NO 12
<211> LENGTH: 172
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        polynucleotide

<400> SEQUENCE: 12
caggattcct acaggaagca agtagtaatt gatggagaaa cctgtctctt ggatattctc      60
gacacagsag gtcaagagga gtacagtgca atgagggacc agtacatgag gactggggag      120
ggctttcttt gtgtatttgc cataaataat actaaatcat ttgaagatat tc                    172

<210> SEQ ID NO 13
<211> LENGTH: 207
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        polynucleotide

<400> SEQUENCE: 13
acaggaagca agtagtaatt gatggagaaa cctgtctctt ggatattctc gacacagcag      60
gtcamgagga gtacagtgca atgagggacc agtacatgag gactggggag ggctttcttt      120
gtgtatttgc cataaataat actaaatcat ttgaagatat tcaccattat aggtggggtt      180
aaattgaata taataagctg acattaa                                          207

```

-continued

<210> SEQ ID NO 14
<211> LENGTH: 169
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 14

tattaacctt atgtgtgaca tgttctaata tagtcacatt ttcattatTT ttattataag 60
gcctgctgaa aatgactgaa tataaacttg tggtagttgg agctrgtggc gtaggcaaga 120
gtgccttgac gatacagcta attcagaatc attttgtgga cgaatatga 169

<210> SEQ ID NO 15
<211> LENGTH: 171
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 15

tattaacctt atgtgtgaca tgttctaata tagtcacatt ttcattatTT ttattataag 60
gcctgctgaa aatgactgaa tataaacttg tggtagttgg agctrgtggc gtaggcaaga 120
gtgccttgac gatacagcta attcagaatc attttgtgga cgaatatgat c 171

<210> SEQ ID NO 16
<211> LENGTH: 230
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 16

cattatTTTT attataaggc ctgctgaaaa tgactgaata taaacttgTg gtagttggag 60
ctggTgrcgt aggcaagagt gccttgacga tacagctaatt tcagaatcat tttgtggacg 120
aatatgatcc aacaatagag gtaaattcttg ttttaatatg catattactg gtgcaggacc 180
attccttgat acagataaag gtttctctga ccattttcat gagtacttat 230

<210> SEQ ID NO 17
<211> LENGTH: 210
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 17

attccttacag aaaacaagtg gttatagatg gtgaaacctg tttgttggaC atactggata 60
cagctggama agaagagtac agtgccatga gagaccaata catgaggaca ggcaaggct 120
tcctctgtgt atttgccatc aataatagca agtcatttgc ggatattaac ctctacaggt 180
actaggagca ttattttctc tgaaggatg 210

<210> SEQ ID NO 18
<211> LENGTH: 206

-continued

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 18

ttacagaaaa caagtgggta tagatgggta aacctgtttg ttggacatac tggatacagc 60
tggacawgaa gagtacagtg ccatgagaga ccaatacatg aggacaggcg aaggcttcct 120
ctgtgtatatt gccatcaata atagcaagtc atttgcggat attaacctct acaggtacta 180
ggagcattat tttctctgaa aggatg 206

<210> SEQ ID NO 19
<211> LENGTH: 162
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 19

tggtttccaa caggttcttg ctggtgtgaa atgactgagt acaaactggt ggtggttgga 60
gcagrtggtg ttgggaaaaa cgcactgaca atccagctaa tccagaacca cttttagat 120
gaatatgata ccaccataga ggtgaggccc agtggtagcc cg 162

<210> SEQ ID NO 20
<211> LENGTH: 158
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 20

tttccaacag gttcttgctg gtgtgaaatg actgagtaca aactggtggt ggttgagca 60
ggtgrtggtg ggaaaagcgc actgacaatc cagctaatacc agaaccactt ttagatgaa 120
tatgatccca ccatagaggt gaggccagc gtagccc 158

<210> SEQ ID NO 21
<211> LENGTH: 189
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 21

ggcaccgcg tccgcgccat ggccatctac aagcagtcac agcacatgac ggaggttggt 60
aggcrtgcc cccaccatga gcgctgctca gatagcgtg gtagcagct ggggctggag 120
agacgacagg gctggttgcc cagggtcccc aggcctctga ttcctcactg attgctctta 180
ggtctggcc 189

<210> SEQ ID NO 22
<211> LENGTH: 165
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic

-continued

polynucleotide

<400> SEQUENCE: 22

```
cctcctcagc atcttatccg agtgggaagga aatttgctg tggagtattt ggatgacaga      60
aacactttty gacatagtggt ggtggtgccc tatgagccgc ctgaggtctg gtttgcaact      120
ggggtctctg ggaggagggg ttaaggggtg ttgtcagtgg ccctc                        165
```

<210> SEQ ID NO 23

<211> LENGTH: 183

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 23

```
cttgggcctg tgttatctcc taggttggt ctgactgtac caccatccac tacaactaca      60
tgtgtaacag tkcctgcatg ggcggcatga accggaggcc catcctcacc atcatcacac      120
tggaagactc caggtcagga gccacttgcc accctgcaca ctggcctgct gtgccccagc      180
ctc                                                                183
```

<210> SEQ ID NO 24

<211> LENGTH: 163

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 24

```
taggttggt ctgactgtac caccatccac tacaactaca tgtgtaacag ttctgcatg      60
ggcrgcatga accggaggcc catcctcacc atcatcacac tggaagactc caggtcagga      120
gccacttgcc accctgcaca ctggcctgct gtgccccagc ctc                    163
```

<210> SEQ ID NO 25

<211> LENGTH: 163

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 25

```
ctgactgtac caccatccac tacaactaca tgtgtaacag ttctgcatg ggcggcatga      60
acyggaggcc catcctcacc atcatcacac tggaagactc caggtcagga gccacttgcc      120
accctgcaca ctggcctgct gtgccccagc ctctgcttgc ctc                    163
```

<210> SEQ ID NO 26

<211> LENGTH: 162

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 26

```
tgactgtacc accatccact acaactacat gtgtaacagt tcctgcatgg ggcggcatgaa      60
```

-continued

```
ccrgaggccc atcctcacca tcacacact ggaagactcc aggtcaggag ccacttgcca 120
ccctgcacac tggcctgctg tgccccagcc tctgettgc tc 162
```

```
<210> SEQ ID NO 27
<211> LENGTH: 176
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      polynucleotide
```

```
<400> SEQUENCE: 27
cctcttgett ctcttttctt atcctgagta gtggaatct actgggacgg aacagctttg 60
aggtygtgt ttgtgcctgt cctgggagag accggcgcac agaggaagag aatctccgca 120
agaaagggga gcctcaccac gagctgcccc caggagcac taagcgaggt aagcaa 176
```

```
<210> SEQ ID NO 28
<211> LENGTH: 601
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
```

```
<400> SEQUENCE: 28
agaaaataac taagggaagg aggaaagtgg ggaggaagga agaacagtgt gaagacaatg 60
gcctgaaaac tgaaaaagtc tgttaaagtt aattatcagt ttttgagtcc aagaactggc 120
tttgctactt tctgtaagtt tctaatttac tgaataagca tgaaaaagat tgctttgagg 180
aatggttata aacacattct tagagcatag taagcagtag ggagtaacaa aataaactg 240
attagaatac ttactctac ttaattaatc aatcatattt agtttgactc accttcccag 300
hacettctag ttctttctta tctttcagtg ctgtccaga caacattttc atttcaacaa 360
ctcctgctat tgcaatgatg ggtacaattg ctaagagtaa cagtgttagt tgccaacccat 420
agatgaagga tataattatt cctgtcccaa gatttgctat attctgggta attacagcaa 480
gcctggaacc tatagcctgc aaacaaaac aaattagaga aattttaaaa atattatctt 540
cacaactcat gcttctatct tctgaaaact caccttcag agactatatt cattatttta 600
t 601
```

```
<210> SEQ ID NO 29
<211> LENGTH: 27
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      primer
```

```
<400> SEQUENCE: 29
atatttagtt tgactcacct tcccagc 27
```

```
<210> SEQ ID NO 30
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      primer
```

```
<400> SEQUENCE: 30
accactcacc tttcccagc 19
```

-continued

<210> SEQ ID NO 31
<211> LENGTH: 27
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 31

atatttagtt tgactcacct tcccaga 27

<210> SEQ ID NO 32
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 32

accactcacc tttcccaga 19

<210> SEQ ID NO 33
<211> LENGTH: 27
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 33

atatttagtt tgactcacct tcccagt 27

<210> SEQ ID NO 34
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 34

accactcacc tttcccagt 19

<210> SEQ ID NO 35
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 35

tggacaagca ctgaaaga 18

<210> SEQ ID NO 36
<211> LENGTH: 27
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 36

-continued

gcaggagttg ttgaaatgaa aatgttg 27

<210> SEQ ID NO 37
<211> LENGTH: 163
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 37

tgactgaata taaacttggt gtagttggag ctgstggcgt aggcaagagt gccttgacga 60

tacagctaata tcagaatcat ttgtggacg aatatgatcc aacaatagag gtaaatcttg 120

ttttaatatg catattactg gtgcaggacc attctttgat aca 163

<210> SEQ ID NO 38
<211> LENGTH: 163
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 38

tgactgaata taaacttggt gtagttggag ctsgtggcgt aggcaagagt gccttgacga 60

tacagctaata tcagaatcat ttgtggacg aatatgatcc aacaatagag gtaaatcttg 120

ttttaatatg catattactg gtgcaggacc attctttgat aca 163

<210> SEQ ID NO 39
<211> LENGTH: 163
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 39

tgactgaata taaacttggt gtagttggag ctgrtggcgt aggcaagagt gccttgacga 60

tacagctaata tcagaatcat ttgtggacg aatatgatcc aacaatagag gtaaatcttg 120

ttttaatatg catattactg gtgcaggacc attctttgat aca 163

<210> SEQ ID NO 40
<211> LENGTH: 163
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 40

tgactgaata taaacttggt gtagttggag ctrgtggcgt aggcaagagt gccttgacga 60

tacagctaata tcagaatcat ttgtggacg aatatgatcc aacaatagag gtaaatcttg 120

ttttaatatg catattactg gtgcaggacc attctttgat aca 163

<210> SEQ ID NO 41
<211> LENGTH: 163
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 41

tgactgaata taaacttgtg gtagttggag ctggtgrcgt aggcaagagt gccttgacga 60
tacagctaata tcagaatcat ttgtggacg aatatgatcc aacaatagag gtaaattcttg 120
ttttaatatg catattactg gtgcaggacc attctttgat aca 163

<210> SEQ ID NO 42
<211> LENGTH: 164
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (34)..(35)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 42

gtgagtttgt attaaaaggt actggtggag tatnngatag tgtattaacc ttatgtgtga 60
catgttctaa tatagtcaca ttttcattat ttttattata aggcctgctg aaaatgactg 120
aatataaaact tgtggtagtt ggagctkgtg gcgtaggcaa gagt 164

<210> SEQ ID NO 43
<211> LENGTH: 111
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 43

atatattcttc atgaagacct cacagtaaaa ataggtgatt ttggtctagc tacagwgaaa 60
tctcgatgga gtgggtccca tcagtttgaa cagttgtctg gatccatttt g 111

<210> SEQ ID NO 44
<211> LENGTH: 111
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (76)..(76)
<223> OTHER INFORMATION: a, c, t, g, unknown or other
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (83)..(83)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 44

gtatccacat cctcttcttc aggattgcct ttaccactca gagaaggagc tgtggragtg 60
gcaccagaat ggattncaga gtnccagtaa gactgttgct gccagtgact a 111

<210> SEQ ID NO 45
<211> LENGTH: 111
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (89)..(89)
<223> OTHER INFORMATION: a, c, t, g, unknown or other
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (96)..(96)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 45

caggacttgg gaggtatcca catcctcttc ctcaggattg cctttaccac tcagayaagg 60
agctgtggta gtggcaccag aatggattnc agagtncagg taagactgtt g 111

<210> SEQ ID NO 46
<211> LENGTH: 111
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 46

cccacgtgtg ccgcctgtctg ggcattctgcc tcacctccac cgtgcagctc atcaygcagc 60
tcatgccctt cggctgcctc ctggactatg tccgggaaca caaagacaat a 111

<210> SEQ ID NO 47
<211> LENGTH: 111
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 47

ttactttgcc tcctcttgca tggattcttt tctcttccgc acccagcagt ttggcckgcc 60
caaaatctgt gatcttgaca tgetgcggtg ttttcaccag tacgttctctg g 111

<210> SEQ ID NO 48
<211> LENGTH: 111
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (63)..(63)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 48

aggaagcaag tagtaattga tggagaaacc tgtctcttgg atattctcga cacagsaggt 60
cangaggagt acagtgcaat gagggaccag tacatgagga ctggggaggg c 111

<210> SEQ ID NO 49
<211> LENGTH: 111
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

-continued

<400> SEQUENCE: 49

aagtagtaat tgaatggagaa acctgtctct tggatattct cgacacagca ggcamgagg 60
agtacagtgc aatgaggggac cagtacatga ggactgggga gggctttctt t 111

<210> SEQ ID NO 50

<211> LENGTH: 111

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 50

ccacaaaatg attctgaatt agctgtatcg tcaaggcact cttgcctacg ccacragctc 60
caactaccac aagtttatat tcagtcattt tcagcaggcc ttataataaa a 111

<210> SEQ ID NO 51

<211> LENGTH: 111

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 51

tccacaaaat gattctgaat tagctgtatc gtcaaggcac tcttgctac gccarcagct 60
ccaactacca caagtttata ttcatgcatt ttcagcaggc cttataataa a 111

<210> SEQ ID NO 52

<211> LENGTH: 111

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 52

attataaggc ctgctgaata tgactgaata taaacttggt gtagttggag ctgggtgrcgt 60
aggcaagagt gccttgacga tacagctaata tcagaatcat tttgtggacg a 111

<210> SEQ ID NO 53

<211> LENGTH: 111

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 53

aacaagtggg tatagatggg gaaacctggt tggtggacat actggatata gctggamaag 60
aagagtacag tgccatgaga gaccaatata tgaggacagg cgaaggcttc c 111

<210> SEQ ID NO 54

<211> LENGTH: 111

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 54

-continued

caagtgggta tagatgggtga aacctgtttg ttggacatac tggatacagc tggacawgaa 60
gagtacagtg ccatgagaga ccaatacatg aggacaggcg aaggcttcct c 111

<210> SEQ ID NO 55
<211> LENGTH: 111
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 55

atatctcttc atgaagacct cacagtaaaa ataggtgatt ttggtctagc tacagwgaaa 60
tctcgatgga gtgggtccca tcagtttgaa cagttgtctg gatccatttt g 111

<210> SEQ ID NO 56
<211> LENGTH: 111
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (76)..(76)
<223> OTHER INFORMATION: a, c, t, g, unknown or other
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (83)..(83)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 56

gtatccacat cctcttcttc aggattgcct ttaccactca gagaaggagc tgtggragtg 60
gcaccagaat ggattnca ga gtnccaggtaa gactgttgct gccagtgact a 111

<210> SEQ ID NO 57
<211> LENGTH: 111
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (89)..(89)
<223> OTHER INFORMATION: a, c, t, g, unknown or other
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (96)..(96)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 57

caggacttgg gaggtatcca catcctcttc ctcaggattg cctttaccac tcagayaagg 60
agctgtggta gtggcaccag aatggattnc agagtncagg taagactggt g 111

<210> SEQ ID NO 58
<211> LENGTH: 111
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 58

-continued

cccacgtgtg cgcctgtgtg ggcattctgcc tcacctccac cgtgcagctc atcaygcagc 60

tcattgccctt cggtgcctc ctggactatg tccgggaaca caaagacaat a 111

<210> SEQ ID NO 59

<211> LENGTH: 111

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 59

ttactttgcc tccttttgcg tggatttctt tctcttccgc acccagcagt ttggcckgcc 60

caaaatctgt gatcttgaca tgctgcggtg ttttcaccag tacgttctg g 111

<210> SEQ ID NO 60

<211> LENGTH: 111

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<220> FEATURE:

<221> NAME/KEY: modified_base

<222> LOCATION: (63)..(63)

<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 60

aggaagcaag tagtaattga tggagaaacc tgtctcttgg atattctcga cacagsaggt 60

cangaggagt acagtgcaat gagggaccag tacatgagga ctggggaggg c 111

<210> SEQ ID NO 61

<211> LENGTH: 111

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 61

aagtagtaat tgatggagaa acctgtctct tggatattct cgacacagca ggctcagagg 60

agtacagtgc aatgagggac cagtacatga ggactgggga gggctttctt t 111

<210> SEQ ID NO 62

<211> LENGTH: 111

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 62

ccacaaaatg attctgaatt agctgtatcg tcaaggcact cttgcctacg ccacragctc 60

caactaccac aagtttatat tcagtcattt tcagcaggcc ttataataaa a 111

<210> SEQ ID NO 63

<211> LENGTH: 111

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic

-continued

polynucleotide

<400> SEQUENCE: 63

tccacaaaat gattctgaat tagctgtatc gtcaaggcac tcttgccctac gccarcagct 60

ccaactacca caagtttata ttcagtcatt ttcagcaggc cttataataa a 111

<210> SEQ ID NO 64

<211> LENGTH: 111

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 64

attataaggc ctgctgaaaa tgactgaata taaacttggtg gtagtgggag ctgggtgroggt 60

aggcaagagt gccttgacga tacagctaatt tcagaatcat tttgtggacg a 111

<210> SEQ ID NO 65

<211> LENGTH: 111

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 65

aacaagtgggt tatagatgggt gaaacctggtt tggtggacat actggataca gctggamaag 60

aagagtacag tgccatgaga gaccaatata tgaggacagg cgaaggcttc c 111

<210> SEQ ID NO 66

<211> LENGTH: 111

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 66

caagtgggta tagatgggtga aacctgtttg ttggacatac tggatacagc tggacawgaa 60

gagtacagtg ccatgagaga ccaatacatg aggacaggcg aaggcttcct c 111

<210> SEQ ID NO 67

<211> LENGTH: 111

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 67

atatctcttc atgaagacct cacagtaaaa ataggtgatt ttggtctagc tacagwgaaa 60

tctcgatgga gtgggtccca tcagtttgaa cagttgtctg gatccatttt g 111

<210> SEQ ID NO 68

<211> LENGTH: 111

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

-continued

```
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (76)..(76)
<223> OTHER INFORMATION: a, c, t, g, unknown or other
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (83)..(83)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 68

gtatccacat cctcttcctc aggattgcct ttaccactca gagaaggagc tgtggragtg      60
gcaccagaat ggattncaga gtncaggtaa gactgttgct gccagtgact a              111

<210> SEQ ID NO 69
<211> LENGTH: 111
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      polynucleotide
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (89)..(89)
<223> OTHER INFORMATION: a, c, t, g, unknown or other
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (96)..(96)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 69

caggacttgg gaggtatcca catcctcttc ctccaggattg cctttaccac tcagayaagg      60
agctgtggta gtggcaccag aatggattnc agagtncagg taagactgtt g              111

<210> SEQ ID NO 70
<211> LENGTH: 111
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      polynucleotide

<400> SEQUENCE: 70

cccacgtgtg ccgcctgtgt ggcattgtgc tcacctccac cgtgcagctc atcaygcagc      60
tcatgccctt cggctgcctc ctggactatg tccgggaaca caaagacaat a              111

<210> SEQ ID NO 71
<211> LENGTH: 111
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      polynucleotide

<400> SEQUENCE: 71

ttactttgcc tcctttctgca tggattcttt tctcttccgc acccagcagt ttggccygcc      60
caaaatctgt gatcttgaca tgctgcggtg ttttcaccag tacgttctctg g              111

<210> SEQ ID NO 72
<211> LENGTH: 111
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      polynucleotide
```

-continued

<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (63)..(63)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 72

aggaagcaag tagtaattga tggagaaacc tgtctcttgg atattctcga cacagkaggt 60
cangaggagt acagtgcaat gagggaccag tacatgagga ctggggaggg c 111

<210> SEQ ID NO 73
<211> LENGTH: 111
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 73

aagtagtaat tgatggagaa acctgtctct tggatattct cgacacagca ggtcamgagg 60
agtacagtgc aatgagggac cagtacatga ggactgggga gggctttctt t 111

<210> SEQ ID NO 74
<211> LENGTH: 111
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 74

ccacaaaaatg attctgaatt agctgtatcg tcaaggcact cttgcctacg ccacragctc 60
caactaccac aagtttatat tcagtcattt tcagcaggcc ttataataaa a 111

<210> SEQ ID NO 75
<211> LENGTH: 111
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 75

tccacaaaaat gattctgaat tagctgtatc gtcaaggcac tcttgcttac gccarcagct 60
ccaactacca caagtttata ttcagtcatt ttcagcaggc cttataataa a 111

<210> SEQ ID NO 76
<211> LENGTH: 111
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 76

attataaggc ctgctgaaaa tgactgaata taaacttggt gtagttggag ctgggtgrcgt 60
aggcaagagt gccttgacga tacagctaata tcagaatcat tttgtggacg a 111

<210> SEQ ID NO 77
<211> LENGTH: 111
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 77

aacaagtggg tatagatggg gaaacctgtt tgttgacat actggataga gctggamaag 60
aagagtacag tgccatgaga gaccaataga tgaggacagg cgaaggcttc c 111

<210> SEQ ID NO 78
<211> LENGTH: 111
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 78

caagtgggta tagatgggta aacctgtttg ttggacatac tggatacagc tggacawgaa 60
gagtacagtg ccatgagaga ccaatacatg aggacaggcg aaggcttcct c 111

<210> SEQ ID NO 79
<211> LENGTH: 290
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 79

gtactgggtg agtatttgat agtgatttaa ccttatgtgt gacatgttct aatatagtca 60
cattttcatt atttttatta taaggcctgc tgaaaatgac tgaatataaa cttgtggtag 120
ttggagctgg tggcgtaggc aagagtgcct tgacgataca gctaattcag aatcattttg 180
tggacgaata tgatccaaca atagaggtaa atcttgtttt aatatgcata ttactgggtgc 240
aggaccattc ttgatacag ataaaggttt ctctgaccat tttcatgagt 290

<210> SEQ ID NO 80
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 80

cccgtgctc ctgtgc 16

<210> SEQ ID NO 81
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 81

acccaacgca caacttagcg 20

<210> SEQ ID NO 82
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic

-continued

primer

<400> SEQUENCE: 82

ccgccttga tggatggaga ttac 25

<210> SEQ ID NO 83
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 83

gcacccttgg tgaaacaatt gttg 24

<210> SEQ ID NO 84
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 84

cgccgaactg gaagtatttt gacag 25

<210> SEQ ID NO 85
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 85

tatccctcac agagagaagg gag 23

<210> SEQ ID NO 86
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 86

gcttgcaatg gctccaacc 19

<210> SEQ ID NO 87
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 87

ggagccattg caagccaag 19

<210> SEQ ID NO 88
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 88
tggcaagagc gatgagtact c 21

<210> SEQ ID NO 89
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 89
ctctcaaaag gtaactgccc actta 25

<210> SEQ ID NO 90
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 90
ctcctgtgct gtcacc 16

<210> SEQ ID NO 91
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 91
cttagcggca cgcac 15

<210> SEQ ID NO 92
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 92
ggagatttac gcaatgtg 18

<210> SEQ ID NO 93
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 93
acaattgttg agggggg 17

<210> SEQ ID NO 94
<211> LENGTH: 20

-continued

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 94
aagtattttg acagctttac 20

<210> SEQ ID NO 95
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 95
cccggtgct cctgtgt 17

<210> SEQ ID NO 96
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 96
gccgaacgca caacttagca 20

<210> SEQ ID NO 97
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 97
cccgcttga tggatggaga tttat 25

<210> SEQ ID NO 98
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 98
tgcacccttg gtgaacaat tgttc 25

<210> SEQ ID NO 99
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 99
cgccgaactg gaagtatttt gacat 25

-continued

<210> SEQ ID NO 100
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 100

atggccctga taagggaga 19

<210> SEQ ID NO 101
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 101

agggcggtgc tcgag 15

<210> SEQ ID NO 102
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 102

tgtcactcta cccagtaaa 19

<210> SEQ ID NO 103
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 103

agaatctgcc aattacctag a 21

<210> SEQ ID NO 104
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 104

acaagtagct gataataaat g 21

<210> SEQ ID NO 105
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 105

gctcctgtgt tgtcacc 17

-continued

<210> SEQ ID NO 106
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 106

cttagcagca cgcacc 16

<210> SEQ ID NO 107
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 107

tggagattta tgcaatgtg 19

<210> SEQ ID NO 108
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 108

acaattgttc agggggg 17

<210> SEQ ID NO 109
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 109

aagtattttg acatctttac 20

<210> SEQ ID NO 110
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 110

gcctgatttt ggtctagcta caga 24

<210> SEQ ID NO 111
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 111

-continued

cgcgagaagg agctgtggt 19

<210> SEQ ID NO 112
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 112

ccgtgccttt accactcaga g 21

<210> SEQ ID NO 113
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 113

gcgcgtgcag ctcacac 18

<210> SEQ ID NO 114
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 114

cggcagcagt ttggccc 17

<210> SEQ ID NO 115
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 115

gccggatatt ctcgacacag c 21

<210> SEQ ID NO 116
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 116

gcggacacag caggtcaa 18

<210> SEQ ID NO 117
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

-continued

<400> SEQUENCE: 117

gcggacacag caggtcaa

18

<210> SEQ ID NO 118

<211> LENGTH: 18

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 118

cgccttgccct acgccact

18

<210> SEQ ID NO 119

<211> LENGTH: 16

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 119

cgctgcctac gccacg

16

<210> SEQ ID NO 120

<211> LENGTH: 17

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 120

gcgttgcccta cgccacc

17

<210> SEQ ID NO 121

<211> LENGTH: 19

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 121

cgcctcttgc ctacgccat

19

<210> SEQ ID NO 122

<211> LENGTH: 18

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 122

gcgtcttgcc tacgccag

18

<210> SEQ ID NO 123

<211> LENGTH: 18

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

-continued

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 123

gcgtcttgcc tacgccac 18

<210> SEQ ID NO 124
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 124

cgcgtagttg gagctggtga 20

<210> SEQ ID NO 125
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 125

ccgatactgg atacagctgg aa 22

<210> SEQ ID NO 126
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 126

ccgtggatac agctggacaa 20

<210> SEQ ID NO 127
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 127

ccgggtggtt ggagcaga 18

<210> SEQ ID NO 128
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 128

ccgggttga gcaggtga 18

<210> SEQ ID NO 129
<211> LENGTH: 18
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 129

ccgggagggtt gtgaggca 18

<210> SEQ ID NO 130
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 130

gccggatgac agaaacactt ttc 23

<210> SEQ ID NO 131
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 131

gcgtacaact acatgtgtaa cagtg 25

<210> SEQ ID NO 132
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 132

gccttcctgc atgggca 17

<210> SEQ ID NO 133
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 133

gccggcggca tgaacc 16

<210> SEQ ID NO 134
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 134

gccgcggcat gaacca 16

<210> SEQ ID NO 135

-continued

<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 135

gccgccaaca gctttgaggt gc 22

<210> SEQ ID NO 136
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 136

ccggattttg gtctagctac agt 23

<210> SEQ ID NO 137
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 137

gccagaagga gctgtggc 18

<210> SEQ ID NO 138
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 138

ccgtgccttt accactcaga a 21

<210> SEQ ID NO 139
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 139

gcgcgtgcag ctcatcat 18

<210> SEQ ID NO 140
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 140

cggcagcagt ttggcca 17

-continued

<210> SEQ ID NO 141
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 141

cgctggatat tctcgacaca gg 22

<210> SEQ ID NO 142
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 142

gcggacacag caggtcac 18

<210> SEQ ID NO 143
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 143

gcggacacag caggtcat 18

<210> SEQ ID NO 144
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 144

gcgttgcccta cgccacc 17

<210> SEQ ID NO 145
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 145

gcgttgcccta cgccacc 17

<210> SEQ ID NO 146
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 146

-continued

gcgttgcccta cgccaca 17

<210> SEQ ID NO 147
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 147

gcgtcttgcc tacgccac 18

<210> SEQ ID NO 148
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 148

gcgtcttgcc tacgccac 18

<210> SEQ ID NO 149
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 149

gcgtcttgcc tacgcca 18

<210> SEQ ID NO 150
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 150

gcctagtgg agctggtgg 19

<210> SEQ ID NO 151
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 151

ccgatactgg atacagctgg ac 22

<210> SEQ ID NO 152
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

-continued

<400> SEQUENCE: 152

cgccctggata cagctggaca t

21

<210> SEQ ID NO 153

<211> LENGTH: 17

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 153

cgcggtggtg gagcagg

17

<210> SEQ ID NO 154

<211> LENGTH: 17

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 154

cgcggttgag caggtgg

17

<210> SEQ ID NO 155

<211> LENGTH: 17

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 155

ggcgaggttg tgaggcg

17

<210> SEQ ID NO 156

<211> LENGTH: 24

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 156

gcctggatga cagaaacact tttt

24

<210> SEQ ID NO 157

<211> LENGTH: 26

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 157

ggcctacaac tacatgtgta acagtt

26

<210> SEQ ID NO 158

<211> LENGTH: 16

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic

-continued

primer

<400> SEQUENCE: 158

gcctcctgca tgggcg 16

<210> SEQ ID NO 159
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 159

gccggcggca tgaact 16

<210> SEQ ID NO 160
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 160

gccgcggcat gaaccg 16

<210> SEQ ID NO 161
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 161

cgcgaacagc tttgaggtgt 20

<210> SEQ ID NO 162
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 162

cccgtggtct agctacaga 19

<210> SEQ ID NO 163
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 163

ccccaaggag ctgtggt 17

<210> SEQ ID NO 164
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 164

cgcccccttta ccaactcagag 20

<210> SEQ ID NO 165
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 165

gggggatgcag ctcacac 18

<210> SEQ ID NO 166
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 166

ggcgcagttt ggccc 15

<210> SEQ ID NO 167
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 167

cccgattctc gacacagc 18

<210> SEQ ID NO 168
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 168

cgcccacagc aggtcaa 17

<210> SEQ ID NO 169
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 169

tgcccacagc aggtcaa 17

<210> SEQ ID NO 170
<211> LENGTH: 15

-continued

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 170
gccgcctacg ccact 15

<210> SEQ ID NO 171
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 171
agcccctacg ccacg 15

<210> SEQ ID NO 172
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 172
accccctacg ccacc 15

<210> SEQ ID NO 173
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 173
ggctgcctac gccat 15

<210> SEQ ID NO 174
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 174
gggctgccta cgccag 16

<210> SEQ ID NO 175
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 175
ggctgcctac gccac 15

-continued

<210> SEQ ID NO 176
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 176

ccccgttgga gctggtga 18

<210> SEQ ID NO 177
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 177

cgggtggata cagctggaa 19

<210> SEQ ID NO 178
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 178

gggcgataca gctggacaa 19

<210> SEQ ID NO 179
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 179

gccctgggtg gagcaga 17

<210> SEQ ID NO 180
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 180

cccgttgag caggtga 17

<210> SEQ ID NO 181
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 181

gccaggttgt gaggca 16

-continued

<210> SEQ ID NO 182
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 182

tgccctgaca gaaacacttt tc 22

<210> SEQ ID NO 183
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 183

cgccctacat gtgtaacagt g 21

<210> SEQ ID NO 184
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 184

ccgcctgcat gggca 15

<210> SEQ ID NO 185
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 185

cccgcggcat gaacc 15

<210> SEQ ID NO 186
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 186

ccccggcatg aacca 15

<210> SEQ ID NO 187
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 187

-continued

cgccagcttt gaggtgc 17

<210> SEQ ID NO 188
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 188

cccgtggtct agctacagt 19

<210> SEQ ID NO 189
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 189

ccccaggagc tgtggc 16

<210> SEQ ID NO 190
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 190

cgccccctta ccactcagaa 20

<210> SEQ ID NO 191
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 191

ggggagtgca gctcatcat 19

<210> SEQ ID NO 192
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 192

ggcgcagttt ggcca 15

<210> SEQ ID NO 193
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

-continued

<400> SEQUENCE: 193

cccgattctc gacacagg

18

<210> SEQ ID NO 194

<211> LENGTH: 17

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 194

tgcccacagc aggtcac

17

<210> SEQ ID NO 195

<211> LENGTH: 17

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 195

cgcccacagc aggtcat

17

<210> SEQ ID NO 196

<211> LENGTH: 14

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 196

gcccctacgc cacc

14

<210> SEQ ID NO 197

<211> LENGTH: 15

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 197

agcccctacg ccacc

15

<210> SEQ ID NO 198

<211> LENGTH: 15

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 198

accgcctacg ccaca

15

<210> SEQ ID NO 199

<211> LENGTH: 16

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

-continued

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 199

gggctgccta cgccac 16

<210> SEQ ID NO 200

<211> LENGTH: 15

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 200

ggctgcctac gccac 15

<210> SEQ ID NO 201

<211> LENGTH: 16

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 201

gggctgccta cgccaa 16

<210> SEQ ID NO 202

<211> LENGTH: 17

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 202

ccccttgagg ctggtgg 17

<210> SEQ ID NO 203

<211> LENGTH: 18

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 203

cgggggatac agctggac 18

<210> SEQ ID NO 204

<211> LENGTH: 19

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 204

gggcgataca gctggacat 19

<210> SEQ ID NO 205

<211> LENGTH: 16

<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 205

gcccggttgg agcagg 16

<210> SEQ ID NO 206
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 206

cccttgagc aggtgg 16

<210> SEQ ID NO 207
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 207

gccaggttgt gaggcg 16

<210> SEQ ID NO 208
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 208

tgccctgaca gaaacacttt tt 22

<210> SEQ ID NO 209
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 209

cgccctacat gtgtaacagt t 21

<210> SEQ ID NO 210
<211> LENGTH: 14
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 210

ccgctgcatg ggcg 14

<210> SEQ ID NO 211

-continued

<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 211

cccgcggcat gaact 15

<210> SEQ ID NO 212
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 212

ccccggcatg aaccg 15

<210> SEQ ID NO 213
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 213

cgccagcttt gaggtgt 17

<210> SEQ ID NO 214
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 214

agcctcaatt cttaccatcc acaaaa 26

<210> SEQ ID NO 215
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 215

catggaacca gacagaaaag cg 22

<210> SEQ ID NO 216
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 216

gtcactggca gcaacagtct ta 22

-continued

<210> SEQ ID NO 217
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 217

tgggagccaa tattgtcttt gtgtt 25

<210> SEQ ID NO 218
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 218

aggaacgtac tggtgaaaac acc 23

<210> SEQ ID NO 219
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 219

ccctccccag tcctcatgta 20

<210> SEQ ID NO 220
<211> LENGTH: 41
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 220

tcttcaaatg atttagtatt atttatggca aatacacaaa g 41

<210> SEQ ID NO 221
<211> LENGTH: 41
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 221

tcttcaaatg atttagtatt atttatggca aatacacaaa g 41

<210> SEQ ID NO 222
<211> LENGTH: 28
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 222

-continued

tgtgtgacat gttctaatat agtcacat 28

<210> SEQ ID NO 223
<211> LENGTH: 28
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 223

tgtgtgacat gttctaatat agtcacat 28

<210> SEQ ID NO 224
<211> LENGTH: 28
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 224

tgtgtgacat gttctaatat agtcacat 28

<210> SEQ ID NO 225
<211> LENGTH: 28
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 225

tgtgtgacat gttctaatat agtcacat 28

<210> SEQ ID NO 226
<211> LENGTH: 28
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 226

tgtgtgacat gttctaatat agtcacat 28

<210> SEQ ID NO 227
<211> LENGTH: 28
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 227

tgtgtgacat gttctaatat agtcacat 28

<210> SEQ ID NO 228
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

-continued

<400> SEQUENCE: 228

ggctctgcac cagtaatatg ca 22

<210> SEQ ID NO 229

<211> LENGTH: 29

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 229

gcaaatgact tgctattatt gatggcaaa 29

<210> SEQ ID NO 230

<211> LENGTH: 29

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 230

gcaaatgact tgctattatt gatggcaaa 29

<210> SEQ ID NO 231

<211> LENGTH: 24

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 231

agtggttctg gattagctgg attg 24

<210> SEQ ID NO 232

<211> LENGTH: 24

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 232

gtggttctgg attagctgga ttgt 24

<210> SEQ ID NO 233

<211> LENGTH: 18

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 233

gcaaccagcc ctgtcgtc 18

<210> SEQ ID NO 234

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic

-continued

primer

<400> SEQUENCE: 234

agaccccggt tgcaaaccag 20

<210> SEQ ID NO 235
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 235

ctggagtctt ccagtgtgat gatg 24

<210> SEQ ID NO 236
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 236

ctggagtctt ccagtgtgat gat 23

<210> SEQ ID NO 237
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 237

tgtgcagggt ggcaagt 17

<210> SEQ ID NO 238
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 238

tgtgcagggt ggcaagt 17

<210> SEQ ID NO 239
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 239

ctttcttgcg gagattctct tcct 24

<210> SEQ ID NO 240
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 240

cagacaactg ttcaaactg 19

<210> SEQ ID NO 241
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 241

ctggcagcaa cagtct 16

<210> SEQ ID NO 242
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 242

tagtggcacc agaattg 17

<210> SEQ ID NO 243
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 243

cccttcgggt gcctcc 16

<210> SEQ ID NO 244
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 244

cagcatgtca agatcac 17

<210> SEQ ID NO 245
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 245

ctggtccttc attgcac 17

<210> SEQ ID NO 246
<211> LENGTH: 16

-continued

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 246

ccctccccag tcctca 16

<210> SEQ ID NO 247
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 247

ccctccccag tcctca 16

<210> SEQ ID NO 248
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 248

cctgctgaaa atgac 15

<210> SEQ ID NO 249
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 249

cctgctgaaa atgac 15

<210> SEQ ID NO 250
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 250

cctgctgaaa atgac 15

<210> SEQ ID NO 251
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 251

cctgctgaaa atgac 15

-continued

<210> SEQ ID NO 252
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 252

cctgctgaaa atgac 15

<210> SEQ ID NO 253
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 253

cctgctgaaa atgac 15

<210> SEQ ID NO 254
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 254

tcgtccacaa aatgattc 18

<210> SEQ ID NO 255
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 255

ccttcgcctg tcctcatg 18

<210> SEQ ID NO 256
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 256

ccttcgcctg tcctcatg 18

<210> SEQ ID NO 257
<211> LENGTH: 14
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 257

cagtgcgctt ttcc 14

-continued

<210> SEQ ID NO 258
<211> LENGTH: 14
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 258

cagtgcgctt ttcc 14

<210> SEQ ID NO 259
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 259

ctgctcacca tcgctatc 18

<210> SEQ ID NO 260
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 260

cctcaggcgg ctcatag 17

<210> SEQ ID NO 261
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 261

atgggcctcc gggtcat 17

<210> SEQ ID NO 262
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 262

accggaggcc catcct 16

<210> SEQ ID NO 263
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 263

-continued

cacactggaa gactcc 16

<210> SEQ ID NO 264
 <211> LENGTH: 16
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 probe

<400> SEQUENCE: 264

cacactggaa gactcc 16

<210> SEQ ID NO 265
 <211> LENGTH: 15
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 probe

<400> SEQUENCE: 265

ctgtgcgcg gtctc 15

<210> SEQ ID NO 266
 <211> LENGTH: 16
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 probe

<400> SEQUENCE: 266

gctacagaga aatctc 16

<210> SEQ ID NO 267
 <211> LENGTH: 17
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 probe

<400> SEQUENCE: 267

gagctgtggt agtggca 17

<210> SEQ ID NO 268
 <211> LENGTH: 16
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 probe

<400> SEQUENCE: 268

cactcagaga aggagc 16

<210> SEQ ID NO 269
 <211> LENGTH: 16
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 probe

-continued

<400> SEQUENCE: 269

catcacgcag ctcatg 16

<210> SEQ ID NO 270

<211> LENGTH: 16

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 270

agtttgcccc gcccaa 16

<210> SEQ ID NO 271

<211> LENGTH: 17

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 271

ctcgacacag caggcca 17

<210> SEQ ID NO 272

<211> LENGTH: 18

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 272

gcaggccaag aggagta 18

<210> SEQ ID NO 273

<211> LENGTH: 18

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 273

gcaggccaag aggagta 18

<210> SEQ ID NO 274

<211> LENGTH: 14

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 274

cgccactagc tcca 14

<210> SEQ ID NO 275

<211> LENGTH: 16

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

-continued

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 275

ccacgagctc caacta 16

<210> SEQ ID NO 276

<211> LENGTH: 14

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 276

acgccaccag ctcc 14

<210> SEQ ID NO 277

<211> LENGTH: 13

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 277

cgccatcagc tcc 13

<210> SEQ ID NO 278

<211> LENGTH: 13

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 278

cgccagcagc tcc 13

<210> SEQ ID NO 279

<211> LENGTH: 14

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 279

cgccaccagc tcca 14

<210> SEQ ID NO 280

<211> LENGTH: 15

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 280

gctggtgacg taggc 15

<210> SEQ ID NO 281

<211> LENGTH: 18

<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 281

gctggaaaag aagagtac 18

<210> SEQ ID NO 282
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 282

cagctggaca agaagag 17

<210> SEQ ID NO 283
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 283

ttggagcaga tgggtgtt 17

<210> SEQ ID NO 284
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 284

tggagcaggt gatgtt 16

<210> SEQ ID NO 285
<211> LENGTH: 13
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 285

tgaggcactg ccc 13

<210> SEQ ID NO 286
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 286

gaaacacttt tcgacatagt gtggtg 26

<210> SEQ ID NO 287

-continued

<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 287

tgtgtaacag tgcctgcatg 20

<210> SEQ ID NO 288
<211> LENGTH: 14
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 288

gcatgggcag catg 14

<210> SEQ ID NO 289
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 289

gcatgaaccg gaggc 15

<210> SEQ ID NO 290
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 290

catgaaccag aggcc 15

<210> SEQ ID NO 291
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 291

ttgaggtgcg tggtttgtg 18

<210> SEQ ID NO 292
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 292

gctacagtga aatctc 16

-continued

<210> SEQ ID NO 293
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 293

agctgtggca gtggca 16

<210> SEQ ID NO 294
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 294

cactcagaaa aggagc 16

<210> SEQ ID NO 295
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 295

tcatcatgca gctcatg 17

<210> SEQ ID NO 296
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 296

agtttggcca gcccaa 16

<210> SEQ ID NO 297
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 297

ctcgacacag gaggtca 17

<210> SEQ ID NO 298
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 298

-continued

cagggtcacga ggagtac 17

<210> SEQ ID NO 299
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 299

gcagggtcatg aggagtac 18

<210> SEQ ID NO 300
<211> LENGTH: 13
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 300

cgccaccagc tcc 13

<210> SEQ ID NO 301
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 301

ccaccagctc caacta 16

<210> SEQ ID NO 302
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 302

acgccacaag ctccaa 16

<210> SEQ ID NO 303
<211> LENGTH: 13
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 303

cgccaccagc tcc 13

<210> SEQ ID NO 304
<211> LENGTH: 13
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

-continued

<400> SEQUENCE: 304

cgccaccagc tcc 13

<210> SEQ ID NO 305

<211> LENGTH: 15

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 305

cgccaacagc tccaa 15

<210> SEQ ID NO 306

<211> LENGTH: 15

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 306

gctggtggcg taggc 15

<210> SEQ ID NO 307

<211> LENGTH: 18

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 307

gctggacaag aagagtac 18

<210> SEQ ID NO 308

<211> LENGTH: 17

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 308

cagctggaca tgaagag 17

<210> SEQ ID NO 309

<211> LENGTH: 16

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 309

tggagcaggt ggtgtt 16

<210> SEQ ID NO 310

<211> LENGTH: 16

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic

-continued

probe

<400> SEQUENCE: 310

tggagcaggt ggtgtt 16

<210> SEQ ID NO 311
<211> LENGTH: 13
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 311

tgaggcgtg ccc 13

<210> SEQ ID NO 312
<211> LENGTH: 27
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 312

agaaacactt tttgacatag tgtggtg 27

<210> SEQ ID NO 313
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 313

atgtgtaaca gttcctgcat g 21

<210> SEQ ID NO 314
<211> LENGTH: 14
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 314

gcatgggcgg catg 14

<210> SEQ ID NO 315
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 315

gcatgaactg gaggcc 16

<210> SEQ ID NO 316
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 316
catgaaccgg aggcc 15

<210> SEQ ID NO 317
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 317
tttgagggtgt gtgtttgtgc 20

<210> SEQ ID NO 318
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 318
ccctggtagt tggagctgg 19

<210> SEQ ID NO 319
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 319
cgctgtggta gttggagctg 20

<210> SEQ ID NO 320
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 320
gcccggtagt tggagctgg 19

<210> SEQ ID NO 321
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 321
cgccgtggta gttggagctg 20

<210> SEQ ID NO 322
<211> LENGTH: 19

-continued

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 322
ccccagttgg agctggtgg 19

<210> SEQ ID NO 323
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 323
gccttgcccta cgccaca 17

<210> SEQ ID NO 324
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 324
gcctcttgcc tacgccaa 18

<210> SEQ ID NO 325
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 325
aaagaatggt cctgcaccag taa 23

<210> SEQ ID NO 326
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 326
ggtcctgcac cagtaatatg ca 22

<210> SEQ ID NO 327
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 327
aaagaatggt cctgcaccag taa 23

-continued

<210> SEQ ID NO 328
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 328
aaagaatggt cctgcaccag taa 23

<210> SEQ ID NO 329
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 329
aaagaatggt cctgcaccag taa 23

<210> SEQ ID NO 330
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 330
gagtttgat taaaaggtagc tggtaggag 29

<210> SEQ ID NO 331
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 331
gagtttgat taaaaggtagc tggtaggag 29

<210> SEQ ID NO 332
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 332
ccctggtagt tggagctgc 19

<210> SEQ ID NO 333
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 333
cgctgtgga gttggagctc 20

-continued

<210> SEQ ID NO 334
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 334
gcccggtagt tggagctga 19

<210> SEQ ID NO 335
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 335
cgccgtggta gttggagcta 20

<210> SEQ ID NO 336
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 336
ccccagttgg agctggtga 19

<210> SEQ ID NO 337
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 337
gccttgccca cgccacc 17

<210> SEQ ID NO 338
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 338
gcctcttgcc tacgccac 18

<210> SEQ ID NO 339
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 339

-continued

atgcatatta aaacaagatt tac 23

<210> SEQ ID NO 340
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 340

aacaatagag gtaaatcttg 20

<210> SEQ ID NO 341
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 341

atgcatatta aaacaagatt tac 23

<210> SEQ ID NO 342
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 342

atgcatatta aaacaagatt tac 23

<210> SEQ ID NO 343
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 343

atgcatatta aaacaagatt tac 23

<210> SEQ ID NO 344
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 344

accttatgtg tgacatgttc 20

<210> SEQ ID NO 345
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

-continued

<400> SEQUENCE: 345

accttatgtg tgacatgttc

20

<210> SEQ ID NO 346

<211> LENGTH: 16

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 346

tggagctgct ggcgta

16

<210> SEQ ID NO 347

<211> LENGTH: 16

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 347

tggagctcgt ggcgta

16

<210> SEQ ID NO 348

<211> LENGTH: 17

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 348

ttggagctga tggcgta

17

<210> SEQ ID NO 349

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 349

tagttggagc tagtggcgta

20

<210> SEQ ID NO 350

<211> LENGTH: 17

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 350

ttggagctgg tgacgta

17

<210> SEQ ID NO 351

<211> LENGTH: 14

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

-continued

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 351

cgccaccagc tcca 14

<210> SEQ ID NO 352

<211> LENGTH: 15

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 352

ctacgccacc agctc 15

<210> SEQ ID NO 353

<211> LENGTH: 16

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 353

tggagctggt ggcgta 16

<210> SEQ ID NO 354

<211> LENGTH: 16

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 354

tggagctggt ggcgta 16

<210> SEQ ID NO 355

<211> LENGTH: 16

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 355

tggagctggt ggcgta 16

<210> SEQ ID NO 356

<211> LENGTH: 16

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 356

tggagctggt ggcgta 16

<210> SEQ ID NO 357

<211> LENGTH: 16

<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 357

tggagctggt ggcgta 16

<210> SEQ ID NO 358
<211> LENGTH: 14
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 358

cgccacaagc tcca 14

<210> SEQ ID NO 359
<211> LENGTH: 14
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 359

tacgccaaaca gctc 14

<210> SEQ ID NO 360
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 360

tgattttggt ctagctacag a 21

<210> SEQ ID NO 361
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 361

gagaaggagc tgtggt 16

<210> SEQ ID NO 362
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 362

tgcctttacc actcagag 18

<210> SEQ ID NO 363

-continued

<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 363

cgtgcagctc atcac 15

<210> SEQ ID NO 364
<211> LENGTH: 14
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 364

cagcagtttg gccc 14

<210> SEQ ID NO 365
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 365

ggatattctc gacacagc 18

<210> SEQ ID NO 366
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 366

gacacagcag gtcaa 15

<210> SEQ ID NO 367
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 367

cttgctacg ccaact 15

<210> SEQ ID NO 368
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 368

ctcttgccca cgccat 16

-continued

<210> SEQ ID NO 369
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 369

gtagttggag ctggtga 17

<210> SEQ ID NO 370
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 370

atactggata cagctggaa 19

<210> SEQ ID NO 371
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 371

tggatacagc tggacaa 17

<210> SEQ ID NO 372
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 372

gattttggtc tagctacagt 20

<210> SEQ ID NO 373
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 373

agaaggagct gtggc 15

<210> SEQ ID NO 374
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 374

-continued

tgcctttacc actcagaa 18

<210> SEQ ID NO 375
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 375

cgtgcagctc atcat 15

<210> SEQ ID NO 376
<211> LENGTH: 14
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 376

cagcagtttg gccca 14

<210> SEQ ID NO 377
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 377

tggatattct cgacacagg 19

<210> SEQ ID NO 378
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 378

gacacagcag gtcac 15

<210> SEQ ID NO 379
<211> LENGTH: 14
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 379

ttgcctacgc cacc 14

<210> SEQ ID NO 380
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

-continued

<400> SEQUENCE: 380

tcttgccctac gccac 15

<210> SEQ ID NO 381

<211> LENGTH: 16

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 381

tagttggagc tggtagg 16

<210> SEQ ID NO 382

<211> LENGTH: 19

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 382

atactggata cagctggac 19

<210> SEQ ID NO 383

<211> LENGTH: 18

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 383

ctggatacag ctggacat 18

<210> SEQ ID NO 384

<211> LENGTH: 24

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 384

gcctgatttt ggtctagcta caga 24

<210> SEQ ID NO 385

<211> LENGTH: 19

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 385

cgcgagaagg agctgtggt 19

<210> SEQ ID NO 386

<211> LENGTH: 21

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic

-continued

primer

<400> SEQUENCE: 386

ccgtgccttt accactcaga g 21

<210> SEQ ID NO 387
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 387

gcgcgtgcag ctcatcac 18

<210> SEQ ID NO 388
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 388

cggcagcagt ttggccc 17

<210> SEQ ID NO 389
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 389

gccggatatt ctcgacacag c 21

<210> SEQ ID NO 390
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 390

gcggacacag caggtcaa 18

<210> SEQ ID NO 391
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 391

cgccttgct acgccact 18

<210> SEQ ID NO 392
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 392
cgctctttgc ctacgccat 19

<210> SEQ ID NO 393
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 393
cgcgtagttg gagctggtga 20

<210> SEQ ID NO 394
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 394
ccgatactgg atacagctgg aa 22

<210> SEQ ID NO 395
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 395
ccgtggatac agctggacaa 20

<210> SEQ ID NO 396
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 396
ccggattttg gtctagctac agt 23

<210> SEQ ID NO 397
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 397
gccagaagga gctgtggc 18

<210> SEQ ID NO 398
<211> LENGTH: 21

-continued

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 398
ccgtgccttt accactcaga a 21

<210> SEQ ID NO 399
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 399
gcgcgtgcag ctcacatc 18

<210> SEQ ID NO 400
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 400
cggcagcagt ttggcca 17

<210> SEQ ID NO 401
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 401
cgctggatat tctcgacaca gg 22

<210> SEQ ID NO 402
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 402
gcggacacag caggtcac 18

<210> SEQ ID NO 403
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 403
gcgttgccca cgccacc 17

-continued

<210> SEQ ID NO 404
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 404
gcgtcttgcc tacgccac 18

<210> SEQ ID NO 405
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 405
gcctagttgg agctggtgg 19

<210> SEQ ID NO 406
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 406
ccgatactgg atacagctgg ac 22

<210> SEQ ID NO 407
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 407
cgcttgata cagctggaca t 21

<210> SEQ ID NO 408
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 408
agcctcaatt cttaccatcc acaaaa 26

<210> SEQ ID NO 409
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 409
catggaacca gacagaaaag cg 22

-continued

<210> SEQ ID NO 410
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 410

gtcactggca gcaacagtct ta 22

<210> SEQ ID NO 411
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 411

tgggagccaa tattgtcttt gtgtt 25

<210> SEQ ID NO 412
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 412

aggaacgtac tggtgaaaac acc 23

<210> SEQ ID NO 413
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 413

ccctcccag tctcatgta 20

<210> SEQ ID NO 414
<211> LENGTH: 41
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 414

tcttcaaatg atttagtatt atttatggca aatacacaaa g 41

<210> SEQ ID NO 415
<211> LENGTH: 28
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 415

-continued

tgtgtgacat gttctaatat agtcacat 28

<210> SEQ ID NO 416
<211> LENGTH: 28
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 416

tgtgtgacat gttctaatat agtcacat 28

<210> SEQ ID NO 417
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 417

ggctcctgcac cagtaatatg ca 22

<210> SEQ ID NO 418
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 418

gcaaatgact tgctattatt gatggcaaa 29

<210> SEQ ID NO 419
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 419

gcaaatgact tgctattatt gatggcaaa 29

<210> SEQ ID NO 420
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 420

cagacaactg ttcaaactg 19

<210> SEQ ID NO 421
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

-continued

<400> SEQUENCE: 421

ctggcagcaa cagtct

16

<210> SEQ ID NO 422

<211> LENGTH: 17

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 422

tagtggcacc agaattgg

17

<210> SEQ ID NO 423

<211> LENGTH: 16

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 423

cccttcggtc gcctcc

16

<210> SEQ ID NO 424

<211> LENGTH: 17

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 424

cagcatgtca agatcac

17

<210> SEQ ID NO 425

<211> LENGTH: 17

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 425

ctggtccttc attgcac

17

<210> SEQ ID NO 426

<211> LENGTH: 16

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 426

ccctccccag tctcca

16

<210> SEQ ID NO 427

<211> LENGTH: 15

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

-continued

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 427

cctgctgaaa atgac 15

<210> SEQ ID NO 428

<211> LENGTH: 15

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 428

cctgctgaaa atgac 15

<210> SEQ ID NO 429

<211> LENGTH: 18

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 429

tcgtccacaa aatgattc 18

<210> SEQ ID NO 430

<211> LENGTH: 18

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 430

ccttcgcctg tctcatg 18

<210> SEQ ID NO 431

<211> LENGTH: 18

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 431

ccttcgcctg tctcatg 18

<210> SEQ ID NO 432

<211> LENGTH: 14

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 432

ctacagagaa atct 14

<210> SEQ ID NO 433

<211> LENGTH: 14

<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 433

ctgtggtagt ggca 14

<210> SEQ ID NO 434
<211> LENGTH: 14
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 434

ctcagagaag gagc 14

<210> SEQ ID NO 435
<211> LENGTH: 13
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 435

atcacgcagc tca 13

<210> SEQ ID NO 436
<211> LENGTH: 11
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 436

ggcccgccca a 11

<210> SEQ ID NO 437
<211> LENGTH: 14
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 437

gacacagcag gtca 14

<210> SEQ ID NO 438
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 438

caggtcaaga ggagta 16

<210> SEQ ID NO 439

-continued

<211> LENGTH: 13
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 439

gccactagct cca 13

<210> SEQ ID NO 440
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 440

gccatcagct cc 12

<210> SEQ ID NO 441
<211> LENGTH: 13
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 441

tggtgacgta ggc 13

<210> SEQ ID NO 442
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 442

gctggaaaag aagag 15

<210> SEQ ID NO 443
<211> LENGTH: 14
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 443

ctggacaaga agag 14

<210> SEQ ID NO 444
<211> LENGTH: 14
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 444

ctacagtga atct 14

-continued

<210> SEQ ID NO 445
<211> LENGTH: 13
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 445

tgtggcagtg gca 13

<210> SEQ ID NO 446
<211> LENGTH: 14
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 446

ctcagaaaag gagc 14

<210> SEQ ID NO 447
<211> LENGTH: 14
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 447

catcatgcag ctca 14

<210> SEQ ID NO 448
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 448

tggccagccc aa 12

<210> SEQ ID NO 449
<211> LENGTH: 14
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 449

gacacaggag gtca 14

<210> SEQ ID NO 450
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 450

-continued

aggtcacgag gagta 15

<210> SEQ ID NO 451
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 451

ccaccagctc ca 12

<210> SEQ ID NO 452
<211> LENGTH: 11
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 452

ccaccagctc c 11

<210> SEQ ID NO 453
<211> LENGTH: 13
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 453

tggtggcgta ggc 13

<210> SEQ ID NO 454
<211> LENGTH: 14
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 454

ctggacaaga agag 14

<210> SEQ ID NO 455
<211> LENGTH: 14
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 455

ctggacatga agag 14

<210> SEQ ID NO 456
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

-continued

<400> SEQUENCE: 456

gtgattttgg tctagctaca ga

22

<210> SEQ ID NO 457

<211> LENGTH: 19

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 457

tcagagaagg agctgtggt

19

<210> SEQ ID NO 458

<211> LENGTH: 21

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 458

gattgccttt accactcaga g

21

<210> SEQ ID NO 459

<211> LENGTH: 18

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 459

caccgtgcag ctcacac

18

<210> SEQ ID NO 460

<211> LENGTH: 18

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 460

caccagcag ttggccc

18

<210> SEQ ID NO 461

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 461

ttggatattc tcgacacagc

20

<210> SEQ ID NO 462

<211> LENGTH: 18

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic

-continued

primer

<400> SEQUENCE: 462

ctcgacacag caggtcaa

18

<210> SEQ ID NO 463

<211> LENGTH: 18

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 463

actcttgcc acgccact

18

<210> SEQ ID NO 464

<211> LENGTH: 18

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 464

cactcttgcc tacgccat

18

<210> SEQ ID NO 465

<211> LENGTH: 19

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 465

tggtagttgg agctgggtga

19

<210> SEQ ID NO 466

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 466

catactggat acagctggaa

20

<210> SEQ ID NO 467

<211> LENGTH: 21

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 467

atactggata cagctggaca a

21

<210> SEQ ID NO 468

<211> LENGTH: 22

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 468

gtgattttgg tctagctaca gt 22

<210> SEQ ID NO 469
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 469

tcagagaagg agctgtggc 19

<210> SEQ ID NO 470
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 470

gattgccttt accactcaga a 21

<210> SEQ ID NO 471
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 471

caccgtgcag ctcacat 18

<210> SEQ ID NO 472
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 472

caccagcag ttggcca 18

<210> SEQ ID NO 473
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 473

cttggatatt ctcgacacag g 21

<210> SEQ ID NO 474
<211> LENGTH: 19

-continued

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 474
tctcgacaca gcaggtcac 19

<210> SEQ ID NO 475
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 475
actcttgcct acgccacc 18

<210> SEQ ID NO 476
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 476
cactcttgcc tacgccac 18

<210> SEQ ID NO 477
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 477
tggtagtgg agctggtgg 19

<210> SEQ ID NO 478
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 478
catactggat acagctggac 20

<210> SEQ ID NO 479
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 479
tactggatac agctggacat 20

-continued

<210> SEQ ID NO 480
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 480
agcctcaatt cttaccatcc acaaaa 26

<210> SEQ ID NO 481
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 481
catggaacca gacagaaaag cg 22

<210> SEQ ID NO 482
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 482
gtcactggca gcaacagtct ta 22

<210> SEQ ID NO 483
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 483
tgggagccaa tattgtcttt gtgtt 25

<210> SEQ ID NO 484
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 484
aggaacgtac tggtgaaaac acc 23

<210> SEQ ID NO 485
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 485
ccctcccag tcctcatgta 20

-continued

<210> SEQ ID NO 486
<211> LENGTH: 41
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 486

tcttcaaatg atttagtatt atttatggca aatacacaaa g 41

<210> SEQ ID NO 487
<211> LENGTH: 28
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 487

tgtgtgacat gttctaatat agtcacat 28

<210> SEQ ID NO 488
<211> LENGTH: 28
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 488

tgtgtgacat gttctaatat agtcacat 28

<210> SEQ ID NO 489
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 489

ggtcctgcac cagtaatatg ca 22

<210> SEQ ID NO 490
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 490

gcaaatgact tgctattatt gatggcaaa 29

<210> SEQ ID NO 491
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 491

-continued

gcaaatgact tgctattatt gatggcaaa 29

<210> SEQ ID NO 492
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 492

cagacaactg ttcaaactg 19

<210> SEQ ID NO 493
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 493

ctggcagcaa cagtct 16

<210> SEQ ID NO 494
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 494

tagtggcacc agaattgg 17

<210> SEQ ID NO 495
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 495

cccttcggct gcctcc 16

<210> SEQ ID NO 496
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 496

cagcatgtca agatcac 17

<210> SEQ ID NO 497
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

-continued

<400> SEQUENCE: 497

ctgggtccctc attgcac

17

<210> SEQ ID NO 498

<211> LENGTH: 16

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 498

ccctccccag tctctca

16

<210> SEQ ID NO 499

<211> LENGTH: 15

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 499

cctgctgaaa atgac

15

<210> SEQ ID NO 500

<211> LENGTH: 15

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 500

cctgctgaaa atgac

15

<210> SEQ ID NO 501

<211> LENGTH: 18

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 501

tcgtccacaa aatgattc

18

<210> SEQ ID NO 502

<211> LENGTH: 18

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 502

ccttcgcctg tctcatg

18

<210> SEQ ID NO 503

<211> LENGTH: 18

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

-continued

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 503

ccttcgectg tctcatg 18

<210> SEQ ID NO 504

<211> LENGTH: 21

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 504

gctacagaga aatctcgatg g 21

<210> SEQ ID NO 505

<211> LENGTH: 17

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 505

gagctgtggt agtggca 17

<210> SEQ ID NO 506

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 506

ttaccactca gagaaggagc 20

<210> SEQ ID NO 507

<211> LENGTH: 17

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 507

gctcatcacg cagctca 17

<210> SEQ ID NO 508

<211> LENGTH: 16

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 508

ttggcccgcc caaaat 16

<210> SEQ ID NO 509

<211> LENGTH: 18

<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 509

gacacagcag gtcacgag 18

<210> SEQ ID NO 510
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 510

cagcagggtca agaggagtac 20

<210> SEQ ID NO 511
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 511

ctacgccact agctccaac 19

<210> SEQ ID NO 512
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 512

ctacgccatc agctcc 16

<210> SEQ ID NO 513
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 513

ctggtgacgt aggcaag 17

<210> SEQ ID NO 514
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 514

ggatacagct ggaaaagaag ag 22

<210> SEQ ID NO 515

-continued

<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 515

agctggacaa gaagagtaca 20

<210> SEQ ID NO 516
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 516

tagctacagt gaaatctcga tg 22

<210> SEQ ID NO 517
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 517

gagctgtggc agtgg 15

<210> SEQ ID NO 518
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 518

ttaccactca gaaaaggagc t 21

<210> SEQ ID NO 519
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 519

cagctcatca tgcagctcat 20

<210> SEQ ID NO 520
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 520

gtttggccag cccaaaatc 19

-continued

<210> SEQ ID NO 521
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 521

gacacaggag gtcaggag 18

<210> SEQ ID NO 522
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 522

cagcagggtca cgaggagta 19

<210> SEQ ID NO 523
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 523

taagccacca gtccaa 17

<210> SEQ ID NO 524
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 524

taagccacca gtcaca 16

<210> SEQ ID NO 525
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 525

ctgggtggcgt aggcaag 17

<210> SEQ ID NO 526
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 526

-continued

atacagctgg acaagaagag t 21

<210> SEQ ID NO 527
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 527

agctggacat gaagagtaca 20

<210> SEQ ID NO 528
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 528

cccccttttg tctagctaca ga 22

<210> SEQ ID NO 529
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 529

gcccgaagga gctgtggt 18

<210> SEQ ID NO 530
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 530

cgccccttta ccactcagag 20

<210> SEQ ID NO 531
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 531

gcgggtgcag ctcacac 18

<210> SEQ ID NO 532
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

-continued

<400> SEQUENCE: 532

cgggagcagt ttggccc

17

<210> SEQ ID NO 533

<211> LENGTH: 21

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 533

gcccgatatt ctcgacacag c

21

<210> SEQ ID NO 534

<211> LENGTH: 18

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 534

gcggacacag caggtcaa

18

<210> SEQ ID NO 535

<211> LENGTH: 16

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 535

cggtgcctac gccact

16

<210> SEQ ID NO 536

<211> LENGTH: 17

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 536

cgccttgccct acgccat

17

<210> SEQ ID NO 537

<211> LENGTH: 19

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 537

gccacgttgg agctggtga

19

<210> SEQ ID NO 538

<211> LENGTH: 21

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic

-continued

```

    primer

<400> SEQUENCE: 538

cgccactgga tacagctgga a                21

<210> SEQ ID NO 539
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
    primer

<400> SEQUENCE: 539

ccgcggatac agctggacaa                20

<210> SEQ ID NO 540
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
    primer

<400> SEQUENCE: 540

cccccttttg tctagctaca gt            22

<210> SEQ ID NO 541
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
    primer

<400> SEQUENCE: 541

gcccgaagga gctgtggc                18

<210> SEQ ID NO 542
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
    primer

<400> SEQUENCE: 542

cgcccccttta ccactcagaa                20

<210> SEQ ID NO 543
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
    primer

<400> SEQUENCE: 543

gcgggtgcag ctcacat                18

<210> SEQ ID NO 544
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence

```

-continued

<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 544

cgggagcagt ttggcca 17

<210> SEQ ID NO 545
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 545

gccccgatatt ctcgacacag g 21

<210> SEQ ID NO 546
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 546

gcggacacag caggtcac 18

<210> SEQ ID NO 547
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 547

cggtgcctac gccacc 16

<210> SEQ ID NO 548
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 548

cgccttgctt acgccac 17

<210> SEQ ID NO 549
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 549

gcccagttgg agctgggtgg 19

<210> SEQ ID NO 550
<211> LENGTH: 21

-continued

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 550
cgccactgga tacagctgga c 21

<210> SEQ ID NO 551
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 551
ccgcggatac agctggacat 20

<210> SEQ ID NO 552
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 552
agcctcaatt cttaccatcc acaaaa 26

<210> SEQ ID NO 553
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 553
catggaacca gacagaaaag cg 22

<210> SEQ ID NO 554
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 554
gtcactggca gcaacagtct ta 22

<210> SEQ ID NO 555
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 555
tgggagccaa tattgtcttt gtgtt 25

-continued

<210> SEQ ID NO 556
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 556
aggaacgtac tggtgaaaac acc 23

<210> SEQ ID NO 557
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 557
ccctccccag tctctcatgta 20

<210> SEQ ID NO 558
<211> LENGTH: 41
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 558
tcttcaaatg atttagtatt atttatggca aatacacaaa g 41

<210> SEQ ID NO 559
<211> LENGTH: 28
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 559
tgtgtgacat gttctaatat agtcacat 28

<210> SEQ ID NO 560
<211> LENGTH: 28
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 560
tgtgtgacat gttctaatat agtcacat 28

<210> SEQ ID NO 561
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 561
ggctcctgcac cagtaatatg ca 22

-continued

<210> SEQ ID NO 562
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 562

gcaaatgact tgctattatt gatggcaaa 29

<210> SEQ ID NO 563
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 563

gcaaatgact tgctattatt gatggcaaa 29

<210> SEQ ID NO 564
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 564

cagacaactg ttcaaaactg 19

<210> SEQ ID NO 565
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 565

ctggcagcaa cagtct 16

<210> SEQ ID NO 566
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 566

tagtggcacc agaattgg 17

<210> SEQ ID NO 567
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 567

-continued

cccttcgggt gcctcc 16

<210> SEQ ID NO 568
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 568

cagcatgtca agatcac 17

<210> SEQ ID NO 569
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 569

ctggtcctc attgac 17

<210> SEQ ID NO 570
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 570

ccctccccag tctca 16

<210> SEQ ID NO 571
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 571

cctgctgaaa atgac 15

<210> SEQ ID NO 572
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 572

cctgctgaaa atgac 15

<210> SEQ ID NO 573
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

-continued

<400> SEQUENCE: 573

tcgtccacaa aatgattc

18

<210> SEQ ID NO 574

<211> LENGTH: 18

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 574

ccttcgcctg tcctcatg

18

<210> SEQ ID NO 575

<211> LENGTH: 18

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 575

ccttcgcctg tcctcatg

18

<210> SEQ ID NO 576

<211> LENGTH: 14

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 576

ctacagagaa atct

14

<210> SEQ ID NO 577

<211> LENGTH: 14

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 577

ctgtggtagt ggca

14

<210> SEQ ID NO 578

<211> LENGTH: 14

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 578

ctcagagaag gagc

14

<210> SEQ ID NO 579

<211> LENGTH: 13

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

-continued

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 579

atcacgcgacg tca 13

<210> SEQ ID NO 580

<211> LENGTH: 11

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 580

ggcccgccca a 11

<210> SEQ ID NO 581

<211> LENGTH: 14

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 581

gacacagcag gtca 14

<210> SEQ ID NO 582

<211> LENGTH: 16

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 582

cagggtcaaga ggagta 16

<210> SEQ ID NO 583

<211> LENGTH: 13

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 583

gccactagct cca 13

<210> SEQ ID NO 584

<211> LENGTH: 12

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 584

gccatcagct cc 12

<210> SEQ ID NO 585

<211> LENGTH: 13

<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 585

tggtgacgta ggc 13

<210> SEQ ID NO 586
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 586

gctggaaaag aagag 15

<210> SEQ ID NO 587
<211> LENGTH: 14
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 587

ctggacaaga agag 14

<210> SEQ ID NO 588
<211> LENGTH: 14
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 588

ctacagtga atct 14

<210> SEQ ID NO 589
<211> LENGTH: 13
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 589

tgtggcagtg gca 13

<210> SEQ ID NO 590
<211> LENGTH: 14
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 590

ctcagaaaag gagc 14

<210> SEQ ID NO 591

-continued

<211> LENGTH: 14
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 591

catcatgcag ctca 14

<210> SEQ ID NO 592
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 592

tggccagccc aa 12

<210> SEQ ID NO 593
<211> LENGTH: 14
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 593

gacacaggag gtca 14

<210> SEQ ID NO 594
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 594

aggtcacgag gagta 15

<210> SEQ ID NO 595
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 595

ccaccagctc ca 12

<210> SEQ ID NO 596
<211> LENGTH: 11
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 596

ccaccagctc c 11

-continued

<210> SEQ ID NO 597
<211> LENGTH: 13
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 597

tggtggcgta ggc 13

<210> SEQ ID NO 598
<211> LENGTH: 14
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 598

ctggacaaga agag 14

<210> SEQ ID NO 599
<211> LENGTH: 14
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 599

ctggacatga agag 14

<210> SEQ ID NO 600
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 600

cccccttttg tctagctaca ga 22

<210> SEQ ID NO 601
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer
<220> FEATURE:
<223> OTHER INFORMATION: Description of Combined DNA/RNA Molecule: Synthetic
primer

<400> SEQUENCE: 601

gcccgaagga gctgtggu 18

<210> SEQ ID NO 602
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

-continued

<400> SEQUENCE: 602

cgcccccttta ccaactcagag

20

<210> SEQ ID NO 603

<211> LENGTH: 18

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 603

gcgggtgcag ctcacac

18

<210> SEQ ID NO 604

<211> LENGTH: 17

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 604

cgggagcagt ttggccc

17

<210> SEQ ID NO 605

<211> LENGTH: 21

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 605

gcccgatatt ctcgacacag c

21

<210> SEQ ID NO 606

<211> LENGTH: 26

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 606

agcctcaatt cttaccatcc acaaaa

26

<210> SEQ ID NO 607

<211> LENGTH: 22

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 607

catggaacca gacagaaaag cg

22

<210> SEQ ID NO 608

<211> LENGTH: 22

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

-continued

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 608

gtcactggca gcaacagtct ta 22

<210> SEQ ID NO 609

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 609

tgggagccaa tattgtcttt gtgtt 25

<210> SEQ ID NO 610

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 610

aggaacgtac tggtgaaaac acc 23

<210> SEQ ID NO 611

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 611

ccctccccag tcctcatgta 20

<210> SEQ ID NO 612

<211> LENGTH: 14

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 612

ctacagagaa atct 14

<210> SEQ ID NO 613

<211> LENGTH: 14

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 613

ctgtggtagt ggca 14

<210> SEQ ID NO 614

<211> LENGTH: 14

<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 614

ctcagagaag gagc 14

<210> SEQ ID NO 615
<211> LENGTH: 13
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 615

atcacgcagc tca 13

<210> SEQ ID NO 616
<211> LENGTH: 11
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 616

ggcccgccca a 11

<210> SEQ ID NO 617
<211> LENGTH: 14
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 617

gacacagcag gtca 14

<210> SEQ ID NO 618
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer
<220> FEATURE:
<223> OTHER INFORMATION: Description of Combined DNA/RNA Molecule: Synthetic primer

<400> SEQUENCE: 618

ccccttttgg tctagctaca gu 22

<210> SEQ ID NO 619
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 619

gcccgaagga gctgtggc 18

-continued

<210> SEQ ID NO 620
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 620
cgcccccttta ccactcagaa 20

<210> SEQ ID NO 621
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer
<220> FEATURE:
<223> OTHER INFORMATION: Description of Combined DNA/RNA Molecule: Synthetic primer

<400> SEQUENCE: 621
gcgggtgcag ctcatcau 18

<210> SEQ ID NO 622
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 622
cgggagcagt ttggcca 17

<210> SEQ ID NO 623
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 623
gcccgatatt ctcgacacag g 21

<210> SEQ ID NO 624
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 624
cagacaactg ttcaaactg 19

<210> SEQ ID NO 625
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic

-continued

```

probe

<400> SEQUENCE: 625

ctggcagcaa cagtct                                16

<210> SEQ ID NO 626
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      probe

<400> SEQUENCE: 626

tagtggcacc agaattg                                17

<210> SEQ ID NO 627
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      probe

<400> SEQUENCE: 627

cccttcggct gcctcc                                16

<210> SEQ ID NO 628
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      probe

<400> SEQUENCE: 628

cagcatgtca agatcac                                17

<210> SEQ ID NO 629
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      probe

<400> SEQUENCE: 629

ctggtccctc attgcac                                17

<210> SEQ ID NO 630
<211> LENGTH: 14
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      probe

<400> SEQUENCE: 630

ctacagtgaa atct                                    14

<210> SEQ ID NO 631
<211> LENGTH: 13
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence

```

-continued

```

<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      probe

<400> SEQUENCE: 631

tgtggcagtg gca                                     13

<210> SEQ ID NO 632
<211> LENGTH: 14
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      probe

<400> SEQUENCE: 632

ctcagaaaag gagc                                     14

<210> SEQ ID NO 633
<211> LENGTH: 14
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      probe

<400> SEQUENCE: 633

catcatgcag ctca                                     14

<210> SEQ ID NO 634
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      probe

<400> SEQUENCE: 634

tggccagccc aa                                       12

<210> SEQ ID NO 635
<211> LENGTH: 14
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      probe

<400> SEQUENCE: 635

gacacaggag gtca                                     14

```

We claim:

1. A method for detecting a first allelic variant of a target sequence in a nucleic acid sample suspected of comprising at least a second allelic variant of the target sequence, comprising:

- a) forming a first reaction mixture by combining:
 - i) the nucleic acid sample;
 - ii) a first allele-specific primer, wherein an allele-specific nucleotide portion of the first allele-specific primer is complementary to the first allelic variant of the target sequence;

- iii) a first allele-specific blocker probe that is complementary to a region of the target sequence comprising the second allelic variant, wherein said region encompasses a position corresponding to the binding position of the allele-specific nucleotide portion of the first allele-specific primer, and wherein the first allele-specific blocker probe comprises a minor groove binder;
- iv) a first locus-specific primer that is complementary to a region of the target sequence that is 3' from the first allelic variant and on the opposite strand; and
- v) a first detector probe;

- b) carrying out an amplification reaction on the first reaction mixture using the first locus-specific primer and the first allele-specific primer to form a first amplicon; and
 - c) detecting the first amplicon by detecting a change in a detectable property of the first detector probe, thereby detecting the first allelic variant of the target gene in the nucleic acid sample.
2. The method of claim 1, further comprising using the change in a detectable property of the first detector probe to quantitate the first allelic variant.
3. The method of claim 1, further comprising:
- d) forming a second reaction mixture by combining:
 - i) the nucleic acid sample;
 - ii) a second allele-specific primer, wherein an allele-specific nucleotide portion of the second allele-specific primer is complementary to the second allelic variant of the target sequence;
 - iii) a second allele-specific blocker probe that is complementary to a region of the target sequence comprising the first allelic variant, wherein said region encompasses a position corresponding to the binding position of the allele-specific nucleotide portion of the second allele-specific primer, and wherein the second allele-specific blocker probe comprises a minor groove binder;
 - iv) a second locus-specific primer that is complementary to a region of the target sequence that is 3' from the second allelic variant and on the opposite strand; and
 - v) a second detector probe;
 - e) carrying out an amplification reaction on the second reaction mixture using the second allele-specific primer and the locus-specific primer, to form a second amplicon; and
 - f) detecting the second amplicon by detecting a change in a detectable property of the detector probe, thereby detecting the second allelic variant of the target gene in the nucleic acid sample.
4. The method of claim 3, further comprising comparing the change in a detectable property of the first detector probe in the first reaction mixture to the change in a detectable property of the second detector probe in the second reaction mixture.
5. The method of claim 1 or 3, wherein said first, second or first and second allele-specific primer and/or said first, second, or first and second allele-specific blocker probe comprises at least one modified base.
6. The method of claim 5, wherein said modified base is an 8-aza-7-deaza-dN (ppN) base analog, where N is adenine (A), cytosine (C), guanine (G), or thymine (T).
7. The method of claim 5, wherein said modified base is a locked nucleic acid (LNA) base.
8. The method of claim 5, wherein said modified base is a fdU or iso dC base.
9. The method of claim 5, wherein said modified base is any modified base that increases the T_m between matched and mismatched target sequences or nucleotides.
10. The method of claim 5, wherein said modified base is located at (a) the 3'-end, (b) the 5'-end, (c) at an internal position or at any combination of (a), (b) or (c) within said allele-specific primer and/or allele-specific blocker probe.
11. The method of claim 5, wherein the specificity of said detecting is improved by the inclusion of said modified base in said first, second or first and second allele-specific primer and/or said first, second, or first and second allele-specific blocker probe as compared to when it is not.
12. The method of claim 11, wherein said improvement is at least 2 fold.
13. The method of claim 1 or 3, wherein the specificity of said detecting is improved by at least 2 fold as compared to the specificity of detecting an allelic variant in a nucleic acid sample using ASB-PCR methods.
14. The method of claim 1 or 3, wherein said carrying out an amplification reaction comprises a 2-stage cycling protocol.
15. The method of claim 14, wherein the number of cycles in the first stage of said 2-stage cycling protocol comprises fewer cycles than the number of cycles used in the second stage.
16. The method of claim 14, wherein said number of cycles in the first stage is about 90% fewer cycles than said number of cycles in the second stage.
17. The method of claim 14, wherein said number of cycles in the first stage is between 3-7 cycles and said number of cycles in the second stage is between 42-48 cycles.
18. The method of claim 14, wherein the annealing/extension temperature used during the first cycling stage of said 2-stage cycling protocol is between 1-3° C. lower than the annealing/extension temperature used during the second stage.
19. The method of claim 14, wherein said annealing/extension temperature used during the first cycling stage of said 2-stage cycling protocol is between 56-59° C. and said annealing/extension temperature used during said second stage is between 60-62° C.
20. The method of claim 1, wherein said step (a) is preceded by a pre-amplification step.
21. The method of claim 20, wherein said pre-amplification step comprises a multiplex amplification reaction that uses at least two complete sets of allele-specific primers and locus-specific primers, wherein each set is suitable or operative for amplifying a specific polynucleotide of interest.
22. The method of claim 21, wherein the products of said multiplex amplification reaction are divided into secondary single-plex amplification reactions, wherein each single-plex amplification reaction contains at least one primer set previously used in said multiplex reaction.
23. The method of claim 21, wherein said multiplex amplification reaction further comprises a plurality of allele-specific blocker probes.
24. The method of claim 21, wherein said multiplex amplification reaction is carried out for a number of cycles suitable to keep the reaction within the linear phase of amplification.
25. A reaction mixture comprising:
- a) a nucleic acid molecule;
 - b) an allele-specific primer, wherein an allele-specific nucleotide portion of the allele-specific primer is complementary to a first allelic variant of a target sequence;
 - c) an allele-specific blocker probe that is complementary to a region of the target sequence comprising a second allelic variant, wherein said region encompasses a position corresponding to the binding position of the allele-specific nucleotide portion of the allele-specific primer, and wherein the allele-specific blocker probe comprises a minor groove binder;

- d) a locus-specific primer that is complementary to a region of the target sequence that is 3' from the first allelic variant and on the opposite strand; and
- e) a detector probe.

26. The reaction mixture of claim 25, wherein said allele-specific primer and/or said allele-specific blocker probe comprises at least one modified base.

27. The reaction mixture of claim 26, wherein said modified base is an 8-aza-7-deaza-dN (ppN) base analog, where N is adenine (A), cytosine (C), guanine (G), or thymine (T).

28. The reaction mixture of claim 26, wherein said modified base is a locked nucleic acid (LNA) base.

29. The reaction mixture of claim 26, wherein said modified base is a fdU or iso dC base.

30. The reaction mixture of claim 26, wherein said modified base is any modified base that increases the T_m between matched and mismatched target sequences or nucleotides.

31. The reaction mixture of claim 26, wherein said modified base is located at (a) the 3'-end, (b) the 5'-end, (c) at an internal position or at any combination of (a), (b) or (c) within said allele-specific primer and/or allele-specific blocker probe.

32. A composition comprising:

- a) a first allele-specific primer, wherein an allele-specific nucleotide portion of the first allele-specific primer is complementary to the first allelic variant of a target sequence;
- b) a first allele-specific blocker probe that is complementary to a region of the target sequence comprising the second allelic variant, wherein said region encompasses a position corresponding to the binding position of the allele-specific nucleotide portion of the first allele-specific primer, and wherein the first allele-specific blocker probe comprises a minor groove binder;

33. The composition of claim 32, further comprising a locus-specific primer that is complementary to a region of the target sequence that is 3' from the first allelic variant and on the opposite strand.

34. The composition of claim 32 or 33, wherein said first allele-specific primer and/or said first allele-specific blocker probe comprises at least one modified base.

35. The composition of claim 34, wherein said modified base is an 8-aza-7-deaza-dN (ppN) base analog, where N is adenine (A), cytosine (C), guanine (G), or thymine (T).

36. The composition of claim 34, wherein said modified base is a locked nucleic acid (LNA) base.

37. The composition of claim 34, wherein said modified base is a fdU or iso dC base.

38. The composition of claim 34, wherein said modified base is any modified base that increases the T_m between matched and mismatched target sequences or nucleotides.

39. The composition of claim 34, wherein said modified base is located at (a) the 3'-end, (b) the 5'-end, (c) at an internal position or at any combination of (a), (b) or (c) within said allele-specific primer and/or allele-specific blocker probe.

40. A kit comprising, two or more containers comprising the following components independently distributed in one of the two or more containers:

- a) a first allele-specific primer, wherein an allele-specific nucleotide portion of the first allele-specific primer is complementary to the first allelic variant of a target sequence; and
- b) a first allele-specific blocker probe that is complementary to a region of the target sequence comprising the

second allelic variant, wherein said region encompasses a position corresponding to the binding position of the allele-specific nucleotide portion of the first allele-specific primer, and wherein the first allele-specific blocker probe comprises a minor groove binder.

41. The kit of claim 40, further comprising a locus-specific primer that is complementary to a region of the target sequence that is 3' from the first allelic variant and on the opposite strand.

42. The kit of claim 41, wherein said first allele-specific primer and/or first allele-specific blocker probe comprises at least one modified base.

43. The kit of claim 42, wherein said modified base is a 8-aza-7-deaza-dN (ppN) base analog, where N is adenine (A), cytosine (C), guanine (G), or thymine (T).

44. The kit of claim 42, wherein said modified base is a locked nucleic acid (LNA) base.

45. The kit of claim 42, wherein said modified base is any modified base that increases the T_m between matched and mismatched target sequences and/or nucleotides.

46. The kit of claim 42, wherein said allele-specific blocker probe comprises an MGB moiety at the 3'-end, the 5'-end and/or at an internal position within said allele-specific blocker probe.

47. A method for detecting a first allelic variant in a target sequence in a nucleic acid sample, comprising:

- a) forming a reaction mixture comprising:
 - i) a nucleic acid sample;
 - ii) an allele-specific primer, wherein an allele-specific nucleotide portion of the allele-specific primer is complementary to the first allelic variant of the target sequence;
 - iii) an allele-specific blocker probe that is complementary to a region of the target sequence comprising a second allelic variant, wherein said region encompasses a position corresponding to the binding position of the allele-specific nucleotide portion of the allele-specific primer, and wherein the allele-specific blocker probe comprises a blocking moiety;
 - iv) a locus-specific primer that is complementary to a region of the target sequence that is 3' from the first allelic variant and on the opposite strand; and
 - v) a detector probe;
- b) PCR amplifying the target sequence using a 2-stage cycling protocol, comprising:
 - i) a first amplification step comprising a first number of cycles run at a first annealing/extension temperature; and
 - ii) a second amplification step comprising a second number of cycles run at a second annealing/extension temperature,

wherein the first number of cycles is fewer than the second number of cycles and the first annealing/extension temperature is lower than said second annealing/extension temperature.

c) detecting a change in a detectable property of the detector probe in the amplified products of the target sequence produced by step (b), thereby detecting the first allelic variant of the target gene in the nucleic acid sample.

48. The method of claim 47, wherein said number of cycles in the first step is about 90% fewer cycles than said number of cycles in the second step.

49. The method of claim **14**, wherein said number of cycles in the first step is between 3-7 cycles and said number of cycles in the second step is between 42-48 cycles.

50. The method of claim **14**, wherein the annealing/extension temperature used during the first step of said 2-stage cycling protocol is between 1-3° C. lower than the annealing/extension temperature used during the second step.

51. The method of claim **14**, wherein said annealing/extension temperature used during the first step of said 2-stage cycling protocol is between 56-59° C. and said annealing/extension temperature used during said second step is between 60-62° C.

52. A method for detecting an allelic variant of a target sequence in a nucleic acid sample, comprising:

- a) forming in a single vessel a first reaction mixture comprising:
 - i) a nucleic acid sample; and
 - ii) at least two sets of primers wherein each set comprises (a) a first allele-specific primer and a second allele-specific primer, wherein the allele-specific nucleotide portion of said first and second allele-specific primers is complementary to a first allele and a second allele, respectively, of a given SNP within a target sequence; and (b) a locus-specific primer that is complementary to a region of said target sequence that is 3' from the first and second alleles and on the opposite strand; wherein the at least two sets of primers is each specific for a different target sequence;

b) amplifying said different target sequences using a number of cycles suitable to keep the reaction within a linear phase;

c) dividing the amplified products of step (b) into at least two separate vessels;

d) adding to each of the divided products of step (c)

- i) at least one set of primers used in step (a),
- ii) an allele-specific blocker probe that is complementary to a region of said target sequence comprising a second allelic variant, wherein said region encompasses a position corresponding to the binding position of the allele-specific nucleotide portion of the first allele-specific primer, and wherein the first allele-specific blocker probe comprises a minor groove binder; and

iv) a detector probe

to form a second and a third reaction mixture in the at least two separate vessels;

e) amplifying said target sequence in said second and third reaction mixtures; and

f) detecting a change in a detectable property of the detector probe in each of the amplified products of said target sequence produced by step (e), thereby detecting the allelic variant of the target gene in the nucleic acid sample.

53. The method of claim **52**, wherein said first reaction mixture is a multiplex reaction and said second and third reaction mixtures are single-plex reactions.

* * * * *