



US009012224B2

(12) **United States Patent**
Bowles et al.

(10) **Patent No.:** **US 9,012,224 B2**
(45) **Date of Patent:** **Apr. 21, 2015**

(54) **CHIMERIC VECTORS**

(75) Inventors: **Dawn E. Bowles**, Durham, NC (US);
Chengwen Li, Chapel Hill, NC (US);
Joseph E. Rabinowitz, Secane, PA
(US); **Josh Grieger**, Chapel Hill, NC
(US); **Mavis Agbandje-McKenna**,
Gainesville, FL (US); **Richard Jude**
Samulski, Chapel Hill, NC (US)

(73) Assignees: **The University of North Carolina at**
Chapel Hill, Chapel Hill, NC (US); **The**
University of Florida Research
Foundation, Gainesville, FL (US)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 1145 days.

(21) Appl. No.: **12/983,547**

(22) Filed: **Jan. 3, 2011**

(65) **Prior Publication Data**

US 2011/0104119 A1 May 5, 2011

Related U.S. Application Data

(63) Continuation of application No. 11/793,430, filed as
application No. PCT/US2005/045552 on Dec. 15,
2005, now Pat. No. 7,892,809.

(60) Provisional application No. 60/636,126, filed on Dec.
15, 2004.

(51) **Int. Cl.**
C12N 15/861 (2006.01)
C12N 15/86 (2006.01)

(52) **U.S. Cl.**
CPC **C12N 15/86** (2013.01); **C12N 2750/14122**
(2013.01); **C12N 2750/14143** (2013.01); **C12N**
2750/14144 (2013.01)

(58) **Field of Classification Search**
None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

5,139,941 A 8/1992 Muzyczka et al.
5,436,146 A 7/1995 Shenk et al.
5,478,745 A 12/1995 Samulski et al.
5,589,377 A 12/1996 Lebkowski et al.
5,622,856 A 4/1997 Natsoulis
5,658,785 A 8/1997 Johnson
5,681,731 A 10/1997 Lebkowski et al.
5,753,500 A 5/1998 Shenk et al.
5,756,283 A 5/1998 Wilson et al.
5,773,289 A 6/1998 Samulski et al.
5,780,280 A 7/1998 Lebkowski et al.
5,780,447 A 7/1998 Nienhuis
5,786,211 A 7/1998 Johnson
5,834,441 A 11/1998 Philip et al.
5,843,742 A 12/1998 Natsoulis et al.
5,846,528 A 12/1998 Podsakoff et al.
5,846,546 A 12/1998 Hurwitz et al.
5,856,152 A 1/1999 Wilson et al.

5,858,351 A 1/1999 Podsakoff et al.
5,858,775 A 1/1999 Johnson
5,861,171 A 1/1999 Philip et al.
5,861,314 A 1/1999 Philip et al.
5,863,541 A 1/1999 Samulski et al.
5,866,552 A 2/1999 Wilson et al.
5,866,696 A 2/1999 Carter et al.
5,869,305 A 2/1999 Samulski et al.
5,871,982 A 2/1999 Wilson et al.
5,872,005 A 2/1999 Wang et al.
5,874,304 A 2/1999 Zolotukhin et al.
5,874,556 A 2/1999 Lupton et al.
5,882,652 A 3/1999 Valdes et al.
5,905,040 A 5/1999 Mazzara et al.
5,916,563 A 6/1999 Young et al.
5,922,315 A 7/1999 Roy
5,945,335 A 8/1999 Colosi
5,952,221 A 9/1999 Kurtzman et al.
5,962,274 A 10/1999 Parks
5,962,313 A 10/1999 Podsakoff et al.
6,001,371 A 12/1999 Young et al.
6,156,303 A 12/2000 Russell et al.
6,468,524 B1 10/2002 Chiorini et al.
6,468,771 B1 10/2002 Einerhand et al.
6,491,907 B1 12/2002 Rabinowitz
6,759,237 B1 7/2004 Wilson et al.
6,984,517 B1 1/2006 Chiorini et al.

(Continued)

FOREIGN PATENT DOCUMENTS

AU 199942205 6/2003
WO WO 95/28493 10/1995

(Continued)

OTHER PUBLICATIONS

Weinberg et al., *Neuropharmacology* vol. 69 (2013) pp. 82-88.*
Rodino-Klapac et al., *Curr Neurol Neurosci Rep* vol. 13 (2013) p.
332.*
Ortolano et al., *Recent Patents on Endocrine, Metabolic & Immune*
Drug Discovery vol. 6 (2012) pp. 47-66.*
Agbandje et al., "Structure Determination of Feline Panleukopenia
Virus Empty Particles" *Proteins: Structure, Function, and Genetics*
16:155-171 (1993).
Alexander et al., "Transfer of Contaminants in Adeno-Associated
Virus Vector Stocks Can Mimic Transduction and Lead to Artifactual
Results" *Human Gene Therapy* 8:1911-1920 (Nov. 1, 1997).
Anderson, "Human Gene Therapy," *Nature* 392: 25-30 (1998).

(Continued)

Primary Examiner — Jim Ketter

(74) *Attorney, Agent, or Firm* — Myers Bigel Sibley &
Sajovec, P.C.

(57) **ABSTRACT**

The present invention is based, in part, on the discovery that
parvovirus (including AAV) capsids can be engineered to
incorporate small, selective regions from other parvoviruses
that confer desirable properties. The inventors have discov-
ered that in some cases as little as a single amino acid inser-
tion or substitution from a first parvovirus (e.g., an AAV) into
the capsid structure of another parvovirus (e.g., an AAV) to
create a chimeric parvovirus is sufficient to confer one or
more of the desirable properties of the first parvovirus to the
resulting chimeric parvovirus and/or to confer a property that
is not exhibited by the first parvovirus or is present to a lesser
extent.

15 Claims, 14 Drawing Sheets

(56)

References Cited

U.S. PATENT DOCUMENTS

7,172,893	B2	2/2007	Rabinowitz et al.
7,252,997	B1	8/2007	Hallek et al.
7,259,151	B2	8/2007	Arbetman et al.
2003/0215422	A1	11/2003	Chiorini et al.
2004/0057931	A1	3/2004	Wilson et al.
2004/0057932	A1	3/2004	Wilson et al.
2004/0057933	A1	3/2004	Wilson et al.
2004/0086490	A1	5/2004	Chiorini et al.
2005/0255089	A1	11/2005	Chiorini et al.
2006/0188483	A1	8/2006	Rabinowitz et al.
2006/0188484	A1	8/2006	Rabinowitz et al.

FOREIGN PATENT DOCUMENTS

WO	WO 96/00587	1/1996
WO	WO96/36364	11/1996
WO	WO 97/05266	2/1997
WO	WO 97/38723	10/1997
WO	WO 98/09524	3/1998
WO	WO 98/11244	3/1998
WO	WO 98/32842	7/1998
WO	WO 98/41240	9/1998
WO	WO 99/61601	12/1999
WO	WO 99/67393	12/1999
WO	WO 00/28004	5/2000
WO	WO 00/28061	5/2000
WO	WO 01/05991	1/2001
WO	WO 01/05990	2/2001
WO	WO 01/68888 A2	9/2001
WO	WO 01/68888 A3	9/2001
WO	WO 01/83692	11/2001

OTHER PUBLICATIONS

Antonietti et al.; "Characterization of the Cell Type-Specific Determinant in the Genome of Minute Virus of Mice," *Journal of Virology* 62:2 552-557 (Feb. 1988).

Ball-Goodrich et al.; Two Amino Acid Substitutions within the Capsid Are Coordinately Required for Acquisition of Fibrotropism by the Lymphotropic Strain of Minute Virus of Mice, *Journal of Virology* 66:6 3415-3423 (Jun. 1992).

Bartlett et al.; "Genetics and Biology of Adeno-Associated Virus," *Viral Vectors* 55-73 (1995).

Bloom et al.; "Characterization of Chimeric Full-Length Molecular Clones of Aleutian Mink Disease Parvovirus (ADV): Identification of a Determinant Governing Replication of ADV in Cell Culture," *Journal of Virology* 59:76-5988 (Oct. 1993).

Bowles et al.; "Marker Rescue of Adeno-Associated Virus (AAV) Capsid Mutants: a Novel Approach for Chimeric AAV Production," *Journal of Virology*, vol. 77:1, pp. 423-432 (Jan. 2003). XP002975999.

Brown et al.; Chimeric Parvovirus B19 Capsids for the Presentation of Foreign Epitopes, *Virology* 198 477-488 (1994).

Burger et al.; "Recombinant AAV Viral Vectors Pseudotyped with Viral Capsids from Serotypes 1,2, and 5 Display Differential Efficiency and Cell Tropism after Delivery to Different Regions of the Central Nervous System", *Molecular Therapy*, vol. 10:2, pp. 302-317 (Aug. 2004). XP004552211.

Chang et al.; Multiple Amino Acids in the Capsid Structure of Canine Parvovirus Coordinately Determine the Canine Host Range and Specific Antigenic and Hemagglutination Properties, *Journal of Virology* 66:12 6858-6867 (Dec. 1992).

Chapman et al.; *Structure, Sequence, and Function Correlations Among Parvoviruses*, *Virology* 194:491-508 (1993).

Chiorini et al.; Adeno-Associated Virus (AAV) Type 5 Rep Protein Cleaves a Unique Terminal Resolution Site Compared with Other AAV Serotypes, *Journal of Virology* 73:5 4293-4298 (May 1999).

Chiorini et al.; Cloning and Characterization of Adeno-Associated virus Type 5, *Journal of Virology* 73:2 1309-1319 (Feb. 1999).

Chiorini et al.; Cloning of Adeno-Associated Virus Type 4 (AAV4) and Generation of Recombinant AAV4 Particles, *Journal of Virology* 71:9 6823-6833 (Sep. 1997).

Conway et al.; "Recombinant Adeno-Associated Virus Type 2 Replication and Packaging Is Entirely Supported by a Herpes Simplex Virus Type 1 Amplicon Expressing Rep and Cap," *Journal of Virology* 71:11 8780-8789 (Nov. 1997).

Dang et al. Gene therapy and translational cancer research. *Clin. Can. Res.* 5:471-474, 1999.

Douar et al. "Deleterious effect of peptide insertions in a permissive site of the AAV2 capsid" *Virology* 309:203-208 (2003).

Du et al.; "Differential Myocardial Gene Delivery by Recombinant Serotype-Specific Adeno-associated Viral Vectors", *Molecular Therapy*, vol. 10:3, pp. 604-608 (Sep. 2004). XP004660600.

Extended European Search Report and Search Opinion (8 pages) corresponding to European Application No. 05826631.3; Dated: Mar. 30, 2009.

Faust et al. "Universal purification of AAV serotypes 1-5 modified to contain a heparin binding epitope" *Molecular Therapy* 9:S36-S36 (2004).

Fu et al.; Viral sequences enable efficient and tissue-specific expression of transgenes in *Xenopus*, *Nature Biotechnology* 16 253-257 (Mar. 1998).

Gao et al.; "Clades of Adeno-Associated Viruses Are Widely Disseminated in Human Tissues," *Journal of Virology* 78:12 6381-6388 (Jun. 2004).

Gao et al.; "Novel adeno-associated viruses from rhesus monkeys as vectors for human gene therapy," *PNAS* 99:18 11854-11859 (Sep. 3, 2002).

Gao et al.; *High-Titer Adeno-Associated Viral Vectors from a Rep/Cap Cell Line and Hybrid Shuttle Virus*, *Human Gene Therapy* 9:2353-2362 (Nov. 1, 1998).

Gardiner et al.; "Mapping of the Fibrotropic and Lymphotropic Host Range Determinants of the Parvovirus Minute Virus of Mice," *Journal of Virology* 62:8 2605-2613 (Aug. 1988).

Girod et al.; Genetic capsid modifications allow efficient re-targeting of adeno-associated virus type 2, *Nature Medicine* 5:9 1052-1056 (Sep. 1999).

Goldman et al.; Targeted Gene Delivery to Kaposi's Sarcoma Cells via the Fibroblast Growth Factor Receptor, *Cancer Research* 57 1447-1451 (Apr. 15, 1997).

Hauck et al.; "Characterization of Tissue Tropism Determinants of Adeno-Associated Virus Type 1," *Journal of Virology* 77:4 2768-2774 (Feb. 2003).

Hermonat et al.; Genetics of Adeno-Associated Virus: Isolation and Preliminary Characterization of Adeno-Associated Virus Type 2 Mutants, *Journal of Virology* 51:2 329-339 (Aug. 1984).

Hodgson, C.P. Advances in vector systems for gene therapy. *Exp. Opin. Ther. Patents* 5:459-468, 1995.

Horiuchi et al.; *Mapping of Determinants of the Host Range for Canine Cells in the Genome of Canine Parvovirus Using Canine Parvovirus/Mink Enteritis Virus Chimeric Viruses*, *Journal of General Virology* 75:1319-1328 (1994).

Li et al.; Role for Highly Regulated rep Gene Expression in Adeno-Associated Virus Vector Production, *Journal of Virology* 71:7 5236-5243 (Jul. 1997).

Lieber et al.; "AAV display-homing in on the target," *Nature Biotechnology* 21:9 1011-1013 (Sep. 2003).

Llamas-Saiz et al.; "Structure Determination of Minute Virus of Mice" *Acta Crysta* D53 93-102 (1997).

Marshall, E. Gene therapy's growing pains. *Science* 269:1050-1055, 1995.

Maxwell et al.; Targeting a Feline Parvovirus to Human Tumor Cells (abstract), Cold Spring Harbor Laboratory, Vector Targeting Strategies for Therapeutic Gene Delivery meeting (Mar. 11-14, 1999) p. 87.

McCarty et al.; "Integration of Adeno-Associated Virus (AAV) and Recombinant AAV Vectors," *Annu. Rev. Genet.* 38:819-45 (2004).

Miyamura et al.; Parvovirus particles as platforms for protein presentation, *Proc. Natl. Acad. Sci. USA* 91 8507-8511 (Aug. 1994).

Moskalenko et al.; Epitope Mapping of Human Anti-Adeno-Associated Virus Type 2 Neutralizing Antibodies: Implications for Gene Therapy and Virus Structure, *Journal of Virology* 74:4 1761-1766 (Feb. 2000).

Muller et al.; "Random peptide libraries displayed on adeno-associated virus to select for targeted gene therapy vectors" *Nature Biotechnology* 21:9 1040-1046 (Sep. 2003).

(56)

References Cited

OTHER PUBLICATIONS

- Muralidhar et al.; Site-Directed Mutagenesis of Adeno-Associated Virus Type 2 Structural Protein Initiation Codons: Effects on Regulation of Synthesis and Biological Activity, *Journal of Virology* 68:1 170-176 (Jan. 1994).
- Muramatsu et al.; Nucleotide Sequencing and Generation of an Infectious Clone of Adeno-Associated Virus 3, *Virology* 221 208-217 (1996).
- Parrish et al.; "Canine Host Range and a Specific Epitope Map along with Variant Sequences in the Capsid Protein Gene of Canine Parvovirus and Related Feline, Mink, and Raccoon Parvoviruses," *Virology* 166:293-307 (1988).
- Parrish et al.; "Rapid Antigenic-Type Replacement and DNA Sequence Evolution of Canine Parvovirus," *Journal of Virology* 65:12 6544-6552 (Dec. 1991).
- Ponnazhagan et al.; Recombinant Human Parvovirus B19 Vectors: Erythroid Cell-Specific Delivery and Expression of Transduced Genes, *Journal of Virology* 72:6 5224-5230 (Jun. 1998).
- Rabinowitz et al.; "Cross-Dressing the Virion: the Transcapsidation of Adeno-Associated Virus Serotypes Functionally Defines Subgroups", *Journal of Virology*, vol. 78:9, pp. 4421-4432 (May 2004). XP002519618.
- Rabinowitz et al.; "Insertional Mutagenesis of AAV2 Capsid and the Production of Recombinant Virus," *Virology* 265: 274-285 (1999).
- Rabinowitz et al.; *Adeno-Associated Virus Expression Systems for Gene Transfer*, *Current Opinion in Biotechnology* 9:5 470-475 (Oct. 1998).
- Rabinowitz et al.; Targeted Adeno-Associated Virus (abstract), Cold Spring Harbor Laboratory, Vector Targeting Strategies for Therapeutic Gene Delivery meeting (Mar. 11-14, 1999) p. 82.
- Romano et al. Latest developments in gene transfer technology: Achievements, perspectives, and controversies over therapeutic applications. *Stem cells* 18:19-39, 2000.
- Ruffing et al.; Mutations in the carboxy terminus of adeno-associated virus 2 capsid proteins affect viral infectivity: lack of an RGD integrin-binding motif, *Journal of General Virology* 75 3385-3392 (1994).
- Rutledge et al.; Infectious Clones and Vectors Derived from Adeno-Associated Virus (AAV) Serotypes Other Than AAV Type 2, *Journal of Virology* 72:1 309-319 (Jan. 1998).
- Sedlik et al.; Recombinant parvovirus-like particles as an antigen carrier: A novel nonreplicative exogenous antigen to elicit protective antiviral cytotoxic T cells, *Proc. Natl. Acad. Sci, USA* 94 7503-7508 (Jul. 1997).
- Simpson et al.; "The Structure of an insect parvovirus (*Galleria mellonella* densovirus) at 3.7 Å resolution" *Structure* 6:11 1355-1367 (1998).
- Smuda et al.; Adeno-Associated Viruses Having Nonsense Mutations in the Capsid Genes: Growth in Mammalian Cells Containing an Inducible Amber Suppressor, *Virology* 184 310-318 (1991).
- Spitzer et al.; "Tropic determinant for canine parvovirus and feline panleukopenia virus functions through the capsid protein VP2," *Journal of General Virology* 78: 925-928 (1997).
- Spitzer et al.; Species specificity for transduction of cultured cells by a recombinant LuIII rodent parvovirus genome encapsidated by canine parvovirus or feline panleukopenia virus, *Journal of General Virology* 77 1787-1792 (1996).
- Srivastava et al.; Construction of a recombinant human parvovirus B19: Adeno-associated virus 2 (AAV) DNA inverted terminal repeats are functional in an AAV-B19 hybrid virus, *Proc. Natl. Acad. Sci. USA* 86 8078-8082 (Oct. 1989).
- Tsao et al.; "The Three-Dimensional Structure of Canine Parvovirus and Its Functional Implications," *Science* 251:1456-1464 (Mar. 22, 1991).
- Urabe et al.; "Insect Cells as a Factory to Produce Adeno-Associated Virus Type 2 Vectors," *Human Gene Therapy* 13:1935-1943 (Nov. 1, 2002).
- Verma et al. Gene therapy-promises, problems and prospects. *Nature* 389:239-242, 1997.
- Wang et al.; "Recombinant AAV serotype 1 transduction efficiency and tropism in the murine brain," *Gene Therapy* 10, 1528-1534 (2003).
- Wang et al.; "Rescue and Replication of Adeno-Associated Virus Type 2 as well as Vector DNA Sequences from Recombinant Plasmids Containing Deletions in the Viral Inverted Terminal Repeats: Selective Encapsidation of Viral Genomes in Progeny Virions" *J. Virol.* 70(3):1668-1667 (1996).
- Warrington et al.; "Adeno-Associated Viruses Type 2 VP2 Capsid Protein Is Nonessential and Can Tolerate Large Peptide Insertions at Its N Terminus" *Journal of Virology* 78:12 6595-6609 (Jun. 2004).
- Wu et al.; "The Canine Parvovirus Empty Capsid Structure" *J. Mol. Biol.* 233: 231-244 (1993).
- Wu et al.; "Mutational Analysis of the Adeno-Associated Virus Type 2 (AAV2) Capsid Gene and Construction of AAV2 Vectors with Altered Tropism", *Journal of Virology*, vol. 74:18, pp. 8635-8647 (Sep. 2000). XP001028219.
- Xiao et al.; "Gene Therapy Vectors Based on Adeno-Associated Virus Type 1," *Journal of Virology* 73(5): 3994-4003 (May 1999).
- Xiao et al.; Efficient Long-Term Gene Transfer into Muscle Tissue of Immunocompetent Mice by Adeno-Associated Virus Vector, *Journal of Virology* 70:11 8098-8108 (Nov. 1996).
- Xiao et al.; Production of high-titer recombinant adeno-associated virus vectors in the absence of helper adenovirus, *J. Virol.* 72:3 2224 (15 pp.) (Mar. 1998).
- Yang et al.; "Virus-Mediated Transduction of Murine Retina with Adeno-Associated Virus: Effects of Viral Capsid and Genome Size," *Journal of Virology* 76:15 7651-7660 (Aug. 2002).
- Yang et al.; Development of Novel Cell Surface CD34-Targeted Recombinant Adenoassociated Virus Vectors for Gene Therapy, *Human Gene Therapy* 9 1929-1937 (Sep. 1, 1998).
- Japanese Patent Application No. 2007-546926, filed Apr. 20, 2009; Office Action mailed Aug. 16, 2011.

* cited by examiner

		1	10	20	30	40	50	60	70																																																														
AAV1	(1)	MAADG	YLP	PDW	LED	NL	SE	GIRE	W	D	L	K	P	G	A	P	K	P	K	A	N	Q	K	Q	D	D	G	R	G	L	V	L	P	G	Y	K	Y	L	P	G	F	N	G	L	D	K	G	E	P	V	N	A	A	D																	
AAV7	(1)	MAADG	YLP	PDW	LED	NL	SE	GIRE	W	D	L	K	P	G	A	P	K	P	K	A	N	Q	K	Q	D	D	G	R	G	L	V	L	P	G	Y	K	Y	L	P	G	F	N	G	L	D	K	G	E	P	V	N	A	A	D																	
AAV2	(1)	MAADG	YLP	PDW	LED	NL	SE	GIRE	W	D	L	K	P	G	A	P	P	P	P	K	P	A	E	R	H	K	D	S	R	G	L	V	L	P	G	Y	K	Y	L	P	G	F	N	G	L	D	K	G	E	P	V	N	E	A	D																
AAV8	(1)	MAADG	YLP	PDW	LED	NL	SE	GIRE	W	D	L	K	P	G	A	P	K	P	K	A	N	Q	K	Q	D	D	G	R	G	L	V	L	P	G	Y	K	Y	L	P	G	F	N	G	L	D	K	G	E	P	V	N	A	A	D																	
AAV1	(71)	AALEH	D	K	A	Y	D	Q	Q	L	K	A	G	D	N	P	Y	L	R	N	H	A	D	A	E	F	Q	E	R	L	Q	E	D	T	S	F	G	N	L	G	R	A	V	F	Q	A	K	R	V	L	E	P	L	G	L	V	E	E	G	A	K	T	A	P							
AAV7	(71)	AALEH	D	K	A	Y	D	Q	Q	L	K	A	G	D	N	P	Y	L	R	N	H	A	D	A	E	F	Q	E	R	L	Q	E	D	T	S	F	G	N	L	G	R	A	V	F	Q	A	K	R	V	L	E	P	L	G	L	V	E	E	G	A	K	T	A	P							
AAV2	(71)	AALEH	D	K	A	Y	D	Q	Q	L	K	A	G	D	N	P	Y	L	R	N	H	A	D	A	E	F	Q	E	R	L	Q	E	D	T	S	F	G	N	L	G	R	A	V	F	Q	A	K	R	V	L	E	P	L	G	L	V	E	E	P	V	K	T	A	P							
AAV8	(71)	AALEH	D	K	A	Y	D	Q	Q	L	K	A	G	D	N	P	Y	L	R	N	H	A	D	A	E	F	Q	E	R	L	Q	E	D	T	S	F	G	N	L	G	R	A	V	F	Q	A	K	R	V	L	E	P	L	G	L	V	E	E	G	A	K	T	A	P							
AAV1	(141)	G	K	R	P	V	E	Q	S	P	O	E	-	P	D	S	S	S	G	I	G	K	T	G	Q	Q	P	A	K	K	R	L	N	F	G	Q	T	G	D	S	E	S	V	P	D	P	Q	P	L	G	E	P	P	A	P	A	V	G	E	T	T	M	A	S	G	G	G	A			
AAV7	(141)	A	K	K	R	P	V	E	S	P	O	R	S	-	P	D	S	S	G	I	G	K	K	G	Q	Q	P	A	K	K	R	L	N	F	G	Q	T	G	D	S	E	S	V	P	D	P	Q	P	L	G	E	P	P	A	P	S	S	V	G	S	G	T	V	A	A	G	G	A			
AAV2	(141)	G	K	K	R	P	V	E	H	S	P	V	E	-	P	D	S	S	G	I	G	K	A	G	Q	Q	P	A	K	K	R	L	N	F	G	Q	T	G	D	A	S	V	P	D	P	Q	P	L	G	Q	P	P	A	P	S	S	V	G	I	G	T	N	I	M	A	T	G	S	G	A	
AAV8	(141)	G	K	K	R	P	V	E	S	P	O	R	S	-	P	D	S	S	G	I	G	K	K	G	Q	Q	P	A	K	K	R	L	N	F	G	Q	T	G	D	S	E	S	V	P	D	P	Q	P	L	G	E	P	P	A	P	S	S	V	G	I	G	T	N	I	M	A	A	G	G	A	
AAV1	(210)	P	M	A	D	N	E	G	A	D	G	V	G	N	A	S	G	N	W	H	C	D	S	T	W	L	G	D	R	V	I	T	T	S	T	R	T	W	A	L	P	T	Y	N	N	H	L	Y	K	O	I	S	S	A	S	T	G	-	A	S	N	D	N	H	Y	F	G	Y	S	T	
AAV7	(211)	P	M	A	D	N	E	G	A	D	G	V	G	N	A	S	G	N	W	H	C	D	S	T	W	L	G	D	R	V	I	T	T	S	T	R	T	W	A	L	P	T	Y	N	N	H	L	Y	K	O	I	S	S	E	I	A	G	-	S	T	N	D	N	T	Y	F	G	Y	S	T	
AAV2	(210)	P	M	A	D	N	E	G	A	D	G	V	G	N	S	S	G	N	W	H	C	D	S	T	W	M	G	D	R	V	I	T	T	S	T	R	T	W	A	L	P	T	Y	N	N	H	L	Y	K	O	I	S	S	G	A	-	-	S	N	D	N	H	Y	F	G	Y	S	T			
AAV8	(211)	P	M	A	D	N	E	G	A	D	G	V	G	S	S	G	N	W	H	C	D	S	T	W	L	G	D	R	V	I	T	T	S	T	R	T	W	A	L	P	T	Y	N	N	H	L	Y	K	O	I	S	N	G	T	S	G	G	A	T	N	D	N	T	Y	F	G	Y	S	T		
AAV1	(279)	P	W	G	Y	F	D	F	N	R	F	H	C	H	F	S	P	R	D	W	Q	R	L	I	N	N	N	W	G	F	R	P	K	R	L	N	F	K	L	F	N	I	Q	V	K	E	V	T	I	N	D	G	V	T	T	I	A	N	N	L	T	S	T	V	Q	V	E	S	D	S	E
AAV7	(280)	P	W	G	Y	F	D	F	N	R	F	H	C	H	F	S	P	R	D	W	Q	R	L	I	N	N	N	W	G	F	R	P	K	R	L	F	K	L	F	N	I	Q	V	K	E	V	T	I	N	D	G	V	T	T	I	A	N	N	L	T	S	T	I	Q	V	E	S	D	S	E	
AAV2	(278)	P	W	G	Y	F	D	F	N	R	F	H	C	H	F	S	P	R	D	W	Q	R	L	I	N	N	N	W	G	F	R	P	K	R	L	N	F	K	L	F	N	I	Q	V	K	E	V	T	I	N	D	G	V	T	T	I	A	N	N	L	T	S	T	V	Q	V	E	S	D	S	E
AAV8	(281)	P	W	G	Y	F	D	F	N	R	F	H	C	H	F	S	P	R	D	W	Q	R	L	I	N	N	N	W	G	F	R	P	K	R	L	S	F	K	L	F	N	I	Q	V	K	E	V	T	I	N	E	C	T	K	T	I	A	N	N	L	T	S	T	I	Q	V	E	S	D	S	E
AAV1	(349)	Y	Q	L	P	Y	V	L	G	S	A	H	Q	G	C	L	P	P	F	P	A	D	V	F	M	I	P	Q	Y	G	Y	L	T	L	N	G	S	Q	A	V	G	R	S	S	F	C	Y	C	L	E	Y	F	P	S	Q	M	L	R	T	G	N	N	F	T	F	S	Y	T	F	F	E
AAV7	(350)	Y	Q	L	P	Y	V	L	G	S	A	H	Q	G	C	L	P	P	F	P	A	D	V	F	M	I	P	Q	Y	G	Y	L	T	L	N	G	S	Q	S	V	G	R	S	S	F	C	Y	C	L	E	Y	F	P	S	Q	M	L	R	T	G	N	N	F	T	F	S	Y	T	F	F	E
AAV2	(348)	Y	Q	L	P	Y	V	L	G	S	A	H	Q	G	C	L	P	P	F	P	A	D	V	F	M	I	P	Q	Y	G	Y	L	T	L	N	G	S	Q	A	V	G	R	S	S	F	C	Y	C	L	E	Y	F	P	S	Q	M	L	R	T	G	N	N	F	T	F	S	Y	T	F	F	E
AAV8	(351)	Y	Q	L	P	Y	V	L	G	S	A	H	Q	G	C	L	P	P	F	P	A	D	V	F	M	I	P	Q	Y	G	Y	L	T	L	N	G	S	Q	A	V	G	R	S	S	F	C	Y	C	L	E	Y	F	P	S	Q	M	L	R	T	G	N	N	F	T	F	S	Y	T	F	F	E

FIG. 1A

AAV1	(419)	421	430	440	450	460	470	480	490										
AAV7	(420)	VPFHSS	YAHSQS	LDRLMN	PLIDQY	LYYINR	TONQ	SGSKQ	NKDL	LLFSR	GSPAGMS	V	OPKNW	LP	GP	CYRQ	Q		
AAV2	(418)	VPFHSS	YAHSQS	LDRLMN	PLIDQY	LYYIAR	TQSN	PGGT	AGNREL	QFYQ	GGPST	MAED	QAKNW	LP	GP	CFRQ	Q		
AAV8	(421)	VPFHSS	YAHSQS	LDRLMN	PLIDQY	LYYISRT	NTTP	SGTT	QSR	LQFSQ	AGASD	IRDD	QSRNW	LP	GP	CYRQ	Q		
		VPFHSS	YAHSQS	LDRLMN	PLIDQY	LYYISRT	QTT	GGT	ANTQ	LGF	SQGGP	NTMAN	QAKNW	LP	GP	CYRQ	Q		
AAV1	(488)	491	500	510	520	530	540	550	560										
AAV7	(490)	RVSKT	KTQNNNS	NFTWTG	ASKYNL	NGRESI	INPG	TAMASH	KDDE	DKFFP	MSGVM	IFGK	ES	AGAS	N	TALDN			
AAV2	(487)	RVSKT	LDQNNNS	NFAWTG	ATKYHL	NGRNS	SLVNP	GVAMATH	KDDE	DRFFP	SSGV	LIFG	KT	GATN	K	TTLN			
AAV8	(490)	RVSKT	SAQNNNS	SEYSWT	GATKYHL	NGRDS	SLVNP	GPAMASH	KDDE	KFFPQ	SGV	LIFG	KQ	SEK	T	NVDIEK			
		RVSTT	TGQNNNS	NFAWTAG	TKYHL	NGRNS	LANPG	GLAMATH	KDDE	ERFFP	SGIL	LIFG	KQ	AA	R	NADYS			
AAV1	(558)	561	570	580	590	600	610	620	630										
AAV7	(559)	VMITDEE	EIRKATNP	VATERF	GTVA	VNFQSS	STD	PATG	DVHAM	GALP	GMVWQ	DRD	VYLQ	GP	IAWAKI	PHTDG			
AAV2	(557)	VLMTNEE	EIRP	TNPV	ATEEY	GIVSSN	LQAA	NTAA	QTVVN	Q	GALP	GMVWQ	NRD	VYLQ	GP	IAWAKI	PHTDG		
AAV8	(560)	VMITDEE	EIRKATNP	VATERF	GTVA	VNFQSS	STD	PATG	DVHAM	GALP	GMVWQ	DRD	VYLQ	GP	IAWAKI	PHTDG			
		VMLTSEE	EIRKATNP	VATERF	GTVA	VNFQSS	STD	PATG	DVHAM	GALP	GMVWQ	DRD	VYLQ	GP	IAWAKI	PHTDG			
AAV1	(628)	631	640	650	660	670	680	690	700										
AAV7	(629)	HFHPS	PLMGGF	GLKN	PPPPQ	ILIKNT	TPVPAN	PPAE	TSATK	FASFI	TOYST	GQVS	VEI	E	WELQ	KENSKR	WNP		
AAV2	(627)	NFHPS	PLMGGF	GLKH	PPPPQ	ILIKNT	TPVPAN	PPAE	TSATK	FASFI	TOYST	GQVS	VEI	E	WELQ	KENSKR	WNP		
AAV8	(630)	NFHPS	PLMGGF	GLKH	PPPPQ	ILIKNT	TPVPAN	PPAE	TSATK	FASFI	TOYST	GQVS	VEI	E	WELQ	KENSKR	WNP		
		NFHPS	PLMGGF	GLKH	PPPPQ	ILIKNT	TPVPAD	PPAE	TSATK	FASFI	TOYST	GQVS	VEI	E	WELQ	KENSKR	WNP		
AAV1	(698)	701	710	720	730	740													
AAV7	(699)	EVQY	TSN	YAKS	ANVDF	TVD	NNGL	YTEP	RP	IG	TR	YL	TR	PL	-				
AAV2	(697)	EIQY	TSN	TEK	Q	TG	VD	FA	V	SSQ	GV	YSE	PR	PI	G	TR	YL	TR	NL
AAV8	(700)	EIQY	TSN	IN	K	SV	NVDF	TVD	NNGL	YTEP	RP	IG	TR	YL	TR	PL	-		

FIG. 1B

AAV1	1	.10	.20	.30	.40	.50	.60	.70
AAV2		MAADGYLPDWLEDNLSEGI	REWWDLKPGAPKPKANQO	KODDGRGLVLP	GGKYLGP	FNGLDKGE	PVNAADA	
		MAADGYLPDWLEDTLSEGI	ROWWKLKPGPPPKPAER	HRDDSRGLVLP	GGKYLGP	FNGLDKGE	PVNEADA	
AAV1	71	.80	.90	.100	.110	.120	.130	.140
AAV2		AALEHDKAYDQOLKAGDNP	YLRYNHADA	AEFOERLQEDTS	FGGNLGRAVFO	AKKRVL	EPGLGLVEEGAKTAP	
		AALEHDKAYDQOLKAGDNP	YLRYNHADA	AEFOERLQEDTS	FGGNLGRAVFO	AKKRVL	EPGLGLVEEPVKTAP	
AAV1	141	.150	.160	.170	.180	.190	.200	.210
AAV2		GKKRPVEQSPQEPDSSSGI	GKTGOQPAKRRLN	FGQTGDSESV	PDQPLGEP	PATPA	AVGPTTMA	SGGGAP
		GKKRPVEHSPVEPDSSSGT	GKAGQOPARKRLN	FGQTGDADSV	PDQPLGQPPA	PSGLGNTMA	TGSGAP	
AAV1	211	.220	.230	.240	.250	.260	.270	.280
AAV2		MADNNEGADGVGNASGN	WHCDSTWLGDRVIT	TSTRTWALPT	YNNHLYKQISS	ASGTASND	NHYFGYSTPW	
		MADNNEGADGVGNSSGN	WHCDSTWLGDRVIT	TSTRTWALPT	YNNHLYKQISS	SGA-SND	NHYFGYSTPW	
AAV1	281	.290	.300	.310	.320	.330	.340	.350
AAV2		GYFDENRFCHFS	SPRDWORLINNNW	GERPKRLNFKL	FNIOVKEVT	TNDGVT	TIANNLTST	VOVFS
		GYFDENRFCHFS	SPRDWORLINNNW	GERPKRLNFKL	FNIOVKEVT	QNDGVT	TIANNLTST	VOVFTDSEYQ
AAV1	351	.360	.370	.380	.390	.400	.410	.420
AAV2		LPYVLGSAHQCCLP	PPADVFMTPOYGYL	TLNNGSOAVGR	SSFYCYCLEY	FP	QMLRTGNN	FTFSYTFE
		LPYVLGSAHQCCLP	PPADVFMTPOYGYL	TLNNGSOAVGR	SSFYCYCLEY	FP	QMLRTGNN	FTFSYTFE
AAV1	421	.430	.440	.450	.460	.470	.480	.490
AAV2		FHSSYAHSQSLDR	LMNPLIDQYLYL	NRTONQSGSAA	QNKDLLFSR	GSPAGMSV	QPKNWL	PGPCYRQ
		FHSSYAHSQSLDR	LMNPLIDQYLYL	NRTONQSGSAA	QNKDLLFSR	GSPAGMSV	QPKNWL	PGPCYRQ
AAV1	491	.500	.510	.520	.530	.540	.550	.560
AAV2		SKTKTDNNNSNFT	WTGASKYNLNGR	ESIINPGTAM	ASHKDDDEK	FFPMSG	VMIFGKE	SAGASNTALDNVM
		SKTSADNNNSEY	SWTGATKYHLN	GRDSLVPNPG	PAMASHKDD	DEEKFFPQ	SGVLIFGKQ	SEKTNVDIEKVM
AAV1	561	.570	.580	.590	.600	.610	.620	.630
AAV2		ITDEEEIKATNP	VATERFGTVAVN	FQSSSTDPAT	GDVHAMGAL	PGMVWQ	DRDVLQ	GPIWAKIPH
		ITDEEEIRTN	VPATEQYGSVSTN	LQGRNRQAATA	DVNTQGVLP	GMVWQ	DRDVLQ	GPIWAKIPH
AAV1	631	.640	.650	.660	.670	.680	.690	.700
AAV2		HPSPIMGGFGLK	NPPPOILLIKNT	VPANPPAE	FSATKFA	SFTTQYST	QVSVEI	IEWELQKENS
		HPSPIMGGFGLK	NPPPOILLIKNT	VPANPPAE	FSATKFA	SFTTQYST	QVSVEI	IEWELQKENS
AAV1	701	.710	.720	.730				
AAV2		VOYTSNYAKS	ANVDFTVDN	NGLYTEPR	PIGTRYL	TRPL		
		IQYTSNYKSV	NVDFTVDN	NGVSEPR	PIGTRYL	TRNL		

FIG. 1C




FIG. 2A




FIG. 2B




FIG. 3A




FIG. 3B







FIG. 3C



**FIG. 5****FIG. 6**

(220)	220	230	240	250	260	270	280	290
AAV1 CAP (217)	GADGVGNASGNWHCDSTWLGDRVITTTSTRTWALPTYNHLYKQISSAS--TGASNDNHFGYSTPWGYFDENRF							
AAV2 CAP (217)	GADGVGNSSGNWHCDSTWMGDRVITTTSTRTWALPTYNHLYKQISSQS--GASNDNHFGYSTPWGYFDENRF							
AAV3A CAP (217)	GADGVGNSSGNWHCDSQWLIGDRVITTTSTRTWALPTYNHLYKQISSQS--GASNDNHFGYSTPWGYFDENRF							
AAV3B CAP (217)	GADGVGNSSGNWHCDSQWLIGDRVITTTSTRTWALPTYNHLYKQISSQS--GASNDNHFGYSTPWGYFDENRF							
AAV4 CAP (211)	GADGVGNASGDWHCDSTWSEGHVITTTSTRTWVLPYNNHLYKRLGESL-----QSNYNGFSTPWGYFDENRF							
AAV5 CAP (207)	GADGVGNASGDWHCDSTWMGDRVVTKSTRTWVLPYNNHLYKQISSAS--VDGSNANAYFGYSTPWGYFDENRF							
AAV6 CAP (217)	GADGVGNASGNWHCDSTWLGDRVITTTSTRTWALPTYNHLYKQISSAS--TGASNDNHFGYSTPWGYFDENRF							
AAV7 CAP (218)	GADGVGNASGNWHCDSTWLGDRVITTTSTRTWALPTYNHLYKQISSAS--AGSTNDNTYFGYSTPWGYFDENRF							
AAV8 CAP (218)	GADGVGSSGNWHCDSTWLGDRVITTTSTRTWALPTYNHLYKQISSAS--TSGGATNDNTYFGYSTPWGYFDENRF							
AAV9 CAP (217)	GADGVGSSGNWHCDSQWLIGDRVITTTSTRTWALPTYNHLYKQISSAS--TSGGSSNDNAYFGYSTPWGYFDENRF							

FIG. 7




FIG. 8

1 50
263, 265 (1) MAADGYLPDWLEDLTLSEGIRQWWKLKPGPPPKPAERHKDDSRGLVLPGY
AAV2CAP (1) MAADGYLPDWLEDLTLSEGIRQWWKLKPGPPPKPAERHKDDSRGLVLPGY

51 100
263, 265 (51) KYLGPFNGLDKGEFVNEADAAALEHDKAYDRQLDSGDNPYLKYNHADAEP
AAV2CAP (51) KYLGPFNGLDKGEFVNEADAAALEHDKAYDRQLDSGDNPYLKYNHADAEP

101 150
263, 265 (101) QERLKEDTSFGGNLGRAVFQAKKRVLEPLGLVEEPVKTAPGKKRPVEHSP
AAV2CAP (101) QERLKEDTSFGGNLGRAVFQAKKRVLEPLGLVEEPVKTAPGKKRPVEHSP

151 200
263, 265 (151) VEPDSSSGTGKAGQQPARKRLNFGQTGDADSVDPQPLGQPPAAPSGLGT
AAV2CAP (151) VEPDSSSGTGKAGQQPARKRLNFGQTGDADSVDPQPLGQPPAAPSGLGT

201 250
263, 265 (201) NTMATGSGAPMADNNEGADGVGNSSGNWHCDSTWMGDRVITTSTRTWALP
AAV2CAP (201) NTMATGSGAPMADNNEGADGVGNSSGNWHCDSTWMGDRVITTSTRTWALP

251 * * 300
263, 265 (251) TYNNHLYKQISSASTGASNDNHYFGYSTPWGYFDFNRFHCHFSPRDWQRL
AAV2CAP (251) TYNNHLYKQISSQS-GASNDNHYFGYSTPWGYFDFNRFHCHFSPRDWQRL

301 350
263, 265 (301) INNNWGFPRKRLNFKLFNIQVKEVTQNDGTTTIANNLTSTVQVFTDSEYQ
AAV2CAP (300) INNNWGFPRKRLNFKLFNIQVKEVTQNDGTTTIANNLTSTVQVFTDSEYQ

351 400
263, 265 (351) LPYVLGSAHQGCLPPFPADVFMVPQYGYLTLNNGSQAVGRSSFYCLEYFP
AAV2CAP (350) LPYVLGSAHQGCLPPFPADVFMVPQYGYLTLNNGSQAVGRSSFYCLEYFP

401 450
263, 265 (401) SQMLRTGNNFTFSYTFEDVPPHSSYAHSQSLDRLMNPLIDQYLYLSRTN
AAV2CAP (400) SQMLRTGNNFTFSYTFEDVPPHSSYAHSQSLDRLMNPLIDQYLYLSRTN

451 500
263, 265 (451) TPSGTTTQSRLOFSQAGASDIRDQSRNWLPGPCYRQQRVSKTSADNNNSE
AAV2CAP (450) TPSGTTTQSRLOFSQAGASDIRDQSRNWLPGPCYRQQRVSKTSADNNNSE

501 550
263, 265 (501) YSWTGATKYHLNGRDSLVPNGPAMASHKDDEEKFFPQSGVLI FGKQGSEK
AAV2CAP (500) YSWTGATKYHLNGRDSLVPNGPAMASHKDDEEKFFPQSGVLI FGKQGSEK

551 600
263, 265 (551) TNVDIEKVMITDEEEIRT'NPNVATEQYGSVSTNLQRGNRQAATADVNTQG
AAV2CAP (550) TNVDIEKVMITDEEEIRT'NPNVATEQYGSVSTNLQRGNRQAATADVNTQG

601 650
263, 265 (601) VLPGMVWQDRDVYLQGPWAKIPHTDGHFHPSPLMGGFGLKHPPQILIK
AAV2CAP (600) VLPGMVWQDRDVYLQGPWAKIPHTDGHFHPSPLMGGFGLKHPPQILIK

651 700
263, 265 (651) NTPVPANPSTTFSAAKFASFITQYSTGQVSVEIEWELQKENS KRWNPEIQ
AAV2CAP (650) NTPVPANPSTTFSAAKFASFITQYSTGQVSVEIEWELQKENS KRWNPEIQ

701 736
263, 265 (701) YTSNYNKS VNVDFTVDTNGVYSEPRPIGTRYLTRNL
AAV2CAP (700) YTSNYNKS VNVDFTVDTNGVYSEPRPIGTRYLTRNL

FIG. 9

	1	50
AAV3bCAP	(1) MAADGYLPDWLEDNLSEGIREWALKPGVPQPKANQQHQDNRRGLVLPGY	
263,265	(1) MAADGYLPDWLEDNLSEGIREWALKPGVPQPKANQQHQDNRRGLVLPGY	
	51	100
AAV3bCAP	(51) KYLGPGNGLDKGEPVNEADAAALEHDKAYDQQLKAGDNPYLKYNHADAEF	
263,265	(51) KYLGPGNGLDKGEPVNEADAAALEHDKAYDQQLKAGDNPYLKYNHADAEF	
	101	150
AAV3bCAP	(101) QERLQEDTSFGGNLGRAVFQAKKRILEPLGLVEEAAKTAPGKKRPVDQSP	
263,265	(101) QERLQEDTSFGGNLGRAVFQAKKRILEPLGLVEEAAKTAPGKKRPVDQSP	
	151	200
AAV3bCAP	(151) QEPDSSSGVGKSGKQPARKRLNFGQTGDSESVDPDPQLGEPPAAPTSLSGS	
263,265	(151) QEPDSSSGVGKSGKQPARKRLNFGQTGDSESVDPDPQLGEPPAAPTSLSGS	
	201	250
AAV3bCAP	(201) NTMASGGGAPMADNNEGADGVGNSSGNWHCDSQWLGD RVITTSTRTWALP	
263,265	(201) NTMASGGGAPMADNNEGADGVGNSSGNWHCDSQWLGD RVITTSTRTWALP	
	251	300
AAV3bCAP	(251) TYNNHLYKQISSQS-GASNDNHYFGYSTPWGYFDNRFHCHFSPRDWQRL	
263,265	(251) TYNNHLYKQISSASTGASNDNHYFGYSTPWGYFDNRFHCHFSPRDWQRL	
	301	350
AAV3bCAP	(300) INNNWGFRPKKLSFKLFNIQVKEVTQNDGTTTIANNLTSTVQVFTDSEYQ	
263,265	(301) INNNWGFRPKKLSFKLFNIQVKEVTQNDGTTTIANNLTSTVQVFTDSEYQ	
	351	400
AAV3bCAP	(350) LPYVLGSAHQGCLPPFPADVFMVPQYGYLTLNNGSQAVGRSSFYCLEYFP	
263,265	(351) LPYVLGSAHQGCLPPFPADVFMVPQYGYLTLNNGSQAVGRSSFYCLEYFP	
	401	450
AAV3bCAP	(400) SQMLRTGNNFQFSYTFEDVFPFHSSYAHSQSLDRLMNPLIDQYLYYLNRTQ	
263,265	(401) SQMLRTGNNFQFSYTFEDVFPFHSSYAHSQSLDRLMNPLIDQYLYYLNRTQ	
	451	500
AAV3bCAP	(450) GTTSGTTNQSRLLFSQAGPQSMQSLQARNWLPGPCYRQQRLSKTANDNNNS	
263,265	(451) GTTSGTTNQSRLLFSQAGPQSMQSLQARNWLPGPCYRQQRLSKTANDNNNS	
	501	550
AAV3bCAP	(500) NFPWTAASKYHLNGRDSLVPNGPAMASHKDDEEKFFPMHGNIIFGKEGTT	
263,265	(501) NFPWTAASKYHLNGRDSLVPNGPAMASHKDDEEKFFPMHGNIIFGKEGTT	
	551	600
AAV3bCAP	(550) ASNAELDNVMITDEEEIRTTPVATEQYGTVANNLQSSNTAPTTRTVNDQ	
263,265	(551) ASNAELDNVMITDEEEIRTTPVATEQYGTVANNLQSSNTAPTTRTVNDQ	
	601	650
AAV3bCAP	(600) GALPGMVWQDRDVYLQGPIWAKIPHTDGHFHPSPLMGGFGLKHPPPQIMI	
263,265	(601) GALPGMVWQDRDVYLQGPIWAKIPHTDGHFHPSPLMGGFGLKHPPPQIMI	
	651	700
AAV3bCAP	(650) KNTFPVPANPPTTFSPAKFASFITQYSTGQVSVEIEWELQKENS KRWNPEI	
263,265	(651) KNTFPVPANPPTTFSPAKFASFITQYSTGQVSVEIEWELQKENS KRWNPEI	
	701	737
AAV3bCAP	(700) QYTSNYNKS VNVDFTVD TNGVYSEPRPIGTRYLTRNL	
263,265	(701) QYTSNYNKS VNVDFTVD TNGVYSEPRPIGTRYLTRNL	

FIG. 10

	1		50
2.5CAP	(1)	MAADGYLPDWLEDTLSEGIQWWKLKPGPPPPKPAERHKDDSRGLVLPGY	
AAV2CAP	(1)	MAADGYLPDWLEDTLSEGIQWWKLKPGPPPPKPAERHKDDSRGLVLPGY	
	51		100
2.5CAP	(51)	KYLGPFGNLDKGEPVNEADAAALEHDKAYDRQLDSGDNPYLKYNHADAEF	
AAV2CAP	(51)	KYLGPFGNLDKGEPVNEADAAALEHDKAYDRQLDSGDNPYLKYNHADAEF	
	101		150
2.5CAP	(101)	QERLKEDTSFGGNLGRAVFQAKKRVLEPLGLVEEPVKTAPGKKRPVEHSP	
AAV2CAP	(101)	QERLKEDTSFGGNLGRAVFQAKKRVLEPLGLVEEPVKTAPGKKRPVEHSP	
	151		200
2.5CAP	(151)	VEPDSSSGTGKAGQQPARKRLNFGQTGDADSVDPDQPLGQPPAAPSGGLGT	
AAV2CAP	(151)	VEPDSSSGTGKAGQQPARKRLNFGQTGDADSVDPDQPLGQPPAAPSGGLGT	
	201		250
2.5CAP	(201)	NTMATGSGAPMADNNEGADGVGNSSGNWHCDSTWMGDRVITTSTRTWALP	
AAV2CAP	(201)	NTMATGSGAPMADNNEGADGVGNSSGNWHCDSTWMGDRVITTSTRTWALP	
	251	* *	300
2.5CAP	(251)	TYNNHLYKQISSASTGASNDNHYFGYSTPWGYFDNRFHCHFSPRDWQRL	
AAV2CAP	(251)	TYNNHLYKQISSQS-GASNDNHYFGYSTPWGYFDNRFHCHFSPRDWQRL	
	301		350
2.5CAP	(301)	INNNWGFRPKRLNFKLFNIQVKEVTQNDGTTIANNLSTVQVFTDSEYQ	
AAV2CAP	(300)	INNNWGFRPKRLNFKLFNIQVKEVTQNDGTTIANNLSTVQVFTDSEYQ	
	351		400
2.5CAP	(351)	LPYVLGSAHQGCLPPFPADVFMVPQYGYLTLNNGSQAVGRSSFYCLEYFP	
AAV2CAP	(350)	LPYVLGSAHQGCLPPFPADVFMVPQYGYLTLNNGSQAVGRSSFYCLEYFP	
	401		450
2.5CAP	(401)	SQMLRTGNNFTFSYTFEDVPFHSSYAHSQSLDRLMNPLIDQYLYLSRTN	
AAV2CAP	(400)	SQMLRTGNNFTFSYTFEDVPFHSSYAHSQSLDRLMNPLIDQYLYLSRTN	
	451		500
2.5CAP	(451)	TPSGTTTQSRLOFSQAGASDIRDQSRNWLPGPCYRQQRVSKTSADNNNSE	
AAV2CAP	(450)	TPSGTTTQSRLOFSQAGASDIRDQSRNWLPGPCYRQQRVSKTSADNNNSE	
	501		550
2.5CAP	(501)	YSWTGATKYHLNGRDSLVPNGPAMASHKDDEEKFFPQSGVLIFGKQGSEK	
AAV2CAP	(500)	YSWTGATKYHLNGRDSLVPNGPAMASHKDDEEKFFPQSGVLIFGKQGSEK	
	551		600
2.5CAP	(551)	TNVDIEKVMITDEEEIRTTNPVATEQYGSVSTNLQRGNRQAATADVNTQG	
AAV2CAP	(550)	TNVDIEKVMITDEEEIRTTNPVATEQYGSVSTNLQRGNRQAATADVNTQG	
	601		650
2.5CAP	(601)	VLPGMVWQDRDVYLQGPIWAKIPHTDGHFHPSPLMGGFGLKHPPPQILIK	
AAV2CAP	(600)	VLPGMVWQDRDVYLQGPIWAKIPHTDGHFHPSPLMGGFGLKHPPPQILIK	
	651		700
2.5CAP	(651)	NTPVPANPSTTFSAAKFASFITQYSTGQVSVEIEWELQKENS KRWNPEIQ	
AAV2CAP	(650)	NTPVPANPSTTFSAAKFASFITQYSTGQVSVEIEWELQKENS KRWNPEIQ	
	701	* * *	736
2.5CAP	(701)	YTSNYAKSANVDFTVDNNGVYSEPRPIGTRYLTRNL	
AAV2CAP	(700)	YTSNYNKS NVNVDFTVDNNGVYSEPRPIGTRYLTRNL	

FIG. 11




FIG. 12




FIG. 13




FIG. 14

1

CHIMERIC VECTORS**CROSS-REFERENCE TO RELATED APPLICATIONS**

This application is a continuation application of, and claims priority to U.S. patent application Ser. No. 11/793,430 filed Aug. 22, 2007, now U.S. Pat. No. 7,892,809, which is a 35 U.S.C. §371 national phase application of International Application No. PCT/US2005/045552, filed Dec. 15, 2005, which claims priority to U.S. Provisional Application No. 60/636,126, filed Dec. 15, 2004, the disclosures of each of which are incorporated herein by reference in their entireties.

STATEMENT OF GOVERNMENT SUPPORT

This invention was made with government support under Grant No. HL066973 awarded by the National Institutes of Health. The government has certain rights in the invention.

FIELD OF THE INVENTION

The present invention relates to novel chimeric viral vectors and methods of making and administering the same.

BACKGROUND OF THE INVENTION

It is becoming clear that vectors based upon adeno-associated virus (AAV) are the vectors of choice for certain gene therapy applications such as muscle delivery. The utilization of AAV vectors in such protocols is based on the advantageous properties of AAV. These properties include lack of pathogenicity and pathology, ease of preparation and purification, long term expression in many tissues including the muscle, and lack of a detrimental cell-mediated immune response.

AAV serotype 2 (AAV2) is the best studied of the AAV isolates. Over the past decade, inroads have been made in the evaluation of the tissue tropism of alternative AAV serotypes. These studies have shown that distinct AAV serotypes may be better suited for particular applications. In this regard, serotypes 1, 6 and 7 are the most promising for delivery to skeletal muscle. For example, as compared with AAV2, AAV1 can be administered at lower dosages (i.e., fewer particles) and can express the transgene at earlier time points and at higher levels of expression.

The purification schemes for AAV2 are well defined. Less streamlined are the purification parameters for some of the other AAV serotypes. It would be desirable to engineer a variant of AAV2 that exhibits advantageous properties, such as the enhanced muscle tropism of AAV1, 6 and 7, but still maintains its ease of purification. Increasing the range of available AAV vectors will also address additional concerns related to re-administration and immune responses.

Accordingly, it would be desirable to have available a broader array of AAV vectors.

SUMMARY OF THE INVENTION

The present invention provides chimeric virus vectors that have been designed to exhibit one or more properties of interest (e.g., enhanced tissue tropism). For example, the inventors have identified the key amino acid of AAV1 responsible for enhanced in vivo transduction and selectively engineered this amino acid into the backbone of AAV2 and AAV3b. In particular embodiments, the chimeric viruses of the invention have enhanced transduction capability (e.g.,

2

transduction of skeletal muscle, cardiac muscle, glial cells, astrocytes, liver, retina and/or lung, etc.), enhanced levels of transgene expression and/or earlier onset of transgene expression. The chimeric virus can also have a reduced transduction capability with respect to one or more cells or tissues (e.g., liver), which can be desirable in terms of targeting the vector to the target tissue of interest and reducing dosage of vector to be administered.

With respect to chimeric viruses based on AAV2 these chimeras can be designed to retain one or more of the desirable properties of this serotype (such as ease of purification and known safety), while exhibiting some of the advantageous properties of other AAV such as AAV1, AAV6 and/or AAV7 (including enhanced transduction of skeletal muscle), and/or other properties of interest that are not seen in other AAV or are present to a lesser extent in other AAV. Further, in particular embodiments, the chimeric virus has a different immunological profile than one or both of the parent viruses (i.e., is only weakly or not at all recognized by neutralizing antisera or antibodies against the parent virus), thereby allowing for administration to subjects that have antibodies directed against the parent virus or repeat administration following administration of another serotype.

Accordingly, as one aspect the invention provides a chimeric virus vector comprising:

- (a) a chimeric AAV capsid comprising a selective amino acid insertion following amino acid position 264 in an AAV2 capsid subunit or a corresponding change in a capsid subunit from other AAV; and
- (b) a nucleic acid comprising an AAV terminal repeat sequence and a heterologous nucleic acid sequence; wherein the nucleic acid is packaged within the chimeric AAV capsid.

As another aspect, the invention also provides chimeric virus vector comprising:

- (a) a chimeric AAV capsid comprising:
 - (i) a selective amino acid substitution of an alanine for glutamine at amino acid position 263 in an AAV2 capsid subunit;
 - (ii) a selective amino acid insertion of a threonine following amino acid position 264 in the AAV2 capsid subunit (e.g., immediately following amino acid position 264);
- (b) a nucleic acid comprising an AAV terminal repeat sequence and a heterologous nucleic acid sequence; wherein the nucleic acid is packaged within the chimeric AAV capsid.

The invention further provides a chimeric virus vector comprising:

- (a) a chimeric AAV capsid comprising:
 - (i) a selective amino acid substitution of an alanine for glutamine at amino acid position 263 in an AAV3b capsid subunit;
 - (ii) a selective amino acid insertion of a threonine following amino acid position 264 in the AAV3b capsid subunit (e.g., immediately following amino acid position 264);
- (b) a nucleic acid comprising an AAV terminal repeat sequence and a heterologous nucleic acid sequence; wherein the nucleic acid is packaged within the chimeric AAV capsid.

As still another aspect, the invention provides chimeric virus vector comprising:

- (a) a chimeric AAV capsid comprising a selective amino acid substitution at amino acid position 450 in an AAV2 capsid subunit or a corresponding change in a capsid subunit from other AAV;

3

(b) a nucleic acid comprising an AAV terminal repeat sequence and a heterologous nucleic acid sequence; wherein the nucleic acid is packaged within the chimeric AAV capsid.

As yet another aspect, the invention provides a chimeric virus vector comprising:

(a) a chimeric AAV capsid comprising a selective amino acid substitution at amino acid position 457 in an AAV2 capsid subunit or a corresponding change in a capsid subunit from other AAV;

(b) a nucleic acid comprising an AAV terminal repeat sequence and a heterologous nucleic acid sequence; wherein the nucleic acid is packaged within the chimeric AAV capsid.

Also provided are pharmaceutical formulations comprising the chimeric virus vectors of the invention.

The invention also provides methods of administering a nucleic acid to a cell comprising contacting the cell with a chimeric virus vector or pharmaceutical formulation of the invention.

As yet a further aspect, the invention provides methods of delivering a nucleic acid to a subject comprising administering to the subject a chimeric virus vector or pharmaceutical formulation of the invention.

The invention also provides for the use of the chimeric virus vectors of the invention in the manufacture of a medicament for the treatment of disease.

These and other aspects of the invention are set forth in more detail in the description of the invention below.

BRIEF DESCRIPTION OF THE DRAWINGS

FIGS. 1A-1B. Rationale of the triage approach. Four AAV serotypes (AAV1, AAV2, AAV7, and AAV8) were subjected to a multiple amino acid alignment using the Align program in the Vector NTI software suite. Amino acids similar in AAV1 (SEQ ID NO:1) and AAV7 (SEQ ID NO:2) and distinct from AAV2 (SEQ ID NO:3) and AAV8 (SEQ ID NO:4) or positions that contained amino acids that were different in the 4 serotypes were pinpointed as potential candidates. This led to the 36 potential candidates to engineer into AAV2. Elimination of amino acids that could not be modeled onto the crystal structure and molecular modeling of candidates onto the surface of the capsid led to the testing of the amino acids shown in FIG. 1C.

FIG. 1C. Alignment of AAV1 (SEQ ID NO:1) and AAV2 (SEQ ID NO:3) using the Align program in Vector NTI software package. The amino acids tested in the triage approach are shown with arrows. The circled amino acids correspond to those amino acids previously identified in Hauck et al., (2003) *J. Virology* 77:2768-2774).

FIG. 2A. Changing the amino acid of AAV2 at the 324, 328 positions did not enhance muscle transduction. Amino acid positions 324, 328 of AAV2 were changed to that of AAV1. 1×10^{10} viral genome-containing particles were injected into each gastrocnemius of male Balb/c mice. Each mouse was imaged at 7, 14, 28, and 42 days post injection. The virus used in this experiment was purified using cesium chloride gradients.

FIG. 2B. Quantitation of light emission from muscle of AAV2 versus 324, 328 variant. The amount of light emitted from the experiment depicted in FIG. 2A was calculated using CMIR_image software. The regions of interest (ROI) from each leg were defined and used to calculate total photons emitted. Data is represented as an average of all 6 limbs.

FIG. 3A. Changing the amino acids of AAV2 to create the 2.5 variant greatly enhanced the muscle tropism of this vari-

4

ant. 1×10^{10} viral genome-containing particles were injected into each gastrocnemius of male Balb/c mice. Each mouse was imaged at 7, 14, 28, 42, and 95 days post injection. The virus used in this experiment was purified using heparin HPLC. The amount of light emitted from each animal was calculated using CMIR_image software. The regions of interest (ROI) from each leg were defined and used to calculate total photons emitted. Data is represented as an average of all 6 limbs.

FIG. 3B. Locations of the amino acid positions on the AAV2 capsid monomer. Depicted is a 3D ribbon structure of the AAV2 capsid monomer. The locations of the changes made in the 2.5 variant are indicated by arrows. The top arrow points to the 263, 265 regions, and the 3 bottom arrows point to the 709, 712, 720 amino acids.

FIG. 3C. The position of the 2.5 amino acids on the AAV2 pentamer. The location of the 2.5 variant amino acids is indicated by the circle. Although 2 of the amino acids lie on one portion of the individual subunit and 3 other amino acids lie on another portion of the same subunit, when 2 subunits come together to form the AAV capsid, these 5 amino acids lie in very close proximity at the 2-fold axis of symmetry.

FIG. 4. 263, 265 amino acids confer enhanced muscle tropism. Subsets (263, 265; 709, 712, 720) of the 2.5 variant were generated. 1×10^{10} viral genome-containing particles were injected into each gastrocnemius of male Balb/c mice. Each mouse was imaged at 7, 14, 28, and 42 days post injection. The virus used in this experiment was purified using cesium chloride gradients. The 263, 265 variant exhibit enhanced muscle tropism similar to the 2.5 variant, whereas the 709, 712, 720 variant exhibited a muscle tropism profile similar to AAV2.

FIG. 5. The threonine insertion at position 265 is responsible for the majority of the enhanced muscle tropism of 2.5. Subsets (263 or 265) of the 263, 265 variant were generated. 1×10^{10} viral genome-containing particles were injected into each gastrocnemius of male Balb/c mice. Each mouse was imaged at 7, 14, 28, 42, and 100 days post injection. The virus used in this experiment was purified using cesium chloride gradients. The 265 variant exhibited enhanced muscle tropism similar to the 263, 265 variant whereas the 263 variant did not exhibit enhanced muscle tropism.

FIG. 6. The 454, 461 variant exhibited enhanced muscle tropism at later time points post injection. 1×10^{10} viral genome-containing particles of AAV2 or 454, 461 were injected into each gastrocnemius of male Balb/c mice. Each mouse was imaged at 7, 14, 21, 28, and 42 days post injection. The virus used in this experiment was purified using cesium chloride gradients. 454, 461 muscle transduction was similar to AAV2 at early time points (day 7 and 14) but expression was better than AAV2 at later time points (days 21-42).

FIG. 7. Identification of the 263, 265 region of other AAV serotypes as an area of natural insertions. The capsid sequences of AAV serotypes 1 (SEQ ID NO:5), 2 (SEQ ID NO:6), 3a (SEQ ID NO:7), 3b (SEQ ID NO:7), 4 (SEQ ID NO:8), 5 (SEQ ID NO:9), 6 (SEQ ID NO:5), 7 (SEQ ID NO:10), 8 (SEQ ID NO:11) and 9 (SEQ ID NO:12) were subjected to multiple sequence alignment using the Align program in the Vector NTI software package. Shown in the box are the amino acids of the other serotypes corresponding to the amino acids identified in AAV1 as responsible for enhanced muscle tropism.

FIG. 8. Heparin binding profile of AAV variants. Equivalent particles of each AAV variant were applied to heparin agarose type 1 and allowed to bind. The columns were washed with PBS, followed by elution in sodium chloride. The number of particles present in the flow thru, washes and

5

elutions were determined via dot blot hybridization. Data is depicted as percentage of unbound particles (wash and flow thru) and bound (elution).

FIG. 9. Amino acid alignment of the capsid sequence of the 263, 265 mutant (SEQ ID NO:13) in an AAV2 background with the capsid sequence of AAV2 (SEQ ID NO:3). Amino acid 1 is numbered with respect to the VP1 sequence.

FIG. 10. Amino acid alignment of the capsid sequence of the 263, 265 mutant (SEQ ID NO:14) in an AAV3b background with the capsid sequence of AAV3b (SEQ ID NO:15). Amino acid 1 is numbered with respect to the VP1 sequence.

FIG. 11. Amino acid alignment of the capsid sequence of the 2.5 mutant (SEQ ID NO:16) in an AAV2 background with the capsid sequence of AAV2 (SEQ ID NO:3). Amino acid 1 is numbered with respect to the VP1 sequence.

FIG. 12. Luciferase activity over time (days 3, 7, 14 and 21) in mice injected with equivalent genome containing particles (1×10^{10}) of AAV1, AAV2, AAV3b, or SASTG as determined by in vivo live animal imaging.

FIG. 13. Human Factor IX (hFIX) levels detected by ELISA in sera from mice treated with an AAV2 or chimeric 2.5 vector carrying the hFIX transgene.

FIG. 14. Green fluorescent protein (GFP) transgene expression in neuronal (left arrow) and non-neuronal (right arrow) cells in the cortex of mouse brain after administration of a chimeric 2.5 vector carrying the GFP transgene.

DETAILED DESCRIPTION OF THE INVENTION

The present invention is based, in part, on the discovery that parvovirus (including AAV) capsids can be engineered to incorporate small, selective regions from other parvoviruses that confer desirable properties. The inventors have discovered that in some cases even a single amino acid insertion or substitution from a first parvovirus (e.g., an AAV) into the capsid structure of another parvovirus (e.g., an AAV) to create a chimeric parvovirus is sufficient to confer one or more of the desirable properties of the first parvovirus to the resulting chimeric parvovirus and/or to confer other properties that are not present in the first parvovirus or are present to a lesser extent.

The present invention will now be described with reference to the accompanying drawings, in which preferred embodiments of the invention are shown. This invention may, however, be embodied in different forms and should not be construed as limited to the embodiments set forth herein. Rather, these embodiments are provided so that this disclosure will be thorough and complete, and will fully convey the scope of the invention to those skilled in the art.

Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. The terminology used in the description of the invention herein is for the purpose of describing particular embodiments only and is not intended to be limiting of the invention. All publications, patent applications, patents, and other references mentioned herein are incorporated by reference in their entirety.

The designation of all amino acid positions in the AAV capsid subunits in the description of the invention and the appended claims is with respect to VP1 capsid subunit numbering.

Except as otherwise indicated, standard methods known to those skilled in the art may be used for the construction of rAAV constructs, modified capsid proteins, packaging vectors expressing the parvovirus rep and/or cap sequences, and transiently and stably transfected packaging cells. Such tech-

6

niques are known to those skilled in the art. See, e.g., SAMBROOK et al., *MOLECULAR CLONING: A LABORATORY MANUAL* 2nd Ed. (Cold Spring Harbor, N.Y., 1989); F. M. AUSUBEL et al. *CURRENT PROTOCOLS IN MOLECULAR BIOLOGY* (Green Publishing Associates, Inc. and John Wiley & Sons, Inc., New York).

Definitions.

The following terms are used in the description herein and the appended claims:

As used in the description of the invention and the appended claims, the singular forms "a," "an" and "the" are intended to include the plural forms as well, unless the context clearly indicates otherwise.

The term "parvovirus" as used herein encompasses the family Parvoviridae, including autonomously-replicating parvoviruses and dependoviruses. The autonomous parvoviruses include members of the genera *Parvovirus*, *Erythrovirus*, *Densovirus*, *Iteravirus*, and *Contravirus*. Exemplary autonomous parvoviruses include, but are not limited to, minute virus of mouse, bovine parvovirus, canine parvovirus, chicken parvovirus, feline panleukopenia virus, feline parvovirus, goose parvovirus, H1 parvovirus, muscovy duck parvovirus, and B19 virus. Other autonomous parvoviruses are known to those skilled in the art. See, e.g., BERNARD N. FIELDS et al., *VIROLOGY*, volume 2, chapter 69 (4th ed., Lippincott-Raven Publishers).

The genus *Dependovirus* contains the adeno-associated viruses (AAV), including but not limited to, AAV type 1, AAV type 2, AAV type 3 (including types 3A and 3B), AAV type 4, AAV type 5, AAV type 6, AAV type 7, AAV type 8, AAV type 9, AAV type 10, AAV type 11, avian AAV, bovine AAV, canine AAV, equine AAV, and ovine AAV or any other AAV now known or later discovered. See, e.g., BERNARD N. FIELDS et al., *VIROLOGY*, volume 2, chapter 69 (4th ed., Lippincott-Raven Publishers). Recently, a number of new AAV serotypes and clades have been identified (see, e.g., Gao et al., (2004) *J. Virology* 78:6381-6388 and Table 1).

The genomic sequences of the various serotypes of AAV and the autonomous parvoviruses, as well as the sequences of the terminal repeats (TRs), Rep proteins, and capsid subunits are known in the art. Such sequences may be found in the literature or in public databases such as GenBank. See, e.g., GenBank Accession Numbers NC 002077, NC 001401, NC 001729, NC 001863, NC 001829, NC 001862, NC 000883, NC 001701, NC 001510, AF063497, U89790, AF043303, AF028705, AF028704, J02275, J01901, J02275, X01457, AF288061, AH009962, AY028226, AY028223, NC 001358, NC 001540, AF513851, AF513852, AY530579, AY631965, AY631966; the disclosures of which are incorporated herein in their entirety. See also, e.g., Srivistava et al., (1983) *J. Virology* 45:555; Chiorini et al., (1998) *J. Virology* 71:6823; Chiorini et al., (1999) *J. Virology* 73:1309; Bantel-Schaal et al., (1999) *J. Virology* 73:939; Xiao et al., (1999) *J. Virology* 73:3994; Muramatsu et al., (1996) *Virology* 221:208; Shade et al., (1986) *J. Virol.* 58:921; Gao et al., (2002) *Proc. Nat. Acad. Sci. USA* 99:11854; international patent publications WO 00/28061, WO 99/61601, WO 98/11244; U.S. Pat. No. 6,156,303; the disclosures of which are incorporated herein in their entirety. See also Table 1. An early description of the AAV1, AAV2 and AAV3 terminal repeat sequences is provided by Xiao, X., (1996), "Characterization of Adeno-associated virus (AAV) DNA replication and integration," Ph.D. Dissertation, University of Pittsburgh, Pittsburgh, Pa. (incorporated herein in its entirety).

7

TABLE 1

Complete Genomes	GenBank Accession Number
Adeno-associated virus 1	NC_002077, AF063497
Adeno-associated virus 2	NC_001401
Adeno-associated virus 3	NC_001729
Adeno-associated virus 3B	NC_001863
Adeno-associated virus 4	NC_001829
Adeno-associated virus 5	Y18065, AF085716
Adeno-associated virus 6	NC_001862
Avian AAV ATCC VR-865	AY186198, AY629583, NC_004828
Avian AAV strain DA-1	NC_006263, AY629583
Bovine AAV	NC_005889, AY388617
Clade A	
AAV1	NC_002077, AF063497
AAV6	NC_001862
Hu.48	AY530611
Hu.43	AY530606
Hu.44	AY530607
Hu.46	AY530609
Clade B	
Hu. 19	AY530584
Hu. 20	AY530586
Hu. 23	AY530589
Hu22	AY530588
Hu24	AY530590
Hu21	AY530587
Hu27	AY530592
Hu28	AY530593
Hu. 29	AY530594
Hu63	AY530624
Hu64	AY530625
Hu13	AY530578
Hu56	AY530618
Hu57	AY530619
Hu49	AY530612
Hu58	AY530620
Hu34	AY530598
Hu35	AY530599
AAV2	NC_001401
Hu45	AY530608
Hu47	AY530610
Hu51	AY530613
Hu52	AY530614
Hu T41	AY695378
Hu S17	AY695376
Hu T88	AY695375
Hu T71	AY695374
Hu T70	AY695373
Hu T40	AY695372
Hu T32	AY695371
Hu T17	AY695370
Hu LG15	AY695377
Clade C	
Hu9	AY530629
Hu10	AY530576
Hu11	AY530577
Hu53	AY530615
Hu55	AY530617
Hu54	AY530616
Hu7	AY530628
Hu18	AY530583
Hu15	AY530580
Hu16	AY530581
Hu25	AY530591
Hu60	AY530622
Ch5	AY243021
Hu3	AY530595
Hu1	AY530575
Hu4	AY530602
Hu2	AY530585
Hu61	AY530623

8

TABLE 1-continued

Complete Genomes	GenBank Accession Number
Clade D	
Rh62	AY530573
Rh48	AY530561
Rh54	AY530567
Rh55	AY530568
Cy2	AY243020
AAV7	AF513851
Rh35	AY243000
Rh37	AY242998
Rh36	AY242999
Cy6	AY243016
Cy4	AY243018
Cy3	AY243019
Cy5	AY243017
Rh13	AY243013
Clade E	
Rh38	AY530558
Hu66	AY530626
Hu42	AY530605
Hu67	AY530627
Hu40	AY530603
Hu41	AY530604
Hu37	AY530600
Rh40	AY530559
Rh2	AY243007
Bb1	AY243023
Bb2	AY243022
Rh10	AY243015
Hu17	AY530582
Hu6	AY530621
Rh25	AY530557
Pi2	AY530554
Pi1	AY530553
Pi3	AY530555
Rh57	AY530569
Rh50	AY530563
Rh49	AY530562
Hu39	AY530601
Rh58	AY530570
Rh61	AY530572
Rh52	AY530565
Rh53	AY530566
Rh51	AY530564
Rh64	AY530574
Rh43	AY530560
AAV8	AF513852
Rh8	AY242997
Rh1	AY530556
Clade F	
Hu14 (AAV9)	AY530579
Hu31	AY530596
Hu32	AY530597
Clonal Isolate	
AAV5	Y18065, AF085716
AAV 3	NC_001729
AAV 3B	NC_001863
AAV4	NC_001829
Rh34	AY243001
Rh33	AY243002
Rh32	AY243003

The term "tropism" as used herein refers to preferential entry of the virus into certain cell or tissue types or preferential interaction with the cell surface that facilitates entry into certain cell or tissue types, optionally and preferably followed by expression (e.g., transcription and, optionally, translation) of sequences carried by the viral genome in the cell, e.g., for a recombinant virus, expression of the heterologous nucleotide sequence(s). Those skilled in the art will appreciate that transcription of a heterologous nucleic acid sequence from the viral genome may not be initiated in the absence of trans-

acting factors, e.g., for an inducible promoter or otherwise regulated nucleic acid sequence. In the case of a rAAV genome, gene expression from the viral genome may be from a stably integrated provirus, from a non-integrated episome, as well as any other form in which the virus nucleic acid may take within the cell.

As used herein, “transduction” of a cell by AAV or a chimeric virus particle means the transfer of genetic material into the cell by the incorporation of nucleic acid into the AAV or chimeric virus particle and subsequent transfer into the cell.

As used herein, a “recipient” virus or virus capsid is the parent virus or virus capsid into which the modification is introduced to produce the chimera. A “donor” virus or virus capsid as used herein is a parent virus or virus capsid from which the modification is taken and transferred into the recipient to produce the chimera.

Unless indicated otherwise, “enhanced transduction” or “enhanced tropism,” or similar terms, by the chimeric virus vectors and capsids of the invention means that there is an increase in transduction or tropism as compared with the parent virus that acted as a recipient and into which the modification was introduced to produce the chimera and/or the chimeric virus vector or capsid exhibits “enhanced transduction” or “enhanced tropism,” or similar terms, as compared with the donor parent virus, i.e., there is an increase in transduction or tropism as compared with the parent virus that acted as a donor and from which the modification was taken and introduced into the recipient to produce the chimera. In particular embodiments, the chimeric virus or capsid has enhanced tropism or transduction for muscle cells (including skeletal muscle, diaphragm muscle and/or cardiac muscle), liver, cells of the eye (including retina, retinal pigment epithelium and/or cornea), brain cells (including glial cells, astrocytes, neurons and/or oligodendrocytes), lung, epithelium cells (including gut and/or respiratory epithelial cells), dendritic cells, pancreatic cells (including islet cells), bone cells (e.g., bone marrow stem cells), hematopoietic stem cells, spleen cells, keratinocytes, fibroblasts, endothelial cells, prostate cells, progenitor cells, or germ cells.

Similarly, unless indicated otherwise, by “reduced transduction” or “reduced tropism” or similar terms by the chimeric virus vectors and capsids of the invention, it is meant that there is a decrease in transduction or tropism as compared with the parent virus that acted as a recipient and into which the modification was introduced to produce the chimera and/or the chimeric virus vector or capsid exhibits “reduced transduction” or “reduced tropism,” or similar terms, as compared with the donor parent virus, i.e., there is a decrease in transduction or tropism as compared with the parent virus that acted as a donor and from which the modification was taken and introduced into the recipient to produce the chimera. In particular embodiments, the chimeric virus or capsid has reduced tropism or transduction for muscle cells (including skeletal muscle, diaphragm muscle and/or cardiac muscle), liver, cells of the eye (including retina, retinal pigment epithelium and/or cornea), brain cells (including glial cells, astrocytes, neurons and/or oligodendrocytes), lung, epithelium cells (including gut and/or respiratory epithelial cells), dendritic cells, pancreatic cells (including islet cells), bone cells (e.g., bone marrow stem cells), hematopoietic stem cells, spleen cells, keratinocytes, fibroblasts, endothelial cells, prostate cells, progenitor cells, or germ cells.

As used herein, the term “polypeptide” encompasses both peptides and proteins, unless indicated otherwise.

A “polynucleotide” is a sequence of nucleotide bases, and may be RNA, DNA or DNA-RNA hybrid sequences (includ-

ing both naturally occurring and non-naturally occurring nucleotide), but are preferably either single or double stranded DNA sequences.

As used herein, an “isolated” polynucleotide (e.g., an “isolated DNA” or an “isolated RNA”) means a polynucleotide separated or substantially free from at least some of the other components of the naturally occurring organism or virus, for example, the cell or viral structural components or other polypeptides or nucleic acids commonly found associated with the polynucleotide.

Likewise, an “isolated” polypeptide means a polypeptide that is separated or substantially free from at least some of the other components of the naturally occurring organism or virus, for example, the cell or viral structural components or other polypeptides or nucleic acids commonly found associated with the polypeptide.

A “therapeutic polypeptide” is a polypeptide that can alleviate or reduce symptoms that result from an absence or defect in a protein in a cell or subject. Alternatively, a “therapeutic polypeptide” is a polypeptide that otherwise confers a benefit to a subject, e.g., anti-cancer effects or improvement in transplant survivability.

By the terms “treat,” “treating” or “treatment of” (or grammatically equivalent terms) it is meant that the severity of the subject’s condition is reduced or at least partially improved or ameliorated and/or that some alleviation, mitigation or decrease in at least one clinical symptom is achieved and/or there is a delay in the progression of the condition and/or prevention or delay of the onset of a disease or disorder. Thus, the terms “treat,” “treating” or “treatment of” (or grammatically equivalent terms) refer to both prophylactic and therapeutic regimens.

A “heterologous nucleotide sequence” or “heterologous nucleic acid” is typically a sequence that is not naturally occurring in the virus. Generally, the heterologous nucleic acid or nucleotide sequence comprises an open reading frame that encodes a polypeptide or nontranslated RNA.

As used herein, the term “vector” or “delivery vector” may refer to a parvovirus (e.g., AAV) particle that functions as a nucleic acid delivery vehicle, and which comprises viral DNA (i.e., the vector genome) packaged within a parvovirus (e.g., AAV) capsid. Alternatively, in some contexts, the term “vector” may be used to refer to the vector genome/vDNA in the absence of the capsid.

As used herein, a “recombinant parvovirus vector genome” is a parvovirus genome (i.e., vDNA) that comprises at least one terminal repeat (e.g., two terminal repeats) and one or more heterologous nucleotide sequences. A “recombinant parvovirus particle” comprises a recombinant parvovirus vector genome packaged within a parvovirus capsid.

A “rAAV vector genome” or “rAAV genome” is an AAV genome (i.e., vDNA) that comprises at least one terminal repeat (e.g., two terminal repeats) and one or more heterologous nucleotide sequences. rAAV vectors generally require only the 145 base terminal repeat(s) (TR(s)) in cis to generate virus. All other viral sequences are dispensable and may be supplied in trans (Muzyczka, (1992) *Curr. Topics Microbiol. Immunol.* 158:97). Typically, the rAAV vector genome will only retain the minimal TR sequence(s) so as to maximize the size of the transgene that can be efficiently packaged by the vector. The structural and non-structural protein coding sequences may be provided in trans (e.g., from a vector, such as a plasmid, or by stably integrating the sequences into a packaging cell). The rAAV vector genome optionally comprises two AAV TRs, which generally will be at the 5’ and 3’

ends of the heterologous nucleotide sequence(s), but need not be contiguous thereto. The TRs can be the same or different from each other.

A “rAAV particle” comprises a rAAV vector genome packaged within an AAV capsid.

A “parvovirus terminal repeat” may be from any parvovirus, including autonomous parvoviruses and AAV (all as defined above). An “AAV terminal repeat” may be from any AAV, e.g., serotypes 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 and 11. The term “terminal repeat” includes synthetic sequences that function as an AAV inverted terminal repeat, such as the “double-D sequence” as described in U.S. Pat. No. 5,478,745 to Samulski et al., the disclosure of which is incorporated in its entirety herein by reference. The AAV terminal repeats need not have a wild-type terminal repeat sequence (e.g., a wild-type sequence may be altered by insertion, deletion, truncation or missense mutations), as long as the terminal repeat mediates the desired functions, e.g., replication, virus packaging, integration, and/or provirus rescue, and the like.

The capsid structure of autonomous parvoviruses and AAV are described in more detail in BERNARD N. FIELDS et al., VIROLOGY, volume 2, chapters 69 & 70 (4th ed., Lippincott-Raven Publishers).

The chimeric virus vector of the invention can further be a “targeted” virus vector (e.g., having a directed tropism) and/or a “hybrid” parvovirus (i.e., in which the rAAV genome and viral capsid are from different parvoviruses) as described in international patent publication WO 00/28004 and Chao et al., (2000) *Molecular Therapy* 2:619. In particular embodiments, the rAAV genome and virus capsid are from different AAV.

In particular embodiments, all of the subunits of the virus capsid are derived from the same AAV capsid protein backbone. In other embodiments, the virus capsid comprises capsid proteins that are derived from different AAV backbones.

The chimeric viruses of the invention can further be duplexed parvovirus particles as described in international patent publication WO 01/92551.

Accordingly, as used herein, the terms “chimeric parvovirus” and “chimeric AAV” encompass hybrid, targeted and duplexed virus particles, as well as other modified forms of parvoviruses and AAV.

Chimeric Virus Vectors.

Selected regions within the AAV capsid can confer desirable properties to other AAV including, but not limited to, enhanced transduction ability (e.g., enhanced transduction of skeletal muscle cells), higher expression of a transgene delivered by the recombinant genome and/or earlier onset of transgene expression. These selected regions can be transferred or “engineered” into other AAV capsids to confer the property(ies) of interest to the resulting chimeric particle. To illustrate, in the case of AAV2, the inventors have discovered that with a single amino acid insertion from AAV1, AAV2 can acquire the enhanced transduction capability of AAV1, while retaining the ease of purification (i.e., retains the ability to bind heparin) and the known safety features of AAV2. Further, in particular embodiments, the resulting chimeric virus has a different immunological profile than the parent virus (e.g., is weakly or not at all recognized by neutralizing antiserum to the parent virus), thereby allowing for repeat administration to subjects that have developed antibodies against the parent virus. For example, the chimeric 2.5 vector described herein is only weakly recognized by neutralizing antisera against AAV2.

In the present studies, the inventors have compared the linear amino acid sequences of two serotypes with high levels

of skeletal muscle transduction (AAV1 and AAV7) with two that are less effective in transducing skeletal muscle (AAV2 and AAV8). Although these four serotypes overall exhibit a high degree of sequence similarity, there are regions of divergence and the characteristics of these four serotypes as delivery vectors are distinct. Using sequence analysis, a number of candidate amino acids in AAV 1 and/or AAV7 were identified for incorporation into AAV2 to determine if they would confer the enhanced skeletal muscle transduction of AAV1/7 to AAV2. This initial pool of candidate positions was narrowed considerably based on analysis of the known crystal structure of AAV2, e.g., to identify amino acid positions that are located in regions that are likely to affect the biology of the virus. This smaller group of amino acids was then substituted and/or inserted into AAV2, either individually or in combination and several of the resulting mutants demonstrated the enhanced skeletal muscle transduction of AAV1. Corresponding modifications can readily be introduced into other AAV based on the present disclosure. A “corresponding” modification can be an insertion and/or a substitution and/or a deletion. For example, a modification that is an insertion in AAV2 may be a substitution mutation in another AAV. Further, in still other AAV, a deletion mutation may bring an amino acid into the desired position within the capsid subunits. “Corresponding” modifications in other AAV will be apparent to those skilled in the art.

According to the present invention, a “selective” amino acid change(s) is introduced into the virus capsid. This approach is in contrast to previous work with whole subunit or large domain swaps between AAV serotypes (see, e.g., international patent publication WO 00/28004 and Hauck et al., (2003) *J. Virology* 77:2768-2774). A “selective” amino acid change results in the insertion and/or substitution and/or deletion of less than about 20, 18, 15, 12, 10, 9, 8, 7, 6, 5, 4 or 3 contiguous amino acids. In particular embodiments, only two contiguous amino acids or even point mutations (i.e., one amino acid) are inserted and/or substituted into and/or deleted from the capsid subunits.

One or more selective amino acid changes can be introduced at different positions within the capsid subunits. For example, in particular embodiments, two, three, four, five, six, seven, eight, nine, ten or more selective amino acid changes can be introduced into the AAV capsid subunits.

It will be understood that the term “chimeric virus vector” or “chimeric capsid” excludes those virus vectors or capsids that have the indicated amino acids at the specified positions in their native state (e.g., in the recipient virus and/or in the wild-type virus).

The invention contemplates that the chimeric viruses of the invention can be produced by modifying the capsids of any AAV now known or later discovered. Further, the recipient parent AAV that is to be modified can be one of the characterized AAV, e.g., AAV2, AAV3a or 3b, AAV4, AAV5, AAV8, AAV9, AAV10 or AAV11 (see also Table 1). Alternatively, the recipient parent virus may already have modifications/alterations as compared with the naturally occurring viruses (i.e., is derived from a wild-type AAV, e.g., AAV2, AAV3a, AAV3b, AAV4, AAV5, AAV6, AAV7, AAV8, AAV9, AAV10 and/or AAV11 or any other AAV now known or later discovered). Such viruses are also within the scope of the present invention. For example, the recipient virus can be an AAV that is derived from AAV8, but has a heparan sulfate-binding domain (e.g., from the AAV2 capsid) incorporated therein or can be an AAV that is modified to contain a poly-His sequence to facilitate purification. As another illustrative example, the AAV can be derived from any of the known serotypes or clades, but have a peptide targeting sequence incorporated

therein. As yet another possibility, the AAV capsid can comprise capsid subunits from different serotypes. Thus, in particular embodiments, the recipient virus comprises a capsid from an AAV serotype or clade that has been modified to comprise sequences that are not from that serotype or clade (e.g., are exogenous to the wild-type virus). Further, a donor parent AAV from a modification that is to be transferred into a recipient is taken can be one of the characterized AAV, e.g., AAV2, AAV3a or 3b, AAV4, AAV5, AAV8, AAV9, AAV10 or AAV11, but is not so limited.

In representative embodiments, the invention provides a chimeric virus vector comprising: (a) a chimeric AAV capsid comprising a selective amino acid insertion following any of amino acid position 260, 261, 262, 263, 264, 265, 266, 267 and/or 268 (e.g., following amino acid position 264) in one or more of the AAV2 capsid subunits (VP1 numbering) or a corresponding change in one or more capsid subunits from other AAV; and (b) a nucleic acid comprising an AAV terminal repeat sequence and a heterologous nucleic acid sequence; wherein the nucleic acid is packaged within the chimeric AAV capsid. In particular embodiments, the chimeric virus vector has enhanced transduction or tropism, an altered immunological profile, enhanced transgene expression and/or earlier onset of transgene expression as compared with the recipient parent rAAV vector and/or a donor parent rAAV vector. By "following amino acid position X" it is intended that the insertion immediately follows the indicated amino acid position (for example, "following amino acid position 264" indicates a point insertion at position 265 or a larger insertion, e.g., from positions 265 to 268, etc.) or is within two or three amino acids of the indicated amino acid position.

The designation of all amino acid positions in the description of the invention and the appended claims is with respect to VP1 numbering. It will be understood by those skilled in the art that the modifications described herein can result in modifications in all of the VP1, VP2 and VP3 capsid subunits. In particular embodiments, the modifications of the invention are found in one, two or three of the subunits of the AAV capsid.

Those skilled in the art will appreciate that for some AAV the corresponding modification will be an insertion and/or a substitution, depending on whether the corresponding amino acid positions are partially or completely present in the virus or, alternatively, are completely absent. Likewise, when modifying AAV other than AAV2, the specific amino acid position(s) may be different than the position in AAV2. The corresponding amino acid position(s) will be readily apparent to those skilled in the art using well-known sequence alignment techniques (see, e.g., FIG. 7).

In particular embodiments, the modifications described herein can be introduced into the capsid subunit(s) at the position that corresponds to the position of the amino acid(s) of interest in the AAV1, AAV6 and/or AAV7 capsid subunit(s) or any other AAV from which the modification is derived. Based on crystal structure analysis, it will be clear that in some instances the insertion/substitution can be moved 1, 2, 3, 4 or 5 or even more amino acids in either direction and still confer the desired characteristic. Because many of the modifications are made in loop structures, those skilled in the art will understand that the result of the modification can be driven more by physical presentation on the surface of the virion than the specific amino acid position.

Amino acids to be substituted and/or inserted according to the present invention can be any naturally occurring amino acids, modified forms thereof or synthetic amino acids.

In representative embodiments, the insertion and/or substitution and/or deletion in the capsid subunit(s) results in the insertion, substitution and/or repositioning of an amino acid that maintains the hydrophilic loop structure in that region and/or an amino acid that alters the configuration of the loop structure, a charged amino acid, or an amino acid that can be phosphorylated or sulfated or otherwise acquire a charge by post-translational modification (e.g., glycosylation) following any of amino acid position 260, 261, 262, 263, 264, 265, 266, 267 and/or 268 (e.g., following position 264) in an AAV2 capsid subunit(s) or a corresponding change in a capsid subunit(s) of another AAV. In particular embodiments, the chimeric virus vector has enhanced transduction for one or more cell types compared with the recipient parent rAAV vector and/or a donor parent rAAV vector. Suitable amino acids include aspartic acid, glutamic acid, lysine, arginine, threonine, serine, tyrosine, glycine, alanine, proline, asparagine or glutamine. In particular embodiments, a threonine is inserted or substituted into the capsid subunit.

According to this aspect of the invention, in particular embodiments the chimeric virus vector comprises a chimeric AAV capsid comprising an amino acid insertion following amino acid position 264 in an AAV2, AAV3a or AAV3b capsid subunit(s) or in the corresponding position in an AAV2, AAV3a or AAV3b capsid that has been modified to comprise non-AAV2, AAV3a or AAV3b sequences, respectively (i.e., is derived from AAV2, AAV3a or AAV3b). The amino acid corresponding to position 264 in an AAV2 (or AAV3a or AAV3b) capsid subunit(s) will be readily identifiable in the starting virus that has been derived from AAV2 (or AAV3a or AAV3b), which can then be further modified according to the present invention.

In particular embodiments, the chimeric virus vectors can comprise a chimeric AAV2, AAV3a or AAV3b capsid comprising an amino acid insertion following amino acid position 264 in an AAV2, AAV3a or AAV3b capsid subunit(s). Suitable amino acid insertions are described above. Illustrative examples of corresponding mutations in other AAV are shown in Table 2 (Position 2). Amino acid substitutions and insertions are as described above. In particular embodiments, the insertion or substitution is a threonine (excepting AAV1, AAV6 and other AAV that have a threonine at this position).

TABLE 2

Serotype	Position 1	Position 2
AAV1	A263X	T265X
AAV2	Q263X	-265X
AAV3a	Q263X	-265X
AAV3b	Q263X	-265X
AAV4	S257X	-259X
AAV5	G253X	V255X
AAV6	A263X	T265X
AAV7	E264X	A266X
AAV8	G264X	S266X
AAV9	S263X	S265X

Where, (X) → mutation to any amino acid (-) → insertion of any amino acid
Note:

Position 2 inserts are indicated by the site of insertion

The invention further provides chimeric virus vectors comprising a chimeric AAV capsid comprising one or more selective amino acid insertions and/or substitutions selected from the group consisting of:

- a selective amino acid insertion following any of amino acid position 260, 261, 262, 263, 264, 265, 266, 267 and/or 268 (e.g., following amino acid position 264) in

15

- an AAV2 capsid subunit(s) or a corresponding change in a capsid subunit(s) from other AAV;
- (b) a selective amino acid substitution at an amino acid from position 260 to 266 (e.g., position 263) in an AAV2 capsid subunit(s) or a corresponding change in a capsid subunit(s) from other AAV;
 - (c) a selective amino acid substitution at amino acid position 705 in an AAV2 capsid subunit(s) or a corresponding change in a capsid subunit(s) from other AAV;
 - (d) a selective amino acid substitution at amino acid position 708 in an AAV2 capsid subunit(s) or a corresponding change in a capsid subunit(s) from other AAV;
 - (e) a selective amino acid substitution at amino acid position 716 in an AAV2 capsid subunit(s) or a corresponding change in a capsid subunit(s) from other AAV;
 - (f) a selective amino acid substitution at an amino acid from position 447 to 453 (e.g., position 450) in an AAV2 capsid subunit(s) or a corresponding change in a capsid subunit(s) from other AAV; and
 - (g) a selective amino acid substitution at an amino acid from position 454 to 460 (e.g., position 457) in an AAV2 capsid subunit(s) or a corresponding change in a capsid subunit(s) from other AAV.

These modifications can be used alone or in any combination with each other and in any combination with any of the other modifications described herein.

In particular embodiments, the chimeric virus vector has enhanced transduction, enhanced transgene expression, earlier onset of transgene expression and/or an altered immunological profile as compared with the recipient parent rAAV vector and/or a donor parent rAAV vector.

In representative embodiments, the chimeric virus vector comprises a chimeric AAV2 capsid comprising an amino acid substitution at amino acid position 263 in an AAV2 capsid subunit(s) or in the corresponding position in an AAV2 capsid that has been modified to comprise non-AAV2 sequences. Alternatively, the chimeric virus vector can comprise a chimeric AAV3b capsid comprising an amino acid substitution at amino acid position 263 in an AAV3b capsid subunit(s) or in the corresponding position in an AAV3b capsid that has been modified to comprise non-AAV3b sequences.

Corresponding modifications can be made in other AAV. For example, in still other embodiments, the substitution is at amino acid position 263 of AAV1, a substitution at amino acid position 263 in AAV3a or AAV3b, or at the corresponding position(s) in an AAV capsid from any of the foregoing serotypes that has been modified to comprise exogenous sequences. Nonlimiting examples of corresponding mutations are shown in Table 2 (Position 1). Suitable amino acid substitutions are described below. In particular embodiments, an alanine is substituted into the capsid subunit(s) at the indicated position (excepting AAV1 and AAV6 and other AAV that have an alanine at this position).

Suitable amino acid substitutions include but are not limited to substitutions of small nonpolar amino acids such as alanine, glycine, valine, leucine or isoleucine or even proline, asparagine, serine or threonine into the capsid subunits (e.g., substitution of these amino acids for the glutamine found at position 263 in the AAV2, AAV3a or AAV3b capsid subunits). In particular embodiments, an alanine is substituted into the capsid subunit(s).

Another chimeric virus vector of the invention comprises a chimeric AAV2 capsid comprising an amino acid substitution at amino acid position 705 in an AAV2 capsid subunit(s) or at the corresponding position in an AAV2 capsid subunit that has been modified to comprise non-AAV2 sequences.

16

Corresponding changes can be made in other AAV. For example, in other embodiments, the invention provides for a chimeric virus vector having an amino acid substitution at amino acid position 706 of an AAV1 capsid subunit(s), a substitution at amino acid position 706 of an AAV3a capsid subunit(s), a substitution at amino acid position 706 of an AAV3b capsid subunit(s), a substitution at amino acid position 707 of an AAV7 capsid subunit(s), a substitution at amino acid position 708 of an AAV8 capsid subunit(s), or a substitution at amino acid position 706 of an AAV9 capsid subunit(s); or at the corresponding position(s) in an AAV capsid from any of the foregoing serotypes that has been modified to comprise exogenous sequences or at the corresponding position(s) of any other AAV.

In representative embodiments of the invention described in the previous two paragraphs, suitable amino acid substitutions include but are not limited to substitutions of serine, threonine, tyrosine, proline, glutamine, alanine, glycine, valine, leucine or isoleucine (e.g., substituted for the asparagine at this position in an AAV2 capsid subunit(s)). In particular embodiments, alanine is substituted into the capsid subunit(s).

A further chimeric virus vector according to the invention comprises a chimeric AAV2 capsid comprising an amino acid substitution at amino acid position 708 in an AAV2 capsid subunit(s) or at the corresponding position in an AAV2 capsid subunit(s) that has been modified to comprise non-AAV2 sequences.

Corresponding modifications can be made in other AAV. In some embodiments, the invention provides for a chimeric virus vector having an amino acid substitution at amino acid position 709 of an AAV1 capsid subunit(s), an amino acid substitution at amino acid position 709 of an AAV3a capsid subunit(s), an amino acid substitution at amino acid position 709 of an AAV3b capsid subunit(s), an amino acid substitution at amino acid position 710 of an AAV7 capsid subunit(s), an amino acid substitution at amino acid position 711 of an AAV8 capsid subunit(s), or an amino acid substitution at amino acid position 709 of an AAV9 capsid subunit(s); or at the corresponding position in an AAV capsid subunit(s) from any of the foregoing AAV that has been modified to comprise exogenous sequences or at a corresponding position in any other AAV.

In representative embodiments of the invention described in the previous two paragraphs, the substitution results in a substitution of serine, threonine, tyrosine, proline, asparagine, glutamine, alanine, glycine, leucine or isoleucine into the capsid subunit(s) (e.g., for the valine at this position in an AAV2 capsid subunit(s)). One exemplary substitution is a substitution of alanine into the capsid subunit(s).

The invention also provides a chimeric virus vector comprising a chimeric AAV2 capsid comprising an amino acid substitution at amino acid position 716 in an AAV2 capsid subunit(s) or at the corresponding position in an AAV2 capsid that has been modified to comprise non-AAV2 sequences.

Corresponding modifications can be made in other AAV. In particular embodiments, the invention provides for a chimeric virus vector having an amino acid substitution at amino acid position 717 of an AAV1 capsid subunit(s), an amino acid substitution at amino acid position 717 of an AAV3a capsid subunit(s), an amino acid substitution at amino acid position 717 of an AAV3b capsid subunit(s), an amino acid substitution at amino acid position 718 of an AAV7 capsid subunit(s), an amino acid substitution at amino acid position 719 of an AAV8 capsid subunit(s), or an amino acid substitution at amino acid position 717 of an AAV9 capsid subunit(s); or at the corresponding position in an AAV capsid subunit(s) from

17

any of the foregoing AAV that has been modified to comprise exogenous sequences or at the corresponding position in any other AAV.

In particular aspects of the embodiments described in the previous two paragraphs, the substitution is a substitution of an amino acid that cannot be phosphorylated for an amino acid that can be phosphorylated (e.g., threonine). Alternatively, the substitution can be a substitution of a serine, tyrosine, glycine, alanine, proline, valine, leucine, isoleucine, asparagine or glutamine into the capsid subunit (e.g., substitution of the threonine at this position in AAV2). One particular chimeric virus comprises an asparagine substituted at this position into the capsid subunit(s).

A further chimeric virus vector according to the invention is a chimeric virus vector comprising a chimeric AAV2 capsid comprising an amino acid substitution at any of amino acid positions 447, 448, 449, 450, 451, 452 and/or 453 (e.g., at position 450) in an AAV2 capsid subunit(s) or at the corresponding position in an AAV2 capsid subunit(s) that has been modified to comprise non-AAV2 sequences or at the corresponding position of any other AAV. In particular embodiments, the substitution is a substitution of an amino acid that cannot be phosphorylated for an amino acid that can be phosphorylated (e.g., threonine). Alternatively, the substitution can be a substitution of a serine, tyrosine, glycine, alanine, proline, valine, leucine, isoleucine, asparagine or glutamine into the capsid subunit (e.g., substitution of the threonine at position 450 in AAV2). One particular chimeric virus comprises an asparagine substituted at this position into the capsid subunits.

Also provided is a chimeric virus vector comprising a chimeric AAV2 capsid comprising an amino acid substitution at any of amino acid position 454, 455, 456, 457, 458, 459 and/or 460 (e.g., at position 457) in an AAV2 capsid subunit(s) or at the corresponding position in an AAV2 capsid subunit(s) that has been modified to comprise non-AAV2 sequences or at the corresponding position of any other AAV. Suitable amino acid substitutions include substitution of a glycine, alanine, valine, lysine, isoleucine, proline, serine, threonine or asparagine into the capsid subunit(s). In representative embodiments, an asparagine is substituted into the capsid subunit (e.g., an asparagine is substituted for the glutamine found at position 457 in the AAV2 capsid subunits).

One non-limiting example of a chimeric virus vector of the invention comprises: (a) a chimeric AAV capsid (e.g., AAV2 capsid) comprising: (i) a selective amino acid insertion following amino acid position 264 in an AAV2 capsid subunit(s) or a corresponding change in a capsid subunit(s) from other AAV; and (ii) a selective amino acid substitution at amino acid position 263 in the AAV2 capsid subunit(s) or a corresponding change in the capsid subunit(s) from other AAV; and (b) a nucleic acid comprising an AAV TR sequence and a heterologous nucleic acid sequence; wherein the nucleic acid is packaged within the chimeric AAV capsid. In particular embodiments, the chimeric virus has enhanced transduction, enhanced transgene expression, earlier transgene expression and/or an altered immunological profile as compared with the recipient parent rAAV vector and/or a donor parent rAAV vector. For example, the chimeric virus vector can comprise a chimeric AAV2, AAV3a or AAV3b capsid in which a threonine has been inserted following amino acid position 264 in an AAV2 or AAV3b capsid subunit(s) and an alanine has been substituted for glutamine at amino acid position 263 in the AAV2, AAV3a or AAV3b capsid subunit(s); or these changes can be made at the corresponding positions in an AAV2, AAV3a or AAV3b capsid that has been modified to comprise

18

non-AAV2, non-AAV3a or non-AAV3b sequences, respectively, or in the corresponding positions in any other AAV. The sequence of the 263, 265 variant described herein is shown in AAV2 (FIG. 9) and AAV3b (FIG. 10) backgrounds and compared with the sequence of the recipient parent virus.

Another exemplary chimeric virus vector of the invention comprises (a) a chimeric AAV capsid e.g., (AAV2 capsid) comprising: (i) a selective amino acid insertion following amino acid position 264 in an AAV2 capsid subunit(s) or a corresponding change in a capsid subunit(s) from other AAV; and (ii) a selective amino acid substitution at amino acid position 263 in the AAV2 capsid subunit(s) or a corresponding change in the capsid subunit(s) from other AAV; (iii) a selective amino acid substitution at amino acid position 705 in the AAV2 capsid subunit(s) or a corresponding change in the capsid subunit(s) from other AAV; (iv) a selective amino acid substitution at amino acid position 708 in the AAV2 capsid subunit(s) or a corresponding change in the capsid subunit(s) from other AAV; and (v) a selective amino acid substitution at amino acid position 716 in the AAV2 capsid subunit(s) or a corresponding change in the capsid subunit(s) from other AAV; and (b) a nucleic acid comprising an AAV TR sequence and a heterologous nucleic acid sequence; wherein the nucleic acid is packaged within the chimeric AAV capsid. In particular embodiments, the chimeric virus has enhanced transduction, enhanced transgene expression, earlier transgene expression and/or an altered immunological profile as compared with the recipient rAAV vector and/or a donor parent rAAV vector.

According to this embodiment, the chimeric virus vector can comprise a chimeric AAV capsid (e.g., AAV2 capsid) comprising:

- (a) a threonine insertion following amino acid position 264 in an AAV2 capsid subunit(s);
- (b) an alanine for glutamine substitution at amino acid position 263 in the AAV2 capsid subunit(s);
- (c) an alanine for asparagine substitution at amino acid position 705 in the AAV2 capsid subunit(s);
- (d) an alanine for valine substitution at amino acid position 708 in the AAV2 capsid subunit(s); and
- (e) an asparagine for threonine substitution at amino acid position 716 in the AAV2 capsid subunit(s);

or these changes can be made at the corresponding positions in an AAV2 capsid that has been modified to comprise non-AAV2 sequences or at the corresponding positions in other AAV.

In particular embodiments, the chimeric virus vector can be the chimeric 2.5 variant disclosed herein. The sequence of the 2.5 mutant as compared with AAV2 is shown in FIG. 11.

The invention further provides a chimeric virus vector comprising:

- (a) a chimeric AAV capsid (e.g., AAV2 capsid) comprising:
 - (i) a selective amino acid substitution at amino acid position 450 in an AAV2 capsid subunit(s) or a corresponding change in a capsid subunit(s) of other AAV; and
 - (ii) a selective amino acid substitution at amino acid position 457 in the AAV2 capsid subunit(s) or a corresponding change in a capsid subunit(s) of other AAV; and
- (b) a nucleic acid comprising an AAV TR sequence and a heterologous nucleic acid sequence; wherein the nucleic acid is packaged within the chimeric AAV capsid.

In particular embodiments, the chimeric viral vector has enhanced transduction, enhanced transgene expression, earlier transgene expression and/or an altered immunological

19

profile as compared with the recipient parent virus vector and/or a donor parent rAAV vector.

According to particular embodiments, a chimeric AAV2 capsid comprises (i) an asparagine for threonine substitution at amino acid position 450 in an AAV2 capsid subunit(s); and (ii) an asparagine for glutamine substitution at amino acid position 457 in the AAV2 capsid subunit(s); or these changes can be made at the corresponding positions in an AAV2 capsid subunit(s) that has been modified to comprise non-AAV2 sequences or at the corresponding positions of any other AAV.

Another exemplary chimeric virus vector of the invention comprises:

- (a) a chimeric AAV capsid (e.g., AAV2 capsid) comprising:
 - (i) a selective amino acid insertion following amino acid position 264 in an AAV2 capsid subunit(s) or a corresponding change in a capsid subunit(s) of other AAV;
 - (ii) a selective amino acid substitution at amino acid position 450 in the AAV2 capsid subunit(s) or a corresponding change in a capsid subunit(s) of other AAV; and
 - (iii) a selective amino acid substitution at amino acid position 457 in the AAV2 capsid subunit(s) or a corresponding change in a capsid subunit(s) of other AAV; and
- (b) a nucleic acid comprising an AAV TR sequence and a heterologous nucleic acid sequence; wherein the nucleic acid is packaged within the chimeric AAV capsid.

In particular embodiments, the chimeric viral vector has enhanced transduction, enhanced transgene expression, earlier transgene expression and/or an altered immunological profile as compared with the recipient parent virus vector and/or a donor parent rAAV vector.

According to particular aspects of this embodiment of the invention, the chimeric virus vector comprises a chimeric AAV2 capsid comprising:

- (a) a threonine insertion following amino acid position 264 in the AAV2 capsid subunit(s);
- (b) an asparagine for threonine substitution at amino acid position 450 in the AAV2 capsid subunit(s); and
- (c) an asparagine for glutamine substitution at amino acid position 457 in the AAV2 capsid subunit(s);

or these changes can be made at the corresponding positions in an AAV2 capsid that has been modified to comprise non-AAV2 sequences or at the corresponding positions of any other AAV.

The chimeric virus vectors of the invention generally comprise a rAAV vector genome, i.e., the AAV vector genome comprises one or more heterologous nucleic acids that encode a polypeptide or a non-translated RNA. Heterologous nucleic acids are discussed in more detail hereinbelow. Chimeric Capsids.

The present invention further encompasses chimeric virus capsids essentially as described above, i.e., in the absence of a rAAV vector genome.

The chimeric virus capsids can be used as "capsid vehicles," as has been described in U.S. Pat. No. 5,863,541 (the disclosure of which is incorporated by reference herein in its entirety). Molecules that can be packaged by the chimeric virus capsids and transferred into a cell include heterologous DNA, RNA, polypeptides, small organic molecules, or combinations of the same. Heterologous molecules are defined as those that are not naturally found in an AAV infection, i.e., those not encoded by a wild-type AAV genome. Further, therapeutically useful molecules can be associated with the outside of the chimeric virus capsid for transfer of the molecules into host target cells. Such associated molecules can

20

include DNA, RNA, small organic molecules, carbohydrates, lipids and/or polypeptides. In one embodiment of the invention the therapeutically useful molecule is covalently linked (i.e., conjugated or chemically coupled) to the capsid proteins. Methods of covalently linking molecules are known by those skilled in the art.

The chimeric virus capsids of the invention also find use in raising antibodies against the novel capsid structures. As a further alternative, an exogenous amino acid sequence may be inserted into the parvovirus capsid for antigen presentation to a cell, e.g., for administration to a subject to produce an immune response to the exogenous amino acid sequence.

According to some embodiments, chimeric virus capsids can be administered to a subject concurrently (e.g., within minutes or hours of each other) with an AAV vector or chimeric virus vector according to the invention. Further, the invention provides compositions and pharmaceutical formulations comprising the inventive chimeric virus capsids and an AAV vector or chimeric virus vector of the invention.

The invention also provides nucleic acids (optionally, isolated nucleic acids) encoding the chimeric virus capsids and capsid subunits of the invention. Further provided are vectors comprising the nucleic acids, and cells (in vivo or in culture) comprising the nucleic acids and/or vectors of the invention. Such nucleic acids, vectors and cells can be used, for example, as reagents (e.g., helper packaging constructs or packaging cells) for the production of chimeric virus capsids or vectors as described herein.

Methods of Producing Chimeric Virus Vectors.

The present invention further provides methods of producing the inventive chimeric virus vector.

In one particular embodiment, the present invention provides a method of producing a recombinant chimeric virus vector, comprising providing to a cell, (a) a rAAV template comprising (i) one or more heterologous nucleotide sequences, and (ii) packaging signal sequences sufficient for the encapsidation of the AAV template into chimeric virus particles, and (b) AAV sequences sufficient for replication and encapsidation of the rAAV template into chimeric viral particles (e.g., the AAV rep and chimeric cap sequences comprising one or more selective insertions, deletions and/or substitutions from other AAV therein). The rAAV template and AAV replication and capsid sequences are provided under conditions such that recombinant chimeric virus particles comprising the rAAV template packed within the chimeric capsid are produced in the cell. The method can further comprise the step of collecting the virus particles from the cell. Virus particles may be collected from the medium and/or by lysing the cells.

The cell is typically a cell that is permissive for AAV viral replication. Any suitable cell known in the art may be employed. Mammalian cells are preferred. Also preferred are trans-complementing packaging cell lines that provide functions deleted from a replication-defective helper virus, e.g., 293 cells or other E1a trans-complementing cells.

The AAV replication and capsid sequences may be provided by any method known in the art. Current protocols typically express the AAV rep/cap genes on a single plasmid. The AAV replication and packaging sequences need not be provided together, although it may be convenient to do so. The AAV rep and/or cap sequences may be provided by any viral or non-viral vector. For example, the rep/cap sequences may be provided by a hybrid adenovirus or herpesvirus vector (e.g., inserted into the E1a or E3 regions of a deleted adenovirus vector). EBV vectors may also be employed to express the AAV cap and rep genes. One advantage of this method is that EBV vectors are episomal, yet will maintain a high copy

number throughout successive cell divisions (i.e., are stably integrated into the cell as extra-chromosomal elements, designated as an "EBV based nuclear episome," see Margolski, (1992) *Curr. Top. Microbiol. Immun.* 158:67).

As a further alternative, the rep/cap sequences may be stably carried (episomal or integrated) within a cell.

Typically, and preferably, the AAV rep/cap sequences will not be flanked by the AAV packaging sequences (e.g., AAV ITRs), to prevent rescue and/or packaging of these sequences.

The rAAV template can be provided to the cell using any method known in the art. For example, the rAAV template may be supplied by a non-viral (e.g., plasmid) or viral vector. In particular embodiments, the rAAV template is supplied by a herpesvirus or adenovirus vector (e.g., inserted into the E1a or E3 regions of a deleted adenovirus). As another illustration, Palombo et al., (1998) *J. Virology* 72:5025, describe a baculovirus vector carrying a reporter gene flanked by the AAV ITRs. EBV vectors may also be employed to deliver the rAAV template, as described above with respect to the rep/cap genes.

In another representative embodiment, the rAAV template is provided by a replicating rAAV virus. In still other embodiments, an AAV provirus is stably integrated into the chromosome of the cell.

To obtain maximal virus titers, helper virus functions (e.g., adenovirus or herpesvirus) essential for a productive AAV infection are generally provided to the cell. Helper virus sequences necessary for AAV replication are known in the art. Typically, these sequences are provided by a helper adenovirus or herpesvirus vector. Alternatively, the adenovirus or herpesvirus sequences can be provided by another non-viral or viral vector, e.g., as a non-infectious adenovirus miniplasmid that carries all of the helper genes required for efficient AAV production as described by Ferrari et al., (1997) *Nature Med.* 3:1295, and U.S. Pat. Nos. 6,040,183 and 6,093,570.

Further, the helper virus functions may be provided by a packaging cell with the helper genes integrated in the chromosome or maintained as a stable extrachromosomal element. It is preferred that these helper virus sequences cannot be packaged into AAV virions, e.g., are not flanked by AAV ITRs.

Those skilled in the art will appreciate that it may be advantageous to provide the AAV replication and capsid sequences and the helper virus sequences (e.g., adenovirus sequences) on a single helper construct. This helper construct may be a non-viral or viral construct, but is preferably a hybrid adenovirus or hybrid herpesvirus comprising the AAV rep/cap genes.

In one particular preferred embodiment, the AAV rep/cap sequences and the adenovirus helper sequences are supplied by a single adenovirus helper vector. This vector further contains the rAAV template. The AAV rep/cap sequences and/or the rAAV template may be inserted into a deleted region (e.g., the E1a or E3 regions) of the adenovirus.

In a further embodiment, the AAV rep/cap sequences and the adenovirus helper sequences are supplied by a single adenovirus helper vector. The rAAV template is provided as a plasmid template.

In another illustrative embodiment, the AAV rep/cap sequences and adenovirus helper sequences are provided by a single adenovirus helper vector, and the rAAV template is integrated into the cell as a provirus. Alternatively, the rAAV template is provided by an EBV vector that is maintained within the cell as an extrachromosomal element (e.g., as an EBV based nuclear episome).

In a further exemplary embodiment, the AAV rep/cap sequences and adenovirus helper sequences are provided by a

single adenovirus helper. The rAAV template is provided as a separate replicating viral vector. For example, the rAAV template may be provided by a rAAV particle or a second recombinant adenovirus particle.

According to the foregoing methods, the hybrid adenovirus vector typically comprises the adenovirus 5' and 3' cis sequences sufficient for adenovirus replication and packaging (i.e., the adenovirus terminal repeats and PAC sequence). The AAV rep/cap sequences and, if present, the rAAV template are embedded in the adenovirus backbone and are flanked by the 5' and 3' cis sequences, so that these sequences may be packaged into adenovirus capsids. As described above, it is preferred that the adenovirus helper sequences and the AAV rep/cap sequences are not flanked by the AAV packaging sequences (e.g., the AAV ITRs), so that these sequences are not packaged into the AAV virions.

Herpesvirus may also be used as a helper virus in AAV packaging methods. Hybrid herpesviruses encoding the AAV Rep protein(s) may advantageously facilitate for more scalable AAV vector production schemes. A hybrid herpes simplex virus type I (HSV-1) vector expressing the AAV-2 rep and cap genes has been described (Conway et al., (1999) *Gene Therapy* 6:986 and WO 00/17377, the disclosures of which are incorporated herein in their entireties).

As a further alternative, the virus vectors of the invention can be produced in insect cells using baculovirus vectors to deliver the rep/cap genes and rAAV template as described by Urabe et al., (2002) *Human Gene Therapy* 13:1935-43.

Other methods of producing AAV use stably transformed packaging cells (see, e.g., U.S. Pat. No. 5,658,785).

AAV vector stocks free of contaminating helper virus may be obtained by any method known in the art. For example, AAV and helper virus may be readily differentiated based on size. AAV may also be separated away from helper virus based on affinity for a heparin substrate (Zolotukhin et al. (1999) *Gene Therapy* 6:973). Preferably, deleted replication-defective helper viruses are used so that any contaminating helper virus is not replication competent. As a further alternative, an adenovirus helper lacking late gene expression may be employed, as only adenovirus early gene expression is required to mediate packaging of AAV virus. Adenovirus mutants defective for late gene expression are known in the art (e.g., ts100K and ts149 adenovirus mutants).

The inventive packaging methods may be employed to produce high titer stocks of chimeric virus particles. Preferably, the virus stock has a titer of at least about 10^5 transducing units (tu)/ml, more preferably at least about 10^6 to/ml, more preferably at least about 10^7 tu/ml, yet more preferably at least about 10^8 to/ml, yet more preferably at least about 10^9 to/ml, still yet more preferably at least about 10^{10} tu/ml. Recombinant Parvovirus Vectors.

The parvovirus vectors of the present invention are useful for the delivery of nucleic acids to cells in vitro, ex vivo, and in vivo. In particular, the parvovirus vectors can be advantageously employed to deliver or transfer nucleic acids to animal, more preferably mammalian, cells.

Any heterologous nucleotide sequence(s) (as defined above) may be delivered in the chimeric virus vectors of the present invention. Nucleic acids of interest include nucleic acids encoding polypeptides, preferably therapeutic (e.g., for medical or veterinary uses) or immunogenic (e.g., for vaccines) polypeptides.

Therapeutic polypeptides include, but are not limited to, cystic fibrosis transmembrane regulator protein (CFTR), dystrophin (including the protein product of dystrophin minigenes, see, e.g. Vincent et al., (1993) *Nature Genetics* 5:130; U.S. Patent Publication No. 2003017131, the disclosures of

which are incorporated herein in their entireties by reference), utrophin (Tinsley et al., (1996) *Nature* 384:349), clotting factors (e.g., Factor VIII, Factor IX, Factor X, etc.), erythropoietin, angiotensin, endostatin, catalase, tyrosine hydroxylase, superoxide dismutase, leptin, the LDL receptor, neprilysin, lipoprotein lipase, ornithine transcarbamylase, β -globin, α -globin, spectrin, α_1 -antitrypsin, adenosine deaminase, hypoxanthine guanine phosphoribosyl transferase, β -glucocerebrosidase, sphingomyelinase, lysosomal hexosaminidase A, branched-chain keto acid dehydrogenase, RP65 protein, cytokines (e.g., α -interferon, β -interferon, interferon- γ , interleukins-1 through -14, granulocyte-macrophage colony stimulating factor, lymphotoxin, and the like), peptide growth factors, neurotrophic factors and hormones (e.g., somatotropin, insulin, insulin-like growth factors including IGF-1 and IGF-2, platelet derived growth factor, epidermal growth factor, fibroblast growth factor, nerve growth factor, neurotrophic factor-3 and -4, brain-derived neurotrophic factor, glial derived growth factor, transforming growth factor- α and - β , and the like), bone morphogenic proteins (including RANKL and VEGF), lysosomal proteins, anti-apoptotic gene products, glutamate receptors, lymphokines, soluble CD4, Fc receptors, T cell receptors, ApoE, ApoC, protein phosphatase inhibitor 1 (I-1), phospholamban, serca2a, lysosomal acid α -glucosidase, α -galactosidase A, barkct, β 2-adrenergic receptor, β 2-adrenergic receptor kinase (BARK), phosphoinositide-3 kinase (PI3 kinase), calsarcin, a sarcoglycan (e.g., α , β , γ), receptors (e.g., the tumor necrosis growth factor- α soluble receptor), anti-inflammatory factors such as IRAP, monoclonal antibodies (including single chain monoclonal antibodies; an exemplary Mab is the herceptin Mab). Other illustrative heterologous nucleotide sequences encode suicide gene products (e.g., thymidine kinase, cytosine deaminase, diphtheria toxin, and tumor necrosis factors such as TNF- α), proteins conferring resistance to a drug used in cancer therapy, tumor suppressor gene products (e.g., p53, Rb, Wt-1), TRAIL, FAS-ligand, and any other polypeptide that has a therapeutic effect in a subject in need thereof.

Heterologous nucleotide sequences encoding polypeptides include those encoding reporter polypeptides (e.g., an enzyme). Reporter polypeptides are known in the art and include, but are not limited to, Green Fluorescent Protein, β -galactosidase, alkaline phosphatase, luciferase, and chloramphenicol acetyltransferase gene.

Alternatively, the heterologous nucleic acid may encode an antisense nucleic acid, a ribozyme (e.g., as described in U.S. Pat. No. 5,877,022), RNAs that effect spliceosome-mediated trans-splicing (see, Puttaraju et al., (1999) *Nature Biotech.* 17:246; U.S. Pat. No. 6,013,487; U.S. Pat. No. 6,083,702), interfering RNAs (RNAi) including small interfering RNAs (siRNA) that mediate gene silencing (see, Sharp et al., (2000) *Science* 287:2431) or other non-translated RNAs, such as "guide" RNAs (Gorman et al., (1998) *Proc. Nat. Acad. Sci. USA* 95:4929; U.S. Pat. No. 5,869,248 to Yuan et al.), and the like. Exemplary untranslated RNAs include RNAi against the multiple drug resistance (MDR) gene product (e.g., to treat tumors and/or for administration to the heart to prevent damage by chemotherapy), RNAi against myostatin (Duchenne muscular dystrophy), to treat or RNAi against VEGF (e.g., to treat tumors).

The parvovirus vector may also comprise a heterologous nucleotide sequence that shares homology with and recombines with a locus on the host chromosome. This approach may be utilized to correct a genetic defect in the host cell.

The present invention also provides parvovirus vectors that express an immunogenic polypeptide, e.g., for vaccination. The nucleic acid may encode any immunogen of interest

known in the art including, but are not limited to, immunogens from human immunodeficiency virus, influenza virus, gag proteins, tumor antigens, cancer antigens, bacterial antigens, viral antigens, and the like.

The use of parvoviruses as vaccines is known in the art (see, e.g., Miyamura et al., (1994) *Proc. Nat. Acad. Sci. USA* 91:8507; U.S. Pat. No. 5,916,563 to Young et al. U.S. Pat. No. 5,905,040 to Mazzara et al., U.S. Pat. No. 5,882,652, U.S. Pat. No. 5,863,541 to Samulski et al.; the disclosures of which are incorporated herein in their entireties by reference). The antigen may be presented in the parvovirus capsid. Alternatively, the antigen may be expressed from a heterologous nucleic acid introduced into a recombinant vector genome.

An immunogenic polypeptide, or immunogen, may be any polypeptide suitable for protecting the subject against a disease, including but not limited to microbial, bacterial, protozoal, parasitic, fungal and viral diseases. For example, the immunogen may be an orthomyxovirus immunogen (e.g., an influenza virus immunogen, such as the influenza virus hemagglutinin (HA) surface protein or the influenza virus nucleoprotein gene, or an equine influenza virus immunogen), or a lentivirus immunogen (e.g., an equine infectious anemia virus immunogen, a Simian Immunodeficiency Virus (SIV) immunogen, or a Human Immunodeficiency Virus (HIV) immunogen, such as the HIV or SIV envelope GP160 protein, the HIV or SIV matrix/capsid proteins, and the HIV or SIV gag, pol and env genes products). The immunogen may also be an arenavirus immunogen (e.g., Lassa fever virus immunogen, such as the Lassa fever virus nucleocapsid protein gene and the Lassa fever envelope glycoprotein gene), a poxvirus immunogen (e.g., vaccinia, such as the vaccinia L1 or L8 genes), a flavivirus immunogen (e.g., a yellow fever virus immunogen or a Japanese encephalitis virus immunogen), a filovirus immunogen (e.g., an Ebola virus immunogen, or a Marburg virus immunogen, such as NP and GP genes), a bunyavirus immunogen (e.g., RVFV, CCHF, and SFS viruses), or a coronavirus immunogen (e.g., an infectious human coronavirus immunogen, such as the human coronavirus envelope glycoprotein gene, or a porcine transmissible gastroenteritis virus immunogen, or an avian infectious bronchitis virus immunogen). The immunogen may further be a polio immunogen, herpes immunogen (e.g., CMV, EBV, HSV immunogens) mumps immunogen, measles immunogen, rubella immunogen, diphtheria toxin or other diphtheria immunogen, pertussis antigen, hepatitis (e.g., hepatitis A, hepatitis B or hepatitis C) immunogen, or any other vaccine immunogen known in the art.

Alternatively, the immunogen may be any tumor or cancer cell antigen. Preferably, the tumor or cancer antigen is expressed on the surface of the cancer cell. Exemplary cancer and tumor cell antigens are described in S. A. Rosenberg, (1999) *Immunity* 10:281. Other illustrative cancer and tumor antigens include, but are not limited to: BRCA1 gene product, BRCA2 gene product, gp100, tyrosinase, GAGE-1/2, BAGE, RAGE, NY-ESO-1, CDK-4, β -catenin, MUM-1, Caspase-8, KIAA0205, HPVE, SART-1, PRAME, p15, melanoma tumor antigens (Kawakami et al., (1994) *Proc. Natl. Acad. Sci. USA* 91:3515; Kawakami et al., (1994) *J. Exp. Med.*, 180:347; Kawakami et al., (1994) *Cancer Res.* 54:3124) including MART-1 (Coulie et al., (1991) *J. Exp. Med.* 180:35), gp100 (Wick et al., (1988) *J. Cutan. Pathol.* 4:201) and MAGE antigen, MAGE-1, MAGE-2 and MAGE-3 (Van der Bruggen et al., (1991) *Science*, 254:1643); CEA, TRP-1, TRP-2, P-15 and tyrosinase (Brichard et al., (1993) *J. Exp. Med.* 178:489); HER-2/neu gene product (U.S. Pat. No. 4,968,603), CA 125, LK26, FB5 (endosialin), TAG 72, AFP, CA19-9, NSE, DUPAN-2, CA50, SPan-1, CA72-4, HCG, STN (sialyl Tn anti-

gen), c-erbB-2 proteins, PSA, L-CanAg, estrogen receptor, milk fat globulin, p53 tumor suppressor protein (Levine, (1993) *Ann. Rev. Biochem.* 62:623); mucin antigens (international patent publication WO 90/05142); telomerases; nuclear matrix proteins; prostatic acid phosphatase; papilloma virus antigens; and antigens associated with the following cancers: melanomas, adenocarcinoma, thymoma, lymphoma, sarcoma, lung cancer, liver cancer, colon cancer, non-Hodgkin's lymphoma, Hodgkin's lymphoma, leukemias, uterine cancer, breast cancer, prostate cancer, ovarian cancer, cervical cancer, bladder cancer, kidney cancer, pancreatic cancer and others (see, e.g., Rosenberg, (1996) *Ann. Rev. Med.* 47:481-91).

Alternatively, the heterologous nucleotide sequence may encode any polypeptide that is desirably produced in a cell in vitro, ex vivo, or in vivo. For example, the parvovirus vectors may be introduced into cultured cells and the expressed gene product isolated therefrom.

It will be understood by those skilled in the art that the heterologous nucleotide sequence(s) of interest may be operably associated with appropriate control sequences. For example, the heterologous nucleic acid may be operably associated with expression control elements, such as transcription/translation control signals, origins of replication, polyadenylation signals, internal ribosome entry sites (IRES), promoters, enhancers, and the like.

Those skilled in the art will appreciate that a variety of promoter/enhancer elements may be used depending on the level and tissue-specific expression desired. The promoter/enhancer may be constitutive or inducible, depending on the pattern of expression desired. The promoter/enhancer may be native or foreign and can be a natural or a synthetic sequence. By foreign, it is intended that the transcriptional initiation region is not found in the wild-type host into which the transcriptional initiation region is introduced.

Promoter/enhancer elements that are native to the target cell or subject to be treated are most preferred. Also preferred are promoters/enhancer elements that are native to the heterologous nucleic acid sequence. The promoter/enhancer element is chosen so that it will function in the target cell(s) of interest. Mammalian promoter/enhancer elements are also preferred. The promoter/enhancer element may be constitutive or inducible.

Inducible expression control elements are preferred in those applications in which it is desirable to provide regulation over expression of the heterologous nucleic acid sequence(s). Inducible promoters/enhancer elements for gene delivery are preferably tissue-specific promoter/enhancer elements, and include muscle specific (including cardiac, skeletal and/or smooth muscle), neural tissue specific (including brain-specific), eye (including retina-specific and cornea-specific), liver specific, bone marrow specific, pancreatic specific, spleen specific, and lung specific promoter/enhancer elements. Other inducible promoter/enhancer elements include hormone-inducible and metal-inducible elements. Exemplary inducible promoters/enhancer elements include, but are not limited to, a Tet on/off element, a RU486-inducible promoter, an ecdysone-inducible promoter, a rapamycin-inducible promoter, and a metallothionein promoter.

In embodiments wherein which the heterologous nucleic acid sequence(s) will be transcribed and then translated in the target cells, specific initiation signals are generally required for efficient translation of inserted protein coding sequences. These exogenous translational control sequences, which may include the ATG initiation codon and adjacent sequences, can be of a variety of origins, both natural and synthetic.

Gene Transfer Technology.

The parvovirus vectors according to the present invention also provide a means for delivering heterologous nucleotide sequences into a broad range of cells, including dividing and non-dividing cells. The parvovirus vectors may be employed to deliver a nucleotide sequence of interest to a cell in vitro, e.g., to produce a polypeptide in vitro or for ex vivo gene therapy. The vectors are additionally useful in a method of delivering a nucleotide sequence to a subject in need thereof, e.g., to express an immunogenic or therapeutic polypeptide. In this manner, the polypeptide may thus be produced in vivo in the subject. The subject may be in need of the polypeptide because the subject has a deficiency of the polypeptide, or because the production of the polypeptide in the subject may impart some therapeutic effect, as a method of treatment or otherwise, and as explained further below.

In general, the parvovirus vectors of the present invention may be employed to deliver any foreign nucleic acid with a biological effect to treat or ameliorate the symptoms associated with any disorder related to gene expression. Alternatively, the invention can be used to treat any disease state for which it is beneficial to deliver a therapeutic polypeptide. Illustrative disease states include, but are not limited to: cystic fibrosis (cystic fibrosis transmembrane regulator protein) and other diseases of the lung, hemophilia A (Factor VIII), hemophilia B (Factor IX), thalassemia (β -globin), anemia (erythropoietin) and other blood disorders, Alzheimer's disease (GDF; neprilysin), multiple sclerosis (β -interferon), Parkinson's disease (glial-cell line derived neurotrophic factor [GDNF]), Huntington's disease (RNAi to remove repeats), amyotrophic lateral sclerosis, epilepsy (galanin, neurotrophic factors), and other neurological disorders, cancer (endostatin, angiostatin, TRAIL, FAS-ligand, cytokines including interferons; RNAi including RNAi against VEGF or the multiple drug resistance gene product), diabetes mellitus (insulin), muscular dystrophies including Duchenne (dystrophin, mini-dystrophin, insulin-like growth factor I, a sarcoglycan [e.g., α , β , γ], RNAi against myostatin) and Becker, Gaucher disease (glucocerebrosidase), Hurler's disease (α -L-iduronidase), adenosine deaminase deficiency (adenosine deaminase), glycogen storage diseases (e.g., Fabry disease [α -galactosidase] and Pompe disease [lysosomal acid α -glucosidase]) and other metabolic defects, congenital emphysema (α 1-antitrypsin), Lesch-Nyhan Syndrome (hypoxanthine guanine phosphoribosyl transferase), Niemann-Pick disease (sphingomyelinase), Tays Sachs disease (lysosomal hexosaminidase A), Maple Syrup Urine Disease (branched-chain keto acid dehydrogenase), retinal degenerative diseases (and other diseases of the eye and retina; e.g., PDGF for macular degeneration), diseases of solid organs such as brain (including Parkinson's Disease [GDNF], astrocytomas [endostatin, angiostatin and/or RNAi against VEGF], glioblastomas [endostatin, angiostatin and/or RNAi against VEGF]), liver, kidney, heart including congestive heart failure or peripheral artery disease (PAD) (e.g., by delivering protein phosphatase inhibitor I (I-1), phospholamban, serca2a, zinc finger proteins that regulate the phospholamban gene, Barkct, β 2-adrenergic receptor, β 2-adrenergic receptor kinase (BARK), phosphoinositide-3 kinase (PI3 kinase), calsarcin, etc.), arthritis (insulin-like growth factors), joint disorders (insulin-like growth factors), intimal hyperplasia (e.g., by delivering enos, inos), improve survival of heart transplants (superoxide dismutase), AIDS (soluble CD4), muscle wasting (insulin-like growth factor I), kidney deficiency (erythropoietin), anemia (erythropoietin), arthritis (anti-inflammatory factors such as IRAP and TNF α soluble receptor), hepatitis (α -interferon), LDL receptor deficiency (LDL

receptor), hyperammonemia (ornithine transcarbamylase), Krabbe's disease (galactocerebrosidase), Batten's disease, spinal cerebral ataxias including SCA1, SCA2 and SCA3, phenylketonuria (phenylalanine hydroxylase), autoimmune diseases, and the like. The invention can further be used following organ transplantation to increase the success of the transplant and/or to reduce the negative side effects of organ transplantation or adjunct therapies (e.g., by administering immunosuppressant agents or inhibitory nucleic acids to block cytokine production). As another example, bone morphogenic proteins (including RANKL and/or VEGF) can be administered with a bone allograft, for example, following a break or surgical removal in a cancer patient.

Alternatively, a gene transfer vector may be administered that encodes any other therapeutic polypeptide.

In particular embodiments, an AAV2-derived and/or AAV3b derived vector comprising the 263 and/or 265 mutations according to the present invention is used to deliver a nucleic acid of interest as described herein to a tumor or to skeletal muscle, cardiac muscle, astrocytes and/or glial cells, for example, to treat a disorder associated with any of these cells or tissues such as Parkinson's disease, astrocytomas, glioblastomas, muscular dystrophy, heart disease (including PAD and congestive heart failure), and the like.

Gene transfer has substantial potential use in understanding and providing therapy for disease states. There are a number of inherited diseases in which defective genes are known and have been cloned. In general, the above disease states fall into two classes: deficiency states, usually of enzymes, which are generally inherited in a recessive manner, and unbalanced states, which may involve regulatory or structural proteins, and which are typically inherited in a dominant manner. For deficiency state diseases, gene transfer could be used to bring a normal gene into affected tissues for replacement therapy, as well as to create animal models for the disease using antisense mutations. For unbalanced disease states, gene transfer could be used to create a disease state in a model system, which could then be used in efforts to counteract the disease state. Thus parvovirus vectors produced according to the methods of the present invention permit the treatment of genetic diseases. As used herein, a disease state is treated by partially or wholly remedying the deficiency or imbalance that causes the disease or makes it more severe. The use of site-specific recombination of nucleic sequences to cause mutations or to correct defects is also possible.

The parvovirus vectors according to the present invention may also be employed to provide an antisense nucleic acid or RNAi to a cell in vitro or in vivo. Expression of the antisense nucleic acid or RNAi in the target cell diminishes expression of a particular protein by the cell. Accordingly, antisense nucleic acids or RNAi may be administered to decrease expression of a particular protein in a subject in need thereof. Antisense nucleic acids or RNAi may also be administered to cells in vitro to regulate cell physiology, e.g., to optimize cell or tissue culture systems.

Further, the parvovirus vectors according to the instant invention find further use in diagnostic and screening methods, whereby a gene of interest is transiently or stably expressed in a cell culture system, or alternatively, a transgenic animal model. The invention can also be practiced to deliver a nucleic acid for the purposes of protein production, e.g., for laboratory, industrial or commercial purposes. Delivery of Immunogenic Polypeptides.

As a further aspect, parvovirus vectors of the present invention may be used to produce an immune response in a subject. According to this embodiment, a parvovirus vector comprising a nucleotide sequence encoding an immunogen may be

administered to a subject, and an active immune response is mounted by the subject against the immunogen. Immunogens are as described hereinabove. Preferably, a protective immune response is elicited.

Alternatively, the parvovirus vector may be administered to a cell ex vivo and the altered cell is administered to the subject. The heterologous nucleotide sequence is introduced into the cell, and the cell is administered to the subject, where the heterologous nucleotide sequence encoding the immunogen is preferably expressed and induces an immune response in the subject against the immunogen. In particular embodiments, the cell is an antigen presenting cell (e.g., a dendritic cell).

An "active immune response" or "active immunity" is characterized by "participation of host tissues and cells after an encounter with the immunogen. It involves differentiation and proliferation of immunocompetent cells in lymphoreticular tissues, which lead to synthesis of antibody or the development of cell-mediated reactivity, or both." Herbert B. Herscovitz, *Immunophysiology: Cell Function and Cellular Interactions in Antibody Formation*, in *IMMUNOLOGY: BASIC PROCESSES* 117 (Joseph A. Bellanti ed., 1985). Alternatively stated, an active immune response is mounted by the host after exposure to immunogens by infection or by vaccination. Active immunity can be contrasted with passive immunity, which is acquired through the "transfer of preformed substances (antibody, transfer factor, thymic graft, interleukin-2) from an actively immunized host to a non-immune host." Id.

A "protective" immune response or "protective" immunity as used herein indicates that the immune response confers some benefit to the subject in that it prevents or reduces the incidence of disease. Alternatively, a protective immune response or protective immunity may be useful in the treatment of disease, in particular cancer or tumors (e.g., by causing regression of a cancer or tumor and/or by preventing metastasis and/or by preventing growth of metastatic nodules). The protective effects may be complete or partial, as long as the benefits of the treatment outweigh any disadvantages thereof.

According to the foregoing methods of inducing an immune response in a subject, it is preferred that the parvovirus vector carrying the heterologous nucleotide sequence is administered in an immunogenically effective amount, as described herein.

The parvovirus vectors of the present invention may also be administered for cancer immunotherapy by administration of a parvovirus vector expressing cancer cell antigens (or an immunologically similar molecule) or any other immunogen that produces an immune response against a cancer cell. To illustrate, an immune response may be produced against a cancer cell antigen in a subject by administering a parvovirus vector comprising a heterologous nucleotide sequence encoding the cancer cell antigen, for example to treat a patient with cancer. The parvovirus vector may be administered to a subject in vivo or by using ex vivo methods, as described herein.

As used herein, the term "cancer" encompasses tumor-forming cancers. Likewise, the term "cancerous tissue" encompasses tumors. A "cancer cell antigen" encompasses tumor antigens.

The term "cancer" has its understood meaning in the art, for example, an uncontrolled growth of tissue that has the potential to spread to distant sites of the body (i.e., metastasize). Exemplary cancers include, but are not limited to, leukemias, lymphomas, colon cancer, renal cancer, liver cancer, breast cancer, lung cancer, prostate cancer, ovarian cancer,

melanoma, and the like. Preferred are methods of treating and preventing tumor-forming cancers.

The term "tumor" is also understood in the art, for example, as an abnormal mass of undifferentiated cells within a multicellular organism. Tumors can be malignant or benign. Preferably, the methods disclosed herein are used to prevent and

treat malignant tumors.

Cancer cell antigens according to the present invention have been described hereinabove. By the terms "treating cancer" or "treatment of cancer," it is intended that the severity of the cancer is reduced or the cancer is prevented or at least partially eliminated. Preferably, these terms indicate that metastasis of the cancer is prevented or reduced or at least partially eliminated. It is further preferred that these terms indicate that growth of metastatic nodules (e.g., after surgical removal of a primary tumor) is prevented or reduced or at least partially eliminated. By the terms "prevention of cancer" or "preventing cancer" it is intended that the methods at least partially eliminate or reduce the incidence or onset of cancer. Alternatively stated, the onset of cancer in the subject may be slowed, controlled, decreased in likelihood or probability, or delayed.

In particular embodiments, cells may be removed from a subject with cancer and contacted with parvovirus particles according to the instant invention. The modified cell is then administered to the subject, whereby an immune response against the cancer cell antigen is elicited. This method is particularly advantageously employed with immunocompromised subjects that cannot mount a sufficient immune response in vivo (i.e., cannot produce enhancing antibodies in sufficient quantities).

It is known in the art that immune responses may be enhanced by immunomodulatory cytokines (e.g., α -interferon, β -interferon, γ -interferon, ω -interferon, τ -interferon, interleukin-1 α , interleukin-1 β , interleukin-2, interleukin-3, interleukin-4, interleukin 5, interleukin-6, interleukin-7, interleukin-8, interleukin-9, interleukin-10, interleukin-11, interleukin 12, interleukin-13, interleukin-14, interleukin-18, B cell Growth factor, CD40 Ligand, tumor necrosis factor- α , tumor necrosis factor- β , monocyte chemoattractant protein-1, granulocyte-macrophage colony stimulating factor, and lymphotoxin). Accordingly, immunomodulatory cytokines (preferably, CTL inductive cytokines) may be administered to a subject in conjunction with the parvovirus vectors.

Cytokines may be administered by any method known in the art. Exogenous cytokines may be administered to the subject, or alternatively, a nucleotide sequence encoding a cytokine may be delivered to the subject using a suitable vector, and the cytokine produced in vivo.

Subjects, Pharmaceutical Formulations, and Modes of Administration.

Parvovirus vectors according to the present invention find use in both veterinary and medical applications. Suitable subjects include both avians and mammals. The term "avian" as used herein includes, but is not limited to, chickens, ducks, geese, quail, turkeys, pheasant, parrots, parakeets. The term "mammal" as used herein includes, but is not limited to, humans, non-human primates, bovines, ovines, caprines, equines, felines, canines, lagomorphs, etc. Human subjects include neonates, infants, juveniles, and adults. In particular embodiments, the subject has antibodies against one or more AAV (e.g., AAV serotypes such as AAV1 and/or AAV2). In other embodiments, the subject has previously been administered a different AAV vector (e.g., an immunologically distinct AAV). Optionally, the subject is "in need of" the methods of the present invention, e.g., because the subject has or is believed at risk for a disorder including those described

herein or that would benefit from the delivery of a nucleic acid including those described herein.

In particular embodiments, the present invention provides a pharmaceutical composition comprising a virus particle or virus capsid of the invention in a pharmaceutically acceptable carrier and, optionally, other medicinal agents, pharmaceutical agents, stabilizing agents, buffers, carriers, adjuvants, diluents, etc. For injection, the carrier will typically be a liquid. For other methods of administration, the carrier may be either solid or liquid. For inhalation administration, the carrier will be respirable, and will preferably be in solid or liquid particulate form.

By "pharmaceutically acceptable" it is meant a material that is not toxic or otherwise undesirable, i.e., the material may be administered to a subject without causing any undesirable biological effects.

One aspect of the present invention is a method of transferring a nucleotide sequence to a cell in vitro. The virus particles may be introduced to the cells at the appropriate multiplicity of infection according to standard transduction methods appropriate for the particular target cells. Titers of virus to administer can vary, depending upon the target cell type and number, and the particular virus vector, and can be determined by those of skill in the art without undue experimentation. Preferably, at least about 10^3 infectious units, more preferably at least about 10^5 infectious units are introduced to the cell.

The cell(s) to be introduced the parvovirus vector may be of any type, including but not limited to neural cells (including cells of the peripheral and central nervous systems, in particular, brain cells such as neurons, oligodendrocytes, glial cells, astrocytes), lung cells, cells of the eye (including retinal cells, retinal pigment epithelium, and corneal cells), epithelial cells (e.g., gut and respiratory epithelial cells), muscle cells, dendritic cells, pancreatic cells (including islet cells), hepatic cells, myocardial cells, bone cells (e.g., bone marrow stem cells), hematopoietic stem cells, spleen cells, keratinocytes, fibroblasts, endothelial cells, prostate cells, germ cells, and the like. Alternatively, the cell may be any progenitor cell. As a further alternative, the cell can be a stem cell (e.g., neural stem cell, liver stem cell). As still a further alternative, the cell may be a cancer or tumor cell. Moreover, the cells can be from any species of origin, as indicated above.

The parvovirus vectors may be introduced to cells in vitro for the purpose of administering the modified cell to a subject. In particular embodiments, the cells have been removed from a subject, the parvovirus vector is introduced therein, and the cells are then replaced back into the subject. Methods of removing cells from subject for treatment ex vivo, followed by introduction back into the subject are known in the art (see, e.g., U.S. Pat. No. 5,399,346; the disclosure of which is incorporated herein in its entirety). Alternatively, the recombinant parvovirus vector is introduced into cells from another subject, into cultured cells, or into cells from any other suitable source, and the cells are administered to a subject in need thereof.

Suitable cells for ex vivo gene therapy are as described above. Dosages of the cells to administer to a subject will vary upon the age, condition and species of the subject, the type of cell, the nucleic acid being expressed by the cell, the mode of administration, and the like. Typically, at least about 10^2 to about 10^8 or about 10^3 to about 10^6 cells will be administered per dose in a pharmaceutically acceptable carrier. In particular embodiments, the cells transduced with the parvovirus vector are administered to the subject in a therapeutically effective amount in combination with a pharmaceutical carrier.

31

A "therapeutically effective" amount as used herein is an amount that is sufficient to provide some improvement or benefit to the subject. Alternatively stated, a "therapeutically effective" amount is an amount that will provide some alleviation, mitigation, or decrease in at least one clinical symptom in the subject. Those skilled in the art will appreciate that the therapeutic effects need not be complete or curative, as long as some benefit is provided to the subject.

In some embodiments, cells that have been transduced with a parvovirus vector may be administered to elicit an immunogenic response against the delivered polypeptide (e.g., expressed as a transgene or in the capsid). Typically, a quantity of cells expressing an immunogenically effective amount of the polypeptide in combination with a pharmaceutically acceptable carrier is administered. An "immunogenically effective amount" is an amount that is sufficient to evoke an active immune response in the subject to which the pharmaceutical formulation is administered. Preferably, the dosage is sufficient to produce a protective immune response (as defined above). The degree of protection conferred need not be complete or permanent, as long as the benefits of administering the immunogenic polypeptide outweigh any disadvantages thereof.

A further aspect of the invention is a method of administering the parvovirus particles or capsids of the invention to subjects. Administration of the parvovirus particles or capsids of the present invention to a human subject or an animal in need thereof can be by any means known in the art for administering virus vectors. Preferably, the parvovirus vector is delivered in a therapeutically effective dose in a pharmaceutically acceptable carrier.

The parvovirus vectors of the invention can further be administered to elicit an immunogenic response (e.g., as a vaccine). Typically, vaccines of the present invention comprise an immunogenically effective amount of virus in combination with a pharmaceutically acceptable carrier. Preferably, the dosage is sufficient to produce a protective immune response (as defined above). The degree of protection conferred need not be complete or permanent, as long as the benefits of administering the immunogenic polypeptide outweigh any disadvantages thereof. Subjects and immunogens are as described above.

Dosages of the parvovirus particles to be administered to a subject will depend upon the mode of administration, the disease or condition to be treated, the individual subject's condition, the particular virus vector, and the nucleic acid to be delivered, and can be determined in a routine manner. Exemplary doses for achieving therapeutic effects are virus titers of at least about 10^5 , 10^6 , 10^7 , 10^8 , 10^9 , 10^{10} , 10^{11} , 10^{12} , 10^{13} , 10^{14} , 10^{15} transducing units or more, preferably about 10^8 - 10^{13} transducing units, yet more preferably 10^{12} transducing units.

In particular embodiments, more than one administration (e.g., two, three, four or more administrations) may be employed to achieve the desired level of gene expression over a period of various intervals, e.g., daily, weekly, monthly, yearly, etc.

Exemplary modes of administration include oral, rectal, transmucosal, topical, intranasal, inhalation (e.g., via an aerosol), buccal (e.g., sublingual), vaginal, intrathecal, intraocular, transdermal, in utero (or in ovo), parenteral (e.g., intravenous, subcutaneous, intradermal, intramuscular [including administration to skeletal, diaphragm and/or cardiac muscle], intradermal, intrapleural, intracerebral, and intraarticular), topical (e.g., to both skin and mucosal surfaces, including airway surfaces, and transdermal administration, and the like, as well as direct tissue or organ injection (e.g., to liver, skel-

32

etal muscle, cardiac muscle, diaphragm muscle or brain). Administration can also be to a tumor (e.g., in or a near a tumor or a lymph node). The most suitable route in any given case will depend on the nature and severity of the condition being treated and on the nature of the particular vector that is being used.

In particular embodiments, a chimeric vector according to the present invention that comprises a 263 and/or 265 mutation as described herein in an AAV2 or AAV3b backbone is administered to skeletal muscle, cardiac muscle and/or brain (e.g., to treat muscular dystrophy, heart disease (e.g., PAD or congestive heart failure), Parkinson's disease, astrocytomas, glioblastomas) or to a tumor.

Injectables can be prepared in conventional forms, either as liquid solutions or suspensions, solid forms suitable for solution or suspension in liquid prior to injection, or as emulsions. Alternatively, one may administer the virus in a local rather than systemic manner, for example, in a depot or sustained-release formulation. Further, the virus particle can be delivered dried to a surgically implantable matrix such as a bone graft substitute, a suture, a stent, and the like (e.g., as described in U.S. Patent Publication No. US-2004-0013645-A1).

Pharmaceutical compositions suitable for oral administration can be presented in discrete units, such as capsules, cachets, lozenges, or tablets, each containing a predetermined amount of the composition of this invention; as a powder or granules; as a solution or a suspension in an aqueous or non-aqueous liquid; or as an oil-in-water or water-in-oil emulsion. Oral delivery can be performed by complexing a composition of the present invention to a carrier capable of withstanding degradation by digestive enzymes in the gut of an animal. Examples of such carriers include plastic capsules or tablets, as known in the art. Such formulations are prepared by any suitable method of pharmacy, which includes the step of bringing into association the composition and a suitable carrier (which may contain one or more accessory ingredients as noted above). In general, the pharmaceutical composition according to embodiments of the present invention are prepared by uniformly and intimately admixing the composition with a liquid or finely divided solid carrier, or both, and then, if necessary, shaping the resulting mixture. For example, a tablet can be prepared by compressing or molding a powder or granules containing the composition, optionally with one or more accessory ingredients. Compressed tablets are prepared by compressing, in a suitable machine, the composition in a free-flowing form, such as a powder or granules optionally mixed with a binder, lubricant, inert diluent, and/or surface active/dispersing agent(s). Molded tablets are made by molding, in a suitable machine, the powdered compound moistened with an inert liquid binder.

Pharmaceutical compositions suitable for buccal (sub-lingual) administration include lozenges comprising the composition of this invention in a flavored base, usually sucrose and acacia or tragacanth; and pastilles comprising the composition in an inert base such as gelatin and glycerin or sucrose and acacia.

Pharmaceutical compositions of this invention suitable for parenteral administration can comprise sterile aqueous and non-aqueous injection solutions of the composition of this invention, which preparations are preferably isotonic with the blood of the intended recipient. These preparations can contain anti-oxidants, buffers, bacteriostats and solutes, which render the composition isotonic with the blood of the intended recipient. Aqueous and non-aqueous sterile suspensions, solutions and emulsions can include suspending agents and thickening agents. Examples of non-aqueous solvents are

33

propylene glycol, polyethylene glycol, vegetable oils such as olive oil, and injectable organic esters such as ethyl oleate. Aqueous carriers include water, alcoholic/aqueous solutions, emulsions or suspensions, including saline and buffered media. Parenteral vehicles include sodium chloride solution, Ringer's dextrose, dextrose and sodium chloride, lactated Ringer's, or fixed oils. Intravenous vehicles include fluid and nutrient replenishers, electrolyte replenishers (such as those based on Ringer's dextrose), and the like. Preservatives and other additives may also be present such as, for example, antimicrobials, anti-oxidants, chelating agents, and inert gases and the like.

The compositions can be presented in unit/dose or multi-dose containers, for example, in sealed ampoules and vials, and can be stored in a freeze-dried (lyophilized) condition requiring only the addition of the sterile liquid carrier, for example, saline or water-for-injection immediately prior to use.

Extemporaneous injection solutions and suspensions can be prepared from sterile powders, granules and tablets of the kind previously described. For example, an injectable, stable, sterile composition of this invention in a unit dosage form in a sealed container can be provided. The composition can be provided in the form of a lyophilizate, which can be reconstituted with a suitable pharmaceutically acceptable carrier to form a liquid composition suitable for injection into a subject. The unit dosage form can be from about 1 µg to about 10 grams of the composition of this invention. When the composition is substantially water-insoluble, a sufficient amount of emulsifying agent, which is physiologically acceptable, can be included in sufficient quantity to emulsify the composition in an aqueous carrier. One such useful emulsifying agent is phosphatidyl choline.

Pharmaceutical compositions suitable for rectal administration are preferably presented as unit dose suppositories. These can be prepared by admixing the composition with one or more conventional solid carriers, such as for example, cocoa butter and then shaping the resulting mixture.

Pharmaceutical compositions of this invention suitable for topical application to the skin preferably take the form of an ointment, cream, lotion, paste, gel, spray, aerosol, or oil. Carriers that can be used include, but are not limited to, petroleum jelly, lanoline, polyethylene glycols, alcohols, transdermal enhancers, and combinations of two or more thereof. In some embodiments, for example, topical delivery can be performed by mixing a pharmaceutical composition of the present invention with a lipophilic reagent (e.g., DMSO) that is capable of passing into the skin.

Pharmaceutical compositions suitable for transdermal administration can be in the form of discrete patches adapted to remain in intimate contact with the epidermis of the subject for a prolonged period of time. Compositions suitable for transdermal administration can also be delivered by iontophoresis (see, for example, *Pharmaceutical Research* 3:318 (1986)) and typically take the form of an optionally buffered aqueous solution of the composition of this invention. Suitable formulations can comprise citrate or bis/tris buffer (pH 6) or ethanol/water and can contain from 0.1 to 0.2M active ingredient.

The parvovirus vectors disclosed herein may be administered to the lungs of a subject by any suitable means, but are preferably administered by administering an aerosol suspension of respirable particles comprised of the parvovirus vectors, which the subject inhales. The respirable particles may be liquid or solid. Aerosols of liquid particles comprising the parvovirus vectors may be produced by any suitable means, such as with a pressure-driven aerosol nebulizer or an ultra-

34

sonic nebulizer, as is known to those of skill in the art. See, e.g., U.S. Pat. No. 4,501,729. Aerosols of solid particles comprising the parvovirus vectors may likewise be produced with any solid particulate medicament aerosol generator, by techniques known in the pharmaceutical art.

Having described the present invention, the same will be explained in greater detail in the following examples, which are included herein for illustration purposes only, and which are not intended to be limiting to the invention.

EXAMPLE 1

Triage Approach

Materials and Methods

Plasmid and DNA Mutagenesis

Starting plasmids for these experiments were pxx1, pxx2, pxx3 (Rabinowitz et al, (2001) *J. Virology* 76:781-801). All plasmid mutagenesis was performed using either the Quik Change Multi Site Mutagenesis or the Quik Change Site Directed Mutagenesis kits (both from Stratagene). Accuracy of the nucleotide changes was verified by DNA sequence analyses followed by DNA subcloning to eliminate any unwanted artifacts generated by the mutagenesis approach. In certain instances, subcloning was performed using the newly generated mutant to generate another mutant.

Recombinant Virus Production

All recombinant AAV (rAAV) viruses were generated using the standard triple transfection method (Haberman et al. 1999. Production of recombinant adeno-associated viral vectors and use in in vitro and in vivo administration, p. 4.17.1-4.17.25. In J. Crawley, et al. (eds.), *Current protocols in neuroscience*, vol. 1. John Wiley & Sons, Inc., New York, N.Y.). rAAV was purified in this study using standard cesium chloride gradients or using heparin column chromatography (Id.). The method used for virus purification is indicated in the corresponding figure. The physical titer of the different viral preparations was evaluated using dot blot hybridization. The preparations being compared in one experiment were evaluated on the same dot blots to control for variation between different dot blots.

Animal Imaging

1×10^{10} viral genome containing particles (vg) were injected into the gastrocnemius of 6-week-old male BALB/c mice. A total of 6 limbs were injected for each virus type using 25 µl of virus. Animals were imaged at different days post injection using the Roper Scientific Imaging (Princeton Instruments). Briefly, animals were anesthetized and injected IP with luciferin substrate. Ten minutes post-injection the animals were placed in the chamber and light emission was determined. The average number of total pixels per region of interest was determined using the CMIR_Image software (Center for Molecular Imaging Research, Mass. General) and plotted over time.

Heparin Binding Experiments

Batch binding of rAAV to heparin agarose was performed as described previously (Rabinowitz, (2004) *J. Virology* 78:4421-4432). Briefly, equivalent particles of rAAV virions were applied to heparin agarose type 1 (H-6508, Sigma, St. Louis, Mo.) in 1xPBS, allowed to bind for one hour at room temperature, centrifuged at low speed for 2 minutes, and supernatant (flow through) was removed. Six washes of five bed-volumes of PBS 1 mM MgCl₂ were performed, followed by a three-step elution of five bed-volumes of PBS 1 mM MgCl₂ containing 0.5 M NaCl (step 1), 1.0 M NaCl (step 2), or 1.5 M NaCl (step 3). The number of rAAV particles present in the washes and the 3-step elution was determined by dot

blot hybridization.

Results

Rationale of the Triage Approach

It is clear in the literature that different serotypes of AAV exhibit differing abilities to transduce different tissues. One well-documented example is the ability of AAV1 to preferentially transduce skeletal muscle. The amino acid sequences of AAV1 and AAV2 are 83% identical. One previous study exchanged large regions of the AAV2 capsid with the corresponding amino acids of AAV1 and examined the ability of the resultant vectors to transduce skeletal muscle (Hauck et al., (2003) *J. Virology* 77:2768-2774). They identified 2 amino acids that appeared to confer enhanced skeletal muscle transduction by AAV2, albeit not as well as AAV1. The present studies examined the phenomenon in a more detailed manner. Employing bio-informatic and structure-function analysis, key amino acids residues in the AAV capsid proteins were identified that account for enhanced AAV1 transduction for testing in the context of majority AAV2 capsid. This approach avoids the need to make all the different changes of amino acids between the 2 serotypes and all of the different permutations. Further, this approach took advantage of the crystal structure of AAV2, amino acid sequence alignments, and known properties of the various AAV serotypes. Alignments were generated of these very similar capsid proteins from the different serotypes. Amino acids in common between AAV1 and AAV7 capsid proteins and distinct from AAV2 and AAV8 capsid proteins were identified. Thirty-six amino acid candidates were identified via this triage approach (FIG. 1A). Based on modeling of the crystal structure, the 12 best candidates were chosen to test for enhanced muscle transduction (FIG. 1B).

Some AAV Variants Exhibited No Difference from AAV2 while Others are Much Better than AAV2.

The different AAV variants were evaluated for their ability to transduce skeletal muscle following injection of 1×10^{10} genome containing viral particles into the gastrocnemius muscle of BALB/c mice. Some virus variants did not exhibit any difference in the capability to transduce skeletal muscle (FIGS. 2a and b). Although the 324, 328 positions are located on loops predicted to lie on the capsid surface, changing of these amino acids to the corresponding amino acids in AAV1 did nothing to improve the capability of this virus to transduce skeletal muscle (FIGS. 2A and 2B).

One variant (2.5) exhibited much higher muscle transduction than AAV2 (FIG. 3A). This variant had 5 amino acid changes that were located in the 2-fold axis of symmetry (FIG. 3B). Although it appears that groups of these 5 amino acids lie on different parts of the same subunit (2 on one side and 3 on another), they are in very close proximity when 2 of the subunits come together (FIG. 3C).

To investigate whether all 5 amino acids were necessary for this enhanced muscle tropism we made the 263, 265 variants as well as the 709, 712, 720 variants. These variants were tested in similar imaging experiments (FIG. 4). The 709, 712, 720 variant exhibited a muscle transduction similar to AAV2, while the 263, 265 variant exhibited an enhanced muscle transduction similar to what was observed with the 2.5 variant containing all 5 amino acid changes. A 263 and a separate 265 variant were constructed and tested in similar experiments (FIG. 5). These experiments revealed that the amino acid mostly responsible for enhanced AAV1 transduction is located at the 265 position. This is essentially an insertion of a threonine in the AAV2 capsid sequence.

The same insertion has been made at positions 263 and 265 in the AAV3b backbone (FIG. 10).

Interestingly, the 263, 265 region is divergent amongst the serotypes (FIG. 7 boxed region). There are additional amino acid insertions in this area compared to AAV2.

Another variant that exhibits higher muscle transduction than AAV2 is the 454, 461 variant (FIG. 6). Different combinations of 265 with the 454, 461 changes are assessed to determine whether further enhancement can be observed by combining these variants.

Finally, we examined the ability of the different variants to be purified by heparin (FIG. 8). All of the variants tested exhibited heparin binding profiles similar to AAV2.

EXAMPLE 2

Immunological Profile

Similar to other non-enveloped viruses, high doses of AAV generate neutralizing antibody that prevents repeated dosing. With the advent of new serotypes, repeat administration is possible. To explore the ability to avoid a pre-existing immune response to AAV2, the chimeric 2.5 vector was tested for transgene expression in vitro after exposure to serum from animals pre-exposed to different AAV serotypes (1, 2, and 2.5 respectively).

To generate animals with a robust immune response to AAV virion shell, 4×10^{10} particles of AAV serotype 1, 2, and 2.5 vector were independently injected intramuscularly in C57BL/6 mice. Four weeks post-injection, blood was isolated and serum collected. Serum from these animals was then used in a neutralizing antibody assay using 293 cells and AAV specific serotype vectors expressing GFP as a reporter gene. In this assay, serum is sequentially diluted and then mixed with a known amount of serotype specific vector (1×10^8 particles) at 4° C. for 2 hr. This mixture of serum and vector is then added to 293 cells in 24-well plates in the presence of adenovirus helper virus at a multiplicity of infection of 5. Under these conditions, green fluorescent protein (GFP) expression is a measure of serotype-specific vector ability to infect cells in the presence of neutralizing antibodies. The neutralizing antibody titer is then calculated as the highest dilution where GFP expression is 50% or less than control vector (without pre-mixture with serotype specific serum).

As seen in Table 3, animals pre-exposed to AAV1 could neutralize AAV 1 GFP transduction with dilutions as high as 1:1000. However this serotype 1 specific neutralizing antibody required more mouse serum to neutralize AAV chimeric 2.5 (1:100 dilution). More importantly, this observation was true for mouse sera obtained from animals pre-exposed to AAV serotype 2 virion shells. In this assay, only after sera were diluted 1:10,000 was observed 50% GFP transduction observed when compared to AAV2 control. However, for chimeric 2.5, 50% GFP transduction was observed with only 1:100 dilution of this mouse serum. Since only 0.6% of the amino acid changes differ from AAV2 in this chimeric vector, these alterations had profound effects on the ability of pre-existing AAV2 neutralizing antibody to recognize the AAV 2.5 capsid shell. Animals pre-exposed to 2.5 and then assayed for neutralizing activity against AAV 1, 2, and 2.5 yielded expected results (see Table 3) with highest dilution required for the 2.5 vector (1:8000) followed by 1:1000 for AAV 2 and 1:100 for AAV serotype 1 respectively. The conclusion from these studies is that the 5 amino acid alteration in chimeric 2.5 although small in number (0.6% total amino acids) was sufficient to significantly affect the immune profile for this virion when challenged with neutralizing antibodies specific for AAV2.

Based on these studies, the 2.5 vector would be suitable for transducing individuals pre-exposed to AAV1, AAV2 or both, thereby providing greater versatility in available vectors. For example, this chimeric vector would allow for re-administration in animals and patients pre-exposed to AAV2. In addition, this chimeric vector demonstrates that selected amino acids can be changed in the AAV2, amino acid capsid sequence that significantly alter immune response. Based on these results, similar alterations at similar capsid coding regions for other AAV serotypes would also result in unique immune profiles allowing one to easily generate a large number of alternative AAV variants for repeat administration without having to change to a new serotype capsid each time.

TABLE 3

The Neutralizing Antibody Assay for AAV2.5 virus		
Sera	Target virus	NA titer
AAV1	AAV1	1:1000
	AAV2.5	1:100
AAV2	AAV2	1:10000
	AAV2.5	1:100
AAV2.5	AAV1	1:100
	AAV2	1:1000
	AAV2.5	1:8000

EXAMPLE 3

263, 265 Mutations in an AAV3b Backbone
(SASTG) Confers Enhanced Skeletal Muscle
Transduction

These amino acids (263, 265) from AAV1 were also engineered into the corresponding positions of the AAV3b capsid, a serotype closely related to AAV2 and one that also does not transduce skeletal muscle well, to generate a vector designated as SASTG. Mice were injected with equivalent genome containing particles (1×10^{10}) of AAV1, AAV2, AAV3b, or SASTG and were assessed for luciferase activity over time (days 3, 7, 14, and 21) using in vivo live animal imaging (FIG. 12).

As expected, both AAV2 and AAV3 do not transduce skeletal muscle as efficiently as AAV1. Expression from the AAV2 and AAV3b injected mice could not be detected at day 3; whereas expression from AAV1 and, importantly, from SASTG could be detected at day 3. The expression pattern of SASTG paralleled that of AAV1 at all time points post injection rather than its AAV3b parent.

EXAMPLE 4

Transduction of Brain and Liver

Six- to eight-week-old male C57bl/6 mice were utilized to determine efficiency of AAV2 and the 2.5 vector transduction in liver. The mice were anesthetized using 300 μ L 2.5% Avertin, and 1×10^{11} particles of AAV2 and 2.5 vector carrying the human Factor IX (hFIX) transgene virus were dissolved in 250 μ L PBS and injected slowly through the portal vein. The vectors were duplexed virus particles as described in international patent publication WO 01/92551. After 1 and 6 weeks, 100 μ L of blood from each mouse was collected from the tail vein using heparin-coated capillary glass tubes. Serum was collected by centrifuging the blood sample at 4° C., 8000 rpm for 20 min. Sera were stored at -80° C. until tested. Expression of hFIX in the serum was tested by standard ELISA methods. Serial dilutions of normal human serum with hFIX levels of 5 μ g/mL were used as a standard. Using this assay, it was found that the 2.5 vector has a reduced ability to transduce liver as compared with the parent AAV2 virus (FIG. 13). This experiment demonstrates that in addition to gaining muscle tropism, the 2.5 vector has lost liver specific tropism characteristic of AAV2.

In another experiment, the duplexed 2.5 vector was injected into the cortex region of the mouse brain (FIG. 14) using a green fluorescent protein (GFP) reporter transgene cassette and conditions previously established for AAV 2 and assayed for neuron specific transduction. It is well-established that AAV1 and AAV2 are specific for neuronal transduction. As shown in FIG. 14, the 2.5 vector transduces neurons (arrow on the left) as well as non-neuronal cells (arrow on right). Based on morphology, these cells appear to be astrocytes.

The sum of these experiments when testing the 2.5 vector for tissue-specific transduction in vivo demonstrates that in addition to gaining tissue-specific tropism (e.g., muscle, skeletal or cardiac derived from the AAV serotype 1 parent), and losing cell type specific transduction (e.g., liver-hepatocyte-specific transduction of recipient AAV2), these vectors have gained a new tropism (non-neuronal/astrocytes) that is not present in either the donor parent (AAV1) or recipient parent capsid (AAV2) and is totally unique to the chimeric 2.5 vector.

The foregoing is illustrative of the present invention, and is not to be construed as limiting thereof. The invention is defined by the following claims, with equivalents of the claims to be included therein.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 16

<210> SEQ ID NO 1

<211> LENGTH: 736

<212> TYPE: PRT

<213> ORGANISM: adeno-associated virus 1

<400> SEQUENCE: 1

Met Ala Ala Asp Gly Tyr Leu Pro Asp Trp Leu Glu Asp Asn Leu Ser
1 5 10 15

Glu Gly Ile Arg Glu Trp Trp Asp Leu Lys Pro Gly Ala Pro Lys Pro
20 25 30

Lys Ala Asn Gln Gln Lys Gln Asp Asp Gly Arg Gly Leu Val Leu Pro

-continued

35					40					45					
Gly	Tyr	Lys	Tyr	Leu	Gly	Pro	Phe	Asn	Gly	Leu	Asp	Lys	Gly	Glu	Pro
50					55					60					
Val	Asn	Ala	Ala	Asp	Ala	Ala	Ala	Leu	Glu	His	Asp	Lys	Ala	Tyr	Asp
65					70					75					80
Gln	Gln	Leu	Lys	Ala	Gly	Asp	Asn	Pro	Tyr	Leu	Arg	Tyr	Asn	His	Ala
				85					90					95	
Asp	Ala	Glu	Phe	Gln	Glu	Arg	Leu	Gln	Glu	Asp	Thr	Ser	Phe	Gly	Gly
			100					105					110		
Asn	Leu	Gly	Arg	Ala	Val	Phe	Gln	Ala	Lys	Lys	Arg	Val	Leu	Glu	Pro
		115						120					125		
Leu	Gly	Leu	Val	Glu	Glu	Gly	Ala	Lys	Thr	Ala	Pro	Gly	Lys	Lys	Arg
		130						135					140		
Pro	Val	Glu	Gln	Ser	Pro	Gln	Glu	Pro	Asp	Ser	Ser	Ser	Gly	Ile	Gly
145									150				155		160
Lys	Thr	Gly	Gln	Gln	Pro	Ala	Lys	Lys	Arg	Leu	Asn	Phe	Gly	Gln	Thr
				165					170					175	
Gly	Asp	Ser	Glu	Ser	Val	Pro	Asp	Pro	Gln	Pro	Leu	Gly	Glu	Pro	Pro
			180						185				190		
Ala	Thr	Pro	Ala	Ala	Val	Gly	Pro	Thr	Thr	Met	Ala	Ser	Gly	Gly	Gly
			195						200				205		
Ala	Pro	Met	Ala	Asp	Asn	Asn	Glu	Gly	Ala	Asp	Gly	Val	Gly	Asn	Ala
		210							215				220		
Ser	Gly	Asn	Trp	His	Cys	Asp	Ser	Thr	Trp	Leu	Gly	Asp	Arg	Val	Ile
225									230				235		240
Thr	Thr	Ser	Thr	Arg	Thr	Trp	Ala	Leu	Pro	Thr	Tyr	Asn	Asn	His	Leu
				245					250					255	
Tyr	Lys	Gln	Ile	Ser	Ser	Ala	Ser	Thr	Gly	Ala	Ser	Asn	Asp	Asn	His
			260						265				270		
Tyr	Phe	Gly	Tyr	Ser	Thr	Pro	Trp	Gly	Tyr	Phe	Asp	Phe	Asn	Arg	Phe
		275							280				285		
His	Cys	His	Phe	Ser	Pro	Arg	Asp	Trp	Gln	Arg	Leu	Ile	Asn	Asn	Asn
									295				300		
Trp	Gly	Phe	Arg	Pro	Lys	Arg	Leu	Asn	Phe	Lys	Leu	Phe	Asn	Ile	Gln
305									310				315		320
Val	Lys	Glu	Val	Thr	Thr	Asn	Asp	Gly	Val	Thr	Thr	Ile	Ala	Asn	Asn
				325					330					335	
Leu	Thr	Ser	Thr	Val	Gln	Val	Phe	Ser	Asp	Ser	Glu	Tyr	Gln	Leu	Pro
				340					345				350		
Tyr	Val	Leu	Gly	Ser	Ala	His	Gln	Gly	Cys	Leu	Pro	Pro	Phe	Pro	Ala
		355							360				365		
Asp	Val	Phe	Met	Ile	Pro	Gln	Tyr	Gly	Tyr	Leu	Thr	Leu	Asn	Asn	Gly
				370					375				380		
Ser	Gln	Ala	Val	Gly	Arg	Ser	Ser	Phe	Tyr	Cys	Leu	Glu	Tyr	Phe	Pro
385									390				395		400
Ser	Gln	Met	Leu	Arg	Thr	Gly	Asn	Asn	Phe	Thr	Phe	Ser	Tyr	Thr	Phe
				405					410					415	
Glu	Glu	Val	Pro	Phe	His	Ser	Ser	Tyr	Ala	His	Ser	Gln	Ser	Leu	Asp
				420					425					430	
Arg	Leu	Met	Asn	Pro	Leu	Ile	Asp	Gln	Tyr	Leu	Tyr	Tyr	Leu	Asn	Arg
				435					440					445	
Thr	Gln	Asn	Gln	Ser	Gly	Ser	Ala	Gln	Asn	Lys	Asp	Leu	Leu	Phe	Ser
				450					455					460	

-continued

Arg Gly Ser Pro Ala Gly Met Ser Val Gln Pro Lys Asn Trp Leu Pro
 465 470 475 480
 Gly Pro Cys Tyr Arg Gln Gln Arg Val Ser Lys Thr Lys Thr Asp Asn
 485 490 495
 Asn Asn Ser Asn Phe Thr Trp Thr Gly Ala Ser Lys Tyr Asn Leu Asn
 500 505 510
 Gly Arg Glu Ser Ile Ile Asn Pro Gly Thr Ala Met Ala Ser His Lys
 515 520 525
 Asp Asp Glu Asp Lys Phe Phe Pro Met Ser Gly Val Met Ile Phe Gly
 530 535 540
 Lys Glu Ser Ala Gly Ala Ser Asn Thr Ala Leu Asp Asn Val Met Ile
 545 550 555 560
 Thr Asp Glu Glu Glu Ile Lys Ala Thr Asn Pro Val Ala Thr Glu Arg
 565 570 575
 Phe Gly Thr Val Ala Val Asn Phe Gln Ser Ser Ser Thr Asp Pro Ala
 580 585 590
 Thr Gly Asp Val His Ala Met Gly Ala Leu Pro Gly Met Val Trp Gln
 595 600 605
 Asp Arg Asp Val Tyr Leu Gln Gly Pro Ile Trp Ala Lys Ile Pro His
 610 615 620
 Thr Asp Gly His Phe His Pro Ser Pro Leu Met Gly Gly Phe Gly Leu
 625 630 635 640
 Lys Asn Pro Pro Pro Gln Ile Leu Ile Lys Asn Thr Pro Val Pro Ala
 645 650 655
 Asn Pro Pro Ala Glu Phe Ser Ala Thr Lys Phe Ala Ser Phe Ile Thr
 660 665 670
 Gln Tyr Ser Thr Gly Gln Val Ser Val Glu Ile Glu Trp Glu Leu Gln
 675 680 685
 Lys Glu Asn Ser Lys Arg Trp Asn Pro Glu Val Gln Tyr Thr Ser Asn
 690 695 700
 Tyr Ala Lys Ser Ala Asn Val Asp Phe Thr Val Asp Asn Asn Gly Leu
 705 710 715 720
 Tyr Thr Glu Pro Arg Pro Ile Gly Thr Arg Tyr Leu Thr Arg Pro Leu
 725 730 735

<210> SEQ ID NO 2

<211> LENGTH: 737

<212> TYPE: PRT

<213> ORGANISM: adeno-associated virus 7

<400> SEQUENCE: 2

Met Ala Ala Asp Gly Tyr Leu Pro Asp Trp Leu Glu Asp Asn Leu Ser
 1 5 10 15
 Glu Gly Ile Arg Glu Trp Trp Asp Leu Lys Pro Gly Ala Pro Lys Pro
 20 25 30
 Lys Ala Asn Gln Gln Lys Gln Asp Asn Gly Arg Gly Leu Val Leu Pro
 35 40 45
 Gly Tyr Lys Tyr Leu Gly Pro Phe Asn Gly Leu Asp Lys Gly Glu Pro
 50 55 60
 Val Asn Ala Ala Asp Ala Ala Ala Leu Glu His Asp Lys Ala Tyr Asp
 65 70 75 80
 Gln Gln Leu Lys Ala Gly Asp Asn Pro Tyr Leu Arg Tyr Asn His Ala
 85 90 95
 Asp Ala Glu Phe Gln Glu Arg Leu Gln Glu Asp Thr Ser Phe Gly Gly

-continued

100							105					110				
Asn	Leu	Gly	Arg	Ala	Val	Phe	Gln	Ala	Lys	Lys	Arg	Val	Leu	Glu	Pro	
		115					120					125				
Leu	Gly	Leu	Val	Glu	Glu	Gly	Ala	Lys	Thr	Ala	Pro	Ala	Lys	Lys	Arg	
		130				135					140					
Pro	Val	Glu	Pro	Ser	Pro	Gln	Arg	Ser	Pro	Asp	Ser	Ser	Thr	Gly	Ile	
145					150					155					160	
Gly	Lys	Lys	Gly	Gln	Gln	Pro	Ala	Arg	Lys	Arg	Leu	Asn	Phe	Gly	Gln	
				165					170					175		
Thr	Gly	Asp	Ser	Glu	Ser	Val	Pro	Asp	Pro	Gln	Pro	Leu	Gly	Glu	Pro	
			180					185					190			
Pro	Ala	Ala	Pro	Ser	Ser	Val	Gly	Ser	Gly	Thr	Val	Ala	Ala	Gly	Gly	
		195					200					205				
Gly	Ala	Pro	Met	Ala	Asp	Asn	Asn	Glu	Gly	Ala	Asp	Gly	Val	Gly	Asn	
		210				215					220					
Ala	Ser	Gly	Asn	Trp	His	Cys	Asp	Ser	Thr	Trp	Leu	Gly	Asp	Arg	Val	
225					230					235					240	
Ile	Thr	Thr	Ser	Thr	Arg	Thr	Trp	Ala	Leu	Pro	Thr	Tyr	Asn	Asn	His	
				245					250					255		
Leu	Tyr	Lys	Gln	Ile	Ser	Ser	Glu	Thr	Ala	Gly	Ser	Thr	Asn	Asp	Asn	
			260					265					270			
Thr	Tyr	Phe	Gly	Tyr	Ser	Thr	Pro	Trp	Gly	Tyr	Phe	Asp	Phe	Asn	Arg	
		275					280					285				
Phe	His	Cys	His	Phe	Ser	Pro	Arg	Asp	Trp	Gln	Arg	Leu	Ile	Asn	Asn	
		290				295					300					
Asn	Trp	Gly	Phe	Arg	Pro	Lys	Lys	Leu	Arg	Phe	Lys	Leu	Phe	Asn	Ile	
305					310					315					320	
Gln	Val	Lys	Glu	Val	Thr	Thr	Asn	Asp	Gly	Val	Thr	Thr	Ile	Ala	Asn	
				325					330					335		
Asn	Leu	Thr	Ser	Thr	Ile	Gln	Val	Phe	Ser	Asp	Ser	Glu	Tyr	Gln	Leu	
			340					345					350			
Pro	Tyr	Val	Leu	Gly	Ser	Ala	His	Gln	Gly	Cys	Leu	Pro	Pro	Phe	Pro	
		355					360					365				
Ala	Asp	Val	Phe	Met	Ile	Pro	Gln	Tyr	Gly	Tyr	Leu	Thr	Leu	Asn	Asn	
		370				375					380					
Gly	Ser	Gln	Ser	Val	Gly	Arg	Ser	Ser	Phe	Tyr	Cys	Leu	Glu	Tyr	Phe	
385					390					395					400	
Pro	Ser	Gln	Met	Leu	Arg	Thr	Gly	Asn	Asn	Phe	Glu	Phe	Ser	Tyr	Ser	
			405						410					415		
Phe	Glu	Asp	Val	Pro	Phe	His	Ser	Ser	Tyr	Ala	His	Ser	Gln	Ser	Leu	
			420					425					430			
Asp	Arg	Leu	Met	Asn	Pro	Leu	Ile	Asp	Gln	Tyr	Leu	Tyr	Tyr	Leu	Ala	
		435				440						445				
Arg	Thr	Gln	Ser	Asn	Pro	Gly	Gly	Thr	Ala	Gly	Asn	Arg	Glu	Leu	Gln	
		450				455					460					
Phe	Tyr	Gln	Gly	Gly	Pro	Ser	Thr	Met	Ala	Glu	Gln	Ala	Lys	Asn	Trp	
465					470					475					480	
Leu	Pro	Gly	Pro	Cys	Phe	Arg	Gln	Gln	Arg	Val	Ser	Lys	Thr	Leu	Asp	
				485					490					495		
Gln	Asn	Asn	Asn	Ser	Asn	Phe	Ala	Trp	Thr	Gly	Ala	Thr	Lys	Tyr	His	
			500					505					510			
Leu	Asn	Gly	Arg	Asn	Ser	Leu	Val	Asn	Pro	Gly	Val	Ala	Met	Ala	Thr	
		515					520					525				

-continued

His Lys Asp Asp Glu Asp Arg Phe Phe Pro Ser Ser Gly Val Leu Ile
 530 535 540
 Phe Gly Lys Thr Gly Ala Thr Asn Lys Thr Thr Leu Glu Asn Val Leu
 545 550 555 560
 Met Thr Asn Glu Glu Glu Ile Arg Pro Thr Asn Pro Val Ala Thr Glu
 565 570 575
 Glu Tyr Gly Ile Val Ser Ser Asn Leu Gln Ala Ala Asn Thr Ala Ala
 580 585 590
 Gln Thr Gln Val Val Asn Asn Gln Gly Ala Leu Pro Gly Met Val Trp
 595 600 605
 Gln Asn Arg Asp Val Tyr Leu Gln Gly Pro Ile Trp Ala Lys Ile Pro
 610 615 620
 His Thr Asp Gly Asn Phe His Pro Ser Pro Leu Met Gly Gly Phe Gly
 625 630 635 640
 Leu Lys His Pro Pro Pro Gln Ile Leu Ile Lys Asn Thr Pro Val Pro
 645 650 655
 Ala Asn Pro Pro Glu Val Phe Thr Pro Ala Lys Phe Ala Ser Phe Ile
 660 665 670
 Thr Gln Tyr Ser Thr Gly Gln Val Ser Val Glu Ile Glu Trp Glu Leu
 675 680 685
 Gln Lys Glu Asn Ser Lys Arg Trp Asn Pro Glu Ile Gln Tyr Thr Ser
 690 695 700
 Asn Thr Glu Lys Gln Thr Gly Val Asp Phe Ala Val Asp Ser Gln Gly
 705 710 715 720
 Val Tyr Ser Glu Pro Arg Pro Ile Gly Thr Arg Tyr Leu Thr Arg Asn
 725 730 735
 Leu

<210> SEQ ID NO 3
 <211> LENGTH: 735
 <212> TYPE: PRT
 <213> ORGANISM: adeno-associated virus 2

<400> SEQUENCE: 3

Met Ala Ala Asp Gly Tyr Leu Pro Asp Trp Leu Glu Asp Thr Leu Ser
 1 5 10 15
 Glu Gly Ile Arg Gln Trp Trp Lys Leu Lys Pro Gly Pro Pro Pro
 20 25 30
 Lys Pro Ala Glu Arg His Lys Asp Ser Arg Gly Leu Val Leu Pro
 35 40 45
 Gly Tyr Lys Tyr Leu Gly Pro Phe Asn Gly Leu Asp Lys Gly Glu Pro
 50 55 60
 Val Asn Glu Ala Asp Ala Ala Ala Leu Glu His Asp Lys Ala Tyr Asp
 65 70 75 80
 Arg Gln Leu Asp Ser Gly Asp Asn Pro Tyr Leu Lys Tyr Asn His Ala
 85 90 95
 Asp Ala Glu Phe Gln Glu Arg Leu Lys Glu Asp Thr Ser Phe Gly Gly
 100 105 110
 Asn Leu Gly Arg Ala Val Phe Gln Ala Lys Lys Arg Val Leu Glu Pro
 115 120 125
 Leu Gly Leu Val Glu Glu Pro Val Lys Thr Ala Pro Gly Lys Lys Arg
 130 135 140
 Pro Val Glu His Ser Pro Val Glu Pro Asp Ser Ser Ser Gly Thr Gly
 145 150 155 160

-continued

Lys Ala Gly Gln Gln Pro Ala Arg Lys Arg Leu Asn Phe Gly Gln Thr
 165 170 175
 Gly Asp Ala Asp Ser Val Pro Asp Pro Gln Pro Leu Gly Gln Pro Pro
 180 185 190
 Ala Ala Pro Ser Gly Leu Gly Thr Asn Thr Met Ala Thr Gly Ser Gly
 195 200 205
 Ala Pro Met Ala Asp Asn Asn Glu Gly Ala Asp Gly Val Gly Asn Ser
 210 215 220
 Ser Gly Asn Trp His Cys Asp Ser Thr Trp Met Gly Asp Arg Val Ile
 225 230 235 240
 Thr Thr Ser Thr Arg Thr Trp Ala Leu Pro Thr Tyr Asn Asn His Leu
 245 250 255
 Tyr Lys Gln Ile Ser Ser Gln Ser Gly Ala Ser Asn Asp Asn His Tyr
 260 265 270
 Phe Gly Tyr Ser Thr Pro Trp Gly Tyr Phe Asp Phe Asn Arg Phe His
 275 280 285
 Cys His Phe Ser Pro Arg Asp Trp Gln Arg Leu Ile Asn Asn Asn Trp
 290 295 300
 Gly Phe Arg Pro Lys Arg Leu Asn Phe Lys Leu Phe Asn Ile Gln Val
 305 310 315 320
 Lys Glu Val Thr Gln Asn Asp Gly Thr Thr Thr Ile Ala Asn Asn Leu
 325 330 335
 Thr Ser Thr Val Gln Val Phe Thr Asp Ser Glu Tyr Gln Leu Pro Tyr
 340 345 350
 Val Leu Gly Ser Ala His Gln Gly Cys Leu Pro Pro Phe Pro Ala Asp
 355 360 365
 Val Phe Met Val Pro Gln Tyr Gly Tyr Leu Thr Leu Asn Asn Gly Ser
 370 375 380
 Gln Ala Val Gly Arg Ser Ser Phe Tyr Cys Leu Glu Tyr Phe Pro Ser
 385 390 395 400
 Gln Met Leu Arg Thr Gly Asn Asn Phe Thr Phe Ser Tyr Thr Phe Glu
 405 410 415
 Asp Val Pro Phe His Ser Ser Tyr Ala His Ser Gln Ser Leu Asp Arg
 420 425 430
 Leu Met Asn Pro Leu Ile Asp Gln Tyr Leu Tyr Tyr Leu Ser Arg Thr
 435 440 445
 Asn Thr Pro Ser Gly Thr Thr Thr Gln Ser Arg Leu Gln Phe Ser Gln
 450 455 460
 Ala Gly Ala Ser Asp Ile Arg Asp Gln Ser Arg Asn Trp Leu Pro Gly
 465 470 475 480
 Pro Cys Tyr Arg Gln Gln Arg Val Ser Lys Thr Ser Ala Asp Asn Asn
 485 490 495
 Asn Ser Glu Tyr Ser Trp Thr Gly Ala Thr Lys Tyr His Leu Asn Gly
 500 505 510
 Arg Asp Ser Leu Val Asn Pro Gly Pro Ala Met Ala Ser His Lys Asp
 515 520 525
 Asp Glu Glu Lys Phe Phe Pro Gln Ser Gly Val Leu Ile Phe Gly Lys
 530 535 540
 Gln Gly Ser Glu Lys Thr Asn Val Asp Ile Glu Lys Val Met Ile Thr
 545 550 555 560
 Asp Glu Glu Glu Ile Arg Thr Thr Asn Pro Val Ala Thr Glu Gln Tyr
 565 570 575

-continued

Gly Ser Val Ser Thr Asn Leu Gln Arg Gly Asn Arg Gln Ala Ala Thr
 580 585 590
 Ala Asp Val Asn Thr Gln Gly Val Leu Pro Gly Met Val Trp Gln Asp
 595 600 605
 Arg Asp Val Tyr Leu Gln Gly Pro Ile Trp Ala Lys Ile Pro His Thr
 610 615 620
 Asp Gly His Phe His Pro Ser Pro Leu Met Gly Gly Phe Gly Leu Lys
 625 630 635 640
 His Pro Pro Pro Gln Ile Leu Ile Lys Asn Thr Pro Val Pro Ala Asn
 645 650 655
 Pro Ser Thr Thr Phe Ser Ala Ala Lys Phe Ala Ser Phe Ile Thr Gln
 660 665 670
 Tyr Ser Thr Gly Gln Val Ser Val Glu Ile Glu Trp Glu Leu Gln Lys
 675 680 685
 Glu Asn Ser Lys Arg Trp Asn Pro Glu Ile Gln Tyr Thr Ser Asn Tyr
 690 695 700
 Asn Lys Ser Val Asn Val Asp Phe Thr Val Asp Thr Asn Gly Val Tyr
 705 710 715 720
 Ser Glu Pro Arg Pro Ile Gly Thr Arg Tyr Leu Thr Arg Asn Leu
 725 730 735

<210> SEQ ID NO 4
 <211> LENGTH: 738
 <212> TYPE: PRT
 <213> ORGANISM: adeno-associated virus 8

<400> SEQUENCE: 4

Met Ala Ala Asp Gly Tyr Leu Pro Asp Trp Leu Glu Asp Asn Leu Ser
 1 5 10 15
 Glu Gly Ile Arg Glu Trp Trp Ala Leu Lys Pro Gly Ala Pro Lys Pro
 20 25 30
 Lys Ala Asn Gln Gln Lys Gln Asp Asp Gly Arg Gly Leu Val Leu Pro
 35 40 45
 Gly Tyr Lys Tyr Leu Gly Pro Phe Asn Gly Leu Asp Lys Gly Glu Pro
 50 55 60
 Val Asn Ala Ala Asp Ala Ala Ala Leu Glu His Asp Lys Ala Tyr Asp
 65 70 75 80
 Gln Gln Leu Gln Ala Gly Asp Asn Pro Tyr Leu Arg Tyr Asn His Ala
 85 90 95
 Asp Ala Glu Phe Gln Glu Arg Leu Gln Glu Asp Thr Ser Phe Gly Gly
 100 105 110
 Asn Leu Gly Arg Ala Val Phe Gln Ala Lys Lys Arg Val Leu Glu Pro
 115 120 125
 Leu Gly Leu Val Glu Glu Gly Ala Lys Thr Ala Pro Gly Lys Lys Arg
 130 135 140
 Pro Val Glu Pro Ser Pro Gln Arg Ser Pro Asp Ser Ser Thr Gly Ile
 145 150 155 160
 Gly Lys Lys Gly Gln Gln Pro Ala Arg Lys Arg Leu Asn Phe Gly Gln
 165 170 175
 Thr Gly Asp Ser Glu Ser Val Pro Asp Pro Gln Pro Leu Gly Glu Pro
 180 185 190
 Pro Ala Ala Pro Ser Gly Val Gly Pro Asn Thr Met Ala Ala Gly Gly
 195 200 205
 Gly Ala Pro Met Ala Asp Asn Asn Glu Gly Ala Asp Gly Val Gly Ser
 210 215 220

-continued

Ser	Ser	Gly	Asn	Trp	His	Cys	Asp	Ser	Thr	Trp	Leu	Gly	Asp	Arg	Val	225	230	235	240
Ile	Thr	Thr	Ser	Thr	Arg	Thr	Trp	Ala	Leu	Pro	Thr	Tyr	Asn	Asn	His	245	250	255	
Leu	Tyr	Lys	Gln	Ile	Ser	Asn	Gly	Thr	Ser	Gly	Gly	Ala	Thr	Asn	Asp	260	265	270	
Asn	Thr	Tyr	Phe	Gly	Tyr	Ser	Thr	Pro	Trp	Gly	Tyr	Phe	Asp	Phe	Asn	275	280	285	
Arg	Phe	His	Cys	His	Phe	Ser	Pro	Arg	Asp	Trp	Gln	Arg	Leu	Ile	Asn	290	295	300	
Asn	Asn	Trp	Gly	Phe	Arg	Pro	Lys	Arg	Leu	Ser	Phe	Lys	Leu	Phe	Asn	305	310	315	320
Ile	Gln	Val	Lys	Glu	Val	Thr	Gln	Asn	Glu	Gly	Thr	Lys	Thr	Ile	Ala	325	330		335
Asn	Asn	Leu	Thr	Ser	Thr	Ile	Gln	Val	Phe	Thr	Asp	Ser	Glu	Tyr	Gln	340	345	350	
Leu	Pro	Tyr	Val	Leu	Gly	Ser	Ala	His	Gln	Gly	Cys	Leu	Pro	Pro	Phe	355	360	365	
Pro	Ala	Asp	Val	Phe	Met	Ile	Pro	Gln	Tyr	Gly	Tyr	Leu	Thr	Leu	Asn	370	375	380	
Asn	Gly	Ser	Gln	Ala	Val	Gly	Arg	Ser	Ser	Phe	Tyr	Cys	Leu	Glu	Tyr	385	390	395	400
Phe	Pro	Ser	Gln	Met	Leu	Arg	Thr	Gly	Asn	Asn	Phe	Gln	Phe	Thr	Tyr	405	410		415
Thr	Phe	Glu	Asp	Val	Pro	Phe	His	Ser	Ser	Tyr	Ala	His	Ser	Gln	Ser	420	425	430	
Leu	Asp	Arg	Leu	Met	Asn	Pro	Leu	Ile	Asp	Gln	Tyr	Leu	Tyr	Tyr	Leu	435	440	445	
Ser	Arg	Thr	Gln	Thr	Thr	Gly	Gly	Thr	Ala	Asn	Thr	Gln	Thr	Leu	Gly	450	455	460	
Phe	Ser	Gln	Gly	Gly	Pro	Asn	Thr	Met	Ala	Asn	Gln	Ala	Lys	Asn	Trp	465	470	475	480
Leu	Pro	Gly	Pro	Cys	Tyr	Arg	Gln	Gln	Arg	Val	Ser	Thr	Thr	Thr	Gly	485	490		495
Gln	Asn	Asn	Asn	Ser	Asn	Phe	Ala	Trp	Thr	Ala	Gly	Thr	Lys	Tyr	His	500	505		510
Leu	Asn	Gly	Arg	Asn	Ser	Leu	Ala	Asn	Pro	Gly	Ile	Ala	Met	Ala	Thr	515	520	525	
His	Lys	Asp	Asp	Glu	Glu	Arg	Phe	Phe	Pro	Ser	Asn	Gly	Ile	Leu	Ile	530	535	540	
Phe	Gly	Lys	Gln	Asn	Ala	Ala	Arg	Asp	Asn	Ala	Asp	Tyr	Ser	Asp	Val	545	550	555	560
Met	Leu	Thr	Ser	Glu	Glu	Glu	Ile	Lys	Thr	Thr	Asn	Pro	Val	Ala	Thr	565	570		575
Glu	Glu	Tyr	Gly	Ile	Val	Ala	Asp	Asn	Leu	Gln	Gln	Gln	Asn	Thr	Ala	580	585		590
Pro	Gln	Ile	Gly	Thr	Val	Asn	Ser	Gln	Gly	Ala	Leu	Pro	Gly	Met	Val	595	600	605	
Trp	Gln	Asn	Arg	Asp	Val	Tyr	Leu	Gln	Gly	Pro	Ile	Trp	Ala	Lys	Ile	610	615	620	
Pro	His	Thr	Asp	Gly	Asn	Phe	His	Pro	Ser	Pro	Leu	Met	Gly	Gly	Phe	625	630	635	640

-continued

Gly Leu Lys His Pro Pro Pro Gln Ile Leu Ile Lys Asn Thr Pro Val
645 650 655

Pro Ala Asp Pro Pro Thr Thr Phe Asn Gln Ser Lys Leu Asn Ser Phe
660 665 670

Ile Thr Gln Tyr Ser Thr Gly Gln Val Ser Val Glu Ile Glu Trp Glu
675 680 685

Leu Gln Lys Glu Asn Ser Lys Arg Trp Asn Pro Glu Ile Gln Tyr Thr
690 695 700

Ser Asn Tyr Tyr Lys Ser Thr Ser Val Asp Phe Ala Val Asn Thr Glu
705 710 715 720

Gly Val Tyr Ser Glu Pro Arg Pro Ile Gly Thr Arg Tyr Leu Thr Arg
725 730 735

Asn Leu

<210> SEQ ID NO 5
 <211> LENGTH: 72
 <212> TYPE: PRT
 <213> ORGANISM: adeno-associated virus 1

<400> SEQUENCE: 5

Gly Ala Asp Gly Val Gly Asn Ala Ser Gly Asn Trp His Cys Asp Ser
1 5 10 15

Thr Trp Leu Gly Asp Arg Val Ile Thr Thr Ser Thr Arg Thr Trp Ala
20 25 30

Leu Pro Thr Tyr Asn Asn His Leu Tyr Lys Gln Ile Ser Ser Ala Ser
35 40 45

Thr Gly Ala Ser Asn Asp Asn His Tyr Phe Gly Tyr Ser Thr Pro Trp
50 55 60

Gly Tyr Phe Asp Phe Asn Arg Phe
65 70

<210> SEQ ID NO 6
 <211> LENGTH: 71
 <212> TYPE: PRT
 <213> ORGANISM: adeno-associated virus 2

<400> SEQUENCE: 6

Gly Ala Asp Gly Val Gly Asn Ser Ser Gly Asn Trp His Cys Asp Ser
1 5 10 15

Thr Trp Met Gly Asp Arg Val Ile Thr Thr Ser Thr Arg Thr Trp Ala
20 25 30

Leu Pro Thr Tyr Asn Asn His Leu Tyr Lys Gln Ile Ser Ser Gln Ser
35 40 45

Gly Ala Ser Asn Asp Asn His Tyr Phe Gly Tyr Ser Thr Pro Trp Gly
50 55 60

Tyr Phe Asp Phe Asn Arg Phe
65 70

<210> SEQ ID NO 7
 <211> LENGTH: 71
 <212> TYPE: PRT
 <213> ORGANISM: adeno-associated virus 3A

<400> SEQUENCE: 7

Gly Ala Asp Gly Val Gly Asn Ser Ser Gly Asn Trp His Cys Asp Ser
1 5 10 15

Gln Trp Leu Gly Asp Arg Val Ile Thr Thr Ser Thr Arg Thr Trp Ala
20 25 30

-continued

Leu Pro Thr Tyr Asn Asn His Leu Tyr Lys Gln Ile Ser Ser Gln Ser
35 40 45

Gly Ala Ser Asn Asp Asn His Tyr Phe Gly Tyr Ser Thr Pro Trp Gly
50 55 60

Tyr Phe Asp Phe Asn Arg Phe
65 70

<210> SEQ ID NO 8
<211> LENGTH: 68
<212> TYPE: PRT
<213> ORGANISM: adeno-associated virus 4

<400> SEQUENCE: 8

Gly Ala Asp Gly Val Gly Asn Ala Ser Gly Asp Trp His Cys Asp Ser
1 5 10 15

Thr Trp Ser Glu Gly His Val Thr Thr Thr Ser Thr Arg Thr Trp Val
20 25 30

Leu Pro Thr Tyr Asn Asn His Leu Tyr Lys Arg Leu Gly Glu Ser Leu
35 40 45

Gln Ser Asn Thr Tyr Asn Gly Phe Ser Thr Pro Trp Gly Tyr Phe Asp
50 55 60

Phe Asn Arg Phe
65

<210> SEQ ID NO 9
<211> LENGTH: 72
<212> TYPE: PRT
<213> ORGANISM: adeno-associated virus 5

<400> SEQUENCE: 9

Gly Ala Asp Gly Val Gly Asn Ala Ser Gly Asp Trp His Cys Asp Ser
1 5 10 15

Thr Trp Met Gly Asp Arg Val Val Thr Lys Ser Thr Arg Thr Trp Val
20 25 30

Leu Pro Ser Tyr Asn Asn His Gln Tyr Arg Glu Ile Lys Ser Gly Ser
35 40 45

Val Asp Gly Ser Asn Ala Asn Ala Tyr Phe Gly Tyr Ser Thr Pro Trp
50 55 60

Gly Tyr Phe Asp Phe Asn Arg Phe
65 70

<210> SEQ ID NO 10
<211> LENGTH: 72
<212> TYPE: PRT
<213> ORGANISM: adeno-associated virus 7

<400> SEQUENCE: 10

Gly Ala Asp Gly Val Gly Asn Ala Ser Gly Asn Trp His Cys Asp Ser
1 5 10 15

Thr Trp Leu Gly Asp Arg Val Ile Thr Thr Ser Thr Arg Thr Trp Ala
20 25 30

Leu Pro Thr Tyr Asn Asn His Leu Tyr Lys Gln Ile Ser Ser Glu Thr
35 40 45

Ala Gly Ser Thr Asn Asp Asn Thr Tyr Phe Gly Tyr Ser Thr Pro Trp
50 55 60

Gly Tyr Phe Asp Phe Asn Arg Phe
65 70

-continued

<210> SEQ ID NO 11
 <211> LENGTH: 73
 <212> TYPE: PRT
 <213> ORGANISM: adeno-associated virus 8

<400> SEQUENCE: 11

```
Gly Ala Asp Gly Val Gly Ser Ser Ser Gly Asn Trp His Cys Asp Ser
1      5      10      15
Thr Trp Leu Gly Asp Arg Val Ile Thr Thr Ser Thr Arg Thr Trp Ala
      20      25      30
Leu Pro Thr Tyr Asn Asn His Leu Tyr Lys Gln Ile Ser Asn Gly Thr
      35      40      45
Ser Gly Gly Ala Thr Asn Asp Asn Thr Tyr Phe Gly Tyr Ser Thr Pro
      50      55      60
Trp Gly Tyr Phe Asp Phe Asn Arg Phe
65      70
```

<210> SEQ ID NO 12
 <211> LENGTH: 73
 <212> TYPE: PRT
 <213> ORGANISM: adeno-associated virus 9

<400> SEQUENCE: 12

```
Gly Ala Asp Gly Val Gly Ser Ser Ser Gly Asn Trp His Cys Asp Ser
1      5      10      15
Gln Trp Leu Gly Asp Arg Val Ile Thr Thr Ser Thr Arg Thr Trp Ala
      20      25      30
Leu Pro Thr Tyr Asn Asn His Leu Tyr Lys Gln Ile Ser Asn Ser Thr
      35      40      45
Ser Gly Gly Ser Ser Asn Asp Asn Ala Tyr Phe Gly Tyr Ser Thr Pro
      50      55      60
Trp Gly Tyr Phe Asp Phe Asn Arg Phe
65      70
```

<210> SEQ ID NO 13
 <211> LENGTH: 736
 <212> TYPE: PRT
 <213> ORGANISM: adeno-associated virus 2

<400> SEQUENCE: 13

```
Met Ala Ala Asp Gly Tyr Leu Pro Asp Trp Leu Glu Asp Thr Leu Ser
1      5      10      15
Glu Gly Ile Arg Gln Trp Trp Lys Leu Lys Pro Gly Pro Pro Pro Pro
      20      25      30
Lys Pro Ala Glu Arg His Lys Asp Asp Ser Arg Gly Leu Val Leu Pro
      35      40      45
Gly Tyr Lys Tyr Leu Gly Pro Phe Asn Gly Leu Asp Lys Gly Glu Pro
      50      55      60
Val Asn Glu Ala Asp Ala Ala Ala Leu Glu His Asp Lys Ala Tyr Asp
65      70      75      80
Arg Gln Leu Asp Ser Gly Asp Asn Pro Tyr Leu Lys Tyr Asn His Ala
      85      90      95
Asp Ala Glu Phe Gln Glu Arg Leu Lys Glu Asp Thr Ser Phe Gly Gly
      100      105      110
Asn Leu Gly Arg Ala Val Phe Gln Ala Lys Lys Arg Val Leu Glu Pro
      115      120      125
Leu Gly Leu Val Glu Glu Pro Val Lys Thr Ala Pro Gly Lys Lys Arg
```

-continued

130	135	140
Pro Val Glu His Ser	Pro Val Glu Pro Asp Ser Ser Ser Gly Thr Gly	
145	150	155 160
Lys Ala Gly Gln Gln	Pro Ala Arg Lys Arg Leu Asn Phe Gly Gln Thr	
	165	170 175
Gly Asp Ala Asp Ser Val Pro Asp Pro Gln Pro Leu Gly Gln Pro Pro		
	180	185 190
Ala Ala Pro Ser Gly Leu Gly Thr Asn Thr Met Ala Thr Gly Ser Gly		
	195	200 205
Ala Pro Met Ala Asp Asn Asn Glu Gly Ala Asp Gly Val Gly Asn Ser		
	210	215 220
Ser Gly Asn Trp His Cys Asp Ser Thr Trp Met Gly Asp Arg Val Ile		
	225	230 235 240
Thr Thr Ser Thr Arg Thr Trp Ala Leu Pro Thr Tyr Asn Asn His Leu		
	245	250 255
Tyr Lys Gln Ile Ser Ser Ala Ser Thr Gly Ala Ser Asn Asp Asn His		
	260	265 270
Tyr Phe Gly Tyr Ser Thr Pro Trp Gly Tyr Phe Asp Phe Asn Arg Phe		
	275	280 285
His Cys His Phe Ser Pro Arg Asp Trp Gln Arg Leu Ile Asn Asn Asn		
	290	295 300
Trp Gly Phe Arg Pro Lys Arg Leu Asn Phe Lys Leu Phe Asn Ile Gln		
	305	310 315 320
Val Lys Glu Val Thr Gln Asn Asp Gly Thr Thr Thr Ile Ala Asn Asn		
	325	330 335
Leu Thr Ser Thr Val Gln Val Phe Thr Asp Ser Glu Tyr Gln Leu Pro		
	340	345 350
Tyr Val Leu Gly Ser Ala His Gln Gly Cys Leu Pro Pro Phe Pro Ala		
	355	360 365
Asp Val Phe Met Val Pro Gln Tyr Gly Tyr Leu Thr Leu Asn Asn Gly		
	370	375 380
Ser Gln Ala Val Gly Arg Ser Ser Phe Tyr Cys Leu Glu Tyr Phe Pro		
	385	390 395 400
Ser Gln Met Leu Arg Thr Gly Asn Asn Phe Thr Phe Ser Tyr Thr Phe		
	405	410 415
Glu Asp Val Pro Phe His Ser Ser Tyr Ala His Ser Gln Ser Leu Asp		
	420	425 430
Arg Leu Met Asn Pro Leu Ile Asp Gln Tyr Leu Tyr Tyr Leu Ser Arg		
	435	440 445
Thr Asn Thr Pro Ser Gly Thr Thr Thr Gln Ser Arg Leu Gln Phe Ser		
	450	455 460
Gln Ala Gly Ala Ser Asp Ile Arg Asp Gln Ser Arg Asn Trp Leu Pro		
	465	470 475 480
Gly Pro Cys Tyr Arg Gln Gln Arg Val Ser Lys Thr Ser Ala Asp Asn		
	485	490 495
Asn Asn Ser Glu Tyr Ser Trp Thr Gly Ala Thr Lys Tyr His Leu Asn		
	500	505 510
Gly Arg Asp Ser Leu Val Asn Pro Gly Pro Ala Met Ala Ser His Lys		
	515	520 525
Asp Asp Glu Glu Lys Phe Phe Pro Gln Ser Gly Val Leu Ile Phe Gly		
	530	535 540
Lys Gln Gly Ser Glu Lys Thr Asn Val Asp Ile Glu Lys Val Met Ile		
	545	550 555 560

-continued

195	200	205
Ala Pro Met Ala Asp Asn Asn Glu Gly Ala Asp Gly Val Gly Asn Ser 210 215 220		
Ser Gly Asn Trp His Cys Asp Ser Gln Trp Leu Gly Asp Arg Val Ile 225 230 235 240		
Thr Thr Ser Thr Arg Thr Trp Ala Leu Pro Thr Tyr Asn Asn His Leu 245 250 255		
Tyr Lys Gln Ile Ser Ser Ala Ser Thr Gly Ala Ser Asn Asp Asn His 260 265 270		
Tyr Phe Gly Tyr Ser Thr Pro Trp Gly Tyr Phe Asp Phe Asn Arg Phe 275 280 285		
His Cys His Phe Ser Pro Arg Asp Trp Gln Arg Leu Ile Asn Asn Asn 290 295 300		
Trp Gly Phe Arg Pro Lys Lys Leu Ser Phe Lys Leu Phe Asn Ile Gln 305 310 315 320		
Val Lys Glu Val Thr Gln Asn Asp Gly Thr Thr Thr Ile Ala Asn Asn 325 330 335		
Leu Thr Ser Thr Val Gln Val Phe Thr Asp Ser Glu Tyr Gln Leu Pro 340 345 350		
Tyr Val Leu Gly Ser Ala His Gln Gly Cys Leu Pro Pro Phe Pro Ala 355 360 365		
Asp Val Phe Met Val Pro Gln Tyr Gly Tyr Leu Thr Leu Asn Asn Gly 370 375 380		
Ser Gln Ala Val Gly Arg Ser Ser Phe Tyr Cys Leu Glu Tyr Phe Pro 385 390 395 400		
Ser Gln Met Leu Arg Thr Gly Asn Asn Phe Gln Phe Ser Tyr Thr Phe 405 410 415		
Glu Asp Val Pro Phe His Ser Ser Tyr Ala His Ser Gln Ser Leu Asp 420 425 430		
Arg Leu Met Asn Pro Leu Ile Asp Gln Tyr Leu Tyr Tyr Leu Asn Arg 435 440 445		
Thr Gln Gly Thr Thr Ser Gly Thr Thr Asn Gln Ser Arg Leu Leu Phe 450 455 460		
Ser Gln Ala Gly Pro Gln Ser Met Ser Leu Gln Ala Arg Asn Trp Leu 465 470 475 480		
Pro Gly Pro Cys Tyr Arg Gln Gln Arg Leu Ser Lys Thr Ala Asn Asp 485 490 495		
Asn Asn Asn Ser Asn Phe Pro Trp Thr Ala Ala Ser Lys Tyr His Leu 500 505 510		
Asn Gly Arg Asp Ser Leu Val Asn Pro Gly Pro Ala Met Ala Ser His 515 520 525		
Lys Asp Asp Glu Glu Lys Phe Phe Pro Met His Gly Asn Leu Ile Phe 530 535 540		
Gly Lys Glu Gly Thr Thr Ala Ser Asn Ala Glu Leu Asp Asn Val Met 545 550 555 560		
Ile Thr Asp Glu Glu Glu Ile Arg Thr Thr Asn Pro Val Ala Thr Glu 565 570 575		
Gln Tyr Gly Thr Val Ala Asn Asn Leu Gln Ser Ser Asn Thr Ala Pro 580 585 590		
Thr Thr Arg Thr Val Asn Asp Gln Gly Ala Leu Pro Gly Met Val Trp 595 600 605		
Gln Asp Arg Asp Val Tyr Leu Gln Gly Pro Ile Trp Ala Lys Ile Pro 610 615 620		

-continued

His Thr Asp Gly His Phe His Pro Ser Pro Leu Met Gly Gly Phe Gly
 625 630 635 640
 Leu Lys His Pro Pro Pro Gln Ile Met Ile Lys Asn Thr Pro Val Pro
 645 650 655
 Ala Asn Pro Pro Thr Thr Phe Ser Pro Ala Lys Phe Ala Ser Phe Ile
 660 665 670
 Thr Gln Tyr Ser Thr Gly Gln Val Ser Val Glu Ile Glu Trp Glu Leu
 675 680 685
 Gln Lys Glu Asn Ser Lys Arg Trp Asn Pro Glu Ile Gln Tyr Thr Ser
 690 695 700
 Asn Tyr Asn Lys Ser Val Asn Val Asp Phe Thr Val Asp Thr Asn Gly
 705 710 715 720
 Val Tyr Ser Glu Pro Arg Pro Ile Gly Thr Arg Tyr Leu Thr Arg Asn
 725 730 735

Leu

<210> SEQ ID NO 15
 <211> LENGTH: 736
 <212> TYPE: PRT
 <213> ORGANISM: adeno-associated virus 3B

<400> SEQUENCE: 15

Met Ala Ala Asp Gly Tyr Leu Pro Asp Trp Leu Glu Asp Asn Leu Ser
 1 5 10 15
 Glu Gly Ile Arg Glu Trp Trp Ala Leu Lys Pro Gly Val Pro Gln Pro
 20 25 30
 Lys Ala Asn Gln Gln His Gln Asp Asn Arg Arg Gly Leu Val Leu Pro
 35 40 45
 Gly Tyr Lys Tyr Leu Gly Pro Gly Asn Gly Leu Asp Lys Gly Glu Pro
 50 55 60
 Val Asn Glu Ala Asp Ala Ala Ala Leu Glu His Asp Lys Ala Tyr Asp
 65 70 75 80
 Gln Gln Leu Lys Ala Gly Asp Asn Pro Tyr Leu Lys Tyr Asn His Ala
 85 90 95
 Asp Ala Glu Phe Gln Glu Arg Leu Gln Glu Asp Thr Ser Phe Gly Gly
 100 105 110
 Asn Leu Gly Arg Ala Val Phe Gln Ala Lys Lys Arg Ile Leu Glu Pro
 115 120 125
 Leu Gly Leu Val Glu Glu Ala Ala Lys Thr Ala Pro Gly Lys Lys Arg
 130 135 140
 Pro Val Asp Gln Ser Pro Gln Glu Pro Asp Ser Ser Ser Gly Val Gly
 145 150 155 160
 Lys Ser Gly Lys Gln Pro Ala Arg Lys Arg Leu Asn Phe Gly Gln Thr
 165 170 175
 Gly Asp Ser Glu Ser Val Pro Asp Pro Gln Pro Leu Gly Glu Pro Pro
 180 185 190
 Ala Ala Pro Thr Ser Leu Gly Ser Asn Thr Met Ala Ser Gly Gly Gly
 195 200 205
 Ala Pro Met Ala Asp Asn Asn Glu Gly Ala Asp Gly Val Gly Asn Ser
 210 215 220
 Ser Gly Asn Trp His Cys Asp Ser Gln Trp Leu Gly Asp Arg Val Ile
 225 230 235 240
 Thr Thr Ser Thr Arg Thr Trp Ala Leu Pro Thr Tyr Asn Asn His Leu
 245 250 255

-continued

Tyr Lys Gln Ile Ser Ser Gln Ser Gly Ala Ser Asn Asp Asn His Tyr
 260 265 270
 Phe Gly Tyr Ser Thr Pro Trp Gly Tyr Phe Asp Phe Asn Arg Phe His
 275 280 285
 Cys His Phe Ser Pro Arg Asp Trp Gln Arg Leu Ile Asn Asn Asn Trp
 290 295 300
 Gly Phe Arg Pro Lys Lys Leu Ser Phe Lys Leu Phe Asn Ile Gln Val
 305 310 315 320
 Lys Glu Val Thr Gln Asn Asp Gly Thr Thr Thr Ile Ala Asn Asn Leu
 325 330 335
 Thr Ser Thr Val Gln Val Phe Thr Asp Ser Glu Tyr Gln Leu Pro Tyr
 340 345 350
 Val Leu Gly Ser Ala His Gln Gly Cys Leu Pro Pro Phe Pro Ala Asp
 355 360 365
 Val Phe Met Val Pro Gln Tyr Gly Tyr Leu Thr Leu Asn Asn Gly Ser
 370 375 380
 Gln Ala Val Gly Arg Ser Ser Phe Tyr Cys Leu Glu Tyr Phe Pro Ser
 385 390 395 400
 Gln Met Leu Arg Thr Gly Asn Asn Phe Gln Phe Ser Tyr Thr Phe Glu
 405 410 415
 Asp Val Pro Phe His Ser Ser Tyr Ala His Ser Gln Ser Leu Asp Arg
 420 425 430
 Leu Met Asn Pro Leu Ile Asp Gln Tyr Leu Tyr Tyr Leu Asn Arg Thr
 435 440 445
 Gln Gly Thr Thr Ser Gly Thr Thr Asn Gln Ser Arg Leu Leu Phe Ser
 450 455 460
 Gln Ala Gly Pro Gln Ser Met Ser Leu Gln Ala Arg Asn Trp Leu Pro
 465 470 475 480
 Gly Pro Cys Tyr Arg Gln Gln Arg Leu Ser Lys Thr Ala Asn Asp Asn
 485 490 495
 Asn Asn Ser Asn Phe Pro Trp Thr Ala Ala Ser Lys Tyr His Leu Asn
 500 505 510
 Gly Arg Asp Ser Leu Val Asn Pro Gly Pro Ala Met Ala Ser His Lys
 515 520 525
 Asp Asp Glu Glu Lys Phe Phe Pro Met His Gly Asn Leu Ile Phe Gly
 530 535 540
 Lys Glu Gly Thr Thr Ala Ser Asn Ala Glu Leu Asp Asn Val Met Ile
 545 550 555 560
 Thr Asp Glu Glu Glu Ile Arg Thr Thr Asn Pro Val Ala Thr Glu Gln
 565 570 575
 Tyr Gly Thr Val Ala Asn Asn Leu Gln Ser Ser Asn Thr Ala Pro Thr
 580 585 590
 Thr Arg Thr Val Asn Asp Gln Gly Ala Leu Pro Gly Met Val Trp Gln
 595 600 605
 Asp Arg Asp Val Tyr Leu Gln Gly Pro Ile Trp Ala Lys Ile Pro His
 610 615 620
 Thr Asp Gly His Phe His Pro Ser Pro Leu Met Gly Gly Phe Gly Leu
 625 630 635 640
 Lys His Pro Pro Pro Gln Ile Met Ile Lys Asn Thr Pro Val Pro Ala
 645 650 655
 Asn Pro Pro Thr Thr Phe Ser Pro Ala Lys Phe Ala Ser Phe Ile Thr
 660 665 670

-continued

Gln Tyr Ser Thr Gly Gln Val Ser Val Glu Ile Glu Trp Glu Leu Gln
 675 680 685
 Lys Glu Asn Ser Lys Arg Trp Asn Pro Glu Ile Gln Tyr Thr Ser Asn
 690 695 700
 Tyr Asn Lys Ser Val Asn Val Asp Phe Thr Val Asp Thr Asn Gly Val
 705 710 715 720
 Tyr Ser Glu Pro Arg Pro Ile Gly Thr Arg Tyr Leu Thr Arg Asn Leu
 725 730 735

 <210> SEQ ID NO 16
 <211> LENGTH: 736
 <212> TYPE: PRT
 <213> ORGANISM: adeno-associated virus 2

 <400> SEQUENCE: 16
 Met Ala Ala Asp Gly Tyr Leu Pro Asp Trp Leu Glu Asp Thr Leu Ser
 1 5 10 15
 Glu Gly Ile Arg Gln Trp Trp Lys Leu Lys Pro Gly Pro Pro Pro Pro
 20 25 30
 Lys Pro Ala Glu Arg His Lys Asp Asp Ser Arg Gly Leu Val Leu Pro
 35 40 45
 Gly Tyr Lys Tyr Leu Gly Pro Phe Asn Gly Leu Asp Lys Gly Glu Pro
 50 55 60
 Val Asn Glu Ala Asp Ala Ala Ala Leu Glu His Asp Lys Ala Tyr Asp
 65 70 75 80
 Arg Gln Leu Asp Ser Gly Asp Asn Pro Tyr Leu Lys Tyr Asn His Ala
 85 90 95
 Asp Ala Glu Phe Gln Glu Arg Leu Lys Glu Asp Thr Ser Phe Gly Gly
 100 105 110
 Asn Leu Gly Arg Ala Val Phe Gln Ala Lys Lys Arg Val Leu Glu Pro
 115 120 125
 Leu Gly Leu Val Glu Glu Pro Val Lys Thr Ala Pro Gly Lys Lys Arg
 130 135 140
 Pro Val Glu His Ser Pro Val Glu Pro Asp Ser Ser Ser Gly Thr Gly
 145 150 155 160
 Lys Ala Gly Gln Gln Pro Ala Arg Lys Arg Leu Asn Phe Gly Gln Thr
 165 170 175
 Gly Asp Ala Asp Ser Val Pro Asp Pro Gln Pro Leu Gly Gln Pro Pro
 180 185 190
 Ala Ala Pro Ser Gly Leu Gly Thr Asn Thr Met Ala Thr Gly Ser Gly
 195 200 205
 Ala Pro Met Ala Asp Asn Asn Glu Gly Ala Asp Gly Val Gly Asn Ser
 210 215 220
 Ser Gly Asn Trp His Cys Asp Ser Thr Trp Met Gly Asp Arg Val Ile
 225 230 235 240
 Thr Thr Ser Thr Arg Thr Trp Ala Leu Pro Thr Tyr Asn Asn His Leu
 245 250 255
 Tyr Lys Gln Ile Ser Ser Ala Ser Thr Gly Ala Ser Asn Asp Asn His
 260 265 270
 Tyr Phe Gly Tyr Ser Thr Pro Trp Gly Tyr Phe Asp Phe Asn Arg Phe
 275 280 285
 His Cys His Phe Ser Pro Arg Asp Trp Gln Arg Leu Ile Asn Asn Asn
 290 295 300
 Trp Gly Phe Arg Pro Lys Arg Leu Asn Phe Lys Leu Phe Asn Ile Gln
 305 310 315 320

-continued

Val	Lys	Glu	Val	Thr	Gln	Asn	Asp	Gly	Thr	Thr	Thr	Ile	Ala	Asn	Asn	
				325					330					335		
Leu	Thr	Ser	Thr	Val	Gln	Val	Phe	Thr	Asp	Ser	Glu	Tyr	Gln	Leu	Pro	
			340					345					350			
Tyr	Val	Leu	Gly	Ser	Ala	His	Gln	Gly	Cys	Leu	Pro	Pro	Phe	Pro	Ala	
		355					360					365				
Asp	Val	Phe	Met	Val	Pro	Gln	Tyr	Gly	Tyr	Leu	Thr	Leu	Asn	Asn	Gly	
	370					375					380					
Ser	Gln	Ala	Val	Gly	Arg	Ser	Ser	Phe	Tyr	Cys	Leu	Glu	Tyr	Phe	Pro	
385					390					395					400	
Ser	Gln	Met	Leu	Arg	Thr	Gly	Asn	Asn	Phe	Thr	Phe	Ser	Tyr	Thr	Phe	
			405						410					415		
Glu	Asp	Val	Pro	Phe	His	Ser	Ser	Tyr	Ala	His	Ser	Gln	Ser	Leu	Asp	
		420						425					430			
Arg	Leu	Met	Asn	Pro	Leu	Ile	Asp	Gln	Tyr	Leu	Tyr	Tyr	Leu	Ser	Arg	
	435						440					445				
Thr	Asn	Thr	Pro	Ser	Gly	Thr	Thr	Thr	Gln	Ser	Arg	Leu	Gln	Phe	Ser	
	450					455					460					
Gln	Ala	Gly	Ala	Ser	Asp	Ile	Arg	Asp	Gln	Ser	Arg	Asn	Trp	Leu	Pro	
465					470				475						480	
Gly	Pro	Cys	Tyr	Arg	Gln	Gln	Arg	Val	Ser	Lys	Thr	Ser	Ala	Asp	Asn	
				485				490						495		
Asn	Asn	Ser	Glu	Tyr	Ser	Trp	Thr	Gly	Ala	Thr	Lys	Tyr	His	Leu	Asn	
		500						505					510			
Gly	Arg	Asp	Ser	Leu	Val	Asn	Pro	Gly	Pro	Ala	Met	Ala	Ser	His	Lys	
	515						520					525				
Asp	Asp	Glu	Glu	Lys	Phe	Phe	Pro	Gln	Ser	Gly	Val	Leu	Ile	Phe	Gly	
	530					535					540					
Lys	Gln	Gly	Ser	Glu	Lys	Thr	Asn	Val	Asp	Ile	Glu	Lys	Val	Met	Ile	
545					550					555					560	
Thr	Asp	Glu	Glu	Glu	Ile	Arg	Thr	Thr	Asn	Pro	Val	Ala	Thr	Glu	Gln	
			565					570						575		
Tyr	Gly	Ser	Val	Ser	Thr	Asn	Leu	Gln	Arg	Gly	Asn	Arg	Gln	Ala	Ala	
		580						585					590			
Thr	Ala	Asp	Val	Asn	Thr	Gln	Gly	Val	Leu	Pro	Gly	Met	Val	Trp	Gln	
	595					600					605					
Asp	Arg	Asp	Val	Tyr	Leu	Gln	Gly	Pro	Ile	Trp	Ala	Lys	Ile	Pro	His	
	610					615					620					
Thr	Asp	Gly	His	Phe	His	Pro	Ser	Pro	Leu	Met	Gly	Gly	Phe	Gly	Leu	
625					630					635					640	
Lys	His	Pro	Pro	Pro	Gln	Ile	Leu	Ile	Lys	Asn	Thr	Pro	Val	Pro	Ala	
			645					650						655		
Asn	Pro	Ser	Thr	Thr	Phe	Ser	Ala	Ala	Lys	Phe	Ala	Ser	Phe	Ile	Thr	
			660					665					670			
Gln	Tyr	Ser	Thr	Gly	Gln	Val	Ser	Val	Glu	Ile	Glu	Trp	Glu	Leu	Gln	
		675				680						685				
Lys	Glu	Asn	Ser	Lys	Arg	Trp	Asn	Pro	Glu	Ile	Gln	Tyr	Thr	Ser	Asn	
	690					695					700					
Tyr	Ala	Lys	Ser	Ala	Asn	Val	Asp	Phe	Thr	Val	Asp	Asn	Asn	Gly	Val	
705					710					715					720	
Tyr	Ser	Glu	Pro	Arg	Pro	Ile	Gly	Thr	Arg	Tyr	Leu	Thr	Arg	Asn	Leu	
			725					730						735		

73

That which is claimed is:

1. A method of administering a nucleic acid to a cell comprising contacting the cell with a chimeric virus vector comprising:

a) a chimeric adeno-associated virus (AAV) capsid comprising a selective amino acid insertion immediately following amino acid position 264 in an AAV2 capsid subunit having the amino acid sequence of (SEQ ID NO:3) or a corresponding change in another AAV capsid subunit; and

b) a nucleic acid comprising an AAV terminal repeat sequence and a heterologous nucleic acid sequence, wherein the nucleic acid is packaged within the chimeric AAV capsid.

2. A method of delivering a nucleic acid to a subject comprising administering to the subject a chimeric virus vector comprising:

a) a chimeric adeno-associated virus (AAV) capsid comprising a selective amino acid insertion immediately following amino acid position 264 in an AAV2 capsid subunit having the amino acid sequence of (SEQ ID NO:3) or a corresponding change in another AAV capsid subunit; and

b) a nucleic acid comprising an AAV terminal repeat sequence and a heterologous nucleic acid sequence, wherein the nucleic acid is packaged within the chimeric AAV capsid.

3. The method of claim 2, wherein the subject is a human subject.

4. The method of claim 2, wherein the subject has or is at risk for a disorder selected from the group consisting of a muscular dystrophy including Duchenne or Becker muscular dystrophy, hemophilia A, hemophilia B, multiple sclerosis, diabetes mellitus, Gaucher disease, Fabry disease, Pompe disease, cancer, arthritis, muscle wasting, heart disease including congenital heart failure or peripheral artery disease, intimal hyperplasia, a neurological disorder including epilepsy, Huntington's disease, Parkinson's disease or Alzheimer's disease, an autoimmune disease, cystic fibrosis, thalassemia, Hurler's disease, Krabbe's disease, phenylketonuria,

74

Batten's disease, spinal cerebral ataxia, LDL receptor deficiency, hyperammonemia, anemia, arthritis, a retinal degenerative disorder including macular degeneration, adenosine deaminase deficiency, and cancer including tumor-forming cancers.

5. The method of claim 2, wherein the chimeric virus vector or pharmaceutical formulation is administered to skeletal muscle.

6. The method of claim 5, wherein the subject has or is at risk for muscular dystrophy.

7. The method of claim 5, wherein the chimeric virus vector comprises a chimeric AAV capsid comprising a selective amino acid insertion following amino acid position 264 in an AAV2 capsid subunit.

8. The method of claim 5, wherein the chimeric virus vector comprises a chimeric AAV capsid comprising a selective amino acid insertion following amino acid position 264 in an AAV3b capsid subunit.

9. The method of claim 2, wherein the chimeric virus vector or pharmaceutical formulation is administered to cardiac muscle.

10. The method of claim 9, wherein the subject has or is at risk for heart disease.

11. The method of claim 10, wherein the subject has or is at risk for congestive heart failure or peripheral artery disease.

12. The method of claim 10, wherein the chimeric virus vector comprises a chimeric AAV capsid comprising a selective amino acid insertion following amino acid position 264 in an AAV3b capsid subunit.

13. The method of claim 2, wherein the chimeric virus vector or pharmaceutical formulation is administered to glial cells and/or astrocytes.

14. The method of claim 13, wherein the subject has or is at risk for Parkinson's Disease, astrocytomas or glioblastoma.

15. The method of claim 13, wherein the chimeric virus vector comprises a chimeric AAV capsid comprising a selective amino acid insertion following amino acid position 264 in an AAV2 capsid subunit.

* * * * *

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 9,012,224 B2
APPLICATION NO. : 12/983547
DATED : April 21, 2015
INVENTOR(S) : Bowles et al.

Page 1 of 1

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

In The Specification

Column 22, Line 47: Please correct "10⁶ to/ml," to read -- 10⁶ tu/ml, --

Column 22, Line 49: Please correct "10⁸ to/ml," to read -- 10⁸ tu/ml, --

Column 22, Lines 49 and 50: Please correct "10⁹ to/ml," to read -- 10⁹ tu/ml, --

Signed and Sealed this
Nineteenth Day of January, 2016



Michelle K. Lee
Director of the United States Patent and Trademark Office