

Tumor Mutational Burden and Efficacy of Nivolumab Monotherapy and in Combination with Ipilimumab in Small-Cell Lung Cancer

Highlights

- This study evaluated the role of tumor mutational burden (TMB) in SCLC
- Efficacy of nivolumab with or without ipilimumab increases with higher TMB in SCLC
- The benefit of nivolumab plus ipilimumab is greatest in patients with high-TMB SCLC
- TMB has a potential role as a biomarker for immunotherapy across lung cancers

Authors

Matthew D. Hellmann,
Margaret K. Callahan,
Mark M. Awad, ..., Han Chang,
William J. Geese, Scott J. Antonia

Correspondence

hellmanm@mskcc.org

In Brief

Hellmann et al. evaluate the impact of tumor mutational burden on the efficacy of nivolumab monotherapy or combination with ipilimumab in patients with small-cell lung cancer (SCLC). They show that treatment efficacy and the increased benefit of the combination are most substantial in SCLC with high tumor mutational burden.



Hellmann et al., 2018, Cancer Cell 33, 853–861
May 14, 2018 © 2018 Elsevier Inc.
<https://doi.org/10.1016/j.ccell.2018.04.001>

CellPress

JHU 2087
Merck Sharp v. Johns Hopkins
IPR2024-00623

Tumor Mutational Burden and Efficacy of Nivolumab Monotherapy and in Combination with Ipilimumab in Small-Cell Lung Cancer

Matthew D. Hellmann,^{1,11,12,*} Margaret K. Callahan,^{1,11} Mark M. Awad,² Emiliano Calvo,³ Paolo A. Ascierto,⁴ Akin Atmaca,⁵ Naiyer A. Rizvi,⁶ Fred R. Hirsch,⁷ Giovanni Selvaggi,⁸ Joseph D. Szustakowski,⁹ Ariella Sasson,⁹ Ryan Golhar,⁹ Patrik Vitazka,⁹ Han Chang,⁹ William J. Geese,⁹ and Scott J. Antonia¹⁰

¹Thoracic Oncology Service, Division of Solid Tumor Oncology, Department of Medicine, Memorial Sloan Kettering Cancer Center, Weill Cornell Medical College, and Parker Center for Cancer Immunotherapy, 885 2nd Avenue, New York, NY 10017, USA

²Department of Medical Oncology, Dana-Farber Cancer Institute, Boston, MA 02215, USA

³START Madrid, Centro Integral Oncológico Clara Campal, Medical Oncology Division, Hospital Universitario Madrid Norte Sanchinarro, Madrid 28050, Spain

⁴Unit of Melanoma, Cancer Immunotherapy and Innovative Therapy, Istituto Nazionale Tumori Fondazione G. Pascale, Naples 80131, Italy

⁵Department of Hematology and Oncology, Krankenhaus Nordwest, UCT-University Cancer Center, Frankfurt am Main 60488, Germany

⁶Division of Hematology and Oncology, Columbia University Medical Center, New York, NY 10032, USA

⁷Department of Medicine, Division of Medical Oncology, University of Colorado Anschutz Medical Campus, Aurora, CO 80045, USA

⁸Clinical Development, Bristol-Myers Squibb, Princeton, NJ 08648, USA

⁹Translational Medicine, Bristol-Myers Squibb, Princeton, NJ 08648, USA

¹⁰Department of Thoracic Oncology, H. Lee Moffitt Cancer Center & Research Institute, Tampa, FL 33612, USA

¹¹These authors contributed equally

¹²Lead Contact

*Correspondence: hellmanm@mskcc.org

<https://doi.org/10.1016/j.ccell.2018.04.001>

SUMMARY

Durable responses and encouraging survival have been demonstrated with immune checkpoint inhibitors in small-cell lung cancer (SCLC), but predictive markers are unknown. We used whole exome sequencing to evaluate the impact of tumor mutational burden on efficacy of nivolumab monotherapy or combined with ipilimumab in patients with SCLC from the nonrandomized or randomized cohorts of CheckMate 032. Patients received nivolumab (3 mg/kg every 2 weeks) or nivolumab plus ipilimumab (1 mg/kg plus 3 mg/kg every 3 weeks for four cycles, followed by nivolumab 3 mg/kg every 2 weeks). Efficacy of nivolumab ± ipilimumab was enhanced in patients with high tumor mutational burden. Nivolumab plus ipilimumab appeared to provide a greater clinical benefit than nivolumab monotherapy in the high tumor mutational burden tertile.

INTRODUCTION

Small-cell lung cancer (SCLC) accounts for 10%–15% of all patients with lung cancer (Alvarado-Luna and Morales-Espinosa, 2016; Herbst et al., 2008), and approximately 75% of patients present with extensive-stage disease. Standard first-line treatment consists of platinum-based chemotherapy, but patients who progress have few effective treatment options and a

routinely poor prognosis (Alvarado-Luna and Morales-Espinosa, 2016; National Comprehensive Cancer Network, 2017). In patients with previously treated SCLC, nivolumab, a fully human immunoglobulin (Ig) G4 programmed death (PD)-1 immune checkpoint inhibitor antibody, showed durable responses and encouraging survival as monotherapy and in combination with ipilimumab, a fully human IgG1 cytotoxic T lymphocyte antigen-4 (CTLA-4) immune checkpoint inhibitor antibody (Antonia

Significance

The identification of predictors of response to immune checkpoint blockade in small-cell lung cancer (SCLC) has been elusive. Expression of tumor programmed death ligand 1, a common biomarker in non-small-cell lung cancer, is uncommon and not predictive in SCLC. We show that efficacy of nivolumab with or without ipilimumab was increased in SCLCs that have high tumor mutational burden. The differential benefit of combination immunotherapy was most substantial in patients with high tumor mutational burden. Our results suggest that tumor mutational burden has a role as a biomarker in SCLC and may help identify patients most likely to benefit from combination immunotherapy. Our findings are a critical contribution in moving toward greater precision in patient selection for treatments in lung cancer.






Figure 1. Flow Diagram of Analyzed Patients

See also Figures S1–S4, Table S1.

RESULTS

Patients and Tumor Mutational Burden

Of all treated patients ($n = 401$), 245 received nivolumab 3 mg/kg every 2 weeks and 156 received nivolumab plus ipilimumab (1 mg/kg plus 3 mg/kg every 3 weeks for four cycles, followed by nivolumab 3 mg/kg every 2 weeks) in the pooled nonrandomized or randomized SCLC cohorts of CheckMate 032 (Figures 1 and S1). Of all treated patients, 61% had sufficient paired tumor and

whole blood samples to attempt whole exome sequencing (WES). WES was successful in 86% of these patients such that overall 211 (53%) of all treated patients were evaluable for efficacy analyses by tumor mutational burden (Figure 1). Baseline patient characteristics were similar between tumor-mutational-burden-evaluable patients and all treated patients across treatment groups (Table 1). Clinical outcomes, including overall survival, progression-free survival, and objective response rate, also were similar between tumor-mutational-burden-evaluable and all treated patients (Figure S2 and Table S1). Overall, the tumor-mutational-burden-evaluable patients were representative of the total trial population.

Tumor mutational burden was defined as the total number of somatic missense mutations, and patients were divided into tertiles, as done previously (Carbone et al., 2017). Tertile boundaries were defined as: low, 0 to <143 mutations; medium, 143 to 247 mutations; and high, ≥ 248 mutations. The distribution of patients by tumor mutational burden was similar between those receiving nivolumab or nivolumab plus ipilimumab (Figure S3A), similar to the distribution seen in an independent previous report (George et al., 2015) of patients with SCLC (Figure S3B), and similar to that of patients with NSCLC treated with nivolumab in the CheckMate 026 study (Carbone et al., 2017) but with a narrower range (Figure S3B). Tumor mutational burden estimated by WES closely correlated with the estimate after *in silico* filtering to the 315 genes in the FoundationOne next-generation sequencing genomic profile (Frampton et al., 2013) (Figure S4).

PD-L1 expression $\geq 1\%$ was uncommon overall, but was evenly distributed between the mutational burden high and medium/low tertiles. There was no association between PD-L1 expression and tumor mutational burden (Table S2) or between PD-L1 expression and objective response rate in the nonrandomized population (Table S3).

The identification of predictors of response to immune checkpoint blockade in SCLC has been elusive. Unlike in some other cancers, including non-small-cell lung cancer (NSCLC) (Reck et al., 2016), tumor PD ligand 1 (PD-L1) expression is uncommon in SCLC ($\sim 18\%$), and responses to nivolumab monotherapy or in combination with ipilimumab have been observed regardless of PD-L1 expression (Antonia et al., 2016). Given the initial signals of benefit with the combination of nivolumab and ipilimumab in SCLC but the potential for greater toxicity compared with nivolumab monotherapy, determining the predictors of response to monotherapy and/or combination therapy is critical (Antonia et al., 2016).

SCLC is characterized by high somatic mutational burden (George et al., 2015; Peifer et al., 2012; Rudin et al., 2012) due to the nearly universal association of SCLC with smoking (Alexandrov et al., 2013; Byers and Rudin, 2015); mutations can affect diverse pathways, including DNA repair mechanisms (Gazdar et al., 2017). Data in other solid tumors, including urothelial carcinoma, melanoma, and NSCLC, have shown an association between high tumor mutational burden and improved efficacy of immune checkpoint blockade monotherapy (Carbone et al., 2017; Galsky et al., 2017; Le et al., 2017; Rizvi et al., 2015; Rosenberg et al., 2016; Snyder et al., 2014); however, this has not previously been examined in patients with SCLC or in patients treated with dual checkpoint blockade. We hypothesized that tumor mutational burden could be associated with efficacy of nivolumab alone and in combination with ipilimumab in previously treated SCLC.

PD-L1 expression $\geq 1\%$ was uncommon overall, but was evenly distributed between the mutational burden high and medium/low tertiles. There was no association between PD-L1 expression and tumor mutational burden (Table S2) or between PD-L1 expression and objective response rate in the nonrandomized population (Table S3).

PD-L1 expression $\geq 1\%$ was uncommon overall, but was evenly distributed between the mutational burden high and medium/low tertiles. There was no association between PD-L1 expression and tumor mutational burden (Table S2) or between PD-L1 expression and objective response rate in the nonrandomized population (Table S3).

PD-L1 expression $\geq 1\%$ was uncommon overall, but was evenly distributed between the mutational burden high and medium/low tertiles. There was no association between PD-L1 expression and tumor mutational burden (Table S2) or between PD-L1 expression and objective response rate in the nonrandomized population (Table S3).

Association of Tumor Mutational Burden and Clinical Outcomes

Within both the nivolumab monotherapy and nivolumab plus ipilimumab treatment groups, objective response rates were higher in those patients with high tumor mutational burden (21.3% and

Table 1. Baseline Characteristics

Characteristic	Nivolumab		Nivolumab Plus Ipilimumab	
	All Treated (n = 245)	Tumor-Mutational-Burden-Evaluable (n = 133)	All Treated (n = 156)	Tumor-Mutational-Burden-Evaluable (n = 78)
Age (years), median (range)	63 (29–83)	63 (29–83)	65 (37–91)	65 (37–80)
Age ≥ 75 years, no. (%)	23 (9)	11 (8)	17 (11)	11 (14)
Male, no. (%)	147 (60)	79 (59)	95 (61)	52 (67)
Smoking status, no. (%)				
Current/former smoker	230 (94)	126 (95)	147 (94)	73 (94)
Never smoker	13 (5)	6 (5)	8 (5)	5 (6)
Unknown	2 (1)	1 (1)	1 (1)	0
ECOG ^a performance status score, no. (%)				
0	73 (30)	42 (32)	49 (31)	23 (29)
1	171 (70)	91 (68)	106 (68)	54 (69)
2	1 (<1)	0	0	0
Unknown	0	0	1 (0.6)	1 (1.3)
Tumor PD-L1 expression, no. (%)				
≥ 1%	24 (10)	17 (13)	19 (12)	8 (10)
<1%	149 (61)	89 (67)	90 (58)	51 (65)
Unknown	72 (29)	27 (20)	47 (30)	19 (24)
Study cohort, no. (%)				
Nonrandomized	98 (40)	50 (38)	61 (39)	25 (32)
Randomized	147 (60)	83 (62)	95 (61)	53 (68)

^aECOG denotes Eastern Cooperative Oncology Group.

46.2%, respectively) than in patients with low (4.8% and 22.2%, respectively) or medium (6.8% and 16.0%, respectively) tumor mutational burden (Figure 2). Within all tumor mutational burden tertiles, the objective response rates were higher among patients treated with nivolumab plus ipilimumab than among those treated with nivolumab monotherapy (Figure 2). Similar associations were seen when tumor mutational burden was grouped by medians or quartile boundaries (Figures S5A and S5B).

We also examined how the association between objective response and tumor mutational burden in SCLC compared with the similar association in NSCLC. The receiver operating characteristics curves of the associations between tumor mutational burden and objective response to nivolumab were similar

in patients with SCLC and NSCLC, suggesting that tumor mutational burden is a predictor of response to nivolumab in both patient populations (Figures S5C and S5D).

Tumor mutational burden was higher among patients with a complete or partial response to either monotherapy or combination therapy than among those with stable disease or progressive disease (Figure 3).

In both treatment groups, estimated 1-year progression-free survival rates were higher in the high tumor mutational burden group (21.2% and 30.0% for nivolumab monotherapy and nivolumab plus ipilimumab, respectively) compared with the low (not calculable and 6.2%, respectively) or medium (3.1% and 8.0%, respectively) tumor mutational burden groups (Figure 4). Similar trends




Figure 2. Objective Response Rate by Tumor Mutational Burden Tertile
See also Figure S5 and Tables S2 and S3.




Figure 3. Tumor Mutational Burden by Best Overall Response in Individual Patients

Black lines in each box denote the median. The bottom and top of each box denote the first and third quartile, respectively. The lower whisker denotes the value at 1.5 times the interquartile range below the 25th percentile or the minimum value of the dataset, whichever value is larger. The upper whisker denotes the value at 1.5 times the interquartile range above the 75th percentile or the maximum value of the dataset, whichever value is smaller.

were observed for overall survival (Figure 5). Within each treatment group, the estimated 1-year overall survival rate was higher in the high tumor mutational burden group (35.2% and 62.4% for nivolumab monotherapy and nivolumab plus ipilimumab, respectively) than in the low (22.1% and 23.4%, respectively) or medium (26.0% and 19.6%, respectively) tumor mutational burden groups. In patients with high tumor mutational burden, 1-year progression-free survival and 1-year overall survival rates were higher with combination therapy than with nivolumab monotherapy; no substantial differences in progression-free and overall survival between treatment groups were observed among patients with lower tumor mutational burden (Figures 4 and 5).

Lastly, we investigated whether there was a prognostic association between tumor mutational burden and overall survival in an independent cohort of patients with SCLC who did not receive immunotherapy (George et al., 2015). We found that there was no prognostic difference in survival based on tumor mutational burden tertile (Figure S6). These data suggest that tumor mutational burden is predictive of improved outcomes specifically in the context of immunotherapy, rather than more generally prognostic in patients with SCLC.

DISCUSSION

This report evaluated the role of tumor mutational burden in SCLC, and its association with outcomes with dual immune checkpoint blockade. We found that patients with high tumor

mutational burden treated with either nivolumab monotherapy or nivolumab plus ipilimumab had improved efficacy compared with those with medium or low tumor mutational burden. Similar to what has been seen in previous reports of nivolumab in NSCLC and urothelial carcinoma (Carbone et al., 2017; Galsky et al., 2017), and ipilimumab in melanoma (Snyder et al., 2014), tumor mutational burden may be a relevant biomarker in SCLC.

The current study suggests that tumor mutational burden may help inform the benefit/risk ratio of nivolumab monotherapy or nivolumab plus ipilimumab for a given patient. Given the initial observation of increased benefit of combination therapy with nivolumab and ipilimumab in SCLC, but also of potentially greater toxicity compared with nivolumab alone, identifying the differential predictors of response to monotherapy or combination therapy is critical (Antonia et al., 2016). Specifically, patients with SCLC and high tumor mutational burden treated with nivolumab plus ipilimumab had outcomes that surpass historical survival expectations for patients with previously treated SCLC. Conversely, although patients with low or medium tumor mutational burden had increased objective response rates with nivolumab plus ipilimumab compared with nivolumab monotherapy, there were no substantial differences between the treatment groups in progression-free or overall survival. Together, these observations suggest a benefit of the combination over nivolumab monotherapy for patients with high tumor mutational burden, while nivolumab alone may be a favorable option for those with medium or low tumor mutational burden. However,




Figure 4. Progression-Free Survival by Treatment and Tumor Mutational Burden Tertile
CI denotes confidence interval.




Figure 5. Overall Survival by Treatment and Tumor Mutational Burden Tertile
CI denotes confidence interval. See also Figure S6.

this was an exploratory analysis and these findings require further evaluation in larger randomized studies.

It was not a foregone conclusion that tumor mutational burden would be informative in SCLC. Given the nearly universal association with smoking in patients with SCLC, tumor mutational burden is high but the range is relatively narrow (Alexandrov et al., 2013). It was not known whether there would be sufficient molecular diversity to identify subgroups with distinct clinical responses to immunotherapy; nevertheless, clinical benefit with immunotherapy was greatest among those with the highest mutational burden. Furthermore, it was speculated that tumor mutational burden analysis may not be feasible given the small biopsies used and necrotic tissue found in SCLC tumors. In this retrospective analysis, 61% of patients had sufficient tumor biopsy material and whole blood sample for WES. As the WES analysis was not pre-planned and tissue requirements for eligibility were relatively minimal, this success rate is perhaps better than expected. In addition, in the vast majority (86%) of these patients, WES and subsequent determination of tumor mutational burden were successful. Overall, this analysis demonstrates that molecular analysis of SCLC for testing of tumor mutational burden is feasible and we would expect that prospective efforts may achieve even higher success rates.

It was also unknown how tumor mutational burden would associate with outcomes in patients treated with nivolumab plus ipilimumab. One hypothesis was that the addition of ipilimumab would broaden the repertoire of anti-tumor T cell clones and diminish the predictive relevance of tumor mutational burden. That does not appear to be the case in SCLC, as tumor mutational burden appears to be a strong biomarker for nivolumab plus ipilimumab. These data may provide insight into molecular determinants of response to combination therapy that may have implications for use in other cancers. Of note, and consistent with the observation in this report, improvement of response to nivolumab plus ipilimumab with high tumor mutational burden also has been observed in patients with NSCLC (Hellmann et al., 2018 [this issue of *Cancer Cell*]). Further work is needed to fully understand the precise immunobiologic mechanisms underlying synergy between PD-1 and CTLA-4 blockade (Wei et al., 2017).

This study is an exploratory analysis and the results presented here will need to be confirmed in larger, prospective datasets. Although WES was performed retrospectively, the clinical and outcome features of the tumor-mutational-burden-evaluable cohort were similar to those of all treated patients and generally typical of patients with SCLC. Inclusion of a control arm would further confirm the predictive (rather than prognostic) nature of tumor mutational burden, although no association of tumor mutational burden and survival was seen in previously published data of SCLC not treated with immunotherapy (George et al., 2015). The association of improved outcomes with higher tumor mutational burden was consistently seen when patients were divided by median, tertile, or quartile. This observation highlights the stability of the association between mutational burden and benefit with immunotherapy in SCLC but also suggests that there is not a singular cut point to enrich for benefit. Further validation and optimization of tumor mutational burden cutoff are warranted, in addition to efforts to refine our understanding of the somatic molecular features that most contribute to immunogenicity (Turajlic et al., 2017). Two ongoing phase 3 trials evalu-

ating the efficacy of nivolumab or nivolumab plus ipilimumab in SCLC (CheckMate 331, NCT02481830, and CheckMate 451, NCT02538666) will assess the association between tumor mutational burden and treatment outcomes in larger datasets.

In conclusion, in patients with SCLC treated with either nivolumab monotherapy or nivolumab plus ipilimumab, efficacy was enhanced among patients with high tumor mutational burden. In the high tumor mutational burden subgroup, the combination of nivolumab and ipilimumab appeared to provide a greater clinical benefit compared with nivolumab monotherapy, with near doubling of estimated 1-year survival rates for the combination versus monotherapy. The progression-free and overall survival seen among patients with high tumor mutational burden treated with nivolumab plus ipilimumab were particularly remarkable for patients with previously treated SCLC. These data, along with existing data on NSCLC (Carbone et al., 2017), suggest that tumor mutational burden has a potential role as a biomarker for immunotherapy across lung cancers.

STAR★METHODS

Detailed methods are provided in the online version of this paper and include the following:

- KEY RESOURCES TABLE
- CONTACT FOR REAGENT AND RESOURCE SHARING
- EXPERIMENTAL MODEL AND SUBJECT DETAILS
 - Patients
 - Trial Design and Treatment
 - Endpoints and Assessments
 - Study Oversight
- METHOD DETAILS
 - PD-L1 Expression
 - Exploratory Biomarker Analysis of Tumor Mutational Burden
 - Molecular Features of Previously Published Small-Cell Lung Cancers
- QUANTIFICATION AND STATISTICAL ANALYSIS
- DATA AND SOFTWARE AVAILABILITY
- ADDITIONAL RESOURCES

SUPPLEMENTAL INFORMATION

Supplemental Information includes six figures and three tables and can be found with this article online at <https://doi.org/10.1016/j.ccell.2018.04.001>.

ACKNOWLEDGMENTS

We thank the patients and their families, as well as the participating study teams. Denmark: Ulrik Lassen (Rigshospitalet); Finland: Petri Bono (Helsinki University Hospital); Germany: Akin Atmaca (Krankenhaus Nordwest GmbH, Frankfurt), Peter Brossart (Universitätsklinikum Bonn), Dirk Jäger (Nationales Centrum für Tumorerkrankungen Heidelberg); Italy: Andrea Ardizzone (Azienda Ospedaliera S. Orsola-Malpighi), Paolo Antonio Ascierto (Istituto Nazionale Tumori Fondazione G. Pascale), Filippo De-Braud (Istituto Nazionale Tumori), Valentina Guarneri (Istituto Oncologico Veneto IOV-IRCCS); Spain: Emiliano Calvo (Hospital De Madrid Norte Sanchinarro), José A. López-Martín (Hospital Universitario 12 De Octubre), Victor Moreno García (Fundación Jiménez Díaz), Noemi Reguart (Hospital Clinic I Provincial); United Kingdom: Ian Chau (Royal Marsden Hospital), Jeff Evans (The Beatson West of Scotland Cancer Centre); United States: Asim Amin (Levine Cancer Institute), Johanna Bendell (Tennessee Oncology PLLC), Scott Antonia (H. Lee Moffitt Cancer Center & Research

Institute, Inc.), Margaret Callahan (Memorial Sloan Kettering Cancer Center), David Chism (Vanderbilt-Ingram Cancer Center), Joseph Eder (Yale University), Jennifer Diamond (University of Colorado Hospital), Ralph Hauke (Nebraska Cancer Specialists), Frederic Kaye (University of Florida), Dung Le (Sidney Kimmel Comprehensive Cancer Center at Johns Hopkins), Michael Morse (Duke Cancer Institute), Rathil Pillai (Emory University – Winship Cancer Institute), Patrick Ott (Dana-Farber Cancer Institute, Massachusetts General Hospital, Brigham & Women's Hospital), Jeffrey Schneider (New York University Winthrop Hospital), Padmanee Sharma (The University of Texas MD Anderson Cancer Center), and Matthew Taylor (Oregon Health & Science University); Dako for collaborative development of the PD-L1 IHC 28-8 pharmDx assay; Andrea Renninger, Rupa Guduru, Marsha Smith, George Green, and Steve Lane for their contributions to experimental design, downstream analysis, and interpretation of the tumor mutational burden data; Peter Szabo for validating data analysis; and Ana Moreno and Soumaya Bendahmane for serving as the protocol managers. Medical writing and editorial assistance was provided by Stefanie Puglielli, PhD, and Beth Burke, PhD, CMPP, of Evidence Scientific Solutions, with funding from Bristol-Myers Squibb. This study was funded by Bristol-Myers Squibb (Princeton, NJ) and ONO Pharmaceutical Company, Ltd. (Osaka, Japan).

AUTHOR CONTRIBUTIONS

Conception or design, S.J.A., M.K.C., H.C., W.J.G., R.G., M.D.H., F.R.H., A.S., G.S., J.D.S., and P.V.; data acquisition, S.J.A., P.A.A., A.A., M.M.A., M.K.C., E.C., M.D.H., and N.A.R.; data analysis/interpretation, all authors; manuscript writing, all authors; final approval of manuscript, all authors.

DECLARATION OF INTERESTS

M.D.H. reports paid consultancy from AstraZeneca/MedImmune, Bristol-Myers Squibb, Genentech, Janssen, Merck, Mirati Therapeutics, Novartis, and Shattuck Labs, research funding from Bristol-Myers Squibb, and a patent filed by Memorial Sloan Kettering related to the use of tumor mutational burden to predict response to immunotherapy (PCT/US2015/062208). M.K.C. reports grants from and employment of a family member by Bristol-Myers Squibb and personal fees for an advisory/consulting role from AstraZeneca/MedImmune, Incyte, Moderna, and Merck. M.M.A. reports a consulting or advisory role for AbbVie, AstraZeneca, Boehringer Ingelheim, Genentech, Merck, and Pfizer. E.C. is employed by START and HM Hospitals Group, where he holds a leadership position. He has ownership at START, Oncoart Associated and International Cancer Consultants. He receives honoraria from HM Hospitals Group, and he is the founder and president of the NPO foundation INTHEOS (Investigational Therapeutics in Oncological Sciences). In the last 2 years, he has been a paid consultant or advisor to Novartis, Nanobiotix, Janssen-Cilag, PsiOxus Therapeutics, Seattle Genetics, EUSA Pharma, Abbvie, Celgene, AstraZeneca, Guidepoint Global, Roche/Genentech, GLG, Pfizer, Servier, and Amcure. He is part of a Speakers' Bureau to Novartis. He has received research funding from AstraZeneca, Novartis, BeiGene, and START, and he has received travel financial support from Roche/Genentech. P.A.A. reports grants for research funding and personal fees for an advisory/consultant role from Array, Bristol-Myers Squibb, and Roche-Genentech. He reports personal fees for an advisory/consultant role from Genmab, Incyte, MedImmune, Merck Serono, MSD, NewLink Genetics, Novartis, Pierre Fabre, AstraZeneca, Syndax, Idera, Sanofi, Sun Pharma, Ultimovacs, Sandoz, and Immunocore. He also received travel support from MSD. A.A. reports travel grants and honoraria for advisory board participation from Bristol-Myers Squibb, Merck Sharp & Dohme, and Roche. N.A.R. reports stock or other ownership from ARMO, Gritstone, and Bellicum. He has a consulting or advisory role for Abbvie, AstraZeneca, Bristol-Myers Squibb, Eli Lilly, EMD Serono, Merck, Neogenomics, Novartis, Pfizer, Regeneron, and Roche. He reports a patent filed by Memorial Sloan Kettering (PCT/US2015/062208) for the use of tumor mutation burden for prediction of immunotherapy efficacy, which is licensed to Personal Genome Diagnostics. F.R.H. has participated in scientific advisory boards for Bristol-Myers Squibb, Merck, Genentech/Roche, Ventana, AstraZeneca, Novartis, Bayer, Loxo, Biocept, HTG, Pfizer, Abbvie, and Helsinn and received research funding (through University of Colorado) from HTG, Bayer, Clovis, Merck, Amgen, Rain Therapeutics, and Genentech. G.S. is employed by

Bristol-Myers Squibb. In addition, he is an inventor on one or more pending patent applications for TMB as a predictive biomarker for immunotherapy. J.D.S. is employed by (and holds shares with) Bristol-Myers Squibb and reports previous employment and stock ownership from Novartis. In addition, he is an inventor on one or more pending patent applications for TMB as a predictive biomarker for immunotherapy. A.S. reports employment by Bristol-Myers Squibb. R.G. reports employment by Bristol-Myers Squibb. P.V. reports employment by Bristol-Myers Squibb. H.C. is employed by Bristol-Myers Squibb. Additionally, he is an inventor on one or more pending patent applications for TMB as a predictive biomarker for immunotherapy. W.J.G. is employed by (and holds shares with) Bristol-Myers Squibb. Additionally, he is an inventor on one or more pending patent applications for TMB as a predictive biomarker for immunotherapy. S.J.A. reports stock or other ownership from Cellular Biomedicine Group and honoraria, a consulting or advisory role, travel, accommodations, and expenses from AstraZeneca, Boehringer Ingelheim, Bristol-Myers Squibb, and Merck. The other authors have no conflicts of interest to declare.

Received: December 14, 2017

Revised: February 12, 2018

Accepted: April 2, 2018

Published: May 3, 2018; corrected online: February 1, 2019

REFERENCES

- Alexandrov, L.B., Nik-Zainal, S., Wedge, D.C., Aparicio, S.A., Behjati, S., Biankin, A.V., Bignell, G.R., Bolli, N., Borg, A., Børresen-Dale, A.L., et al. (2013). Signatures of mutational processes in human cancer. *Nature* 500, 415–421.
- Alvarado-Luna, G., and Morales-Espinosa, D. (2016). Treatment for small cell lung cancer, where are we now?—a review. *Transl. Lung Cancer Res.* 5, 26–38.
- Antonia, S.J., López-Martin, J.A., Bendell, J., Ott, P.A., Taylor, M., Eder, J.P., Jäger, D., Pietanza, M.C., Le, D.T., de Braud, F., et al. (2016). Nivolumab alone and nivolumab plus ipilimumab in recurrent small-cell lung cancer (CheckMate 032): a multicentre, open-label, phase 1/2 trial. *Lancet Oncol.* 17, 883–895.
- Bamford, S., Dawson, E., Forbes, S., Clements, J., Pettett, R., Dogan, A., Flanagan, A., Teague, J., Futreal, P.A., Stratton, M.R., and Wooster, R. (2004). The COSMIC (catalogue of somatic mutations in cancer) database and website. *Br. J. Cancer* 91, 355–358.
- Byers, L.A., and Rudin, C.M. (2015). Small cell lung cancer: where do we go from here? *Cancer* 121, 664–672.
- Carbone, D.P., Reck, M., Paz-Ares, L., Creelan, B., Horn, L., Steins, M., Felip, E., van den Heuvel, M.M., Ciuleanu, T.E., Badin, F., et al. (2017). First-line nivolumab in stage IV or recurrent non-small-cell lung cancer. *N. Engl. J. Med.* 376, 2415–2426.
- Cibulskis, K., Lawrence, M.S., Carter, S.L., Sivachenko, A., Jaffe, D., Sougnez, C., Gabriel, S., Meyerson, M., Lander, E.S., and Getz, G. (2013). Sensitive detection of somatic point mutations in impure and heterogeneous cancer samples. *Nat. Biotechnol.* 31, 213–219.
- Cingolani, P., Platts, A., Wang, L., Coon, M., Nguyen, T., Wang, L., Land, S.J., Lu, X., and Ruden, D.M. (2012). A program for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the genome of *Drosophila melanogaster* strain w1118; iso-2; iso-3. *Fly (Austin)* 6, 80–92.
- Eisenhauer, E.A., Therasse, P., Bogaerts, J., Schwartz, L.H., Sargent, D., Ford, R., Dancey, J., Arbuck, S., Gwyther, S., Mooney, M., et al. (2009). New response evaluation criteria in solid tumours: revised RECIST guideline (version 1.1). *Eur. J. Cancer* 45, 228–247.
- Fisher, S., Barry, A., Abreu, J., Minie, B., Nolan, J., Delorey, T.M., Young, G., Fennell, T.J., Allen, A., Ambrogio, L., et al. (2011). A scalable, fully automated process for construction of sequence-ready human exome targeted capture libraries. *Genome Biol.* 12, R1.
- Frampton, G.M., Fichtenholtz, A., Otto, G.A., Wang, K., Downing, S.R., He, J., Schnall-Levin, M., White, J., Sanford, E.M., An, P., et al. (2013). Development and validation of a clinical cancer genomic profiling test based on massively parallel DNA sequencing. *Nat. Biotechnol.* 31, 1023–1031.

- Galsky, M.D., Sazi, A., Szabo, P.M., Azrilevich, A., Horak, C., Lambert, A., Siefker-Radtke, A., Necchi, A., and Sharma, P. (2017). Impact of tumor mutation burden on nivolumab efficacy in second-line urothelial carcinoma patients: exploratory analysis of the phase II CheckMate 275 study, [abstract 848PD]. *Ann. Oncol.* 28 (suppl 5), v295–v329.
- Gazdar, A.F., Bunn, P.A., and Minna, J.D. (2017). Small-cell lung cancer: what we know, what we need to know and the path forward. *Nat. Rev. Cancer* 17, 725–737.
- George, J., Lim, J.S., Jang, S.J., Cun, Y., Ozretić, L., Kong, G., Leenders, F., Lu, X., Fernández-Cuesta, L., Bosco, G., et al. (2015). Comprehensive genomic profiles of small cell lung cancer. *Nature* 524, 47–53.
- Hellmann, M.D., Antonia, S.J., Ponce, S., Ott, P.A., Calvo, E., Taylor, M., Ready, N., Hann, C.L., De Braud, F., Eder, J.P., et al. (2016). Nivolumab alone or with ipilimumab in recurrent small cell lung cancer (SCLC): 2-year survival and updated analyses from the CheckMate 032 trial, [abstract MA09.05]. *J. Thorac. Oncol.* 12, S393–S394.
- Hellmann, M.D., Nathanson, T., Rizvi, H., Creelan, B.C., Sanchez-Vega, F., Ahuja, A., Ni, A., and Novik, J.B. (2018). Genomic features of response to combination immunotherapy in patients with advanced non-small cell lung cancer. *Cancer Cell*. <https://doi.org/10.1016/j.ccell.2018.03.018>.
- Hellmann, M.D., Ott, P.A., Zugazagoitia, J., Ready, N.E., Hann, C.L., De Braud, F.G., Antonia, S.J., Ascierto, P.A., Moreno, V., Atmaca, A., et al. (2017). Nivolumab (nivo) ± ipilimumab (ipi) in advanced small-cell lung cancer (SCLC): first report of a randomized expansion cohort from CheckMate 032, [abstract 8503]. *J. Clin. Oncol.* 35 (suppl 15), 8503.
- Herbst, R.S., Heymach, J.V., and Lippman, S.M. (2008). Lung cancer. *N. Engl. J. Med.* 359, 1367–1380.
- Le, D.T., Durham, J.N., Smith, K.N., Wang, H., Bartlett, B.R., Aulakh, L.K., Lu, S., Kemberling, H., Wilt, C., Luber, B.S., et al. (2017). Mismatch repair deficiency predicts response of solid tumors to PD-1 blockade. *Science* 357, 409–413.
- Li, H. (2013). Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. *arXiv*, 1303.3997v1 [q-bio.GN].
- Mills, R.E., Luttig, C.T., Larkins, C.E., Beauchamp, A., Tsui, C., Pittard, W.S., and Devine, S.E. (2006). An initial map of insertion and deletion (INDEL) variation in the human genome. *Genome Res.* 16, 1182–1190.
- National Comprehensive Cancer Network (2017). Clinical practice guidelines in Oncology. Small cell lung cancer. Version 1.2018. https://www.nccn.org/professionals/physician_gls/default.aspx
- Peifer, M., Fernández-Cuesta, L., Sos, M.L., George, J., Seidel, D., Kasper, L.H., Plenker, D., Leenders, F., Sun, R., Zander, T., et al. (2012). Integrative genome analyses identify key somatic driver mutations of small-cell lung cancer. *Nat. Genet.* 44, 1104–1110.
- Reck, M., Rodríguez-Abreu, D., Robinson, A.G., Hui, R., Csőszi, T., Fülöp, A., Gottfried, M., Peled, N., Tafreshi, A., Cuffe, S., et al. (2016). Pembrolizumab versus chemotherapy for PD-L1-positive non-small-cell lung cancer. *N. Engl. J. Med.* 375, 1823–1833.
- Rizvi, N.A., Hellmann, M.D., Snyder, A., Kvistborg, P., Makarov, V., Havel, J.J., Lee, W., Yuan, J., Wong, P., Ho, T.S., et al. (2015). Mutational landscape determines sensitivity to PD-1 blockade in non-small cell lung cancer. *Science* 348, 124–128.
- Rosenberg, J.E., Hoffman-Censits, J., Powles, T., van der Heijden, M.S., Balar, A.V., Necchi, A., Dawson, N., O'Donnell, P.H., Balmanoukian, A., Loriot, Y., et al. (2016). Atezolizumab in patients with locally advanced and metastatic urothelial carcinoma who have progressed following treatment with platinum-based chemotherapy: a single-arm, multicentre, phase 2 trial. *Lancet* 387, 1909–1920.
- Rudin, C.M., Durinck, S., Stawiski, E.W., Poirier, J.T., Modrusan, Z., Shames, D.S., Bergbower, E.A., Guan, Y., Shin, J., Guillory, J., et al. (2012). Comprehensive genomic analysis identifies SOX2 as a frequently amplified gene in small-cell lung cancer. *Nat. Genet.* 44, 1111–1116.
- Saunders, C.T., Wong, W.S., Swamy, S., Becq, J., Murray, L.J., and Cheetham, R.K. (2012). Strelka: accurate somatic small-variant calling from sequenced tumor-normal sample pairs. *Bioinformatics* 28, 1811–1817.
- Snyder, A., Makarov, V., Merghoub, T., Yuan, J., Zaretsky, J.M., Desrichard, A., Walsh, L.A., Postow, M.A., Wong, P., Ho, T.S., et al. (2014). Genetic basis for clinical response to CTLA-4 blockade in melanoma. *N. Engl. J. Med.* 371, 2189–2199.
- Turajlic, S., Litchfield, K., Xu, H., Rosenthal, R., McGranahan, N., Reading, J.L., Wong, Y.N.S., Rowan, A., Kanu, N., Al Bakir, M., et al. (2017). Insertion-and-deletion-derived tumour-specific neoantigens and the immunogenic phenotype: a pan-cancer analysis. *Lancet Oncol.* 18, 1009–1021.
- Wang, P.P., Parker, W.T., Branford, S., and Schreiber, A.W. (2016). BAM-matcher: a tool for rapid NGS sample matching. *Bioinformatics* 32, 2699–2701.
- Weber, J.A., Aldana, R., Gallagher, B.D., and Edwards, J.S. (2016). Sentieon DNA pipeline for variant detection - software-only solution, over 20x faster than GATK 3.3 with identical results. *PeerJ PrePrints* 4, e1672.
- Wei, S.C., Levine, J.H., Cogdill, A.P., Zhao, Y., Anang, N.A.S., Andrews, M.C., Sharma, P., Wang, J., Wargo, J.A., Pe'er, D., and Allison, J.P. (2017). Distinct cellular mechanisms underlie anti-CTLA-4 and anti-PD-1 checkpoint blockade. *Cell* 170, 1120–1133.

STAR★METHODS

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Rabbit anti-human PD-L1 antibody	Epitomics Inc, Burlingame, CA	Clone 28-8
Critical Commercial Assays		
Agilent/Dako PD-L1 IHC 28-8 pharmDx	Agilent Technologies, Santa Clara, CA	Code SK005
Allprep DNA/RNA kit	Qiagen, Hilden, Germany	Cat# 80234
QIAamp DNA Blood Midi Kit	Qiagen, Hilden, Germany	Cat# 51185
Agilent SureSelectXT reagent kit	Agilent Technologies, Santa Clara, CA	Cat# 5500-0133
Tapestation	Agilent Technologies, Santa Clara, CA	RRID:SCR_014994
FoundationOne comprehensive genomic profile	Foundation Medicine, Inc, Cambridge, MA	
Deposited Data		
VCF files containing somatic snvs per Tumor-Normal pair	Sentieon's TNSnv software	European Variation Archive Accession Number: PRJEB25807
VCF files containing somatic snvs per Tumor-Normal pair	Strelka software	European Variation Archive Accession Number: PRJEB25808
Oligonucleotides		
SureSelect All Exon v5 bait	Agilent Technologies, Santa Clara, CA	Cat# 5190-6210
Software and Algorithms		
Sentieon's somatic mutation pipeline, version 201704.01	Sentieon, Inc., Mountain View, CA	
The Broad Institute's best practices for somatic mutation calling with MuTect1.3	https://software.broadinstitute.org/gatk/best-practices/	
BWA (version 0.7.15)	Sentieon, Inc., Mountain View, CA	
Sentieon's TNSnv (reimplementation of Mutect1)	Sentieon, Inc., Mountain View, CA	
Strelka (v1.0.15)	https://sites.google.com/site/strelkasomaticvariantcaller/home	
snpEff (v4.1c)	http://snpeff.sourceforge.net/index.html	
Mills' insertions and deletions	https://software.broadinstitute.org/gatk/download/bundle	
Other		
Agencourt AMPure XP beads	Beckman Coulter, Indianapolis, IN	Cat# A63882

CONTACT FOR REAGENT AND RESOURCE SHARING

Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contact, Matthew D. Hellmann, (hellmanm@mskcc.org).

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Patients

The SCLC cohort of this phase 1/2 multicenter, multi-arm, open-label trial included patients with histologically or cytologically confirmed, limited- or extensive-stage SCLC with progression after at least one platinum-based chemotherapy regimen (Figure S1) (Antonia et al., 2016). Inclusion and exclusion criteria have been previously described. Briefly, eligible patients were aged 18 years or older and had an Eastern Cooperative Oncology Group performance status of 0 or 1. Patients were eligible irrespective of PD-L1 expression and platinum sensitivity (relapse ≥ 90 days after chemotherapy) or resistance (relapse < 90 days after or during chemotherapy). Baseline or on-study tumor biopsy was required for biomarker analyses; four patients included in this analysis did not have

baseline tissue but had on-study biopsy that was used for tumor mutational burden testing. Written informed consent was collected from all patients prior to enrollment.

Trial Design and Treatment

Patients analyzed here include those treated as part of both the initial nonrandomized and subsequent randomized SCLC cohorts. Patients received nivolumab (3 mg/kg intravenously every 2 weeks until disease progression or unacceptable toxicity) or nivolumab plus ipilimumab (1 mg/kg plus 3 mg/kg intravenously every 3 weeks for four cycles, followed by nivolumab 3 mg/kg every 2 weeks until disease progression or unacceptable toxicity). Patients were permitted to receive therapy beyond Response Evaluation Criteria In Solid Tumors (RECIST) v1.1–defined progression ([Eisenhauer et al., 2009](#)) if protocol-defined criteria were met.

Here, we report an exploratory analysis of the SCLC cohort based on a database lock of March 30, 2017. Median follow-up times in the nonrandomized cohort were 23.3 months for nivolumab ($n = 98$) and 28.6 months for nivolumab plus ipilimumab ($n = 61$); median follow-up times in the randomized cohort were 10.8 months for nivolumab ($n = 147$) and 11.2 months for nivolumab plus ipilimumab ($n = 95$). Patients in the nonrandomized cohort who were treated with nivolumab 3 mg/kg plus ipilimumab 1 mg/kg ($n = 54$) were not included in the March 2017 database lock (because this regimen was not chosen for subsequent development) and therefore were not available for this analysis.

Endpoints and Assessments

The primary endpoint of the trial was the objective response rate, defined as the proportion of patients with a best overall response of complete response or partial response with nivolumab plus ipilimumab versus nivolumab monotherapy, per RECIST v1.1, as assessed by a blinded, independent central review. Secondary endpoints included overall survival and progression-free survival.

Tumor assessments by radiographic imaging (computed tomography and magnetic resonance imaging) were performed at baseline, every 6 weeks for the first 24 weeks, and every 12 weeks thereafter until disease progression (investigator-assessed per RECIST v1.1) or treatment discontinuation. Survival was monitored continuously while patients were on treatment and every 3 months after treatment discontinuation until death or conclusion of study.

Study Oversight

The study was designed by the academic authors in collaboration with the sponsor (Bristol-Myers Squibb); the sponsor worked jointly with the investigators to collect and analyze data. The study protocol was approved by an institutional review board or ethics committee at each of the participating centers (all participating centers listed in the Acknowledgements). The study was conducted in accordance with the provisions of the Declaration of Helsinki and Good Clinical Practice guidelines as defined by the International Conference on Harmonisation. An independent data and safety monitoring committee provided oversight of safety and efficacy. This study is registered with [ClinicalTrials.gov](#), number NCT01928394.

All authors attest that the study was conducted in accordance with the protocol and vouch for the accuracy and completeness of the data and analyses. All authors signed a confidentiality agreement with the sponsor. Medical writing support, including writing of the first draft, was provided by Beth Burke, Ph.D., CMPP, and Stefanie Puglielli, Ph.D., of Evidence Scientific Solutions, with funding from the sponsor.

METHOD DETAILS

PD-L1 Expression

Tumor PD-L1 expression was retrospectively assessed in pretreatment (archival or fresh) tumor biopsy specimens using an automated immunohistochemical assay (Dako North America, Carpinteria, CA) and a rabbit anti-human PD-L1 antibody (clone 28-8; Epitomics Inc, Burlingame, CA).

Exploratory Biomarker Analysis of Tumor Mutational Burden

Whole Exome Capture and Sequencing

DNA was isolated from fresh or archival formalin-fixed, paraffin-embedded tumor tissue using the Allprep DNA/RNA kit (Qiagen, Hilden, Germany). Germline DNA from whole blood was isolated using the QIAamp DNA Blood Midi Kit (Qiagen, Hilden, Germany) following the manufacturer's instructions.

Genomic DNA (150 ng) was used for library preparation. Genomic DNA was fragmented to approximately 150 bp using the Covaris instrument (Covaris, Woburn, MA), and the fragmented DNA was then purified with Agencourt AMPure XP beads (Beckman Coulter, Indianapolis, IN). The library was prepared using the Agilent SureSelectXT reagent kit (Agilent Technologies, Santa Clara, CA) with the on-bead modifications of [Fisher et al. \(2011\)](#). Briefly, the DNA was blunted and a single "A-tail" was added to each fragment. Truncated PE P5 and P7 adaptors were ligated to the DNA fragments and the fragments were purified with AMPure beads. The purified DNA fragments were amplified by polymerase chain reaction (PCR; six cycles).

A total of 500 ng of enriched library was used in the hybridization and captured with the SureSelect All Exon v5 (Agilent Technologies) bait. Following hybridization, the captured libraries were purified according to the manufacturer's recommendations and amplified by PCR for 11 cycles using a universal primer and a unique index primer specific to each library. The amplified product was checked for quality using the TapeStation (Agilent Technologies) and quantified by qPCR (Kapa Biosystems). Normalized

libraries were pooled and DNA was sequenced on the Illumina HiSeq 2500 using 2 × 100-bp paired-end reads at a plex-level appropriate to the coverage required for the particular application. An average of 93 million reads were sequenced per tumor sample (average of 103 × the mean tumor target coverage), and an average of 94 million reads were sequenced per germline sample (average of 104 × the mean germline target coverage).

Alignment and Assembly

Exome sequence data processing was performed using Sentieon's somatic mutation pipeline, version 201704.01 (Weber et al., 2016), a proprietary reimplement of The Broad Institute's best practices for somatic mutation calling with MuTect1 (Cibulskis et al., 2013). Briefly, BWA (version 0.7.15) (Li, 2013) was used to align FASTQ files to the publicly available human genome reference hg19, followed by removal of duplicates, realignment around insertions and deletions identified by Mills (Mills et al., 2006), and recalibration of base quality scores. Finally, a co-realignment step was run for patients who had matching tumor and blood sequencing data to increase the quality of the somatic mutations identified.

Sequencing Quality Control

The quality of the aligned sequencing data (mean base quality for each flowcell cycle, the base quality score distribution, GC bias metrics, alignment metrics, and insert size metrics) was calculated for each sample with Sentieon's quality control algorithms based on Picard's various alignment metrics. Additional targeted metrics were collected with Picard's CollectHsMetrics (<http://broadinstitute.github.io/picard/>) based on the Agilent SureSelect v5 target bed file. CollectHsMetrics collected the key metrics of the aligned reads, the average coverage, the percentage of bases >20× as well as other metrics. Finally, BMS cohort-matcher tool (<https://github.com/golharam/cohort-matcher>), which utilizes BAM-matcher (Wang et al., 2016), compared the tumor and blood BAMs to ensure that they came from the same patient, in addition to checking for potential sample swaps within the cohort. If any sample (tumor or blood) of a tumor-blood patient pair failed quality control (total reads <45,000,000, mean target coverage <50×, or percentage of bases >20× <80%) or genotype match between tumor and blood samples was <0.85, then the pair was rejected from the final analysis.

Mutation Calling

The somatic mutations were called from the co-realigned BAM file, which contains the tumor and blood alignments from the same patient. Somatic single-nucleotide mutations were called by Sentieon's TNSnv (reimplementation of Mutect1) (Cibulskis et al., 2013; Weber et al., 2016). In addition, both single-nucleotide mutations and insertions/deletions were called with Strelka (v1.0.15) (Saunders et al., 2012). Both these callers have different approaches to identifying the specific variants present in the tumor but not the blood sample.

Annotation and Filtering

Variant Call Formats (VCFs) generated from TNSnv and Strelka were filtered to retain only variants that passed all internal filters and met a minimum quality standard. These variants were then annotated with snpEff (v4.1c) (Cingolani et al., 2012). Single nucleotide mutations generated by TNSnv and Strelka were filtered to exclude common single nucleotide polymorphisms (SNPs) by comparison to 1000 Genomes phase 1 snps and indels (<http://www.internationalgenome.org/>), Mills insertions and deletions (Mills et al., 2006), ExAC (r3.0, <http://exac.broadinstitute.org/>), and dbSNP (release 138, <https://www.ncbi.nlm.nih.gov/projects/SNP/>). If variants found in SNP databases, but also in COSMIC (v67, <http://cancer.sanger.ac.uk/cosmic>) (Bamford et al., 2004), then they were retained.

Determination of Tumor Mutational Burden

The data processing pipeline used to determine tumor mutational burden has been published previously (Carbone et al., 2017). Tumor mutational burden was evaluated in patients with matched tumor and blood samples who had sufficient whole exome sequencing to pass quality control. Tumor mutational burden was defined as the total number of somatic missense mutations identified by either (or both) somatic variant callers after filtering. For patients with more than one tumor sample whose matched pair passed quality control, the average value from all samples was used to calculate tumor mutational burden. Tumor mutational burden testing was conducted in a research laboratory, and the methodology is not a Clinical Laboratory Improvement Amendments–approved clinical diagnostic test.

For most analyses of treatment outcomes by tumor mutational burden, patients were grouped in thirds according to tumor mutational burden (analysis by tumor mutational burden tertiles); additional analyses were based on median or quartiles. A receiver operating characteristic analysis of best overall response by tumor mutational burden as a continuous variable was also performed.

Molecular Features of Previously Published Small-Cell Lung Cancers

Called mutations and clinical outcomes were collected from a cohort of SCLCs examined by whole genome sequencing as described (George et al., 2015). A total of 100 patients had paired tumor and normal sequencing, available molecular data, and known overall survival. To align with methods applied here, only missense mutations were used for quantifying tumor mutational burden.

QUANTIFICATION AND STATISTICAL ANALYSIS

All analyses were performed on the basis of the original treatment assignment, regardless of crossover status. The Kaplan–Meier method was used to estimate survival. The Clopper–Pearson method was used to estimate response rates and their exact 95% confidence intervals. Statistical analyses were performed using R version 3.4.1.

DATA AND SOFTWARE AVAILABILITY

Human sequencing data (WES; from patients who provided full written informed consent to share these data) were deposited into the European Variation Archive (Accession numbers: PRJEB25808 for Strelka and PRJEB25807 for TNsnv).

ADDITIONAL RESOURCES

Clinical trial registry: <https://clinicaltrials.gov/ct2/show/NCT01928394>
1000 Genomes phase 1 snps and indels: <http://www.internationalgenome.org/>
COSMIC (v67): <http://cancer.sanger.ac.uk/cosmic>
ExAC (r3.0): <http://exac.broadinstitute.org/>
dbSNP (release 138): <https://www.ncbi.nlm.nih.gov/projects/SNP/>
Picard's CollectHsMetrics: <http://broadinstitute.github.io/picard/>
BMS cohort-matcher tool: <https://github.com/golharam/cohort-matcher>