

This Site Uses Cookies and Your Privacy Choice Is Important to Us. Choose *Customize My Settings* to make your privacy choices. Choose *Accept All Cookies* to accept third-party cookies. [See our Privacy Policy](#).


[Customize My Settings](#)
[Accept All Cookies](#)

KEYTRUDA[®]
(pembrolizumab) injection 100 mg

 
Cancer Type Menu

Advanced MSI-H/dMMR cancer

IS KEYTRUDA RIGHT FOR YOU?



Related topics

> [Side effects](#)

> [Results from clinical trials](#)

> [Adrienne's patient story](#)

On this page:

[About MSI-H/dMMR](#) ▾

[Test for MSI-H/dMMR](#) ▾

Immunotherapy may be right for you. It's TRU. KEYTRUDA.

What are MSI-H/dMMR cancers?

In normal cells, a system called DNA mismatch repair (MMR) corrects errors that occur during DNA replication. Defects in MMR can lead to microsatellite instability-high (MSI-H), which can be found in many types of cancer. MSI-H or mismatch repair deficient (dMMR) tumors have an accumulation of errors in genetic sequences that are normally repeated (called microsatellites).



How a test for MSI-H/dMMR works

Everyone's cancer is different. To help develop a treatment plan that's right for you, your doctor may order laboratory tests. These tests can evaluate your tumor for different biomarkers, including MSI-H or dMMR. Some biomarkers are used to help select patients that may be eligible for certain treatments, such as KEYTRUDA. Talk to your doctor to see if a laboratory test for the MSI-H or dMMR biomarker is right for you.

1. **A tissue sample (biopsy) of your tumor is taken**, and your doctor orders the test.
2. **Your doctor typically gets the results** in 10 to 20 days.
3. **You talk with your doctor** to discuss treatment options based on your results.
4. **You may begin treatment with KEYTRUDA** if you and your doctor have decided it's right

<https://www.keytruda.com/msi-h/treatment-options>

for you.



Your doctor can help you find out if you're a match for **KEYTRUDA**

A laboratory test for MSI-H or dMMR can help determine whether KEYTRUDA is right for you.

KEYTRUDA may still be an option even though you have tried other cancer treatments.

Questions you should ask your doctor about MSI-H/dMMR testing:

- Will you be testing my tumor for MSI-H/dMMR?
- How would this test be helpful in determining my treatment plan?
- Is KEYTRUDA right for me?



Talk to your doctor about a test for MSI or MMR to see if **KEYTRUDA** could be an option for you

Getting tested for MSI or MMR status can help determine if KEYTRUDA could be an appropriate treatment for you or someone you care about.

KEYTRUDA was studied in these MSI-H/dMMR cancers:

- | | |
|--|--------------------|
| • Colorectal | • Mesothelioma |
| • Endometrial | • Nasopharyngeal |
| • Adrenocortical | • Neuroendocrine |
| • Anal | • Ovarian |
| • Appendiceal cancer | • Pancreatic |
| • Biliary | • Prostate |
| • Bladder | • Renal cell |
| • Brain | • Retroperitoneal |
| • Breast | • Salivary |
| • Cervical | • Sarcoma |
| • Gastric or gastroesophageal junction | • Small cell lung |
| • Head and neck | • Small intestinal |
| | • Testicular |

- Head and neck squamous cell
- Liver
- Vaginal
- Thyroid
- Unknown origin
- Vulvar



Patient support throughout your treatment

Patients who have been prescribed KEYTRUDA for an approved indication can access additional support from KEY+YOU, the patient support program for KEYTRUDA. KEY+YOU helps eligible patients by offering educational resources and support information throughout treatment.

Call 85-KEYTRUDA (855-398-7832) >



Additional educational resources for patients

- > [Questions to ask before starting treatment \(PDF\)](#)
- > [Questions to ask while on treatment \(PDF\)](#)
- > [Track your symptoms \(PDF\)](#)
- > [Frequently asked questions about KEYTRUDA](#)

You also might be interested in:

About KEYTRUDA

> [Adrienne's patient story](#)

Results with KEYTRUDA

> [Results from clinical trials](#)

About KEYTRUDA

> [Side effects](#)

KEYTRUDA
(pembrolizumab) injection 100 mg

FDA-Approved Indications

KEYTRUDA is a prescription medicine used to treat a kind of cancer that is shown by a laboratory test to be a [microsatellite instability-high \(MSI-H\)](#) or a mismatch repair deficient ([dMMR](#)) solid tumor. KEYTRUDA may be used in adults and children to treat:

- cancer that has spread or cannot be removed by surgery (advanced cancer), and
- has progressed following treatment, and you have no satisfactory treatment options.

Chat

Important Safety Information

KEYTRUDA is a medicine that may treat certain cancers by working with your immune system. KEYTRUDA can cause your immune system to attack normal organs and tissues in any area of your body and can affect the way they work. These problems can sometimes become severe or life-threatening and can lead to death. You can have more than one of these problems at the same time. These problems may happen any time during treatment or even after your treatment has ended.

Call or see your health care provider right away if you develop any signs or symptoms of the following problems or if they get worse. These are not all of the signs and symptoms of immune system problems that can happen with KEYTRUDA:

Lung problems: cough, shortness of breath, or chest pain.

Intestinal problems: diarrhea (loose stools) or more frequent bowel movements than usual; stools that are black, tarry, sticky, or have blood or mucus; or severe stomach-area (abdomen) pain or tenderness.

Liver problems: yellowing of your skin or the whites of your eyes; severe nausea or vomiting; pain on the right side of your stomach area (abdomen); dark urine (tea colored); or bleeding or bruising more easily than normal.

Hormone gland problems: headaches that will not go away or unusual headaches; eye sensitivity to light; eye problems; rapid heartbeat; increased sweating; extreme tiredness; weight gain or weight loss; feeling more hungry or thirsty than usual; urinating more often than usual; hair loss; feeling cold; constipation; your voice gets deeper; dizziness or fainting; changes in mood or behavior, such as decreased sex drive, irritability, or forgetfulness.

Kidney problems: decrease in the amount of your urine; blood in your urine; swelling of your ankles; loss of appetite.

Skin problems: rash; itching; skin blistering or peeling; painful sores or ulcers in your mouth or in your nose, throat, or genital area; fever or flu-like symptoms; swollen lymph nodes.

Problems can also happen in other organs and tissues. Signs and symptoms of these problems may include: chest pain; irregular heartbeat; shortness of breath; swelling of ankles; confusion; sleepiness; memory problems; changes in mood or behavior; stiff neck; balance problems; tingling or numbness of the arms or legs; double vision; blurry vision; sensitivity to light; eye pain; changes in eyesight; persistent or severe muscle pain or weakness; muscle cramps; low red blood cells; bruising.

Infusion reactions that can sometimes be severe or life-threatening. Signs and symptoms of infusion reactions may include chills or shaking, itching or rash, flushing, shortness of breath or wheezing, dizziness, feeling like passing out,

fever, and back pain.

Rejection of a transplanted organ or tissue. Your health care provider should tell you what signs and symptoms you should report and they will monitor you, depending on the type of organ or tissue transplant that you have had.

Complications, including graft-versus-host disease (GVHD), in people who have received a bone marrow (stem cell) transplant that uses donor stem cells (allogeneic). These complications can be serious and can lead to death. These complications may happen if you underwent transplantation either before or after being treated with KEYTRUDA. Your health care provider will monitor you for these complications.

Getting medical treatment right away may help keep these problems from becoming more serious. Your health care provider will check you for these problems during treatment with KEYTRUDA. They may treat you with corticosteroid or hormone replacement medicines. They may also need to delay or completely stop treatment with KEYTRUDA if you have severe side effects.

Before you receive KEYTRUDA, tell your health care provider if you have immune system problems such as Crohn's disease, ulcerative colitis, or lupus; have had an organ or tissue transplant, including corneal transplant, or have had or plan to have a bone marrow (stem cell) transplant that uses donor stem cells (allogeneic); have had radiation treatment in your chest area; have a condition that affects your nervous system, such as myasthenia gravis or Guillain-Barré syndrome.

If you are pregnant or plan to become pregnant, tell your health care provider. KEYTRUDA can harm your unborn baby. If you are able to become pregnant, you will be given a pregnancy test before you start treatment. Use effective birth control during treatment with KEYTRUDA and for 4 months after your last dose of KEYTRUDA. Tell them right away if you think you may be pregnant or you become pregnant during treatment with KEYTRUDA.

Tell your health care provider if you are breastfeeding or plan to breastfeed. It is not known if KEYTRUDA passes into your breast milk. Do not breastfeed during treatment with KEYTRUDA and for 4 months after your last dose of KEYTRUDA.

Tell your health care provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Common side effects of KEYTRUDA when used alone include feeling tired; pain, including pain in muscles; rash; diarrhea; fever; cough; decreased appetite; itching; shortness of breath; constipation; bones or joints and stomach-area (abdominal) pain; nausea; and low levels of thyroid hormone.

In children, when KEYTRUDA is used alone, fever, vomiting, headache, stomach area (abdominal) pain, and low levels of white blood cells and red blood cells (anemia) are more common than in adults.

Common side effects of KEYTRUDA when given with certain chemotherapy medicines include feeling tired or weak; nausea; constipation; diarrhea;

decreased appetite; rash; vomiting; cough; trouble breathing; fever; hair loss; inflammation of the nerves that may cause pain, weakness, and paralysis in the arms and legs; swelling of the lining of the mouth, nose, eyes, throat, intestines, or vagina; mouth sores; headache; weight loss; stomach-area (abdominal) pain; joint and muscle pain; trouble sleeping; bleeding, blisters, or rash on the palms of your hands and soles of your feet; urinary tract infection; and low levels of thyroid hormone.

Common side effects of KEYTRUDA when given with chemotherapy with radiation therapy medicines include feeling tired or weak; nausea; constipation; diarrhea; decreased appetite; rash; vomiting; cough; trouble breathing; fever; hair loss; inflammation of the nerves that may cause pain, weakness, and paralysis in the arms and legs; swelling of the lining of the mouth, nose, eyes, throat, intestines, or vagina; mouth sores; headache; weight loss; stomach-area (abdominal) pain; joint and muscle pain; trouble sleeping; bleeding, blisters, or rash on the palms of your hands and soles of your feet; urinary tract infection; and low levels of thyroid hormone.

Common side effects of KEYTRUDA when given with enfortumab vedotin include rash; tingling or numbness of the arms or legs; feeling tired; itching; diarrhea; hair loss; weight loss; decreased appetite; dry eye; nausea; constipation; changes in sense of taste; and urinary tract infection.

Common side effects of KEYTRUDA when given with chemotherapy and bevacizumab include tingling or numbness of the arms or legs; hair loss; low red blood cell count; feeling tired or weak; nausea; low white blood cell count; diarrhea; high blood pressure; decreased platelet count; constipation; joint aches; vomiting; urinary tract infection; rash; low levels of thyroid hormone; and decreased appetite.

Common side effects of KEYTRUDA when given with axitinib include diarrhea; feeling tired or weak; high blood pressure; liver problems; low levels of thyroid hormone; decreased appetite; blisters or rash on the palms of your hands and soles of your feet; nausea; mouth sores or swelling of the lining of the mouth, nose, eyes, throat, intestines, or vagina; hoarseness; rash; cough; and constipation.

These are not all the possible side effects of KEYTRUDA. Talk to your health care provider for medical advice about side effects.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

Please read the accompanying [Medication Guide](#) for KEYTRUDA and discuss it with your doctor. The physician [Prescribing Information](#) also is available.



About KEYTRUDA

- > [How does KEYTRUDA work?](#)
- > [Side effects](#)
- > [How KEYTRUDA is given](#)
- > [Frequently asked questions](#)

Cost Info & Financial Help

- > [Cost info & financial help](#)

Indications & Safety

- > [Indications](#)
- > [Important safety information](#)

Results From Clinical Trials

- > [Results from clinical trials](#)

Patient Support

- > [What to know about treatment with KEYTRUDA](#)

[Privacy Policy](#) • [Terms of Use](#) • [Cookie Preferences](#) • [Accessibility](#)



[View Health Care Professionals Site](#)

Copyright © 2024 Merck & Co., Inc., Rahway, NJ, USA and its affiliates. All rights reserved.

This site is intended only for residents of the United States, its territories, and Puerto Rico.

US-OVA-00921 04/23



Need Help?