



Actor portrayals. OPDIVO or OPDIVO + YERVOY is not approved for people younger than 12 years of age.

Your fight against colorectal cancer (MSI-H/dMMR) may continue here

OPDIVO, alone or with YERVOY, is an FDA-approved immunotherapy option to treat advanced colorectal cancer (MSI-H/dMMR)

Previously treated colorectal cancer

OPDIVO + YERVOY is a treatment option for certain adults and children 12 years and older who have had prior treatments for MSI-H/dMMR colorectal cancer

[Clinical trial results](#) →

[Getting an infusion](#) →

Previously treated colorectal cancer

OPDIVO is a treatment option for certain adults and children 12 years and older who have had prior treatments for MSI-H/dMMR colorectal cancer

[Clinical trial results](#) →

[Getting an infusion](#) →

OPDIVO will not work for everyone. Individual results may vary.

Important Facts About OPDIVO® (nivolumab) and OPDIVO + YERVOY® (ipilimumab)

This is a summary of important information that you need to know about OPDIVO and OPDIVO + YERVOY. Your healthcare team can work with you to help answer any questions you may have about these medications. Keep this information in a safe place so you can refer to it before and during your treatment.

^




Colorectal cancer (MSI-H...

[About OPDIVO](#)

[Patient & Caregiver Support](#)

[Cost & Access](#)



Patient stories

Real patients share their experiences with OPDIVO and OPDIVO-based treatments

[Watch videos](#) →



Important Facts About OPDIVO® (nivolumab) and OPDIVO + YERVOY® (ipilimumab)



This is a summary of important information that you need to know about OPDIVO and OPDIVO + YERVOY. Your healthcare team can work with you to help answer any questions you may have about these medications. Keep this information in a safe place so you can refer to it before and during your treatment.



Colorectal cancer (MSI-H...

[About OPDIVO](#)

[Patient & Caregiver Support](#)

[Cost & Access](#)



Cost information and financial help

Find information about cost and access to OPDIVO

[Learn more](#) →

Important Facts About OPDIVO® (nivolumab) and OPDIVO + YERVOY® (ipilimumab)

This is a summary of important information that you need to know about OPDIVO and OPDIVO + YERVOY. Your healthcare team can work with you to help answer any questions you may have about these medications. Keep this information in a safe place so you can refer to it before and during your treatment.

Look out for the following icons as you read:



Talk to your
healthcare team

Important Facts About OPDIVO® (nivolumab) and OPDIVO + YERVOY® (ipilimumab)



This is a summary of important information that you need to know about OPDIVO and OPDIVO + YERVOY. Your healthcare team can work with you to help answer any questions you may have about these medications. Keep this information in a safe place so you can refer to it before and during your treatment.



What are the serious side effects of OPDIVO and OPDIVO + YERVOY?

A **serious side effect** is a side effect that can sometimes become severe or life-threatening, and can lead to death. They may happen anytime during treatment or even after your treatment has ended. You may have more than one of these problems at the same time.

Call or see your healthcare provider right away if you develop any new or worse signs or symptoms, including:

Lung problems — Things to look out for may include:

- cough
- shortness of breath
- chest pain

Intestinal problems — Things to look out for may include:

- diarrhea (loose stools) or more frequent bowel movements than usual
- stools that are black, tarry, sticky, or have blood or mucus
- severe stomach-area (abdominal) pain or tenderness

Liver problems — Things to look out for may include:

- yellowing of your skin or the whites of your eyes
- severe nausea or vomiting
- pain on the right side of your stomach area (abdomen)
- dark urine (tea colored)
- bleeding or bruising more easily than normal

Hormone gland problems — Things to look out for may include:

- headaches that will not go away or unusual headaches
- eye sensitivity to light
- eye problems
- rapid heartbeat
- increased sweating
- extreme tiredness
- weight gain or weight loss
- feeling more hungry or thirsty than usual
- urinating more often than usual
- hair loss
- feeling cold
- constipation
- your voice gets deeper

Important Facts About OPDIVO® (nivolumab) and OPDIVO + YERVOY® (ipilimumab)



This is a summary of important information that you need to know about OPDIVO and OPDIVO + YERVOY. Your healthcare team can work with you to help answer any questions you may have about these medications. Keep this information in a safe place so you can refer to it before and during your treatment.



- eye pain or redness

Problems can also happen in other organs and tissues. These are not all the signs and symptoms of immune system problems that can happen with OPDIVO and YERVOY. Call or see your healthcare provider right away for any new or worsening signs or symptoms, which may include:

- Chest pain, irregular heartbeat, shortness of breath, swelling of ankles
- Confusion, sleepiness, memory problems, changes in mood or behavior, stiff neck, balance problems, tingling or numbness of the arms or legs
- Double vision, blurry vision, sensitivity to light, eye pain, changes in eyesight
- Persistent or severe muscle pain or weakness, muscle cramps
- Low red blood cells, bruising

Rejection of a transplanted organ or tissue. Your healthcare provider should tell you what signs and symptoms you should report and monitor you depending on the type of organ transplant that you have had.

What are the possible side effects of OPDIVO + YERVOY?

OPDIVO and OPDIVO + YERVOY can cause serious side effects, including:

See the previous section, **“What is the most important information I should know about OPDIVO + YERVOY?”**

Severe infusion reactions — Things to look out for may include:

- chills or shaking
- itching or rash
- flushing
- shortness of breath or wheezing
- dizziness
- feel like passing out
- fever
- back or neck pain



Tell your healthcare team right away if you get these symptoms during an infusion of OPDIVO or YERVOY.

Complications, including graft-versus-host disease (GVHD), of bone marrow (stem cell) transplant that uses donor stem cells (allogeneic). These complications can be serious and can lead to death. These complications may happen if you underwent transplantation either before or after being treated with OPDIVO or YERVOY. Your healthcare provider will monitor you for these complications.

What are the most common side effects?

Important Facts About OPDIVO® (nivolumab) and OPDIVO + YERVOY® (ipilimumab)

This is a summary of important information that you need to know about OPDIVO and OPDIVO + YERVOY. Your healthcare team can work with you to help answer any questions you may have about these medications. Keep this information in a safe place so you can refer to it before and during your treatment.

Indications	Prescribing Information >
Medication Guide >	View Healthcare Professional Site >



Colorectal cancer (MSI-H...

[About OPDIVO](#)

[Patient & Caregiver Support](#)

[Cost & Access](#)

- **vomiting**
- **urinary tract infection**

The most common side effects of OPDIVO when used in combination with YERVOY include:

- **feeling tired**
- **diarrhea**
- **rash**
- **itching**
- **nausea**
- **pain in muscles, bones, and joints**
- **fever**
- **cough**
- **decreased appetite**
- **vomiting**
- **stomach-area (abdominal) pain**
- **shortness of breath**
- **upper respiratory tract infection**
- **headache**
- **low thyroid hormone levels (hypothyroidism)**
- **constipation**
- **decreased weight**
- **dizziness**

The most common side effects of OPDIVO when used in combination with YERVOY and chemotherapy include:

- **feeling tired**
- **pain in muscles, bones, and joints**
- **nausea**
- **diarrhea**
- **rash**
- **decreased appetite**
- **constipation**
- **itching**

The most common side effects of OPDIVO when used in combination with chemotherapy include:

- **nausea**
- **feeling tired**
- **pain in muscles, bones and joints**
- **constipation**
- **decreased appetite**
- **rash**

Important Facts About OPDIVO® (nivolumab) and OPDIVO + YERVOY® (ipilimumab) ^

This is a summary of important information that you need to know about OPDIVO and OPDIVO + YERVOY. Your healthcare team can work with you to help answer any questions you may have about these medications. Keep this information in a safe place so you can refer to it before and during your treatment.



Colorectal cancer (MSI-H...

About OPDIVO Patient & Caregiver Support Cost & Access

- **upper respiratory tract infection**
The most common side effects of OPDIVO when used in combination with fluoropyrimidine and platinum-containing chemotherapy include:
- **nausea**
- **numbness, pain, tingling, or burning in your hands or feet**
- **decreased appetite**
- **feeling tired**
- **constipation**
- **mouth sores**
- **diarrhea**
- **vomiting**
- **stomach-area (abdominal) pain**
- **pain in muscles, bones, and joints**

These are not all the possible side effects. **Talk to your healthcare team** or pharmacist for more information. You are encouraged to report side effects of prescription drugs to the FDA. Call 1-800-FDA-1088.

What should I discuss with my healthcare team before receiving OPDIVO or YERVOY?



Tell your healthcare provider about all your medical conditions, including if you:

- have immune system problems such as Crohn's disease, ulcerative colitis, or lupus
- have received an organ transplant, including corneal transplant
- have received or plan to receive a stem cell transplant that uses donor stem cells (allogeneic)
- have received radiation treatment to your chest area in the past and have received other medicines that are like OPDIVO
- have a condition that affects your nervous system, such as myasthenia gravis or Guillain-Barré syndrome
- are pregnant or plan to become pregnant. OPDIVO and YERVOY can harm your unborn baby.
- are breastfeeding or plan to breastfeed. It is not known if OPDIVO or YERVOY passes into your breast milk. Do not breastfeed during treatment and for 5 months after your last dose of OPDIVO or YERVOY.



Females who are able to become pregnant:

Your healthcare provider should do a pregnancy test before you start receiving OPDIVO or YERVOY.

- You should use an effective method of birth control during your treatment and for 5 months after the last dose of OPDIVO or YERVOY. Talk to your healthcare provider about birth control methods that you can use during this time.

Important Facts About OPDIVO® (nivolumab) and OPDIVO + YERVOY® (ipilimumab)

This is a summary of important information that you need to know about OPDIVO and OPDIVO + YERVOY. Your healthcare team can work with you to help answer any questions you may have about these medications. Keep this information in a safe place so you can refer to it before and during your treatment.

Indications	Prescribing Information >
Medication Guide >	View Healthcare Professional Site >



- Colorectal cancer (MSI-H...
- About OPDIVO
- Patient & Caregiver Support
- Cost & Access

Information provided on this website is not a substitute for talking with your healthcare professional.
Your healthcare professional is the best source of information about your disease.
All individuals depicted are models used for illustrative purposes only.

1506-US-2400728 11/24



[Site Map](#) | [Legal Notice](#) | [Privacy Policy](#) | [Your Privacy Choices](#) | 1-855-OPDIVO-1

© 2024 Bristol-Myers Squibb Company.
OPDIVO®, YERVOY®, Access Support®, A Chance to Live Longer™, and the related logos are trademarks of Bristol-Myers Squibb Company.

This site is intended for U.S. residents 18 years of age or older.

466-US-2400031 03/24

Important Facts About OPDIVO® (nivolumab) and OPDIVO + YERVOY® (ipilimumab)

This is a summary of important information that you need to know about OPDIVO and OPDIVO + YERVOY. Your healthcare team can work with you to help answer any questions you may have about these medications. Keep this information in a safe place so you can refer to it before and during your treatment.