

Pharmacokinetic and Pharmacodynamic (PK/PD) Relationship of Intravenous Ganaxolone in Refractory Status Epilepticus

R. Eugene Ramsay, MD⁴, Aatif M. Husain, MD², Christa B. Swisher, MD², Steven Smith³, Heather Van Heusen³, Julia Tsai, PhD³, Lorianne Masuoka, MD³, Maciej Gasior, MD, PhD³, Henrikas Vaitkevicius, MD¹
¹Neurosciences Institute, Ochsner Health System, New Orleans, LA; ²Dept. of Neurology, Duke University, Durham, NC; ³Marinus Pharmaceuticals, Radnor, PA; ⁴Dept. of Neurology, Brigham and Women's Hospital, Boston, MA

RSE OVERVIEW

Status Epilepticus (SE) is the most life-threatening occurrence within the spectrum of epileptic disorders

- Continuous seizures lasting >5 min for convulsive seizures or >30 minutes for non-convulsive seizures
- Heterogenous patient population with various/unknown etiologies
- Prolonged seizure activity can result in permanent neuronal damage and contribute to the high morbidity and mortality rates associated with SE
- SE becomes more difficult to control as its duration increases and is associated with increased mortality

Established status epilepticus (ESE) is defined as failure of at least one benzodiazepine at adequate doses

RSE defined as failure of at least one 2nd line intravenous (IV) anti-epileptic drug (AED)

Super Refractory Status Epilepticus (SRSE) is SE that continues or recurs 24hrs or more after the onset of IV anesthetics or failure to wean off IV anesthetics

GOALS OF A NEW THERAPY FOR THE TREATMENT OF RSE

Established Status Epilepticus (ESE) Refractory Status Epilepticus (RSE) Super Refractory Status Epilepticus (SRSE)

1st line Benzodiazepine Administered 2nd line IV AED's 3rd line IV Anesthetics Medically induced Coma

Goals of a new treatment for RSE

- Prevent patient progression towards escalation of treatment (IV anesthetics)
- Rapid cessation of SE
- Maintenance of seizure control over study period

GANAXOLONE (GNX) MECHANISM OF ACTION

- GNX is a synthetic analog of endogenous allopregnanolone and a potent positive allosteric modulator of GABA_A-receptors.
- GNX targets unique binding sites on GABA_A-receptors that are not susceptible to tolerance build up (e.g., benzodiazepines)
- GNX acts on both synaptic and extrasynaptic GABA_A-receptors to maximize inhibitory signaling as well as maintain activity when synaptic receptors are down-regulated

Study Design

Objectives: Evaluate safety, tolerability, efficacy, and pharmacokinetics of IV Ganaxolone in RSE patients

RSE Patients → Screening → Treatment Period → Post-treatment Follow-up

Treatment Period:

Loading Dose	Maintenance	Taper
10 mg/kg	2-4 day infusion	18 hour taper

Post-treatment Follow-up: 24 hr, Weeks 2, 3, 4

Endpoints:

- Primary:** number of patients who do not require escalation of treatment with IV anesthetic within the first 24 hours after ganaxolone initiation
- Secondary:** additional efficacy, safety and tolerability

Patient Baseline Characteristics

17 patients were enrolled (8 males, 9 females) with an average age of 57 years old (range: 23-88)

Patient	Dosing Cohort	Etiology	History of Epilepsy	Type of SE	Failed Antiseizure Medications Prior to GNX*
1	Low	Vascular	No	NCSE	LAC, LEV
2	Low	Unknown	Yes	NCSE	PHPT, LEV
3	Low	Vascular	No	NCSE	LOR, LAC, LEV
4	Low	Vascular	No	NCSE	LOR, LAC, LEV
5	Low	Tumor	No	CSE	LOR, LAC, LEV
6	Medium	Vascular	No	NCSE	LOR, LAC, LEV
7	Medium	Drug Overdose / Withdrawal	Yes	CSE	LOR, LEV
8	Medium	Unknown	Yes	CSE → NCSE	LOR, LAC, LEV
9	Medium	Tumor	Yes	NCSE	LAC, LEV, PHT
10	Target	Vascular	Yes	CSE	LOR, LAC, VPA
11	Target	Drug Overdose / Withdrawal	No	CSE	LOR, LAC, LEV
12	Target	Tumor	Yes	NCSE	LOR, LEV, VPA
13	Target	Autoimmune	No	NCSE	LOR, LEV
14	Target	Vascular	No	NCSE	LOR, LAC, LEV, PHT
15	Target	Vascular	Yes	CSE	LOR, LEV
16	Target	Tumor	No	NCSE	LOR, LAC, LEV
17	Target	Autoimmune	No	NCSE	LOR, PHT, LAC, LEV, VPA

NCSE: Non-convulsive status epilepticus, CSE: Convulsive status epilepticus, LAC: Lacosamide, LEV: Levetiracetam, LOR: Lorazepam, PHT: Phenytoin, PHPT: Fosphenytoin, VPA: Valproic Acid
*All prior AEDs were administered at therapeutically relevant doses and within recommended dosing guidelines

PHASE 2 TRIAL OF IV GANAXOLONE IN RSE

Efficacy of IV Ganaxolone

Cohort	No escalation to IV anesthetics within 24 hours from infusion initiation (Primary Endpoint)	Status-free through 24 hours from infusion initiation	No escalation to additional IV AEDs or IV anesthetics for status relapse at any time through 24hr after GNX discontinuation	No SE Relapse at anytime during the 4-wk follow up period	
Target (713 mg/day; n=8)	100% (8 of 8)	88% (7 of 8)	100% (8 of 8)	100% (6 of 6); 1ET, 1died	One patient had status relapse @ Day 1, which resolved during the ganaxolone infusion without treatment escalation
Medium (650 mg/day; n=4)	100% (4 of 4)	100% (4 of 4)	75% (3 of 4)	67% (2 of 3); 1ET	One patient escalated to additional IV AED @ Day 1 for seizure relapse. One patient experienced status relapse @ Day 2 (during taper).
Low (500 mg/day; n=5)	100% (5 of 5)	100% (5 of 5)	60% (3 of 5)	50% (1 of 2); 1died	Two patients escalated to 3rd line therapy for seizure relapse @ Day3

ET: Early Termination

Safety of IV Ganaxolone

10 SAEs in 6 patients (also included in AEs)	50 AEs in 16 patients
2 related in 2 patients • 2 severe sedation	13 Related in 7 subjects • 6 mild (2 hypotension, 2 somnolence, 1 urinary retention, 1 hypercarbia) • 5 moderate (4 somnolence; 1 hypercarbia) • 2 severe (2 sedation)

Intubation

9 patients were not intubated upon enrollment. Of these, 6 remained intubation-free during the entire ganaxolone treatment period

PK/PD Relationship and Rationale for Target Dose

GNX PK and Correlation with EEG

Modeled PK Curves for Dosing Groups

Parameter	Low (500 mg/day)	Medium (650 mg/day)	Target (713 mg/day)
Target acute minimum plasma concentration	~200 ng/mL	~400 ng/mL	~500 ng/mL

Conclusions

- No patients progressed to IV anesthetics during first 24 hours
- Median time to SE cessation = 5 minutes (3.4-14.4 minutes 25th-75th percentile) (n=15 evaluable)
- Durable response throughout study period in target dose cohort
- Patients failed mean of 2.1 second line IV AEDs
- Highly heterogeneous underlying cause of status
- Target patient population and dose identified for Ph. 3 study
- Ganaxolone shows an acceptable safety profile in patients with RSE