

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

TRANSMITTAL FOR POWER OF ATTORNEY TO ONE OR MORE REGISTERED PRACTITIONERS

NOTE: This form is to be submitted with the Power of Attorney by Applicant form (PTO/AIA/82B or equivalent) to identify the application to which the Power of Attorney is directed, in accordance with 37 CFR 1.5. If the Power of Attorney by Applicant form is not accompanied by this transmittal form or an equivalent, the Power of Attorney will not be recognized in the application.

Application Number	Not Yet Assigned
Filing Date	Concurrently Herewith
First Named Inventor	Matthew DURING
Title	METHODS AND COMPOSITIONS FOR TREATMENT OF EPILEPTIC DISORDERS
Art Unit	Not Yet Assigned
Examiner Name	Not Yet Assigned
Attorney Docket Number	2262-29

SIGNATURE of Applicant or Patent Practitioner

Signature	/Jeffrey S. Steen, Reg. #32063/	Date	August 10, 2017
Name	Jeffrey S. Steen	Telephone	(631) 501-5700
Registration Number	32,063		

NOTE: This form must be signed in accordance with 37 CFR 1.33. See 37 CFR 1.4(d) for signature requirements and certifications.

☒ *Total of 1 forms are submitted.

This collection of information is required by 37 CFR 1.31, 1.32 and 1.33. The information is required to obtain or retain a benefit by the public which is to file (and by the USPTO to process) an application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.11 and 1.14. This collection is estimated to take 3 minutes to complete, including gathering, preparing, and submitting the completed application form to the USPTO. Time will vary depending upon the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, P.O. Box 1450, Alexandria, VA 22313-1450. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. **SEND TO: Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.**

If you need assistance in completing the form, call 1-800-PTO-9199 and select option 2.

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

Application Data Sheet 37 CFR 1.76		Attorney Docket Number	2262-29
		Application Number	
Title of Invention	METHODS AND COMPOSITIONS FOR TREATMENT OF EPILEPTIC DISORDERS		
The application data sheet is part of the provisional or nonprovisional application for which it is being submitted. The following form contains the bibliographic data arranged in a format specified by the United States Patent and Trademark Office as outlined in 37 CFR 1.76. This document may be completed electronically and submitted to the Office in electronic format using the Electronic Filing System (EFS) or the document may be printed and included in a paper filed application.			

Secrecy Order 37 CFR 5.2

☐ Portions or all of the application associated with this Application Data Sheet may fall under a Secrecy Order pursuant to 37 CFR 5.2 (Paper filers only. Applications that fall under Secrecy Order may not be filed electronically.)

Inventor Information:

Inventor	1				Remove	
Legal Name						
Prefix	Given Name	Middle Name	Family Name	Suffix		
	Matthew		DURING			
Residence Information (Select One) • US Residency Non US Residency Active US Military Service						
City	Weston	State/Province	CT	Country of Residence	US	
Mailing Address of Inventor:						
Address 1	14 Cedar Lane					
Address 2						
City	Weston	State/Province	CT			
Postal Code	06883	Country	US			
All Inventors Must Be Listed - Additional Inventor Information blocks may be generated within this form by selecting the Add button.						
Add						

Correspondence Information:

Enter either Customer Number or complete the Correspondence Information section below. For further information see 37 CFR 1.33(a).			
☐ An Address is being provided for the correspondence information of this application.			
Customer Number	31554		
Email Address	docket@cdfslaw.com	Add Email	Remove Email

Application Information:

Title of the Invention	METHODS AND COMPOSITIONS FOR TREATMENT OF EPILEPTIC DISORDERS		
Attorney Docket Number	2262-29	Small Entity Status Claimed	☐
Application Type	Nonprovisional		
Subject Matter	Utility		
Total Number of Drawing Sheets (if any)	4	Suggested Figure for Publication (if any)	
Filing By Reference :			

Application Data Sheet 37 CFR 1.76		Attorney Docket Number	2262-29
		Application Number	
Title of Invention	METHODS AND COMPOSITIONS FOR TREATMENT OF EPILEPTIC DISORDERS		

Only complete this section when filing an application by reference under 35 U.S.C. 111(c) and 37 CFR 1.57(a). Do not complete this section if application papers including a specification and any drawings are being filed. Any domestic benefit or foreign priority information must be provided in the appropriate section(s) below (i.e., "Domestic Benefit/National Stage Information" and "Foreign Priority Information").

For the purposes of a filing date under 37 CFR 1.53(b), the description and any drawings of the present application are replaced by this reference to the previously filed application, subject to conditions and requirements of 37 CFR 1.57(a).

Application number of the previously filed application	Filing date (YYYY-MM-DD)	Intellectual Property Authority or Country

Publication Information:

☐ Request Early Publication (Fee required at time of Request 37 CFR 1.219)

☐ **Request Not to Publish.** I hereby request that the attached application not be published under 35 U.S.C. 122(b) and certify that the invention disclosed in the attached application **has not and will not** be the subject of an application filed in another country, or under a multilateral international agreement, that requires publication at eighteen months after filing.

Representative Information:

Representative information should be provided for all practitioners having a power of attorney in the application. Providing this information in the Application Data Sheet does not constitute a power of attorney in the application (see 37 CFR 1.32). Either enter Customer Number or complete the Representative Name section below. If both sections are completed the customer Number will be used for the Representative Information during processing.

Please Select One:	☒ Customer Number	☐ US Patent Practitioner	☐ Limited Recognition (37 CFR 11.9)
Customer Number	B1554		

Domestic Benefit/National Stage Information:

This section allows for the applicant to either claim benefit under 35 U.S.C. 119(e), 120, 121, 365(c), or 386(c) or indicate National Stage entry from a PCT application. Providing this information in the application data sheet constitutes the specific reference required by 35 U.S.C. 119(e) or 120, and 37 CFR 1.78.

When referring to the current application, please leave the application number blank.

Prior Application Status	Pending	Remove	
Application Number	Continuity Type	Prior Application Number	Filing Date (YYYY-MM-DD)
	Claims benefit of provisional	62373589	2016-08-11
Prior Application Status	Pending	Remove	
Application Number	Continuity Type	Prior Application Number	Filing Date (YYYY-MM-DD)
	Claims benefit of provisional	62490293	2017-04-26
Additional Domestic Benefit/National Stage Data may be generated within this form by selecting the Add button.			Add

Application Data Sheet 37 CFR 1.76		Attorney Docket Number	2262-29
		Application Number	
Title of Invention	METHODS AND COMPOSITIONS FOR TREATMENT OF EPILEPTIC DISORDERS		

Foreign Priority Information:

This section allows for the applicant to claim priority to a foreign application. Providing this information in the application data sheet constitutes the claim for priority as required by 35 U.S.C. 119(b) and 37 CFR 1.55. When priority is claimed to a foreign application that is eligible for retrieval under the priority document exchange program (PDX)ⁱ the information will be used by the Office to automatically attempt retrieval pursuant to 37 CFR 1.55(i)(1) and (2). Under the PDX program, applicant bears the ultimate responsibility for ensuring that a copy of the foreign application is received by the Office from the participating foreign intellectual property office, or a certified copy of the foreign priority application is filed, within the time period specified in 37 CFR 1.55(g)(1).

			Remove
Application Number	Country ⁱ	Filing Date (YYYY-MM-DD)	Access Code ⁱ (if applicable)
Additional Foreign Priority Data may be generated within this form by selecting the Add button.			Add

Statement under 37 CFR 1.55 or 1.78 for AIA (First Inventor to File) Transition Applications

☐ This application (1) claims priority to or the benefit of an application filed before March 16, 2013 and (2) also contains, or contained at any time, a claim to a claimed invention that has an effective filing date on or after March 16, 2013. NOTE: By providing this statement under 37 CFR 1.55 or 1.78, this application, with a filing date on or after March 16, 2013, will be examined under the first inventor to file provisions of the AIA.

Authorization to Permit Access:

☒ Authorization to Permit Access to the Instant Application by the Participating Offices
--

Application Data Sheet 37 CFR 1.76		Attorney Docket Number	2262-29
		Application Number	
Title of Invention	METHODS AND COMPOSITIONS FOR TREATMENT OF EPILEPTIC DISORDERS		

If checked, the undersigned hereby grants the USPTO authority to provide the European Patent Office (EPO), the Japan Patent Office (JPO), the Korean Intellectual Property Office (KIPO), the World Intellectual Property Office (WIPO), and any other intellectual property offices in which a foreign application claiming priority to the instant patent application is filed access to the instant patent application. See 37 CFR 1.14(c) and (h). This box should not be checked if the applicant does not wish the EPO, JPO, KIPO, WIPO, or other intellectual property office in which a foreign application claiming priority to the instant patent application is filed to have access to the instant patent application.

In accordance with 37 CFR 1.14(h)(3), access will be provided to a copy of the instant patent application with respect to: 1) the instant patent application-as-filed; 2) any foreign application to which the instant patent application claims priority under 35 U.S.C. 119(a)-(d) if a copy of the foreign application that satisfies the certified copy requirement of 37 CFR 1.55 has been filed in the instant patent application; and 3) any U.S. application-as-filed from which benefit is sought in the instant patent application.

In accordance with 37 CFR 1.14(c), access may be provided to information concerning the date of filing this Authorization.

Applicant Information:

Providing assignment information in this section does not substitute for compliance with any requirement of part 3 of Title 37 of CFR to have an assignment recorded by the Office.

Applicant	1	Remove
If the applicant is the inventor (or the remaining joint inventor or inventors under 37 CFR 1.45), this section should not be completed. The information to be provided in this section is the name and address of the legal representative who is the applicant under 37 CFR 1.43; or the name and address of the assignee, person to whom the inventor is under an obligation to assign the invention, or person who otherwise shows sufficient proprietary interest in the matter who is the applicant under 37 CFR 1.46. If the applicant is an applicant under 37 CFR 1.46 (assignee, person to whom the inventor is obligated to assign, or person who otherwise shows sufficient proprietary interest) together with one or more joint inventors, then the joint inventor or inventors who are also the applicant should be identified in this section.		
Clear		
Assignee	Legal Representative under 35 U.S.C. 117	Joint Inventor
Person to whom the inventor is obligated to assign.		Person who shows sufficient proprietary interest
If applicant is the legal representative, indicate the authority to file the patent application, the inventor is:		

Name of the Deceased or Legally Incapacitated Inventor :			
If the Applicant is an Organization check here. ☒			
Organization Name	Ovid Therapeutics Inc.		
Mailing Address Information For Applicant:			
Address 1	1460 Broadway		
Address 2			
City	New York	State/Province	NY
Country	US	Postal Code	10036
Phone Number		Fax Number	

Application Data Sheet 37 CFR 1.76		Attorney Docket Number	2262-29
		Application Number	
Title of Invention	METHODS AND COMPOSITIONS FOR TREATMENT OF EPILEPTIC DISORDERS		
Email Address			
Additional Applicant Data may be generated within this form by selecting the Add button.		Add	

Assignee Information including Non-Applicant Assignee Information:

Providing assignment information in this section does not substitute for compliance with any requirement of part 3 of Title 37 of CFR to have an assignment recorded by the Office.

Assignee	1			
Complete this section if assignee information, including non-applicant assignee information, is desired to be included on the patent application publication. An assignee-applicant identified in the "Applicant Information" section will appear on the patent application publication as an applicant. For an assignee-applicant, complete this section only if identification as an assignee is also desired on the patent application publication.				
				Remove
If the Assignee or Non-Applicant Assignee is an Organization check here. ☐				
Prefix	Given Name	Middle Name	Family Name	Suffix
Mailing Address Information For Assignee including Non-Applicant Assignee:				
Address 1				
Address 2				
City		State/Province		
Country ⁱ		Postal Code		
Phone Number		Fax Number		
Email Address				
Additional Assignee or Non-Applicant Assignee Data may be generated within this form by selecting the Add button.				Add

Signature:

NOTE: This form must be signed in accordance with 37 CFR 1.33. See 37 CFR 1.4 for signature requirements and certifications.					
Signature	/Jeffrey S. Steen, Reg. #32063/			Date (YYYY-MM-DD)	2017-08-10
First Name	Jeffrey S.	Last Name	Steen	Registration Number	32063
Additional Signature may be generated within this form by selecting the Add button.				Add	

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

Application Data Sheet 37 CFR 1.76		Attorney Docket Number	2262-29
		Application Number	
Title of Invention	METHODS AND COMPOSITIONS FOR TREATMENT OF EPILEPTIC DISORDERS		

This collection of information is required by 37 CFR 1.76. The information is required to obtain or retain a benefit by the public which is to file (and by the USPTO to process) an application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.14. This collection is estimated to take 23 minutes to complete, including gathering, preparing, and submitting the completed application data sheet form to the USPTO. Time will vary depending upon the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, P.O. Box 1450, Alexandria, VA 22313-1450. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. **SEND TO: Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.**

Privacy Act Statement

The Privacy Act of 1974 (P.L. 93-579) requires that you be given certain information in connection with your submission of the attached form related to a patent application or patent. Accordingly, pursuant to the requirements of the Act, please be advised that: (1) the general authority for the collection of this information is 35 U.S.C. 2(b)(2); (2) furnishing of the information solicited is voluntary; and (3) the principal purpose for which the information is used by the U.S. Patent and Trademark Office is to process and/or examine your submission related to a patent application or patent. If you do not furnish the requested information, the U.S. Patent and Trademark Office may not be able to process and/or examine your submission, which may result in termination of proceedings or abandonment of the application or expiration of the patent.

The information provided by you in this form will be subject to the following routine uses:

1. The information on this form will be treated confidentially to the extent allowed under the Freedom of Information Act (5 U.S.C. 552) and the Privacy Act (5 U.S.C. 552a). Records from this system of records may be disclosed to the Department of Justice to determine whether the Freedom of Information Act requires disclosure of these records.
2. A record from this system of records may be disclosed, as a routine use, in the course of presenting evidence to a court, magistrate, or administrative tribunal, including disclosures to opposing counsel in the course of settlement negotiations.
3. A record in this system of records may be disclosed, as a routine use, to a Member of Congress submitting a request involving an individual, to whom the record pertains, when the individual has requested assistance from the Member with respect to the subject matter of the record.
4. A record in this system of records may be disclosed, as a routine use, to a contractor of the Agency having need for the information in order to perform a contract. Recipients of information shall be required to comply with the requirements of the Privacy Act of 1974, as amended, pursuant to 5 U.S.C. 552a(m).
5. A record related to an International Application filed under the Patent Cooperation Treaty in this system of records may be disclosed, as a routine use, to the International Bureau of the World Intellectual Property Organization, pursuant to the Patent Cooperation Treaty.
6. A record in this system of records may be disclosed, as a routine use, to another federal agency for purposes of National Security review (35 U.S.C. 181) and for review pursuant to the Atomic Energy Act (42 U.S.C. 218(c)).
7. A record from this system of records may be disclosed, as a routine use, to the Administrator, General Services, or his/her designee, during an inspection of records conducted by GSA as part of that agency's responsibility to recommend improvements in records management practices and programs, under authority of 44 U.S.C. 2904 and 2906. Such disclosure shall be made in accordance with the GSA regulations governing inspection of records for this purpose, and any other relevant (i.e., GSA or Commerce) directive. Such disclosure shall not be used to make determinations about individuals.
8. A record from this system of records may be disclosed, as a routine use, to the public after either publication of the application pursuant to 35 U.S.C. 122(b) or issuance of a patent pursuant to 35 U.S.C. 151. Further, a record may be disclosed, subject to the limitations of 37 CFR 1.14, as a routine use, to the public if the record was filed in an application which became abandoned or in which the proceedings were terminated and which application is referenced by either a published application, an application open to public inspections or an issued patent.
9. A record from this system of records may be disclosed, as a routine use, to a Federal, State, or local law enforcement agency, if the USPTO becomes aware of a violation or potential violation of law or regulation.




FIG. 1

Study Timeline
 (all animals pretreated with Li-Cl (127 mg/kg, i.p.)
 24 hours prior to Study Day)





FIG. 2



VEH (0% protected)

FIG. 3




FIG. 4




FIG. 5





FIG. 6A



Significant Main Effect of Treatment: $F_{(3,23)}=3.123$, $p = 0.045$
 Dunnett's post-hoc test VEH vs. GBD: $p = 0.0545$

FIG. 6B




FIG. 7

2262-29

ABSTRACT

Use of allosteric modulators and/or gaboxadol for the treatment of epileptic disorders in a subject in need thereof.

WHAT IS CLAIMED IS:

1. A method of treating an epileptic disorder comprising administering to a patient in need thereof a pharmaceutical composition comprising an allosteric modulator.
2. The method of claim 1, wherein the epileptic disorder is selected from the group consisting of epilepsy, epilepsy with generalized tonic-clonic seizures, epilepsy with myoclonic absences, frontal lobe epilepsy, temporal lobe epilepsy, Landau-Kleffner Syndrome, Ohtahara syndrome, Rasmussen's syndrome, West's syndrome, Lennox-Gastaut syndrome (LGS), Rett syndrome, CDKL5 disorder, childhood absence epilepsy, essential tremor, Dravet syndrome, Doose syndrome, acute repetitive seizures, benign rolandic epilepsy, status epilepticus, refractory status epilepticus, super-refractory status epilepticus (SRSE), PCDH19 pediatric epilepsy, increased seizure activity or breakthrough seizures.
3. The method of claim 1, wherein the epileptic disorder is status epilepticus.
4. The method of claim 1, wherein the epileptic disorder is acute repetitive seizure.
5. The method of claim 1, wherein the epileptic disorder is Lennox-Gastaut syndrome.
6. The method of claim 1, wherein the composition provides reduction in the frequency of seizures, the severity of seizures, or a combination thereof in a patient diagnosed with an epileptic disorder.
7. The method of claim 1, wherein the allosteric modulator is selected from the group consisting of a neurosteroid, a benzodiazepine and a potassium channel opener.
8. The method of claim 1, wherein the allosteric modulator is ganaxolone.

9. The method of claim 1, wherein the allosteric modulator is allopregnanolone.
10. The method of claim 1, wherein the allosteric modulator is a benzodiazepine.
11. The method of claim 10, wherein the benzodiazepine is selected from the group consisting of midazolam, clobazam, clonazepam, diazepam, lorazepam, flurazepam, and lorazepam.
12. The method of claim 1, wherein the allosteric modulator is a potassium channel opener.
13. The method of claim 12, wherein the potassium channel opener is retigabine or flupirtine.
14. The method of claim 1, wherein the patient is further administered gaboxadol or a pharmaceutically acceptable salt thereof.
15. The method of claim 1, wherein the allosteric modulator is administered once, twice, three times daily, or every other day.
16. The method of claim 1, further comprising administering a compound selected from the group consisting of acetazolamide, carbamazepine, eslicarbazepine acetate, ethosuximide, gabapentin, lacosamide, lamotrigine, leviteracetam, nitrazepam, oxcarbazepine, perampanel, piracetam, phenobarbital, phenytoin, pregabalin, primidone, rufinamide, sodium valproate, stiripentol, tiagabine, topiramate, vigabatrin, and zonisamide.
17. The method of claim 1 wherein the pharmaceutical composition is a parenteral formulation.

18. A method of treating an epileptic disorder comprising administering to a patient in need thereof a pharmaceutical composition comprising gaboxadol or a pharmaceutically acceptable salt thereof.
19. The method of claim 18, wherein the epileptic disorder is selected from the group consisting of epilepsy, epilepsy with generalized tonic-clonic seizures, epilepsy with myoclonic absences, frontal lobe epilepsy, temporal lobe epilepsy, Landau-Kleffner Syndrome, Ohtahara syndrome, Rasmussen's syndrome, West's syndrome, Lennox-Gastaut syndrome (LGS), Rett syndrome, CDKL5 disorder, childhood absence epilepsy, essential tremor, Dravet syndrome, Doose syndrome, acute repetitive seizures, benign rolandic epilepsy, status epilepticus, refractory status epilepticus, super-refractory status epilepticus (SRSE), PCDH19 pediatric epilepsy, increased seizure activity or breakthrough seizures.
20. The method of claim 18, wherein the epileptic disorder is status epilepticus.
21. The method of claim 18, wherein the epileptic disorder is acute repetitive seizure.
22. The method of claim 18, wherein the epileptic disorder is Lennox-Gastaut syndrome.
23. The method of claim 18, wherein the composition provides reduction in the frequency of seizures, the severity of seizures, or a combination thereof in a patient diagnosed with an epileptic disorder.
24. The method of claim 18, wherein the patient is further administered an allosteric modulator.
25. The method of claim 24, wherein the allosteric modulator is selected from the group consisting of a neurosteroid, a benzodiazepine and a potassium channel opener.

26. The method of claim 24, wherein the allosteric modulator is ganaxolone.
27. The method of claim 24, wherein the allosteric modulator is allopregnanolone.
28. The method of claim 24, wherein the allosteric modulator is a benzodiazepine.
29. The method of claim 28, wherein the benzodiazepine is selected from the group consisting of midazolam, clobazam, clonazepam, diazepam, lorazepam, flurazepam, and lorazepam.
30. The method of claim 24, wherein the allosteric modulator is a potassium channel opener.
31. The method of claim 30, wherein the potassium channel opener is retigabine or flupirtine.
32. The method of claim 18, wherein the gaboxadol or a pharmaceutically acceptable salt thereof is administered once, twice, three times daily, or every other day.
33. The method of claim 32, wherein the gaboxadol or a pharmaceutically acceptable salt thereof is administered immediately after the onset of a seizure.
34. The method of claim 32, wherein the gaboxadol or a pharmaceutically acceptable salt thereof is administered after the onset of a seizure warning sign.
35. The method of claim 18, further comprising administering a compound selected from the group consisting of acetazolamide, carbamazepine, eslicarbazepine acetate, ethosuximide, gabapentin, lacosamide, lamotrigine, leviteracetam, nitrazepam, oxcarbazepine, perampanel, piracetam, phenobarbital, phenytoin, pregabalin, primidone, rufinamide, sodium valproate, stiripentol, tiagabine, topiramate, vigabatrin, and zonisamide.

36. The method of claim 18 wherein the pharmaceutical composition is a parenteral formulation.
37. A pharmaceutical composition for parenteral administration comprising about 0.005 µg/ml to about 500 µg/ml gaboxadol or a pharmaceutically acceptable salt thereof.
38. A pharmaceutical composition for parenteral administration comprising about 0.05 mg to about 100 mg gaboxadol or a pharmaceutically acceptable salt thereof.
39. A pharmaceutical composition for parenteral administration according to claims 37 or 38 wherein the gaboxadol or a pharmaceutically acceptable salt thereof is present at a molarity of less than about 10.0 M.
40. A pharmaceutical composition for parenteral administration according to claims 37 or 38 wherein the solubility of gaboxadol or salt thereof in the composition is between about 1 mg/mL to about 50 mg/mL.
41. A pharmaceutical composition for parenteral administration according to claims 37 or 38 wherein the pharmaceutical composition exhibits a time of maximum plasma concentration (T_{max}) for gaboxadol of about 1 hour or more after administration of the parenteral dosage form.
42. A pharmaceutical composition for parenteral administration according to claims 37 or 38 wherein parenteral administration of the composition provides an *in vivo* plasma profile of gaboxadol comprising a mean $AUC_{0-\infty}$ of more than about 25 ng·hr/ml.
43. A pharmaceutical composition for parenteral administration according to claims 37 or 38 wherein parenteral administration of the composition provides an *in vivo* plasma profile of gaboxadol comprising a mean C_{max} of less than about 10000 ng/ml.

44. A pharmaceutical composition for parenteral administration according to claims 37 or 38 wherein parenteral administration exhibits a pharmacokinetic profile comprising a T_{max} at about 1 to about 120 minutes after administration of the parenteral composition; followed by a plasma drug concentration of at least 50% C_{max} for a duration of about 90 to about 360 minutes.
45. A pharmaceutical composition for parenteral administration according to claims 37 or 38 wherein the composition is contained in a bag, a glass vial, a plastic vial, or a bottle.
46. A pharmaceutical composition for parenteral administration according to claims 37 or 38 wherein the composition is aqueous.
47. A pharmaceutical composition for parenteral administration according to claims 37 or 38 wherein the composition is ready-to-use.
48. A pharmaceutical composition for parenteral administration according to claims 37 or 38 wherein the composition is sufficiently soluble.
49. A pharmaceutical composition for parenteral administration according to claims 37 or 38 wherein the composition further comprises an excipient selected from the group consisting of a buffering agent, solubilizing agent, tonicity agent, antioxidant, chelating agent, antimicrobial agent and preservative.
50. A pharmaceutical composition for parenteral administration according to claims 37 or 38 wherein the composition further comprises an excipient which is present at a weight percent (w/v) of less than about 10%.
51. A pharmaceutical composition for parenteral administration according to claims 37 or 38 wherein the composition further comprises an excipient which is present at a weight percent (w/v) between about 0.01% to about 10%.

52. A pharmaceutical composition for parenteral administration according to claims 37 or 38 wherein the excipient is present in a molar ratio of the excipient to gaboxadol or pharmaceutically acceptable salt thereof of about 0.1:1 to about 0.25:1.
53. A pharmaceutical composition for parenteral administration according to claims 37 or 38 wherein the composition comprises an excipient which includes a stabilizing amount of a buffering agent.
54. A pharmaceutical composition for parenteral administration according to claims 37 or 38 wherein the composition is at a pH of about 4 to about 8.
55. A pharmaceutical composition for parenteral administration according to claims 37 or 38 wherein the composition is at a pH of about 6 to about 8.
56. A pharmaceutical composition for parenteral administration according to claims 37 or 38 wherein composition is an aqueous solution and the pH of the aqueous solution is from about 6.8 to about 7.8.
57. A pharmaceutical composition for parenteral administration according to claims 37 or 38 wherein the composition further comprises sodium chloride at a concentration of between about 0.01 and about 2.0 weight percent.
58. A pharmaceutical composition for parenteral administration according to claims 37 or 38 wherein the composition further comprises sodium chloride at a concentration of about 0.9 weight percent.
59. A pharmaceutical composition for parenteral administration according to claims 37 or 38 wherein the composition is formulated as a total volume selected from the group consisting of 20 ml, 50 ml, and 100 ml.

60. A pharmaceutical composition for parenteral administration according to claims 37 or 38 wherein the composition is prepared for subcutaneous, intramuscular, transdermal, or intravenous administration.

61. A pharmaceutical composition for parenteral administration according to claims 37 or 38 wherein the composition comprises an excipient which is a stabilizing amount of a solubilizing agent.

62. A pharmaceutical composition for parenteral administration according to claims 37 or 38 wherein the composition shows no detectable chemical degradation after one month at 40°C.

63. A pharmaceutical composition for parenteral administration according to claims 37 or 38 wherein the composition can be stored at ambient conditions and remains clear and colorless for at least about 12 weeks.

METHODS AND COMPOSITIONS FOR TREATMENT OF EPILEPTIC DISORDERS

CROSS REFERENCE TO RELATED APPLICATIONS

- 5 This application claims benefit of and priority to U.S. Provisional Application No. 62/373,589, filed August 11, 2016 and U.S. Provisional Application No. 62/490,293, filed April 26, 2017, which are incorporated herein by reference in their respective entireties.

TECHNICAL FIELD

- 10 Methods of using allosteric modulators and/or gaboxadol or pharmaceutically acceptable salts thereof for the treatment of epileptic disorders in a subject in need thereof.

BACKGROUND

- Allosteric modulators such as neurosteroids (*e.g.*, ganaxolone, allopregnanolone),
15 benzodiazapines (*e.g.*, diazepam) and potassium channel openers (*e.g.*, ritigabine) have been used in the treatment of epilepsy. However, treatment with these agents is often limited to patients that do not respond to traditional medications. For example, allopregnanolone is currently in development for the treatment of super refractory status epilepticus. In addition, diazepam is currently marketed (Diastat®) for use in emergency situations to stop cluster
20 seizures in people who are taking other medications to treat epilepsy.

- Gaboxadol (4,5,6,7-tetrahydroisoxazolo [5,4-c] pyridine-3-ol) (THIP)) is described in EP Patent No. 0000338, in EP Patent No. 0840601, and in U.S. Patent Nos. 4,278,676, 4,362,731, 4,353,910, and WO 2005/094820. Gaboxadol is a selective GABA_A receptor agonist with a preference for δ -subunit containing GABA_A receptors. In the early 1980s gaboxadol was the
25 subject of a series of pilot studies that tested its efficacy as an analgesic and anxiolytic, as well as a treatment for tardive dyskinesia, Huntington's disease, Alzheimer's disease, and spasticity. In the 1990s gaboxadol moved into late stage development for the treatment of insomnia but failed to show significant effects in sleep onset and sleep maintenance in a three-month efficacy study. Additionally, patients with a history of drug abuse who received gaboxadol experienced a steep
30 increase in psychiatric adverse events. As a result of these negative results the development of gaboxadol was terminated.

Parenteral dosage forms are intended for administration as an injection or infusion. Common injection types are intravenous (into a vein), subcutaneous (under the skin), and intramuscular (into muscle). Infusions typically are given by intravenous route. Parenteral formulations often include excipients to enhance or maintain active ingredient solubility (solubilizers) and/or stability (buffers, antioxidants, chelating agents, cryo- and lyoprotectants). Excipients also are important in parenteral formulations to assure safety (antimicrobial preservatives), minimize pain and irritation upon injection (tonicity agents), and control or prolong drug delivery (polymers). However, excipients may also produce negative effects such as loss of drug solubility, activity, and/or stability.

There remains a need in the art for safe and effective methods and pharmaceutical compositions that provide epileptic treatment. Accordingly, this disclosure provides pharmaceutical compositions and methods that may be used in applications of epileptic disorders, such as status epilepticus.

SUMMARY

Methods are provided for treatment of epileptic disorders including epilepsy, epilepsy with generalized tonic-clonic seizures, epilepsy with myoclonic absences, frontal lobe epilepsy, temporal lobe epilepsy, Landau-Kleffner Syndrome, Ohtahara syndrome, Rasmussen's syndrome, West's syndrome, Lennox-Gastaut syndrome (LGS), Rett syndrome, CDKL5 disorder, childhood absence epilepsy, essential tremor, Dravet syndrome, Doose syndrome, acute repetitive seizures, benign rolandic epilepsy, status epilepticus, refractory status, epilepticus, super-refractory status epilepticus (SRSE), PCDH19 pediatric epilepsy, increased seizure activity or breakthrough seizures (increased seizure activity, also called serial or cluster seizures) and sodium channel protein type 1 subunit alpha (Scn1a)-related disorders by administering to a patient in need thereof a pharmaceutical composition containing an allosteric modulator. Allosteric modulators include one or more of neurosteroids, benzodiazapines, and potassium channel openers. In embodiments, methods of treating epileptic disorders are provided which include administering to a patient in need thereof a pharmaceutical composition including an allosteric modulator in combination with gaboxadol or a pharmaceutically acceptable salt thereof.

Parenteral formulations of gaboxadol or pharmaceutically acceptable salts thereof are provided herein. Methods of treating epileptic disorders, including status epilepticus, with parenteral formulations of gaboxadol or a pharmaceutically acceptable salt thereof are provided. In embodiments, parenteral formulations including gaboxadol or a pharmaceutically acceptable salt thereof, alone or in combination with an allosteric modulator, are administered to a patient in
 5 need thereof to treat an epileptic disorder.

In embodiments, methods are provided for treatment of epileptic disorders including status epilepticus, benign rolandic epilepsy (BRE), intractable childhood epilepsy (ICE), childhood absence epilepsy (CAE), juvenile myoclonic epilepsy (JME), infantile spasms (or
 10 West syndrome), Dravet syndrome and Lennox-Gastaut syndrome (LGS) by administering to a patient in need thereof a pharmaceutical composition including an allosteric modulator either alone or in combination with gaboxadol or a pharmaceutically acceptable salt thereof. In embodiments, methods are provided for treatment of epileptic disorders including status epilepticus, benign rolandic epilepsy (BRE), intractable childhood epilepsy (ICE), childhood
 15 absence epilepsy (CAE), juvenile myoclonic epilepsy (JME), infantile spasms (or West syndrome), Dravet syndrome and Lennox-Gastaut syndrome (LGS) by administering to a patient in need thereof a pharmaceutical composition including gaboxadol or a pharmaceutically acceptable salt thereof either alone or in combination with an allosteric modulator.

In embodiments, methods are provided for treatment of epileptic disorders characterized
 20 as a sodium channel protein type 1 subunit alpha (Scn1a)-related disorder. Scn1a-related disorders include generalized epilepsy with febrile seizures plus, intractable childhood epilepsy with generalized tonic-clonic seizures, intractable infantile partial seizures, myoclonic-astatic epilepsy, severe myoclonic epilepsy in infancy, simple febrile seizures, Dravet syndrome, Lennox-Gastaut syndrome (LGS), infantile spasms, and vaccine-related encephalopathy and
 25 seizures. In embodiments, methods are provided for treatment of a sodium channel protein type 1 subunit alpha (Scn1a)-related disorder by administering to a patient in need thereof a pharmaceutical composition including an allosteric modulator. In embodiments, methods are provided for treatment of a sodium channel protein type 1 subunit alpha (Scn1a)-related disorder by administering to a patient in need thereof a pharmaceutical composition including an
 30 allosteric modulator in combination with gaboxadol or a pharmaceutically acceptable salt thereof. In embodiments, methods are provided for treatment of a sodium channel protein type 1

subunit alpha (Scn1a)-related disorder by administering to a patient in need thereof a pharmaceutical composition including gaboxadol or a pharmaceutically acceptable salt thereof. In embodiments, methods are provided for treatment of a sodium channel protein type 1 subunit alpha (Scn1a)-related disorder by administering to a patient in need thereof a pharmaceutical composition including gaboxadol or a pharmaceutically acceptable salt thereof in combination with an allosteric modulator. In embodiments, methods are provided for treatment of a sodium channel protein type 1 subunit alpha (Scn1a)-related disorder by administering to a patient in need thereof a parenteral formulation including gaboxadol or a pharmaceutically acceptable salt thereof. In embodiments, methods are provided for treatment a sodium channel protein type 1 subunit alpha (Scn1a)-related disorder by administering to a patient in need thereof a pharmaceutical composition including an allosteric modulator in combination with a parenteral formulation including gaboxadol or a pharmaceutically acceptable salt thereof.

In embodiments, a combination of allosteric modulators, *e.g.*, a neurosteroid, benzodiazepine, or potassium channel opener, may be administered to a patient in need thereof. In embodiments, a combination of one or more allosteric modulators and gaboxadol or a pharmaceutically acceptable salt thereof is administered to a patient in need thereof.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 shows both the theoretical and measured solubility of gaboxadol at different pH values.

FIG. 2 is a schematic drawing depicting a timeline for an evaluation study of the ability of allopregnanolone, ganaxolone and gaboxadol to block benzodiazepine resistant status epilepticus in rats.

FIG. 3 is a bar graph depicting the percent protected versus dose with respect to allopregnanolone, ganaxalone, or gaboxadol.

FIG. 4 is a bar graph depicting the 24-hour survival results based on dose with respect to allopregnanolone, ganaxalone, or gaboxadol.

FIG. 5 is a bar graph showing the number of observed seizures versus dose of allopregnanolone, ganaxalone, or gaboxadol.

FIG. 6A is a bar graph showing body weight changes 24 hours post-status epilepticus as a function of percent loss versus dose.

FIG. 6B is a bar graph showing 24 hour body weight loss for 0.5 mg/kg dose groups.

FIG. 7 is a schematic drawing depicting a timeline for a prospective evaluation study of the ability of allopregnanolone, ganaxolone and gaboxadol to synergistically block benzodiazepine resistant status epilepticus in rats.

5

DETAILED DESCRIPTION

Described herein are methods of treating epileptic disorders including epilepsy, epilepsy with generalized tonic-clonic seizures, epilepsy with myoclonic absences, frontal lobe epilepsy, temporal lobe epilepsy, Landau-Kleffner Syndrome, Ohtahara syndrome, Rasmussen's syndrome, infantile spasms (or West syndrome), Lennox-Gastaut syndrome (LGS), Rett syndrome, Dravet syndrome, Doose syndrome, CDKL5 disorder, intractable childhood epilepsy (ICE), childhood absence epilepsy (CAE), juvenile myoclonic epilepsy (JME), essential tremor, acute repetitive seizures, benign rolandic epilepsy, status epilepticus, refractory status epilepticus, super-refractory status epilepticus (SRSE), PCDH19 pediatric epilepsy, increased seizure activity or breakthrough seizures (increased seizure activity; also called serial or cluster seizures). Compositions and methods described herein may be used to treat epileptic disorders characterized as a sodium channel protein type 1 subunit alpha (Scn1A)-related disorder. For example, Scn1A-related disorders include generalized epilepsy with febrile seizures, intractable childhood epilepsy with generalized tonic-clonic seizures, intractable infantile partial seizures, myoclonic-astatic epilepsy, severe myoclonic epilepsy in infancy, simple febrile seizures, Dravet syndrome, Lennox-Gastaut syndrome, infantile spasms, and vaccine-related encephalopathy and seizures. The compositions and methods described herein involve allosteric modulators and/or gaboxadol or a pharmaceutically acceptable salt thereof.

In embodiments, a method of treating an epileptic disorder may include administering to a patient in need thereof a pharmaceutical composition including an allosteric modulator. In embodiments, a method of treating an epileptic disorder may include administering to a patient in need thereof a pharmaceutical composition including gaboxadol or a pharmaceutically acceptable salt thereof. In embodiments, a method of treating an epileptic disorder may include administering to a patient in need thereof gaboxadol or a pharmaceutically acceptable salt thereof, in combination with an allosteric modulator. In embodiments, a method of treating an epileptic disorder may include administering to a patient in need thereof a parenteral

pharmaceutical formulation including an allosteric modulator. In embodiments, a method of treating an epileptic disorder may include administering to a patient in need thereof a parenteral pharmaceutical composition including gaboxadol or a pharmaceutically acceptable salt thereof. In embodiments, a method of treating an epileptic disorder may include administering to a
5 patient in need thereof a parenteral pharmaceutical composition including an allosteric modulator and gaboxadol or a pharmaceutically acceptable salt thereof.

Many pharmaceutical products are administered as a fixed dose, at regular intervals, to achieve therapeutic efficacy. The duration of action is reflected by the product's plasma half-life. Since efficacy is often dependent on sufficient exposure within the central nervous system
10 administration of CNS drugs with a short half-life may require frequent maintenance dosing. Advantageously disclosed herein are methods of treating epileptic disorders by administration of an allosteric modulator. For example, in embodiments, methods of treating an epileptic disorder are provided which include administering to a patient in need thereof a pharmaceutical composition including about 0.05 mg to about 2000 mg of an allosteric modulator wherein the
15 composition provides improvement for more than 6 hours after administration to the patient. Advantageously disclosed herein are methods of treating epileptic disorders by administration of gaboxadol or pharmaceutically acceptable salt thereof. For example, in embodiments, methods of treating an epileptic disorder are provided which include administering to a patient in need thereof a pharmaceutical composition including about 0.05 mg to about 75 mg of gaboxadol or
20 pharmaceutically acceptable salt thereof, wherein the composition provides improvement for more than 6 hours after administration to the patient.

In embodiments, methods of treating an epileptic disorder include administering to a patient in need thereof a pharmaceutical composition including about 0.05 mg to about 50 mg gaboxadol or a pharmaceutically acceptable salt thereof. In embodiments, methods of treating a
25 epileptic disorder include administering to a patient in need thereof a pharmaceutical composition including about 0.1 mg to about 30 mg gaboxadol or a pharmaceutically acceptable salt thereof.

For example, dosages may include amounts of gaboxadol or pharmaceutically acceptable salt thereof in the range of about, *e.g.*, 0.05 mg to 50 mg, 1 mg to 30 mg, 1 mg to 20 mg, 1 mg to
30 15 mg, 0.01 mg to 10 mg, 0.1 mg to 15 mg, 0.1 mg to 30 mg, 0.15 mg to 12.5 mg, or 0.2 mg to 10 mg, with doses of 0.05 mg, 0.1 mg, 0.2 mg, 0.3 mg, 0.4 mg, 0.5 mg, 0.6 mg, 0.7 mg, 0.8 mg,

0.9 mg, 1.5 mg, 1.0 mg, 1.75 mg, 2 mg, 2.5 mg, 2.75 mg, 3 mg, 3.5 mg, 3.75 mg, 4 mg, 4.5 mg, 4.75 mg, 5 mg, 5.5 mg, 6 mg, 6.5 mg, 7 mg, 7.5 mg, 8 mg, 8.5 mg, 9 mg, 10 mg, 11 mg, 12 mg, 15 mg, 20 mg, 25 mg, and 30 mg being specific examples of doses.

Typically, dosages of gaboxadol or a pharmaceutically acceptable salt thereof are administered once or twice daily to a patient in need thereof. The methods and compositions described herein may provide reduced dosing frequency and reduced adverse events and/or increased efficacy. In embodiments, the dosage is about, *e.g.*, 0.05-30 mg/day, 0.1-20 mg/day, or 0.2-15 mg/day, or 0.5-10 mg/day, or 0.75-5 mg/day, for example 0.1 mg/day, 0.2 mg/day, 0.5 mg/day, 0.75 mg/day, 1 mg/day, 1.5 mg/day, 2 mg/day, 3 mg/day, 4 mg/day, 5 mg/day, 6 mg/day, 7 mg/day, 8 mg/day, 9 mg/day, 10 mg/day, 11 mg/day, 12 mg/day, 13 mg/day, 14 mg/day, 15 mg/day, 16 mg/day, 17 mg/day, 18 mg/day, 19 mg/day, 20 mg/day, 21 mg/day, 22 mg/day, 23 mg/day, 24 mg/day, 25 mg/day, 26 mg/day, 27 mg/day, 28 mg/day, 29 mg/day, or 30 mg/day. In embodiments, gaboxadol or a pharmaceutically acceptable salt thereof, or a derivative or analogue thereof is administered at doses of 0.2 mg to 1 mg in infants or 1-20 mg in adults, once daily.

In embodiments, the pharmaceutical compositions include 0.1 mg to 25 mg, 0.1 mg to 20 mg, 0.1 mg to 15 mg, 0.5 mg to 25 mg, 0.5 mg to 20 mg, 0.5 to 15 mg, 1 mg to 25 mg, 1 mg to 20 mg, 1 mg to 15 mg, 1.5 mg to 25 mg, 1.5 mg to 20 mg, 1.5 mg to 15 mg, 2 mg to 25 mg, 2 mg to 20 mg, 2 mg to 15 mg, 2.5 mg to 25 mg, 2.5 mg to 20 mg, 2.5 mg to 15 mg, 3 mg to 25 mg, 3 mg to 20 mg, 3 mg to 15 mg gaboxadol or a pharmaceutically acceptable salt thereof.

In embodiments, the pharmaceutical compositions include 5 mg to 20 mg, 5 mg to 10 mg, 4 mg to 6 mg, 6 mg to 8 mg, 8 mg to 10 mg, 10 mg to 12 mg, 12 mg to 14 mg, 14 mg to 16 mg, 16 mg to 18 mg, or 18 mg to 20 mg gaboxadol or a pharmaceutically acceptable salt thereof.


In embodiments, the pharmaceutical compositions include 0.1 mg, 0.25 mg, 0.5 mg, 1 mg, 2.5 mg, 3 mg, 4 mg, 5 mg, 7 mg, 7.5 mg, 10 mg, 12.5 mg, 15 mg, 17.5 mg, 20 mg gaboxadol or a pharmaceutically acceptable salt thereof or amounts that are multiples of such doses. In embodiments, the pharmaceutical compositions include 2.5 mg, 5 mg, 7.5 mg, 10 mg, 15 mg, or 20 mg gaboxadol or a pharmaceutically acceptable salt thereof.

In embodiments, the total amount of gaboxadol or pharmaceutically acceptable salt thereof and/or gaboxadol administered to a subject in a 24-hour period is 1 mg to 50 mg. In embodiments, the total amount of gaboxadol or pharmaceutically acceptable salt thereof and/or

gaboxadol administered to a subject in a 24-hour period is 1 mg to 20 mg. In embodiments, the total amount of gaboxadol or pharmaceutically acceptable salt thereof and/or gaboxadol administered to a subject in a 24-hour period is 5 mg, 10 mg, 15 mg or 20 mg. In embodiments, the total amount of gaboxadol or a pharmaceutically acceptable salt thereof administered to a subject in a 24-hour period is 1 mg to 50 mg. In embodiments, the subject may be started at a low dose and the dosage is escalated. In this manner, it can be determined if the drug is well tolerated in the subject. Dosages can be lower for children than for adults. In embodiments, a dose of gaboxadol for children can be 0.1 mg/kg to 1 mg/kg.

Allosteric modulators may include a neurosteroid, *e.g.*, ganaxolone or allopregnanolone, a benzodiazepine, *e.g.*, midazolam, clobazam, clonazepam, diazepam, lorazepam, flurazepam, lorazepam etc., or a potassium channel opener, *e.g.*, retigabine or flupirtine.

In embodiments, methods are provided for treating an epileptic disorder by administering ganaxolone to a patient in need thereof. In embodiments, methods are provided for treating an epileptic disorder by administering allopregnanolone to a patient in need thereof. In embodiments methods are provided for treating an epileptic disorder by administering a compound of Formula I:



Formula I

In embodiments, an allosteric modulator or a pharmaceutically acceptable salt thereof is administered at dosages ranging from about 0.001 mg/kg to about 30 mg/kg of body weight of a patient in need thereof, *e.g.*, from about 0.01 mg/kg to 20 mg/kg at least once a day. For example, dosages may include amounts of an allosteric modulator or a pharmaceutically acceptable salt thereof in the range of about, *e.g.*, 1 mg to 30 mg, 1 mg to 25 mg, 1 mg to 20 mg, 1 mg to 15 mg, 1 mg to 10 mg, 0.01 mg to 10 mg, 0.1 mg to 15 mg, 0.15 mg to 12.5 mg, or 0.1 mg to 10 mg, or 0.2 mg to 10 mg, with doses of 0.1 mg, 0.2 mg, 0.3 mg, 0.4 mg, 0.5 mg, 0.6 mg, 0.7 mg, 0.8 mg, 0.9 mg, 1.5 mg, 1.0 mg, 1.75 mg, 2 mg, 2.5 mg, 2.75 mg, 3 mg, 3.5 mg, 3.75 mg, 4 mg, 4.5 mg, 4.75 mg, 5 mg, 5.5 mg, 6 mg, 6.5 mg, 7 mg, 7.5 mg, 8 mg, 8.5 mg, 9 mg, 10 mg, 11 mg, 12 mg, 13 mg, 14 mg, 15 mg, 16 mg, 17 mg, 18 mg, 19 mg, 20 mg, 21 mg, 22 mg,

23 mg, 24 mg, 25 mg, 26 mg, 27 mg 28 mg, 29 mg, and 30 mg being specific examples of doses. For example, dosages may include amounts of an allosteric modulator or a pharmaceutically acceptable salt thereof in the range of about, e.g., 50 mg to 75 mg, 75 mg to 100 mg, 100 mg to 125 mg, 125 mg to 150 mg, 150 mg to 175 mg, 175 mg to 200 mg, 200 mg to 225 mg, 225 mg to 250 mg, 250 mg to 275 mg, 275 mg to 300 mg, 300 mg to 325 mg, 325 mg to 350 mg, 350 mg to 375 mg, 375 mg to 400 mg, 400 mg to 425 mg, 425 mg to 450 mg, 450 mg to 475 mg, 475 mg to 500 mg, 500 mg to 525 mg, 525 mg to 550 mg, 550 mg to 575 mg, 575 mg to 600 mg, 600 mg to 625 mg, 625 mg to 650 mg, 650 mg to 675 mg, 675 mg to 700 mg, 700 mg to 725 mg, 725 mg to 750 mg, 750 mg to 775 mg, 775 mg to 800 mg, 800 mg to 825 mg, 825 mg to 850 mg, 850 mg to 875 mg, 875 mg to 900 mg, 900 mg to 925 mg, 925 mg to 950 mg, 950 mg to 975 mg, 975 mg to 1000 mg, 1000 mg to 1025 mg, 1025 mg to 1050 mg, 1050 mg to 1075 mg, 1075 mg to 1100 mg, 1100 mg to 1125 mg, 1125 mg to 1150 mg, 1150 mg to 1175 mg, 1175 mg to 1200 mg, 1200 mg to 1225 mg, 1225 mg to 1250 mg, 1250 mg to 1275 mg, 1275 mg to 1300 mg, 1300 mg to 1325 mg, 1325 mg to 1350 mg, 1350 mg to 1375 mg, 1375 mg to 1400 mg, 1400 mg to 1425 mg, 1425 mg to 1450 mg, 1450 mg to 1475 mg, 1475 mg to 1500 mg, 1500 mg to 1525 mg, 1525 mg to 1550 mg, 1550 mg to 1575 mg, 1575 mg to 1600 mg, 1600 mg to 1625 mg, 1625 mg to 1650 mg, 1650 mg to 1675 mg, 1675 mg to 1700 mg, 1700 mg to 1725 mg, 1725 mg to 1750 mg, 1750 mg to 1775 mg, 1775 mg to 1800 mg, 1800 mg to 1825 mg, 1825 mg to 1850 mg, 1850 mg to 1875 mg, 1875 mg to 1900 mg, 1900 mg to 1925 mg, 1925 mg to 1950 mg, 1950 mg to 1975 mg, or 1975 mg to 2000 mg, of an allosteric modulator.

Typically, dosages of an allosteric modulator or pharmaceutically acceptable salts thereof are administered once daily, twice daily, three times daily or four times daily to a patient in need thereof. In embodiments, allosteric modulators may be administered once weekly. The methods and compositions described herein may provide reduced dosing frequency and reduced adverse events and/or increased efficacy. In embodiments, the dosage of an allosteric modulator can be about, e.g., 0.1-20 mg/day, or 0.2-15 mg/day, or 0.5-10 mg/day, or 0.75-5 mg/day, for example 0.2 mg/day, 0.5 mg/day, 0.75 mg/day, 1 mg/day, 1.5 mg/day, 2 mg/day, 3 mg/day, 4 mg/day, 5 mg/day, 6 mg/day, 7 mg/day, 8 mg/day, 9 mg/day, or 10 mg/day. In embodiments, a patient can be administered an allosteric modulator in an amount of, e.g., 10 mg to 25 mg/day, 25 mg to 50 mg/day, 50 mg to 75 mg/day, 75 mg to 100 mg/day, 100 mg to 125 mg/day, 125 mg to 150 mg/day, 150 mg to 175 mg/day, 175 mg to 200 mg/day, 200 mg to 225 mg/day, 225 mg to 250

mg/day, 250 mg to 275 mg/day, 275 mg to 300 mg/day, 300 mg to 325 mg/day, 325 mg to 350 mg/day, 350 mg to 375 mg/day, 375 mg to 400 mg/day, 400 mg to 425 mg/day, 425 mg to 450 mg/day, 450 mg to 475 mg/day, 475 mg to 500 mg/day, 500 mg to 525 mg/day, 525 mg to 550 mg/day, 550 mg to 575 mg/day, 575 mg to 600 mg/day, 600 mg to 625 mg/day, 625 mg to 650 mg/day, 650 mg to 675 mg/day, 675 mg to 700 mg/day, 700 mg to 725 mg/day, 725 mg to 750 mg/day, 750 mg to 775 mg/day, 775 mg to 800 mg/day, 800 mg to 825 mg/day, 825 mg to 850 mg/day, 850 mg to 875 mg/day, 875 mg to 900 mg/day, 900 mg to 925 mg/day, 925 mg to 950 mg/day, 950 mg to 975 mg/day, 975 mg to 1000 mg/day, 1000 mg to 1025 mg/day, 1025 mg to 1050 mg/day, 1050 mg to 1075 mg/day, 1075 mg to 1100 mg/day, 1100 mg to 1125 mg/day, 1125 mg to 1150 mg/day, 1150 mg to 1175 mg/day, 1175 mg to 1200 mg/day, 1200 mg to 1225 mg/day, 1225 mg to 1250 mg/day, 1250 mg to 1275 mg/day, 1275 mg to 1300 mg/day, 1300 mg to 1325 mg/day, 1325 mg to 1350 mg/day, 1350 mg to 1375 mg/day, 1375 mg to 1400 mg/day, 1400 mg to 1425 mg/day, 1425 mg to 1450 mg/day, 1450 mg to 1475 mg/day, 1475 mg to 1500 mg/day, 1500 mg to 1525 mg/day, 1525 mg to 1550 mg/day, 1550 mg to 1575 mg/day, 1575 mg to 1600 mg/day, 1600 mg to 1625 mg/day, 1625 mg to 1650 mg/day, 1650 mg to 1675 mg/day, 1675 mg to 1700 mg/day, 1700 mg to 1725 mg/day, 1725 mg to 1750 mg/day, 1750 mg to 1775 mg/day, 1775 mg to 1800 mg/day, 1800 mg to 1825 mg/day, 1825 mg to 1850 mg/day, 1850 mg to 1875 mg/day, 1875 mg to 1900 mg/day, 1900 mg to 1925 mg/day, 1925 mg to 1950 mg/day, 1950 mg to 1975 mg/day, or 1975 mg to 2000 mg/day. In embodiments, an allosteric modulator, or a derivative or analogue thereof can be administered at doses of 0.2 mg to 1 mg in infants or 1-20 mg in adults, once daily.

In embodiments, a method of treating an epileptic disorder such as status epilepticus includes administering ganaxolone or a pharmaceutically acceptable salt thereof to a patient in need thereof. Ganaxolone or a pharmaceutically acceptable salt thereof can be administered in doses ranging from 10 mg/kg to 40 mg/kg, e.g., 11 mg/kg to 39 mg/kg, 12 mg/kg to 38 mg/kg, 13 mg/kg to 37 mg/kg, 14 mg/kg to 36 mg/kg, 15 mg/kg to 35 mg/kg, 16 mg/kg to 34 mg/kg, 17 mg/kg to 33 mg/kg, 18 mg/kg to 32 mg/kg, 19 mg/kg to 31 mg/kg, 20 mg/kg to 30 mg/kg, 21 mg/kg to 29 mg/kg, 22 mg/kg to 28 mg/kg, 23 mg/kg to 27 mg/kg, or 24 mg/kg to 26 mg/kg. In embodiments, ganaxolone doses can be, e.g., 50 mg, 75 mg, 100 mg, 125 mg, 150 mg, 175 mg, 200 mg, 225 mg, 250 mg, 275 mg, 300 mg, 325 mg, 350 mg, 375 mg, 400 mg, 425 mg, 450 mg, 475 mg, 500 mg, 525 mg, 550 mg, 575 mg, 600 mg, 625 mg, 650 mg, 675 mg, 700 mg, 725 mg,

750 mg, 775 mg, 800 mg, 825 mg, 850 mg, 875 mg, 900 mg, 925 mg, 950 mg, 975 mg, 1000 mg, 1225 mg, 1250 mg, 1275 mg, 1300 mg, 1325 mg, 1350 mg, 1375 mg, 1400 mg, 1425 mg, 1450 mg, 1475 mg, 1500 mg, 1525 mg, 1550 mg, 1575 mg, 1600 mg, 1625 mg, 1650 mg, 1675 mg, 1700 mg, 1725 mg, 1750 mg, 1775 mg, 1800 mg, 1825 mg, 1850 mg, 1875 mg, 1900 mg, 1925 mg, 1950 mg, 1975 mg, or 2000 mg.

Ganaxolone or a pharmaceutically acceptable salt thereof can be administered, e.g., once daily, twice daily, three times daily, or four times daily. In embodiments, ganaxolone or a pharmaceutically acceptable salt thereof can be administered once weekly. In embodiments, ganaxolone or a pharmaceutically acceptable salt thereof can be administered parenterally as soon as possible after the onset of seizures. In embodiments, ganaxolone or a pharmaceutically acceptable salt thereof can be administered parenterally in escalating doses as soon as possible after the onset of seizures.

In embodiments, a method of treating an epileptic disorder such as status epilepticus includes administering allopregnanolone or a pharmaceutically acceptable salt thereof to a patient in need thereof. Allopregnanolone or a pharmaceutically acceptable salt thereof can be administered in doses ranging, e.g., from 0.01 mg/kg to 20 mg/kg, 0.02 mg/kg to 19 mg/kg, 0.03 mg/kg to 18 mg/kg, 0.04 mg/kg to 17 mg/kg, 0.05 mg/kg to 16 mg/kg, 0.06 mg/kg to 15 mg/kg, 0.07 mg/kg to 14 mg/kg, 0.08 mg/kg to 14 mg/kg, 0.09 mg/kg to 13 mg/kg, 0.1 mg/kg to 12 mg/kg, 0.2 mg/kg to 11 mg/kg, 0.3 mg/kg to 10 mg/kg, 0.4 mg/kg to 9 mg/kg, 0.5 mg/kg to 8 mg/kg, 0.6 mg/kg to 7 mg/kg, 0.7 mg/kg to 6 mg/kg, 0.8 mg/kg to 5 mg/kg, 0.9 mg/kg to 4 mg/kg, or 1 mg/kg to 3 mg/kg. In embodiments, allopregnanolone doses can be, e.g., 1 mg, 2 mg, 3 mg, 4 mg, 5 mg, 6 mg, 7 mg, 8 mg, 9 mg, 10 mg, 11 mg, 12 mg, 13 mg, 14 mg, 15 mg, 16 mg, 17 mg, 18 mg, 19 mg, or 20 mg.

Allopregnanolone or a pharmaceutically acceptable salt thereof can be administered, e.g., once daily, twice daily, three times daily, or four times daily. In embodiments, allopregnanolone or a pharmaceutically acceptable salt thereof can be administered once weekly. In embodiments, allopregnanolone or a pharmaceutically acceptable salt thereof can be administered parenterally as soon as possible after the onset of seizures. In embodiments, allopregnanolone or a pharmaceutically acceptable salt thereof can be administered parenterally in escalating doses as soon as possible after the onset of seizures.

Methods of treating epileptic disorders by administering to a subject in need thereof an effective amount of gaboxadol or a pharmaceutically acceptable salt thereof, either alone or in combination with, an allosteric modulator or a pharmaceutically acceptable salt, derivative or analogue, or combination thereof, are provided. Methods of treating epileptic disorders by
5 administering to a subject in need thereof an effective amount of an allosteric modulator or a pharmaceutically acceptable salt, derivative or analogue, or combination thereof, either alone or in combination with, gaboxadol or a pharmaceutically acceptable salt thereof, are provided.

An effective amount or therapeutically effective amount can be a dosage sufficient to treat, inhibit, or alleviate one or more symptoms of an epileptic disorder such as reducing the
10 frequency or severity of seizures, reducing behavior abnormalities (or otherwise improving behavior); or to provide a desired pharmacologic and/or physiologic effect, for example, reducing, inhibiting, or reversing one or more of the underlying pathophysiological mechanisms underlying the neurological dysfunction, increasing dopamine levels or signaling, or a combination thereof. The precise dosage will vary according to a variety of factors such as
15 subject-dependent variables (*e.g.*, age, immune system health, clinical symptoms *etc.*). In embodiments, a subject may be started at a low dose and the dosage is escalated. In this manner, it can be determined if the drug is well tolerated in the subject. Dosages can be lower for children than for adults.

In embodiments, the methods described herein are effective to reduce, delay, or prevent
20 one or more other clinical symptoms of an epileptic disorder, such as acute repetitive seizures. For example, the effect of a gaboxadol or a pharmaceutically acceptable salt thereof, and/or an allosteric modulator or a pharmaceutically acceptable salt, derivative or analogue thereof, on a particular symptom, pharmacologic, or physiologic indicator can be compared to an untreated subject, or the condition of the subject prior to treatment. In embodiments, the symptom,
25 pharmacologic, and/or physiologic indicator is measured in a subject prior to treatment, and again one or more times after treatment is initiated. In embodiments, the control is a reference level, or average determined based on measuring the symptom, pharmacologic, or physiologic indicator in one or more subjects that do not have the disease or condition to be treated (*e.g.*, healthy subjects). In embodiments, the effect of the treatment is compared to a conventional
30 treatment that is known the art.

In embodiments, compositions and methods of treatment are provided with low dosages of gaboxadol and/or an allosteric modulator such that the patient is provided one or more beneficial effects related to an epileptic disorder, such as, reduced seizure activity, reduced fatigue, increased mood, increased concentration, increased behavioral control and/or increased cognitive ability. Provided herein are dosing regimens that allow effective treatment of an epileptic disorder with potentially limited or substantially few negative side effects, e.g., convulsions and/or sleep disruption. Accordingly, the methods described herein may provide treatment of an epileptic disorder that may be considered surprising and unexpected. For example, methods are provided herein of treating epileptic disorders in a patient in need thereof which may not cause sleep disruption. In embodiments, methods described herein may provide effective treatment of an epileptic disorder without interrupting Slow Wave Sleep. In embodiments, methods of treating an epileptic disorder without causing insomnia or trouble falling asleep are provided.

In embodiments, the methods described herein may be used to treat epileptic disorders including acute repetitive seizures, Landau-Kleffner Syndrome, Lennox-Gastaut syndrome (LGS) and Dravet syndrome. In embodiments, the methods include treatment of acute repetitive seizure.

In embodiments, the methods described herein may be used to treat epileptic disorders including benign rolandic epilepsy (BRE), intractable childhood epilepsy (ICE), childhood absence epilepsy (CAE), juvenile myoclonic epilepsy (JME), infantile spasms (or West syndrome), generalized epilepsy with febrile seizure plus (GEFS+) and Lennox-Gastaut syndrome (LGS).

In embodiments, the methods described herein may be used to treat a sodium channel protein type 1 subunit alpha (Scn1A)-related disorder. For example Scn1A-related disorders include generalized epilepsy with febrile seizures plus, intractable childhood epilepsy with generalized tonic-clonic seizures, intractable infantile partial seizures, myoclonic-astatic epilepsy, severe myoclonic epilepsy in infancy, simple febrile seizures, Dravet syndrome, Lennox-Gastaut syndrome (LGS), infantile spasms, and vaccine-related encephalopathy and seizures.

The methods described herein may also be effective in subjects experiencing intractable seizures, status epilepticus, akinetic seizures, myoclonic seizures, absence seizures, or severe

myoclonic epilepsy in infancy (SMEI). In embodiments, the disorders are characterized by intractable seizures. Intractable seizures (also referred to as “uncontrolled” or “refractory” seizures) are seizures that cannot be controlled with conventional treatments. For example, the subject can have intractable epilepsy or another disorder characterized by intractable seizures, or a disorder characterized by status epilepticus. Status epilepticus is a condition in which seizures follow one another without recovery of consciousness between them. Accordingly, in embodiments, the disclosed methods are used to treat subjects that would otherwise be resistant to one or more conventional therapies.

The methods described herein may be particularly useful for treating children and infants, and for treating disorders that onset during infancy or childhood. In embodiments, the subject of the disclosed method is a newborn, a baby, a toddler, a preschooler, a school-age child, a tween, or a teenager. In embodiments, the subject is 18 years old or younger, 12 years old or younger, 10 years old or younger, 8 years old or younger, 6 years old or younger, 4 years old or younger, 2 years old or younger, 1 year old or younger. In embodiments, the subject is an adult that is over eighteen years old.

In embodiments, the epileptic disorders are characterized by seizures associated with epilepsy. In embodiments, the seizures are non-epileptic seizures (NES) or dissociative seizures that are distinguished from epilepsy. Non-epileptic seizures include organic non-epileptic seizures and psychogenic seizures.

Epilepsy is a neurological disorder that occurs when nerve cell activity in the brain becomes disrupted, leading to seizures or periods of unusual behavior, sensations and sometimes loss of consciousness. A subject can be said to have epilepsy when having two seizures without an obvious cause. Epilepsy can occur in both adults and children, and can be associated with a specific syndrome. Accordingly, in embodiments, the subject has a childhood epilepsy syndrome such as benign rolandic epilepsy (BRE), childhood absence epilepsy (CAE), juvenile myoclonic epilepsy (JME), infantile spasms (or West syndrome), Dravet syndrome or Lennox-Gastaut syndrome (LGS).

In embodiments, the subject does not experience diagnosable seizures, but exhibits subclinical electrical discharges, which refers to a high rate of seizure-like activity when their brain waves are measured with an electroencephalogram. Epileptic syndromes associated with

these seizure-like discharges include Landau-Kleffner Syndrome, Dravet syndrome and Continuous Spike-wave Activity during Slow-wave Sleep.

In embodiments, the epileptic disorders treated by the methods and compositions described herein include Scn1A-related seizure disorders. Scn1A-related seizure disorders include simple febrile seizures (FS) and generalized epilepsy with febrile seizures plus (GEFS+) at the mild end to Dravet syndrome and intractable childhood epilepsy with generalized tonic-clonic seizures (ICE-GTC) at the severe end. Specific Scn1A-related seizure disorders include, but are not limited to, generalized epilepsy with febrile seizures, intractable childhood epilepsy with generalized tonic-clonic seizures, intractable infantile partial seizures, myoclonic-astatic epilepsy, severe myoclonic epilepsy in infancy, simple febrile seizures, Dravet syndrome, Lennox-Gastaut syndrome (LGS), infantile spasms, and vaccine-related encephalopathy.

In embodiments, the subject has an intellectual epileptic disability (IDD) such as an Autism Spectrum Disorders (ASD). In embodiments, the subject of the disclosed method has epilepsy and an IDD or ASD disorder. Common IDD and ASD that are comorbid with seizures and epilepsy include, but are not limited to, fragile X syndrome (FXS), Rett syndrome (RTT), Angelman syndrome, Prader-Willi syndrome, Velocardiofacial syndrome, Smith-Lemli-Opitz syndrome, neuroligin mutations and "interneuronopathies" resulting from aristaless-related homeobox, X-linked (ARX) and Nueropilin 2 (NRP2) gene mutations.

Also provided herein are methods and compositions for treating epileptic disorders by co-administering to a patient in need thereof an allosteric modulator, and gaboxadol, a derivative thereof, or a pharmaceutically acceptable salt thereof. In embodiments, the methods and compositions described herein include a dosage form including gaboxadol or a pharmaceutically acceptable salt thereof, and an allosteric modulator. In embodiments, the methods and compositions described herein can include a dosage form including gaboxadol or a pharmaceutically acceptable salt thereof, and a separate dosage form including an allosteric modulator or a pharmaceutically acceptable salt thereof.

Gaboxadol or pharmaceutically acceptable salt thereof may be provided as an acid addition salt, a zwitter ion hydrate, zwitter ion anhydrate, hydrochloride or hydrobromide salt, or in the form of the zwitter ion monohydrate. Acid addition salts, include but are not limited to, maleic, fumaric, benzoic, ascorbic, succinic, oxalic, bis-methylenesalicylic, methanesulfonic, ethane-disulfonic, acetic, propionic, tartaric, salicylic, citric, gluconic, lactic, malic, mandelic,

cinnaamic, citraconic, aspartic, stearic, palmitic, itaconic, glycolic, p-amino-benzoic, glutamic, benzene sulfonic or theophylline acetic acid addition salts, as well as the 8-halotheophyllines, for example 8-bromo-theophylline. In other suitable embodiments, inorganic acid addition salts, including but not limited to, hydrochloric, hydrobromic, sulfuric, sulfamic, phosphoric or nitric acid addition salts may be used.

In embodiments, gaboxadol is provided as gaboxadol monohydrate. One skilled in the art will readily understand that the amounts of active ingredient in a pharmaceutical composition will depend on the form of gaboxadol provided. For example, pharmaceutical compositions of including 5.0, 10.0, or 15.0 mg gaboxadol correspond to 5.6, 11.3, or 16.9 mg gaboxadol monohydrate.

In embodiments, gaboxadol is crystalline, such as the crystalline hydrochloric acid salt, the crystalline hydrobromic acid salt, or the crystalline zwitter ion monohydrate. In embodiments, gaboxadol is provided as a crystalline monohydrate.

Deuteration of pharmaceuticals to improve pharmacokinetics (PK), pharmacodynamics (PD), and toxicity profiles, has been demonstrated previously with some classes of drugs. Accordingly the use of deuterium enriched gaboxadol is contemplated and within the scope of the methods and compositions described herein. Deuterium can be incorporated in any position in replace of hydrogen synthetically, according to the synthetic procedures known in the art. For example, deuterium may be incorporated to various positions having an exchangeable proton, such as the amine N--H, via proton-deuterium equilibrium exchange. Thus, deuterium may be incorporated selectively or non-selectively through methods known in the art to provide deuterium enriched gaboxadol. *See Journal of Labeled Compounds and Radiopharmaceuticals* 19(5) 689-702 (1982).

Pharmaceutical compositions herein may be provided with immediate release, delayed release, extended release, or modified release profiles. In embodiments, pharmaceutical compositions with different drug release profiles may be combined to create a two phase or three-phase release profile. For example, pharmaceutical compositions may be provided with an immediate release and an extended release profile. In embodiments, pharmaceutical compositions may be provided with an extended release and delayed release profile. Such composition may be provided as pulsatile formulations, multilayer tablets, or capsules containing tablets, beads, granules, etc. Compositions may be prepared using a pharmaceutically acceptable

“carrier” composed of materials that are considered safe and effective. The “carrier” includes all components present in the pharmaceutical formulation other than the active ingredient or ingredients. The term “carrier” includes, but is not limited to, diluents, binders, lubricants, disintegrants, fillers, and coating compositions.

5 In embodiments, the pharmaceutical compositions described herein are administered once, twice, three times or four times daily, or every other day. In embodiments, a pharmaceutical composition described herein is provided to the patient in the evening. In embodiments, a pharmaceutical composition described herein is provided to the patient once in the evening and once in the morning. In embodiments, a pharmaceutical composition herein is provided as soon as possible after the occurrence of a seizure. In embodiments, a pharmaceutical composition herein is provided continuously.

10 In embodiments, the total amount of allosteric modulator and/or gaboxadol administered to a subject in a 24-hour period is 1 mg to 50 mg. In embodiments, the total amount of allosteric modulator and/or gaboxadol administered to a subject in a 24-hour period is 1 mg to 20 mg. In 15 embodiments, the total amount of allosteric modulator and/or gaboxadol administered to a subject in a 24-hour period is 5 mg, 10 mg, or 15 mg. In embodiments, the total amount of allosteric modulator and/or gaboxadol or a pharmaceutically acceptable salt thereof administered to a subject in a 24-hour period is 1 mg to 50 mg. In embodiments, the subject may be started at a low dose and the dosage is escalated. In this manner, it can be determined if the drug is well 20 tolerated in the subject. Dosages can be lower for children than for adults.

 In embodiments, provided herein are methods of treating an epileptic disorder including administering to a patient in need thereof a pharmaceutical composition including an allosteric modulator and/or gaboxadol wherein the composition provides improvement in at least one symptom of the epileptic disorder. In embodiments, methods of treating epileptic disorders by 25 administering to a subject in need thereof an effective amount of allosteric modulator and/or gaboxadol, or combination thereof, are provided. An effective amount or therapeutically effective amount can be a dosage sufficient to treat, inhibit, or alleviate one or more symptoms of an epileptic disorder such as reducing the frequency or severity of seizures, reducing behavior abnormalities (or otherwise improving behavior); or to provide a desired pharmacologic and/or 30 physiologic effect, for example, reducing, inhibiting, or reversing one or more of the underlying pathophysiological mechanisms underlying the neurological dysfunction, increasing dopamine

levels or signaling, or a combination thereof. The precise dosage will vary according to a variety of factors such as subject-dependent variables (*e.g.*, age, immune system health, clinical symptoms etc.).

5 In embodiments, the methods described herein are effective to reduce, delay, or prevent one or more other clinical symptoms of an epileptic disorder, *e.g.*, epilepsy or Dravet syndrome. For example, the effect of a composition including allosteric modulator and/or gaboxadol on a particular symptom, pharmacologic, or physiologic indicator can be compared to an untreated subject, or the condition of the subject prior to treatment. In embodiments, the symptom, pharmacologic, and/or physiologic indicator is measured in a subject prior to treatment, and
10 again one or more times after treatment is initiated. In embodiments, the control is a reference level, or average determined based on measuring the symptom, pharmacologic, or physiologic indicator in one or more subjects that do not have the disease or condition to be treated (*e.g.*, healthy subjects). In embodiments, the effect of the treatment is compared to a conventional treatment that is known the art.

15 In embodiments, provided herein are methods of treating an epileptic disorder, *e.g.*, status epilepticus, including administering to a patient in need thereof a pharmaceutical composition including an allosteric modulator and/or gaboxadol or pharmaceutically acceptable salt thereof wherein the composition provides improvement in at least one symptom of the epileptic disorder. In embodiments, the methods provided may also surprisingly and unexpectedly reduce or
20 prevent seizures, or symptoms thereof in a subject in need thereof. In embodiments, the methods provided may reduce or prevent one or more different types of seizures. In embodiments, the methods provided may also surprisingly and unexpectedly reduce or prevent seizures, or symptoms thereof in a subject in need thereof. In embodiments, the methods provided may reduce or prevent one or more different types of seizures. Generally, a seizure can include
25 convulsions, repetitive movements, unusual sensations, and combinations thereof. Seizures can be categorized as focal seizures (also referred to as partial seizures) and generalized seizures. Focal seizures affect only one side of the brain, while generalized seizures affect both sides of the brain. Specific types of focal seizures include simple focal seizures, complex focal seizures, and secondarily generalized seizures. Simple focal seizures can be restricted or focused on a
30 particular lobe (*e.g.*, temporal lobe, frontal lobe, parietal lobe, or occipital lobe). Complex focal seizures generally affect a larger part of one hemisphere than simple focal seizures, but

commonly originate in the temporal lobe or the frontal lobe. When a focal seizure spreads from one side (hemisphere) to both sides of the brain, the seizure is referred to as a secondarily generalized seizure. Specific types of generalized seizures include absences (also referred to as petit mal seizures), tonic seizures, atonic seizures, myoclonic seizures, tonic clonic seizures (also referred to as grand mal seizures), and clonic seizures.

In embodiments, methods described herein may reduce the frequency of seizures, reduce the severity of seizures, change the type of seizures (*e.g.*, from a more severe type to a less severe type), or a combination thereof in a subject after treatment compared to the absence of treatment (*e.g.*, before treatment), or compared to treatment with an alternative conventional treatment.

In embodiments, provided herein are methods of treating an epileptic disorder wherein the patient is provided improvement of at least one symptom for more than 4 hours after administration of the pharmaceutical composition to the patient. In embodiments, the improvement of at least one symptom for more than 6 hours after administration of the pharmaceutical composition to the patient is provided in accordance with the present disclosure. In embodiments, improvement of at least one symptom for more than, *e.g.*, 8 hours, 10 hours, 12 hours, 15 hours, 18 hours, 20 hours, or 24 hours after administration of the pharmaceutical composition to the patient is provided in accordance with the present disclosure. In embodiments, improvement in at least one symptom for at least *e.g.*, 8 hours, 10 hours, 12 hours, 15 hours, 18 hours, 20 hours, or 24 hours after administration of the pharmaceutical composition to the patient is provided in accordance with the present disclosure. In embodiments, improvement in at least one symptom for 12 hours after administration of the pharmaceutical composition to the patient is provided in accordance with the present disclosure.

In embodiments, provided herein methods of treating an epileptic disorder including administering a composition herein to a patient in need thereof wherein the composition provides improvement in next day functioning to the patient.

In embodiments, provided herein are methods of treating an epileptic disorder wherein the amount of active substance, *e.g.*, allosteric modulator and/or gaboxadol, within the patient about 4 hours after administration of the pharmaceutical composition is less than about 75% of the administered dose. In embodiments, provided herein are methods wherein the amount of allosteric modulator and/or gaboxadol or pharmaceutically acceptable salt thereof within the

patient about, *e.g.*, 6 hours, 8 hours, 10 hours, 12 hours, 15 hours, or 20 hours after administration of the pharmaceutical composition is less than about 75%.

In embodiments, provided herein are methods of treating an epileptic disorder wherein the amount of active substance, *e.g.*, allosteric modulator and/or gaboxadol, within the patient about 4 hours after administration of the pharmaceutical composition is less than about 80% of the administered dose. In embodiments, provided herein are methods wherein the amount of active substance, *e.g.*, allosteric modulator and/or gaboxadol, within the patient about, *e.g.*, 6 hours, 8 hours, 10 hours, 12 hours, 15 hours, or 20 hours after administration of the pharmaceutical composition is less than about 80% of the administered dose.

In embodiments, provided herein are methods of treating an epileptic disorder wherein the amount of active substance, *e.g.*, allosteric modulator and/or gaboxadol, within the patient about 4 hours after administration of the pharmaceutical composition is between about 65% to about 85% of the administered dose. In embodiments, the amount of active substance, *e.g.*, allosteric modulator and/or gaboxadol, within the patient after about, *e.g.*, 6 hours, 8 hours, 10 hours, 12 hours, 15 hours, or 20 hours after administration of the pharmaceutical composition is between about 65% to about 85% of the administered dose.

In embodiments, provided herein are methods of treating an epileptic disorder including administering to a patient in need thereof a pharmaceutical composition including an active substance, *e.g.*, allosteric modulator and/or gaboxadol, wherein the composition provides an *in vivo* plasma profile having a C_{max} less than about 500 ng/ml. In embodiments, the composition provides improvement for more than 6 hours after administration to the patient.

In embodiments, the composition provides an *in vivo* plasma profile having a C_{max} less than about, *e.g.*, 450 ng/ml, 400 ng/ml, 350 ng/ml, or 300 ng/ml and wherein the composition provides improvement of next day functioning of the patient. In embodiments, the composition provides an *in vivo* plasma profile having a C_{max} less than about, *e.g.*, 250 ng/ml, 200 ng/ml, 150 ng/ml, or 100 ng/ml and wherein the composition provides improvement of next day functioning of the patient.

In embodiments, provided herein are methods of treating an epileptic disorder including administering to a patient in need thereof a pharmaceutical composition wherein the composition provides a consistent *in vivo* plasma profile having a $AUC_{0-\infty}$ of less than about 900 ng•hr/ml. In embodiments, the composition provides improvement in next day functioning of the patient. In

embodiments, the compositions provide an *in vivo* plasma profile having a $AUC_{0-\infty}$ of less than about, *e.g.*, 850 ng•hr/ml, 800 ng•hr/ml, 750 ng•hr/ml, or 700 ng•hr/ml and wherein the composition provides improvement of next day functioning of the patient. In embodiments, the composition provides improvement in one or more symptom for more than 6 hours after administration.

In embodiments, provided herein are methods of treating an epileptic disorder including administering to a patient in need thereof a pharmaceutical composition including an active substance, *e.g.*, allosteric modulator and/or gaboxadol, wherein the composition provides an *in vivo* plasma profile having a $AUC_{0-\infty}$ of less than about, *e.g.*, 650 ng•hr/ml, 600 ng•hr/ml, 550 ng•hr/ml, 500 ng•hr/ml, or 450 ng•hr/ml. In embodiments, wherein the composition provides an *in vivo* plasma profile having a $AUC_{0-\infty}$ of less than about, *e.g.*, 400 ng•hr/ml, 350 ng•hr/ml, 300 ng•hr/ml, 250 ng•hr/ml, or 200 ng•hr/ml. In embodiments, the composition provides an *in vivo* plasma profile having a $AUC_{0-\infty}$ of less than about, *e.g.*, 150 ng•hr/ml, 100 ng•hr/ml, 75 ng•hr/ml, or 50 ng•hr/ml. In embodiments, the composition provides improvement of next day functioning of the patient after administration for more than, *e.g.*, 4 hours, 6 hours, 8 hours, 10 hours, or 12 hours, after administration of the composition to the patient.

In embodiments, provided herein are methods of treating a epileptic disorder including administering to a patient in need thereof a first pharmaceutical composition including gaboxadol or a pharmaceutically acceptable salt thereof and a second pharmaceutical composition including gaboxadol or a pharmaceutically acceptable salt thereof. In some embodiments, the second pharmaceutical composition provides an *in vivo* plasma profile having a mean $AUC_{0-\infty}$ of at least about 20% less than the first pharmaceutical composition.

In embodiments, provided herein are methods of treating an epileptic disorder including administering to a patient in need thereof a first pharmaceutical composition including an allosteric modulator and a second pharmaceutical composition including gaboxadol or a pharmaceutically acceptable salt thereof. In embodiments, the second pharmaceutical composition provides an *in vivo* plasma profile having a mean $AUC_{0-\infty}$ of at least about 20% less than the first pharmaceutical composition.

In embodiments, the first and/or the second pharmaceutical compositions are administered once, twice, three or four times daily, or every other day. In embodiments, the first or the second pharmaceutical composition is provided to the patient in the evening. In

embodiments, the second pharmaceutical composition includes an amount of gaboxadol that is at least one third of the amount of the allosteric modulator in the first pharmaceutical composition. In embodiments, the second pharmaceutical composition includes an amount of gaboxadol that is at least half of the amount of the amount of the allosteric modulator provided in the first pharmaceutical composition.

5 In embodiments, the first and/or the second pharmaceutical composition are provided to the patient once in the evening and once in the morning. In embodiments, the total amount of allosteric modulator administered to a subject in a 24-hour period is 1 mg to 2500 mg. In embodiments, the total amount of allosteric modulator administered to a subject in a 24-hour
10 period is 1 mg/kg to 35 mg/kg. In embodiments, the total amount of gaboxadol or a pharmaceutically acceptable salt thereof administered to a subject in a 24-hour period is 1 mg to 75 mg. In embodiments, the total amount of active substance, *e.g.*, gaboxadol or pharmaceutically acceptable salt thereof, administered to a subject in a 24-hour period is less than about 75 mg, 50 mg, 25 mg, 20 mg, 10 mg, or 5 mg. In embodiments, the total amount of
15 active substance, *e.g.*, gaboxadol or pharmaceutically acceptable salt thereof, administered to a subject in a 24-hour period is less than 15 mg. In embodiments, the total amount of active substance, *e.g.*, allosteric modulator and/or gaboxadol, administered to a subject in a 24-hour period is less than about 2500mg, 2250 mg, 2000 mg, 1750 mg, 1500 mg, 1250 mg, 1000 mg, 750 mg, 500 mg, 250 mg, 200 mg, 175 mg, 150 mg, 125 mg, 100 mg, 75 mg, 50 mg, 25 mg, 20
20 mg, 10 mg, or 5 mg. In embodiments, the total amount of active substance, *e.g.*, allosteric modulator and/or gaboxadol, administered to a subject in a 24-hour period is less than 15 mg.

In embodiments, the first and/or the second pharmaceutical compositions may be provided with immediate release, delayed release, extended release, or modified release profiles. The first and second pharmaceutical compositions may be provided at the same time or separated
25 by an interval of time, *e.g.*, 6 hours, 12 hours etc. In embodiments, the first and the second pharmaceutical compositions may be provided with different drug release profiles to create a two-phase release profile. For example, the first pharmaceutical composition may be provided with an immediate release profile and the second pharmaceutical composition may provide an extended release profile. In embodiments, one or both of the first and second pharmaceutical
30 compositions may be provided with an extended release or delayed release profile. Such compositions may be provided as pulsatile formulations, multilayer tablets or capsules

containing tablets, beads, granules, etc. In some embodiments, the first pharmaceutical composition is an immediate release composition. In embodiments, the second pharmaceutical composition is an immediate release composition. In embodiments, the first and second pharmaceutical compositions are provided as separate immediate release compositions, *e.g.*,
 5 tablets or capsules. In embodiments the first and second pharmaceutical compositions are provided 12 hours apart.

In embodiments, compositions described herein are suitable for parenteral administration, including, *e.g.*, intramuscularly (i.m.), intravenously (i.v.), subcutaneously (s.c.), intraperitoneally (i.p.), or intrathecally (i.t.). Parenteral compositions must be sterile for
 10 administration by injection, infusion or implantation into the body and may be packaged in either single-dose or multi-dose containers.

In embodiments, liquid pharmaceutical compositions for parenteral administration to a subject include an active substance, *e.g.*, allosteric modulator and/or gaboxadol, at a concentration of about 0.005 µg/ml to about 500 µg/ml. In embodiments, the composition
 15 comprises an active substance, *e.g.*, allosteric modulator and/or gaboxadol, at a concentration of, *e.g.*, about 0.005 µg/ml to about 250 µg/ml, about 0.005 µg/ml to about 200 µg/ml, about 0.005 µg/ml to about 150 µg/ml, about 0.005 µg/ml to about 100 µg/ml, or about 0.005 µg/ml to about 50 µg/ml.

In embodiments, the compositions comprises an active substance, *e.g.*, allosteric
 20 modulator and/or gaboxadol, at a concentration of, *e.g.*, about 0.05 µg/ml to about 50 µg/ml, about 0.1 µg/ml to about 50 µg/ml, about 0.05 µg/ml to about 25 µg/ml, about 0.05 µg/ml to about 10 µg/ml, about 0.05 µg/ml to about 5 µg/ml, or about 0.05 µg/ml to about 1 µg/ml. In embodiments, the composition comprises an active substance, *e.g.*, allosteric modulator and/or
 25 gaboxadol, at a concentration of, *e.g.*, about 0.05 µg/ml to about 15 µg/ml, about 0.5 µg/ml to about 10 µg/ml, about 0.5 µg/ml to about 7 µg/ml, about 1 µg/ml to about 10 µg/ml, about 5 µg/ml to about 10 µg/ml, or about 5 µg/ml to about 15 µg/ml. In embodiments, the pharmaceutical compositions for parenteral administration is formulated as a total volume of about, *e.g.*, 10 ml, 20 ml, 25 ml, 50 ml, 100 ml, 200 ml, 250 ml, or 500 ml. In embodiments, the compositions are contained in a bag, a glass vial, a plastic vial, or a bottle.

30 In embodiments, compositions for parenteral administration include about 0.05 mg to about 100 mg active substance, *e.g.*, allosteric modulator and/or gaboxadol. In embodiments, the

pharmaceutical compositions comprise about, *e.g.*, 0.1 mg to 25 mg, 0.1 mg to 20 mg, 0.1 mg to 15 mg, 0.5 mg to 25 mg, 0.5 mg to 20 mg, 0.5 mg to 15 mg, 1 mg to 25 mg, 1 mg to 20 mg, 1 mg to 15 mg, 1.5 mg to 25 mg, 1.5 mg to 20 mg, 1.5 mg to 15 mg, 2 mg to 25 mg, 2 mg to 20 mg, 2 mg to 15 mg, 2.5 mg to 25 mg, 2.5 mg to 20 mg, 2.5 mg to 15 mg, 3 mg to 25 mg, 3 mg to 20 mg, 3 mg to 15 mg active substance, *e.g.*, allosteric modulator and/or gaboxadol.

In embodiments, pharmaceutical compositions include about, *e.g.*, 5 mg to 20 mg, 5 mg to 10 mg, 4 mg to 6 mg, 6 mg to 8 mg, 8 mg to 10 mg, 10 mg to 12 mg, 12 mg to 14 mg, 14 mg to 16 mg, 16 mg to 18 mg, or 18 mg to 20 mg active substance, *e.g.*, allosteric modulator and/or gaboxadol. In embodiments, the pharmaceutical compositions include about, *e.g.*, 0.1 mg, 0.25 mg, 0.5 mg, 1 mg, 2.5 mg, 3 mg, 4 mg, 5 mg, 7 mg, 7.5 mg, 10 mg, 12.5 mg, 15 mg, 17.5 mg, 20 mg active substance, *e.g.*, allosteric modulator and/or gaboxadol, or amounts that are multiples of such doses. The compositions may be contained in a bag, a glass vial, a plastic vial, or a bottle.

In embodiments, pharmaceutical compositions for parenteral administration to a subject include an active substance, *e.g.*, allosteric modulator and/or gaboxadol, at a concentration of about 0.005 mg/ml to about 500 mg/ml. In embodiments, the compositions include an active substance, *e.g.*, allosteric modulator and/or gaboxadol, at a concentration of, *e.g.*, about 0.05 mg/ml to about 50 mg/ml, about 0.1 mg/ml to about 50 mg/ml, about 0.1 mg/ml to about 10 mg/ml, about 0.05 mg/ml to about 25 mg/ml, about 0.05 mg/ml to about 10 mg/ml, about 0.05 mg/ml to about 5 mg/ml, or about 0.05 mg/ml to about 1 mg/ml. In embodiments, the composition includes an active substance, *e.g.*, allosteric modulator and/or gaboxadol, at a concentration of, *e.g.*, about 0.05 mg/ml to about 15 mg/ml, about 0.5 mg/ml to about 10 mg/ml, about 0.25 mg/ml to about 5 mg/ml, about 0.5 mg/ml to about 7 mg/ml, about 1 mg/ml to about 10 mg/ml, about 5 mg/ml to about 10 mg/ml, or about 5 mg/ml to about 15 mg/ml. In embodiments, pharmaceutical compositions for parenteral administration are formulated as a total volume of about, *e.g.*, 10 ml, 20 ml, 25 ml, 50 ml, 100 ml, 200 ml, 250 ml, or 500 ml. In embodiments, the compositions are packages and stored in a bag, a glass vial, a plastic vial, or a bottle.

In embodiments, pharmaceutical compositions herein include an active substance, *e.g.*, allosteric modulator and/or gaboxadol, wherein the active substance is present at a molarity less than about 1.0 M. In embodiments, the active substance, *e.g.*, allosteric modulator and/or gaboxadol, is present at a molarity greater than, *e.g.*, about 0.0001 M about 0.001 M, about 0.01

M, about 0.1 M, about 0.2 M, greater than about 0.5, greater than about 1.0 M, greater than about 1.2 M, greater than about 1.5 M, greater than about 1.75 M, greater than about 2.0 M, or greater than about 2.5 M. In embodiments, the active substance, *e.g.*, allosteric modulator and/or gaboxadol, is present at a molarity of between, *e.g.*, about 0.00001 M to about 0.1 M, about 0.01 to about 0.1 M, about 0.1 M to about 1.0 M, about 1.0 M to about 5.0 M, or about 5.0 M to about 10.0 M. In embodiments, the active substance, *e.g.*, allosteric modulator and/or gaboxadol, is present at a molarity of less than, *e.g.*, about 0.01 M, about 0.1 M, about 1.0 M, about 5.0 M, or about 10.0 M

In embodiments, the solubility of the active substance, *e.g.*, allosteric modulator and/or gaboxadol, in the composition is greater than, *e.g.*, about 10 mg/mL, about 15 mg/mL, about 20 mg/mL, about 25 mg/mL, about 30 mg/mL, about 40 mg/mL, about 50 mg/mL, about 75 mg/mL, about 100 mg/mL, about 150 mg/mL, when measured, for example, in water at 25°C.

In embodiments, the solubility of the active substance, *e.g.*, allosteric modulator and/or gaboxadol, in the composition is between, *e.g.*, about 1 mg/mL to about 50 mg/mL, about 5 mg/mL to about 50 mg/mL, about 10 mg/mL to about 50 mg/mL, about 20 mg/mL to about 50 mg/mL, from about 20 mg/mL to about 30 mg/mL or from about 10 mg/mL to about 45 mg/mL, when measured, for example, in water at 25 C.

In embodiments, a pharmaceutical composition for parenteral administration is provided wherein the pharmaceutical composition is stable for at least six months. In embodiments, the pharmaceutical compositions herein exhibit no more than about 5% decrease in active substance, *e.g.*, allosteric modulator and/or gaboxadol, *e.g.*, 3 months or 6 months. In embodiments, the amount of gaboxadol or pharmaceutically acceptable salt thereof degrades at no more than about, *e.g.*, 2.5%, 1%, 0.5% or 0.1%. In embodiments, the degradation is less than about, *e.g.*, 5%, 2.5%, 1%, 0.5%, 0.25%, 0.1%, for at least six months.

In embodiments, pharmaceutical compositions for parenteral administration are provided wherein the pharmaceutical composition remains soluble. In embodiments, pharmaceutical compositions are provided that are stable, soluble, local site compatible and/or ready-to-use. In embodiments, the pharmaceutical compositions herein are ready-to-use for direct administration to a patient in need thereof.

The parenteral compositions provided herein may comprise one or more excipients, *e.g.*, solvents, solubility enhancers, suspending agents, buffering agents, isotonicity agents, stabilizers

or antimicrobial preservatives. When used, the excipients of the parenteral compositions will not adversely affect the stability, bioavailability, safety, and/or efficacy of allosteric modulator and/or gaboxadol or pharmaceutically acceptable salt(s) used in the composition. Thus, parenteral compositions are provided wherein there is no incompatibility between any of the components of the dosage form.

In embodiments, parenteral compositions of allosteric modulator and/or gaboxadol or a pharmaceutically acceptable salt thereof include a stabilizing amount of at least one excipient. For example, excipients may be selected from the group consisting of buffering agents, solubilizing agents, tonicity agents, antioxidants, chelating agents, antimicrobial agents, and preservative. One skilled in the art will appreciate that an excipient may have more than one function and be classified in one or more defined group.

In embodiments, pharmaceutical compositions include an allosteric modulator and/or gaboxadol, or a pharmaceutically acceptable salt thereof and an excipient wherein the excipient is present at a weight percent (w/v) of less than about, *e.g.*, 10%, 5%, 2.5%, 1%, or 0.5%. In embodiments, the excipient is present at a weight percent between about, *e.g.*, 1.0% to 10%, 10% to 25%, 15% to 35%, 0.5% to 5%, 0.001% to 1%, 0.01% to 1%, 0.1% to 1%, or 0.5% to 1%. In embodiments, the excipient is present at a weight percent between about, *e.g.*, 0.001% to 1%, 0.01% to 1%, 1.0% to 5%, 10% to 15%, or 1% to 15%.

In embodiments pharmaceutical compositions are provided including gaboxadol, or a pharmaceutically acceptable salt thereof, and an excipient wherein the excipient is present in a molar ratio of the excipient to gaboxadol or pharmaceutically acceptable salt of, *e.g.*, about 0.01:1 to about 0.45:1, about 0.1:1 to about 0.15:1, about 0.01:1 to about 0.1:1, and about 0.001:1 to about 0.01:1 are provided. In embodiments, the excipient is present at a molar ratio of the excipient to gaboxadol or pharmaceutically acceptable salt is about 0.0001:1 to about 0.1:1 or about 0.001:1 to about 0.001:1.

In embodiments, pharmaceutical compositions are provided including gaboxadol, or a pharmaceutically acceptable salt thereof and an excipient wherein the excipient comprises a stabilizing amount of a buffering agent. The buffering agent may be used to maintain the pH of the pharmaceutical composition wherein the gaboxadol or pharmaceutically acceptable salt thereof remains soluble, stable, and/or physiologically compatible. For example, in embodiments, the parenteral compositions include a buffering agent wherein the composition

remains stable without significant gaboxadol degradation. In embodiments, the addition of a buffer is desired for controlling the pH to enhance stability without significantly catalyzing or degrading the gaboxadol or salt thereof and/or causing pain to the patient upon infusion.

In embodiments, the buffering agent can be a citrate, phosphate, acetate, tartrate, carbonate, glutamate, lactate, succinate, bicarbonate buffer and combinations thereof. For example, sodium citrate, trisodium citrate anhydrous, trisodium citrate dihydrate, sodium citrate dehydrate, triethanolamine (TRIS), trisodium citrate pentahydrate dihydrate (*i.e.*, trisodium citrate dehydrate), acetic acid, citric acid, glutamic acid, phosphoric acid, may be used as a buffering agent. In embodiments, the buffering agent may be an amino acid, alkali metal, or alkaline earth metal buffer. For example, the buffering agent may be sodium acetate or hydrogen phosphate.

In embodiments, parenteral compositions of an active substance, *e.g.*, allosteric modulator and/or gaboxadol are provided, wherein the pH of the composition is between about 4.0 to about 8.0. In embodiments, the pH of the compositions is between, *e.g.*, about 5.0 to about 8.0, about 6.0 to about 8.0, about 6.5 to about 8.0. In embodiments, the pH of the compositions is between, *e.g.*, about 6.5 to about 7.5, about 7.0 to about 7.8, about 7.2 to about 7.8, or about 7.3 to about 7.6. In embodiments, the pH of the aqueous solution is, *e.g.*, about 6.8, about 7.0, about 7.2, about 7.4, about 7.6, about 7.7, about 7.8, about 8.0, about 8.2, about 8.4, or about 8.6.

In embodiments, pharmaceutical compositions of an active substance, *e.g.*, allosteric modulator and/or gaboxadol, or a pharmaceutically acceptable salt thereof, and an excipient wherein the excipient includes a solubilizing agent. For example, solubilizing agents according to the invention may include, *e.g.*, sodium hydroxide, L-lysine, L-arginine, sodium carbonate, potassium carbonate, sodium phosphate, and/or potassium phosphate. The amount of solubilizing agent in the composition will be sufficient such that the solution remains soluble at all concentrations, *i.e.*, does not turn hazy and/or form precipitates.

In embodiments, provided herein are pharmaceutical compositions of an active substance, *e.g.*, allosteric modulator and/or gaboxadol, or a pharmaceutically acceptable salt thereof, and an excipient wherein the excipient includes a particulate formation inhibitor. A particulate formation inhibitor refers to a compound that has the desired property of inhibiting the formation of particles in parenteral compositions. Particulate formation inhibitors of the invention include ethylenediaminetetraacetic acid (EDTA) and salts thereof, for example,

ethylenediaminetetraacetic acid, calcium disodium salt (preferably as the hydrate);
 ethylenediaminetetraacetic acid, diammonium salt (preferably as the hydrate);
 ethylenediaminetetraacetic acid, dipotassium salt (preferably as the dihydrate);
 ethylenediaminetetraacetic acid, disodium salt (preferably as the dihydrate and, if desired, as the
 5 anhydrous form); ethylenediaminetetraacetic acid, tetrasodium salt (preferably as the hydrate);
 ethylenediaminetetraacetic acid, tripotassium salt (preferably as the dihydrate);
 ethylenediaminetetraacetic acid, trisodium salt (preferably as the hydrate) and
 ethylenediaminetetraacetic acid disodium salt, USP (preferably as the dihydrate). In
 embodiments, pharmaceutical compositions described herein have an effective amount of a
 10 particulate formation inhibitor. In embodiments the excipients may include, e.g., an amino acid,
 urea, alcohol, ascorbic acid, phospholipids, proteins, such as serum albumin, collagen, and
 gelatin; salts such as EDTA or EGTA, and sodium chloride, liposomes, polyvinylpyrrolidone,
 sugars, such as dextran, mannitol, sorbitol, and glycerol, propylene glycol and polyethylene
 glycol (e.g., PEG-4000, PEG-6000), glycerol, glycine, and/or lipids.

15 In embodiments, provided herein are pharmaceutical compositions of an active substance,
 e.g., allosteric modulator and/or gaboxadol, or a pharmaceutically acceptable salt thereof, and an
 excipient wherein the excipient includes a solubilizing agent. For example, solubilizing agents
 may include, but are not limited to, acids, such as carboxylic acids, amino acids. In other
 examples, the solubilizing agents may be saturated carboxylic acids, unsaturated carboxylic
 20 acids, fatty acids, keto acids, aromatic carboxylic acids, dicarboxylic acids, tricarboxylic acids,
 α -hydroxy acids, amino acids, and combinations thereof.

In embodiments, provided herein are pharmaceutical compositions of an active substance,
 e.g., allosteric modulator and/or gaboxadol, or a pharmaceutically acceptable salt thereof, and an
 excipient wherein the excipient includes a solubilizing agent such as formic acid, acetic acid,
 25 propionic acid, butyric acid, valeric acid, caproic acid, enanthic acid, caprylic acid, pelargonic
 acid, capric acid, lauric acid, stearic acid, acrylic acid, docosahexaenoic acid, eicosapentaenoic
 acid, pyruvic acid, benzoic acid, salicylic acid, aldarcic acid, oxalic acid, malonic acid, malic acid,
 succinic acid, glutaric acid, adipic acid, citric acid, lactic acid, alanine, arginine, asparagine,
 aspartic acid, cysteine, glutamine, glycine, histidine, isoleucine, leucine, lysine, methionine,
 30 phenylalanine, proline, serine, threonine, tryptophan, tyrosine, valine, and combinations thereof.

In embodiments, the solubilizing agent is selected from acetic acid, salts thereof, and combinations thereof, (*e.g.*, acetic acid/sodium acetate), citric acid, salts thereof and combinations thereof (*e.g.*, citric acid/sodium citrate), DL arginine, L-arginine and histadine. In
 5 embodiments, the solubilizing agent is DL-arginine. In embodiments, the solubilizing agent is L-arginine. In embodiments, the solubilizing agent is acetic acid/sodium acetate. In embodiments, the solubilizing agent is citric acid/sodium citrate.

In embodiments, provided herein are pharmaceutical compositions of an active substance, *e.g.*, allosteric modulator and/or gaboxadol, or a pharmaceutically acceptable salt thereof, and an
 10 excipient wherein the excipient renders the composition isotonic. Isotonic pharmaceutical compositions herein may be achieved by adding an appropriate quantity of sodium chloride, glucose, laevulose, dextrose, mannitol, or potassium chloride, or calcium chloride, or calcium gluconoglucoheptonate, or mixtures thereof. For example, the excipients may include one or more tonicity agents, such as, *e.g.*, sodium chloride, potassium chloride, glycerin, mannitol, and/or dextrose. Tonicity agents may be used to minimize tissue damage and irritation, reduce
 15 hemolysis of blood cells, and/or prevent electrolyte imbalance. For example, the parenteral compositions may be an aqueous solution including sodium chloride wherein the composition is isotonic. In embodiments, the isotonicizing agent is sodium chloride. In embodiments, the concentration of the isotonicizing agent is between about 0.01 and about 2.0 weight percent. In embodiments, the pharmaceutical compositions may comprise up to about 10% isotonicizing
 20 agent. In embodiments the pharmaceutical compositions may comprise up to about, *e.g.*, 0.25%, 0.5%, 1%, 2.5% isotonicizing agent. In embodiments the amount of isotonicizing agent in the pharmaceutical is between about, *e.g.*, 0.01% to 1%, 0.1% to 1%, 0.25% to 1%, or 0.5% to 1%.

In embodiments, provided herein are pharmaceutical compositions of an active
 25 substance, *e.g.*, allosteric modulator and/or gaboxadol, or a pharmaceutically acceptable salt thereof, and an excipient wherein the excipient includes a free radical antagonist. In embodiments, the free radical antagonist is ascorbic acid, ascorbic acid derivatives, organic compounds having at least one thiol, alkyl polyhydroxylated, and cycloalkyl polyhydroxylated compounds, and combinations thereof.

In embodiments, provided herein are pharmaceutical compositions of an active
 30 substance, *e.g.*, allosteric modulator and/or gaboxadol, or a pharmaceutically acceptable salt thereof, and an excipient wherein the excipient includes a free radical scavenger selected from

thiolglycolic acid, thiolacetic acid, dithiothreitol, reduced glutathion, thiourea, α -thioglycerol, cystein, acetlcystein, mercaptoethane sulfonic acid and combinations thereof.

In embodiments, provided herein are pharmaceutical of an active substance, *e.g.*, allosteric modulator and/or gaboxadol, or a pharmaceutically acceptable salt thereof, and an excipient wherein the excipient includes riboflavin, dithiothreitol, sodium thiosulfate, thiourea, ascorbic acid, methylene blue, sodium metabisulfite, sodium bisulfite, propyl gallate acetylcysteine, phenol, acetone sodium bisulfate, ascorbic acid, ascorbic acid esters, butylhydroxyanisol (BHA), Butylhydroxytoluene (BHT), cysteine, nordihydroguaiaretic acid (NDGA), monothioglycerol, sodium bisulfite, sodium metabisulfate, tocopherols, and/or glutathione.

In embodiments, provided herein are pharmaceutical compositions of an active substance, *e.g.*, allosteric modulator and/or gaboxadol, or a pharmaceutically acceptable salt thereof, and an excipient wherein the excipient includes a preservative. In embodiments, the preservative is selected from benzalkonium chloride, benzethonium chloride, benzyl alcohol, chlorobutanol, chlorocresol, metacresol, Phenol, phenylmercuric nitrate, phenylmercuric acetate, methyl p-hydroxybenzoate, propyl p-hydroxybenzoate, butyl p-hydroxybenzoate, and thimerosal. In other embodiments, the preservative is selected from the group consisting of phenol, meta-cresol, benzyl alcohol, parabens (*e.g.*, methyl, propyl, butyl), benzalkonium chloride, chlorobutanol, thimerosal, phenylmercuric salts (*e.g.*, acetate, borate, or nitrate), and combinations thereof.

In embodiments, the compositions herein include a co-solvent. For example, in some instances the solubility of gaboxadol may be well below the therapeutic dose and therefore a co-solvent system may be used. A co-solvent is a mixture of solvents that may be used to achieve sufficiently high solubility and may increase the stability. For example, co-solvents may be a water-miscible organic solvents, such as ethanol, propylene, glycol, Capmul PG, propylene glycol, glycerin, polyethylene glycol, sorbitol, dimethylacetamide, and/or dimethylsulfoxide (DMSO). In embodiments, the cosolvent may comprise up to about 75% of the pharmaceutical composition. In other embodiments the amount of cosolvent used include up to about, *e.g.*, 1%, 5%, 10%, 15%, 25%, 40%, 50%, of the pharmaceutical composition.

The dosage forms may be prepared, for example, by mixing of an allosteric modulator and/or gaboxadol, or a pharmaceutically acceptable salt thereof, and one or more excipients (*e.g.*, buffering agents, solubilizing agents, tonicity agents, antioxidants, chelating agents,

antimicrobial agents and/or preservatives) in a blender under sterile conditions until a uniform blend is obtained. Pre-sterilized vials may then be filled with an appropriate amount of the sterile blend. The predetermined amount of sterile blend may then be mixed with a solvent, *e.g.*, water, saline, about 5-10% sugar (*e.g.*, glucose, dextrose) solution and combinations thereof prior to
5 administration. In addition, the solution may be frozen and thawed prior to further processing.

The excipients may be used in solid or in solution form. When used in solid form, the excipients and an allosteric modulator and/or gaboxadol, or a pharmaceutically acceptable salt thereof, may be mixed together as described above, and then solvent added prior to parenteral administration. When used in solution form, the allosteric modulator and/or gaboxadol, or a
10 pharmaceutically acceptable salt thereof, may be mixed with a solution of the excipient prior to parenteral administration.

Parenteral solutions including an allosteric modulator and/or gaboxadol, or a pharmaceutically acceptable salt thereof, may be prepared by mixing the required amount of allosteric modulator and/or gaboxadol, or a pharmaceutically acceptable salt thereof, which may
15 be purified prior to use in parenteral fluids such as DSW, distilled water, saline or PEG and adjusting the pH of this solution between 6.8-8. The process may be carried out at room temperature, or to increase concentration, the solution may be warmed appropriately. Other solvents such as PEG 400, 600, polypropylene glycol or other glycols can be used to enhance solubility. The resulting solutions after cooling to room temperature, may be sterilized by known
20 means such as ultrafiltration using, *e.g.*, 0.45 micron filter or ethylene oxide treatment or heating and may be packaged into ampules, vials or pre-filled syringes suitable for dispensing a sterile parenteral formulation.

When administered, the parenteral compositions herein provide a time of maximum plasma concentration (T_{max}) for gaboxadol in human patients of about 1 or more hours (*e.g.*,
25 about 1.5 or more hours). In embodiments, a T_{max} of gaboxadol in human patients ranging from between, *e.g.*, about 1 to about 5 hours, about 1 to about 4 hours, about 1 to about 3 hours, about 1 to about 2 hours. In embodiments, a T_{max} for gaboxadol in human patients of more than about 1.5 is observed. In embodiments, a T_{max} for gaboxadol in human patients of less than about 3 hours is observed. The time of maximum plasma concentration is measured once infusion is
30 complete.

In embodiments herein a dosage form includes from about 1 mg to about 500 mg gaboxadol, wherein parenteral administration (e.g., intramuscular, intravenous, subcutaneous, intraperitoneal, or intrathecal) of the dosage form provides an *in vivo* plasma profile for gaboxadol encompassing a mean $AUC_{0-\infty}$ of more than about 25 ng•hr/ml. In embodiments, single dose administration of the dosage form provides an *in vivo* plasma profile for gaboxadol encompassing a mean $AUC_{0-\infty}$ of more than about, e.g., 50 ng•hr/ml, 75 ng•hr/ml, 150 ng•hr/ml, 250 ng•hr/ml, 500 ng•hr/ml, 1000 ng•hr/ml, or 1500 ng•hr/ml.

In embodiments, the dosage form includes from about 1 mg to about 500 mg gaboxadol, wherein administration of the dosage form provides an *in vivo* plasma profile for gaboxadol encompassing a mean C_{max} of less than about 10000 ng/ml. In embodiments, single dose administration of the compositions provide an *in vivo* plasma profile for gaboxadol of a mean C_{max} of less than about, e.g., 5000 ng/ml, 2500 ng/ml, 1000 ng/ml, 500 ng/ml, 250 ng/ml, or 100 ng/ml.

In embodiments, pharmaceutical compositions for parenteral administration include gaboxadol or a pharmaceutically acceptable salt thereof wherein parenteral administration exhibits a pharmacokinetic profile of a T_{max} at about 1 to about 120 minutes after administration of the parenteral composition; followed by a plasma drug concentration of at least 50% C_{max} for a duration of about 90 to about 360 minutes. In embodiments, parenteral administration of gaboxadol is followed by a plasma drug concentration of at least 50% C_{max} for a duration of, e.g., about 10 to about 60 minutes, about 15 to about 90 minutes, about 30 to about 120 minutes, about 60 to about 180 minutes, about 90 to about 180 minutes.

In embodiments, stable pharmaceutical compositions are provided in unit dosage form in a vial or ampoule suitable for parenteral administration having a therapeutically effective amount of allosteric modulator and/or gaboxadol, or a pharmaceutically acceptable salt thereof, dissolved in sterile water to form a solution wherein the composition is substantially free of any excipient, organic solvent, buffer, acid, base, salt other than allosteric modulator and/or gaboxadol, or a pharmaceutically acceptable salt thereof. In embodiments, the pharmaceutical composition remains sufficiently soluble and is capable of direct administration. In embodiments, the pharmaceutical composition is capable of storage in the absence of an inert atmosphere for at least 6 months.

In embodiments, provided herein are stable pharmaceutical compositions in unit dosage form in a vial or ampoule suitable for parenteral administration having a therapeutically effective amount of allosteric modulator and/or gaboxadol, or a pharmaceutically acceptable salt thereof, dissolved in sterile water to form a solution wherein the composition is free of any excipient, organic solvent, buffer, acid, base, salt other than allosteric modulator and/or gaboxadol, or a pharmaceutically acceptable salt thereof. In embodiments, the pharmaceutical composition remains sufficiently soluble and is capable of direct administration. In embodiments, the pharmaceutical composition is capable of storage in the absence of an inert atmosphere for at least 6 months.

In embodiments, stable pharmaceutical compositions suitable for parenteral administration include allosteric modulator and/or gaboxadol, or a pharmaceutically acceptable salt thereof, in an aqueous solution having an osmolarity between 225 and 350 mOsm/kg and at a pH in the range between 7.0 and 8.0. In embodiments, the aqueous solution has an osmolarity between 270 and 310. In embodiments, the aqueous solution has a pH in the range between 7.2 and 7.8.

In embodiments, provided herein are methods of treating an epileptic disorder including administering to a patient in need thereof a first pharmaceutical and a second pharmaceutical composition including gaboxadol or a pharmaceutically acceptable salt thereof, and/or an allosteric modulator, wherein the second pharmaceutical composition that provides a stable *in vivo* plasma profile having a mean $AUC_{0-\infty}$ of at least about, e.g., 25%, 30%, 35%, 40%, 45% or 50% less than the first pharmaceutical composition. In embodiments, the composition provides improvement of next day functioning of the patient. For example, the composition may provide improvement in one or more symptoms for more than about, e.g., 6 hours, 8 hours, 10 hours, or 12 hours after administration of the first and/or second pharmaceutical composition.

In embodiments, parenteral compositions may be administered as needed, e.g., once, twice, thrice or four or more times daily, or continuously depending on the patient's needs. In embodiments, the parenteral compositions may be administered immediately or as soon thereafter as seizures start. In embodiments, the parenteral compositions may be administered as soon as warning signs of a seizure are exhibited such as auras, unusual smells, usual feelings, etc.

In embodiments, provided herein are methods of treating an epileptic disorder including administering to a patient in need thereof a first pharmaceutical dosage including a sub-

therapeutic dosage of an allosteric modulator wherein the composition provides improvement for more than 6 hours after administration.

In embodiments, provided herein are methods of treating an epileptic disorder including administering to a patient in need thereof a first pharmaceutical dosage including a sub
5 therapeutic dosage of gaboxadol or a pharmaceutically acceptable salt thereof wherein the composition provides improvement for more than 6 hours after administration.

In embodiments, provided herein are methods of treating an epileptic disorder including administering to a patient in need thereof a first pharmaceutical composition including an
10 allosteric modulator and a second pharmaceutical composition including gaboxadol or a pharmaceutically acceptable salt thereof wherein the second pharmaceutical composition provides an *in vivo* plasma profile having a mean $AUC_{0-\infty}$ of less than about 900 ng•hr/ml. In embodiments, the second pharmaceutical composition provides an *in vivo* plasma profile having a $AUC_{0-\infty}$ of less than about, e.g., 800 ng•hr/ml, 750 ng•hr/ml, 700 ng•hr/ml, 650 ng•hr/ml, or 600 ng•hr/ml. In embodiments, the second pharmaceutical composition provides an *in vivo*
15 plasma profile having a $AUC_{0-\infty}$ of less than about, e.g., 550 ng•hr/ml, 500 ng•hr/ml, 450 ng•hr/ml, 400 ng•hr/ml, or 350 ng•hr/ml. In embodiments, the second pharmaceutical composition provides an *in vivo* plasma profile having a $AUC_{0-\infty}$ of less than about, e.g., 300 ng•hr/ml, 250 ng•hr/ml, 200 ng•hr/ml, 150 ng•hr/ml, or 100 ng•hr/ml. In embodiments, the first and second pharmaceutical composition are administered wherein the compositions provide
20 improvement of next day functioning of the patient. In embodiments, the first pharmaceutical composition provides improvement in one or more symptom for more than, e.g., 6 hours, 8 hours or 12 hours after administration of the first pharmaceutical composition.

In embodiments, provided herein are methods of treating an epileptic disorder including administering to a patient in need thereof a first pharmaceutical composition including an
25 allosteric modulator and a second pharmaceutical composition including gaboxadol or a pharmaceutically acceptable salt thereof wherein the first composition provides an *in vivo* plasma profile with a C_{max} that is more than about 50% greater than the C_{max} provided by the administration of the second pharmaceutical composition. As used herein the C_{max} provided by the administration of the second pharmaceutical composition may or may not include the plasma
30 profile contribution of the first pharmaceutical composition. In embodiments, the administration of the second pharmaceutical composition does not include the plasma profile contribution of the

first pharmaceutical composition. In embodiments, the first composition provides an *in vivo* plasma profile having a C_{max} that is more than about *e.g.*, 60%, 70%, 80%, or 90% greater than the C_{max} provided by the administration of the second pharmaceutical composition.

In embodiments, the T_{max} of the first pharmaceutical composition is less than 3 hours. In
5 embodiments, the T_{max} of the first pharmaceutical composition is less than 2.5 hours. In
embodiments, the T_{max} of the first pharmaceutical composition is less than 2 hours. In
embodiments, the T_{max} of the first pharmaceutical composition is less than 1.5 hours. In
embodiments, the T_{max} of the first pharmaceutical composition is less than 1 hour.

In embodiments, the first pharmaceutical composition provides a dissolution of at least
10 about 80% within the first 20 minutes of administration to a patient in need thereof. In
embodiments, the first pharmaceutical composition provides a dissolution of at least about, *e.g.*,
85%, 90% or 95% within the first 20 minutes of administration to a patient in need thereof. In
embodiments, the first pharmaceutical composition provides a dissolution of at least 80% within
the first 10 minutes of administration to a patient in need thereof.

15 In embodiments the first and/or the second pharmaceutical compositions are sub
therapeutic dosages. A sub therapeutic dosage is an amount of active substance, *e.g.*, allosteric
modulator and/or gaboxadol, or a pharmaceutically acceptable salt thereof, that is less than the
amount required for a therapeutic effect. In embodiments, a sub therapeutic dosage is an amount
of allosteric modulator or a pharmaceutically acceptable salt thereof that alone may not provide
20 improvement in at least one symptom of the epileptic disorder but is sufficient to maintain such
improvement. In embodiments, a sub therapeutic dosage is an amount of gaboxadol
pharmaceutically acceptable salt thereof that alone may not provide improvement in at least one
symptom of the epileptic disorder but is sufficient to maintain such improvement. In
embodiments, the methods provide administering a first pharmaceutical composition that
25 provides improvement in at least one symptom of an epileptic disorder and a second composition
that maintains the improvement. In embodiments, after administration of the first pharmaceutical
composition, the second pharmaceutical composition may provide a synergistic effect to improve
at least one symptom of an epileptic disorder. In embodiments the second pharmaceutical
composition may provide a synergistic effect to improve at least one symptom of an epileptic
30 disorder.

In embodiments, provided herein are methods of treating an epileptic disorder including administering to a patient in need thereof a pharmaceutical composition including a first pharmaceutical dosage wherein the composition provides improvement for more than 6 hours after administration and a second pharmaceutical composition including a sub therapeutic dosage of gaboxadol or a pharmaceutically acceptable salt thereof.

Administration of the first and second pharmaceutical compositions may be simultaneous or separated by an interval of time to achieve immediate, intermediate or long-term improvement in at least one symptom. In embodiments, the first and second pharmaceutical composition may be administered 6 hours apart. In embodiments, the first and second pharmaceutical composition may be administered 12 hours apart. In embodiments, the first and second pharmaceutical compositions may administered within, *e.g.*, 15 minutes, 30 minutes, 1 hour, 2 hours, 6 hours, 12 hours, 18 hours, 24 hours etc. In embodiments, the first and second pharmaceutical composition may be administered together. In embodiments, the first and second pharmaceutical compositions may administered separated by at least, *e.g.*, 15 minutes, 30 minutes, 1 hour, 2 hours, 12 hours, 18 hours, 24 hours etc. In embodiments, improvement in at least one symptom of an epileptic disorder for more than 8 hours after administration to the patient is provided. In embodiments, improvement for more than about, *e.g.*, 10 hours, 12 hours, 15 hours, 18 hours, 20 hours, or 24 hours after administration to the patient is provided.

In embodiments, the administration of the first and second pharmaceutical composition may provide a synergistic effect to improve at least one symptom of an epileptic disorder.

In embodiments, the first and/or the second pharmaceutical composition include any of the aforementioned amounts of active substance, *e.g.*, allosteric modulator and/or gaboxadol a pharmaceutically acceptable salt thereof.

In embodiments, the first and/or the second pharmaceutical composition include 5 mg to 15 mg, 5 mg to 10 mg, 4 mg to 6 mg, 6 mg to 8 mg, 8 mg to 10 mg, 10 mg to 12 mg, 12 mg to 14 mg, 14 mg to 16 mg, 16 mg to 18 mg, or 18 mg to 20 mg active substance, *e.g.*, allosteric modulator and/or gaboxadol.

In embodiments, the first and/or the second pharmaceutical composition include 0.1 mg, 0.25 mg, 0.5 mg, 1 mg, 2.5 mg, 3 mg, 4 mg, 5 mg, 7 mg, 7.5 mg, 10 mg, 12.5 mg, 15 mg, 17.5 mg, 20 mg active substance, *e.g.*, allosteric modulator and/or gaboxadol or amounts that are multiples of such doses. In embodiments, the first pharmaceutical compositions include 2.5 mg,

5 mg, 7.5 mg, 10 mg, 15 mg, or 20 mg of an allosteric modulator. In embodiments, the second pharmaceutical compositions include 2.5 mg, 5 mg, 7.5 mg, 10 mg, 15 mg, or 20 mg of an allosteric modulator.

In embodiments, methods of treating epileptic disorders include administration of an
 5 allosteric modulator and/or gaboxadol or a pharmaceutically acceptable salt thereof, in combination with one or more other active compounds. The combination therapies can include administration of the active agents together in the same admixture, or in separate admixtures. In embodiments, the pharmaceutical composition includes two, three, or more active agents. In
 10 embodiments, the combinations result in a more than additive effect on the treatment of the disease or disorder. Thus, treatment is provided of an epileptic disorder with a combination of agents that combined, may provide a synergistic effect that enhances efficacy.

In embodiments, gaboxadol or a pharmaceutically acceptable salt thereof, is administered in combination with conventional therapy for seizures, epilepsy, or one of the other disorders disclosed herein. For example, common conventional therapies for seizures and epilepsy include
 15 antiepileptic drugs and non-antiepileptic drug treatments such as low carbohydrate diet (*e.g.*, ketogenic diets, such as classical diet, medium chain triglyceride (MCT) diet, modified Atkins diet (MAD), and low glycemic index treatment (LGIT)), intravenous immunoglobulin, steroids, elimination diet, vagus nerve stimulation, corticectomy, and multiple subpial transections.

Common antiepileptic and anticonvulsive active compounds that may be used in
 20 combination with an allosteric modulator and/or gaboxadol or a pharmaceutically acceptable salt thereof include, but are not limited to, acetazolamide, eslicarbazepine acetate, ethosuximide, gabapentin, lacosamide, lamotrigine, levetiracetam, nitrazepam, oxcarbazepine, perampanel, piracetam, phenobarbital, phenytoin, pregabalin, primidone, retigabine, rufinamide, sodium valproate, stiripentol, tiagabine, topiramate, vigabatrin, and zonisamide.

25 The disclosed compounds, such as gaboxadol or pharmaceutically acceptable salts thereof, or an allosteric modulator, pharmaceutically acceptable salts, derivatives and/or analogues thereof, can be used individually as a monotherapy as the only active agent. In embodiments, methods are provided of treating epileptic disorders using an allosteric modulator or a pharmaceutically acceptable salt thereof. In embodiments, methods of treating epileptic
 30 disorders include administration of an allosteric modulator, pharmaceutically acceptable salts, derivatives and/or analogues thereof in combination with one or more other active agent, *e.g.*,

allosteric modulator or gaboxadol. The combination therapies can include administration of the active agent, *e.g.*, allosteric modulator or gaboxadol, together in the same admixture, or in separate admixtures. In embodiments, the pharmaceutical composition includes two, three, or more active agents. In embodiments, the combinations result in a more than additive effect on the treatment of the disease or disorder. Thus, treatment is provided of an epileptic disorder with a combination of agents that combined, may provide a synergistic effect that enhances efficacy.

In embodiments, gaboxadol or pharmaceutically acceptable salts thereof, or an allosteric modulator, pharmaceutically acceptable salts, derivatives and/or analogues thereof, or both, is administered in combination with conventional therapy for seizures, epilepsy, or one of the other disorders disclosed herein. For example, common conventional therapies for seizures and epilepsy include antiepileptic drugs and non-antiepileptic drug treatments such as low carbohydrate diet (*e.g.*, ketogenic diets, such as classical diet, medium chain triglyceride (MCT) diet, modified Atkins diet (MAD), and low glycemic index treatment (LGIT)), intravenous immunoglobulin, steroids, elimination diet, vagus nerve stimulation, corticectomy, and multiple subpial transections.

Common antiepileptic and anticonvulsive active compounds that may be used in combination with an allosteric modulator include, but are not limited to, acetazolamide, carbamazepine, clobazam, clonazepam, eslicarbazepine acetate, ethosuximide, gabapentin, lacosamide, lamotrigine, levetiracetam, nitrazepam, oxcarbazepine, perampanel, piracetam, phenobarbital, phenytoin, pregabalin, primidone, retigabine, rufinamide, sodium valproate, stiripentol, tiagabine, topiramate, vigabatrin, and zonisamide.

In embodiments, a co-therapy of an allosteric modulator, a pharmaceutically acceptable salt thereof, or a derivative thereof and gaboxadol or a pharmaceutically acceptable salt thereof is effective to reduce seizure frequency or severity in the subject greater than either compound is administered alone. In embodiments, the co-therapy produces a more than additive result compared to compounds administered individually.

In embodiments, the subject may be started at a low dose and the dosage is escalated. In this manner, it can be determined if the drug is well tolerated in the subject. Dosages can be lower for children than for adults.

In embodiments, such as combination therapies, a dose of gaboxadol for children can be 0.1 mg/kg to 1 mg/kg, and the dose for an allosteric modulator may be 0.01 mg/kg to 0.1mg/kg.

In embodiments, the weight/weight ratio of gaboxadol and an allosteric modulator can be 10-to-1. However, the dosing ratio based on milligrams of active pharmaceutical ingredient (API) can range from 0.1-to-1 to 100-to-1 of gaboxadol-to-an allosteric modulator respectively.

Effective treatment of an epileptic disorder (*e.g.*, acute repetitive seizure) herein may be established by showing reduction in the frequency of seizures (*e.g.*, more than 50%) after a period of time compared with baseline. For example, after a baseline period of 1 month, the patients may be randomly allocated gaboxadol or an allosteric modulator or placebo as add-on therapy to standard therapies, such as valproate and clobazam, during a double-blind period of 2 months. Primary outcome measurements may include the percentage of responders on gaboxadol or an allosteric modulator and on placebo, defined as having experienced at least a 50% reduction of clonic (or tonic-clonic) seizure frequency during the second month of the double-blind period compared with baseline. Patients who present with status epilepticus during the double-blind period may be regarded as non-responders. Secondary outcomes may include the absolute count of clonic (or tonic-clonic) seizures during the second month of the double-blind period (normalized to 30 days, by dividing the raw count by the exact number of days of observation and multiplying by 30) and the percentage of change from baseline.

The effectiveness of gaboxadol and/or an allosteric modulator for the treatment of a disclosed epileptic disorder, *e.g.*, associated with Dravet syndrome or Lennox-Gastaut syndrome, may be established in other controlled studies. For example, a randomized, double-blind, placebo-controlled study consisting of a 4-week baseline period followed by a 3-week titration period and 12-week maintenance period may be used in patients age 2-54 years with a current or prior diagnosis of Dravet syndrome or LGS. Multiple target maintenance doses of gaboxadol and/or an allosteric modulator may be tested according to patient body weight and specific dosing regime. A primary efficacy measure may include the percent reduction in the weekly frequency of drop seizures (atonic, tonic, or myoclonic), also known as drop attacks, from the 4-week baseline period to 12-week maintenance period. Thus, efficacy may be measured as percentage reduction in weekly seizure (*e.g.*, atonic, tonic, or myoclonic) frequency from baseline of, *e.g.*, 0 to <20, 20 to <40, 40 to <60, 60 to <80, 80 to <100.

Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of skill in the art to which the disclosure herein belongs.

The term "about" or "approximately" as used herein means within an acceptable error range for the particular value as determined by one of ordinary skill in the art, which will depend in part on how the value is measured or determined, *i.e.*, the limitations of the measurement system. For example, "about" can mean within 3 or more than 3 standard deviations, per the practice in the art. Alternatively, "about" can mean a range of up to 20%, preferably up to 10%, more preferably up to 5%, and more preferably still up to 1% of a given value. Alternatively, particularly with respect to biological systems or processes, the term can mean within an order of magnitude, preferably within 5-fold, and more preferably within 2-fold, of a value.

As used herein, the term "treating" or "treatment" refers to alleviating, attenuating or delaying the appearance of clinical symptoms of a disease or condition in a subject that may be afflicted with or predisposed to the disease or condition, but does not yet experience or display clinical or subclinical symptoms of the disease or condition. In certain embodiments, treating" or "treatment" may refer to preventing the appearance of clinical symptoms of a disease or condition in a subject that may be afflicted with or predisposed to the disease or condition, but does not yet experience or display clinical or subclinical symptoms of the disease or condition. "Treating" or "treatment" may also refer to inhibiting the disease or condition, *e.g.*, arresting or reducing its epileptic or at least one clinical or subclinical symptom thereof. "Treating" or "treatment" further refers to relieving the disease or condition, *e.g.*, causing regression of the disease or condition or at least one of its clinical or subclinical symptoms. The benefit to a subject to be treated may be statistically significant, mathematically significant, or at least perceptible to the subject and/or the physician. Nonetheless, prophylactic (preventive) and therapeutic treatment are two separate embodiments of the disclosure herein.

"Effective amount" or "therapeutically effective amount" means a dosage sufficient to alleviate one or more symptom of a disorder, disease, or condition being treated, or to otherwise provide a desired pharmacological and/or physiologic effect.

"Improvement" refers to the treatment of symptoms or conditions associated with epileptic disorders, measured relative to at least one symptom or condition of the metabolic disease.

"Improvement in next day functioning" or "wherein there is improvement in next day functioning" refers to improvement after waking from an overnight sleep period wherein the beneficial effect of administration of one or more of gaboxadol or a pharmaceutically acceptable

salt thereof alone, or an allosteric modulator alone, or gaboxadol in combination an allosteric modulator, applies to at least one symptom or condition associated with an epileptic disorder and is discernable, either subjectively by a patient or objectively by an observer, for a period of time, e.g., immediately, 1 hour, 2 hours, hours, 3 hours, 4 hours, 5 hours, 6 hours, 12 hours, 24 hours, etc. after waking.

"Composition", "pharmaceutical composition", "formulation", "pharmaceutical formulation" are used interchangeably herein. "Composition", "pharmaceutical composition", "formulation", "pharmaceutical formulation" encompass dosage forms. Dosage forms can encompass unit doses.

"Pharmaceutically acceptable" refers to molecular entities and compositions that are "generally regarded as safe", e.g., that are physiologically tolerable and do not typically produce an allergic or similar untoward reaction, such as gastric upset and the like, when administered to a human. In embodiments, this term refers to molecular entities and compositions approved by a regulatory agency of the federal or a state government, as the GRAS list under section 204(s) and 409 of the Federal Food, Drug and Cosmetic Act, that is subject to premarket review and approval by the FDA or similar lists, the U.S. Pharmacopeia or another generally recognized pharmacopeia for use in animals, and more particularly in humans.

As used herein, the term "prevention" or "preventing" means to administer a composition to a subject or a system at risk for or having a predisposition for one or more symptoms caused by a disease or disorder to facilitate cessation of a particular symptom of the disease or disorder, a reduction or prevention of one or more symptoms of the disease or disorder, a reduction in the severity of the disease or disorder, the complete ablation of the disease or disorder, stabilization or delay of the epileptic or progression of the disease or disorder.

"Prodrug", as used herein, refers to a pharmacological substance (drug) that is administered to a subject in an inactive (or significantly less active) form. Once administered, the prodrug is metabolized in the body (*in vivo*) into a compound having the desired pharmacological activity.

"Analog" "analogue, and "derivative" are used herein interchangeably and refer to a compound that possesses the same core as the parent compound, but may differ from the parent compound in bond order, the absence or presence of one or more atoms and/or groups of atoms, and combinations thereof. The derivative can differ from the parent compound, for example, in

one or more substituents present on the core, which may include one or more atoms, functional groups, or substructures. In general, a derivative can be imagined to be formed, at least theoretically, from the parent compound via chemical and/or physical processes.

“Stereoisomer”, as used herein, refers to isomeric molecules that have the same
5 molecular formula and sequence of bonded atoms (constitution), but which differ in the three dimensional orientations of their atoms in space. Examples of stereoisomers include enantiomers and diastereomers. As used herein, an enantiomer refers to one of the two mirror-image forms of an optically active or chiral molecule. Diastereomers (or diastereoisomers) are stereoisomers that are not enantiomers (non-superimposable mirror images of each other).
10 Chiral molecules contain a chiral center, also referred to as a stereocenter or stereogenic center, which is any point, though not necessarily an atom, in a molecule bearing groups such that an interchanging of any two groups leads to a stereoisomer. In organic compounds, the chiral center is typically a carbon, phosphorus or sulfur atom, though it is also possible for other atoms to be stereocenters in organic and inorganic compounds. A molecule can have multiple stereocenters,
15 giving it many stereoisomers. In compounds whose stereoisomerism is due to tetrahedral stereogenic centers (*e.g.*, tetrahedral carbon), the total number of hypothetically possible stereoisomers will not exceed 2^n , where n is the number of tetrahedral stereocenters. Molecules with symmetry frequently have fewer than the maximum possible number of stereoisomers. A 50:50 mixture of enantiomers is referred to as a racemic mixture. Alternatively, a mixture of
20 enantiomers can be enantiomerically enriched so that one enantiomer is present in an amount greater than 50%. Enantiomers and/or diastereomers can be resolved or separated using techniques known in the art. “Chirality” also includes axial and planar chirality.

The term “pharmaceutically acceptable salt”, as used herein, refers to derivatives of the compounds defined herein, wherein the parent compound is modified by making acid or base
25 salts thereof. Example of pharmaceutically acceptable salts include but are not limited to mineral or organic acid salts of basic residues such as amines; and alkali or organic salts of acidic residues such as carboxylic acids. The pharmaceutically acceptable salts include the conventional non-toxic salts or the quaternary ammonium salts of the parent compound formed, for example, from non-toxic inorganic or organic acids. Such conventional non-toxic salts
30 include those derived from inorganic acids such as hydrochloric, hydrobromic, sulfuric, sulfamic, phosphoric, and nitric acids; and the salts prepared from organic acids such as acetic,

propionic, succinic, glycolic, stearic, lactic, malic, tartaric, citric, ascorbic, pamoic, maleic, hydroxymaleic, phenylacetic, glutamic, benzoic, salicylic, sulfanilic, 2-acetoxybenzoic, fumaric, tolunesulfonic, naphthalenesulfonic, methanesulfonic, ethane disulfonic, oxalic, and isethionic salts. The pharmaceutically acceptable salts of the compounds can be synthesized from the parent
 5 compound, which contains a basic or acidic moiety, by conventional chemical methods. For example, gaboxadol may be formulated for administration to a patient using pharmaceutically acceptable salts including acid addition salts, a zwitter ion hydrate, zwitter ion anhydrate, hydrochloride or hydrobromide salt, or in the form of the zwitter ion monohydrate. Acid addition salts, include but are not limited to, maleic, fumaric, benzoic, ascorbic, succinic, oxalic,
 10 bis-methylenesalicylic, methanesulfonic, ethane-disulfonic, acetic, propionic, tartaric, salicylic, citric, gluconic, lactic, malic, mandelic, cinnamic, citraconic, aspartic, stearic, palmitic, itaconic, glycolic, p-amino-benzoic, glutamic, benzene sulfonic or theophylline acetic acid addition salts, as well as the 8-halotheophyllines, for example 8-bromo-theophylline. In other suitable embodiments, inorganic acid addition salts, including but not limited to, hydrochloric,
 15 hydrobromic, sulfuric, sulfamic, phosphoric or nitric acid addition salts may be used.

"Excipient" is a substance, other than the active drug substance, *e.g.*, gaboxadol, of a pharmaceutical composition, which has been appropriately evaluated for safety and are included in a drug delivery system to either aid the processing of the drug delivery system during its manufacture; protect; support; enhance stability, bioavailability, or patient acceptability; assist in
 20 product identification; or enhance any other attributes of the overall safety and effectiveness of the drug delivery system during storage or use.

"Stabilizer" or "stabilizing amount" refers to an amount of one or more excipients included in the parenteral compositions that provide sufficient stability but do not adversely affect the bioavailability, safety and/or efficacy of allosteric modulator and/or gaboxadol or
 25 pharmaceutically acceptable salt thereof used in the composition.

"Stable" means that there is substantially no degradation of the gaboxadol or pharmaceutically acceptable salt thereof after a specified period of time, *e.g.*, after 3 months or 6 months.

"Soluble" means that the solution of allosteric modulator and/or gaboxadol or
 30 pharmaceutically acceptable salt thereof does not turn hazy and/or there is substantially no precipitate in the solution

"Sufficiently soluble" means that the particle content is sufficiently low, and the material is sufficiently sterile such that it is useful for parenteral administration. For example, the number of particles in a liquid composition should be, *e.g.*, less than 6,000 10 μ m particles should be present in a volume of 10 ml solvent, preferably less than 10,000, less than 5,000, less than 3,000, less than 1,000, or less than 400 10 μ m particles. In some examples, the number of particles in a liquid composition should be less than 1000, less than 600, or less than 200 25 μ m particles in the 10 ml volume.

"Local site compatible" herein shall mean the composition is tolerant at the site of injection or infusion, thus minimizing side effects, such as local skin irritations or venous irritations, including inflammatory reactions at the infusion site. The parenteral compositions herein may have less side reactions than conventional products, such as skin irritation or phlebitis.

"Purified" as used herein refers to material that has been isolated under conditions that reduce or eliminate the presence of unrelated materials, *i.e.*, contaminants, including native materials from which the material is obtained. As used herein, the term "substantially free" is used operationally, in the context of analytical testing of the material. Preferably, purified material substantially free of contaminants is at least 95% pure; more preferably, at least 97% pure, and more preferably still at least 99% pure. Purity can be evaluated, for example, by chromatography or any other methods known in the art. In embodiments, purified means that the level of contaminants is below a level acceptable to regulatory authorities for safe administration to a human or non-human animal.

"Ready-to-use" with reference to the compositions herein shall mean the preparation in the reconstituted form, with standardized concentration and quality, prefilled in the single-use container, such as glass vials, infusion bags or syringes, ready for direct administration to the patient.

"Direct administration" with reference to the compositions herein shall mean the immediate administration, *i.e.*, without further dilution, premixing with other substances or otherwise changing the composition or formulation of the composition. Such composition is typically directly discharged from an infusion device and administered via a vascular access port or through a central line.

"Dosage" is intended to encompass a formulation expressed in terms of $\mu\text{g/kg/day}$, $\mu\text{g/kg/hr}$, mg/kg/day or mg/kg/hr . The dosage is the amount of an ingredient administered in accordance with a particular dosage regimen. A "dose" is an amount of an agent administered to a mammal in a unit volume or mass, *e.g.*, an absolute unit dose expressed in mg or μg of the agent. The dose depends on the concentration of the agent in the formulation, *e.g.*, in moles per liter (M), mass per volume (m/v), or mass per mass (m/m). The two terms are closely related, as a particular dosage results from the regimen of administration of a dose or doses of the formulation. The particular meaning in any case will be apparent from context.

"Patient" and "subject" are used interchangeably herein and include, but not limited to, primates such as humans, canines, porcine, ungulates, rodents, poultry, and avian.

"Co-administered with", "in combination with", "administered in combination with", "a combination of", "administered along with", or "co-therapy", may be used interchangeably and mean that two or more agents are administered in the course of therapy. The agents may be administered together at the same time or separately in spaced apart intervals. The agents may be administered in a single dosage form or in separate dosage forms.

"PK" refers to the pharmacokinetic profile. C_{max} is defined as the highest plasma drug concentration estimated during an experiment (ng/ml). T_{max} is defined as the time when C_{max} is estimated (min). $\text{AUC}_{0-\infty}$ is the total area under the plasma drug concentration-time curve, from drug administration until the drug is eliminated ($\text{ng}\cdot\text{hr/ml}$ or $\mu\text{g}\cdot\text{hr/ml}$). The area under the curve is governed by clearance. Clearance is defined as the volume of blood or plasma that is totally cleared of its content of drug per unit time (ml/min).

EXAMPLES

The Examples provided herein are included solely for augmenting the disclosure herein and should not be considered to be limiting in any respect.

Example 1

Gaboxadol Plasma Concentration Profiles

The following Example provides the plasma concentration profiles and dose proportionality of gaboxadol monohydrate following single oral doses ranging from 2.5 to 20

mg. The absolute bioavailability of gaboxadol monohydrate capsules ranging from 2.5 to 20 mg is also assessed.

This study was composed of separate groups of 10 healthy adult subjects (at least 4 of each gender) who participated in a 6-period, double-blind, randomized, crossover study designed to access the dose proportionality and absolute bioavailability of 5 single oral doses of gaboxadol across the dose range of 2.5 to 20 mg. The order in which the subjects received the 5 single oral doses of gaboxadol (2.5; 5; 10; 15; and 20 mg) was randomized within Treatment Periods 1 through 5. Each subject was expected to complete all 6 treatment periods and there was a washout of at least 4 days between each treatment period.

Each oral dosing within Treatment Periods consisted of 2 capsules of test drug taken simultaneously at each scheduled dosing. The treatment designations for the orally administered study drugs were as follows: Treatment A - one 2.5 mg gaboxadol capsule and 1 matching placebo capsule; Treatment B - one 5 mg gaboxadol capsule and 1 matching placebo capsule; Treatment C - one 10 mg gaboxadol capsule and 1 matching placebo capsule; Treatment D - one 15 mg gaboxadol capsule and 1 matching placebo capsule; and Treatment E - 20 mg gaboxadol (two 10 mg gaboxadol capsules). Subjects received their study drug after an overnight fast with 240 mL of water in the morning about 8:00 AM. Water was permitted *ad libitum* except within 1 hour prior to and after study drug administration. No food was allowed for 4 hours post dose.

For each subject in each treatment, plasma and urine samples were collected over 16 hours post-dosing for the determination of pharmacokinetic parameters (*e.g.*, AUC, C_{max}, T_{max}, apparent t_{1/2}, cumulative urinary excretion, renal clearance, clearance, and steady-state volume of distribution, as appropriate). AUC and C_{max} for gaboxadol were potency adjusted to facilitate comparison of pharmacokinetic data across studies. Table 1 provides the individual potency-adjusted pharmacokinetic parameters of gaboxadol following single oral doses (2.5, 5, 10, 15, and 20 mg).

Table 1. Pharmacokinetic parameters for gaboxadol following oral and IV administration

Pharmacokinetic parameters for gaboxadol following oral and IV administration							
Parameter	Geometric Mean (N=10)						Slope (90 % CI) ^{††}
	2.5 mg	5 mg	10 mg Oral	10 mg I.V.	15 mg	20 mg	
AUC _{0-∞} (ng·hr/mL)	90	171	346	380	539	669	0.98 (0.95, 1.01)
C _{max} (ng/mL) [†]	61	110	232	212	382	393	0.95 (0.88, 1.02)
T _{max} (hr) [‡]	0.5	0.6	0.5	~	0.5	0.6	
Apparent t _{1/2} (hr) [§]	1.5	1.5	1.6	1.5	1.5	1.6	
CL/F (mL/min) [¶]	461	488	476	438	469	499	
F _e (%)	43	45	53	53	50	53	
CL _R (mL/min)	196	222	250	208	234	265	
F (%) (90% CI) [#]	92% (0.86, 0.97)						
[†] C _{coi} (ng/mL) for 10 mg. IV.							
[‡] Median.							
[§] Harmonic Mean.							
[¶] CL (mL/min) for 10 mg IV.							
[#] Bioavailability relative to 10 mg I.V. reference based on pooled dose-adjusted (to 10 mg) oral AUC _{0-∞} values.							
^{††} Dose proportionality assessment of oral treatments only.							

5

Example 2

Assessment of Residual Effects Resulting from Gaboxadol Administration

This study was a double blind, double-dummy, randomized, active- and placebo- controlled, single dose, 3-period crossover study, followed by an open-label, single-dose, single period study in healthy elderly male and female subjects. Subjects were randomized to each of 3 treatments (Treatments A, B, and C) to be administered in a crossover manner over the first 3 treatment periods. For Treatment A, subjects received a single dose of gaboxadol 10 mg; for Treatment B, subjects received a single dose of flurazepam 30 mg; and for Treatment C, subjects received a single dose of placebo. Doses were administered orally at bedtime on Day 1. Subjects were domiciled from early in the evening of dosing until ~36 hours post-dose (morning of Day 3) during each treatment period. The subjects who participated in treatment periods 1-3 participated in a fourth treatment period. In this period, a single dose of gaboxadol 10 mg (Treatment D) was administered orally in an open-label manner on the morning of Day 1 for PK of gaboxadol. There was at least a 14-day washout between the doses of consecutive treatment

20

periods. Study participants included healthy, elderly male and female subjects between 65 and 80 years of age, with a Mini Mental Status 24, weighing at least 55 kg. All subjects received 10 mg gaboxadol monohydrate capsules and 30 mg flurazepam (provided as 2 x 15 mg capsules), matching placebo was provided for both gaboxadol and flurazepam.

5 The primary endpoints evaluated included pharmacodynamics (measurement of psychomotor performance, memory, attention and daytime sleepiness the following pm dosing), gaboxadol pharmacokinetics, and safety. Gaboxadol (single dose 10 mg) did not show residual effect 9 hours post-dose on the primary endpoints Choice Reaction Time and Critical Flicker Fusion, whereas the active reference Flurazepam (30 mg single dose) showed significant effect
10 on the same tests. In addition, gaboxadol did not show any signs of residual effects on other measurements applied in the study (Multiple Sleep Latency Test (MSLT); Digit symbol substitution test (DSST), Tracking, Memory tests, Body Sway, and Leeds Sleep Evaluation Questionnaire).

15

Example 3

Evaluation of the Ability of Allopregnanolone, Ganaxalone, and Gaboxadol to Block Benzodiazepine-Resistant Status Epilepticus

20 Allopregnanolone, ganaxalone, and gaboxadol are evaluated for acute anticonvulsant efficacy when administered with escalating doses at 30 min after the onset of convulsive status epilepticus, a time-point typically resistant to benzodiazepines. Results obtained with these agents are compared to those obtained from side-by-side studies with vehicle-treated animals.

25 Twenty-four hours prior to administration of the chemoconvulsant pilocarpine, male Sprague Dawley rats (n = 10 time or treatment/group, 100-125 g; Charles River Laboratories) are treated systemically with lithium chloride (127 mg/kg; intraperitoneal (i.p.)). On the next day, the rats receive pilocarpine hydrochloride (50 mg/kg; i.p.) and are monitored carefully for the presence or absence of convulsive seizure activity. Administration of pilocarpine induces behavioral seizures within 5-20 min and any rat not showing convulsive seizure activity within
30 45 min of pilocarpine administration is excluded from further study. On the study day, the ability of each of the investigational compounds (allopregnanolone (ALLO), ganaxalone (GNX), or gaboxadol (GBD)) or vehicle (VEH) (40% hydroxypropyl β -cyclodextrin) to halt convulsive

status epilepticus in the Li-Pilo model of status epilepticus is evaluated with escalating doses administered i.p. 30 min after the first observed convulsive seizure. Throughout the study, the experimenter conducting the behavioral observations is blinded to treatment conditions (i.e., allopregnanolone, ganaxalone, or gaboxadol). All rats are observed and scored for seizure severity for 120 min post drug administration, and any accompanying behavioral effects are also noted by an experimenter blinded to treatment conditions. At the conclusion of the behavioral observation period, a 3 mL injection of lactated Ringer's solution is administered to all surviving rats to replace any SE-induced fluid loss.

The dose of each of the investigational compounds (allopregnanolone, ganaxalone, or gaboxadol) is varied in groups of ten rats until at least two points are established between the limits of 100% protection (no further convulsive seizures after 10 min of drug administration) and 0% protection. The dose of drug required to produce the desired endpoint in 50% of animals (ED50 or TD50) and the 95% confidence interval is calculated by a computer program based on the Probit method (Finney DJ. Probit Analysis. Cambridge University Press. 1971). This dose response evaluation typically requires up to 5 treatment groups per investigational compound, for a total of up to 50 rats per compound. Thus, up to 150 rats can be used for the investigational compounds (allopregnanolone, ganaxalone, or gaboxadol) quantification, and there is also one vehicle treatment group (n = 10), such that the total number of rodents in Study 1 is 160. All animals in the study are retained for 24 hours following completion of the study for assessment of weight change. The doses administered for allopregnanolone, ganaxalone, or gaboxadol are 0.5 mg/kg, 2 mg/kg, 5 mg/kg, 10 mg/kg, or 20 mg/kg.

Pharmacokinetics Sample Collection: Brains and plasma may be collected for assessment from a satellite cohort of rats for each dose (n = 3 rats/dose/compound; up to 45 rats in total). Plasma is collected from trunk blood after 10,000xg centrifugation for 10 min at 4°C. The anticoagulant is lithiumheparin. Brains are snap frozen on dry ice. All samples are stored at -20°C. The testing procedure timeline is set forth in FIG. 2.

Dose-response curves are constructed and expressed as ED50 (95% confidence intervals) calculated for each investigational compound (allopregnanolone, ganaxalone, or gaboxadol) administered 30 min after the first observed convulsive seizure. In cases that the data does not permit the calculation of an ED50 (95% confidence interval) because of lack of efficacy, the highest dose tested is noted. Additional measures of effects, e.g., motor impairment, weight

change post-SE, or survival from SE for 24 hrs are also recorded during the studies for all treatment conditions.

Table 2. Overall survival and protection results

Compound (dose, mg/kg)	Protection (includes PK rats)	24-hour Survival (excludes PK rats)
VEH (40% HPBCD)	0/13 (0%)	8/11 [#] (72%)
ALLO (0.5 mg/kg)	0/13 (0%)	6/10 (60%)
GBD (0.5 mg/kg)	2/13 (15%)	7/10 (70%)
GNX (0.5 mg/kg)	0/13 (0%)	6/10 (60%)
ALLO (2 mg/kg)	0/13 (0%)	7/10 (70%)
GBD (2 mg/kg)	0/13 (0%)	8/10 (80%)
GNX (2 mg/kg)	0/13 (0%)	8/10 (80%)
ALLO (5 mg/kg)	0/13 (0%)	7/9 (72%)
GBD (5 mg/kg)	0/13 (0%)	6/11 [#] (55%)
GNX (5 mg/kg)	0/13 (0%)	7/10 (70%)
ALLO (10 mg/kg)	2/13 (15%)	9/10 (90%)
GBD (10 mg/kg)	0/13 (0%)	10/10 (100%)*
GNX (10 mg/kg)	1/13 (7.7%)	9/10 (90%)
ALLO (20 mg/kg)	1/13 (7.7%)	9/10 (90%)
GBD (20 mg/kg)	0/13 (0%)	7/9 (78%)
GNX (20 mg/kg)	4/13 (30.8%)*	9/10 (90%)

5

*Significantly different from VEH, $p < 0.05$

[#] Spontaneous death during SE in PK group rat 110 min post –SE onset, but sample was collected.

10

FIG. 3 is a bar graph showing the percent protected versus dose with respect to allopregnanolone, ganaxalone, or gaboxadol. FIG. 4 is a bar graph showing the 24-hour survival results based on dose with respect to allopregnanolone, ganaxalone, or gaboxadol. FIG. 5 is a bar graph showing the number of observed seizures versus dose. FIG. 6A is a bar graph showing

body weight changes 24 hours post-status epilepticus as a function of percent loss versus dose. FIG. 6B is a bar graph showing 24 hour body weight loss for 0.5 mg/kg dose groups. Dose-response evaluation of treatment during benzodiazepine-resistant status epilepticus suggests dose-dependent efficacy of ganaxalone (i.p.) – significant ($p < 0.02$) improvement in protection at 20 mg/kg. Potential for inverted U-response profile of gaboxadol (i.p.) – $p = 0.071$ at 0.5 mg/kg. Significant improvements in 24-hour survival are shown for gabaxodol-treated rats (10 mg/kg).

Example 4

- 10 Prospective study for the Characterization of the Ability of Allopregnanolone, Ganaxalone, and Gaboxadol to Block Benzodiazepine-Resistant Status Epilepticus

Initial dose-response studies with intraperitoneal (i.p.) administration of allopregnanolone (ALLO), ganaxalone (GNX), and gaboxadol (GBD) suggests the potential for efficacy against benzodiazepine-resistant status epilepticus (Example 3). This study will assess the potential for synergistic activity of gaboxadol with either allopregnanolone, ganaxalone, or the benzodiazepine, lorazepam (LZP), against benzodiazepine-resistant status epilepticus in the Li-Pilo model in rats. Specifically, gaboxadol at low-doses (e.g. 0.5 mg/kg) will be administered in combination with a fixed dose of either allopregnanolone, ganaxalone, or lorazepam. The doses of allopregnanolone and ganaxalone will be those that were previously found to confer anticonvulsant effect in Example 3 (Table 2). The dose of lorazepam will be 2 mg/kg (Walton and Treiman, 1990. *Neurology* 40: 990-994 1990). The activity of each compound will also be evaluated alone. A vehicle-treatment group will be included. At 30 min post- status epilepticus onset, a single i.p. dose of each dose or dose combination (Table 3) will be administered to rats (n = 13/group). Rats will be observed for presence or absence of further convulsive activity.

Table 3. Investigational Compounds and Combinations to be Evaluated (n = 13 rats / treatment group with n = 3 rats sacrificed for PK analysis).

Compound	Dose (all i.p.)
GBD	0.25 mg/kg
GBD	0.5 mg/kg
ALLO	10 mg/kg
GNX	20 mg/kg
GNX	30 mg/kg
LZP	2 mg/kg
GBD + ALLO	0.5 mg/kg + 10 mg/kg
GBD + GNX	0.5 mg/kg + 20 mg/kg
GBD + LZP	0.5 mg/kg + 2 mg/kg
VEH	40% HPBCD

- 5 Twenty-four hours prior to administration of the chemoconvulsant pilocarpine, male Sprague Dawley rats (n = 13 time or treatment/group; 100-125 g at arrival from Charles River Laboratories) will be treated systemically with lithium chloride (127 mg/kg; i.p.). On the next day, the rats will receive pilocarpine hydrochloride (50 mg/kg; i.p.) and monitored carefully for the presence or absence of convulsive seizure activity. Administration of pilocarpine induces
- 10 behavioral seizures within 5-20 min and any rat not showing convulsive seizure activity within 45 min of pilocarpine administration will be excluded from further study. On the study day, the ability of each of the investigational compounds (allopregnanolone, ganaxalone, or gaboxadol) or vehicle (VEH) to halt convulsive status epilepticus in the Li-Pilo model of status epilepticus will be evaluated as outlined in Table 3 with administration 30 min after the first observed
- 15 convulsive seizure. Throughout the study, the experimenter conducting the behavioral observations will be blinded to treatment conditions (e.g., allopregnanolone, ganaxalone, or gaboxadol, lorazepam, or VEH). All rats will be observed and scored for seizure severity for 120 min post drug administration, and any accompanying behavioral effects will also be noted by an experimenter blinded to treatment conditions. At the conclusion of the behavioral observation

period, a 3 mL injection of lactated Ringer's solution will be administered to all surviving rats to replace any status epilepticus -induced fluid loss.

5 The dose of each of the investigational compounds will be administered to rats (n = 13/compound dose; Table 3), for a total of 130 rats. A 15% non-responder rate is anticipated for rats pre-treated with Li-Cl, but that do not develop convulsive status epilepticus in the 45-minute time period. Thus, up to 150 rats will be used for this study, which includes potential non-responders. All animals in the study will be retained for 24 hours following completion of the study for assessment of weight change, as well as overall behavioral appearance at that time (e.g. lethargic/active). Behavioral appearance will be assessed by an investigator blinded to treatment
10 condition.

Pharmacokinetics Sample Collection: Brains and plasma will be collected for assessment from a cohort of rats for each dose (n = 3 rats/dose/compound). Plasma will be isolated from trunk blood after 10,000xg centrifugation for 10 min at 4°C. The anticoagulant will be lithium-heparin. Brains would be snap frozen on dry ice. The testing procedure timeline is set forth on
15 FIG. 7.

Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments described herein. Such equivalents are intended to be encompassed by the claims.

REMARKS

By this amendment, claims 1-63 have been cancelled and new claims 64-82 have been added.

Early and favorable consideration of this application is respectfully requested.

Dated: August 10, 2017

Respectfully submitted,

/Jeffrey S. Steen, Reg. #32063/

Jeffrey S. Steen
Reg. No.: 32,063
Attorney for Applicants

Carter, DeLuca, Farrell & Schmidt, LLP
445 Broad Hollow Road - Suite 420
Melville, New York 11747
Tel.: (631) 501-5700
Fax: (631) 501-3526

JSS:jd

This listing of claims will replace all prior versions, and listings, of claims in the application:

Listing of Claims:

Claims 1-63 (Cancelled).

64. (New) A method of treating an epileptic disorder comprising administering to a patient in need thereof a pharmaceutical composition comprising gaboxadol or a pharmaceutically acceptable salt thereof.
65. (New) The method of claim 64, wherein the epileptic disorder is selected from the group consisting of epilepsy, epilepsy with generalized tonic-clonic seizures, epilepsy with myoclonic absences, frontal lobe epilepsy, temporal lobe epilepsy, Landau-Kleffner Syndrome, Ohtahara syndrome, Rasmussen's syndrome, West's syndrome, Lennox-Gastaut syndrome (LGS), Rett syndrome, CDKL5 disorder, childhood absence epilepsy, essential tremor, Dravet syndrome, Doose syndrome, acute repetitive seizures, benign rolandic epilepsy, status epilepticus, refractory status epilepticus, super-refractory status epilepticus (SRSE), PCDH19 pediatric epilepsy, increased seizure activity or breakthrough seizures.
66. (New) The method of claim 64, wherein the epileptic disorder is status epilepticus.
67. (New) The method of claim 64, wherein the epileptic disorder is acute repetitive seizure.
68. (New) The method of claim 64, wherein the epileptic disorder is Lennox-Gastaut syndrome.
69. (New) The method of claim 64, wherein the composition provides reduction in the frequency of seizures, the severity of seizures, or a combination thereof in a patient diagnosed with an epileptic disorder.

70. (New) The method of claim 64, wherein the patient is further administered an allosteric modulator.
71. (New) The method of claim 70, wherein the allosteric modulator is selected from the group consisting of a neurosteroid, a benzodiazepine and a potassium channel opener.
72. (New) The method of claim 70, wherein the allosteric modulator is ganaxolone.
73. (New) The method of claim 70, wherein the allosteric modulator is allopregnanolone.
74. (New) The method of claim 70, wherein the allosteric modulator is a benzodiazepine.
75. (New) The method of claim 74, wherein the benzodiazepine is selected from the group consisting of midazolam, clobazam, clonazepam, diazepam, lorazepam, flurazepam, and lorazepam.
76. (New) The method of claim 70, wherein the allosteric modulator is a potassium channel opener.
77. (New) The method of claim 76, wherein the potassium channel opener is retigabine or flupirtine.
78. (New) The method of claim 64, wherein the gaboxadol or a pharmaceutically acceptable salt thereof is administered once, twice, three times daily, or every other day.
79. (New) The method of claim 78, wherein the gaboxadol or a pharmaceutically acceptable salt thereof is administered immediately after the onset of a seizure.
80. (New) The method of claim 78, wherein the gaboxadol or a pharmaceutically acceptable salt thereof is administered after the onset of a seizure warning sign.

81. (New) The method of claim 64, further comprising administering a compound selected from the group consisting of acetazolamide, carbamazepine, eslicarbazepine acetate, ethosuximide, gabapentin, lacosamide, lamotrigine, leviteracetam, nitrazepam, oxcarbazepine, perampanel, piracetam, phenobarbital, phenytoin, pregabalin, primidone, rufinamide, sodium valproate, stiripentol, tiagabine, topiramate, vigabatrin, and zonisamide.
82. (New) The method of claim 64 wherein the pharmaceutical composition is a parenteral formulation.

Docket No.: 2262-29
(PATENT)

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Application of:

Matthew DURING

Confirmation No.: To Be Assigned

Application No.: To Be Assigned

Group Art Unit: To Be Assigned

Filed: Concurrently Herewith

Examiner: To Be Assigned

For: METHODS AND COMPOSITIONS FOR
TREATMENT OF EPILEPTIC DISORDERS

Dated: August 10, 2017

Commissioner for Patents
P.O. Box 1450
Alexandria, VA 22313-1450

Filed via EFS Web

Sir:

PREMINARY AMENDMENT

Prior to the examination of the above-identified patent application, please amend the application as follows:

Amendments to the Claims are reflected at page 2 of this submission.

Remarks begin on page 5 of this paper.

Electronic Patent Application Fee Transmittal				
Application Number:				
Filing Date:				
Title of Invention:		METHODS AND COMPOSITIONS FOR TREATMENT OF EPILEPTIC DISORDERS		
First Named Inventor/Applicant Name:		Matthew During		
Filer:		Jeffrey S. Steen/Julie Dascoli		
Attorney Docket Number:		2262-29		
Filed as Large Entity				
Filing Fees for Track I Prioritized Examination - Nonprovisional Application under 35 USC 111(a)				
Description	Fee Code	Quantity	Amount	Sub-Total in USD(\$)
Basic Filing:				
UTILITY APPLICATION FILING	1011	1	280	280
UTILITY SEARCH FEE	1111	1	600	600
UTILITY EXAMINATION FEE	1311	1	720	720
REQUEST FOR PRIORITIZED EXAMINATION	1817	1	4000	4000
Pages:				
Claims:				
Miscellaneous-Filing:				

Description	Fee Code	Quantity	Amount	Sub-Total in USD(\$)
PUBL. FEE- EARLY, VOLUNTARY, OR NORMAL	1504	1	0	0
PROCESSING FEE, EXCEPT PROV. APPLS.	1830	1	140	140
Petition:				
Patent-Appeals-and-Interference:				
Post-Allowance-and-Post-Issuance:				
Extension-of-Time:				
Miscellaneous:				
Total in USD (\$)				5740

Electronic Acknowledgement Receipt	
EFS ID:	30042761
Application Number:	15673737
International Application Number:	
Confirmation Number:	9025
Title of Invention:	METHODS AND COMPOSITIONS FOR TREATMENT OF EPILEPTIC DISORDERS
First Named Inventor/Applicant Name:	Matthew During
Customer Number:	31554
Filer:	Jeffrey S. Steen/Julie Dascoli
Filer Authorized By:	Jeffrey S. Steen
Attorney Docket Number:	2262-29
Receipt Date:	10-AUG-2017
Filing Date:	
Time Stamp:	14:05:32
Application Type:	Utility under 35 USC 111(a)

Payment information:

Submitted with Payment	yes
Payment Type	CARD
Payment was successfully received in RAM	\$5740
RAM confirmation Number	081117INTEFSW14105800
Deposit Account	
Authorized User	
The Director of the USPTO is hereby authorized to charge indicated fees and credit any overpayment as follows:	

File Listing:					
Document Number	Document Description	File Name	File Size(Bytes)/ Message Digest	Multi Part /.zip	Pages (if appl.)
1		POA.pdf	255123	yes	2
			839856926999091a0e1ca89296767c3c99d322f4		
	Multipart Description/PDF files in .zip description				
	Document Description		Start	End	
	Power of Attorney		2	2	
	Miscellaneous Incoming Letter		1	1	
Warnings:					
Information:					
2	Application Data Sheet	Application_Data_Sheet.pdf	1818971	no	7
			4b1971fb8ea82f5773d0f943ea2396c507ef5a63		
Warnings:					
Information:					
3	TrackOne Request	Cert_and_Request_for_Prioritized_Exam.pdf	140535	no	2
			44aec3588f98ecb08f125564428086ce60cdeb6		
Warnings:					
Information:					
4	Oath or Declaration filed	Executed_Declaration.pdf	62143	no	2
			354fe5ceae51a71f2d66e283ccb761f7d28d3495		
Warnings:					
The page size in the PDF is too large. The pages should be 8.5 x 11 or A4. If this PDF is submitted, the pages will be resized upon entry into the Image File Wrapper and may affect subsequent processing					
Information:					
5		Application_as_filed_810.pdf	3964086	yes	66
			cdad1854351887a0083212e2e0c991e7a41ba9e5		

	Multipart Description/PDF files in .zip description				
	Document Description		Start	End	
	Drawings-only black and white line drawings		63	66	
	Abstract		62	62	
	Claims		54	61	
	Specification		1	53	
Warnings:					
Information:					
6		Preliminary_Amendment.pdf	123645	yes	5
		97e524ec1d74ed70d1ff0af8ff23365d371dc115			
	Multipart Description/PDF files in .zip description				
	Document Description		Start	End	
	Applicant Arguments/Remarks Made in an Amendment		5	5	
	Claims		2	4	
	Preliminary Amendment		1	1	
Warnings:					
Information:					
7	Fee Worksheet (SB06)	fee-info.pdf	40225	no	2
		48b0c3d774310ae5a59fcc844ce986d8c3110fab			
Warnings:					
Information:					
Total Files Size (in bytes):			6404728		

This Acknowledgement Receipt evidences receipt on the noted date by the USPTO of the indicated documents, characterized by the applicant, and including page counts, where applicable. It serves as evidence of receipt similar to a Post Card, as described in MPEP 503.

New Applications Under 35 U.S.C. 111

If a new application is being filed and the application includes the necessary components for a filing date (see 37 CFR 1.53(b)-(d) and MPEP 506), a Filing Receipt (37 CFR 1.54) will be issued in due course and the date shown on this Acknowledgement Receipt will establish the filing date of the application.

National Stage of an International Application under 35 U.S.C. 371

If a timely submission to enter the national stage of an international application is compliant with the conditions of 35 U.S.C. 371 and other applicable requirements a Form PCT/DO/EO/903 indicating acceptance of the application as a national stage submission under 35 U.S.C. 371 will be issued in addition to the Filing Receipt, in due course.

New International Application Filed with the USPTO as a Receiving Office

If a new international application is being filed and the international application includes the necessary components for an international filing date (see PCT Article 11 and MPEP 1810), a Notification of the International Application Number and of the International Filing Date (Form PCT/RO/105) will be issued in due course, subject to prescriptions concerning national security, and the date shown on this Acknowledgement Receipt will establish the international filing date of the application.