



US008901378B2

(12) **United States Patent**
Barbour et al.(10) **Patent No.:** **US 8,901,378 B2**
(45) **Date of Patent:** ***Dec. 2, 2014**(54) **CORN EVENT TC1507 AND METHODS FOR DETECTION THEREOF**(75) Inventors: **Eric Barbour**, Johnston, IA (US);
James Wayne Bing, Ankeny, IA (US);
Guy A. Cardineau, Tempe, AZ (US);
Robert F. Cressman, Jr., Wilmington, DE (US); **Manju Gupta**, Carmel, IN (US); **Mary E. Hartnett Locke**, Mickleton, NJ (US); **David Hondred**, Ankeny, IA (US); **Joseph W. Keaschall**, Clive, IA (US); **Michael G. Koziel**, Raleigh, NC (US); **Terry EuClaire Meyer**, Urbandale, IA (US); **Daniel Moellenbeck**, Granger, IA (US); **Kenneth Edwin Narva**, Carlsbad, CA (US); **Wilas Nirunsuksiri**, Auburn, WA (US); **Stephen W. Ritchie**, Omaha, NE (US); **Marjorie L. Rudert**, Boone, IA (US); **Craig D. Sanders**, Bear, DE (US); **Aihua Shao**, Johnston, IA (US); **Steven Jeffrey Stelman**, San Diego, CA (US); **David S. Stucker**, Johnston, IA (US); **Laura Ann Tagliani**, Zionsville, IN (US); **William M. Van Zante**, Urbandale, IA (US)(73) Assignee: **Pioneer Hi-Bred International, Inc.**, Johnston, IA (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 1071 days.

This patent is subject to a terminal disclaimer.

(21) Appl. No.: **12/914,271**(22) Filed: **Oct. 28, 2010**(65) **Prior Publication Data**

US 2011/0099662 A1 Apr. 28, 2011

Related U.S. Application Data

(60) Division of application No. 12/333,044, filed on Mar. 13, 2009, now Pat. No. 7,989,607, which is a continuation of application No. 11/774,236, filed on Jul. 6, 2007, now Pat. No. 7,514,544, which is a division of application No. 10/837,105, filed on Apr. 30, 2004, now Pat. No. 7,288,643.

(60) Provisional application No. 60/467,772, filed on May 2, 2003.

(51) **Int. Cl.**
A01H 1/00 (2006.01)
C12N 15/82 (2006.01)
C12Q 1/68 (2006.01)(52) **U.S. Cl.**
CPC **C12N 15/8277** (2013.01); **C12Q 1/6895**(2013.01); **C12Q 1/686** (2013.01); **C12N 15/8286** (2013.01); **Y10S 435/975** (2013.01)USPC **800/302**; 800/265; 435/975(58) **Field of Classification Search**

None

See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

5,474,796	A	12/1995	Brennan	
5,877,012	A *	3/1999	Estruch et al.	435/252.3
6,166,302	A *	12/2000	Merlo et al.	800/320.1
6,646,184	B2	11/2003	Hohn et al.	
6,713,259	B2	3/2004	Levine	
6,825,400	B2	11/2004	Behr et al.	
7,288,643	B2 *	10/2007	Barbour et al.	536/23.1
7,417,132	B2 *	8/2008	Barbour et al.	536/23.1
7,435,807	B1 *	10/2008	Barbour et al.	536/23.1
7,449,564	B2 *	11/2008	Barbour et al.	536/23.1
7,514,544	B2 *	4/2009	Barbour et al.	536/23.1
7,989,607	B2 *	8/2011	Barbour et al.	536/23.1
2001/0053519	A1	12/2001	Fodor et al.	
2002/0177526	A1 *	11/2002	Chen et al.	504/100

FOREIGN PATENT DOCUMENTS

WO WO 2004/011601 2/2004

OTHER PUBLICATIONS

Cornejo et al 1993 Plant Molecular Biology 23: 567-581.*

Spencer et al 1992, Plant Molecular Biology 18: 201-210.*

Anonymous: "DAS-01507-1 (TC 1507)", Internet Article, Online Oct. 17, 2002, Internet: aqbios.com/dbase.php?action=showprod&date=TC1507&data=TC1507&format=long> 'retrieved on 1/412005.

Anonymous: "Skov—og Naturstyrelsen" Internet Article, Online 12111/2000, Denmark, Internet: sns.dk/erhvogadm/boeringer/n n1 00/indhold.htm> retried on Jan. 6, 2005.

Database EMBL 'Online! Nov. 18, 2002, "Mus musculus chromosome 5 clone rp23-426e16 strain C57BL16J, complete sequence." retried from EBI, Database accession No. AC084071.

New England Biolabs 1988/99 Catalog (NEB Catalog).

Printout from aqbios.com/docroot/decdocs/04-225-004.pdf.

Printout from epa.gov/fedrgstr/EPA-IMPACT/2001/August/Day-14/20307.htm.

Printout from aqbios.com/docroot/decdocs/02122001/pdf.


Stratagene ("Gene Characterization Kits" 1988).

* cited by examiner

Primary Examiner — David H Kruse(74) *Attorney, Agent, or Firm* — Ballard Spahr LLP(57) **ABSTRACT**

The invention provides DNA compositions that relate to transgenic insect resistant maize plants. Also provided are assays for detecting the presence of the maize TC1507 event based on the DNA sequence of the recombinant construct inserted into the maize genome and the DNA sequences flanking the insertion site. Kits and conditions useful in conducting the assays are provided.

8 Claims, 1 Drawing SheetInari Exhibit 1001
Inari v. Corteva



This bottom diagram shows where primers are located by SID number. The green is flanking, the grey is insert. The red stars are rough (not to scale) locations of primers and the number is the SID.

1

CORN EVENT TC1507 AND METHODS FOR DETECTION THEREOF

CROSS-REFERENCE TO RELATED APPLICATIONS

This application is a divisional of U.S. Ser. No. 12/333,044, filed Mar. 13, 2007, which is a continuation of U.S. Ser. No. 11/774,236 filed Jul. 6, 2007, which is a divisional of 10/837,105, filed Apr. 30, 2004, which claims the benefit of U.S. Provisional Application Ser. No. 60/467,772, filed May 2, 2003, the contents of all of which are herein incorporated by reference in their entirety.

FIELD OF THE INVENTION

The present invention relates to the field of plant molecular biology, specifically the invention relates to a DNA construct for conferring insect resistance to a plant. The invention more specifically relates to an insect resistant corn plant TC1507 and to assays for detecting the presence of corn plant TC1507 DNA in a sample and compositions thereof.

BACKGROUND OF THE INVENTION

This invention relates to the insect resistant corn (*Zea mays*) plant TC1507, also referred to as maize line TC1507 or maize event TC1507, and to the DNA plant expression construct of corn plant TC1507 and the detection of the transgene/flanking insertion region in corn plant TC1507 and progeny thereof.

Corn is an important crop and is a primary food source in many areas of the world. Damage caused by insect pests is a major factor in the loss of the world's corn crops, despite the use of protective measures such as chemical pesticides. In view of this, insect resistance has been genetically engineered into crops such as corn in order to control insect damage and to reduce the need for traditional chemical pesticides. One group of genes which have been utilized for the production of transgenic insect resistant crops are the delta-endotoxins from *Bacillus thuringiensis* (B.t.). Delta-endotoxins have been successfully expressed in crop plants such as cotton, potatoes, rice, sunflower, as well as corn, and have proven to provide excellent control over insect pests. (Perlak, F. J et al. (1990) *Bio/Technology* 8, 939-943; Perlak, F. J. et al. (1993) *Plant Mol. Biol.* 22: 313-321; Fujimoto H. et al. (1993) *Bio/Technology* 11: 1151-1155; Tu et al. (2000) *Nature Biotechnology* 18:1101-1104; PCT publication number WO 01/13731; and Bing J W et al. (2000) Efficacy of Cry1F Transgenic Maize, 14th Biennial International Plant Resistance to Insects Workshop, Fort Collins, Colo.).

The expression of foreign genes in plants is known to be influenced by their location in the plant genome, perhaps due to chromatin structure (e.g., heterochromatin) or the proximity of transcriptional regulatory elements (e.g., enhancers) close to the integration site (Weising et al., *Ann. Rev. Genet* 22:421-477, 1988). At the same time the presence of the transgene at different locations in the genome will influence the overall phenotype of the plant in different ways. For this reason, it is often necessary to screen a large number of events in order to identify an event characterized by optimal expression of an introduced gene of interest. For example, it has been observed in plants and in other organisms that there may be a wide variation in levels of expression of an introduced gene among events. There may also be differences in spatial or temporal patterns of expression, for example, differences in the relative expression of a transgene in various plant

2

tissues, that may not correspond to the patterns expected from transcriptional regulatory elements present in the introduced gene construct. For this reason, it is common to produce hundreds to thousands of different events and screen those events for a single event that has desired transgene expression levels and patterns for commercial purposes. An event that has desired levels or patterns of transgene expression is useful for introgressing the transgene into other genetic backgrounds by sexual outcrossing using conventional breeding methods. Progeny of such crosses maintain the transgene expression characteristics of the original transformant. This strategy is used to ensure reliable gene expression in a number of varieties that are well adapted to local growing conditions.

It would be advantageous to be able to detect the presence of a particular event in order to determine whether progeny of a sexual cross contain a transgene of interest. In addition, a method for detecting a particular event would be helpful for complying with regulations requiring the pre-market approval and labeling of foods derived from recombinant crop plants, for example, or for use in environmental monitoring, monitoring traits in crops in the field, or monitoring products derived from a crop harvest, as well as for use in ensuring compliance of parties subject to regulatory or contractual terms.

It is possible to detect the presence of a transgene by any nucleic acid detection method known in the art including, but not limited to, the polymerase chain reaction (PCR) or DNA hybridization using nucleic acid probes. These detection methods generally focus on frequently used genetic elements, such as promoters, terminators, marker genes, etc., because for many DNA constructs, the coding region is interchangeable. As a result, such methods may not be useful for discriminating between different events, particularly those produced using the same DNA construct or very similar constructs unless the DNA sequence of the flanking DNA adjacent to the inserted heterologous DNA is known. For example, an event-specific PCR assay is described in U.S. Pat. No. 6,395,485 for the detection of elite event GAT-ZM1. Accordingly, it would be desirable to have a simple and discriminative method for the identification of event TC1507.

SUMMARY OF THE INVENTION

This invention relates preferably to methods for producing and selecting an insect resistant monocot crop plant. More specifically, a DNA construct is provided that when expressed in plant cells and plants confers resistance to insects. According to one aspect of the invention, a DNA construct, capable of introduction into and replication in a host cell, is provided that when expressed in plant cells and plants confers insect resistance to the plant cells and plants. The DNA construct is comprised of a DNA molecule named PHI8999A and it includes two transgene expression cassettes. The first expression cassette comprises a DNA molecule which includes the promoter, 5' untranslated exon, and first intron of the maize ubiquitin (Ubi-1) gene (Christensen et al. (1992) *Plant Mol. Biol.* 18:675-689 and Christensen and Quail (1996) *Transgenic Res.* 5:213-218) operably connected to a DNA molecule encoding a B.t. δ -endotoxin identified as Cry1F (U.S. Pat. Nos. 5,188,960 and 6,218,188) operably connected to a DNA molecule comprising a 3' ORF25 transcriptional terminator isolated from *Agrobacterium tumefaciens* (Barker et al. (1983) *Plant Mol. Biol.* 2:335-350). The second transgene expression cassette of the DNA construct comprises a DNA molecule of the cauliflower mosaic virus (CaMV) 35S promoter (Odell J. T. et al. (1985) *Nature* 313: 810-812; Mitsu-hara et al. (1996) *Plant Cell Physiol.* 37: 49-59) operably

3

connected to a DNA molecule encoding a phosphinothricin acetyltransferase (PAT) gene (Wohleben W. et al. (1988) *Gene* 70: 25-37) operably connected to a DNA molecule comprising a 3' transcriptional terminator from (CaMV) 35S (see Mitsuhashi et al. (1996) *Plant Cell Physiol.* 37: 49-59). Plants containing the DNA construct are also provided.

According to another aspect of the invention, compositions and methods are provided for identifying a novel corn plant designated TC1507, which methods are based on primers or probes which specifically recognize the 5' and/or 3' flanking sequence of TC1507. DNA molecules are provided that comprise primer sequences that when utilized in a PCR reaction will produce amplicons unique to the transgenic event TC1507. These molecules may be selected from the group consisting of:

(SEQ ID NO: 1)
5' -GTAGTACTATAGATTATATTATTCGTAGAG-3' ;

(SEQ ID NO: 2)
5' -GCCATACAGAAGCTCAAAATCTTTTCCGGAG-3' ;

(SEQ ID NO: 23)
5' -CTTCAACAAGTGTGACAAA-3' ;

(SEQ ID NO: 3)
5' -TGTGGTGTGTTGTGGCTCTGTCTCTAA-3' ;

(SEQ ID NO: 4)
5' -AGCACCTTTTTCATCTTTTCATATAC-3' ;

(SEQ ID NO: 5)
5' -GACCTCCCCA CAGGCATGAT TGATC-3' ;

and complements thereof. The corn plant and seed comprising these molecules is an aspect of this invention. Further, kits utilizing these primer sequences for the identification of the TC1507 event are provided.

An additional aspect of the invention relates to the specific flanking sequences of TC1507 described herein, which can be used to develop specific identification methods for TC1507 in biological samples. More particularly, the invention relates to the 5' and/or 3' flanking regions of TC1507, SEQ ID NO:21 and SEQ ID NO:22, respectively, which can be used for the development of specific primers and probes. The invention further relates to identification methods for the presence of TC1507 in biological samples based on the use of such specific primers or probes.

According to another aspect of the invention, methods of detecting the presence of DNA corresponding to the corn event TC1507 in a sample are provided. Such methods comprise: (a) contacting the sample comprising DNA with a DNA primer set, that when used in a nucleic acid amplification reaction with genomic DNA extracted from corn event TC1507 produces an amplicon that is diagnostic for corn event TC1507; (b) performing a nucleic acid amplification reaction, thereby producing the amplicon; and (c) detecting the amplicon.

DNA molecules that comprise the novel transgene/flanking insertion region, SEQ ID NO: 26 and SEQ ID NO: 27 and are homologous or complementary to SEQ ID NO: 26 and SEQ ID NO: 27 are an aspect of this invention.

DNA sequences that comprise the novel transgene/flanking insertion region, SEQ ID NO:26 are an aspect of this invention. DNA sequences that comprise a sufficient length of polynucleotides of transgene insert sequence and a sufficient length of polynucleotides of maize genomic and/or flanking sequence from maize plant TC1507 of SEQ ID NO:26 that are

4

useful as primer sequences for the production of an amplicon product diagnostic for maize plant TC1507 are included.

In addition, DNA sequences that comprise the novel transgene/flanking insertion region, SEQ ID NO:27 are provided. DNA sequences that comprise a sufficient length of polynucleotides of transgene insert sequence and a sufficient length of polynucleotides of maize genomic and/or flanking sequence from maize plant TC1507 of SEQ ID NO:27 that are useful as primer sequences for the production of an amplicon product diagnostic for maize plant TC1507 are included.

According to another aspect of the invention, the DNA sequences that comprise at least 11 or more nucleotides of the transgene portion of the DNA sequence of SEQ ID NO:26 or complements thereof, and a similar length of 5' flanking maize DNA sequence of SEQ ID NO:26 or complements thereof are useful as DNA primers in DNA amplification methods. The amplicons produced using these primers are diagnostic for maize event TC1507. Therefore, the invention also includes the amplicons produced by DNA primers homologous or complementary to SEQ ID NO:26.

According to another aspect of the invention, the DNA sequences that comprise at least 11 or more nucleotides of the transgene portion of the DNA sequence of SEQ ID NO:27 or complements thereof, and a similar length of 3' flanking maize DNA sequence of SEQ ID NO:27 or complements thereof are useful as DNA primers in DNA amplification methods. The amplicons produced using these primers are diagnostic for maize event TC1507. Therefore, the invention also includes the amplicons produced by DNA primers homologous or complementary to SEQ ID NO:27.

More specifically, a pair of DNA molecules comprising a DNA primer set, wherein the DNA molecules are identified as SEQ ID NO: 1 or complements thereof and SEQ ID NO: 2 or complements thereof; SEQ ID NO: 2 or complements thereof and SEQ ID NO: 23 or complements thereof; SEQ ID NO: 3 or complements thereof and SEQ ID NO: 5 or complements thereof; SEQ ID NO: 4 or complements thereof and SEQ ID NO: 5 or complements thereof are aspects of the invention.

Further aspects of the invention include the amplicon comprising the DNA molecules of SEQ ID NO: 1 and SEQ ID NO: 2; the amplicon comprising the DNA molecules of SEQ ID NO: 2 and SEQ ID NO: 23; the amplicon comprising the DNA molecules of SEQ ID NO: 3 and SEQ ID NO: 5; and the amplicon comprising the DNA molecules of SEQ ID NO: 4 and SEQ ID NO: 5.

According to another aspect of the invention, methods of detecting the presence of a DNA molecule corresponding to the TC1507 event in a sample, such methods comprising: (a) contacting the sample comprising DNA extracted from a corn plant with a DNA probe, molecule that hybridizes under stringent hybridization conditions with DNA extracted from corn event TC1507 and does not hybridize under the stringent hybridization conditions with a control corn plant DNA; (b) subjecting the sample and probe to stringent hybridization conditions; and (c) detecting hybridization of the probe to the DNA. More specifically, a method for detecting the presence of a DNA molecule corresponding to the TC1507 event in a sample, such methods, consisting of (a) contacting the sample comprising DNA extracted from a corn plant with a DNA probe molecule that consists of sequences that are unique to the event, e.g. junction sequences, wherein said DNA probe molecule hybridizes under stringent hybridization conditions with DNA extracted from corn event TC1507 and does not hybridize under the stringent hybridization conditions with a control corn plant DNA; (b) subjecting the sample and probe to stringent hybridization conditions; and (c) detecting hybridization of the probe to the DNA.

In addition, a kit and methods for identifying event TC1507 in a biological sample which detects a TC1507 specific region within SEQ ID NO: 24 are provided.

DNA molecules are provided that comprise at least one junction sequence of TC1507 selected from the group consisting of SEQ ID NO: 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56 and 57 and complements thereof; wherein a junction sequence spans the junction between heterologous DNA inserted into the genome and the DNA from the corn cell flanking the insertion site, i.e. flanking DNA, and is diagnostic for the TC1507 event.

According to another aspect of the invention, methods of producing an insect resistant corn plant that comprise the steps of: (a) sexually crossing a first parental corn line comprising the expression cassettes of the present invention, which confers resistance to insects, and a second parental corn line that lacks insect resistance, thereby producing a plurality of progeny plants; and (b) selecting a progeny plant that is insect resistant. Such methods may optionally comprise the further step of back-crossing the progeny plant to the second parental corn line to producing a true-breeding corn plant that is insect resistant.

The present invention provides a method of producing a corn plant that is resistant to insects comprising transforming a corn cell with the DNA construct PHI8999A (SEQ ID NO: 25), growing the transformed corn cell into a corn plant, selecting the corn plant that shows resistance to insects, and further growing the corn plant into a fertile corn plant. The fertile corn plant can be self pollinated or crossed with compatible corn varieties to produce insect resistant progeny.

The invention further relates to a DNA detection kit for identifying maize event TC1507 in biological samples. Preferably the kit of the invention comprises a first primer which specifically recognizes the 5' or 3' flanking region of TC1507, and a second primer which specifically recognizes a sequence within the foreign DNA of TC1507, or within the flanking DNA, for use in a PCR identification protocol. The invention also relates to a kit for identifying event TC1507 in biological samples, which kit comprises a specific probe having a sequence which corresponds or is complementary to, a sequence having between 80% and 100% sequence identity with a specific region of event TC1507. Preferably the sequence of the probe corresponds to a specific region comprising part of the 5' or 3' flanking region of event TC1507.

The methods and kits encompassed by the present invention can be used for different purposes such as, but not limited to the following: to identify event TC1507 in plants, plant material or in products such as, but not limited to, food or feed products (fresh or processed) comprising, or derived from plant material; additionally or alternatively, the methods and kits of the present invention can be used to identify transgenic plant material for purposes of segregation between transgenic and non-transgenic material; additionally or alternatively, the methods and kits of the present invention can be used to determine the quality of plant material comprising maize event TC1507. The kits may also contain the reagents and materials necessary for the performance of the detection method.

This invention further relates to the TC1507 corn plant or its parts, including, but not limited to, pollen, ovules, vegetative cells, the nuclei of pollen cells, and the nuclei of egg cells of the corn plant TC1507 and the progeny derived thereof. The corn plant and seed TC1507 from which the DNA primer molecules of the present invention provide a specific amplification product is an aspect of the invention.

The foregoing and other aspects of the invention will become more apparent from the following detailed description and accompanying drawing.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1. Linear map showing the transgenic insert PHI8999A, as well as the sequences flanking the transgenic insert. Note: FIG. 1 is not necessarily to scale.

DETAILED DESCRIPTION

The following definitions and methods are provided to better define the present invention and to guide those of ordinary skill in the art in the practice of the present invention. Unless otherwise noted, terms are to be understood according to conventional usage by those of ordinary skill in the relevant art. Definitions of common terms in molecular biology may also be found in Rieger et al., *Glossary of Genetics: Classical and Molecular*, 5th edition, Springer-Verlag; New York, 1991; and Lewin, *Genes V*, Oxford University Press: New York, 1994. The nomenclature for DNA bases as set forth at 37 CFR 1.822 is used.

As used herein, the term "comprising" means "including but not limited to".

As used herein, the term "corn" means *Zea mays* or maize and includes all plant varieties that can be bred with corn, including wild maize species.

As used herein, the term "TC1507 specific" refers to a nucleotide sequence which is suitable for discriminatively identifying event TC1507 in plants, plant material, or in products such as, but not limited to, food or feed products (fresh or processed) comprising, or derived from plant material.

As used herein, the terms "insect resistant" and "impacting insect pests" refers to effecting changes in insect feeding, growth, and/or behavior at any stage of development, including but not limited to: killing the insect; retarding growth; preventing reproductive capability; and the like.

As used herein, the terms "pesticidal activity" and "insecticidal activity" are used synonymously to refer to activity of an organism or a substance (such as, for example, a protein) that can be measured by numerous parameters including, but not limited to, pest mortality, pest weight loss, pest attraction, pest repellency, and other behavioral and physical changes of a pest after feeding on and/or exposure to the organism or substance for an appropriate length of time. For example "pesticidal proteins" are proteins that display pesticidal activity by themselves or in combination with other proteins.

"Coding sequence" refers to a nucleotide sequence that codes for a specific amino acid sequence. As used herein, the terms "encoding" or "encoded" when used in the context of a specified nucleic acid mean that the nucleic acid comprises the requisite information to guide translation of the nucleotide sequence into a specified protein. The information by which a protein is encoded is specified by the use of codons. A nucleic acid encoding a protein may comprise non-translated sequences (e.g., introns) within translated regions of the nucleic acid or may lack such intervening non-translated sequences (e.g., as in cDNA).

"Gene" refers to a nucleic acid fragment that expresses a specific protein, including regulatory sequences preceding (5' non-coding sequences) and following (3' non-coding sequences) the coding sequence. "Native gene" refers to a gene as found in nature with its own regulatory sequences. "Chimeric gene" refers any gene that is not a native gene, comprising regulatory and coding sequences that are not found together in nature. Accordingly, a chimeric gene may

comprise regulatory sequences and coding sequences that are derived from different sources, or regulatory sequences and coding sequences derived from the same source, but arranged in a manner different than that found in nature. "Endogenous gene" refers to a native gene in its natural location in the genome of an organism. "Foreign" refers to material not normally found in the location of interest. Thus "foreign DNA" may comprise both recombinant DNA as well as newly introduced, rearranged DNA of the plant. A "foreign" gene refers to a gene not normally found in the host organism, but that is introduced into the host organism by gene transfer. Foreign genes can comprise native genes inserted into a non-native organism, or chimeric genes. A "transgene" is a gene that has been introduced into the genome by a transformation procedure. The site in the plant genome where a recombinant DNA has been inserted may be referred to as the "insertion site" or "target site".

As used herein, "insert DNA" refers to the heterologous DNA within the expression cassettes used to transform the plant material while "flanking DNA" can exist of either genomic DNA naturally present in an organism such as a plant, or foreign (heterologous) DNA introduced via the transformation process which is extraneous to the original insert DNA molecule, e.g. fragments associated with the transformation event. A "flanking region" or "flanking sequence" as used herein refers to a sequence of at least 20 base pair, preferably at least 50 base pair, and up to 5000 base pair which is located either immediately upstream of and contiguous with or immediately downstream of and contiguous with the original foreign insert DNA molecule. Transformation procedures leading to random integration of the foreign DNA will result in transformants containing different flanking regions characteristic and unique for each transformant. When recombinant DNA is introduced into a plant through traditional crossing, its flanking regions will generally not be changed. Transformants will also contain unique junctions between a piece of heterologous insert DNA and genomic DNA, or 2 pieces of genomic DNA, or 2 pieces of heterologous DNA. A "junction" is a point where 2 specific DNA fragments join. For example, a junction exists where insert DNA joins flanking DNA. A junction point also exists in a transformed organism where 2 DNA fragments join together in a manner that is modified from that found in the native organism. "Junction DNA" refers to DNA that comprises a junction point.

As used herein, "heterologous" in reference to a nucleic acid is a nucleic acid that originates from a foreign species, or, if from the same species, is substantially modified from its native form in composition and/or genomic locus by deliberate human intervention. For example, a promoter operably linked to a heterologous nucleotide sequence can be from a species different from that from which the nucleotide sequence was derived, or, if from the same species, the promoter is not naturally found operably linked to the nucleotide sequence. A heterologous protein may originate from a foreign species, or, if from the same species, is substantially modified from its original form by deliberate human intervention.

"Regulatory sequences" refer to nucleotide sequences located upstream (5' non-coding sequences), within, or downstream (3' non-coding sequences) of a coding sequence, and which influence the transcription, RNA processing or stability, or translation of the associated coding sequence. Regulatory sequences may include promoters, translation leader sequences, introns, and polyadenylation recognition sequences.

"Promoter" refers to a nucleotide sequence capable of controlling the expression of a coding sequence or functional RNA. In general, a coding sequence is located 3' to a promoter sequence. The promoter sequence consists of proximal and more distal upstream elements, the latter elements are often referred to as enhancers. Accordingly, an "enhancer" is a nucleotide sequence that can stimulate promoter activity and may be an innate element of the promoter or a heterologous element inserted to enhance the level or tissue-specificity of a promoter. Promoters may be derived in their entirety from a native gene, or be composed of different elements derived from different promoters found in nature, or even comprise synthetic nucleotide segments. It is understood by those skilled in the art that different promoters may direct the expression of a gene in different tissues or cell types, or at different stages of development, or in response to different environmental conditions. Promoters that cause a nucleic acid fragment to be expressed in most cell types at most times are commonly referred to as "constitutive promoters". New promoters of various types useful in plant cells are constantly being discovered; numerous examples may be found in the compilation by Okamuro and Goldberg (1989) *Biochemistry of Plants* 15:1-82. It is further recognized that since in most cases the exact boundaries of regulatory sequences have not been completely defined, nucleic acid fragments of different lengths may have identical promoter activity.

The "translation leader sequence" refers to a nucleotide sequence located between the promoter sequence of a gene and the coding sequence. The translation leader sequence is present in the fully processed mRNA upstream of the translation start sequence. The translation leader sequence may affect numerous parameters including, processing of the primary transcript to mRNA, mRNA stability and/or translation efficiency. Examples of translation leader sequences have been described (Turner and Foster (1995) *Mol. Biotechnol.* 3:225-236).

The "3' non-coding sequences" refer to nucleotide sequences located downstream of a coding sequence and include polyadenylation recognition sequences and other sequences encoding regulatory signals capable of affecting mRNA processing or gene expression. The polyadenylation signal is usually characterized by affecting the addition of polyadenylic acid tracts to the 3' end of the mRNA precursor. The use of different 3' non-coding sequences is exemplified by Ingelbrecht et al. (1989) *Plant Cell* 1:671-680.

A "protein" or "polypeptide" is a chain of amino acids arranged in a specific order determined by the coding sequence in a polynucleotide encoding the polypeptide.

A DNA construct is an assembly of DNA molecules linked together that provide one or more expression cassettes. The DNA construct may be a plasmid that is enabled for self replication in a bacterial cell and contains various endonuclease enzyme restriction sites that are useful for introducing DNA molecules that provide functional genetic elements, i.e., promoters, introns, leaders, coding sequences, 3' termination regions, among others; or a DNA construct may be a linear assembly of DNA molecules, such as an expression cassette. The expression cassette contained within a DNA construct comprise the necessary genetic elements to provide transcription of a messenger RNA. The expression cassette can be designed to express in prokaryote cells or eukaryotic cells. Expression cassettes of the present invention are designed to express most preferably in plant cells.

The DNA molecules of the invention are provided in expression cassettes for expression in an organism of interest. The cassette will include 5' and 3' regulatory sequences operably linked to a coding sequence of the invention. "Operably

linked” means that the nucleic acid sequences being linked are contiguous and, where necessary to join two protein coding regions, contiguous and in the same reading frame. Operably linked is intended to indicate a functional linkage between a promoter and a second sequence, wherein the promoter sequence initiates and mediates transcription of the DNA sequence corresponding to the second sequence. The cassette may additionally contain at least one additional gene to be cotransformed into the organism. Alternatively, the additional gene(s) can be provided on multiple expression cassettes or multiple DNA constructs.

The expression cassette will include in the 5' to 3' direction of transcription: a transcriptional and translational initiation region, a coding region, and a transcriptional and translational termination region functional in the organism serving as a host. The transcriptional initiation region (i.e., the promoter) may be native or analogous, or foreign or heterologous to the host organism. Additionally, the promoter may be the natural sequence or alternatively a synthetic sequence. The expression cassettes may additionally contain 5' leader sequences in the expression cassette construct. Such leader sequences can act to enhance translation.

It is to be understood that as used herein the term “transgenic” includes any cell, cell line, callus, tissue, plant part, or plant the genotype of which has been altered by the presence of a heterologous nucleic acid including those transgenics initially so altered as well as those created by sexual crosses or asexual propagation from the initial transgenic. The term “transgenic” as used herein does not encompass the alteration of the genome (chromosomal or extra-chromosomal) by conventional plant breeding methods or by naturally occurring events such as random cross-fertilization, non-recombinant viral infection, non-recombinant bacterial transformation, non-recombinant transposition, or spontaneous mutation.

A transgenic “event” is produced by transformation of plant cells with a heterologous DNA construct(s), including a nucleic acid expression cassette that comprises a transgene of interest, the regeneration of a population of plants resulting from the insertion of the transgene into the genome of the plant, and selection of a particular plant characterized by insertion into a particular genome location. An event is characterized phenotypically by the expression of the transgene. At the genetic level, an event is part of the genetic makeup of a plant. The term “event” also refers to progeny produced by a sexual outcross between the transformant and another variety that include the heterologous DNA. Even after repeated back-crossing to a recurrent parent, the inserted DNA and flanking DNA from the transformed parent is present in the progeny of the cross at the same chromosomal location. The term “event” also refers to DNA from the original transformant comprising the inserted DNA and flanking sequence immediately adjacent to the inserted DNA that would be expected to be transferred to a progeny that receives inserted DNA including the transgene of interest as the result of a sexual cross of one parental line that includes the inserted DNA (e.g., the original transformant and progeny resulting from selfing) and a parental line that does not contain the inserted DNA.

An insect resistant TC1507 corn plant can be bred by first sexually crossing a first parental corn plant consisting of a corn plant grown from the transgenic TC1507 corn plant and progeny thereof derived from transformation with the expression cassettes of the present invention that confers insect resistance, and a second parental corn plant that lacks insect resistance, thereby producing a plurality of first progeny plants; and then selecting a first progeny plant that is resistant to insects; and selfing the first progeny plant, thereby produc-

ing a plurality of second progeny plants; and then selecting from the second progeny plants an insect resistant plant. These steps can further include the back-crossing of the first insect resistant progeny plant or the second insect resistant progeny plant to the second parental corn plant or a third parental corn plant, thereby producing a corn plant that is resistant to insects.

As used herein, the term “plant” includes reference to whole plants, plant organs (e.g., leaves, stems, roots, etc.), seeds, plant cells, and progeny of same. Parts of transgenic plants understood to be within the scope of the invention comprise, for example, plant cells, protoplasts, tissues, callus, embryos as well as flowers, stems, fruits, leaves, and roots originating in transgenic plants or their progeny previously transformed with a DNA molecule of the invention and therefore consisting at least in part of transgenic cells, are also an aspect of the present invention.

As used herein, the term “plant cell” includes, without limitation, seeds, suspension cultures, embryos, meristematic regions, callus tissue, leaves, roots, shoots, gametophytes, sporophytes, pollen, and microspores. The class of plants that can be used in the methods of the invention is generally as broad as the class of higher plants amenable to transformation techniques, including both monocotyledonous and dicotyledonous plants.

“Transformation” refers to the transfer of a nucleic acid fragment into the genome of a host organism, resulting in genetically stable inheritance. Host organisms containing the transformed nucleic acid fragments are referred to as “transgenic” organisms. Examples of methods of plant transformation include *Agrobacterium*-mediated transformation (De Blaere et al. (1987) *Meth. Enzymol.* 143:277) and particle-accelerated or “gene gun” transformation technology (Klein et al. (1987) *Nature* (London) 327:70-73; U.S. Pat. No. 4,945,050, incorporated herein by reference). Additional transformation methods are disclosed below.

Thus, isolated polynucleotides of the present invention can be incorporated into recombinant constructs, typically DNA constructs, which are capable of introduction into and replication in a host cell. Such a construct can be a vector that includes a replication system and sequences that are capable of transcription and translation of a polypeptide-encoding sequence in a given host cell. A number of vectors suitable for stable transfection of plant cells or for the establishment of transgenic plants have been described in, e.g., Pouwels et al., (1985; Supp. 1987) *Cloning Vectors: A Laboratory Manual*, Weissbach and Weissbach (1989) *Methods for Plant Molecular Biology*, (Academic Press, New York); and Flevin et al., (1990) *Plant Molecular Biology Manual*, (Kluwer Academic Publishers). Typically, plant expression vectors include, for example, one or more cloned plant genes under the transcriptional control of 5' and 3' regulatory sequences and a dominant selectable marker. Such plant expression vectors also can contain a promoter regulatory region (e.g., a regulatory region controlling inducible or constitutive, environmentally- or developmentally-regulated, or cell- or tissue-specific expression), a transcription initiation start site, a ribosome binding site, an RNA processing signal, a transcription termination site, and/or a polyadenylation signal.

It is also to be understood that two different transgenic plants can also be mated to produce offspring that contain two independently segregating added, exogenous genes. Selfing of appropriate progeny can produce plants that are homozygous for both added, exogenous genes. Back-crossing to a parental plant and out-crossing with a non-transgenic plant are also contemplated, as is vegetative propagation. Descriptions of other breeding methods that are commonly used for

different traits and crops can be found in one of several references, e.g., Fehr, in *Breeding Methods for Cultivar Development*, Wilcos J. ed., American Society of Agronomy, Madison Wis. (1987).

A "probe" is an isolated nucleic acid to which is attached a conventional detectable label or reporter molecule, e.g., a radioactive isotope, ligand, chemiluminescent agent, or enzyme. Such a probe is complementary to a strand of a target nucleic acid, in the case of the present invention, to a strand of isolated DNA from corn event TC1507 whether from a corn plant or from a sample that includes DNA from the event. Probes according to the present invention include not only deoxyribonucleic or ribonucleic acids but also polyamides and other probe materials that bind specifically to a target DNA sequence and can be used to detect the presence of that target DNA sequence.

"Primers" are isolated nucleic acids that are annealed to a complementary target DNA strand by nucleic acid hybridization to form a hybrid between the primer and the target DNA strand, then extended along the target DNA strand by a polymerase, e.g., a DNA polymerase. Primer pairs of the present invention refer to their use for amplification of a target nucleic acid sequence, e.g., by the polymerase chain reaction (PCR) or other conventional nucleic-acid amplification methods. "PCR" or "polymerase chain reaction" is a technique used for the amplification of specific DNA segments (see, U.S. Pat. Nos. 4,683,195 and 4,800,159; herein incorporated by reference).

Probes and primers are of sufficient nucleotide length to bind to the target DNA sequence specifically in the hybridization conditions or reaction conditions determined by the operator. This length may be of any length that is of sufficient length to be useful in a detection method of choice. Generally, 11 nucleotides or more in length, preferably 18 nucleotides or more, and more preferably 22 nucleotides or more, are used. Such probes and primers hybridize specifically to a target sequence under high stringency hybridization conditions. Preferably, probes and primers according to the present invention have complete DNA sequence similarity of contiguous nucleotides with the target sequence, although probes differing from the target DNA sequence and that retain the ability to hybridize to target DNA sequences may be designed by conventional methods. Probes can be used as primers, but are generally designed to bind to the target DNA or RNA and not be used in an amplification process.

Specific primers can be used to amplify an integration fragment to produce an amplicon that can be used as a "specific probe" for identifying event TC1507 in biological samples. When the probe is hybridized with the nucleic acids of a biological sample under conditions which allow for the binding of the probe to the sample, this binding can be detected and thus allow for an indication of the presence of event TC1507 in the biological sample. Such identification of a bound probe has been described in the art. The specific probe is preferably a sequence which, under optimized conditions, hybridizes specifically to a region within the 5' or 3' flanking region of the event and preferably also comprises a part of the foreign DNA contiguous therewith. Preferably the specific probe comprises a sequence of at least 80%, preferably between 80 and 85%, more preferably between 85 and 90%, especially preferably between 90 and 95%, and most preferably between 95 and 100% identical (or complementary) to a specific region of the event.

Methods for preparing and using probes and primers are described, for example, in *Molecular Cloning: A Laboratory Manual*, 2nd ed., vol. 1-3, ed. Sambrook et al., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. 1989

(hereinafter, "Sambrook et al., 1989"); *Current Protocols in Molecular Biology*, ed. Ausubel et al., Greene Publishing and Wiley-Interscience, New York, 1992 (with periodic updates) (hereinafter, "Ausubel et al., 1992"); and Innis et al., *PCR Protocols: A Guide to Methods and Applications*, Academic Press: San Diego, 1990. PCR primer pairs can be derived from a known sequence, for example, by using computer programs intended for that purpose such as the PCR primer analysis tool in Vector NTI version 6 (Informax Inc., Bethesda Md.); PrimerSelect (DNASTAR Inc., Madison, Wis.); and Primer (Version 0.5, © 1991, Whitehead Institute for Biomedical Research, Cambridge, Mass.). Additionally, the sequence can be visually scanned and primers manually identified using guidelines known to one of skill in the art.

A "kit" as used herein refers to a set of reagents for the purpose of performing the method of the invention, more particularly, the identification of the event TC1507 in biological samples. The kit of the invention can be used, and its components can be specifically adjusted, for purposes of quality control (e.g. purity of seed lots), detection of event TC1507 in plant material, or material comprising or derived from plant material, such as but not limited to food or feed products. "Plant material" as used herein refers to material which is obtained or derived from a plant.

Primers and probes based on the flanking DNA and insert sequences disclosed herein can be used to confirm (and, if necessary, to correct) the disclosed sequences by conventional methods, e.g., by re-cloning and sequencing such sequences. The nucleic acid probes and primers of the present invention hybridize under stringent conditions to a target DNA sequence. Any conventional nucleic acid hybridization or amplification method can be used to identify the presence of DNA from a transgenic event in a sample. Nucleic acid molecules or fragments thereof are capable of specifically hybridizing to other nucleic acid molecules under certain circumstances. As used herein, two nucleic acid molecules are said to be capable of specifically hybridizing to one another if the two molecules are capable of forming an anti-parallel, double-stranded nucleic acid structure.

A nucleic acid molecule is said to be the "complement" of another nucleic acid molecule if they exhibit complete complementarity. As used herein, molecules are said to exhibit "complete complementarity" when every nucleotide of one of the molecules is complementary to a nucleotide of the other. Two molecules are said to be "minimally complementary" if they can hybridize to one another with sufficient stability to permit them to remain annealed to one another under at least conventional "low-stringency" conditions. Similarly, the molecules are said to be "complementary" if they can hybridize to one another with sufficient stability to permit them to remain annealed to one another under conventional "high-stringency" conditions. Conventional stringency conditions are described by Sambrook et al., 1989, and by Haymes et al., In: *Nucleic Acid Hybridization, a Practical Approach*, IRL Press, Washington, D.C. (1985), departures from complete complementarity are therefore permissible, as long as such departures do not completely preclude the capacity of the molecules to form a double-stranded structure. In order for a nucleic acid molecule to serve as a primer or probe it need only be sufficiently complementary in sequence to be able to form a stable double-stranded structure under the particular solvent and salt concentrations employed.

In hybridization reactions, specificity is typically the function of post-hybridization washes, the critical factors being the ionic strength and temperature of the final wash solution. The thermal melting point (T_m) is the temperature (under defined ionic strength and pH) at which 50% of a comple-

mentary target sequence hybridizes to a perfectly matched probe. For DNA-DNA hybrids, the T_m can be approximated from the equation of Meinkoth and Wahl (1984) *Anal. Biochem.* 138:267-284: $T_m = 81.5^\circ \text{C} + 16.6 (\log M) + 0.41 (\% \text{GC}) - 0.61 (\% \text{form}) - 500/L$; where M is the molarity of monovalent cations, $\% \text{GC}$ is the percentage of guanosine and cytosine nucleotides in the DNA, $\% \text{form}$ is the percentage of formamide in the hybridization solution, and L is the length of the hybrid in base pairs. T_m is reduced by about 1°C . for each 1% of mismatching; thus, T_m , hybridization, and/or wash conditions can be adjusted to hybridize to sequences of the desired identity. For example, if sequences with $>90\%$ identity are sought, the T_m can be decreased 10°C . Generally, stringent conditions are selected to be about 5°C . lower than the T_m for the specific sequence and its complement at a defined ionic strength and pH. However, severely stringent conditions can utilize a hybridization and/or wash at 1, 2, 3, or 4°C . lower than the T_m ; moderately stringent conditions can utilize a hybridization and/or wash at 6, 7, 8, 9, or 10°C . lower than the T_m ; low stringency conditions can utilize a hybridization and/or wash at 11, 12, 13, 14, 15, or 20°C . lower than the T_m .

Using the equation, hybridization and wash compositions, and desired T_m , those of ordinary skill will understand that variations in the stringency of hybridization and/or wash solutions are inherently described. If the desired degree of mismatching results in a T_m of less than 45°C . (aqueous solution) or 32°C . (formamide solution), it is preferred to increase the SSC concentration so that a higher temperature can be used. An extensive guide to the hybridization of nucleic acids is found in Tijssen (1993) *Laboratory Techniques in Biochemistry and Molecular Biology—Hybridization with Nucleic Acid Probes*, Part I, Chapter 2 (Elsevier, New York); and Ausubel et al., eds. (1995) *Current Protocols in Molecular Biology*, Chapter 2 (Greene Publishing and Wiley-Interscience, New York). See Sambrook et al. (1989) *Molecular Cloning: A Laboratory Manual* (2d ed., Cold Spring Harbor Laboratory Press, Plainview, N.Y.).

As used herein, a substantially homologous sequence is a nucleic acid molecule that will specifically hybridize to the complement of the nucleic acid molecule to which it is being compared under high stringency conditions. Appropriate stringency conditions which promote DNA hybridization, for example, $6\times$ sodium chloride/sodium citrate (SSC) at about 45°C ., followed by a wash of $2\times \text{SSC}$ at 50°C ., are known to those skilled in the art or can be found in *Current Protocols in Molecular Biology*, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6. Typically, stringent conditions will be those in which the salt concentration is less than about 1.5 M Na ion, typically about 0.01 to 1.0 M Na ion concentration (or other salts) at pH 7.0 to 8.3 and the temperature is at least about 30°C . for short probes (e.g., 10 to 50 nucleotides) and at least about 60°C . for long probes (e.g., greater than 50 nucleotides). Stringent conditions may also be achieved with the addition of a destabilizing agent such as formamide. Exemplary low stringency conditions include hybridization with a buffer solution of 30 to 35% formamide, 1 M NaCl, 1% SDS (sodium dodecyl sulphate) at 37°C ., and a wash in $1\times$ to $2\times \text{SSC}$ ($20\times \text{SSC} = 3.0 \text{ M NaCl}/0.3 \text{ M trisodium citrate}$) at 50 to 55°C . Exemplary moderate stringency conditions include hybridization in 40 to 45% formamide, 1 M NaCl, 1% SDS at 37°C ., and a wash in $0.5\times$ to $1\times \text{SSC}$ at 55 to 60°C . Exemplary high stringency conditions include hybridization in 50% formamide, 1 M NaCl, 1% SDS at 37°C ., and a wash in $0.1\times \text{SSC}$ at 60 to 65°C . In a preferred embodiment, a nucleic acid of the present invention will specifically hybridize to one or more of

the nucleic acid molecules unique to the TC1507 event or complements thereof or fragments of either under moderately stringent conditions.

Methods of alignment of sequences for comparison are well known in the art. Thus, the determination of percent identity between any two sequences can be accomplished using a mathematical algorithm. Non-limiting examples of such mathematical algorithms are the algorithm of Myers and Miller (1988) *CABIOS* 4:11-17; the local homology algorithm of Smith et al. (1981) *Adv. Appl. Math.* 2:482; the homology alignment algorithm of Needleman and Wunsch (1970) *J. Mol. Biol.* 48:443-453; the search-for-similarity-method of Pearson and Lipman (1988) *Proc. Natl. Acad. Sci.* 85:2444-2448; the algorithm of Karlin and Altschul (1990) *Proc. Natl. Acad. Sci. USA* 87:2264, modified as in Karlin and Altschul (1993) *Proc. Natl. Acad. Sci. USA* 90:5873-5877.

Computer implementations of these mathematical algorithms can be utilized for comparison of sequences to determine sequence identity. Such implementations include, but are not limited to: CLUSTAL in the PC/Gene program (available from Intelligenetics, Mountain View, Calif.); the ALIGN program (Version 2.0); the ALIGN PLUS program (version 3.0, copyright 1997); and GAP, BESTFIT, BLAST, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Version 10 (available from Accelrys, 9685 Scranton Road, San Diego, Calif. 92121, USA). Alignments using these programs can be performed using the default parameters.

The CLUSTAL program is well described by Higgins and Sharp, *Gene* 73: 237-244 (1988); Higgins and Sharp, *CABIOS* 5: 151-153 (1989); Corpet, et al., *Nucleic Acids Research* 16: 10881-90 (1988); Huang, et al., *Computer Applications in the Biosciences* 8: 155-65 (1992), and Pearson, et al., *Methods in Molecular Biology* 24: 307-331 (1994). The ALIGN and the ALIGN PLUS programs are based on the algorithm of Myers and Miller (1988) supra. The BLAST programs of Altschul et al. (1990) *J. Mol. Biol.* 215:403 are based on the algorithm of Karlin and Altschul (1990) supra. The BLAST family of programs which can be used for database similarity searches includes: BLASTN for nucleotide query sequences against nucleotide database sequences; BLASTX for nucleotide query sequences against protein database sequences; BLASTP for protein query sequences against protein database sequences; TBLASTN for protein query sequences against nucleotide database sequences; and TBLASTX for nucleotide query sequences against nucleotide database sequences. See, *Current Protocols in Molecular Biology*, Chapter 19, Ausubel, et al., Eds., Greene Publishing and Wiley-Interscience, New York (1995). Alignment may also be performed manually by inspection.

To obtain gapped alignments for comparison purposes, Gapped BLAST (in BLAST 2.0) can be utilized as described in Altschul et al. (1997) *Nucleic Acids Res.* 25:3389. Alternatively, PSI-BLAST (in BLAST 2.0) can be used to perform an iterated search that detects distant relationships between molecules. See Altschul et al. (1997) supra. When utilizing BLAST, Gapped BLAST, PSI-BLAST, the default parameters of the respective programs (e.g., BLASTN for nucleotide sequences, BLASTX for proteins) can be used. See www.ncbi.nlm.nih.gov.

As used herein, "sequence identity" or "identity" in the context of two nucleic acid or polypeptide sequences makes reference to the residues in the two sequences that are the same when aligned for maximum correspondence over a specified comparison window. When percentage of sequence identity is used in reference to proteins it is recognized that residue positions which are not identical often differ by conservative amino acid substitutions, where amino acid residues

are substituted for other amino acid residues with similar chemical properties (e.g., charge or hydrophobicity) and therefore do not change the functional properties of the molecule. When sequences differ in conservative substitutions, the percent sequence identity may be adjusted upwards to correct for the conservative nature of the substitution. Sequences that differ by such conservative substitutions are said to have "sequence similarity" or "similarity". Means for making this adjustment are well known to those of skill in the art. Typically this involves scoring a conservative substitution as a partial rather than a full mismatch, thereby increasing the percentage sequence identity. Thus, for example, where an identical amino acid is given a score of 1 and a non-conservative substitution is given a score of zero, a conservative substitution is given a score between zero and 1. The scoring of conservative substitutions is calculated, e.g., as implemented in the program PC/GENE (Intelligenetics, Mountain View, Calif.).

As used herein, "percentage of sequence identity" means the value determined by comparing two optimally aligned sequences over a comparison window, wherein the portion of the polynucleotide sequence in the comparison window may comprise additions or deletions (i.e., gaps) as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. The percentage is calculated by determining the number of positions at which the identical nucleic acid base or amino acid residue occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison, and multiplying the result by 100 to yield the percentage of sequence identity.

Regarding the amplification of a target nucleic acid sequence (e.g., by PCR) using a particular amplification primer pair, "stringent conditions" are conditions that permit the primer pair to hybridize only to the target nucleic-acid sequence to which a primer having the corresponding wild-type sequence (or its complement) would bind and preferably to produce a unique amplification product, the amplicon, in a DNA thermal amplification reaction.

The term "specific for (a target sequence)" indicates that a probe or primer hybridizes under stringent hybridization conditions only to the target sequence in a sample comprising the target sequence.

As used herein, "amplified DNA" or "amplicon" refers to the product of nucleic acid amplification of a target nucleic acid sequence that is part of a nucleic acid template. For example, to determine whether a corn plant resulting from a sexual cross contains transgenic event genomic DNA from the corn plant of the present invention, DNA extracted from the corn plant tissue sample may be subjected to a nucleic acid amplification method using a DNA primer pair that includes a first primer derived from flanking sequence adjacent to the insertion site of inserted heterologous DNA, and a second primer derived from the inserted heterologous DNA to produce an amplicon that is diagnostic for the presence of the event DNA. Alternatively, the second primer may be derived from the flanking sequence. The amplicon is of a length and has a sequence that is also diagnostic for the event. The amplicon may range in length from the combined length of the primer pairs plus one nucleotide base pair to any length of amplicon producible by a DNA amplification protocol. Alternatively, primer pairs can be derived from flanking sequence on both sides of the inserted DNA so as to produce an amplicon that includes the entire insert nucleotide sequence of the PH18999A expression construct, see FIG. 1, approximately 6.2 Kb in size. A member of a primer pair derived from the

flanking sequence may be located a distance from the inserted DNA sequence, this distance can range from one nucleotide base pair up to the limits of the amplification reaction, or about twenty thousand nucleotide base pairs. The use of the term "amplicon" specifically excludes primer dimers that may be formed in the DNA thermal amplification reaction.

Nucleic acid amplification can be accomplished by any of the various nucleic acid amplification methods known in the art, including the polymerase chain reaction (PCR). A variety of amplification methods are known in the art and are described, inter alia, in U.S. Pat. Nos. 4,683,195 and 4,683,202 and in *PCR Protocols: A Guide to Methods and Applications*, ed. Innis et al., Academic press, San Diego, 1990. PCR amplification methods have been developed to amplify up to 22 Kb of genomic DNA and up to 42 Kb of bacteriophage DNA (Cheng et al., *Proc. Natl. Acad. Sci. USA* 91:5695-5699, 1994). These methods as well as other methods known in the art of DNA amplification may be used in the practice of the present invention. It is understood that a number of parameters in a specific PCR protocol may need to be adjusted to specific laboratory conditions and may be slightly modified and yet allow for the collection of similar results. These adjustments will be apparent to a person skilled in the art.

The amplicon produced by these methods may be detected by a plurality of techniques, including, but not limited to, Genetic Bit Analysis (Nikiforov, et al. *Nucleic Acid Res.* 22:4167-4175, 1994) where a DNA oligonucleotide is designed which overlaps both the adjacent flanking DNA sequence and the inserted DNA sequence. The oligonucleotide is immobilized in wells of a microwell plate. Following PCR of the region of interest (using one primer in the inserted sequence and one in the adjacent flanking sequence) a single-stranded PCR product can be hybridized to the immobilized oligonucleotide and serve as a template for a single base extension reaction using a DNA polymerase and labeled ddNTPs specific for the expected next base. Readout may be fluorescent or ELISA-based. A signal indicates presence of the insert/flanking sequence due to successful amplification, hybridization, and single base extension.

Another detection method is the Pyrosequencing technique as described by Winge (*Innov. Pharma. Tech.* 00: 18-24, 2000). In this method an oligonucleotide is designed that overlaps the adjacent DNA and insert DNA junction. The oligonucleotide is hybridized to a single-stranded PCR product from the region of interest (one primer in the inserted sequence and one in the flanking sequence) and incubated in the presence of a DNA polymerase, ATP, sulfurylase, luciferase, apyrase, adenosine 5' phosphosulfate and luciferin. dNTPs are added individually and the incorporation results in a light signal which is measured. A light signal indicates the presence of the transgene insert/flanking sequence due to successful amplification, hybridization, and single or multi-base extension.

Fluorescence Polarization as described by Chen et al., (*Genome Res.* 9:492-498, 1999) is a method that can be used to detect an amplicon of the present invention. Using this method an oligonucleotide is designed which overlaps the flanking and inserted DNA junction. The oligonucleotide is hybridized to a single-stranded PCR product from the region of interest (one primer in the inserted DNA and one in the flanking DNA sequence) and incubated in the presence of a DNA polymerase and a fluorescent-labeled ddNTP. Single base extension results in incorporation of the ddNTP. Incorporation can be measured as a change in polarization using a fluorometer. A change in polarization indicates the presence of the transgene insert/flanking sequence due to successful amplification, hybridization, and single base extension.

Taqman® (PE Applied Biosystems, Foster City, Calif.) is described as a method of detecting and quantifying the presence of a DNA sequence and is fully understood in the instructions provided by the manufacturer. Briefly, a FRET oligonucleotide probe is designed which overlaps the flanking and insert DNA junction. The FRET probe and PCR primers (one primer in the insert DNA sequence and one in the flanking genomic sequence) are cycled in the presence of a thermostable polymerase and dNTPs. Hybridization of the FRET probe results in cleavage and release of the fluorescent moiety away from the quenching moiety on the FRET probe. A fluorescent signal indicates the presence of the flanking/transgene insert sequence due to successful amplification and hybridization.

Molecular Beacons have been described for use in sequence detection as described in Tyangi et al. (*Nature Biotech.* 14:303-308, 1996). Briefly, a FRET oligonucleotide probe is designed that overlaps the flanking and insert DNA junction. The unique structure of the FRET probe results in it containing secondary structure that keeps the fluorescent and quenching moieties in close proximity. The FRET probe and PCR primers (one primer in the insert DNA sequence and one in the flanking sequence) are cycled in the presence of a thermostable polymerase and dNTPs. Following successful PCR amplification, hybridization of the FRET probe to the target sequence results in the removal of the probe secondary structure and spatial separation of the fluorescent and quenching moieties. A fluorescent signal results. A fluorescent signal indicates the presence of the flanking/transgene insert sequence due to successful amplification and hybridization.

A hybridization reaction using a probe specific to a sequence found within the amplicon is yet another method used to detect the amplicon produced by a PCR reaction.

The present invention is further defined in the following Examples. It should be understood that these Examples, while indicating preferred embodiments of the invention, are given by way of illustration only. From the above discussion and these Examples, one skilled in the art can ascertain the essential characteristics of this invention, and without departing from the spirit and scope thereof, can make various changes and modifications of the invention to adapt it to various usages and conditions. Thus, various modifications of the invention, in addition to those shown and described herein, will be apparent to those skilled in the art from the foregoing description. Such modifications are also intended to fall within the scope of the appended claims.

The disclosure of each reference set forth herein is incorporated herein by reference in its entirety.

EXAMPLES

Example 1

Transformation of Maize by Particle Bombardment and Regeneration of Transgenic Plants Containing the Cry1F Gene

A DNA molecule of 6.2 Kb, designated PHI8999A (see FIG. 1 and SEQ ID NO:25), which includes a first transgene expression cassette comprising the promoter, 5' untranslated exon, and first intron of the maize ubiquitin (Ubi-1) gene (Christensen et al. (1992) *Plant Mol. Biol.* 18:675-689 and Christensen and Quail (1996) *Transgenic Res.* 5:213-218) operably connected to a DNA molecule encoding a *Bacillus thuringiensis* δ -endotoxin identified as Cry1F (U.S. Pat. Nos. 5,188,960 and 6,218,188) operably connected to a DNA molecule comprising a 3' ORF25 transcriptional terminator iso-

lated from *Agrobacterium tumefaciens* (Barker et al. (1983) *Plant Mol. Biol.* 2:335-350), and a second transgene expression cassette comprising a DNA molecule of the cauliflower mosaic virus (CaMV) 35S promoter (Odell J. T. et al. (1985) *Nature* 313: 810-812; Mitsuhashi et al. (1996) *Plant Cell Physiol.* 37:49-59) operably connected to a DNA molecule encoding the selectable marker, phosphinothricin acetyltransferase (PAT) gene (Wohleben W. et al. (1988) *Gene* 70:25-37) operably connected to a DNA molecule comprising a 3' transcriptional terminator from (CaMV) 35S (see Mitsuhashi et al. (1996) *Plant Cell Physiol.* 37:49-59) was used to transform maize embryo tissue.

B.t. Cry1F maize plants were obtained by microprojectile bombardment using the Biolistics® PDS-1000He particle gun manufactured by Bio-Rad, Hercules, Calif.; essentially as described by Klein et al. (1987) *Nature*, UK 327(6117): 70-73. Immature embryos isolated from maize ears, harvested soon after pollination were cultured on callus initiation medium for several days. On the day of transformation, microscopic tungsten particles were coated with purified PHI8999A DNA (SEQ ID NO:25) and accelerated into the cultured embryos, where the insert DNA was incorporated into the cell chromosome. Only insert PHI8999A was used during transformation and no additional plasmid DNA was incorporated into the transformant. After bombardment, embryos were transferred to callus initiation medium containing glufosinate as the selection agent. Individual embryos were kept physically separate during culture, and the majority of explants died on the selective medium.

Those embryos that survived and produced healthy, glufosinate-resistant callus tissue were assigned unique identification codes representing putative transformation events, and continually transferred to fresh selection medium. Plants were regenerated from tissue derived from each unique event and transferred to the greenhouse. Leaf samples were taken for molecular analysis to verify the presence of the transgene by PCR and to confirm expression of the Cry1F protein by ELISA. Plants were then subjected to a whole plant bioassay using European corn borer insects. Positive plants were crossed with inbred lines to obtain seed from the initial transformed plants. A number of lines were evaluated in the field. The TC1507 event was selected from a population of independent transgenic events based on a superior combination of characteristics, including insect resistance and agronomic performance (see Bing J W et al. (2000) Efficacy of Cry1F Transgenic Maize, 14th Biennial International Plant Resistance to Insects Workshop, Fort Collins, Colo., herein incorporated by reference).

Example 2

Identification of Nucleotides Comprising the Flanking Sequence 5' to the Transgenic Insert DNA in *Bacillus thuringiensis* Cry1F Maize Line TC1507

To identify a DNA fragment that included sequence 5' to the PHI8999A insert in event TC1507, Spe I restriction enzyme fragments from event TC1507 genomic DNA were size selected on agarose gels, purified, and screened by Southern analysis to confirm hybridization to a Cry1F probe. Following confirmation of hybridization and fragment size, the fragments of interest were cloned into a pBluescript II SK (+)TM cloning vector to prepare an enriched size selected plasmid based genomic DNA library. A probe homologous to a portion of the Cry1F gene was used to screen the plasmid library for positive clones. A positive clone was identified, purified by additional screening, and confirmed to result in a

19

positive signal when hybridized to the Cry1F probe. Nearly 3 Kb of the Spe I fragment contained in the isolated positive clone was sequenced using a primer walking approach. To initiate the first sequencing run, a primer that binds to a known sequence in the cloning vector DNA was designed to sequence a portion of the DNA of interest. A second sequencing run over the same region using another primer oriented in the reverse direction provided second strand coverage. Primer walking was accomplished by repeatedly using sequence data from previous runs to design new primers that were then used to extend the next round of sequencing further into the DNA of interest until the flanking sequence 5' to the inserted transgenic DNA in maize event TC1507 was obtained. Specific sequence information is provided in Example 4.

Example 3

Confirmation of Flanking Sequence 5' to the B.t. Cry1F Maize Line TC1507 Insert

To confirm the 5' flanking sequence of the B.t. Cry1F maize line TC1507 insert, PCR primer pairs were designed to obtain overlapping PCR products extending from the 5' flanking region into the full-length PHI8999A transgenic insert. PCR

20

products were successfully amplified from B.t. Cry1F maize line TC1507 genomic DNA, isolated, and sequenced for Region 1 through Region 6, shown in Table 1, and confirmed to match the previously determined sequence from the Spe I fragment, described in Example 2. However, the region from by 2358 to by 2829, immediately adjacent and 5' to the start of the full-length insert was recalcitrant to PCR amplification and appeared to be larger than the sequence obtained from the Spe I clone described above. The use of primer pairs flanking this region and the Advantage®-GC 2 Polymerase Mix (BD Biosciences Clontech, Palo Alto, Calif.) was successful in amplifying PCR products from B.t. Cry1F maize line TC1507 genomic DNA for sequencing. The amplification conditions used to produce amplicons with the Advantage®-GC 2 system are shown in Table 10. The DNA primer pairs used to confirm the sequence in the region from by 2358 to 2829 are those listed in SEQ ID NO:1 and SEQ ID NO:2; and SEQ ID NO:2 and SEQ ID NO:23. Sequence from this region is described in Table 1 (Regions 7a, 7b, 7c, and 8).

Example 4

Event TC1507 5' Flanking Sequence

A description of each region is provided in Table 1.

Region 1 Maize genomic (no significant homology) (SEQ ID NO: 28)

```

1  ACTAGTTTCC TAGCCCGCGT CGTGCCCTA CCCACCGAC GTTTATGGAA
51 GGTGCCATT CACGGTTCTT CGTGGCCGCC CCTAAGGATG TAAATGGTCG
101 GTAAATCCG GTAAATTTCC GGTACCGTTT ACCAGATTTT TCCAGCCGTT
151 TTCGGATT A TCGGGATATA CAGAAAACGA GACGAAACG GAATAGGTTT
201 TTTTTCGAAA ACGGTACGGT AACCGGTGAG ACAAACCTAC CGTCCGTTTT
251 CGTATTTCTC GGGAACTCT GGTATATTCC CGTATTTGTC CCGTATTTTC
301 CCGACCCACG GACCTGCCAA TCAACCATCA GCCAGTCAGC CCATCCCCAC
351 AGCTATGGCC CATGGGGCCA TGTGGCCAC ATGCCACGC AACGCAAGGC
401 AGTAAGGCTG GCAGCCTGGC ACGCATTGAC GCATGTGGAC ACACACAGCC
451 GCCGCCTGTT CGTGTTCCTG TGCCGTGTG CGAGACTGTG ACTGCGAGTG
501 GCGGAGTCGG CGAACGGCGA GCGCTCTCCG GAGTCTGGAC TCGCGCTGTG
551 GACAGCGACG CTGTGACGGC GACTCGGCGA AGCCCCAAGC TACCAAGCCC
601 CCAAGTCCCC ATCCATCTCT GCTTCTCTGG TCATCTCCTT CCCCTGGTCG
651 ATCTGCAGGC GCCAGACCG

```

Region 2 Undescribed maize genomic sequence (complement) (SEQ ID NO: 29)

```

670 G CCGAAGCATC ACGAAACGCA CTAAGACCTC
701 GAAGGAGTCA AACCACTCCT CCGAGGCCCT GGGGGCTACA CCCGGCGGGT
751 GCGCTCGCGC GCACCCACCG GAACAAAATG TAACCGAGAA AGGTCGGTCC
801 CTTTGCAAAA AAAGTGCAC AAAAGCCTCC AAGCGAGTAT TAACACTCAC
851 TTTGAGGCTC GGGGGCTAC

```

Region 3 Fragment of maize Huck-1 retrotransposon (SEQ ID NO: 30)

```

870 T GTCGGGGACC ATAATTAGGG GTACCCCAA
901 GACTCCTAAT CTCAGCTGGT AACCCCATC AGCACAAGC TGCAAAGGC
951 TGATGGGTGC GATTAAGTCA AGGCTCGGTC CACTCAAGG ACACGATCTC

```

-continued

1001 GCCTCGCCCG AGCCCAGCCT CGGGCAAGGG CGGCCGACCC CGAGGATTCA
 1051 CGTCTCGCCC GAGGGCCCCC TCAAGCGACG GGCACACCTT CGGCTCGCCC
 1101 GAGGCCCATC CTTCGCCGAG AAGCAACCTT GGCCAGATCG CCACACCGAC
 1151 CGACCGTATC GCAGGAGCAT TTAATGCGAG GATCGCCTGA CACCTTATCC
 1201 TGACGCGCGC TCTTCAGTCG ACAGAGCCGA AGTGACCGCA ATCACTTCGC
 1251 CGCTCCACTG ACCGACCTGA CAAGAAGACA GCGCCGCCTG CGTCGCTCCG
 1301 ACTGCTGTGC CACTCGACAG AGTGAGGCTG ACAGCAGCCA AGTCCGGCCT
 1351 CGGCGCCCAT AGGAAGCTCC GCCTCGCCCG ACCCTAGGGC TCGGACTCGG
 1401 CCTCGGCTCC GGAAGACGAC GAACTACGCT TCGCCCGACC CCAGGGCTTG
 1451 GACTCAGCCT CGGCTCCGGA AGACGACGAA TTCCGCCTCG CCCGACCCCA
 1501 GGGCTCGGAC TCGGCCTCGG CTCCAGAAGA CGACGAACTC CGCCTCGCCC
 1551 GACCCAGGG CTCGGACTCA GCCTCGGCTC CGGAAGACGA CGAACTCCGC
 1601 CTCGCCCAGC CCCAGGGCTC GGAATCAGCC TCGGCCTCAG ACGATGGTCT
 1651 CCGCCTCGCC CGACCCGGGG CTCGGACTCG A

Region 4 Fragment of crylF gene

(SEQ ID NO: 31)

1682 CCTTTCTAT CGGACCTTGT
 1701 CAGATCCTGT CTTCGTCCGA GGAGGCTTTG GCAATCCTCA CTATGTACTC
 1751 GGTCTTAGGG GAGTGGCCTT TCAACAACT GGTACGAATC ACACCCGCAC
 1801 ATTCAAGAAC TCCGGGACCA TTGACTCTCT AGATGAGATA CCACCTCAAG
 1851 ACAACAGCGG CGCACCTTGG AATGACTACT CCCATGTGCT GAATCATGTT
 1901 ACCTTTGTGC GCTGGCCAGG TGAGATCTCA GGTTCCGACT CATGGAGAGC
 1951 ACCAATGTTT TCTTGACGC ATCGTAGCGC TACCCCAACA AACACCATTG
 2001 ATCCAGAGAG AATCAC

Region 5 Fragment of maize chloroplast rpoC2 gene

(SEQ ID NO: 32)

2017 TCAT TCTTCAAGAA CTGCATATCT TGCCGAGATC
 2051 CTCATCCCTA AAGGTACTTG ACAATAGTAT TATTGGAGTC GATACACAAC
 2101 TCACAAAAAA TACAAGAAGT CGACTAGGTG GATTGGTCCG AGTGAAGAGA
 2151 AAAAAAAGCC ATACAGAAGT CAAAATCTTT TCCGGAGATA TTCATTTTCC
 2201 TGAAGAGCGG GATAAGATAT TAGGTGGCAG TTTGATACCA CCAGAAAGAG
 2251 AAAAAAAGA TTCTAAGGAA TCAAAAAAAA GGAAAAATTG GTTTATGTT
 2301 CAACGGAAAA AATTTCTCAA AAGCAAGGAA AAGTATT

Region 6 Fragment of maize chloroplast or ubiZM1(2) promoter

(SEQ ID NO: 33)

2338 GTG GCTATTTATC

2351 TATC

Nucleotides 2355-2358 (CGT) connect Region 6 to Region 7a.

Region 7a Fragment of pat gene

(SEQ ID NO: 34)

2358 GCA GCTGATATGG CCGCGGTTTG TGATATCGTT AACCATTACA
 2401 TTGAGACGTC TACAGTGAAC TTTAGGACAG AGCCACAAAC ACCACAAGAG
 2451 TGGATTGATG ATCTAGAGAG GTTGCAAGAT AGATACCCCTT GGTGTTGTC
 2501 TGAGGTTGAG GGTGTTGTGG CTGGTATTGC TTACGCTGGG CCCTGGAAGG
 2551 CTAGGAAC

-continued

Region 7b Fragment of pat gene (complement)
(SEQ ID NO: 35)
2559 CC TCAACCTCAG CAACCAACCA ATGGTATCTA TCTTGCAACC
2601 TCTCTAGATC ATCAATCCAC TCTTGTGGTG TTTGTGGCTC TGTCTTAAAG
2651 TTCCTGTAG ACGTCTCAAT GTAATGGTTA ACGATATCAC AAACCG
Region 7c Fragment of cry1F gene (complement)
(SEQ ID NO: 36)
2697 AGAG
2701 AAGAGGGATC T
Region 8 Fragment of Polylinker
(SEQ ID NO: 37)
2712 CGAAGCTTC GGCCGGGGCC CATCGATATC CGCGGGCATG
2751 CCTGCAGTGC AGCGTGACCC GGTCGTGCCC CTCTCTAGAG ATAATGAGCA
2801 TTGCATGTCT AAGTTATAAA AAATTACCA
Region 9 Full-length insert of PHI8999A
(SEQ ID NO: 25)

Example 5

Description of the Flanking Sequence 5' to the Insert
in Maize Event TC1507

In order to more fully describe the event TC1507 5' flanking sequence, homology searching was done against the GenBank public databases (release 122, 2/01) using the Basic Local Alignment Search Tool (BLAST). The BLAST program performs sequence similarity searching and is particularly useful for identifying homologs to an unknown sequence. In addition to searching the public databases, pairwise alignments were performed using AlignX (InforMax Inc., Bethesda, Md.) to look for homology between the maize event TC1507 flanking sequence and the PHI8999A transgenic insert. The results of these homology searches are presented in Table 1. The TC1507 5' flanking sequence is numbered with base 1 being the furthest 5' to the insert and base 2830 at the starting point of the full-length PHI8999A trans-

genic insert (see FIG. 1). The percent identity values indicate the percentage of identical matches across the length of the sequences analyzed.

In most cases, similarity searching with the event TC1507 5' flanking sequence resulted in a match to one unique sequence based on a very high percent identity value. Those sequences are identified in Table 1. In addition, there are two regions in the TC1507 5' DNA flanking sequence with high similarity to more than one known sequence. In regions 870-1681 and 2338-2354, the percent identity scores with both sequence fragments are sufficiently high that a single match (homolog) cannot be determined. The two possible homologs for each of these regions are indicated in Table 1.

Highly similar sequences were identified for all but the first 669 base pairs of sequence. Generally, the results of similarity searching indicate high homology with maize genomic sequences 5' to base 1681. The region from base 1682 to the start of the PHI8999A insert at position 2830 contains some fragments associated with the transformation event.

TABLE 1

Sequence summary for event TC1507 insert						
Region	Location in SEQ ID NO: 24	Size bp	% Identity	Homolog	Location in homologous sequence	Description
1	1-669	669	N/A ¹	N/A	N/A	No significant homology detected
2	670-869	200	90.5	AF123535	52432-52632 (complement)	Undescribed maize genomic sequence
3	870-1681	812	89.4	AF050439	1-801	Fragment of maize Huck-1 retrotransposon 5' LTR ²
			86.6	AF050438	1-797	Fragment of maize Huck-1 retrotransposon 3' LTR
4	1682-2016	335	100.0	PHI8999A	3149-3483	Fragment of cry1F gene
5	2017-2337	321	100.0	X86563	29429-29749	Fragment of maize chloroplast rpoC2 gene (RNA polymerase beta-2 subunit)

TABLE 1-continued

Sequence summary for event TC1507 insert						
Region	Location in SEQ ID NO: 24	Size bp	% Identity	Homolog	Location in homologous sequence	Description
6	2338-2354	17	100.0	X86563	97643-97659	Fragment of maize chloroplast trnI gene (tRNA-Ile)
			82.4	PHI8999A	182-197	Fragment of maize ubiZM1(2) promoter
7a	2358-2558	201	100.0	PHI8999A	5320-5475	Fragment of pat gene
7b	2559-2696	138	99	PHI8999A	5336-5518 (complement)	Fragment of pat gene
7c	2697-2711	15	100.0	PHI8999A	2544-2558 (complement)	Fragment of cry1F gene
8	2712-2829	118	100.0	PHI8999A	36-153	Fragment of polylinker region (bases 36-80) and ubiZM1(2) promoter (bases 81-153)
9	2830-9015	6186	100.0	PHI8999A	11-6196	Full-length insert of PHI8999A
10	9016-9565	550	100.0	PHI8999A	3906-4456 (complement)	Inverted ORF25 terminator
11	9566-9693	128	100.0	NC_001666	121851-121978 (complement) & 100759-100886	Fragment of maize chloroplast rps12 rRNA (23S ribosomal RNA)
12	9696-10087	392	99	NC_001666	17091-17483 (complement)	Fragment of maize chloroplast genome
13	10088-10275	188	99	PHI8999A	5333-5520 (complement)	Fragment of pat gene
14	10278-10358	81	100	NC_001666	137122-137202 (complement)	Fragment of maize chloroplast "ORF241"-hypothetical protein gene
15	10359-10612	254	N/A ¹	N/A	N/A	No significant homology detected
16	10613-11361	749	N/A ¹	N/A	N/A	No description available

¹N/A; not applicable²LTR; long terminal repeat

Example 6

Confirmation of the Presence of Regions 1, 2, and 3 in an Unmodified Control Corn Line

PCR analysis was used to determine if Regions 1, 2, and 3 (Table 1) in the 5' flanking region of Event TC1507 are present in an unmodified control corn line used for transformation to produce maize event TC1507 and thus represents a border with corn genomic DNA. Nine different PCR analyses were carried out on genomic DNA prepared from TC1507 and the unmodified control corn line Hi-II (see Armstrong (1994) *The Maize Handbook*, ed. Freeling and Walbot, Springer-Verlag, New York, pp. 663-671, for information on Hi-II) as outlined in Table 2 using the primer sequences shown in Table 3. Two reactions were designed to amplify DNA within Region 1 of the 5' flanking region from by 25 to 324 (Reaction A—300 by amplicon); and from by 25 to 480 (Reaction B—456 by amplicon). The expected amplicons

45 were present in both the Hi-II unmodified corn line and in maize event TC1507. One PCR primer pair, Reaction C, spanned Region 2 to Region 3 of the 5' flanking region from by 759 to 1182 (424 by amplicon) and again produced PCR products of the expected size in both Hi-II and TC1507. Reaction D, spanned Region 1 to Region 3 of the 5' flanking region from by 415 to 1182 (768 by amplicon) and again produced PCR products of the expected size in both Hi-II and TC1507. Reactions E and F were designed as specific primer pairs for the pat gene region of the full-length insert of PHI8999A in TC1507 and thus an amplicon in the unmodified Hi-II corn line is not expected. The results indicate that both Reactions E and F are specific for a maize line transformed with a pat gene region and produce the expected amplicon, whereas no amplicon was produced in the unmodified Hi-II corn line. Reaction G was also designed as a primer pair that would produce an amplicon of 366 by in the maize event TC1507 and no amplicon in the unmodified Hi-II corn line.

Reactions H and I were designed as specific primer pairs for TC1507 that would span the end of the transgenic insert

27

into the 5' flanking region. In both Reactions H and I, the reverse primer was located in the ubiquitin promoter region of the full-length PHI8999A insert (Region 9 in Table 1) and the forward primer was located in Region 5, the rpoC2 gene fragment (see Table 1). Reaction H and Reaction I both produced an amplicon in maize line TC1507 and did not produce an amplicon in the unmodified control corn line. These results indicate that both Reactions H and I are specific for the TC1507 event.

The PCR results show that the undescribed sequence (Region 1) is present in the unmodified corn line Hi-II and that Regions 1, 2 and 3, are contiguous in the unmodified corn line Hi-II. The DNA sequences amplified in Reactions A, B, C, and D are not unique to the 5' flanking region of maize event TC1507 but are also present in the unmodified corn line Hi-II.

28

TABLE 3

PCR primers for sequence 5' to the PHI8999A insert in TC1507 and for regions within the full-length insert of PHI8999A in maize event TC1507				
Reaction	Amplicon Size (bp)	Primer Pair	Primer Sequences 5' to 3'	
A	300	SEQ ID NO: 10	CCCTACCCACCGACGT TTAT	
		SEQ ID NO: 11	TTGATTGGCAGGTCCGTG GGTC	
B	456	SEQ ID NO: 10	CCCTACCCACCGACGT TTAT	

TABLE 2

PCR reactions for sequence 5' to the PHI8999A insert in maize event TC1507 and for regions within the full-length insert of PHI8999A in maize event TC1507					
Reaction	PCR Amplicon Location	Amplicon Size (bp)	Region in TC1507 flanking sequence or PHI8999A insert	Amplicon present in Hi-II	Amplicon present in maize line TC1507
A	25-324 bp in TC1507 flanking sequence	300	Region 1	Yes	Yes
B	25-480 bp in TC1507 flanking sequence	456	Region 1	Yes	Yes
D	415-1182 bp in TC1507 5' flanking sequence	768	Region 1 to Region 3	Yes	Yes
E Not Unique to TC1507	4750-5794 bp in PHI8999A	1045	Region 9 (in full-length insert of PHI8999A 35S promoter to pat gene)	No	Yes
F Not Unique to TC1507	4827-5308 bp in PHI8999A	482	Region 9 (in full-length insert of PHI8999A 35S promoter to pat gene)	No	Yes
Detects cry1F fragment in 5' flanking region	cry1F sequence in 5' flanking sequence and in full-length insert of PHI8999A	366	Spans 335 bp cry1F sequence in 5' flanking sequence and same sequence in the full-length insert	No	Yes
H Unique to TC1507	2158 bp in Region 5 (rpoC2 gene fragment) to 3069 bp in Region 9 (full-length insert of PHI8999A)	912	Region 5 to Region 9 Unique to Insertion Event [SPANS UNIQUE JUNCTION REGIONS]	No	Yes
I Unique to TC1507	2158 bp in Region 5 (rpoC2 gene fragment) to 3001 bp in Region 9 (full-length insert of PHI8999A)	844	Region 5 to Region 9 Unique to Insertion Event [SPANS UNIQUE JUNCTION REGIONS]	No	Yes

29

TABLE 3-continued

PCR primers for sequence 5' to the PHI8999A insert in TC1507 and for regions within the full-length insert of PHI8999A in maize event TC1507				
Reaction	Amplicon Size (bp)	Primer Pair	Primer Sequences 5' to 3'	
		SEQ ID NO: 12	CACAAACGGCACAGAAACA CGAA	
C	424	SEQ ID NO: 13	GCGCACCCACCGGAACAA AATG	
		SEQ ID NO: 14	TCCTCGCATTAAATGCTC CTGC	
D	768	SEQ ID NO: 15	CCTGGCAGCATTGACGC ATGT	
		SEQ ID NO: 14	TCCTCGCATTAAATGCTC CTGC	
E	1045	SEQ ID NO: 6	TAGAGGACCTAACAGAAC TCGCCGT	
		SEQ ID NO: 7	GAGCTGGCAACTCAAAAT CCCTTT	
F	482	SEQ ID NO: 8	AAAATCTTCGTCAACATG GTGGAGC	
		SEQ ID NO: 9	TAATCTCAACTGGTCTCC TCTCCGG	
G	366	SEQ ID NO: 19	GGCTCGGACTCGACCTTT CTAT	
		SEQ ID NO: 20	GCAGTTCTTGAAGAATGA GTGA	
H	912	SEQ ID NO: 1	GTAGTACTATAGATTATA TTATTCGTAGAG	
		SEQ ID NO: 2	GCCATACAGAACTCAAAA TCTTTTCCGGAG	
I	844	SEQ ID NO: 2	GCCATACAGAACTCAAAA TCTTTTCCGGAG	
		SEQ ID NO: 23	CTTCAACAAGGTGACA AA	

Example 7

Flanking Sequence 3' to Inserted Transgenic DNA in Maize Event TC1507

Two separate PCR approaches were used to extend the length of the sequence information 3' to the full-length PHI8999A insert in maize event TC1507. In the first approach PCR primer pairs were designed to amplify a product that spanned the junction between the full-length insert and the inverted ORF25 terminator, see FIG. 1 for a depiction of the inverted ORF25 terminator. A forward primer was located at the end of the full-length PHI8999A insert and a series of reverse primers were located at 100 by intervals in the inverted sequence. In this manner the length of the inverted fragment present in the maize event TC1507 could be determined within a 100 by region based on the successful PCR reactions. This method indicated the inverted fragment contained the majority of the ORF25 terminator but no Cry1F sequence. PCR fragments were isolated and sequenced from this region.

In the second approach PCR primers were designed to walk out into the flanking DNA sequence from the inverted ORF25 terminator region as determined in the PCR experiment

30

described above. Genomic DNA isolated from two to three individual plants of event TC1507 and an unmodified control corn line was digested with various restriction enzymes and then ligated to adaptors specific for the restriction enzyme used for digestion (Universal Genome Walker™ Kit, Clontech Laboratories, Inc. and Devon et al. (1995) *Nucleic Acids Res.* 23:1644-1645). Primary PCR was carried out using an ORF25 terminator specific primer and a primer homologous to the adaptor sequence ligated onto the digested DNA. In order to increase the specificity of the reaction a nested secondary PCR was performed again with another ORF25 terminator specific primer and a secondary primer homologous to the adaptor sequence with the secondary primers being internal to the respective primers used in the primary PCR. Products produced by the nested PCR were analyzed by agarose gel electrophoresis and fragments unique to TC1507 DNA samples were isolated and sequenced. Fragments were amplified from both the ORF25 terminator contained within the full-length insert and from the targeted (inverted) ORF25 terminator on the 3' end of the full-length PHI8999A insert. Fragments from the full-length insert were of a predicted size based on the knowledge of the restriction enzyme sites located in the full-length insert. Fragments produced from the 3' inverted ORF25 terminator appeared as fragments of unexpected size. Sequence analysis of amplified fragments from the 3' inverted ORF25 terminator resulted in flanking DNA sequence of 1043 bp. Resultant sequence from the above series of genome walking experiments was used to design additional primers to walk further out from the insert into the bordering maize genome with a final 3' flanking sequence, of 2346 bp.

In order to describe the TC1507 3' flanking sequence, homology searching was done against the GenBank public databases using the Basic Local Alignment Search Tool (BLAST). The BLAST program performs sequence similarity searching and is particularly useful for identifying homologs to an unknown sequence. In addition to searching the public databases, alignments were performed using SeqMan 4.05™, Martinez and Needleman-Wunsch alignment algorithms (DNASTAR Inc.) to look for homology between the TC1507 3' flanking sequence and the PHI8999A transgenic insert. The results of these homology searches are presented in Table 1. The percent identity values indicate the percentage of identical matches across the length of the sequences analyzed. The results of similarity searching for the 3' flanking sequence indicate high homology with three regions of maize chloroplast DNA, a 188 by fragment of the pat gene, and 254 by of DNA (Region 15, Table 1) with no significant homology. An additional 749 by (Region 16) beyond Region 15 (see Table 1) was also sequenced. No similarity searching results are available for Region 16.

PCR analysis on control and TC1507 genomic DNA determined that the 254 by sequence (Region 15, fragment of maize chloroplast "ORF241") is present in the maize genome. The DNA sequence of Region 15 in the 3' flanking region is not unique to the 3' flanking region of maize event TC1507 but is also present in the unmodified control corn line. The TC1507 3' flanking sequence is presented in Example 8 and diagrammed in FIG. 1.

Sequence of the Region 3' to the End of the
Full-Length Insert DNA in Maize Event TC1507

A description of each region is in Table 1.

Region 10 Fragment of ORF25 Terminator (complement)

(SEQ ID NO: 38)

9016 CTCAC TCCGCTTGAT CTTGGCAAAG ATATTTGACG
9051 CATTATTAG TATGTGTTAA TTTTCATTG CAGTGCAGTA TTTTCTATTC
9101 GATCTTTATG TAATTCGTTA CAATTAATAA ATATTCAAAT CAGATTATTG
9151 ACTGTCATTT GTATCAAATC GTGTTTAATG GATATTTTAA TTATAATATT
9201 GATGATATCT CAATCAAAAC GTAGATAATA ATAATATTTA TTAAATATT
9251 TTGCGTCGCA CAGTGAAAAT CTATATGAGA TTACAAAATA CCGACAACAT
9301 TATTTAAGAA ACATAGACAT TAACCCCTGAG ACTGTTGGAC ATCAACGGGT
9351 AGATTCTTC ATGCATAGCA CCTCATTCTT GGGGACAAAA GCACGGTTTG
9401 GCCGTTCCAT TGCTGCACGA ACGAGCTTTG CTATATCCTC GGGTTGGATC
9451 ATCTCATCAG GTCCAATCAA ATTTGTCCAA GAACTCATGT TAGTCGCAAC
9501 GAAACCGGG CATATGTCGG GTATCTCGAG CTCGCGAAAG CTTGGCTGCA
9551 GGTCGACGGA TCCTT

Region 11 Fragment of maize chloroplast rps12 rRNA gene (complement)

(SEQ ID NO: 39)

9566 CAACA AAAGGGTACC TGTACCCGAA ACCGACACAG
9601 GTGGGTAGGT AGAGAATACC TAGGGGCGCG AGACAACCTCT CTCTAAGGAA
9651 CTCGGCAAAA TAGCCCCGTA ACTTCGGGAG AAGGGGTGCC CCC

Nucleotides 9694-9695 (CG) connect Region 11 to Region 12.

Region 12 Fragment of maize chloroplast genome

(SEQ ID NO: 40)

9696 CTAAC
9701 AATAAACGAA TACGGTTTAT GTATGGATTC CGGTAAATA CCGGTACTCG
9751 ATTTCATAAG AGTCGAATAG GAAGTTAAGA TGAGGGTGGT ATCATCATAA
9801 AAATGGAGTA GTATCCTAAA TTATACTAAT CCACGTATGA TATGTATGCC
9851 TTTCTTATC AACCGGAAGT AGTGCAAAAA AAATTCCTATA CTGCACTGCT
9901 CTCTTTTAC TGAGAAATGC AAAAAAATAA AAGTGAAGTA AGGGTGCCCC
9951 ATAGATATTT GATCTTGCTT CCTGTCCCC CCCCTTTT TTCATCAAAA
10001 ATTTCCATGA AAAAAGAAAA GATGAATTG TCCATTCATT GAACCTAGT
10051 TCGGGACTGA CGGGGCTCGA ACCCGCAGCT TCGCCT

Region 13 Fragment of pat gene (complement)

(SEQ ID NO: 41)

10088 GTT CCTAGCCTTC
10101 CAGGGCCCAG CGTAAGCAAT ACCAGCCACA GCACCCTCAA CCTCAGCAAC
10151 CAACCAAGGG TATCTATCTT GCAACCTCTC TAGATCATCA ATCCACTCTT
10201 GTGGTGTTTG TGGCTCTGTC CTAAAGTTCA CTGTAGACGT CTCAATGTAA
10251 TGGTTAACGA TATCACAAAC CGCGG

- continued

Nucleotides 10276-10277 (AA) connect Region 13 to Region 14.

Region 14 Fragment of maize chloroplast ORF241 (complement)

(SEQ ID NO: 42)

10278 CAC AAGAACGAAA GCACCTTTTC

10301 ATTCTTTTCAT ATACTAGGGG TTTTACTTG GAAAAGACAA TGTTCCATAC

10351 TAAAGGAT

Region 15 Maize genomic (no significant homology)

(SEQ ID NO: 43)

10359 AG CTGCAGAAGC CGCCACCGTC TTGAGGACCT TCCGGGGAGC

10401 CAGACCGGTC GAACCGTGCC TCCACTTGCT AAGGAGAAAG GGAAATCAG

10451 GGCCAGGACA TACGAAGGAG GAGCCAGAAC GAAGATATCC TAAGATACTT

10501 ACTCGCTCCG GGCCATGATC AATCATGCCT GTGGGGAGGT CTCTCGCACC

10551 TCGATCCATG AAGGTACCAC CGAGGTCTGC CCCGCCGCCG GCTTCGGTAC

10601 CGTCCTCGCC TT

Region 16 Maize genomic

(SEQ ID NO: 44)

10613 GGGCGCCC GAGGCACCCG GGGGATGGAC TGCCAGGCG

10651 CAGCCACGAC GACCCAAGGA TCACCTCCT GCGCAGTCGG CACGAGCAAT

10701 AGTTCTCGGG GAACAGGCAG CTTGGCCTGA CTCGCCGGG TCACCTCAAC

10751 TACCTCGGCC GAGGGGTCAA GTACCCCTC AGTCCGCCCC CGCTCTTCGG

10801 ACCGGGACCC CGACGTCCCG GCCCCGGATA CCGACGGCAC CAGCCCGCTC

10851 GGGGGCTGGC TTGACGACCC CTGGCCAGC CTCAGATCTG GGCTGAGGCC

10901 GAGGCAGGCG GCCATGTCGT CGTCTTCATC ATCGTCTTCA TCATCGTCGT

10951 CGTCATCAGG CGTCTCCGGC GACGGCTCCC TTGGGAGCCC CTCCCTCTCC

11001 TGCCGACGAC GAAGCCTTTC CAAGGCATCC CGAGCCACG TCCGCTCGTG

11051 GGCCCGAGCC TTCTTTGCGT CCTTCTCTC CTTCTCTTC TCCGCGGTGA

11101 CCCTCCGCGC AGCTCGGTCC ACCGCATCCT CCGGGACTGG TGGCAGGGAA

11151 GGCTTGATGAT GCCCTACCTC CTGGAGACAG ACGAAAAGTC TCAGCTATGA

11201 GAACCGAGGG CAATCTGACG CAAGAAGGAA GAAGGAGCGG ATACTCACCA

11251 GAGACACGCA CCCGCGATCG GGACGCATTA AGGGCTGGGA AAAAGTGCCG

11301 GCCTCTAATT TCGCTACCGT GCCGTCCACC CACCTGTGGA GGTATCGAT

11351 GGGAAGGGGA A

Example 9

Confirmation of the Presence of Region 15 in the
Unmodified Control Corn Line

PCR analysis was used to determine if the undescribed region of sequence on the end of the 3' flanking sequence (Region 15 in Table 1) is present in the unmodified control corn line used for transformation to produce maize event TC1507 and thus represents a border with corn genomic DNA. Successful PCR amplification of Region 15 in both maize line TC1507 and the unmodified Hi-II control corn line revealed that Region 15 was indeed present in corn genomic DNA. Five different PCR analyses were carried out on genomic DNA prepared from TC1507 and the unmodified Hi-II control corn line as outlined in Table 7 below using the primer sequences shown in Table 8. Three reactions were designed to amplify DNA within Region 15 of the 3' flanking region; Reaction L—producing a 175 by amplicon, Reaction

M—producing a 134 by amplicon, and Reaction N—producing a 107 by amplicon. The expected amplicons were present in both the unmodified control corn line and in maize line TC1507. Reactions J and K were designed as specific primer pairs for TC1507 that would span the end of the insert into the 3' flanking region. In Reaction J, the forward primer was located in the pat gene fragment on the 3' end of the full-length PHI8999A insert (Region 13 in Table 1) and the reverse primer was located in the undefined Region 15. In Reaction K the forward primer was located in the chloroplast hypothetical protein gene on the 3' end of the full-length insert (Region 14 in Table 1) and the reverse primer was located in the undefined Region 15. Both Reaction J and Reaction K produced an amplicon in maize line TC1507 and did not produce an amplicon in the unmodified control corn line. The results indicate that both Reactions J and K are specific for the TC1507 event.

The PCR results indicate that the undescribed sequence (Region 15) of the 3' flanking sequence of TC1507 is also

35

present in genomic DNA isolated from the unmodified Hi-II control corn line. The DNA sequences amplified in Reactions L, M, and N are not unique to the 3' flanking region of TC1507 but are also present in the unmodified control corn line.

TABLE 7

PCR reactions for sequence 3' to the PHI8999A insert in maize event TC1507				
Reaction	Amplicon Size (bp)	Region in TC1507 3' flanking sequence	Amplicon present in Control	Amplicon present in maize line TC1507
J	342	Region 13 (pat gene fragment) to Region 15	No	Yes
K	252	Region 14 (chloroplast gene) to Region 15	No	Yes
L	175	Region 15	Yes	Yes
M	134	Region 15	Yes	Yes
N	107	Region 15	Yes	Yes

TABLE 8

PCR primers for sequence 3' to the PHI8999A insert in maize event TC1507				
Reaction	Amplicon Size (bp)	Primer Pair	Primer Sequences 5' to 3'	
J	342	SEQ ID NO: 3	TGTGGTGTGTTGTGGCTCTGT CCTAA	
		SEQ ID NO: 5	GACCTCCCCACAGGCATGAT TGATC	
K	252	SEQ ID NO: 4	AGCACCTTTTCATTCTTTCA TATAC	
		SEQ ID NO: 5	GACCTCCCCACAGGCATGAT TGATC	
L	175	SEQ ID NO: 16	AAGCCGCCACCGTCTTGAGG ACCTT	
		SEQ ID NO: 5	GACCTCCCCACAGGCATGAT TGATC	
M	134	SEQ ID NO: 17	GTCGAACCGTGCCTCCACTT GCTAA	
		SEQ ID NO: 5	GACCTCCCCACAGGCATGAT TGATC	
N	107	SEQ ID NO: 18	AGAAAGGGAAAATCAGGGCC AGGAC	
		SEQ ID NO: 5	GACCTCCCCACAGGCATGAT TGATC	

Example 10

PCR Primers

DNA event specific primer pairs were used to produce an amplicon diagnostic for TC1507. These event primer pairs include, but are not limited to, SEQ ID NO: 1 and SEQ ID NO: 2; SEQ ID NO: 2 and SEQ ID NO: 23; SEQ ID NO: 3 and SEQ ID NO: 5; and SEQ ID NO: 4 and SEQ ID NO: 5. In addition to these primer pairs, any primer pair derived from SEQ ID NO: 26 and SEQ ID NO: 27 that when used in a DNA amplification reaction produces a DNA amplicon diagnostic for TC1507 is an aspect of the present invention. The amplification conditions for this analysis are illustrated in Table 9, however, any modification of these methods that use DNA

36

primers or complements thereof to produce an amplicon DNA molecule diagnostic for TC1507 is within the ordinary skill of the art. The preferred amplification conditions for reactions utilizing the PCR primers identified in SEQ ID NOS: 1, 2, and 23 are illustrated in Table 10. In addition, control primer pairs, which include SEQ ID NOS: 10 and 11; SEQ ID NOS: 10 and 12; SEQ ID NOS: 13 and 14; SEQ ID NOS: 14 and 15; SEQ ID NOS: 5 and 16; SEQ ID NOS: 5 and 17; and SEQ ID NOS: 5 and 18; for amplification of an endogenous corn gene are included as internal standards for the reaction conditions. Also included are primer pairs that will produce an amplicon in transgenic events containing a pat gene (SEQ ID NOS: 6 and 7; SEQ ID NOS: 8 and 9), and a primer pair that will produce an amplicon in transgenic events containing a cry1F gene (SEQ ID NOS: 19 and 20).

The analysis of plant tissue DNA extracts to test for the presence of the TC1507 event should include a positive tissue DNA extract control (a DNA sample known to contain the transgenic sequences). A successful amplification of the positive control demonstrates that the PCR was run under conditions which allow for the amplification of target sequences. A negative, or wild-type, DNA extract control in which the template DNA provided is either genomic DNA prepared from a non-transgenic plant, or is a non-TC1507 transgenic plant, should also be included. Additionally a negative control that contains no template corn DNA extract will be a useful gauge of the reagents and conditions used in the PCR protocol.

Additional DNA primer molecules of sufficient length can be selected from SEQ ID NO: 26 and SEQ ID NO: 27 by those skilled in the art of DNA amplification methods, and conditions optimized for the production of an amplicon that may differ from the methods shown in Table 9 or Table 10 but result in an amplicon diagnostic for event TC1507. The use of these DNA primer sequences with modifications to the methods shown in Table 9 and Table 10 are within the scope of the invention. The amplicon wherein at least one DNA primer molecule of sufficient length derived from SEQ ID NO: 26 and SEQ ID NO: 27 that is diagnostic for event TC1507 is an aspect of the invention. The amplicon wherein at least one DNA primer of sufficient length derived from any of the genetic elements of PHI8999A that is diagnostic for event TC1507 is an aspect of the invention. The assay for the TC1507 amplicon can be performed by using a Stratagene Robocycler, MJ Engine, Perkin-Elmer 9700, or Eppendorf Mastercycler Gradient thermocycler, or by methods and apparatus known to those skilled in the art.

TABLE 9

PCR Conditions: Conditions: Kit used: Perkin-Elmer AmpliTaq Gold kit	
Volume	Component
5 µl	template (10 ng/µl)
4 µl	2 µl each primer (10 µM)
2 µl	10X PCR Gold Buffer
2 µl	25 mM MgCl ₂
2 µl	50X dNTP's (10 mM)

37

TABLE 9-continued

PCR Conditions: Conditions: Kit used: Perkin-Elmer AmpliTaq Gold kit	
Volume	Component
0.1 μ l	Amplitaq Gold Polymerase
4.9 μ l	H ₂ O
20 μ l	Total

Cycling Parameters
GeneAmp $\otimes$ PCR System 9700
9 min 92° C.
30 cycles:
94° C. 30 sec
60° C. 30 sec
72° C. 1 min
7 min 72° C.
Hold 4° C.

TABLE 10

PCR Conditions used with the Advantage $\otimes$ -GC 2 Polymerase Mix: Conditions: Kit used: Advantage $\otimes$ -GC 2 Polymerase Mix	
Volume	Component
5 μ l	template (10 ng/ μ l)
5 μ l	2.5 μ l each primer (10 μ M)
10 μ l	5x GC2 Buffer
10 μ l	GC melt (1.0M final conc.)

38

TABLE 10-continued

PCR Conditions used with the Advantage $\otimes$ -GC 2 Polymerase Mix: Conditions: Kit used: Advantage $\otimes$ -GC 2 Polymerase Mix	
Volume	Component
1.5 μ l	50X dNTP's (10 mM)
1.0 μ l	Advantage GC2 Polymerase
17.5 μ l	H ₂ O
50 μ l	Total

Cycling Parameters
GeneAmp $\otimes$ PCR System 9700
5 min 94° C.
35 cycles:
94° C. 1 min
60° C. 2 min
72° C. 3 min
7 min 72° C.
Hold 4° C.

Having illustrated and described the principles of the present invention, it should be apparent to persons skilled in the art that the invention can be modified in arrangement and detail without departing from such principles. We claim all modifications that are within the spirit and scope of the appended claims.

All publications and published patent documents cited in this specification are incorporated herein by reference to the same extent as if each individual publication or patent application was specifically and individually indicated to be incorporated by reference.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 57

<210> SEQ ID NO 1

<211> LENGTH: 30

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Event specific primer sequence designed for TC1507

<400> SEQUENCE: 1

gtagtactat agattatatt attcgtagag

30

<210> SEQ ID NO 2

<211> LENGTH: 30

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Event specific primer sequence designed for TC1507

<400> SEQUENCE: 2

gccatacaga actcaaaatc ttttccggag

30

<210> SEQ ID NO 3

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Event specific primer sequence designed for TC1507

<400> SEQUENCE: 3

-continued

tgtgggtgtt gtggctctgt cctaa 25

<210> SEQ ID NO 4
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Event specific primer for TC1507

<400> SEQUENCE: 4

agcacctttt cattctttca tatac 25

<210> SEQ ID NO 5
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Genomic DNA primer sequence

<400> SEQUENCE: 5

gacctcccca caggcatgat tgatc 25

<210> SEQ ID NO 6
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer in full length insert, 35S promoter to
pat gene

<400> SEQUENCE: 6

tagaggacct aacagaactc gccgt 25

<210> SEQ ID NO 7
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer in full length insert, 35S promoter to
pat gene

<400> SEQUENCE: 7

gagctggcaa ctcaaaatcc cttt 24

<210> SEQ ID NO 8
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer in full length insert, 35S promoter to
pat gene

<400> SEQUENCE: 8

aaaatcttcg tcaacatggt ggagc 25

<210> SEQ ID NO 9
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer in full length insert, 35S promoter to
pat gene

<400> SEQUENCE: 9

taatctcaac tggctcctc tccgg 25

-continued

<210> SEQ ID NO 10
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer - Zea mays genomic DNA

<400> SEQUENCE: 10

cccctacccc accgacgttt at 22

<210> SEQ ID NO 11
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer - Zea mays genomic DNA

<400> SEQUENCE: 11

ttgattggca ggtccgtggg tc 22

<210> SEQ ID NO 12
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer - Zea mays genomic DNA

<400> SEQUENCE: 12

cacaacggca cagaaacacg aa 22

<210> SEQ ID NO 13
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer - Zea mays genomic DNA

<400> SEQUENCE: 13

gcgcacccac cggaacaaaa tg 22

<210> SEQ ID NO 14
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer - Zea mays genomic DNA

<400> SEQUENCE: 14

tcctcgcat aaatgctcct gc 22

<210> SEQ ID NO 15
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer - Zea mays genomic DNA

<400> SEQUENCE: 15

cctggcacgc attgacgcat gt 22

<210> SEQ ID NO 16
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer - Zea mays genomic DNA

-continued

<400> SEQUENCE: 16

aagccgccac cgtcttgagg acctt

25

<210> SEQ ID NO 17

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Primer - Zea mays genomic DNA

<400> SEQUENCE: 17

gtcgaaccgt gcctccactt gctaa

25

<210> SEQ ID NO 18

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Primer - Zea mays genomic DNA

<400> SEQUENCE: 18

agaaagggaa aatcagggcc aggac

25

<210> SEQ ID NO 19

<211> LENGTH: 22

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Cry1F sequence primer

<400> SEQUENCE: 19

ggctcggact cgacctttct at

22

<210> SEQ ID NO 20

<211> LENGTH: 22

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Cry1F sequence primer

<400> SEQUENCE: 20

gcagttcttg aagaatgagt ga

22

<210> SEQ ID NO 21

<211> LENGTH: 2829

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: 5' flanking sequence of event TC1507

<400> SEQUENCE: 21

actagtttcc tagcccgct cgtgcccta cccaccgac gtttatggaa ggtgccattc

60

cacggttctt cgtggccgcc cctaaggatg taaatggctg gtaaaatccg gtaaatttcc

120

ggtaccgttt accagatttt tccagccgtt ttcggattta tcgggatata cagaaaacga

180

gacggaaacg gaataggttt tttttcgaaa acggtacggt aaacggtgag acaaacttac

240

cgtccgtttt cgtattttctc gggaaactct ggtatattcc cgtatttgtc ccgtattttc

300

ccgacccacg gacctgcaa tcaaccatca gccagtcagc ccatcccccac agctatggcc

360

catggggcca tgttgccac atgccacgc aacgcaaggc agtaaggctg gcagcctggc

420

acgcattgac gcatgtggac acacacagcc gccgcctgtt cgtgtttctg tgccgttgtg

480

cgagactgtg actgcagtg gcggagtcgg cgaacggcga ggcgtctccg gagtctggac

540

-continued

tgcggctgtg gacagcgacg ctgtgacggc gactcggcga agccccaagc taccaagccc	600
ccaagtcccc atccatctct gcttctcttg tcatctcctt cccctggtcg atctgcaggc	660
gccagaccgg ccgaagcatc acgaaacgca ctaagacctc gaaggagtca aaccactcct	720
ccgaggcctc gggggctaca cccggcgggt gcgctcgcgc gcaccacccg gaacaaaatg	780
taaccgagaa aggtcggctc ccttgcaaaa aaagtgcgac aaaagcctcc aagcgagtat	840
taacactcac tttgaggtc gggggctact gtccgggacc ataattagggt gtaccccaaa	900
gactcctaata ctcagctggt aacccccatc agcacaaaag tgcaaggccc tgatgggtgc	960
gattaagtca aggtcgggtc cactcaagggt acacgatctc gcctcgcccc agcccagcct	1020
cgggcaagggt cggccgaccc cgaggattca cgtctcgccc gagggccccc tcaagcgacg	1080
ggcacacctt cggtcgcgcc gagggccatt cttcgccgag aagcaacctt ggccagatcg	1140
ccacaccgac cgaccgtatc gcaggagcat ttaatgcgag gatcgctga cacttatcc	1200
tgacgcgcgc tcttcagtcg acagagccga agtgaccgca atcacttcgc cgctccactg	1260
accgacctga caagaagaca ggcgccctg cgtcgctccg actgctgtgc cactcgacag	1320
agtgaggctg acagcagcca agtcggcct cgggcgccat aggaagctcc gcctcgcccc	1380
accctagggtc tcggactcgg cctcggtcc ggaagacgac gaactacgct tcgcccgcac	1440
ccagggtctg gactcagcct cggtccgga agacgacgaa ttcgcctcg cccgacccca	1500
gggtcggac tcggcctcgg ctccagaaga cgacgaactc cgctcgccc gacccagggt	1560
ctcgactca gcctcggtc cggaagacga cgaactccgc ctgcccgcac cccagggtc	1620
ggactcagcc tcggcctcag acgatgggtc ccgcctcgcc cgacccgggg ctcgactcg	1680
acctttctat cggaccttgt cagatcctgt cttcgtccga ggaggcttg gcaatcctca	1740
ctatgtactc ggtcttaggg gagtgccct tcaacaaact ggtacgaatc acaccgcac	1800
attcaggaac tccgggacca ttgactctct agatgagata ccacctcaag acaacagcgg	1860
cgcaccttgg aatgactact cccatgtgct gaatcatgtt acctttgtgc gctggccagg	1920
tgagatctca ggttcgact catggagagc accaatgttc tcttgacgc atcgtagcgc	1980
tacccccaca aacaccattg atccagagag aatcactcat tcttcaagaa ctgcatact	2040
tgccgagatc ctcatcccta aaggacttg acaatagtat tattggagtc gatacacaa	2100
tcacaaaaaa tacaagaagt cgactagggt gattgggtcc agtgaagaga aaaaaagcc	2160
atacagaact caaaatcttt tccggagata ttcattttcc tgaagaggcg gataagatat	2220
taggtggcag tttgatacca ccagaaagag aaaaaaaga ttctaaggaa tcaaaaaaa	2280
ggaaaaattg ggtttatgtt caacggaaaa aatttctcaa aagcaaggaa aagtattgtg	2340
gctatttate tatccgtgca gctgatatgg ccgcggtttg tgatatcgtt aaccattaca	2400
ttgagacgtc tacagtgaac tttaggacag agccacaaac accacaagag tggattgatg	2460
atctagagag gttcaagat agataccctt ggttggttgc tgagggttag ggtgtgtggt	2520
ctggtattgc ttacgctggg ccttgaagg ctaggaaacc tcaacctcag caaccaacca	2580
atggtatcta tcttgcaacc tctctagatc atcaatccac tcttggtgtg tttgtggctc	2640
tgtcctaaag ttcactgtag acgtctcaat gtaatgggtta acgatatcac aaaccgagag	2700
aagagggtac tcgaagcttc ggccggggcc catcgatata cgcgggcatg cctgcagtgc	2760
agcgtgaccc ggtcgtgccc ctctctagag ataatgagca ttgcatgtct aagttataaa	2820
aaattacca	2829

-continued

```

<210> SEQ ID NO 22
<211> LENGTH: 2346
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: 3' flanking sequence of event TC1507

<400> SEQUENCE: 22

ctcactccgc ttgatcttgg caaagatatt tgacgcattt attagtatgt gttaattttc    60
atttgcagtg cagtattttc tattcgatct ttatgtaatt cggtacaatt aataaatatt    120
caaatcagat tattgactgt catttgtatc aaatcgtgtt taatggatat ttttattata    180
atattgatga tatctcaatc aaaacgtaga taataataat atttatttaa ttttttgcg    240
tcgcacagtg aaaatctata tgagattaca aaataccgac aacattattt aagaaacata    300
gacattaacc ctgagactgt tggacatcaa cgggtagatt ccttcacgca tagcacctca    360
ttcttgggga caaaagcacg gtttggccgt tccattgctg cacgaacgag ctttgcata    420
tcctcggggt ggatcatctc atcagggtcca atcaaatgtg tccaagaact catgttagtc    480
gcaacgaaac cggggcatat gtcgggtatc tcgagctcgc gaaagcttgg ctgcaggtcg    540
acggatcctt caacaaaagg gtacctgtac ccgaaaccga cacagggtggg taggtagaga    600
atacctaggg gcgcgagaca actctctcta aggaactcgg caaaatagcc ccgtaacttc    660
gggagaaggg gtgccccccg ctaacaataa acgaatacgg tttatgtatg gattccggta    720
aaataccggt actcgatttc ataagagtcg aataggaagt taagatgagg gtggtatcat    780
cataaaaatg gagtagtata ctaaattata ctaatccacg tatgatatgt atgcctttcc    840
ttatcaaccg gaagtagtgc aaaaaaatt ctatactgca ctgctctctt ttactgaga    900
aatgcaaaaa aataaaagtg aagtaagggt gcccataga tatttgatct tgccctctgt    960
ccccccccc cttttttcat caaaaatttc catgaaaaaa gaaaagatga atttgtccat    1020
tcattgaacc ctagtctggg actgacgggg ctgcgaaccg cagcttcgcg ctgttcctag    1080
ccttcagggy cccagcgtaa gcaataccag ccacagcacc ctcaacctca gcaaccaacc    1140
aagggtatct atcttgcaac ctctctagat catcaatcca ctcttggtgt gtttgtggct    1200
ctgtcctaaa gttcactgta gacgtctcaa tgtaatggtt aacgatatca caaacgcggg    1260
aacacaagaa cgaaagcacc ttttcattct ttcataact aggggttttt acttgaaaaa    1320
gacaatgttc catactaaag gatagctgca gaagccgcca ccgtcttgag gaccttcggg    1380
ggagccagac cggtcgaacc gtgcctccac ttgctaagga gaaagggaaa atcagggcca    1440
ggacatacga aggaggagcc agaacgaaga tatcctaaga tacttactcg ctccgggcca    1500
tgatcaatca tgctgtggg gaggtctctc gcacctcgat ccatgaaggt accaccgagg    1560
tctgccccgc cgccggcttc ggtaccgtcc tcgccttggg cgcccgaggc acccggggga    1620
tggactgccc aggcgcagcc acgacgaccc aaggatcacc ctctgcgca gtcggcacga    1680
gcaatagttc tcggggaaca ggcagcttgg cctgaactccc cggggtcacc tcaactacct    1740
cggccgaggg gtcaagtacc cctcagtcg gcccccgctc ttcggaacgg gaccccgacg    1800
tccccggccc ggataccgac ggcaccagcc cgctcggggg ctggcttgac gacccctggc    1860
ccagcctcag atctgggctg aggcggaggc aggcggccat gtctgtctct tcatcatcgt    1920
cttcatcctc gtctgtctca tcaggcgtct ccggcgacgg ctcccttggg agccctctcc    1980
tctctgccc acgacgaagc ctttccaagg catcccagac ccacgtccgc tcgtgggccc    2040
gagccttctt tgcgtccttc ttctccttc tcttctccgc ggtgacctc cgcgcagctc    2100

```

-continued

```

ggtcaccgc atcctccggg actggtggca gggaaggctt gtgatgccct acctcctgga 2160
gacagacgaa aagtctcagc tatgagaacc gagggcaatc tgacgcaaga aggaagaagg 2220
agcggatact caccagagac acgcaccgc gatcgggacg cattaagggc tgggaaaaag 2280
tgccggcctc taatttcgct accgtgccgt ccaccacct gtggaggtea tcgatgggaa 2340
ggggaa 2346

```

```

<210> SEQ ID NO 23
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Event specific primer sequence designed for
TC1507

```

```

<400> SEQUENCE: 23

```

```

cttcaaacia gtgtgacaaa 20

```

```

<210> SEQ ID NO 24
<211> LENGTH: 11361
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: The sequence represents the transgenic insert
in maize line TC1507 as well as the sequence flanking
the insertion sites.

```

```

<400> SEQUENCE: 24

```

```

actagtttcc tagcccgct cgtgcccta cccaccgac gtttatggaa ggtgccattc 60
cacggttctt cgtggcgcc cctaaggatg taaatggctg gtaaaatccg gtaaatcttc 120
ggtaccgttt accagatttt tccagccgtt ttcggattta tcgggatata cagaaaacga 180
gacggaaacg gaataggttt tttttcgaaa acggtacggt aaacggtgag acaaacttac 240
cgtccgtttt cgtatttctc gggaaactct ggtatatcc cgtatttgc ccgtattttc 300
ccgacccacg gacctgcaa tcaaccatca gccagtcagc ccacccccc agctatggcc 360
catggggcca tgttggccac atgccacgc aacgcaaggc agtaaggctg gcagcctggc 420
acgcattgac gcattgtggac acacacagcc gccgcctgtt cgtgtttctg tgccgttgtg 480
cgagactgtg actgcgagtg gcggagtcgg cgaacggcga ggcgtctccg gactctggac 540
tgcggtgtg gacagcgacg ctgtgacggc gactcggcga agccccaagc taccaagccc 600
ccaagtcccc atccatctct gcttctctgg tcatctcctt cccctggtcg atctgcaggc 660
gccagaccgg ccgaagcatc acgaaacgca ctaagacctc gaaggagtca aaccactcct 720
ccgaggcctc gggggctaca cccggcggtt gcgctcgcgc gcacccaccg gaacaaaatg 780
taaccgagaa aggtcggtcc ccttgcaaaa aaagtgcgac aaaagcctcc aagcgagtat 840
taacactcac tttgaggctc gggggctact gtcggggacc ataattaggg gtaccccaaa 900
gactcctaata ctacgtggtt aacccccatc agcacaaggc tgcaaaggcc tgatgggtgc 960
gattaagtca aggtcctggt cactcaaggg acacgatctc gcctcgcccc agccagcct 1020
cgggcaaggg cggccgaccc cgaggattca cgtctcgccc gagggcccc tcaagcgacg 1080
ggcacacett cggtcgcgcc gagggccatt cttcgccgag aagcaacett ggccagatcg 1140
ccacaccgac cgaccgtatc gcaggagcat ttaatgcgag gatcgccctga caccctatcc 1200
tgacgcgcgc tcttcagtcg acagagccga agtgaccgca atcacttcgc cgctccactg 1260
accgacctga caagaagaca gcgccgctg cgctcgctccg actgctgtgc cactcgacag 1320

```

-continued

agtgaggctg	acagcagcca	agtcgggctt	cgggcgccat	aggaagctcc	gcctcgccc	1380
accctagggc	tgggactcgg	cctcggtctc	ggaagacgac	gaactacgct	tcgcccagcc	1440
ccagggtctg	gactcagcct	cggtccgga	agacgacgaa	ttccgcctcg	cccgacccca	1500
gggtcggac	tcggcctcgg	ctccagaaga	cgacgaactc	cgctcgccc	gacccaggg	1560
ctcgactca	gcctcggtc	cggaagacga	cgaactccgc	ctcgcccgac	cccagggtc	1620
ggactcagcc	tcggcctcag	acgatggtct	cgcctcgcc	cgacccgggg	ctcgactcg	1680
acctttctat	cggaacctgt	cagatcctgt	cttcgtccga	ggaggctttg	gcaatcctca	1740
ctatgtactc	ggtcttaggg	gagtggcctt	tcaacaaact	ggtacgaatc	acacccgcac	1800
attcaggaac	tccgggacca	ttgactctct	agatgagata	ccacctcaag	acaacagcgg	1860
cgcaccttgg	aatgactact	cccatgtgct	gaatcatggt	acctttgtgc	gctggccagg	1920
tgagatctca	ggttcgact	catggagagc	accaatgttc	tcttggaagc	atcgtagcgc	1980
tacccccaca	aacaccattg	atccagagag	aatcactcat	tcttcaagaa	ctgcatactt	2040
tgcgagatc	ctcatcccta	aaggtagctg	acaatagtat	tattggagtc	gatacacaa	2100
tcacaaaaaa	tacaagaagt	cgactaggtg	gattgggtccg	agtgaagaga	aaaaaaagcc	2160
atacagaact	caaaatcttt	tccggagata	ttcattttcc	tgaagaggcg	gataagatat	2220
taggtggcag	tttgatacca	ccagaagag	aaaaaaaaga	ttctaaggaa	tcaaaaaaaa	2280
ggaaaaattg	ggtttatgtt	caacggaaaa	aattttctca	aagcaaggaa	aagtattgtg	2340
gctatttatc	tatcgtgca	gctgatatgg	cgcgggtttg	tgatatcggt	aaccattaca	2400
ttgagacgtc	tacagtgaac	tttaggacag	agccacaaac	accacaagag	tggtattgatg	2460
atctagagag	ggtgcaagat	agataccctt	ggttggttgc	tgagggttag	ggtggtgtgg	2520
ctggtattgc	ttacgctggg	ccttggaagg	ctaggaaccc	tcaacctcag	caaccaacca	2580
atggtatcta	tcttgcaacc	tctctagatc	atcaatccac	tcttggtgtg	tttgtggctc	2640
tgtcctaaag	ttcactgtag	acgtctcaat	gtaatgggta	acgatatcac	aaaccgagag	2700
aagagggatc	tcgaagcttc	ggcgggggcc	catcgatata	cgcgggcatg	cctgcagtgc	2760
agcgtgaccc	ggtcgtgccc	ctctctagag	ataatgagca	ttgcatgtct	aagtataaaa	2820
aaattaccac	aactggaaga	gcggttacct	ggaccgaagc	ttcgccgggg	gcccacgat	2880
atccgcgggc	atgcctgcag	tgacgcgtga	cccgtctgtg	cccctctcta	gagataatga	2940
gcattgcagt	tctaagttat	aaaaaattac	cacatatatt	ttttgtcaca	cttggttgaa	3000
gtgcagttta	tctatcttta	tacatatatt	taaactttac	tctacgaata	atataatcta	3060
tagtactaca	ataatatcag	tgttttagag	aatcatataa	atgaacagtt	agacatggtc	3120
taaaggacaa	ttgagtattt	tgacaacagg	actctacagt	tttatctttt	tagtgtgcat	3180
gtgttctcct	ttttttttgc	aaatagcttc	acctatataa	tacttcatcc	attttattag	3240
tacatccatt	tagggttttag	ggttaatggg	ttttatagac	taattttttt	agtacatcta	3300
ttttattcta	tttttagctc	taaattaaga	aaactaaaac	tctatttttag	tttttttatt	3360
taataattta	gatataaaat	agaataaaat	aaagtgacta	aaaattaaac	aaataccctt	3420
taagaaatta	aaaaaactaa	ggaacattt	ttcttgtttc	gagtagataa	tgccagcctg	3480
ttaaacgccg	tcgacgagtc	taacggacac	caaccagcga	accagcagcg	tcgctcgagg	3540
ccaagcgaag	cagacggcac	ggcatctctg	tcgctgcctc	tggacccctc	tcgagagttc	3600
cgtccaccgg	ttggacttgc	tccgctgtcg	gcataccagaa	attgcgtggc	ggagcggcag	3660

-continued

acgtgagccg	gcacggcagg	cggcctcctc	ctcctctcac	ggcacggcag	ctacggggga	3720
ttcctttccc	accgctcctt	cgctttccct	tctcgcgccg	ccgtaataaa	tagacacccc	3780
ctccacaccc	tctttcccca	acctcgtgtt	gttcgggagc	cacacacaca	caaccagatc	3840
tcccccaaat	ccacccgctg	gcacctccgc	ttcaaggtag	gccgctcgtc	ctcccccccc	3900
ccccctctct	accttctcta	gatcggcggt	cgggtccatg	gttagggccc	ggtagttcta	3960
cttctgttca	tgtttgtgtt	agatccgtgt	ttgtgttaga	tccgtgctgc	tagcgttcgt	4020
acacggatgc	gacctgtacg	tcagacacgt	tctgattgct	aacttgccag	tgtttctctt	4080
tggggaatcc	tgggatggct	ctagccgttc	cgcagacggg	atcgatttca	tgattttttt	4140
tgtttcgttg	catagggttt	ggtttgcctt	tttctttat	ttcaatatat	gccgtgcact	4200
tgtttgctcg	gtcatctttt	catgcttttt	tttgtcttgg	ttgtgatgat	gtggtctggt	4260
tgggcggtcg	ttctagatcg	gagtagaatt	ctgtttcaaa	ctacctgggtg	gattttattaa	4320
ttttggatct	gtatgtgtgt	gccatacata	ttcatagtta	cgaattgaag	atgatggatg	4380
gaaatatcga	tctaggatag	gtatacatgt	tgatgcgggt	tttactgatg	catatacaga	4440
gatgcttttt	gttcgcttgg	ttgtgatgat	gtggtgtggt	tgggcggtcg	ttcatcgtt	4500
ctagatcgga	gtagaatact	gtttcaaact	acctgggtga	tttattaatt	ttggaactgt	4560
atgtgtgtgt	catacatctt	catagttacg	agttaaagat	ggatggaaat	atcgatctag	4620
gatagggtata	catgttgatg	tgggtttttac	tgatgcatat	acatgatggc	atatgcagca	4680
tctattcata	tgctctaacc	ttgagtacct	atctattata	ataaacaagt	atgttttata	4740
attattttga	tcttgatata	cttgatgat	ggcatatgca	gcagctatat	gtggattttt	4800
ttagccctgc	cttcatacgc	tattttattg	cttggtactg	tttcttttgt	cgatgctcac	4860
cctgttgttt	ggtgttactt	ctgcaggctg	actctagagg	atccaacaat	ggagaacaac	4920
atacagaatc	agtgcgctcc	ctacaactgc	ctcaacaatc	ctgaagtaga	gattctcaac	4980
gaagagaggt	cgactggcag	attgccgtta	gacatctccc	tgctccctac	acgtttcctg	5040
ttgtctgagt	ttgttccagg	tgtgggagtt	gcgtttggcc	tcttcgacct	catctggggc	5100
ttcatcactc	catctgattg	gagcctcttt	cttctccaga	ttgaacagtt	gattgaacaa	5160
aggattgaga	ccttggaag	gaatcgggcc	atcactaccc	ttcgtggcct	agcagacagc	5220
tatgatagct	acattgaagc	actaagagag	tgggaagcca	atcctaacaa	tgcccaactg	5280
agagaagatg	tgcgatatcg	ctttgctaac	acagatgatg	ctttgatcac	agccatcaac	5340
aacttcaccc	ttaccagctt	cgagatccct	cttctctcgg	tctatgttca	agctgctaac	5400
ctgcacttgt	cactactgcg	cgacgctgtg	tcgtttgggc	aaggttgggg	actggacata	5460
gctactgtca	acaatcacta	caacagactc	atcaatctga	ttcatcgata	cacgaaacat	5520
tgtttgata	cctacaatca	gggattggag	aaactgagag	gtactaacac	tcgccaatgg	5580
gccagggtta	atcagttcag	gagagacctt	acacttactg	tgtagacat	agttgctctc	5640
tttccgaact	acgatgttcg	tacctatccg	attcaaacgt	catcccaact	tacaaggag	5700
atctacacca	gttcagtcct	tgaagactct	ccagtttctg	cgaacatacc	caatggtttc	5760
aacagggctg	agtttgaggt	cagaccaccc	catctcatgg	acttcatgaa	ctctttgttt	5820
gtgactgcag	agactgttag	atcccaaaact	gtgtggggag	gacacttagt	tagctcacgc	5880
aacacggctg	gcaatcgtat	caactttcct	agttacgggg	tcttcaatcc	cgggggcgcc	5940
atctggattg	cagatgaaga	tccacgtcct	ttctatcgga	ccttgtcaga	tcctgtcttc	6000
gtccgaggag	gctttggcaa	tcctcactat	gtactcggtc	ttaggggagt	ggcctttcaa	6060

-continued

caaactggta cgaatcacac ccgcacattc aggaactccg ggaccattga ctctctagat	6120
gagataccac ctcaagacaa cagcggcgca ccttggaatg actactccca tgtgctgaat	6180
catgttacct ttgtgcgctg gccaggtgag atctcaggtt ccgactcatg gagagcacca	6240
atgttctctt ggacgcacg tagcgctacc cccacaaaca ccattgatcc agagagaatc	6300
actcagattc ccttggtgaa ggcacacaca cttcagtcag gaactacagt tgtaagaggg	6360
ccgggggttca cgggaggaga cattcttcga cgcactagtg gaggaccatt cgcgtacacc	6420
attgtcaaca tcaatgggca acttccccaa aggtatcgtg ccaggatagc ctatgcctct	6480
actaccaatc taagaatcta cgttacggtt gcaggtgaac ggatctttgc tggtcagttc	6540
aacaagacaa tggataccgg tgatccactt acattccaat cttctctcta cgccactatc	6600
aacaccgcgt tcacctttcc aatgagccag agcagtttca cagtaggtgc tgataccttc	6660
agttcaggca acgaagtgtta cattgacagg tttagattga ttccagttac tgccacactc	6720
gagtaaggat ccgtcgacct gcagccaagc ttctcgcgagc tcgagatccc cgacatatgc	6780
cccggtttcg ttgcgactaa catgagttct tggacaaatt tgattggacc tgatgagatg	6840
atccaacccg aggatatagc aaagctcgtt cgtgcagcaa tggaaaggcc aaaccgtgct	6900
tttgtcccca agaatgaggt gctatgcatg aaggaatcta cccgttgatg tccaacagtc	6960
tcagggttaa tgtctatgta tcttaataa tgtgtgcggt attttgaat ctcatataga	7020
ttttcactgt gcgacgcaaa aatattaaat aaatattatt attatctacg ttttgattga	7080
gatatcatca atattataat aaaaatatcc attaaacacg atttgataca aatgacagtc	7140
aataatctga tttgaatatt tattaattgt aacgaattac ataaagatcg aatagaaaat	7200
actgcactgc aaatgaaaat taacacatac taataaatgc gtcaaatac tttgccaaga	7260
tcaagcggag tgagggcctc atatccggtc tcagttacaa gcacgggtac cccgaagcgc	7320
gtccaccaaa tgccctcgac atagatgccg ggctcgacgc tgaggacatt gcctaccttg	7380
agcatggtct cagcgccggc tttaagctca atcccatccc aatctgaata tcctatcccg	7440
cgcccagtc cgtgtaagaa cgggtctgtc catccacctc tgttgggaat tccggtcagg	7500
gtcacctttg tccaccaaga tggaaactgc gccgcggacc gaattcccat ggagtcaaag	7560
attcaaatag aggacctaac agaactcgcc gtaaagactg gcgaacagtt catacagagt	7620
ctcttacgac tcaatgacaa gaagaaaaat ttctgcaaca tgggtggagca cgacacgctt	7680
gtctactcca aaaatatcaa agatacagtc tcagaagacc aaagggcaat tgagactttt	7740
caacaaaggg taatatccgg aaacctctc ggattccatt gccagctat ctgtcacttt	7800
attgtgaaga tagtgaaaa ggaaggtggc tcctacaaat gccatcattg cgataaagga	7860
aaggccatcg ttgaagatgc ctctgccgac agtgggtcca aagatggacc cccacccacg	7920
aggagcatcg tggaaaaaga agacgttcca accacgtctt caaagcaagt ggattgatgt	7980
gatatctcca ctgacgtaag ggatgacgca caatccccact atccttcgca agacccttcc	8040
tctatataag gaagttcatt tcatttgag aggacagggg acccggggat ccaccatgtc	8100
tccggagagg agaccagttg agattaggcc agctacagca gctgatatgg ccgcggtttg	8160
tgatcctgtt aaccattaca ttgagacgtc tacagtgaac tttaggacag agccacaaac	8220
accacaagag tggattgatg atctagagag gttgcaagat agataccctt ggttggttgc	8280
tgaggttgag ggtgttgtgg ctggtattgc ttacgctggg ccctggaagg ctaggaaacg	8340
ttacgattgg acagttgaga gtactgttta cgtgtcacat aggcatacaa ggttgggcct	8400

-continued

aggatccaca ttgtacacac atttgcttaa gtctatggag gcgcaagggt ttaagtctgt	8460
ggttgctgtt ataggccttc caaacgatcc atctgttagg ttgcatgagg ctttgggata	8520
cacagcccggt ggtacattgc gcgcagctgg atacaagcat ggtggatggc atgatgttg	8580
tttttgcaa agggattttg agttgccagc tcctccaagg ccagttaggc cagttacca	8640
gatctgagtc gacctgcagg catgcccgt gaaatcacca gtctctctct acaaactctat	8700
ctctctctat aataatgtgt gagtagttcc cagataaggg aattagggtt cttatagggt	8760
ttcgctcatg tgttgagcat ataagaaacc cttagtatgt atttgattt gtaaaatact	8820
tctatcaata aaatttctaa ttctaaaaac caaaatccag tggcgagctc gaattcgagc	8880
tcgagcccggt gtggatctc tagagtcgac ctgcagaagc ttcggtccgg cgcgcctcta	8940
gttgaagaca cgttcatgtc ttcatcgtaa gaagacactc agtagtcttc ggccagaatg	9000
gcctaactca aggcctctac tccgcttgat cttggcaaag atatttgacg catttattag	9060
tatgtgttaa ttttcatttg cagtgcagta ttttctattc gatctttatg taattcggtta	9120
caattaataa atattcaaat cagattattg actgtcattt gtatcaaatc gtgtttaatg	9180
gatatcttta ttataatatt gatgatattc caatcaaac gtagataata ataatactta	9240
tttaataatt ttgcgtcgca cagtgaatct ctatatgaga ttacaaaata ccgacaacat	9300
tatttaagaa acatagacat taacctgag actgttgac atcaacgggt agattccttc	9360
atgcatagca cctcattctt ggggacaaaa gcacgggttg gccgttccat tgctgcacga	9420
acgagctttg ctatatctc ggggttgatc atctcatcag gtccaatcaa atttgtccaa	9480
gaactcatgt tagtcgcaac gaaaccgggg catatgtcgg gtatctcgag ctgcgaaag	9540
cttggtcgca ggtcgacgga tccttcaaca aaagggtagc tgtaccgaa accgacacag	9600
gtgggttagt agagaatacc taggggcgcg agacaactct ctctaaggaa ctcgcaaaa	9660
tagccccgta acttcgggag aaggggtgcc ccccgctaac aataaacgaa tacggtttat	9720
gtatggattc cggtaaaata ccggtactcg atttcataag agtcgaatag gaagttaaga	9780
tgagggtggt atcatcataa aaatggagta gtatcctaaa ttataactaat ccacgtatga	9840
tatgtatgcc tttccttacc aaccggaagt agtgcaaaaa aaattctata ctgcaactgt	9900
ctctttttac tgagaaatgc aaaaaataa aagtgaagta aggggtcccc atagatattt	9960
gatcttgctt cctgtcccc cccccctttt ttcatacaaa atttccatga aaaaagaaaa	10020
gatgaatttg tccattcatt gaaccctagt tcgggactga cggggctcga acccgagct	10080
tcgcctgtt cctagccttc cagggcccg cgtaagcaat accagccaca gcaccctcaa	10140
cctcagcaac caaccaagggt tatctatctt gcaacctctc tagatcatca atccactctt	10200
gtggtgtttg tggctctgtc ctaaagtcca ctgtagacgt ctcaatgtaa tggttaacga	10260
tatcacaaac cgcggaacac aagaacgaaa gcacctttt attctttcat atactagggg	10320
tttttacttg gaaaagacaa tgttcatac taaaggatag ctgcagaagc cgccaccgtc	10380
ttgaggacct tccggggagc cagaccggtc gaaccgtgcc tccacttgct aaggagaaag	10440
ggaaaatcag ggccaggaca tacgaaggag gagccagaac gaagatatcc taagatactt	10500
actcgctccg ggccatgac aatcatgcct gtggggaggt ctctcgacc tcgatccatg	10560
aaggtagcac cgaggctgc cccgcgcgcg gcttcgggtac cgtcctcgcc ttgggcgcgc	10620
gaggcaccgg gggatggac tgcccaggcg cagccacgac gacccaagga tcaccctcct	10680
gcgcagtcgg cacgagcaat agttctcggg gaacaggcag cttggcctga ctccccgggg	10740
tcacctcaac tacctcgcc gaggggtcaa gtacccctc agtcgcgcgc cgtctctcgg	10800

-continued

```

accgggaccc cgacgtcccg gccccggata cgcacggcac cagcccgcgc gggggctggc 10860
ttgacgaccc ctggcccagc ctcagatctg ggctgaggcc gaggcaggcg gccatgtcgt 10920
cgtcttcacg atcgtcttca tcacgtcgtc cgtcatcagg cgtctccggc gacggctccc 10980
ttgggagccc ctcctctccc tgcgcacgac gaagccttcc caaggcatcc cgagcccacg 11040
tccgctcgtg ggcccagacc ttctttgctg ccttcttctc cttcctcttc tccgcgggtga 11100
ccctccgcgc agctcgggtc accgcaccc cggggactgg tggcagggaa ggcttgtgat 11160
gccctacccc ctggagacag acgaaaagtc tcagctatga gaaccgaggg caatctgacg 11220
caagaaggaa gaaggagcgg atactcacca gagacacgca cccgcgatcg ggacgcatta 11280
agggctggga aaaagtgcgc gcctctaatt tcgctaccgt gccgtccacc cacctgtgga 11340
ggtcacgatg gggaagggga a                                     11361

```

<210> SEQ ID NO 25

<211> LENGTH: 6186

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: The sequence represents the DNA molecule used to transform maize line TC1507. This sequence represents insert PHI8999A.

<400> SEQUENCE: 25

```

caactggaag agcgggtacc cggaccgaag cttcggcccg ggcccacga tatccgcggg 60
catgcctgca gtgcagcgtg acccggtcgt gccctctct agagataatg agcattgcat 120
gtctaagtta taaaaatta ccacatattt tttttgtcac acttgtttga agtgcagttt 180
atctatcttt atacatatat ttaacttta ctctacgaat aatataatct atagtactac 240
aataatatca gtgttttaga gaatcatata aatgaacagt tagacatggt ctaaaggaca 300
attgagtatt ttgacaacag gactctacag ttttatcttt ttagtgtgca tgtgttctcc 360
tttttttttg caaatagctt cacctatata atacttcac cattttatta gtacatccat 420
ttagggttta ggggtaatgg tttttataga ctaatttttt tagtacatct attttattct 480
attttagcct ctaaatgaag aaaactaaaa ctctatttta gtttttttat ttaataattt 540
agataaaaa tagaataaaa taaagtgact aaaaattaaa caaataccct ttaagaaatt 600
aaaaaaacta aggaacatt tttctgtgtt cgagtagata atgccagcct gttaaacgcc 660
gtcgacgagt ctaacggaca ccaaccagcg aaccagcagc gtcgcgtcgg gccaagcgaa 720
gcagacggca cggcatctct gtcgctgcct ctggaccctc ctcgagagtt ccgctccacc 780
gttggaactg ctccgctgtc ggcattccga aattgcgtgg cggagcggca gacgtgagcc 840
ggcacggcag ggggcctcct cctcctctca cggcacggca gctacggggg attcctttcc 900
caccgtcctc tcgctttccc ttctcgcgcc gccgtaataa atagacaccc cctccacacc 960
ctctttcccc aacctcgtgt tgttcggagc gcacacacac acaaccagat ctcccccaaa 1020
tccaccgcgc ggcacctccg cttcaaggta cgcgcgtcgt cctccccccc cccccctctc 1080
taccttctct agatcggcgt tccgggccat ggtagggcc cggtagttct acttctgttc 1140
atgtttgtgt tagatccgtg tttgtgttag atccgtgctg ctacggttcg tacacggatg 1200
cgacctgtac gtcagacacg ttctgattgc taacttgcca gtgtttctct ttggggaatc 1260
ctgggatggc tctagccgtt ccgcagacgg gatcgatttc atgatttttt ttgtttcgtt 1320
gcatagggtt tggtttgccc ttttcttta tttcaatata tgccgtgcac ttgtttgtcg 1380

```

-continued

ggatcatcttt	tcatgctttt	ttttgtcttg	gttgtgatga	tgtgggtctgg	ttgggcgggc	1440
gttctagatc	ggagtagaat	tctgtttcaa	actacctggt	ggattttatta	atthttggatc	1500
tgtatgtgtg	tgccatacat	attcatagtt	acgaattgaa	gatgatggat	ggaaatatcg	1560
atctaggata	ggtatacatg	ttgatgcggg	ttttactgat	gcatatacag	agatgctttt	1620
tgttcgcttg	gttgtgatga	tgtgggtgtg	ttgggcgggc	gttcattcgt	tctagatcgg	1680
agtagaatc	tgthttcaac	tacctggtgt	atthttaaat	tttggaaactg	tatgtgtgtg	1740
tcatacatct	tcatagttac	gagthtaaga	tggatggaaa	tatcgatcta	ggataggtat	1800
acatgttgat	gtgggtthta	ctgatgcata	tacatgatgg	catatgcagc	atctattcat	1860
atgctctaac	cttgagtacc	tatctattat	aataaacaag	tatgtthtat	aattatthtg	1920
atcttgatat	acttggatga	tggcatatgc	agcagctata	tgtggatttt	tttagccttg	1980
ccttcatacg	ctatthtttt	gcttggtact	gtthctthtg	togatgctca	ccctgttggt	2040
tgggtgttact	tctgcaggtc	gactctagag	gatccaacaa	tggagaacaa	catacagaat	2100
cagtgcgtcc	cctacaactg	cctcaacaat	cctgaagtag	agattctcaa	cgaagagagg	2160
tcgactggca	gattgcggtt	agacatctcc	ctgtccctta	cacgtthctt	gttgtctgag	2220
thtgttccag	gtgtgggagt	tgcgtthggc	ctcttcgacc	tcatctgggg	cttcactact	2280
ccatctgatt	ggagcctctt	tcttctccag	attgaacagt	tgattgaaca	aaggattgag	2340
accttggaac	ggaatcgggc	catcactacc	cttcgtggct	tagcagacag	ctatgagatc	2400
tacattgaag	cactaagaga	gtgggaagcc	aatcctaaca	atgccaact	gagagaagat	2460
gtgcgtatac	gctthgtctaa	cacagatgat	gctthgatca	cagccatcaa	caacttcacc	2520
cttaccagct	tcgagatccc	tcttctctcg	gtctatgttc	aagctgctaa	cctgcacttg	2580
tcactactgc	gcgacgtgtg	gtcgtthggg	caaggthggg	gactggacat	agctactgtc	2640
aacaatcact	acaacagact	catcaatctg	attcatcgat	acacgaacaa	ttgtthggat	2700
acctacaatc	agggattgga	gaacctgaga	ggtactaaca	ctcgccaatg	ggccagggtc	2760
aatcagthta	ggagagacct	tacacttact	gtgttagaca	tagthgtctt	ctthccgaac	2820
tacgatgttc	gtacctatcc	gattcaaacg	tcattcccaac	ttacaaggga	gatctacacc	2880
agthcagtca	ttgaagactc	tccagthttct	gcgaacatac	ccaatggtht	caacagggtc	2940
gagthttggag	tcagaccacc	ccatctcatg	gacttcatga	actctthgtt	tgtgactgca	3000
gagactgtta	gatcccaaac	tgtgtgggga	ggacacttag	ttagctcacg	caacacggct	3060
ggcaatcgta	tcaactthtc	tagttacggg	gtcttcaatc	ccgggggcgc	catctggatt	3120
gcagatgaag	atccacgtcc	thctatcgg	acctgtcag	atcctgtctt	cgtccgagga	3180
ggctthtgca	atcctcacta	tgtactcgg	cttaggggag	tggcctthca	acaaactggt	3240
acgaatcaca	ccgcacatt	caggaaactc	gggaccattg	actctctaga	tgagatacca	3300
cctcaagaca	acagcggcgc	accttggaat	gactactccc	atgtgtgtaa	tcatgttacc	3360
thtgtgcgt	ggccagggtga	gatctcagg	tccgactcat	ggagagcacc	aatgttctct	3420
tggacgcato	gtagcgttac	ccccacaaac	accttgatc	cagagagaat	cactcagatt	3480
cccttggtga	aggcacacac	acttcagtca	ggaactacag	ttgtaagagg	gccgggggtc	3540
acgggaggag	acattctctg	acgcactagt	ggaggaccat	tcgcgtacac	cattgtcaac	3600
atcaatgggc	aacttcccca	aaggatcgt	gccaggatac	gctatgcctc	tactaccaat	3660
ctaagaatct	acgttacgg	tgcaggtgaa	cggatctthg	ctggtcagtt	caacaagaca	3720
atggataccg	gtgatccact	tacattccaa	tctthctcct	acgccactat	caacaccgcg	3780

-continued

ttcacctttc caatgagcca gagcagtttc acagtaggtg ctgatacctt cagttcaggc	3840
aacgaagtgt acattgacag gtttgagttg attccagtta ctgccacact cgagtaagga	3900
tccgtcgacc tgcagccaag cttttcgca gctcgagatc cccgacatat gccccggtt	3960
cgttcgact aacatgagtt cttggacaaa tttgattgga cctgatgaga tgatccaacc	4020
cgaggatata gcaaagctcg ttcgtgcagc aatggaacgg ccaaaccgtg cttttgtccc	4080
caagaatgag gtgctatgca tgaaggaatc tacccgttga tgtccaacag tctcagggtt	4140
aatgtctatg tatcttaaat aatgtgtcg gtattttgta atctcatata gattttcact	4200
gtgcgacgca aaaatattaa ataaatatta ttattatcta cgttttgatt gagatatcat	4260
caatattata ataaaaatat ccattaaaca cgatttgata caaatgacag tcaataatct	4320
gatttgaata tttattaatt gtaacgaatt acataaagat cgaatagaaa atactgcact	4380
gcaaatgaaa attaacacat actaataaat gcgtcaaata tctttgcaa gatcaagcgg	4440
agtgaaggcc tcatatccgg tctcagttac aagcacggta tcccgaagc gcgctccacc	4500
aatgcctcg acatagatgc cgggctcgac gctgaggaca ttgcctacct tgagcatggt	4560
ctcagcgccg gctttaagct caatccatc ccaatctgaa tatcctatcc cgcgcccagt	4620
cgggtgtaag aacgggtctg tccatccacc tctgttgga attccggtcc gggtcacctt	4680
tgtccaccaa gatggaactg cggccgcgga ccgaattccc atggagtcaa agattcaaat	4740
agaggaccta acagaactcg ccgtaaagac tggcgaacag ttcatacaga gtctcttacg	4800
actcaatgac aagaagaaaa tcttcgtcaa catggtggag cagcacgc ttgtctactc	4860
caaaaatata aaagatacag tctcagaaga ccaaagggca attgagactt tcaacaaaag	4920
ggtaatatcc ggaaacctcc tcggattcca ttgccagct atctgtcact ttattgtgaa	4980
gatagtggaa aaggaagggtg gctcctacaa atgccatcat tgcgataaag gaaaggccat	5040
cgttgaagat gcctctgcg acagtgggtcc caaagatgga cccccacca cgaggagcat	5100
cgtggaaaaa gaagacgttc caaccacgtc ttcaaagcaa gtggattgat gtgatcttc	5160
cactgacgta agggatgacg cacaatccca ctatccttcg caagaccctt cctctatata	5220
aggaagtta tttcatttgg agaggacagg gtaccgggg atccaccatg tctccggaga	5280
ggagaccagt tgagattagg ccagctacag cagctgatat ggccgcggtt tgtgatatcg	5340
ttaaccatta cattgagacg tctacagtga actttaggac agagccacaa acaccacaag	5400
agtggattga tgatctagag aggttgcaag atagataccc ttggttggtt gctgagggtg	5460
aggggtgtgt ggctggtatt gcttacgctg ggccctggaa ggctaggaa gcttacgatt	5520
ggacagttga gactactgtt tacgtgtcac ataggcatca aaggttgggc ctaggatcca	5580
cattgtacac acatttgctt aagtctatgg aggcgcaagg ttttaagtct gtggttgctg	5640
ttataggcct tccaaacgat ccattctgta ggttgcatga ggctttggga tacacagccc	5700
gggttacatt gcgcgcagct ggatacaagc atgggtggatg gcatgatgtt ggtttttggc	5760
aaagggattt tgagttgcca gctcctccaa ggccagttag gccagttacc cagatctgag	5820
tcgacctgca ggcatgccgc tgaatcacc agtctctctc taaaaatcta tctctctcta	5880
taataatgtg tgagtagttc ccagataagg gaattagggt tcttataggg ttctgctcat	5940
gtgttgagca tataagaaac ccttagtatg tatttgtatt tgtaaaatac ttctatcaat	6000
aaaatttcta attcctaaaa ccaaaatcca gtggcgagct cgaattcgag ctcgagcccc	6060
ggtggatcct ctagagtoga cctgcagaag cttcggtccg gcgcgcctct agttgaagac	6120

-continued

```
acgttcacgt cttcatcgta agaagacact cagtagtctt cggccagaat ggctaactc 6180
aaggcc 6186
```

```
<210> SEQ ID NO 26
<211> LENGTH: 3830
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Sequence that represents part of the PHI8999A
insert as well as flanking sequence 5' to the
insert.
```

```
<400> SEQUENCE: 26
```

```
actagtttcc tagcccggt cgtgcccta cccaccgac gtttatggaa ggtgccattc 60
cacggttctt cgtggccgcc cctaaggatg taaatggctg gtaaaatccg gtaaatttcc 120
ggtaccgttt accagatttt tccagccgtt ttcggattta tcgggatata cagaaaacga 180
gacggaaacg gaataggttt tttttcgaaa acggtacggt aaacggtgag aaaaacttac 240
cgtccgtttt cgtattttctc gggaaactct ggtatattcc cgtatttgct ccgtattttc 300
ccgacccacg gacctgcaa tcaacatca gccagtcagc ccacccccc agctatggcc 360
catggggcca tgttgccac atgccacgc aacgcaaggc agtaaggctg gcagcctggc 420
acgcattgac gcattgtggc acacacagcc gccgcctgtt cgtgtttctg tgccgttgtg 480
cgagactgtg actgcagtg gcggagtcgg cgaacggcga ggcgtctccg gactctggac 540
tcggctgtg gacagcgac ctgtgacggc gactcggcga agccccaagc taccaagccc 600
ccaagtcccc atccatctct gcttctctgg tcatctcctt cccctggctg atctgcaggc 660
gccagaccgg ccgaagcatc acgaaacgca ctaagacctc gaaggagtca aaccactcct 720
ccgaggcctc gggggctaca cccggcgggt gcgctcgcgc gcacccaccg gaacaaaatg 780
taaccgagaa aggtcggctc ccttgcaaaa aaagtgcgac aaaagcctcc aagcgagtat 840
taacactcac tttgagctc gggggctact gtcggggacc ataattaggg gtaccccaaa 900
gactccta atcagctggt aacccccatc agcacaaaagc tgcaaaaggc tgatgggtgc 960
gattaagtca aggtcggctc cactcaaggc acacgatctc gcctcgccc agcccagcct 1020
cgggcaaggc cggccgaccc cgaggattca cgtctcgccc gagggcccc tcaagcgacg 1080
ggcacacctt cggtcgcgcc gagggccatt cttcgccgag aagcaacctt ggccagatcg 1140
ccacaccgac cgaccgtatc gcaggagcat ttaatgcgag gatcgctga caccttatcc 1200
tgacgcgcgc tcttcagtcg acagagccga agtgaccga atcacttcgc cgtccactg 1260
accgacctga caagaagaca gcgccgctg cgtcgctccg actgctgtgc cactcgacag 1320
agtgaggctg acagcagcca agtcggcct cgggcgccat aggaagctcc gcctcgccc 1380
accctagggc tcggactcgg cctcgctcc ggaagacgac gaactacgct tcgcccagcc 1440
ccagggttg gactcagcct cggtccgga agacgacgaa ttccgctcg cccgacccca 1500
gggctcggac tcggcctcgg ctccagaaga cgacgaactc cgctcgccc gaccccaggg 1560
ctcgactca gcctcgctc cggaagacga cgaactccgc ctgcccgcac cccagggtc 1620
ggactcagcc tcggcctcag acgatggtct ccgcctcgcc cgacccgggg ctcgactcg 1680
acctttctat cggacctgtg cagatcctgt cttcgctcca ggaggcttg gcaatcctca 1740
ctatgtactc ggtcttaggg gactggcctt tcaacaaact ggtacgaatc acaccgcac 1800
attcaggaac tccgggacca ttgactctct agatgagata ccacctcaag acaacagcgg 1860
cgacaccttg aatgactact cccatgtgct gaatcatgtt acctttgtgc gctggccagg 1920
```

-continued

tgagatctca ggttccgact catggagagc accaatgttc tcttgagcgc atcgtagcgc	1980
tacccccaca aacaccattg atccagagag aatcactcat tcttcaagaa ctgcatactt	2040
tgccgagatc ctcaccccta aaggctactg acaatagtat tattggagtc gatacacaa	2100
tcacaaaaaa tacaagaagt cgactagggtg gattgggtccg agtgaagaga aaaaaaagcc	2160
atacagaact caaaatcttt tccggagata ttcattttcc tgaagaggcg gataagatat	2220
taggtggcag tttgatacca ccagaaagag aaaaaaaga ttctaaggaa tcaaaaaaa	2280
ggaaaaattg ggtttatgtt caacggaaaa aatttctcaa aagcaaggaa aagtattgtg	2340
gctatttate tatccgtgca gctgatatgg ccgcggtttg tgatatacgtt aaccattaca	2400
ttgagacgtc tacagtgaac tttaggacag agccacaaac accacaagag tggattgatg	2460
atctagagag gttgcaagat agataccctt ggttgggttc tgagggtgag ggtgtgtg	2520
ctggtattgc ttacgtggg ccttgaaggt ctaggaaccc tcaacctcag caaccaacca	2580
atggtatcta tcttgcaacc tctctagatc atcaatccac tcttggtgtg tttgtggctc	2640
tgctctaaag ttcactgtag acgtctcaat gtaatggta acgatatcac aaaccgagag	2700
aagaggatc tcgaagcttc ggccggggcc catcgatata cgccggcatg cctgcagtgc	2760
agcgtgaccc ggtcgtgccc ctctctagag ataatgagca ttgcatgtct aagttataaa	2820
aaattaccac aactggaaga gcgggttacc ggaccgaagc ttcggccggg gcccatcgat	2880
atcccggggc atgcctgcag tgcagcgtga cccggtcgtg cccctctcta gagataatga	2940
gcattgcagt tctaagttat aaaaaattac cacatatatt tttgtgcaca cttgtttgaa	3000
gtgcagttta tctatcttta tacatatatt taaactttac tctacgaata atataatcta	3060
tagtactaca ataatacag tgttttagag aatcatataa atgaacagtt agacatggtc	3120
taaaggacaa ttgagtattt tgacaacagg actctacagt tttatcttt tagtgtgcat	3180
gtgttctcct ttttttttgc aaatagcttc acctatataa tacttcatcc attttattag	3240
tacatccatt tagggtttag ggtaaatggt ttttatagac taatttttt agtacatcta	3300
ttttattcta ttttagctc taaattaaga aaactaaaac tctatttttag tttttttatt	3360
taataattta gatataaaat agaataaaat aaagtgacta aaaattaaac aaataccctt	3420
taagaaatta aaaaaactaa ggaaacattt ttcttgtttc gagtagataa tgccagcctg	3480
ttaaacgccg tcgacgagtc taacggacac caaccagcga accagcagcg tcgctcggg	3540
ccaagcgaag cagacggcac ggcactctctg tcgctgcctc tggaccctc tcgagagttc	3600
cgctccaccg ttggacttgc tccgtgtctg gcattccagaa attgcgtggc ggagcggcag	3660
acgtgagccg gcacggcagg cggcctctc ctctctcac ggcacggcag ctacggggga	3720
ttctttccc accgctcctt cgttttccct tctcgcctc ccgtaataaa tagacacccc	3780
ctccacccc tctttcccca acctcgtgtt gtccggagcg cacacacaca	3830

<210> SEQ ID NO 27

<211> LENGTH: 3347

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Sequence that represents part of the PHI8999A
insert as well as flanking sequence 3' to the
insert.

<400> SEQUENCE: 27

cccactatcc ttcgcaagac ccttctctta tataaggaag ttcatttcat ttggagagga	60
---	----

-continued

cagggtagcc	ggggatccac	catgtctccg	gagaggagac	cagttgagat	taggccagct	120
acagcagctg	atatggccgc	ggtttgtgat	atcgtaaac	attacattga	gacgtctaca	180
gtgaacttta	ggacagagcc	acaaacacca	caagagtgga	ttgatgatct	agagagggtg	240
caagatagat	acccttggtt	ggttgctgag	gttgaggggtg	ttgtggctgg	tattgcttac	300
gctgggccct	ggaaggctag	gaacgcttac	gattggacag	ttgagagtac	tgtttacgtg	360
tcacataggc	atcaaagggt	gggcctagga	tccacattgt	acacacattt	gcttaagtct	420
atggaggcgc	aagggtttta	gtctgtggtt	gctgttatag	gccttccaaa	cgatccatct	480
gttaggttgc	atgaggcttt	gggatacaca	gcccggggta	cattgcgcgc	agctggatac	540
aagcatgggt	gatggcatga	tgttggtttt	tggcaaagg	attttgagtt	gccagctcct	600
ccaaggccag	ttaggccagt	taccagatc	tgagtcgacc	tgcaggcatg	cccgtgaaa	660
tcaccagtct	ctctctacaa	atctatctct	ctctataata	atgtgtgagt	agttcccaga	720
taagggaatt	agggttctta	tagggtttcg	ctcatgtgtt	gagcatataa	gaaaccctta	780
gtatgtat	gtatttgtaa	aatacttcta	tcaataaaat	ttctaattcc	taaaacccaa	840
atccagtggc	gagctcgaat	tcgagctcga	gcccgggtgg	atcctctaga	gtcgacctgc	900
agaagcttcg	gtccggcgcg	cctctagttg	aagacacgtt	catgtcttca	tcgtaagaag	960
acactcagta	gtcttcggcc	agaatggcct	aactcaaggc	cctcactccg	cttgatcttg	1020
gcaaagatat	ttgacgcatt	tattagtatg	tgttaatttt	catttgagct	gcagtatttt	1080
ctattcgatc	tttatgtaat	tcgttacaat	taataaatat	tcaaatcaga	ttattgactg	1140
tcatttgtat	caaatacgtg	ttaatggata	tttttattat	aatattgatg	atatctcaat	1200
caaaacgtag	ataataataa	tattttattt	atatttttgc	gtcgcacagt	gaaaatctat	1260
atgagattac	aaaataccga	caacattatt	taagaacat	agacattaac	cctgagactg	1320
ttggacatca	acgggtagat	tccttcacgc	atagcacctc	attcttgggg	acaaaagcac	1380
ggtttggccg	ttccattgct	gcacgaacga	gctttgctat	atcctcgggt	tggatcatct	1440
catcagggtc	aatcaatttt	gtccaagaac	tcattgttagt	cgcaacgaaa	ccggggcata	1500
tgtcgggtat	ctcgagctcg	cgaaagcttg	gctgcaggtc	gacggatcct	tcaacaaaag	1560
ggtacctgta	cccgaaccg	acacaggtgg	gtaggtagag	aatacctagg	ggcgcgagac	1620
aactctctct	aaggaaactcg	gcaaaatagc	cccgtaaact	cgggagaagg	ggtgcccccc	1680
gctaacaata	aacgaatacg	gtttatgtat	ggattccggt	aaaataccgg	tactcgattt	1740
cataagatgc	gaataggaag	ttaagatgag	ggtggtatca	tcataaaaat	ggagtagtat	1800
cctaaattat	actaatccac	gtatgatatg	tatgccttcc	cttatcaacc	ggaagtagtg	1860
caaaaaaaat	tctatactgc	actgctctct	ttttactgag	aatgcaaaa	aaataaaagt	1920
gaagtaagg	tgccccatag	atatttgatc	ttgcctcctg	ccccccccc	ccttttttca	1980
tcaaaaattt	ccatgaaaaa	agaaaagatg	aatttgtcca	ttcattgaac	cctagttcgg	2040
gactgacggg	gctcgaaccc	gcagcttccg	cctgttccta	gccttcagg	gccagcgta	2100
agcaatacca	gccacagcac	cctcaacctc	agcaaccaac	caagggtatc	tatcttgcaa	2160
cctctctaga	tcataaatcc	actcttggtg	tgtttgtggc	tctgtcctaa	agttcactgt	2220
agacgtctca	atgtaatggt	taacgatatc	acaaaccgcg	gaacacaaga	acgaaagcac	2280
cttttcattc	tttcatatac	taggggtttt	tacttggaag	agacaatggt	ccatactaaa	2340
ggatagctgc	agaagccgcc	accgtcttga	ggaccttccg	gggagccaga	ccggtcgaac	2400
cgtgcctcca	cttgctaagg	agaaagggaa	aatcagggcc	aggacatacg	aaggaggagc	2460

-continued

```

cagaacgaag atatacctaag atacttactc gctccggggc atgatcaatc atgcctgtgg 2520
ggaggtctct cgcacctcga tccatgaagg taccaccgag gtctgccccg ccgccggctt 2580
cggtaaccgtc ctgccttgg gcgcccgagg caccgggggg atggactgcc caggcgcagc 2640
cacgacgacc caaggatcac cctcctgcgc agtcggcacg agcaatagtt ctcggggaac 2700
aggcagcttg gcctgactcc ccgggggtcac ctcaactacc tcggccgagg ggtcaagtac 2760
cccctcagtc cgcccccgct ctccggaccg ggaccccgac gtcccgggcc cgataaccga 2820
cggcaccagc ccgctcgggg gctggcttga cgacccctgg ccagcctca gatctgggct 2880
gaggccgagg caggcgccca tgcgtcgtc ttcacatcgc tcttcacatc cgtcgtcgtc 2940
atcaggcgctc tccggcgacg gctcccttgg gagccctcc ctctcctgcc gacgacgaag 3000
cctttccaag gcataccgag ccacgctccg ctctggggcc cgagccttct ttgcgtcctt 3060
cttctccttc ctcttctccg cgttgacctc ccgcgcagct cggtcaccgc cctcctccgg 3120
gactggtggc aggggaaggct tgtgatgcc taccctcctg agacagacga aaagtctcag 3180
ctatgagaac cgagggcaat ctgacgcaag aaggagaag gagcggatac tcaccagaga 3240
cacgcacccg cgatcgggac gcattaaggg ctgggaaaaa gtgccggcct ctaatttcgc 3300
taccgtgccg tccaccacc tgtggaggtc atcgatggga aggggaa 3347

```

```

<210> SEQ ID NO 28
<211> LENGTH: 669
<212> TYPE: DNA
<213> ORGANISM: Zea mays

```

```

<400> SEQUENCE: 28

```

```

actagtttcc tagcccgct cgtgcccta cccaccgac gtttatggaa ggtgccattc 60
cacggttctt cgtggcgcc cctaaggatg taaatggctg gtaaaatccg gtaaatcttc 120
ggtaccgttt accagatttt tccagccgtt ttcggattta tcgggatata cagaaaaacga 180
gacggaacg gaataggttt tttttcgaaa acggtacggt aaacggtgag acaaacttac 240
cgtccgtttt cgtatttctc gggaaactct ggtatatccc cgtatttgc ccgtattttc 300
ccgacccacg gacctgcaa tcaaccatca gccagtcagc ccatcccccac agctatggcc 360
catggggcca tgttggccac atgccacgc aacgcaaggc agtaaggctg gcagcctggc 420
acgcattgac gcattgtggc acacacagcc gccgcctgtt cgtgtttctg tgccgttgtg 480
cgagactgtg actgcgagt gcggagtcgg cgaacggcga ggcgtctccg gactctggac 540
tgccgtgtg gacagcgac ctgtgacggc gactcggcga agcccaagc taccaagccc 600
ccaagtcccc atccatctct gcttctctg tcatctcctt cccctggctg atctgcaggc 660
gccagaccg 669

```

```

<210> SEQ ID NO 29
<211> LENGTH: 200
<212> TYPE: DNA
<213> ORGANISM: Zea mays

```

```

<400> SEQUENCE: 29

```

```

gccgaagcat cacgaaacgc actaagacct cgaaggagtc aaaccactcc tccgaggcct 60
cgggggctac acccggcggg tgcgtcgcg cgcacccacc ggaacaaaat gtaaccgaga 120
aaggctcggtc cccttgcaaa aaaagtgcga caaagcctc caagcgagta ttaacctca 180
ctttgaggct cgggggctac 200

```

-continued

```

<210> SEQ ID NO 30
<211> LENGTH: 812
<212> TYPE: DNA
<213> ORGANISM: Zea mays
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (0)...(0)
<223> OTHER INFORMATION: Fragment of maize Huck-1 retrotransposon

<400> SEQUENCE: 30

tgtcggggac cataattagg ggtaccccca agactcctaa tctcagctgg taacccccat    60
cagcacaaaag ctgcaaaggc ctgatgggtg cgattaagtc aaggctcggc cactcaagg    120
gacacgatct cgctcgccc gagccagcc tcgggcaagg gcggccgacc ccgaggattc    180
acgtctcgcc cgaggggccc ctcaagcgac gggcacacct tcggctcgcc cgaggcccat    240
tcttcgccga gaagcaacct tggccagatc gccacaccga ccgaccgtat cgcaggagca    300
tttaatgcga ggatgcctg acaccttata ctgacgcgct ctcttcagtc gacagagccg    360
aagtgaaccg aatcacttcg ccgctccact gaccgacctg acaagaagac agcgccgcct    420
gcgtcgctcc gactgctgtg ccactcgaca gactgaggct gacagcagcc aagtcgggcc    480
tcgggcgcca taggaagtc cgctcgccc gacctagggt ctgggactcg gcctcggtc    540
cggaagacga cgaactacgc ttgccccgac cccagggtt ggactcagcc tcggctcgg    600
aagacgacga attccgctc gcccgacccc agggctcgga ctgggctcg gctccagaag    660
acgacgaact ccgctcgcc cgacccaggt gctcggaact agcctcggt ccggaagacg    720
acgaactccg cctcgcccga ccccgaggct cggactcagc ctcggcctca gacgatggtc    780
tccgcctcgc ccgacccggg gctcggaact ga                                812

```

```

<210> SEQ ID NO 31
<211> LENGTH: 335
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Represents part of PHI8999A insert sequence -
        fragment of Cry1F gene

```

```

<400> SEQUENCE: 31

cctttctatc ggacctgtc agatcctgtc ttgctccgag gagccttgg caatcctcac    60
tatgtactcg gtcttagggg agtggccttt caacaaactg gtacgaatca caccgcaca    120
ttcaggaact ccggggacct tgactctcta gatgagatac cacctcaaga caacagcggc    180
gcaccttgga atgactactc ccatgtgctg aatcatgtta cctttgtgct ctggccaggt    240
gagatctcag gttccgactc atggagagca ccaatgttct cttggacgca tcgtagcgct    300
acccccacaa acaccattga tccagagaga atcac                                335

```

```

<210> SEQ ID NO 32
<211> LENGTH: 321
<212> TYPE: DNA
<213> ORGANISM: Zea mays
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (0)...(0)
<223> OTHER INFORMATION: Fragment of maize chloroplast rpoC2 gene

```

```

<400> SEQUENCE: 32

tcattcttca agaactgcat atcttgccga gatcctcatc cctaaaggta cttgacaata    60
gtattattgg agtcgatata caactcacia aaaatacaag aagtcgacta ggtggattgg    120

```


-continued

<400> SEQUENCE: 37

```
cgaagcttcg gccggggccc atcgatatcc gcgggcatgc ctgcagtgc gcgtagcccg    60
gtcgtgcccc tctctagaga taatgagcat tgcattgtcta agttataaaa aattacca    118
```

<210> SEQ ID NO 38

<211> LENGTH: 550

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Represents part of PHI8999A insert
sequence-fragment of ORF25 terminator (complement)

<400> SEQUENCE: 38

```
ctcactccgc ttgatcttgg caaagatatt tgacgcattt attagtatgt gttaattttc    60
atttgcagtg cagtattttc tattcgatct ttatgtaatt cgttacaatt aataaatatt    120
caaatcagat tattgactgt catttgtatc aaatcgtgtt taatggatat ttttattata    180
atattgatga tatctcaatc aaaacgtaga taataataat atttatttaa ttttttgcg    240
tcgcacagtg aaaatctata tgagattaca aaataccgac aacattatct aagaaacata    300
gacattaacc ctgagactgt tggacatcaa cggttagatt ctttcattgca tagcacctca    360
ttcttgggga caaaagcacg gtttggccgt tccattgctg cacgaacgag ctttgcata    420
tcctcggggt ggatcatctc atcagggtcca atcaaatttg tccaagaact catgttagtc    480
gcaacgaaac cggtgcatat gtcgggtatc tcgagctcgc gaaagcttgg ctgcaggtcg    540
acggatcctt                                     550
```

<210> SEQ ID NO 39

<211> LENGTH: 128

<212> TYPE: DNA

<213> ORGANISM: Zea mays

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (0)...(0)

<223> OTHER INFORMATION: Fragment of maize chloroplast rps12 rRNA gene
(complement)

<400> SEQUENCE: 39

```
caacaaaagg gtacctgtac ccgaaaccga cacaggtggg taggtagaga atacctaggg    60
gcgcgagaca actctctcta aggaactcgg caaaatagcc ccgtaacttc gggagaaggg    120
gtgcccc                                     128
```

<210> SEQ ID NO 40

<211> LENGTH: 392

<212> TYPE: DNA

<213> ORGANISM: Zea mays

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (0)...(0)

<223> OTHER INFORMATION: Fragment of maize chloroplast genome

<400> SEQUENCE: 40

```
ctaacaataa acgaatacgg tttatgtatg gattccggta aaataccggt actcgatttc    60
ataagagtcg aataggaagt taagatgagg gtggtatcat cataaaaaat gagtagtatc    120
ctaaattata ctaatccacg tatgatatgt atgcctttcc ttatcaaccg gaagtagtgc    180
aaaaaaaatt ctatactgca ctgctctctt tttactgaga aatgcaaaaa aataaaagtg    240
aagtaagggg gcccocataga tttttgatct tgcoctctgt ccccccccc cttttttcat    300
```

-continued

caaaaatttc catgaaaaaa gaaaagatga atttgtccat tcattgaacc ctagtccggg	360
actgacgggg ctcgaaccgc cagcttccgc ct	392

<210> SEQ ID NO 41
 <211> LENGTH: 188
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Represents part of PHI8999A insert
 sequence-fragment of pat gene (complement)

<400> SEQUENCE: 41	
gttcctagcc ttccagggcc cagcgtaagc aataccagcc acagcaccct caacctcagc	60
aaccaaccaa gggatatctat ctgtcaacct ctctagatca tcaatccact ctgtgggtgt	120
ttgtggctct gtcctaaagt tcaactgtaga cgtctcaatg taatgggttaa cgatatcaca	180
aaccgcgg	188

<210> SEQ ID NO 42
 <211> LENGTH: 81
 <212> TYPE: DNA
 <213> ORGANISM: Zea mays
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (0)...(0)
 <223> OTHER INFORMATION: Fragment of maize chloroplast ORF241
 (complement)

<400> SEQUENCE: 42	
cacaagaacg aaagcacctt ttcattcttt catatactag gggtttttac ttggaaaaga	60
caatgttcca tactaaagga t	81

<210> SEQ ID NO 43
 <211> LENGTH: 254
 <212> TYPE: DNA
 <213> ORGANISM: Zea mays

<400> SEQUENCE: 43	
agctgcagaa gccgccaccg tcttgaggac cttccgggga gccagaccgg tcgaaccgtg	60
cctccacttg ctaaggagaa agggaaaatc agggccagga catacgaagg aggagccaga	120
acgaagatat cctaagatac ttactcgctc cgggccatga tcaatcatgc ctgtggggag	180
gtctctcgca cctcgatcca tgaaggtagc accgaggtct gcccgcgcgc cggttcggt	240
accgtcctcg cctt	254

<210> SEQ ID NO 44
 <211> LENGTH: 749
 <212> TYPE: DNA
 <213> ORGANISM: Zea mays

<400> SEQUENCE: 44	
gggcgcccga ggcacccggg ggatggactg cccaggcgca gccacgacga cccaaggatc	60
accctcctgc gcagtcggca cgagcaatag ttctcgggga acaggcagct tggcctgact	120
ccccggggtc acctcaacta cctcggccga ggggtcaagt accccctcag tccgcccccg	180
ctcttcggac cgggaccccg acgtccggc cccggatacc gacggcacca gccgctcgg	240
gggctggctt gacgaccctt ggcccagcct cagatctggg ctgaggccga ggcaggcggc	300
catgtcgtcg tcttcacat cgtcttcac atcgtcgtcg tcatcaggcg tctccggcga	360
cggctccctt gggagccctt cctctcctg ccgacgacga agcctttcca aggcaccccg	420

-continued

```

agccccacgtc cgctcgtggg cccgagcctt ctttgcgtec ttcttctcct tcctcttctc 480
cgcggtgacc ctccgcgcag ctccggtccac cgcctcctcc gggactgggtg gcaggggaagg 540
cttgtgatgc cctacctcct ggagacagac gaaaagtctc agctatgaga accgagggca 600
atctgacgca agaaggaaga aggagcggat actcaccaga gacacgcacc cgcgatcggg 660
acgcattaag ggctgggaaa aagtgccggc ctctaatttc gctaccgtgc cgtccaccca 720
cctgtggagg tcctcgatgg gaaggggaa 749

```

```

<210> SEQ ID NO 45
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Biolistic transformation event 5' flanking
sequence; junction between regions 3 and 4

```

```

<400> SEQUENCE: 45

```

```

tcggactcga ctttctatc 20

```

```

<210> SEQ ID NO 46
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Biolistic transformation event 5' flanking
sequence; junction between regions 4 and 5

```

```

<400> SEQUENCE: 46

```

```

agagaatcac tcattcttca 20

```

```

<210> SEQ ID NO 47
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Biolistic transformation event 5' flanking
sequence; junction between regions 5 and 6

```

```

<400> SEQUENCE: 47

```

```

gaaaagtatt gtggctatct 20

```

```

<210> SEQ ID NO 48
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Biolistic transformation event 5' flanking
sequence; junction between regions 6 and 7a

```

```

<400> SEQUENCE: 48

```

```

tctcaaggcc gcagctgata 20

```

```

<210> SEQ ID NO 49
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Biolistic transformation event 5' flanking
sequence; junction between regions 7a and 7b

```

```

<400> SEQUENCE: 49

```

```

ggctaggaac cctcaacctc 20

```

-continued

<210> SEQ ID NO 50
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Biolistic transformation event 5' flanking
sequence; junction between regions 7b and 7c

<400> SEQUENCE: 50

tcacaaaccg agagaagagg 20

<210> SEQ ID NO 51
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Biolistic transformation event 5' flanking
sequence; junction between regions 7c and 8

<400> SEQUENCE: 51

agagggatct cgaagcttcg 20

<210> SEQ ID NO 52
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Biolistic transformation event 5' flanking
sequence; junction between regions 8 and 9

<400> SEQUENCE: 52

aaaattacca caactggaag 20

<210> SEQ ID NO 53
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Biolistic transformation event 3' flanking
sequence; junction between regions 9 and 10

<400> SEQUENCE: 53

agctatgttt ctactccgc 20

<210> SEQ ID NO 54
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Biolistic transformation event 3' flanking
sequence; junction between regions 10 and 11

<400> SEQUENCE: 54

acggatcctt caacaaaagg 20

<210> SEQ ID NO 55
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Biolistic transformation event 3' flanking
sequence; junction between regions 11 and 12

<400> SEQUENCE: 55

gtgcccccg ctaacaataa 20

<210> SEQ ID NO 56

-continued

```

<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Biolistic transformation event 3' flanking
sequence; junction between regions 12 and 13

<400> SEQUENCE: 56

gcttcgcgct gttcctagcc                                20

<210> SEQ ID NO 57
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Biolistic transformation event 3' flanking
sequence; junction between regions 13 and 14

<400> SEQUENCE: 57

aaaccgcgga acacaagaac                                20

```

What is claimed is:

1. A corn plant comprising a DNA construct, said DNA construct comprising a first and a second expression cassette, wherein said first expression cassette in operable linkage comprises

- (a) a maize ubiquitin promoter;
- (b) a 5' untranslated exon of a maize ubiquitin gene;
- (c) a maize ubiquitin intron;
- (d) a Cry1F encoding DNA molecule; and
- (e) a 3' ORF25 transcriptional terminator; and said second expression cassette comprising in operable linkage
 - (i) a CaMV 35S promoter;
 - (ii) a pat encoding DNA molecule; and
 - (iii) a 3' transcriptional terminator from (CaMV) 35S comprising the nucleic acid sequence of positions 5843 to 6032 of SEQ ID NO: 25;
 and wherein the first and second expression cassettes are flanked by SEQ ID NO: 26 at the 5' end and SEQ ID NO: 27 at the 3' end.

2. The corn plant of claim 1, wherein the corn plant is a seed.

3. A method of producing an insect resistant corn plant comprising:

- (a) breeding with an insect resistant corn plant comprising in its genome DNA the DNA construct of claim 1 and
- (b) selecting a progeny of (a) by analyzing the genome of the progeny for the presence of the DNA construct of claim 1.

4. The method of claim 3, wherein breeding comprises crossing the insect resistant corn plant of (a) with a second plant.

5. The method of claim 3, wherein the insect resistant corn plant of (a) is the pollen parent for the crossing.

6. The method of claim 4, wherein the second plant is the pollen parent for the crossing.

7. The method of claim 3, wherein the progeny comprises the insect resistance of the insect resistant corn plant of (a).

8. The corn plant of claim 1, wherein the DNA construct comprises the polynucleotide of SEQ ID NO: 24.

* * * * *