

05-05-03

60467772 .050203

PTO/SB/16 (10-01)

Approved for use through 10/31/2002. OMB 0651-0032

Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

**PROVISIONAL APPLICATION FOR PATENT COVER SHEET**

This is a request for filing a PROVISIONAL APPLICATION FOR PATENT under 37 CFR 1.53 (c).

Express Mail Label No. EL870385389 US

**INVENTOR(S)**

| Given Name (first and middle [if any]) | Family Name or Surname | Residence<br>(City and either State or Foreign Country) |
|----------------------------------------|------------------------|---------------------------------------------------------|
| Mary E.                                | Hartnett Locke         | Mickleton, NJ                                           |

☐ Additional inventors are being named on the \_\_\_\_\_ separately numbered sheets attached hereto**TITLE OF THE INVENTION (500 characters max)**

Corn Event TC1507 and Methods for Detection Thereof

**CORRESPONDENCE ADDRESS**

Direct all correspondence to:

☒ Customer Number

27310

Place Customer Number  
Bar Code Label here

OR

Type Customer Number here

☐ Firm or  
Individual Name

Address

Address

City

State

ZIP

Country

Telephone

Fax

**ENCLOSED APPLICATION PARTS (check all that apply)**☒ Specification Number of Pages

51

☐ CD(s), Number☒ Drawing(s) Number of Sheets

1

☒ Other (specify) Sequence Listing and  
Statement regarding Sequence Listing☒ Application Data Sheet. See 37 CFR 1.76**METHOD OF PAYMENT OF FILING FEES FOR THIS PROVISIONAL APPLICATION FOR PATENT**☐ Applicant claims small entity status. See 37 CFR 1.27.☐ A check or money order is enclosed to cover the filing fees☒ The Commissioner is hereby authorized to charge filing  
fees or credit any overpayment to Deposit Account Number:

16-1852

FILING FEE  
AMOUNT (\$)

160.00

☐ Payment by credit card. Form PTO-2038 is attached.

The invention was made by an agency of the United States Government or under a contract with an agency of the United States Government.

☒ No.☐ Yes, the name of the U.S. Government agency and the Government contract number are: \_\_\_\_\_

Respectfully submitted,

SIGNATURE

Date

5/2/03

TYPED or PRINTED NAME

Amy L. Cigan

REGISTRATION NO.  
(if appropriate)

45,494

Docket Number:

1523P

TELEPHONE

515-248-4896

**USE ONLY FOR FILING A PROVISIONAL APPLICATION FOR PATENT**

This collection of information is required by 37 CFR 1.51. The information is used by the public to file (and by the PTO to process) a provisional application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.14. This collection is estimated to take 8 hours to complete, including gathering, preparing, and submitting the complete provisional application to the PTO. Time will vary depending upon the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, Washington, D.C., 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Box Provisional Application, Assistant Commissioner for Patents, Washington, D.C. 20231.

Inari Exhibit 1005  
Inari v. Corteva

**Certificate of Mailing under 37 CFR 1.10****Express Mailing Label Number EL870385389US**

I hereby certify that this correspondence is being deposited with the United States Postal Service with sufficient postage as "Express Mail Post Office to Addressee" Service in an envelope addressed to:

Mail Stop Provisional Patent Application  
Commissioner for Patents  
P.O. Box 1450  
Alexandria, VA 22313-1450

on May 2, 2003.

  
\_\_\_\_\_  
Signature

\_\_\_\_\_  
Tracy Westphal  
Typed or printed name of person signing Certificate

Note: Each paper must have its own certificate of mailing, or this certificate must identify each submitted paper.

1. Provisional Application for Patent Cover Sheet (1 orig + 1 copy)
2. Specification (1 copy)
3. Fig. 1 (1 copy)
4. Application Data Sheet (1 copy)
5. Statement to Support Filing and Submission in Accordance with  
37 CFR 1.821 through 1.825 (1 copy)
6. Sequence Listing (1 paper copy + 1 diskette)

## IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

First Named Inventor : Mary E. Hartnett Locke  
Application Number :  
Filing Date : May 2, 2003  
Title : Corn Event TC1507 and Methods for Detection  
Thereof  
  
Group/Art Unit :  
Examiner :  
  
Attorney Docket Number : 1523P

Mail Stop Sequence  
Commissioner for Patents  
P.O. Box 1450  
Alexandria, VA 22313-1450

**STATEMENT TO SUPPORT FILING AND SUBMISSION  
IN ACCORDANCE WITH 37 CFR 1.821 THROUGH 1.825**

- ☒ I hereby state that the contents of the paper and computer readable copies of the Sequence Listing, submitted in accordance with 37 CFR 1.821(c) and (e), respectively, are the same.
- ☐ I hereby state that the submission filed in accordance with 37 CFR 1.821(g) does not include new matter.
- ☐ I hereby state that the submission filed in accordance with 37 CFR 1.821(h) does not include new matter or go beyond the disclosure in the international application as filed.
- ☐ I hereby state that the amendments, made in accordance with 37 CFR 1.825(a), included in the substitute sheet(s) of the Sequence Listing are supported in the application, as filed, at pages \_\_\_\_\_. I hereby state that the substitute sheet(s) of the Sequence Listing does (do) not include new matter.
- ☐ I hereby state that the substitute copy of the computer readable form, submitted in accordance with 37 CFR 1.825(b), is the same as the amended Sequence Listing.

☐ I hereby state that the substitute copy of the computer readable form, submitted in accordance with 37 CFR 1.825(d), is identical to that originally filed.

Respectfully submitted,



Amy L. Cigan  
Agent for Applicant(s)  
Registration No. 45,494

PIONEER HI-BRED INTERNATIONAL, INC.  
Corporate Intellectual Property  
7100 N.W. 62<sup>nd</sup> Avenue  
P.O. Box 1000  
Johnston, Iowa 50131-1000  
Phone: (515) 248-4896  
Facsimile: (515) 334-6883

Application Data Sheet**Application Information**

Application Type:: Provisional  
Subject Matter::  
Suggested Classification::  
Suggested Group Art Unit::  
CD-ROM or CD-R?::  
Number of CD disks::  
Number of copies of CDs::  
Sequence submission?:: paper  
Computer Readable Form  
(CRF)?:: Yes  
Number of copies of CRF:: 01  
Title:: Corn Event TC1507 and Methods for Detection  
Thereof  
Attorney Docket Number:: 1523P  
Request for Early Publication?::  
Request for Non-Publication?::  
Suggested Drawing Figure::  
Total Drawing Sheets:: 1  
Small Entity::  
Petition included?::  
Secrecy Order in Parent Appl.?::

**Applicant Information:**

Applicant Authority type:: Inventor  
Primary Citizenship Country:: US  
Status:: Full Capacity  
Given Name:: Mary  
Middle Name:: E.  
1 Initial 5/1/2003

Family Name:: Hartnett Locke  
City of Residence:: Mickleton  
State or Province of Residence:: NJ  
Country of Residence:: US  
Street of mailing address:: 31 Millwood Drive  
City of mailing address:: Mickleton  
State or Province of mailing address:: NJ  
Postal or Zip Code of mailing address:: 08056

**Correspondence Information**

Correspondence Customer Number:: 27310

**Representative Information**

Representative Customer Number:: 27310

**Domestic Priority Information**

| Application:: | Continuity Type:: | Parent Application:: | Parent Filing Date:: |
|---------------|-------------------|----------------------|----------------------|
|               |                   |                      |                      |
|               |                   |                      |                      |
|               |                   |                      |                      |

**Assignee Information**

Assignee Name:: Pioneer Hi-Bred International, Inc.

## CORN EVENT TC1507 AND METHODS FOR DETECTION THEREOF

### 5 FIELD OF THE INVENTION

The present invention relates to the field of plant molecular biology, specifically the invention relates to a DNA construct for conferring insect resistance to a plant. The invention more specifically relates to an insect resistant corn plant TC1507 and to assays for detecting the presence of corn plant TC1507 DNA in a sample and compositions thereof.

### 10 BACKGROUND OF THE INVENTION

This invention relates to the insect resistant corn (*Zea mays*) plant TC1507, also referred to as maize line TC1507 or maize event TC1507, and to the DNA plant expression construct of corn plant TC1507 and the detection of the transgene/flanking insertion region in corn plant TC1507 and progeny thereof.

15 Corn is an important crop and is a primary food source in many areas of the world. Damage caused by insect pests is a major factor in the loss of the world's corn crops, despite the use of protective measures such as chemical pesticides. In view of this, insect resistance has been genetically engineered into crops such as corn in order to control insect damage and to reduce the need for traditional chemical pesticides. One group of genes which have been  
20 utilized for the production of transgenic insect resistant crops are the delta-endotoxins from *Bacillus thuringiensis* (*B.t.*). Delta-endotoxins have been successfully expressed in crop plants such as cotton, potatoes, rice, sunflower, as well as corn, and have proven to provide excellent control over insect pests. (Perlak, F.J *et al.* (1990) *Bio/Technology* 8, 939-943; Perlak, F.J. *et al.* (1993) *Plant Mol. Biol.* 22: 313-321; Fujimoto H. *et al.* (1993) *Bio/Technology* 11: 1151-  
25 1155; Tu *et al.* (2000) *Nature Biotechnology* 18:1101-1104; PCT publication number WO 01/13731; and Bing JW *et al.* (2000) Efficacy of Cry1F Transgenic Maize, 14<sup>th</sup> Biennial International Plant Resistance to Insects Workshop, Fort Collins, CO).

The expression of foreign genes in plants is known to be influenced by their location in the plant genome, perhaps due to chromatin structure (e.g., heterochromatin) or the proximity  
30 of transcriptional regulatory elements (e.g., enhancers) close to the integration site (Weising *et al.*, *Ann. Rev. Genet* 22:421-477, 1988). At the same time the presence of the transgene at different locations in the genome will influence the overall phenotype of the plant in different ways. For this reason, it is often necessary to screen a large number of events in order to

identify an event characterized by optimal expression of an introduced gene of interest. For example, it has been observed in plants and in other organisms that there may be a wide variation in levels of expression of an introduced gene among events. There may also be differences in spatial or temporal patterns of expression, for example, differences in the relative  
5 expression of a transgene in various plant tissues, that may not correspond to the patterns expected from transcriptional regulatory elements present in the introduced gene construct. For this reason, it is common to produce hundreds to thousands of different events and screen those events for a single event that has desired transgene expression levels and patterns for commercial purposes. An event that has desired levels or patterns of transgene expression is  
10 useful for introgressing the transgene into other genetic backgrounds by sexual outcrossing using conventional breeding methods. Progeny of such crosses maintain the transgene expression characteristics of the original transformant. This strategy is used to ensure reliable gene expression in a number of varieties that are well adapted to local growing conditions.

It would be advantageous to be able to detect the presence of a particular event in order  
15 to determine whether progeny of a sexual cross contain a transgene of interest. In addition, a method for detecting a particular event would be helpful for complying with regulations requiring the pre-market approval and labeling of foods derived from recombinant crop plants, for example, or for use in environmental monitoring, monitoring traits in crops in the field, or monitoring products derived from a crop harvest, as well as for use in ensuring compliance of  
20 parties subject to regulatory or contractual terms.

It is possible to detect the presence of a transgene by any nucleic acid detection method known in the art including, but not limited to, the polymerase chain reaction (PCR) or DNA hybridization using nucleic acid probes. These detection methods generally focus on frequently used genetic elements, such as promoters, terminators, marker genes, etc., because  
25 for many DNA constructs, the coding region is interchangeable. As a result, such methods may not be useful for discriminating between different events, particularly those produced using the same DNA construct or very similar constructs unless the DNA sequence of the flanking DNA adjacent to the inserted heterologous DNA is known. For example, an event-specific PCR assay is described in U.S. Patent No. 6,395,485 for the detection of elite event GAT-ZM1.  
30 Accordingly, it would be desirable to have a simple and discriminative method for the identification of event TC1507.

# SUMMARY OF THE INVENTION

This invention relates preferably to methods for producing and selecting an insect resistant monocot crop plant. More specifically, a DNA construct is provided that when expressed in plant cells and plants confers resistance to insects. According to one aspect of the invention, a DNA construct, capable of introduction into and replication in a host cell, is provided that when expressed in plant cells and plants confers insect resistance to the plant cells and plants. The DNA construct is comprised of a DNA molecule named PHI8999A and it includes two transgene expression cassettes. The first expression cassette comprises a DNA molecule which includes the promoter, 5' untranslated exon, and first intron of the maize ubiquitin (Ubi-1) gene (Christensen *et al.* (1992) *Plant Mol. Biol.* 18:675-689 and Christensen and Quail (1996) *Transgenic Res.* 5:213-218) operably connected to a DNA molecule encoding a *B.t.*  $\delta$ -endotoxin identified as Cry1F (U.S. Pat. Nos. 5,188,960 and 6,218,188) operably connected to a DNA molecule comprising a 3' ORF25 transcriptional terminator isolated from *Agrobacterium tumefaciens* (Barker *et al.* (1983) *Plant Mol. Biol.* 2:335-350). The second transgene expression cassette of the DNA construct comprises a DNA molecule of the cauliflower mosaic virus (CaMV) 35S promoter (Odell J.T. *et al.* (1985) *Nature* 313: 810-812; Mitsuhashi *et al.* (1996) *Plant Cell Physiol.* 37: 49-59) operably connected to a DNA molecule encoding a phosphinothricin acetyltransferase (PAT) gene (Wohlleben W. *et al.* (1988) *Gene* 70: 25-37) operably connected to a DNA molecule comprising a 3' transcriptional terminator from (CaMV) 35S (see Mitsuhashi *et al.* (1996) *Plant Cell Physiol.* 37: 49-59). Plants containing the DNA construct are also provided.

According to another aspect of the invention, compositions and methods are provided for identifying a novel corn plant designated TC1507, which methods are based on primers or probes which specifically recognize the 5' and/or 3' flanking sequence of TC1507. DNA molecules are provided that comprise primer sequences that when utilized in a PCR reaction will produce amplicons unique to the transgenic event TC1507. These molecules may be selected from the group consisting of:

- 5'-GTAGTACTATAGATTATATTATTCGTAGAG-3' (SEQ ID NO: 1);
- 5'-GCCATACAGAACTCAAAATCTTTCCGGAG-3' (SEQ ID NO: 2);
- 5'-CTTCAAACAAGTGTGACAAA-3' (SEQ ID NO: 23 );
- 5'-TGTGGTGTGTTGTGGCTCTGTCCTAA-3' (SEQ ID NO: 3);

5'-AGCACCTTTTCATTCTTTCATATAC-3' (SEQ ID NO: 4);

5'-GACCTCCCCA CAGGCATGAT TGATC-3' (SEQ ID NO: 5);

and complements thereof. The corn plant and seed comprising these molecules is an aspect of this invention. Further, kits utilizing these primer sequences for the identification of the

5 TC1507 event are provided.

An additional aspect of the invention relates to the specific flanking sequences of TC1507 described herein, which can be used to develop specific identification methods for TC1507 in biological samples. More particularly, the invention relates to the 5' and/or 3' flanking regions of TC1507, SEQ ID NO:21 and SEQ ID NO:22, respectively, which can be  
10 used for the development of specific primers and probes. The invention further relates to identification methods for the presence of TC1507 in biological samples based on the use of such specific primers or probes.

According to another aspect of the invention, methods of detecting the presence of DNA corresponding to the corn event TC1507 in a sample are provided. Such methods  
15 comprise: (a) contacting the sample comprising DNA with a DNA primer set, that when used in a nucleic acid amplification reaction with genomic DNA extracted from corn event TC1507 produces an amplicon that is diagnostic for corn event TC1507; (b) performing a nucleic acid amplification reaction, thereby producing the amplicon; and (c) detecting the amplicon.

DNA molecules that comprise the novel transgene/flanking insertion region, SEQ ID  
20 NO: 26 and SEQ ID NO: 27 and are homologous or complementary to SEQ ID NO: 26 and SEQ ID NO: 27 are an aspect of this invention.

DNA sequences that comprise the novel transgene/flanking insertion region, SEQ ID NO:26 are an aspect of this invention. DNA sequences that comprise a sufficient length of polynucleotides of transgene insert sequence and a sufficient length of polynucleotides of  
25 maize genomic and/or flanking sequence from maize plant TC1507 of SEQ ID NO:26 that are useful as primer sequences for the production of an amplicon product diagnostic for maize plant TC1507 are included.

In addition, DNA sequences that comprise the novel transgene/flanking insertion region, SEQ ID NO:27 are provided. DNA sequences that comprise a sufficient length of  
30 polynucleotides of transgene insert sequence and a sufficient length of polynucleotides of maize genomic and/or flanking sequence from maize plant TC1507 of SEQ ID NO:27 that are

useful as primer sequences for the production of an amplicon product diagnostic for maize plant TC1507 are included.

According to another aspect of the invention, the DNA sequences that comprise at least 11 or more nucleotides of the transgene portion of the DNA sequence of SEQ ID NO:26 or complements thereof, and a similar length of 5' flanking maize DNA sequence of SEQ ID NO:26 or complements thereof are useful as DNA primers in DNA amplification methods. The amplicons produced using these primers are diagnostic for maize event TC1507. Therefore, the invention also includes the amplicons produced by DNA primers homologous or complementary to SEQ ID NO:26.

According to another aspect of the invention, the DNA sequences that comprise at least 11 or more nucleotides of the transgene portion of the DNA sequence of SEQ ID NO:27 or complements thereof, and a similar length of 3' flanking maize DNA sequence of SEQ ID NO:27 or complements thereof are useful as DNA primers in DNA amplification methods. The amplicons produced using these primers are diagnostic for maize event TC1507. Therefore, the invention also includes the amplicons produced by DNA primers homologous or complementary to SEQ ID NO:27.

More specifically, a pair of DNA molecules comprising a DNA primer set, wherein the DNA molecules are identified as SEQ ID NO: 1 or complements thereof and SEQ ID NO: 2 or complements thereof; SEQ ID NO: 2 or complements thereof and SEQ ID NO: 23 or complements thereof; SEQ ID NO: 3 or complements thereof and SEQ ID NO: 5 or complements thereof; SEQ ID NO: 4 or complements thereof and SEQ ID NO: 5 or complements thereof are aspects of the invention.

Further aspects of the invention include the amplicon comprising the DNA molecules of SEQ ID NO: 1 and SEQ ID NO: 2; the amplicon comprising the DNA molecules of SEQ ID NO: 2 and SEQ ID NO: 23; the amplicon comprising the DNA molecules of SEQ ID NO: 3 and SEQ ID NO: 5; and the amplicon comprising the DNA molecules of SEQ ID NO: 4 and SEQ ID NO: 5.

According to another aspect of the invention, methods of detecting the presence of a DNA molecule corresponding to the TC1507 event in a sample, such methods comprising: (a) contacting the sample comprising DNA extracted from a corn plant with a DNA probe, molecule that hybridizes under stringent hybridization conditions with DNA extracted from corn event TC1507 and does not hybridize under the stringent hybridization conditions with a

control corn plant DNA; (b) subjecting the sample and probe to stringent hybridization conditions; and (c) detecting hybridization of the probe to the DNA. More specifically, a method for detecting the presence of a DNA molecule corresponding to the TC1507 event in a sample, such methods, consisting of (a) contacting the sample comprising DNA extracted from  
 5 a corn plant with a DNA probe molecule that consists of sequences that are unique to the event, e.g. junction sequences, wherein said DNA probe molecule hybridizes under stringent hybridization conditions with DNA extracted from corn event TC1507 and does not hybridize under the stringent hybridization conditions with a control corn plant DNA; (b) subjecting the sample and probe to stringent hybridization conditions; and (c) detecting hybridization of the  
 10 probe to the DNA.

In addition, a kit and methods for identifying event TC1507 in a biological sample which detects a TC1507 specific region within SEQ ID NO: 24 are provided.

DNA molecules are provided that comprise at least one junction sequence of TC1507 selected from the group consisting of SEQ ID NO:45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56  
 15 and 57 and complements thereof; wherein a junction sequence spans the junction between heterologous DNA inserted into the genome and the DNA from the corn cell flanking the insertion site, i.e. flanking DNA, and is diagnostic for the TC1507 event.

According to another aspect of the invention, methods of producing an insect resistant corn plant that comprise the steps of: (a) sexually crossing a first parental corn line comprising  
 20 the expression cassettes of the present invention, which confers resistance to insects, and a second parental corn line that lacks insect resistance, thereby producing a plurality of progeny plants; and (b) selecting a progeny plant that is insect resistant. Such methods may optionally comprise the further step of back-crossing the progeny plant to the second parental corn line to producing a true-breeding corn plant that is insect resistant.

The present invention provides a method of producing a corn plant that is resistant to  
 25 insects comprising transforming a corn cell with the DNA construct PHI8999A (SEQ ID NO:25), growing the transformed corn cell into a corn plant, selecting the corn plant that shows resistance to insects, and further growing the corn plant into a fertile corn plant. The fertile corn plant can be self pollinated or crossed with compatible corn varieties to produce insect  
 30 resistant progeny.

The invention further relates to a DNA detection kit for identifying maize event TC1507 in biological samples. Preferably the kit of the invention comprises a first primer

which specifically recognizes the 5' or 3' flanking region of TC1507, and a second primer which specifically recognizes a sequence within the foreign DNA of TC1507, or within the flanking DNA, for use in a PCR identification protocol. The invention also relates to a kit for identifying event TC1507 in biological samples, which kit comprises a specific probe having a  
 5 sequence which corresponds or is complementary to, a sequence having between 80% and 100% sequence identity with a specific region of event TC1507. Preferably the sequence of the probe corresponds to a specific region comprising part of the 5' or 3' flanking region of event TC1507.

The methods and kits encompassed by the present invention can be used for different  
 10 purposes such as, but not limited to the following: to identify event TC1507 in plants, plant material or in products such as, but not limited to, food or feed products (fresh or processed) comprising, or derived from plant material; additionally or alternatively, the methods and kits of the present invention can be used to identify transgenic plant material for purposes of segregation between transgenic and non-transgenic material; additionally or alternatively, the  
 15 methods and kits of the present invention can be used to determine the quality of plant material comprising maize event TC1507. The kits may also contain the reagents and materials necessary for the performance of the detection method.

This invention further relates to the TC1507 corn plant or its parts, including, but not limited to, pollen, ovules, vegetative cells, the nuclei of pollen cells, and the nuclei of egg cells  
 20 of the corn plant TC1507 and the progeny derived thereof. The corn plant and seed TC1507 from which the DNA primer molecules of the present invention provide a specific amplicon product is an aspect of the invention.

The foregoing and other aspects of the invention will become more apparent from the following detailed description and accompanying drawing.

#### BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1. Linear map showing the transgenic insert PHI8999A, as well as the sequences flanking the transgenic insert.

#### DETAILED DESCRIPTION

The following definitions and methods are provided to better define the present invention and to guide those of ordinary skill in the art in the practice of the present invention.

Unless otherwise noted, terms are to be understood according to conventional usage by those of ordinary skill in the relevant art. Definitions of common terms in molecular biology may also be found in Rieger *et al.*, *Glossary of Genetics: Classical and Molecular*, 5<sup>th</sup> edition, Springer-Verlag; New York, 1991; and Lewin, *Genes V*, Oxford University Press: New York,

5 1994. The nomenclature for DNA bases as set forth at 37 CFR 1.822 is used.

As used herein, the term “comprising” means “including but not limited to”.

As used herein, the term “corn” means *Zea mays* or maize and includes all plant varieties that can be bred with corn, including wild maize species.

As used herein, the term “TC1507 specific” refers to a nucleotide sequence which is  
10 suitable for discriminatively identifying event TC1507 in plants, plant material, or in products such as, but not limited to, food or feed products (fresh or processed) comprising, or derived from plant material.

As used herein, the terms “insect resistant” and “impacting insect pests” refers to effecting changes in insect feeding, growth, and/or behavior at any stage of development,  
15 including but not limited to: killing the insect; retarding growth; preventing reproductive capability; and the like.

As used herein, the terms “pesticidal activity” and “insecticidal activity” are used synonymously to refer to activity of an organism or a substance (such as, for example, a protein) that can be measured by numerous parameters including, but not limited to, pest  
20 mortality, pest weight loss, pest attraction, pest repellency, and other behavioral and physical changes of a pest after feeding on and/or exposure to the organism or substance for an appropriate length of time. For example “pesticidal proteins” are proteins that display pesticidal activity by themselves or in combination with other proteins.

“Coding sequence” refers to a nucleotide sequence that codes for a specific amino acid  
25 sequence. As used herein, the terms “encoding” or “encoded” when used in the context of a specified nucleic acid mean that the nucleic acid comprises the requisite information to guide translation of the nucleotide sequence into a specified protein. The information by which a protein is encoded is specified by the use of codons. A nucleic acid encoding a protein may comprise non-translated sequences (e.g., introns) within translated regions of the nucleic acid  
30 or may lack such intervening non-translated sequences (e.g., as in cDNA).

“Gene” refers to a nucleic acid fragment that expresses a specific protein, including regulatory sequences preceding (5' non-coding sequences) and following (3' non-coding

sequences) the coding sequence. "Native gene" refers to a gene as found in nature with its own regulatory sequences. "Chimeric gene" refers any gene that is not a native gene, comprising regulatory and coding sequences that are not found together in nature. Accordingly, a chimeric gene may comprise regulatory sequences and coding sequences that are derived from different sources, or regulatory sequences and coding sequences derived from the same source, but arranged in a manner different than that found in nature. "Endogenous gene" refers to a native gene in its natural location in the genome of an organism. "Foreign" refers to material not normally found in the location of interest. Thus "foreign DNA" may comprise both recombinant DNA as well as newly introduced, rearranged DNA of the plant. A "foreign" gene refers to a gene not normally found in the host organism, but that is introduced into the host organism by gene transfer. Foreign genes can comprise native genes inserted into a non-native organism, or chimeric genes. A "transgene" is a gene that has been introduced into the genome by a transformation procedure. The site in the plant genome where a recombinant DNA has been inserted may be referred to as the "insertion site" or "target site".

As used herein, "insert DNA" refers to the heterologous DNA within the expression cassettes used to transform the plant material while "flanking DNA" can exist of either genomic DNA naturally present in an organism such as a plant, or foreign (heterologous) DNA introduced via the transformation process which is extraneous to the original insert DNA molecule, e.g. fragments associated with the transformation event. A "flanking region" or "flanking sequence" as used herein refers to a sequence of at least 20 base pair, preferably at least 50 base pair, and up to 5000 base pair which is located either immediately upstream of and contiguous with or immediately downstream of and contiguous with the original foreign insert DNA molecule. Transformation procedures leading to random integration of the foreign DNA will result in transformants containing different flanking regions characteristic and unique for each transformant. When recombinant DNA is introduced into a plant through traditional crossing, its flanking regions will generally not be changed. Transformants will also contain unique junctions between a piece of heterologous insert DNA and genomic DNA, or 2 pieces of genomic DNA, or 2 pieces of heterologous DNA. A "junction" is a point where 2 specific DNA fragments join. For example, a junction exists where insert DNA joins flanking DNA. A junction point also exists in a transformed organism where 2 DNA fragments join together in a manner that is modified from that found in the native organism. "Junction DNA" refers to DNA that comprises a junction point.

As used herein, “heterologous” in reference to a nucleic acid is a nucleic acid that originates from a foreign species, or, if from the same species, is substantially modified from its native form in composition and/or genomic locus by deliberate human intervention. For example, a promoter operably linked to a heterologous nucleotide sequence can be from a species different from that from which the nucleotide sequence was derived, or, if from the same species, the promoter is not naturally found operably linked to the nucleotide sequence. A heterologous protein may originate from a foreign species, or, if from the same species, is substantially modified from its original form by deliberate human intervention.

“Regulatory sequences” refer to nucleotide sequences located upstream (5’ non-coding sequences), within, or downstream (3’ non-coding sequences) of a coding sequence, and which influence the transcription, RNA processing or stability, or translation of the associated coding sequence. Regulatory sequences may include promoters, translation leader sequences, introns, and polyadenylation recognition sequences.

“Promoter” refers to a nucleotide sequence capable of controlling the expression of a coding sequence or functional RNA. In general, a coding sequence is located 3’ to a promoter sequence. The promoter sequence consists of proximal and more distal upstream elements, the latter elements are often referred to as enhancers. Accordingly, an “enhancer” is a nucleotide sequence that can stimulate promoter activity and may be an innate element of the promoter or a heterologous element inserted to enhance the level or tissue-specificity of a promoter.

Promoters may be derived in their entirety from a native gene, or be composed of different elements derived from different promoters found in nature, or even comprise synthetic nucleotide segments. It is understood by those skilled in the art that different promoters may direct the expression of a gene in different tissues or cell types, or at different stages of development, or in response to different environmental conditions. Promoters that cause a nucleic acid fragment to be expressed in most cell types at most times are commonly referred to as “constitutive promoters”. New promoters of various types useful in plant cells are constantly being discovered; numerous examples may be found in the compilation by Okamuro and Goldberg (1989) *Biochemistry of Plants* 15:1-82. It is further recognized that since in most cases the exact boundaries of regulatory sequences have not been completely defined, nucleic acid fragments of different lengths may have identical promoter activity.

The “translation leader sequence” refers to a nucleotide sequence located between the promoter sequence of a gene and the coding sequence. The translation leader sequence is

present in the fully processed mRNA upstream of the translation start sequence. The translation leader sequence may affect numerous parameters including, processing of the primary transcript to mRNA, mRNA stability and/or translation efficiency. Examples of translation leader sequences have been described (Turner and Foster (1995) *Mol. Biotechnol.* 3:225-236).

The “3’ non-coding sequences” refer to nucleotide sequences located downstream of a coding sequence and include polyadenylation recognition sequences and other sequences encoding regulatory signals capable of affecting mRNA processing or gene expression. The polyadenylation signal is usually characterized by affecting the addition of polyadenylic acid tracts to the 3’ end of the mRNA precursor. The use of different 3’ non-coding sequences is exemplified by Ingelbrecht *et al.* (1989) *Plant Cell* 1:671-680.

A “protein” or “polypeptide” is a chain of amino acids arranged in a specific order determined by the coding sequence in a polynucleotide encoding the polypeptide.

A DNA construct is an assembly of DNA molecules linked together that provide one or more expression cassettes. The DNA construct may be a plasmid that is enabled for self replication in a bacterial cell and contains various endonuclease enzyme restriction sites that are useful for introducing DNA molecules that provide functional genetic elements, i.e., promoters, introns, leaders, coding sequences, 3’ termination regions, among others; or a DNA construct may be a linear assembly of DNA molecules, such as an expression cassette. The expression cassette contained within a DNA construct comprise the necessary genetic elements to provide transcription of a messenger RNA. The expression cassette can be designed to express in prokaryote cells or eukaryotic cells. Expression cassettes of the present invention are designed to express most preferably in plant cells.

The DNA molecules of the invention are provided in expression cassettes for expression in an organism of interest. The cassette will include 5’ and 3’ regulatory sequences operably linked to a coding sequence of the invention. “Operably linked” means that the nucleic acid sequences being linked are contiguous and, where necessary to join two protein coding regions, contiguous and in the same reading frame. Operably linked is intended to indicate a functional linkage between a promoter and a second sequence, wherein the promoter sequence initiates and mediates transcription of the DNA sequence corresponding to the second sequence. The cassette may additionally contain at least one additional gene to be cotransformed into the

organism. Alternatively, the additional gene(s) can be provided on multiple expression cassettes or multiple DNA constructs.

The expression cassette will include in the 5' to 3' direction of transcription: a transcriptional and translational initiation region, a coding region, and a transcriptional and translational termination region functional in the organism serving as a host. The transcriptional initiation region (*i.e.*, the promoter) may be native or analogous, or foreign or heterologous to the host organism. Additionally, the promoter may be the natural sequence or alternatively a synthetic sequence. The expression cassettes may additionally contain 5' leader sequences in the expression cassette construct. Such leader sequences can act to enhance translation.

It is to be understood that as used herein the term "transgenic" includes any cell, cell line, callus, tissue, plant part, or plant the genotype of which has been altered by the presence of a heterologous nucleic acid including those transgenics initially so altered as well as those created by sexual crosses or asexual propagation from the initial transgenic. The term "transgenic" as used herein does not encompass the alteration of the genome (chromosomal or extra-chromosomal) by conventional plant breeding methods or by naturally occurring events such as random cross-fertilization, non-recombinant viral infection, non-recombinant bacterial transformation, non-recombinant transposition, or spontaneous mutation.

A transgenic "event" is produced by transformation of plant cells with a heterologous DNA construct(s), including a nucleic acid expression cassette that comprises a transgene of interest, the regeneration of a population of plants resulting from the insertion of the transgene into the genome of the plant, and selection of a particular plant characterized by insertion into a particular genome location. An event is characterized phenotypically by the expression of the transgene. At the genetic level, an event is part of the genetic makeup of a plant. The term "event" also refers to progeny produced by a sexual outcross between the transformant and another variety that include the heterologous DNA. Even after repeated back-crossing to a recurrent parent, the inserted DNA and flanking DNA from the transformed parent is present in the progeny of the cross at the same chromosomal location. The term "event" also refers to DNA from the original transformant comprising the inserted DNA and flanking sequence immediately adjacent to the inserted DNA that would be expected to be transferred to a progeny that receives inserted DNA including the transgene of interest as the result of a sexual

cross of one parental line that includes the inserted DNA (e.g., the original transformant and progeny resulting from selfing) and a parental line that does not contain the inserted DNA.

An insect resistant TC1507 corn plant can be bred by first sexually crossing a first parental corn plant consisting of a corn plant grown from the transgenic TC1507 corn plant and  
 5 progeny thereof derived from transformation with the expression cassettes of the present invention that confers insect resistance, and a second parental corn plant that lacks insect resistance, thereby producing a plurality of first progeny plants; and then selecting a first progeny plant that is resistant to insects; and selfing the first progeny plant, thereby producing a plurality of second progeny plants; and then selecting from the second progeny plants an insect  
 10 resistant plant. These steps can further include the back-crossing of the first insect resistant progeny plant or the second insect resistant progeny plant to the second parental corn plant or a third parental corn plant, thereby producing a corn plant that is resistant to insects.

As used herein, the term "plant" includes reference to whole plants, plant organs (e.g., leaves, stems, roots, etc.), seeds, plant cells, and progeny of same. Parts of transgenic plants  
 15 understood to be within the scope of the invention comprise, for example, plant cells, protoplasts, tissues, callus, embryos as well as flowers, stems, fruits, leaves, and roots originating in transgenic plants or their progeny previously transformed with a DNA molecule of the invention and therefore consisting at least in part of transgenic cells, are also an aspect of the present invention.

20 As used herein, the term "plant cell" includes, without limitation, seeds, suspension cultures, embryos, meristematic regions, callus tissue, leaves, roots, shoots, gametophytes, sporophytes, pollen, and microspores. The class of plants that can be used in the methods of the invention is generally as broad as the class of higher plants amenable to transformation techniques, including both monocotyledonous and dicotyledonous plants.

25 "Transformation" refers to the transfer of a nucleic acid fragment into the genome of a host organism, resulting in genetically stable inheritance. Host organisms containing the transformed nucleic acid fragments are referred to as "transgenic" organisms. Examples of methods of plant transformation include *Agrobacterium*-mediated transformation (De Blaere *et al.* (1987) *Meth. Enzymol.* 143:277) and particle-accelerated or "gene gun" transformation  
 30 technology (Klein *et al.* (1987) *Nature* (London) 327:70-73; U.S. Patent No. 4,945,050, incorporated herein by reference). Additional transformation methods are disclosed below.

Thus, isolated polynucleotides of the present invention can be incorporated into recombinant constructs, typically DNA constructs, which are capable of introduction into and replication in a host cell. Such a construct can be a vector that includes a replication system and sequences that are capable of transcription and translation of a polypeptide-encoding sequence in a given host cell. A number of vectors suitable for stable transfection of plant cells or for the establishment of transgenic plants have been described in, e.g., Pouwels *et al.*, (1985; Supp. 1987) *Cloning Vectors: A Laboratory Manual*, Weissbach and Weissbach (1989) *Methods for Plant Molecular Biology*, (Academic Press, New York); and Flevin *et al.*, (1990) *Plant Molecular Biology Manual*, (Kluwer Academic Publishers). Typically, plant expression vectors include, for example, one or more cloned plant genes under the transcriptional control of 5' and 3' regulatory sequences and a dominant selectable marker. Such plant expression vectors also can contain a promoter regulatory region (e.g., a regulatory region controlling inducible or constitutive, environmentally- or developmentally-regulated, or cell- or tissue-specific expression), a transcription initiation start site, a ribosome binding site, an RNA processing signal, a transcription termination site, and/or a polyadenylation signal.

It is also to be understood that two different transgenic plants can also be mated to produce offspring that contain two independently segregating added, exogenous genes. Selfing of appropriate progeny can produce plants that are homozygous for both added, exogenous genes. Back-crossing to a parental plant and out-crossing with a non-transgenic plant are also contemplated, as is vegetative propagation. Descriptions of other breeding methods that are commonly used for different traits and crops can be found in one of several references, e.g., Fehr, in *Breeding Methods for Cultivar Development*, Wilcos J. ed., American Society of Agronomy, Madison Wis. (1987).

A "probe" is an isolated nucleic acid to which is attached a conventional detectable label or reporter molecule, e.g., a radioactive isotope, ligand, chemiluminescent agent, or enzyme. Such a probe is complementary to a strand of a target nucleic acid, in the case of the present invention, to a strand of isolated DNA from corn event TC1507 whether from a corn plant or from a sample that includes DNA from the event. Probes according to the present invention include not only deoxyribonucleic or ribonucleic acids but also polyamides and other probe materials that bind specifically to a target DNA sequence and can be used to detect the presence of that target DNA sequence.

“Primers” are isolated nucleic acids that are annealed to a complementary target DNA strand by nucleic acid hybridization to form a hybrid between the primer and the target DNA strand, then extended along the target DNA strand by a polymerase, e.g., a DNA polymerase. Primer pairs of the present invention refer to their use for amplification of a target nucleic acid sequence, e.g., by the polymerase chain reaction (PCR) or other conventional nucleic-acid amplification methods. “PCR” or “polymerase chain reaction” is a technique used for the amplification of specific DNA segments (see, U.S. Patent Nos. 4,683,195 and 4,800,159; herein incorporated by reference).

Probes and primers are of sufficient nucleotide length to bind to the target DNA sequence specifically in the hybridization conditions or reaction conditions determined by the operator. This length may be of any length that is of sufficient length to be useful in a detection method of choice. Generally, 11 nucleotides or more in length, preferably 18 nucleotides or more, and more preferably 22 nucleotides or more, are used. Such probes and primers hybridize specifically to a target sequence under high stringency hybridization conditions.

Preferably, probes and primers according to the present invention have complete DNA sequence similarity of contiguous nucleotides with the target sequence, although probes differing from the target DNA sequence and that retain the ability to hybridize to target DNA sequences may be designed by conventional methods. Probes can be used as primers, but are generally designed to bind to the target DNA or RNA and not be used in an amplification process.

Specific primers can be used to amplify an integration fragment to produce an amplicon that can be used as a “specific probe” for identifying event TC1507 in biological samples. When the probe is hybridized with the nucleic acids of a biological sample under conditions which allow for the binding of the probe to the sample, this binding can be detected and thus allow for an indication of the presence of event TC1507 in the biological sample. Such identification of a bound probe has been described in the art. The specific probe is preferably a sequence which, under optimized conditions, hybridizes specifically to a region within the 5’ or 3’ flanking region of the event and preferably also comprises a part of the foreign DNA contiguous therewith. Preferably the specific probe comprises a sequence of at least 80%, preferably between 80 and 85%, more preferably between 85 and 90%, especially preferably between 90 and 95%, and most preferably between 95 and 100% identical (or complementary) to a specific region of the event.

Methods for preparing and using probes and primers are described, for example, in *Molecular Cloning: A Laboratory Manual*, 2<sup>nd</sup> ed., vol. 1-3, ed. Sambrook *et al.*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. 1989 (hereinafter, "Sambrook *et al.*, 1989"); *Current Protocols in Molecular Biology*, ed. Ausubel *et al.*, Greene Publishing and Wiley-Interscience, New York, 1992 (with periodic updates) (hereinafter, "Ausubel *et al.*, 1992"); and Innis *et al.*, *PCR Protocols: A Guide to Methods and Applications*, Academic Press: San Diego, 1990. PCR primer pairs can be derived from a known sequence, for example, by using computer programs intended for that purpose such as the PCR primer analysis tool in Vector NTI version 6 (Informax Inc., Bethesda MD); PrimerSelect (DNASTAR Inc., Madison, WI); and Primer (Version 0.5, © 1991, Whitehead Institute for Biomedical Research, Cambridge, Mass.). Additionally, the sequence can be visually scanned and primers manually identified using guidelines known to one of skill in the art.

A "kit" as used herein refers to a set of reagents for the purpose of performing the method of the invention, more particularly, the identification of the event TC1507 in biological samples. The kit of the invention can be used, and its components can be specifically adjusted, for purposes of quality control (e.g. purity of seed lots), detection of event TC1507 in plant material, or material comprising or derived from plant material, such as but not limited to food or feed products. "Plant material" as used herein refers to material which is obtained or derived from a plant.

Primers and probes based on the flanking DNA and insert sequences disclosed herein can be used to confirm (and, if necessary, to correct) the disclosed sequences by conventional methods, e.g., by re-cloning and sequencing such sequences. The nucleic acid probes and primers of the present invention hybridize under stringent conditions to a target DNA sequence. Any conventional nucleic acid hybridization or amplification method can be used to identify the presence of DNA from a transgenic event in a sample. Nucleic acid molecules or fragments thereof are capable of specifically hybridizing to other nucleic acid molecules under certain circumstances. As used herein, two nucleic acid molecules are said to be capable of specifically hybridizing to one another if the two molecules are capable of forming an anti-parallel, double-stranded nucleic acid structure.

A nucleic acid molecule is said to be the "complement" of another nucleic acid molecule if they exhibit complete complementarity. As used herein, molecules are said to exhibit "complete complementarity" when every nucleotide of one of the molecules is complementary

to a nucleotide of the other. Two molecules are said to be “minimally complementary” if they can hybridize to one another with sufficient stability to permit them to remain annealed to one another under at least conventional “low-stringency” conditions. Similarly, the molecules are said to be “complementary” if they can hybridize to one another with sufficient stability to permit them to remain annealed to one another under conventional “high-stringency” conditions. Conventional stringency conditions are described by Sambrook *et al.*, 1989, and by Haymes *et al.*, In: *Nucleic Acid Hybridization, a Practical Approach*, IRL Press, Washington, D.C. (1985), departures from complete complementarity are therefore permissible, as long as such departures do not completely preclude the capacity of the molecules to form a double-stranded structure. In order for a nucleic acid molecule to serve as a primer or probe it need only be sufficiently complementary in sequence to be able to form a stable double-stranded structure under the particular solvent and salt concentrations employed.

In hybridization reactions, specificity is typically the function of post-hybridization washes, the critical factors being the ionic strength and temperature of the final wash solution. The thermal melting point ( $T_m$ ) is the temperature (under defined ionic strength and pH) at which 50% of a complementary target sequence hybridizes to a perfectly matched probe. For DNA-DNA hybrids, the  $T_m$  can be approximated from the equation of Meinkoth and Wahl (1984) *Anal. Biochem.* 138:267-284:  $T_m = 81.5^\circ\text{C} + 16.6 (\log M) + 0.41 (\%GC) - 0.61 (\% \text{ form}) - 500/L$ ; where M is the molarity of monovalent cations, %GC is the percentage of guanosine and cytosine nucleotides in the DNA, % form is the percentage of formamide in the hybridization solution, and L is the length of the hybrid in base pairs.  $T_m$  is reduced by about  $1^\circ\text{C}$  for each 1% of mismatching; thus,  $T_m$ , hybridization, and/or wash conditions can be adjusted to hybridize to sequences of the desired identity. For example, if sequences with >90% identity are sought, the  $T_m$  can be decreased  $10^\circ\text{C}$ . Generally, stringent conditions are selected to be about  $5^\circ\text{C}$  lower than the  $T_m$  for the specific sequence and its complement at a defined ionic strength and pH. However, severely stringent conditions can utilize a hybridization and/or wash at 1, 2, 3, or  $4^\circ\text{C}$  lower than the  $T_m$ ; moderately stringent conditions can utilize a hybridization and/or wash at 6, 7, 8, 9, or  $10^\circ\text{C}$  lower than the  $T_m$ ; low stringency conditions can utilize a hybridization and/or wash at 11, 12, 13, 14, 15, or  $20^\circ\text{C}$  lower than the  $T_m$ .

Using the equation, hybridization and wash compositions, and desired  $T_m$ , those of ordinary skill will understand that variations in the stringency of hybridization and/or wash

5    *Molecular Biology—Hybridization with Nucleic Acid Probes*, Part I, Chapter 2 (Elsevier, New York); and Ausubel *et al.*, eds. (1995) *Current Protocols in Molecular Biology*, Chapter 2 (Greene Publishing and Wiley-Interscience, New York). See Sambrook *et al.* (1989) *Molecular Cloning: A Laboratory Manual* (2d ed., Cold Spring Harbor Laboratory Press, Plainview, New York).

Typically, stringent conditions will be those in which the salt concentration is less than about 1.5 M Na ion, typically about 0.01 to 1.0 M Na ion concentration (or other salts) at pH 7.0 to 8.3 and the temperature is at least about 30°C for short probes (e.g., 10 to 50 nucleotides) and at least about 60°C for long probes (e.g., greater than 50 nucleotides). Stringent conditions may also be achieved with the addition of a destabilizing agent such as formamide. Exemplary low stringency conditions include hybridization with a buffer solution of 30 to 35% formamide, 1 M NaCl, 1% SDS (sodium dodecyl sulphate) at 37°C, and a wash in 1X to 2X SSC (20X SSC = 3.0 M NaCl/0.3 M trisodium citrate) at 50 to 55°C. Exemplary moderate stringency conditions include hybridization in 40 to 45% formamide, 1 M NaCl, 1% SDS at 37°C, and a wash in 0.5X to 1X SSC at 55 to 60°C. Exemplary high stringency conditions include hybridization in 50% formamide, 1 M NaCl, 1% SDS at 37°C, and a wash in 0.1X SSC at 60 to 65°C. In a preferred embodiment, a nucleic acid of the present invention will specifically hybridize to one or more of the nucleic acid molecules unique to the TC1507 event or complements thereof or fragments of either under moderately stringent conditions.

algorithm of Myers and Miller (1988) *CABIOS* 4:11-17; the local homology algorithm of Smith *et al.* (1981) *Adv. Appl. Math.* 2:482; the homology alignment algorithm of Needleman and Wunsch (1970) *J. Mol. Biol.* 48:443-453; the search-for-similarity-method of Pearson and Lipman (1988) *Proc. Natl. Acad. Sci.* 85:2444-2448; the algorithm of Karlin and Altschul (1990) *Proc. Natl. Acad. Sci. USA* 87:2264, modified as in Karlin and Altschul (1993) *Proc. Natl. Acad. Sci. USA* 90:5873-5877.

Computer implementations of these mathematical algorithms can be utilized for comparison of sequences to determine sequence identity. Such implementations include, but are not limited to: CLUSTAL in the PC/Gene program (available from Intelligenetics, Mountain View, California); the ALIGN program (Version 2.0); the ALIGN PLUS program (version 3.0, copyright 1997); and GAP, BESTFIT, BLAST, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Version 10 (available from Accelrys, 9685 Scranton Road, San Diego, CA 92121, USA). Alignments using these programs can be performed using the default parameters.

The CLUSTAL program is well described by Higgins and Sharp, *Gene* 73: 237-244 (1988); Higgins and Sharp, *CABIOS* 5: 151-153 (1989); Corpet, *et al.*, *Nucleic Acids Research* 16: 10881-90 (1988); Huang, *et al.*, *Computer Applications in the Biosciences* 8: 155-65 (1992), and Pearson, *et al.*, *Methods in Molecular Biology* 24: 307-331 (1994). The ALIGN and the ALIGN PLUS programs are based on the algorithm of Myers and Miller (1988) *supra*. The BLAST programs of Altschul *et al.* (1990) *J. Mol. Biol.* 215:403 are based on the algorithm of Karlin and Altschul (1990) *supra*. The BLAST family of programs which can be used for database similarity searches includes: BLASTN for nucleotide query sequences against nucleotide database sequences; BLASTX for nucleotide query sequences against protein database sequences; BLASTP for protein query sequences against protein database sequences; TBLASTN for protein query sequences against nucleotide database sequences; and TBLASTX for nucleotide query sequences against nucleotide database sequences. See, *Current Protocols in Molecular Biology*, Chapter 19, Ausubel, *et al.*, Eds., Greene Publishing and Wiley-Interscience, New York (1995). Alignment may also be performed manually by inspection.

To obtain gapped alignments for comparison purposes, Gapped BLAST (in BLAST 2.0) can be utilized as described in Altschul *et al.* (1997) *Nucleic Acids Res.* 25:3389. Alternatively, PSI-BLAST (in BLAST 2.0) can be used to perform an iterated search that

detects distant relationships between molecules. See Altschul *et al.* (1997) *supra*. When utilizing BLAST, Gapped BLAST, PSI-BLAST, the default parameters of the respective programs (e.g., BLASTN for nucleotide sequences, BLASTX for proteins) can be used. See [www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov).

5 As used herein, “sequence identity” or “identity” in the context of two nucleic acid or polypeptide sequences makes reference to the residues in the two sequences that are the same when aligned for maximum correspondence over a specified comparison window. When percentage of sequence identity is used in reference to proteins it is recognized that residue positions which are not identical often differ by conservative amino acid substitutions, where  
 10 amino acid residues are substituted for other amino acid residues with similar chemical properties (e.g., charge or hydrophobicity) and therefore do not change the functional properties of the molecule. When sequences differ in conservative substitutions, the percent sequence identity may be adjusted upwards to correct for the conservative nature of the substitution. Sequences that differ by such conservative substitutions are said to have  
 15 “sequence similarity” or “similarity”. Means for making this adjustment are well known to those of skill in the art. Typically this involves scoring a conservative substitution as a partial rather than a full mismatch, thereby increasing the percentage sequence identity. Thus, for example, where an identical amino acid is given a score of 1 and a non-conservative substitution is given a score of zero, a conservative substitution is given a score between zero  
 20 and 1. The scoring of conservative substitutions is calculated, e.g., as implemented in the program PC/GENE (Intelligenetics, Mountain View, California).

As used herein, “percentage of sequence identity” means the value determined by comparing two optimally aligned sequences over a comparison window, wherein the portion of the polynucleotide sequence in the comparison window may comprise additions or deletions  
 25 (i.e., gaps) as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. The percentage is calculated by determining the number of positions at which the identical nucleic acid base or amino acid residue occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison, and  
 30 multiplying the result by 100 to yield the percentage of sequence identity.

Regarding the amplification of a target nucleic acid sequence (e.g., by PCR) using a particular amplification primer pair, “stringent conditions” are conditions that permit the primer

pair to hybridize only to the target nucleic-acid sequence to which a primer having the corresponding wild-type sequence (or its complement) would bind and preferably to produce a unique amplification product, the amplicon, in a DNA thermal amplification reaction.

The term “specific for (a target sequence)” indicates that a probe or primer hybridizes  
5 under stringent hybridization conditions only to the target sequence in a sample comprising the target sequence.

As used herein, “amplified DNA” or “amplicon” refers to the product of nucleic acid amplification of a target nucleic acid sequence that is part of a nucleic acid template. For example, to determine whether a corn plant resulting from a sexual cross contains transgenic  
10 event genomic DNA from the corn plant of the present invention, DNA extracted from the corn plant tissue sample may be subjected to a nucleic acid amplification method using a DNA primer pair that includes a first primer derived from flanking sequence adjacent to the insertion site of inserted heterologous DNA, and a second primer derived from the inserted heterologous DNA to produce an amplicon that is diagnostic for the presence of the event DNA.

15 Alternatively, the second primer may be derived from the flanking sequence. The amplicon is of a length and has a sequence that is also diagnostic for the event. The amplicon may range in length from the combined length of the primer pairs plus one nucleotide base pair to any length of amplicon producible by a DNA amplification protocol. Alternatively, primer pairs can be derived from flanking sequence on both sides of the inserted DNA so as to produce an  
20 amplicon that includes the entire insert nucleotide sequence of the PHI8999A expression construct, see FIG. 1, approximately 6.2 Kb in size. A member of a primer pair derived from the flanking sequence may be located a distance from the inserted DNA sequence, this distance can range from one nucleotide base pair up to the limits of the amplification reaction, or about twenty thousand nucleotide base pairs. The use of the term “amplicon” specifically excludes  
25 primer dimers that may be formed in the DNA thermal amplification reaction.

Nucleic acid amplification can be accomplished by any of the various nucleic acid amplification methods known in the art, including the polymerase chain reaction (PCR). A variety of amplification methods are known in the art and are described, inter alia, in U.S. Pat. Nos. 4,683,195 and 4,683,202 and in *PCR Protocols: A Guide to Methods and Applications*,  
30 ed. Innis *et al.*, Academic press, San Diego, 1990. PCR amplification methods have been developed to amplify up to 22 Kb of genomic DNA and up to 42 Kb of bacteriophage DNA (Cheng *et al.*, *Proc. Natl. Acad. Sci. USA* 91:5695-5699, 1994). These methods as well as other

methods known in the art of DNA amplification may be used in the practice of the present invention. It is understood that a number of parameters in a specific PCR protocol may need to be adjusted to specific laboratory conditions and may be slightly modified and yet allow for the collection of similar results. These adjustments will be apparent to a person skilled in the art.

5           The amplicon produced by these methods may be detected by a plurality of techniques, including, but not limited to, Genetic Bit Analysis (Nikiforov, *et al. Nucleic Acid Res.* 22:4167-4175, 1994) where a DNA oligonucleotide is designed which overlaps both the adjacent flanking DNA sequence and the inserted DNA sequence. The oligonucleotide is immobilized in wells of a microwell plate. Following PCR of the region of interest (using one primer in the  
10 inserted sequence and one in the adjacent flanking sequence) a single-stranded PCR product can be hybridized to the immobilized oligonucleotide and serve as a template for a single base extension reaction using a DNA polymerase and labeled ddNTPs specific for the expected next base. Readout may be fluorescent or ELISA-based. A signal indicates presence of the insert/flanking sequence due to successful amplification, hybridization, and single base  
15 extension.

          Another detection method is the Pyrosequencing technique as described by Winge (*Innov. Pharma. Tech.* 00: 18-24, 2000). In this method an oligonucleotide is designed that overlaps the adjacent DNA and insert DNA junction. The oligonucleotide is hybridized to a single-stranded PCR product from the region of interest (one primer in the inserted sequence  
20 and one in the flanking sequence) and incubated in the presence of a DNA polymerase, ATP, sulfurylase, luciferase, apyrase, adenosine 5' phosphosulfate and luciferin. dNTPs are added individually and the incorporation results in a light signal which is measured. A light signal indicates the presence of the transgene insert/flanking sequence due to successful amplification, hybridization, and single or multi-base extension.

25           Fluorescence Polarization as described by Chen *et al.*, (*Genome Res.* 9:492-498, 1999) is a method that can be used to detect an amplicon of the present invention. Using this method an oligonucleotide is designed which overlaps the flanking and inserted DNA junction. The oligonucleotide is hybridized to a single-stranded PCR product from the region of interest (one primer in the inserted DNA and one in the flanking DNA sequence) and incubated in the  
30 presence of a DNA polymerase and a fluorescent-labeled ddNTP. Single base extension results in incorporation of the ddNTP. Incorporation can be measured as a change in polarization using a fluorometer. A change in polarization indicates the presence of the transgene

insert/flanking sequence due to successful amplification, hybridization, and single base extension.

Taqman® (PE Applied Biosystems, Foster City, Calif.) is described as a method of detecting and quantifying the presence of a DNA sequence and is fully understood in the instructions provided by the manufacturer. Briefly, a FRET oligonucleotide probe is designed which overlaps the flanking and insert DNA junction. The FRET probe and PCR primers (one primer in the insert DNA sequence and one in the flanking genomic sequence) are cycled in the presence of a thermostable polymerase and dNTPs. Hybridization of the FRET probe results in cleavage and release of the fluorescent moiety away from the quenching moiety on the FRET probe. A fluorescent signal indicates the presence of the flanking/transgene insert sequence due to successful amplification and hybridization.

Molecular Beacons have been described for use in sequence detection as described in Tyangi *et al.* (*Nature Biotech.* 14:303-308, 1996). Briefly, a FRET oligonucleotide probe is designed that overlaps the flanking and insert DNA junction. The unique structure of the FRET probe results in it containing secondary structure that keeps the fluorescent and quenching moieties in close proximity. The FRET probe and PCR primers (one primer in the insert DNA sequence and one in the flanking sequence) are cycled in the presence of a thermostable polymerase and dNTPs. Following successful PCR amplification, hybridization of the FRET probe to the target sequence results in the removal of the probe secondary structure and spatial separation of the fluorescent and quenching moieties. A fluorescent signal results. A fluorescent signal indicates the presence of the flanking/transgene insert sequence due to successful amplification and hybridization.

A hybridization reaction using a probe specific to a sequence found within the amplicon is yet another method used to detect the amplicon produced by a PCR reaction.

The present invention is further defined in the following Examples. It should be understood that these Examples, while indicating preferred embodiments of the invention, are given by way of illustration only. From the above discussion and these Examples, one skilled in the art can ascertain the essential characteristics of this invention, and without departing from the spirit and scope thereof, can make various changes and modifications of the invention to adapt it to various usages and conditions. Thus, various modifications of the invention, in addition to those shown and described herein, will be apparent to those skilled in the art from

the foregoing description. Such modifications are also intended to fall within the scope of the appended claims.

The disclosure of each reference set forth herein is incorporated herein by reference in its entirety.

5

## EXAMPLES

### Example 1. Transformation of Maize by Particle Bombardment and Regeneration of Transgenic Plants Containing the Cry1F Gene

- 10 A DNA molecule of 6.2 Kb, designated PHI8999A (see FIG. 1 and SEQ ID NO:25), which includes a first transgene expression cassette comprising the promoter, 5' untranslated exon, and first intron of the maize ubiquitin (Ubi-1) gene (Christensen *et al.* (1992) *Plant Mol. Biol.* 18:675-689 and Christensen and Quail (1996) *Transgenic Res.* 5:213-218) operably connected to a DNA molecule encoding a *Bacillus thuringiensis*  $\delta$ -endotoxin identified as
- 15 Cry1F (U.S. Pat. Nos. 5,188,960 and 6,218,188) operably connected to a DNA molecule comprising a 3' ORF25 transcriptional terminator isolated from *Agrobacterium tumefaciens* (Barker *et al.* (1983) *Plant Mol. Biol.* 2:335-350), and a second transgene expression cassette comprising a DNA molecule of the cauliflower mosaic virus (CaMV) 35S promoter (Odell J.T. *et al.* (1985) *Nature* 313: 810-812; Mitsuhashi *et al.* (1996) *Plant Cell Physiol.* 37:49-59)
- 20 operably connected to a DNA molecule encoding the selectable marker, phosphinothricin acetyltransferase (PAT) gene (Wohllleben W. *et al.* (1988) *Gene* 70:25-37) operably connected to a DNA molecule comprising a 3' transcriptional terminator from (CaMV) 35S (see Mitsuhashi *et al.* (1996) *Plant Cell Physiol.* 37:49-59) was used to transform maize embryo tissue.
- 25 *B.t.* Cry1F maize plants were obtained by microprojectile bombardment using the Biolistics® PDS-1000He particle gun manufactured by Bio-Rad, Hercules, CA; essentially as described by Klein *et al.* (1987) *Nature*, UK 327(6117):70-73. Immature embryos isolated from maize ears, harvested soon after pollination were cultured on callus initiation medium for several days. On the day of transformation, microscopic tungsten particles were coated with
- 30 purified PHI8999A DNA (SEQ ID NO:25) and accelerated into the cultured embryos, where the insert DNA was incorporated into the cell chromosome. Only insert PHI8999A was used during transformation and no additional plasmid DNA was incorporated into the transformant.

After bombardment, embryos were transferred to callus initiation medium containing glufosinate as the selection agent. Individual embryos were kept physically separate during culture, and the majority of explants died on the selective medium.

Those embryos that survived and produced healthy, glufosinate-resistant callus tissue were assigned unique identification codes representing putative transformation events, and continually transferred to fresh selection medium. Plants were regenerated from tissue derived from each unique event and transferred to the greenhouse. Leaf samples were taken for molecular analysis to verify the presence of the transgene by PCR and to confirm expression of the Cry1F protein by ELISA. Plants were then subjected to a whole plant bioassay using European corn borer insects. Positive plants were crossed with inbred lines to obtain seed from the initial transformed plants. A number of lines were evaluated in the field. The TC1507 event was selected from a population of independent transgenic events based on a superior combination of characteristics, including insect resistance and agronomic performance (see Bing JW *et al.* (2000) Efficacy of Cry1F Transgenic Maize, 14<sup>th</sup> Biennial International Plant Resistance to Insects Workshop, Fort Collins, CO, herein incorporated by reference).

Example 2. Identification of nucleotides comprising the flanking sequence 5' to the transgenic insert DNA in *Bacillus thuringiensis* Cry1F maize line TC1507

To identify a DNA fragment that included sequence 5' to the PHI8999A insert in event TC1507, Spe I restriction enzyme fragments from event TC1507 genomic DNA were size selected on agarose gels, purified, and screened by Southern analysis to confirm hybridization to a Cry1F probe. Following confirmation of hybridization and fragment size, the fragments of interest were cloned into a pBluescript II SK (+)<sup>TM</sup> cloning vector to prepare an enriched size selected plasmid based genomic DNA library. A probe homologous to a portion of the Cry1F gene was used to screen the plasmid library for positive clones. A positive clone was identified, purified by additional screening, and confirmed to result in a positive signal when hybridized to the Cry1F probe. Nearly 3 Kb of the Spe I fragment contained in the isolated positive clone was sequenced using a primer walking approach. To initiate the first sequencing run, a primer that binds to a known sequence in the cloning vector DNA was designed to sequence a portion of the DNA of interest. A second sequencing run over the same region using another primer oriented in the reverse direction provided second strand coverage. Primer walking was accomplished by repeatedly using sequence data from previous runs to design new

primers that were then used to extend the next round of sequencing further into the DNA of interest until the flanking sequence 5' to the inserted transgenic DNA in maize event TC1507 was obtained. Specific sequence information is provided in Example 4.

5 Example 3. Confirmation of flanking sequence 5' to the *B.t.* Cry1F maize line TC1507 insert

To confirm the 5' flanking sequence of the *B.t.* Cry1F maize line TC1507 insert, PCR primer pairs were designed to obtain overlapping PCR products extending from the 5' flanking region into the full-length PHI8999A transgenic insert. PCR products were successfully  
10 amplified from *B.t.* Cry1F maize line TC1507 genomic DNA, isolated, and sequenced for Region 1 through Region 6, shown in Table 1, and confirmed to match the previously determined sequence from the Spe I fragment, described in Example 2. However, the region from bp 2358 to bp 2829, immediately adjacent and 5' to the start of the full-length insert was recalcitrant to PCR amplification and appeared to be larger than the sequence obtained from the  
15 Spe I clone described above. The use of primer pairs flanking this region and the Advantage®-GC 2 Polymerase Mix (BD Biosciences Clontech, Palo Alto, CA) was successful in amplifying PCR products from *B.t.* Cry1F maize line TC1507 genomic DNA for sequencing. The amplification conditions used to produce amplicons with the Advantage®-GC 2 system are shown in Table 10. The DNA primer pairs used to confirm the sequence in the region from bp  
20 2358 to 2829 are those listed in SEQ ID NO:1 and SEQ ID NO:2; and SEQ ID NO:2 and SEQ ID NO:23. Sequence from this region is described in Table 1 (Regions 7a, 7b, 7c, and 8).

Example 4. Event TC1507 5' flanking sequence. A description of each region is provided in Table 1.

25

**Region 1 (SEQ ID NO: 28) Maize genomic (no significant homology)**

1 ACTAGTTTCC TAGCCCGCGT CGTGCCCTA CCCCACCGAC GTTTATGGAA  
51 GGTGCCATTC CACGGTCTT CGTGCCGCC CCTAAGGATG TAAATGGTCG  
101 GTAAAATCCG GTAAATTTCC GGTACCGTTT ACCAGATTTT TCCAGCCGTT  
30 151 TTCGGATTTA TCGGGATATA CAGAAAACGA GACGGAAACG GAATAGGTTT  
201 TTTTTCGAAA ACGGTACGGT AAACGGTGAG ACAAACCTAC CGTCCGTTTT  
251 CGTATTTCTC GGGAAACTCT GGTATATTCC CGTATTGTG CCGTATTTTC  
301 CCGACCCACG GACCTGCCAA TCAACCATCA GCCAGTCAGC CCATCCCCAC  
351 AGCTATGGCC CATGGGGCCA TGTTGGCCAC ATGCCACGC AACGCAAGGC  
35 401 AGTAAGGCTG GCAGCCTGGC ACGCATTGAC GCATGTGGAC ACACACAGCC  
451 GCCGCTGTT CGTGTCTCTG TGCCGTTGTG CGAGACTGTG ACTGCGAGTG  
501 GCGGAGTCGG CGAACGGCGA GCGTCTCCG GAGTCTGGAC TGCGGCTGTG

551 GACAGCGACG CTGTGACGGC GACTCGGCGA AGCCCCAAGC TACCAAGCCC  
 601 CCAAGTCCCC ATCCATCTCT GCTTCTCTGG TCATCTCCTT CCCCTGGTCG  
 651 ATCTGCAGGC GCCAGACCG

5 **Region 2 (SEQ ID NO: 29) Undescribed maize genomic sequence (complement)**

670 G CCGAAGCATC ACGAAACGCA CTAAGACCTC  
 701 GAAGGAGTCA AACCCTCCT CCGAGGCCTC GGGGGCTACA CCCGGCGGGT  
 751 GCGCTCGCGC GCACCCACCG GAACAAAATG TAACCGAGAA AGGTCGGTCC  
 801 CCTTGCAAAA AAAGTGCGAC AAAAGCCTCC AAGCGAGTAT TAACACTCAC  
 10 851 TTTGAGGCTC GGGGGCTAC

**Region 3 (SEQ ID NO: 30) Fragment of maize Huck-1 retrotransposon**

870 T GTCGGGGACC ATAATTAGGG GTACCCCCAA  
 901 GACTCCTAAT CTCAGCTGGT AACCCCCATC AGCACAAAGC TGCAAAGGCC  
 15 951 TGATGGGTGC GATTAAGTCA AGGCTCGGTC CACTCAAGGG ACACGATCTC  
 1001 GCCTCGCCCG AGCCCAGCCT CGGGCAAGGG CGGCCGACCC CGAGGATTCA  
 1051 CGTCTCGCCC GAGGGCCCCC TCAAGCGACG GGCACACCTT CGGCTCGCCC  
 1101 GAGGCCATT CTTCGCCGAG AAGCAACCTT GGCCAGATCG CCACACCGAC  
 1151 CGACCGTATC GCAGGAGCAT TTAATGCGAG GATCGCCTGA CACCTTATCC  
 20 1201 TGACGCGCGC TCTTCAGTCG ACAGAGCCGA AGTGACCGCA ATCACTTCGC  
 1251 CGCTCCACTG ACCGACCTGA CAAGAAGACA GCGCCGCCTG CGTCGCTCCG  
 1301 ACTGCTGTGC CACTCGACAG AGTGAGGCTG ACAGCAGCCA AGTCCGGCCT  
 1351 CGGGCGCCAT AGGAAGCTCC GCCTCGCCCG ACCCTAGGGC TCGGACTCGG  
 1401 CCTCGGCTCC GGAAGACGAC GAACTACGCT TCGCCCGACC CCAGGGCTTG  
 25 1451 GACTCAGCCT CGGCTCCGGA AGACGACGAA TTCCGCCTCG CCCGACCCCA  
 1501 GGGCTCGGAC TCGGCCTCGG CTCGAGAAGA CGACGAACTC CGCCTCGCCC  
 1551 GACCCAGGG CTCGGACTCA GCCTCGGCTC CGAAGACGA CGAACTCCGC  
 1601 CTCGCCCGAC CCCAGGGCTC GGACTCAGCC TCGGCCTCAG ACGATGGTCT  
 1651 CCGCCTCGCC CGACCCGGGG CTCGGACTCG A

30

**Region 4 (SEQ ID NO: 31) Fragment of *cry1F* gene**

1682 CCTTTCTAT CGGACCTTGT  
 1701 CAGATCCTGT CTTCGTCCGA GGAGGCTTTG GCAATCCTCA CTATGTACTC  
 1751 GGTCTTAGGG GAGTGGCCTT TCAACAACT GGTACGAATC ACACCCGCAC  
 35 1801 ATTCAGGAAC TCCGGGACCA TTGACTCTCT AGATGAGATA CCACCTCAAG  
 1851 ACAACAGCGG CGCACCTTGG AATGACTACT CCCATGTGCT GAATCATGTT  
 1901 ACCTTTGTGC GCTGGCCAGG TGAGATCTCA GGTTCGACT CATGGAGAGC  
 1951 ACCAATGTTT TCTTGACGC ATCGTAGCGC TACCCCAACA AACACCATTG  
 2001 ATCCAGAGAG AATCAC

40

**Region 5 (SEQ ID NO: 32) Fragment of maize chloroplast *rpoC2* gene**

2017 TCAT TCTTCAAGAA CTGCATATCT TGCCGAGATC  
 2051 CTCATCCCTA AAGGTACTTG ACAATAGTAT TATTGGAGTC GATACACAAC  
 2101 TCACAAAAAA TACAAGAAGT CGACTAGGTG GATTGGTCCG  
 45 AGTGAAGAGA  
 2151 AAAAAAAGCC ATACAGAAGT CAAAATCTTT TCCGGAGATA TTCATTTTCC

- 2201 TGAAGAGGCG GATAAGATAT TAGGTGGCAG TTTGATACCA  
CCAGAAAGAG  
2251 AAAAAAAGA TTCTAAGGAA TCAAAAAA GGAAAAATTG  
GGTTTATGTT  
5 2301 CAACGGAAAA AATTTCTCAA AAGCAAGGAA AAGTATT
- Region 6 (SEQ ID NO: 33) Fragment of maize chloroplast or *ubiZM1(2)* promoter**  
2338 GTG GCTATTTATC  
2351 TATC  
10
- Nucleotides 2355-2358 (CGT) connect Region 6 to Region 7a.**
- Region 7a (SEQ ID NO: 34) Fragment of *pat* gene**  
2358 GCA GCTGATATGG CCGCGGTTTG TGATATCGTT AACCATTACA  
15 2401 TTGAGACGTC TACAGTGAAC TTAGGACAG AGCCACAAAC ACCACAAGAG  
2451 TGGATTGATG ATCTAGAGAG GTTGCAAGAT AGATACCCTT GGTTGGTTGC  
2501 TGAGGTTGAG GGTGTTGTGG CTGGTATTGC TTACGCTGGG CCCTGGAAGG  
2551 CTAGGAAC
- 20 **Region 7b (SEQ ID NO: 35) Fragment of *pat* gene (complement)**  
2559 CC TCAACCTCAG CAACCAACCA ATGGTATCTA TCTTGCAACC  
2601 TCTCTAGATC ATCAATCCAC TCTTGTTGGTG TTTGTGGCTC TGTCCTAAAG  
2651 TTCCTGTAG ACGTCTCAAT GTAATGGTTA ACGATATCAC AAACCG
- 25 **Region 7c (SEQ ID NO: 36) Fragment of *cry1F* gene (complement)**  
2697 AGAG  
2701 AAGAGGGATC T
- Region 8 (SEQ ID NO: 37) Fragment of Polylinker**  
30 2712 CGAAGCTTC GGCCGGGGCC CATCGATATC CGCGGGCATG  
2751 CCTGCAGTGC AGCGTGACCC GGTCGTGCCC CTCTCTAGAG ATAATGAGCA  
2801 TTGCATGTCT AAGTTATAAA AAATTACCA
- 35 **Region 9 (SEQ ID NO: 25) Full-length insert of PHI8999A**

Example 5. Description of the flanking sequence 5' to the insert in maize event TC1507

- 40 In order to more fully describe the event TC1507 5' flanking sequence, homology  
searching was done against the GenBank public databases (release 122, 2/01) using the Basic  
Local Alignment Search Tool (BLAST). The BLAST program performs sequence similarity  
searching and is particularly useful for identifying homologs to an unknown sequence. In  
addition to searching the public databases, pairwise alignments were performed using AlignX  
(InforMax Inc., Bethesda, MD) to look for homology between the maize event TC1507

flanking sequence and the PHI8999A transgenic insert. The results of these homology searches are presented in Table 1. The TC1507 5' flanking sequence is numbered with base 1 being the furthest 5' to the insert and base 2830 at the starting point of the full-length PHI8999A transgenic insert (see FIG. 1). The percent identity values indicate the percentage of identical matches across the length of the sequences analyzed.

In most cases, similarity searching with the event TC1507 5' flanking sequence resulted in a match to one unique sequence based on a very high percent identity value. Those sequences are identified in Table 1. In addition, there are two regions in the TC1507 5' DNA flanking sequence with high similarity to more than one known sequence. In regions 870-1681 and 2338-2354, the percent identity scores with both sequence fragments are sufficiently high that a single match (homolog) cannot be determined. The two possible homologs for each of these regions are indicated in Table 1.

Highly similar sequences were identified for all but the first 669 base pairs of sequence. Generally, the results of similarity searching indicate high homology with maize genomic sequences 5' to base 1681. The region from base 1682 to the start of the PHI8999A insert at position 2830 contains some fragments associated with the transformation event.

**Table 1. Sequence summary for event TC1507 insert**

| Region | Location in SEQ ID NO:24 | Size bp | % Identity       | Homolog  | Location in homologous sequence | Description                                                  |
|--------|--------------------------|---------|------------------|----------|---------------------------------|--------------------------------------------------------------|
| 1      | 1-669                    | 669     | N/A <sup>1</sup> | N/A      | N/A                             | No significant homology detected                             |
| 2      | 670-869                  | 200     | 90.5             | AF123535 | 52432-52632 (complement)        | Undescribed maize genomic sequence                           |
| 3      | 870-1681                 | 812     | 89.4             | AF050439 | 1-801                           | Fragment of maize Huck-1 retrotransposon 5' LTR <sup>2</sup> |
|        |                          |         | 86.6             | AF050438 | 1-797                           | Fragment of maize Huck-1 retrotransposon 3' LTR              |

| Region | Location in SEQ ID NO:24 | Size bp | % Identity | Homolog   | Location in homologous sequence            | Description                                                                              |
|--------|--------------------------|---------|------------|-----------|--------------------------------------------|------------------------------------------------------------------------------------------|
| 4      | 1682-2016                | 335     | 100.0      | PHI8999A  | 3149-3483                                  | Fragment of <i>cry1F</i> gene                                                            |
| 5      | 2017-2337                | 321     | 100.0      | X86563    | 29429-29749                                | Fragment of maize chloroplast <i>rpoC2</i> gene (RNA polymerase beta-2 subunit)          |
| 6      | 2338-2354                | 17      | 100.0      | X86563    | 97643-97659                                | Fragment of maize chloroplast <i>trnI</i> gene (tRNA-Ile)                                |
|        |                          |         | 82.4       | PHI8999A  | 182-197                                    | Fragment of maize <i>ubiZM1(2)</i> promoter                                              |
| 7a     | 2358-2558                | 201     | 100.0      | PHI8999A  | 5320-5475                                  | Fragment of <i>pat</i> gene                                                              |
| 7b     | 2559-2696                | 138     | 99         | PHI8999A  | 5336-5518 (complement)                     | Fragment of <i>pat</i> gene                                                              |
| 7c     | 2697-2711                | 15      | 100.0      | PHI8999A  | 2544-2558 (complement)                     | Fragment of <i>cry1F</i> gene                                                            |
| 8      | 2712-2829                | 118     | 100.0      | PHI8999A  | 36-153                                     | Fragment of polylinker region (bases 36-80) and <i>ubiZM1(2)</i> promoter (bases 81-153) |
| 9      | 2830-9015                | 6186    | 100.0      | PHI8999A  | 11 – 6196                                  | Full-length insert of PHI8999A                                                           |
| 10     | 9016-9565                | 550     | 100.0      | PHI8999A  | 3906-4456 (complement)                     | Inverted ORF25 terminator                                                                |
| 11     | 9566-9693                | 128     | 100.0      | NC_001666 | 121851-121978 (complement) & 100759-100886 | Fragment of maize chloroplast <i>rps12</i> rRNA (23S ribosomal RNA)                      |
| 12     | 9696-10087               | 392     | 99         | NC_001666 | 17091-17483 (complement)                   | Fragment of maize chloroplast genome                                                     |
| 13     | 10088-10275              | 188     | 99         | PHI8999A  | 5333-5520 (complement)                     | Fragment of <i>pat</i> gene                                                              |

| Region | Location in SEQ ID NO:24 | Size bp | % Identity       | Homolog   | Location in homologous sequence | Description                                                        |
|--------|--------------------------|---------|------------------|-----------|---------------------------------|--------------------------------------------------------------------|
| 14     | 10278-10358              | 81      | 100              | NC_001666 | 137122-137202 (complement)      | Fragment of maize chloroplast "ORF241" – hypothetical protein gene |
| 15     | 10359-10612              | 254     | N/A <sup>1</sup> | N/A       | N/A                             | No significant homology detected                                   |
| 16     | 10613-11361              | 749     | N/A <sup>1</sup> | N/A       | N/A                             | No description available                                           |

<sup>1</sup> N/A; not applicable

<sup>2</sup> LTR; long terminal repeat

5

Example 6: Confirmation of the presence of Regions 1, 2, and 3 in an unmodified control corn line

10 PCR analysis was used to determine if Regions 1, 2, and 3 (Table 1) in the 5' flanking region of Event TC1507 are present in an unmodified control corn line used for transformation to produce maize event TC1507 and thus represents a border with corn genomic DNA. Nine different PCR analyses were carried out on genomic DNA prepared from TC1507 and the unmodified control corn line Hi-II (see Armstrong (1994) *The Maize Handbook*, ed. Freeling and Walbot, Springer-Verlag, New York, pp. 663-671, for information on Hi-II) as outlined in

15 Table 2 using the primer sequences shown in Table 3. Two reactions were designed to amplify DNA within Region 1 of the 5' flanking region from bp 25 to 324 (Reaction A - 300 bp amplicon); and from bp 25 to 480 (Reaction B - 456 bp amplicon). The expected amplicons were present in both the Hi-II unmodified corn line and in maize event TC1507. One PCR primer pair, Reaction C, spanned Region 2 to Region 3 of the 5' flanking region from bp 759 to

20 1182 (424 bp amplicon) and again produced PCR products of the expected size in both Hi-II and TC1507. Reaction D, spanned Region 1 to Region 3 of the 5' flanking region from bp 415 to 1182 (768 bp amplicon) and again produced PCR products of the expected size in both Hi-II and TC1507. Reactions E and F were designed as specific primer pairs for the *pat* gene region of the full-length insert of PHI8999A in TC1507 and thus an amplicon in the unmodified Hi-II

25 corn line is not expected. The results indicate that both Reactions E and F are specific for a

maize line transformed with a *pat* gene region and produce the expected amplicon, whereas no amplicon was produced in the unmodified Hi-II corn line. Reaction G was also designed as a primer pair that would produce an amplicon of 366 bp in the maize event TC1507 and no amplicon in the unmodified Hi-II corn line.

- 5 Reactions H and I were designed as specific primer pairs for TC1507 that would span the end of the transgenic insert into the 5' flanking region. In both Reactions H and I, the reverse primer was located in the ubiquitin promoter region of the full-length PHI8999A insert (Region 9 in Table 1) and the forward primer was located in Region 5, the *rpoC2* gene fragment (see Table 1). Reaction H and Reaction I both produced an amplicon in maize line  
10 TC1507 and did not produce an amplicon in the unmodified control corn line. These results indicate that both Reactions H and I are specific for the TC1507 event.

- The PCR results show that the undescribed sequence (Region 1) is present in the unmodified corn line Hi-II and that Regions 1, 2 and 3, are contiguous in the unmodified corn line Hi-II. The DNA sequences amplified in Reactions A, B, C, and D are not unique to the 5'  
15 flanking region of maize event TC1507 but are also present in the unmodified corn line Hi-II.

**Table 2. PCR reactions for sequence 5' to the PHI8999A insert in maize event TC1507 and for regions within the full-length insert of PHI8999A in maize event TC1507**

| Reaction | PCR Amplicon Location                     | Amplicon Size (bp) | Region in TC1507 flanking sequence or PHI8999A insert | Amplicon present In Hi-II | Amplicon present in maize line TC1507 |
|----------|-------------------------------------------|--------------------|-------------------------------------------------------|---------------------------|---------------------------------------|
| A        | 25 – 324 bp in TC1507 flanking sequence   | 300                | Region 1                                              | Yes                       | Yes                                   |
| B        | 25 – 480 bp in TC1507 flanking sequence   | 456                | Region 1                                              | Yes                       | Yes                                   |
| C        | 759 – 1182 bp in TC1507 flanking sequence | 424                | Region 2 to Region 3                                  | Yes                       | Yes                                   |

20

| Reaction                                                        | PCR Amplicon Location                                                                                     | Amplicon Size (bp) | Region in TC1507 flanking sequence or PHI8999A insert                                                  | Amplicon present In Hi-II | Amplicon present in maize line TC1507 |
|-----------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------|--------------------|--------------------------------------------------------------------------------------------------------|---------------------------|---------------------------------------|
| <b>D</b>                                                        | 415 – 1182 bp in TC1507 5' flanking sequence                                                              | 768                | Region 1 to Region3                                                                                    | Yes                       | Yes                                   |
| <b>E</b><br>Not Unique to TC1507                                | 4750 – 5794 bp in PHI8999A                                                                                | 1045               | Region 9 (in full-length insert of PHI8999A 35S promoter to <i>pat</i> gene)                           | No                        | Yes                                   |
| <b>F</b><br>Not Unique to TC1507                                | 4827 - 5308 bp in PHI8999A                                                                                | 482                | Region 9 (in full-length insert of PHI8999A 35S promoter to <i>pat</i> gene)                           | No                        | Yes                                   |
| <b>G</b><br>Detects <i>cry1F</i> fragment in 5' flanking region | <i>cry1F</i> sequence in 5' flanking sequence and in full-length insert of PHI8999A                       | 366                | Spans 335 bp <i>cry1F</i> sequence in 5' flanking sequence and same sequence in the full-length insert | No                        | Yes                                   |
| <b>H</b><br>Unique to TC1507                                    | 2158 bp in Region 5 ( <i>rpoC2</i> gene fragment) to 3069 bp in Region 9 (full-length insert of PHI8999A) | 912                | Region 5 to Region 9<br>Unique to Insertion Event<br>[SPANS UNIQUE JUNCTION REGIONS]                   | No                        | Yes                                   |
| <b>I</b><br>Unique to TC1507                                    | 2158 bp in Region 5 ( <i>rpoC2</i> gene fragment) to 3001 bp in Region 9 (full-length insert of PHI8999A) | 844                | Region 5 to Region 9<br>Unique to Insertion Event<br>[SPANS UNIQUE JUNCTION REGIONS]                   | No                        | Yes                                   |

**Table 3. PCR primers for sequence 5' to the PHI8999A insert in TC1507 and for regions within the full-length insert of PHI8999A in maize event TC1507**

| Reaction | Amplicon Size (bp) | Primer Pair  | Primer Sequences<br>5' to 3'       |
|----------|--------------------|--------------|------------------------------------|
| A        | 300                | SEQ ID NO:10 | CCCCTACCCACCGACGTTTAT              |
|          |                    | SEQ ID NO:11 | TTGATTGGCAGGTCCGTGGGTC             |
| B        | 456                | SEQ ID NO:10 | CCCCTACCCACCGACGTTTAT              |
|          |                    | SEQ ID NO:12 | CACAACGGCACAGAAACACGAA             |
| C        | 424                | SEQ ID NO:13 | GCGCACCCACCGGAACAAAATG             |
|          |                    | SEQ ID NO:14 | TCCTCGCATTAAATGCTCCTGC             |
| D        | 768                | SEQ ID NO:15 | CCTGGCACGCATTGACGCATGT             |
|          |                    | SEQ ID NO:14 | TCCTCGCATTAAATGCTCCTGC             |
| E        | 1045               | SEQ ID NO:6  | TAGAGGACCTAACAGAACTCGCCGT          |
|          |                    | SEQ ID NO:7  | GAGCTGGCAACTCAAAATCCCTTT           |
| F        | 482                | SEQ ID NO:8  | AAAATCTTCGTCAACATGGTGGAGC          |
|          |                    | SEQ ID NO:9  | TAATCTCAACTGGTCTCCTCTCCGG          |
| G        | 366                | SEQ ID NO:19 | GGCTCGGACTCGACCTTTCTAT             |
|          |                    | SEQ ID NO:20 | GCAGTTCTTGAAGAATGAGTGA             |
| H        | 912                | SEQ ID NO:1  | GTAGTACTATAGATTATATTATTCGT<br>AGAG |
|          |                    | SEQ ID NO:2  | GCCATACAGAACTCAAAATCTTTTCC<br>GGAG |
| I        | 844                | SEQ ID NO:2  | GCCATACAGAACTCAAAATCTTTTCC<br>GGAG |
|          |                    | SEQ ID NO:23 | CTTCAAACAAGTGTGACAAA               |

10 Example 7. Flanking sequence 3' to inserted transgenic DNA in maize event TC1507

Two separate PCR approaches were used to extend the length of the sequence information 3' to the full-length PHI8999A insert in maize event TC1507. In the first approach PCR primer pairs were designed to amplify a product that spanned the junction between the full-length insert and the inverted ORF25 terminator, see FIG. 1 for a depiction of the inverted ORF25 terminator. A forward primer was located at the end of the full-length PHI8999A insert and a series of reverse primers were located at 100 bp intervals in the inverted sequence. In this manner the length of the inverted fragment present in the maize event TC1507 could be determined within a 100 bp region based on the successful PCR reactions. This method

indicated the inverted fragment contained the majority of the ORF25 terminator but no Cry1F sequence. PCR fragments were isolated and sequenced from this region.

In the second approach PCR primers were designed to walk out into the flanking DNA sequence from the inverted ORF25 terminator region as determined in the PCR experiment described above. Genomic DNA isolated from two to three individual plants of event TC1507 and an unmodified control corn line was digested with various restriction enzymes and then ligated to adaptors specific for the restriction enzyme used for digestion (Universal Genome Walker™ Kit, Clontech Laboratories, Inc. and Devon *et al.* (1995) *Nucleic Acids Res.* 23:1644-1645). Primary PCR was carried out using an ORF25 terminator specific primer and a primer homologous to the adaptor sequence ligated onto the digested DNA. In order to increase the specificity of the reaction a nested secondary PCR was performed again with another ORF25 terminator specific primer and a secondary primer homologous to the adaptor sequence with the secondary primers being internal to the respective primers used in the primary PCR. Products produced by the nested PCR were analyzed by agarose gel electrophoresis and fragments unique to TC1507 DNA samples were isolated and sequenced. Fragments were amplified from both the ORF25 terminator contained within the full-length insert and from the targeted (inverted) ORF25 terminator on the 3' end of the full-length PHI8999A insert. Fragments from the full-length insert were of a predicted size based on the knowledge of the restriction enzyme sites located in the full-length insert. Fragments produced from the 3' inverted ORF25 terminator appeared as fragments of unexpected size. Sequence analysis of amplified fragments from the 3' inverted ORF25 terminator resulted in flanking DNA sequence of 1043 bp. Resultant sequence from the above series of genome walking experiments was used to design additional primers to walk further out from the insert into the bordering maize genome with a final 3' flanking sequence, of 2346 bp.

In order to describe the TC1507 3' flanking sequence, homology searching was done against the GenBank public databases using the Basic Local Alignment Search Tool (BLAST). The BLAST program performs sequence similarity searching and is particularly useful for identifying homologs to an unknown sequence. In addition to searching the public databases, alignments were performed using SeqMan 4.05™, Martinez and Needleman-Wunsch alignment algorithms (DNASTAR Inc.) to look for homology between the TC1507 3' flanking sequence and the PHI8999A transgenic insert. The results of these homology searches are presented in Table 1. The percent identity values indicate the percentage of identical matches

across the length of the sequences analyzed. The results of similarity searching for the 3' flanking sequence indicate high homology with three regions of maize chloroplast DNA, a 188 bp fragment of the *pat* gene, and 254 bp of DNA (Region 15, Table 1) with no significant homology. An additional 749 bp (Region 16) beyond Region 15 (see Table 1) was also sequenced. No similarity searching results are available for Region 16.

PCR analysis on control and TC1507 genomic DNA determined that the 254 bp sequence (Region 15, fragment of maize chloroplast "ORF241") is present in the maize genome. The DNA sequence of Region 15 in the 3' flanking region is not unique to the 3' flanking region of maize event TC1507 but is also present in the unmodified control corn line. The TC1507 3' flanking sequence is presented in Example 8 and diagrammed in FIG. 1.

Example 8. Sequence of the region 3' to the end of the full-length insert DNA in maize event TC1507. A description of each region is in Table 1.

**Region 10 (SEQ ID NO: 38) Fragment of ORF25 Terminator (complement)**

9016 CTCAC TCCGCTTGAT CTTGGCAAAG ATATTTGACG  
9051 CATTATTAG TATGTGTTAA TTTTCATTG CAGTGCAGTA TTTTCTATTC  
9101 GATCTTTATG TAATTCGTTA CAATTAATAA ATATTCAAAT CAGATTATTG  
9151 ACTGTCATTT GTATCAAATC GTGTTAATG GATATTTTTA TTATAATATT  
9201 GATGATATCT CAATCAAAAC GTAGATAATA ATAATATTTA TTAAATATTT  
9251 TTGCGTCGCA CAGTGAAAAT CTATATGAGA TTACAAAATA CCGACAACAT  
9301 TATTTAAGAA ACATAGACAT TAACCCTGAG ACTGTTGGAC ATCAACGGGT  
9351 AGATTCCTTC ATGCATAGCA CCTCATTCTT GGGGACAAAA GCACGGTTTG  
9401 GCCGTTCCAT TGCTGCACGA ACGAGCTTTG CTATATCCTC GGGTTGGATC  
9451 ATCTCATCAG GTCCAATCAA ATTTGTCCAA GAACTCATGT TAGTCGCAAC  
9501 GAAACCGGGG CATATGTCGG GTATCTCGAG CTCGCGAAAAG CTTGGCTGCA  
9551 GGTCGACGGA TCCTT

**Region 11 (SEQ ID NO: 39) Fragment of maize chloroplast *rps12* rRNA gene (complement)**

9566 CAACA AAAGGGTACC TGTACCCGAA ACCGACACAG  
9601 GTGGGTAGGT AGAGAATACC TAGGGGCGCG AGACAACTCT CTCTAAGGAA  
9651 CTCGGCAAAA TAGCCCCGTA ACTTCGGGAG AAGGGGTGCC CCC

**Nucleotides 9694-9695 (CG) connect Region 11 to Region 12.**

**Region 12 (SEQ ID NO: 40) Fragment of maize chloroplast genome**

9696 CTAAC  
9701 AATAAACGAA TACGGTTTAT GTATGGATTC CGGTAAAATA CCGGTACTCG  
9751 ATTCATAAG AGTCGAATAG GAAGTTAAGA TGAGGGTGGT ATCATCATAA  
9801 AAATGGAGTA GTATCCTAAA TTATACTAAT CCACGTATGA TATGTATGCC  
9851 TTTCCTTATC AACCGGAAGT AGTGCAAAAA AAATTCTATA CTGCACTGCT

9901 CTCTTTTAC TGAGAAATGC AAAAAAATAA AAGTGAAGTA AGGGTGCCCC  
 9951 ATAGATATTT GATCTTGCTT CCTGTCCCCC CCCCCCTTTT TTCATCAAAA  
 10001 ATTTCCATGA AAAAAGAAAA GATGAATTG TCCATTCATT GAACCCTAGT  
 10051 TCGGGACTGA CGGGGCTCGA ACCCGCAGCT TCCGCCT

5

**Region 13 (SEQ ID NO: 41) Fragment of *pat* gene (complement)**

10088 GTT CCTAGCCTTC  
 10101 CAGGGCCCAG CGTAAGCAAT ACCAGCCACA GCACCCTCAA  
 CCTCAGCAAC  
 10151 CAACCAAGGG TATCTATCTT GCAACCTCTC TAGATCATCA ATCCACTCTT  
 10201 GTGGTGTGTTG TGGCTCTGTC CTAAAGTTCA CTGTAGACGT CTCAATGTAA  
 10251 TGGTTAACGA TATCACAAC CGCGG

10

**Nucleotides 10276-10277 (AA) connect Region 13 to Region 14.**

15

**Region 14 (SEQ ID NO: 42) Fragment of maize chloroplast ORF241 (complement)**

10278 CAC AAGAACGAAA GCACCTTTTC  
 10301 ATTCTTTCAT ATACTAGGGG TTTTACTTG GAAAAGACAA TGTTCCATAC  
 10351 TAAAGGAT

20

**Region 15 (SEQ ID NO: 43) Maize genomic (no significant homology)**

10359 AG CTGCAGAAGC CGCCACCGTC TTGAGGACCT TCCGGGGAGC  
 10401 CAGACCGGTC GAACCGTGCC TCCACTTGCT AAGGAGAAAG  
 GGAAAATCAG  
 10451 GGCCAGGACA TACGAAGGAG GAGCCAGAAC GAAGATATCC  
 TAAGATACTT  
 10501 ACTCGCTCCG GGCCATGATC AATCATGCCT GTGGGGAGGT CTCTCGCACC  
 10551 TCGATCCATG AAGGTACCAC CGAGGTCTGC CCCGCCGCCG GCTTCGGTAC  
 10601 CGTCCTCGCC TT

25

30

**Region 16 (SEQ ID NO: 44) Maize genomic**

10613 GGGCGCCC GAGGCACCCG GGGGATGGAC TGCCCAGGCG  
 10651 CAGCCACGAC GACCCAAGGA TCACCCTCCT GCGCAGTCGG  
 CACGAGCAAT  
 10701 AGTTCTCGGG GAACAGGCAG CTTGGCCTGA CTCCCCGGGG TCACCTCAAC  
 10751 TACCTCGGCC GAGGGGTCAA GTACCCCTC AGTCCGCCCC CGCTCTTCGG  
 10801 ACCGGGACCC CGACGTCCCG GCCCCGGATA CCGACGGCAC CAGCCCGCTC  
 10851 GGGGGCTGGC TTGACGACCC CTGGCCAGC CTCAGATCTG GGCTGAGGCC  
 10901 GAGGCAGGCG GCCATGTCGT CGTCTTCATC ATCGTCTTCA TCATCGTCGT  
 10951 CGTCATCAGG CGTCTCCGGC GACGGCTCCC TTGGGAGCCC CTCCCTCTCC  
 11001 TGCCGACGAC GAAGCCTTTC CAAGGCATCC CGAGCCCACG TCCGCTCGTG  
 11051 GGCCCGAGCC TTCTTTGCGT CCTTCTTCTC CTTCCTCTTC TCCGCGGTGA  
 11101 CCCTCCGCGC AGCTCGGTCC ACCGCATCCT CCGGGACTGG TGGCAGGGAA  
 11151 GGCTTGTGAT GCCCTACCTC CTGGAGACAG ACGAAAAGTC TCAGCTATGA  
 11201 GAACCGAGGG CAATCTGACG CAAGAAGGAA GAAGGAGCGG  
 ATACTACCA

45

11251 GAGACACGCA CCCGCGATCG GGACGCATTA AGGGCTGGGA  
 AAAAGTGCCG  
 11301 GCCTCTAATT TCGCTACCGT GCCGTCCACC CACCTGTGGA GGTCATCGAT  
 11351 GGGAAGGGGA A

5

#### Example 9. Confirmation of the presence of Region 15 in the unmodified control corn line

PCR analysis was used to determine if the undescribed region of sequence on the end of the 3' flanking sequence (Region 15 in Table 1) is present in the unmodified control corn line used for transformation to produce maize event TC1507 and thus represents a border with corn genomic DNA. Successful PCR amplification of Region 15 in both maize line TC1507 and the unmodified Hi-II control corn line revealed that Region 15 was indeed present in corn genomic DNA. Five different PCR analyses were carried out on genomic DNA prepared from TC1507 and the unmodified Hi-II control corn line as outlined in Table 7 below using the primer sequences shown in Table 8. Three reactions were designed to amplify DNA within Region 15 of the 3' flanking region; Reaction L – producing a 175 bp amplicon, Reaction M – producing a 134 bp amplicon, and Reaction N – producing a 107 bp amplicon. The expected amplicons were present in both the unmodified control corn line and in maize line TC1507. Reactions J and K were designed as specific primer pairs for TC1507 that would span the end of the insert into the 3' flanking region. In Reaction J, the forward primer was located in the *pat* gene fragment on the 3' end of the full-length PHI8999A insert (Region 13 in Table 1) and the reverse primer was located in the undefined Region 15. In Reaction K the forward primer was located in the chloroplast hypothetical protein gene on the 3' end of the full-length insert (Region 14 in Table 1) and the reverse primer was located in the undefined Region 15. Both Reaction J and Reaction K produced an amplicon in maize line TC1507 and did not produce an amplicon in the unmodified control corn line. The results indicate that both Reactions J and K are specific for the TC1507 event.

The PCR results indicate that the undescribed sequence (Region 15) of the 3' flanking sequence of TC1507 is also present in genomic DNA isolated from the unmodified Hi-II control corn line. The DNA sequences amplified in Reactions L, M, and N are not unique to the 3' flanking region of TC1507 but are also present in the unmodified control corn line.

35

Table 7. PCR reactions for sequence 3' to the PHI8999A insert in maize event TC1507

| Reaction | Amplicon Size (bp) | Region in TC1507 3' flanking sequence              | Amplicon present in Control | Amplicon present in maize line TC1507 |
|----------|--------------------|----------------------------------------------------|-----------------------------|---------------------------------------|
| J        | 342                | Region 13 ( <i>pat</i> gene fragment) to Region 15 | No                          | Yes                                   |
| K        | 252                | Region 14 (chloroplast gene) to Region 15          | No                          | Yes                                   |
| L        | 175                | Region 15                                          | Yes                         | Yes                                   |
| M        | 134                | Region 15                                          | Yes                         | Yes                                   |
| N        | 107                | Region 15                                          | Yes                         | Yes                                   |

5 Table 8. PCR primers for sequence 3' to the PHI8999A insert in maize event TC1507

| Reaction | Amplicon Size (bp) | Primer Pair  | Primer Sequences 5' to 3'  |
|----------|--------------------|--------------|----------------------------|
| J        | 342                | SEQ ID NO:3  | TGTGGTGTGTTGTGGCTCTGTCCTAA |
|          |                    | SEQ ID NO:5  | GACCTCCCCACAGGCATGATTGATC  |
| K        | 252                | SEQ ID NO:4  | AGCACCTTTTCATTCTTTCATATAC  |
|          |                    | SEQ ID NO:5  | GACCTCCCCACAGGCATGATTGATC  |
| L        | 175                | SEQ ID NO:16 | AAGCCGCCACCGTCTTGAGGACCTT  |
|          |                    | SEQ ID NO:5  | GACCTCCCCACAGGCATGATTGATC  |
| M        | 134                | SEQ ID NO:17 | GTCGAACCGTGCCTCCACTTGTAA   |
|          |                    | SEQ ID NO:5  | GACCTCCCCACAGGCATGATTGATC  |
| N        | 107                | SEQ ID NO:18 | AGAAAGGGAAAATCAGGGCCAGGAC  |
|          |                    | SEQ ID NO:5  | GACCTCCCCACAGGCATGATTGATC  |

#### Example 10. PCR Primers

DNA event specific primer pairs were used to produce an amplicon diagnostic for TC1507. These event primer pairs include, but are not limited to, SEQ ID NO: 1 and SEQ ID NO: 2; SEQ ID NO: 2 and SEQ ID NO: 23; SEQ ID NO: 3 and SEQ ID NO: 5; and SEQ ID NO: 4 and SEQ ID NO: 5. In addition to these primer pairs, any primer pair derived from SEQ ID NO: 26 and SEQ ID NO: 27 that when used in a DNA amplification reaction produces a DNA amplicon diagnostic for TC1507 is an aspect of the present invention. The amplification conditions for this analysis are illustrated in Table 9, however, any modification of these methods that use DNA primers or complements thereof to produce an amplicon DNA molecule diagnostic for TC1507 is within the ordinary skill of the art. The preferred amplification conditions for reactions utilizing the PCR primers identified in SEQ ID NOS: 1, 2, and 23 are illustrated in Table 10. In addition, control primer pairs, which include SEQ ID NOS: 10 and 11; SEQ ID NOS: 10 and 12; SEQ ID NOS: 13 and 14; SEQ ID NOS: 14 and 15; SEQ ID NOS: 5 and 16; SEQ ID NOS: 5 and 17; and SEQ ID NOS: 5 and 18; for amplification of an endogenous corn gene are included as internal standards for the reaction conditions. Also included are primer pairs that will produce an amplicon in transgenic events containing a *pat* gene (SEQ ID NOS: 6 and 7; SEQ ID NOS: 8 and 9), and a primer pair that will produce an amplicon in transgenic events containing a *cry1F* gene (SEQ ID NOS: 19 and 20).

The analysis of plant tissue DNA extracts to test for the presence of the TC1507 event should include a positive tissue DNA extract control (a DNA sample known to contain the transgenic sequences). A successful amplification of the positive control demonstrates that the PCR was run under conditions which allow for the amplification of target sequences. A negative, or wild-type, DNA extract control in which the template DNA provided is either genomic DNA prepared from a non-transgenic plant, or is a non-TC1507 transgenic plant, should also be included. Additionally a negative control that contains no template corn DNA extract will be a useful gauge of the reagents and conditions used in the PCR protocol.

Additional DNA primer molecules of sufficient length can be selected from SEQ ID NO: 26 and SEQ ID NO: 27 by those skilled in the art of DNA amplification methods, and conditions optimized for the production of an amplicon that may differ from the methods shown in Table 9 or Table 10 but result in an amplicon diagnostic for event TC1507. The use of these DNA primer sequences with modifications to the methods shown in Table 9 and Table 10 are within the scope of the invention. The amplicon wherein at least one DNA primer

molecule of sufficient length derived from SEQ ID NO: 26 and SEQ ID NO: 27 that is diagnostic for event TC1507 is an aspect of the invention. The amplicon wherein at least one DNA primer of sufficient length derived from any of the genetic elements of PHI8999A that is diagnostic for event TC1507 is an aspect of the invention. The assay for the TC1507 amplicon can be performed by using a Stratagene Robocycler, MJ Engine, Perkin-Elmer 9700, or Eppendorf Mastercycler Gradient thermocycler, or by methods and apparatus known to those skilled in the art.

Table 9. PCR Conditions:

| Conditions: |                                    |
|-------------|------------------------------------|
| Kit used:   | Perkin-Elmer AmpliTAQ Gold kit     |
| Volume      | Component                          |
| 5 $\mu$ l   | template (10 ng/ $\mu$ l)          |
| 4 $\mu$ l   | 2 $\mu$ l each primer (10 $\mu$ M) |
| 2 $\mu$ l   | 10X PCR Gold Buffer                |
| 2 $\mu$ l   | 25 mM MgCl <sub>2</sub>            |
| 2 $\mu$ l   | 50X dNTP's (10 mM)                 |
| 0.1 $\mu$ l | Amplitaq Gold Polymerase           |
| 4.9 $\mu$ l | H <sub>2</sub> O                   |
| 20 $\mu$ l  | Total                              |

Cycling Parameters  
GeneAmp® PCR System 9700

9 min 92°C  
30 cycles:  
94°C 30 sec  
60°C 30 sec  
72°C 1 min  
7 min 72°C  
Hold 4°C

Table 10. PCR Conditions used with the Advantage<sup>®</sup>-GC 2 Polymerase Mix:

| Conditions:                          |                                             |
|--------------------------------------|---------------------------------------------|
| Kit used:                            | Advantage <sup>®</sup> -GC 2 Polymerase Mix |
| Volume                               | Component                                   |
| 5 µl                                 | template (10 ng/µl)                         |
| 5 µl                                 | 2.5 µl each primer (10 µM)                  |
| 10 µl                                | 5x GC2 Buffer                               |
| 10 µl                                | GC melt (1.0 M final conc.)                 |
| 1.5 µl                               | 50X dNTP's (10 mM)                          |
| 1.0 µl                               | Advantage GC2 Polymerase                    |
| 17.5 µl                              | H <sub>2</sub> O                            |
| 50 µl                                | Total                                       |
| Cycling Parameters                   |                                             |
| GeneAmp <sup>®</sup> PCR System 9700 |                                             |
| 5 min 94°C                           |                                             |
| <u>35 cycles:</u>                    |                                             |
| 94°C 1 min                           |                                             |
| 60°C 2 min                           |                                             |
| 72°C 3 min                           |                                             |
| 7 min 72°C                           |                                             |
| Hold 4°C                             |                                             |

Having illustrated and described the principles of the present invention, it should be  
 5 apparent to persons skilled in the art that the invention can be modified in arrangement and  
 detail without departing from such principles. We claim all modifications that are within the  
 spirit and scope of the appended claims.

All publications and published patent documents cited in this specification are  
 incorporated herein by reference to the same extent as if each individual publication or patent  
 10 application was specifically and individually indicated to be incorporated by reference.

## WHAT IS CLAIMED IS:

1. An isolated DNA molecule comprising a nucleotide sequence identified as SEQ ID NO:21.
2. An isolated DNA molecule comprising a nucleotide sequence identified as SEQ ID NO:22.
3. An isolated DNA molecule comprising a nucleotide sequence identified as SEQ ID NO:24.
4. An isolated DNA molecule comprising a nucleotide sequence identified as SEQ ID NO:26.
5. An isolated DNA molecule comprising a nucleotide sequence identified as SEQ ID NO:27.
6. A kit for identifying event TC1507 in a biological sample which detects a TC1507 specific region, said kit comprising at least a first primer, which recognizes a sequence within SEQ ID NO:21 or within SEQ ID NO:22.
7. The kit of claim 6, further comprising at least a second primer which recognizes a sequence within SEQ ID NO:25.
8. The kit of claim 6, further comprising at least a second primer which recognizes a sequence within SEQ ID NO:22.
9. The kit of claim 6, wherein the first primer recognizes a sequence within SEQ ID NO:21.
10. The kit of claim 6, wherein the first primer recognizes a sequence within SEQ ID NO:22.
11. The kit of claim 7, wherein said at least first and second primers comprise the sequence of SEQ ID NO:1 and SEQ ID NO:2, respectively.
12. The kit of claim 7, wherein said at least first and second primers comprise the sequence of SEQ ID NO:2 and SEQ ID NO:23, respectively.
13. The kit of claim 8, wherein said at least first and second primers comprise the sequence of SEQ ID NO:3 and SEQ ID NO:5, respectively.
14. The kit of claim 8, wherein said at least first and second primers comprise the sequence of SEQ ID NO:4 and SEQ ID NO:5, respectively.
15. A DNA detection kit specific for junction DNA of maize event TC1507 and its progeny comprising at least one DNA molecule of a sufficient length of contiguous DNA

- polynucleotides to function in a DNA detection method, that is homologous or complementary to SEQ ID NO: 26.
16. A DNA detection kit specific for junction DNA of maize event TC1507 and its progeny comprising at least one DNA molecule of a sufficient length of contiguous DNA polynucleotides to function in a DNA detection method, that is homologous or complementary to SEQ ID NO: 27.
  17. A kit for identifying event TC1507 in a biological sample, said kit comprising a specific probe comprising a sequence which hybridizes with SEQ ID NO:21 and SEQ ID NO:25, contiguous therewith.
  18. A kit for identifying event TC1507 in a biological sample, said kit comprising a specific probe comprising a sequence which hybridizes with SEQ ID NO:22 and SEQ ID NO:25, contiguous therewith.
  19. A DNA detection kit comprising: at least one DNA molecule of a sufficient length of contiguous nucleotides homologous or complementary to SEQ ID NO:26 or SEQ ID NO:27 that functions as a DNA primer or probe specific for maize event TC1507 and its progeny.
  20. A DNA construct comprising: a first and second expression cassette, wherein said first expression cassette in operable linkage comprises
    - (i) a maize ubiquitin promoter;
    - (ii) a 5' untranslated exon of a maize ubiquitin gene;
    - (iii) a maize ubiquitin intron;
    - (iv) a Cry1F encoding DNA molecule; and
    - (v) a 3' ORF25 transcriptional terminator; and
 said second expression cassette comprising in operable linkage
    - (a) a CaMV 35S promoter;
    - (b) a *pat* encoding DNA molecule; and
    - (c) a 3' transcriptional terminator from (CaMV) 35S.
  21. A plant comprising the DNA construct of claim 20.
  22. A plant of claim 21, wherein said plant is a corn plant.
  23. A method for identifying event TC1507 in a biological sample, comprising detecting a TC1507 specific region with a probe or first primer which specifically recognizes a sequence within SEQ ID NO:21 or SEQ ID NO:22.

24. The method of claim 23, further comprising amplifying a DNA fragment from a nucleic acid present in said biological sample using a polymerase chain reaction with at least two primers, wherein said first primer recognizes a sequence within SEQ ID NO:21 or SEQ ID NO:22, and a second primer recognizes a sequence within SEQ ID NO:22 or SEQ ID NO: 25.
25. The method of claim 24, wherein said first primer recognizes a sequence within SEQ ID NO:21 and said second primer recognizes a sequence within SEQ ID NO: 25.
26. The method of claim 23, wherein said first primer recognizes a sequence within SEQ ID NO:22 and a second primer recognizes a sequence within SEQ ID NO:22.
27. The method of claim 25, wherein said first and second primers comprise the sequence of SEQ ID NO:2 and SEQ ID NO:1 respectively.
28. The method of claim 25, wherein said first and second primers comprise the sequence of SEQ ID NO:2 and SEQ ID NO:23 respectively.
29. The method of claim 26, wherein said first and second primers comprise the sequence of SEQ ID NO:3 and SEQ ID NO:5 respectively.
30. The method of claim 26, wherein said first and second primers comprise the sequence of SEQ ID NO:4 and SEQ ID NO:5 respectively.
31. The method of claim 27, comprising amplifying a fragment of about 912 bp using a TC1507 PCR identification protocol.
32. The method of claim 28, comprising amplifying a fragment of about 844 bp using a TC1507 PCR identification protocol.
33. The method of claim 29, comprising amplifying a fragment of about 342 bp using a TC1507 PCR identification protocol.
34. The method of claim 30, comprising amplifying a fragment of about 252 bp using a TC1507 PCR identification protocol.
35. A method of detecting the presence of maize event TC1507 or progeny thereof in a biological sample, comprising:
  - (a) extracting a DNA sample from said biological sample;
  - (b) providing DNA primer molecules SEQ ID NO:1 and SEQ ID NO:2 ;
  - (c) providing DNA amplification reaction conditions;
  - (d) performing said DNA amplification reaction, thereby producing a DNA amplicon molecule; and

- (e) detecting said DNA amplicon molecule, wherein the detection of said DNA amplicon molecule in said DNA amplification reaction indicates the presence of maize event TC1507.
36. A method of detecting the presence of maize event TC1507 or progeny thereof in a biological sample, comprising:
- (a) extracting a DNA sample from said biological sample;
  - (b) providing DNA primer molecules SEQ ID NO:2 and SEQ ID NO:23 ;
  - (c) providing DNA amplification reaction conditions;
  - (d) performing said DNA amplification reaction, thereby producing a DNA amplicon molecule; and
  - (e) detecting said DNA amplicon molecule, wherein the detection of said DNA amplicon molecule in said DNA amplification reaction indicates the presence of maize event TC1507.
37. A method of detecting the presence of maize event TC1507 or progeny thereof in a biological sample, comprising:
- (a) extracting a DNA sample from said biological sample;
  - (b) providing DNA primer molecules SEQ ID NO:3 and SEQ ID NO:5 ;
  - (c) providing DNA amplification reaction conditions;
  - (d) performing said DNA amplification reaction, thereby producing a DNA amplicon molecule; and
  - (e) detecting said DNA amplicon molecule, wherein the detection of said DNA amplicon molecule in said DNA amplification reaction indicates the presence of maize event TC1507.
38. A method of detecting the presence of maize event TC1507 or progeny thereof in a biological sample, comprising:
- (a) extracting a DNA sample from said biological sample;
  - (b) providing DNA primer molecules SEQ ID NO:4 and SEQ ID NO:5 ;
  - (c) providing DNA amplification reaction conditions;
  - (d) performing said DNA amplification reaction, thereby producing a DNA amplicon molecule; and

(e) detecting said DNA amplicon molecule, wherein the detection of said DNA amplicon molecule in said DNA amplification reaction indicates the presence of maize event TC1507.

39. An isolated DNA molecule comprising the amplicon produced by the method of claim 35.
40. An isolated DNA molecule comprising the amplicon produced by the method of claim 36.
41. An isolated DNA molecule comprising the amplicon produced by the method of claim 37.
42. An isolated DNA molecule comprising the amplicon produced by the method of claim 38.
43. A method of detecting the presence of DNA corresponding to the TC1507 event in a sample, the method comprising:
  - a. contacting the sample comprising maize DNA with a polynucleotide probe that hybridizes under stringent hybridization conditions with DNA from maize event TC1507 and does not hybridize under said stringent hybridization conditions with a non-TC1507 maize plant DNA;
  - b. subjecting the sample and probe to stringent hybridization conditions; and
  - c. detecting hybridization of the probe to the DNA, wherein detection of hybridization indicates the presence of the TC1507 event.
44. An isolated DNA nucleotide primer sequence comprising a sequence selected from the group consisting of SEQ ID NO:1, 2, 3, 4, 5, 23, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56 and 57 or its complement.
45. An isolated DNA nucleotide primer sequence of claim 44 comprising a sequence selected from the group consisting of SEQ ID NO:1, 2, 3, 4, 5 and 23 or its complement.
46. A pair of DNA molecules comprising: a first DNA molecule and a second DNA molecule, wherein the DNA molecules are of a sufficient length of contiguous nucleotides of SEQ ID NO:26 or its complement to function as DNA primers or probes diagnostic for DNA extracted from a TC1507 corn plant or progeny thereof.

47. A pair of DNA molecules comprising: a first DNA molecule and a second DNA molecule, wherein the DNA molecules are of a sufficient length of contiguous nucleotides of SEQ ID NO:27 or its complement to function as DNA primers or probes diagnostic for DNA extracted from a TC1507 corn plant or progeny thereof.
48. An isolated DNA molecule comprising a junction sequence comprising a sequence selected from the group consisting of SEQ ID NO: 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56 and 57 and complements thereof.
49. A method for confirming seed purity, comprising detection of a TC1507 specific region with a specific primer or probe which specifically recognizes a sequence within SEQ ID NO:21 or SEQ ID NO:22, in a seed sample.
50. A method for screening seeds for the presence of event TC1507, comprising detection of a TC1507 specific region with a specific primer or probe which specifically recognizes a sequence within SEQ ID NO:21 or SEQ ID NO:22 in a sample of a seed lot.
51. An insect resistant corn plant, or parts thereof, wherein DNA having at least one nucleotide sequence selected from the group consisting of SEQ ID NO: 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56 and 57 and complements thereof forms part of the plant's genome.
52. A descent plant of the insect resistant corn plant of claim 51 wherein DNA having at least one nucleotide sequence selected from the group consisting of SEQ ID NO: 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56 and 57 and complements thereof forms part of the plant's genome.
53. Seed of a plant of claim 51 or 52.
54. A method of producing an insect resistant corn plant comprising breeding with a plant of claim 51 or 52, and selecting progeny by analyzing for at least one nucleotide sequence selected from the group consisting of SEQ ID NO: 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56 and 57 and complements thereof.
55. An isolated DNA sequence comprising a sequence selected from the group consisting of DNA having at least one nucleotide sequence selected from the group consisting of SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 23, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56 and 57 and complements thereof.

56. A pair of isolated DNA sequences, each comprising at least ten nucleotides and which when used together in a DNA amplification procedure will produce an amplicon diagnostic for event TC1507.
57. The pair of isolated DNA sequences of claim 56 wherein each is chosen from SEQ ID NO:26.
58. The pair of isolated DNA sequences of claim 56 wherein each is chosen from SEQ ID NO:27.
59. A method of detecting the presence of the TC1507 event insertion in corn tissue comprising:
  - (a) selecting a primer pair each comprising at least ten nucleotides from SEQ ID NO:26 or SEQ ID NO:27 wherein each member of the pair is on opposite sides of a sequence diagnostic for said TC1507 event insertion;
  - (b) contacting a sample of said corn tissue with said primer pair;
  - (c) performing DNA amplification and analyzing for an amplicons.
60. The method of claim 59 wherein said primer pair is selected from the group consisting of SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 23, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56 and 57 and complements thereof.
61. A method of detecting the presence of the TC1507 event insertion in corn tissue comprising:
  - (a) contacting a sample of said corn tissue with a polynucleotide probe that hybridizes under stringent hybridization conditions with one or more DNA sequence selected from the group consisting of SEQ ID NO: 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56 and 57 and complements thereof;
  - (b) subjecting said sample and probe to stringent hybridization conditions; and
  - (c) analyzing for hybridization of the probe.
62. A DNA detection kit comprising a polynucleotide probe that hybridizes under stringent hybridization conditions with one or more DNA sequences selected from the group consisting of SEQ ID NO: 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56 and 57 and complements thereof.

63. A DNA detection kit comprising a primer pair each comprising at least 10 nucleotides from SEQ ID NO:26 and SEQ ID NO:27, wherein each is on opposite sides of a sequence diagnostic for the TC1507 event insertion.
64. The DNA detection kit of claim 61 wherein said primer pair is selected from the group consisting of SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 23, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56 and 57 and complements thereof.
65. A kit for identifying event TC1507 in a biological sample which detects TC1507 specific region within SEQ ID NO: 24.
66. A method for identifying TC1507 in a biological sample which detects a TC1507 specific region within SEQ ID NO: 24.

# ABSTRACT OF THE DISCLOSURE

The invention provides DNA compositions that relate to transgenic insect resistant maize plants. Also provided are assays for detecting the presence of the maize TC1507 event  
5 based on the DNA sequence of the recombinant construct inserted into the maize genome and the DNA sequences flanking the insertion site. Kits and conditions useful in conducting the assays are provided.




FIG. 1

SEQUENCE LISTING

<110> Mary E. Hartnett Locke

<120> Corn Event TC1507 and Methods for  
Detection Thereof

<130> 1523P

<160> 57

<170> FastSEQ for Windows Version 4.0

<210> 1  
<211> 30  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Event specific primer sequence designed for TC1507

<400> 1  
gtagtactat agattatatt attcgtagag 30

<210> 2  
<211> 30  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Event specific primer sequence designed for TC1507

<400> 2  
gccatacaga actcaaaatc ttttcggag 30

<210> 3  
<211> 25  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Event specific primer designed for TC1507.

<400> 3  
tgtggtgttt gtggctctgt cctaa 25

<210> 4  
<211> 25  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Event specific primer for TC1507.

<400> 4  
agcacctttt cattctttca tatac 25

<210> 5  
 <211> 25  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Genomic DNA primer sequence  
  
 <400> 5  
 gacctcccca caggcatgat tgatc 25  
  
 <210> 6  
 <211> 25  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Primer in full length insert, 35S promoter to pat  
 gene  
  
 <400> 6  
 tagaggacct aacagaactc gccgt 25  
  
 <210> 7  
 <211> 24  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Primer in full length insert, 35S promoter to pat  
 gene  
  
 <400> 7  
 gagctggcaa ctcaaaatcc cttt 24  
  
 <210> 8  
 <211> 25  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Primer in full length insert, 35S promoter to pat  
 gene  
  
 <400> 8  
 aaaatcttcg tcaacatggt ggagc 25  
  
 <210> 9  
 <211> 25  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Primer in full length insert, 35S promoter to pat  
 gene

<400> 9  
taatctcaac tggctctctc tccgg 25

<210> 10  
<211> 22  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Primer - Zea mays genomic DNA

<400> 10  
cccctacccc accgacgttt at 22

<210> 11  
<211> 22  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Primer - Zea mays genomic DNA

<400> 11  
ttgattggca ggtccgtggg tc 22

<210> 12  
<211> 22  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Primer - Zea mays genomic DNA.

<400> 12  
cacaacggca cagaaacacg aa 22

<210> 13  
<211> 22  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Primer - Zea mays genomic DNA.

<400> 13  
gcgcacccac cggaacaaaa tg 22

<210> 14  
<211> 22  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Primer - Zea mays genomic DNA.

<400> 14  
tcctcgcat aaatgctcct gc 22

<210> 15  
 <211> 22  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Primer - Zea mays genomic DNA.  
  
 <400> 15  
 cctggcacgc attgacgcat gt 22  
  
 <210> 16  
 <211> 25  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Primer - Zea mays genomic DNA.  
  
 <400> 16  
 aagccgccac cgtcttgagg acctt 25  
  
 <210> 17  
 <211> 25  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Primer - Zea mays genomic DNA.  
  
 <400> 17  
 gtcgaaccgt gcctccactt gctaa 25  
  
 <210> 18  
 <211> 25  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Primer - Zea mays genomic DNA.  
  
 <400> 18  
 agaaagggaa aatcagggcc aggac 25  
  
 <210> 19  
 <211> 22  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> CrylF sequence primer  
  
 <400> 19  
 ggctcggact cgacctttct at 22  
  
 <210> 20

<211> 22  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> CrylF sequence primer

<400> 20  
gcagttcttg aagaatgagt ga

22

<210> 21  
<211> 2829  
<212> DNA  
<213> 5' flanking sequence of event TC1507

<400> 21  
actagtttcc tagcccgct cgtgcccta cccaccgac gtttatggaa ggtgccattc 60  
cacggttctt cgtggccgcc cctaaggatg taaatggctg gtaaaatccg gtaaatcttc 120  
ggtaccggtt accagatttt tccagccgtt ttcggattta tcgggatata cagaaaacga 180  
gacggaaacg gaatagggtt tttttcgaaa acggtacggt aaacggtgag acaaaacttac 240  
cgtccggttt cgtattttctc gggaaactct ggtatatcc cgtattttgtc ccgtattttc 300  
ccgacccacg gacctgccaa tcaaccatca gccagtcagc ccatccccac agctatggcc 360  
catggggcca tgttgccac atgccacgc aacgcaaggc agtaaggctg gcagcctggc 420  
acgcattgac gcatgtggac acacacagcc gccgcctgtt cgtgtttctg tgccgttgtg 480  
cgagactgtg actgcgagtg gcggagtcgg cgaacggcga ggctctccg gactctggac 540  
tgccgctgtg gacagcgagc ctgtgacggc gactcggcga agcccaagc taccagccc 600  
ccaagtcccc atccatctct gcttctctgg tcatctcctt ccctggctg atctgcaggc 660  
gccagaccgg ccgaagcatc acgaaacgca ctaagacctc gaaggagtca aaccactcct 720  
ccgaggcctc gggggctaca cccggcgggt gcgctcgcgc gcacccaccg gaacaaaatg 780  
taaccgagaa aggtcggtcc ccttgcaaaa aaagtgcgac aaaagcctcc aagcgagtat 840  
taacactcac tttgaggtc gggggctact gtcggggacc ataattaggg gtaccccaa 900  
gactccta atctcagctgtt aaccccatc agcacaagc tgcaaaaggc tgatgggtgc 960  
gattaagtca aggtcgggtc cactcaaggg acacgatctc gcctcgccc agccagcct 1020  
cgggcaaggg cggccgaccc cgaggattca cgtctcgccc gagggccccc tcaagcgacg 1080  
ggcacacctt cggctcggcc gaggccatt ctccgcccag aagcaacctt ggccagatcg 1140  
ccacaccgac cgaccgtatc gcaggagcat ttaatgcgag gatcgccctga caccttatcc 1200  
tgacgcgcgc tcttcagtcg acagagcga agtgaccgca atcacttcgc cgctccactg 1260  
accgacctga caagaagaca gcgcgcctg cgtcgctccg actgctgtgc cactcgacag 1320  
agtgaaggctg acagacgcca agtcggcct cgggcgccaat aggaagctcc gcctcgccc 1380  
accctagggc tcggactcgg cctcggctcc ggaagacgac gaactacgct tcgcccagcc 1440  
ccagggcttg gactcagcct cggctccgga agacgacgaa ttccgcctcg cccgacccca 1500  
gggctcggac tcggcctcgg ctccagaaga cgacgaactc cgctcgccc gacccaggg 1560  
ctcggactca gcctcggtc cggaagacga cgaactccgc ctgcccagc cccagggtc 1620  
ggactcagcc tcggcctcag acgatgtct cgcctcggc cgacccggg ctcgactcg 1680  
acctttctat cggacctgtg cagatcctgt ctctgctcga ggaggctttg gcaatcctca 1740  
ctatgtactc ggtcttaggg gactggcctt tcaacaaact ggtacgaatc acaccgcac 1800  
attcaggaac tccgggacca ttgactctct agatgagata ccacctcaag acaacagcgg 1860  
cgcaccttg aatgactact cccatgtgct gaatcatgtt acctttgtgc gctggccagg 1920  
tgagatctca ggttcgact catggagagc accaatgttc tcttgagcgc atcgtagcgc 1980  
tacccccaca aacaccattg atccagagag aatcactcat tcttcaagaa ctgcatatct 2040  
tgccgagatc tctatcccta aaggtaactg acaatagat tattggagtc gatacacaac 2100  
tcacaaaaaa tacaagaagt cgactaggtg gattgggtccg agtgaagaga aaaaaaagcc 2160  
atacagaact caaaatcttt tccggagata ttcatcttcc tgaagaggcg gataagatat 2220  
taggtggcag tttgatacca ccagaaaag aaaaaaaaga ttctaaggaa tcaaaaaaaa 2280  
ggaaaaattg ggtttatgtt caacggaaaa aatttctcaa aagcaaggaa aagtattgtg 2340  
gctatctatc tatccgtgca gctgataggg ccgcggtttg tgatatcggt aaccattaca 2400  
ttgagacgtc tacagtgaac tttaggacag agccacaaac accacaagag tggattgatg 2460

```

atctagagag gttgcaagat agataccctt ggttggttgc tgagggttgag ggtggttggtg 2520
ctgggtattgc ttacgctggg ccctggaagg ctaggaaccc tcaacctcag caaccaacca 2580
atggtatcta tcttgcaacc tctctagatc atcaatccac tcttggtgtg tttgtggctc 2640
tgtcctaaag ttactgttag acgtctcaat gtaatggtta acgatatcac aaaccgagag 2700
aagagggatc tcgaagcttc ggccggggcc catcgatata cgcgggcatg cctgcagtgc 2760
agcgtgaccc ggtcgtgccc ctctctagag ataatgagca ttgcatgtct aagttataaa 2820
aaattacca 2829

```

<210> 22

<211> 2346

<212> DNA

<213> 3' flanking sequence of event TC1507

<400> 22

```

ctcactccg ttagctcttg caaagatatt tgacgcattt attagtatgt gttaattttc 60
atttgcagt cagtattttc tattcgatct ttatgtaatt cgttacaatt aataaatatt 120
caaatcagat tattgactgt cattttgatc aaatcgtgtt taatggatat ttttattata 180
atattgatga tatctcaatc aaaacgtaga taataataat atttatttaa tttttttgcg 240
tcgcacagtg aaaatctata tgagattaca aaataccgac aacattattt aagaaacata 300
gacattaacc ctgagactgt tggacatcaa cgggtagatt ccttcattgca tagcacctca 360
ttcttgggga caaaagcacg gtttgccgtg tccattgctg cacgaacgag ctttgctata 420
tcctcggggt ggatcatctc atcaggtoaa atcaaatttg tccaagaact catgttagtc 480
gcaacgaaac cggggcatat gtcgggtatc tcgagctcgc gaaagcttgg ctgcaggctc 540
acggatcctt caacaaaagg gtacctgtac ccgaaaccga cacagggtgg taggtagaga 600
atacctaggg gcgcgagaca actctctcta aggaactcgg caaaatagcc ccgtaacttc 660
gggagaaggg gtgccccccg ctaacaataa acgaatacgg ttatgtatg gattccggta 720
aaataccggt actcgatttc ataagagtcg aatagggaagt taagatgagg gtggtatcat 780
cataaaaaat gagtagtata ctaaatataa ctaatccacg tatgatattg atgcctttcc 840
ttatcaaccg gaagttagtc aaaaaaaatt ctatactgca ctgctctctt tttactgaga 900
aatgcaaaaa aataaaagtg aagtaagggt gccccataga tatttgatct tgcctcctgt 960
cccccccccc cttttttcat caaaaatttc catgaaaaaa gaaaagatga atttgtccat 1020
tcattgaacc ctagtctggg actgacgggg ctcgaaccgc cagcttccgc ctgttccatg 1080
ccttccaggg cccagcgtaa gcaataccag ccacagcacc ctcaacctca gcaaccaacc 1140
aagggtatct atcttgcaac ctctctagat catcaatcca ctcttggtg gtttggtggc 1200
ctgtcctaaa gtctactgta gacgtctcaa tgtaatggtt aacgatata caaacgcg 1260
aacacaagaa cgaaaagcacc ttttcattct tcatataact aggggttttt acttgaaaa 1320
gacaatgttc catactaaag gatagctgca gaagccgcca cgtcttgag gaccttccg 1380
ggagccagac cggtcgaacc gtgcctccac ttgctaagga gaaagggaat atcagggcc 1440
ggacatacga aggaggagcc agaacgaaga tatectaaga tacttactcg ctccgggcc 1500
tgatcaatca tgctgtggg gaggtctctc gcacctcgat ccataaagg accaccgag 1560
tctgccccgc cgccgcttc ggtaccgtcc tcgccttggg cgcgcgagc acccggggg 1620
tggactgcc aggcgcagcc acgacgacc aaggatcacc ctctgcca gtcggcacg 1680
gcaatagttc tcggggaaca ggcagcttgg cctgactccc cggggtcacc tcaactacct 1740
cggccgaggg gtcaagtacc cctcagtc gcccccgtc ttcggaccg gaccccgac 1800
tcccggcccc ggataccgac ggcaccagcc cgctcgggg ctggttgac gaccctggc 1860
ccagcctcag atctgggctg aggcgaggg aggcggccat gtcgtcgtct tcatcatcgt 1920
cttcatcatc gtcgtcgtca tcaggcgtct ccggcgacgg ctcccttgg agccctccc 1980
tctcctgcc acgacgaagc ctttccaagg catcccagc ccacgtccgc tcgtgggcc 2040
gagccttctt tgcgtccttc ttctccttc tcttctccgc ggtgacctc cgcgcagctc 2100
ggtccaccgc atcctccggg actggtggca gggaaggctt gtgatgccct acctcctgga 2160
gacagacgaa aagtctcagc tatgagaacc gagggcaatc tgacgcaaga aggaagaagg 2220
agcggatact caccagagac acgcaccgc gatcgggacg cattaaggc tgggaaaaag 2280
tgccggcctc taatttcgct accgtgccgt ccaccacct gtggaggtca tcgatgggaa 2340
ggggaa 2346

```

<210> 23

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Event specific primer sequence designed for TC1507

<400> 23

cttcaaacaa gtgtgacaaa

20

<210> 24

<211> 11361

<212> DNA

<213> Artificial Sequence

<220>

<223> The sequence represents the transgenic insert in  
maize line TC1507 as well as the sequence flanking  
the insertion sites.

<400> 24

```

actagtttcc tagcccgct cgtgccccta cccaccgac gtttatggaa ggtgccattc 60
cacggttctt cgtggccgcc cctaaggatg taaatggctg gtaaatccg gtaaatcttc 120
ggtaccgttt accagatttt tccagccgtt ttcggattta tcgggatata cagaaaacga 180
gacggaaacg gaatagggtt tttttcgaaa acgggtacgg aaacgggtgag acaaaacttac 240
cgtccgtttt cgtattttctc gggaaactct ggtatatctc cgtatttgte ccgtattttc 300
ccgacccacg gacctgccaa tcaaccatca gccagtcagc ccatccccac agctatggcc 360
catggggcca tgttgccac atgcccacgc aacgcaaggc agtaaggctg gcagcctggc 420
acgcattgac gcatgtggac acacacagcc gccgcctgtt cgtgtttctg tgcggtgtg 480
cgagactgtg actgcgagtg gccgagtcgg cgaacggcga gccgtctccg gactctggac 540
tgcggctgtg gacagcgacg ctgtgacggc gactcggcga agccccaagc taccagccc 600
ccaagtcccc atccatctct gtttctctgg tcatctcctt cccctggctg atctgcaggc 660
gccagaccgg ccgaagcatc acgaaacgca ctaagacctc gaaggagtca aaccactcct 720
ccgaggcctc gggggctaca cccggcgggt gcgctcgcgc gcacccaccg gaacaaaaatg 780
taaccgagaa aggtcggctc ctttgcaaaa aaagtgcgac aaaagcctcc aagcgagtat 840
taacactcac tttgaggtc gggggctact gtccgggacc ataattagggt gtaccccaaa 900
gactccta atctcagctgtt aaccccatc agcacaaagc tgcaaggcc tgatgggtgc 960
gattaagtca aggtcgggtc cactcaaggg acacgatctc gctcgcgcc agccagcct 1020
cgggcaaggg cggccgaccc cgaggattca cgtctcgcgc gagggccccc tcaagcgacg 1080
ggcacacctt cggctcgccc gaggccatt ctccgcccag aagcaacctt gccagatcg 1140
ccacaccgac cgaccgtatc gcaggagcat ttaatgcgag gatcgectga caccttatcc 1200
tgacgcgcgc tcttcagtcg acagagccga agtgaccgca atcacttcgc cgtccactg 1260
accgacctga caagaagaca gcgcgcctg cgtcgcctcg actgctgtgc cactcgacag 1320
agtgaggctg acagcagcca agtcggcct cgggcgccat aggaagctcc gcctcgccc 1380
accctagggc tcggactcgg cctcggctcc ggaagacgac gaactacgt tcgcccagcc 1440
ccagggtctg gactcagcct cggctccgga agacgacgaa ttccgectcg cccgacccca 1500
gggctcggac tcggcctcgg ctccagaaga cgacgaaact cgcctcgccc gacccaggg 1560
ctcggactca gcctcggctc cggaagacga cgaactccgc ctgcgccgac cccagggtc 1620
ggactcagcc tcggcctcag acgatggtct ccgcctcgcc cgaccgggg ctggactcg 1680
acctttctat cggaccttgt cagatcctgt cttcgtccga ggaggcttg gcaatcctca 1740
ctatgtactc ggtcttaggg gactggcctt tcaacaaact ggtacgaatc acaccgcac 1800
attcaggaac tccgggacca ttgactctct agatgagata ccacctcaag acaacagcgg 1860
cgcaccttg aatgactact cccatgtgct gaatcatgtt acctttgtgc gctggccagg 1920
tgagatctca ggttcgact catggagagc accaatgttc tcttgacgc atcgtagcgc 1980
tacccccaca aacaccattg atccagagag aatcactcat tcttcaagaa ctgcatact 2040
tgccgagatc ctatcccta aaggtacttg acaatagtat tattggagtc gatacacaac 2100
tcacaaaaaa tacaagaagt cgactagggt gattggtccg agtgaagaga aaaaaagcc 2160
atacagaact caaaatcttt tccggagata ttcatcttcc tgaagaggcg gataagatat 2220
taggtggcag tttgatacca ccagaaagag aaaaaaaga ttctaaggaa tcaaaaaaaa 2280
ggaaaaattg ggtttatgtt caacggaaaa aatttctcaa aagcaaggaa aagtattgtg 2340

```

```

gctatattatc tatccgtgca gctgatattgg ccgcggtttg tgatatcggt aaccattaca 2400
ttgagacgtc tacagtgaac tttaggacag agccacaaac accacaagag tggattgatg 2460
atctagagag gttgcaagat agataccctt ggttggttgc tgagggtttag ggtgttggg 2520
ctggtattgc ttacgctggg ccctggaagg cttagaacc tcaacctcag caaccaacca 2580
atggtatcta tcttgcaacc tctctagatc atcaatccac tcttggtgtg tttgtggctc 2640
tgtcctaaag ttactgttag acgtctcaat gtaatgggta acgatatcac aaaccgagag 2700
aagagggatc tcgaagcttc ggccggggcc catcgatc cgcgggcatg cctgcagtgc 2760
agcgtgaccc ggtcgtgccc ctctctagag ataatgagca ttgcatgtct aagtataaaa 2820
aaattaccac aactggaaga gcggttacc ggaccgaagc ttcggccggg gcccatcgat 2880
atccgcgggc atgcctgcag tgcagcgtga ccggtcgtg cccctctcta gagataatga 2940
gcattgcatg tctaagttat aaaaaattac cacatatttt ttttgtcaca cttgtttgaa 3000
gtgcagttta tctatcttta tacatatatt taaactttac tctacgaata atataatcta 3060
tagtactaca ataatatcag tgttttagag aatcatataa atgaacagtt agacatggtc 3120
taaaagcaaa ttgagtattt tgacaacagg actctacagt tttatctttt tagtgtgcat 3180
gtgttctcct ttttttttgc acctatataa tacttcatcc attttattag 3240
tacatccatt tagggttttag ggttaaatggt tttttagac taattttttt agtacatcta 3300
ttttattcta ttttagcctc taaattaaga aaactaaaac tctatttttag tttttttatt 3360
taataattta gatataaaat agaataaaat aaagtgacta aaaattaaac aaataccctt 3420
taagaaatta aaaaaactaa ggaacatttt ttcttgtttc gagtagataa tgccagcctg 3480
ttaacgcgcg tcgacgagtc taacggacac caaccagcga accagcagcg tcgcgtcggg 3540
ccaagcgaag cagacggcac ggcatctctg tcgctgcctc tggaccctc tcgagagttc 3600
cgctccaccg ttggaacttc tccgctgtcg gcatccagaa attgcgtggc ggagcggcag 3660
acgtgagccg gcacggcagg cggcctcctc ctectctcac ggcacggcag ctacggggga 3720
ttcctttccc accgctcctt cgttttccct tccctgcccc cgtaataaaa tagacacccc 3780
ctccacaccc tctttcccca acctcgtgtt gttcggagcg cacacacaca caaccagatc 3840
tcccccaaat ccaccgctcg gcacctccgc tccaaggtag gccgctcgtc ctcccccccc 3900
ccccctctct acctctctta gatcggcgtt ccggtccatg gttagggccc ggtagttcta 3960
cttctgttca tgtttgtgtt agatccgtgt ttgtgttaga tccgtgctgc tagcgttcgt 4020
acacggatgc gacctgtacg tcagacacgt tctgattgct aacttgccag tgtttctctt 4080
tggggaatcc tgggatggct ctacccgttc cgcagacggg atcgatttca tgattttttt 4140
tgtttcggtg cataggggtt ggtttgccct tttcctttat ttcaatataat gccgtgcaact 4200
tgtttgtcgg gtcacttttt catgcttttt tttgtccttg ttgtgatgat gtggtctggt 4260
tgggcggtcg ttctagatcg gagtagaatt ctgtttcaaa ctacctggtg gatttattaa 4320
ttttggatct gtatgtgtgt gccatacata ttcatagtta cgaattgaag atgatggatg 4380
gaaatatcga tctaggatag gtatacatgt tgatgcgggt ttactgatg catatacaga 4440
gatgcttttt gttcgcttgg ttgtgatgat gtggtgtggt tgggcggtcg ttcatctggt 4500
ctagatcgga gtgaataact gtttcaaact acctggtgta tttattaatt ttggaactgt 4560
atgtgtgtgt catagctctt catagttacg agtttaagat ggatggaaat atcgatctag 4620
gataggatata catgttgatg tgggttttac tgatgcata acatgatggc atatgcagca 4680
tctattcata tgctctaacc ttgagtacct atctattata ataaacaagt atgttttata 4740
attattttga tcttgatata cttggatgat ggcataatga gcagctatat gtggattttt 4800
ttagccctgc cttcatacgc tatttatttg cttggtactg tttcttttgc cgatgctcac 4860
cctgttgttt ggtgttactt ctgcaggctc actctagagg atccaacaat ggagaacaac 4920
atacagaatc agtgcgtccc ctacaactgc ctcaacaatc ctgaagtaga gattctcaac 4980
gaagagaggt cgactggcag attgcccgtta gacatctccc tgtcccttac acgtttcctg 5040
ttgtctgagt ttgttccagg tgtgggagtt gcgtttggcc tcttcgacct catctggggc 5100
ttcatcactc catctgattg gagcctcttt cttctccaga ttgaacagtt gattgaacaa 5160
aggattgaga ccttggaag gaatcgggcc atcactaccc ttcgtggctt agcagacagc 5220
tatgagatct acattgaagc actaagagag tgggaagcca atcctaaca tgcccaactg 5280
agagaagatg tgcgtatagc ctttgctaac acagatgatg ctttgatcac agccatcaac 5340
aacttcaccc ttaccagctt cgagatccct cttctctcgg tctatgttca agctgctaac 5400
ctgcacttgt cactactgcg cgacgctgtg tcgtttgggc aaggttgggg actggacata 5460
gctactgtca acaatcacta caacagactc atcaatctga ttcactgata cacgaaacat 5520
tgtttggata cctacaatca gggattggag aacctgagag gtactaacac tcgccaatgg 5580
gccaggttca atcagttcag gagagacctt acacttactg tgtagacat agttgctctc 5640
tttccgaact acgagtgtcg tacctatccg attcaaacgt catcccaact tacaagggag 5700
atctacacca gttcagtcac tgaagactct ccagtttctg cgaacatacc caatggtttc 5760
aacagggtcg agtttgagat cagaccaccc catctcatgg acttcatgaa ctctttgttt 5820

```

```

gtgactgcag agactgttag atcccaaact gtgtggggag gacacttagt tagctcacgc 5880
aacacggctg gcaatcgtat caactttcct agttacgggg tcttcaatcc cgggggcgcc 5940
atctggattg cagatgaaga tccacgtcct ttctatcgga ccttgtcaga tcctgtcttc 6000
gtccgaggag gctttggcaa tctcactat gtactcggtc ttaggggagt ggcctttcaa 6060
caaaactggt cgaatcacac ccgcacattc aggaactccg ggaccattga ctctctagat 6120
gagataccac ctcaagacaa cagcggcgca ccttggaatg actactccca tgtgctgaat 6180
catgttacct ttgtgcgtg gccaggtgag atctcaggtt ccgactcatg gagagcacca 6240
atgttctctt ggacgcatcg tagcgctacc cccacaaaca ccattgatcc agagagaatc 6300
actcagattc ccttggtgaa ggcacacaca cttcagtcag gaactacagt tgtaagaggg 6360
ccgggggttca cgggaggaga cattcttcga cgcactagtg gaggaccatt cgcgtacacc 6420
attgtcaaca tcaatgggca acttccccaa aggtatcgtg ccaggatacg ctatgcctct 6480
actaccaatc taagaatcta cgttacggtt gcaggtgaac ggatctttgc tggtcagttc 6540
aacaagacaa tggataccgg tgatccactt acattccaat ctttctccta cgccactatc 6600
aacaccgctg tcacctttcc aatgagccag agcagtttca cagtaggtgc tgataccttc 6660
agttcaggca acgaagtgtg cattgacagg tttagattga ttccagttac tgccacactc 6720
gagtaaggat ccgtcgacct gcagccaagc tttcgcgagc tcgagatccc cgacatatgc 6780
cccggtttgc ttgcgactaa catgagttct tggacaaatt tgattggacc tgatgagatg 6840
atccaaccgg aggatatagc aaagctcgtt cgtgcagcaa tggaacggcc aaaccgtgct 6900
tttgtcccca agaattgaggt gctatgcatg aaggaatcta cccgttgatg tccaacagtc 6960
tcagggttaa tgtctatgta tcttaataaa tgttgtcggt attttgtaat ctcatataga 7020
ttttcactgt gcgacgcaaa aatattaaat aaatattatt attatctacg ttttgattga 7080
gatatcatca atattataat aaaaatatcc attaaacacg atttgataca aatgacagtc 7140
aataatctga tttgaatatt tattaattgt aacgaattac ataaagatcg aatagaaaat 7200
actgcactgc aaatgaaaat taacacatac taataaatgc gtcaaatac tttgccaaga 7260
tcaagcggag tgagggcctc atatccggtc tcagttacaa gcacgggtatc cccgaagcgc 7320
gctccaccaa tgccctcgac atagatgccg ggtcgcagcg tgaggacatt gcctaccttg 7380
agcatggtct cagcgcgggc tttaaagctc atcccacccc aatctgaata tcctatcccc 7440
cgcccgatcc ggtgtaagaa cgggtctgtc catccacctc tgttgggaat tccggtccgg 7500
gtcacctttg tccaccaaga tggaaactgc gccgcggacc gaattcccat ggagtcгааг 7560
attcaaatag aggacctaac agaactcgcc gtaaagactg gcgaacagtt catacagagt 7620
ctcttacgac tcaatgacaa gaagaaaatc ttcgtcaaca tgggtggagca cgacacgctt 7680
gtctactcca aaaatatcaa agatacagtc tcagaagacc aaagggcaat tgagactttt 7740
caacaaaggg taatatccgg aaacctctc ggattccatt gccagctat ctgtcacttt 7800
attgtgaaga tagtgaaaaa ggaaggtggc tctacaaat gccatcattg cgataaagga 7860
aaggccatcg ttgaagatgc ctctgccgac agtgggtcca aagatggacc cccacccacg 7920
aggagcatcg tggaaaaaga agacgttcca accacgtctt caaagcaagt ggattgatgt 7980
gatattctca ctgacgtaag ggatgacgca caatcccact atccttcgca agacccttcc 8040
tctatataag gaagttcatt tcatltggag aggacagggg acccggggat ccacctatgc 8100
tccggagagg agaccagttg agattaggcc agctacagca gctgatatgg ccgcgggttg 8160
tgatatcgtt aaccattaca ttgagacgtc tacagtgaac tttaggacag agccacaaac 8220
accacaagag tggattgatg atctagagag gttgcaagat agataccctt ggttggttgc 8280
tgaggttgag ggtgttggtg ctggtattgc ttacgctggg cctggaagg ctaggaacgc 8340
ttacgatttg acagttgaga gtactgttta cgtgtcacat aggcatacaa ggttgggcct 8400
aggatccaca ttgtacacac atttgcttaa gtctatggag gcgcaagggt ttaagtctgt 8460
ggttgctgtt ataggccttc caaacgatcc atctgttagg ttgcatgagg ctttgggata 8520
cacagcccgg ggtacattgc gcgcagctgg atacaagcat ggtggatggc atgatgttgg 8580
tttttggcaa agggattttg agttgccagc tectccaagg ccagttaggc cagttacca 8640
gatctgagtc gacctgcagg catgcccgtc gaaatcacca gtctctctc acaaatctat 8700
ctctctctat aataatgtgt gagtagttcc cagataaggg aattagggtt ctatagggt 8760
ttcgctcatg tgttgagcat ataagaaacc cttagtatgt atttgattt gtaaaatact 8820
tctatcaata aaatttctaa ttctaaaac caaaatccag tggcgagctc gaattcgagc 8880
tcgagcccgg gtggatcctc ctgcagaagc ctgcagaagc agtagtcttc gcccagaatg 8940
gttgaagaca cgttcatgtc ttcacgttaa gaagacactc agtagtcttc gcccagaatg 9000
gcctaactca aggcctcac tccgcttgat cttggcaaaг atatttgacg catttattag 9060
tatgtgttaa ttttcatttg cagtgcatga actgtcattt gtatcaaac gtgtttaatg 9120
caattaataa atattcaaat gatgatattc caatcaaaac gtagataata ataatttta 9180
gatattttta ttataatatt ctatatgaga ttacaaaata ccgacaacat 9240
tttaatattt ttgcgtcgca cagtgaaaat ctatatgaga ttacaaaata ccgacaacat 9300

```

```

tatttaagaa acatagacat taaccctgag actgttggac atcaacgggt agattccttc 9360
atgcatagca cctcattctt ggggacaaaa gcacgggtttg gccgttccat tgctgcacga 9420
acgagctttg ctatatactc ggggttggatc atctcatcag gtccaatcaa atttgtccaa 9480
gaactcatgt tagtcgcaac gaaaccgggg catatgtcgg gtatctcgag ctgcgcaaaa 9540
cttggtgca ggtcgacgga tccttcaaca aaagggtacc tgtaccgaa accgacacag 9600
gtgggttagt agagaatacc tagggcgcg agacaactct ctctaaggaa ctgcgcaaaa 9660
tagccccgta acttcgggag aaggggtgcc ccccgctaac aataaacgaa tacggtttat 9720
gtatggattc cggtaaaata ccggtactcg atttcataag agtcgaatag gaagttaaga 9780
tgagggtggt atcatcataa aaatggagta gtatcctaaa ttatactaata ccacgtatga 9840
tatgtatgcc ttctcttata aaccggaagt agtgcaaaaa aaattctata ctgcactgct 9900
ctctttttac tgagaaatgc aaaaaataa aagtgaagta agggtgcccc atagatatatt 9960
gatcttgcc cctgtcccc ccccccttt ttcatcaaaa atttccatga aaaaagaaaa 10020
gatgaatttg tccattcatt gaaccctagt tcgggactga cggggctcga acccgagct 10080
tccgctgtt cctagccttc cagggcccg cgtaagcaat accagccaca gcaccctcaa 10140
cctcagcaac caaccaaggg tatctatctt gcaacctctc tagatcatca atccactctt 10200
gtggtgtttg tggctctgtc cttaaagttca ctgtagacgt ctcaatgtaa tggttaacga 10260
tatcacaac cgcggaacac aagaacgaaa gcacctttt attctttcat atactagggg 10320
tttttacttg gaaaagacaa tgttccatac taaaggatag ctgcagaagc cgccaccgct 10380
ttgaggacct tccggggagc cagaccggtc gaaccgtgcc tccacttgct aaggagaaaag 10440
ggaaaatcag ggccaggaca tacgaaggag gagccagaac gaagatatcc taagataact 10500
actcgctccg ggccatgatc aatcatgcct gtggggaggt ctctcgacc tcgatccatg 10560
aaggtaccac cgaggtctgc cccgcccgcg gcttcggtac cgtcctcgcc ttgggcgccc 10620
gaggcaccgc ggggatggac tgcccaggcg cagccacgac gaccaagga tcaccctcct 10680
gcgcagtcg cagcagcaat agttctcggg gaacaggcag cttggcctga ctccccggg 10740
tcacctcaac tacctcgcc gaggggtcaa gtacccctc agtcggccc cgctcttcg 10800
accgggaccc cgacgtccc gccccggata ccgacggcac cagcccgtc gggggctggc 10860
ttgacgaccc ctggcccagc ctcatgctgt ggtcagggcc gaggcaggcg gccatgtcgt 10920
cgtcttcac atcgtcttca tcctcgtcgt cgtcatcagg cgtctcggc gacggtccc 10980
ttgggagccc ctccctctcc tgccgacgac gaagccttcc caaggcatcc cgagcccacg 11040
tccgctcgtg ggcccagacc ttctttgcgt ccttcttctc ctctctctc tccgcggtga 11100
ccctccgcgc agctcgggtc accgcaccc cccgggactg tggcaggga ggcttgtgat 11160
gcctacctc ctggagacag acgaaaagtc tcagctatga gaaccgaggg caatctgacg 11220
caagaaggaa taaggagcgg atactacca gagacacga cccgcgatcg ggacgcatta 11280
agggtcggga aaaagtgcg gcctctaatt tcgctaccgt gccgtccacc cacctgtgga 11340
ggtcatcgat gggaaggga a 11361

```

<210> 25

<211> 6235

<212> DNA

<213> Artificial Sequence

<220>

<223> The sequence represents the DNA molecule used to transform maize line TC1507. This sequence represents insert PHI8999A.

<400> 25

```

aaacgctctt caactggaag agcggttacc cggaccgaag cttcggccgg ggcccatcga 60
tatccgcggg catgcctgca gtgcagcgtg acccggtcgt gcccctctct agagataatg 120
agcattgcat gtctaagtta taaaaaatta ccacatattt ttttgtcac acttggttga 180
agtgcagttt atctatcttt atacatatat ttaaacttta ctctacgaat aatataatct 240
atagtactac aataatatca gtgttttaga gaatcatata aatgaacagt tagacatggt 300
ctaaggaca attgagtatt ttgacaacag gactctacag ttttatcttt ttagtgtgca 360
tgtgttctcc tttttttttg caaatagctt cacctatata ataacttcac cattttatta 420
gtacatccat ttagggttta gggtaaatgg tttttataga ctaatttttt tagtacatct 480
attttattct attttagcct ctaaattaag aaaactaaaa ctctatttta gtttttttat 540

```

|             |            |             |             |            |             |      |
|-------------|------------|-------------|-------------|------------|-------------|------|
| ttaataaattt | agatataaaa | tagaataaaa  | taaagtgact  | aaaaattaaa | caaataccct  | 600  |
| ttaagaaatt  | aaaaaaacta | aggaaacatt  | tttcttgttt  | cgagtagata | atgccagcct  | 660  |
| gttaaaccgc  | gtcgacgagt | ctaaccgaca  | ccaaccagcg  | aaccagcagc | gtcgcgtcgg  | 720  |
| gccaaagcaa  | gcagacggca | cggcatctct  | gtcgcgtcct  | ctggaccctt | ctcgagagtt  | 780  |
| ccgctccacc  | gttggacttg | ctccgctgtc  | ggcatccaga  | aattgcgtgg | cggagcggca  | 840  |
| gacgtgagcc  | ggcacggcag | gcggcctcct  | cctcctctca  | cggcacggca | gctacggggg  | 900  |
| attcctttcc  | caccgctcct | tcgctttccc  | ttcctcgccc  | gccgtaataa | atagacaccc  | 960  |
| cctccacacc  | ctctttcccc | aacctcgtgt  | tggtcggagc  | gcacacacac | acaaccagat  | 1020 |
| ctcccccaaa  | tccaccctgc | ggcacctcgg  | cttcaaggta  | cgccgctcgt | cctccccccc  | 1080 |
| ccccctctc   | taccttctct | agatcggcgt  | tccggtccat  | ggttagggcc | cggtagttct  | 1140 |
| acttctgttc  | atgtttgtgt | tagatccgtg  | tttgtgttag  | atccgtgctg | ctagcgttcg  | 1200 |
| tacacggatg  | cgacctgtac | gtcagacacg  | ttctgattgc  | taacttgcca | gtgtttctct  | 1260 |
| ttgggggaatc | ctgggatggc | tctagccgtt  | ccgcagacgg  | gatcgatttc | atgatttttt  | 1320 |
| ttgtttcggt  | gcatagggtt | tggtttgccc  | ttttccttta  | tttcaatata | tgccgtgcac  | 1380 |
| ttgtttgtcg  | ggtcatcttt | tcattgctttt | ttttgtcttg  | gttgtgatga | tgtgggtctg  | 1440 |
| ttgggcgggtc | gttctagatc | ggagtagaat  | tctgtttcaa  | actacctggt | ggatttatta  | 1500 |
| attttggtatc | tgatgtgtgt | tgccatacat  | attcatagtt  | acgaattgaa | gatgatggat  | 1560 |
| ggaatatatcg | atctaggata | ggtatacatg  | ttgatcgagg  | ttttactgat | gcataatacag | 1620 |
| agatgctttt  | tggtcgcttg | gttgtgatga  | tgtgggtgtg  | ttgggcggtc | gttcattcgt  | 1680 |
| tctagatcgg  | agtagaatac | tggttcaaac  | tacctggtgt  | atttattaat | tttggaactg  | 1740 |
| tatgtgtgtg  | tcatacatct | tcatagtta   | gagtttaaga  | tggttgga   | tatcgatcta  | 1800 |
| ggaataggtat | acatgttgat | gtgggtttta  | ctgatgcata  | tacatgatgg | catatgcagc  | 1860 |
| atctattcat  | atgctctaac | cttgagtacc  | tatctattat  | aataaacaag | tatgttttat  | 1920 |
| aattattttg  | atcttgatat | acttgatga   | tgccatatgc  | agcagctata | tgtggatttt  | 1980 |
| tttagccctg  | ccttcatacg | ctattttatt  | gcttggtact  | gtttcttttg | tcgatgctca  | 2040 |
| ccctgttgtt  | tggtgttact | tctgcaggtc  | gactctagag  | gatccaaaca | tggaagaaca  | 2100 |
| catacagaat  | cagtgcgtcc | cctacaactg  | cctcaacaat  | cctgaagtag | agattctcaa  | 2160 |
| cgaagagagg  | tcgactggca | gattgccggt  | agacatctcc  | ctgtccctta | cacgtttcct  | 2220 |
| gttgtctgag  | tttgtccag  | gtgtgggagt  | tgcggttggc  | ctcttcgacc | tcactcgggg  | 2280 |
| cttcactact  | ccatctgatt | ggagcctctt  | tcttctccag  | attgaacagt | tgattgaaca  | 2340 |
| aaggattgag  | accttgga   | ggaatcgggc  | catcactacc  | cttcgtggct | tagcagacag  | 2400 |
| ctatgagatc  | tacattgaag | cactaagaga  | gtgggaagcc  | aatcctaaca | atgcccaact  | 2460 |
| gagagaagat  | gtgctgatac | gctttgctaa  | cacagatgat  | gctttgatca | cagccatcaa  | 2520 |
| caacttcacc  | cttaccagct | tcgagatccc  | tcttctctcg  | gtctatgttc | aagctgctaa  | 2580 |
| cctgcacttg  | tcactactgc | gcgacgtgtg  | gtcggttggg  | caaggttggg | gactggacat  | 2640 |
| agctactgtc  | aacaatcact | acaacagact  | catcaatctg  | attcatcgat | acacgaaaca  | 2700 |
| ttgtttggat  | acctacaatc | agggattgga  | gaacctgaga  | ggtactaaca | ctcgccaatg  | 2760 |
| ggccagggttc | aatcagttca | ggagagacct  | tacacttact  | gtgttagaca | tagttgctct  | 2820 |
| ctttccgaac  | tacgatgttc | gtacctatcc  | gattcaaacg  | tcacccaac  | ttacaaggga  | 2880 |
| gatctacacc  | agttcagtc  | ttgaagactc  | tccagtttct  | gcgaacatac | ccaatgggtt  | 2940 |
| caacagggct  | gagtttgag  | tcagaccacc  | ccatctcatg  | gacttcatga | actctttgtt  | 3000 |
| tgtgactgca  | gagactgtta | gatcccaaac  | tgtgtgggga  | ggacacttag | ttagctcacg  | 3060 |
| caacacggct  | ggcaatcgta | tcaactttcc  | tagttacggg  | gtcttcaatc | ccgggggcgc  | 3120 |
| catctggatt  | gcagatgaag | atccacgtcc  | tttctatcgg  | accttgtcag | atcctgtctt  | 3180 |
| cgtccgagga  | ggctttggca | atcctcacta  | tgtactcggg  | cttaggggag | tgccctttca  | 3240 |
| acaaactggt  | acgaatcaca | ccgcacattc  | caggaaactcc | gggaccattg | actctctaga  | 3300 |
| tgagatacca  | cctcaagaca | acagcggcgc  | accttgggat  | gactactccc | atgtgctgaa  | 3360 |
| tcattgttacc | tttgtgcgct | ggccagggtga | gatctcaggt  | tccgactcat | ggagagcacc  | 3420 |
| aatgtttctct | tggaagcgtc | gtagcgtctc  | ccccacaaac  | accattgatc | cagagagaat  | 3480 |
| cactcagatt  | cccttggtga | aggcacacac  | acttcagtca  | ggaactacag | ttgtaagagg  | 3540 |
| gccgggggttc | acgggaggag | acattctctc  | acgcactagt  | ggaggaccat | tcgcgtacac  | 3600 |
| cattgtcaac  | atcaatgggc | aacttcccca  | aaggtatcgt  | gccaggatac | gctatgcctc  | 3660 |
| tactaccaat  | ctaagaatct | acgttacggg  | tgacgggtga  | cggatctttg | ctgggtcagtt | 3720 |
| caacaagaca  | atggataccg | gtgatccact  | tacattccaa  | tctttctcct | acgccactat  | 3780 |
| caacaccgcg  | ttcacccttc | caatgagcca  | gagcagtttc  | acagtaggtg | ctgatacctt  | 3840 |
| cagttcaggc  | aacgaagtgt | acattgacag  | gtttgagttg  | attccagtta | ctgccacact  | 3900 |
| cgaagtaagga | tccgtcgacc | tgacgccaag  | cttttcgcga  | gctcgagatc | ccgacatat   | 3960 |
| gccccgggtt  | cgttgcgact | aacatgagtt  | cttggaacaa  | tttgattgga | cctgatgaga  | 4020 |

```

tgatccaacc cgaggatata gcaaagctcg ttcgtgcagc aatggaacgg ccaaaccgtg 4080
cttttgtccc caagaatgag gtgctatgca tgaaggaatc taccggttga tgtccaacag 4140
tctcagggtt aatgtctatg tatcttaaat aatgtttgtc gtatttttga atctcatata 4200
gatttttact gtgcgacgca aaaatattaa ataaatatta ttattatcta cgttttgatt 4260
gagatatcat caatattata ataaaaatat ccattaaaca cgatttgata caaatgacag 4320
tcaataatct gatttgaata tttattaatt gtaacgaatt acataaagat cgaatagaaa 4380
atactgcact gcaaatgaaa attaacacat actaataaat gcgtcaaata tctttgccaa 4440
gatcaagcgg agtgagggcc tcatatccgg tctcagttac aagcacggta tccccgaagc 4500
gcgctccacc aatgccctcg acatagatgc cgggctcgac gctgaggaca ttgcctacct 4560
tgagcatggt ctcagcgccg gctttaagct caatcccatc ccaatctgaa tatcctatcc 4620
cgcgcccagt ccggtgtaag aacgggtctg tccatccacc tctgttgagg attccggtcc 4680
gggtcacctt tgtccaccaa gatggaactg cggccgcgga ccgaattccc atggagtcaa 4740
agattcaaat agaggacctt acagaactcg ccgtaaagac tggcgaacag ttcatacaga 4800
gtctcttacg actcaatgac aagaagaaaa tcttcgtcaa catggtggag cagcacgc 4860
ttgtctactc caaaaatata aaagatacag tctcagaaga ccaaaggcca attgagactt 4920
ttcaacaaag ggtaatatcc ggaaacctcc tcggattcca ttgcccagct atctgtcact 4980
ttattgtgaa gatagtggaa aagggaaggtg gctcctacaa atgccatcat tgcgataaag 5040
gaaaggccat cgttgaagat gcctctgcgg acagtggtcc caaagatgga cccccaccca 5100
cgaggagcat cgtggaaaaa gaagacgttc caaccacgtc ttcaaagcaa gtggattgat 5160
gtgatattct cactgacgta agggatgacg cacaatccca ctatccttcg caagaccctt 5220
cctctatata aggaagttca tttcatttgg agaggacagg gtaccggggg atccaccatg 5280
tctcgggaga ggagaccagt tgagattagg ccagctacag cagctgatat ggccgcggtt 5340
tgtgatattc ttaaccatta cattgagacg tctacagtga actttaggac agagccacaa 5400
acaccacaag agtggtattga tgatctagag aggttgcaag atagataccc ttggttggtt 5460
gctgaggttg aggggtgtgt ggctggtatt gcttacgctg ggccctggaa ggctaggaac 5520
gcttacgatt ggacagttga gagtactggt tacgtgtcac ataggcatca aaggttgggc 5580
ctaggatcca cattgtacac acatttgctt aagtctatgg aggcgcaagg ttttaagtct 5640
gtggttgctg ttataggcct tccaaacgat ccatctgtta ggttgcata ggctttggga 5700
tacacagccc ggggtacatt gcgcgcagct ggatacaagc atggtggatg gcatgatgtt 5760
ggtttttggc aaagggattt tgagttgcca gctcctccaa ggccagttag gccagttacc 5820
cagatctgag tcgacctgca ggcagccgcg tgaaatcacc agtctctctc taaaaatcta 5880
tctctctcta taataatgtg tgagtgttc ccagataagg gaattagggg tcttataggg 5940
tttcgctcat gtgttgagca tataagaaac ccttagtatg tatttgattt tgtaaaatac 6000
ttctatcaat aaaatttcta attcctaaaa ccaaaatcca gtggcgagct cgaattcgag 6060
ctcgagcccg ggtggatcct ctgaggtcga cctgcagaag ctcggtccg gcgcgcctct 6120
agttagaac acgttcatgt cttcatcgta agaagacact cagtagtctt cggccagaat 6180
ggcctaactc aaggccatcg tggcctcttg ctcttcagga tgaagagcta tgtttt 6235

```

<210> 26

<211> 3830

<212> DNA

<213> Artificial Sequence

<220>

<223> Sequence that represents part of the PHI8999A  
insert as well as flanking sequence 5' to the  
insert.

<400> 26

```

actagtttcc tagcccgct cgtgcccta cccaccgac gtttatggaa ggtgccattc 60
cacggttctt cgtggccgcc cctaaggatg taaatggctg gtaaaatccg gtaaaatttc 120
ggtaccgttt accagatttt tcagccggtt ttcggattta tcgggatata cagaaaacga 180
gacggaaacg gaataggttt tttttcgaaa acggtagcgt aaacggtgag acaaaacttac 240
cgtccgtttt cgtatttctc gggaaactca ggtatatccc cgtattttgc ccgtattttc 300
ccgaccacag gacctgccaa tcaaccatca gccagtcagc ccatccccc agctatggcc 360
catggggcca tgttgccac atgccacgc aacgcaaggc agtaaggctg gcagcctggc 420

```

```

acgcattgac gcatgtggac acacacagcc gccgcctgtt cgtgtttctg tgccgttgtg 480
cgagactgtg actgcgagtg gccgagtcgg cgaacggcga ggctgtctcg gactctggac 540
tgcggtgtg gacagcgacg ctgtgacggc gactcggcga agccccaagc taccaagccc 600
ccaagtcccc atccatctct gcttctcttg tcactctcct cccttggtcg atctgcaggc 660
gccagaccgg ccgaagcadc acgaaacgca ctaagacctc gaaggagtca aaccactcct 720
ccgagggcctc gggggctaca cccggcgggt gcgtctcgcg gcacccaccg gaacaaaatg 780
taaccgagaa aggtcggtcc ccttgcaaaa aaagtgcgac aaaagcctcc aagcgagtat 840
taacactcac tttgaggctc gggggctact gtccggggacc ataattaggg gtaccccca 900
gactccta at ctcagctggg aacccccatc agcacaaaagc tgcaaaaggc tgatgggtgc 960
gattaagtca aggtcctggg cactcaaggg acacgatctc gcctcgccc agcccagcct 1020
cgggcaaggg cggccgaccc cgaggattca cgtctcgccc gagggccccc tcaagcgacg 1080
ggcacacctt cggctcgccc gagggccatt ctctcgccgag aagcaacctt ggccagatcg 1140
ccacaccgac cgaccgtatc gcaggagcat ttaatgcgag gatcgccctga caccttatcc 1200
tgacgcgcgc tcttcagtcg acagagccga agtgaccgca atcacttcgc cgtccactg 1260
accgacctga caagaagaca gcgcgcctg cgtcgctccg actgctgtgc cactcgacag 1320
agtgaggtcg acagcagcca agtcggcctc cgggcgcctc aggaagctcc gcctcgccc 1380
acctagggc tcgactcgg cctcggtctc ggaagacgac gaactacgct tcgcccagc 1440
ccagggtctg gactcagcct cggctccgga agacgacgaa tccgcctcg cccgacccca 1500
gggctcggac tcggcctcgg ctccagaaga cgacgaactc cgctcgccc gacccaggg 1560
ctcggactca cctcggctc cggaagacga cgaactccgc ctgcccagc cccagggtc 1620
ggactcagcc tcggcctcag acgatggtct ccgctcgcc cgacccggg ctcgactcg 1680
acctttctat cggacctgtg cagatcctgt ctctgctcga ggaggtttg gcaatcctca 1740
ctatgtactc ggtcttaggg gagtggcctt tcaacaaact ggtacgaatc acaccgcac 1800
attcaggaac tccgggacca ttgactctct agatgagata ccacctcaag acaacagcg 1860
cgcaccttg aatgactact cccatgtgct gaatcatgtt acctttgtgc gctggccagg 1920
tgagatctca ggttcogact catggagagc accaatgttc tcttgagcg atctgagcg 1980
tacccccaca aacaccattg atccagagag aatcactcat tctcaagaa ctgcatact 2040
tgccgagatc ctcatcccta aaggtacttg acaatagtat tattggagt gatacaca 2100
tcacaaaaaa tacaagaagt cgactagggt gattggtccg agtgaagaga aaaaaagcc 2160
atacagaact caaaatcttt tccggagata ttcatcttcc tgaagaggcg gataagatat 2220
taggtggcag tttgatacca ccagaaagag aaaaaaaga ttctaaggaa tcaaaaaaa 2280
ggaaaaattg ggtttatgtt caacggaaaa aatttctcaa aagcaaggaa aagtattgtg 2340
gctatttact tatccgtgca gctgatagc cgcgggtttg tgatatcgtt aaccattaca 2400
ttgagacgct tacagtgaac tttaggacag agccacaaac accacaagag tggattgatg 2460
atctagagag gttgcaagat agataccctt ggttggttgc tgaggttgag ggtgttgtg 2520
ctggtattgc ttacgtggg ccctggaagg ctaggaaccc tcaacctcag caaccaacca 2580
atggtatcta tcttgaacc tctctagatc atcaatccac tcttgtggtg tttgtggctc 2640
tgtcttaag ttcatgtag acgtctcaat gtaatggtta acgatatcac aaaccgagag 2700
aagagggatc tcgaagcttc ggccggggcc catcgatcgc cgcgggcatg cctgcagtgc 2760
agcgtgaccc ggtcgtgccc ctctctagag ataatgagca ttgcatgtct aagttataaa 2820
aaattaccac aactggaaga gccgttaccg ggaccgaagc ttcggccggg gcccatcgat 2880
atccgcgggc atgcctgcag tgcagcgtga cccggtcgtg cccctctcta gagataatga 2940
gcattgcatg tctaagttat aaaaaattac cacatatatt tttgtcaca cttgtttgaa 3000
gtgcagttta tctatcttta tacatatatt taaactttac tctacgaata atataatcta 3060
tagtactaca ataatacag tgttttagag aatcatataa atgaacagtt agacatggct 3120
taaaggacaa ttgagtattt tgacaacagg actctacagt tttatctttt tagtgtgcat 3180
gtgttctcct ttttttttgc aaatagcttc acctatataa tacttcatcc attttattag 3240
tacatccatt tagggtttag ggttaatggt ttttatagac taattttttt agtacatcta 3300
ttttattcta ttttagcctc taaattaaga aaactaaaac tctatttttag tttttttatt 3360
taataattta gatataaaat agaataaaat aaagtgacta aaaattaaac aaataccctt 3420
taagaattaa aaaaaactaa ggaaacattt ttcttgttcc gagtagataa tgccagcctg 3480
ttaaacgcgc tcgacgagtc taacggacac caaccagcga accagcagcg tcgcgtcggg 3540
ccaagcgaag cagacggcac ggcatctctg tcgctgcctc tggacccctc tcgagagttc 3600
cgctccaccg ttggacttgc tccgctgtcg gcacccagaa attcggtggc ggagcggcag 3660
acgtgagccg gcacggcagg cggcctcctc ctctctcac ggcacggcag ctacggggga 3720
ttcctttccc accgctcctt cgctttccct tcctcgccc cgtataaaa tagacacccc 3780
ctccacaccc tctttcccca acctcgtgtt gttcggagcg cacacacaca 3830

```

<210> 27  
 <211> 3347  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Sequence that represents part of the PHI8999A  
 insert as well as flanking sequence 3' to the  
 insert.

```
<400> 27
cccactatcc ttcgcaagac ccttcctcta tataaggaag ttcatttcat ttggagagga 60
cagggtagcc ggggatccac catgtctccg gagaggagac cagttgagat taggccagct 120
acagcagctg atatggccgc ggtttgtgat atcgtaaac attacattga gacgtctaca 180
gtgaacttta ggacagagcc acaaacacca caagagtggg ttgatgatct agagaggttg 240
caagatagat acccttggtt ggttgctgag gttgaggggt ttgtggctgg tattgcttac 300
gctgggccct ggaaggctag gaacgcttac gattggacag ttgagagtac tgtttacgtg 360
tcacatagga atcaaaaggt gggcctagga tccacattgt acacacattt gcttaagtct 420
atggaggcgc aagggtttta gtctgtggtt gctgttatag gccttccaaa cgatccatct 480
gttaggttgc atgaggcttt gggatacaca gcccggggta cattgcgcgc agctggatac 540
aagcatgggt gatggcatga tgttggtttt tggcaaaggg attttgagtt gccagctcct 600
ccaaggccag ttaggccagt taccagatc tgagtcgacc tgcaggcatg cccgctgaaa 660
tcaccagtct ctctctacaa atctatctct ctctataata atgtgtgagt agttcccaga 720
taagggaatt agggttctta tagggtttcg ctcatgtgtt gagcatataa gaaaccctta 780
gtatgttatt gtatttgtaa aatacttcta tcaataaaat ttctaattcc taaaaccaa 840
atccagtggc gagctcgaat tcgagctcga gcccggttgg atcctctaga gtcgacctgc 900
agaagcttgc gtccgcgcgc cctctagttg aagacacgtt catgtcttca tcgtaagaag 960
acactcagta gtcttcggcc agaatggcct aactcaaggc cctcactccg cttgatcttg 1020
gcaaagatat ttgacgcatt tattagtatg tgttaatttt catttgcatg gcagtatttt 1080
ctattcagtc tttatgtaat tcgttacaat taataaatat tcaaatcaga ttattgactg 1140
tcatttgtat caaatcgtgt ttaatggata tttttattat aatattgatg atatctcaat 1200
caaaacgtga ataataataa tatttattta atatttttgc gtcgcacagt gaaaaatctat 1260
atgagattac aaaataaccga caacattatt taagaaacat agacattaac cctgagactg 1320
ttggacatca acgggtagat tccttcatgc atagcacctc attcttgggg acaaaagcac 1380
ggtttggccg ttccattgct gcacgaacga gctttgctat atcctcgggt tggatcatct 1440
catcaggtcc aatcaaattt gtccaagAAC tcattgttagt cgcaacgaaa ccggggcata 1500
tgtcgggtat ctgcagctcg cgaaagcttg gctgcaggtc gacggatcct tcaacaaaag 1560
ggtagctgta cccgaaaccg acacaggtgg gtaggtagag aatacctagg ggcgcgagac 1620
aactctctct aaggaaactcg gcaaaatagc cccgtaactt cgggagaagg ggtgcccccc 1680
gctaacaata aacgaatacg gtttatgtat ggattccggt aaaatacccg tactcgattt 1740
cataagagtc gaataggaag ttaagatgag ggtgggtatca tcataaaaaat ggagtagtat 1800
cctaaattat actaatccac gtatgatatg tatgcctttc cttatcaacc ggaagtagtg 1860
caaaaaaaat tctatactgc actgctctct ttttactgag aaatgcaaaa aaataaaagt 1920
gaagtaaggg tgccccatag atatttgatc ttgcctctcg tcccccccc ccttttttca 1980
tcaaaaattt ccatgaaaaa agaaaagatg aatttgtcca ttcatgaaac cctagtctcg 2040
gactgacggg gctcgaaccc gcagcttccg cctgttctta gccttccagg gccagcgta 2100
agcaatacca gccacagcac cctcaacctc agcaaccaac caagggtatc tatcttgcaa 2160
cctctctaga tcatcaatcc actcttgttg tgtttgtggc tctgtcctaa agttcactgt 2220
agacgtctca atgtaatggt taacgatatc acaaacgcgc gaacacaaga acgaaagcac 2280
cttttcattc tttcatatac taggggtttt tacttggaag agacaatgtt ccatactaaa 2340
ggatagctgc agaagccgcc accgtcttga ggaccttccg gggagccaga ccggctgaac 2400
cgtgcctcca ctgtctaag agaaaggga aatcagggcc aggacatacg aaggaggagc 2460
cagaacgaag atatcctaag atacttactc gctccgggcc atgatcaatc atgcctgtgg 2520
ggaggtctct cgcacctcga tccatgaagg taccaccgag gtctgccccg ccgccggctt 2580
cggtagcgtc ctgccttggt gcgccgaggt caccgggggg atggactgcc caggcgagc 2640
cacgacgacc caaggatcac cctcctgcgc agtcggcacg agcaatagtt ctcggggaac 2700
aggcagcttg cctgactcc ccggggtcac ctcaactacc tcggccgagg ggtcaagtac 2760
cccctcagtc cgccccgcct cttcggaccg ggaccccgac gtcccggccc cggataccga 2820
```

```

cggcaccagc ccgctcgggg gctggcttga cgacccctgg cccagcctca gatctgggct 2880
gaggccgagg caggcgccca tgtcgtcgtc ttcctcatcg tcttcatcat cgtcgtcgtc 2940
atcaggcgctc tccggcgacg gctcccttgg gagccctcc ctctcctgcc gacgacgaag 3000
cctttccaag gcatcccgag cccacgtccg ctcggtggcc cgagccttct ttgctcctt 3060
cttctccttc ctcttctccg cgggtgacct ccgcgcagct cggtcaccg catcctccg 3120
gactgggtggc aggggaaggct tgtgatgcc tacctcctgg agacagacga aaagtctcag 3180
ctatgagaac cgagggcaat ctgacgcaag aaggaagaag gagcggatac tcaccagaga 3240
cacgcaccgc cgatcgggac gcattaaggc ctgggaaaaa gtgccggcct ctaatttcgc 3300
taccgtgccg tccacccacc tgtggaggtc atcgatggga aggggaa 3347

```

<210> 28  
 <211> 669  
 <212> DNA  
 <213> Zea mays

```

<400> 28
actagtttcc tagcccgct cgtgcccta cccaccgac gtttatggaa ggtgccattc 60
cacggttctt cgtggccgcc cctaaggatg taaatggctg gtaaaatccg gtaaatcttc 120
ggtagcgttt accagatttt tccagccgtt ttcggattta tcgggatata cagaaaacga 180
gacggaaacg gaataggttt tttttcgaaa acggtacggt aaacgggtgag acaaaacttac 240
cgtccgtttt cgtattttctc gggaaactct ggtatatctc cgtattttgc ccgtattttc 300
ccgaccacgc gacctgccaa tcaaccatca gccagtcagc ccacccccac agctatggcc 360
catggggcca tgttggccac atgccacgc aacgcaagge agtaaggctg gcagcctggc 420
acgcattgac gcattgtggc acacacagcc gccgcctgtt cgtgtttctg tgccgttgtg 480
cgagactgtg actgcgagtg gcggagtcgg cgaacggcga ggctgtctcg gactctggac 540
tgccgtgtgt gacagcgacg ctgtgacggc gactcggcga agccccaagc taccaagccc 600
ccaagtcccc atccatctct gcttctcttg tcatctcctt ccctgggtcg atctgcaggc 660
gccagaccg
669

```

<210> 29  
 <211> 200  
 <212> DNA  
 <213> Zea mays

```

<400> 29
gccgaagcat cagcaaacgc actaagacct cgaaggagtc aaaccactcc tccgaggcct 60
cgggggctac accggcgagg tgcgtcgcg cgcaccacc ggaacaaaat gtaaccgaga 120
aaggtcggtc cccttgcaaa aaaagtgcga caaagcctc caagcgagta ttaacactca 180
ctttgaggct cgggggctac
200

```

<210> 30  
 <211> 812  
 <212> DNA  
 <213> Zea mays

<220>  
 <223> Fragment of maize Huck-1 retrotransposon

```

<400> 30
tgtcggggac cataattagg ggtaccccca agactcctaa tctcagctgg taacccccat 60
cagcaciaaag ctgcaaaggc ctgatgggtg cgattaagtc aaggtcgggt ccactcaagg 120
gacacgatct cgcctcgccc gageccagcc tcgggcaagg gcggccgacc ccgaggattc 180
acgtctcgcc cgaggggccc ctcaagcgac ggccacacct tcggctcgcc cgaggcccat 240
tcttcgccga gaagcaacct tggccagatc gccacaccga ccgaccgtat cgcaggagca 300
tttaatgcga ggatcgccctg acaccttatc ctgacgcgcg ctcttcagtc gacagagccg 360
aagtgaaccg aatcacttcg ccgctccact gaccgacctg acaagaagac agcgcgcgct 420
gcgtcgctcc gactgctgtg ccactcgaca gagtgggctg gacagcagcc aagtccggcc 480
tcgggcgcca taggaagctc cgcctcgccc gacctaggg ctcggaactc gcctcggtc 540

```

```

cggaagacga cgaactacgc ttgccccgac cccagggcct ggactcagcc tcggctccgg 600
aagacgacga attccgcctc gcccgacccc agggctcgga ctggcctcg gctccagaag 660
acgacgaact ccgcctcgcc cgaccccagg gctcggactc agcctcggt ccggaagacg 720
acgaactccg cctcgcccgga ccccagggct cggactcagc ctggcctca gacgatggtc 780
tccgcctcgc ccgacccggg gctcggactc ga 812

```

<210> 31  
 <211> 335  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Represents part of PHI8999A insert sequence -  
 fragment of cry1F gene

```

<400> 31
cctttctatc ggaccttgtc agatcctgtc ttcgtccgag gaggcttttg caatcctcac 60
tatgtactcg gtcttagggg agtggccttt caacaaactg gtacgaatca caccgcaca 120
ttcaggaact ccgggaccat tgactctcta gatgagatac cacctcaaga caacagcggc 180
gcaccttgga atgactactc ccatgtgctg aatcatgtta cctttgtgcg ctggccaggt 240
gagatctcag gttccgactc atggagagca ccaatgttct cttggacgca tcgtagcgtc 300
acccccacaa acaccattga tccagagaga atcac 335

```

<210> 32  
 <211> 321  
 <212> DNA  
 <213> Zea mays

<220>  
 <223> Fragment of maize chloroplast rpoC2 gene

```

<400> 32
tcattcttca agaactgcat atcttgccga gatcctcacc cctaaaggta cttgacaata 60
gtattatttg agtcgataca caactcacia aaaatacaag aagtcgacta ggtggatttg 120
tccgagtga gagaaaaaaa agccatacag aactcaaaat cttttccgga gatattcatt 180
ttctgaaga ggcggataag atattaggtg gcagtttgat accaccagaa agagaaaaaa 240
aagattctaa ggaatcaaaa aaaaggaaaa attgggttta tgttcaacgg aaaaaatttc 300
tcaaaagcaa ggaaaagtat t 321

```

<210> 33  
 <211> 17  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Represents part of PHI8999A insert  
 sequence-fragment of ubi2M1(2) promoter; also a  
 fragment of the maize chloroplast trn1 gene

```

<400> 33
gtggctatct atctatc 17

```

<210> 34  
 <211> 201  
 <212> DNA  
 <213> Artificial Sequence

<220>

<223> Represents part of PHI8999A insert sequence -  
fragment of pat gene

<400> 34  
gcagctgata tggccgcggt ttgtgatata gttaaccatt acattgagac gtctacagtg 60  
aacttttagga cagagccaca aacaccacaa gagggtgattg atgatctaga gaggttgcaa 120  
gatagatacc cttggttggg tgctgaggtt gaggggtgtg tggctggat tgcttacgct 180  
gggcctgga aggctaggaa c 201

<210> 35  
<211> 138  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Represents part of PHI8999A insert sequence -  
fragment of pat gene (complement)

<400> 35  
cctcaacctc agcaaccaac caatggtatc tatcttgcaa cctctctaga tcatcaatcc 60  
actcttgagg tgtttgtggc tctgtcctaa agttcactgt agacgtctca atgtaagtgt 120  
taacgatata acaaaccg 138

<210> 36  
<211> 15  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Represents part of PHI8999A insert sequence -  
fragment of crylF gene (complement)

<400> 36  
agagaagagg gatct 15

<210> 37  
<211> 118  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Represents part of PHI8999A insert sequence -  
fragment of polylinker

<400> 37  
cgaagcttcg gccggggccc atcgatatcc gcgggcatgc ctgcagtga gcgtgaccg 60  
gtcgtgcccc tctctagaga taatgagcat tgcatgtcta agttataaaa aattacca 118

<210> 38  
<211> 550  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Represents part of PHI8999A insert sequence -  
fragment of ORF25 terminator (complement)

<400> 38

```
ctcactccgc ttgatcttgg caaagatatt tgacgcattt attagtatgt gttaattttc 60
atttgcagtg cagtattttc tattcgatct ttatgtaatt cgttacaatt aataaatatt 120
caaatcagat tattgactgt catttgtatc aaatcgtgtt taatggatat ttttattata 180
atattgatga tatctcaatc aaaacgtaga taataataat atttatttaa tatttttgcg 240
tcgcacagtg aaaatctata tgagattaca aaataccgac aacattattt aagaaacata 300
gacattaacc ctgagactgt tggacatcaa cgggtagatt ccttcatgca tagcacctca 360
ttcttgggga caaaagcacg gtttggccgt tccattgctg cacgaacgag ctttgctata 420
tcctcgggtt ggatcatctc atcagggtcca atcaaatttg tccaagaact catgttagtc 480
gcaacgaaac cggggcatat gtcgggtatc tcgagctcgc gaaagcttgg ctgcaggctc 540
acggatcctt                                     550
```

<210> 39  
 <211> 128  
 <212> DNA  
 <213> Zea mays

<220>  
 <223> Fragment of maize chloroplast rps12 rRNA gene  
 (complement)

```
<400> 39
caacaaaagg gtacctgtac ccgaaaccga cacagggtgg taggtagaga atacctaggg 60
gcgcgagaca actctctcta aggaactcgg caaaatagcc ccgtaacttc gggagaaggg 120
gtgcccc
```

<210> 40  
 <211> 392  
 <212> DNA  
 <213> Zea mays

<220>  
 <223> Fragment of maize chloroplast genome

```
<400> 40
ctaacaataa acgaatacgg tttatgtatg gattccggta aaataccggt actcgatttc 60
ataagagtcg aataggaagt taagatgagg gtggtatcat cataaaaaatg gagtagtata 120
ctaaattata ctaatccacg tatgatatgt atgcctttcc ttatcaaccg gaagtagtgc 180
aaaaaaaatt ctatactgca ctgctctctt ttactgaga aatgcacaaa aataaaagtg 240
aagtaagggt gcccataga ttttgatctt tgctctctgt cccccccccc cttttttcat 300
caaaaatttc catgaaaaaa gaaaagatga atttgtccat tcattgaacc ctagtctggg 360
actgacgggg ctgcaacccg cagcttccgc ct                                     392
```

<210> 41  
 <211> 188  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Represents part of PHI8999A insert sequence -  
 fragment of pat gene (complement)

```
<400> 41
gttcctagcc ttccagggcc cagcgtaagc aataccagcc acagcaccct caacctcagc 60
aaccaaccaa gggatcttat cttgcaacct ctctagatca tcaatccact cttgtggtgt 120
ttgtggtctt gtctctaaagt tcaactgtaga cgtctcaatg taatgggtta cgatatacaca 180
aaccgcgg
```

<210> 42

<211> 81  
<212> DNA  
<213> Zea mays

<220>  
<223> Fragment of maize chloroplast ORF241 (complement)

<400> 42  
cacaagaacg aaagcacctt ttcattcttt catatactag gggtttttac ttggaaaaga 60  
caatgttcca tactaaagga t 81

<210> 43  
<211> 254  
<212> DNA  
<213> Zea mays

<400> 43  
agctgcagaa gccgccaccg tcttgaggac cttccgggga gccagaccgg tcgaaccgtg 60  
cctccacttg ctaaggagaa agggaaaatc agggccagga catacgaagg aggagccaga 120  
acgaagatat cctaagatac ttactcgctc cgggccatga tcaatcatgc ctgtggggag 180  
gtctctcgca cctcgatcca tgaaggtacc accgaggtct gccccgccgc cggcttcggt 240  
accgtcctcg cctt 254

<210> 44  
<211> 749  
<212> DNA  
<213> Zea mays

<400> 44  
ggcgccccga ggcaccocgg ggatggactg cccaggcgca gccacgacga cccaaggatc 60  
accctcctgc gcagtcggca cgagcaatag ttctcgggga acaggcagct tggcctgact 120  
ccccggggtc acctcaacta cctcggccga ggggtcaagt accccctcag tccgcccccg 180  
ctcttcggac cgggaccccg acgtcccggc cccggatacc gacggcacca gcccgctcgg 240  
gggctggcct gacgacccct ggcccagcct cagatctggg ctgaggccga ggcaggcggc 300  
catgtcgtcg tcttcatcat cgtcttcatc atcgtcgtcg tcatcaggcg tctccggcga 360  
cggctccctt gggagccctt ccctctcctg ccgacgacga agcctttcca aggcatccc 420  
agcccaagtc cgtcgtggg cccgagcctt ctttgcgtcc ttcttctcct tctcttctc 480  
cgcggtgacc ctccgcgcag ctccgtccac cgcacccctc gggactgggtg gcagggaagg 540  
cttgtgatgc cctacctcct ggagacagac gaaaagtctc agctatgaga accgagggca 600  
atctgacgca agaaggaaga aggagcggat actcaccaga gacacgcacc cgcgatcggg 660  
acgcattaag ggctgggaaa aagtgccggc ctctaatttc gctaccgtgc cgtccaccca 720  
cctgtggagg tcatcgatgg gaaggggaa 749

<210> 45  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> 5' event flanking sequence; junction between  
regions 3 and 4

<400> 45  
tcggactcga cctttctatc 20

<210> 46  
<211> 20  
<212> DNA

<213> Artificial Sequence

<220>

<223> 5' event flanking sequence; junction between  
regions 4 and 5

<400> 46

agagaatcac tcattcttca

20

<210> 47

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> 5' event flanking sequence; junction between  
regions 5 and 6

<400> 47

gaaaagtatt gtggctatatt

20

<210> 48

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> 5' event flanking sequence; junction between  
regions 6 and 7a

<400> 48

atctatccgt gcagctgata

20

<210> 49

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> 5' event flanking sequence; junction between  
regions 7a and 7b

<400> 49

ggctaggaac cctcaacctc

20

<210> 50

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> 5' event flanking sequence; junction between  
regions 7b and 7c

<400> 50

tcacaaaccg agagaagagg

20

<210> 51

20

<211> 20  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> 5' event flanking sequence; junction between  
           regions 7c and 8  
  
 <400> 51  
 agagggatct cgaagcttcg 20  
  
 <210> 52  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> 5' event flanking sequence; junction between  
           regions 8 and 9  
  
 <400> 52  
 aaaattacca caactggaag 20  
  
 <210> 53  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> 3' event flanking sequence; junction between  
           regions 9 and 10  
  
 <400> 53  
 agctatgttt ctcaactccgc 20  
  
 <210> 54  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> 3' event flanking sequence; junction between  
           regions 10 and 11  
  
 <400> 54  
 acggatcctt caacaaaagg 20  
  
 <210> 55  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> 3' event flanking sequence; junction between  
           regions 11 and 12  
  
 <400> 55  
 gtgcccccg ctaacaataa 20

6046772 050203

<210> 56  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> 3' event flanking sequence; junction between  
regions 12 and 13

<400> 56  
gcttcgcct gttcctagcc 20

<210> 57  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> 3' event flanking sequence; junction between  
regions 13 and 14

<400> 57  
caaaccgcgg cacaagaacg 20