



US 20060037095A9

(19) **United States**(10) **Pub. No.: US 2006/0037095 A9**(12) **Patent Application Publication**
Barbour et al.(48) **Pub. Date: Feb. 16, 2006**
CORRECTED PUBLICATION(54) **CORN EVENT TC1507 AND METHODS FOR
DETECTION THEREOF**(21) Appl. No.: **10/837,105**(75) Inventors: **Eric Barbour**, Johnston, IA (US);
James Wayne Bing, Ankeny, IA (US);
Guy A. Cardineau, Tempe, AZ (US);
Robert F. Cressman JR., Wilmington,
DE (US); **Manju Gupta**, Carmel, IN
(US); **Mary E. Hartnett Locke**,
Mickleton, NJ (US); **David Hondred**,
Ankeny, IA (US); **Joseph W.**
Keaschall, Clive, IA (US); **Michael G.**
Koziel, Raleigh, NC (US); **Terry**
EuClaire Meyer, Urbandale, IA (US);
Daniel Moellenbeck, Granger, IA (US);
Kenneth Edwin Narva, Carlsbad, CA
(US); **Wilas Nirunsuksiri**, Auburn, WA
(US); **Steven W. Ritchie**, Omaha, NE
(US); **Marjorie L. Rudert**, Boone, IA
(US); **Craig D. Sanders**, Bear, DE
(US); **Aihua Shao**, Johnston, IA (US);
Steven Jeffrey Stelman, San Diego,
CA (US); **David S. Stucker**, Johnston,
IA (US); **Laura Ann Tagliani**,
Zionsville, IN (US); **William M. Van**
Zante, Urbandale, IA (US)(22) Filed: **Apr. 30, 2004****Prior Publication Data**(15) Correction of US 2005/0039226 A1 Feb. 17, 2005
See Claims 15, 20, and 23

(65) US 2005/0039226 A1 Feb. 17, 2005

Related U.S. Application Data(60) Provisional application No. 60/467,772, filed on May
2, 2003.**Publication Classification**(51) **Int. Cl.**
A01H 1/00 (2006.01)
C12N 15/82 (2006.01)
A01H 5/00 (2006.01)
C12Q 1/68 (2006.01)
C07H 21/04 (2006.01)
(52) **U.S. Cl.** **800/278**; 800/320.1; 536/23.2;
435/6

Correspondence Address:

PIONEER HI-BRED INTERNATIONAL, INC.
7250 N.W. 62ND AVENUE
P.O. BOX 552
JOHNSTON, IA 50131-0552 (US)(73) Assignees: **Dow AgroSciences LLC**; **E.I. duPont**
deNemours and Company; **Pioneer Hi-**
Bred International, Inc.(57) **ABSTRACT**The invention provides DNA compositions that relate to
transgenic insect resistant maize plants. Also provided are
assays for detecting the presence of the maize TC1507 event
based on the DNA sequence of the recombinant construct
inserted into the maize genome and the DNA sequences
flanking the insertion site. Kits and conditions useful in
conducting the assays are provided.




FIG. 1

CORN EVENT TC1507 AND METHODS FOR DETECTION THEREOF

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the benefit of U.S. Provisional Application No. 60/467,772, filed May 2, 2003.

FIELD OF THE INVENTION

[0002] The present invention relates to the field of plant molecular biology, specifically the invention relates to a DNA construct for conferring insect resistance to a plant. The invention more specifically relates to an insect resistant corn plant TC1507 and to assays for detecting the presence of corn plant TC1507 DNA in a sample and compositions thereof.

BACKGROUND OF THE INVENTION

[0003] This invention relates to the insect resistant corn (*Zea mays*) plant TC1507, also referred to as maize line TC1507 or maize event TC1507, and to the DNA plant expression construct of corn plant TC1507 and the detection of the transgene/flanking insertion region in corn plant TC1507 and progeny thereof.

[0004] Corn is an important crop and is a primary food source in many areas of the world. Damage caused by insect pests is a major factor in the loss of the world's corn crops, despite the use of protective measures such as chemical pesticides. In view of this, insect resistance has been genetically engineered into crops such as corn in order to control insect damage and to reduce the need for traditional chemical pesticides. One group of genes which have been utilized for the production of transgenic insect resistant crops are the delta-endotoxins from *Bacillus thuringiensis* (*B.t.*). Delta-endotoxins have been successfully expressed in crop plants such as cotton, potatoes, rice, sunflower, as well as corn, and have proven to provide excellent control over insect pests. (Perlak, F. J et al. (1990) *Bio/Technology* 8, 939-943; Perlak, F. J. et al. (1993) *Plant Mol. Biol.* 22: 313-321; Fujimoto H. et al. (1993) *Bio/Technology* 11: 1151-1155; Tu et al. (2000) *Nature Biotechnology* 18:1101-1104; PCT publication number WO 01/13731; and Bing J W et al. (2000) Efficacy of Cry1F Transgenic Maize, 14th Biennial International Plant Resistance to Insects Workshop, Fort Collins, Colo.).

[0005] The expression of foreign genes in plants is known to be influenced by their location in the plant genome, perhaps due to chromatin structure (e.g., heterochromatin) or the proximity of transcriptional regulatory elements (e.g., enhancers) close to the integration site (Weising et al., *Ann. Rev. Genet* 22:421-477, 1988). At the same time the presence of the transgene at different locations in the genome will influence the overall phenotype of the plant in different ways. For this reason, it is often necessary to screen a large number of events in order to identify an event characterized by optimal expression of an introduced gene of interest. For example, it has been observed in plants and in other organisms that there may be a wide variation in levels of expression of an introduced gene among events. There may also be differences in spatial or temporal patterns of expression, for example, differences in the relative expression of a transgene in various plant tissues, that may not correspond to the patterns expected from transcriptional regulatory elements

present in the introduced gene construct. For this reason, it is common to produce hundreds to thousands of different events and screen those events for a single event that has desired transgene expression levels and patterns for commercial purposes. An event that has desired levels or patterns of transgene expression is useful for introgressing the transgene into other genetic backgrounds by sexual outcrossing using conventional breeding methods. Progeny of such crosses maintain the transgene expression characteristics of the original transformant. This strategy is used to ensure reliable gene expression in a number of varieties that are well adapted to local growing conditions.

[0006] It would be advantageous to be able to detect the presence of a particular event in order to determine whether progeny of a sexual cross contain a transgene of interest. In addition, a method for detecting a particular event would be helpful for complying with regulations requiring the pre-market approval and labeling of foods derived from recombinant crop plants, for example, or for use in environmental monitoring, monitoring traits in crops in the field, or monitoring products derived from a crop harvest, as well as for use in ensuring compliance of parties subject to regulatory or contractual terms.

[0007] It is possible to detect the presence of a transgene by any nucleic acid detection method known in the art including, but not limited to, the polymerase chain reaction (PCR) or DNA hybridization using nucleic acid probes. These detection methods generally focus on frequently used genetic elements, such as promoters, terminators, marker genes, etc., because for many DNA constructs, the coding region is interchangeable. As a result, such methods may not be useful for discriminating between different events, particularly those produced using the same DNA construct or very similar constructs unless the DNA sequence of the flanking DNA adjacent to the inserted heterologous DNA is known. For example, an event-specific PCR assay is described in U.S. Pat. No. 6,395,485 for the detection of elite event GAT-ZM1. Accordingly, it would be desirable to have a simple and discriminative method for the identification of event TC1507.

SUMMARY OF THE INVENTION

[0008] This invention relates preferably to methods for producing and selecting an insect resistant monocot crop plant. More specifically, a DNA construct is provided that when expressed in plant cells and plants confers resistance to insects. According to one aspect of the invention, a DNA construct, capable of introduction into and replication in a host cell, is provided that when expressed in plant cells and plants confers insect resistance to the plant cells and plants. The DNA construct is comprised of a DNA molecule named PH18999A and it includes two transgene expression cassettes. The first expression cassette comprises a DNA molecule which includes the promoter, 5' untranslated exon, and first intron of the maize ubiquitin (Ubi-1) gene (Christensen et al. (1992) *Plant Mol. Biol.* 18:675-689 and Christensen and Quail (1996) *Transgenic Res.* 5:213-218) operably connected to a DNA molecule encoding a *B.t.* 6-endotoxin identified as Cry1F (U.S. Pat. Nos. 5,188,960 and 6,218,188) operably connected to a DNA molecule comprising a 3' ORF25 transcriptional terminator isolated from *Agrobacterium tumefaciens* (Barker et al. (1983) *Plant Mol. Biol.* 2:335-350). The second transgene expression cassette of the

DNA construct comprises a DNA molecule of the cauliflower mosaic virus (CaMV) ³⁵S promoter (Odell J. T. et al. (1985) *Nature* 313: 810-812; Mitsuhashi et al. (1996) *Plant Cell Physiol.* 37: 49-59) operably connected to a DNA molecule encoding a phosphinothricin acetyltransferase (PAT) gene (Wohlleben W. et al. (1988) *Gene* 70: 25-37) operably connected to a DNA molecule comprising a 3' transcriptional terminator from (CaMV) ³⁵S (see Mitsuhashi et al. (1996) *Plant Cell Physiol.* 37: 49-59). Plants containing the DNA construct are also provided.

[0009] According to another aspect of the invention, compositions and methods are provided for identifying a novel corn plant designated TC1507, which methods are based on primers or probes which specifically recognize the 5' and/or 3' flanking sequence of TC1507. DNA molecules are provided that comprise primer sequences that when utilized in a PCR reaction will produce amplicons unique to the transgenic event TC1507. These molecules may be selected from the group consisting of:

5'-GTAGTACTATAGATTATATTATTCGTA (SEQ ID NO: 1)
GAG-3';

5'-GCCATACAGAACTCAAAATCTTTCCG (SEQ ID NO: 2)
GAG-3';

5'-CTTCAACAAGTGTGACAAA-3'; (SEQ ID NO: 23)

5'-TGTTGGTGTGTTGGCTCTGTCTCTAA-3'; (SEQ ID NO: 3)

5'-AGCACCTTTTCATTCTTCATATAC-3'; (SEQ ID NO: 4)

5'-GACCTCCCCA CAGGCATGAT TGATC-3'; (SEQ ID NO: 5)

and complements thereof. The corn plant and seed comprising these molecules is an aspect of this invention. Further, kits utilizing these primer sequences for the identification of the TC1507 event are provided.

[0010] An additional aspect of the invention relates to the specific flanking sequences of TC1507 described herein, which can be used to develop specific identification methods for TC1507 in biological samples. More particularly, the invention relates to the 5' and/or 3' flanking regions of TC1507, SEQ ID NO:21 and SEQ ID NO:22, respectively, which can be used for the development of specific primers and probes. The invention further relates to identification methods for the presence of TC1507 in biological samples based on the use of such specific primers or probes.

[0011] According to another aspect of the invention, methods of detecting the presence of DNA corresponding to the corn event TC1507 in a sample are provided. Such methods comprise: (a) contacting the sample comprising DNA with a DNA primer set, that when used in a nucleic acid amplification reaction with genomic DNA extracted from corn event TC1507 produces an amplicon that is diagnostic for corn event TC1507; (b) performing a nucleic acid amplification reaction, thereby producing the amplicon; and (c) detecting the amplicon.

[0012] DNA molecules that comprise the novel transgene/flanking insertion region, SEQ ID NO: 26 and SEQ ID NO: 27 and are homologous or complementary to SEQ ID NO: 26 and SEQ ID NO: 27 are an aspect of this invention.

[0013] DNA sequences that comprise the novel transgene/flanking insertion region, SEQ ID NO:26 are an aspect of this invention. DNA sequences that comprise a sufficient length of polynucleotides of transgene insert sequence and a sufficient length of polynucleotides of maize genomic and/or flanking sequence from maize plant TC1507 of SEQ ID NO:26 that are useful as primer sequences for the production of an amplicon product diagnostic for maize plant TC1507 are included.

[0014] In addition, DNA sequences that comprise the novel transgene/flanking insertion region, SEQ ID NO:27 are provided. DNA sequences that comprise a sufficient length of polynucleotides of transgene insert sequence and a sufficient length of polynucleotides of maize genomic and/or flanking sequence from maize plant TC1507 of SEQ ID NO:27 that are useful as primer sequences for the production of an amplicon product diagnostic for maize plant TC1507 are included.

[0015] According to another aspect of the invention, the DNA sequences that comprise at least 11 or more nucleotides of the transgene portion of the DNA sequence of SEQ ID NO:26 or complements thereof, and a similar length of 5' flanking maize DNA sequence of SEQ ID NO:26 or complements thereof are useful as DNA primers in DNA amplification methods. The amplicons produced using these primers are diagnostic for maize event TC1507. Therefore, the invention also includes the amplicons produced by DNA primers homologous or complementary to SEQ ID NO:26.

[0016] According to another aspect of the invention, the DNA sequences that comprise at least 11 or more nucleotides of the transgene portion of the DNA sequence of SEQ ID NO:27 or complements thereof, and a similar length of 3' flanking maize DNA sequence of SEQ ID NO:27 or complements thereof are useful as DNA primers in DNA amplification methods. The amplicons produced using these primers are diagnostic for maize event TC1507. Therefore, the invention also includes the amplicons produced by DNA primers homologous or complementary to SEQ ID NO:27.

[0017] More specifically, a pair of DNA molecules comprising a DNA primer set, wherein the DNA molecules are identified as SEQ ID NO: 1 or complements thereof and SEQ ID NO: 2 or complements thereof; SEQ ID NO: 2 or complements thereof and SEQ ID NO: 23 or complements thereof; SEQ ID NO: 3 or complements thereof and SEQ ID NO: 5 or complements thereof; SEQ ID NO: 4 or complements thereof and SEQ ID NO: 5 or complements thereof are aspects of the invention.

[0018] Further aspects of the invention include the amplicon comprising the DNA molecules of SEQ ID NO: 1 and SEQ ID NO: 2; the amplicon comprising the DNA molecules of SEQ ID NO: 2 and SEQ ID NO: 23; the amplicon comprising the DNA molecules of SEQ ID NO: 3 and SEQ ID NO: 5; and the amplicon comprising the DNA molecules of SEQ ID NO: 4 and SEQ ID NO: 5.

[0019] According to another aspect of the invention, methods of detecting the presence of a DNA molecule corresponding to the TC1507 event in a sample, such methods comprising: (a) contacting the sample comprising DNA extracted from a corn plant with a DNA probe, molecule that hybridizes under stringent hybridization conditions with DNA extracted from corn event TC1507 and does not

hybridize under the stringent hybridization conditions with a control corn plant DNA; (b) subjecting the sample and probe to stringent hybridization conditions; and (c) detecting hybridization of the probe to the DNA. More specifically, a method for detecting the presence of a DNA molecule corresponding to the TC1507 event in a sample, such methods, consisting of (a) contacting the sample comprising DNA extracted from a corn plant with a DNA probe molecule that consists of sequences that are unique to the event, e.g. junction sequences, wherein said DNA probe molecule hybridizes under stringent hybridization conditions with DNA extracted from corn event TC1507 and does not hybridize under the stringent hybridization conditions with a control corn plant DNA; (b) subjecting the sample and probe to stringent hybridization conditions; and (c) detecting hybridization of the probe to the DNA.

[0020] In addition, a kit and methods for identifying event TC1507 in a biological sample which detects a TC1507 specific region within SEQ ID NO: 24 are provided.

[0021] DNA molecules are provided that comprise at least one junction sequence of TC1507 selected from the group consisting of SEQ ID NO:45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56 and 57 and complements thereof; wherein a junction sequence spans the junction between heterologous DNA inserted into the genome and the DNA from the corn cell flanking the insertion site, i.e. flanking DNA, and is diagnostic for the TC1507 event.

[0022] According to another aspect of the invention, methods of producing an insect resistant corn plant that comprise the steps of: (a) sexually crossing a first parental corn line comprising the expression cassettes of the present invention, which confers resistance to insects, and a second parental corn line that lacks insect resistance, thereby producing a plurality of progeny plants; and (b) selecting a progeny plant that is insect resistant. Such methods may optionally comprise the further step of back-crossing the progeny plant to the second parental corn line to producing a true-breeding corn plant that is insect resistant.

[0023] The present invention provides a method of producing a corn plant that is resistant to insects comprising transforming a corn cell with the DNA construct PH18999A (SEQ ID NO:25), growing the transformed corn cell into a corn plant, selecting the corn plant that shows resistance to insects, and further growing the corn plant into a fertile corn plant. The fertile corn plant can be self pollinated or crossed with compatible corn varieties to produce insect resistant progeny.

[0024] The invention further relates to a DNA detection kit for identifying maize event TC1507 in biological samples. Preferably the kit of the invention comprises a first primer which specifically recognizes the 5' or 3' flanking region of TC1507, and a second primer which specifically recognizes a sequence within the foreign DNA of TC1507, or within the flanking DNA, for use in a PCR identification protocol. The invention also relates to a kit for identifying event TC1507 in biological samples, which kit comprises a specific probe having a sequence which corresponds or is complementary to, a sequence having between 80% and 100% sequence identity with a specific region of event TC1507. Preferably the sequence of the probe corresponds to a specific region comprising part of the 5' or 3' flanking region of event TC1507.

[0025] The methods and kits encompassed by the present invention can be used for different purposes such as, but not limited to the following: to identify event TC1507 in plants, plant material or in products such as, but not limited to, food or feed products (fresh or processed) comprising, or derived from plant material; additionally or alternatively, the methods and kits of the present invention can be used to identify transgenic plant material for purposes of segregation between transgenic and non-transgenic material; additionally or alternatively, the methods and kits of the present invention can be used to determine the quality of plant material comprising maize event TC1507. The kits may also contain the reagents and materials necessary for the performance of the detection method.

[0026] This invention further relates to the TC1507 corn plant or its parts, including, but not limited to, pollen, ovules, vegetative cells, the nuclei of pollen cells, and the nuclei of egg cells of the corn plant TC1507 and the progeny derived thereof. The corn plant and seed TC1507 from which the DNA primer molecules of the present invention provide a specific amplicon product is an aspect of the invention.

[0027] The foregoing and other aspects of the invention will become more apparent from the following detailed description and accompanying drawing.

BRIEF DESCRIPTION OF THE DRAWINGS

[0028] **FIG. 1.** Linear map showing the transgenic insert PH18999A, as well as the sequences flanking the transgenic insert.

DETAILED DESCRIPTION

[0029] The following definitions and methods are provided to better define the present invention and to guide those of ordinary skill in the art in the practice of the present invention. Unless otherwise noted, terms are to be understood according to conventional usage by those of ordinary skill in the relevant art. Definitions of common terms in molecular biology may also be found in Rieger et al., *Glossary of Genetics: Classical and Molecular*, 5th edition, Springer-Verlag; New York, 1991; and Lewin, *Genes V*, Oxford University Press: New York, 1994. The nomenclature for DNA bases as set forth at 37 CFR 1.822 is used.

[0030] As used herein, the term “comprising” means “including but not limited to”.

[0031] As used herein, the term “corn” means *Zea mays* or maize and includes all plant varieties that can be bred with corn, including wild maize species.

[0032] As used herein, the term “TC1507 specific” refers to a nucleotide sequence which is suitable for discriminatively identifying event TC1507 in plants, plant material, or in products such as, but not limited to, food or feed products (fresh or processed) comprising, or derived from plant material.

[0033] As used herein, the terms “insect resistant” and “impacting insect pests” refers to effecting changes in insect feeding, growth, and/or behavior at any stage of development, including but not limited to: killing the insect; retarding growth; preventing reproductive capability; and the like.

[0034] As used herein, the terms “pesticidal activity” and “insecticidal activity” are used synonymously to refer to activity of an organism or a substance (such as, for example, a protein) that can be measured by numerous parameters including, but not limited to, pest mortality, pest weight loss, pest attraction, pest repellency, and other behavioral and physical changes of a pest after feeding on and/or exposure to the organism or substance for an appropriate length of time. For example “pesticidal proteins” are proteins that display pesticidal activity by themselves or in combination with other proteins.

[0035] “Coding sequence” refers to a nucleotide sequence that codes for a specific amino acid sequence. As used herein, the terms “encoding” or “encoded” when used in the context of a specified nucleic acid mean that the nucleic acid comprises the requisite information to guide translation of the nucleotide sequence into a specified protein. The information by which a protein is encoded is specified by the use of codons. A nucleic acid encoding a protein may comprise non-translated sequences (e.g., introns) within translated regions of the nucleic acid or may lack such intervening non-translated sequences (e.g., as in cDNA).

[0036] “Gene” refers to a nucleic acid fragment that expresses a specific protein, including regulatory sequences preceding (5' non-coding sequences) and following (3' non-coding sequences) the coding sequence. “Native gene” refers to a gene as found in nature with its own regulatory sequences. “Chimeric gene” refers any gene that is not a native gene, comprising regulatory and coding sequences that are not found together in nature. Accordingly, a chimeric gene may comprise regulatory sequences and coding sequences that are derived from different sources, or regulatory sequences and coding sequences derived from the same source, but arranged in a manner different than that found in nature. “Endogenous gene” refers to a native gene in its natural location in the genome of an organism. “Foreign” refers to material not normally found in the location of interest. Thus “foreign DNA” may comprise both recombinant DNA as well as newly introduced, rearranged DNA of the plant. A “foreign” gene refers to a gene not normally found in the host organism, but that is introduced into the host organism by gene transfer. Foreign genes can comprise native genes inserted into a non-native organism, or chimeric genes. A “transgene” is a gene that has been introduced into the genome by a transformation procedure. The site in the plant genome where a recombinant DNA has been inserted may be referred to as the “insertion site” or “target site”.

[0037] As used herein, “insert DNA” refers to the heterologous DNA within the expression cassettes used to transform the plant material while “flanking DNA” can exist of either genomic DNA naturally present in an organism such as a plant, or foreign (heterologous) DNA introduced via the transformation process which is extraneous to the original insert DNA molecule, e.g. fragments associated with the transformation event. A “flanking region” or “flanking sequence” as used herein refers to a sequence of at least 20 base pair, preferably at least 50 base pair, and up to 5000 base pair which is located either immediately upstream of and contiguous with or immediately downstream of and contiguous with the original foreign insert DNA molecule. Transformation procedures leading to random integration of the foreign DNA will result in transformants containing

different flanking regions characteristic and unique for each transformant. When recombinant DNA is introduced into a plant through traditional crossing, its flanking regions will generally not be changed. Transformants will also contain unique junctions between a piece of heterologous insert DNA and genomic DNA, or 2 pieces of genomic DNA, or 2 pieces of heterologous DNA. A “junction” is a point where 2 specific DNA fragments join. For example, a junction exists where insert DNA joins flanking DNA. A junction point also exists in a transformed organism where 2 DNA fragments join together in a manner that is modified from that found in the native organism. “Junction DNA” refers to DNA that comprises a junction point.

[0038] As used herein, “heterologous” in reference to a nucleic acid is a nucleic acid that originates from a foreign species, or, if from the same species, is substantially modified from its native form in composition and/or genomic locus by deliberate human intervention. For example, a promoter operably linked to a heterologous nucleotide sequence can be from a species different from that from which the nucleotide sequence was derived, or, if from the same species, the promoter is not naturally found operably linked to the nucleotide sequence. A heterologous protein may originate from a foreign species, or, if from the same species, is substantially modified from its original form by deliberate human intervention.

[0039] “Regulatory sequences” refer to nucleotide sequences located upstream (5' non-coding sequences), within, or downstream (3' non-coding sequences) of a coding sequence, and which influence the transcription, RNA processing or stability, or translation of the associated coding sequence. Regulatory sequences may include promoters, translation leader sequences, introns, and polyadenylation recognition sequences.

[0040] “Promoter” refers to a nucleotide sequence capable of controlling the expression of a coding sequence or functional RNA. In general, a coding sequence is located 3' to a promoter sequence. The promoter sequence consists of proximal and more distal upstream elements, the latter elements are often referred to as enhancers. Accordingly, an “enhancer” is a nucleotide sequence that can stimulate promoter activity and may be an innate element of the promoter or a heterologous element inserted to enhance the level or tissue-specificity of a promoter. Promoters may be derived in their entirety from a native gene, or be composed of different elements derived from different promoters found in nature, or even comprise synthetic nucleotide segments. It is understood by those skilled in the art that different promoters may direct the expression of a gene in different tissues or cell types, or at different stages of development, or in response to different environmental conditions. Promoters that cause a nucleic acid fragment to be expressed in most cell types at most times are commonly referred to as “constitutive promoters”. New promoters of various types useful in plant cells are constantly being discovered; numerous examples may be found in the compilation by Okamuro and Goldberg (1989) *Biochemistry of Plants* 15:1-82. It is further recognized that since in most cases the exact boundaries of regulatory sequences have not been completely defined, nucleic acid fragments of different lengths may have identical promoter activity.

[0041] The “translation leader sequence” refers to a nucleotide sequence located between the promoter sequence of a

gene and the coding sequence. The translation leader sequence is present in the fully processed mRNA upstream of the translation start sequence. The translation leader sequence may affect numerous parameters including, processing of the primary transcript to mRNA, mRNA stability and/or translation efficiency. Examples of translation leader sequences have been described (Turner and Foster (1995) *Mol. Biotechnol.* 3:225-236).

[0042] The “3’ non-coding sequences” refer to nucleotide sequences located downstream of a coding sequence and include polyadenylation recognition sequences and other sequences encoding regulatory signals capable of affecting mRNA processing or gene expression. The polyadenylation signal is usually characterized by affecting the addition of polyadenylic acid tracts to the 3’ end of the mRNA precursor. The use of different 3’ non-coding sequences is exemplified by Ingelbrecht et al. (1989) *Plant Cell* 1:671-680.

[0043] A “protein” or “polypeptide” is a chain of amino acids arranged in a specific order determined by the coding sequence in a polynucleotide encoding the polypeptide.

[0044] A DNA construct is an assembly of DNA molecules linked together that provide one or more expression cassettes. The DNA construct may be a plasmid that is enabled for self replication in a bacterial cell and contains various endonuclease enzyme restriction sites that are useful for introducing DNA molecules that provide functional genetic elements, i.e., promoters, introns, leaders, coding sequences, 3’ termination regions, among others; or a DNA construct may be a linear assembly of DNA molecules, such as an expression cassette. The expression cassette contained within a DNA construct comprise the necessary genetic elements to provide transcription of a messenger RNA. The expression cassette can be designed to express in prokaryote cells or eukaryotic cells. Expression cassettes of the present invention are designed to express most preferably in plant cells.

[0045] The DNA molecules of the invention are provided in expression cassettes for expression in an organism of interest. The cassette will include 5’ and 3’ regulatory sequences operably linked to a coding sequence of the invention. “Operably linked” means that the nucleic acid sequences being linked are contiguous and, where necessary to join two protein coding regions, contiguous and in the same reading frame. Operably linked is intended to indicate a functional linkage between a promoter and a second sequence, wherein the promoter sequence initiates and mediates transcription of the DNA sequence corresponding to the second sequence. The cassette may additionally contain at least one additional gene to be cotransformed into the organism. Alternatively, the additional gene(s) can be provided on multiple expression cassettes or multiple DNA constructs.

[0046] The expression cassette will include in the 5’ to 3’ direction of transcription: a transcriptional and translational initiation region, a coding region, and a transcriptional and translational termination region functional in the organism serving as a host. The transcriptional initiation region (i.e., the promoter) may be native or analogous, or foreign or heterologous to the host organism. Additionally, the promoter may be the natural sequence or alternatively a synthetic sequence. The expression cassettes may additionally contain 5’ leader sequences in the expression cassette construct. Such leader sequences can act to enhance translation.

[0047] It is to be understood that as used herein the term “transgenic” includes any cell, cell line, callus, tissue, plant part, or plant the genotype of which has been altered by the presence of a heterologous nucleic acid including those transgenics initially so altered as well as those created by sexual crosses or asexual propagation from the initial transgenic. The term “transgenic” as used herein does not encompass the alteration of the genome (chromosomal or extra-chromosomal) by conventional plant breeding methods or by naturally occurring events such as random cross-fertilization, non-recombinant viral infection, non-recombinant bacterial transformation, non-recombinant transposition, or spontaneous mutation.

[0048] A transgenic “event” is produced by transformation of plant cells with a heterologous DNA construct(s), including a nucleic acid expression cassette that comprises a transgene of interest, the regeneration of a population of plants resulting from the insertion of the transgene into the genome of the plant, and selection of a particular plant characterized by insertion into a particular genome location. An event is characterized phenotypically by the expression of the transgene. At the genetic level, an event is part of the genetic makeup of a plant. The term “event” also refers to progeny produced by a sexual outcross between the transformant and another variety that include the heterologous DNA. Even after repeated back-crossing to a recurrent parent, the inserted DNA and flanking DNA from the transformed parent is present in the progeny of the cross at the same chromosomal location. The term “event” also refers to DNA from the original transformant comprising the inserted DNA and flanking sequence immediately adjacent to the inserted DNA that would be expected to be transferred to a progeny that receives inserted DNA including the transgene of interest as the result of a sexual cross of one parental line that includes the inserted DNA (e.g., the original transformant and progeny resulting from selfing) and a parental line that does not contain the inserted DNA.

[0049] An insect resistant TC1507 corn plant can be bred by first sexually crossing a first parental corn plant consisting of a corn plant grown from the transgenic TC1507 corn plant and progeny thereof derived from transformation with the expression cassettes of the present invention that confers insect resistance, and a second parental corn plant that lacks insect resistance, thereby producing a plurality of first progeny plants; and then selecting a first progeny plant that is resistant to insects; and selfing the first progeny plant, thereby producing a plurality of second progeny plants; and then selecting from the second progeny plants an insect resistant plant. These steps can further include the back-crossing of the first insect resistant progeny plant or the second insect resistant progeny plant to the second parental corn plant or a third parental corn plant, thereby producing a corn plant that is resistant to insects.

[0050] As used herein, the term “plant” includes reference to whole plants, plant organs (e.g., leaves, stems, roots, etc.), seeds, plant cells, and progeny of same. Parts of transgenic plants understood to be within the scope of the invention comprise, for example, plant cells, protoplasts, tissues, callus, embryos as well as flowers, stems, fruits, leaves, and roots originating in transgenic plants or their progeny previously transformed with a DNA molecule of the invention and therefore consisting at least in part of transgenic cells, are also an aspect of the present invention.

[0051] As used herein, the term “plant cell” includes, without limitation, seeds, suspension cultures, embryos, meristematic regions, callus tissue, leaves, roots, shoots, gametophytes, sporophytes, pollen, and microspores. The class of plants that can be used in the methods of the invention is generally as broad as the class of higher plants amenable to transformation techniques, including both monocotyledonous and dicotyledonous plants.

[0052] “Transformation” refers to the transfer of a nucleic acid fragment into the genome of a host organism, resulting in genetically stable inheritance. Host organisms containing the transformed nucleic acid fragments are referred to as “transgenic” organisms. Examples of methods of plant transformation include *Agrobacterium*-mediated transformation (De Blaere et al. (1987) *Meth. Enzymol.* 143:277) and particle-accelerated or “gene gun” transformation technology (Klein et al. (1987) *Nature* (London) 327:70-73; U.S. Pat. No. 4,945,050, incorporated herein by reference). Additional transformation methods are disclosed below.

[0053] Thus, isolated polynucleotides of the present invention can be incorporated into recombinant constructs, typically DNA constructs, which are capable of introduction into and replication in a host cell. Such a construct can be a vector that includes a replication system and sequences that are capable of transcription and translation of a polypeptide-encoding sequence in a given host cell. A number of vectors suitable for stable transfection of plant cells or for the establishment of transgenic plants have been described in, e.g., Pouwels et al., (1985; Supp. 1987) *Cloning Vectors: A Laboratory Manual*, Weissbach and Weissbach (1989) *Methods for Plant Molecular Biology*, (Academic Press, New York); and Flevin et al., (1990) *Plant Molecular Biology Manual*, (Kluwer Academic Publishers). Typically, plant expression vectors include, for example, one or more cloned plant genes under the transcriptional control of 5' and 3' regulatory sequences and a dominant selectable marker. Such plant expression vectors also can contain a promoter regulatory region (e.g., a regulatory region controlling inducible or constitutive, environmentally- or developmentally-regulated, or cell- or tissue-specific expression), a transcription initiation start site, a ribosome binding site, an RNA processing signal, a transcription termination site, and/or a polyadenylation signal.

[0054] It is also to be understood that two different transgenic plants can also be mated to produce offspring that contain two independently segregating added, exogenous genes. Selfing of appropriate progeny can produce plants that are homozygous for both added, exogenous genes. Back-crossing to a parental plant and out-crossing with a non-transgenic plant are also contemplated, as is vegetative propagation. Descriptions of other breeding methods that are commonly used for different traits and crops can be found in one of several references, e.g., Fehr, in *Breeding Methods for Cultivar Development*, Wilcos J. ed., American Society of Agronomy, Madison Wis. (1987).

[0055] A “probe” is an isolated nucleic acid to which is attached a conventional detectable label or reporter molecule, e.g., a radioactive isotope, ligand, chemiluminescent agent, or enzyme. Such a probe is complementary to a strand of a target nucleic acid, in the case of the present invention, to a strand of isolated DNA from corn event TC1507 whether from a corn plant or from a sample that includes

DNA from the event. Probes according to the present invention include not only deoxyribonucleic or ribonucleic acids but also polyamides and other probe materials that bind specifically to a target DNA sequence and can be used to detect the presence of that target DNA sequence.

[0056] “Primers” are isolated nucleic acids that are annealed to a complementary target DNA strand by nucleic acid hybridization to form a hybrid between the primer and the target DNA strand, then extended along the target DNA strand by a polymerase, e.g., a DNA polymerase. Primer pairs of the present invention refer to their use for amplification of a target nucleic acid sequence, e.g., by the polymerase chain reaction (PCR) or other conventional nucleic acid amplification methods. “PCR” or “polymerase chain reaction” is a technique used for the amplification of specific DNA segments (see, U.S. Pat. Nos. 4,683,195 and 4,800,159; herein incorporated by reference).

[0057] Probes and primers are of sufficient nucleotide length to bind to the target DNA sequence specifically in the hybridization conditions or reaction conditions determined by the operator. This length may be of any length that is of sufficient length to be useful in a detection method of choice. Generally, 11 nucleotides or more in length, preferably 18 nucleotides or more, and more preferably 22 nucleotides or more, are used. Such probes and primers hybridize specifically to a target sequence under high stringency hybridization conditions. Preferably, probes and primers according to the present invention have complete DNA sequence similarity of contiguous nucleotides with the target sequence, although probes differing from the target DNA sequence and that retain the ability to hybridize to target DNA sequences may be designed by conventional methods. Probes can be used as primers, but are generally designed to bind to the target DNA or RNA and not be used in an amplification process.

[0058] Specific primers can be used to amplify an integration fragment to produce an amplicon that can be used as a “specific probe” for identifying event TC1507 in biological samples. When the probe is hybridized with the nucleic acids of a biological sample under conditions which allow for the binding of the probe to the sample, this binding can be detected and thus allow for an indication of the presence of event TC1507 in the biological sample. Such identification of a bound probe has been described in the art. The specific probe is preferably a sequence which, under optimized conditions, hybridizes specifically to a region within the 5' or 3' flanking region of the event and preferably also comprises a part of the foreign DNA contiguous therewith. Preferably the specific probe comprises a sequence of at least 80%, preferably between 80 and 85%, more preferably between 85 and 90%, especially preferably between 90 and 95%, and most preferably between 95 and 100% identical (or complementary) to a specific region of the event.

[0059] Methods for preparing and using probes and primers are described, for example, in *Molecular Cloning: A Laboratory Manual*, 2nd ed., vol. 1-3, ed. Sambrook et al., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. 1989 (hereinafter, “Sambrook et al., 1989”); *Current Protocols in Molecular Biology*, ed. Ausubel et al., Greene Publishing and Wiley-Interscience, New York, 1992 (with periodic updates) (hereinafter, “Ausubel et al., 1992”); and Innis et al., *PCR Protocols: A Guide to Methods and*

Applications, Academic Press: San Diego, 1990. PCR primer pairs can be derived from a known sequence, for example, by using computer programs intended for that purpose such as the PCR primer analysis tool in Vector NTI version 6 (Informax Inc., Bethesda Md.); PrimerSelect (DNASTAR Inc., Madison, Wis.); and Primer (Version 0.5, © 1991, Whitehead Institute for Biomedical Research, Cambridge, Mass.). Additionally, the sequence can be visually scanned and primers manually identified using guidelines known to one of skill in the art.

[0060] A “kit” as used herein refers to a set of reagents for the purpose of performing the method of the invention, more particularly, the identification of the event TC1507 in biological samples. The kit of the invention can be used, and its components can be specifically adjusted, for purposes of quality control (e.g. purity of seed lots), detection of event TC1507 in plant material, or material comprising or derived from plant material, such as but not limited to food or feed products. “Plant material” as used herein refers to material which is obtained or derived from a plant.

[0061] Primers and probes based on the flanking DNA and insert sequences disclosed herein can be used to confirm (and, if necessary, to correct) the disclosed sequences by conventional methods, e.g., by re-cloning and sequencing such sequences. The nucleic acid probes and primers of the present invention hybridize under stringent conditions to a target DNA sequence. Any conventional nucleic acid hybridization or amplification method can be used to identify the presence of DNA from a transgenic event in a sample. Nucleic acid molecules or fragments thereof are capable of specifically hybridizing to other nucleic acid molecules under certain circumstances. As used herein, two nucleic acid molecules are said to be capable of specifically hybridizing to one another if the two molecules are capable of forming an anti-parallel, double-stranded nucleic acid structure.

[0062] A nucleic acid molecule is said to be the “complement” of another nucleic acid molecule if they exhibit complete complementarity. As used herein, molecules are said to exhibit “complete complementarity” when every nucleotide of one of the molecules is complementary to a nucleotide of the other. Two molecules are said to be “minimally complementary” if they can hybridize to one another with sufficient stability to permit them to remain annealed to one another under at least conventional “low-stringency” conditions. Similarly, the molecules are said to be “complementary” if they can hybridize to one another with sufficient stability to permit them to remain annealed to one another under conventional “high-stringency” conditions. Conventional stringency conditions are described by Sambrook et al., 1989, and by Haymes et al., In: *Nucleic Acid Hybridization, a Practical Approach*, IRL Press, Washington, D.C. (1985), departures from complete complementarity are therefore permissible, as long as such departures do not completely preclude the capacity of the molecules to form a double-stranded structure. In order for a nucleic acid molecule to serve as a primer or probe it need only be sufficiently complementary in sequence to be able to form a stable double-stranded structure under the particular solvent and salt concentrations employed.

[0063] In hybridization reactions, specificity is typically the function of post-hybridization washes, the critical factors

being the ionic strength and temperature of the final wash solution. The thermal melting point (T_m) is the temperature (under defined ionic strength and pH) at which 50% of a complementary target sequence hybridizes to a perfectly matched probe. For DNA-DNA hybrids, the T_m can be approximated from the equation of Meinkoth and Wahl (1984) *Anal. Biochem.* 138:267-284: $T_m = 81.5^\circ \text{C.} + 16.6 (\log M) + 0.41 (\% \text{ GC}) - 0.61 (\% \text{ form}) - 500/L$; where M is the molarity of monovalent cations, % GC is the percentage of guanosine and cytosine nucleotides in the DNA, % form is the percentage of formamide in the hybridization solution, and L is the length of the hybrid in base pairs. T_m is reduced by about 1°C. for each 1% of mismatching; thus, T_m , hybridization, and/or wash conditions can be adjusted to hybridize to sequences of the desired identity. For example, if sequences with >90% identity are sought, the T_m can be decreased 10°C. Generally, stringent conditions are selected to be about 5°C. lower than the T_m for the specific sequence and its complement at a defined ionic strength and pH. However, severely stringent conditions can utilize a hybridization and/or wash at 1, 2, 3, or 4°C. lower than the T_m ; moderately stringent conditions can utilize a hybridization and/or wash at 6, 7, 8, 9, or 10°C. lower than the T_m ; low stringency conditions can utilize a hybridization and/or wash at 11, 12, 13, 14, 15, or 20°C. lower than the T_m .

[0064] Using the equation, hybridization and wash compositions, and desired T_m , those of ordinary skill will understand that variations in the stringency of hybridization and/or wash solutions are inherently described. If the desired degree of mismatching results in a T_m of less than 45°C. (aqueous solution) or 32°C. (formamide solution), it is preferred to increase the SSC concentration so that a higher temperature can be used. An extensive guide to the hybridization of nucleic acids is found in Tijssen (1993) *Laboratory Techniques in Biochemistry and Molecular Biology—Hybridization with Nucleic Acid Probes*, Part I, Chapter 2 (Elsevier, New York); and Ausubel et al., eds. (1995) *Current Protocols in Molecular Biology*, Chapter 2 (Greene Publishing and Wiley-Interscience, New York). See Sambrook et al. (1989) *Molecular Cloning: A Laboratory Manual* (2d ed., Cold Spring Harbor Laboratory Press, Plainview, N.Y.).

[0065] As used herein, a substantially homologous sequence is a nucleic acid molecule that will specifically hybridize to the complement of the nucleic acid molecule to which it is being compared under high stringency conditions. Appropriate stringency conditions which promote DNA hybridization, for example, 6× sodium chloride/sodium citrate (SSC) at about 45°C. , followed by a wash of 2×SSC at 50°C. , are known to those skilled in the art or can be found in *Current Protocols in Molecular Biology*, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6. Typically, stringent conditions will be those in which the salt concentration is less than about 1.5 M Na ion, typically about 0.01 to 1.0 M Na ion concentration (or other salts) at pH 7.0 to 8.3 and the temperature is at least about 30°C. for short probes (e.g., 10 to 50 nucleotides) and at least about 60°C. for long probes (e.g., greater than 50 nucleotides). Stringent conditions may also be achieved with the addition of a destabilizing agent such as formamide. Exemplary low stringency conditions include hybridization with a buffer solution of 30 to 35% formamide, 1 M NaCl, 1% SDS (sodium dodecyl sulphate) at 37°C. , and a wash in 1× to 2×SSC (20×SSC=3.0 M NaCl/0.3 M trisodium citrate) at 50 to 55°C. Exemplary

moderate stringency conditions include hybridization in 40 to 45% formamide, 1 M NaCl, 1% SDS at 37° C., and a wash in 0.5× to 1×SSC at 55 to 60° C. Exemplary high stringency conditions include hybridization in 50% formamide, 1 M NaCl, 1% SDS at 37° C., and a wash in 0.1×SSC at 60 to 65° C. In a preferred embodiment, a nucleic acid of the present invention will specifically hybridize to one or more of the nucleic acid molecules unique to the TC1507 event or complements thereof or fragments of either under moderately stringent conditions.

[0066] Methods of alignment of sequences for comparison are well known in the art. Thus, the determination of percent identity between any two sequences can be accomplished using a mathematical algorithm. Non-limiting examples of such mathematical algorithms are the algorithm of Myers and Miller (1988) *CABIOS* 4:11-17; the local homology algorithm of Smith et al. (1981) *Adv. Appl. Math.* 2:482; the homology alignment algorithm of Needleman and Wunsch (1970) *J. Mol. Biol.* 48:443-453; the search-for-similarity-method of Pearson and Lipman (1988) *Proc. Natl. Acad. Sci.* 85:2444-2448; the algorithm of Karlin and Altschul (1990) *Proc. Natl. Acad. Sci. USA* 87:2264, modified as in Karlin and Altschul (1993) *Proc. Natl. Acad. Sci. USA* 90:5873-5877.

[0067] Computer implementations of these mathematical algorithms can be utilized for comparison of sequences to determine sequence identity. Such implementations include, but are not limited to: CLUSTAL in the PC/Gene program (available from Intelligenetics, Mountain View, Calif.); the ALIGN program (Version 2.0); the ALIGN PLUS program (version 3.0, copyright 1997); and GAP, BESTFIT, BLAST, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Version 10 (available from Accelrys, 9685 Scranton Road, San Diego, Calif. 92121, USA). Alignments using these programs can be performed using the default parameters.

[0068] The CLUSTAL program is well described by Higgins and Sharp, *Gene* 73: 237-244 (1988); Higgins and Sharp, *CABIOS* 5: 151-153 (1989); Corpet, et al., *Nucleic Acids Research* 16: 10881-90 (1988); Huang, et al., *Computer Applications in the Biosciences* 8: 155-65 (1992), and Pearson, et al., *Methods in Molecular Biology* 24: 307-331 (1994). The ALIGN and the ALIGN PLUS programs are based on the algorithm of Myers and Miller (1988) supra. The BLAST programs of Altschul et al. (1990) *J. Mol. Biol.* 215:403 are based on the algorithm of Karlin and Altschul (1990) supra. The BLAST family of programs which can be used for database similarity searches includes: BLASTN for nucleotide query sequences against nucleotide database sequences; BLASTX for nucleotide query sequences against protein database sequences; BLASTP for protein query sequences against protein database sequences; TBLASTN for protein query sequences against nucleotide database sequences; and TBLASTX for nucleotide query sequences against nucleotide database sequences. See, *Current Protocols in Molecular Biology*, Chapter 19, Ausubel, et al., Eds., Greene Publishing and Wiley-Interscience, New York (1995). Alignment may also be performed manually by inspection.

[0069] To obtain gapped alignments for comparison purposes, Gapped BLAST (in BLAST 2.0) can be utilized as described in Altschul et al. (1997) *Nucleic Acids Res.*

25:3389. Alternatively, PSI-BLAST (in BLAST 2.0) can be used to perform an iterated search that detects distant relationships between molecules. See Altschul et al. (1997) supra. When utilizing BLAST, Gapped BLAST, PSI-BLAST, the default parameters of the respective programs (e.g., BLASTN for nucleotide sequences, BLASTX for proteins) can be used. See www.ncbi.nlm.nih.gov.

[0070] As used herein, "sequence identity" or "identity" in the context of two nucleic acid or polypeptide sequences makes reference to the residues in the two sequences that are the same when aligned for maximum correspondence over a specified comparison window. When percentage of sequence identity is used in reference to proteins it is recognized that residue positions which are not identical often differ by conservative amino acid substitutions, where amino acid residues are substituted for other amino acid residues with similar chemical properties (e.g., charge or hydrophobicity) and therefore do not change the functional properties of the molecule. When sequences differ in conservative substitutions, the percent sequence identity may be adjusted upwards to correct for the conservative nature of the substitution. Sequences that differ by such conservative substitutions are said to have "sequence similarity" or "similarity". Means for making this adjustment are well known to those of skill in the art. Typically this involves scoring a conservative substitution as a partial rather than a full mismatch, thereby increasing the percentage sequence identity. Thus, for example, where an identical amino acid is given a score of 1 and a non-conservative substitution is given a score of zero, a conservative substitution is given a score between zero and 1. The scoring of conservative substitutions is calculated, e.g., as implemented in the program PC/GENE (Intelligenetics, Mountain View, Calif.).

[0071] As used herein, "percentage of sequence identity" means the value determined by comparing two optimally aligned sequences over a comparison window, wherein the portion of the polynucleotide sequence in the comparison window may comprise additions or deletions (i.e., gaps) as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. The percentage is calculated by determining the number of positions at which the identical nucleic acid base or amino acid residue occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison, and multiplying the result by 100 to yield the percentage of sequence identity.

[0072] Regarding the amplification of a target nucleic acid sequence (e.g., by PCR) using a particular amplification primer pair, "stringent conditions" are conditions that permit the primer pair to hybridize only to the target nucleic-acid sequence to which a primer having the corresponding wild-type sequence (or its complement) would bind and preferably to produce a unique amplification product, the amplicon, in a DNA thermal amplification reaction.

[0073] The term "specific for (a target sequence)" indicates that a probe or primer hybridizes under stringent hybridization conditions only to the target sequence in a sample comprising the target sequence.

[0074] As used herein, "amplified DNA" or "amplicon" refers to the product of nucleic acid amplification of a target nucleic acid sequence that is part of a nucleic acid template.

For example, to determine whether a corn plant resulting from a sexual cross contains transgenic event genomic DNA from the corn plant of the present invention, DNA extracted from the corn plant tissue sample may be subjected to a nucleic acid amplification method using a DNA primer pair that includes a first primer derived from flanking sequence adjacent to the insertion site of inserted heterologous DNA, and a second primer derived from the inserted heterologous DNA to produce an amplicon that is diagnostic for the presence of the event DNA. Alternatively, the second primer may be derived from the flanking sequence. The amplicon is of a length and has a sequence that is also diagnostic for the event. The amplicon may range in length from the combined length of the primer pairs plus one nucleotide base pair to any length of amplicon producible by a DNA amplification protocol. Alternatively, primer pairs can be derived from flanking sequence on both sides of the inserted DNA so as to produce an amplicon that includes the entire insert nucleotide sequence of the PH18999A expression construct, see **FIG. 1**, approximately 6.2 Kb in size. A member of a primer pair derived from the flanking sequence may be located a distance from the inserted DNA sequence, this distance can range from one nucleotide base pair up to the limits of the amplification reaction, or about twenty thousand nucleotide base pairs. The use of the term "amplicon" specifically excludes primer dimers that may be formed in the DNA thermal amplification reaction.

[0075] Nucleic acid amplification can be accomplished by any of the various nucleic acid amplification methods known in the art, including the polymerase chain reaction (PCR). A variety of amplification methods are known in the art and are described, inter alia, in U.S. Pat. Nos. 4,683,195 and 4,683,202 and in *PCR Protocols: A Guide to Methods and Applications*, ed. Innis et al., Academic press, San Diego, 1990. PCR amplification methods have been developed to amplify up to 22 Kb of genomic DNA and up to 42 Kb of bacteriophage DNA (Cheng et al., *Proc. Natl. Acad. Sci. USA* 91:5695-5699, 1994). These methods as well as other methods known in the art of DNA amplification may be used in the practice of the present invention. It is understood that a number of parameters in a specific PCR protocol may need to be adjusted to specific laboratory conditions and may be slightly modified and yet allow for the collection of similar results. These adjustments will be apparent to a person skilled in the art.

[0076] The amplicon produced by these methods may be detected by a plurality of techniques, including, but not limited to, Genetic Bit Analysis (Nikiforov, et al. *Nucleic Acid Res.* 22:4167-4175, 1994) where a DNA oligonucleotide is designed which overlaps both the adjacent flanking DNA sequence and the inserted DNA sequence. The oligonucleotide is immobilized in wells of a microwell plate. Following PCR of the region of interest (using one primer in the inserted sequence and one in the adjacent flanking sequence) a single-stranded PCR product can be hybridized to the immobilized oligonucleotide and serve as a template for a single base extension reaction using a DNA polymerase and labeled ddNTPs specific for the expected next base. Readout may be fluorescent or ELISA-based. A signal indicates presence of the insert/flanking sequence due to successful amplification, hybridization, and single base extension.

[0077] Another detection method is the Pyrosequencing technique as described by Winge (*Innov. Pharma. Tech.* 00: 18-24, 2000). In this method an oligonucleotide is designed that overlaps the adjacent DNA and insert DNA junction. The oligonucleotide is hybridized to a single-stranded PCR product from the region of interest (one primer in the inserted sequence and one in the flanking sequence) and incubated in the presence of a DNA polymerase, ATP, sulfurylase, luciferase, apyrase, adenosine 5' phosphosulfate and luciferin. dNTPs are added individually and the incorporation results in a light signal which is measured. A light signal indicates the presence of the transgene insert/flanking sequence due to successful amplification, hybridization, and single or multi-base extension.

[0078] Fluorescence Polarization as described by Chen et al., (*Genome Res.* 9:492-498, 1999) is a method that can be used to detect an amplicon of the present invention. Using this method an oligonucleotide is designed which overlaps the flanking and inserted DNA junction. The oligonucleotide is hybridized to a single-stranded PCR product from the region of interest (one primer in the inserted DNA and one in the flanking DNA sequence) and incubated in the presence of a DNA polymerase and a fluorescent-labeled ddNTP. Single base extension results in incorporation of the ddNTP. Incorporation can be measured as a change in polarization using a fluorometer. A change in polarization indicates the presence of the transgene insert/flanking sequence due to successful amplification, hybridization, and single base extension.

[0079] Taqman® (PE Applied Biosystems, Foster City, Calif.) is described as a method of detecting and quantifying the presence of a DNA sequence and is fully understood in the instructions provided by the manufacturer. Briefly, a FRET oligonucleotide probe is designed which overlaps the flanking and insert DNA junction. The FRET probe and PCR primers (one primer in the insert DNA sequence and one in the flanking genomic sequence) are cycled in the presence of a thermostable polymerase and dNTPs. Hybridization of the FRET probe results in cleavage and release of the fluorescent moiety away from the quenching moiety on the FRET probe. A fluorescent signal indicates the presence of the flanking/transgene insert sequence due to successful amplification and hybridization.

[0080] Molecular Beacons have been described for use in sequence detection as described in Tyangi et al. (*Nature Biotech.* 14:303-308, 1996). Briefly, a FRET oligonucleotide probe is designed that overlaps the flanking and insert DNA junction. The unique structure of the FRET probe results in it containing secondary structure that keeps the fluorescent and quenching moieties in close proximity. The FRET probe and PCR primers (one primer in the insert DNA sequence and one in the flanking sequence) are cycled in the presence of a thermostable polymerase and dNTPs. Following successful PCR amplification, hybridization of the FRET probe to the target sequence results in the removal of the probe secondary structure and spatial separation of the fluorescent and quenching moieties. A fluorescent signal results. A fluorescent signal indicates the presence of the flanking/transgene insert sequence due to successful amplification and hybridization.

[0081] A hybridization reaction using a probe specific to a sequence found within the amplicon is yet another method used to detect the amplicon produced by a PCR reaction.

[0082] The present invention is further defined in the following Examples. It should be understood that these Examples, while indicating preferred embodiments of the invention, are given by way of illustration only. From the above discussion and these Examples, one skilled in the art can ascertain the essential characteristics of this invention, and without departing from the spirit and scope thereof, can make various changes and modifications of the invention to adapt it to various usages and conditions. Thus, various modifications of the invention, in addition to those shown and described herein, will be apparent to those skilled in the art from the foregoing description. Such modifications are also intended to fall within the scope of the appended claims.

[0083] The disclosure of each reference set forth herein is incorporated herein by reference in its entirety.

EXAMPLES

Example 1

Transformation of Maize by Particle Bombardment and Regeneration of Transgenic Plants Containing the Cry1F Gene

[0084] A DNA molecule of 6.2 Kb, designated PH18999A (see FIG. 1 and SEQ ID NO:25), which includes a first transgene expression cassette comprising the promoter, 5' untranslated exon, and first intron of the maize ubiquitin (Ubi-1) gene (Christensen et al. (1992) *Plant Mol. Biol.* 18:675-689 and Christensen and Quail (1996) *Transgenic Res.* 5:213-218) operably connected to a DNA molecule encoding a *Bacillus thuringiensis* 6-endotoxin identified as Cry1F (U.S. Pat. Nos. 5,188,960 and 6,218,188) operably connected to a DNA molecule comprising a 3' ORF25 transcriptional terminator isolated from *Agrobacterium tumefaciens* (Barker et al. (1983) *Plant Mol. Biol.* 2:335-350), and a second transgene expression cassette comprising a DNA molecule of the cauliflower mosaic virus (CaMV) ³⁵S promoter (Odell J. T. et al. (1985) *Nature* 313: 810-812; Mitsuhashi et al. (1996) *Plant Cell Physiol.* 37:49-59) operably connected to a DNA molecule encoding the selectable marker, phosphinothricin acetyltransferase (PAT) gene (Wohlleben W. et al. (1988) *Gene* 70:25-37) operably connected to a DNA molecule comprising a 3' transcriptional terminator from (CaMV) ³⁵S (see Mitsuhashi et al. (1996) *Plant Cell Physiol.* 37:49-59) was used to transform maize embryo tissue.

[0085] *B.t.* Cry1F maize plants were obtained by micro-projectile bombardment using the Biolistics® PDS-1000He particle gun manufactured by Bio-Rad, Hercules, Calif.; essentially as described by Klein et al. (1987) *Nature*, UK 327(6117):70-73. Immature embryos isolated from maize ears, harvested soon after pollination were cultured on callus initiation medium for several days. On the day of transformation, microscopic tungsten particles were coated with purified PH18999A DNA (SEQ ID NO:25) and accelerated into the cultured embryos, where the insert DNA was incorporated into the cell chromosome. Only insert PH18999A was used during transformation and no additional plasmid DNA was incorporated into the transformant. After bombardment, embryos were transferred to callus initiation medium containing glufosinate as the selection agent. Individual embryos were kept physically separate during culture, and the majority of explants died on the selective medium.

[0086] Those embryos that survived and produced healthy, glufosinate-resistant callus tissue were assigned unique identification codes representing putative transformation events, and continually transferred to fresh selection medium. Plants were regenerated from tissue derived from each unique event and transferred to the greenhouse. Leaf samples were taken for molecular analysis to verify the presence of the transgene by PCR and to confirm expression of the Cry1F protein by ELISA. Plants were then subjected to a whole plant bioassay using European corn borer insects. Positive plants were crossed with inbred lines to obtain seed from the initial transformed plants. A number of lines were evaluated in the field. The TC1507 event was selected from a population of independent transgenic events based on a superior combination of characteristics, including insect resistance and agronomic performance (see Bing J W et al. (2000) Efficacy of Cry1F Transgenic Maize, 14th Biennial International Plant Resistance to Insects Workshop, Fort Collins, Colo., herein incorporated by reference).

Example 2

Identification of Nucleotides Comprising the Flanking Sequence 5' to the Transgenic Insert DNA in *Bacillus thuringiensis* Cry1F Maize Line TC1507

[0087] To identify a DNA fragment that included sequence 5' to the PH18999A insert in event TC1507, Spe I restriction enzyme fragments from event TC1507 genomic DNA were size selected on agarose gels, purified, and screened by Southern analysis to confirm hybridization to a Cry1F probe. Following confirmation of hybridization and fragment size, the fragments of interest were cloned into a pBluescript II SK (+)TM cloning vector to prepare an enriched size selected plasmid based genomic DNA library. A probe homologous to a portion of the Cry1F gene was used to screen the plasmid library for positive clones. A positive clone was identified, purified by additional screening, and confirmed to result in a positive signal when hybridized to the Cry1F probe. Nearly 3 Kb of the Spe I fragment contained in the isolated positive clone was sequenced using a primer walking approach. To initiate the first sequencing run, a primer that binds to a known sequence in the cloning vector DNA was designed to sequence a portion of the DNA of interest. A second sequencing run over the same region using another primer oriented in the reverse direction provided second strand coverage. Primer walking was accomplished by repeatedly using sequence data from previous runs to design new primers that were then used to extend the next round of sequencing further into the DNA of interest until the flanking sequence 5' to the inserted transgenic DNA in maize event TC1507 was obtained. Specific sequence information is provided in Example 4.

Example 3

Confirmation of Flanking Sequence 5' to the *B.t.* Cry1F Maize Line TC1507 Insert

[0088] To confirm the 5' flanking sequence of the *B.t.* Cry1F maize line TC1507 insert, PCR primer pairs were designed to obtain overlapping PCR products extending from the 5' flanking region into the full-length PH18999A transgenic insert. PCR products were successfully amplified from *B.t.* Cry1F maize line TC1507 genomic DNA, isolated,

and sequenced for Region 1 through Region 6, shown in Table 1, and confirmed to match the previously determined sequence from the Spe I fragment, described in Example 2. However, the region from bp 2358 to bp 2829, immediately adjacent and 5' to the start of the full-length insert was recalcitrant to PCR amplification and appeared to be larger than the sequence obtained from the Spe I clone described above. The use of primer pairs flanking this region and the Advantage®-GC 2 Polymerase Mix (BD Biosciences Clontech, Palo Alto, Calif.) was successful in amplifying PCR products from *B.t.* Cry1F maize line TC1507 genomic DNA for sequencing. The amplification conditions used to pro-

duce amplicons with the Advantage®-GC 2 system are shown in Table 10. The DNA primer pairs used to confirm the sequence in the region from bp 2358 to 2829 are those listed in SEQ ID NO: 1 and SEQ ID NO:2; and SEQ ID NO:2 and SEQ ID NO:23. Sequence from this region is described in Table 1 (Regions 7a, 7b, 7c, and 8).

Example 4

Event TC1507 5' Flanking Sequence.

[0089] A description of each region is provided in Table 1.

Region 1 (SEQ ID NO: 28) Maize genomic (no significant homology)

```

1  ACTAGTTTCC TAGCCCGCGT CGTGCCCTTA CCCCACCGAC GTTTATGGAA
51  GGTGCCATTC CACGGTTCTT CGTGGCCGCC CCTAAGGATG TAAATGGTCG
101 GTAAAATCCG GTAAATTTCG GGTACCGTTT ACCAGATTTT TCCAGCCGTT
151 TTCGGATTTA TCGGGATATA CAGAAAACGA GACGGAAACG GAATAGGTTT
201 TTTTTCGAAA ACGGTACGGT AAACGGTGAG ACAAACCTAC CGTCCGTTTT
251 CGTATTCTTC GGGAAACTCT GGTATATTCC CGTATTGTGC CCGTATTTTC
301 CCGACCCACG GACCTGCCAA TCAACCATCA GCCAGTCAGC CCATCCCCAC
351 AGCTATGGCC CATGGGGCCA TGTGGCCAC ATGCCACGC AACGCAAGGC
401 AGTAAGGCTG GCAGCCTGGC ACGCATTGAC GCATGTGGAC ACACACAGCC
451 GCCGCCTGTT CGTGTCTCTG TGCCGTTGTG CGAGACTGTG ACTGCGAGTG
501 GCGGAGTCGG CGAACGGCGA GCGCTCTCCG GAGTCTGGAC TCGGGCTGTG
551 GACAGCGACG CTGTGACGGC GACTCGGCGA AGCCCCAAGC TACCAAGCCC
601 CCAAGTCCCC ATCCATCTCT GCTTCTCTGG TCATCTCCTT CCCCTGGTCG
651 ATCTGCAGGC GCCAGACCG

```

Region 2 (SEQ ID NO: 29) Undescribed maize genomic sequence (complement)

```

670 G CCGAAGCATC ACGAAACGCA CTAAGACCTC
701 GAAGGAGTCA AACCACTCCT CCGAGGCCTC GGGGGCTACA CCCGGCGGGT
751 GCGCTCGCGC GCACCCACCG GAACAAAATG TAACCGAGAA AGGTGCGTCC
801 CCTTGCAAAA AAAGTGCGAC AAAAGCCTCC AAGCGAGTAT TAACACTCAC
851 TTTGAGGCTC GGGGGCTAC

```

Region 3 (SEQ ID NO: 30) Fragment of maize Huck-1 retrotransposon

```

870 T GTCGGGACG ATAATTAGGG GTACCCCAAA
901 GACTCCTAAT CTCAGCTGGT AACCCCATC AGCACAAAGC TGCAAAGGCC
951 TGATGGGTGC GATTAAGTCA AGGCTCGGTC CACTCAAGGG ACACGATCTC
1001 GCCTCGCCCG AGCCCAGCCT CGGGCAAGGG CGGCCGACCC CGAGGATTCA
1051 CGTCTCGCCC GAGGGCCCC TCAAGCGACG GGCACACCTT CGGCTCGCCC
1101 GAGGCCCATC CTTGCGCGAG AAGCAACCTT GGCCAGATCG CCACACCGAC
1151 CGACCGTATC GCAGGAGCAT TTAATGCGAG GATCGCCTGA CACCTTATCC
1201 TGACGCGCGC TCTTCAGTCG ACAGAGCCGA AGTGACCGCA ATCACTTCGC

```

-continued

1251 CGCTCCACTG ACCGACCTGA CAAGAAGACA GCGCCGCCTG CGTCGCTCCG
 1301 ACTGCTGTGC CACTCGACAG AGTGAGGCTG ACAGCAGCCA AGTCCGGCCT
 1351 CGGGCGCCAT AGGAAGCTCC GCCTCGCCCG ACCCTAGGGC TCGGACTCGG
 1401 CCTCGGCTCC GGAAGACGAC GAACTACGCT TCGCCCGACC CCAGGGCTTG
 1451 GACTCAGCCT CGGCTCCGGA AGACGACGAA TTCCGCCTCG CCCGACCCCA
 1501 GGGCTCGGAC TCGGCCTCGG CTCCAGAAGA CGACGAACTC CGCCTCGCCC
 1551 GACCCAGGG CTCGGACTCA GCCTCGGCTC CGGAAGACGA CGAACTCCGC
 1601 CTCGCCCGAC CCCAGGGCTC GGAATCAGCC TCGGCCTCAG ACGATGGTCT
 1651 CCGCCTCGCC CGACCCGGGG CTCGGACTCG A

Region 4 (SEQ ID NO: 31) Fragment of *cry1F* gene
 1682 CCTTTCTAT CGGACCTTGT

1701 CAGATCCTGT CTTCGTCCGA GGAGGCTTTG GCAATCCTCA CTATGTACTC
 1751 GGTCTTAGGG GAGTGGCCTT TCAACAAACT GGTACGAATC ACACCCGCAC
 1801 ATTCAGGAAC TCCGGGACCA TTGACTCTCT AGATGAGATA CCACCTCAAG
 1851 ACAACAGCGG CGCACCTTGG AATGACTACT CCCATGTGCT GAATCATGTT
 1901 ACCTTTGTGC GCTGGCCAGG TGAGATCTCA GGTTCGCACT CATGGAGAGC
 1951 ACCAATGTTT TCTTGACGC ATCGTAGCGC TACCCCCACA AACACCATTG
 2001 ATCCAGAGAG AATCAC

Region 5 (SEQ ID NO: 32) Fragment of maize chloroplast
rpoC2 gene

2017 TCAT TCTTCAAGAA CTGCATATCT TGCCGAGATC
 2051 CTCATCCCTA AAGGTACTTG ACAATAGTAT TATGGAGTC GATACACAAC
 2101 TCACAAAAAA TACAAGAAGT CGACTAGGTG GATTGGTCCG AGTGAAGAGA
 2151 AAAAAAAGCC ATACAGAACT CAAAATCTTT TCCGGAGATA TTCATTTTCC
 2201 TGAAGAGGCG GATAAGATAT TAGGTGGCAG TTTGATACCA CCAGAAAGAG
 2251 AAAAAAAGA TTCTAAGGAA TCAAAAAAAA GGAAAAATTG GGTTTATGTT
 2301 CAACGAAAA AATTTCTCAA AAGCAAGGAA AAGTATT

Region 6 (SEQ ID NO: 33) Fragment of maize chloroplast
 or *ubizM1(2)* promoter

2338 GTG GCTATTTATC

2351 TATC

Nucleotides 2355-2358 (CGT) connect Region 6 to Region 7a.

Region 7a (SEQ ID NO: 34) Fragment of *pat* gene

2358 GCA GCTGATATGG CCGCGGTTG TGATATCGTT AACCATTACA
 2401 TTGAGACGTC TACAGTGAAC TTTAGGACAG AGCCACAAAC ACCACAAGAG
 2451 TGGATTGATG ATCTAGAGAG GTTGCAAGAT AGATACCCTT GGTGTTGTC
 2501 TGAGGTTGAG GGTTGTTGG CTGGTATTGC TTACGCTGGG CCCTGGAAGG
 2551 CTAGGAAC

Region 7b (SEQ ID NO: 35) Fragment of *pat* gene (complement)
 (complement)

2559 CC TCAACCTCAG CAACCAACCA ATGGTATCTA TCTTGCAACC
 2601 TCTCTAGATC ATCAATCCAC TCTTGTGGTG TTTGTGGCTC TGTCTAAAG
 2651 TTCCTGTAG ACGTCTCAAT GTAATGGTTA ACGATATCAC AAACCG

-continued

Region 7c (SEQ ID NO: 36) Fragment of cry1F gene
(complement)
2697 AGAG

2701 AAGAGGGATC T

Region 8 (SEQ ID NO: 37) Fragment of Polylinker
2712 CGAAGCTTC GGCCGGGGCC CATCGATATC CGCGGGCATG

2751 CCTGCAGTGC AGCGTGACCC GGTCGTGCCC CTCTCTAGAG ATAATGAGCA

2801 TTGCATGTCT AAGTTATAAA AAATTACCA

Region 9 (SEQ ID NO: 25) Full-length insert of PHI8999A

Example 5

Description of the Flanking Sequence 5' to the
Insert in Maize Event TC1507

[0090] In order to more fully describe the event TC1507 5' flanking sequence, homology searching was done against the GenBank public databases (release 122, 2/01) using the Basic Local Alignment Search Tool (BLAST). The BLAST program performs sequence similarity searching and is particularly useful for identifying homologs to an unknown sequence. In addition to searching the public databases, pairwise alignments were performed using AlignX (InforMax Inc., Bethesda, Md.) to look for homology between the maize event TC1507 flanking sequence and the PHI8999A transgenic insert. The results of these homology searches are presented in Table 1. The TC1507 5' flanking sequence is numbered with base 1 being the furthest 5' to the insert and base 2830 at the starting point of the full-length PHI8999A transgenic insert (see **FIG. 1**). The percent identity values indicate the percentage of identical matches across the length of the sequences analyzed.

[0091] In most cases, similarity searching with the event TC1507 5' flanking sequence resulted in a match to one unique sequence based on a very high percent identity value. Those sequences are identified in Table 1. In addition, there are two regions in the TC1507 5' DNA flanking sequence with high similarity to more than one known sequence. In regions 870-1681 and 2338-2354, the percent identity scores with both sequence fragments are sufficiently high that a single match (homolog) cannot be determined. The two possible homologs for each of these regions are indicated in Table 1.

[0092] Highly similar sequences were identified for all but the first 669 base pairs of sequence. Generally, the results of similarity searching indicate high homology with maize genomic sequences 5' to base 1681. The region from base 1682 to the start of the PHI8999A insert at position 2830 contains some fragments associated with the transformation event.

TABLE 1

Sequence summary for event TC1507 insert						
Region	Location in SEQ ID NO: 24	Size bp	% Identity	Homolog	Location in homologous sequence	Description
1	1-669	669	N/A ¹	N/A	N/A	No significant homology detected
2	670-869	200	90.5	AF123535	52432-52632 (complement)	Undescribed maize genomic sequence
3	870-1681	812	89.4	AF050439	1-801	Fragment of maize Huck-1 retrotransposon 5' LTR ²
			86.6	AF050438	1-797	Fragment of maize Huck-1 retrotransposon 3' LTR
4	1682-2016	335	100.0	PHI8999A	3149-3483	Fragment of cry1F gene
5	2017-2337	321	100.0	X86563	29429-29749	Fragment of maize chloroplast rpoC2 gene (RNA polymerase beta-2 subunit)
6	2338-2354	17	100.0	X86563	97643-97659	Fragment of maize chloroplast

TABLE 1-continued

Sequence summary for event TC1507 insert						
Region	Location in SEQ ID NO: 24	Size bp	% Identity	Homolog	Location in homologous sequence	Description
			82.4	PHI8999A	182–197	trnI gene (tRNA-Ile)
7a	2358–2558	201	100.0	PHI8999A	5320–5475	Fragment of maize ubiZM1(2) promoter
7b	2559–2696	138	99	PHI8999A	5336–5518 (complement)	Fragment of pat gene
7c	2697–2711	15	100.0	PHI8999A	2544–2558 (complement)	Fragment of cry1F gene
8	2712–2829	118	100.0	PHI8999A	36–153	Fragment of polylinker region (bases 36–80) and ubiZM1(2) promoter (bases 81–153)
9	2830–9015	6186	100.0	PHI8999A	11–6196	Full-length insert of PHI8999A
10	9016–9565	550	100.0	PHI8999A	3906–4456 (complement)	Inverted ORF25 terminator
11	9566–9693	128	100.0	NC_001666	121851–121978 (complement) & 100759–100886	Fragment of maize chloroplast rps12 rRNA (23S ribosomal RNA)
12	9696–10087	392	99	NC_001666	17091–17483 (complement)	Fragment of maize chloroplast genome
13	10088–10275	188	99	PHI8999A	5333–5520 (complement)	Fragment of pat gene
14	10278–10358	81	100	NC_001666	137122–137202 (complement)	Fragment of maize chloroplast “ORF241” - hypothetical protein gene
15	10359–10612	254	N/A ¹	N/A	N/A	No significant homology detected
16	10613–11361	749	N/A ¹	N/A	N/A	No description available

¹N/A; not applicable²LTR; long terminal repeat

Example 6

Confirmation of the Presence of Regions 1, 2, and 3 in an Unmodified Control Corn Line

[0093] PCR analysis was used to determine if Regions 1, 2, and 3 (Table 1) in the 5' flanking region of Event TC1507 are present in an unmodified control corn line used for transformation to produce maize event TC1507 and thus represents a border with corn genomic DNA. Nine different PCR analyses were carried out on genomic DNA prepared from TC1507 and the unmodified control corn line Hi-II (see Armstrong (1994) *The Maize Handbook*, ed. Freeling and Walbot, Springer-Verlag, New York, pp. 663-671, for information on Hi-II) as outlined in Table 2 using the primer sequences shown in Table 3. Two reactions were designed to amplify DNA within Region 1 of the 5' flanking region from bp 25 to 324 (Reaction A-300 bp amplicon); and from bp 25 to 480 (Reaction B—456 bp amplicon). The expected amplicons were present in both the Hi-II unmodified corn line and in maize event TC1507. One PCR primer pair, Reaction C, spanned Region 2 to Region 3 of the 5' flanking region from

bp 759 to 1182 (424 bp amplicon) and again produced PCR products of the expected size in both Hi-II and TC1507. Reaction D, spanned Region 1 to Region 3 of the 5' flanking region from bp 415 to 1182 (768 bp amplicon) and again produced PCR products of the expected size in both Hi-II and TC1507. Reactions E and F were designed as specific primer pairs for the pat gene region of the full-length insert of PHI8999A in TC1507 and thus an amplicon in the unmodified Hi-II corn line is not expected. The results indicate that both Reactions E and F are specific for a maize line transformed with a pat gene region and produce the expected amplicon, whereas no amplicon was produced in the unmodified Hi-II corn line. Reaction G was also designed as a primer pair that would produce an amplicon of 366 bp in the maize event TC1507 and no amplicon in the unmodified Hi-II corn line.

[0094] Reactions H and I were designed as specific primer pairs for TC1507 that would span the end of the transgenic insert into the 5' flanking region. In both Reactions H and I, the reverse primer was located in the ubiquitin promoter

region of the full-length PH18999A insert (Region 9 in Table 1) and the forward primer was located in Region 5, the rpoC2 gene fragment (see Table 1). Reaction H and Reaction I both produced an amplicon in maize line TC1507 and did not produce an amplicon in the unmodified control corn line. These results indicate that both Reactions H and I are specific for the TC1507 event.

[0095] The PCR results show that the undescribed sequence (Region 1) is present in the unmodified corn line Hi-II and that Regions 1, 2 and 3, are contiguous in the unmodified corn line Hi-II. The DNA sequences amplified in Reactions A, B, C, and D are not unique to the 5' flanking region of maize event TC1507 but are also present in the unmodified corn line Hi-II.

TABLE 2

PCR reactions for sequence 5' to the PH18999A insert in maize event TC1507 and for regions within the full-length insert of PH18999A in maize event TC1507					
Reaction	PCR Amplicon Location	Amplicon Size (bp)	Region in TC1507 flanking sequence or PH18999A insert	Amplicon present In Hi-II	Amplicon present in maize line TC1507
A	25–324 bp in TC1507 flanking sequence	300	Region 1	Yes	Yes
B	25–480 bp in TC1507 flanking sequence bp in TC1507 flanking sequence	456	Region 1 3	Yes	Yes
D	415–1182 bp in TC1507 5' flanking sequence	768	Region 1 to Region3	Yes	Yes
E Not Unique to TC1507	4750–5794 bp in PH18999A	1045	Region 9 (in full- length insert of PH18999A 35S promoter to pat gene)	No	Yes
F Not Unique to TC1507	4827–5308 bp in PH18999A	482	Region 9 (in full- length insert of PH18999A 35S promoter to pat gene)	No	Yes
Detects cry1F fragment in 5' flanking region	cry1F sequence in 5' flanking sequence and in full-length insert of PH18999A	366	Spans 335 bp cry1F sequence in 5' flanking sequence and same sequence in the full-length insert	No	Yes
H Unique to TC1507	2158 bp in Region 5 (rpoC2 gene fragment) to 3069 bp in Region 9 (full-length insert of PH18999A	912	Region 5 to Region 9 Unique to Insertion Event [SPANS UNIQUE JUNCTION REGIONS]	No	Yes
I Unique to TC1507	2158 bp in Region 5 (rpoC2 gene fragment) to 3001 bp in Region 9 (full-length insert of PH18999A)	844	Region 5 to Region 9 Unique to Insertion Event [SPANS UNIQUE JUNCTION REGIONS]	No	Yes

[0096]

TABLE 3

PCR primers for sequence 5' to the PHI8999A insert in TC1507 and for regions within the full-length insert of PHI8999A in maize event TC1507				
Reaction	Amplicon Size (bp)	Primer Pair	Primer Sequences 5' to 3'	
A	300	SEQ ID NO:10	CCCCTACCCACCGACGTTTAT	
		SEQ ID NO:11	TTGATTGGCAGGTCCGTGGGTC	
B	456	SEQ ID NO:10	CCCCTACCCACCGACGTTTAT	
		SEQ ID NO:12	CACAAACGGCACAGAAACACGAA	
C	424	SEQ ID NO:13	GCGCACCCACCGAACAATG	
		SEQ ID NO:14	TCCTCGATTAAATGCTCCTGC	
D	768	SEQ ID NO:15	CCTGGCAGCATTCGCGCATGT	
		SEQ ID NO:14	TCCTCGATTAAATGCTCCTGC	
E	1045	SEQ ID NO:6	TAGAGGACCTAACAGAACTCGCCGT	
		SEQ ID NO:7	GAGCTGGCAACTCAAAATCCCTTT	
F	482	SEQ ID NO:8	AAAATCTTCGTCAACATGGTGGAGC	
		SEQ ID NO:9	TAATCTCAACTGGTCTCCTCTCCGG	
G	366	SEQ ID NO:19	GGCTCGGACTCGACCTTTCTAT	
		SEQ ID NO:20	GCAGTTCTTGAAGAATGAGTGA	
H	912	SEQ ID NO:1	GTAGTACTATAGATTATATTATTCGTAGAG	
		SEQ ID NO:2	GCCATACAGAACTCAAAATCTTTTCCGGAG	
I	844	SEQ ID NO:2	GCCATACAGAACTCAAAATCTTTTCCGGAG	
		SEQ ID NO:23	CTTCAAACAAGTGTGACAAA	

Example 7

Flanking Sequence 3' to Inserted Transgenic DNA in Maize Event TC1507

[0097] Two separate PCR approaches were used to extend the length of the sequence information 3' to the full-length PH18999A insert in maize event TC1507. In the first approach PCR primer pairs were designed to amplify a product that spanned the junction between the full-length insert and the inverted ORF25 terminator, see FIG. 1 for a depiction of the inverted ORF25 terminator. A forward primer was located at the end of the full-length PH18999A insert and a series of reverse primers were located at 100 bp intervals in the inverted sequence. In this manner the length of the inverted fragment present in the maize event TC1507 could be determined within a 100 bp region based on the successful PCR reactions. This method indicated the inverted fragment contained the majority of the ORF25 terminator but no Cry1F sequence. PCR fragments were isolated and sequenced from this region.

[0098] In the second approach PCR primers were designed to walk out into the flanking DNA sequence from the inverted ORF25 terminator region as determined in the PCR experiment described above. Genomic DNA isolated from two to three individual plants of event TC1507 and an unmodified control corn line was digested with various restriction enzymes and then ligated to adaptors specific for the restriction enzyme used for digestion (Universal Genome Walker™ Kit, Clontech Laboratories, Inc. and Devon et al. (1995) *Nucleic Acids Res.* 23:1644-1645). Primary PCR was carried out using an ORF25 terminator

specific primer and a primer homologous to the adaptor sequence ligated onto the digested DNA. In order to increase the specificity of the reaction a nested secondary PCR was performed again with another ORF25 terminator specific primer and a secondary primer homologous to the adaptor sequence with the secondary primers being internal to the respective primers used in the primary PCR. Products produced by the nested PCR were analyzed by agarose gel electrophoresis and fragments unique to TC1507 DNA samples were isolated and sequenced. Fragments were amplified from both the ORF25 terminator contained within the full-length insert and from the targeted (inverted) ORF25 terminator on the 3' end of the full-length PH18999A insert. Fragments from the full-length insert were of a predicted size based on the knowledge of the restriction enzyme sites located in the full-length insert. Fragments produced from the 3' inverted ORF25 terminator appeared as fragments of unexpected size. Sequence analysis of amplified fragments from the 3' inverted ORF25 terminator resulted in flanking DNA sequence of 1043 bp. Resultant sequence from the above series of genome walking experiments was used to design additional primers to walk further out from the insert into the bordering maize genome with a final 3' flanking sequence, of 2346 bp.

[0099] In order to describe the TC1507 3' flanking sequence, homology searching was done against the GenBank public databases using the Basic Local Alignment Search Tool (BLAST). The BLAST program performs sequence similarity searching and is particularly useful for identifying homologs to an unknown sequence. In addition to searching the public databases, alignments were per-

formed using SeqMan 4.05™, Martinez and Needleman-Wunsch alignment algorithms (DNASTAR Inc.) to look for homology between the TC1507 3' flanking sequence and the PHI8999A transgenic insert. The results of these homology searches are presented in Table 1. The percent identity values indicate the percentage of identical matches across the length of the sequences analyzed. The results of similarity searching for the 3' flanking sequence indicate high homology with three regions of maize chloroplast DNA, a 188 bp fragment of the *pat* gene, and 254 bp of DNA (Region 15, Table 1) with no significant homology. An additional 749 bp (Region 16) beyond Region 15 (see Table 1) was also sequenced. No similarity searching results are available for Region 16.

[0100] PCR analysis on control and TC1507 genomic DNA determined that the 254 bp sequence (Region 15, fragment of maize chloroplast “ORF241”) is present in the maize genome. The DNA sequence of Region 15 in the 3' flanking region is not unique to the 3' flanking region of maize event TC1507 but is also present in the unmodified control corn line. The TC1507 3' flanking sequence is presented in Example 8 and diagrammed in FIG. 1.

Example 8

Sequence of the Region 3' to the End of the Full-Length-Insert DNA in maize Event TC1507.

[0101] A description of each region is in Table 1.

Region 10 (SEQ ID NO: 38) Fragment of ORF25 Terminator (complement)

```
9016 CTCAC TCCGCTTGAT CTTGGCAAAG ATATTTGACG

9051 CATTTATTAG TATGTGTAA TTTTCATTG CAGTGCAGTA TTTTCTATTC

9101 GATCTTTATG TAATTCGTTA CAATTAATAA ATATTCAAAT CAGATTATTC

9151 ACTGTCATTG GTATCAAATC GTGTTAATG GATATTTTGA TTATAATATT

9201 GATGATATCT CAATCAAAC GTAGATAATA ATAATATTTA TTTAATATTT

9251 TTGCGTCGCA CAGTGAAAAT CTATATGAGA TTACAAAATA CCGACAACAT

9301 TATTTAAGAA ACATAGACAT TAACCTGAG ACTGTTGGAC ATCAACGGGT

9351 AGATTCTTC ATGCATAGCA CCTCATTCTT GGGGACAAAA GCACGGTTTG

9401 GCCGTTCCAT TGCTGCACGA ACGAGCTTTG CTATATCCTC GGGTTGGATC

9451 ATCTCATCAG GTCCAATCAA ATTTGTCCAA GAACTCATGT TAGTCGCAAC

9501 GAAACCGGG CATATGTCGG GTATCTCGAG CTCGCGAAAG CTTGGCTGCA

9551 GGTCGACGGA TCCTT
```

Region 11 (SEQ ID NO: 39) Fragment of maize chloroplast *rps12* rRNA gene (complement)

```
9566 CAACA AAAGGGTACC TGTACCCGAA ACCGACACAG

9601 GTGGGTAGGT AGAGAATACC TAGGGGCGCG AGACAACTCT CTCTAAGGAA

9651 CTCGGCAAAA TAGCCCCGTA ACTTCGGGAG AAGGGTGCC CCC
```

Nucleotides 9694–9695 (CG) connect Region 11 to Region 12.

Region 12 (SEQ ID NO: 40) Fragment of maize chloroplast genome

```
9696 CTAAC

9701 AATAAACGAA TACGGTTTAT GTATGGATTC CGGTAAAATA CCGGTACTCG

9751 ATTTTCATAAG AGTCGAATAG GAAGTTAAGA TGAGGGTGGT ATCATCATAA

9801 AAATGGAGTA GTATCCTAAA TTATACTAAT CCACGTATGA TATGTATGCC

9851 TTTCTTATC AACCGGAAGT AGTGCAAAAA AAATTCATATA CTGCACTGCT

9901 CTCTTTTAC TGAGAAATGC AAAAAAATAA AAGTGAAGTA AGGGTGCCCC

9951 ATAGATATTT GATCTTGCTT CCTGTCCCCC CCCCCCTTTT TTCATCAAAA

10001 ATTTCCATGA AAAAAAGAAA GATGAATTG TCCATTCAAT GAACCTAGT

10051 TCGGGACTGA CGGGGCTCGA ACCCGCAGCT TCCGCCT
```

-continued

Region 13 (SEQ ID NO: 41) Fragment of *pat* gene (complement)

10088 GTT CCTAGCCTTC
10101 CAGGGCCGAG CGTAAGCAAT ACCAGCCACA GCACCCTCAA CCTCAGCAAC
10151 CAACCAAGGG TATCTATCTT GCAACCTCTC TAGATCATCA ATCCACTCTT
10201 GTGGTGTTTG TGGCTCTGTC CTAAAGTTCA CTGTAGACGT CTCAATGTAA
10251 TGGTTAACGA TATCACAAAC CGCGG

Nucleotides 10276–10277 (AA) connect Region 13 to Region 14.

Region 14 (SEQ ID NO: 42) Fragment of maize chloroplast ORF241 (complement)

10278 CAC AAGAACGAAA GCACCTTTTC
10301 ATTCTTTCAT ATACTAGGGG TTTTACTTG GAAAAGACAA TGTTCATAC
10351 TAAAGGAT

Region 15 (SEQ ID NO: 43) Maize genomic (no significant homology)

10359 AG CTGCAGAAGC CGCCACCGTC TTGAGGACCT TCCGGGGAGC
10401 CAGACCGGTC GAACCGTGCC TCCACTTGCT AAGGAGAAAG GGAATCAG
10451 GGCCAGGACA TACGAAGGAG GAGCCAGAAC GAAGATATCC TAAGATACTT
10501 ACTCGCTCCG GGCCATGATC AATCATGCCT GTGGGAGGT CTCTCGCACC
10551 TCGATCCATG AAGGTACCAC CGAGGTCTGC CCCGCCGCCG GCTTCGGTAC
10601 CGTCCTCGCC TT

Region 16 (SEQ ID NO: 44) Maize genomic

10613 GGGCGCCC GAGGCACCG GGGATGGAC TGCCAGGCG
10651 CAGCCACGAC GACCAAGGA TCACCCTCCT GCGCAGTCGG CACGAGCAAT
10701 AGTTCTCGGG GAACAGGCAG CTGGCCTGA CTCCCCGGG TCACCTCAAC
10751 TACCTCGGCC GAGGGTCAA GTACCCCTC AGTCCGCCC CGCTCTTCGG
10801 ACCGGGACCC CGACGTCCC GCCCCGATA CCGACGGCAC CAGCCCGCTC
10851 GGGGCTGGC TTGACGACC CTGGCCAGC CTCAGATCTG GGCTGAGGCC
10901 GAGCAGGCG GCCATGTCGT CGTCTTCATC ATCGTCTTCA TCATCGTCGT
10951 CGTCATCAGG CGTCTCCGGC GACGGCTCCC TTGGGAGCCC CTCCTCTCC
11001 TGCCGACGAC GAAGCCTTTC CAAGGCATCC CGAGCCACG TCCGCTCGTG
11051 GGCCGAGGCC TTCTTTGCGT CCTTCTTCTC CTTCTCTTC TCCGCGGTGA
11101 CCCTCCGCGC AGCTCGGTCC ACCGCATCCT CCGGACTGG TGGCAGGGAA
11151 GGCTTGATGAT GCCCTACCTC CTGGAGACAG ACGAAAAGTC TCAGCTATGA
11201 GAACCGAGGG CAATCTGACG CAAGAAGGAA GAAGGAGCGG ATACTACCA
11251 GAGACACGCA CCCGCGATCG GGACGCATTA AGGGCTGGGA AAAAGTGCCG
11301 GCCTCTAATT TCGCTACCGT GCCGTCCACC CACCTGTGGA GGTCATCGAT
11351 GGAAGGGGA A

Example 9

Confirmation of the Presence of Region 15 in the
Unmodified Control Corn Line

[0102] PCR analysis was used to determine if the undescribed region of sequence on the end of the 3' flanking sequence (Region 15 in Table 1) is present in the unmodified control corn line used for transformation to produce maize event TC1507 and thus represents a border with corn genomic DNA. Successful PCR amplification of Region 15 in both maize line TC1507 and the unmodified Hi-II control corn line revealed that Region 15 was indeed present in corn genomic DNA. Five different PCR analyses were carried out on genomic DNA prepared from TC1507 and the unmodified Hi-II control corn line as outlined in Table 7 below using the primer sequences shown in Table 8. Three reactions were designed to amplify DNA within Region 15 of the 3' flanking region; Reaction L—producing a 175 bp amplicon, Reaction M—producing a 134 bp amplicon, and Reaction N—producing a 107 bp amplicon. The expected amplicons were present in both the unmodified control corn line and in maize line TC1507.

[0103] Reactions J and K were designed as specific primer pairs for TC1507 that would span the end of the insert into the 3' flanking region. In Reaction J, the forward primer was located in the pat gene fragment on the 3' end of the full-length PH18999A insert (Region 13 in Table 1) and the reverse primer was located in the undefined Region 15. In Reaction K the forward primer was located in the chloroplast hypothetical protein gene on the 3' end of the full-length insert (Region 14 in Table 1) and the reverse primer was located in the undefined Region 15. Both Reaction J and Reaction K produced an amplicon in maize line TC1507 and did not produce an amplicon in the unmodified control corn line. The results indicate that both Reactions J and K are specific for the TC1507 event.

[0104] The PCR results indicate that the undescribed sequence (Region 15) of the 3' flanking sequence of TC1507 is also present in genomic DNA isolated from the unmodified Hi-II control corn line. The DNA sequences amplified in Reactions L, M, and N are not unique to the 3' flanking region of TC1507 but are also present in the unmodified control corn line.

TABLE 7

PCR reactions for sequence 3' to the PH18999A insert in maize event TC1507				
Reaction	Amplicon Size (bp)	Region in TC1507 3' flanking sequence	Amplicon present in Control	Amplicon present in maize line TC1507
J	342	Region 13 (pat gene fragment) to Region 15	No	Yes
K	252	Region 14 (chloroplast gene) to Region 15	No	Yes
L	175	Region 15	Yes	Yes
M	134	Region 15	Yes	Yes
N	107	Region 15	Yes	Yes

[0105]

TABLE 8

PCR primers for sequence 3' to the PH18999A insert in maize event TC1507				
Reaction	Amplicon Size (bp)	Primer Pair	Primer Sequences 5' to 3'	
J	342	SEQ ID NO:3	TGTGGTGTTCGTGGCTCTG	
		SEQ ID NO:5	TCCTAA GACCTCCCCACAGGCATGA TTGATC	
K	252	SEQ ID NO:4	AGCACCTTTTCATTCTTTC	
		SEQ ID NO:5	ATATAC GACCTCCCCACAGGCATGA TTGATC	
L	175	SEQ ID NO:16	AAGCCGCCACCGTCTTGAG	
		SEQ ID NO:5	GACCTT GACCTCCCCACAGGCATGA TTGATC	
M	134	SEQ ID NO:17	GTCGAACCGTGCTCCACT	
		SEQ ID NO:5	TGCTAA GACCTCCCCACAGGCATGA TTGATC	
N	107	SEQ ID NO:18	AGAAAGGGAAAATCAGGGC	
		SEQ ID NO:5	CAGGAC GACCTCCCCACAGGCATGA TTGATC	

Example 10. PCR Primers

[0106] DNA event specific primer pairs were used to produce an amplicon diagnostic for TC1507. These event primer pairs include, but are not limited to, SEQ ID NO: 1 and SEQ ID NO: 2; SEQ ID NO: 2 and SEQ ID NO: 23; SEQ ID NO: 3 and SEQ ID NO: 5; and SEQ ID NO: 4 and SEQ ID NO: 5. In addition to these primer pairs, any primer pair derived from SEQ ID NO: 26 and SEQ ID NO: 27 that when used in a DNA amplification reaction produces a DNA amplicon diagnostic for TC1507 is an aspect of the present invention. The amplification conditions for this analysis are illustrated in Table 9, however, any modification of these methods that use DNA primers or complements thereof to produce an amplicon DNA molecule diagnostic for TC1507 is within the ordinary skill of the art. The preferred amplification conditions for reactions utilizing the PCR primers identified in SEQ ID NOS: 1, 2, and 23 are illustrated in Table 10. In addition, control primer pairs, which include SEQ ID NOS: 10 and 11; SEQ ID NOS: 10 and 12; SEQ ID NOS: 13 and 14; SEQ ID NOS: 14 and 15; SEQ ID NOS: 5 and 16; SEQ ID NOS: 5 and 17; and SEQ ID NOS: 5 and 18; for amplification of an endogenous corn gene are included as internal standards for the reaction conditions. Also included are primer pairs that will produce an amplicon in transgenic events containing a pat gene (SEQ ID NOS: 6 and 7; SEQ ID NOS: 8 and 9), and a primer pair that will produce an amplicon in transgenic events containing a cry1F gene (SEQ ID NOS: 19 and 20).

[0107] The analysis of plant tissue DNA extracts to test for the presence of the TC1507 event should include a positive tissue DNA extract control (a DNA sample known to contain the transgenic sequences). A successful amplification of the positive control demonstrates that the PCR was run under

conditions which allow for the amplification of target sequences. A negative, or wild-type, DNA extract control in which the template DNA provided is either genomic DNA prepared from a non-transgenic plant, or is a non-TC1507 transgenic plant, should also be included. Additionally a negative control that contains no template corn DNA extract will be a useful gauge of the reagents and conditions used in the PCR protocol.

[0108] Additional DNA primer molecules of sufficient length can be selected from SEQ ID NO: 26 and SEQ ID NO: 27 by those skilled in the art of DNA amplification methods, and conditions optimized for the production of an amplicon that may differ from the methods shown in Table 9 or Table 10 but result in an amplicon diagnostic for event TC1507. The use of these DNA primer sequences with modifications to the methods shown in Table 9 and Table 10 are within the scope of the invention. The amplicon wherein at least one DNA primer molecule of sufficient length derived from SEQ ID NO: 26 and SEQ ID NO: 27 that is diagnostic for event TC1507 is an aspect of the invention. The amplicon wherein at least one DNA primer of sufficient length derived from any of the genetic elements of PHI8999A that is diagnostic for event TC1507 is an aspect of the invention. The assay for the TC1507 amplicon can be performed by using a Stratagene Robocycler, MJ Engine, Perkin-Elmer 9700, or Eppendorf Mastercycler Gradient thermocycler, or by methods and apparatus known to those skilled in the art.

TABLE 9

PCR Conditions:	
Conditions:	Perkin-Elmer AmpliTaq Gold kit
Kit used:	Component
Volume	
5 μ l	template (10 ng/ μ l)
4 μ l	2 μ l each primer (10 μ M)
2 μ l	10X PCR Gold Buffer
2 μ l	25 mM MgCl ₂
2 μ l	50X dNTP's (10 mM)
0.1 μ l	Amplitaq Gold Polymerase
4.9 μ l	H ₂ O
20 μ l	Total

Cycling Parameters
GeneAmp ® PCR System 9700
9 min 92° C.
30 cycles:
94° C. 30 sec
60° C. 30 sec
72° C. 1 min
7 min 72° C.
Hold 4° C.

[0109]

TABLE 10

PCR Conditions used with the Advantage ®-GC 2 Polymerase Mix:	
Conditions:	Advantage ®-GC 2 Polymerase Mix
Kit used:	Component
Volume	
5 μ l	template (10 ng/ μ l)
5 μ l	2.5 μ l each primer (10 μ M)
10 μ l	5x GC2 Buffer
10 μ l	GC melt (1.0 M final conc.)
1.5 μ l	50X dNTP's (10 mM)
1.0 μ l	Advantage GC2 Polymerase
17.5 μ l	H ₂ O
50 μ l	Total

Cycling Parameters
GeneAmp ® PCR System 9700
5 min 94° C.
35 cycles:
94° C. 1 min
60° C. 2 min
72° C. 3 min
7 min 72° C.
Hold 4° C.

[0110] Having illustrated and described the principles of the present invention, it should be apparent to persons skilled in the art that the invention can be modified in arrangement and detail without departing from such principles. We claim all modifications that are within the spirit and scope of the appended claims.

[0111] All publications and published patent documents cited in this specification are incorporated herein by reference to the same extent as if each individual publication or patent application was specifically and individually indicated to be incorporated by reference.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 57

<210> SEQ ID NO 1

<211> LENGTH: 30

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Event specific primer sequence designed for

-continued

TC1507

<400> SEQUENCE: 1

gtagtactat agattatatt attcgtagag 30

<210> SEQ ID NO 2

<211> LENGTH: 30

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Event specific primer sequence designed for TC1507

<400> SEQUENCE: 2

gccatacaga actcaaaatc ttttccggag 30

<210> SEQ ID NO 3

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Event specific primer designed for TC1507.

<400> SEQUENCE: 3

tgtgtgtgtt gtggctctgt cctaa 25

<210> SEQ ID NO 4

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Event specific primer for TC1507.

<400> SEQUENCE: 4

agcacctttt cattctttca tatac 25

<210> SEQ ID NO 5

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Genomic DNA primer sequence

<400> SEQUENCE: 5

gacctcccca caggcatgat tgatc 25

<210> SEQ ID NO 6

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Primer in full length insert, 35S promoter to pat gene

<400> SEQUENCE: 6

tagaggacct aacagaactc gccgt 25

<210> SEQ ID NO 7

<211> LENGTH: 24

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Primer in full length insert, 35S promoter to pat gene

-continued

<400> SEQUENCE: 7

gagctggcaa ctcaaatcc cttt

24

<210> SEQ ID NO 8

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Primer in full length insert, 35S promoter to
pat gene

<400> SEQUENCE: 8

aaaatcttcg tcaacatggt ggagc

25

<210> SEQ ID NO 9

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Primer in full length insert, 35S promoter to
pat gene

<400> SEQUENCE: 9

taatctcaac tggctctcctc tccgg

25

<210> SEQ ID NO 10

<211> LENGTH: 22

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Primer - Zea mays genomic DNA

<400> SEQUENCE: 10

cccctacccc accgacgttt at

22

<210> SEQ ID NO 11

<211> LENGTH: 22

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Primer - Zea mays genomic DNA

<400> SEQUENCE: 11

ttgattggca ggtccgtggg tc

22

<210> SEQ ID NO 12

<211> LENGTH: 22

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Primer - Zea mays genomic DNA.

<400> SEQUENCE: 12

cacaacggca cagaaacacg aa

22

<210> SEQ ID NO 13

<211> LENGTH: 22

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Primer - Zea mays genomic DNA.

<400> SEQUENCE: 13

-continued

gcgcacccac cggaacaaaa tg 22

<210> SEQ ID NO 14
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer - Zea mays genomic DNA.

<400> SEQUENCE: 14

tcctcgcatc aaatgctcct gc 22

<210> SEQ ID NO 15
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer - Zea mays genomic DNA.

<400> SEQUENCE: 15

cctggcacgc attgacgcac gt 22

<210> SEQ ID NO 16
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer - Zea mays genomic DNA.

<400> SEQUENCE: 16

aagccgccac cgtcttgagg acctt 25

<210> SEQ ID NO 17
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer - Zea mays genomic DNA.

<400> SEQUENCE: 17

gtcgaaccgt gcctccactt gctaa 25

<210> SEQ ID NO 18
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer - Zea mays genomic DNA.

<400> SEQUENCE: 18

agaaagggaa aatcagggcc aggac 25

<210> SEQ ID NO 19
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Cry1F sequence primer

<400> SEQUENCE: 19

ggctcggact cgacctttct at 22

-continued

<210> SEQ ID NO 20
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Cry1F sequence primer

<400> SEQUENCE: 20

gcagttcttg aagaatgagt ga 22

<210> SEQ ID NO 21
<211> LENGTH: 2829
<212> TYPE: DNA
<213> ORGANISM: 5' flanking sequence of event TC1507

<400> SEQUENCE: 21

actagtttcc tagcccgct cgtgcccta cccaccgac gtttatggaa ggtgccattc 60
cacggttctt cgtggccgcc cctaaggatg taaatggctg gtaaaatccg gtaaatcttc 120
ggtaccgttt accagatttt tccagccgtt ttcggattta tcgggatata cagaaaacga 180
gacggaaacg gaatagggtt tttttcgaaa acggtacggt aaacggtgag acaaacttac 240
cgtccgtttt cgtattttctc gggaaactct ggtatatctc cgtatttgtc ccgtattttc 300
ccgacccacg gacctgcaa tcaaccatca gccagtcagc ccatccccac agctatggcc 360
catggggcca tgttggccac atgccacgc aacgcaaggc agtaaggctg gcagcctggc 420
acgcattgac gcattgtggc acacacagcc gccgcctgtt cgtgtttctg tgccgttggtg 480
cgagactgtg actgcgagtg gcggagtcgg cgaacggcga ggcgtctccg gagtctggac 540
tgccgctgtg gacagcgacg ctgtgacggc gactcggcga agccccagc taccaagccc 600
ccaagtcccc atccatctct gtttctctg tcatctcctt cccctggctg atctgcaggc 660
gccagaccgg ccgaagcatc acgaaacgca ctaagacctc gaaggagtca aaccactcct 720
ccgaggcctc gggggctaca cccggcgggt gcgctcgcgc gcaccacccg gaacaaaatg 780
taaccgagaa aggtcgttcc ccttgcaaaa aaagtgcgac aaaagcctcc aagcgagtat 840
taaacctcac tttgaggctc gggggctact gtcggggacc ataattaggg gtaccccaaa 900
gactccta atctagctggt aacccccatc agcacaagc tgcaaaggcc tgatgggtgc 960
gattaagtca aggtcgtgct cactcaaggg acacgatctc gcctcgcgc agccagcct 1020
cgggcaaggg cgcccgaccc cgaggattca cgtctcgcgc gagggcccc tcaagcgacg 1080
ggcacacctt cggtcgcgc gagggccatt ctctcgcgc aagcaacctt ggccagatcg 1140
ccacaccgac cgaccgtatc gcaggagcat ttaatgcgag gatcgctga caccttatcc 1200
tgacgcgcgc tcttcagtcg acagagccga agtgaccgca atcacttcgc cgctccactg 1260
accgacctga caagaagaca gcgcgcctg cgtcgcctcg actgctgtgc cactcgacag 1320
agtgaaggctg acagcagcca agtcggcct cgggcgccat aggaagctcc gcctcgcgc 1380
accctagggc tcggactcgg cctcggctcc ggaagacgac gaactacgct tcgcccagcc 1440
ccagggttg gactcagcct cggtccgga agacgacgaa ttccgcctcg cccgacccca 1500
gggctcggac tcggcctcgg ctccagaaga cgacgaactc cgcctcgcgc gacccaggg 1560
ctcggactca gcctcggctc cggaagacga cgaactccgc ctgcgccgac cccagggtc 1620
ggactcagcc tcggcctcag acgatggctt ccgcctcgc cgacccgggg ctcgactcg 1680
acctttctat cggaacctgt cagatcctgt ctctgtccga ggaggcttg gcaatcctca 1740

-continued

```

ctatgtactc ggtcttaggg gagtggcctt tcaacaaact ggtacgaatc acaccgcac 1800
attcaggaac tccgggacca ttgactctct agatgagata ccacctcaag acaacagcgg 1860
cgcaccttg aatgactact cccatgtgct gaatcatgtt acctttgtgc gctggccagg 1920
tgagatctca ggttccgact catggagagc accaatgttc tcttgagcgc atcgtagcgc 1980
tacccccaca aacaccattg atccagagag aatcactcat tcttcaagaa ctgcatactt 2040
tgccgagatc ctcatcccta aaggtagctg acaatagtat tattggagtc gatacacaa 2100
tcacaaaaaa tacaagaagt cgactagggtg gattgggtccg agtgaagaga aaaaaagcc 2160
atacagaact caaaatcttt tccggagata ttcattttcc tgaagaggcg gataagatat 2220
taggtggcag tttgatacca ccagaaagag aaaaaaaga ttctaaggaa tcaaaaaaa 2280
ggaaaaattg ggtttatgtt caacggaaaa aatttctcaa aagcaaggaa aagtattgtg 2340
gctattttatc tatccgtgca gctgatattg ccgcgggttg tgatatcgtt aaccattaca 2400
ttgagacgtc tacagtgaac tttagacag agccacaaac accacaagag tggattgatg 2460
atctagagag gttgcaagat agataccctt ggttggttgc tgagggtgag ggtgtgtggy 2520
ctggtattgc ttacgctggg ccttggagg cttaggaaccc tcaacctcag caaccaacca 2580
atggtatcta tcttgcaacc tctctagatc atcaatccac tcttgtggtg tttgtggctc 2640
tgtcctaaag ttcactgtag acgtctcaat gtaatggta acgatatcac aaaccgagag 2700
aagagggatc tcgaagcttc ggcggggcc catcgatata cgcgggcatg cctgcagtgc 2760
agcgtgaccc ggtcgtgccc ctctctagag ataatgagca ttgcatgtct aagttataaa 2820
aaattacca 2829

```

<210> SEQ ID NO 22

<211> LENGTH: 2346

<212> TYPE: DNA

<213> ORGANISM: 3' flanking sequence of event TC1507

<400> SEQUENCE: 22

```

ctcactccgc ttgatcttgg caaagatatt tgacgcattt attagtatgt gttaattttc 60
atttgcagtg cagtattttc tattcgatct ttatgtaatt cgttacaatt aataaatatt 120
caaatcagat tattgactgt catttgtatc aaatcgtgtt taatggatat ttttattata 180
atattgatga tatctcaatc aaaacgtaga taataataat atttatttaa ttttttgcg 240
tcgcacagtg aaaatctata tgagattaca aaataccgac aacattatct aagaaacata 300
gacattaacc ctgagactgt tggacatcaa cgggtagatt ccttcoatga tagcacctca 360
ttcttgggga caaaagcacg gtttggccgt tccattgtcg cacgaacgag ctttgcata 420
tcctcggggt ggatcatctc atcagggtcca atcaaatctg tccaagaact catgttagtc 480
gcaacgaaac cggggcatat gtcgggtatc tcgagctcgc gaaagcttgg ctgcaggctc 540
acggatcctt caacaaaagg gtacctgtac ccgaaaccga cacagggtgg taggtagaga 600
atacctaggg gcgcgagaca actctctcta aggaactcgg caaaatagcc ccgtaacttc 660
gggagaaggg gtgccccccg ctaacaataa acgaatacgg tttatgtatg gattccggta 720
aaataccggt actcgatttc ataagagtcg aataggaagt taagatgagg gtggtatcat 780
cataaaaatg gagtagtatc ctaaattata ctaatccacg tatgatattg atgcctttcc 840
ttatcaaccg gaagtagtgc aaaaaaaatt ctatactgca ctgctctctt tttactgaga 900

```

-continued

```

aatgcaaaaa aataaaagtg aagtaagggt gcccacataga tatttgatct tgccctcctgt    960
ccccccccc cttttttcat caaaaatttc catgaaaaaa gaaaagatga atttgccat    1020
tcattgaacc ctagttcggg actgacgggg ctgcaacccg cagcttcgcg ctgttcctag    1080
ccttcacagg cccagcgtaa gcaataccag ccacagcacc ctcaacctca gcaaccaacc    1140
aagggtatct atcttgcaac ctctctagat catcaatcca ctcttggtgt gtttggtgct    1200
ctgtcctaaa gttcactgta gacgtctcaa tgtaatggtt aacgatatca caaacgcgg    1260
aacacaagaa cgaaagcacc ttttcattct ttcataact aggggttttt acttggaana    1320
gacaatgttc cactactaaag gatagctgca gaagccgcca ccgtcttgag gaccttcgg    1380
ggagccagac cggctgaacc gtgcctccac ttgctaagga gaaagggaaa atcagggcca    1440
ggacatacga aggaggagcc agaacgaaga taccctaaga tacttactcg ctccgggcca    1500
tgatcaatca tgccctgtgg gaggtctctc gcacctcgat ccataaagg accaccgagg    1560
tctgccccgc cgccggcttc ggtaccgtcc tcgccttggg cgcccgaggc acccggggga    1620
tggactgccc aggcgcagcc acgacgaccc aaggatcacc ctccctgcga gtcggcacga    1680
gcaatagttc tcggggaaca ggcagcttgg cctgactccc cggggtcacc tcaactacct    1740
cgcccgaggg gtcaagtacc ccctcagtcc gcccccgctc tcgggaccgg gaccccgacg    1800
tcccggcccc ggataccgac ggcaccagcc cgctcggggg ctggcttgac gacccctggc    1860
ccagcctcag atctgggctg aggcggaggc aggcggccat gtcgtcgtct tcacatcgt    1920
cttcatcatc gtcgtcgtca tcaggcgtct ccggcgacgg ctcccttggg agccctccc    1980
tctcctgcg acgacgaagc ctttccaagg catcccgagc ccacgtccgc tcgtggggcc    2040
gagccttctt tgctccttc ttctccttc tcttctccgc ggtgacctc cgcgcagctc    2100
ggtccaccgc atcctccggg actggtggca gggaaggctt gtgatgccct acctcctgga    2160
gacagacgaa aagtctcagc tatgagaacc gagggcaatc tgacgcaaga aggaagaagg    2220
agcggatact caccagagac acgcacccgc gatcgggacg cattaagggc tgggaaaaag    2280
tgccggcctc taatttcgct accgtgccgt ccacccacct gtggaggtca tcgatgggaa    2340
ggggaa                                           2346

```

```

<210> SEQ ID NO 23
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Event specific primer sequence designed for
TC1507

```

```

<400> SEQUENCE: 23

```

```

cttcaaacaa gtgtgacaaa                                           20

```

```

<210> SEQ ID NO 24
<211> LENGTH: 11361
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: The sequence represents the transgenic insert
in maize line TC1507 as well as the sequence flanking the
insertion sites.

```

```

<400> SEQUENCE: 24

```

-continued

actagtttcc tagcccgct cgtgcccta cccaccgac gtttatggaa ggtgccattc	60
cacggttctt cgtggccgcc cctaaggatg taaatggctg gtaaatccg gtaaatttcc	120
ggtaccgttt accagatttt tccagccgtt ttcggattta tcgggatata cagaaaacga	180
gacggaaaacg gaatagggtt tttttcgaaa acggtacggt aaacggtgag acaaacttac	240
cgtccgtttt cgtattttctc gggaaactct ggtatattcc cgtatttgc ccgtattttc	300
ccgacccacg gacctgcaa tcaacatca gccagtcagc ccatcccccac agctatggcc	360
catggggcca tgttgccac atgccacgc aacgcaaggc agtaaggctg gcagcctggc	420
acgcattgac gcattgtggac acacacagcc gccgcctgtt cgtgtttctg tgccgttgtg	480
cgagactgtg actgcagtg gcggagtcg cgaacggcga ggctctccg gactctggac	540
tgcggctgtg gacagcgac ctgtgacggc gactcggcga agccccaagc taccaagccc	600
ccaagtcccc atccatctct gcttctctgg tcatctcctt cccctggtcg atctgcaggc	660
gccagaccgg ccgaagcatc acgaaacgca ctaagacctc gaaggagtca aaccactcct	720
ccgagggctc gggggctaca cccggcggt gcgctcgcgc gcacccaccg gaacaaaatg	780
taaccgagaa aggtcggtcc ccttgcaaaa aaagtgcgac aaaagcctcc aagcgagtat	840
taacactcac tttgaggctc gggggctact gtcggggacc ataattaggg gtaccccaaa	900
gactccta atcagctggt aacccccatc agcacaagc tgcaaaaggc tgatgggtgc	960
gattaagtca aggtcggtc cactcaaggc acacgatctc gcctcgccc agcccagcct	1020
cgggcaaggc cgccgaccc cgaggattca cgtctcgccc gagggcccc tcaagcgacg	1080
ggcacacctt cggtcggccc gagggccatt cttcgccgag aagcaacctt ggccagatcg	1140
ccacaccgac cgaccgtatc gcaggagcat ttaatgcgag gatcgccga caccttatcc	1200
tgacgcgcgc tcttcagtcg acagagccga agtgaccgca atcaattcgc cgtccactcg	1260
accgacctga caagaagaca gcgccgctg cgtcgtccg actgctgtgc cactcgacag	1320
agtgaggctg acagcagcca agtcggcct cgggcgccat aggaagctcc gcctcgccc	1380
accctagggc tcggactcg cctcggtcc ggaagacgac gaactacgct tcgcccagcc	1440
ccagggttg gactcagcct cggtccgga agacgacgaa ttccgctcg cccgacccca	1500
gggtcggac tcggcctcg ctccagaaga cgacgaactc cgctcgccc gacccaggg	1560
ctcgactca gcctcggtc cggaagacga cgaactcgc ctgcccgcac cccagggtc	1620
ggactcagcc tcggcctcag acgatggtct ccgcctcgc cgacccggg ctcgactcg	1680
acctttctat cggacctgt cagatcctgt cttcgtccga ggaggcttg gcaatcctca	1740
ctatgtactc ggtcttaggg gagtggcctt tcaacaaact ggtacgaatc acaccgcac	1800
attcaggaac tccgggacca ttgactctct agatgagata ccacctcaag acaacagcgg	1860
cgcaccttg aatgactact cccatgtgct gaatcatgtt acctttgtgc gctggccagg	1920
tgagatctca ggttcgact catggagagc accaatgttc tcttgagcgc atcgtagcgc	1980
tacccccaca aacaccattg atccagagag aatcactcat tcttcaagaa ctgcatatct	2040
tgccgagatc ctcatcccta aaggctactg acaatagtat tattggagtc gatacacaac	2100
tcacaaaaaa tacaagaagt cgactagggt gattgggtccg agtgaagaga aaaaaagcc	2160
atacagaact caaaatcttt tccggagata ttcattttcc tgaagaggcg gataagatat	2220
taggtggcag tttgatacca ccagaagag aaaaaaaga ttctaaggaa tcaaaaaaa	2280

-continued

ggaaaaattg ggtttatggt caacggaaaa aattttctcaa aagcaaggaa aagtattgtg	2340
gctattttatc tatcogtgca gctgatatgg ccgcggtttg tgatatacgtt aaccattaca	2400
ttgagacgtc tacagtgaac tttaggacag agccacaaac accacaagag tggattgatg	2460
atctagagag gttgcaagat agataccctt ggttggttgc tgaggttgag ggtgtgtgtg	2520
ctggtattgc ttacgctggg ccctggaagg ctaggaaccc tcaacctcag caaccaacca	2580
atggtatcta tcttgcaacc tctctagatc atcaatccac tcttggtgtg tttgtggctc	2640
tgctcctaaag ttactgtag acgtctcaat gtaatggtta acgatatac aaaccgagag	2700
aagagggatc tcgaagcttc ggcgggggcc catcgatata cgcgggcatg cctgcagtgc	2760
agcgtgaccc ggtcgtgccc ctctctagag ataatgagca ttgcatgtct aagttataaa	2820
aaattaccac aactggaaga gcggttaccg ggaccgaagc ttcggccggg gcccatcgat	2880
atccgcgggc atgcctgcag tgcagcgtga cccggtcgtg cccctctcta gagataatga	2940
gcattgcatg tctaagttat aaaaaattac cacatatttt tttgtgcaca cttgtttgaa	3000
gtgcagttta tctatcttta tacatatatt taaactttac tctacgaata atataatcta	3060
tagtactaca ataatatcag tgttttagag aatcatataa atgaacagtt agacatggtc	3120
taaaggacaa ttgagtattt tgacaacagg actctacagt tttatctttt tagtgtgcat	3180
gtgttctcct ttttttttgc aaatagcttc acctatataa tacttcatcc attttattag	3240
tacatccatt tagggtttag ggttaatggt ttttatagac taattttttt agtacatcta	3300
ttttattcta ttttagcttc taaattaaga aaactaaac tctatttttag tttttttatt	3360
taataattta gatataaaat agaataaaat aaagtacta aaaattaaac aaataccctt	3420
taagaaatta aaaaaactaa ggaacattt ttcttgtttc gagtagataa tgccagcctg	3480
ttaaacgcg tcgacgagtc taacggacac caaccagcga accagcagcg tcgctgcggg	3540
ccaagcgaag cagacggcac ggcattctctg tcgctgcctc tggaccctc tcgagagttc	3600
cgctccaccg ttggacttgc tccgctgtcg gcatccagaa attgctgtgc ggagcggcag	3660
acgtgagcgg gcacggcagg cggcctcctc ctccctctcac ggcacggcag ctacggggga	3720
ttcctttccc accgctcctt cgctttccct tcctcgcccg ccgtaataaa tagacacccc	3780
ctccacaccc tctttcccca acctcgtgtt gttcggagcg cacacacaca caaccagatc	3840
tcccccaaat ccaccgctcg gcacctccgc ttcaaggtag gccgctcgtc ccccccccc	3900
ccccctctct accttctcta gatcggcgtt ccggtccatg gttagggcc ggtagtctta	3960
cttctgttca tgtttgtgtt agatccgtgt ttgtgttaga tccgtgctgc tagcgttcgt	4020
acacggatgc gacctgtacg tcagacacgt tctgattgct aacttgccag tgtttctctt	4080
tggggaatcc tgggatggct ctagccgttc cgcagacggg atcgatttca tgattttttt	4140
tgtttcgttg catagggttt ggtttgccct tttcctttat ttcaatatat gccgtgcact	4200
tgtttgtcgg gtcacttttt catgcttttt tttgtcttgg ttgtgatgat gtggtctggt	4260
tgggcggctg ttctagatcg gagtagaatt ctgtttcaaa ctacctggtg gatattataa	4320
ttttggatct gtatgtgtgt gccatacata ttcatagtta cgaattgaag atgatggatg	4380
gaaatatcga tctaggatag gtatacatgt tgatgcgggt tttactgatg catatacaga	4440
gatgcttttt gtccgcttgg ttgtgatgat gtggtgtggt tgggcggctg ttcattcgtt	4500
ctagatcgga gtagaatact gtttcaaact acctggtgta tttattaatt ttggaactgt	4560

-continued

atgtgtgtgt	catacatctt	catagttacg	agtttaagat	ggatggaaat	atcgatctag	4620
gataggtata	catgttgatg	tgggttttac	tgatgcatat	acatgatggc	atatgcagca	4680
tctattcata	tgctctaacc	ttgagtacct	atctattata	ataaacaagt	atgttttata	4740
attattttga	tcttgatata	cttggtgatg	ggcatatgca	gcagctatat	gtggattttt	4800
ttagccctgc	cttcatacgc	tattttattg	cttggtactg	tttcttttgt	cgatgctcac	4860
cctgttgttt	ggtgttactt	ctgcaggctg	actctagagg	atccaacaat	ggagaacaac	4920
atacagaatc	agtgcgctcc	ctacaactgc	ctcaacaatc	ctgaagtaga	gattctcaac	4980
gaagagaggt	cgactggcag	attgcccgtta	gacatctccc	tgtcccttac	acgtttctctg	5040
ttgtctgagt	ttgttccagg	tgtgggagtt	gcgtttggcc	tcttcgacct	catctggggc	5100
ttcatcactc	catctgattg	gagcctcttt	cttctccaga	ttgaacagtt	gattgaacaa	5160
aggattgaga	ccttggaaag	gaatcggggc	atcactaccc	ttcgtggctt	agcagacagc	5220
tatgagatct	acattgaagc	actaagagag	tgggaagcca	atcctaacaa	tgcccaactg	5280
agagaagatg	tgcgatacag	ctttgctaac	acagatgatg	ctttgatcac	agccatcaac	5340
aaactcacc	ttaccagctt	cgagatccct	cttctctcgg	tctatgttca	agctgctaac	5400
ctgcacttgt	cactactgcg	cgacgctgtg	tcgtttgggc	aaggttgggg	actggacata	5460
gctactgtca	acaatcacta	caacagactc	atcaatctga	ttcatcgata	cacgaaacat	5520
tgtttgata	cctacaatca	gggattggag	aacctgagag	gtactaacac	tcgccaatgg	5580
gccaggttca	atcagttcag	gagagacctt	acacttactg	tgtagacat	agttgctctc	5640
tttccgaact	acgatgttcg	tacctatccg	attcaaacgt	catcccaact	tacaagggag	5700
atctacacca	gttcagtcct	tgaagactct	ccagtttctg	cgaacatacc	caatgggttc	5760
aacagggctg	agtttgaggt	cagaccaccc	catctcatgg	acttcatgaa	ctctttgttt	5820
gtgactgcag	agactgttag	atcccaaact	gtgtggggag	gacacttagt	tagctcacgc	5880
aacacggctg	gcaatcgtat	caactttcct	agttacgggg	tcttcaatcc	cgggggcgcc	5940
atctggattg	cagatgaaga	tccacgtcct	ttctatcgga	ccttgtcaga	tcctgtcttc	6000
gtccgaggag	gctttggcaa	tcctcactat	gtactcggtc	ttaggggagt	ggcctttcaa	6060
caaactggta	cgaatcacac	ccgcacattc	aggaaactcc	ggaccattga	ctctctagat	6120
gagataccac	ctcaagacaa	cagcggcgca	ccttggaatg	actactccca	tgtgctgaat	6180
catgttacct	ttgtgcgctg	gccaggtgag	atctcaggtt	ccgactcatg	gagagcacca	6240
atgttctctt	ggacgcatcg	tagcgctacc	cccacaaaca	ccattgatcc	agagagaatc	6300
actcagattc	ccttgggtgaa	ggcacacaca	cttcagtcag	gaactacagt	tgtaagaggg	6360
ccgggggttca	cgggaggaga	cattcttcga	cgcactagtg	gaggaccatt	cgcgtacacc	6420
attgtcaaca	tcaatgggca	acttcccaa	aggatcgtg	ccaggatagc	ctatgcctct	6480
actaccaatc	taagaatcta	cgttacggtt	gcaggtgaac	ggatctttgc	tggtcagttc	6540
aacaagacaa	tggataccgg	tgatccactt	acattccaat	ctttctccta	cgcactatc	6600
aacaccgcgt	tcacctttcc	aatgagccag	agcagtttca	cagtaggtgc	tgataccttc	6660
agttcaggca	acgaagtgtg	cattgacagg	tttgagttga	ttccagttac	tgccacactc	6720
gagtaaggat	ccgtcgacct	gcagccaagc	tttcgcgagc	tcgagatccc	cgacatatgc	6780
cccggtttctg	ttgcgactaa	catgagttct	tggacaaaatt	tgattggacc	tgatgagatg	6840

-continued

atccaacccg	aggatatagc	aaagctcgtt	cgtgcagcaa	tggaacggcc	aaaccgtgct	6900
tttgtcccca	agaatgaggt	gctatgcatg	aaggaatcta	cccgttgatg	tccaacagtc	6960
tcagggttaa	tgtctatgta	tcttaataa	tgttgctcgt	atthttgtaat	ctcatataga	7020
ttttcactgt	gcgacgcaaa	aatattaaat	aatattatt	attatctacg	ttttgattga	7080
gatatcatca	atattataat	aaaaatatcc	attaaacacg	atthtgatata	aatgacagtc	7140
aataatctga	tttgaatatt	tattaattgt	aacgaattac	ataaagatcg	aatagaaaat	7200
actgcactgc	aatgaaaat	taacacatac	taataaatgc	gtcaaatatc	tttgccaaga	7260
tcaagcggag	tgagggcctc	atatccggtc	tcagttacaa	gcacggatc	cccgaagcgc	7320
gctccaccaa	tgccctcgac	atagatgccg	ggctcgacgc	tgaggacatt	gcctaccttg	7380
agcatgggtc	cagcgccggc	tttaagctca	atcccatccc	aatctgaata	tcctatcccg	7440
cgccacgtcc	gggtgaagaa	cgggtctgtc	catccacctc	tgthgggaat	tcgggtccgg	7500
gtcacctttg	tccaccaaga	tggaactgcg	gccgcggacc	gaattcccat	ggagtcaaa	7560
attcaaatag	aggacctaac	agaactcgcc	gtaaagactg	gcgaacagtt	catacagagt	7620
ctcttacgac	tcaatgacaa	gaagaaaatc	ttcgtcaaca	tggtggagca	cgacacgctt	7680
gtctactcca	aaaatatcaa	agatacagtc	tcagaagacc	aaagggcaat	tgagactttt	7740
caacaaaggg	taatatccgg	aaacctcctc	ggattccatt	gccacgtat	ctgtcacttt	7800
attgtgaaga	tagtggaaaa	ggaaggtggc	tcctacaaat	gccatcattg	cgataaagga	7860
aaggccatcg	ttgaagatgc	ctctgccgac	agtggtccca	aagatggacc	cccacccacg	7920
aggagcatcg	tggaaaaaga	agacgttcca	accacgtctt	caaagcaagt	ggattgatgt	7980
gatatctcca	ctgacgtaag	ggatgacgca	caatcccat	atccttcgca	agacccttcc	8040
tctatataag	gaagttcatt	tcatttgag	aggacaggt	accggggat	ccaccatgtc	8100
tcgggagagg	agaccagttg	agattaggcc	agctacagca	gctgatattg	ccgcggtttg	8160
tgatatcgtt	aaccattaca	ttgagacgtc	tacagtgaac	tttaggacag	agccacaaac	8220
accacaagag	tggattgatg	atctagagag	gttgcaagat	agataccott	ggttggttgc	8280
tgaggttgag	gggtgtgtg	ctggatttgc	ttacgctggg	ccctggaag	ctaggaaacgc	8340
ttacgatttg	acagttgaga	gtactgttta	cgtgtcacat	aggcatcaaa	ggttgggcct	8400
aggatccaca	ttgtacacac	atttgcttaa	gtctatggag	gcgcaaggtt	ttaagtctgt	8460
ggttgctgtt	ataggccttc	caaacgatcc	atctgttagg	ttgcatgagg	ctttgggata	8520
cacagcccg	ggtagattgc	gcgcagctgg	atacaagcat	ggtaggatggc	atgatgttgg	8580
tttttgcaa	agggattttg	agttgccagc	tcctccaagg	ccagttaggc	cagttaccca	8640
gatctgagtc	gacctgcagg	catgcccgct	gaaatcacca	gtctctctct	acaaatctat	8700
ctctctctat	aataatgtgt	gagtagttcc	cagataagg	aattaggggt	cttatagggt	8760
ttcgtctatg	tgttgagcat	ataagaaacc	cttagtatgt	atthgtatth	gtaaaatact	8820
tctatcaata	aaattttctaa	ttcctaaaac	caaaatccag	tggcgagctc	gaattcgagc	8880
tcgagcccg	gtggatctc	tagagtcgac	ctgcagaagc	ttcgggtccg	cgcgctctta	8940
gttgaagaca	cgttcatgtc	ttcatcgtaa	gaagacactc	agtagtcttc	ggccagaatg	9000
gcctaactca	aggccctcac	tccgcttgat	cttgccaaag	atatttgacg	cattttattag	9060
tatgtgttaa	ttttcatttg	cagtgcagta	ttttctattc	gatctttatg	taattcgtta	9120

-continued

caattaataa atattcaaat cagattattg actgtcattt gtatcaaatac gtgtttaatg	9180
gatatatttta ttataatatt gatgatattt caatcaaaac gtagataata ataataattta	9240
tttaataattt ttgcgtcgca cagtgtaaaat ctatatgaga ttacaaaata ccgacaacat	9300
tatttaagaa acatagacat taaccctgag actgttggac atcaacgggt agattccttc	9360
atgcatagca cctcattctt ggggacaaaa gcacggtttg gccgttccat tgctgcacga	9420
acgagccttg ctatatcctc ggggttgatc atctcatcag gtccaatcaa atttgtccaa	9480
gaactcatgt tagtcgcaac gaaaccgggg catatgtcgg gtatctcgag ctgcgaaaag	9540
cttggtcgca ggtcgacgga tccttcaaca aaagggtacc tgtaccgaa accgacacag	9600
gtgggtaggt agagaatacc tagggcgcg agacaactct ctctaaggaa ctcgcaaaaa	9660
tagccccgta acttcgggag aagggtgcc ccccgctaac aataaacgaa tacggtttat	9720
gtatggattc cggtaaaata ccggtactcg atttcataag agtcgaatag gaagttaaga	9780
tgagggtggt atcatcataa aaatggagta gtatcctaaa ttataactaat ccacgtatga	9840
tatgtatgcc ttctcttacc aaccggaagt agtgcaaaaa aaattctata ctgcactgct	9900
ctctttttac tgagaaatgc aaaaaataa aagtgaagta agggtgcccc atagatattt	9960
gatcttgctt cctgtccccc cccccctttt ttcatacaaa atttcctatga aaaaagaaaa	10020
gatgaatttg tccattcatt gaaccctagt tcgggactga cggggctcga acccgagct	10080
tccgcctgtt cctagccttc cagggccag cgtaagcaat accagccaca gcaccctcaa	10140
cctcagcaac caaccaaggg tatctatctt gcaacctctc tagatcatca atccactctt	10200
gtggtgtttg tggtctgttc cttaaagtca ctgtagacgt ctcaatgtaa tggttaacga	10260
tatcacaaac cgcggaacac aagaacgaaa gcaccttttc attctttcat atactagggg	10320
tttttacttg gaaaagacaa tgttcatac taaaggatag ctgcagaagc cgcacccgtc	10380
ttgaggacct tccggggagc cagaccggtc gaaccgtgcc tccacttgct aaggagaaa	10440
ggaaaatcag ggccaggaca tacgaaggag gagccagaac gaagatatcc taagatactt	10500
actcgtctcg ggccatgac aatcatgcct gtggggagggt ctctcgacc tcgatccatg	10560
aaggtaccac cgaggctgc cccgcgcgcg gcttcggtac cgtcctcgcc ttgggcgccc	10620
gaggcaccgc ggggatggac tgcccaggcg cagccacgac gacccaagga tcaccctcct	10680
gcgcagtcgg cagcagcaat agttctcggg gaacaggcag cttggcctga ctccccggg	10740
tcacctcaac tacctcggcc gaggggtcaa gtacccctc agtcgcgccc cgtctctcgg	10800
accgggaccc cgacgtcccg gccccggata ccgacggcac cagcccgctc gggggctggc	10860
ttgacgaccc ctggcccagc ctcatatctg ggctgaggcc gaggcaggcg gccatgtcgt	10920
cgtcttcatc atcgtcttca tcacgtcgtc cgtcatcagg cgtctccggc gacggctccc	10980
ttgggagccc ctccctctcc tgccgacgac gaagcctttc caaggcatcc cgagcccacg	11040
tccgctcgtg ggcccagacc ttctttgctt ccttcttctc cttcctcttc tccgcgggtga	11100
ccctccgcgc agctcgggtc accgcaccc cggggactgg tggcagggaa ggcttgtgat	11160
gccctacctc ctggagacag acgaaaagtc tcagctatga gaaccgagg caatctgacg	11220
caagaaggaa gaaggagcgg atactacca gagacacgca cccgcgatcg ggacgcatta	11280
agggctggga aaaagtgcgc gcctctaatt tcgtaccgt gccgtccacc cactgtgga	11340
ggatcatcat gggaaggga a	11361

-continued

<210> SEQ ID NO 25
<211> LENGTH: 6186
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: The sequence represents the DNA molecule used to transform maize line TC1507. This sequence represents insert PHI8999A.

<400> SEQUENCE: 25

caactggaag agcgggttacc cggaccgaag cttcggccgg ggcccatcga tatccgcggg 60
catgcctgca gtgcagcgtg acccggtcgt gccctctct agagataatg agcattgcat 120
gtctaagtta taaaaaatta ccacatattt tttttgtcac acttgtttga agtgcagttt 180
atctatcttt atacatatat ttaaaacttta ctctacgaat aatataatct atagtactac 240
aataatatca gtgtttttaga gaatcatata aatgaacagt tagacatggt cttaaaggaca 300
attgagtatt ttgacaacag gactctacag ttttatcttt ttagtgtgca tgtgttctcc 360
tttttttttg caaatagctt cacctatata atacttcato cattttatta gtacatccat 420
ttagggttta ggggtaatgg tttttataga ctaatttttt tagtacatct attttattct 480
atttttagcct ctaaataaag aaaactaaaa ctctatttta gtttttttat ttaataattt 540
agatataaaa tagaataaaa taaagtgact aaaaattaaa caaataccct ttaagaaatt 600
aaaaaaacta aggaacattt tttctgtgtt cgagtagata atgccagcct gttaaacgcc 660
gtcgacgagt ctaacggaca ccaaccagcg aaccagcagc gtcgcgtcgg gccaagcgaa 720
gcagacggca cggcatctct gtcgctgcct ctggaccctt ctcgagagtt ccgctccacc 780
gttggtactg ctccgctgtc ggcatccaga aattgcgtgg cggagcggca gacgtgagcc 840
ggcacggcag gcggcctcct cctcctctca cggcacggca gctacggggg attcctttcc 900
caccgctcct tcgctttccc ttcctcgccc gccgtaataa atagacaccc cctccacacc 960
ctctttcccc aacctcgtgt tgttcggagc gcacacacac acaaccagat ctccccaaaa 1020
tccacccgct ggcaacctcg ctccaaggta cgcgcgtcgt cctccccccc cccccctctc 1080
taccttctct agatggcggt tccggtccat ggttagggcc cggtagttct acttctgttc 1140
atgtttgtgt tagatccgtg tttgtgttag atccgtgctg ctagcgttcg tacacggatg 1200
cgacctgtac gtcagacacg ttctgattgc taacttgcca gtgtttctct ttggggaatc 1260
ctgggatggc tctagccgtt ccgcagacgg gatcgatttc atgatTTTTT ttgtttcgtt 1320
gcatagggtt tggtttgccc ttttccttta tttcaatata tgccgtgcac ttgtttgtcg 1380
ggtcaccttt tcatgctttt ttttgccttg gttgtgatga tgtggctctg ttgggcggtc 1440
gttctagatc ggagtagaat tctgtttcaa actacctggt ggatttatta attttggatc 1500
tgtatgtgtg tgccatacat attcatagtt acgaattgaa gatgatggat ggaaatcgcg 1560
atctaggata ggtatacatg ttgatcgagg ttttactgat gcatatacag agatgctttt 1620
tgttcgcttg gttgtgatga tgtggtgtgg ttgggcggtc gttcattcgt tctagatcgg 1680
agtagaatac tgtttcaaac tacctgggtg atttattaat tttggaactg tatgtgtgtg 1740
tcatacatct tcatagttac gagtttaaga tggatggaaa tatcgatcta ggataggtat 1800
acatgttgat gtgggtttta ctgatgcata tacatgatgg catatgcagc atctattcat 1860
atgctctaac cttgagtacc tatctattat aataaacaag tatgttttat aattattttg 1920

-continued

atcttgatat	acttggatga	tggcatatgc	agcagctata	tgtggatttt	tttagccctg	1980
ccttcatacg	ctatttat	gcttggtagt	gtttcttttg	tcgatgctca	ccctgttgtt	2040
tgggtgttact	tctgcaggtc	gactctagag	gatccaacaa	tggagaacaa	catacagaat	2100
cagtgcgtcc	cctacaactg	cctcaacaat	cctgaagtag	agattctcaa	cgaagagagg	2160
tcgactggca	gattgccgtt	agacatctcc	ctgtccctta	cacgtttcct	gttgtctgag	2220
tttgttccag	gtgtgggagt	tgcgtttggc	ctcttcgacc	tcctctgggg	cttcactcact	2280
ccatctgatt	ggagcctcct	tcttctccag	attgaacagt	tgattgaaca	aaggattgag	2340
acottggaaa	ggaatcgggc	catcactacc	cttcgtggct	tagcagacag	ctatgagatc	2400
tacattgaag	cactaagaga	gtgggaagcc	aatcctaaca	atgcccaact	gagagaagat	2460
gtgcgtatac	gctttgctaa	cacagatgat	gctttgatca	cagccatcaa	caacttcacc	2520
cttaccagct	tcgagatccc	tcttctctcg	gtctatgttc	aagctgctaa	cctgcacttg	2580
tcactactgc	gcgacgtgt	gtcgtttggg	caaggttggg	gactggacat	agctactgtc	2640
aacaatcact	acaacagact	catcaatctg	attcatcgat	acacgaaaca	ttgtttggat	2700
acctacaatc	agggattgga	gaacctgaga	ggtactaaca	ctcgccaatg	ggccagggtc	2760
aatcagttca	ggagagacct	tacacttact	gtgttagaca	tagttgctct	ctttccgaac	2820
tacgatgttc	gtacctatcc	gattcaaacg	tcattcccaac	ttacaaggga	gatctacacc	2880
agttcagtca	ttgaagactc	tccagtttct	gcgaacatac	ccaatgggtt	caacagggtc	2940
gagtttggag	tcagaccacc	ccatctcatg	gacttcatga	actctttgtt	tgtgactgca	3000
gagactgtta	gatcccaaac	tgtgtgggga	ggacacttag	ttagctcacg	caacacggct	3060
ggcaatcgta	tcaactttcc	tagttacggg	gtcttcaatc	ccgggggcgc	catctggatt	3120
gcagatgaag	atccacgtcc	tttctatcgg	acottgtcag	atcctgtcct	cgtccgagga	3180
ggctttggca	atcctcacta	tgtactcggg	cttaggggag	tggcctttca	acaaactggt	3240
acgaatcaca	cccgcacatt	cagggaactcc	gggaccattg	actctctaga	tgagatacca	3300
cctcaagaca	acagcgggcg	acottggaat	gactactccc	atgtgctgaa	tcattgttacc	3360
tttgtgcgct	ggccagggtg	gatctcaggt	tccgactcat	ggagagcacc	aatgttctct	3420
tggacgcato	gtagcgctac	ccccacaaac	accattgato	cagagagaat	cactcagatt	3480
ccottggtga	aggcacacac	acttcagtca	ggaactacag	ttgtaagagg	gccgggggtc	3540
acgggaggag	acattctctg	acgcactagt	ggaggaccat	tcgcgtacac	cattgtcaac	3600
atcaatgggc	aacttcccca	aaggatcgt	gccaggatag	gctatgcctc	tactaccaat	3660
ctaagaatct	acgttacggg	tgcagggtgaa	cggatctttg	ctggtcagtt	caacaagaca	3720
atggataccg	gtgatccact	tacattccaa	tctttctcct	acgccactat	caacaccgcg	3780
ttcacccttc	caatgagcca	gagcagtttc	acagtaggtg	ctgatacctt	cagttcaggc	3840
aacgaagtgt	acattgacag	gtttgagttg	attccagtta	ctgccacact	cagtaagga	3900
tccgtcgacc	tgcagccaag	cttttcgcga	gctcgagatc	cccacatat	gccccgggtt	3960
cgttgcgact	aacatgagtt	cttgacaaaa	tttgattgga	cctgatgaga	tgatccaacc	4020
cagagatata	gcaaagctcg	ttcgtgcagc	aatggaacgg	ccaaaccgtg	cttttgtccc	4080
caagaatgag	gtgctatgca	tgaaggaaac	taccggttga	tgtccaacag	tctcagggtt	4140
aatgtctatg	tatcttaaat	aatgttgtcg	gtattttgta	atctcatata	gattttcact	4200

-continued

gtgcgacgca	aaaatattaa	ataaatatta	ttattatcta	cgttttgatt	gagatatcat	4260
caatattata	ataaaaatat	ccattaaaca	cgatttgata	caaatagacag	tcaataatct	4320
gatttgaata	tttattaatt	gtaacgaatt	acataaagat	cgaatagaaa	atactgcact	4380
gcaaatgaaa	attaacacat	actaataaat	gcgtcaaata	tctttgccaa	gatcaagcgg	4440
agtgagggcc	tcatatccgg	tctcagttac	aagcacggta	tcccgaagc	gcgctccacc	4500
aatgccctcg	acatagatgc	cgggctcgac	gctgaggaca	ttgcctacct	tgagcatggt	4560
ctcagcgccg	gctttaagct	caatcccac	ccaatctgaa	tatcctatcc	cgcgcccagt	4620
ccggtgtaag	aacgggtctg	tccatccacc	tctgttggga	attccggtcc	gggtcacctt	4680
tgtccaccaa	gatggaactg	cggccgcgga	ccgaattccc	atggagtcaa	agattcaaat	4740
agaggaccta	acagaactcg	ccgtaaagac	tggcgaacag	ttcatacaga	gtctcttacg	4800
actcaatgac	aagaagaaaa	tcttcgtcaa	catggtggag	cacgacacgc	ttgtctactc	4860
caaaaatatc	aaagatacag	tctcagaaga	caaagggca	attgagactt	ttcaacaaag	4920
ggtaatatcc	ggaaacctcc	tcggattcca	ttgccagct	atctgtcact	ttattgtgaa	4980
gatagtggaa	aagggaagtg	gctcctacaa	atgccatcat	tgcgataaag	gaaaggccat	5040
cgttgaagat	gcctctgccg	acagtgggtc	caaagatgga	ccccaccca	cgaggagcat	5100
cgtggaaaaa	gaagacgttc	caaccacgtc	ttcaaagcaa	gtggattgat	gtgatattct	5160
cactgacgta	agggatgacg	cacaatccca	ctatccttcg	caagaccctt	cctctatata	5220
aggaagtcca	tttcatttgg	agaggacag	gtaccgggg	atccaccatg	tctccggaga	5280
ggagaccagt	tgagattagg	ccagctacag	cagctgatat	ggccgcgggt	tgtgatatcg	5340
ttaaccatta	cattgagacg	tctacagtga	actttaggac	agagccacaa	acaccacaag	5400
agtggattga	tgatctagag	aggttgcaag	atagataccc	ttggttggtt	gctgaggttg	5460
aggggtgtgt	ggctggtatt	gcttacgctg	ggccctggaa	ggctaggaac	gcttacgatt	5520
ggacagttga	gagtactggt	tacgtgtcac	ataggcatca	aaggttgggc	ctaggatcca	5580
cattgtacac	acatttgctt	aagtctatgg	aggcgcaagg	ttttaagtct	gtggttgctg	5640
ttataggcct	tccaaacgat	ccatctgtta	ggttgcatga	ggctttggga	tacacagccc	5700
gggttacatt	gcgcgcagct	ggatacaagc	atggtggatg	gcatgatggt	ggtttttggc	5760
aaagggattt	tgagttgcc	gtcctcccaa	ggccagttag	gccagttacc	cagatctgag	5820
tcgacctgca	ggcatgccgc	tgaatcacc	agtctctctc	tacaaatcta	tctctctcta	5880
taataatgtg	tgagtagttc	ccagataaag	gaattagggt	tcttatagg	tttcgctcat	5940
gtgttgagca	tataagaaac	ccttagtatg	tatttgtatt	tgtaaaatac	ttctatcaat	6000
aaaatttcta	attcctaaaa	ccaaaatcca	gtggcgagct	cgaattcgag	ctcgagcccc	6060
ggtggtacct	ctagagtcga	cctgcagaag	cttcgggtccg	gcgcgcctct	agttgaagac	6120
acgttcatgt	cttcacgta	agaagacact	cagtagtctt	cggccagaat	ggcctaactc	6180
aaggcc						6186

<210> SEQ ID NO 26

<211> LENGTH: 3830

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Sequence that represents part of the PHI8999A insert as well as flanking sequence 5' to the

-continued

insert.

<400> SEQUENCE: 26

actagtttcc tagcccgcgt cgtgccccta cccaccgcac gtttatggaa ggtgccattc	60
cacggttctt cgtggccgcc cctaaggatg taaatggtcg gtaaaatccg gtaaatttcc	120
ggtaccgttt accagatttt tccagccgtt ttcggattta tcgggatata cagaaaacga	180
gacggaaacg gaataggttt tttttcgaaa acggtacggt aaacggtgag acaaacttac	240
cgtccgtttt cgtattttctc gggaaactct ggtatatattcc cgtatttgtc ccgtattttc	300
ccgacccacg gacctgccaa tcaaccatca gccagtcagc ccatccccc acgtatggcc	360
catggggcca tgttggccac atgccacgc aacgcaaggc agtaaggctg gcagcctggc	420
acgcattgac gcattgtggac acacacagcc gccgcctgtt cgtgtttctg tgccgttggtg	480
cgagactgtg actgcgagtg gcggagtcgg cgaacggcga ggcgtctccg gagtctggac	540
tcgggctgtg gacagcgacg ctgtgacggc gactcggcga agccccaagc taccaagccc	600
ccaagtcccc atccatctct gcttctctgg tcatctcctt cccctggtcg atctgcaggc	660
gccagaccgg ccgaagcatc acgaaacgca ctaagacctc gaaggagtca aaccactcct	720
ccgaggcctc gggggctaca cccggcgggt gcgctcgcgc gcacccaccg gaacaaaatg	780
taaccgagaa aggtcggctc ccttgcaaaa aaagtgcgac aaaagcctcc aagcgagtat	840
taaacctcac tttgaggctc gggggctact gtcggggacc ataattaggg gtacccccaa	900
gactcctaata ctacgtcgtt aacccccatc agcacaaaag tgcaaaggcc tgatgggtgc	960
gattaagtca aggtcggctc cactcaaggg acacgatctc gcctcgccc agcccagcct	1020
cgggcaaggg cgcccgaccc cgaggattca cgtctcgccc gagggccccc tcaagcgacg	1080
ggcacacctt cggctcgccc gagggccatt ctctcgccgag aagcaacctt ggccagatcg	1140
ccacaccgac cgaccgtatc gcaggagcat ttaatgcgag gatcgctga cacttatcc	1200
tgacgcgcgc tcttcagtcg acagagccga agtgaccgca atcacttcgc cgctccactg	1260
accgacctga caagaagaca gcgccgctg cgtcgctccg actgctgtgc cactcgacag	1320
agtgaggctg acagcagcca agtcggcctt cgggcgccat aggaagctcc gcctcgccc	1380
accctagggc tcggactcgg cctcggtcc ggaagacgac gaactacgct tcgcccgacc	1440
ccagggttg gactcagcct cggctccgga agacgacgaa ttccgcctcg cccgacccca	1500
gggctcggac tcggcctcgg ctccagaaga cgacgaactc cgcctcgccc gacccaggg	1560
ctcggactca gcctcggctc cggaagacga cgaactccgc ctcgcccgac cccagggtc	1620
ggactcagcc tcggcctcag acgatgggtc cgcctcgcc cgacccgggg ctcgactcg	1680
acctttctat cggaccttgt cagatcctgt ctctgctcca ggaggctttg gcaatcctca	1740
ctatgtactc ggtcttaggg gagtgccctt tcaacaaact ggtacgaatc acaccgcac	1800
attcaggaac tccgggacca ttgactctct agatgagata ccacctcaag acaacagcgg	1860
cgcaccttg aatgactact cccatgtgct gaatcatgtt acctttgtgc gctggccagg	1920
tgagatctca ggttcgact catggagagc accaatgttc tcttgagcgc atcgtagcgc	1980
tacccccaca aacaccattg atccagagag aatcactcat tcttcaagaa ctgcatact	2040
tgccgagatc ctcatcccta aaggtacttg acaatagtat tattggagtc gatacacaac	2100
tcacaaaaaa tacaagaagt cgactagggt gattggtccg agtgaagaga aaaaaagcc	2160

-continued

atacagaact caaaatcttt tccggagata ttcattttcc tgaagaggcg gataagatat	2220
taggtggcag tttgatacca ccagaaagag aaaaaaaga ttctaaggaa tcaaaaaaa	2280
ggaaaaattg ggtttatgtt caacggaaaa aatttctcaa aagcaaggaa aagtattgtg	2340
gctattttatc tatccgtgca gctgatatgg ccgcgggttg tgatatcgtt aaccattaca	2400
ttgagacgtc tacagtgaac tttaggacag agccacaaac accacaagag tggattgatg	2460
atctagagag gttgcaagat agataccctt ggttggttgc tgagggtgag ggtgtgtg	2520
ctggtattgc ttacgtggg ccttgaagg ctaggaaccc tcaacctcag caaccaacca	2580
atggtatcta tcttgcaacc tctctagatc atcaatccac tcttgtggtg tttgtggctc	2640
tgctctaaag ttcactgtag acgtctcaat gtaatggta acgatatcac aaaccgagag	2700
aagagggatc tcgaagcttc ggccggggcc catcgatata cgccggcatg cctgcagtgc	2760
agcgtgaccc ggtcgtgccc ctctctagag ataatgagca ttgcatgtct aagttataaa	2820
aaattaccac aactggaaga gcggttaccc ggaccgaagc ttcggccggg gcccatcgat	2880
atccgcgggc atgcctgcag tgcagcgtga cccggtcgtg cccctctcta gagataatga	2940
gcattgcatg tctaagttat aaaaaattac cacatatctt ttttgtcaca cttgtttgaa	3000
gtgcagttta tctatcttta tacatatatt taaactttac tctacgaata atataatcta	3060
tagtactaca ataatatcag tgttttagag aatcatataa atgaacagtt agacatggtc	3120
taaaggacaa ttgagtattt tgacaacagg actctacagt tttatctttt tagtgtgcat	3180
gtgttctcct ttttttttgc aaatagcttc acctatataa tacttcatcc attttattag	3240
tacatccatt tagggtttag ggtaaatggt ttttatagac taattttttt agtacatcta	3300
ttttattcta ttttagcctc taaattaaga aaactaaaac tctatttttag tttttttatt	3360
taataattta gatataaaat agaataaaat aaagtgacta aaaattaaac aaataccctt	3420
taagaaatta aaaaaactaa ggaacattt ttcttgtttc gagtagataa tgccagcctg	3480
ttaaacgccg tcgacgagtc taacggacac caaccagcga accagcagcg tcgctcgagg	3540
ccaagcgaag cagacggcac ggcattctctg tcgctgcctc tggaccctc tcgagagttc	3600
cgtccaccg ttggacttgc tccgtgtcgc gcattccagaa attcgtggc ggagcggcag	3660
acgtgagccg gcacggcagg cggcctctc ctccctctcac ggcacggcag ctacggggga	3720
ttcctttccc accgctcctt cgctttccct tccctgcccg ccgtaataaa tagacacccc	3780
ctccacaccc tctttcccca acctcgtgtt gttcggagcg cacacacaca	3830

<210> SEQ ID NO 27

<211> LENGTH: 3347

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Sequence that represents part of the PHI8999A insert as well as flanking sequence 3' to the insert.

<400> SEQUENCE: 27

cccactatcc ttcgcaagac ccttcctcta tataaggaag ttcatttcat ttggagagga	60
caggggtaccc ggggatccac catgtctccg gagaggagac cagttgagat taggccagct	120
acagcagctg atatggccgc ggtttgtgat atcgtaaac attacattga gacgtctaca	180
gtgaacttta ggacagagcc acaaacacca caagagtgga ttgatgatct agagaggttg	240

-continued

caagatagat acccttggtt ggttgctgag gttgaggggtg ttgtggctgg tattgcttac	300
gctgggccct ggaaggctag gaacgcttac gattggacag ttgagagtac tgtttaoctg	360
tcacataggc atcaaaggtt gggcctagga tccacattgt acacacattt gcttaagtct	420
atggaggcgc aaggttttaa gtctgtggtt gctgttatag gccttccaaa cgatccatct	480
gttaggttgc atgaggcttt gggatacaca gcccggggta cattgcgcgc agctggatac	540
aagcatggtg gatggcatga tgttggtttt tggcaaaggg attttgagtt gccagctcct	600
ccaaggccag ttaggccagt taccagatc tgagtcgacc tgcaggcatg cccgctgaaa	660
tcaccagtct ctctctacaa atctatctct ctctataata atgtgtgagt agttcccaga	720
taagggaatt aggggttcta tagggtttcg ctcatgtgtt gagcatataa gaaaccctta	780
gtatgtatgtt gtatttgtta aatacttcta tcaataaaat ttctaattcc taaaacccaa	840
atccagtggc gagctcgaat tcgagctcga gcccggtggg atcctctaga gtcgacctgc	900
agaagcttcg gtccggcgcg cctctagttg aagacacgtt catgtcttca tcgtaagaag	960
acactcagta gtcttcggcc agaatggcct aactcaaggc cctcactccg cttgatcttg	1020
gcaaagatat ttgacgcatt tattagtatg tgttaatttt catttgcagt gcagtatgtt	1080
ctattcgcgc tttatgtaat tcgttacaat taataaatat tcaaatcaga ttattgactg	1140
tcatttgtat caaatcgtgt ttaatggata tttttattat aatattgatg atatctcaat	1200
caaaacgtag ataataataa tattttattt atatttttgc gtgcacagc gaaaatctat	1260
atgagattac aaaataccga caacattatt taagaaacat agacattaac cctgagactg	1320
ttggacatca acgggtagat tccttcacgc atagcacctc attcttgggg acaaaagcac	1380
ggtttgcccg ttccattgct gcacgaacga gctttgctat atcctcgggt tggatcatct	1440
catcagggtc aatcaaatgt gtccaagaac tcatgttagt cgcaacgaaa ccggggcata	1500
tgctgggtat ctcgagctcg cgaagcctt gctgcaggtc gacggatcct tcaacaaaag	1560
ggtacctgta ccgaaaccg acacaggtgg gtaggtagag aatacctagg ggcgcgagac	1620
aactctctct aaggaaactc gcaaaatagc cccgtaactt cgggagaagg ggtgcccccc	1680
gctaacaata aacgaatagc gtttatgtat ggattccggt aaaataccgg tactcgattt	1740
cataagagtc gaataggaag ttaagatgag ggtggtatca tcataaaaat ggagtagtat	1800
cctaaattat actaatccac gtatgatatg tatgccttct cttatcaacc ggaagtagtg	1860
caaaaaaaat tctatactgc actgctctct ttttactgag aaatgcaaaa aaataaaagt	1920
gaagtaaggg tgcccatag atatttgatc ttgcctcctg tccccccccc ctttttttca	1980
tcaaaaaatt ccatgaaaaa agaaaagatg aatttgtcca ttcatgaaac cctagttcgg	2040
gactgacggg gctcgaaccc gcagcttccg cctgttccta gccttccagg gccagcgta	2100
agcaatacca gccacagcac cctcaacctc agcaaccaac caagggtatc tatcttgcaa	2160
cctctctaga tcatcaatcc actcttggtg tgtttgtggc tctgtcctaa agttcactgt	2220
agacgtctca atgtaatggt taacgatatc acaaaccgcg gaacacaaga acgaaagcac	2280
cttttcatcc tttcatatac taggggtttt tacttggaag agacaatgtt ccatactaaa	2340
ggatagctgc agaagccgcc accgtcttga ggaccttccg gggagccaga ccggtcgaac	2400
cgtgcctcca cttgctaagg agaaagggaa aatcagggcc aggcatacag aaggaggagc	2460
cagaacgaag atatcctaag atacttactc gctccgggcc atgatcaatc atgcctgtgg	2520

-continued

ggaggtctct cgcacctcga tccatgaagg taccaccgag gtctgccccg ccgccggctt	2580
cggtaccgtc ctgccttgg gcgccgagg caccggggg atggactgcc caggcgcagc	2640
cacgacgacc caaggatcac cctcctgcgc agtcggcacg agcaatagtt ctcggggaac	2700
aggcagcttg gcctgactcc ccggggtcac ctcaactacc tcggccgagg ggtcaagtac	2760
ccctcagtc cgcctccgct ctctggaccg ggaccccgac gtcccgcccc cggataccga	2820
cggcaccagc ccgctcgggg gctggcttga cgacccctgg ccagcctca gatctgggct	2880
gaggccgagg caggcgccca tgcgtcgtc ttcatcatcg tcttcatcat cgtcgtcgtc	2940
atcaggcgtc tccggcgacg gctcccttgg gagccctcc ctctcctgcc gacgacgaag	3000
cctttccaag gcaccccgag ccacagtcct ctctggggcc cgagccttct ttgcgtcctt	3060
cttctccttc ctcttctccg cggtagacct ccgcgagct cggtcaccg cctcctccgg	3120
gactggtggc aggggaaggct tgtgatgcc tacctcctgg agacagacga aaagtctcag	3180
ctatgagaac cgagggcaat ctgacgcaag aagggaagag gagcggatac tcaccagaga	3240
cacgcacccg cgatcgggac gcattaaggg ctgggaaaaa gtgccggcct ctaatttcgc	3300
taccgtgccg tccaccacc tgtggaggtc atcgatggga aggggaa	3347

<210> SEQ ID NO 28

<211> LENGTH: 669

<212> TYPE: DNA

<213> ORGANISM: Zea mays

<400> SEQUENCE: 28

actagtttcc tagcccgct cgtgccccta cccaccgac gtttatggaa ggtgccattc	60
cacggttctt cgtggccgcc cctaaggatg taaatggctg gtaaaatccg gtaaatcttc	120
ggtaccgttt accagatttt tccagcgtt ttcggattta tcgggatata cagaaaacga	180
gacggaaacg gaataggttt tttttcgaac acggtacggt aaacggtgag acaaacttac	240
cgtccgtttt cgtatttctc gggaaactct ggtatattcc cgtatttgct ccgtattttc	300
ccgacccacg gacctgcaa tcaaccatca gccagtcagc ccatccccc acgtatggcc	360
catggggcca tgttggccac atgccacgc aacgcaaggc agtaaggctg gcagcctggc	420
acgcattgac gcatgtggac acacacagcc gccgcctgtt cgtgtttctg tgccgttgtg	480
cgagactgtg actgcagtg gcggagtcgg cgaacggcga ggctctccg gactctggac	540
tgccgctgtg gacagcgacg ctgtgacggc gactcggcga agccccaagc taccaagccc	600
ccaagtcccc atccatctct gcttctctgg tcatctcctt cccctggctg atctgcaggc	660
gccagaccg	669

<210> SEQ ID NO 29

<211> LENGTH: 200

<212> TYPE: DNA

<213> ORGANISM: Zea mays

<400> SEQUENCE: 29

gccgaagcat cacgaaacgc actaagacct cgaaggagtc aaaccactcc tccgaggcct	60
cgggggctac acccgggcgg tgccgtcgcg cgcacccacc ggaacaaaat gtaaccgaga	120
aaggctcggt cccttgcaaa aaaagtgcga caaaagcctc caagcgagta ttaacactca	180
ctttgaggct cgggggctac	200

-continued

```

<210> SEQ ID NO 30
<211> LENGTH: 812
<212> TYPE: DNA
<213> ORGANISM: Zea mays
<220> FEATURE:
<223> OTHER INFORMATION: Fragment of maize Huck-1 retrotransposon

<400> SEQUENCE: 30
tgtcggggac cataattagg ggtaccccca agactcctaa tctcagctgg taaccccat    60
cagcacaaag ctgcaaaggc ctgatgggtg cgattaagtc aaggctcggc cactcaagg    120
gacacgatct cgcctcgccc gagccagcc tcgggcaagg gcggccgacc ccgaggattc    180
acgtctcgcc cgaggggccc ctcaagcgac gggcacacct tcggctcgcc cgaggcccat    240
tcttcgccga gaagcaacct tggccagatc gccacaccga ccgaccgtat cgcaggagca    300
tttaatgcga ggatcgctg acaccttatt ctgacgcgcg ctcttcagtc gacagagccg    360
aagtgaaccg aatcacttcg ccgctccact gaccgacctg acaagaagac agcgccgcct    420
gcgtcgctcc gactgctgtg cactcgaca gagtgaggct gacagcagcc aagtccggcc    480
tcgggcgcca taggaagtc cgcctcgccc gaccctaggg ctcggaactc gcctcggtc    540
cggaagacga cgaactacgc ttcgcccgc cccagggtt ggactcagcc tcggctccgg    600
aagacgacga attccgcctc gcccgacccc agggctcgga ctcggcctcg gctccagaag    660
acgacgaact ccgcctcgcc cgacccaggg gctcggactc agcctcggtt ccggaagacg    720
acgaactccg cctcgccga cccaggggt cggactcagc ctcggcctca gacgatggtc    780
tccgcctcgc ccgacccggg gctcggactc ga                                812

```

```

<210> SEQ ID NO 31
<211> LENGTH: 335
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Represents part of PHI8999A insert sequence -
        fragment of cryIF gene

```

```

<400> SEQUENCE: 31
cctttctatc ggacctgtc agatcctgtc ttcgtccgag gaggttttg caatcctcac    60
tatgtactcg gtcttagggg agtggccttt caacaaactg gtacgaatca caccgcaca    120
ttcaggaact ccgggacat tgactctcta gatgagatac cacctcaaga caacagcggc    180
gcaccttgga atgactactc ccatgtgctg aatcatgtta cctttgtgcg ctggccaggc    240
gagatctcag gttccgactc atggagagca ccaatgttct cttggacgca tcgtagcgct    300
acccccacaa acaccattga tccagagaga atcac                                335

```

```

<210> SEQ ID NO 32
<211> LENGTH: 321
<212> TYPE: DNA
<213> ORGANISM: Zea mays
<220> FEATURE:
<223> OTHER INFORMATION: Fragment of maize chloroplast rpoC2 gene

```

```

<400> SEQUENCE: 32
tcattcttca agaactgcat atcttgccga gatcctcatc cctaaaggta cttgacaata    60
gtattattgg agtcgataca caactcacia aaaatacaag aagtcgacta ggtggattgg    120
tccgagtga gagaaaaaaa agccatacag aactcaaat cttttccgga gatattcatt    180

```

-continued

```

ttcctgaaga ggcggataag atattaggtg gcagtttgat accaccagaa agagaaaaaa    240
aagattctaa ggaatcaaaa aaaaggaaaa attgggttta tgttcaacgg aaaaaatttc    300
tcaaaagcaa ggaaaagtat t                                              321

```

```

<210> SEQ ID NO 33
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Represents part of PHI8999A insert
sequence-fragment of ubiZM1(2) promoter; also a
fragment of the maize chloroplast trn1 gene

```

```

<400> SEQUENCE: 33
gtggctatct atctatc                                              17

```

```

<210> SEQ ID NO 34
<211> LENGTH: 201
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Represents part of PHI8999A insert sequence -
fragment of pat gene

```

```

<400> SEQUENCE: 34
gcagctgata tggccgcggt ttgtgatatc gttaaccatt acattgagac gtctacagtg    60
aaccttagga cagagccaca aacaccacaa gagtggattg atgatctaga gaggttgcaa    120
gatagatacc cttggttggt tgctgaggtt gaggggtgtg tggctggtat tgcttacgct    180
gggccctgga aggctaggaa c                                          201

```

```

<210> SEQ ID NO 35
<211> LENGTH: 138
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Represents part of PHI8999A insert sequence -
fragment of pat gene (complement)

```

```

<400> SEQUENCE: 35
cctcaacctc agcaaccaac caatggtatc tatcttgcaa cctctctaga tcatcaatcc    60
actcttgtag tgtttgtggc tctgtcctaa agttcactgt agacgtctca atgtaatggt    120
taacgatatc acaaaccg                                              138

```

```

<210> SEQ ID NO 36
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Represents part of PHI8999A insert sequence -
fragment of cry1F gene (complement)

```

```

<400> SEQUENCE: 36
agagaagagg gatct                                              15

```

```

<210> SEQ ID NO 37
<211> LENGTH: 118
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

```

-continued

<223> OTHER INFORMATION: Represents part of PHI8999A insert sequence -
fragment of polylinker

<400> SEQUENCE: 37

```
cgaagcttcg gccggggccc atcgatatcc gcgggcatgc ctgcagtga gcgtgacccg      60
gtcgtgcccc tctctagaga taatgagcat tgcattgtcta agttataaaa aattacca      118
```

<210> SEQ ID NO 38

<211> LENGTH: 550

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Represents part of PHI8999A insert sequence -
fragment of ORF25 terminator (complement)

<400> SEQUENCE: 38

```
ctcactccgc ttgatcttgg caaagatatt tgacgcattt attagtatgt gttaattttc      60
atttgcagtg cagtattttc tattcgatct ttatgtaatt cgttacaatt aataaatatt      120
caaatcagat tattgactgt catttgtatc aaatcgtgtt taatggatat ttttattata      180
atattgatga tatctcaatc aaaacgtaga taataataat atttatttaa tttttttgcg      240
tcgcacagtg aaaatctata tgagattaca aaataccgac aacattatct aagaaacata      300
gacattaacc ctgagactgt tggacatcaa cgggtagatt ccttcattga tagcacctca      360
ttcttgggga caaaagcacg gtttggccgt tccattgctg cacgaacgag ctttgcata      420
tcctcggggt ggatcatctc atcagggtcca atcaaatttg tccaagaact catgttagtc      480
gcaacgaaac cggggcatat gtcgggtatc tcgagctcgc gaaagcttgg ctgcaggtcg      540
acggatcctt                                     550
```

<210> SEQ ID NO 39

<211> LENGTH: 128

<212> TYPE: DNA

<213> ORGANISM: Zea mays

<220> FEATURE:

<223> OTHER INFORMATION: Fragment of maize chloroplast rps12 rRNA gene
(complement)

<400> SEQUENCE: 39

```
caacaaaagg gtacctgtac ccgaaaccga cacagggtgg taggtagaga atacctaggg      60
gcgcgagaca actctctcta aggaactcgg caaaatagcc ccgtaacttc gggagaaggg      120
gtgcccccc                                     128
```

<210> SEQ ID NO 40

<211> LENGTH: 392

<212> TYPE: DNA

<213> ORGANISM: Zea mays

<220> FEATURE:

<223> OTHER INFORMATION: Fragment of maize chloroplast genome

<400> SEQUENCE: 40

```
ctaacaataa acgaatacgg tttatgtatg gattccggta aaataccggt actcgatttc      60
ataagagtcg aataggaagt taagatgagg gtggtatcat cataaaaatg gagtagtatc      120
ctaaattata ctaatccacg tatgatattg atgcctttcc ttatcaaccg gaagtagtgc      180
aaaaaaaatt ctatactgca ctgctctctt tttactgaga aatgcaaaaa aataaaagtg      240
aagtaagggt gcccataga tatttgatct tgccctcctgt cccccccccc cttttttcat      300
```

-continued

caaaaatttc catgaaaaaa gaaaagatga atttgtccat tcattgaacc ctagttcggg 360
actgacgggg ctcgaaccgc cagcttcgc ct 392

<210> SEQ ID NO 41
<211> LENGTH: 188
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Represents part of PHI8999A insert sequence -
fragment of pat gene (complement)

<400> SEQUENCE: 41
gttcctagcc ttccagggcc cagcgtaagc aataccagcc acagcaccct caacctcagc 60
aaccaaccaa gggatatctat cttgcaacct ctctagatca tcaatccact cttgtgggtgt 120
ttgtggctct gtcctaaagt tcaactgtaga cgtctcaatg taatgggtta cgatatcaca 180
aaccgcgg 188

<210> SEQ ID NO 42
<211> LENGTH: 81
<212> TYPE: DNA
<213> ORGANISM: Zea mays
<220> FEATURE:
<223> OTHER INFORMATION: Fragment of maize chloroplast ORF241
(complement)

<400> SEQUENCE: 42
cacaagaacg aaagcacctt ttcattcttt catatactag gggtttttac ttggaaaaga 60
caatgttcca tactaaagga t 81

<210> SEQ ID NO 43
<211> LENGTH: 254
<212> TYPE: DNA
<213> ORGANISM: Zea mays

<400> SEQUENCE: 43
agctgcagaa gccgccaccg tcttgaggac cttccgggga gccagaccgg tcgaaccgtg 60
cctccacttg ctaaggagaa agggaaaatc agggccagga catacgaagg aggagccaga 120
acgaagatat cctaagatag ttactcgctc cgggccatga tcaatcatgc ctgtggggag 180
gtctctcgca cctcgatcca tgaaggtagc accgaggtct gcccgcgcgc cggtctcggt 240
accgtcctcg cctt 254

<210> SEQ ID NO 44
<211> LENGTH: 749
<212> TYPE: DNA
<213> ORGANISM: Zea mays

<400> SEQUENCE: 44
gggggcccca ggcaccggg ggatggactg cccaggcgca gccacgacga cccaaggatc 60
accctcctgc gcagtcggca cgagcaatag ttctcgggga acaggcagct tggcctgact 120
ccccggggtc acctcaacta cctcggccga ggggtcaagt acccctcag tccgccccgc 180
ctcttcggac cgggacccgc acgtcccgc cccgatacc gacggcacca gccgcctcgc 240
gggctggctt gacgaccctt ggcccagcct cagatctggg ctgaggccga ggcaggcggc 300
catgtcgtcg tottcatcat cgtotcatc atcgtcgtcg tcatcaggcg tctccggcga 360

-continued

```

cggctccctt gggagccctt ccctctcttg cgcacgacga agcctttcca aggcattccc 420
agcccacgtc cgctcgtggg cccgagcctt ctttgcgtcc ttcttctcct tctctttctc 480
cgcggtgacc ctccgcgcag ctccgtccac cgcattctcc gggactggtg gcagggaagg 540
cttgtgatgc cctacctcct ggagacagac gaaaagtctc agctatgaga accgagggca 600
atctgacgca agaaggaaga aggagcggat actcaccaga gacacgcacc cgcgatcggg 660
acgcattaag ggctgggaaa aagtgccggc ctctaatttc gctaccgtgc cgtccacca 720
cctgtggagg tcacgatgg gaaggggaa 749

```

```

<210> SEQ ID NO 45
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: 5' event flanking sequence; junction between
        regions 3 and 4

```

```

<400> SEQUENCE: 45

```

```

tcggactcga cctttctatc 20

```

```

<210> SEQ ID NO 46
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: 5' event flanking sequence; junction between
        regions 4 and 5

```

```

<400> SEQUENCE: 46

```

```

agagaatcac tcattcttca 20

```

```

<210> SEQ ID NO 47
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: 5' event flanking sequence; junction between
        regions 5 and 6

```

```

<400> SEQUENCE: 47

```

```

gaaaagtatt gtggctatct 20

```

```

<210> SEQ ID NO 48
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: 5' event flanking sequence; junction between
        regions 6 and 7a

```

```

<400> SEQUENCE: 48

```

```

tctcaaggcc gcagctgata 20

```

```

<210> SEQ ID NO 49
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: 5' event flanking sequence; junction between
        regions 7a and 7b

```

-continued

<400> SEQUENCE: 49

ggctaggaac cctcaacctc

20

<210> SEQ ID NO 50

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: 5' event flanking sequence; junction between regions 7b and 7c

<400> SEQUENCE: 50

tcacaaaccg agagaagagg

20

<210> SEQ ID NO 51

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: 5' event flanking sequence; junction between regions 7c and 8

<400> SEQUENCE: 51

agagggatct cgaagcttcg

20

<210> SEQ ID NO 52

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: 5' event flanking sequence; junction between regions 8 and 9

<400> SEQUENCE: 52

aaaattacca caactggaag

20

<210> SEQ ID NO 53

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: 3' event flanking sequence; junction between regions 9 and 10

<400> SEQUENCE: 53

agctatgttt ctcaactcgc

20

<210> SEQ ID NO 54

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: 3' event flanking sequence; junction between regions 10 and 11

<400> SEQUENCE: 54

acggatcctt caacaaaagg

20

<210> SEQ ID NO 55

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: 3' event flanking sequence; junction between

-continued

```

regions 11 and 12

<400> SEQUENCE: 55
gtgcccccg ctaacaataa                                20

<210> SEQ ID NO 56
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: 3' event flanking sequence; junction between
regions 12 and 13

<400> SEQUENCE: 56
gcttcgcgct gttcctagcc                                20

<210> SEQ ID NO 57
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: 3' event flanking sequence; junction between
regions 13 and 14

<400> SEQUENCE: 57
aaaccgcgga acacaagaac                                20

```

What is claimed is:

1. An isolated DNA molecule comprising a nucleotide sequence identified as SEQ ID NO:21.
2. An isolated DNA molecule comprising a nucleotide sequence identified as SEQ ID NO:22.
3. An isolated DNA molecule comprising a nucleotide sequence identified as SEQ ID NO:24.
4. An isolated DNA molecule comprising a nucleotide sequence identified as SEQ ID NO:26.
5. An isolated DNA molecule comprising a nucleotide sequence identified as SEQ ID NO:27.
6. A kit for identifying event TC1507 in a biological sample which detects a TC1507 specific region, said kit comprising at least a first primer, which recognizes a sequence within SEQ ID NO:21 or within SEQ ID NO:22.
7. The kit of claim 6, further comprising at least a second primer which recognizes a sequence within SEQ ID NO:25.
8. The kit of claim 6, further comprising at least a second primer which recognizes a sequence within SEQ ID NO:22.
9. The kit of claim 6, wherein the first primer recognizes a sequence within SEQ ID NO:21.
10. The kit of claim 6, wherein the first primer recognizes a sequence within SEQ ID NO:22.
11. The kit of claim 7, wherein said at least first and second primers comprise the sequence of SEQ ID NO: 1 and SEQ ID NO:2, respectively.
12. The kit of claim 7, wherein said at least first and second primers comprise the sequence of SEQ ID NO:2 and SEQ ID NO:23, respectively.
13. The kit of claim 8, wherein said at least first and second primers comprise the sequence of SEQ ID NO:3 and SEQ ID NO:5, respectively.

14. The kit of claim 8, wherein said at least first and second primers comprise the sequence of SEQ ID NO:4 and SEQ ID NO:5, respectively.

15. A DNA detection kit specific for junction DNA of maize event TC1507 and its progeny comprising at least one DNA molecule of a sufficient length of contiguous DNA polynucleotides to function in a DNA detection method, that is homologous or complementary to SEQ ID NO: 26.

16. A DNA detection kit specific for junction DNA of maize event TC1507 and its progeny comprising at least one DNA molecule of a sufficient length of contiguous DNA polynucleotides to function in a DNA detection method, that is homologous or complementary to SEQ ID NO: 27.

17. A kit for identifying event TC1507 in a biological sample, said kit comprising a specific probe comprising a sequence which hybridizes with SEQ ID NO:21 and SEQ ID NO:25, contiguous therewith.

18. A kit for identifying event TC1507 in a biological sample, said kit comprising a specific probe comprising a sequence which hybridizes with SEQ ID NO:22 and SEQ ID NO:25, contiguous therewith.

19. A DNA detection kit comprising: at least one DNA molecule of a sufficient length of contiguous nucleotides homologous or complementary to SEQ ID NO:26 or SEQ ID NO:27 that functions as a DNA primer or probe specific for maize event TC1507 and its progeny.

20. A DNA construct comprising: a first and second expression cassette, wherein said first expression cassette in operable linkage comprises

- (a) a maize ubiquitin promoter;
- (b) a 5' untranslated exon of a maize ubiquitin gene;
- (c) a maize ubiquitin intron;

- (d) a Cry1F encoding DNA molecule; and
 - (e) a 3' ORF25 transcriptional terminator; and
- said second expression cassette comprising in operable linkage
- (i) a CaMV 35S promoter;
 - (ii) a pat encoding DNA molecule; and
 - (iii) a 3' transcriptional terminator from (CaMV) 35S.
- 21.** A plant comprising the DNA construct of claim 20.
- 22.** A plant of claim 21, wherein said plant is a corn plant.
- 23.** A method for identifying event TC1507 in a biological sample, comprising detecting a TC1507 specific region with a probe or first primer which specifically recognizes a sequence within SEQ ID NO:21 or SEQ ID NO:22.
- 24.** The method of claim 23, further comprising amplifying a DNA fragment from a nucleic acid present in said biological sample using a polymerase chain reaction with at least two primers, wherein said first primer recognizes a sequence within SEQ ID NO:21 or SEQ ID NO:22, and a second primer recognizes a sequence within SEQ ID NO:22 or SEQ ID NO: 25.
- 25.** The method of claim 24, wherein said first primer recognizes a sequence within SEQ ID NO:21 and said second primer recognizes a sequence within SEQ ID NO: 25.
- 26.** The method of claim 23, wherein said first primer recognizes a sequence within SEQ ID NO:22 and a second primer recognizes a sequence within SEQ ID NO:22.
- 27.** The method of claim 25, wherein said first and second primers comprise the sequence of SEQ ID NO:2 and SEQ ID NO:1 respectively.
- 28.** The method of claim 25, wherein said first and second primers comprise the sequence of SEQ ID NO:2 and SEQ ID NO:23 respectively.
- 29.** The method of claim 26, wherein said first and second primers comprise the sequence of SEQ ID NO:3 and SEQ ID NO:5 respectively.
- 30.** The method of claim 26, wherein said first and second primers comprise the sequence of SEQ ID NO:4 and SEQ ID NO:5 respectively.
- 31.** The method of claim 27, comprising amplifying a fragment of about 912 bp using a TC1507 PCR identification protocol.
- 32.** The method of claim 28, comprising amplifying a fragment of about 844 bp using a TC1507 PCR identification protocol.
- 33.** The method of claim 29, comprising amplifying a fragment of about 342 bp using a TC1507 PCR identification protocol.
- 34.** The method of claim 30, comprising amplifying a fragment of about 252 bp using a TC1507 PCR identification protocol.
- 35.** A method of detecting the presence of maize event TC1507 or progeny thereof in a biological sample, comprising:
- (a) extracting a DNA sample from said biological sample;
 - (b) providing DNA primer molecules SEQ ID NO: 1 and SEQ ID NO:2;
 - (c) providing DNA amplification reaction conditions;
 - (d) performing said DNA amplification reaction, thereby producing a DNA amplicon molecule; and

- (e) detecting said DNA amplicon molecule, wherein the detection of said DNA amplicon molecule in said DNA amplification reaction indicates the presence of maize event TC1507.

36. A method of detecting the presence of maize event TC1507 or progeny thereof in a biological sample, comprising:

- (a) extracting a DNA sample from said biological sample;
- (b) providing DNA primer molecules SEQ ID NO:2 and SEQ ID NO:23;
- (c) providing DNA amplification reaction conditions;
- (d) performing said DNA amplification reaction, thereby producing a DNA amplicon molecule; and
- (e) detecting said DNA amplicon molecule, wherein the detection of said DNA amplicon molecule in said DNA amplification reaction indicates the presence of maize event TC1507.

37. A method of detecting the presence of maize event TC1507 or progeny thereof in a biological sample, comprising:

- (a) extracting a DNA sample from said biological sample;
- (b) providing DNA primer molecules SEQ ID NO:3 and SEQ ID NO:5;
- (c) providing DNA amplification reaction conditions;
- (d) performing said DNA amplification reaction, thereby producing a DNA amplicon molecule; and
- (e) detecting said DNA amplicon molecule, wherein the detection of said DNA amplicon molecule in said DNA amplification reaction indicates the presence of maize event TC1507.

38. A method of detecting the presence of maize event TC1507 or progeny thereof in a biological sample, comprising:

- (a) extracting a DNA sample from said biological sample;
- (b) providing DNA primer molecules SEQ ID NO:4 and SEQ ID NO:5;
- (c) providing DNA amplification reaction conditions;
- (d) performing said DNA amplification reaction, thereby producing a DNA amplicon molecule; and
- (e) detecting said DNA amplicon molecule, wherein the detection of said DNA amplicon molecule in said DNA amplification reaction indicates the presence of maize event TC1507.

39. An isolated DNA molecule comprising the amplicon produced by the method of claim 35.

40. An isolated DNA molecule comprising the amplicon produced by the method of claim 36.

41. An isolated DNA molecule comprising the amplicon produced by the method of claim 37.

42. An isolated DNA molecule comprising the amplicon produced by the method of claim 38.

43. A method of detecting the presence of DNA corresponding to the TC1507 event in a sample, the method comprising:

- (a) contacting the sample comprising maize DNA with a polynucleotide probe that hybridizes under stringent

hybridization conditions with DNA from maize event TC1507 and does not hybridize under said stringent hybridization conditions with a non-TC1507 maize plant DNA;

(b) subjecting the sample and probe to stringent hybridization conditions; and

(c) detecting hybridization of the probe to the DNA, wherein detection of hybridization indicates the presence of the TC1507 event.

44. An isolated DNA nucleotide primer sequence comprising a sequence selected from the group consisting of SEQ ID NO:1, 2, 3, 4, 5, 23, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56 and 57 or its complement.

45. An isolated DNA nucleotide primer sequence of claim 44 comprising a sequence selected from the group consisting of SEQ ID NO: 1, 2, 3, 4, 5 and 23 or its complement.

46. A pair of DNA molecules comprising: a first DNA molecule and a second DNA molecule, wherein the DNA molecules are of a sufficient length of contiguous nucleotides of SEQ ID NO:26 or its complement to function as DNA primers or probes diagnostic for DNA extracted from a TC1507 corn plant or progeny thereof.

47. A pair of DNA molecules comprising: a first DNA molecule and a second DNA molecule, wherein the DNA molecules are of a sufficient length of contiguous nucleotides of SEQ ID NO:27 or its complement to function as DNA primers or probes diagnostic for DNA extracted from a TC1507 corn plant or progeny thereof.

48. An isolated DNA molecule comprising a junction sequence comprising a sequence selected from the group consisting of SEQ ID NO: 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56 and 57 and complements thereof.

49. A method for confirming seed purity, comprising detection of a TC1507 specific region with a specific primer or probe which specifically recognizes a sequence within SEQ ID NO:21 or SEQ ID NO:22, in a seed sample.

50. A method for screening seeds for the presence of event TC1507, comprising detection of a TC1507 specific region with a specific primer or probe which specifically recognizes a sequence within SEQ ID NO:21 or SEQ ID NO:22 in a sample of a seed lot.

51. An insect resistant corn plant, or parts thereof, wherein DNA having at least one nucleotide sequence selected from the group consisting of SEQ ID NO: 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56 and 57 and complements thereof forms part of the plant's genome.

52. A descent plant of the insect resistant corn plant of claim 51 wherein DNA having at least one nucleotide sequence selected from the group consisting of SEQ ID NO: 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56 and 57 and complements thereof forms part of the plant's genome.

53. Seed of a plant of claim 51 or 52.

54. A method of producing an insect resistant corn plant comprising breeding with a plant of claim 51 or 52, and selecting progeny by analyzing for at least one nucleotide sequence selected from the group consisting of SEQ ID NO: 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56 and 57 and complements thereof.

55. An isolated DNA sequence comprising a sequence selected from the group consisting of DNA having at least one nucleotide sequence selected from the group consisting

of SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 23, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56 and 57 and complements thereof.

56. A pair of isolated DNA sequences, each comprising at least ten nucleotides and which when used together in a DNA amplification procedure will produce an amplicon diagnostic for event TC1507.

57. The pair of isolated DNA sequences of claim 56 wherein each is chosen from SEQ ID NO:26.

58. The pair of isolated DNA sequences of claim 56 wherein each is chosen from SEQ ID NO:27.

59. A method of detecting the presence of the TC1507 event insertion in corn tissue comprising:

(a) selecting a primer pair each comprising at least ten nucleotides from SEQ ID NO:26 or SEQ ID NO:27 wherein each member of the pair is on opposite sides of a sequence diagnostic for said TC1507 event insertion;

(b) contacting a sample of said corn tissue with said primer pair;

(c) performing DNA amplification and analyzing for an amplicons.

60. The method of claim 59 wherein said primer pair is selected from the group consisting of SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 23, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56 and 57 and complements thereof.

61. A method of detecting the presence of the TC1507 event insertion in corn tissue comprising:

(a) contacting a sample of said corn tissue with a polynucleotide probe that hybridizes under stringent hybridization conditions with one or more. DNA sequence selected from the group consisting of SEQ ID NO: 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56 and 57 and complements thereof;

(b) subjecting said sample and probe to stringent hybridization conditions; and

(c) analyzing for hybridization of the probe.

62. A DNA detection kit comprising a polynucleotide probe that hybridizes under stringent hybridization conditions with one or more DNA sequences selected from the group consisting of SEQ ID NO: 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56 and 57 and complements thereof.

63. A DNA detection kit comprising a primer pair each comprising at least 10 nucleotides from SEQ ID NO:26 and SEQ ID NO:27, wherein each is on opposite sides of a sequence diagnostic for the TC1507 event insertion.

64. The DNA detection kit of claim 61 wherein said primer pair is selected from the group consisting of SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 23, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56 and 57 and complements thereof.

65. A kit for identifying event TC1507 in a biological sample which detects TC1507 specific region within SEQ ID NO: 24.

66. A method for identifying TC1507 in a biological sample which detects a TC1507 specific region within SEQ ID NO: 24.