



US 20020177526A1

(19) **United States**(12) **Patent Application Publication** (10) **Pub. No.: US 2002/0177526 A1****Chen et al.**(43) **Pub. Date: Nov. 28, 2002**(54) **INSECTICIDAL SEED COATING****Publication Classification**(76) Inventors: **Yuguang Chen**, Lakeville, MN (US);
Kevin Mark Turnblad, Urbandale, IA
(US)(51) **Int. Cl.⁷** **A01H 5/00**; A01N 25/00;
A01N 25/26(52) **U.S. Cl.** **504/100**; 800/320.1

Correspondence Address:

SYNGENTA BIOTECHNOLOGY, INC.**PATENT DEPARTMENT****3054 CORNWALLIS ROAD****P.O. BOX 12257****RESEARCH TRIANGLE PARK, NC****27709-2257 (US)**(57) **ABSTRACT**

The present invention relates to an insecticidal coating for a corn seed comprising one or more binders selected from the group consisting of polymers and copolymers of polyvinyl acetate, methyl cellulose, polyvinyl alcohol, vinylidene chloride, acrylic, cellulose, polyvinylpyrrolidone and polysaccharide and an insecticide and filler wherein the binder forms a matrix for the insecticide and filler resulting in a substantially non-phytotoxic seed coating; methods for producing said coating and the seed treated with the above described coating.

(21) Appl. No.: **10/097,186**(22) Filed: **Mar. 13, 2002****Related U.S. Application Data**

(63) Continuation of application No. 09/259,930, filed on Mar. 1, 1999, now abandoned, which is a continuation-in-part of application No. 08/881,178, filed on Jun. 23, 1997, now Pat. No. 5,876,739, which is a continuation-in-part of application No. 08/664,052, filed on Jun. 13, 1996, now Pat. No. 5,849,320.

Additionally, the invention concerns a method of controlling soil-borne insect pests, particularly from the order of Coleoptera or Lepidoptera, comprising a) applying to a corn seed an insecticidal coating as described above, and transforming the corn seed with one or more transgenes coding for one or more proteins that confer pesticidal resistance to the corn seed.

INSECTICIDAL SEED COATING

BACKGROUND OF THE INVENTION

[0001] Numerous technologies are currently utilized to treat seeds with the desired purpose of enhancing crop performance. These treatments include inter alia the coating, pelleting and/or film overcoating of seeds. A range of fungicide treatments as dusts, liquids and slurries have long been used to control soil and seed-borne diseases in vegetable seeds. However, the use of insecticides for vegetable seed treatment has lagged behind that of fungicide seed treatments, and moreover, seed treatment of agronomic crops is even less advanced than that for vegetable seed.

[0002] Recent developments in seed treatment technology have focused on the use of techniques to deliver pesticides to seeds. Film-coating has been studied as a means of delivering insecticides such as benfuracarb, chlorpyrifos, chlorfenvinphos and others. In the past, the concentration of these insecticides and other active ingredients present in the seed coating was limited due to direct phytotoxic effect of the insecticide on the seed. Using the instant seed coating of the present invention solves the problem of direct insecticide phytotoxicity to the seed.

SUMMARY OF THE INVENTION

[0003] The present invention relates to an insecticidal coating for a seed comprising one or more binders selected from the group consisting of polymers and copolymers of polyvinyl acetate, vinyl acetate/ethylene, methyl vinyl ether, malic anhydride, methyl cellulose, polyvinyl alcohol, vinylidene chloride, acrylic, cellulose, polyvinylpyrrolidone and polysaccharide, one or more filler(s), and an insecticidally effective amount of an insecticide, preferably an organophosphate, phenyl pyrazole, pyrethoid, oxadiazine or imidacloprid insecticide, most preferably tefluthrin and/or an oxadiazine or combinations thereof, wherein the binder forms a matrix for the insecticide and filler resulting in a substantially non-phytotoxic seed coating.

[0004] In a preferred embodiment, the invention concerns an insecticidal coating for a corn seed comprising a binder in an amount from about 0.01 to about 15% of the weight of the seed wherein said binder is a vinyl acetate-ethylene copolymer or polymer or copolymer of vinylidene chloride, optionally in combination with methyl vinyl ether/malic anhydride copolymer, a filler in an amount of up to about 70% of the weight of the seed, an insecticide in an amount from about 0.005 to about 50% of the weight of the seed and selected from the group consisting of imidacloprid, terbufos, chlorpyrifos, fipronil, tefluthrin, chloroethoxyfos, tebufirimfos, and mixtures thereof, wherein said binder forms a matrix for the insecticide and filler(s) resulting in a substantially non-phytotoxic seed coating.

[0005] In another preferred embodiment the invention relates to a method of controlling soil-borne insect pests, such as corn root worm, comprising applying to a seed an insecticidal coating as described above wherein said insecticide is applied to the seed in an insecticidally effective amount to control insect pests.

[0006] Additionally the invention concerns a method of preparing a coated seed comprising a) mixing one or more binders with an insecticide selected from the group consist-

ing of imidacloprid, terbufos, chlorpyrifos, tefluthrin, fipronil, chloroethoxyfos, tebufirimfos, and mixtures thereof, wherein the binder serves as a matrix for the insecticide; b) applying the mixture to a seed; c) allowing the mixture to dry on said seed and d) applying a film overcoating to the seed.

[0007] In still another embodiment the invention is concerned with a method of protecting emerging seedlings of a crop plant from one or more insect organisms in the seed growing soil environment which comprises applying to the seeds of the crop plant an insecticidally effective amount of an insecticidal coating as described herein above.

[0008] The seed growing environment as used herein includes that area directly surrounding the seed and young roots.

[0009] Frequently when pesticides are applied to seeds, the pesticide may cause physiological damage to the seed or seedling due to phytotoxic effects of the concentrated pesticide in or around the seed. The present invention is novel in that pesticides may be applied in a polymer matrix at pesticidally effective concentrations without having an adverse phytotoxic effect on the emerging seedling. In particular it has been found that the use of the coating of the present invention is effective in the control of insect pests, such as Coleoptera and Lepidoptera insects.

[0010] It is a most important object of the present invention to provide a seed coated with a coating which protects the seed or the emerging seedling from physiological damage potentially caused by the insecticidal ingredient of the coating.

[0011] A still further object of the invention is to provide a hybrid corn seed optionally transformed with gene encoding a pesticidally active protein, coated with an insecticide which aids in the control of Coleopteran pests, such as Diabrotica insects, and particularly corn rootworm larvae.

[0012] Another object of the invention is to provide a seed which is coated with an insecticide whereby the use of the insecticide in the coating increases the safety of using said insecticide by reducing operator exposure.

[0013] Still another object of the invention is to provide an insecticidal seed coating wherein the insecticide is encapsulated within a matrix that provides for the controlled release of the insecticide over a prolonged period of time.

[0014] Yet another aim of the invention is to improve the flowability of seed through a seed planter which will ultimately result in more uniform seed planting.

DETAILED DESCRIPTION OF THE INVENTION

[0015] The present invention relates to improved seed coatings which are superior to conventional seed coatings comprising insecticidal components. The invention also relates to the use of insecticides in seed coatings wherein the insecticide has heretofore been ineffective against certain insect pests, however, may be formulated in the seed coating of the present invention in sufficiently high amounts to be effective against those same insect pests without causing unacceptable phytotoxicity to the seed.

[0016] As used herein the term coated seed means a seed that has been subjected to a procedure whereby the seed is treated with one or more adhering coating layers.

[0017] The coating described herein includes two main components: a) one or more binders and b) an insecticide. The binder serves as a matrix for the insecticide and is preferably present in the seed coating in an amount sufficient to prevent or reduce the levels of phytotoxicity caused by the insecticide.

[0018] The binder component of the coating is composed preferably of an adhesive polymer that may be natural or synthetic and is without phytotoxic effect on the seed to be coated. The binder may be selected from polyvinyl acetates, polyvinyl acetate copolymers (-ethylene), methyl vinyl ether, malic anhydride, polyvinyl alcohols, polyvinyl alcohol copolymers, celluloses, including ethylcelluloses and methylcelluloses, hydroxymethylcelluloses, hydroxypropylcellulose, hydroxymethylpropylcelluloses, polyvinylpyrrolidones, dextrans, maltodextrins, polysaccharides, fats, oils, proteins, gum arabics, shellacs, vinylidene chloride, vinylidene chloride copolymers, calcium lignosulfonates, acrylic copolymers, starches, polyvinylacrylates, zeins, gelatin, carboxymethylcellulose, chitosan, polyethylene oxide, acrylimide polymers and copolymers, polyhydroxyethyl acrylate, methylacrylimide monomers, alginate, ethylcellulose, polychloroprene and syrups or mixtures thereof. Preferred binders include polymers and copolymers of vinyl acetate, methyl vinyl ether, malic anhydride, methyl cellulose, polyvinyl alcohol, vinylidene chloride, acrylic, cellulose, polyvinylpyrrolidone and polysaccharide. The above-identified polymers include those known in the art and for example some are identified as Rhoplex™ B-60A, Methocel™ A15LV, Methocel™ E15LV, Cellosize™ QP, Air-Flex™ 500, Daratak™ SP 1090, Elvanol™ 85-30, Gantrez™ (AN139) 10%, Rhoplex™ AC-33-NP, Rhoplex™ B-85 and Vinamul™ 18132. Particularly preferred classes of polymers include polymers and copolymers of vinylidene chloride, methyl vinyl ether/malic anhydride copolymers and vinyl acetate-ethylene copolymers.

[0019] The amount of binder in the coating will be in the range of about 0.01 to 15% of the weight of the seed. A preferred range will be about 0.1 to 10.0% of the weight of the seed.


[0020] The binder must be chosen so that it serves as a matrix for the insecticide. While the binders disclosed above may all be useful as a matrix, the specific binder will depend on the properties of the insecticide. The term "matrix" as used herein means a continuous solid phase of one or more binder compounds and contains vacancies, voids or spaces occupied by the insecticide and filler. The term matrix is given a broad meaning and includes what may be viewed as a matrix system, a reservoir system or a microencapsulated system. In general a matrix system consists of an insecticide and filler uniformly dispersed within a polymer, while a reservoir system consists of a separate insecticide phase, insecticide particles or droplets physically dispersed within a surrounding, rate limiting polymeric phase. Microencapsulation includes the coating of small particles or droplets of liquids. The term microencapsulation has not only been applied to coated particles or droplets but also to dispersions in a solid matrix. Without being limited to the specific encapsulating system (matrix, reservoir or microencapsulated) the term matrix is meant to be inclusive of the above listed systems.

[0021] The invention further contemplates the use of fillers, such as absorbent or inert fillers, in the insecticidal coating. It has been discovered that the use of fillers in the coating is particularly effective for protecting the seed during stress conditions. Fillers for such formulations are known in the art and may include woodflours, clays, activated carbon, sugars, diatomaceous earth, cereal flours, fine-grain inorganic solids, calcium carbonate and the like. Clays and inorganic solids which may be used include calcium, bentonite, kaolin, china clay, talc, perlite, mica, vermiculite, silicas, quartz powder, montmorillonite and mixtures thereof. Sugars which may be used include dextrin and maltodextrin. Cereal flours include: wheat flour, oat flour and barley flour. Preferred fillers include diatomaceous earth, perlite, silica and calcium carbonates and mixtures thereof. For example, a product containing diatomaceous earth and amorphous silica such as that manufactured by Celite Corporation (Celite™) or bentonite (Actisil™) are most preferred. One skilled in the art will appreciate that this is a non-exhaustive list of materials and that other recognized filler materials may be used depending on the seed to be coated and the insecticide used in the coating.

[0022] The filler is chosen so that it will provide a proper microclimate for the seed, for example the filler is used to increase the loading rate of the active ingredient and to adjust the control-release of the active ingredient. A filler aids in the production or process of coating the seed. The effect varies, because in some instances formulated insecticidal compounds will comprise a filler. The amount of filler used may vary, but generally the weight of the filler components will be in the range of about 0.005 to 70% of the seed weight, more preferably about 0.01 to 50% and most preferably about 0.1 to 15%. The filler may be supplied in the coating of the invention with the insecticidal component. The specific examples as described herein utilize readily available commercial formulations of known insecticides wherein filler material is included in the formulation.

[0023] Suitable insecticides include those selected from azoles, for example, triazoles, azines, pyrethroids, organophosphates, caramoyloximes, pyrroles, pyrazoles, pyridines, amidines, halogenated hydrocarbons, and carbamates and combinations and derivatives thereof. Particularly suitable classes of insecticides include insect growth regulators (e.g., Mimic™) organophosphates (e.g., Fortress™), phenylpyrazoles and pyrethroids. Preferred insecticides are those known as terbufos, chlorpyrifos, fipronil (Regent™), chloroethoxyfos, tefluthrin (Force™), fiproles, phenoxycarb, diofenolan (Aware™), pymetrozine (Chess™), carbofuran, tebufirimfos, and imidacloprid (Gaucho™, Confidor™, Admire™), including imidacloprid analogs, such as (substituted or unsubstituted) nitro-, oxo-, thio- or cyano-substituted-guanidines, enamines, and oxadiazines. More preferred are imidacloprid and imidacloprid analogs, for example, those insecticidal compounds disclosed in U.S. Pat. No. 5,034,524, terbufos, chlorpyrifos, fipronil, tefluthrin, chloroethoxyfos, tebufirimfos, and mixtures thereof, for example, oxadiazines and tefluthrin. Most preferred are tefluthrin or an oxadiazine, or a combination thereof.

[0024] It has also been discovered that the seed treatment of this invention demonstrates applicability to oxadiazine compounds of the general formula (I) (See U.S. Pat. No. 5,852,0120):



[0025] in which

[0026] A is an unsubstituted or mono- to tetrasubstituted, aromatic or non-aromatic, monocyclic or bicyclic heterocyclic radical, where one to two of the substituents of A can be selected from the group consisting of halo-C₁-C₃alkyl, cyclopropyl, halocyclopropyl, C₂-C₃alkenyl, C₂-C₃alkynyl, halo-C₂-C₃alkenyl, halo-C₂-C₃alkynyl, halo-C₁-C₃alkoxy, C₁-C₃alkylthio, halo-C₁-C₃alkylthio, allyloxy, propargyloxy, allythio, propargylthio, haloallyloxy, haloallylthio, cyano and nitro, and one to four of the substituents of A can be selected from the group consisting of C₁-C₃alkyl, C₁-C₃alkoxy and halogen;


[0027] R is hydrogen, C₁-C₆alkyl, phenyl-C₁-C₄alkyl, C₃-C₆cycloalkyl, C₂-C₆alkenyl or C₂-C₆alkynyl; and

[0028] X is N—NO₂ or N—CN,


[0029] or, if appropriate, a tautomer thereof, in each case in free form or in salt form.

[0030] An especially preferred oxadiazine insecticide of the general formula (I) is the compound

[0031] wherein A is



[0032] and X is N—NO₂ which yields the compound of formula II:



[0033] The compound of formula II may be used in combination with other pesticides, for example, with tefluthrin. Other combinations of pesticides, for example, pyrethroids and organophosphates, are also suitable.

[0034] Also included are insect growth regulators for example, methoprene and hydroprene. These are well known to those skilled in the art.

[0035] The amount of the insecticide in the coating will vary depending on the type of seed and particular active ingredient, but in general will range from about 0.005 to 50% of the weight of the seed. A preferred percent range for the insecticide is about 0.01 to 40%. In particular, the more preferred range for chlorpyrifos will be about 0.5 to 20.0%, a most preferred range will be about 1.0 to 15%; the more preferred range for tefluthrin will be about 0.01 to 10.0%, a most preferred range will be about 0.05 to 5.0%; the more preferred range for tebupirimfos will be about 0.01 to 5.0%, a most preferred range from about 0.05 to 3.0%; the more preferred range for fipronil will be about 0.01 to 10.0%, and a most preferred range will be about 0.05 to 5.0%; and the more preferred range for imidacloprid and imidacloprid analog insecticides will be about 0.01 to 10%, and a most preferred range will be about 0.05 to 5%.

[0036] As one skilled in the art can appreciate the exact amount will vary depending of the size of the seed to be coated. The insecticide of the coating must not inhibit germination of the seed and should be efficacious during that time in the target insects life cycle which causes injury to a crop plant. One skilled in the art will appreciate that this time will vary depending on the target insect among other factors. In general the coating will be efficacious for approximately 0 to 120 days after sowing. The coating of the present invention will contain an amount of insecticide that is insecticidally effective. An insecticidally effective amount as used herein means that amount of insecticide that will kill insect pests in the larvae or pupal stage of growth or will consistently reduce or retard the amount of damage produced by insect pests.

[0037] When the insecticidal component used for the coating is an oily type formulation and little or no filler is present, it may be necessary to hasten the drying process by drying the formulation. This optional procedure may be accomplished by means well known to those skilled in the art and includes the addition of calcium carbonate, kaolin or bentonite clay, perlite, diatomaceous earth or any adsorbent material added preferably concurrently with the insecticidal coating layer to absorb the oil or excess moisture. The amount of calcium carbonate or related compounds to dry charge the solution will be in the range of about 0.5 to 10.0% (w/w).

[0038] The coatings formed by the invention are capable of effecting a slow rate of release of the insecticide by diffusion or movement through the matrix to the surrounding medium.

[0039] Virtually any crop seed can be treated in accordance with the invention, such as cereals, vegetables, ornamentals, and fruits. Particular crop seeds are selected from the group of corn (sweet and field), soybean, wheat, barley, oats, rice, cotton, sunflower, alfalfa, sorghum, rapeseed, sugarbeet, Brassica spp., tomato, bean, carrot, tobacco and flower seed, for example, pansy, impatiens, petunia and geranium. The most preferred seeds include corn and soybean.

[0040] Preferred target pest organisms include soil borne insects of the order:

[0041] Coleoptera particularly *Diabrotica* sp. (Western, Southern and Northern corn rootworm), for example, *D. balteata*, *D. virgifera*, *D. undecimpunctata*, and *D. longicornis*; *Melanotus* sp. (Corn wireworm); *Phyllophaga* sp. (White grubs, Wireworms, and False wireworms); *Limoni* sp. (Sugarbeet wireworms) and *Agrotis* sp. (Wheat wireworms, White grubs and Seed maggots):

[0042] Lepidoptera particularly *Peridroma* sp. (Varigated cutworm); *Euxoa* sp. (Army cutworm); and *Agrotis* sp. (Black cutworm):

[0043] Diptera particularly *Hylemya* sp. (Seedcorn maggot) and *Tetanops* sp. (Sugarbeet root maggot):

[0044] Homoptera particularly *Pemphigus* sp. (Sugarbeet root aphid, Cutworm, and White grub) and *Aphis* sp. (Corn root aphid).

[0045] The insecticidal coating is particularly useful in accommodating high insecticide loads, as are required to treat typically refractory pests, such as corn root worm, while at the same time preventing unacceptable phytotoxicity due to the increased insecticide load.

[0046] Most preferred target organisms include in addition to *Diabrotica* sp, *Agrotis ypsilon*; *melanotus cribrulosus*; *Hylemya cilicrura*; *Agonoderus lecontei*; *Feltia subgothica* and *Phyllophaga rugosa*.

[0047] In other embodiments a plasticizer may be used in the insecticidal solution. Plasticizers are typically used to make the film that is formed by the insecticidal coating layer more flexible, improve adhesion, spreadability and improve the speed during processing. The improved film flexibility is important to minimize chipping, breakage or flaking during handling or sowing processes. Many plasticizers may be used however, most preferred plasticizers include polyethylene glycol, glycerol, butylbenzylphthalate, glycol benzoates and related compounds. The range of the percent of plasticizer in the insecticidal coating layer will be in the range of about 0.1 to about 20%.

[0048] In addition to the insecticidal coating layer, the seed may be treated with one or more of the following ingredients: other pesticides including fungicides and herbicides; herbicidal safeners; fertilizers and/or biocontrol agents. These ingredients may be added as a separate layer or alternatively may be added in the insecticidal coating layer.

[0049] Furthermore, the seed may also be genetically transformed, e.g., with genes coding for proteins having insecticide, herbicide and/or fungicide resistance properties. It is preferable to transform the seed to protect it from the indigenous pests that cause the greatest harm. For corn root worm, for example, the seed may be transformed by a gene encoding an active Bt (*Bacillus thuringiensis*) protein or active Bt fragment or synthetic or semi-synthetic polypeptide replicate thereof, e.g., a cry 1A(b) protein, preferably encoded by a plant optimized gene as described in U.S. Pat. Nos. 5,165,136 and 5,689,032 as well as WO 98/04984, the disclosures of which are incorporated herein by reference, or a gene encoding a vegetative insecticidal protein (VIP), preferably a plant optimized and/or synthetic gene, such as

disclosed in WO 98/44137, WO 96/10083, and U.S. Pat. No. 5,872,212, the disclosures of which are incorporated herein by reference.

[0050] Where both the seed coating and the transformed seed are active against the same target insect, the combination is useful (i) in a method for enhancing activity against the target insect and (ii) in a method for preventing development of resistance to a transgenic insecticidal protein by providing a second mechanism of action against the target insect. Thus, the invention provides a method of enhancing activity against or preventing development of resistance in a target insect, e.g., a Lepidopteran or Coleopteran pest, for example Black Cutworm or Corn Rootworm, comprising applying the seed coating of the invention to a seed comprising one or more transgenes expressing an insecticidal protein or combination of proteins, preferably a protein or proteins derived from or having the insecticidal activity of the insecticidal proteins from *Photobacterium* or *Bacillus*, such as a delta endotoxin from *Bacillus thuringiensis* (especially cry1Ab, cry1Ac, cry1F, cry9c, cry2A, or cry 6A) or a vegetative insecticidal protein from *Bacillus thuringiensis* or *Bacillus cereus* (especially VIP3).

[0051] Even where the transformed seed is active against a different target insect, the seed coating is useful to expand the range of insect control, for example by adding control of Coleopteran pests such as Corn Rootworm to a transformed seed having a transgene which expresses a protein which is active against Lepidopteran pests, such as Black Cutworm, Fall Armyworm and European Corn Borer, but not against Coleopteran pests.

[0052] The invention thus further provides seed coated with the seed coating of the invention, wherein said seed comprises one or more transgenes expressing an insecticidal protein or combination of proteins, preferably a protein or proteins derived from or having the insecticidal activity of the insecticidal proteins from *Photobacterium* or *Bacillus*, such as a delta endotoxin from *Bacillus thuringiensis* (especially cry1Ab, cry1Ac, cry1F, cry9c, cry2A, or cry6A) or a vegetative insecticidal protein (VIP) from *Bacillus thuringiensis* or *Bacillus cereus* (especially VIP3).

[0053] Suitable pesticides include those listed herein and those listed in The Pesticide Manual, 9th Ed., Editor, Charles Worthing, published by the British Crop Protection Council and hereby incorporated by reference.

[0054] A fungicide may be applied to the seed prior to the coating layer described herein. The application of a fungicide as a dust, slurry or the like is a well known practice in the art and is not considered a coating layer within the meaning of the term used herein. Suitable examples of fungicides include Captan (N-(trichloromethyl)thio-4-cyclohexane-1,2-dicarboximide); Thiram (tetramethylthioperoxydicarbonic diamide); Metalaxyl (methyl N-(2,6-dimethylphenyl)-N-(methoxyacetyl)-DL-alaninate; Fludioxonil (4-(2,2-difluoro-1,3-benzodioxol-4-yl)-1H-pyrrol-3-carbonitrile; and Oxadixyl (N-(2,6-dimethylphenyl)-2-methoxy-N-(2-oxo-3-oxazolidinyl) acetamide. One skilled in the art will be aware of other beneficial fungicides suitable for combating harmful pathogens which are not only a problem for a particular locale where the coated seed is to be grown but also suitable for the protection of seeds in storage before planting.

[0055] The amount of fungicide to be added will vary due to the strength of its active ingredient, but in general may

range from about 0.001 to about 10% of the weight of the seed and preferably from about 0.01 to 2.0%. However, for a particular situation the amounts may be greater or less.

[0056] Suitable herbicides include those selected from carbamates, thiocarbamates, acetamides, particularly chloroacetamides, triazines, dinitroanilines, glycerol ethers, pyridazinones, uracils, phenoxy, ureas, and benzoic acids and derivatives. Suitable safeners include for example, benzoxazine, benzhydryl derivatives, N,N-diallyl dichloroacetamide, various dihaloacyl, oxazolidinyl and thiazolidinyl compounds, ethanone, naphthalic anhydride compounds, and oxime derivatives.

[0057] Suitable biocontrol agents are bacteria of the genera *Rhizobium*, *Bacillus*, *Pseudomonas*, and *Serratia*, fungi of the genera *Trichoderma*, *Glomus*, and *Gliocladium* and mycorrhizal fungi.

[0058] The above compounds are listed as examples and are not intended to be an exhaustive list of compounds that can be used in the insecticidal coating layer or in additional other coating layers.

[0059] Conventional means of coating may be used for carrying out the coating of the invention. Additionally, various coating machines are available to one skilled in the art. Three well known techniques include the use of drum coaters, and fluidized bed techniques. Other methods, such as spouted beds may also be useful. The seeds may be presized prior to coating. After coating the seeds are dried and then optionally sized by transfer to a sizing machine. These machines are known in the art for example, a typical machine used when sizing seed corn in the industry.

[0060] Film-forming compositions for enveloping coated seeds are well known in the art, and a film overcoating can be optionally applied to the coated seeds of the present invention. The film overcoat protects the coating layers and optionally allows for easy identification of the treated seeds. In general, additives are dissolved or dispersed in a liquid adhesive, usually a polymer into or with which seeds are dipped or sprayed before drying. Alternatively a powder adhesive can be used. Various materials are suitable for overcoating including but not limited to, methyl cellulose, hydroxypropylmethylcellulose, dextrin, gums, waxes, vegetable or paraffin oils; water soluble or water disperse polysaccharides and their derivatives such as alginates, starch, and cellulose; and synthetic polymers such as polyethylene oxide, polyvinyl alcohol and polyvinylpyrrolidone and their copolymers and related polymers and mixtures of these.

[0061] Further materials may be added to the overcoat including optionally plasticizers, colorants, brighteners and surface active agents such as, dispersants, emulsifiers and flow agents including for example, calcium stearate, talc and vermiculite. Additionally pesticides, such as fungicides may be added to the film coat, however, it has been observed that fungicides added initially to the seed gives better results than when added with the overcoat. Fluidized bed and drum film coating techniques described above can be employed for film coating.

[0062] The overcoating film is in the range of approximately 0.01% to about 20% of the weight of the seed it is applied to. The preferred range will be about 0.01% to about 10.0% and a most preferred range will be about 0.01 to

5.0%. A preferred solution will vary depending on the specific active ingredient. A preferred overcoating film may include methyl cellulose, hydroxypropylmethylcellulose, polyvinyl acetate, PEG and mixtures thereof. Additionally, depending on the type of insecticide the overcoat film may have incorporated therein a bird repellent compound. These compounds are known in the art and include for example; anthraquinone, methyl anthranilate, capsaicin and oxygenated tetracyclic triterpenoid compounds (also referred to as cucurbitacins) and disclosed in U.S. Pat. No. 5,292,533. Moreover these compounds may be added to the insecticidal coating layer as well.

[0063] Depending on the germplasm and crop seed to be coated other benefits may be seen by subjecting the seeds to additional treatments prior to coating such as priming or treatment with growth regulators and biocontrol agents.

[0064] In seed priming, the seeds are exposed to enough moisture to enhance germination, but not enough to cause premature germination. Examples of priming procedures are known in the art and include drum priming and steep priming.

[0065] Growth regulators which may be added prior to coating includes auxins, cytokinins, gibberellins, ethylene, abscisic acid, IAA, AA, IBA, ethephon, acetamide, maleic, hydrazide, diminozide and like compounds. Biological treatment may also be used to enhance seed performance and help in the control of harmful organisms.

[0066] Seeds treated with the coating of the invention have the following advantages: They possess less risk with respect to operator handling and exposure because of reduced dust exposure. Pesticides can be applied in a uniform way and loss of pesticides during transport and handling is prevented. Phytotoxicity is reduced to the seeds and developing seedlings and therefore there is no adverse effect on germination, seedling emergence and seedling development. Additionally, placement of pesticides, particularly insecticides, closer to the insect soil feeding zone may improve insect control.

[0067] Specific embodiments of the invention are described hereinafter by way of example only.

EXAMPLE 1

[0068] Seeds Treated with an Insecticidal Coating Layer:

[0069] a. Five pounds maize seed pretreated with Captan 400 fungicide and Pro-Ized Red colorant, are coated in an Vector LDCS coating machine. The seeds are warmed for is approximately 5 minutes in a rotating pan at approximately 20 r.p.m. with an air flow of 31-32 c.f.m. and an air pressure of 20 p.s.i. A 992 gram insecticide solution comprising 183 grams of 50% chlorpyrifos (the active ingredient in Lorsban™ 50-SL) and 50% premixed inert filler, 47 gram Airflex™ 500 (vinyl acetate-ethylene), and 762 grams water is sprayed onto the seed until all the solution is utilized. Inlet air temperature is approximately 50° C. and may be periodically adjusted to maintain seed temperature about equal to or less than 35° C. or germination may be effected.

[0070] b. Seed is treated as described above however, a 334.0 gram insecticide solution comprising 51.87 grams of 27.9% tefluthrin (the active ingredient in Force™ 30CS), 21.0 grams Vinamul™ 18132 (vinyl acetate-ethylene), 30 grams of Celite™ 266 and 231.0 grams water is applied to the seed.

[0071] c.

[0072] (1) Seed is treated as described above however, a 900 gram insecticide solution comprising 15.5 grams of 92% tebuirimphos (the active ingredient in Mat™ 7484), 90 grams of Celite™ 266, 22 grams Airflex™ 500 (vinyl-acetate-ethylene), and 772 grams water is applied to the seed.

[0073] (2) Seed is treated as described in c. (1), except no Celite™ 266 is added to the solution. A total of 30 grams of calcium carbonate is applied to the seed during the coating process as a drying charge.

[0074] d. Seed is treated as described above however, a 293 gram insecticide solution comprising 11.4 grams of 80% fipronil (the active ingredient in Regent™), 22.6 grams Celite™ 266, 9 grams Airflex™ 500 and 250 grams water is applied to the seed.

[0075] Lorsban™ 50-SL; active ingredient is=0,0-dimethyl 0-(3,5,6 trichloro-2-pyridyl) phosphorothioate, Dow Elanco.

[0076] Force™ 30CS; active ingredient is=[1 α , 3 α (Z)]-($\pm$)-(2,3,5,6-tetrafluoro-4-methylphenyl)methyl 3-(2-chloro-3,3,3-trifluoro-1-propenyl)-2,2-dimethylcyclopropanecarboxylate, Zeneca, Inc.

[0077] Mat™ 7484; active ingredient=o-[2-(1,1-dimethylethyl)-5-pyrimidinyl-o-ethyl o-(1-methylethyl) phosphorothioate, Bayer.

[0078] Regent™ 80WDG; active ingredient=5-amino-1-(2,6-dichloro-4-trifluoromethylphenyl)-3-cyano-4-trifluoromethane sulphonyl pyrazole, Rhone-Poulenc.

EXAMPLE 2

[0079] Overcoating of Coated Seeds:

[0080] To each of the seed examples described in Example 1 is added a film overcoat. The same machine and set points

are used for the overcoat. The machine is set at an inlet temperature of 50° C. which is periodically adjusted to maintain seed temperature about equal to or less than 35° C. A solution comprised of 438 grams water, 2.2 grams Methocel K100M, 0.36 grams PEG 8000, 5.8 grams Pro-Ized red colorant 0.73 grams GP Hytech Calcium stearate 5899 and 22.2 grams Airflex is sprayed on the seed for 15 minutes.

EXAMPLE 3

[0081] Seed Germination and Field Emergence Results:

[0082] Corn hybrid seed germination is tested under testing rules of Association of Official Seed Analysts (AOSA). Both paper towel and soil are used for separate germination tests. In addition, seedling shoots and roots were separated after 7 to 10 days of germination. Dry weight of shoots and roots is recorded to indicate seedling growth. For field emergence test, corn hybrid seed is planted in Iowa, Minnesota, Wisconsin, Nebraska, Illinois, and Ohio in a split plot design with two rows per treatment and four replications. Emergence counts are taken every other day beginning with the first day plants emerge for each treatment and continue until 21 days after emergence. The % emergence is determined for all locations for each treatment and is exhibited in Table 1. This Table demonstrates that seed coating does not adversely affect seedling emergence. The treatments are described as follows:

[0083] Treatment 1(a): Seed is treated as described in Example 1(a);

[0084] Treatment 1(b): Seed is treated with chlorpyrifos at the same rate as described in Example 1(a), except that no binder (Airflex™ 500) is used.

[0085] Treatment 2: Seed is treated as described in Example 1(b);

[0086] Treatment 3: Seed is treated as described in Example 1(c)(1);

[0087] Treatment 4: Seed is treated as described in Example 1(d);

[0088] Treatment 5: Seed is treated with Captan 400 fungicide;

[0089] Treatment 6: Seed is treated as described in Example 2;

TABLE 1

Treatment	% Germination				Shoots (mg)	Roots (mg)	% Corn Emergence
	Paper	Soil					
		4 Days	7 Days				
1(a)	99	95	95		10.2	12.55	88
1(b)		70	80*				
2	98				—	—	81
3	98				13.5	11.6	87
4	99				13.1	12.95	86
5	99				—	—	87
6	99	95	97		11.67	13.14	88

EXAMPLE 4

[0090] Effect on Phytotoxicity of Using Filler in the Coating

[0091] A high resolution saturated cold test was used to test the effect of filler on the reduction of phytotoxicity. In this example seeds were germinated at 10° C. with low oxygen, and a water saturated condition for 5 days, and then transferred to 25° C. to finish the germination process. The % saturated cold germination and % field emergence is exhibited in Table 2.

TABLE 2

Hybrid Seed	Treatment	% Sat. Cold Germ	% Corn Emergence
N4242	C (1)	72	88
N4242	C (2)	66	84
N7590	C (1)	71	
N7590	C (2)	54	

The results indicate that the formulation containing a filler (C (1)) provides an improvement in % germination and emergence.

EXAMPLE 5

[0092] Corn Rootworm Efficacy Studies:

[0093] Hybrid corn seeds with and without an insecticidal coating are planted in Nebraska, Iowa, Minnesota, Wisconsin and Illinois to determine the efficacy of various coated seed treatments in standard tests. All seeds contain a Captan pretreatment whether or not a coating is applied. Additionally, an overcoat layer is applied to all coated seed which comprises the coating described in Example 2. Banded granular insecticides are applied at the manufactures label rates using granular insecticide applicators. Plot size for each treatment is two ten foot rows spaced 30 inches apart with four replications of each treatment at each location. Non-diapausing 1st instar Western Corn Rootworm larvae are side dressed on both sides of the corn plants in the row at a rate of 800 eggs per foot of row. Application of the insect larvae is made when the corn seedlings are approximately at the two leaf stage. The placement of the larvae is approximately two inches to the side of the corn plants and two inches below the soil surface. Plots are maintained with normal cultural practices until the corn plants reach the flowering stage. At late brown silk stage, 10 sequential roots from each plot are obtained. The corn stalks are cut at approximately 12 inches above the soil line. Each corn root is washed and after washing the roots are rated on a scale of 1-6 using the Iowa Root Rating Method developed at Iowa State University (Hills and Peters, 1971) and used commonly throughout the corn belt. In the rating, 1=no damage or only a few minor feeding scars; 2=feeding scars evident but no roots eaten off to within 1½ inch of the plant; 3=several roots eaten off to within 1½ inch of the plant but never the equivalent of an entire node of roots is destroyed; 4=one root node completely destroyed; 5=two root nodes completely destroyed and 6=three or more root nodes destroyed. A destroyed root is defined as a root that has been pruned to within 1½ inch of the base. Pruned roots do not have to originate from a single node, but all pruned roots must equal the equivalent of a full node to count as a destroyed node. The mean root rating results are reported in Table 2 below. In addition, a growth chamber and a greenhouse bioassay was used to determine the effect of insecti-

cidal seed coating on corn rootworm control. For growth chamber assay, corn seed was planted in a 16 oz cup with garden Jiffymix, and allowed to grow for two weeks. Twenty neonate corn rootworm larvae are placed in the Jiffymix near corn roots and allowed to feed for approximately 2 weeks. The larvae are then screened out and the % mortality of the larvae is determined. Each corn root is washed and after washing the roots are rated on a scale of 1-6 using the Iowa Root Rating Method developed at Iowa State University (Hills and Peters, 1971). Some of the results are indicated below in Table 3. The greenhouse assay use 2 gallon pot with 20 mesh corn field soil. Forty corn rootworm larvae are placed in each pot when corn plant was at 4 to 5-leaf stage, and allowed to feed for 2 to 3 weeks. Each corn root is washed and after washing the roots are rated on a scale of 1-6 using the Iowa Root Rating Method developed at Iowa State University (Hills and Peters, 1971). Some of the results are indicated below in Table 3.

TABLE 3

Corn Rootworm Field Efficacy Results	
Treatment	Root Rating
Captan only	4.7
Overcoating only	4.5
Example 1(a)	3.43
Example 1(b)	3.0
Example 1(c)(2)	3.2
Example 1(d)	3.55
Lorsban™ 15 G soil T-band*	3.0
Aztec™ 2.1 G soil T-band*	2.8
Force™ 1.5 G soil T-band*	2.6

Lorsban 15 G = 15% 0,0-diethyl-0-(3,5,6-trichloro-2-pyridinyl)phosphorothioate, DowElanco.
 Aztec 2.1 G = 2% 0-[2-(1,1-dimethylethyl)-5-pyrimidinyl-o-ethyl o -(1-methylethyl) phosphorothioate and 0.1% cyano(4-fluoro-3-phenoxyphenyl)methyl 3-(2,2-dichloroethyl)-2,2-dimethylcyclopropanecarboxylate, Bayer.
 Force 1.5 G = 1.5% [1α, 3α(Z)]-(±)-(2,3,5,6-tetrafluoro-4-methylphenyl)methyl 3-(2-chloro-3,3,3-trifluoro-1-propenyl)-2,2-dimethylcyclopropanecarboxylate, Zeneca, Inc.

[0094]

TABLE 4

Corn Rootworm Bioassay Results			
Treatment	Growth Chamber		Greenhouse
	% Mortality	Root Rating	Root Rating
Captan only	27	5.57	4.02
Example 1(a)	87	3.55	3.15
Example 1(c)	96	1.44	2.15
Example 1(d)	80	3.75	—

EXAMPLE 6

[0095] Corn Rootworm Efficacy Studies:

[0096] Maize seed is treated as described in Examples 1 and 2. A combination of methyl cellulose and polyvinyl alcohol is used as a binder and Celite 266 is used as filler for insecticide solution, which contains the compound of formula II as the insecticide. Bioassay methods are as described in Example 4. Root rating was based on the Iowa State 1-6 scale.

TABLE 5

Corn Rootworm Bioassay Results		
Treatment (mg/seed)	Growth Chamber (16 oz cup) Root Rating	Greenhouse (2 gal pot) Root Rating
Control	5.69	4.54
Tefluthrin 50% (1.0)	2.30	3.52
Formula II		
(0.01)	5.30	
(0.03)	4.30	
(0.3)	2.70	
(0.6)	2.10	2.73
(0.8)	2.30	
(1.0)	2.10	
(1.2)		3.08
(1.5)		2.96
(2.0)		2.90
Raze*/Formula II		
(0.01)	2.48	2.63
(0.3)	1.87	2.34

*Raze™ is a tefluthrin seed treatment formulation (0.13 mg/seed) marketed by Wilber Ellis for wireworm control. Raze™ seed treatment alone did not show any significant activity against CRW.

EXAMPLE 7

[0097] The following seed coating (hereinafter "Formula X") was prepared and applied to a corn seed at a dose rate of 62.5%:

TABLE 6

COATING COMPONENT	GRAMS/1000 SEEDS
Force™ (tefluthrin) 29% a.i.	4.00
Proised red™ (colorant)	0.10
Actisil™ (bentonite)	2.00
Airflex™ 500 (vinyl acetate/ethylene copolymer emulsion)	0.75
Gantrez™ AN 139 @ 10% (methyl vinyl ether/maleic anhydride copolymer)**	2.50
Water	1.65

This coating formulation included a 75/25 (talc/mica WG325) dry charge of 0.25 g.

**Also known as Agrimer VEMA AN 990™.

EXAMPLE 8

[0098] The following film coating was prepared and applied to a corn seed:

TABLE 7

COATING COMPONENT	GRAMS/1000 SEEDS
Maxim™ (fludioxonil)/Apron™ (metalaxyl)	0.030
Proised red™ (colorant)	0.400
Airflex™ 500 (vinyl acetate/ethylene copolymer emulsion)	0.500
Gantrez™ 10% (methyl vinyl ether/maleic anhydride copolymer)	0.500
Afflair™ (mica/titanium dioxide)	0.300
calcium stearate	0.015

TABLE 7-continued

COATING COMPONENT	GRAMS/1000 SEEDS
titanium dioxide	0.250
polyethylene glycol 8000	0.010
Water	4.500

This coating formulation included 0.25 g of a 75/25 dry charge.

EXAMPLE 9

[0099] The following seed coating was applied to a corn seed at a concentration of 1.0 mg a.i./seed:

TABLE 8

COATING COMPONENT	GRAMS/1000 SEEDS
compound of formula II	4.839
Gantrez AN139 @ 10%	4.247
Airflex 500	1.314
Actisil	1.676
Colorant	0.200
Water	6.399

EXAMPLES 10-12

[0100] Following the procedure set forth in Example 5, field trials were conducted at three separate locations, two in Minnesota (Tables 9 and 11) and one in Illinois (Table 10). In addition to the use of the Root Rating Method, roots were evaluated in terms of "Consistency", i.e., the percentage of the corn plant roots having a root rating of less than 3.0 (above which, the roots have been, at least in part, eaten off by corn root worm larvae).

TABLE 9

TREATMENT	ROOT RATING	CONSISTENCY
Control	5.28	0.0
Formula X (50.0%)	2.97	66.7
Formula X (62.5%)	3.25	48.3
Formula X (75%)	2.84	76.7
Formula X (50%) w/Reax*	3.68	30.0
Formula X (62.5%) w/Reax	3.36	51.7
Formula X (75.0%) w/Reax	3.28	46.7
Aztec™ 2.1 G T-Band	2.55	96.7
Counter™ 20 G T-Band	3.38	56.7
Force™ 3G T-Band	3.34	62.1
Lorsban 15G T-Band	2.90	80.0

*Reax (Reax 85A™) is a sulfonated kraft lignin (lignosulfonic acid, sodium salt)

[0101]

TABLE 10

TREATMENT	ROOT RATING	CONSISTENCY
Control	4.00	32.5
Formula X (50.0%)	2.33	90.0
Formula X (62.5%)	2.60	90.0
Formula X (75%)	2.68	82.5
Formula X (50%) w/Reax	2.80	87.5
Formula X 62.5% w/Reax	2.78	90.0
Formula X (75.0%) w/Reax	2.78	80.0
Aztec™ 2.1G T-Band	3.05	80.0
Counter™ 20G T-Band	3.15	72.5

TABLE 10-continued

TREATMENT	ROOT RATING	CONSISTENCY
Force TM TM 3G T-Band	2.75	87.5
Lorsban 15G T-Band	3.60	50.0

[0102]

TABLE 11

TREATMENT	ROOT RATING	CONSISTENCY
Control	3.42	46.2
Formula X (50.0%)	2.32	97.5
Formula X (62.5%)	2.56	87.2
Formula X (75%)	2.59	87.2
Formula X (50%) w/Reax*	2.43	87.5
Formula X (62.5%) w/Reax	2.58	92.3
Formula X (75.0%) w/Reax	2.21	100.0
Aztec TM 2.1G T-Band	2.28	100.0
Counter TM 20G T-Band	2.67	85.0
Force TM 3G T-Band	2.56	94.9
Lorsban 15G T-Band	2.60	95.0

EXAMPLES 13-14

[0103] IR Corn/IST Trials

[0104] In order to evaluate the effect against black cutworm that the seed coating of the invention would have on corn seed transformed with an insect resistance gene compared to a) non-coated and b) non transformed seed, three field trials were conducted, two in Illinois and one in Missouri, with three different corn events, either transformed with VIP3 or non-transformed. The effect was measured in terms of percentage of corn plants a) cut by the insect and b) killed by the insect

TABLE 12

TREATMENT	Black Cutworm Trials Percent Dead Plants		
	Trial 1	Trial 2	Trial 3
VIP3A* Event 1	1.67	0.0	2
VIP3A* Event 1 + Formula X	0.00	0.0	2
Non-transformed Event 1	6.66	0.0	19
Non-transformed Event 1 + Formula X	0.00	0.0	10
VIP3A* Event 2	0.00	0.0	4
VIP3A* Event 2 + Formula X	0.00	0.0	2
Non-transformed Event 2	1.67	1.6	30
Non-transformed Event 2 + Formula X	3.33	0.0	20
VIP3A* Event 3	3.34	1.0	5
VIP3A* Event 3 + Formula X	1.67	0.0	9
Non-transformed Event 3	6.92	0.0	22

*Vegetative insecticidal protein (VIP3 transgene)

[0105]

TABLE 13

TREATMENT	Black Cutworm Trials Percent Cut Plants		
	Trial 1	Trial 2	Trial 3
VIP3A* Event 1	1.67	6.98	8
VIP3A* Event 1 + Formula X	1.67	5.56	9

TABLE 13-continued

TREATMENT	Black Cutworm Trials Percent Cut Plants		
	Trial 1	Trial 2	Trial 3
Non-transformed Event 1	10.00	11.21	36
Non-transformed Event 1 + Formula X	1.67	12.65	22
VIP3A* Event 2	1.67	5.94	5
VIP3A* Event 2 + Formula X	3.34	3.22	8
Non-transformed Event 2	6.67	10.15	45
Non-transformed Event 2 + Formula X	5.00	10.36	29
VIP3A* Event 3	6.67	26.03	22
VIP3A* Event 3 + Formula X	5.00	2.88	28
Non-transformed Event 3	12.43	16.00	45

*Vegetative insecticidal protein (VIP3 transgene)

What is claimed is:

1. An insecticidal coating for a corn seed comprising:

a) one or more binders selected from the group consisting of polymers and copolymers of polyvinyl acetate, methyl cellulose, polyvinyl alcohol, methyl vinyl ether, malic anhydride, vinylidene chloride, acrylic, cellulose, polyvinylpyrrolidone and polysaccharide;

b) an insecticidally effective amount of an insecticide comprising tefluthrin or an oxadiazine or a combination thereof; and

c) one or more fillers wherein the binder forms a matrix for the insecticide and filler and is present in an amount effective to prevent or reduce the phytotoxic effect on the seed caused by the insecticide.

2. The coating of claim 1 further comprising a film overcoat.

3. The coating of claim 1 wherein the filler is an absorbent filler selected from the group consisting of bentonite, diatomaceous earth, perlite, silica, calcium carbonates, and mixtures thereof.


4. Corn seed coated with the coating according to claim 1.

5. The corn seed according to claim 4 comprising one or more stably incorporated transgenes that code on expression for one or more proteins that confer pesticidal resistance to the seed.

6. The corn seed according to claim 5 wherein the transgenes comprise genes which code on expression for pesticidally active proteins selected from the group consisting of crylAb, crylAc, crylF, cry9c, cry2A, cry6A and VIP3.

7. The coating of claim 1 wherein the insecticide further comprises one or more insecticides selected from the group consisting of imidacloprid, an organophosphate, phenylpyrazole or pyrethroid other than tefluthrin, or combinations thereof.

8. The coating of claim 1 wherein the insecticide is an oxadiazine compound of the general formula (I)



(I)

in which


A is an unsubstituted or mono- to tetrasubstituted, aromatic or non-aromatic, monocyclic or bicyclic heterocyclic radical, where one to two of the substituents of A can be selected from the group consisting of halo-C₁-C₃alkyl, cyclopropyl, halocyclopropyl, C₂-C₃alkenyl, C₂-C₃alkynyl halo-C₂-C₃alkenyl, halo-C₂-C₃alkynyl, halo-C₁-C₃alkoxy, C₁-C₃alkylthio, halo-C₁-C₃alkylthio, allyloxy, propargyloxy, allythio, propargylthio, haloallyloxy, haloallylthio, cyano and nitro, and one to four of the substituents of A can be selected from the group consisting of C₁-C₃alkyl, C₁-C₃alkoxy and halogen;

R is hydrogen, C₁-C₆alkyl, phenyl-C₁-C₄alkyl, C₃-C₆cycloalkyl, C₂-C₆alkenyl or C₂-C₆alkynyl; and

X is N—NO₂ or N—CN,

or, if appropriate, a tautomer thereof, in each case in free form or in salt form, terbufos, chlorpyrifos, fipronil, tefluthrin, chloroethoxyfos or tebufipirimfos, or combinations thereof.

9. The coating of claim 8 wherein the insecticide is the compound of formula II



(II)

and optionally tefluthrin.

10. The coated corn seed of claim 4 further comprising a fungicide.

11. The coating of claim 1 further comprising a plasticizer.

12. The coating of claim 1 further comprising a bird repellent compound.

13. The coating of claim 1 wherein the filler comprises a mixture of diatomaceous earth and silica.

14. The coating of claim 1 wherein the binder is in the range of 0.01 to 15% of the weight of the corn seed, the filler is in the range of 0.01 to 50% of the weight of the corn seed, and the insecticide is in the range of 0.005 to 50% of the weight of the corn seed.

15. An insecticidal coating for a corn seed comprising a binder in an amount from about 0.01 to about 15% of the weight of the seed wherein said binder is a vinyl acetate-ethylene copolymer or polymer or copolymer of vinylidene

chloride, a filler in an amount from about 0.01 to about 50% of the weight of the seed wherein said filler is diatomaceous earth and amorphous silica, and an insecticidally effective amount of a compound effective against corn root worm or its larvae selected from compounds of the group consisting of the compounds of formula II, imidacloprid, tefluthrin, and combinations thereof, and in the range from about 0.005 to about 50% of the weight of the seed.

16. A method of controlling insect pests from damaging a crop plant comprising applying to a seed an insecticidal coating according to claim 1.

17. A method of controlling insect pests from damaging a crop plant comprising applying to a seed an insecticidal coating according to claim 7.

18. A method of controlling insect pests from damaging a corn plant comprising transforming the corn plant or corn seed with one or more transgenes that code on expression for one or more proteins that confer pesticidal resistance to the seed, and applying to the corn seed the insecticidal coating according to claim 1.

19. The method according to claim 18 wherein the insect is a Lepidopteran or Coleopteran insect.

20. The method according to claim 19 wherein the insect is corn root worm.

21. A method of preparing coated corn seeds comprising:

a) mixing a binder selected from vinyl acetate-ethylene copolymers or polymers or co-polymers of vinylidene chloride or mixtures thereof, and a filler selected from bentonite, diatomaceous earth, perlite, silica, calcium carbonates and mixtures thereof, with a terbufos, oxadiazine, chlorpyrifos, tefluthrin, fipronil or tebufipirimfos insecticide;

b) applying the mixture to a seed;

c) allowing the mixture to dry on said seed; and

d) applying a film overcoating to the seed.

22. A seed coated according to the method of claim 21.

23. A method of protecting emerging seedlings of a crop plant from one or more insect organisms in the seed growing soil environment which comprises applying to the seeds of the crop plant an insecticidally effective amount of a coating according to claim 1.

24. The method of claim 23 wherein the crop plant is a corn plant transformed with a transgene coding for a protein that confers pesticidal resistance to the seed.

25. The method of claim 23 wherein the insect is a Lepidopteran or Coleopteran insect.

26. The method of claim 25 wherein the insect is corn root worm.

27. A method for reducing the resistance of a target insect to a transgenic insecticidal protein comprising:

a) transforming a corn seed with one or more transgenes that code on expression for a protein or combination of proteins having insecticidal activity; and

b) coating the seed with the seed coating according to claim 1.

28. The method according to claim 27 wherein the one or more transgenes comprise genes that code on expression for proteins selected from the group consisting of cry1Ab, cry1Ac, cry1F, cry9c, cry2A, cry6A and VIP3.

* * * * *