



# JOURNAL *of the* **AMERICAN** **OIL CHEMISTS'** **SOCIETY**

FORMERLY OIL AND SOAP



JUNE 1976 • 221-462 • VOL. 53, NO. 6

COMPLETE PROCEEDINGS

**World Conference**

on

**Oilseed and Vegetable Oil  
Processing Technology**

Amsterdam, The Netherlands  
March 1-5, 1976

ANALYTICAL CHEMISTRY  
AND METHODOLOGY OF LIPIDS  
PHYSICAL AND CHEMICAL  
PROPERTIES OF FATS

SOAPS AND DETERGENTS

FATTY ACIDS  
AND FATTY CHEMICALS  
DRYING OILS AND  
PROTECTIVE COATINGS

LIPID BIOCHEMISTRY  
AND NUTRITION  
EDIBLE FATS AND OILS

SOLVENT EXTRACTION  
FAT AND OIL PROCESSING

JAOCA7 53(6) 221-426(1976)

CARGILL EXHIBIT 1008  
Cargill v. Bunge Loders Crokiaan

# The JOURNAL of the AMERICAN OIL CHEMISTS' SOCIETY

Formerly Published As

Chemists' Section, Cotton Oil Press, 1917 to 1924; Journal of the  
Oil and Fat Industries, 1924 to 1931; Oil & Soap, 1932 to 1947

June, 1976 • Volume 53, No. 6

## AOCS OFFICERS

**PRESIDENT** . . . . . F.B. White  
Manager, International Operations, Process Plants  
Division, Foster Wheeler Corp., Livingston, NJ.

**VICE PRESIDENT** . . . . . T.H. Applewhite  
Manager, Edible Oil Products Laboratory, Kraftco  
Corp., Research and Development, Glenview, IL.

**SECRETARY** . . . . . G.A. Jacobson  
Division Head, Basic Research Department, Campbell  
Institute for Food Research, Camden, NJ.

**TREASURER** . . . . . D. Firestone  
Research Chemist, Division of Chemistry and Physics,  
Bureau of Foods, Food and Drug Administration,  
Washington, DC.

## JOURNAL STAFF

**EDITOR** . . . . . A.R. Baldwin

### ASSOCIATE EDITORS

|                 |                |                |
|-----------------|----------------|----------------|
| T.H. Applewhite | M.E. Ginn      | W.M. Linfield  |
| N.R. Artman     | L.A. Goldblatt | E.G. Perkins   |
| W.J. Beach      | S.F. Herb      | E.E. Rice      |
| G.C. Cavanagh   | T.J. Jacks     | J.A. Robertson |
| S.S. Chang      | E. Jungerman   | W.H. Tallent   |
| J.C. Cowan      |                | I.A. Wolf      |

**MANAGING EDITOR** . . . . . J.C. Lyon

**PRODUCTION MANAGER** . . . . . D.I. Peters

**STAFF EDITOR** . . . . . M.F. Grzyb

**ABSTRACTS EDITOR** . . . . . S. Koritala

**BOOK REVIEW EDITOR** . . . . . J.F. Gerecht

**INDEX EDITOR** . . . . . M.W. Formo

**ADVERTISING DIRECTOR** . . . . . K.T. Zilch

**TECHNICAL CORRESPONDENT** . . . . . J.G. Endres

**CIRCULATION MANAGER** . . . . . J. Nelson

Published monthly by  
THE AMERICAN OIL CHEMISTS' SOCIETY  
508 South Sixth St., Champaign, Illinois 61820  
Phone: (217) 359-2344

Second class postage paid at  
Champaign, Illinois

Copyright 1976 by  
THE AMERICAN OIL CHEMISTS' SOCIETY

Subscriptions are \$40 for one year, \$77 for two  
years, and \$111 for three years. Foreign postage is \$3  
extra per year. Air mail postage is \$30 extra per year.  
Claims for copies lost in the mails must be received  
within 30 days (90 days foreign) of the date of issue.  
Notice of change of address must be received two  
weeks before date of issue.

The American Oil Chemists' Society assumes no  
responsibility for statements or opinions of contribu-  
tors to its columns.

The Journal is indexed in the Applied Science and  
Technology Index of the H.W. Wilson Company, New  
York.

**NEWS**—Submit to AOCS six weeks preceding month  
of issue.

**MANUSCRIPT PUBLICATION**—Submit to Dr. A.R.  
Baldwin, Cargill, Inc., Cargill Building, Minneap-  
olis, Minn. 55402.

### ADVERTISING INSTRUCTIONS AND DEADLINES

**ADVERTISING MANAGER** . . . . . C. Sharp  
508 South Sixth Street, Champaign, Illinois 61820  
Telephone . . . . . Champaign (217) 359-8028

Closing date 12th of month preceding date of  
issue. Insertion orders received after closing will be  
accepted by the production manager at advertiser's  
risk. AOCS cannot guarantee positioning or proofs for  
ads received after closing. Cancellation date 12th of  
preceding month. Ship mechanical material direct to  
Kowa Graphics Incorporated, 510 N. Hickory Street,  
Champaign, Illinois 61820.

## News

- 298A Introduction to this World Conference Proceedings Issue of JAOCS
- 302A Cosponsoring and Supporting Organizations
- 306A Steering Committee and Program Committee
- 310A Conference Officers
- 312A Local Committee
- 314A Session Chairpersons
- 316A Pictorial Review of the Conference
- 333A Support for Participants from Developing Countries
- 336A AOCS Official Referee Chemists: A Long-Valued Society Professional Service
- 340A AOCS Uniform Methods and Smalley Committees Oversee Vital Services to Industry
- 348A Exhibitors: Products and Services
- 364A List of Delegates
- 378A Classified Advertising & Advertisers' Index

## Technical

- 221-222 Welcoming Address, by Dr. I. Samkalden
- 222 Call to Order, by T.H. Applewhite and C.J.M. Meershoek
- 223 Introduction to the Technical Meeting, by T.H. Applewhite

### SESSION I — ECONOMIC CONSIDERATIONS

- 224-231 World Supply and Demand Situation for Oilseeds, Oils, and Meals, by S. Mielke
- 232-234 Referee Chemists and Official Methods, by W.E. Link
- 235-237 International Arbitration, by E. Faure
- 238-240 Effect of Shipping on Quality of Seeds, Meals, Fats, and Oils, by M.R. Spencer

### SESSION II — PROCESSING VEGETABLE OIL-BEARING MATERIALS

- 241-247 Characteristics and Composition of Vegetable Oil-bearing Materials, by A. Langstraat
- 248-250 Loading, Unloading, Storage, Drying, and Cleaning of Vegetable Oil-bearing Materials, by E.H. Gustafson
- 251-255 Oil Mill Energy Sources and Power Balances in a Palm Oil Operation, by B.E. Fonade
- 256-258 Continuous Processing of Palm Fruit, by B. de Ramecourt
- 259-260 Processing Peanuts and Cottonseed, by E.J. Garcia M.
- 261-264 Processing High Oil Content Seeds in Continuous Screw Presses, by J.A. Ward
- 265-270 Current Equipment for Mechanical Oil Extraction, by L.H. Tindale and S.R. Hill-Haas
- 271-274 Cleaning, Cracking, Dehulling, Decorticating, and Flaking of Oil-bearing Materials, by J.P. Galloway
- 275-278 Batch and Continuous Solvent Extraction, by E. Bernardini
- 279-282 Continuous Solvent Extraction of Soybeans and Cottonseed, by R.P. Hutchins

- 283-285 Continuous Solvent Extraction of Sunflower Seed, Groundnuts, Palmkernels, Rapeseed, and Copra, by W. Stein and F.-W. Glaser
- 286-290 Survey of Current Solvent Extraction Equipment, by E.D. Milligan
- 291-292 Energy Conservation in Solvent Extraction Plants, by H.P.J. Jongeneelen
- 293-294 Safety in Solvent Extraction Plants – USA, by J.E. Heilman
- 295-298 Safety in Solvent Extraction Plants – Europe, by C.E.M. Critchfield
- 299-301 Processing Effects on Oil Quality, by J.S.R. Ohlson
- 302-304 Protein Products and Hulls for Animal Foods, by R. Roberts
- 304-309 Whole Seed Processing by Extrusion Cooking, by E. Nielsen
- 310-315 Oilseed Flour for Human Food, by F.G. Bastiaens
- 316-320 Standards for Protein Based Foods in Developing Countries, by K.T. Achaya
- 321-324 Oilseed Vegetable Protein Concentrates, by D.W. Johnson
- 325-326 Texturized Protein Products, by R. Brian
- 327-331 New Technology in Oilseed Proteins, by K.W. Becker and E.A. Tiernan

### SESSION III – PROCESSING VEGETABLE FATS AND OILS

- 332-333 Crude Oil Handling and Storage, by J.P. Burkhalter
- 334-337 Special Problems in Small Scale Operations, by G.M. Neumunz
- 338 Small Scale Processing of Vegetable Oils, by C. Baumann
- 339-341 Bleaching Practices in Europe, by H.B.W. Patterson
- 342-343 Bleaching Practices in the U.S., by E.H. Goebel
- 344-346 Degumming, Refining, Bleaching, and Deodorization Theory, by J.C. Cowan
- 347-352 Degumming and Refining Practices in the U.S., by R.A. Carr
- 353-357 Degumming and Refining Practices in Europe, by B. Braae
- 358-360 Steam Refining, by F.E. Sullivan
- 361-363 Miscella Refining, by G.C. Cavanagh
- 364-369 Deodorization 1975, by C.T. Zehnder
- 370-381 Automation in Vegetable Oil Refineries, by A.J. Duff

### SESSION IV – FATS AND OILS MODIFICATION

- 382-389 Hydrogenation of Edible Oils, by J.W.E. Coenen
- 390-392 Interesterification of Edible Oils, by H.H. Hustedt
- 393-396 Fractionation and Winterization of Edible Fats and Oils, by H.P. Kreulen
- 397-399 Blending, Chilling, and Tempering of Margarines and Shortenings, by A.J. Haighton
- 400-407 Food Emulsifiers: Surface Activity, Edibility, Manufacture, Composition, and Application, by J.B. Lauridsen

### SESSION V – FINISHED PRODUCTS

- 408-409 Finished Oil Handling and Storage in the U.S., by L.M. Wright
- 410-413 Finished Oil Handling and Storage in Europe, by G.M.R. Johansson
- 414-416 Salad and Cooking Oils, by B. Lesieur
- 417-420 Margarine Oils, Shortenings, and Vanaspati, by K.-F. Gander
- 421-424 Specialty Fats, by F.R. Paulicka
- 425-427 Lecithin Production and Properties, by W. Van Nieuwenhuyzen
- 428-429 Utilization of Lecithin, by P.M. Scocca
- 430-436 Antioxidants for Vegetable Oils, by E.R. Sherwin

### SESSION VI – OIL PROCESSING WASTES AND BY-PRODUCTS

- 437-442 Use or Disposal of By-Products and Spent Material from the Vegetable Oil Processing Industry in the U.S., by K.S. Watson and C.H. Meierhoefer
- 443-445 Use or Disposal of By-Products and Spent Material from the Vegetable Oil Processing Industry in Europe, by C.H. Svensson
- 446-448 Liquid Waste Treatment in the Vegetable Oil Processing Industry – European Practices, by G. Choffel
- 449-458 Liquid Waste Treatment in the Vegetable Oil Processing Industry – U.S. Practices, by G.N. McDermott
- 459-460 Reflections on the Conference – European View, by J. Pasman
- 461 Reflections on the Conference – U.S. View, by T.H. Applewhite

# Interesterification of Edible Oils

H.H. HUSTEDT, Chemical and Technical Director, Noble & Thörl GmbH, Hamburg-Harburg, West Germany



## ABSTRACT

Types of interesterification discussed are (a) interchange between a fat and free fatty acids, in which the most important reaction is the introduction of acids of low mol wt into a fat with higher fatty acids; (b) interchange between a fat and an alcohol, e.g., with glycerol, in order to produce emulsifiers like monoglycerides; (c) rearrangement of fatty acid radicals in triglycerides, the so-called transesterification which in recent years has taken on the same importance as hydrogenation or fractionation. In natural fats, the fatty acid radicals are not usually randomly distributed but become so by rearrangement; the distinctive physical properties of natural fats and oils can be changed within limits by this transesterification. Well-known examples are cocoa butter, palm oil, and lard. More important is the transesterification of a mixture of different fats and oils; e.g., the combination of hydrogenation and interesterification allows the production of a solid fat with high linoleic acid content. The composition of glycerides after random interesterification can be calculated by formulas.

Distinct from random is such directed interesterification. This is done by working at low temperatures that glycerides with higher melting point crystallize from the reaction mixture. Directed interesterification can be combined with fractionation, for instance, to get a higher yield of liquid fraction from palm oil than is obtained by fractionation alone.

The transesterification process can be performed in a batch or continuously. A small amount of metallic sodium or sodium ethylate is used as catalyst, which is destroyed by water or acid and removed after the reaction.

Since 1950, the importance of the interesterification process has grown continuously, and today it is used as frequently as hydrogenation in many areas.

The term *interesterification* refers to that reaction of fats and oils in which fatty acid esters react with other esters or fatty acids to produce new esters by an interchange of fatty acid groups. It is this part which I will describe, and no reference will be made to reactions of fatty acid esters with alcohols.

Since the fatty acid distribution in fats can be changed




FIG. 1. Minimum distribution of fatty acids. S = saturated and U = unsaturated fatty acids.

by interesterification, I will first briefly discuss the possible distributions of the fatty acids in triglyceride molecules. For this purpose, I will assume the simple case that we have only two different types of fatty acids, namely, saturated (S) and unsaturated (U). The minimum distribution as shown in Figure 1 is the simplest possibility. For instance, a composition which consists of 1/3 saturated and 2/3 unsaturated fatty acids is considered to be a mixture of only two types of glycerides, of which 1/3 is  $GS_3$  (glyceride with three saturated acids) and 2/3 is  $GU_3$  (glycerides with three unsaturated acids). Another simple distribution is the even distribution (Fig. 2). In the example of 1/3 S and 2/3 U, there is only one type  $GSU_2$ . In another example of 1/2 S and 1/2 U, the composition is 1/2  $GSU_2$  and 1/2  $GS_2U$ .

Random distribution is most frequent because in a liquid phase the normal ester rearrangement or esterification of acids with glycerine proceeds at random with the attainment of an equilibrium composition, which in our simple case of only two types of fatty acids can be calculated by using the formulas shown in Figure 3. Figure 4 shows the curves of these formulas. For instance, 50% S and 50% U give 12.5%  $GS_3$ , 12.5%  $GU_3$ , 37.5%  $GSU_2$ , and 37.5%  $GS_2U$ .

At this point, reference should be made to isomerism. In




FIG. 2. Even distribution of fatty acids. S = saturated and U = unsaturated fatty acids.

|                                                                   |                                                                            |
|-------------------------------------------------------------------|----------------------------------------------------------------------------|
| $\begin{matrix} S \\ S \\ S \end{matrix}$                         | Percent triglyceride $GS_3 = \frac{S \cdot S \cdot S}{10000}$              |
| $\begin{matrix} U \\ U \\ U \end{matrix}$                         | Percent triglyceride $GU_3 = \frac{U \cdot U \cdot U}{10000}$              |
| $\begin{matrix} S & U \\ U & S \end{matrix}$                      | Percent triglyceride $GSU_2 = 3 \cdot \frac{S \cdot U \cdot U}{10000}$     |
| $\begin{matrix} U & S \\ S & U \end{matrix}$                      | Percent triglyceride $GUS_2 = 3 \cdot \frac{U \cdot S \cdot S}{10000}$     |
| $\begin{matrix} A & A & B \\ B & C & A \\ C & B & C \end{matrix}$ | Percent triglyceride $GABC = 6 \cdot \frac{A \cdot B \cdot C}{10000}$ etc. |

FIG. 3. Formulas for calculating the random distribution of fatty acids. s,u,a,b,c = molar percentages of the fatty acids S (saturated), U (unsaturated), A, B, and C; G = glycerine.




FIG. 4. Random distribution of fatty acids. S = saturated and U = unsaturated fatty acids.




FIG. 5. Glyceride structure of cocoa butter before and after random interesterification.

the glyceride  $GS_2U$ , the unsaturated acid could be located in two different positions: the middle  $\beta$ -position (2) or the  $\alpha$ -position (1). After interesterification of an ester, the random distribution occurs whereby in the molecule type  $GS_2U$  2/3 of the unsaturated acid appears in position 1 and 1/3 in position 2.

Cocoa butter is a typical example in this respect (Fig. 5). The fatty acid composition consists of ca. 2/3 saturated acids (stearic and palmitic acids) and 1/3 unsaturated acids. The glyceride composition of the natural cocoa butter may be called an even distribution because it contains mainly glycerides with one unsaturated acid in position 2 and two saturated acids in position 1. This is the explanation for the unusual physical behavior of cocoa butter. By interesterification cocoa butter will lose these typical properties because the fat then has a random distribution with only a slight amount (12.5 %) of the original cocoa butter glycerides. Therefore, cocoa butter cannot be produced by interesterification or by esterification of the correct fatty acid composition.

Before uses of the interesterification process can be discussed, some mention must be made of the technological procedure. Interesterification may proceed in two ways: random or directed. Both procedures may be carried out in a discontinuous or in a continuous way.

When random rearrangement is preferred, the fat is heated to ca. 80 C in the presence of a catalyst and with rapid stirring for half an hour. The most common catalysts are sodium methylate or sodium-metal. It is essential that the fat contain no substances which destroy the catalyst. The amount of catalyst required depends upon several factors. If the water content is < 0.01%, the free fatty acid level below 0.05%, and the peroxide content is also very low, the addition of only 0.1% of catalyst is sufficient. Therefore, the fat must be well refined, dried, and heated to




FIG. 6. Solid fat index of palm oil before and after interesterifications. S = saturated and U = unsaturated fatty acids.

150 C under a nitrogen blanket before the catalyst is added. Experience indicates that the addition of each additional 0.1% catalyst results in a loss of ca. 1% neutral fat.

When the directed interesterification process is applied, some of the fatty acids or their glycerides are removed from the equilibrium reaction. The procedure commonly used is to continuously distill the acids with low mol wt from the fat, such as a mixture of coconut oil with an excess of free fatty acids, or, even more important, the interesterification is carried out at such low temperatures that higher melting glycerides which are formed by the interesterification get separated from the residual liquid phase after crystallization. During rearrangement and continuous crystallization, the formation of an increasing amount of high melting glycerides occurs. This does not result in random interesterification but, more or less, the minimum distribution of fatty acids in which a mixture of glycerides exists containing saturated and unsaturated acids as mentioned above.

Since the rate of the reaction is slow due to the low temperature, and since the fats tend to supercool, the reaction time is longer than for nondirected interesterification reactions. The reaction time required could be 1 day, and the process is, of course, expensive. Sodium methylate or a liquid alloy of sodium-potassium is used as catalyst. At the end of the reaction, the catalyst must, of course, be inactivated with water or dilute acid before the fat will be heated or the solids will be separated.

In many cases, the directed interesterification increases the melting point, since the melting point of a fat depends to some degree upon the content of saturated glycerides. By this method it is possible to convert a liquid oil into a plastic product with the consistency of a shortening. Directed interesterified sunflower oil has a melting point which is slightly below 20 C and a solid fat index of 10 at 0 C and of 5 at 15 C. If this fat is blended with 5% hard-stock of hydrogenated oil, a solid fat with > 60% essential fatty acids can be produced.

A winterized oil which contains no linolenic acid can be produced by hydrogenation and directed interesterification followed by a fractionation. The yield of the liquid phase from directed interesterified palm oil is higher than from untreated oil, as shown in Figure 6. The soap, which is formed by destroying the catalyst with water, can be used in the fractionating step as tenside.

Since the distribution of fatty acids in most of the natural fats and oils occurs not at random, the nondirected interesterification also changes the distinctive physical properties of the fats, such as the melting point, the solid fat index, the crystallization tendency, the texture, etc. The graininess of ordinary lard is due to large crystals of disaturated glycerides (OPS), in which the palmitic acid is in the 2 (middle)-position and the oleic acid in the 1-position (OPS). Random interesterification reduces the amount of




FIG. 7. Solidification curves of sunflower and coconut oils interesterified with hard fats.

these glycerides. In this case, directed interesterification will eliminate the graininess and simultaneously create a hardstock, so that soft lard will become an improved shortening.

A hard butter with a low melting point (32 C) and also with sharp melting characteristics is obtained by random interesterification of a hydrogenated lauric oil with the melting point of 40-42 C.

Use of interesterification of fat and oil mixtures in the industry is very important and is extremely versatile. It is impossible to describe all the possible applications. However, I would like to explain some typical examples.

One principal advantage of interesterification is that use of this process allows the combination of the properties of different fats and oils. Many of these interesterified fat mixtures are used in margarine and shortening.

Lauric oils, such as palmkernel and coconut oil, have low melting points and short plastic ranges; a margarine with high content of these fats will be very hard if kept in the refrigerator but will melt partly at room temperature. If it becomes desirable to use a large percentage of coconut oil in margarine—and, according to the present low price of coconut oil, one must—the glycerides with three lauric acid moieties must be removed or decreased. This is possible by




FIG. 8. Solid fat index (%). Hydr. Co = hydrogenated coconut oil.

interesterifying the coconut oil with another type of fat—for instance, with palm oil. Thereby, a margarine may be produced which consists of 60% of the interesterified palm oil-coconut oil mixture, which is then blended with 40% of an oil, e.g., sunflower oil.

Intesterification is especially useful when it is desired to combine solid and liquid fats. Many patents relate to fats or margarines having a high polyunsaturated (linoleic) fatty acid content. In these products, the solid fat fraction used could be animal fats, hydrogenated fats, palm oil, or synthetic tristearin. The effect of interesterification can be seen in Figure 7. This shows the solidification curves of a mixture of a hydrogenated oil having a melting point of 70 C and a liquid oil before and after interesterification. The congeal point decreases from 50 C to 31 C.

The solid fat indices of an interesterified hardstock of hydrogenated soybean oil and coconut oil blended with 80% liquid oil are shown in Figure 8. Increasing the percentage of oil before and simultaneously decreasing the percentage after interesterification causes the melting point and the solid fat index to drop. This blended fat represents a good margarine having a high content of essential fatty acids and a low content of *trans*-double bonds.