

2005

Volume 14 Number 4

Asia Pacific journal of clinical nutrition.
DUP-General Collection
W1 AS139BG
v. 14, no. 4
2005

Asia Pacific Journal of Clinical Nutrition

Editor-in-Chief

Mark Wahlqvist, Melbourne

Co-Editors

Akira Okada, Osaka

Duo Li, Hangzhou



PROPERTY OF THE
NATIONAL
LIBRARY OF
MEDICINE

INCLUDED IN
MEDLINE
and
CURRENT CONTENTS

HEC PRESS

www.healthyeatingclub.org/APJCN/

CARGILL EXHIBIT 1015
Cargill v. Bunge Loders Croklaan

Asia Pacific Journal of Clinical Nutrition

2005, HEC PRESS, Melbourne, Australia ISSN 0964-7058

Asia Pacific Journal of Clinical Nutrition

THE JOURNAL OF THE ASIA PACIFIC CLINICAL NUTRITION SOCIETY

APJCN 2005: Volume 14 (Number 4) : 294-440 ISSN 0964-7058

EDITORS:

Professor Mark Wahlqvist MD, Asia Pacific Health and Nutrition Centre, Monash Asia Institute, 8th Floor, Menzies Building, Monash University, Wellington Road, Clayton, Melbourne, Victoria 3800, AUSTRALIA. Fax: (+61) 3 9905 8146; Email: Mark.Wahlqvist@adm.monash.edu.au.

Professor Akira Okada MD, President, Osaka Medical Center and Research Institute for Maternal and Child Health, 840 Murodo-cho, Izumi City, Osaka, 594-1101, JAPAN. Fax: (+81) 725 56 5682; Email: aokada@mch.pref.osaka.jp.

Professor Duo Li MD, Professor of Nutrition & Food Safety, Dept of Food Science & Nutrition, Zhejiang University, 268 Kaixuan Road, Hangzhou 310029, CHINA. Fax: (+86) 571 86971024; Email: duoli@zju.edu.cn

Managing Editor: Dr Antigone Kouris-Blazos, HEC PRESS, PO Box 4121, McKinnon, Melbourne, Victoria 3204, AUSTRALIA; Fax: (+61) 3 95154544. Email: info@healthyeatingclub.org

Editorial Office: Asia Pacific Journal of Clinical Nutrition, Asia Pacific Health and Nutrition Centre, Monash Asia Institute, 8th Floor, Menzies Building, Monash University, Wellington Road, Clayton, Melbourne, Victoria 3800, AUSTRALIA
Fax: (+61) 3 9905 8146. Email: apjcn@ozemail.com.au. Administration officer: Wendy Yu

Asia Pacific Journal of Clinical Nutrition (APJCN) is a peer-reviewed journal published for the Asia Pacific Clinical Nutrition Society (APCNS). The journal publishes original research reports, short communications, reviews, and letters to the editors. Letters to the editors may be either very short articles (500 words) containing new material, case reports, or comments on previous papers or on other topics of current interest. News, book reviews and other items may also be included. All articles are peer-reviewed by at least two researchers expert in the field of the submitted paper. For detailed instructions concerning the submission of manuscripts, please refer to the Instructions to Authors at the back of this issue or visit the journal web page, listed below.

The aims of the *Asia Pacific Clinical Nutrition Society (APCNS)* are to promote the education and training of clinical nutritionists in the region and to enhance the practice of human nutrition and related disciplines in their application to health and the prevention of disease.

The journal is also an organ of *The Nutrition Society of New Zealand* and of *The Nutrition Society of Australia*, whose members receive *APJCN* as part of their society membership. *The International Union of Nutritional Sciences (IUNS)* is acknowledged as a non-financial co-sponsor of the journal.

Abstracting and Indexing Services: The Journal is indexed by Australasian Medical Index, CAB Abstracts, Chemical Abstracts Service, Current Contents/Clinical Medicine, Current Opinion, Index Medicus/MEDLINE, ISI Alerting Services, Science Citation Index, SciSearch and SUBIS Current Awareness in Biomedicine. The Journal is also indexed selectively in APAIS.

Business Correspondence and Advertisement Booking: All matters relating to orders, subscriptions, back issues, offprints, advertisement booking and general enquiries should be addressed to: HEC PRESS, PO Box 4121, McKinnon, Melbourne, Victoria 3204, Australia; Mobile: +61 0408551702; Fax: +61(0)395154544; info@healthyeatingclub.org; <http://www.healthyeatingclub.org>

Subscription Rates: The journal is published four times per year and the prices per annum for 2005 are as follows:

Personal: Hard copy (including on-line access) AUS\$135;
On-line access only AUS\$120.

Institution: Hard copy (including on-line access) AUS\$400;
On-line access only AUS\$170.


The journal is despatched to subscribers outside Australia by surface air-lifted delivery.

Copyright and Photocopying: © 2005 HEC PRESS, Healthy Eating Club Pty Ltd. Authorization to photocopy items for internal or personal use, or the internal or personal use of specific clients, is granted by HEC PRESS, Healthy Eating Club Pty Ltd for libraries and other users registered with the Copyright Clearance Center (CCC) Transactional Reporting Service, provided that the base fee of US\$15 per copy is paid directly to: CCC, 222 Rosewood Drive, Danvers, MA 01923, USA. This consent does not extend to other kinds of copying, such as copying for general distribution, for advertising or promotional purposes, for creating new collective works or for resale. Special requests should be addressed to the Publisher.

APJCN online & CD-ROM

Issues prior to 2004: Full papers are available free-of-charge via the website <http://www.healthyeatingclub.org>. Volumes 1-13 can also be purchased on CD-ROM for AU\$190 (to order CD-ROM use APJCN subscription form on last page of this issue). **Issues after 2003:** subscribers have on-line access to full papers on the website www.healthyeatingclub.org (password access).

Disclaimer: The Publisher, the Society and the Editors cannot be held responsible for errors or any consequences arising from the use of information contained in this journal; the views and opinions expressed do not necessarily reflect those of the Publisher, the Society or the Editors, neither does the publication of advertisements constitute any endorsement by the Publisher, the Society or the Editors of the products advertised.



Asia Pacific Clinical Nutrition Society

www.apcns.org

President: Professor Prasong Tienboon, Thailand

Immediate Past President: Professor Widjaja Lukito, Indonesia

Vice President: Professor Zhu-Ming Jiang, China

Secretary: Dr Irene Darmadi, Australia

Treasurer: Professor Mark Wahlqvist, Australia

Councillor: Professor Kazuo Kondo, Japan

Webmaster: *Dr Antigone Kouris-Blazos, Australia*

Asia Pacific Clinical Nutrition Society is established to create links between clinical nutritionists in the Asia Pacific region. The Society seeks thereby to promote the contribution of nutritionists to the health of the populations in Asia Pacific countries. Asia Pacific Clinical Nutrition Society will encourage continuing nutrition and training in the region so as to promote the highest possible level of research and practical innovation. *Asia Pacific Journal of Clinical Nutrition*, along with the sponsorship by Asia Pacific Clinical Nutrition Society of regional and local clinical nutrition meetings, is expected to assist greatly in the achievement of these aims.

Membership of Asia Pacific Clinical Nutrition Society, for which there is a modest annual fee and which includes on-line access to *Asia Pacific Journal of Clinical Nutrition*, is open to all clinical nutritionists in the region. The Society will consider application for membership based on submission of a *curriculum vitae* and a statement of support from one of the officers and the individuals listed below. The Asia Pacific Clinical Nutrition Society representatives should state: 'On behalf of Asia Pacific Clinical Nutrition Society I support the application of [Name and address] for membership in the Society'. After the applicant has obtained this endorsement from the representative it should be forwarded with the applicant's *curriculum vitae* for consideration and acknowledgement to: The Secretary, Asia Pacific Clinical Nutrition Society, Asia Pacific Health and Nutrition Centre, Monash Asia Institute, 8th Floor, Menzies Building, Monash University, Wellington Road, Clayton, Victoria 3168, Australia.

Contents:

Editorial

Population-based studies of nutrition and health in Asia Pacific elderly
MEEI-SHYUAN LEE AND MARK L WAHLQVIST

294

Review Articles

Globalisation, food and health in Pacific Island countries
ROBERT G HUGHES AND MARK A LAWRENCE

298

Antiarrhythmic effects of n-3 polyunsaturated fatty acids
WANWARANG WONGCHAROEN AND NIPON CHATTIPAKORN

307

Dietary fat and the prevention of chronic disease
MARK L WAHLQVIST

313

Original Articles

Antioxidant properties of palm fruit extracts
NAGENDRAN BALASUNDRAM, TAN YEW AI, RAVIGADEVI SAMBANTHAMURTHI, KALYANA SUNDRAM
AND SAMIR SAMMAN

319

Effect of palm oil on blood pressure, endothelial function and oxidative stress
MOHAMED A BAYORH, IMAD KABUKHALAF AND AGABA A GANAFA

325

Dietary red palm oil supplementation protects against the consequences of global ischemia
in the isolated perfused rat heart

340

AJ ESTERHUYSE, EF DU TOIT AND J VAN ROOYEN

Palm oil *versus* hydrogenated soybean oil: Effects on serum lipids and plasma haemostatic variables
JAN I PEDERSEN, HANNE MÜLLER, INGEBJØRG SELJEFLØT AND BENTE KIRKHUS

348

A comparison between tocopherol and tocotrienol effects on gastric parameters in rats exposed to stress
MF NUR AZLINA, MI NAFEEZA AND BAK KHALID

358

Intracellular mechanisms mediating tocotrienol-induced apoptosis in neoplastic mammary
epithelial cells
PAUL W SYLVESTER AND SUMIT SHAH

366

Effects of tocotrienols on cell viability and apoptosis in normal murine liver cells (BNL CL.2) and
liver cancer cells (BNL IME A.7R.1), *in vitro*
CHAN HOOI HAR AND CHAN KOK KEONG

374

A manufacturer's perspective on selected palm-based products
NEILO CARR AND W FRASER HOGG

381

Quality of margarine: fats selection and processing parameters
MAT SAHRI MISKANDAR, YAAKOB CHE MAN, MOHD SURIA AFFANDI YUSOFF AND RUSSLY ABD RAHMAN

387

Interesterified palm products as alternatives to hydrogenation
NOR AINI IDRIS AND NOOR LIDA HABI MAT DIAN

396

Performance of phytochemical antioxidant systems in refined-bleached-deodorized palm olein during frying
IRWANDI JASWIR, YAAKOB B CHE MAN AND TORLA H HASSAN

402

Palm oil and palm olein frying applications
RAZALI ISMAIL

414

Preventive action of food seasoning spices mixture on fructose-induced lipid abnormalities
RAMU SUGANTHI, SHANMUGANATHAN RAJAMANI, MAMBAKKAM KATCHAPESWARAN RAVICHANDRAN AND
CARANI VENKATRAMAN ANURADHA

420

Mother's milk and hydrogen peroxide
EBTEHAL ALI ABAS AL-KERWI, ALI HMAL-HASHIMI AND ATHRAA M SALMAN

428

Anti-schistosomal activity of colostral and mature camel milk on *Schistosoma mansoni* infected mice
AMANY S MAGHRABY, MAHMOUD A MOHAMED AND AHMED M ABDEL-SALAM

432

News & Views

Future Events

439

Author Index

440



Original Article

Interesterified palm products as alternatives to hydrogenation

Nor Aini Idris¹ and Noor Lida Habi Mat Dian MSc¹

Malaysian Palm Oil Board, No. 6, Persiaran Institusi, Bandar Baru Bangi, 43000 Kajang, Selangor, Malaysia

Interesterification is one of the processes used to modify the physico-chemical characteristics of oils and fats. Interesterification is an acyl-rearrangement reaction on the glycerol molecule. On the other hand, hydrogenation involves addition of hydrogen to the double bonds of unsaturated fatty acids. Due to health implications of *trans* fatty acids, which are formed during hydrogenation, the industry needs to find alternatives to hydrogenated fats. This paper discusses some applications of interesterified fats, with particular reference to interesterified palm products, as alternatives to hydrogenation. Some physico-chemical properties of interesterified fats used in shortenings are discussed. With interesterification, more palm stearin can be incorporated in vanaspati. For confectionary fats and infant formulations, enzymatic interesterification has been employed.

Key Words: Interesterification, palm products, palm oil, hydrogenation, margarine

Introduction

Interesterification is one of the four modification processes to alter the physico-chemical characteristics of oils and fats, the others being blending, fractionation and hydrogenation. Interesterification is an acyl-arrangement reaction. Natural oils and fats have a specific distribution of fatty acids. In most oils and fats, the middle, or 2-position, of the triacylglycerol molecules are preferentially occupied by an unsaturated fatty acid, such as linoleic or linolenic acid. During interesterification, the distribution of fatty acids is randomized on the glycerol backbone. This leads to an altered triacylglycerol composition and increased triacylglycerol species. Consequently, this will affect the physical characteristics of the oil or fat, including melting and crystallization.

In hydrogenation, hydrogen is added to the double bonds of unsaturated fatty acids. But in partial hydrogenation, some of the saturated acids are isomerized into *trans* fatty acids from their natural *cis* configuration. In the *cis* configuration, the two carbon moieties are on the same side of the double bond and on opposite sides in the *trans* configuration (Fig. 1). The *cis* configuration is a bent molecule, whereas the *trans* configuration is a straight chain.

Trans fatty acids have similar melting points to those of the corresponding saturated fatty acids and are very important contributors to the functional properties of hydrogenated products. Hydrogenation converts the liquid oil into a semi-solid or solid fat. The margarine firmness is increased by hydrogenation of the base stock due to the formation of saturated and *trans* fatty acids¹, however, *trans* fatty acids are now considered a risk factor for cardiovascular diseases.² Due to the health implications, the oils and fats industry is moving towards zero or low *trans* products. For example, in Europe, the levels of *trans* fatty

acids in margarine products have been lowered.³ The industry, therefore, needs to find alternatives to hydrogenated fats. Being naturally semi-solid, palm oil is an obvious choice to replace hydrogenated fats in solid fat food formulations. The industry is now relying more on interesterification to produce low or zero *trans* fats.⁴ Liquid




Figure 1. Molecular structure of *cis* and *trans* isomers of C18:1⁵ (Reproduced with permission from Elsevier)

Correspondence address: Nor Aini Idris, Malaysian Palm Oil Board, No. 6, Persiaran Institusi, Bandar Baru Bangi, 43000 Kajang, Selangor, Malaysia

Tel: 603-89282434; Fax: 603-89259446

E-mail: aini@mpob.gov.my

Accepted 30th June 2005

palm products such as palm olein can be interesterified or use in solid fat formulations. Interesterified palm stearin or its inter-esterified blends with other oils or fats can be used in several food applications.

Interesterification

Interesterification was first used in the US in 1940's to improve the poor creaming properties of lard. Unmodified lard is a beta-tending fat, becoming grainy due to the uniform nature of its triglycerides. Most of the palmitic acid in lard is located in the 2-position of the glyceride molecule. In palm oil, the palmitic acid is located in the 1, 3 positions. Interesterification eliminates the graininess in lard as the position of palmitic acid is randomized, resulting in the formation of more desirable beta prime crystals.

Interesterification can be carried out chemically or enzymatically. A chemical, such as sodium methoxide, is used as a catalyst in chemical interesterification which produces complete positional randomization of the acyl groups in the triacylglycerols. On the other hand, enzymatic interesterification uses microbial lipases as the catalyst. Each type of interesterification has its advantages and disadvantages. The advantages of chemical interesterification over the enzymatic reaction include cost recovery and low initial investment as the catalysts

are much cheaper than lipases. The process has been around for a long time, with the industrial procedures and equipment available. Enzymatic interesterification is more specific, requires less severe reaction conditions and produces less waste than chemical interesterification. In view of the current concern over *trans* fatty acids, MPOB have been investigating several *trans* fatty acid-free products - shortenings, margarines, vanaspati, ice-cream and confectionaries. For this paper, only selected products will be discussed with work done elsewhere cited.

Table 1 shows the fatty acid compositions of commercial shortenings from five countries compared with two interesterified palm-based shortenings. The levels of *trans* acids is expressed as elaidic acid (C18:1t). The shortening from the USA contained the highest *trans* acids at 15.1%, followed by that from Turkey with 11.5% and (closely behind) that from Canada at 11.0%. All these shortenings are based on hydrogenated oils. The South African shortening contained negligible *trans* acids with a fatty acid profile similar to that of the shortening from Malaysia. It is, therefore, likely that it contained high amounts of palm oil products. Malaysian shortenings are usually based on palm oil and palm stearin. Like commercial Malaysian shortenings, the shortening based on interesterified palm olein contained high palmitic and oleic acids [C16:0 and C18:1 (n-9), respectively].

Table 1. Fatty acid composition of shortenings from different countries and shortenings based on interesterified (IE) palm olein (POo) and an interesterified blend of palm stearin (POs) with cottonseed oil (CSO)^a

Fatty acid (%)	USA	Canada	Turkey	South Africa	Malaysia	IE POo	IE POs:CSO
12:0	0	0	0.8	0.7	0.2	0.2	0
14:0	0.1	0.3	0.9	1.4	1.1	1.1	0
16:0	12.0	16.1	24.8	52.7	45.4	39.9	38.2
16:1	0.1	0.1	0.2	0.1	0.2	0.1	0.1
18:0	11.0	9.9	6.2	4.9	4.6	4.1	4.7
18:1t	15.1	11.0	11.5	0.8	0.1	0	0
18:1(n-9)	40.0	32.7	27.7	29.5	37.0	43.2	27.1
18:2tt,ct	4.0	2.7	0.5	0.2	0.3	0.1	0.2
18:2(n-6)	21.0	22.4	24.7	7.3	10.2	11.0	26.0
18:3(n-3)	1.3	1.5	1.5	0.1	0.3	0	0.1
20:0	0.4	0.5	0.2	0.4	0.4	0	0.5
20:1	0	0.3	0	0.1	0	0	0
22:0	0	0.3	0	0	0	0	0
24:0	0	0.1	0	0	0	0	0
Others	1.7	0.1	0.9	1.7	0	0	0

^aThe fatty acids were analysed as methyl esters using a Hewlett-Packard GC System HP 6890 Model G1530A (Wilmington, DE); Column: Fused silica capillary column, 60 m x 0.25 mm, film thickness 0.25 μ m; Oven temperature: 185°C; Injector temperature: 240°C; Detector temperature: 240°C; Carrier gas: He, 0.8 ml/min. Quantification of the peak was carried out using an integrator built into the system.

It also contained a moderate amount of linoleic acid (C18:2). The shortening based on inter-esterified palm stearin and cottonseed oil contained high palmitic acid and considerable amounts of oleic and linoleic acids.

Applications of interesterification in product formulation

Interesterification is an effective way to increase the slip melting point (SMP) and solid fat content (SFC) of palm olein:tallow blends.⁶ After interesterification, there were changes in the triglyceride compositions of the blends (Table 2) - lower C50 and C52 and higher C48 and C54. The creaming performance improved in the interesterified palm olein : tallow blends, as did the baking performance (Table 3).

Beta prime crystals are desirable in margarines and in shortenings for icing and cakes. It is important that the fat crystals are able to remain in beta prime form during storage to maintain the textural quality and functional properties. Studies at MPOB have shown products made from hydrogenated palm oil to be stable in the beta prime form, even after two months from production. Some beta crystals were found in products made from unmodified palm oil after the same storage time, but beta prime crystals still predominated. Beta crystals were predominant in direct blends of palm stearin and sunflower oil. With interesterification, more beta prime crystals were formed in this blend.

Table 2. Slip melting point and triglyceride composition of direct and interesterified (IE) blends of palm olein (POo) and tallow (T)

Sample	POo:T 70:30	IEPOo:T 70:30	POo:T 60:40	IEPOo:T 60:40	Commercial
Slip melting point (°C)					
	33.8	39.7	33.9	35.5	43.8
Triglyceride composition (wt %) by carbon number					
46	0.9	1.9	1.0	1.0	1.9
48	4.6	9.1	5.0	5.6	9.1
50	35.4	27.6	33.6	32.9	20.6
52	41.9	36.0	45.7	44.5	36.4
54	11.9	15.5	14.0	15.0	19.9
56	0.4	0.3	0.5	0.5	0

Table 3. Shortening baking test for blends of palm olein (POo) and tallow (T)

Sample	Specific cake volume (cm ³ /g)
POo:T 70:30	2.33
IE POo:T 70:30	2.38
POo:T 60:40	2.46
IE POo:T 60:40	2.49
Commercial	2.30

Margarines and shortenings

The slow crystallizing property of palm oil may lead to a post-hardening in products such as margarine and shortening. Interesterification can ameliorate the problem. Table 4 shows the consistencies of several palm-based shortenings during storage. The shortening made from hydrogenated palm oil was the firmest. For cake making, a softer shortening is preferred (yield value of 200 to 500g/cm²). The softer shortening acts as a lubricant and aids in the mixing of ingredients. The interesterified palm oil shortening was softer than that made from unmodified palm oil. Similarly, the product based on interesterified palm stearin:sunflower oil was generally softer than the non-interesterified one. The changes in consistency during storage were also minimal in the interesterified products.

Table 4. Consistency measurements^a of several palm based shortenings

Shortening	PO	HPO	IEPO	POs:SFO	IE POs:SFO
Day					
15	250	1120	190	224	240
30	260	1290	200	441	274
60	300	1370	220	444	365

^aIn terms of yield value (g/cm²). PO, palm oil; HPO, hydrogenated palm oil; IEPO, interesterified palm oil; POs, palm stearin; SFO, sunflower oil.

Vanaspatti

Vanaspatti, or vegetable ghee, is a very popular cooking fat in India and Pakistan. In India, vanaspatti is made from hydrogenated vegetable oils. The oils commonly used are palm, soybean, rapeseed and cottonseed. Malaysian manufacturers also produce vanaspatti, mainly for export to Middle Eastern countries. The Malaysian products are usually from direct blends of palm oil and palm stearin. *Trans* free vanaspatti can be produced using palm stearin and other liquid oils either by direct blending or by employing interesterification. Our studies at MPOB have shown that more palm stearin can be incorporated in vanaspatti with interesterification.⁷ Figure 2 shows the SFC profiles of some of the palm-based vanaspatti compared with a commercial hydrogenated vanaspatti from India. The vanaspattis based on interesterified palm products had good characteristics, comparable to those in the commercial product. This is proof that interesterified palm products are good alternatives to hydrogenated other oils.

Confectionary products

In confectionary, cocoa butter is an important raw material. Although cocoa butter confers excellent eating qualities, products containing it are subject to bloom, difficult to temper and may melt in hot climates. Furthermore, the inconsistent supply and high price of cocoa butter have encouraged the search for alternatives. These alternatives are fats generally classified into three types, based on their chemical composition and compatibility with cocoa




Figure 2. Solid fat content profiles of vanaspati formulations compared to commercial ghee and vanaspati. A = commercial ghee; B = IE PO; C = POs:SBO; D = IE POs:SBO; E = HPO and F = commercial vanaspati

butter. Cocoa butter equivalents (CBE) have chemical and physical properties compatible with those of cocoa butter, and can be used to supplement cocoa butter in confectionery products. Cocoa butter substitutes (CBS) are generally lauric fats incompatible with cocoa butter. Cocoa butter replacers (CBR) are partially compatible with cocoa butter. CBRs are primarily non-lauric fats which have properties in between those of CBEs and CBSs, and are sometimes referred to as non-lauric cocoa butter substitutes. Among these three principal types of alternative fats, CBE is the most and CBS the least expensive. Typically, the substitutes cost only one-third to one-fourth of cocoa butter, making them more economically attractive to consumers.⁸

There are a number of fats suitable for total or partial replacement of cocoa butter. Palm oil is an important source oil in the development of substitutes or equivalents. Undurraga *et al.*⁹ used enzymatic interesterification of palm mid fraction with stearic acid in a solvent free system with a commercially available enzyme to produce CBE. In an earlier study, Chong *et al.*¹⁰ used a solvent-free lipase catalyzed inter-esterification of palm olein with stearic acid, followed by solvent fractionation procedures to produce palm-based cocoa butter-like fat. The yield of cocoa butter-like fat was approximately 25%

Table 5. Composition of a fraction of interesterified palm olein obtained by different solvent fractionation procedures (A, B, C).

Fraction	Triglycerides ^a (%)		
	POP	POS	SOS
F2(A)	15.3	44.1	29.6
F2(B)	16.0	38.7	23.1
F2(C)	13.7	38.5	24.8
Cocoa butter	18.9	41.3	29.7

^aP, palmitate; S, stearate; O, oleate. (Reproduced with permission from American Oil Chemists' Society).

of the weight of the original palm olein. Table 5 shows the triacylglycerol compositions of a fraction of interesterified palm olein obtained by different solvent fractionation procedures.

Infant formula

Another important application of interesterification is infant formula. The fatty acid composition of human milk is often used as the basis for infant formula to create a close semblance to the milk.¹¹ Human milk fat has triacylglycerols which fatty acid structure, composition and distribution are specific. Its unique characteristics include, firstly, two long-chain polyunsaturated fatty acids (PUFAs) - arachidonic (C20:4, n-6) and docosahexaenoic (C22:6, n-3) - and, secondly, abundance of a saturated fatty acid - palmitic (C16:0) - predominantly at the 2-position of the triacylglycerol.¹²

To obtain a similar fatty acid composition and positional distribution as in human milk fat, oils and fats may be modified by interesterification. Lipid compositions for infant formula, which mimics human milk, as well as the processes for preparation of the compositions by interesterification, either chemical or enzymatic, are described in various international patents. In one, a mixture of lipids containing palm oil enriched with palmitic acid (for example, from palm stearin), a vegetable oil high in linoleic and alpha-linolenic fatty acids, an oil for arachidonic acid and another for docosahexaenoic acid in defined proportions are interesterified to obtain the required fatty acid composition with random distribution of the fatty acid residues between the 1-, 2- and 3- positions of the triacylglycerols. The interesterification can be enzymatic (catalyzed by a non-regiospecific lipase) or chemical. By the reaction, the non-random distribution of fatty acids on the triacylglycerols of the natural lipids is converted to random distribution, with the fatty acids rearranged equally at the three positions. The randomized mixture was then interesterified with a mixture of free fatty acids high in medium chain fatty acids and oleic acid using a 1,3-regiospecific lipase. This step selectively replaces palmitic acid at the 1- and 3- positions with other fatty acids, in particular, with medium chain fatty acids and oleic but not with PUFAs. Many scientists have reported that enzymatic interesterification using 1,3-specific lipase is the most effective way to produce triacylglycerols similar to those found in human milk.¹³⁻¹⁷

References

1. Mensink RP, Katan MB. Trans monounsaturated fatty acids in nutrition and their impact on serum lipoprotein levels in man. *Progress in Lipid Research* 1993; 32: 111-122.
2. Reddy SY, Jeyarani T. Trans-free bakery shortenings from mango kernel and mahua fats by fractionation and blending. *J Am Oil Chemists Soc* 2001; 78: 635-640.
3. Michels K, Sacks F. Trans fatty acids in European margarines. *N Engl J Med* 1995; 332: 541-542.
4. Yang T, Xu, X. Enzymatic modification of palm oils: Useful products with potential processes. *Proceedings of the 2001 PIPOC International Palm Oil Congress*, Malaysian Palm Oil Board, K. Lumpur. 2001
5. Ghotra BS, Dyal SD, Narine SS. Lipid shortenings: A review. *Food Research International* 2002; 35:1015-1048.

6. Osman A, Nor Aini I. Physical and chemical properties of shortenings from palm oil:tallow and palm olein:tallow blends with and without interesterification. *J Palm Oil Research* 1999; 11:1-10.
7. Nor Aini I, Yusoff MSA, Mohd Arif S. Palm-based trans free vanaspati. PORIM Information Series No. 66; 1997.
8. Nalur SC. Palm kernel oil blends. United States Patent 1999; No. 5, 932, 275.
9. Undurragaa D, Markovits A, Erazob S. Cocoa butter equivalent through enzymatic interesterification of palm oil mid fraction. *Process Biochemistry* 2001; 36: 933-939.
10. Chong CN, Hoh YM, Wang CW. Fractionation procedures for obtaining cocoa butter-like fat from enzymatically interesterified palm olein. *JAOCS* 1992; 69:137-140.
11. Jensen RG. Human milk lipids as a model for infant formula. *Lipid Technol* 1998; 3: 34 -38.
12. Wang J, Bertholet R, Ducret P, Fleith M. Lipid composition for infant formula and method of preparation. United States Patent 2000; No. 6,034,130.
13. Akoh CC, Lee LN. Enzymatic synthesis of position-specific low-calorie structured lipid. *JAOCS*, 1997; 74: 1409 -1413.
14. Akoh CC, Moussata CO. Lipase-catalyzed modification of borage oil: incorporation of capric and eicosapentaenoic acids to form structured lipids. *JAOCS* 1998; 75: 697-701.
15. Xu X, Balchen S, Hoy CE, Adler-Nissen J. Pilot batch production of specific-structured lipids by lipase-catalyzed interesterification: preliminary study on incorporation and acyl migration. *JAOCS* 1998; 75:301-308.
16. Xu X, Mu H, Hoy CE, Adler-Nissen J. Production of specifically structured lipids by enzymatic interesterification in a pilot enzyme bed reactor: process optimization by response surface methodology. *Fett/Lipid*, 1999; 101: 207-214.
17. Shimada Y, Sakai N, Sugihara A, Fujita H, Honda Y, Tominaga Y. Large-scale purification of linoleic acid by selective esterification using *Rhizopus delemar* lipase. *JAOCS* 1998; 75:1539-1543.