

and amplification agents sufficient to process at least about 384 samples. Any one of the samples may comprise a single cell. The nucleic acid amplification may result in a measurement of about 1, 2, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 200, 300, 400, 500, 600, 700, 800, 900 or 1000 targeted nucleic acids in a sample. The nucleic acid amplification may result in a measurement of about 1000 targeted nucleic acids in a sample. The nucleic acid amplification may result in a measurement of about 100 targeted nucleic acids in a sample. The nucleic acid amplification may result in a measurement of about 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 100% of total nucleic acids in single cells. The nucleic acid amplification may result in a global measurement of all nucleic acid sequences in single cells. The nucleic acid amplification may result in a measurement of targeted nucleic acid sequences in single cells by sequencing. The nucleic acid amplification may result in a measurement of targeted nucleic acid sequences in single cells by an array.

[00402] The amplification agents may comprise a fixed panel of primers. The amplification agents may comprise at least one pair of custom primers. The amplification agents may comprise at least one pair of control primers. The amplification agents may comprise at least one pair of housekeeping gene primers. The amplification agents suitable may comprise a PCR master mix. The kit may further comprise instructions for primer design and optimization. The kit may further comprise a microwell plate, wherein the microwell plate may comprise at least one well in which no more than one bead is distributed. The kit may further comprise one or more additional containers. The one or more additional containers may comprise one or more additional plurality of sample tags. The plurality of one or more additional sample tags in the one or more additional containers are different from the first plurality of sample tags in the first container. The one or more additional containers may comprise one or more additional molecular identifier labels. The one or more additional molecular identifier labels of the one or more additional containers are the same as the one or more additional molecular identifier labels of the second container.

[00403] The methods and kits disclosed herein may comprise the use of one or more pipette tips and/or containers (*e.g.*, tubes, vials, multiwell plates, microwell plates, eppendorf tubes, glass slides, beads). In some instances, the pipet tips are low binding pipet tips. Alternatively, or additionally, the containers may be low binding containers. Low binding pipet tips and low binding containers may have reduced leaching and/or subsequent sample degradation associated with silicone-based tips and non-low binding containers. Low binding pipet tips and low binding containers may have reduced sample binding as compared to non-low

binding pipet tips and containers. Examples of low binding tips include, but are not limited to, Corning® DeckWorks™ low binding tips and Avant Premium low binding graduated tips. A non-limiting list of low-binding containers include Corning® Costar® low binding microcentrifuge tubes and Cosmobrand low binding PCR tubes and microcentrifuge tubes.

[00404] Any of the kits disclosed herein can further comprise software. For example, a kit can comprise software for analyzing sequences, such as barcodes or target sequences. For example, a kit can comprise software for analyzing sequences, such as barcodes or target sequences for counting unique target molecules, such as unique target molecules from a single cell. For example, a kit can comprise software for analyzing sequences, such as barcodes or target sequences for counting unique target molecules, such as unique target molecules from a gene, such as a gene from a single cell.

Microwells and microwell arrays

[00405] In some instances, the methods of the disclosure provide for contacting a solid support comprising a conjugated oligonucleotide with a cell. The contacting step may be performed on a surface. Exemplary surfaces may include a microwell, a tube, a flask, and chip. In some instances, the surface comprises a microwell. In some instances, the microwell is part of a microwell array.

[00406] The microwells of a microwell array may be of a size and shape capable of containing at least 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 or more cells per microwell. The microwells may be of a size and shape capable of containing at most 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 or more cells per microwell. The microwells of a microwell array may be of a size and shape capable of containing at least 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 or more solid supports per microwell. The microwells may be of a size and shape capable of containing at most 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 or more solid supports per microwell. A microwell may comprise at most one cell and one solid support. A microwell may comprise at most one cell and two solid supports. A microwell may comprise at least one cell and at most one solid support. A microwell may comprise at least one cell and at most two solid supports.

[00407] Microwells on the microwell array may be arranged horizontally. The microwells may be arranged vertically. The microwells may be arranged with equal or near equal spacing. The microwell array may have markers associated with one or more microwells. For example, the microwells of the microwell array may be divided into groups each comprised of a prescribed number of microwells. These groups may be provided on the principal surface of the substrate. Markers may be provided so that the position of each group may be

determined. A marker may be detectable by the naked eye. A marker may be a marker that requires optics to see (*e.g.*, fluorescent marker, emission marker, UV marker).

[00408] A microwell array may comprise at least about 96, 384, 1000, 5000, 10000, 15000, 100000, 150000, 500000, 1000000, or 5000000 or more microwells. A microwell array may comprise at most about 96, 384, 1000, 5000, 10000, 15000, 100000, 150000, 500000, 1000000, or 5000000 or more microwells.

[00409] The shape of the microwell may be cylindrical. The shape of the microwell may be noncylindrical, such as a polyhedron comprised of multiple faces (for example, a parallelepiped, hexagonal column, or octagonal column), an inverted cone, an inverted pyramid (inverted triangular pyramid, inverted square pyramid, inverted pentagonal pyramid, inverted hexagonal pyramid, or an inverted polygonal pyramid with seven or more angles). The microwell may comprise a shape combining two or more of these shapes. For example, it may be partly cylindrical, with the remainder having the shape of an inverted cone. The shape of the microwell may be one in which a portion of the top of an inverted cone or inverted pyramid is cut off. The mouth of the microwell may be on the top of the microwell or the bottom of the microwell. The bottom of the microwell may be flat, but curved surfaces (*e.g.*, convex or concave) are also possible. The shape and size of the microwell may be determined in consideration of the type of cell and/or solid substrate (*e.g.*, shape, size) to be stored in the microwell.

[00410] The diameter of the microwell may refer to the largest circle that may be inscribed in the planar shape of the microwell. The diameter of the microwell may be at least about 0.1, 0.5, 1, 2, or 3-fold or more the diameter of the cell and/or solid support to be contained in the microwell. The diameter of the microwell may be at most about 0.1, 0.5, 1, 2, or 3-fold or more the diameter of the cell and/or solid support to be contained in the microwell. The diameter of the microwell may be at least about 10, 20, 30, 40, or 50% or more the diameter of the solid support. The diameter of the microwell may be at most about 10, 20, 30, 40, or 50% or more the diameter of the solid support. The diameter of the microwell may be at least about 5, 10, 15, 20, 25, 30, 35, 40, 45, or 50 or more micrometers. The diameter of the microwell may be at most about 5, 10, 15, 20, 25, 30, 35, 40, 45, or 50 or more micrometers. The diameter of the microwell is about 25 micrometers. In some instances, the diameter of the microwell is about 30 micrometers. In some instances, the diameter of the microwell is about 28 micrometers.

[00411] The difference between the microwell volume and the solid support volume may be at least about $1 \times 10^{(-14)} \text{ m}^3$, $1.5 \times 10^{(-14)} \text{ m}^3$, $1.7 \times 10^{(-14)} \text{ m}^3$, $2.0 \times 10^{(-14)} \text{ m}^3$, $2.5 \times 10^{(-14)} \text{ m}^3$, or

$3.0 \times 10^{(-14)} \text{ m}^3$ or more. The difference between the microwell volume and the solid support volume may be at most about $1 \times 10^{(-14)} \text{ m}^3$, $1.5 \times 10^{(-14)} \text{ m}^3$, $1.7 \times 10^{(-14)} \text{ m}^3$, $2.0 \times 10^{(-14)} \text{ m}^3$, $2.5 \times 10^{(-14)} \text{ m}^3$, or $3.0 \times 10^{(-14)} \text{ m}^3$ or more. The difference between the microwell volume and the solid support volume may be at least about $1 \times 10^{(-11)} \text{ L}$, $1.5 \times 10^{(-11)} \text{ L}$, $1.7 \times 10^{(-11)} \text{ L}$, $2.0 \times 10^{(-11)} \text{ L}$, $2.5 \times 10^{(-11)} \text{ L}$, or $3.0 \times 10^{(-11)} \text{ L}$ or more. The difference between the microwell volume and the solid support volume may be at most about $1 \times 10^{(-11)} \text{ L}$, $1.5 \times 10^{(-11)} \text{ L}$, $1.7 \times 10^{(-11)} \text{ L}$, $2.0 \times 10^{(-11)} \text{ L}$, $2.5 \times 10^{(-11)} \text{ L}$, or $3.0 \times 10^{(-11)} \text{ L}$ or more. **FIG. 7** illustrates exemplary statistics about the volume of the microwell, the solid support, and the differences between the microwell and the solid support volumes.

[00412] The depth of the microwell may be at least about 0.1, 0.5, 1, 2, 3, 4, or 5-fold or more the diameter of the cell and/or solid support to be contained in the microwell. The depth of the microwell may be at most about 0.1, 0.5, 1, 2, 3, 4, or 5-fold or more the diameter of the cell and/or solid support to be contained in the microwell. The depth of the microwell may be at least about 10, 20, 30, 40, or 50% or more the depth of the solid support. The depth of the microwell may be at most about 10, 20, 30, 40, or 50% or more the depth of the solid support. The depth of the microwell may be at least about 5, 10, 15, 20, 25, 30, 35, 40, 45, or 50 or more micrometers. The depth of the microwell may be at most about 5, 10, 15, 20, 25, 30, 35, 40, 45, or 50 or more micrometers. The depth of the microwell may be about 30 micrometers. The depth of the microwell may be about 28 micrometers. The microwell may be flat, or substantially flat.

[00413] A microwell array may comprise spacing between the wells. The spacing between the wells may be at least about 5, 10, 25, 20, 25, 30, 35, 40, 45, or 50 or more micrometers. The spacing between the wells may be at most about 5, 10, 25, 20, 25, 30, 35, 40, 45, or 50 or more micrometers. The spacing between the wells may be about 15 micrometers. The spacing between the wells may be about 25 micrometers.

[00414] There may be differences in the height of dips and rises at any position on the inner wall of a microwell. By creating dips and rises on a portion of the inner wall of a well that has been treated for smoothness, functionality may be added to the well. The inner wall of a microwell may be smoothed by etching. The degree of vacuum in the etching device, the type of etching gas, the etching steps, and the like may be suitably selected. For example, smoothing of the inner wall of a microwell may be conducted by wet etching or by combining a hot oxidation step with oxide film etching. The inner wall of the microwell may be functionalized (*e.g.*, functionalized with an oligonucleotide, a reactive group, a functional group).

[00415] The microwell array may be made of silicon, metal (*e.g.*, aluminum, stainless steel, copper, nickel, chromium, and titanium), PDMS (elastomer), glass, polypropylene, agarose, gelatin, pluronic (*e.g.*, pluronic F127), plastics (*e.g.*, plastics that are naturally hydrophilic, such as PMMA), plastics (*e.g.*, PP, COP, COC) and elastomer (*e.g.*, PDMS) that are hydrophobic but may be treated to be made hydrophilic), hydrogels (*e.g.*, polyacrylamide, alginate), or resin (*e.g.*, polyimide, polyethylene, vinyl chloride, polypropylene, polycarbonate, acrylic, and polyethylene terephthalate). The microwell array may be made of a material that is hydrophobic. The microwell array may be made of a material that is hydrophobic but coated to be made hydrophilic (*e.g.*, by oxygen plasma treatment). The microwell array may be made of a material that is hydrophilic but coated to be made hydrophobic.

[00416] A microwell array may be assembled. Microwell array assembly may comprise obtaining a silicon wafer with patterning (*e.g.*, patterned posts made with SU8 photoresist) and incubating it with PDMS material to create arrays of wells through soft lithography (*e.g.*, at 80 C for a few hours). For example, uncured PDMS may be liquid. Uncured PDMS may fill gaps between posts. When PDMS is cured by heat, it may become solid, thereby generating the array of wells. An optical adhesive (*e.g.*, NOA81/NOA63) may be applied to the PDMS material (*e.g.*, using UV light) to create an array of posts (*e.g.*, a plurality of arrays). The application may be performed for at least about 1 second, 2 seconds, 3 seconds, 4 seconds, 5 seconds, 6 seconds, 7 seconds, 8 seconds, 9 seconds, 10 seconds or 1 minute, 2 minutes, 3 minutes, 4 minutes, or 5 or more minutes. A layer comprising agarose may be applied to the optical adhesive. The agarose layer may be at least about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10% or more agarose. The agarose layer may be most about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10% or more agarose. The agarose layer may be about 5% agarose. The agarose layer may be set on Gelbond film, or any hydrophilic substrate that the agarose may adhere to. The incubation of the agarose layer on the optical surface may be at least about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more minutes. The incubation of the agarose layer on the optical surface may be at most about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more minutes.

[00417] In some instances, the methods of the disclosure may use a surface that may not comprise microwells. The surface may be glass, plastic, metal. The surface may be coated with solid supports, extracellular matrix, polymers. The surface may not comprise wells. The surface may comprise solid supports spatially arranged to limit molecular diffusion. The methods of the disclosure of capturing cells and/or cell contents may occur on a flat surface.

The methods of the disclosure of capturing cells and/or cell contents may occur in a suspension.

Cells and samples

[00418] The cell and of the disclosure may be a cell from an animal (*e.g.*, human, rat, pig, horse, cow, dog, mouse). In some instances, the cell is a human cell. The cell may be a fetal human cell. The fetal human cell may be obtained from a mother pregnant with the fetus. The cell may be a cell from a pregnant mother. The cell may be a cell from a vertebrate, invertebrate, fungi, archae, or bacteria. The cell may be from a multicellular tissue (*e.g.*, an organ (*e.g.*, brain, liver, lung, kidney, prostate, ovary, spleen, lymph node, thyroid, pancreas, heart, skeletal muscle, intestine, larynx, esophagus, and stomach), a blastocyst). The cell may be a cell from a cell culture. The cell may be a HeLa cell, a K562 cell, a Ramos cell, a hybridoma, a stem cell, an undifferentiated cell, a differentiated cell, a circulating cell, a CHO cell, a 3T3 cell, and the like.

[00419] In some instances, the cell is a cancerous cell. Non-limiting examples of cancer cells may include a prostate cancer cell, a breast cancer cell, a colon cancer cell, a lung cancer cell, a brain cancer cell, and an ovarian cancer cell. In some instances, the cell is from a cancer (*e.g.*, a circulating tumor cell). Non-limiting examples of cancers may include, adenoma, adenocarcinoma, squamous cell carcinoma, basal cell carcinoma, small cell carcinoma, large cell undifferentiated carcinoma, chondrosarcoma, and fibrosarcoma.

[00420] In some instances, the cell is a rare cell. A rare cell can be a circulating tumor cell (CTC), circulating epithelial cell (CEC), circulating stem cell (CSC), stem cells, undifferentiated stem cells, cancer stem cells, bone marrow cells, progenitor cells, foam cells, fetal cells, mesenchymal cells, circulating endothelial cells, circulating endometrial cells, trophoblasts, immune system cells (host or graft), connective tissue cells, bacteria, fungi, or pathogens (for example, bacterial or protozoa), microparticles, cellular fragments, proteins and nucleic acids, cellular organelles, other cellular components (for example, mitochondria and nuclei), and viruses.

[00421] In some instances, the cell is from a tumor. In some instances, the tumor is benign or malignant. The tumor cell may comprise a metastatic cell. In some instances, the cell is from a solid tissue that comprises a plurality of different cell types (*e.g.*, different genotypes).

[00422] The cell may comprise a virus, bacterium, fungus, and parasite. Viruses may include, but are not limited to, DNA or RNA animal viruses (*e.g.*, Picornaviridae (*e.g.*, polioviruses), Reoviridae (*e.g.*, rotaviruses), Togaviridae (*e.g.*, encephalitis viruses, yellow fever virus, rubella virus), Orthomyxoviridae (*e.g.*, influenza viruses), Paramyxoviridae (*e.g.*, respiratory

syncytial virus, measles virus, mumps virus, parainfluenza virus), Rhabdoviridae (*e.g.*, rabies virus), Coronaviridae, Bunyaviridae, Flaviviridae, Filoviridae, Arenaviridae, Bunyaviridae and Retroviridae (*e.g.*, human T cell lymphotropic viruses (HTLV), human immunodeficiency viruses (HIV), Papovaviridae (*e.g.*, papilloma viruses), Adenoviridae (*e.g.*, adenovirus), Herpesviridae (*e.g.*, herpes simplex viruses), and Poxviridae (*e.g.*, variola viruses)).

[00423] Exemplary bacteria that may be used in the methods of the disclosure may include Actinomeduræ, Actinomyces israelii, Bacillus anthracis, Bacillus cereus, Clostridium botulinum, Clostridium difficile, Clostridium perfringens, Clostridium tetani, Corynebacterium, Enterococcus faecalis, Listeria monocytogenes, Nocardia, Propionibacterium acnes, Staphylococcus aureus, Staphylococcus epiderm, Streptococcus mutans, Streptococcus pneumoniae and the like. Gram negative bacteria include, but are not limited to, Afipia felis, Bacteriodes, Bartonella bacilliformis, Bortadella pertussis, Borrelia burgdorferi, Borrelia recurrentis, Brucella, Calymmatobacterium granulomatis, Campylobacter, Escherichia coli, Francisella tularensis, Gardnerella vaginalis, Haemophilus aegyptius, Haemophilus ducreyi, Haemophilus influenzae, Helicobacter pylori, Legionella pneumophila, Leptospira interrogans, Neisseria meningitidis, Porphyromonas gingivalis, Providencia stuartii, Pseudomonas aeruginosa, Salmonella enteritidis, Salmonella typhi, Serratia marcescens, Shigella boydii, Streptobacillus moniliformis, Streptococcus pyogenes, Treponema pallidum, Vibrio cholerae, Yersinia enterocolitica, Yersinia pestis and the like. Other bacteria may include Mycobacterium avium, Mycobacterium leprae, Mycobacterium tuberculosis, Bartonella henselae, Chlamydia psittaci, Chlamydia trachomatis, Coxiella burnetii, Mycoplasma pneumoniae, Rickettsia akari, Rickettsia prowazekii, Rickettsia rickettsii, Rickettsia tsutsugamushi, Rickettsia typhi, Ureaplasma urealyticum, Diplococcus pneumoniae, Ehrlichia chaffeensis, Enterococcus faecium, Meningococci and the like.

[00424] Exemplary fungi to be used in the methods of the disclosure may include, but are not limited to Aspergilli, Candida, Candida albicans, Coccidioides immitis, Cryptococci, and combinations thereof.

[00425] Exemplary parasites to be used in the methods of the disclosure may include, but are not limited to, Balantidium coli, Cryptosporidium parvum, Cyclospora cayatanensis, Encephalitozoa, Entamoeba histolytica, Enterocytozoon bienersi, Giardia lamblia, Leishmaniae, Plasmodium, Toxoplasma gondii, Trypanosomae, trapezoidal amoeba, worms (*e.g.*, helminthes), particularly parasitic worms including, but not limited to, Nematoda

(roundworms, *e.g.*, whipworms, hookworms, pinworms, ascarids, filarids and the like), Cestoda (*e.g.*, tapeworms).

[00426] The sample of the disclosure may be a sample from an animal (*e.g.*, human, rat, pig, horse, cow, dog, mouse). In some instances, the sample is a human sample. The sample may be a fetal human sample. The fetal human sample may be obtained from a mother pregnant with the fetus. The sample may be a sample from a pregnant mother. The sample may be a sample from a vertebrate, invertebrate, fungi, archae, or bacteria. The sample may be from a multicellular tissue (*e.g.*, an organ (*e.g.*, brain, liver, lung, kidney, prostate, ovary, spleen, lymph node, thyroid, pancreas, heart, skeletal muscle, intestine, larynx, esophagus, and stomach), a blastocyst). The sample may be a cell from a cell culture.

[00427] The sample may comprise a plurality of cells. The sample may comprise a plurality of the same type of cell. The sample may comprise a plurality of different types of cells. The sample may comprise a plurality of cells at the same point in the cell cycle and/or differentiation pathway. The sample may comprise a plurality of cells at different points in the cell cycle and/or differentiation pathway. A sample may comprise a plurality of samples.

[00428] The plurality of samples may comprise at least 5, 10, 20, 30, 40, 50, 60, 70, 80, 90 or 100 or more samples. The plurality of samples may comprise at least about 100, 200, 300, 400, 500, 600, 700, 800, 900 or 1000 or more samples. The plurality of samples may comprise at least about 1000, 2000, 3000, 4000, 5000, 6000, 7000, 8000 samples, 9000, or 10,000 samples, or 100,000 samples, or 1,000,000 or more samples. The plurality of samples may comprise at least about 10,000 samples.

[00429] The one or more nucleic acids in the first sample may be different from one or more nucleic acids in the second sample. The one or more nucleic acids in the first sample may be different from one or more nucleic acids in a plurality of samples. The one or more nucleic acids may comprise a length of at least about 1 nucleotide, 2 nucleotides, 5 nucleotides, 10 nucleotides, 20 nucleotides, 50 nucleotides, 100 nucleotides, 200 nucleotides, 300 nucleotides, 500 nucleotides, 1000 nucleotides, 2000 nucleotides, 3000 nucleotides, 4000 nucleotides, 5000 nucleotides, 10,000 nucleotides, 100,000 nucleotides, 1,000,000 nucleotides.

[00430] The first sample may comprise one or more cells and the second sample may comprise one or more cells. The one or more cells of the first sample may be of the same cell type as the one or more cells of the second sample. The one or more cells of the first sample may be of a different cell type as one or more different cells of the plurality of samples. The cell type may be chondrocyte, osteoclast, adipocyte, myoblast, stem cell, endothelial cell or

smooth muscle cell. The cell type may be an immune cell type. The immune cell type may be a T cell, B cell, thrombocyte, dendritic cell, neutrophil, macrophage or monocyte.

[00431] The plurality of samples may comprise one or more malignant cell. The one or more malignant cells may be derived from a tumor, sarcoma or leukemia.

[00432] The plurality of samples may comprise at least one bodily fluid. The bodily fluid may comprise blood, urine, lymphatic fluid, saliva. The plurality of samples may comprise at least one blood sample.

[00433] The plurality of samples may comprise at least one cell from one or more biological tissues. The one or more biological tissues may be a bone, heart, thymus, artery, blood vessel, lung, muscle, stomach, intestine, liver, pancreas, spleen, kidney, gall bladder, thyroid gland, adrenal gland, mammary gland, ovary, prostate gland, testicle, skin, adipose, eye or brain.

[00434] The biological tissue may comprise an infected tissue, diseased tissue, malignant tissue, calcified tissue or healthy tissue.

[00435] The plurality of samples may be from one or more sources. The plurality of samples may be from two or more sources. The plurality of samples may be from one or more subjects. The plurality of samples may be from two or more subjects. The plurality of samples may be from the same subject. The one or more subjects may be from the same species. The one or more subjects may be from different species. The one or more subjects may be healthy. The one or more subjects may be affected by a disease, disorder or condition. The plurality of samples may comprise cells of an origin selected from a mammal, bacteria, virus, fungus or plant. The one or more samples may be from a human, horse, cow, chicken, pig, rat, mouse, monkey, rabbit, guinea pig, sheep, goat, dog, cat, bird, fish, frog and fruit fly.

[00436] The plurality of samples may be obtained concurrently. The plurality of samples may be obtained at the same time. The plurality of samples may be obtained sequentially. The plurality of samples may be obtained over a course of years, 100 years, 10 years, 5 years, 4 years, 3 years, 2 years or 1 year of obtaining one or more different samples. One or more samples may be obtained within about one year of obtaining one or more different samples. One or more samples may be obtained within 12 months, 11 months, 10 months, 9 months, 8 months, 7 months, 6 months, 4 months, 3 months, 2 months or 1 month of obtaining one or more different samples. One or more samples may be obtained within 30 days, 28 days, 26 days, 24 days, 21 days, 20 days, 18 days, 17 days, 16 days, 15 days, 14 days, 13 days, 12 days, 11 days, 10 days, 9 days, 8 days, 7 days, 6 days, 5 days, 4 days, 3 days, 2 days or one day of obtaining one or more different samples. One or more samples may be obtained within about 24 hours, 22 hours, 20 hours, 18 hours, 16 hours, 14 hours, 12 hours, 10 hours, 8 hours,

6 hours, 4 hours, 2 hours or 1 hour of obtaining one or more different samples. One or more samples may be obtained within about 60sec, 45sec, 30sec, 20sec, 10sec, 5 sec, 2sec or 1 sec of obtaining one or more different samples. One or more samples may be obtained within less than one second of obtaining one or more different samples.

Target Molecules

[00437] The methods and kits disclosed herein may be used in the stochastic labeling of molecules. Such molecules include, but are not limited to, polynucleotides and polypeptides. As used herein, the terms “polynucleotide” and “nucleic acid molecule” refers to a polymeric form of nucleotides of any length, either ribonucleotides, deoxyribonucleotides, locked nucleic acids (LNA) or peptide nucleic acids (PNAs), that comprise purine and pyrimidine bases, or other natural, chemically or biochemically modified, non-natural, or derivatized nucleotide bases. A “polynucleotide” or “nucleic acid molecule” may consist of a single nucleotide or base pair. Alternatively, the “polynucleotide” or “nucleic acid molecule” comprises two or more nucleotides or base pairs. For example, the “polynucleotide” or “nucleic acid molecule” comprises at least about 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 30, 40, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500, 600, 700, 800, 900, or 1000 nucleotides or base pairs. In another example, the polynucleotide comprises at least about 1500, 2000, 2500, 3000, 3500, 4000, 4500, 5000, 5500, 6000, 6500, 7000, 7500, 8000, 8500, 9000, 9500, or 10000 nucleotides or base pairs. The backbone of the polynucleotide may comprise sugars and phosphate groups, as may typically be found in RNA or DNA, or modified or substituted sugar or phosphate groups. A polynucleotide may comprise modified nucleotides, such as methylated nucleotides and nucleotide analogs. The sequence of nucleotides may be interrupted by non-nucleotide components. Thus the terms nucleoside, nucleotide, deoxynucleoside and deoxynucleotide generally include analogs such as those described herein. These analogs are those molecules having some structural features in common with a naturally occurring nucleoside or nucleotide such that when incorporated into a nucleic acid or oligonucleoside sequence, they allow hybridization with a naturally occurring nucleic acid sequence in solution. Typically, these analogs are derived from naturally occurring nucleosides and nucleotides by replacing and/or modifying the base, the ribose or the phosphodiester moiety. The changes may be tailor made to stabilize or destabilize hybrid formation or enhance the specificity of hybridization with a complementary nucleic acid sequence as desired. In some instances, the molecules are DNA, RNA, or DNA-RNA hybrids. The molecules may be single-stranded or double-stranded. In some instances, the molecules are RNA molecules, such as mRNA, rRNA, tRNA, ncRNA,

lncRNA, siRNA, microRNA or miRNA. The RNA molecules may be polyadenylated. Alternatively, the mRNA molecules are not polyadenylated. Alternatively, the molecules are DNA molecules. The DNA molecules may be genomic DNA. The DNA molecules may comprise exons, introns, untranslated regions, or any combination thereof. In some instances, the molecules are a panel of molecules.

[00438] The methods and kits disclosed herein may be used to stochastically label individual occurrences of identical or nearly identical molecules and/or different molecules. In some instances, the methods and kits disclosed herein may be used to stochastically label identical or nearly identical molecules (*e.g.*, molecules comprise identical or nearly identical sequences). For example, the molecules to be labeled comprise at least about 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity. The nearly identical molecules may differ by less than about 100, 90, 80, 70, 60, 50, 40, 30, 25, 20, 15, 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 nucleotide or base pair. The plurality of nucleic acids in one or more samples of the plurality of samples may comprise two or more identical sequences. At least about 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%, or 100% of the total nucleic acids in one or more of the plurality of samples may comprise the same sequence. The plurality of nucleic acids in one or more samples of the plurality of samples may comprise at least two different sequences. At least about 5%, 10 %, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 100% of the total nucleic acids in one or more of the plurality of samples may comprise at least two different sequences. In some instances, the molecules to be labeled are variants of each other. For example, the molecules to be labeled may contain single nucleotide polymorphisms or other types of mutations. In another example, the molecules to be labeled are splice variants. In some instances, at least one molecule is stochastically labeled. In other instances, at least 2, 3, 4, 5, 6, 7, 8, 9, or 10 identical or nearly identical molecules are stochastically labeled. Alternatively, at least 20, 30, 40, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500, 600, 700, 800, 900, or 1000 identical or nearly identical molecules are stochastically labeled. In other instances, at least 1500; 2,000; 2500; 3,000; 3500; 4,000; 4500; 5,000; 6,000; 7,000; 8,000; 9,000; or 10000 identical or nearly identical molecules are stochastically labeled. In other instances; at least 15,000; 20,000; 25,000; 30,000; 35,000; 40,000; 45,000; 50,000; 60,000; 70,000; 80,000; 90,000; or 100,000 identical or nearly identical molecules are stochastically labeled.

[00439] In other instances, the methods and kits disclosed herein may be used to stochastically label different molecules. For example, the molecules to be labeled comprise less than 75%, 70%, 65%, 60%, 55%, 50%, 45%, 40%, 35%, 30%, 25%, 20%, 15%, 10%, 5%, 4%, 3%, 2%, 1% sequence identity. The different molecules may differ by at least about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 40, 50, 60, 70, 80, 90, 100 or more nucleotides or base pairs. In some instances, at least one molecule is stochastically labeled. In other instances, at least 2, 3, 4, 5, 6, 7, 8, 9, or 10 different molecules are stochastically labeled. Alternatively, at least 20, 30, 40, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500, 600, 700, 800, 900, or 1000 different molecules are stochastically labeled. In other instances, at least 1500; 2,000; 2500; 3,000; 3500; 4,000; 4500; 5,000; 6,000; 7,000; 8,000; 9,000; or 10000 different molecules are stochastically labeled. In other instances; at least 15,000; 20,000; 25,000; 30,000; 35,000; 40,000; 45,000; 50,000; 60,000; 70,000; 80,000; 90,000; or 100,000 different molecules are stochastically labeled.

[00440] The different molecules to be labeled may be present in the sample at different concentrations or amounts. For example, the concentration or amount of one molecule is greater than the concentration or amount of another molecule in the sample. In some instances, the concentration or amount of at least one molecule in the sample is at least about 1.5, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, or 100 or more times greater than the concentration or amount of at least one other molecule in the sample. In some instances, the concentration or amount of at least one molecule in the sample is at least about 1000 or more times greater than the concentration or amount of at least one other molecule in the sample. In another example, the concentration or amount of one molecule is less than the concentration or amount of another molecule in the sample. The concentration or amount of at least one molecule in the sample may be at least about 1.5, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, or 100 or more times less than the concentration or amount of at least one other molecule in the sample. The concentration or amount of at least one molecule in the sample may be at least about 1000 or more times less than the concentration or amount of at least one other molecule in the sample.

[00441] In some instances, the molecules to be labeled are in one or more samples. The molecules to be labeled may be in two or more samples. The two or more samples may contain different amounts or concentrations of the molecules to be labeled. In some instances, the concentration or amount of one molecule in one sample may be greater than the concentration or amount of the same molecule in a different sample. For example, a blood sample might contain a higher amount of a particular molecule than a urine sample.

Alternatively, a single sample is divided into two or more subsamples. The subsamples may contain different amounts or concentrations of the same molecule. The concentration or amount of at least one molecule in one sample may be at least about 1.5, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, or 100 or more times greater than the concentration or amount of the same molecule in another sample. Alternatively, the concentration or amount of one molecule in one sample may be less than the concentration or amount of the same molecule in a different sample. For example, a heart tissue sample might contain a higher amount of a particular molecule than a lung tissue sample. The concentration or amount of at least one molecule in one sample may be at least about 1.5, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, or 100 or more times less than the concentration or amount of the same molecule in another sample. In some instances, the different concentrations or amounts of a molecule in two or more different samples is referred to as sample bias.

[00442] The methods and kits disclosed herein may be used for the analysis of two or more molecules from two or more samples. The two or more molecules may comprise two or more polypeptides. The method may comprise determining the identity of two or more labeled polypeptides. Determining the identity of two or more labeled polypeptides may comprise mass spectrometry. The method may further comprise combining the labeled polypeptides of the first sample with the labeled polypeptides of the second sample. The labeled polypeptides may be combined prior to determining the number of different labeled polypeptides. The method may further comprise combining the first sample-tagged polypeptides and the second sample-tagged polypeptides. The first sample-tagged polypeptides and the second sample-tagged polypeptides may be combined prior to contact with the plurality of molecular identifier labels. Determining the number of different labeled polypeptides may comprise detecting at least a portion of the labeled polypeptide. Detecting at least a portion of the labeled polypeptide may comprise detecting at least a portion of the sample tag, molecular identifier label, polypeptide, or a combination thereof.

[00443] As used herein, the term “polypeptide” refers to a molecule comprising at least one peptide. In some instances, the polypeptide consists of a single peptide. Alternatively, the polypeptide comprises two or more peptides. For example, the polypeptide comprises at least about 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 30, 40, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500, 600, 700, 800, 900, or 1000 peptides. Examples of polypeptides include, but are not limited to, amino acids, proteins, peptides, hormones, oligosaccharides,

lipids, glycolipids, phospholipids, antibodies, enzymes, kinases, receptors, transcription factors, and ligands.

Subjects

[00444] The methods and kits disclosed herein may comprise use of a cell or sample from one or more subjects. A subject may be a human or a non-human subject. A subject may be living. A subject may be dead. A subject may be a human that is under the care of a caregiver (*e.g.*, medical professional). A subject may be suspected of having a disease. A subject may have a disease. A subject may have symptoms of a disease. A subject may be a subject that provides one or more samples. A subject may be a mammal, reptile, amphibian, and/or bird. A subject may be a non-human primate.

Enzymes

[00445] The methods and kits disclosed herein may comprise one or more enzymes. Examples of enzymes include, but are not limited to ligases, reverse transcriptases, polymerases, and restriction nucleases. In some instances, attachment of the oligonucleotide tag to the molecules comprises the use of one or more ligases. Examples of ligases include, but are not limited to, DNA ligases such as DNA ligase I, DNA ligase III, DNA ligase IV, and T4 DNA ligase, and RNA ligases such as T4 RNA ligase I and T4 RNA ligase II.

[00446] The methods and kits disclosed herein may further comprise the use of one or more reverse transcriptases. In some instances, the reverse transcriptase is a HIV-1 reverse transcriptase, M-MLV reverse transcriptase, AMV reverse transcriptase, and telomerase reverse transcriptase. In some instances, the reverse transcriptase is M-MLV reverse transcriptase.

[00447] In some instances, the methods and kits disclosed herein comprise the use of one or more polymerases. Examples of polymerases include, but are not limited to, DNA polymerases and RNA polymerases. In some instances, the DNA polymerase is a DNA polymerase I, DNA polymerase II, DNA polymerase III holoenzyme, and DNA polymerase IV. Commercially available DNA polymerases include, but are not limited to, Bst 2.0 DNA Polymerase, Bst 2.0 WarmStart™ DNA Polymerase, Bst DNA Polymerase, Sulfolobus DNA Polymerase IV, Taq DNA Polymerase, 9°N™m DNA Polymerase, Deep VentR™ (exo-) DNA Polymerase, Deep VentR™ DNA Polymerase, Hemo KlenTaq™, LongAmp® Taq DNA Polymerase, OneTaq® DNA Polymerase, Phusion® DNA Polymerase, Q5™ High-Fidelity DNA Polymerase, Terminator™ γ DNA Polymerase, Terminator™ DNA Polymerase, Terminator™ II DNA Polymerase, Terminator™ III DNA Polymerase, VentR® DNA Polymerase, VentR® (exo-) DNA Polymerase, Bsu DNA Polymerase, phi29

DNA Polymerase, T4 DNA Polymerase, T7 DNA Polymerase, Terminal Transferase, Titanium® Taq Polymerase, KAPA Taq DNA Polymerase and KAPA Taq Hot Start DNA Polymerase.

[00448] Alternatively, the polymerase is an RNA polymerases such as RNA polymerase I, RNA polymerase II, RNA polymerase III, E. coli Poly(A) polymerase, phi6 RNA polymerase (RdRP), Poly(U) polymerase, SP6 RNA polymerase, and T7 RNA polymerase.

[00449] In some instances, the methods and kits disclosed herein comprise one or more restriction enzymes. Restriction enzymes include type I, type II, type III, and type IV restriction enzymes. In some instances, Type I enzymes are complex, multi-subunit, combination restriction-and-modification enzymes that cut DNA at random far from their recognition sequences. Generally, type II enzymes cut DNA at defined positions close to or within their recognition sequences. They may produce discrete restriction fragments and distinct gel banding patterns. Type III enzymes are also large combination restriction-and-modification enzymes. They often cleave outside of their recognition sequences and may require two such sequences in opposite orientations within the same DNA molecule to accomplish cleavage; they rarely give complete digests. In some instances, type IV enzymes recognize modified, typically methylated DNA and may be exemplified by the McrBC and Mrr systems of E. coli.

Additional Reagents

[00450] The methods and kits disclosed herein may comprise the use of one or more reagents. Examples of reagents include, but are not limited to, PCR reagents, ligation reagents, reverse transcription reagents, enzyme reagents, hybridization reagents, sample preparation reagents, and reagents for nucleic acid purification and/or isolation.

[00451] The methods and kits disclosed herein may comprise the use of one or more buffers. Examples of buffers include, but are not limited to, wash buffers, ligation buffers, hybridization buffers, amplification buffers, and reverse transcription buffers. In some instances, the hybridization buffer is a commercially available buffer, such as TMAC Hyb solution, SSPE hybridization solution, and ECONOTM hybridization buffer. The buffers disclosed herein may comprise one or more detergents.

[00452] The methods and kits disclosed herein may comprise the use of one or more carriers. Carriers may enhance or improve the efficiency of one or more reactions disclosed herein (*e.g.*, ligation reaction, reverse transcription, amplification, hybridization). Carriers may decrease or prevent non-specific loss of the molecules or any products thereof (*e.g.*, labeled-molecule, labeled-cDNA molecule, labeled-amplicon). For example, the carrier may

decrease non-specific loss of a labeled-molecule through absorption to surfaces. The carrier may decrease the affinity of the molecule, labeled-molecule, or any product thereof to a surface or substrate (*e.g.*, container, eppendorf tube, pipet tip). Alternatively, the carrier may increase the affinity of the molecule or any product thereof to a surface or substrate (*e.g.*, bead, array, glass, slide, chip). Carriers may protect the molecule or any product thereof from degradation. For example, carriers may protect an RNA molecule or any product thereof from ribonucleases. Alternatively, carriers may protect a DNA molecule or any product thereof from a DNase. Examples of carriers include, but are not limited to, nucleic acid molecules such as DNA and/or RNA, or polypeptides. Examples of DNA carriers include plasmids, vectors, polyadenylated DNA, and DNA oligonucleotides. Examples of RNA carriers include polyadenylated RNA, phage RNA, phage MS2 RNA, E.coli RNA, yeast RNA, yeast tRNA, mammalian RNA, mammalian tRNA, short polyadenylated synthetic ribonucleotides and RNA oligonucleotides. The RNA carrier may be a polyadenylated RNA. Alternatively, the RNA carrier may be a non-polyadenylated RNA. In some instances, the carrier is from a bacteria, yeast, or virus. For example, the carrier may be a nucleic acid molecule or a polypeptide derived from a bacteria, yeast or virus. For example, the carrier is a protein from *Bacillus subtilis*. In another example, the carrier is a nucleic acid molecule from *Escherichia coli*. Alternatively, the carrier is a nucleic acid molecule or peptide from a mammal (*e.g.*, human, mouse, goat, rat, cow, sheep, pig, dog, or rabbit), avian, amphibian, or reptile.

[00453] The methods and kits disclosed herein may comprise the use of one or more control agents. Control agents may include control oligos, inactive enzymes, non-specific competitors. Alternatively, the control agents comprise bright hybridization, bright probe controls, nucleic acid templates, spike-in controls, PCR amplification controls. The PCR amplification controls may be positive controls. In other instances, the PCR amplification controls are negative controls. The nucleic acid template controls may be of known concentrations. The control agents may comprise one or more labels.

[00454] Spike-in controls may be templates that are added to a reaction or sample. For example, a spike-in template may be added to an amplification reaction. The spike-in template may be added to the amplification reaction any time after the first amplification cycle. In some instances, the spike-in template is added to the amplification reaction after the 2nd, 3rd, 4th, 5th, 6th, 7th, 8th, 9th, 10th, 11th, 12th, 13th, 14th, 15th, 20th, 25th, 30th, 35th, 40th, 45th, or 50th amplification cycle. The spike-in template may be added to the amplification reaction any time before the last amplification cycle. The spike-in template may comprise one or more nucleotides or nucleic acid base pairs. The spike-in template may

comprise DNA, RNA, or any combination thereof. The spike-in template may comprise one or more labels.

Detectable labels

[00455] The methods, kits, and compositions disclosed herein may further comprise a detectable label. The terms “detectable label”, “tag” or “label” may be used interchangeably and refer to any chemical moiety attached to a molecule (*e.g.*, nucleotide, nucleotide polymer, or nucleic acid binding factor, molecular barcode). The chemical moiety may be covalently attached the molecule. The chemical moiety may be non-covalently attached to the molecule. The molecular barcodes, sample tags and molecular identifier labels may further comprise a detectable label, tag or label. Preferably, the label is detectable and renders the nucleotide or nucleotide polymer detectable to the practitioner of the invention. Detectable labels that may be used in combination with the methods disclosed herein include, for example, a fluorescent label, a chemiluminescent label, a quencher, a radioactive label, biotin, pyrene moiety, gold, or combinations thereof. Non-limiting example of detectable labels include luminescent molecules, fluorochromes, fluorescent quenching agents, colored molecules, radioisotopes or scintillants.

[00456] In some instances, the methods disclosed herein further comprise attaching one or more detectable labels to the molecular barcode, molecular identifier label, the sample tag, the labeled nucleic acid or any product thereof (*e.g.*, labeled-amplicon). The methods may comprise attaching two or more detectable labels to the molecular barcode, molecular identifier label, the sample tag or the labeled nucleic acid. Alternatively, the method comprises attaching at least about 3, 4, 5, 6, 7, 8, 9, or 10 detectable labels to the molecular barcode, molecular identifier label, the sample tag or the labeled nucleic acid. In some instances, the detectable label is a CyTM label. The CyTM label is a Cy3 label. Alternatively, or additionally, the detectable label is biotin. In some embodiments the detectable label is attached to a probe which binds to the molecular barcode, molecular identifier label, the sample tag or the labeled nucleic acid. This may occur, for example, after the nucleic acid or labeled nucleic acid has been hybridized to an array. In one example the nucleic acid or labeled nucleic acid is bound to partners on an array. After the binding, a probe which may bind the labeled nucleic acid is bound to the molecules on the array. This process may be repeated with multiple probes and labels to decrease the likelihood that a signal is the result of nonspecific binding of a label or nonspecific binding of the molecule to the array.

[00457] A donor acceptor pair may be used as the detectable labels. Either the donor or acceptor may be attached to a probe that binds a nucleic acid. The probe may be, for example,

a nucleic acid probe that may bind to a nucleic acid or the labeled nucleic acid. The corresponding donor or acceptor may be added to cause a signal.

[00458] In some instances, the detectable label is a Freedom dye, Alexa Fluor® dye, CyTM dye, fluorescein dye, or LI-COR IRDyes®. In some instances, the Freedom dye is fluorescein (6-FAMTM, 6-carboxyfluorescein), MAX (NHS Ester), TYETM 563, TEX 615, TYETM 665, TYE 705. The detectable label may be an Alexa Fluor dye. Examples of Alexa Fluor® dyes include Alexa Fluor® 488 (NHS Ester), Alexa Fluor® 532 (NHS Ester), Alexa Fluor® 546 (NHS Ester), Alexa Fluor® 594 (NHS Ester), Alexa Fluor® 647 (NHS Ester), Alexa Fluor® 660 (NHS Ester), or Alexa Fluor® 750 (NHS Ester). Alternatively, the detectable label is a CyTM dye. Examples of CyTM dyes include, but are not limited to, Cy2, Cy3, Cy3B, Cy3.5, Cy5, Cy5.5, and Cy7. In some instances, the detectable label is a fluorescein dye. Non-limiting examples of fluorescein dyes include 6-FAMTM (Azide), 6-FAMTM (NHS Ester), Fluorescein dT, JOE (NHS Ester), TETTM, and HEXTM. In some instances, the detectable label is a LI-COR IRDyes®, such as 5' IRDye® 700, 5' IRDye® 800, or IRDye® 800CW (NHS Ester). In some instances, the detectable label is TYETM 563. Alternatively, the detectable label is Cy3.

[00459] The detectable label may be Rhodamine dye. Examples of rhodamine dyes include, but are not limited to, Rhodamine GreenTM-X (NHS Ester), TAMRATM, TAMRATM (NHS Ester), Rhodamine RedTM-X(NHS Ester), ROXTM (NHS Ester), and 5'TAMRATM (Azide). In other instances, the detectable label is a WellRED Dye. WellRED Dyes include, but are not limited to, WellRED D4 dye, WellRED D3 dye, and WellRED D2 dye. In some instances, the detectable label is Texas Red®-X (NHS Ester), Lightcycler® 640 (NHS Ester), or Dy 750 (NHS Ester).

[00460] In some instances, detectable labels include a linker molecule. Examples of linker molecules include, but are not limited to, biotin, avidin, streptavidin, HRP, protein A, protein G, antibodies or fragments thereof, Grb2, polyhistidine, Ni²⁺, FLAG tags, myc tags. Alternatively, detectable labels include heavy metals, electron donors/acceptors, acridinium esters, dyes and calorimetric substrates. In other instances, detectable labels include enzymes such as alkaline phosphatase, peroxidase and luciferase.

[00461] A change in mass may be considered a detectable label, as is the case of surface plasmon resonance detection. The skilled artisan would readily recognize useful detectable labels that are not mentioned herein, which may be employed in the operation of the present invention.

[00462] In some instances, detectable labels are used with primers. For example, the universal primer is labeled with the detectable label (*e.g.*, Cy3 labeled universal primer, fluorophore labeled universal primer). Alternatively, the target specific primer is labeled with the detectable label (*e.g.*, TYE 563-labeled target specific primer). In other instances, detectable labels are used with the sample tags or molecular identifier labels. For example, the oligonucleotide tag is labeled with a detectable label (*e.g.*, biotin-labeled oligonucleotide tag). In other instances, detectable labels are used with the nucleic acid template molecule. Detectable labels may be used to detect the labeled-molecules or labeled-amplicons.

Alternatively, detectable labels are used to detect the nucleic acid template molecule.

[00463] In some instances, the detectable label is attached to the primer, molecular barcode, sample tag, molecular identifier label, labeled-molecule, labeled-amplicon, probe, HCR probe, and/or non-labeled nucleic acid. Methods for attaching the detectable label to the primer, oligonucleotide tag, labeled-molecule, labeled-amplicon, and/or non-labeled nucleic acid include, but are not limited to, chemical labeling and enzymatic labeling. In some instances, the detectable label is attached by chemical labeling. In some embodiments, chemical labeling techniques comprise a chemically reactive group. Non-limiting examples of reactive groups include amine-reactive succinimidyl esters such as NHS-fluorescein or NHS-rhodamine, amine-reactive isothiocyanate derivatives including FITC, and sulfhydryl-reactive maleimide-activated fluorophores such as fluorescein-5-maleimide. In some embodiments, reaction of any of these reactive dyes with another molecule results in a stable covalent bond formed between a fluorophore and the linker and/or agent. In some embodiments, the reactive group is isothiocyanates. In some embodiments, a label is attached to an agent through the primary amines of lysine side chains. In some embodiments, chemical labeling comprises a NHS-ester chemistry method.

[00464] Alternatively, the detectable label is attached by enzymatic labeling. Enzymatic labeling methods may include, but are not limited to, a biotin acceptor peptide/biotin ligase (AP/Bir A), acyl carrier protein/phosphopantetheine transferase (ACP/PPTase), human O⁶-alkylguanine transferase (hAGT), Q-tag/transglutaminase (TGase), aldehyde tag/formylglycine-generating enzyme, mutated prokaryotic dehalogenase (HaloTagTM), and farnesylation motif/protein farnesyltransferase (PFTase) methods. Affinity labeling may include, but is not limited to, noncovalent methods utilizing dihydrofolate reductase (DHFR) and Phe36Val mutant of FK506-binding protein 12 (FKBP12(F36V)), and metal-chelation methods.

[00465] Crosslinking reagents may be used to attach a detectable label to the primer, oligonucleotide tag, labeled-molecule, labeled-amplicon, and/or non-labeled nucleic acid. In some instances, the crosslinking reagent is glutaraldehyde. Glutaraldehyde may react with amine groups to create crosslinks by several routes. For example, under reducing conditions, the aldehydes on both ends of glutaraldehyde couple with amines to form secondary amine linkages.

[00466] In some instances, attachment of the detectable label to the primer, oligonucleotide tag, labeled-molecule, labeled-amplicon, and/or non-labeled nucleic acid comprises periodate-activation followed by reductive amination. In some instances, Sulfo-SMCC or other heterobifunctional crosslinkers are used to conjugate the detectable to the primer, oligonucleotide tag, labeled-molecule, labeled-amplicon, and/or non-labeled nucleic acid. For example, Sulfo-SMCC is used to conjugate an enzyme to a drug. In some embodiments, the enzyme is activated and purified in one step and then conjugated to the drug in a second step. In some embodiments, the directionality of crosslinking is limited to one specific orientation (*e.g.*, amines on the enzyme to sulfhydryl groups on the antibody).

Diseases/Conditions

[00467] Disclosed herein are methods, kits and compositions for diagnosing, monitoring, and/or prognosing a status or outcome of a disease or condition in a subject. Generally, the method comprises (a) stochastically labeling two or more molecules from two or more samples to produce two or more labeled nucleic acids; (b) detecting and/or quantifying the two or more labeled nucleic acids; and (c) diagnosing, monitoring, and/or prognosing a status or outcome of a disease or condition in a subject based on the detecting and/or quantifying of the two or more labeled nucleic acids. The method may further comprise determining a therapeutic regimen. The two or more of samples may comprise one or more samples from a subject suffering from a disease or condition. The two or more samples may comprise one or more samples from a healthy subject. The two or more samples may comprise one or more samples from a control.

[00468] Monitoring a disease or condition may further comprise monitoring a therapeutic regimen. Monitoring a therapeutic regimen may comprise determining the efficacy of a therapeutic regimen. In some instances, monitoring a therapeutic regimen comprises administering, terminating, adding, or altering a therapeutic regimen. Altering a therapeutic regimen may comprise increasing or reducing the dosage, dosing frequency, or mode of administration of a therapeutic regimen. A therapeutic regimen may comprise one or more

therapeutic drugs. The therapeutic drugs may be an anticancer drug, antiviral drug, antibacterial drug, antipathogenic drug, or any combination thereof.

Cancer

[00469] In some instances, the disease or condition is a cancer. The molecules to be stochastically labeled may be from a cancerous cell or tissue. In some instances, the cancer is a sarcoma, carcinoma, lymphoma or leukemia. Sarcomas are cancers of the bone, cartilage, fat, muscle, blood vessels, or other connective or supportive tissue. Sarcomas include, but are not limited to, bone cancer, fibrosarcoma, chondrosarcoma, Ewing's sarcoma, malignant hemangioendothelioma, malignant schwannoma, bilateral vestibular schwannoma, osteosarcoma, soft tissue sarcomas (*e.g.*, alveolar soft part sarcoma, angiosarcoma, cystosarcoma phylloides, dermatofibrosarcoma, desmoid tumor, epithelioid sarcoma, extraskelatal osteosarcoma, fibrosarcoma, hemangiopericytoma, hemangiosarcoma, Kaposi's sarcoma, leiomyosarcoma, liposarcoma, lymphangiosarcoma, lymphosarcoma, malignant fibrous histiocytoma, neurofibrosarcoma, rhabdomyosarcoma, and synovial sarcoma).

[00470] Carcinomas are cancers that begin in the epithelial cells, which are cells that cover the surface of the body, produce hormones, and make up glands. By way of non-limiting example, carcinomas include breast cancer, pancreatic cancer, lung cancer, colon cancer, colorectal cancer, rectal cancer, kidney cancer, bladder cancer, stomach cancer, prostate cancer, liver cancer, ovarian cancer, brain cancer, vaginal cancer, vulvar cancer, uterine cancer, oral cancer, penile cancer, testicular cancer, esophageal cancer, skin cancer, cancer of the fallopian tubes, head and neck cancer, gastrointestinal stromal cancer, adenocarcinoma, cutaneous or intraocular melanoma, cancer of the anal region, cancer of the small intestine, cancer of the endocrine system, cancer of the thyroid gland, cancer of the parathyroid gland, cancer of the adrenal gland, cancer of the urethra, cancer of the renal pelvis, cancer of the ureter, cancer of the endometrium, cancer of the cervix, cancer of the pituitary gland, neoplasms of the central nervous system (CNS), primary CNS lymphoma, brain stem glioma, and spinal axis tumors. In some instances, the cancer is a skin cancer, such as a basal cell carcinoma, squamous cell carcinoma, melanoma, nonmelanoma, or actinic (solar) keratosis.

[00471] In some instances, the cancer is a lung cancer. Lung cancer may start in the airways that branch off the trachea to supply the lungs (bronchi) or the small air sacs of the lung (the alveoli). Lung cancers include non-small cell lung carcinoma (NSCLC), small cell lung carcinoma, and mesothelioma. Examples of NSCLC include squamous cell carcinoma, adenocarcinoma, and large cell carcinoma. The mesothelioma may be a cancerous tumor of the lining of the lung and chest cavity (pleura) or lining of the abdomen (peritoneum). The

mesothelioma may be due to asbestos exposure. The cancer may be a brain cancer, such as a glioblastoma.

[00472] Alternatively, the cancer may be a central nervous system (CNS) tumor. CNS tumors may be classified as gliomas or nongliomas. The glioma may be malignant glioma, high grade glioma, diffuse intrinsic pontine glioma. Examples of gliomas include astrocytomas, oligodendrogliomas (or mixtures of oligodendroglioma and astrocytoma elements), and ependymomas. Astrocytomas include, but are not limited to, low-grade astrocytomas, anaplastic astrocytomas, glioblastoma multiforme, pilocytic astrocytoma, pleomorphic xanthoastrocytoma, and subependymal giant cell astrocytoma.

Oligodendrogliomas include low-grade oligodendrogliomas (or oligoastrocytomas) and anaplastic oligodendrogliomas. Nongliomas include meningiomas, pituitary adenomas, primary CNS lymphomas, and medulloblastomas. In some instances, the cancer is a meningioma.

[00473] The leukemia may be an acute lymphocytic leukemia, acute myelocytic leukemia, chronic lymphocytic leukemia, or chronic myelocytic leukemia. Additional types of leukemias include hairy cell leukemia, chronic myelomonocytic leukemia, and juvenile myelomonocytic leukemia.

[00474] Lymphomas are cancers of the lymphocytes and may develop from either B or T lymphocytes. The two major types of lymphoma are Hodgkin's lymphoma, previously known as Hodgkin's disease, and non-Hodgkin's lymphoma. Hodgkin's lymphoma is marked by the presence of the Reed-Sternberg cell. Non-Hodgkin's lymphomas are all lymphomas which are not Hodgkin's lymphoma. Non-Hodgkin lymphomas may be indolent lymphomas and aggressive lymphomas. Non-Hodgkin's lymphomas include, but are not limited to, diffuse large B cell lymphoma, follicular lymphoma, mucosa-associated lymphatic tissue lymphoma (MALT), small cell lymphocytic lymphoma, mantle cell lymphoma, Burkitt's lymphoma, mediastinal large B cell lymphoma, Waldenström macroglobulinemia, nodal marginal zone B cell lymphoma (NMZL), splenic marginal zone lymphoma (SMZL), extranodal marginal zone B cell lymphoma, intravascular large B cell lymphoma, primary effusion lymphoma, and lymphomatoid granulomatosis.

Pathogenic Infection

[00475] In some instances, the disease or condition is a pathogenic infection. The molecules to be stochastically labeled may be from a pathogen. The pathogen may be a virus, bacterium, fungi, or protozoan. In some instances, the pathogen may be a protozoan, such as *Acanthamoeba* (e.g., *A. astronyxis*, *A. castellanii*, *A. culbertsoni*, *A. hatchetti*, *A. polyphaga*,

A. rhyodes, *A. healyi*, *A. divionensis*), *Brachiola* (*e.g.*, *B. connori*, *B. vesicularum*), *Cryptosporidium* (*e.g.*, *C. parvum*), *Cyclospora* (*e.g.*, *C. cayetanensis*), *Encephalitozoon* (*e.g.*, *E. cuniculi*, *E. hellem*, *E. intestinalis*), *Entamoeba* (*e.g.*, *E. histolytica*), *Enterocytozoon* (*e.g.*, *E. bieneusi*), *Giardia* (*e.g.*, *G. lamblia*), *Isospora* (*e.g.*, *I. belli*), *Microsporidium* (*e.g.*, *M. africanum*, *M. ceylonensis*), *Naegleria* (*e.g.*, *N. fowleri*), *Nosema* (*e.g.*, *N. algerae*, *N. ocularum*), *Pleistophora*, *Trachipleistophora* (*e.g.*, *T. anthropophthera*, *T. hominis*), and *Vittaforma* (*e.g.*, *V. corneae*). The pathogen may be a fungus, such as, *Candida*, *Aspergillus*, *Cryptococcus*, *Histoplasma*, *Pneumocystis*, and *Stachybotrys*.

[00476] The pathogen may be a bacterium. Exemplary bacteria include, but are not limited to, *Bordetella*, *Borrelia*, *Brucella*, *Campylobacter*, *Chlamydia*, *Chlamydophila*, *Clostridium*, *Corynebacterium*, *Enterococcus*, *Escherichia*, *Francisella*, *Haemophilus*, *Helicobacter*, *Legionella*, *Leptospira*, *Listeria*, *Mycobacterium*, *Mycoplasma*, *Neisseria*, *Pseudomonas*, *Rickettsia*, *Salmonella*, *Shigella*, *Staphylococcus*, *Streptococcus*, *Treponema*, *Vibrio*, or *Yersinia*.

[00477] The virus may be a reverse transcribing virus. Examples of reverse transcribing viruses include, but are not limited to, single stranded RNA-RT (ssRNA-RT) virus and double-stranded DNA –RT (dsDNA-RT) virus. Non-limiting examples of ssRNA-RT viruses include retroviruses, alpharetrovirus, betaretrovirus, gammaretrovirus, deltaretrovirus, epsilon retrovirus, lentivirus, spuma virus, metavirus, and pseudoviruses. Non-limiting examples of dsDNA-RT viruses include hepadenavirus and caulimovirus. The virus can be a DNA virus. The virus can be a RNA virus. The DNA virus may be a double-stranded DNA (dsDNA) virus. In some instances, the dsDNA virus is an adenovirus, herpes virus, or pox virus. Examples of adenoviruses include, but are not limited to, adenovirus and infectious canine hepatitis virus. Examples of herpes viruses include, but are not limited to, herpes simplex virus, varicella-zoster virus, cytomegalovirus, and Epstein-Barr virus. A non-limiting list of pox viruses includes smallpox virus, cow pox virus, sheep pox virus, monkey pox virus, and vaccinia virus. The DNA virus may be a single-stranded DNA (ssDNA) virus. The ssDNA virus may be a parvovirus. Examples of parvoviruses include, but are not limited to, parvovirus B19, canine parvovirus, mouse parvovirus, porcine parvovirus, feline panleukopenia, and Mink enteritis virus.

[00478] The virus can be a RNA virus. The RNA virus may be a double-stranded RNA (dsRNA) virus, (+) sense single-stranded RNA virus ((+)ssRNA) virus, or (-) sense single-stranded ((-) ssRNA) virus. A non-limiting list of dsRNA viruses include reovirus, orthoreovirus, cypovirus, rotavirus, bluetongue virus, and phytoreovirus. Examples of (+)

ssRNA viruses include, but are not limited to, picornavirus and togavirus. Examples of picornaviruses include, but are not limited to, enterovirus, rhinovirus, hepatovirus, cardiovirus, aphthovirus, poliovirus, parechovirus, erbovirus, kobuvirus, teschovirus, and coxsackie. In some instances, the togavirus is a rubella virus, Sindbis virus, Eastern equine encephalitis virus, Western equine encephalitis virus, Venezuelan equine encephalitis virus, Ross River virus, O'nyong'nyong virus, Chikungunya, or Semliki Forest virus. A non-limiting list of (-) ssRNA viruses include orthomyxovirus and rhabdovirus. Examples of orthomyxoviruses include, but are not limited to, influenzavirus a, influenzavirus B, influenzavirus C, isavirus, and thogotovirus. Examples of rhabdoviruses include, but are not limited to, cytorhabdovirus, dichorhabdovirus, ephemerovirus, lyssavirus, novirhabdovirus, and vesiculovirus.

Fetal disorders

[00479] In some instances, the disease or condition is pregnancy. The methods and kits disclosed herein may comprise diagnosing a fetal condition in a pregnant subject. The methods and kits disclosed herein may comprise identifying fetal mutations or genetic abnormalities. The molecules to be stochastically labeled may be from a fetal cell or tissue. Alternatively, or additionally, the molecules to be labeled may be from the pregnant subject.

[00480] The methods and kits disclosed herein may be used in the diagnosis, prediction or monitoring of autosomal trisomies (*e.g.*, Trisomy 13, 15, 16, 18, 21, or 22). In some cases the trisomy may be associated with an increased chance of miscarriage (*e.g.*, Trisomy 15, 16, or 22). In other cases, the trisomy that is detected is a liveborn trisomy that may indicate that an infant will be born with birth defects (*e.g.*, Trisomy 13 (Patau Syndrome), Trisomy 18 (Edwards Syndrome), and Trisomy 21 (Down Syndrome)). The abnormality may also be of a sex chromosome (*e.g.*, XXY (Klinefelter's Syndrome), XYY (Jacobs Syndrome), or XXX (Trisomy X)). The molecule(s) to be labeled may be on one or more of the following chromosomes: 13, 18, 21, X, or Y. For example, the molecule is on chromosome 21 and/or on chromosome 18, and/or on chromosome 13.

[00481] Further fetal conditions that may be determined based on the methods and kits disclosed herein include monosomy of one or more chromosomes (X chromosome monosomy, also known as Turner's syndrome), trisomy of one or more chromosomes (13, 18, 21, and X), tetrasomy and pentasomy of one or more chromosomes (which in humans is most commonly observed in the sex chromosomes, *e.g.*, XXXX, XXYY, XXXY, XYYY, XXXXX, XXXXY, XXXYY, XYYYY and XXYYY), monoploidy, triploidy (three of every

chromosome, *e.g.*, 69 chromosomes in humans), tetraploidy (four of every chromosome, *e.g.*, 92 chromosomes in humans), pentaploidy and multiploidy.

[00482] Further disclosed herein is a method of forensic analysis comprising any of the above described methods. Forensic scientists may use nucleic acids in various samples (*e.g.*, blood, semen, skin, saliva, hair) found at a crime scene to identify the presence of an individual at the scene, such as a perpetrator. This process is formally termed DNA profiling, but may also be called “genetic fingerprinting.” For example, DNA profiling comprises measuring and comparing the lengths of variable sections of repetitive DNA, such as short tandem repeats and minisatellites, in various samples and people. This method is usually an extremely reliable technique for matching a DNA sample from a person with DNA in a sample found at the crime scene. However, identification may be complicated if the scene is contaminated with DNA from several people. In this instance, as well as in other forensic applications, it may be advantageous to obtain absolute quantification of nucleic acids from a single cell or small number of cells.

[00483] In some instances, the disease or condition is an immune disorder. An immune disorder can be an inflammatory disorder, an autoimmune disorder, irritable bowel syndrome or ulcerative colitis. Examples of autoimmune diseases include Chrohn's disease, lupus, and Graves' disease.

[00484] In some instances, the disease or disorder is a neorlogical condition or disorder. A neorlogical condition or disorder can be Acquired Epileptiform Aphasia, Acute Disseminated Encephalomyelitis, Adrenoleukodystrophy, Agenesis of the corpus callosum, Agnosia, Aicardi syndrome, Alexander disease, Alpers' disease, Alternating hemiplegia, Alzheimer's disease, Amyotrophic lateral sclerosis (see Motor Neuron Disease), Anencephaly, Angelman syndrome, Angiomatosis, Anoxia, Aphasia, Apraxia, Arachnoid cysts, Arachnoiditis, Arnold-Chiari malformation, Arteriovenous malformation, Asperger's syndrome, Ataxia Telangiectasia, Attention Deficit Hyperactivity Disorder, Autism, Auditory processing disorder, Autonomic Dysfunction, , Back Pain, Batten disease, Behcet's disease, Bell's palsy, Benign Essential Blepharospasm, Benign Focal Amyotrophy, Benign Intracranial Hypertension, Bilateral frontoparietal polymicrogyria, Binswanger's disease, Blepharospasm, Bloch-Sulzberger syndrome, Brachial plexus injury, Brain abscess, Brain damage, Brain injury, Brain tumor, Brown-Sequard syndrome, Canavan disease, Carpal tunnel syndrome (CTS), Causalgia, Central pain syndrome, Central pontine myelinolysis, Centronuclear myopathy, Cephalic disorder, Cerebral aneurysm, Cerebral arteriosclerosis, Cerebral atrophy, Cerebral gigantism, Cerebral palsy, Charcot-Marie-Tooth disease, Chiari malformation,

Chorea, Chronic inflammatory demyelinating polyneuropathy (CIDP), Chronic pain, Chronic regional pain syndrome, Coffin Lowry syndrome, Coma, including Persistent Vegetative State, Complex I deficiency syndrome, Complex I deficiency syndrome, Complex II deficiency syndrome, Complex III deficiency syndrome, Complex IV/COX deficiency syndrome, Complex V deficiency syndrome, Congenital facial diplegia, Corticobasal degeneration, Cranial arteritis, Craniosynostosis, Creutzfeldt-Jakob disease, Cumulative trauma disorders, Cushing's syndrome, Cytomegalic inclusion body disease (CIBD), Cytomegalovirus Infection, , Dandy-Walker syndrome, Dawson disease, Deficiency of mitochondrial NADH dehydrogenase component of Complex I, De Morsier's syndrome, Dejerine-Klumpke palsy, Dejerine-Sottas disease, Delayed sleep phase syndrome, Dementia, Dermatomyositis, Neurological Dyspraxia, Diabetic neuropathy, Diffuse sclerosis, Dysautonomia, Dyscalculia, Dysgraphia, Dyslexia, Dystonia, , Early infantile epileptic encephalopathy, Empty sella syndrome, Encephalitis, Encephalocele, Encephalotrigeminal angiomatosis, Encopresis, Epilepsy, Erb's palsy, Erythromelalgia, Essential tremor, , Fabry's disease, Fahr's syndrome, Fainting, Familial spastic paralysis, Febrile seizures, Fisher syndrome, Friedreich's ataxia, FART Syndrome, Gaucher's disease, Gerstmann's syndrome, Giant cell arteritis, Giant cell inclusion disease, Globoid cell Leukodystrophy, Gray matter heterotopia, Guillain-Barre syndrome, HTLV-1 associated myelopathy, Hallervorden-Spatz disease, Head injury, Headache, Hemifacial Spasm, Hereditary Spastic Paraplegia, Heredopathia atactica polyneuritiformis, Herpes zoster oticus, Herpes zoster, Hirayama syndrome, Holoprosencephaly, Huntington's disease, Hydranencephaly, Hydrocephalus, Hypercortisolism, Hypoxia, Immune-Mediated encephalomyelitis, Inclusion body myositis, Incontinentia pigmenti, Infantile phytanic acid storage disease, Infantile Refsum disease, Infantile spasms, Inflammatory myopathy, Intracranial cyst, Intracranial hypertension, Joubert syndrome, Kearns-Sayre syndrome, Kennedy disease, Kinsbourne syndrome, Klippel Feil syndrome, Krabbe disease, , Kufor-Rakeb syndrome, Kugelberg-Welander disease, Kuru, Lafora disease, Lambert-Eaton myasthenic syndrome, Landau-Kleffner syndrome, Lateral medullary (Wallenberg) syndrome, Learning disabilities, Leigh's disease, Lennox-Gastaut syndrome, Lesch-Nyhan syndrome, Leukodystrophy, Lewy body dementia, Lissencephaly, Locked-In syndrome, Lou Gehrig's disease, Lumbar disc disease, Lyme disease - Neurological Sequelae, Machado-Joseph disease (Spinocerebellar ataxia type 3), Macrencephaly, Maple Syrup Urine Disease, Megalencephaly, Melkersson-Rosenthal syndrome, Menieres disease, Meningitis, Menkes disease, Metachromatic leukodystrophy, Microcephaly, Migraine, Miller Fisher syndrome, Mini-Stroke, Mitochondrial disease,

Mitochondrial dysfunction, Mitochondrial Myopathies, Mitochondrial Respiratory Chain Complex I Deficiency, Mobius syndrome, Monomelic amyotrophy, Motor Neuron Disease, Motor skills disorder, Moyamoya disease, Mucopolysaccharidoses, Multi-Infarct Dementia, Multifocal motor neuropathy, Multiple sclerosis, Multiple system atrophy with postural hypotension, Muscular dystrophy, Myalgic encephalomyelitis, Myasthenia gravis, Myelinoclastic diffuse sclerosis, Myoclonic Encephalopathy of infants, Myoclonus, Myopathy, Myotubular myopathy, Myotonia congenita, NADH-coenzyme Q reductase deficiency, NADH:Q(1) oxidoreductase deficiency, Narcolepsy, Neurofibromatosis, Neuroleptic malignant syndrome, Neurological manifestations of AIDS, Neurological sequelae of lupus, Neuromyotonia, Neuronal ceroid lipofuscinosis, Neuronal migration disorders, Niemann-Pick disease, Non 24-hour sleep-wake syndrome, Nonverbal learning disorder, , O'Sullivan-McLeod syndrome, Occipital Neuralgia, Occult Spinal Dysraphism Sequence, Ohtahara syndrome, Olivopontocerebellar atrophy, Opsoclonus myoclonus syndrome, Optic neuritis, Orthostatic Hypotension, Overuse syndrome, oxidative phosphorylation disorders, Palinopsia, Paresthesia, Parkinson's disease, Paramyotonia Congenita, Paraneoplastic diseases, Paroxysmal attacks, Parry-Romberg syndrome (also known as Rombergs Syndrome), Pelizaeus-Merzbacher disease, Periodic Paralysis, Peripheral neuropathy, Persistent Vegetative State, Pervasive neurological disorders, Photic sneeze reflex, Phytanic Acid Storage disease, Pick's disease, Pinched Nerve, Pituitary Tumors, PMG, Polio, Polymicrogyria, Polymyositis, Porencephaly, Post-Polio syndrome, Postherpetic Neuralgia (PHN), Postinfectious Encephalomyelitis, Postural Hypotension, Prader-Willi syndrome, Primary Lateral Sclerosis, Prion diseases, Progressive Hemifacial Atrophy also known as Rombergs_Syndrome, Progressive multifocal leukoencephalopathy, Progressive Sclerosing Poliodystrophy, Progressive Supranuclear Palsy, Pseudotumor cerebri, Ramsay-Hunt syndrome (Type I and Type II), Rasmussen's encephalitis, Reflex sympathetic dystrophy syndrome, Refsum disease, Repetitive motion disorders, Repetitive stress injury, Restless legs syndrome, Retrovirus-associated myelopathy, Rett syndrome, Reye's syndrome, Rombergs_Syndrome, Rabies, Saint Vitus dance, Sandhoff disease, Schytsophrenia, Schilder's disease, Schizencephaly, Sensory Integration Dysfunction, Septo-optic dysplasia, Shaken baby syndrome, Shingles, Shy-Drager syndrome, Sjogren's syndrome, Sleep apnea, Sleeping sickness, Sniatiation, Sotos syndrome, Spasticity, Spina bifida, Spinal cord injury, Spinal cord tumors, Spinal muscular atrophy, Spinal stenosis, Steele-Richardson-Olszewski syndrome, see Progressive Supranuclear Palsy, Spinocerebellar ataxia, Stiff-person syndrome, Stroke, Sturge-Weber syndrome, Subacute sclerosing

panencephalitis, Subcortical arteriosclerotic encephalopathy, Superficial siderosis, Sydenham's chorea, Syncope, Synesthesia, Syringomyelia, Tardive dyskinesia, Tay-Sachs disease, Temporal arteritis, Tethered spinal cord syndrome, Thomsen disease, Thoracic outlet syndrome, Tic Douloureux, Todd's paralysis, Tourette syndrome, Transient ischemic attack, Transmissible spongiform encephalopathies, Transverse myelitis, Traumatic brain injury, Tremor, Trigeminal neuralgia, Tropical spastic paraparesis, Trypanosomiasis, Tuberous sclerosis, Vasculitis including temporal arteritis, Von Hippel-Lindau disease (VHL), Viliuisk Encephalomyelitis (VE), Wallenberg's syndrome, Werdnig-Hoffman disease, West syndrome, Whiplash, Williams syndrome, Wilson's disease, X-Linked Spinal and Bulbar Muscular Atrophy, or Zellweger syndrome.

Definitions

[00485] Where a range of values is provided, it is understood that each intervening value, to the tenth of the unit of the lower limit unless the context clearly dictates otherwise, between the upper and lower limits of that range is also specifically disclosed. Each smaller range between any stated value or intervening value in a stated range and any other stated or intervening value in that stated range is encompassed within the invention. The upper and lower limits of these smaller ranges may independently be included or excluded in the range, and each range where either, neither or both limits are included in the smaller ranges is also encompassed within the invention, subject to any specifically excluded limit in the stated range. Where the stated range includes one or both of the limits, ranges excluding either or both of those included limits are also included in the invention.

[00486] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, some potential and preferred methods and materials are now described. All publications mentioned herein are incorporated herein by reference to disclose and describe the methods and/or materials in connection with which the publications are cited. It is understood that the present disclosure supersedes any disclosure of an incorporated publication to the extent there is a contradiction.

[00487] As will be apparent to those of skill in the art upon reading this disclosure, each of the individual embodiments described and illustrated herein has discrete components and features which may be readily separated from or combined with the features of any of the other several embodiments without departing from the scope or spirit of the present

invention. Any recited method can be carried out in the order of events recited or in any other order which is logically possible.

[00488] It must be noted that as used herein and in the appended claims, the singular forms “a”, “an”, and “the” include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to “a cell” includes a plurality of such cells and reference to “the peptide” includes reference to one or more peptides and equivalents thereof, *e.g.*, polypeptides, known to those skilled in the art, and so forth.

[00489] As used herein, the term “label” may refer to a unique oligonucleotide sequence that may allow a corresponding nucleic acid base and/or nucleic acid sequence to be identified. In some embodiments, the nucleic acid base and/or nucleic acid sequence may be located at a specific position on a larger polynucleotide sequence (*e.g.*, a polynucleotide attached to a bead).

[00490] As used herein, the term “hybridization” may refer to the process in which two single-stranded polynucleotides bind non-covalently to form a stable double-stranded polynucleotide. The term “hybridization” may also refer to triple-stranded hybridization. The resulting (usually) double-stranded polynucleotide is a “hybrid” or “duplex.”

[00491] As used herein, “nucleoside” may include natural nucleosides, such as 2′-deoxy and 2′-hydroxyl forms. “Analog” in reference to nucleosides may include synthetic nucleosides comprising modified base moieties and/or modified sugar moieties, or the like. Analogs may be capable of hybridization. Analogs may include synthetic nucleosides designed to enhance binding properties, reduce complexity, increase specificity, and the like. Exemplary types of analogs may include oligonucleotide phosphoramidates (referred to herein as “amidates”), peptide nucleic acids (referred to herein as “PNAs”), oligo-2′-O-alkylribonucleotides, polynucleotides containing C-5 propynylpyrimidines, and locked nucleic acids (LNAs).

[00492] As used herein, the terms “nucleic acid molecule,” “nucleic acid sequence,” “nucleic acid fragment,” “oligonucleotide,” “oligonucleotide fragment” and “polynucleotide” may be used interchangeably and may be intended to include, but are not limited to, polymeric forms of nucleotides that may have various lengths, either deoxyribonucleotides or ribonucleotides, or analogs thereof. Nucleic acid molecules may include single stranded DNA (ssDNA), double stranded DNA (dsDNA), single stranded RNA (ssRNA) and double stranded RNA (dsRNA). Different nucleic acid molecules may have different three-dimensional structures, and may perform various functions. Non-limiting examples of nucleic acid molecules may include a gene, a gene fragment, a genomic gap, an exon, an intron, intergenic DNA (including, without limitation, heterochromatic DNA), messenger RNA

(mRNA), transfer RNA, ribosomal RNA, ribozymes, small interfering RNA (siRNA), miRNA, small nucleolar RNA (snoRNA), cDNA, recombinant polynucleotides, branched polynucleotides, plasmids, vectors, isolated DNA of a sequence, isolated RNA of a sequence, nucleic acid probes, and primers.

[00493] Oligonucleotides may refer to a linear polymer of natural or modified nucleosidic monomers linked by phosphodiester bonds or analogs thereof. An “oligonucleotide fragment” refers to an oligonucleotide sequence that has been cleaved into two or more smaller oligonucleotide sequences. Oligonucleotides may be natural or synthetic. Oligonucleotides may include deoxyribonucleosides, ribonucleosides, and non-natural analogs thereof, such as anomeric forms thereof, peptide nucleic acids (PNAs), and the like. Oligonucleotides may be capable of specifically binding to a target genome by way of a regular pattern of monomer-to-monomer interactions, such as Watson-Crick type of base pairing, base stacking, Hoogsteen or reverse Hoogsteen types of base pairing, or the like. Oligonucleotides and the term “polynucleotides” may be used interchangeably herein.

[00494] Whenever an oligonucleotide is represented by a sequence of letters, such as “ATGCCTG,” it may be understood that the nucleotides are in 5′ to 3′ order from left to right and that “A” denotes deoxyadenosine, “C” denotes deoxycytidine, “G” denotes deoxyguanosine, “T” denotes deoxythymidine, and “U” denotes the ribonucleoside, uridine, unless otherwise noted.

[00495] Oligonucleotides may include one or more non-standard nucleotide(s), nucleotide analog(s) and/or modified nucleotides. Examples of modified nucleotides may include, but are not limited to diaminopurine, 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xantine, 4-acetylcytosine, 5-(carboxyhydroxymethyl)uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5′-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-D46-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid (v), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, oligonucleotide phosphoramidates (referred to herein as “amidates”), peptide nucleic acids (referred to herein as “PNAs”), oligo-2′-O-alkylribonucleotides,

polynucleotides containing C-5 propynylpyrimidines, locked nucleic acids (LNAs), 2,6-diaminopurine and the like. Nucleic acid molecules may also be modified at the base moiety (*e.g.*, at one or more atoms that typically are available to form a hydrogen bond with a complementary nucleotide and/or at one or more atoms that are not typically capable of forming a hydrogen bond with a complementary nucleotide), sugar moiety or phosphate backbone.

[00496] As used herein, a “sample” may refer to a single cell or many cells. Nucleic acid molecules may be obtained from one or more samples. A sample may comprise a single cell type or a combination of two or more cell types. A sample may include a collection of cells that perform a similar function such as those found, for example, in a tissue. A sample may comprise one or more tissues. Examples of tissues may include, but are not limited to, epithelial tissue (*e.g.*, skin, the lining of glands, bowel, skin and organs such as the liver, lung, kidney), endothelium (*e.g.*, the lining of blood and lymphatic vessels), mesothelium (*e.g.*, the lining of pleural, peritoneal and pericardial spaces), mesenchyme (*e.g.*, cells filling the spaces between the organs, including fat, muscle, bone, cartilage and tendon cells), blood cells (*e.g.*, erythrocytes, granulocytes, neutrophils, eosinophils, basophils, monocytes, T-lymphocytes (also known as T-cells), B-lymphocytes (also known as B-cells), plasma cells, megakaryocytes and the like), neurons, germ cells (*e.g.*, spermatozoa, oocytes), amniotic fluid cells, placenta, stem cells and the like. A sample may be obtained from one or more of single cells in culture, metagenomic samples, embryonic stem cells, induced pluripotent stem cells, cancer samples, tissue sections, and biopsies, or any combination thereof.

[00497] As used herein, the term “organism” may include, but is not limited to, a human, a non-human primate, a cow, a horse, a sheep, a goat, a pig, a dog, a cat, a rabbit, a mouse, a rat, a gerbil, a frog, a toad, a fish (*e.g.*, *Danio rerio*) a roundworm (*e.g.*, *C. elegans*) and any transgenic species thereof. The term “organism” may also include, but is not limited to, a yeast (*e.g.*, *S. cerevisiae*) cell, a yeast tetrad, a yeast colony, a bacterium, a bacterial colony, a virion, virosome, virus-like particle and/or cultures thereof, and the like.

[00498] As used herein, the term “attach,” “conjugate,” and “couple” may be used interchangeably and may refer to both covalent interactions and noncovalent interactions.

[00499] While preferred embodiments of the present invention have been shown and described herein, it will be obvious to those skilled in the art that such embodiments are provided by way of example only. Numerous variations, changes, and substitutions will now occur to those skilled in the art without departing from the invention. It should be understood that various alternatives to the embodiments of the invention described herein may be

employed in practicing the invention. It is intended that the following claims define the scope of the invention and that methods and structures within the scope of these claims and their equivalents be covered thereby.

EXAMPLES

EXAMPLE 1: Enzymatic Split-Pool Synthesis

[00500] In this example, an enzymatic split-pool synthesis method was used to produce oligonucleotide coupled beads. As shown in **FIG. 2A**, a set of oligonucleotides was added to each well of a first plate. An oligonucleotide in a set of oligonucleotides comprises a 5' amine, universal sequence, cell label and a linker. The 5' amine, universal sequence and linker are the same for each set of oligonucleotides. The universal sequence and linker are different from each other. However, the cell label is different for each set of oligonucleotides. Thus, each well has a different cell label.

In Step 1 of the enzymatic split-pool synthesis, oligonucleotide-coupled beads were synthesized by adding a single bead to each well and performing 1-Ethyl-3-(3-dimethylaminopropyl)carbodiimide (EDC) coupling reactions. The oligonucleotides beads resulting from Step 1 comprise a bead coupled to multiple oligonucleotides. The oligonucleotide comprises a 5'-amine, universal sequence, cellular label 1, and linker 1 (see **FIG. 2A**). The oligonucleotides on the same bead are the same. However, oligonucleotides on a first bead are different from oligonucleotides on a second bead.

[00501] In Step 2 of the enzymatic split-pool synthesis, multiple washes were performed to remove uncoupled oligonucleotides. Once the uncoupled oligonucleotides were removed, the oligonucleotide-coupled beads were pooled (see **FIG. 2A**). The oligonucleotide coupled beads resulting from Step 2 comprise a bead coupled to multiple single stranded oligonucleotides. The single stranded oligonucleotide comprises a 5' amine, universal sequence, cell label 1 and linker 1. Each oligonucleotide on a bead is identical. However, each bead comprises a different oligonucleotide. The oligonucleotides coupled to the different beads differ by the cell label 1 sequence.

[00502] As shown in **FIG. 2B**, a set of oligonucleotides was added to each well of a second plate. An oligonucleotide in a set of oligonucleotides comprises a first linker, cell label, and a second linker. The first and second linkers are the same for each set of oligonucleotides. The first and second linkers are different from each other. However, the cell label is different for each set of oligonucleotides. Thus, each well has a different cell label.

[00503] In Step 3 of the enzymatic split-pool synthesis, the oligonucleotide coupled beads that were pooled in Step 2 were split into the wells of the second plate. Because the first

linker of the oligonucleotides in the wells of the second plate are complementary to the linker of the oligonucleotides coupled to the beads, primer extension using Klenow large fragment was performed to couple the oligonucleotides from the second plate to the oligonucleotide coupled beads from Step 2. The oligonucleotides coupled beads resulting from Step 3 comprise a bead coupled to multiple double stranded oligonucleotides. The double stranded oligonucleotide comprises a 5' amine, universal sequence, cell label 1, linker 1, cell label 2, and linker 2 (see **FIG. 2B**).

[00504] In Step 4 of the enzymatic split-pool synthesis, multiple washes were performed to remove uncoupled oligonucleotides and the Klenow large fragment enzymes. The second plate was heated to denature the double stranded oligonucleotides, and the oligonucleotide coupled beads were pooled (see **FIG. 2B**). The oligonucleotide coupled beads resulting from Step 4 comprise a bead coupled to multiple single stranded oligonucleotides. The single stranded oligonucleotide comprises a 5' amine, universal sequence, cell label 1, linker 1, cell label 2, and linker 2. Each oligonucleotide on a bead is identical. However, each bead comprises a different oligonucleotide. The oligonucleotides coupled to the different beads differ by the combined cell label sequences. For example, a first bead may comprise oligonucleotides comprising a first cell label of cell label A and second cell label of cell label C and a second bead may comprise oligonucleotides comprising a first cell label of cell label C and a second cell label of cell label D. Thus, the first bead and the second bead may comprise the same cell label (in this case, cell label C), however, the combined cell label sequences of the first bead and the second bead are different (*e.g.*, for the first bead, the combined cell label sequence is cell label A + cell label C; for the second bead, the combined cell label sequence is cell label C + cell label A). In other instances, two beads may comprise oligonucleotides comprising different cell labels. For example, a first bead may comprise oligonucleotides comprising cell label A and cell label B and a second bead may comprise oligonucleotides comprising cell label C and cell label D. In this instance, both of the cell labels of the first bead are different from both of the cell labels of the second bead.

[00505] As shown in **FIG. 2C**, a set of oligonucleotides was added to each well of a third plate. An oligonucleotide in a set of oligonucleotides comprises a linker, cell label, molecular label, and an oligodT. The linker and oligodT sequences are the same for each set of oligonucleotides. However, the cell label is different for each set of oligonucleotides. Thus, each well has a different cell label. In addition, the molecular label is different for oligonucleotides within a set. Thus, a single well contains a plurality of oligonucleotides with

the same cell label, but different molecular labels. The oligonucleotides from different wells may contain the same molecular label.

[00506] In Step 5 of the enzymatic split-pool synthesis, the oligonucleotide coupled beads that were pooled in Step 4 were split into the wells of the third plate. Because the linker of the oligonucleotides in the wells of the second plate are complementary to the second linker of the oligonucleotides coupled to the beads, primer extension using Klenow large fragment was performed to couple the oligonucleotides from the third plate to the oligonucleotide coupled beads from Step 4. The oligonucleotides coupled beads resulting from Step 5 comprise a bead coupled to multiple double stranded oligonucleotides. The double stranded oligonucleotide comprises a 5' amine, universal sequence, cell label 1, linker 1, cell label 2, linker 2, cell label 3, molecular label and oligodT (see **FIG. 2C**).

[00507] In Step 6 of the enzymatic split-pool synthesis, multiple washes were performed to remove uncoupled oligonucleotides and the Klenow large fragment enzymes. The third plate was heated to denature the double stranded oligonucleotides, and the oligonucleotide coupled beads were pooled (see **FIG. 2C**). The oligonucleotide coupled beads resulting from Step 4 comprise a bead coupled to multiple single stranded oligonucleotides. The single stranded oligonucleotide comprises a 5' amine, universal sequence, cell label 1, linker 1, cell label 2, linker 2, cell label 3, molecular label and oligodT. The multiple single stranded oligonucleotides on a single bead may be differentiated by the molecular label. The cell label portions of the multiple oligonucleotides on a single bead are identical. Each bead comprises different oligonucleotides. The oligonucleotides coupled to the different beads differ by the cell label sequences. The molecular label on the oligonucleotides from different beads may be the same. The molecular label on the oligonucleotides from different beads may be different. Two or more beads may differ by the combined cell label sequences. For example, a first bead may comprise an oligonucleotide comprising cell label A, cell label B and cell label C and a second bead may comprise an oligonucleotide comprising cell label B, cell label D and cell label A. In this instance, the first and second bead both contain cell label B, however the two other cell labels are different. Thus, two or more beads may comprise oligonucleotides differing by at least one cell label. Two or more beads may comprise oligonucleotides differing by at least two cell labels. Two or more beads may comprise oligonucleotides differing by at least three cell labels. However, a bead may comprise an oligonucleotide comprising two or more identical cell labels. For example, a bead may comprise an oligonucleotide comprising cell label A, cell label A and cell label D. A bead may comprise oligonucleotides comprising at least three identical cell labels. For example, a bead may

comprise an oligonucleotide comprising cell label A, cell label A and cell label A. A bead may comprise oligonucleotides comprising three non-identical cell labels. For example, a bead may comprise an oligonucleotide comprising cell label A, cell label D and cell label E. A bead may comprise at least two oligonucleotides comprising at least two different molecular labels. For example, a bead may comprise a first oligonucleotide comprising molecular label A and a second oligonucleotide comprising molecular label D. However, a bead may comprise multiple copies of an oligonucleotide comprising a first molecular label. Thus, a bead may comprise at least two oligonucleotides comprising the same molecular label. For example, a bead may comprise a first oligonucleotide comprising molecular label A and a second oligonucleotide comprising molecular label A. At least 30% of the oligonucleotides on a bead may comprise different molecular labels. At least 40% of the oligonucleotides on a bead may comprise different molecular labels. At least 50% of the oligonucleotides on a bead may comprise different molecular labels. At least 60% of the oligonucleotides on a bead may comprise different molecular labels. Less than 30% of the oligonucleotides on a bead may comprise the same molecular label. Less than 20% of the oligonucleotides on a bead may comprise the same molecular label. Less than 15% of the oligonucleotides on a bead may comprise the same molecular label. Less than 10% of the oligonucleotides on a bead may comprise the same molecular label. Less than 5% of the oligonucleotides on a bead may comprise the same molecular label.

[00508] The enzymatic split-pool synthesis technique may be performed on multiple plates or plates with a greater number of wells to produce a larger number of oligonucleotide coupled beads. The use of three separate cell label portions may increase the diversity of the total cell label portions on the beads. With 96 different sequence options for each cell label portion, 884,736 different cell label combinations may be created.

EXAMPLE 2: Comparison of amplification in tube and microwell

[00509] The disclosure provides a method for capturing cells. About 5,000 Ramos cells were captured on a microwell array comprising microwells of about 30 micron in diameter. Some cells were not captured. The control for the experiment was an equivalent number of cells captured in a tube. Both the cells in the tube and the cells in the microwell array were lysed. The nucleic acid was allowed to hybridize to a conjugated bead. Real time PCR of GAPDH and RPL19 genes was performed.

[00510] **FIG. 9** shows the results of the real time PCR amplification. The yield from the microwell was larger than the yield from the nucleic acid in the tube, indicating that the

hybridization of the nucleic acid to the oligonucleotide was more effective in the microwell than the tube (compare grey bar and white bar, respectively).

EXAMPLE 3: Comparison of amplification of second synthesized strand and synthesis on bead

[00511] Cells were obtained and lysed as described in Example 1. RPL19, TUBB, and GAPDH were amplified either off the second strand synthesized off the solid supports, or direct on the solid supports using a universal primer. **FIG. 10** shows, amplification directly on the solid supports (**FIG. 10**) yielded less off-target amplification than amplification not directly off a solid support. GAPDH and TUBB amplifications produced correctly sized products regardless of method (the left lane of each triplet in **FIG. 10** corresponds to solid support plus lysate in tube format, the middle lane of each triplet corresponds to solid supports from the microwell, and the right lane of each triplet corresponds to solid supports plus purified nucleic acid). The RPL19 product had minimal off-target amplification products, but only produced a strong product when purified nucleic acid was used with the solid support. These experiments indicate that amplification directly on the beads produces less off-target amplification products than amplification using a second strand synthesized off the solid support.

EXAMPLE 4: Multiplex analysis of target nucleic acids

[00512] Cells are obtained and lysed as described in Example 1. Target nucleic acids are hybridized to the solid support comprising oligonucleotides. A plurality of copies of the target nucleic acid are hybridized to a target binding region comprising an oligodT sequence. The plurality of copies of the target nucleic acid are reverse transcribed using reverse transcriptase. Reverse transcription incorporates the features of the oligonucleotide to which the copy of the target nucleic acid was hybridized (*e.g.*, the molecular label, the cellular label, and the universal label). The plurality of copies of the target nucleic acid are amplified using PCR. The amplified copies of the target nucleic acid are sequenced. The sequenced target nucleic acids are counted to determine the copy number of the target nucleic acid in the cell. The counting is performed by counting the number of different molecular labels for each of the same sequence read of target nucleic acid. In this way, amplification bias may be diminished.

EXAMPLE 5: Evaluating efficacy of split-pool synthesis to produce beads with clonal copies of one cell label combination

[00513] In this example, the efficacy of split-pool synthesis to produce beads with clonal copies of one cell label combination was evaluated. Oligonucleotide coupled beads were

synthesized by the enzymatic split-pool synthesis method as described in Example 1. 250 ng of total RNA was purified from Ramos cells, which is equivalent to RNA from 25,000 cells. The total RNA was contacted with 35,000 oligonucleotide coupled beads, resulting in hybridization of mRNA to the oligonucleotide coupled beads. cDNA synthesis was performed on the mRNA hybridized to the oligonucleotide coupled beads. Samples comprising 18, 175, and 1750 beads were used for further analysis. PCR amplification reactions using GAPDH-specific primers and IGJ-specific primers were performed on the cDNA bound to the beads from the 18-, 175- and 1750-bead samples. The cDNA molecules attached to the beads were sequenced. FIG. 11A-I show graphical representations of the sequencing results. For FIG. 11A-C, the number of reads per bead is plotted on the y-axis and the unique barcode (*e.g.*, cell label combination) is plotted on the x-axis for the 18-bead, 175-bead and 1750-bead samples, respectively. For FIG. 11D-F, the number of unique molecules per bead is plotted on the y-axis and the unique barcode (*e.g.*, cell label combination) is plotted on the x-axis for the 18-bead, 175-bead and 1750-bead samples, respectively. For FIG. 11G-I, the number of unique molecules per bead is plotted on the y-axis and the unique barcode is plotted on the x-axis for the 18-bead, 175-bead and 1750-bead samples, respectively. The results for FIG. 11G-I are sorted by the total number of molecules. The median number of unique molecules per bead for the various samples is shown in Table 1. Numerical values for the sequencing results are shown in Table 2. For FIG. 11J-L, the number of unique barcode (bc) combination using the index is plotted on the y-axis and the barcode (bc) segment index is plotted on the x-axis for the cell label 1, cell label 2, and cell label 3 for the 1750-bead sample, respectively. The barcode (bc) refers to the cell label (*e.g.*, bc segment1 = cell label part 1). As shown in FIG. 11J-L, the presence of almost all 96 barcodes within each segment was detected by sequencing. These results demonstrate the success of the enzymatic split-pool synthesis method to produce beads with clonal copies of one cell label combination.

Table 1			
	Median number of unique molecules		
	18 beads	175 beads	1750 beads
IGJ	78	85	40
GAPDH	22	45	25

Table 2			
Expected # of beads	17.5	175	1750
Total number of reads	58321	60308	133043
>=8 match in constant 1	56385	57615	123349
>=8 match in constant 2	54117	55187	115126
>=8 match in constant 1 & 2	54114	55185	115107
Perfect match in all 3 sub-barcode	38585	46066	95217
Perfect match in gene (40bp)	29968	33775	72260
Total number of unique barcode combination	239	407	1654
% useful reads	51.388%	56.00%	54.31%
Number of unique barcode combinations > 20 read	5	26	288

EXAMPLE 6. Single cell RNA labeling using oligonucleotide coupled beads

[00514] In this example, the efficacy of single cell RNA labeling using oligonucleotide coupled beads was evaluated. Three cell samples were prepared as follows:

	Sample 1: K562 only	Sample 2: Ramos only	Sample 3: Ramos + K562 mixture
Number of microwells	~10000	~10000	~10000
Number of Ramos cells	0	5000	3750
Number of K562 cells	1000	0	2500

[00515] The cell suspension of the samples was added to the top of a microwell and cells were allowed to settle into the wells of the microwell array. Cells not captured by the microwell array were washed away in a phosphate buffered saline (PBS) bath.

Oligonucleotide coupled beads, as prepared by the enzymatic split-pool synthesis method described in Example 1, were added to the microwell array. The oligonucleotide coupled bead comprises a magnetic bead with a plurality of oligonucleotides. Each oligonucleotide on the bead comprises a 5' amine, universal sequence, cell label 1, linker 1, cell label 2, linker 2, cell label 3, molecular label, and oligodT. For each oligonucleotide on the same bead, the sequences of the oligonucleotides are identical except for the molecular label. For oligonucleotides on different beads, the cell label 1, 2, and 3 combinations are different. Approximately 5-6 beads were added per well of the microwell array. In some instances, for every 10 wells, 50 beads may be deposited on the array, with 0-2 beads falling into each well. The beads were allowed to settle into the wells and uncaptured beads were washed away in a PBS bath. A magnet was placed underneath the microwell array. Cells were lysed by the addition of cold lysis buffer. The array and magnet were placed on a cold aluminum block for 5 minutes. mRNA from the lysed cells were hybridized to the oligonucleotides coupled to the beads. The array was washed with excess lysis buffer to remove unbound mRNA. The beads were retrieved from the wells by placing a magnet on top of the microwell array. The

retrieved beads were washed. cDNA synthesis was performed on the beads using Superscript III at 50 °C for 50 minutes on a rotor. Non-extended oligodT from the oligonucleotides on the beads were removed by ExoI treatment conducted at 37 °C for 30 minutes on a rotor. Gene-specific PCR amplification was conducted on the cDNA. The genes selected for the gene-specific PCR were cell-type specific and are shown in Table 3. The PCR amplified products were sequenced. Sequencing statistics are shown in Table 4. Figures 12A-C show a histogram of the sequencing results for the K562-only sample, Ramos-only sample, and K562 + Ramos mixture sample, respectively. For FIG. 12A-C, the unique molecule per barcode plotted on the y-axis and the unique bc combination index, sorted by read per bc plotted on the x-axis.

Table 3		
Number	Gene	Cell-type
1	CD74	Ramos-specific
2	CD79a	Ramos-specific
3	IGJ	Ramos-specific
4	TCL1A	Ramos-specific
5	SEPT9	Ramos-specific
6	CD27	Ramos-specific
7	CD41	K562-specific
8	GYPA	K562-specific
9	GATA1	K562-specific
10	GATA2	K562-specific
11	HBG1	K562-specific
12	GAPDH	common

Table 4			
	Sample 1: K562 only	Sample 2: Ramos only	Sample 3: Ramos + K562 mixture
Number of Ramos cells	0	5000	3750
Number of K562 cells	1000	0	2500
Total number of reads	717718	1329189	2399025
>=8 match in constant 1	657911	1201081	2026726
>=8 match in constant 2	581581	1071364	1513466
>=8 match in constant 1 & 2	581508	1071153	1513102
Perfect match in all 3 sub-barcodes	481564	862348	1248073
Perfect match in gene (40bp)	283463	575713	1004338
% useful reads	39.50%	43.31%	41.86%
Total number of unique barcode combination	8501	29647	28783
Number of unique barcode combinations >	145	1072	768

30 molecule			
Capture efficiency	0.145	0.2144	0.12288

[00516] Single cell labeling was used to determine the copy number for the single-cell type samples (*e.g.*, K562-only sample, Ramos-only sample). FIG. 12D-E shows a graph of the copy number for genes listed in Table 3 for the Ramos-only cell sample and K562-only cell sample, respectively. For FIG. 12D-E, the number of molecules per barcode (bc) combination is plotted on the y-axis and the unique barcode combination, sorted by total number of molecules per bc combination is plotted on the x-axis. The results shown in Figures 12D-E were based on sequencing data from beads with > 30 total number of unique molecules. These results demonstrate that the proportion of molecules per amplicon per bead matches expectations for the cell type. For the K562-only cell sample, the skew of the number of molecules is more severe and it appears that HBG1, which is highly abundant in this cell type, has a variable copy number. However, GAPDH copy number appears to be constant even though the total number of molecules per bead is skewed. The copy number for the individual genes are shown in FIG. 12F-I. For FIG. 12F-G, the copy number is represented as copy per bead or single cell for Ramos-only cells and K562-only cells, respectively. For FIG. 12H-I, the copy number is represented as relative abundance per bead or single cell for Ramos-only cells and K562-only cells, respectively.

[00517] Single cell labeling was used to determine the cell type of single cells in the K562+Ramos mixture sample. Sequencing results from 100 unique barcode combinations with the most abundant molecules were analyzed to evaluate the efficacy of single cell labeling to determine the cell type of single cells in the K562+Ramos mixture sample. FIG. 12J-M show graphs of the number of unique molecules per gene (y-axis) for the beads with the 100 unique barcode combinations. The numbers on the x-axis refer to the gene (see Table 3). FIG. 12J-M clearly depict general gene expression patterns for the K562 and Ramos cells. FIG. 12N-O show enlarged graphs of two beads that depict the general pattern of gene expression profiles for the two cell types. FIG. 12N shows the general pattern of gene expression profile for K562-like cells and FIG. 12O shows the general pattern of gene expression profile for Ramos-like cells. FIG. 12P shows a scatter plot of results based on principal component analysis of gene expression profile of 768 beads with > 30 molecules per bead from the K562+Ramos mixture sample. Component 1, which is plotted on the x-axis, separates the two cell types. Component 2, which is plotted on the y-axis, separates K562 cells with high and low HBG1 copy number. Each dot on the scatter plot represents one

unique barcode combination, which is equivalent to one bead or one cell. Based on the principal component analysis, 409 beads corresponded to K562 cells and 347 beads corresponded to Ramos cells. The copy number of the genes from Table 3 was determined for the K562-like and Ramos-like cell types. FIG. 12Q-R show histograms of the copy number per amplicon per bead for the K562-like cells (beads on the left of the first principal component based on FIG. 12P) and Ramos-like cells (beads on the right of the first principal component based on FIG. 12P), respectively. For FIG. 12Q-R, number of per bc combination is on the y-axis and unique barcode combination, sorted by total number of molecules per bc combination is on the x-axis. FIG. 12S-T show the copy number per bead or single cell of the individual genes for the K562-like cells (beads on the left of the first principal component based on FIG. 12P) and Ramos-like cells (beads on the right of the first principal component based on FIG. 12P), respectively. Table 5 shows the mean copy number per bead for the single cell and mixture samples.

Table 5				
Gene	Single cell type samples		K562+Ramos mixture sample	
	K562-only	Ramos-only	K562-like	Ramos-like
CD74	0.00	39.95	0.10	7.50
CD79a	0.02	30.97	0.84	18.88
IGJ	0.03	42.43	0.81	27.76
TCL1A	0.01	31.78	0.71	19.44
SEPT9	0.88	3.89	1.35	1.52
CD27	0.00	5.31	0.03	1.30
CD41	0.61	0.00	0.47	0.01
GYPA	1.92	0.00	0.73	0.02
GATA2	1.38	0.00	0.60	0.04
GATA1	0.94	0.00	1.04	0.04
HBG1	201.09	0.00	72.27	1.37
GAPDH	51.77	39.13	44.94	13.53
GAPDH read redundancy	2.04	1.47	7.67	7.22

EXAMPLE 7. Evaluating cross-talk between beads

[00518] In this example, the cross-talk between beads was evaluated. Samples comprising mixtures of mouse EL4 cells and Ramos cells were prepared as follows:

	High density	Low density
Number of microwells	~10000	~10000
Number of mouse EL4 cells	2500	1500
Number of Ramos cells	3750	1500

[00519] The cell suspension of the samples was added to the top of a microwell and cells were allowed to settle into the wells of the microwell array. Cells not captured by the microwell array were washed away in a phosphate buffered saline (PBS) bath. Oligonucleotide coupled beads, as prepared by the enzymatic split-pool synthesis method described in Example 1, were added to the microwell array. The oligonucleotide coupled bead comprises a magnetic bead with a plurality of oligonucleotides. Each oligonucleotide on the bead comprises a 5' amine, universal sequence, cell label 1, linker 1, cell label 2, linker 2, cell label 3, molecular label, and oligodT. For each oligonucleotide on the same bead, the sequences of the oligonucleotides are identical except for the molecular label. For oligonucleotides on different beads, the cell label 1, 2, and 3 combinations are different. Approximately 5-6 beads were added per well of the microwell array. The beads were allowed to settle into the wells and uncaptured beads were washed away in a PBS bath. A magnet was placed underneath the microwell array. Cells were lysed by the addition of cold lysis buffer. The array and magnet were placed on a cold aluminum block for 5 minutes. mRNA from the lysed cells were hybridized to the oligonucleotides coupled to the beads. The array was washed with excess lysis buffer to remove unbound mRNA. The beads were retrieved from the wells by placing a magnet on top of the microwell array. The retrieved beads were washed. cDNA synthesis was performed on the beads using Superscript III at 50 °C for 50 minutes on a rotor. Non-extended oligodT from the oligonucleotides on the beads were removed by ExoI treatment conducted at 37 °C for 30 minutes on a rotor. Gene-specific PCR amplification was conducted on the cDNA. The genes selected for the gene-specific PCR were cell-type specific and are shown in Table 6.

Table 6		
Number	Gene	Cell-type
1	HS_CD74	human
2	HS_CD79a	human
3	HS_IGJ	human
4	HS_TCL1A	human
5	HS_SEPT9	human
6	HS_CD27	human
7	HS_GAPDH	human
8	MM_B2M	mouse
9	MM_ACTM	mouse
10	MM_HPRT	mouse
11	MM_SHDA	mouse

[00520] The PCR amplified products were sequenced. Sequencing statistics are shown in Table 7.

Table 7		
	Low density	High density
Number of Ramos cells	15000	3750
Number of mouse cells	1500	2500
Total number of reads	2391780	4038217
>=8 match in constant 1	2162945	3651643
>=8 match in constant 2	1981835	3356493
>=8 match in constant 1 & 2	1981626	3355787
Perfect match in all 3 sub-barcodes	1645994	2790879
Perfect match in gene (40bp)	1083013	2171930
% useful reads	45%	54%
Total number of unique barcode combination	16695	36595
Number of unique barcode combinations > 30 molecule	80	281
Capture efficiency	0.03	0.04

[00521] Gene expression profiles for 100 unique barcode combinations with the most abundant molecules were determined for the high density and low density samples. The gene expression profiles were generated based on the sequencing results. FIG. 13A shows graphs of the gene expression profile for 35 of the 100 unique barcode combinations from the high density sample. For FIG. 13A, the number of unique molecules is on the y-axis and the gene reference number is on the x-axis (see Table 6 for genes corresponding to the gene reference number). FIG. 13A clearly depicts general gene expression patterns for the mouse and Ramos cells. FIG. 13B-C show scatter plots of results based on principal component analysis of gene expression profile of the high density sample and low density sample, respectively.

Component 1, which is plotted on the x-axis, separates the two cell types. Component 2, which is plotted on the y-axis, indicates variability in gene expression within the Ramos cell population. Each dot on the scatter plot represents one unique barcode combination, which is equivalent to one bead or one cell. Based on the principal component analysis of the high density sample, 144 beads corresponded to the mouse cells and 132 beads corresponded to Ramos cells. Based on the principal component analysis of the low density sample, 52 beads corresponded to the mouse cells and 27 beads corresponded to Ramos cells.

[00522] Once the cell types were determined, cross-talk between the beads was assessed by detecting the genes from Table 6 in the different cell types. FIG. 13D-E depict graphs of the read per barcode (bc) combination (y-axis) versus the unique barcode combination, sorted by the total number of molecules per bc combination (x-axis) for Ramos-like cells and mouse-

like cells from the high density sample, respectively. FIG. 13F-G depict graphs of the number of molecules per bc combination (y-axis) versus the unique barcode combination, sorted by the total number of molecules per bc combination (x-axis) for Ramos-like cells and mouse-like cells from the high density sample, respectively. FIG. 13H-I depict graphs of the read per barcode (bc) combination (y-axis) versus the unique barcode combination, sorted by the total number of molecules per bc combination (x-axis) for Ramos-like cells and mouse-like cells from the low density sample, respectively. FIG. 13J-K depict graphs of the number of molecules per bc combination (y-axis) versus the unique barcode combination, sorted by the total number of molecules per bc combination (x-axis) for Ramos-like cells and mouse-like cells from the low density sample, respectively. Table 8 shows the average fold coverage or read redundancy per unique molecule for the low and high density samples.

Table 8				
Gene	Low density		High density	
	Ramos-like cells	Mouse-like cells	Ramos-like cells	Mouse-like cells
HS_CD74	29.75	3.17	23.75	2.15
HS_CD79a	47.2	4.09	42.30	2.67
HS_IGJ	29.65	1.39	30.23	2.4
HS_TCL1A	45.74	2.26	39.00	4.13
HS_SEPT9	11.85	1.00	12.75	1.18
HS_CD27	37.99	1.00	32.12	1.10
HS_GAPDH	19.97	1.55	17.37	2.57
MM_B2M	1.21	31.98	3.05	31.48
MM_ACTM	1.05	29.08	1.90	28.38
MM_HPRT	1.02	39.96	1.03	43.65
MM_SHDA	1.00	39.60	1.02	29.60

[00523] The results in Table 8 show that average fold coverage per unique molecule was much higher for human genes than mouse genes in Ramos cells, and vice versa.

[00524] As a control, a mixture of mouse and human cells were lysed in a tube, converted to cDNA synthesis with the beads, and the cDNA was sequenced. FIG. 4XL shows a graphical representation of the sequencing results. As expected, a large number of unique barcode (bc) combinations was observed, and most beads only had one to two copies total.

[00525] These results demonstrate that there was minimal cross-talk between beads and that the cross-talk may be identified bioinformatically.

EXAMPLE 8. Single cell nucleic acid library production

[00526] The oligonucleotide conjugated supports disclosed herein may be used to produce single cell nucleic acid libraries. In this example, single cell nucleic acid libraries are produced by adding a cell sample to a surface (*e.g.*, grid) that has the oligonucleotide conjugated supports. An oligonucleotide conjugated support comprises a plurality of oligonucleotides conjugated to a bead. An oligonucleotide comprises (a) a cell label region comprising at least two distinct regions connected by a linker; and (b) a molecular label region. Two or more oligonucleotides on a bead comprise identical cell label regions. Two or more oligonucleotides on a bead comprise two or more different molecular label regions. Two or more oligonucleotides on two or more different beads comprise two or more different cell label regions. Thus, each cell associated with an oligonucleotide conjugated support has a different cell label region. The concentration of cells in the cell sample is sufficiently dilute to enable association of one or fewer cells to one oligonucleotide conjugated support on the surface. Cells are lysed using a lysis buffer. mRNAs from a cell are hybridized to the oligonucleotides of the oligonucleotide conjugated support. Thus, all mRNAs from a cell are labeled with oligonucleotides comprising identical cell label regions. Two or more mRNAs from a cell are labeled with two or more oligonucleotides comprising two or more different molecular label regions. A magnet is applied to the surface to purify the oligonucleotide conjugated solid supports from the surface. The oligonucleotide conjugated solid supports may be individually purified from the surface. The mRNAs hybridized to the oligonucleotides on the oligonucleotide conjugated solid support are reverse transcribed to produce labeled cDNA. The labeled cDNA comprise a reverse complement of the mRNA and a copy of the oligonucleotide that the mRNA was hybridized to. The labeled cDNA are amplified by PCR to produce labeled amplicons. The labeled cDNA and/or labeled amplicons may be removed from the bead by restriction enzyme digestion. A library of nucleic acids from the single cell is produced from the labeled amplicons.

[00527] Alternatively, the oligonucleotide conjugated solid supports are purified together. Reverse transcription of the mRNA may be performed on the combined oligonucleotide conjugated solid supports. Because mRNAs from different cells are labeled with oligonucleotides comprising different cell label regions, the cell label regions may be used to determine which cell the labeled cDNA or labeled amplicons originated from. Thus, a library of nucleic acids from a plurality cells may be produced, wherein the identity of the cell from which the labeled amplicon originated from may be determined by the cell label region.

[00528] Single cell nucleic acid libraries may also be produced by contacting the cells with an agent prior to lysing the cell. The agent may be an antigen, drug, cell, toxin, etc. Thus,

specialized single cell nucleic libraries may be produced. Analysis of the nucleic acid libraries may be used to generate single cell drug expression profiles. Signal transduction pathways on a single cell level may also be determined from these nucleic acid libraries. The nucleic acid libraries may also be used to determine the effects of antigens on specific cell types.

EXAMPLE 9. Single cell expression profiling

[00529] The oligonucleotide conjugated supports disclosed herein may be used to determine the expression profile of single cells. In this example, a cell sample comprising a mixture of cells is contacted with a plurality of antibodies. A subset of the cells is purified using flow cytometry. The subset of cells is added to a microwell array. A plurality of oligonucleotide conjugated supports is added to the microwell array. An oligonucleotide conjugated support comprises a plurality of oligonucleotides coupled to a nanoparticle. An oligonucleotide comprises (a) a cell label region comprising three distinct sequences connected by two predetermined sequences; and (b) a molecular label region. Two or more oligonucleotides on a nanoparticle comprise identical cell label regions. Two or more oligonucleotides on a nanoparticle comprise two or more different molecular label regions. Two or more oligonucleotides on two or more different nanoparticles comprise two or more different cell label regions. Thus, each cell associated with an oligonucleotide conjugated support has a different cell label region.

[00530] A magnet is applied to the microwell array and the cells that are not associated with an oligonucleotide conjugated support are washed away. A sponge comprising a lysis buffer is placed on top of the microwell array, thereby lysing the cells.

[00531] mRNAs from the lysed cells hybridize to the oligonucleotides on the bead. The mRNAs are reverse transcribed to produce labeled cDNA. The labeled cDNA comprise a reverse complement of the mRNA and a copy of the oligonucleotide that the mRNA was hybridized to. The labeled cDNA are amplified by PCR to produce labeled amplicons. The labeled amplicons are sequenced. Because each mRNA from a cell is labeled with the same cell label and mRNAs from different cells are labeled with different cell labels, the sequence information of the labeled amplicons is used to generate single cell expression profiles.

EXAMPLE 10: Immunophenotyping by single cell sequencing

[00532] A blood sample was collected from a subject and peripheral blood mononuclear cells (PMBCs) were isolated from the blood sample. PMBCs were cultured in RPMI1640 medium and placed in an incubator overnight. The PMBCs were washed multiple times in PBS to remove the serum. Approximately 7000 PMBCs were deposited onto a microwell

array with 32,400 wells. Thus, most wells on the microwell array contained no cells and some wells on the cell contained only 1 cell. Oligonucleotide-conjugated beads were applied to the microwell array. Each oligonucleotide-conjugated bead contained approximately 1 billion oligonucleotides attached to a bead. Each oligonucleotide attached to the bead contained a 5' amine, universal sequence, three-part cellular label (*e.g.*, three cell label sections connected by two linkers), molecular label, and oligodT. Each bead contained a unique three-part cellular label, which is a result of the unique combination of the three cell label sections. All of the oligonucleotides on a single bead contained the same three-part cellular label.

Oligonucleotides from different beads contained different three-part cellular labels. Each well contained 1 or fewer oligonucleotide-conjugated bead. A cell lysis reagent was applied to the microwell array, resulting in lysis of the cells. Polyadenylated molecules (*e.g.*, mRNA) from the cell hybridized to the oligodT sequence of the oligonucleotides from the oligonucleotide-conjugated beads. The polyadenylated molecules that were hybridized to the oligonucleotides from the oligonucleotide-conjugated bead were reverse transcribed with SuperScript II at 42°C at 90 minutes on a rotor. The oligonucleotide from the oligonucleotide-conjugated bead served as a primer for first strand cDNA synthesis. A SMART oligo was incorporated in the cDNA synthesis such that the superscript II may add the complement of the SMART oligo sequence to the 3' end of the cDNA when it reaches the end. The cDNA synthesis reaction produces a bead conjugated to unextended oligonucleotides (*e.g.*, oligonucleotides that were not attached to the polyadenylated molecule from the cell) and the extended oligonucleotides (*e.g.*, oligonucleotides that were attached to the polyadenylated molecule and comprise a polyadenylated molecule/cDNA hybrid).

[00533] The beads are combined and the oligonucleotides comprising the polyadenylated molecule/cDNA hybrid were amplified. Multiplex PCR was performed to amplify a panel of 98 genes (see Table 9) from the cDNA on the beads. Primers for the multiplexed PCR comprised a first gene specific primer that was designed to sit approximately 500 base pairs from the 3' end of the mRNA and a nested gene-specific primer that was designed to sit approximately 300 base pairs from the 3' end of the mRNA. Primers for the multiplex PCR were designed to require no significant complementarity in the last 6 bases of the primers in the panel. If complementarity was detected in the multiplex PCR primers, then the primers were manually replaced. The multiplex PCR reaction comprised the following steps: 1) 15 cycles of first gene specific PCR (KAPA multiplex mix, 50nM of each primer –first gene specific primer and universal primer that is complementary to the universal sequence of the oligonucleotide-conjugated bead), Ampure clean up (0.7x bead to template ratio), 15 cycles

of nested gene specific PCR (KAPA multiplex mix, 50nM of each primer –nested gene-specific primer and universal primer that is complementary to the universal sequence of the oligonucleotide-conjugated bead), Ampure clean up (0.7x bead to template ratio), 8 cycles of final PCR to add full length Illumina adaptor (KAPA HiFi ReadyMix), and Ampure clean up (1x bead to template ratio).

Table 9. Gene Panel							
Cell type	Gene	Cell type	Gene	Cell type	Gene	Cell type	Gene
B cell	PAX5	monocytes	CD14	naïve	CD62L (SELL)	Th17	IL17A
	CD19	classical monocytes	S100A12	Naïve Th	CD45RA		IL17F
	CD20		CCR2		THPOK/ZBTB7B		IL21
	BCMA/TNFRSF17		SELL/CD62L (L-selectin)	Naïve Tc	RUNX3		IL22
	BAFF	nonclassical monocytes	CD16/FCGR3B	memory	CD45RO/PTPRC		CCL20
	TCL1A		CX3CR1		CD44		IL23R
	TACI		ITGAL	Central Memory CD8+/CD4+	CCR7		RORa/RORA
B naïve	IGHD	conventional myeloid DC	CD1b	Effector Memory CD8+/CD4+ Th1	TXK	Follicular T helper	RORgamat/RORC
	IGHM		FOXP1		MBD2		OX40L/TNFSF4/CD252
B memory	CD27		CD209/DC-SIGN		BCL6		CXCR5
	CD38	plasmacytoid dendritic cell (rare)	CD1e	Activated T	BLIMP1	Activated T and B	SLAM/SLAMF1
	CD24		CCL17		CXCR3		ICOS
	AICDA		DTNA		IFNGR1		SAP/SH2D1A
	CD95	rare myeloid dendritic cell (0.02%)	CLEC4C/CD303		IL12RB2		CD69
B transitional	CD10		CD141/TM		IFN gamma		CD30
B reg	IL10	NKT		Th2	IL33R/IL1RL1	Toll-like receptors (innate)	TLR1
	RASD1		PLZF/ZBTB16		IL4R		TLR2
	AMPD1		SLAMF1		CCR4		TLR3
NK	SDC1 (CD138)	T cell	CD3 (CD3D)	Treg	CRTH2/PTGDR2		TLR4
	OSBPL5		CD3 (CD3E)		IL4		TLR5
	CD56/NCAM 1		CD8 (CD8A)		IL5		TLR6
	IGFBP7	Cytotoxic T	CD8 (CD8B)		CD25		TLR7
	KIR2DS5		PRF1 (perforin)		FOXP1		TLR8
	KIR2DS2		EOMES		TGFbeta		TLR9
	RAB4B	Helper T	CD4		IL10		TLR10

[00534] The amplified products were sequenced. The sequence reads with 150bp were aligned to entire mRNA sequences of the 98 genes listed in Table 9 using Bowtie2. The results of the sequence alignment (see Table 10) demonstrate that the multiplex PCR reaction resulted in highly specific products. FIG. 14 shows a graph depicting the genes on the X-axis and the log10 of the number of reads. 16 genes of the 98 genes were not present. Absence of these genes may be due to the fact that some of the genes target rare cells that may not be

present in this blood sample. Overall, approximately, 84% of the genes from the 98 gene panel were detected.

Table 10	
total	6357075
aligned 0 times	703616
aligned exactly 1 time	5584201
aligned > 1 time	69258
% aligned exactly once	88%

[00535] Table 11 shows the results of the overall sequencing statistics. For Read1, the total read1 match criteria required a perfect match to the three-part cellular label (*e.g.*, cell barcode) and at most 1 mismatch to the linkers.

Table 11	
total num read	6357075
total read1 match criteria	4384245
read2 also align	3943667
% read2 align	89.95%
number of unique cell bc	31129
read count per unique bc > 100	3228
read count per unique bc > 50	3721
% useful reads	62.04%

[00536] FIG. 15A shows a graph of the distribution of genes detected per three-part cell label (*e.g.*, cell barcode). FIG. 15B shows a graph of the distribution of unique molecules detected per bead (expressing the gene panel).

[00537] Cell clustering analysis was performed to determine whether the sequencing results could be used to analyze cell populations based on the cell barcode. SPADE (a minimum spanning tree algorithm developed by the Nolan lab for CyTOF data) was used to cluster cells based on the presence/absence of 17 genes. For a gene to be considered present, the average sequencing redundancy for the gene has to be greater than 5 fold. After sequence filtering, there were approximately 500 unique cell barcodes (*e.g.*, cell labels) associated with greater than 20 unique molecules. Each unique cell barcode corresponds to a single cell. Based on the genes that were associated with a unique cell barcode, cells were clustered into cell types. Table 12 shows a list of genes that may be used to definitively identify a cell type. Thus, cell barcodes that are associated with CD20, IGHM, TCL1A and CD24 were designated as B-cells, whereas cell barcodes that are associated with CD8A, CD3D, CD3E,

CD4 and CD62L were designated as T-cells. The remaining genes from Table 9 were mapped to the cell clusters. FIG. 16 depicts the cell clusters based on the genes associated with a cell barcode. The size of the cluster is proportionate to the number of cells that were assigned to the cluster. The results shown in FIG. 16 demonstrate that the combination of cell and molecular barcoding may be used to uniquely label copies of molecules from a single cell, which may enable immunophenotyping by single cell sequencing. In addition to clustering PMBCs into the major cell types based on the genes listed in Table 12, the 98 gene panel may also be used to identify clusters of sub-types of the major cell types. Table 13 shows the frequency of each major cell type detected by single cell sequencing. As shown in Table 13, with the exception of CD8⁺ T cells, the percentage of each cell type corresponded to the normal cell percentage range. A slightly higher percentage of CD8⁺ T cells was observed in the PMBC sample. Using the cell clusters based on FIG. 16, expression profiles of additional genes from the 98 gene panel were used to further analyze the cell clusters.

Table 12	
Major cell types	Genes
B cells	CD20, IGHM, TCL1A, CD24
T cells	CD8A, CD3D, CD3E, CD4, CD62L
NKT cells	ZBTB16
Dendritic cells	CD209
Natural Killer cells	KIR2DS5, KIR2DS2, CD16
Monocytes	CD16, CD14, CCR2, S100A12, CD62L

Table 13			
Cell type	# cells	percentage	normal range
monocytes	67	13.3%	10-30%
NK	85	16.9%	up to 15%
B	47	9.3%	up to 15%
CD8	210	41.7%	5-30%
CD4	94	18.7%	25-60%
total assigned cluster	503	100.0%	

[00538] FIG. 17A-D show the analysis of monocyte specific markers. FIG. 17E shows the cell cluster depicted in FIG. 16. FIG. 17A shows the cell expression profile for CD14, which is a monocyte specific marker. “Hot colors” (e.g., red) represent high gene expression and “cool colors” (e.g., blue) represent low gene expression. As shown in FIG. 17A, CD14 is

highly expressed in the monocyte population and had low to no expression in the other cell types. The cell expression profile for CD16 which is known to be present in both monocytes and NK is shown in FIG. 17B. As shown in FIG. 17B, the monocyte and NK cell clusters had high expression of CD16, whereas the other cell types had low to no expression. CCR2 and S100A12 are known to be highly expressed in monocytes. The CCR2 and S100A12 monocyte-specific expression was also demonstrated in the cell expression profiles shown in FIG. 17C and D, respectively. However, the expression of CCR2 and S100A12 separated into two branches of monocyte cells. The other cell types exhibited low to no expression of CCR2 and S100A12.

[00539] FIG. 18A-B show the analysis of the T cell specific markers. FIG. 18C shows the cell cluster depicted in FIG. 16. FIG. 18A shows the cell expression profile for CD3D which is a chain of the CD3 molecule. CD3 is a pan T cell marker. FIG. 18A shows that CD3D is highly expressed in two branches of CD8⁺ T cells and moderately expressed in a third branch of CD8⁺ T cells. However, CD3D is not highly expressed in CD4⁺ T cells. Also, the other cell types have low to no expression of CD3D. FIG. 18B shows the cell expression profile for CD3E which is a chain of the CD3 molecule. FIG. 18B shows that CD3D is highly expressed in CD4⁺ T cells. Different branches of CD8⁺ T cells exhibit high to moderate expression of CD3D. Little to no expression of CD3D is observed in the other cell types.

[00540] FIG. 19A-B show the analysis of the CD8⁺ T cell specific markers. FIG. 19C shows the cell cluster depicted in FIG. 16. FIG. 19A shows the cell expression profile for CD8A which is a chain of the CD8 molecule. As shown in FIG. 19A, different branches of CD8⁺ T cells have various levels of CD8A expression, with some branches having high expression, other branches having moderate expression and one branch exhibiting low to no expression of CD8A. High CD8A expression was observed in a branch of the CD16⁺ NK cells. It has been reported in the literature that up to 80% of NK cells express CD8. Little to no CD8A expression was observed in the other cell types. FIG. 19B shows the cell expression profile for CD8B which is a chain of the CD8 model. As shown in FIG. 19B, different branches of CD8⁺ T cells have various levels of CD8B expression, with one branch having high expression, some branches having moderate expression and two branches exhibiting low to no expression of CD8B. High CD8B expression was also observed in a branch of the CD16⁺ NK cells. Little to no CD8B expression was observed in the other cell types.

[00541] FIG. 20A shows the analysis of CD4⁺ T cell specific markers. FIG. 20B shows the cell cluster depicted in FIG. 16. FIG. 20A shows the expression profile for CD4. Moderate expression of CD4 was observed in a subset of cells in the CD4⁺ T cell cluster and high

expression of CD4 was observed in a branch of the monocyte cluster. It has previously been documented in the literature that monocytes also express CD4. Moderate to low expression of CD4 was observed in a branch of CD8+ T-cells and in NK cells. Low to no expression of CD4 was observed in the other cell types.

[00542] FIG. 21A-D show the analysis of Natural Killer (NK) cell specific markers. FIG. 20E shows the cell cluster depicted in FIG. 16. FIG. 20A shows the expression profile for KIR2DS2. All of the cell types exhibited little to no KIR2DS2 expression. FIG. 20B shows the expression profile for KIR2DS5. Killer immunoglobulin receptors (KIRs) are known to be expressed in NK cells and a subset of T cells. High expression of KIR2DS5 was observed in 2 branches of NK cells and moderate to low expression of KIR2DS5 was observed in one branch of NK cells. Moderate to high expression of KIR2DS5 was observed in 2 branches of CD8+ T cells. Low to no expression of KIR2DS5 was observed in all other cell types. OSBPL5 and IGFBP7 are known to be highly expressed in NK cells. FIG. 20C shows the expression profile for OSBPL5. OSBPL5 was highly expressed in one branch of NK cells. Moderate to low expression of OSBPL5 was observed in a branch of B cells. Low to no expression of OSBPL5 was observed in all other cell types. FIG. 20D shows the expression profile for IGFBP7. High expression of IGFBP7 was observed in two branches of NK cells and one branch of monocytes. Moderate expression of IGFBP7 was observed in one branch of B cells. Low to no expression of IGFBP7 was observed in all other cell types.

[00543] FIG. 22A-E show the analysis of B cell specific markers. FIG. 22F shows the cell cluster depicted in FIG. 16. FIG. 22A shows the expression profile for IGHM CH4. IGHM CH4 was highly expressed in one branch of B cells and moderately expressed in the second branch of B cells. Low to no expression of IGHM CH4 was observed in all other cell types. FIG. 22B shows the expression profile for PAX5. PAX5 was highly expressed in one branch of B cells. Low to no expression of PAX5 was observed in all other cell types. FIG. 22C shows the expression profile for CD20. CD20 was highly expressed in one branch of B cells. Low to no expression of CD20 was observed in all other cell types. FIG. 22D shows the expression profile for TCL1A. Low to no expression of TCL1A was observed in all other cell types. FIG. 22E shows the expression profile for IGHD CH2. IGHD CH2 was highly expressed in one branch of B cells. Low to no expression of IGHD CH2 was observed in all other cell types.

[00544] FIG. 23A-F show the analysis of Toll-like receptors. Toll-like receptors are mainly expressed by monocytes and some B cells. FIG. 23G shows the cell cluster depicted in FIG. 16. FIG. 23A shows the expression profile for TLR1. One branch of monocytes exhibited

high expression of TLR1 and two branches of monocytes exhibited moderate expression of TLR1. Low to no expression of TLR1 was observed in all other cell types. FIG. 23B shows the expression profile for TLR4. One branch of monocytes exhibited high expression of TLR4. Moderate TLR4 expression was observed in two branches of monocytes and one branch of NK cells. Low to no expression of TLR4 was observed in all other cell types. FIG. 23C shows the expression profile for TLR7. High expression of TLR7 was observed in one branch of monocytes and moderate expression of TLR7 was observed in one branch of NK cells. Low to no expression of TLR7 was observed in all other cell types. FIG. 23D shows the expression profile for TLR2. High expression of TLR2 was observed in one branch of B cells. Low to no expression of TLR2 was observed in all other cell types. FIG. 23E shows the expression profile for TLR3. High expression of TLR3 was observed in one branch of B cells. Low to no expression of TLR3 was observed in all other cell types. FIG. 23F shows the expression profile for TLR8. High expression of TLR8 was observed in three branches of monocytes. Moderate to low expression of TLR8 was observed in two branches of monocytes and one branch of NK cells. Low to no expression of TLR8 was observed in all other cell types.

[00545] These results demonstrate that massively parallel single cell sequencing may successfully identify major cell types in PMBCs. The sequencing results also determined that some cell markers that are used in FACs for identifying cell types do not have high mRNA expression (*e.g.*, CD56 for NK cells, CD19 for B cells). In addition, many of the genes in the gene panel were expressed across multiple cell types. These expression profiles may be used to subtype cells within a major cell type (*e.g.*, activated cell versus resting cell, etc.).

EXAMPLE 11: Identifying rare cells in a population

[00546] In this experiment, massively parallel single cell sequencing is used to identify cancer cells from a mixture of cancer and non-cancer cells. Ramos (Burkitt lymphoma) cells were spiked into a population of CD19⁺ B cells that were isolated from a healthy individual. The concentration of the Ramos cells in the mixed population was about 4-5%.

Approximately 7000 normal B cells and 300 Ramos cells were deposited on a microwell array with 25,200 wells. Thus, most wells on the microwell array contained no cells and some wells on the cell contained only 1 cell. Oligonucleotide-conjugated beads were applied to the microwell array. Each oligonucleotide-conjugated bead contained approximately 1 billion oligonucleotides attached to a bead. Each oligonucleotide attached to the bead contained a 5' amine, universal sequence, three-part cellular label (*e.g.*, three cell label sections connected by two linkers), molecular label, and oligodT. Each bead contained a unique three-part

cellular label, which is a result of the unique combination of the three cell label sections. All of the oligonucleotides on a single bead contained the same three-part cellular label. Oligonucleotides from different beads contained different three-part cellular labels. Each well contained 1 or fewer oligonucleotide-conjugated bead. A cell lysis reagent was applied to the microwell array, resulting in lysis of the cells. Polyadenylated molecules (*e.g.*, mRNA) from the cell hybridized to the oligodT sequence of the oligonucleotides from the oligonucleotide-conjugated beads. The polyadenylated molecules that were hybridized to the oligonucleotides from the oligonucleotide-conjugated bead were reverse transcribed with SuperScript II at 42°C at 90 minutes on a rotor. The oligonucleotide from the oligonucleotide-conjugated bead served as a primer for first strand cDNA synthesis. A SMART oligo was incorporated in the cDNA synthesis such that the superscript II may add the complement of the SMART oligo sequence to the 3' end of the cDNA when it reaches the end. The cDNA synthesis reaction produces a bead conjugated to unextended oligonucleotides (*e.g.*, oligonucleotides that were not attached to the polyadenylated molecule from the cell) and the extended oligonucleotides (*e.g.*, oligonucleotides that were attached to the polyadenylated molecule and comprise a polyadenylated molecule/cDNA hybrid).

[00547] The beads are combined and the oligonucleotides comprising the polyadenylated molecule/cDNA hybrid were amplified. Multiplex PCR was performed to amplify a panel of 111 genes from the cDNA on the beads. The 111 genes represent markers for different subsets of B cells. Primers for the multiplexed PCR comprised a first gene specific primer that was designed to sit approximately 500 base pairs from the 3' end of the mRNA and a nested gene-specific primer that was designed to sit approximately 300 base pairs from the 3' end of the mRNA. Primers for the multiplex PCR were designed to require no significant complementarity in the last 6 bases of the primers in the panel. If complementarity was detected in the multiplex PCR primers, then the primers were manually replaced. The multiplex PCR reaction comprised the following steps: 1) 15 cycles of first gene specific PCR (KAPA multiplex mix, 50nM of each primer –first gene specific primer and universal primer that is complementary to the universal sequence of the oligonucleotide-conjugated bead), Ampure clean up (0.7x bead to template ratio), 15 cycles of nested gene specific PCR (KAPA multiplex mix, 50nM of each primer –nested gene-specific primer and universal primer that is complementary to the universal sequence of the oligonucleotide-conjugated bead), Ampure clean up (0.7x bead to template ratio), 8 cycles of final PCR to add full length Illumina adaptor (KAPA HiFi ReadyMix), and Ampure clean up (1x bead to template ratio).

[00548] The amplified products were sequenced. The sequence reads comprising 150bp were aligned to entire mRNA sequences of the 111 genes (Table 17) using Bowtie2. The results of the sequence alignment (see Table 14) demonstrate that the multiplex PCR reaction resulted in highly specific products. FIG. 24 depicts a graph of the genes versus the log10 of the number of reads. 24 of the 111 genes were not present. At least two of the genes, RAG1 and RAG2 which are involved in VDJ recombination and should be present only in pre-B cells, should not be present. A few of the absent genes are specific for plasma cells, which are very rarely preserved in frozen cells.

Table 17.							
CD19	AURKB	FOXP1	CCND3	TLR1	FOXP3	CXCL12	GNAI2
CD27	CD81	MCL1	IL12A	TLR2	LAG3	CCL3	RGS1
CD138	CD80	IFNB1	IFNG	TLR3	CD73	CCL14	CD5
CD38	CD23a	BLNK	TNFA	TLR4	CD70	CCL20	CD22
CD24	CD44	CD40LG	IL2	TLR5	CCR7	CCL18	PIK3CD
CD10	LEF1	IGBP1	IL4	TLR6	CD45RA	TCL1A	DOCK8
CD95	CXCR5	IRF4	IL6	TLR7	PDCD1	TAC1	CD11b
CD21	PRKCB	CD79a	BAFF	TLR8	MYC	AICDA	FCGR2B
CXCR3	PRKCD	LTA	IGHE	TLR9	CD25	FCRL4	CD72
CD40	CD20	HDAC5	IGHD	TLR10	FCAMR	BCL2	BCL11B
CD69	CD30	RAG1	IGHM	GAPDH	CCND2	FASLG	CD86
CD1c	CD30L	RAG2	IGHA	CD9	MKI67	BCL6	TBX21
IL10	BAFFR	CD1d	IGHG1	CD11c	IL21R	IGHG2	PRDM1
IL4R	CMRF-35H	TGFB1	IGHG4	IL6R	HLA-DRA	IGHG3	

Table 14	
total	5711013
aligned 0 times	504775
aligned exactly 1 time	5203308
aligned > 1 time	2930
% aligned exactly once	91.6%

[00549] Table 15 shows the results of the overall sequencing statistics. For Read1, the total read1 match criteria required a perfect match to the three-part cellular label (*e.g.*, cell barcode) and at most 1 mismatch to the linkers.

Table 15	
total num read	5711013
total read1 match criteria	3795915
read2 also align	3495392
% read2 align	92%
number of unique cell bc	40764
read count per unique bc > 100	3313
read count per unique bc > 50	4154
% useful reads	61%

[00550] FIG. 25A-D shows graphs of the molecular barcode versus the number of reads or log10 of the number of reads for two genes. FIG. 25A shows a graph of the molecular barcode (sorted by abundance) versus the number of reads for CD79. FIG. 25B shows a graph of the molecular barcode (sorted by abundance) versus the log10 of the number of reads for CD79. FIG. 25C shows a graph of the molecular barcode (sorted by abundance) versus the number of reads for GAPDH. FIG. 25D shows a graph of the molecular barcode (sorted by abundance) versus the log10 of the number of reads for GAPDH.

[00551] 856 cells were retained for analysis. FIG. 26A shows a graph of the number of genes in the panel expressed per cell barcode versus the number of unique cell barcodes/single cell. FIG. 26B shows a histogram of the number of unique molecules detected per bead versus frequency of the number of cells per unique cell barcode carrying a given number of molecules. A small subset of cells showed distinctly higher number of mRNA molecules and number of genes expressed from the 111 gene panel (see circled sections in FIG. 26A-B). FIG. 26C shows a histogram of the number of unique GAPDH molecules detected per bead versus frequency of the number of cells/unique cell barcode carrying a given number of molecules.

[00552] Principal component analysis (PCA) was used to generate a scatterplot of cells. FIG. 27 shows a scatterplot of the 856 cells. PCA identified the small subset of cells with a different gene expression pattern than the majority of cells. The subset of cells contained 18 cells, which is approximately 2% of all of the cells analyzed. This percentage is similar to the percentage of Ramos cells that was spiked into the population.

[00553] Ramos cells are derived from follicular B cells and strongly express B cell differentiation markers CD20, CD22, CD19, CD10 and BCL6. Ramos cells also express IgM and overexpress c-myc. FIG. 28 shows a heat map of expression of the top 100 (in terms of the total number of molecules detected). The subset of cells (18 cells) that express much

higher levels of mRNA also strongly express genes that are known markers for Ramos cells (*e.g.*, CD10, Bcl-6, CD22, C-my, and IgM).

[00554] These results demonstrate that massively parallel single cell sequencing successfully identified small subsets (as low as 2%) of abnormal cell types in a cell suspension. Massively parallel single cell sequencing may be used in cancer diagnostics (*e.g.*, biopsy/circulating tumor cells). Since cancer cells are larger in size and carry more mRNA, they may be easily differentiated from normal cells.

EXAMPLE 12: Massively parallel single cell whole genome and multiplex amplification of gDNA targets using RESOLVE

[00555] FIG. 29 shows a workflow for this example. As shown in FIG. 29, a cell suspension is applied to a microwell array (2901). The number of cells in the cell suspension is less than the number of wells in the microwell array, such that application of the cell suspension to the microwell array results in a well in the microwell array containing 1 or fewer cells.

Oligonucleotide-conjugated beads (2905) are applied to the microwell array. An oligonucleotide-conjugated bead (2905) contains a bead (2910) attached to an oligonucleotide comprising a 5' amine (2915), universal primer sequence (2920), cell label (2925), molecular label (2930) and randomer (2935). The oligonucleotide-conjugated bead contains approximately 1 billion oligonucleotides. An oligonucleotide contains a 5' amine, universal primer sequence, cell label, molecular label, and randomer. Each oligonucleotide on a single bead contains the same cell label. However, two or more oligonucleotides on a single bead may contain two or more different molecular labels. A bead may contain multiple copies of oligonucleotides containing the same molecular label.

[00556] After the oligonucleotide-conjugated beads are added to the microwell array, a cell lysis buffer is applied to the array surface. As shown in FIG. 29, the genomic DNA (2945) from the cell hybridizes to the randomer sequence (2935) of the oligonucleotide-conjugated beads (2940). A neutralization buffer is added to the array surface. A DNA polymerase (*e.g.*, Phi29) and dNTPs are added to the array surface. The randomer sequence (2935) acts as a primer for amplification of the genomic DNA, thereby produce a gDNA-conjugated bead (2555). The gDNA-conjugated bead (2955) contains an oligonucleotide comprising a 5' amine (2915), universal primer sequence (2920), cell label (2925), molecular label (2925), randomer (2935) and copy of the genomic DNA (2955). The original genomic DNA (2945) is hybridized to the randomer (2935) and the copy of the genomic DNA (2955). For a single bead, there are multiple different genomic DNA molecules attached to the oligonucleotides.

[00557] As shown in FIG. 29, the gDNA-conjugated beads (2950) from the wells are combined into an eppendorf tube (2960). The genomic DNA on the gDNA MDA mix containing randomers, dNTPs and a DNA polymerase (*e.g.*, Phi29) is added to the eppendorf tube containing the combined gDNA-conjugated beads. The labeled genomic DNA is further amplified to yield labeled amplicons (2965) in solution. A labeled amplicon (2965) comprises a universal primer sequence (2920), cell label (2925), molecular label (2930), randomer (2935), and copy of the genomic DNA (2955). The labeled amplicons are sheared to smaller pieces of approximately 1kb or less. Alternatively, the labeled amplicons may be fragmented by to Tagmentation (Nextera). Shearing or fragmenting the labeled amplicons results in labeled-fragments (2980) and unlabeled fragments (2985). The labeled fragment (2980) contains the universal primer sequence (2920), cell label (2925), molecular label (2930), randomer (2935), and fragment of the copy of the genomic DNA (2955). Adaptors (2970, 2975) are added to the fragments. The universal primer sequence may be used to select for labeled fragments (2980) via hybridization pulldown or PCR using the universal primer sequence and a primer against one of the adaptors (2970, 2975).

[00558] The labeled fragments may be sequenced. Sequence reads comprising a sequence of the cell label, molecular label and genomic fragment may be used to identify cell populations from the cell suspension. Principal component analysis may be used to generate scatterplots of the cells based on known cell markers. Alternatively, or additionally, SPADE may be used to produce cell cluster plots. A computer software program may be used to generate a list comprising a cell label and the molecular labels and genomic fragments associated with the cell label.

EXAMPLE 13: Massively parallel sequencing to identify cells in a heterogeneous population

[00559] The experimental workflow for this example is shown in FIG. 30. As shown in FIG. 30, a mixed population of cells was stochastically dispersed onto a microwell array. In this example, the mixed population of cells comprises a mixture of Ramos cells and K562 cells. The cell suspension comprises a low concentration of cells such that each microwell in the array contains 1 or fewer cells. After the cells were applied to the microwell array, a plurality of oligonucleotide conjugated beads was stochastically dispersed onto the microwell array. The oligonucleotide bead contains a plurality of oligonucleotides comprising a 5' amine, universal primer sequence, cell label, molecular label, and oligodT. The cell labels of the plurality of oligonucleotides from a single bead are identical. A single bead may comprise multiple oligonucleotides comprising the same molecular label. In addition, a single bead may comprise multiple oligonucleotides comprising different molecular labels. A cell label of

an oligonucleotide conjugated to a first bead is different from a cell label of an oligonucleotide conjugated to a second bead. Thus, the cell label may be used to differentiate two or more oligonucleotide conjugated beads. The cells were lysed and the RNA molecules from a single cell were attached to the oligonucleotide conjugated beads in the same well. FIG. 30 shows the attachment of the polyA sequence of a RNA to the oligodT sequence of the oligonucleotide. After attachment of the RNA molecules from the individual cells to the oligonucleotide conjugated beads in the same well, the beads were combined into a single sample. A cDNA synthesis reaction was carried out on the beads in the single sample. FIG. 30 shows the product of the cDNA synthesis comprises a bead attached to an oligonucleotide, the oligonucleotide comprising the 5' amine, universal primer sequence, cell label, molecular label, oligodT and a copy of the RNA molecule. For simplicity, only one oligonucleotide is depicted in FIG. 30, however, in this example, each oligonucleotide conjugated bead comprises approximately 1 billion oligonucleotides. As shown in FIG. 30, multiplexed PCR was performed with the beads in the single sample using a universal primer that hybridized to the universal primer sequence and a gene-specific primer that hybridized to the copy of the RNA molecule. The gene-specific primers were designed to bind to Ramos-specific genes or K562-specific genes from the gene panel shown in Table 16. As a control, a GAPDH gene-specific primer was also used in the multiplexed PCR reaction. Lastly, next-generation sequencing was used to sequence the amplified products. The sequencing reads included information pertaining to the cell label, molecular label and the gene. Using principal component analysis, a scatter plot of the cells was constructed based on the sequencing information pertaining to the cell label, molecular label and the gene. Analogous to how FACs is used to sort cells and scatter plots based on the surface markers is used to group cells, the cell label is used to identify genes from a single cell and the molecular label is used to determine the quantity of the genes. This combined information is then used to relate the gene expression profile individual cells. As shown in FIG. 31A, massively parallel single cell sequencing with cell and molecular labels was able to successfully identify the two cell populations (K562 and Ramos cells) in the mixed cell population.

Table 16			
Gene	Cell	Gene	Cell
CD74	Ramos specific	CD41	K562 specific
CD79a	Ramos specific	GYPA	K562 specific
IGJ	Ramos specific	GATA2	K562 specific
TCL1A	Ramos specific	GATA1	K562 specific
SEPT9	Ramos specific	HBG1	K562 specific
CD27	Ramos specific	GAPDH	Common

EXAMPLE 14: Massively parallel single cell sequencing with principal component analysis

[00560] In this example, mRNA molecules from individual cells were stochastically labeled with oligonucleotide conjugated beads in parallel. PBMCs were isolated from blood and frozen at 80°C in RPMI1640 plus FBS and DMSO. The PMBCs were thawed and washed three times with PBS. A PBMC sample comprising a mixture of cell types (4000 total cells) was stochastically applied to an agarose microwell array. The agarose microwell array contained 37,500 cells. A mixture of 150,000 oligonucleotide conjugated beads was stochastically applied to the microwell array via a PDMS gasket that surrounded the microwell array. The oligonucleotide conjugated bead is depicted in FIG. 1. For simplicity, only one oligonucleotide is shown to be attached to the bead, however, the oligonucleotide conjugated beads contained approximately 1 billion oligonucleotides.

[00561] Cells were lysed by placing the microwell array on a cold block for 10 minutes and by applying lysis buffer to the array surface. Once the cells in the wells were lysed, the mRNA molecules from the single cells were attached to the oligonucleotide conjugated bead via the oligodT sequence. A magnet was applied to the array and the array was washed twice with wash buffer.

[00562] The beads with the attached mRNA molecules were combined into an eppendorf tube. The mRNA molecules attached to the beads were reverse transcribed to produce cDNA. The following cDNA synthesis mixture was prepared as follows:

Component	Volume (uL)
Water	8
dNTP (10mM)	2
5x first strand buffer	4
MgCl ₂	2.4
SuperRase In	1
SMART oligo (50uM)	0.4
0.1M DTT	1
100x BSA	0.2
SSII	1
total	20

[00563] The cDNA synthesis mixture was added to the eppendorf tube containing the beads with the attached mRNA molecules. The eppendorf tube was incubated at 40°C for 90 minutes on a rotor. The cDNA synthesis reaction occurred on the beads. After 90 minutes, a magnet was applied to the tube and the cDNA mix was removed and replaced with the following ExoI reaction mixture:

Component	Volume (uL)
ExoI buffer	2
water	17
ExoI	1

[00564] The tubes were incubated at 37°C for 30 minutes on a rotor. The tubes were then transferred to a thermal cycler for 15 minutes at 80°C. After incubating the tube at 80°C for 15 minutes, 70 microliters of TE+Tween20 was added to the tube. A magnet was applied to the tube and the buffer was removed. The beads were then resuspended in 50 microliters TE+Tween20.

[00565] The cDNA attached to the beads were amplified by real-time PCR using the following amplification mixture:

Component	Volume (uL)
2x iTaq mix	10
GAPDH ILMN (10uM)	0.6
ILR2 (10uM)	0.6
bead	2
water	6.8
total	20

[00566] The labeled cDNA amplicons were sequenced to detect the cell label, molecular index, and gene. Sequencing reads were aligned to the cell label, then the gene, and lastly the molecular label. A cell label associated with 4 or more genes or associated with 10 or more unique transcript molecules, with each unique transcript molecule sequenced more than once,

was designated a cell. Principal component analysis with all of genes from Table 9 detected was used to identify the set of genes that had the greatest contribution to the variation in data. 632 single cells were used in the principal component analysis. Based on the sequencing results, 81 out of the 98 genes were detected.

[00567] FIG. 32 shows a principal component analysis plot for GAPDH expression. As shown in FIG. 32, two cell clusters were observed based on the location of the principal component space.

[00568] FIG. 33A-F shows the principal component analysis (PCA) for monocyte associated genes. FIG. 33A shows the PCA for CD16. FIG. 33B shows the PCA for CCRvarA. FIG. 33C shows the PCA for CD14. FIG. 33D shows the PCA for S100A12. FIG. 33E shows the PCA for CD209. FIG. 33F shows the PCA for IFNGR1.

[00569] FIG. 34A-B shows the principal component analysis (PCA) for pan-T cell markers (CD3). FIG. 34A shows the PCA for CD3D and FIG. 34B shows the PCA for CD3E.

[00570] FIG. 35A-E shows the principal component analysis (PCA) for CD8 T cell associated genes. FIG. 35A shows the PCA for CD8A. FIG. 35B shows the PCA for EOMES. FIG. 35C shows the PCA for CD8B. FIG. 35D shows the PCA for PRF1. FIG. 35E shows the PCA for RUNX3.

[00571] FIG. 36A-C shows the principal component analysis (PCA) for CD4 T cell associated genes. FIG. 36A shows the PCA for CD4. FIG. 36B shows the PCA for CCR7. FIG. 36C shows the PCA for CD62L.

[00572] FIG. 37A-F shows the principal component analysis (PCA) for B cell associated genes. FIG. 37A shows the PCA for CD20. FIG. 37B shows the PCA for IGHD. FIG. 37C shows the PCA for PAX5. FIG. 37D shows the PCA for TCL1A. FIG. 37E shows the PCA for IGHM. FIG. 37F shows the PCA for CD24.

[00573] FIG. 38A-C shows the principal component analysis (PCA) for Natural Killer cell associated genes. FIG. 38A shows the PCA for KIR2DS5. FIG. 38B shows the PCA for CD16. FIG. 38C shows the PCA for CD62L.

[00574] Based on the principal component analyses, monocytes and lymphocytes formed two distinct clusters on PC1. B, T, and NK cells formed another cluster that resided as a continuum in the cluster along PC2. FIG. 39 shows the PCA analysis of GAPDH expression with annotations for the cell types and cell subtypes. FIG. 40 depicts a heat map that shows the correlation in gene expression profile between cells. Along the diagonal starting with the left upper corner, the cells are monocytes, naive CD4 T cells, naive CD8 T cells, cytotoxic CD8 T cells, NK cells, and B cells. FIG. 41 shows another version of a heat map

demonstrating the correlation between gene expression and cell type. FIG. 42 shows a heat map demonstrating the correlation in gene expression profile between genes.

EXAMPLE 15: Uncovering cellular heterogeneity by digital gene expression cytometry

[00575] An approach for gene expression cytometry is presented combining next-generation sequencing with stochastic barcoding of single cells. Thousands of cells were deposited randomly onto an array of approximately 150,000 microwells. A library of beads bearing cell- and transcript-barcoding capture probes was added so that each cell is partitioned alongside a bead with a unique cell barcode. Following cell lysis, mRNAs were hybridized to beads, and were pooled for reverse transcription, amplification, and sequencing. The digital gene expression profile for each cell was reconstructed when barcoded transcripts were counted and assigned to the cell of origin. We applied the technology to dissect the human hematopoietic system into cell sub-populations, and to characterize the heterogeneous response of immune cells to *in vitro* stimulation. Furthermore, the high sensitivity of the method was demonstrated by the detection of rare cells, such as antigen-specific T cells, and tumor cells in a high background of normal cells.

Introduction

[00576] Understanding cellular diversity and function in a large collection of cells requires the measurement of specific genes or proteins expressed by individual cells. Flow cytometry is well established for measuring protein expression of single cells, yet mRNA expression measurements are typically conducted in bulk samples, obscuring individual cell contributions. While single cell mRNA expression measurements using microtiter plates or commercial microfluidic chips have recently been reported (1-5), these approaches are extremely low-throughput and difficult to scale. Because of these limitations, most studies to date are restricted in both the number of cells interrogated and the number of conditions explored.

[00577] Here, we have developed a highly scalable approach that enables routine, digital gene expression profiling of thousands of single cells across an arbitrary number of genes. Microscale engineering and combinatorial chemistry were used to label all mRNA molecules in a cell with a unique cellular barcode in a massively parallel manner. In addition, each transcript copy within a cell was tagged with a molecular barcode, allowing absolute digital gene expression measurements (6). Tagged mRNA molecules from all cells were pooled, amplified, and sequenced. The digital gene expression profile of each cell was reconstructed using the cell and molecular barcodes on each sequence. This highly scalable technology enables gene expression cytometry, which we term CytoSeq. We have applied the technique

to multiparameter genetic classification of the hematopoietic system and demonstrated its use for studying cellular heterogeneity and detecting rare cells in a population.

Results

CytoSeq

[00578] The procedure was outlined in **FIG. 43A**. A cell suspension was first loaded onto a microfabricated surface with up to 150,000 microwells. Each 30 micron diameter microwell has a volume of ~20 picoliters. The number of cells was adjusted so that only ~1 out of 10 or more wells receives a cell. The cells settled within the wells by gravity.

[00579] Magnetic beads were loaded onto the microwell array to saturation, such that a bead sat partially on top of, or adjacent to, each cell within a well. The dimension of the bead was chosen such that each microwell may hold only one bead. Each magnetic bead carried approximately one billion oligonucleotide templates with the structure outlined in **FIG. 43B**. Each oligonucleotide displayed a universal priming site, followed by a cell label, a molecular label, and a capture sequence of oligo(dT). All the oligonucleotides on each bead have the same cell label but contain a diversity of molecular labels. We have devised a combinatorial split-pool method to synthesize beads with a diversity of close to one million. The probability of having two single cells being tagged with the same cell label was low (on the order of 10^{-4}) because only ~10% of the wells were occupied by a single cell. Similarly, the diversity of the molecular labels on a single bead was on the order of 10^4 , and the likelihood of two transcript molecules of the same gene in the same cell being tagged with the same molecular label was also low.

[00580] Lysis buffer was applied onto the surface of the microwell array and diffuses into the microwells. The poly(dA) tailed mRNA molecules released from a cell hybridize to the oligo(dT) on the 3' end of the oligonucleotides on the bead. Because the cell was adjacent to the bead, under the high salt conditions of the lysis buffer and high local concentration of mRNA (tens of nanomolar), mRNA molecules were captured on the bead.

[00581] After lysis and hybridization, all beads were collected from the microwell array into a tube using a magnet. From this point forward, all reactions were carried out in a single tube. cDNA synthesis was performed on the beads using conventional protocols (Methods). The cDNA molecules derived from each cell were covalently attached to their corresponding bead, each tagged on the 5' end with a cell label and a molecular label. Nested multiplex PCRs were carried out to amplify genes of interest (**FIG. 55**). Because the mRNA from each cell had been copied onto a bead as cDNA, the beads may be repeatedly amplified and analyzed, for example, for a different set of genes.

[00582] Sequencing of the amplicons revealed the cell label, the molecular label, and the gene identity (**FIG. 55**). Computational analysis grouped the reads based on the cell label, and collapsed the reads with the same molecular label and gene sequence into a single entry to suppress any amplification bias. The use of molecular label enabled us to measure the absolute number of molecules per gene per cell, and therefore allowed the direct comparison of cellular expression level across biological samples that may have undergone different depths of sequencing.

Identification of distinct cell types in controlled cell mixtures

[00583] In order to measure the ability of the method to separate two cell types, a ~1:1 mixture of K562 and Ramos cells was loaded onto the microwell array with 10,000 wells. Approximately 6000 cells were used to capture ~1000 cells. A panel of 12 genes was selected and amplified from the beads. The panel consists of 5 genes specific for K562 (myelogenous leukemia) cells, 6 genes specific for Ramos (follicular lymphoma) cells, and the housekeeping gene GAPDH (**Table 18**). With approximately 1000 cells captured on a 10,000-well array each with a single bead, only 10% of the beads should carry mRNA and one should in theory observe only a maximum of 1000 unique cell labels in the sequencing data. Indeed, we found 768 cell labels that were associated with significant number of reads after data filtering (see Methods for filtering criteria). As a comparison, we carried out bulk cell lysis and mRNA capture in a microcentrifuge tube with similar number of cells and beads, and observed a large number of cell labels with mostly only one read associated with each cell label. This demonstrates that the microwell array was effective in confining hybridization of mRNA from a single cell to the bead in the same well.

[00584] The gene expression profile of each of the 768 single cells was clustered using principal component analysis (PCA) (**FIG. 31A**). The first principal component (PC) clearly separated the single cells into two major clusters based on the cell type. The genes that contributed to the positive side of the first principal component were those that are specific to Ramos, while the genes that contributed to the negative side of the same principal component were those that are specific to K562. This successful clustering of cells into groups based on their specific expression showed that inter-well contamination, if any, was negligible. The second principal component highlighted the high degree of variability in fetal hemoglobin (HBG1) within the K562 cells, which had been observed previously (7).

Table 18		
Gene	Outer Primer	Nested Primer with Common 5' Flanking Sequence
CD41	CCCCTGGAAGAAGATGATGA	CAGACGTGTGCTCTTCCGATCTTTCTCCAACAAGTTGCCTCC
GYPD	GAGGAAATGAAGCCAAACA CA	CAGACGTGTGCTCTTCCGATCTAATCGTGACCTTAAAGGCCC
GATA1	TTAGCCACCTCATGCCTTTC	CAGACGTGTGCTCTTCCGATCTCTACTGTGGTGGCTCCGCT
GATA2	GGAGGAGGATTGTGCTGATG	CAGACGTGTGCTCTTCCGATCTGTGTCCGCATAAGAAAAAGAA TC
HBG1	GCAAGAAGGTGCTGACTTCC	CAGACGTGTGCTCTTCCGATCTCTGCATGTGGATCCTGAGAA
CD27	CTGCAGTCCCATCCTCTTGT	CAGACGTGTGCTCTTCCGATCTGATGAGGTGGAGAGTGGGAA
IGJ	GGACATAACAGACTTGAA GCA	CAGACGTGTGCTCTTCCGATCTCAATCCATTTTGTAACTGAACC TT
TCL1A	AAGCCTCTGGGTCAGTGGT	CAGACGTGTGCTCTTCCGATCTTGGAAAAGGGATAGAGGTTGG
CD74	TAGACAGATCCCCGTTCCCTG	CAGACGTGTGCTCTTCCGATCTACAGGGAGAAGGGATAACCC
SEPT9	CAGCATCCCAGCCTTGAG	CAGACGTGTGCTCTTCCGATCTCCTCAATGGCCTTTTGCTAC
CD79a	CCTCTAAACTGCCCCACCTC	CAGACGTGTGCTCTTCCGATCTCCTTAATCGCTGCCTCTAGG
GAPDH	CACATGGCCUCCAAGGAGU H AA	CAGACGTGTGCTCTTCCGATCTCAGCAAGAGCACAAAGAGGAA

[00585] In another experiment, we spiked in Ramos (Burkitt lymphoma) cells at a few percentage into primary B cells from a healthy individual. A panel of 111 genes (**Table 22**) was designed to represent different states of B cells. 1198 single cells were analyzed. A small group of the population, constituting 18 single cells (~1.5% of the population), was found to have a distinct gene expression pattern as compared to the rest (**FIG. 31B**). The genes that were preferentially expressed by this group are known to be associated with Burkitt lymphoma, such as MYC and IgM, as well as B cell differentiation markers (CD10, CD20, CD22, BCL6) that are expressed specifically by follicular B cells, which are the subset of B cells that Burkitt lymphoma originates (**FIG. 31C and 31D**). In addition, this group of cells carried higher level of CCND3 and GAPDH, as well as an overall higher mRNA content, as determined by the total number of unique mRNA molecules detected based on analyzing the molecular indices (**FIG. 31B**). This finding was consistent with the fact that lymphoma cells are physically larger than the primary B cells in normal individuals, and that they are rapidly proliferating and producing larger amount of transcripts.

Simultaneous identification of multiple cell types in human PBMCs

[00586] While the controlled experiments involved artificial mixtures of two distinct types of cells, most naturally occurring biological samples contain diverse populations with

numerous cell types and states with more subtle differences in gene expression profile. A prominent example is blood. We carried out an experiment in which we aimed to simultaneously identify all of the major cell types in human peripheral blood mononuclear cells (PBMCs), including monocytes, NK cells, and the different T and B cell subsets, by measuring the expression profile of a panel of 98 genes (**Table 19**) that are specific to each of the major cell type. Unlike traditional immunophenotyping that is limited mostly to surface protein markers, we included genes that encode cytokines, transcription factors, and intracellular proteins of various cellular functions in addition to surface proteins. We analyzed with PCA the digital gene expression profile of 632 single PBMCs using 81 genes present (**FIG. 32-39**). The first principal component clearly separated monocytes and lymphocytes into two orthogonal clusters, as evidenced by the expression of CD16a, CD14, S100A12, and CCR2 in one cluster, and lymphocyte associated genes in the other. The different subtypes of lymphocytes lay in a continuum along the second principal component, with B cells (expressing IgM, IgD, TCL1A, CD20, CD24, PAX5) at one end, naïve T cells (expressing CD4, CCR7, CD62L) in the middle, and cytotoxic T cells (expressing CD8A, CD8B, EOMES, PRF1) at the other end. Natural killer cells that express killer-like immunoglobulin receptor, CD16a, and perforin (PRF1) lay in the space between monocytes and cytotoxic T cells. We also observed that GAPDH, an indicator of cellular metabolism, was expressed at highest levels in monocytes and lowest in B cells, which are presumably mostly resting. Correlation analysis of gene expression profile across cells reiterated observations with PCA and revealed additional smaller subsets of cells within each major cell type (**FIG. 40A-B**). A replicate experiment of the same PBMC sample with 731 cells yielded largely similar segregation and cell type frequency (**FIG. 41**).

Table 22		
Gene	Outer Primer	Nested Primer with Common 5' Flanking Sequence
CD19	GCAGGGTCCCAGTCCTATG	CAGACGTGTGCTCTTCCGATCTCCAATCATGAGGAAGATGCA
CD27	TCCAGGAGGATTACCGAAAA	CAGACGTGTGCTCTTCCGATCTCCATCCAAGGGAGAGTGAGA
CD138	AATGGCAAAGGAAGGTGGAT	CAGACGTGTGCTCTTCCGATCTGCAGACACCTTGGACATCCT
CD38	AGATCTGAGCCAGTCGCTGT	CAGACGTGTGCTCTTCCGATCTTGGTGCAGAGCTGAAGATTTT
CD24	AAAAGTGGGCTTGATTCTGC	CAGACGTGTGCTCTTCCGATCTTTTGTTCGCATGGTCACAC
CD10	ATATTCCTTTGGGCCTCTGC	CAGACGTGTGCTCTTCCGATCTTCAAGTTTGGGTCTGTGCTG
CD95	CCCCGAAAATGTTCAATAA	CAGACGTGTGCTCTTCCGATCTTGCTCTTGTCCATACCCCCA
CD21	TAGCTTCCTCCTCTGGTGGT	CAGACGTGTGCTCTTCCGATCTTTTGCCTTTCCATAATCACTCA
CXCR3	CTGGCTCTCCCCAATATCCT	CAGACGTGTGCTCTTCCGATCTGCTCTGAGGACTGCACCACTT
CD40	GTGGTGTGGGGTATGTTT	CAGACGTGTGCTCTTCCGATCTATACACAGATGCCATTGCA
CD69	AGACAGGTCTTTTCGATGG	CAGACGTGTGCTCTTCCGATCTTGTGCAATATGTGATGTGGC
CD1c	TTGAGACAGGCACATACAGCTT	CAGACGTGTGCTCTTCCGATCTTGTCTTCTCAATCTGTCCA
IL10	CCCCAACCACTTCATTCTTG	CAGACGTGTGCTCTTCCGATCTTCAATTCCTCTGGGAATGTT

IL4R	TGCCTAGAGGTGCTCATTCA	CAGACGTGTGCTCTTCCGATCTGTTGATGCTGGAGGCAGAAT
IL21R	AGCCTGGGTACAGATCAAG	CAGACGTGTGCTCTTCCGATCTAGGTAGGAGGGTGGATGGAG
IL6R	CCAGCACCAGGGAGTTTCTA	CAGACGTGTGCTCTTCCGATCTAGGAAAGGATTGGAACAGCA
CXCL12	GGGTTTCAGGTTCCAATCAG	CAGACGTGTGCTCTTCCGATCTTTTGTAACTTTTGAAGGCA
CCL3	GTGAGGAGTGGGTCCAGAAA	CAGACGTGTGCTCTTCCGATCTAGTGGGAGGAGCAGGAG
CCL14	CCATTCCCTTCTTCTCCTC	CAGACGTGTGCTCTTCCGATCTTACCTACAAGATCCCGCGTC
CCL20	TTGGACATAGCCCAAGAACA	CAGACGTGTGCTCTTCCGATCTTGTGCCTCACTGGACTTGTCTC
CCL18	ACCTGAAGCTGAATGCCTGA	CAGACGTGTGCTCTTCCGATCTCTGGAGGCCACCTCTTCTAA
TCL1A	GGTAAACACGCCTGCAAAC	CAGACGTGTGCTCTTCCGATCTCAGGACTCAGAAGCCTCTGG
TACI	CAACAAAGCACAGTGTTAAATGAA	CAGACGTGTGCTCTTCCGATCTTGTGTCTAGCTACTGCGGAAA
AICDA	TGAGCAGATCCACAGGAAAA	CAGACGTGTGCTCTTCCGATCTGAAATGGAGTCTCAAAGCTTCA
FCRL4	TCCCAACTACGCTGATTTGA	CAGACGTGTGCTCTTCCGATCTGACCAAAAGGAATGTGTGGG
BCL2	TGCAAGAGTGACAGTGGATTG	CAGACGTGTGCTCTTCCGATCTTCAACCAAGGTTTGCTTTTGT
FASLG	AGAGGCTGAAAGAGGCCAAT	CAGACGTGTGCTCTTCCGATCTAATATGGGTTGCATTTGGTCA
BCL6	AAATCTGCAGAAGGAAAAATGTG	CAGACGTGTGCTCTTCCGATCTAGTTTTCAATGATGGGCGAG
AURKB	GCTCAAGGGAGAGCTGAAGA	CAGACGTGTGCTCTTCCGATCTGACTACCTGCCCCAGAGAT
CD81	GTGGCGTGATGAGTGGAGA	CAGACGTGTGCTCTTCCGATCTCACTCGCCCAGAGACTCAG
CD80	GCACATCTCATGGCAGCTAA	CAGACGTGTGCTCTTCCGATCTGCTTCACAAACCTTGCTCCT
CD23a	ACATTTTCTGCCACCCAAAC	CAGACGTGTGCTCTTCCGATCTAACAGCACCTCTCCAGATG
CD44	GCCTGGTAGAATTGGCTTTTC	CAGACGTGTGCTCTTCCGATCTTTTTGTAGCCAACATTCAATCAA
LEF1	CAATTGGCAGCCCTATTTCA	CAGACGTGTGCTCTTCCGATCTGTTGAGCAGACTGGTTTGCA
CXCR5	CCGTGAGGATGTCACTCAGA	CAGACGTGTGCTCTTCCGATCTACGAGGAAGCCCTAAGACGT
PRKCB	TTGAGCCTGGGGTGTAAAGAC	CAGACGTGTGCTCTTCCGATCTGTCTTCCAGATTACGGTG
PRKCD	GAGCACCTCCTGGAAGATTG	CAGACGTGTGCTCTTCCGATCTTAAGCACCAGTGGGACTGTG
CD20	TAGGAGCAGGCCTGAGAAAA	CAGACGTGTGCTCTTCCGATCTGATTCTCTCCAAACCCATG
CD30	TGTTTTGGGGAAAGTTGGAG	CAGACGTGTGCTCTTCCGATCTCTGTTTGCCCAAGTGTGTG
CD30L	TGCAACCCAACTGTGTGTTA	CAGACGTGTGCTCTTCCGATCTTTTACCAACTGTTCTCTGAGC
BAFFR	GCCCTGAGCAACAATAGCAG	CAGACGTGTGCTCTTCCGATCTTTCAGCTCTTCACTCCAGCA
CMRF-35H	AGGAAAAGATGTGGCTCACG	CAGACGTGTGCTCTTCCGATCTGGAGTTGGGGAGAACTGTCA
PRDM1	TCGAATAATCCAGGGAAACC	CAGACGTGTGCTCTTCCGATCTACCAAAGCATCACGTTGACA
HLA-DRA	GGCTTTACAAAGCTGGCAAT	CAGACGTGTGCTCTTCCGATCTTATGCCTCTTCGATTGCTCC
GNAI2	CCTTGAGTGTGTCTGCGTGT	CAGACGTGTGCTCTTCCGATCTCCACAGAATTGGGTTCCAAG
RGS1	AACTGGGAAGGCCAGGTAAC	CAGACGTGTGCTCTTCCGATCTTGTCTTCAAATTGCCATTGC
CD5	CTTTCTCCACGCCATTTGAT	CAGACGTGTGCTCTTCCGATCTACTAGGATATGGGGTGGGCT
CD22	GGGATCTGCTCGTCATCATT	CAGACGTGTGCTCTTCCGATCTGTTTCTGCCTCTGAGGGAAA
PIK3CD	GCGTGCGCGTTATTTATTTA	CAGACGTGTGCTCTTCCGATCTTGTCTGGGGAAGGCAAGTTA
DOCK8	GCAGTCAGCCAGAAATCACA	CAGACGTGTGCTCTTCCGATCTTTTTCTCCTCTCTGGGACCA
CD11b	TGAAAAGTCTCCCTTTCCAGA	CAGACGTGTGCTCTTCCGATCTCCTTCAGACACATTCCAGCC
FCGR2B	GGAGAGGAGAGATGGGGATT	CAGACGTGTGCTCTTCCGATCTGAGTGAGTGCCCTTTTCTT
CD72	CTCATGCCAACAGAACCTG	CAGACGTGTGCTCTTCCGATCTTGACCCACACCTGACACTTC
BCL11B	TCGTGGAACACAGGCAAAC	CAGACGTGTGCTCTTCCGATCTTTGCATTTGTACTGGCAAGG
CD86	TCAAGGCAACCAGAGGAAAC	CAGACGTGTGCTCTTCCGATCTACTAAGGGATGGGGCAGTCT
TBX21	ACCTTTTCGTTGGCATGTGT	CAGACGTGTGCTCTTCCGATCTTCAGGGAAAGGACTCACCTG
FOXP1	ATGCTGAAGGCATTTCTTGG	CAGACGTGTGCTCTTCCGATCTCTGTGAGCATGGTGCTTCAT
MCL1	GAGGGGAGTGGTGGGTTTAT	CAGACGTGTGCTCTTCCGATCTCAAAAGGGAAAGGGAGGATT
IFNB1	AGGGGAAAACATGAGCAG	CAGACGTGTGCTCTTCCGATCTTCACTGTGCCTGGACCATAG
BLNK	TTGGGCAGAAAGAAAAATGG	CAGACGTGTGCTCTTCCGATCTCAAAAGATTCCACCAGACTGAA

CD40LG	CCTCCCCCAGTCTCTCTTCT	CAGACGTGTGCTCTTCCGATCTGAGTCAGGCCGTTGCTAGTC
IGBP1	GGCTGATCTTCCCAACAC	CAGACGTGTGCTCTTCCGATCTACGAGGGCAAAGATGCTAAA
IRF4	ATTCCCGTGTGCTTCAAAC	CAGACGTGTGCTCTTCCGATCTAGAAGTCCAGCAGGTAGGA
CD79a	CACTTCCCTGGGACATTCTC	CAGACGTGTGCTCTTCCGATCTCACTCTTCTCCAGGCCAG
LTA	TGATGTCTGTCTGGCTGAGG	CAGACGTGTGCTCTTCCGATCTCCACACACAGAGGAAGAGCA
HDAC5	CCAGCCTGTAGGAAACCAAA	CAGACGTGTGCTCTTCCGATCTCTCCTTCTATCTCCAGGGCC
RAG1	GGATGCAGGTGGTTTTGAT	CAGACGTGTGCTCTTCCGATCTCATTGTACCCATTTTACATTTTCTT
RAG2	CAAACTTAAACACCCAGAAGC	CAGACGTGTGCTCTTCCGATCTATAACAATTCGGCAGTTGGC
CD1d	GAACCAGTTTCTCTCTGTGC	CAGACGTGTGCTCTTCCGATCTAAGATGTGGAGGCTGTTGCT
TGFB1	GACTGCGGATCTCTGTGTCA	CAGACGTGTGCTCTTCCGATCTTCTGCACTATTCTTTGCC
CD9	TCAGTATGATCTTGTGCTGTGCT	CAGACGTGTGCTCTTCCGATCTTACCCATGAAGATTGGTGGG
CD11c	CACAGCATGAGAGGCTCTGT	CAGACGTGTGCTCTTCCGATCTTCTCAGTTCCGATTTCCAG
FOXP3	TCAGGATCTGAGGTCCCAAC	CAGACGTGTGCTCTTCCGATCTTCACTGTGTATCTCACGCA
LAG3	AGAGCTGTCTAGCCAGGTG	CAGACGTGTGCTCTTCCGATCTTGGTGTCTTTCTCTGTCTCC
CD73	CTTAACGTGGGAGTGGAACC	CAGACGTGTGCTCTTCCGATCTGTGTGCAAATGGCAGCTAGA
CD70	TCTCAGCTTCCACCAAGGTT	CAGACGTGTGCTCTTCCGATCTTCACTGGGACACTTTTGCCT
CCR7	CAGGGGAGAGTGTGGTGT	CAGACGTGTGCTCTTCCGATCTGACATGCACTCAGCTCTTGG
CD45RA	TGCATAGTTCCCATGTTAAATCC	CAGACGTGTGCTCTTCCGATCTTACCAGGAATGGATGTCGCT
PDCD1	ACATCCTACGGTCCCAAGGT	CAGACGTGTGCTCTTCCGATCTGCAGAAAGTGCAGGCACCTA
MYC	TGCATGATCAAATGCAACCT	CAGACGTGTGCTCTTCCGATCTTGGACTTTGGGCATAAAAGA
CD25	AAATCACGGCAGTTTTCAGC	CAGACGTGTGCTCTTCCGATCTCTCATCTGTGCACTCTCCCC
FCAMR	GTGGGAAGAGAAGCTGATGC	CAGACGTGTGCTCTTCCGATCTTCAAGCATTATCCACGTCCA
CCND2	TGTGATGCCATATCAAGTCCA	CAGACGTGTGCTCTTCCGATCTTCAAGTATGCGAAAAGGTTTTT
MKI67	AGCCTCTCTTGGGCTTTCTT	CAGACGTGTGCTCTTCCGATCTGTTTTCCCTGCCTGGAACCTT
CCND3	CTTTGCTGCTGAAGGCTCAT	CAGACGTGTGCTCTTCCGATCTACAAGTGGTGGTAACCTGG
IL12A	TGCTTCTAAAAAGCGAGGT	CAGACGTGTGCTCTTCCGATCTGAAGTGGGAGGGGAAAGA
IFNG	GCAGCCAACCTAAGCAAGAT	CAGACGTGTGCTCTTCCGATCTATCCAGTACTGCCGTTTTG
TNFA	GAATGCTGCAGGACTTGAGA	CAGACGTGTGCTCTTCCGATCTACTTCCTTGAGACACGGAGC
IL2	ACCCAGGGACTTAATCAGCA	CAGACGTGTGCTCTTCCGATCTGCTGATGAGACAGCAACCATT
IL4	GACATCTTTGCTGCCTCCA	CAGACGTGTGCTCTTCCGATCTATGAGAAGGACACTCGCTGC
IL6	TTAAGGAGTTCCTGCAGTCCA	CAGACGTGTGCTCTTCCGATCTTCCACTGGGCACAGAACTTA
BAFF	TCCTTCGCTTTGCTTGTCTT	CAGACGTGTGCTCTTCCGATCTAGGTGGAAAAATAGATGCCAGTC
IGHE	CCCGGAAGTCTATGCGTTT	CAGACGTGTGCTCTTCCGATCTAGGACATCTCGGTGCAGTG
IGHD	TGTGTGAGGTGTCTGGCTTC	CAGACGTGTGCTCTTCCGATCTAGGAGCACCACGTTCTGG
IGHM	CCCGGAGAAGTATGTGACCA	CAGACGTGTGCTCTTCCGATCTGTACTTCGCCACAGCATC
IGHA	CTGAACGAGCTGGTGACG	CAGACGTGTGCTCTTCCGATCTAGTACCTGACTTGGGCATCC
IGHG1	CAAGGGCCCATCGGTCTT	CAGACGTGTGCTCTTCCGATCTTTGTGACAAAACACACATGC
IGHG4	CAAGGGCCCATCGGTCTT	CAGACGTGTGCTCTTCCGATCTCAAATATGGTCCCCCATGC
IGHG2	CAAGGGCCCATCGGTCTT	CAGACGTGTGCTCTTCCGATCTGCAAATGTTGTGTGAGTGC
IGHG3	CAAGGGCCCATCGGTCTT	CAGACGTGTGCTCTTCCGATCTACCCCACTTGGTGACACAAC
TLR1	CCATTCCGCAGTACTCCATT	CAGACGTGTGCTCTTCCGATCTAAGGAAAAGAGCAAACGTGG
TLR2	TTGGTTGACTTCATGGATGC	CAGACGTGTGCTCTTCCGATCTGGAACAGCACAAATGAACCTAA
TLR3	CATCATGCAGTTCAACAAGC	CAGACGTGTGCTCTTCCGATCTATGCACTCTGTTTGCGAAGA
TLR4	GGGTGTGTTTCCATGTCTCA	CAGACGTGTGCTCTTCCGATCTTTGAAAGTGTGTGTGTCGGC
TLR5	TCAGGCTGTTGCATGAAGAA	CAGACGTGTGCTCTTCCGATCTGTATGCCCTTGCTGGACCTA
TLR6	ATGCGCAGTAAAACTCGTG	CAGACGTGTGCTCTTCCGATCTTACAGTTCCACGCTGAGCTG
TLR7	GCCTGTACTTTAGCTGGGTA	CAGACGTGTGCTCTTCCGATCTAAGGTGTTTGTGCCATTTGG
TLR8	GGTGAGCTCTGATTGCTTCA	CAGACGTGTGCTCTTCCGATCTTATCAGGAGGCAGGGATCAC

TLR9	GACCGGGTCAGTGGTCTCT	CAGACGTGTGCTCTTCCGATCTGGTGATCCTGAGCCCTGAC
TLR10	TGCAGTGAGCTGAGATCGAG	CAGACGTGTGCTCTTCCGATCTATGGAAAACATCCTCATGGC
GAPDH	CACATGGCCUCCAAGGAGUAA	CAGACGTGTGCTCTTCCGATCTCAGCAAGAGCACAAAGAGGAA

Table 19

Gene	Outer Primer	Nested Primer with Common 5' Flanking Sequence
CD19	GCAGGGTCCCAGTCCTATG	CAGACGTGTGCTCTTCCGATCTCCAATCATGAGGAAGATGCA
CD20	TAGGAGCAGGCCTGAGAAAA	CAGACGTGTGCTCTTCCGATCTGATTCTCTCCAAACCCATG
BAFF	TCCTTCGCTTTGCTTGCTT	CAGACGTGTGCTCTTCCGATCTAGGTGGAAAAATAGATGCCAGTC
TCL1A	GGTAAACACGCCTGCAAAAC	CAGACGTGTGCTCTTCCGATCTCAGGACTCAGAAGCCTCTGG
TACI	CAACAAAGCACAGTGTTAAATGAA	CAGACGTGTGCTCTTCCGATCTTGCTCAGCTACTGCGGAAA
IGHD	TGTGTGAGGTGTCTGGCTTC	CAGACGTGTGCTCTTCCGATCTAGGAGCACCACGTTCTGG
IGHM	CCCGGAGAAGTATGTGACCA	CAGACGTGTGCTCTTCCGATCTGTACTTCGCCACAGCATC
CD27	TCCAGGAGGATTACCGAAAA	CAGACGTGTGCTCTTCCGATCTCCATCCAAGGGAGAGTGAGA
CD38	AGATCTGAGCCAGTCGCTGT	CAGACGTGTGCTCTTCCGATCTTGGTGCAGAGCTGAAGATTTT
CD24	AAAAGTGGGCTTGATTCTGC	CAGACGTGTGCTCTTCCGATCTTTTGTTCGCATGGTCACAC
AICDA	TGAGCAGATCCACAGGAAAA	CAGACGTGTGCTCTTCCGATCTGAAATGGAGTCTCAAAGCTTCA
CD95	CCCCCGAAAATGTTCAATAA	CAGACGTGTGCTCTTCCGATCTTGCTCTTGTCATACCCCCA
CD10	ATATTCTTTGGGCCTCTGC	CAGACGTGTGCTCTTCCGATCTTCAAGTTTGGGTCTGTGCTG
IL10	CCCCAACCACTTCATTCTTG	CAGACGTGTGCTCTTCCGATCTTCAATTCTCTGGGAATGTT
CD138	AATGGCAAAGGAAGGTGGAT	CAGACGTGTGCTCTTCCGATCTGCAGACACCTTGACATCCT
CD45RA	TGCATAGTTCCCATGTTAAATCC	CAGACGTGTGCTCTTCCGATCTTACCAGGAATGGATGTCGCT
BCL6	AAATCTGCAGAAGGAAAAATGTG	CAGACGTGTGCTCTTCCGATCTAGTTTTCAATGATGGCGAG
PRDM1	TCGAATAATCCAGGGAAACC	CAGACGTGTGCTCTTCCGATCTACCAAAAGCATCAGTTGACA
CXCR3	CTGGCTCTCCCCAATATCCT	CAGACGTGTGCTCTTCCGATCTGCTCTGAGGACTGCACCATT
IFNG	GCAGCCAACCTAAGCAAGAT	CAGACGTGTGCTCTTCCGATCTATCCAGTTACTGCCGTTTG
IL4R	TGCCTAGAGGTGCTCATTCA	CAGACGTGTGCTCTTCCGATCTGTTGATGCTGGAGGCAGAAT
IL4	GACATCTTTGCTGCCTCCA	CAGACGTGTGCTCTTCCGATCTATGAGAAGGACACTCGCTGC
CCL20	TTGGACATAGCCCAAGAACA	CAGACGTGTGCTCTTCCGATCTTGCTGCTCACTGGACTTGTC
CD25	AAATCACGGCAGTTTTCAGC	CAGACGTGTGCTCTTCCGATCTCTCATCTGTGCACTCTCCCC
FOXP1	ATGCTGAAGGCATTTCTTGG	CAGACGTGTGCTCTTCCGATCTCTGTGAGCATGGTGCTTCAT
TGFB1	GACTGCGGATCTCTGTGCA	CAGACGTGTGCTCTTCCGATCTTCTGCACTATTCTTTGCC
CXCR5	CCGTGAGGATGTCACTCAGA	CAGACGTGTGCTCTTCCGATCTACGAGGAAGCCCTAAGACGT
CD69	AGACAGGTCTTTTCGATGG	CAGACGTGTGCTCTTCCGATCTTGTCGAATATGTGATGTGGC
CD30	TGTTTTGGGGAAAGTTGGAG	CAGACGTGTGCTCTTCCGATCTGTTTGCCCAAGTGTGTTG
PDCD1	ACATCCTACGGTCCCAAGGT	CAGACGTGTGCTCTTCCGATCTGCAGAAGTGCAGGCACCTA
LAG3	AGAGCTGTCTAGCCCAGGTG	CAGACGTGTGCTCTTCCGATCTTGGTGTCTTTCTCTGCTCC
PAX5	TGACGTGTGTTGCTTTTGTG	CAGACGTGTGCTCTTCCGATCTACTTGGGAGAAAACAGGGGT
TNFRSF17	GCTTCCACTCCCAGCTATG	CAGACGTGTGCTCTTCCGATCTTGCTTTGAGTGCTACGGAGA
RASD1	GGGGGAGGGATGTGAAGTTA	CAGACGTGTGCTCTTCCGATCTATCTTGCTGTGATTCCGGG
AMPD1	ACAGATGACCCAATGCAATTC	CAGACGTGTGCTCTTCCGATCTGAGCACCTGTGATATGTGCG
OSBPL5	AGACCGATGCACAGTCTTCC	CAGACGTGTGCTCTTCCGATCTTTCACGTCTGGCCTCAGTC
CD56	GGAGCACTCAAGTGTGACGA	CAGACGTGTGCTCTTCCGATCTTTTCTATGGAGCCTTCCGA
IGFBP7	CATCCAATTCCAAGGACAG	CAGACGTGTGCTCTTCCGATCTGGTGAAGGTGCCGAGCTATA
KIR2DS5	GCTCTTCTCAAACACGAA	CAGACGTGTGCTCTTCCGATCTCACACTCCTTTGCTTAGCCC
KIR2DS2	TCCTCACACCAGCAATCTGA	CAGACGTGTGCTCTTCCGATCTCACTCCTTTGCTTAGCCAC
RAB4B	CCAGCTCACCTGTTCTCCAG	CAGACGTGTGCTCTTCCGATCTGAATCCCGTACCTGCTGCT
CD14	CTAAAGGACTGCCAGCCAAG	CAGACGTGTGCTCTTCCGATCTATAACCTGACACTGGACGGG

S100A12	CACATTCCTGTGCATTGAGG	CAGACGTGTGCTCTTCCGATCTATACTCAGTTCGGAAGGGGC
CCR2variantB	GAAGGAGGGAGACATGAGCA	CAGACGTGTGCTCTTCCGATCTACTGGTCCTTAGCCCCATCT
CD62L	TCAGTTGGCTGACTTCCACA	CAGACGTGTGCTCTTCCGATCTTAGTTGGGGGTTTTGCTG
CD16	TCTTGGCCAGGGTAGTAAGAA	CAGACGTGTGCTCTTCCGATCTGTCAGTTCCAATGAGGTGGG
CX3CR1	CGTCCAGACCTTGTTACAC	CAGACGTGTGCTCTTCCGATCTCCACAAATAGTGCTCGCTTTC
CD1b	TAGAGGGCCAGGACATCATC	CAGACGTGTGCTCTTCCGATCTTTGCTCCTTTTGCTATGCCT
FOXQ1	TGCTATTGACCGATGCTTCA	CAGACGTGTGCTCTTCCGATCTGCAACGGGCTACAGCTTTAT
CD209	GCTCTTGTTCTTGCCGTTTT	CAGACGTGTGCTCTTCCGATCTGAGTCCCTCAGTGGAGCAAG
CD1e	CACAAGCACATTCATCTCTTCC	CAGACGTGTGCTCTTCCGATCTATTAGGGCCAGCTTCATAA
CCL17	TACTTCAAGGGAGCCATTCC	CAGACGTGTGCTCTTCCGATCTTTGTAAGTGTGCAGGGCAG
DTNA	AGCAACGTGGAGTCAGTCTGT	CAGACGTGTGCTCTTCCGATCTCTCACCTTCTCTTGCCCTTGG
CLEC4C	TTATTTTCTGGGGCTGTCAGA	CAGACGTGTGCTCTTCCGATCTCATTCTGGCACTCAGGTGAA
ZBTB16	TGATCAAGCACCTGAGAACG	CAGACGTGTGCTCTTCCGATCTTACCAGTGCAACCATCTGCAC
SLAMF1	TGCAAAACCCAGAAGCTAAAA	CAGACGTGTGCTCTTCCGATCTGTTCTGTGCAAAATGGCATTTC
CD3D	AGAGCTGTGTGGAGCTGGAT	CAGACGTGTGCTCTTCCGATCTGGAGTCTTCTGCTTTGCTGG
CD3E	GCCCTCTTGCCAGGATATTT	CAGACGTGTGCTCTTCCGATCTGCATGTAAGTTGTCCCCCAT
CD8A	CTGGCCTCTGCTCAACTAGC	CAGACGTGTGCTCTTCCGATCTATGGTACAAGCAATGCCTGC
CD8B	CAGCCTCAAGGGGAAGGTAT	CAGACGTGTGCTCTTCCGATCTTGCTTAACCCATGGATCCTG
PRF1	CCCTGCAGTCACAGCTACAC	CAGACGTGTGCTCTTCCGATCTTCAGGGCTGGTCTTTTAGGA
EOMES	TGGGATAATGTAAACTGGTGCT	CAGACGTGTGCTCTTCCGATCTCATCCCCATGATATTTGGGA
CD4	AGCTAGCCTGAGAGGGAACC	CAGACGTGTGCTCTTCCGATCTTCTCCAGACCATTACAGGAC
THPOK	GGCTCTGCCTTGCACTATTT	CAGACGTGTGCTCTTCCGATCTCTCTTCTCCCTTCCATGC
RUNX3	TAAGGCCCAAAGTGGGTACA	CAGACGTGTGCTCTTCCGATCTTAGGAAGCACAGGAAAGGA
CD45RO	ACCCTCTCTCCCTCCCTTTC	CAGACGTGTGCTCTTCCGATCTTAGTTGGCTATGCTGGCATG
CD44	GCCTGGTAGAATTGGCTTTTC	CAGACGTGTGCTCTTCCGATCTTTTGTAGCCAACATTCATTCAA
CCR7	CAGGGGAGAGTGTGGTGTTT	CAGACGTGTGCTCTTCCGATCTACTCAGCTCTTGGCTCCACT
TXK	ACATCAAGCTCCATTGTTTCG	CAGACGTGTGCTCTTCCGATCTTTTGCTGCACTCTTTGTAGG
MBD2	GCCTGGCACGTAATAGCTTG	CAGACGTGTGCTCTTCCGATCTAGGAAAGAAATGCCCTTGGT
IFNGR1	GAGGATGTGTGGCATTITCA	CAGACGTGTGCTCTTCCGATCTGGTTTCTAGGTGAGCAGGTG
IL12RB2	AGCAGGCTGTACACAGCAGA	CAGACGTGTGCTCTTCCGATCTGACACTAGGCACATTGGCTG
IL33R	ACTGTGCCCTCATCCAGAAC	CAGACGTGTGCTCTTCCGATCTAACGACGCCAAGGTGATACT
CCR4	TGGTGAAATGCAGAGTCAATG	CAGACGTGTGCTCTTCCGATCTTCAGGAGGAAGGCTTACACC
CRTH2	TGAATTTTGCTTGGTGGATG	CAGACGTGTGCTCTTCCGATCTTGTCAGTGGAAGAAGCAGATG
IL5	CAGTGAGAATGAGGGCCAAG	CAGACGTGTGCTCTTCCGATCTGAATGAGGGCCAAGAAAGAG
IL17A	AAAATGAAACCCTCCCCAAA	CAGACGTGTGCTCTTCCGATCTTCTTTGGAGATTAAGGCC
IL17F	CTGCATCAATGCTCAAGGAA	CAGACGTGTGCTCTTCCGATCTCCAAGGCTGCTCTGTTTCTT
IL21	AAATCAAGCTCCCAAGGTCA	CAGACGTGTGCTCTTCCGATCTTGTAATGACTTGGTCCCTG
IL22	ATGCCCCAAAGCGATTTTT	CAGACGTGTGCTCTTCCGATCTCAAAGGAAACCAATGCCACT
IL23R	TCCCTCATTGAAAGATGCAA	CAGACGTGTGCTCTTCCGATCTTAGAATCATTAGGCCAGGCG
RORA	TGCAAGCCATTATGGGAAT	CAGACGTGTGCTCTTCCGATCTCCTTGGGTTTTCTTTTCAATTC
RORC	ATTTCCATGGTGCTCCAGTC	CAGACGTGTGCTCTTCCGATCTAGAGAAGCAGAAGTCGCTCG
OX40L	CTGCTGGCCCTGTACCTG	CAGACGTGTGCTCTTCCGATCTCTCCACCCTGGCCAAGAT
ICOS	TTCAGCTGACTTGGACAACCT	CAGACGTGTGCTCTTCCGATCTGGACAACCTGACTGGCTTTG
SH2D1A	GGGTGTTGGTGAACCTGGTT	CAGACGTGTGCTCTTCCGATCTTTAATATGGATGCCGTGGG
CCR2variantA	GTTGCCAGTGTGTTTCTGA	CAGACGTGTGCTCTTCCGATCTAACCAGGCAACTTGGGAAC
TLR1	CCATTCCGCAGTACTCCATT	CAGACGTGTGCTCTTCCGATCTAAGGAAAAGACAAACGTGG
TLR2	TTGGTTGACTTCATGGATGC	CAGACGTGTGCTCTTCCGATCTGGAAACAGCACAAATGAACTTAA
TLR3	CATCATGCAGTTCAACAAGC	CAGACGTGTGCTCTTCCGATCTATGCACTCTGTTTGCGAAGA

TLR4	GGGTGTGTTTCCATGTCTCA	CAGACGTGTGCTCTTCCGATCTTTGAAAGTGTGTGTGTCGGC
TLR5	TCAGGCTGTTGCATGAAGAA	CAGACGTGTGCTCTTCCGATCTGTATGCCCTTGCTGGACCTA
TLR6	ATGCGCAGTAAAACTCGTG	CAGACGTGTGCTCTTCCGATCTTACAGTCCACGCTGAGCTG
TLR7	GCCTGTACTTTCAGCTGGGTA	CAGACGTGTGCTCTTCCGATCTAAGGTGTTTGTGCCATTGG
TLR8	GGTGAGCTCTGATTGCTTCA	CAGACGTGTGCTCTTCCGATCTTATCAGGAGGCAGGGATCAC
TLR9	GACCGGGTCAGTGGTCTCT	CAGACGTGTGCTCTTCCGATCTGGTGATCCTGAGCCCTGAC
TLR10	TGCAGTGAGCTGAGATCGAG	CAGACGTGTGCTCTTCCGATCTATGGAAACATCCTCATGGC
GAPDH	CACATGGCCUCCAAGGAGUAA	CAGACGTGTGCTCTTCCGATCTCAGCAAGAGCACAAGAGGAA

Studying diversity of response of human T cells to in vitro stimulus

[00587] When examining the gene expression pattern of a bulk sample, the observed pattern was contributed by both the sample's cell composition and the expression level of each gene in each cell type or subtype. These two effects cannot be deconvoluted by bulk analysis but only with large-scale single cell analysis. To illustrate, we utilized our platform to study the variability of response of human T cells to an in vitro stimulus.

[00588] We purified CD3⁺ T cells by negative selection from a blood donor and stimulated them with anti-CD28/anti-CD3 beads for 6 hours, and performed experiments with the stimulated and a separate aliquot of unstimulated cells. We designed a panel of 93 genes (**Table 20**) that encompassed surface proteins, cytokines, chemokines, and effector molecules expressed by the different T cell subsets. A total of 3517 and 1478 single cells were analyzed for the stimulated and unstimulated samples, respectively.

Table 20		
Gene	Outer Primer	Nested Primer with Common 5' Flanking Sequence
GAPDH	GACTTCAACAGCGACACCCA	CAGACGTGTGCTCTTCCGATCTGCCCTCAACGACCACTTTGT
CD3D	GAAAACGCATCCTGGACCCA	CAGACGTGTGCTCTTCCGATCTTGATGTCATTGCCACTCTGCT
CD3E	AAGTTGTCCCCATCCCAA	CAGACGTGTGCTCTTCCGATCTCTGGGGATGGACTGGGTAAT
CD8A	ACTGCTGTCCCAACATGCA	CAGACGTGTGCTCTTCCGATCTATGCCTGCCATTGGAGAGAA
CD8B	CCACCATCTTTCAGGTTGC	CAGACGTGTGCTCTTCCGATCTGCTGTCCAGTTCCCAGAAGG
CD4	CTGGGAGAGGGGGTAGCTAG	CAGACGTGTGCTCTTCCGATCTACCACTTCCCTCAGTCCCAA
FOXP3	ACAGAAGCAGCGTCAGTACC	CAGACGTGTGCTCTTCCGATCTGGGTCTCTTGAGTCCCGTG
CCR7	GGGGAGAGTGTGGTGTTC	CAGACGTGTGCTCTTCCGATCTCTTGGCTCCACTGGGATG
CD5	ATCAATGGTCCAAGCCGCAT	CAGACGTGTGCTCTTCCGATCTAGGTCACAGATCTTCCCCCG
IL32	CTTTCAGTCCTACGGAGCC	CAGACGTGTGCTCTTCCGATCTTGCTCTGAACCCCAATCCTC
CD28	ACCATCACAGGCATGTTCT	CAGACGTGTGCTCTTCCGATCTTGATGACCTGGCTTGCC
SELL	GCATCTCATGAGTGCCAAGC	CAGACGTGTGCTCTTCCGATCTCCTGCCCCAGACCTTTTATC
CD27	TGCAGAGCCTTGTGTTACA	CAGACGTGTGCTCTTCCGATCTCGTGACAGAGTGCCCTTTTCG
GZMB	AGGTGAAGATGACAGTGCAAG	CAGACGTGTGCTCTTCCGATCTAGGCCCTCTTGTTGTGAACA
GZMA	GGAACCATGTGCCAAGTTGC	CAGACGTGTGCTCTTCCGATCTCCTTTGTTGTGCGAGGGTGT
GZMH	AGTGTGCTGACAGTGACAGA	CAGACGTGTGCTCTTCCGATCTCCAAAGAAGACACAGACCGGT
GZMK	TTGCCACAAAGCCTGGAATC	CAGACGTGTGCTCTTCCGATCTAAAGCAACCTTGTCCCGCCT
PRF1	GGAGTCCAGCGAATGACGTC	CAGACGTGTGCTCTTCCGATCTCATGGCCACGTTGTATTGT
NKG2D	CAACACCCAGGGGATCAGTG	CAGACGTGTGCTCTTCCGATCTCCACCCTCCACAGGAAATTG

LAG3	AGCTGTACCAGGGGGAGAG	CAGACGTGTGCTCTTCCGATCTCTTTGGAGAAGACAGTGCGCA
CD160	GGAAGACAGCCAGATCCAGTG	CAGACGTGTGCTCTTCCGATCTTTGTGCAGACCAAGAGCACC
CD244	GGGCTGAGAATGAGGCAGTT	CAGACGTGTGCTCTTCCGATCTGGAAGCGACAAGGGTGAAC
EOMES	ACTTAACAGCTGCAGGGGC	CAGACGTGTGCTCTTCCGATCTACTAACTTGAACCGTGTTTAAGG
TBX21	TTATAACCATCAGCCCGCCA	CAGACGTGTGCTCTTCCGATCTAGAAAAGGGGCTGGAAGGG
PRDM1	ACCAAAGCATCACGTTGACAT	CAGACGTGTGCTCTTCCGATCTACATGTGAATGTTGAGCCCA
IRF4	CTCTTCAGCATCCCCGTAC	CAGACGTGTGCTCTTCCGATCTGCCCCAAATGAAAGCTTGA
ZNF683	GGAGAGCGTCCATTCCAGTG	CAGACGTGTGCTCTTCCGATCTATCCACCTGAAGCTGCACC
ZBED2	AATGTACCAGCCAGTCAGCG	CAGACGTGTGCTCTTCCGATCTGGTTTTGGTGGAGCTGACGA
CD30	TTTACTCATCGGGCAGCCAC	CAGACGTGTGCTCTTCCGATCTTGTGGCCAGTGTGTGTC
CD69	GCTGTAGACAGGTCTTTTCG	CAGACGTGTGCTCTTCCGATCTAGTGTGGAAAATGTGCAATATGTG
HLA-DRA	GGGTCTGGTGGGCATCATTA	CAGACGTGTGCTCTTCCGATCTGCCTCTTCGATTGCTCCGTA
CD38	AGGTCAATGCCAGAGACGGA	CAGACGTGTGCTCTTCCGATCTATCAGCATACCTTTATTGTGATCTATC
TNFRSF9	TGGCATGTGAGTCATTGCTC	CAGACGTGTGCTCTTCCGATCTTTTGTGTGAGGGGCGGAT
MKI67	TACTTTTTCGCCTCCCAGGG	CAGACGTGTGCTCTTCCGATCTTCTGCCCCACCAAGATCAT
BIRC5	TGCCACGGCCTTTCCTTAA	CAGACGTGTGCTCTTCCGATCTTTGTCTAAGTGAACCGCCT
FOSL1	CTCCTGACAGAAGGTGCCAC	CAGACGTGTGCTCTTCCGATCTGGTGATTGGACCAGGCCATT
MCL1	GACTGGCTACGTAGTTCGGG	CAGACGTGTGCTCTTCCGATCTTTTGCTTAGAAGGATGGCGC
MYC	AGCTACGGAACCTCTGTGCG	CAGACGTGTGCTCTTCCGATCTCAACCTTGGCTGAGCTTGA
TYMS	TCAGTCTTTAGGGGTTGGGC	CAGACGTGTGCTCTTCCGATCTATGTGCATTTCAATCCCACGTAC
CDCA7	CCAGTCTAGTTCTGGGCAGG	CAGACGTGTGCTCTTCCGATCTATGTAAACCATTGCTGTGCCATT
UHRF1	CCAGTCTTCTGACACCGG	CAGACGTGTGCTCTTCCGATCTCCAAAGTTTGACGCTATACC
SAP30	ACCAACCAGACCAGGACTTA	CAGACGTGTGCTCTTCCGATCTTCACTAGGAGACGTGGAATTG
CX3CR1	CACCCGTCCAGACCTTGT	CAGACGTGTGCTCTTCCGATCTTGTCTTCTTAACGTTAGACCAC
BCL2	TGCAAGAGTGACAGTGGAATTG	CAGACGTGTGCTCTTCCGATCTGCTGATATTCTGCAACACTGTACA
BCL6	TGCTCTACGGTGCCTTTT	CAGACGTGTGCTCTTCCGATCTGTAGGCAGACACAGGGACTT
FASLG	CCTCAAGGGGGACTGTCTTTC	CAGACGTGTGCTCTTCCGATCTGCATATCCTGAGCCATCGGT
FAS	ATTGCTGGTAGAGACCCCA	CAGACGTGTGCTCTTCCGATCTCCCCATTTCCCCGATGT
CCL4	CCCAGCCAGCTGTGGTATTC	CAGACGTGTGCTCTTCCGATCTTGAACTGAACTGAGCTGCT
IFNG	CTAGGCAGCCAACCTAAGCA	CAGACGTGTGCTCTTCCGATCTCCTGCAATCTGAGCCAGTGC
TNF	AGTGGACCTTAGGCCTTCCT	CAGACGTGTGCTCTTCCGATCTGGCTCAGACATGTTTTCCGTG
IL2	TCACTTAAGACCCAGGGACTT	CAGACGTGTGCTCTTCCGATCTAAGCATCATCTCAACACTGACTT
IL4	ACCATGAGAAGGACACTCGC	CAGACGTGTGCTCTTCCGATCTCGGGCTTGAATTCCTGTCT
IL6	CGGCAAATGTAGCATGGGC	CAGACGTGTGCTCTTCCGATCTGGAAGTGGCTATGCAGTTTG
IL1A	GGCATCCTCCACAATAGCAGA	CAGACGTGTGCTCTTCCGATCTGCATTTTGGTCCAAGTTGTGC
IL1B	CTTAAAGCCCGCCTGACAGA	CAGACGTGTGCTCTTCCGATCTACATTCTGATGAGCAACCGC
IL3	ACAGACGACTTTGAGCCTCG	CAGACGTGTGCTCTTCCGATCTATTTACCTTTCTGCGGC
IL13	GGAGCCAAGGGTTCAGAGAC	CAGACGTGTGCTCTTCCGATCTTGCTACCTCACTGGGGTCT
IL31	GGCCATCTCTTCTTTTCGGA	CAGACGTGTGCTCTTCCGATCTGTGTGGGAACCTGCGCGT
IL24	CTACCCCATCATCCCTTTC	CAGACGTGTGCTCTTCCGATCTGCCAGTGAGACTGTGTGT
IL26	TACTGACGGCATGTTAGGTG	CAGACGTGTGCTCTTCCGATCTTGTGTGTGGAGTGGGATGTG
LTA	AGGCAGGGAGGGGACTATTT	CAGACGTGTGCTCTTCCGATCTGGAGAAACAGAGACAGGCC
IL5	GCAGTGAGAATGAGGGCCA	CAGACGTGTGCTCTTCCGATCTAGGCATACTGACACTTTGCC
CSF2	AGCCAGTCCAGGAGTGAGAC	CAGACGTGTGCTCTTCCGATCTGGCCACACTGACCCTGATAC
IL21	CCCAAGGTCAAGATCGCCAC	CAGACGTGTGCTCTTCCGATCTCTGCCAGCTCCAGAAGATGT
IL22	TGGGAAGCCAACTCCATCAT	CAGACGTGTGCTCTTCCGATCTGGAACCAATGCCACTTTTGT
IL17A	GCCTTCAAGACTGAACACCGA	CAGACGTGTGCTCTTCCGATCTGCCCCCTCAGAGATCAACAGAC
IL17F	TTGGAGAAGGTGCTGGTGAC	CAGACGTGTGCTCTTCCGATCTTACCCAGTGCTCTGCAAC

TGFB1	TATTCCTTTGCCCGGCATCA	CAGACGTGTGCTCTTCCGATCTACCTTGGGCACTGTTGAAGT
CCL20	ACTTGACATCATGGAGGGT	CAGACGTGTGCTCTTCCGATCTTCCATAAGCTATTTGGTTTAGTGC
IL12A	GGTCCCTCCAAACCGTTGTC	CAGACGTGTGCTCTTCCGATCTGAACTAGGGAGGGGAAAGAAG
CXCL12	TGGGAGTTGATCGCCTTTCC	CAGACGTGTGCTCTTCCGATCTCTCATTCTGAAGGAGCCCAT
CCL3	TGGACTGGTTGTTGCCAAAC	CAGACGTGTGCTCTTCCGATCTCTCTGAGAGTTCCCCTGTCC
CCL14	TTCCTCCTCATCACCATCGC	CAGACGTGTGCTCTTCCGATCTCTTACCACCCCTCAGAGTGC
CCL18	GAAGCTGAATGCCTGAGGGG	CAGACGTGTGCTCTTCCGATCTGTCCCATCTGCTATGCCCA
CCL17	GAGTGCTGCCTGGAGTACTT	CAGACGTGTGCTCTTCCGATCTCTCACCCAGACTCCTGACT
IL12B	GCTATGGTGAGCCGTGATTG	CAGACGTGTGCTCTTCCGATCTTCCCTACCCCCACCTCTCTA
CXCR3	GACCTCAGAGGCCTCCTACT	CAGACGTGTGCTCTTCCGATCTCCAATATCCTCGCTCCCGG
IL33R	TTCAGGACTCCCTCCAGCAT	CAGACGTGTGCTCTTCCGATCTAGGTACCAAAATGCCTGTGCC
IL4R	TGAACCTCAGGGAGGGTGGT	CAGACGTGTGCTCTTCCGATCTTCCCTCGTATGCATGGAACCC
CCR4	CCAAAGGGAAGAGTGCAGGG	CAGACGTGTGCTCTTCCGATCTATTCTGTATAACACTCATATCTTTGCC
IL23R	AGAATCATTAGGCCAGGCGTG	CAGACGTGTGCTCTTCCGATCTCTGGCCAATATGCTGAAACCC
IL21R	ATTTGAGGCTGCAGTGAGCT	CAGACGTGTGCTCTTCCGATCTAGACAAGAGCTGGCTCACCT
CXCR5	CCTCCCCAGCCTTTGATCAG	CAGACGTGTGCTCTTCCGATCTTCCCTCGAAGCTGGGTAATC
IL6R	CCAGCACCAAGGAGTTTCTA	CAGACGTGTGCTCTTCCGATCTACAGCATGTCAAGGCTGT
CXCL13	AGGCAGATGGAACCTGAGCC	CAGACGTGTGCTCTTCCGATCTGCATTGGAAGATCCCCAGACTT
LIF	TCCCCATCGTCTCTTGTCT	CAGACGTGTGCTCTTCCGATCTTTGCCGGCTCTCCAGAGTA
PTPRCv1 (CD45RA)	GTTCCCATGTTAAATCCCATTCAT	CAGACGTGTGCTCTTCCGATCTTACCAGGAATGGATGTCGCTAATCA
PTPRCv2 (CD45RO)	ACCTCTCTCCCTCCCTTTC	CAGACGTGTGCTCTTCCGATCTTAGTTGGCTATGCTGGCATG
IL10	CCCCAACCACTTCATTCTTG	CAGACGTGTGCTCTTCCGATCTTTCAATTCTCTGGGAATGTT
CD40LG	CCTCCCCCAGTCTCTCTTCT	CAGACGTGTGCTCTTCCGATCTGAGTCAGGCCGTTGCTAGTC

[00589] In the unstimulated sample, PCA analysis revealed two major subsets of cells. A closer look at the genes enriched in each subset showed that one subset represented CD8⁺ cells with expression of CD8A, CD8B, NKG2D, GZMA, GZMH, GZMK, and EOMES, and the other subset represented CD4⁺ cells with expression of CD4, CCR7 and SELL (**FIG. 44A and FIG. 45**).

[00590] In the stimulated sample, two branches of cells were immediately clear on the PCA plot (**FIG. 44B and FIG. 46A-D**). The first principal component represented the degree of response of individual cells to stimulant in terms of varying level of expression of IFNG, TNF, CD69, and GAPDH. Expression of CCL3, CCL4, and GZMB, which are cytokines and effector molecules associated with cytotoxic T cells, and LAG3, a marker associated with exhausted cells, was localized to cells in the upper branch. Expression of IL2, LTA, CD40LG, and CCL20, which are cytokines associated with helper T cells, was localized to the lower branch. Other genes that have been known to be upregulated in activated T cells, including ZBED2, IL4R, PRDM1, TBX21, MYC, FOSL1, CSF2, TNFRSF9, BCL2 and FASLG, were expressed in various degrees in a smaller number of cells (**FIG. 46A-D**). Most

of these cytokines, effector molecules, and transcription factors were not expressed or were expressed at very low levels by cells in the unstimulated sample. While most of the cells that responded within this short period of stimulation were presumably memory cells, we observed a small population of cells that produced lower level of IL2 and not other cytokines nor effector molecules, and may represent naïve cells (**FIG. 44B, arrow**).

[00591] To fully appreciate the heterogeneity in response, we clustered the cells based on a pair-wise correlation coefficient. While the two main groups of CD4 and CD8 cells were obvious, there was considerable diversity within each set in terms of the combination and level of activated genes expressed (**FIG. 47 and FIG. 48**).

[00592] We observed that there were a few cytokines, namely IL4, IL5, IL13, IL17F, IL22, LIF, IL3, and IL21, that were upregulated by a few hundred or more folds in the stimulated sample as a whole as compared to the unstimulated one, but were contributed only by a few cells in the sample (**FIG. 44C**). Subsets of these cytokines were expressed by the same cells (**FIG. 49A-C**). For instance, the same single cell contributed to most of the counts of IL17F and IL22, which were signatures for Th17 cells. Another 7 cells expressed various combinations of IL4, IL5, IL13, which were signatures of Th2 cells, and expressed various combinations of them. Such observation highlights the importance of large-scale single cell analysis, especially when the contribution to overall expression changes was derived from a rare subpopulation.

[00593] We repeated the same stimulation experiment with T cells from a second blood donor and analyzed the profile of 669 and 595 single cells in the stimulated and unstimulated sample, respectively. While the overall level of activation was lower (smaller magnitude in terms of change in expression) in this individual (possibly indicating inter-individual variability to stimulation), we observed the same trends in PCA analysis, as well as heterogeneity in individual cell's response to stimulus (**FIG. 48**).

Identification of rare antigen specific T cells

[00594] We demonstrated the utility of our platform to identify rare cells using the model of antigen specific cells in CD8+ T cell population. We exposed fresh blood of the same two blood donors who were seropositive for cytomegalovirus (CMV) to CMV pp65 peptide pool. A separate untreated blood aliquot of each donor served as negative control. We subsequently isolated CD8+ T cells and analyzed the response of stimulated and unstimulated cells on our platform. We obtained data from 2274, 2337, 581, and 253 cells in donor 2's CMV stimulated and unstimulated, and donor 1's CMV stimulated and unstimulated samples, respectively.

[00595] Except for donor 1's negative control that yielded relatively small number of cells to form obvious clusters in clustering analysis, all the rest of the samples showed two main groups of cells (**FIG. 50A, 51 and 52**). Cells in one group expressed naïve cell and central memory associated markers *SELL*, *CCR7*, and *CD27*, while cells in the other group expressed effector memory cell (*CCL4*, *CX3CR1*, *CXCR3*) and effector cell associated genes (*EOMES*, *GZMA*, *GZMB*, *GZMH*, *TBX21*, *ZNF683*). There was a distinct small subset of cells that occupy space in between the two branches and express granzyme K (*GZMK*), as well as another subset of HLA-DRA expressing cells. The differential expression of the different types of granzymes has previously been reported (8). Our results recapitulated those observed in previous CyTOF experiments with CD8+ T cells (9).

[00596] While a considerable proportion of cells seemed to respond to the exposure to the antigen via expression *CD69* and *MYC* (**FIG. 52**), we found only a few cells that expressed *IFNG*, a signature cytokine for activated antigen specific cell. Most of the *IFNG* expressing cells were also among those cells carried the most total detected transcript molecules in the gene panel, an indication of active cell state, and belong to the effector memory/effector cell cluster (**FIG. 50B and 53**). We identified 5 out of 581 (0.86%) and 2 out of 2274 (0.09%) cells in donors 1 and 2 respectively that were likely to be CMV specific based on *IFNG* expression and overall transcription level. Among those cells, there was substantial amount of heterogeneity in terms of combinations and levels of effector molecules (*e.g.*, granzymes) and cytokines (*e.g.*, *IFNG*, *IL2*, *CCL3*, *CCL4*, *TNF*, *CSF2*, *IL4*) expressed (**FIG. 54**). Interesting, the single cell that expressed most transcripts in donor 2 expressed both *IL6* and *IL1B* but not *IFNG*.

Discussion

[00597] In this example, we presented highly scalable mRNA cytometry that used a recursive Poisson strategy to isolate single cells, to uniquely barcode cellular content, and to barcode individual molecules for quantitative analysis. We have shown that we may simultaneously identify and count transcript molecules belonging to each cell in a sample containing a few thousands cells. Further, we have demonstrated to use of this technique to characterize individual cells based on their expression profiles in naturally occurring heterogeneous systems, and detection of rare cells in a large background population.

[00598] The throughput and simplicity of CytoSeq presents a major advance over existing approaches involving microtiter plates or microfluidic chips for sequencing based measurement of gene expression of single cells. Because the experimental procedure is simple and reagent consumption per cell is low (in the nanoliter range), it enables one to

readily carry out single cell analysis for large number of cells across multiple conditions. In this study alone, we performed gene expression profiling of a total of ~14,600 single cell across 12 experiments, which would be costly and time-consuming if carried out by existing approaches. The number of cells measured by CytoSeq may be further scaled up simply by increasing the size of the microwell array and the library size of the barcoded beads, which is readily achieved by combinatorial synthesis. In addition, there is no restriction on the uniformity of cell sizes, thus allowing direct analysis of complex samples containing cells with a variety of cell sizes and shapes, such as PBMCs shown in this example, without any pre-sorting.

[00599] CytoSeq data resembled those of flow cytometry (FC), but with important differences. First, CytoSeq offers more versatility in terms of the number and type of gene products studied. Unlike flow cytometry that is confined mostly to a handful of surface proteins and requires optimally binding antibodies, CytoSeq allowed measurement of any transcribed mRNAs via nucleic acid amplification techniques. Optimal primer design and assay conditions enable us to routinely achieve ~88% mapped rate via multiplex PCR for an arbitrarily chosen panel of 100 or more genes (**Table 21**). Additionally, the entire transcriptome of each single cell in the sample may also be measured via universal amplification of the bead bound cDNA, although one has to be mindful with the relatively low efficiency of commonly used universal amplification techniques (7) and the high sequencing depth required for measuring the whole transcriptome across thousands of cells.

[00600] Second, in contrast to flow cytometry that relies on the kinetics of antibody binding, CytoSeq provides digital, absolute readout of gene expression level through molecular indexing. It has higher sensitivity and specificity to a single rare cell event because the detection was achieved by the co-expression of large number of genes specific to the rare cells. It therefore consumes much smaller amount of sample as compared to flow cytometry that requires certain number of events in order to form reliable clusters for gating.

[00601] Our data illustrates the importance of single cell versus bulk analysis. For instance, we showed scenarios where the most highly expressed genes in a sample of thousands of cells as whole were contributed by only one or a few cells. Most importantly, our experiments illustrate the importance of examining both large number of cells and large number of genes in single cell gene expression studies, an ability that is extremely limiting in prior approaches. The availability of such a tool for the routine measurement of expression across thousands of single cells in a biological sample may help accelerate the understanding of complex biological systems and drive novel applications in clinical diagnostics, such as

circulating tumor cell analysis and immune responses monitoring. We envision that our massive parallel single cell barcoding regime may also be adopted to measure the genome, as well as the genome and the transcriptome simultaneously, for studying single cell genome instability in areas such as cancer biology and neuroscience.

Table 21							
Experiment	total number of reads	number of reads with exactly 1 match to gene in panel	% reads aligned to one gene in the panel	number of reads with exact match to a cell barcode and alignment to one gene	% read after gene and barcode alignment	number of unique cell barcodes that satisfy filtering criteria	number of reads associated with those cell barcodes
K562 + Ramos	2399025	2154454	90%	1175715	49%	768	859470
Primary B+ Ramos	5711013	5203308	91%	3495392	61%	1198	2868577
PBMC	1270214	1105687	87%	803151	63%	632	670576
PBMC replicate	3927672	3468538	88%	2459367	63%	731	1920956
Donor 1 antiCD3/antiCD28 stimulated	3529898	3249998	92%	2122416	60%	3517	1466000
Donor 1 antiCD3/antiCD28 negative control	1557996	1292211	83%	939094	60%	1478	719351
Donor 2 antiCD3/antiCD28 stimulated	606865	552877	91%	403943	67%	669	246234
Donor 2 antiCD3/antiCD28 negative control	332951	283723	85%	205762	62%	595	86866
Donor 1 CMV stimulated	1064648	958410	90%	697057	65%	581	401629
Donor 1 CMV negative control	619957	547259	88%	406801	66%	253	192605
Donor 2 CMV stimulated	1902977	1692734	89%	1229667	65%	2274	688296
Donor 2 CMV negative control	1671419	1346637	81%	977344	58%	2337	715453

Synthesis of bead library

[00602] Beads were manufactured by Cellular Research, Inc. using a split-pool combinatorial approach. Briefly, twenty-micron magnetic beads functionalized with carboxyl groups were distributed into a 96 tubes containing oligos with 5' amine, followed by a universal sequence, first part of the cell label that is different for different tubes, and a linker sequence. The oligos were covalently coupled onto the beads by carbodiimide chemistry. Beads were pooled and split into a second set of 96 tubes containing oligos with a second

linker sequence on the 5' end, followed by the second part of the cell label that is different for different tubes, and complementary sequence to the first linker. Oligos on the beads were extended by DNA polymerase upon hybridization to oligos in solution via the first linker. Beads were pooled and split into a third set of 96 tubes containing oligos with oligo(dA) on the 5' end, followed by a randomer sequence that serves as the molecular label, the third part of the cell label, and a complementary sequence to the second linker. Oligos on the beads were extended by DNA polymerase upon hybridization to oligos in solution via the second linker. The final bead library has a size of 96x96x96 (884,736) cell labels.

Fabrication of microwell array

[00603] Microwell arrays were fabricated using standard photolithography. Arrays of pillars were patterned on photoresist on silicon wafer. PDMS was poured onto the wafer to create arrays of microwells. Replicas of the wafer were made with NOA63 optical adhesive using PDMS microwell array as template. Agarose (5%, type IX-A, Sigma) microwell arrays were casted from the NOA63 replica before each experiment.

Sample preparation

[00604] K562 and Ramos cells were cultured in RPMI-1640 with 10% FBS and 1x antibiotic-antimycotic. Primary B cells from a healthy donor were purchased from Sanguine Biosciences. PBMCs from a healthy donor were isolated from fresh whole blood in sodium heparin tube acquired from the Stanford Blood Center using Lymphoprep solution (StemCell).

T cell stimulation

[00605] Heparinized whole blood of two CMV seropositive blood donors was obtained from the Stanford Blood Center. For CMV stimulation, 1ml of whole blood was stimulated with CMV pp65 peptide pool diluted in PBS (Miltenyi Biotec) at a final concentration of 1.81 µg/ml for 6 hours at 37C. A separate aliquot of whole blood of each donor was incubated with PBS as negative controls. CD8⁺ T cells were isolated using RosetteSep cocktail (StemCell) and subsequently deposited onto microwell arrays. For anti-CD3/anti-CD28 stimulation, T cells from the same two donors were isolated from whole blood using RosetteSep T cell enrichment cocktail and resuspended in RPMI-1640 with 10% FBS and 1x antibiotic-antimycotic. One aliquot of cells from each donor was incubated with Dynabeads Human T-Activator CD3/CD28 (Life Technologies) at ~1:1 bead to cell ratio at 37C for 6 hours. A separate aliquot of cells from each donor were placed in incubator with no stimulation and served as negative control.

Single cell capture

[00606] Single cell suspension was pipetted on to the microwell array at a density of ~1 cell per 10 microwells. After washing to remove uncaptured cells, magnetic beads were loaded at a density of ~5 beads per well to saturate the microwell array. After washing to remove excess beads, cold lysis buffer (0.1M Tris-HCl pH 7.5, 0.5M LiCl, 1% LiSDS, 10mM EDTA, 5mM DTT) was pipetted over the surface of the microwell array. After 10 minutes of incubation on a slide magnet, beads were retrieved from the microwell array. Beads were collected in a microcentrifuge tube, and washed twice with wash A buffer (0.1M Tris-HCl, 0.5M LiCl, 1mM EDTA) and once with wash B buffer (20mM Tris-HCl pH 7.5, 50mM KCl, 3mM MgCl₂). From this point forward, all reactions were carried out in a single tube.

cDNA synthesis

[00607] Washed beads were resuspended in 40µL RT mix (1x First Strand buffer, 1µL SuperRase Inhibitor, 1µL SuperScript II or SuperScript III, 3mM additional MgCl₂, 1mM dNTP, 0.2ug/µL BSA) in a microcentrifuge tube placed on a rotor in a hybridization oven at temperatures 50C for 50 minutes (when using SuperScript III for the early experiment with K562 and Ramos cells) or 42C for 90 minutes (when using Superscript II for the rest of the experiments). Beads were treated with 1 µL of ExoI (NEB) in 20 µL of 1x ExoI buffer at 37 °C for 30 minutes, and 80 °C for 15 minutes.

Multiplex PCR and sequencing

[00608] Each gene panel contained two sets of gene specific primers designed by Primer3. A custom MATLAB script was written to select PCR primers such that there was minimal 3' end complementarity across the primers within the set. Primers in each panel are listed in **Table 21**. The amplification scheme is shown in **FIG. 55**. PCR were performed with the beads with KAPA Fast Multiplex Kit, with 50nM of each gene specific primer in the first primer set and 400nM universal primer, in a volume of 100µL or 200µL, with the following cycling protocol: 3min at 95C; 15 cycles of 15s at 95C, 60s at 60C, 90s at 72C; 5min at 72C. Magnetic beads were recovered and PCR products were purified with 0.7x Ampure XP. Half of the purified products were used for the next round of nested PCR with the second primer set using the same KAPA kit and cycling protocol. After clean up with 0.7x Ampure XP, 1/10th of the product was input into a final PCR reaction whereby the full-length Illumina adaptors were appended (1x KAPA HiFi Ready Mix, 200nM of P5, 200nM of P7. 95C 5min; 8 cycles of 98C 15s, 60C 30s, 72C 30s; 72C 5min).

Data analysis

[00609] Sequencing of library was performed on Illumina MiSeq instrument with 150x2 bp chemistry at a median depth of 1.6 million reads per sample. Sequencing revealed the cell label, the molecular label, and the gene of each read (**FIG. 55**). The assignment of gene of each read was done with the alignment software 'bowtie' (ref). The cell and molecular labels of each read were analyzed using custom MATLAB scripts. Reads were grouped first by cell label, then by gene and molecular label. To calculate the number of unique molecules per gene per cell, the molecular labels of reads with the same cell label and gene assignment were clustered. Edit distance greater than 1 base was considered as a unique cluster, and thus a unique transcript molecule. A table containing digital gene expression information of each cell was constructed for each sample - each row in the table represented a unique cell label, each column represented a gene, and each entry in the table represented the count of unique molecules within a gene per cell label. The table was filtered such that unique molecules that were sequenced only once (i.e. redundancy = 1) were removed. Subsequently, cells with a sum of unique molecules less than 10 or with co-expression of 4 or less genes in the panel were removed. The filtered table was then used for clustering analysis. Principal component analysis and hierarchical clustering was performed on log-transformed transcript count (with pseudocount of 1 added) with built-in functions in MATLAB.

[00610] References cited in Example 15, all of which are incorporated by reference in their entireties:

- [00611]** 1. A. K. Shalek *et al.*, Single-cell transcriptomics reveals bimodality in expression and splicing in immune cells. *Nature* 498, 236 (Jun 13, 2013).
- [00612]** 2. S. C. Bendall *et al.*, Single-cell mass cytometry of differential immune and drug responses across a human hematopoietic continuum. *Science* 332, 687 (May 6, 2011).
- [00613]** 3. A. R. Wu *et al.*, Quantitative assessment of single-cell RNA-sequencing methods. *Nature methods* 11, 41 (Jan, 2014).
- [00614]** 4. B. Treutlein *et al.*, Reconstructing lineage hierarchies of the distal lung epithelium using single-cell RNA-seq. *Nature* 509, 371 (May 15, 2014).
- [00615]** 5. S. Islam *et al.*, Characterization of the single-cell transcriptional landscape by highly multiplex RNA-seq. *Genome research* 21, 1160 (Jul, 2011).
- [00616]** 6. G. K. Fu, J. Hu, P. H. Wang, S. P. Fodor, Counting individual DNA molecules by the stochastic attachment of diverse labels. *Proceedings of the National Academy of Sciences of the United States of America* 108, 9026 (May 31, 2011).

[00617] 7. G. K. Fu, J. Wilhelmy, D. Stern, H. C. Fan, S. P. Fodor, Digital encoding of cellular mRNAs enabling precise and absolute gene expression measurement by single-molecule counting. *Analytical chemistry* 86, 2867 (Mar 18, 2014).

[00618] 8. K. Bratke, M. Kuepper, B. Bade, J. C. Virchow, Jr., W. Luttmann, Differential expression of human granzymes A, B, and K in natural killer cells and during CD8⁺ T cell differentiation in peripheral blood. *European journal of immunology* 35, 2608 (Sep, 2005).

[00619] 9. E. W. Newell, N. Sigal, S. C. Bendall, G. P. Nolan, M. M. Davis, Cytometry by time-of-flight shows combinatorial cytokine expression and virus-specific cell niches within a continuum of CD8⁺ T cell phenotypes. *Immunity* 36, 142 (Jan 27, 2012).

EXAMPLE 16: Development of Single Cell Quantification Protocol

[00620] FIG. 56 depicts a general workflow for the quantification of RNA molecules in a sample. In this example, the total number of RNA molecules in the sample was equivalent to the total number of RNA molecules in a single cell. As shown in Step 1 of FIG. 56, RNA molecules (110) were reverse transcribed to produce cDNA molecules (105) by the stochastic hybridization of a set of molecular identifier labels (115) to the polyA tail region of the RNA molecules. The molecular identifier labels (115) comprised an oligodT region (120), label region (125), and universal PCR region (130). The set of molecular identifier labels contained 960 different types of label regions.

Part I. Reverse Transcription and Labeling of RNA molecules

[00621] An RNA sample was prepared by mixing the following:

Genes	number of RNA molecules
Lys (spike-in control)	456
Phe (spike-in control)	912
Thr (spike-in control)	1824
Dap (spike-in control)	6840
Kan (spike-in control)	7352
Lymphocyte cell line RNA	10 pg (1 cell equivalent)
MS2 carrier (no polyA)	6×10^{11}

[00622] RNA molecules were labeled by preparing in an eppendorf tube a labeling mix as follows:

	Amount (μL)
RNA sample	2
ms2 RNA 1μg/μL	1
10mM dNTP	1
960 dT oligos pool (set#4) 10μM	0.4
water	9.1

[00623] Note: dT oligos pool (set #4) refers to the set of molecular identifier labels.

[00624] The molecular identifier labels were hybridized to the RNA molecules by incubation at 65°C for 5 minutes. The labeling mix was stored on ice for at least 1 minute.

[00625] The labeled RNA molecules were reverse transcribed by the addition of the reverse transcription mix as described below:

	Amount (μL)
5X first strand buffer	4
0.1M DTT	1
suprase-in 20u/μL	0.5
superscript III RT	1

[00626] Once the reverse transcription mix was added to the eppendorf tube containing labeling mix reaction, the reverse transcription reaction was conducted by incubating the sample at 37°C for 5 minutes, followed by incubation at 50°C for 30 minutes, and lastly incubation at 75°C for 15 minutes. Reverse transcription of the labeled RNA molecules produced labeled cDNA molecules (170).

[00627] Once the RNA molecules were reverse transcribed and labeled, excess oligos were removed from the sample by Ampure bead purification (Step 2 of FIG. 1). Ampure bead purification was performed by adding 20 μL of ampure beads to the eppendorf tube containing the reverse transcribed and labeled RNA molecules and incubating the tube at room temperature for 5 minutes, The beads were washed twice with 70% ethanol to remove the excess oligos. Once the excess oligos were removed by the ethanol washes, 20 μL of 10mM Tris was added to the tube containing the bead-bound labeled cDNA molecules.

[00628] As shown in Step 3 of FIG. 56, the labeled cDNA molecules (170) were amplified by multiplex PCR. Custom amplification of the labeled cDNA molecules was performed by using a custom forward primer (F1, 135 in FIG. 1) and a universal PCR primer (140). Table 23 lists the 96 different custom forward primers that were used to amplify 96 different genes to produce labeled amplicons (180) in a single reaction volume.

[00629] In order to optimize multiplex PCR reactions, 3 multiplex PCR reactions mixtures were prepared. Multiplex PCR reaction 1 was prepared as follows:

	Reaction 1
	Amount (μ L)
10X titanium	5
10mM dNTP	1.5
water	35.5
1 μ M each F1 primer pool	5
PCR004 10 μ M	1
purified cDNA	1
Titanium polymerase	1

[00630] The reaction condition for Multiplex PCR reaction was 1 cycle at 94°C for 2min, followed by 25 cycles of 94°C for 30 sec, 57°C for 60 sec, and 68°C for 1 min, then 1 cycle of 68°C for 7 min and 1 hold cycle at 4°C.

[00631] Multiplex PCR reactions 2 and 3 were prepared as follows:

	Reaction 2	Reaction 3
	Amount (μ L)	Amount (μ L)
2X Qiagen Multiplex mix	25	25
1 μ M each F1 primer pool	5	5
PCR004 10 μ M	1	1
Q solution		5
water	18	13
purified cDNA	1	1

[00632] The multiplex PCR reaction condition for Reactions 2 and 3 was 1 cycle at 95°C for 15min, followed by 25 cycles of 94°C for 30 sec, 57°C for 90 sec, and 72°C for 1 min, then 1 cycle of 68°C for 7 min and 1 hold cycle at 4°C.

[00633] The F1 primer pools contained the following primers:

F1 PCR Primers	Sequence
100611KanF2	CTGCCTCGGTGAGTTTTCTC
Lys_L_269	CTTCCCGTTACGGTTTTGAC
phe_L_177	AAAACCGGATTAGGCCATTA
thr_L_332	TCTCGTCATGACCGAAAAAG
dap_L_276	CAACGCCTACAAAAGCCAGT

[00634] Kan, Phe and Dap control genes were selectively amplified by nested PCR. Nested PCR amplification reactions were prepared as follows:

Multiplex PCR Rxn # →	1	2	3	1	2	3	1	2	3
PCR Rxn # →	1	2	3	4	5	6	7	8	9
	μ L	μ L	μ L	μ L	μ L	μ L	μ L	μ L	μ L
10x Taq	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5
10mM dNTP	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5
water	22.25	22.25	22.25	22.25	22.25	22.25	22.25	22.25	22.25

Cy3 PCR004 10 μ M	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5
KanF3_5P 5 μ M	1	1	1						
Phe_L_215 5 μ M				1	1	1			
Dap_L_290 5 μ M							1	1	1
Multiplex PCR Rxn	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5
USB tag	0.25	0.25	0.25	0.25	0.25	0.25	0.25	0.25	0.25

[00635] Note: The multiplex PCR reaction used for PCR reactions 1, 4, and 7 was multiplex PCR reaction # 1. The multiplex PCR reaction used for PCR reactions 2, 5 and 8 was multiplex PCR reaction # 2. The multiplex PCR reaction used for PCR reactions 3, 6 and 9 was multiplex PCR reaction # 3.

[00636] The primers used for nested PCR are disclosed as follows:

Nested PCR primer	Sequence
KanF4_5P	/5Phos/GTGGCAAAGCAAAAGTTCAA
Phe_L_215	TGAGAAAGCGTTTGATGATGTA
Dap_L_290	GCCAGTTTATCCCGTCAAAG

[00637] The PCR amplification reaction condition for Reactions 1-9 was 1 cycle of 94 °C for 2 min, 30 cycles of 94 °C for 20 sec, 55 °C for 20 sec, and 72 °C for 20 sec, then 1 cycle at 72 °C for 4 min and 1 hold cycle at 4 °C.

[00638] The 4 μ L of PCR products of PCR amplification reactions 1-9 were run on an agarose gel. As shown in FIG. 58A, Reactions 1-3 showed the presence of the Kan control gene, Reactions 4-6 showed the presence of the Phe control gene, and Reactions 7-9 showed the presence of the Dap control gene.

[00639] The PCR products from PCR reactions 1-9 were prepared for hybridization to an Applied Microarray Inc. (AMI) array. Hybridization mixtures were prepared as follows:

	μ L
PCR product	20
Wash A (6X SSPE+0.01% Triton X-100)	55
Cy3 Oligo (760pM)	1

[00640] The hybridization mixtures 1-9, corresponding to the mixtures containing PCR products from PCR reactions 1-9, respectively, were denatured at 95°C for 5 minutes and then placed at 4°C. The hybridization mixtures were transferred to an AMI array slide and incubated overnight at 37°C.

[00641] After the overnight hybridization, the AMI array slide was washed and then scanned. Theoretical and actual measurements and percent accuracy are depicted below:

Hybridization mixtures # →	1	2	3	4	5	6	7	8	9
Multiplex PCR condition (Rxn #)	Titani um (1)	Qiagen (2)	Qiagen +Q (3)	Titani um (1)	Qiage n (2)	Qiagen +Q (3)	Titani um (1)	Qiage n (2)	Qiagen +Q (3)
Theoretical measurement	3676 (bioanalyzer)			912			6840		
Actual measurement	1826	1740	2116	299	235	251	1165	1077	172
% detection	49.7	47.3	57.6	32.8	25.8	27.5	17	15.7	2.5

[00642] Note: The theoretical measurement is based on detection of 100% of the Kan, Phe and Dap control genes.

[00643] PCR products from Reaction 2 were purified by Ampure purification. Ampure purification was performed as follows:

	μL
F1 PCR products from X01 sample 2	30
Ampure beads	30

[00644] Ampure purification reactions were incubated at room temperature for 5 min and then washed in 70% ethanol. Purified PCR products were eluted from the beads in 30 μL of water. The concentration of the PCR products was 6 ng/ μL as determined by a Nanodrop spectrometer.

Part II: Library Preparation Protocol

[00645] PCR products purified from the X01 sample 2 (see Example 1) were used to prepare a DNA library. An F2 primer pool was created from mixing the following primers:

F2 PCR Primers	Sequence
Lys_L_269	CTTCCCGTTACGGTTTTGAC
phe_L_177	AAAACCGGATTAGGCCATTA
thr_L_332	TCTCGTCATGACCGAAAAAG
dap_L_276	CAACGCCTACAAAAGCCAGT

[00646] An F2 primer mix was prepared by mixing the following

F2 primer mix	μL
water	750
F2 primer pool 1uM each/100uM total	100

[00647] The F2 primer mix was incubated at 95°C for 3 min and then stored on ice. The following ligation mix was added to the F2 primer mix to produce an F2 primer ligation mix:

Ligation mix	μL
10X DNA ligase buffer NEB	100
T4 PNK USB	50

[00648] The F2 primer ligation mix was incubated at 37°C for 1 hour, followed by an incubation at 65°C for 20 min. The F2 PCR primers were ethanol precipitated and the concentration of the primer pool was determined by a Nanodrop spectrophotometer. The F2 primer pool was resuspended to produce a final concentration of 1uM each/100uM total.

[00649] As shown in Step 4 of FIG. 56, the labeled amplicons (180) were amplified by multiplex PCR. 96 different custom forward primers (F2, 145 in FIG. 1) and a universal PCR primer (140) were used to amplify the labeled amplicons (X01 sample 2 from Example 1) in a single reaction volume. Table 24 lists the 96 different custom forward primers.

[00650] The multiplex PCR reaction was prepared as follows:

Multiplex PCR mix	μL
2X Qiagen Multiplex mix	25
1μM each F2 primer pool kinase	5
PCR004 5'P 10μM	1
water	18
purified first PCR X01 sample 2	1

[00651] The multiplex PCR condition was 1 cycle at 95°C for 15min, followed by 18 cycles of 94°C for 30 sec, 57°C for 90 sec, and 72°C for 1 min, then 1 cycle of 68°C for 7 min and 1 hold cycle at 4°C. The multiplexed amplicons were purified by Ampure purification and eluted with 50 μL of water. The concentration of the amplicons was determined to be 30 ng/μL by a Nanodrop spectrophotometer. 5 μL of the amplicons was run on an agarose gel (FIG. 58B).

As shown in Step 5 of FIG. 56, adaptors (150, 155) were ligated to the labeled amplicons (180) to produce adaptor labeled amplicons (190). Adaptor labeled amplicons were produced as follows:

Adaptor mix	μL
10X T4 ligase USB	10
water	60
purified nested PCR product	10
annealed, pooled 96 ABC adaptors 50μM	10
T4 DNA ligase (3μl neb hc, 7 μl usb)	10

[00652] The adaptor mix was incubated at 16°C for 4 hours. The adaptor labeled amplicons were purified by Ampure purification and eluted in 20 μL of 10mM Tris.

[00653] The purified adaptor labeled amplicons were gap-repaired and PCR amplified as follows:

Fill-in and PCR mix	μL
10x thermoPol buffer	5
10mM dNTP	1.5
water	32
CR P1 10μM	3
CR IDX D1 10μM	3
purified adaptor labeled amplicons	5
Vent exo- 2u/μL	0.5

[00654] The PCR condition was 1 cycle of 72°C for 2 min, followed by 94°C for 1 min, 12 cycles of 94°C for 15 sec, 60°C for 15 sec and 72°C for 30 sec, 1 cycle of 72°C for 4 min and 1 hold cycle at 4°C. The PCR products were purified by Ampure purification and eluted in 30 μL of TE. The concentration of the purified PCR product was 22ng/ μL (83nM) as determined by Nanodrop spectroscopy. 5 μL of the PCR purified products were run on a 1% agarose gel (FIG. 58B)

[00655] *Part III. Sequencing of the adaptor labeled amplicon library*

[00656] The adaptor labeled amplicon library was sequenced using a MiSeq Sequencer.

[00657] A sequence mapping summary is shown below:

	Require Perfect Match	Allow 1bp mismatch
Total Read Pairs	7,724,955	
# of RNA with universal primer and polyA match	2,499,444 (32%)	4,716,378 (61%)
# of RNA mapped to targets (96 genes)	2,373,700	4,489,485

[00658] As shown in the sequence mapping summary above, many reads were lost due to the stringent polyA matching criteria. FIG. 59 shows the reads and counts across all detected genes.

[00659] Sequencing reads were also used to quantify specific genes. FIG. 61-62 depict a plot of the reads observed per label detected (RPLD) for various genes. Conventional rpkm values are also shown in the plots depicted in FIG. 61-62. FIG. 59 summarizes a comparison of RPLD and RPKM for various genes.

[00660] FIG. 63 depicts a plot of total reads (labels) versus rpld for various genes.

[00661] The data represented in FIGS. 4, 7 and 8 are also shown in numerical form in Table 25.

[00662] FIG. 64 depicts a plot of RPKM for undetected genes.

[00663] The quantity of the spike-in controls in the adaptor labeled amplicon library was determined by MiSeq sequencing. Results from MiSeq sequencing of the spike-in controls are shown in the table below.

Spike-in Control	input N (mfg)	Reads	Labels (K)
Dap	6840	1,920,503	893
Phe	912	470,738	859
Thr	1824	410,664	847
Lys	456	282,174	847
Kan	7352	24	23

[00664] In the table above, input N refers to the original number of the spike-in control; Reads refers to the total number of read pairs; and Labels (K) refers to the number of different labels detected by sequencing. FIG. 60A-D depicts a plot of the reads observed per label detected (RPLD) for Lys, Phe, Thr, and Dap spike-in controls, respectively. FIG. 60E depicts a plot of Reads versus Input.

Table 23	
Name	Sequence
NM_144646.3F1	TTGACTTTGCCTTGGAGAGC
NR_015342.1F1	TTTTTCTTACAGTGTCTTGGCATA
NM_000193.2F1	CGTGACCCTAAGCGAGGAG
NM_001777.3F1	TTTGCAGTGATTTGAAGACCA
NM_000600.3F1	GGCATTCCTTCTTCTGGTCA
NM_021127.2F1	CTGGGCTATATACAGTCCTCAAA
NM_004318.3F1	GGGGTGATTATGACCAGTTGA
NM_002467.4F1	TGCATGATCAAATGCAACCT
NM_001773.2F1	TCTTCCGAAAAATCCTCTTCC
NM_001770.5F1	CTGGGGTCCCAGTCCTATG
NM_001718.4F1	TGTACTGGGAAGGCAATTTCA
NR_023920.1F1	GAGCCGCTGGGGTACTC
NM_000267.3F1	CAGTTAGTTGCTGCACATGGA
NM_000633.2F1	TTGCATTTCTTTTGGGGAAG
NM_000314.4F1	GTCATGCATGCAGATGGAAG
NM_021151.3F1	GCTGCAGTGAGCTGTGATGT
NM_002415.1F1	GTTCTCTCCGAGCTCACC
NM_004985.3F1	TCCGAAAGTTTCCAATTCCA
NM_005375.2F1	TTGTTTGGGAGACTCTGCATT
NM_000555.3F1	GACCCCACTTGGACTGGTAG
NM_001668.3F1	GTGATCTTGATTGCGGCTTT
NM_025237.2F1	GGGGGAAAAACTACAAGTGC
NM_021117.3F1	TGATTCCTTTTCCTGCCTGT
NM_016316.2F1	AAAAACCTCCAGGCCAGACT
NM_021975.3F1	AATCAAAATAACGCCCCAGA
NM_004333.4F1	TTGCTAAAAATTGGCAGAGC
NM_001621.4F1	TTGTTAAGTGCCAAACAAAGGA
NM_005239.5F1	AAGCTGGGAAGAGCAAAGC
NM_000485.2F1	AGGACAGAGGGTGGTCGTC
NM_004048.2F1	TGAGTGCTGTCTCCATGTTG

NM_001657.2F1	CCTCACAGCTGTTGCTGTTATT
NM_012238.4F1	AAAACACCCAGCTAGGACCA
NM_002055.4F1	AACTGAGGCACGAGCAAAGT
NM_002392.4F1	GCTTTATGGGTGGATGCTGA
NM_001625.3F1	ATAATATCGCCAGCCTCAGC
NM_002110.3F1	TCCAGAGTGTGCTGGATGAC
NM_002943.3F1	TGCAAGCCATTTATGGGAAT
NM_000059.3F1	TGGAATGAGGTCTCTTAGTACAGTT
NM_018136.4F1	TCCCAGAAACACCTGTAAGGA
NM_003467.2F1	TGTCTAGGCAGGACCTGTGG
NM_004958.3F1	AGTGATGCTGCGACTCACAC
NM_006139.3F1	GGCTCAGAAAGTCTCTCTTTCC
NM_002693.2F1	CTCCCAAACCTCAGGCTTTCA
NM_001080432.2F1	AAAGCGCTGGGATTACAGG
NM_005954.2F1	CGTCCAGTTGCTTGGAGAAG
NM_024865.2F1	AATAACCTTGGCTGCCGTCT
NM_001905.2F1	GGGAATTCTCAGTGCCAACT
NM_002046.4F1	GCATCCTGGGCTACACTGAG
NM_002253.2F1	TGCTGGGAACAATGACTATAAGA
NM_002356.5F1	GCCTAAAACACTTTGGGTGGT
NM_000189.4F1	GGGTGCCCACAAAATAGAGA
NM_000546.5F1	GAGACTGGGTCTCGCTTTGT
NM_152860.1F1	TGGGGAAGGCTTTCTCTAGG
NM_016231.4F1	TTCAACTTGAGTGATCTGAGCTG
NM_000518.4F1	TATGGGCAACCCTAAGGTGA
NM_000905.3F1	CGCTGCGACACTACATCAAC
NM_005038.2F1	TGGAGTCTTGCTCTGTCACC
NM_000041.2F1	ACGAGGTGAAGGAGCAGGT
NM_005957.4F1	CGATGCCTTTGGGTAGAGAG
NR_002785.2F1	ACTGATCGTCCAAGGACTGG
NM_000321.2F1	AAAAAGAAATCTGGTCTTGTTAGAAAA
NM_152756.3F1	TTGAAAAGTGGTAAAGGAATTGTGA

NM_000610.3F1	CACCAAGAATTGATTTTGTAGCC
NR_033314.1F1	AAAAATGGGGGAAAATGGTG
NM_017460.5F1	CATGGTTGAAACCCCATCTC
NR_002196.1F1	TTCAAAGCCTCCACGACTCT
NM_000591.3F1	GCTGGAACAGGTGCCTAAAG
NM_000106.5F1	CCCTAAGGGAACGACACTCA
NM_138712.3F1	ACCTGCTACAAGCCCTGGA
NM_004304.4F1	GGATCCCTAAGACCGTGGAG
NM_000754.3F1	CCACCTCAGAGGCTCCA
NM_000492.3F1	TGCTGTATTTTAAAAGAATGATTATGA
NM_000444.4F1	GTAGCTGGGACGCTGGTTTA
NM_002463.1F1	ATTCCCTTCCCCCTACAAGA
NM_000552.3F1	CCTGAGTGCAACGACATCAC
NM_005430.3F1	GGGGGAACCAGCAGAAAT
NM_003150.3F1	GACCTAGGGCGAGGGTTC
NM_000388.3F1	AATTCCTGAAGCCAGATCCA
NM_007294.3F1	AAAATGTTTATTGTTGTAGCTCTGG
NM_005933.3F1	TTTCAAGAGCTCAACAGATGACA
NM_002343.3F1	GACTGCCCGGACAAGTTTT
NM_000376.2F1	GAGAAGGTGCCCCAAAATG
NM_002462.3F1	AGCCACTGGACTGACGACTT
NM_021005.3F1	GGAGGACTAGTGAGGGAGGTG
NM_012343.3F1	GGCAAGTGATGTGGCAATTA
NM_001741.2F1	GTTGGAGCACCTGGAAAGAA
NM_014417.4F1	ATGCCTGCCTCACCTTCAT
NM_014009.3F1	ACAGGGGCACTGTCAACAC
NM_006908.4F1	AAAAATCATGTGTTGCAGCTTT
NM_005228.3F1	TGCTTTCACAACATTTGCAG
NM_013994.2F1	AATGTTTCCTTGTGCCTGCT
NM_000639.1F1	ATATCCTGAGCCATCGGTGA
NM_002701.4F1	TTTTGGTACCCAGGCTATG
NM_000268.3F1	ACCCCGTGGCATTACATAAC

NM_003140.1F1	CTTCCAGGAGGCACAGAAAT
NM_000551.3F1	CTAACCTGGGCGACAGAGTG

Table 24

Name	Sequence
NM_144646.3F2	ATATTTGGACATAACAGACTTGGAA
NR_015342.1F2	TGCTGACTTTTAAAATAAGTGATTCTG
NM_000193.2F2	GCGGCAGAGTAGCCCTAAC
NM_001777.3F2	TGGGCTATTTCTATTGCTGCT
NM_000600.3F2	AATGGAAAGTGGCTATGCAG
NM_021127.2F2	GGTTGTAGTCACTTTAGATGGAAAA
NM_004318.3F2	TTTGTTTGACTTTGAGCACCA
NM_002467.4F2	AATGTTTCTCTGTAAATATTGCCATT
NM_001773.2F2	CACCCCCATATGGTCATAGC
NM_001770.5F2	AGCACCAGGTGATCCTCAG
NM_001718.4F2	TGTTTTGCTGTAACATTGAAGGA
NR_023920.1F2	TAATGCCACAGTGGGGATG
NM_000267.3F2	GGGCCTAAACTTTGGCAGTT
NM_000633.2F2	TTTACCTTCCATGGCTCTTTT
NM_000314.4F2	GCCTTACTCTGATTCAGCCTCTT
NM_021151.3F2	CGTAACAAAATTCATTGTGGTGT
NM_002415.1F2	AGAACCGCTCCTACAGCAAG
NM_004985.3F2	GTGCTTTCTTTTGTGGGACA
NM_005375.2F2	GGGAGTTCTGCATTTGATCC
NM_000555.3F2	TGGGTCAGAGGACTTCAAGG
NM_001668.3F2	AGGGTTCTGATCACATTGCAC
NM_025237.2F2	CTGCAGGACTGGTCGTTTTT
NM_021117.3F2	AGGGCAGGGTAGAGAGGGTA
NM_016316.2F2	TTCTTCCATGCGGAGAAATC
NM_021975.3F2	CATGGCTGAAGGAAACCAGT
NM_004333.4F2	TTGCCAGCTATCACATGTCC
NM_001621.4F2	TCTTTTCCTGTACCAGGTTTTTC

NM_005239.5F2	TGACTGGGAACATCTTGCTG
NM_000485.2F2	TGGCACCTGTACCCTTCTTC
NM_004048.2F2	TTCAATCTCTTGCACTCAAAGC
NM_001657.2F2	TGGAGTCACTGCCAAGTCAT
NM_012238.4F2	TTTGCATGATGTTTGTGTGC
NM_002055.4F2	GCACCCACTCTGCTTTGACT
NM_002392.4F2	ACCATGTAGCCAGCTTTCAA
NM_001625.3F2	GCAACTGGGCATGAGTACCT
NM_002110.3F2	CCACACCCCCTTCCTACTC
NM_002943.3F2	AGTCTGCTTATTTCCAGCTGTTT
NM_000059.3F2	TCCTGTTCAAAAGTCAGGATGA
NM_018136.4F2	AAATCACAAATCCCCTGCAA
NM_003467.2F2	CTGAACATTCCAGAGCGTGT
NM_004958.3F2	CAGTGGGACCACCCTCACT
NM_006139.3F2	TCTGTAGATGACCTGGCTTGC
NM_002693.2F2	TCAGAACCAAGATGCCAACA
NM_001080432.2F2	CATGACCCAGCCTATGGTTT
NM_005954.2F2	ACCTCCTGCAAGAAGAGCTG
NM_024865.2F2	TTGGGAGGCTTTGCTTATTTT
NM_001905.2F2	CTGGGAAACACTCCTTGCA
NM_002046.4F2	CAACGAATTTGGCTACAGCA
NM_002253.2F2	CAAAGGTCATAATGCTTTCAGC
NM_002356.5F2	TTTGACGTATCTTTTCATCCAA
NM_000189.4F2	TGTTGTTGGTTTCCAAAAAGG
NM_000546.5F2	GCCAACTTTTGCATGTTTTG
NM_152860.1F2	CCCAAGCTGATCTGGTGGT
NM_016231.4F2	TGCTGTGAAAGAAACAAACATTG
NM_000518.4F2	GCACGTGGATCCTGAGAACT
NM_000905.3F2	CCAGCCCAGAGACACTGATT
NM_005038.2F2	CACGCCCAGCTAATTTTTGT
NM_000041.2F2	CCTGGTGGAAGACATGCAG
NM_005957.4F2	TCACACCTGTAATCCCAGCA

NR_002785.2F2	CAGAGCTCCGCCTCATTAGT
NM_000321.2F2	TCCATTTTCATCATTGTTTCTGC
NM_152756.3F2	TGGTGTTTGTAGGTCACCTGAACA
NM_000610.3F2	AACATGGTCCATTACCTTTATG
NR_033314.1F2	AGAGCGAGACTCCGTCTCAA
NM_017460.5F2	AGTGAGCTGAGATTGCACCA
NR_002196.1F2	AGACGGCCTTGAGTCTCAGT
NM_000591.3F2	GGGAATCCCTTCCTGGTC
NM_000106.5F2	CTTCCTGCCTTTCTCAGCAG
NM_138712.3F2	TGCAGGTGATCAAGAAGACG
NM_004304.4F2	GGTTTGTAGCATGGGTTCAT
NM_000754.3F2	CCAGCCCACTCCTATGGAT
NM_000492.3F2	AAACTGGGACAGGGGAGAAC
NM_000444.4F2	TTTGGGTAGGTGACCTGCTT
NM_002463.1F2	TCACTGAACGAATGAGTGCTG
NM_000552.3F2	ACGATGTGCAGGACCAGTG
NM_005430.3F2	AATTTGCACTGAAACGTGGA
NM_003150.3F2	CTGTTGTGGCCATTAAAGAA
NM_000388.3F2	TTCCCTCCAGCAGTGGTATT
NM_007294.3F2	CACCAGGAAGGAAGCTGTTG
NM_005933.3F2	TTTCCTTGTGTTCTTCCAAGC
NM_002343.3F2	TCGCAGGCATTACTAATCTGAA
NM_000376.2F2	CTCTGGCTGGCTAACTGGAA
NM_002462.3F2	AGAGCCCCACCCTCAGAT
NM_021005.3F2	TGTGCAGAGTTCTCCATCTGA
NM_012343.3F2	TGCCTGTTACAAATATCAAGGAA
NM_001741.2F2	TTTCCCTTCTTGCATCCTTC
NM_014417.4F2	TGTGACCACTGGCATTTCATT
NM_014009.3F2	CTCACACACACGGCCTGTTA
NM_006908.4F2	CACTTGACCAATACTGACCCTCT
NM_005228.3F2	GTGTGTGCCCTGTAAACCTGA
NM_013994.2F2	CCACTTCCCACTTGCAGTCT

NM_000639.1F2	TGTGTGTGTGTGTGTGTGTGTGT
NM_002701.4F2	TCTCCCATGCATTCAAACCTG
NM_000268.3F2	TCTAAGTGTTCTCACTGACAGG
NM_003140.1F2	TACTCTGCAGCGAAGTGCAA
NM_000551.3F2	CCAAGATCACACCATTGCAC
NM_144646.3F2	ATATTTGGACATAACAGACTTGGAA

Table 25			
Gene	Number of reads	Number of labels	reads per kb/million (RPKM)
APOE	1585	408	0.2
APRT	11280	56	103.4
AREG	147	102	0.0
ASPM	4683	53	4.4
B2M	209362	698	3891.7
BBC3	8	1	0.0
BCL2	3627	27	33.2
BDNF	12778	116	0.3
CD19	38	5	43.1
CD44	6789	47	8.1
COMT	2828	16	10.7
CTPS1	3998	15	25.4
CXCR4	10547	54	19.2
CYP3A4	80982	267	0.1
DCX	28	24	0.0
ETS2	6	5	0.0
FASLG	3182	565	0.2
FTO	8877	58	11.3
GAPDH	227129	661	3870.8
HCK	294	2	2.4
HK2	593	2	12.7
IGJ	119449	454	438.9
KDR	2	2	0.0
KRAS	64	31	6.8
LTF	126	90	0.0
MARCKS	1563	12	36.9
MIF	17775	89	760.4
MLL	72	9	2.6
MTHFR	4854	282	3.9
MX1	100701	285	119.0
MX2	2145	13	45.2
MYB	18361	100	2.8
MYC	6859	27	130.5
NF1	4	1	3.7
NNT	15673	78	14.1

PMAIP1	50604	244	126.9
POLG	5163	46	7.4
POU5F1	1924	12	1.0
PPID	27354	303	39.0
PTEN	20884	109	51.6
RAC1	12454	67	44.8
RB1	1420	14	46.1
RELA	3893	26	17.9
RICTOR	898	5	5.2
RORA	954	7	0.1
SOST	1	1	0.0
SP7	1	1	0.0
STAT3	706	28	16.9
TP53	900	34	14.6
VHL	11576	106	0.0

[00665] While preferred embodiments of the present invention have been shown and described herein, it will be obvious to those skilled in the art that such embodiments are provided by way of example only. Numerous variations, changes, and substitutions will now occur to those skilled in the art without departing from the invention. It should be understood that various alternatives to the embodiments of the invention described herein may be employed in practicing the invention.

[00666] The publications discussed herein are provided solely for their disclosure prior to the filing date of the present application. Nothing herein is to be construed as an admission that the present invention is not entitled to antedate such publication by virtue of prior invention. Further, the dates of publication provided may be different from the actual publication dates which may need to be independently confirmed.

EMBODIMENTS

[00667] Disclosed herein are methods for analyzing molecules in two or more samples. The method may comprise: a) producing a plurality of sample-tagged nucleic acids by: i) contacting a first sample comprising a plurality of nucleic acids with a plurality of first sample tags to produce a plurality of first sample-tagged nucleic acids; and ii) contacting a second sample comprising a plurality of nucleic acids with a plurality of second sample tags to produce a plurality of second sample-tagged nucleic acids, wherein the plurality of second sample tags are different from the first sample tags; b) contacting the plurality of sample-tagged nucleic acids with a plurality of molecular identifier labels to produce a plurality of labeled nucleic acids; and c) detecting at least one of the labeled nucleic acids, thereby determining a count of a plurality of nucleic acids in a plurality of samples. One or more of

the plurality of samples may comprise a single cell or cell lysate. One or more of the plurality of samples may consist of a single cell. The sample tag may comprise a cellular label that identifies the cell from which the labeled nucleic acids originated from. The plurality of samples consisting of a single cell may be from one or more sources. The sample tag may comprise a sample index region that identifies the source of the single cell. The molecular identifier labels may be referred to as a molecular label. One or more of the plurality of samples may comprise fewer than 1,000,000 cells. One or more of the plurality of samples may comprise fewer than 100,000 cells. One or more of the plurality of samples may comprise fewer than 10,000 cells. One or more of the plurality of samples may comprise fewer than 1,000 cells. One or more of the plurality of the samples may comprise fewer than 100 cells. One or more of the plurality of samples may comprise a cell lysate.

[00668] Alternatively, the method for analyzing molecules in a plurality of samples may comprise: a) producing a plurality of labeled nucleic acids comprising: i) contacting a first sample with a first plurality of sample tags, wherein the first plurality of sample tags comprise identical nucleic acid sequences; ii) contacting the first sample with a first plurality of molecular identifier labels comprising different nucleic acid sequences, thereby producing a plurality of first-labeled nucleic acids; iii) contacting a second sample with a second plurality of sample tags, wherein the second plurality of sample tags comprise identical nucleic acid sequences; iv) contacting the second sample with a second plurality of molecular identifier labels comprising different nucleic acid sequences, thereby producing a plurality of second-labeled nucleic acids, wherein the plurality of labeled nucleic acids comprises the plurality of first-labeled nucleic acids and the second-labeled nucleic acids; and b) determining a number of different labeled nucleic acids, thereby determining a count of a plurality of nucleic acids in a plurality of samples. The sample tag may comprise a cellular label that identifies the cell from which the labeled nucleic acids originated from. The sample tag may comprise a sample index region that identifies the source of the single cell. The molecular identifier labels may be referred to as a molecular label.

[00669] Alternatively, the method for analyzing molecules in a plurality of samples may comprise: a) contacting a plurality of samples comprising two or more different nucleic acids with a plurality of sample tags and a plurality of molecular identifier labels to produce a plurality of labeled nucleic acids, wherein: i) the plurality of labeled nucleic acids comprise two or more nucleic acids attached to two or more sample tags and two or more molecular identifier labels; ii) the sample tags attached to nucleic acids from a first sample of the plurality of samples are different from the sample tags attached to nucleic acid molecules

from a second sample of the plurality of samples; and iii) two or more identical nucleic acids in the same sample are attached to two or more different molecular identifier labels; and b) detecting at least a portion of the labeled nucleic acids, thereby determining a count of two or more different nucleic acids in the plurality of samples. The sample tag may comprise a cellular label that identifies the cell from which the labeled nucleic acids originated from. The sample tag may comprise a sample index region that identifies the source of the single cell. The molecular identifier labels may be referred to as a molecular label.

[00670] Further disclosed herein are methods for analyzing molecules in a plurality of samples comprising: a) contacting a first plurality of molecules from a first sample of a plurality of samples with a first set of molecular barcodes to produce a first plurality of labeled molecules, wherein a molecular barcode of the first plurality of molecular barcodes comprises a label region and a sample index region; b) contacting a second plurality of molecules from a second sample of the plurality of samples with a second set of molecular barcodes to produce a second plurality of labeled molecules, wherein a molecular barcodes of the second plurality of molecular barcodes comprises a label region and a sample index region, and wherein the first plurality of molecular barcodes and the second plurality of molecular barcodes differ at least by the sample index region of the molecular barcodes; and c) detecting at least a portion of two or more molecules of the first plurality of labeled molecules and at least a portion of two or more molecules of the second plurality of labeled molecules, thereby determining a count of the two or more molecules in the plurality of samples. The first plurality of molecules may comprise nucleic acid molecules. The second plurality of molecules may comprise nucleic acid molecules. The label region may be referred to as a molecular label. The molecular barcode may further comprise a cellular label. In instances in which a sample of the plurality of samples consists of a single cell, the sample index region may refer to the cellular label.

[00671] Disclosed herein is a method of selecting a custom primer, comprising: a) a first pass, wherein primers chosen comprise: i) no more than three sequential guanines, no more than three sequential cytosines, no more than four sequential adenines, and no more than four sequential thymines; ii) at least 3, 4, 5, or 6 nucleotides that are guanines or cytosines; and iii) a sequence that does not easily form a hairpin structure; b) a second pass, comprising: i) a first round of choosing a plurality of sequences that have high coverage of all transcripts; and ii) one or more subsequent rounds, selecting a sequence that has the highest coverage of remaining transcripts and a complementary score with other chosen sequences of no more

than 4; and c) adding sequences to a picked set until a coverage saturates or a total number of customer primers is less than or equal to about 96.

[00672] Further disclosed herein is a method for producing a labeled molecule library comprising: a) producing a plurality of sample-tagged nucleic acids by: i) contacting a first sample comprising a plurality of nucleic acids with a plurality of first sample tags to produce a plurality of first sample-tagged nucleic acids; and ii) contacting a second sample comprising a plurality of nucleic acids with a plurality of second sample tags to produce a plurality of second sample-tagged nucleic acids, wherein the plurality of first sample tags are different from the second sample tags; and b) contacting the plurality of sample-tagged nucleic acids with a plurality of molecular identifier labels to produce a plurality of labeled nucleic acids, thereby producing a labeled nucleic acid library.

[00673] Disclosed herein are kits for use in analyzing molecules in a plurality of samples. The kit may comprise: a) two or more sets of molecular barcodes, wherein a molecular barcode of the set of one or more molecular barcodes comprise a sample index region and a label region, wherein (i) the sample index regions of the molecular barcodes of a set of molecular barcodes is the same; and (ii) the sample index regions of a first set of molecular barcodes are different from the sample index regions of a second set of molecular barcodes; and b) a plurality of beads. The two or more sets of molecular barcodes may be attached to the plurality of beads. The two or more sets of molecular barcodes may be conjugated to the bead. The label region may be referred to as a molecular label. The molecular barcode may further comprise a cellular label. In instances in which a sample of the plurality of samples consists of a single cell, the sample index region may refer to a cellular label.

[00674] The kit for analyzing molecules in a plurality of samples may comprise: a) a first container comprising a first plurality of molecular barcodes, wherein: (i) a molecular barcode comprises a sample index region and a label region; (ii) the sample index regions of at least about 80% of the total number of molecular barcodes of the first plurality of molecular barcodes are identical; and (iii) the label regions of two or more barcodes of the first plurality of molecular barcodes are different; and (b) a second container comprising a second plurality of molecular barcodes, wherein: (i) a molecular barcode comprises a sample index region and a label region; (ii) the sample index regions of at least about 80% of the total number of molecular barcodes of the first plurality of molecular barcodes are identical; and (iii) the label regions of two or more barcodes of the first plurality of molecular barcodes are different; wherein the sample index regions of the first plurality of molecular barcodes is different from the sample index regions of the second plurality of molecular barcodes. . The label region

may be referred to as a molecular label. The molecular barcode may further comprise a cellular label. In instances in which a sample of the plurality of samples consists of a single cell, the sample index region may refer to a cellular label.

[00675] Alternatively, the kit for analyzing molecules in a plurality of samples comprises: a) a first container comprising a first plurality of sample tags, wherein the plurality of sample tags comprises the same nucleic acid sequence; and b) a second container comprising a first plurality of molecular identifier labels, wherein the plurality of molecular identifier labels comprises two or more different nucleic acid sequences. The label region may be referred to as a molecular label. In instances in which a sample of the plurality of samples consists of a single cell, the sample tag may refer to a cellular label. The kit may further comprise a third container comprising a first plurality of cellular labels, wherein the plurality of cellular labels comprises two or more different nucleic acid sequences.

[00676] The kits and methods disclosed herein may comprise one or more sets of molecular barcodes. The kits and methods disclosed herein may comprise one or more molecular barcodes. The molecular barcodes may comprise a sample index region, molecular label region, cellular label region, or a combination thereof. At least two molecular barcodes of a set of molecular barcodes may comprise two or more different label regions. Label regions of two or more molecular barcodes of two or more sets of molecular barcodes may be identical. Two or more sets of molecular barcodes may comprise molecular barcodes comprising the same label region. In instances in which a sample of the plurality of samples consists of a single cell, the sample tag may refer to a cellular label.

[00677] The molecular barcodes disclosed herein may comprise a sample index region. The sample index region of molecular barcodes of two or more sets of molecular barcodes may be different. The sample index region may comprise one or more nucleotides. Two or more sequences of sample index regions of two or more different sets of molecular barcodes may be less than about 90%, 85%, 80%, 75%, 70%, 65%, 60%, 55%, 50%, 45%, 40%, 35%, 30%, 25%, 20%, 15%, 10%, or 5% homologous. Two or more sequences of sample index regions of two or more different sets of molecular barcodes may be less than about 80% homologous. Two or more sequences of sample index regions of two or more different sets of molecular barcodes may be less than about 60% homologous. Two or more sequences of sample index regions of two or more different sets of molecular barcodes may be less than about 40% homologous. Two or more sequences of sample index regions of two or more different sets of molecular barcodes may be less than about 20% homologous.

[00678] The molecular barcodes disclosed herein may comprise a cellular label. The cellular label of molecular barcodes of two or more sets of molecular barcodes may be different. The cellular label may comprise one or more nucleotides. Two or more sequences of cellular labels of two or more different sets of molecular barcodes may be less than about 90%, 85%, 80%, 75%, 70%, 65%, 60%, 55%, 50%, 45%, 40%, 35%, 30%, 25%, 20%, 15%, 10%, or 5% homologous. Two or more sequences of cellular labels of two or more different sets of molecular barcodes may be less than about 80% homologous. Two or more sequences of cellular labels of two or more different sets of molecular barcodes may be less than about 60% homologous. Two or more sequences of cellular labels of two or more different sets of molecular barcodes may be less than about 40% homologous. Two or more sequences of cellular labels of two or more different sets of molecular barcodes may be less than about 20% homologous.

[00679] The molecular barcode disclosed herein may further comprise a universal PCR region. The molecular barcode may further comprise a target-specific region. The molecular barcode may comprise one or more nucleotides. The label region may comprise one or more nucleotides. The sample index region may comprise one or more nucleotides. The universal PCR region may comprise one or more nucleotides. The target-specific region may comprise one or more nucleotides.

[00680] The kits and methods disclosed herein may comprise one or more sets of sample tags. The kits and methods disclosed herein may comprise one or more sample tags. The sample tags may comprise a sample index region. The sample index region of the sample tags of a first set of sample tags may be different from the sample index region of the sample tags of a second set of sample tags. The sample index region may comprise one or more nucleotides. Two or more sequences of sample index regions of two or more different sets of sample tags may be less than about 90%, 85%, 80%, 75%, 70%, 65%, 60%, 55%, 50%, 45%, 40%, 35%, 30%, 25%, 20%, 15%, 10%, or 5% homologous. Two or more sequences of sample index regions of two or more different sets of sample tags may be less than about 80% homologous. Two or more sequences of sample index regions of two or more different sets of sample tags may be less than about 60% homologous. Two or more sequences of sample index regions of two or more different sets of sample tags may be less than about 40% homologous. Two or more sequences of sample index regions of two or more different sets of sample tags may be less than about 20% homologous.

[00681] The kits and methods disclosed herein may comprise one or more sets of molecular identifier labels. The kits and methods disclosed herein may comprise one or more molecular

identifier labels. The molecular identifier labels may comprise a label region. The label regions of two or more molecular identifier labels of a set of molecular identifier labels may be different. The label region may comprise one or more nucleotides. A sequence of label regions of two or more molecular identifier labels of a set of molecular identifier labels may be less than about 90%, 85%, 80%, 75%, 70%, 65%, 60%, 55%, 50%, 45%, 40%, 35%, 30%, 25%, 20%, 15%, 10%, or 5% homologous. A sequence of label regions of two or more molecular identifier labels of a set of molecular identifier labels may be less than about 80% homologous. A sequence of label regions of two or more molecular identifier labels of a set of molecular identifier labels may be less than about 60% homologous. A sequence of label regions of two or more molecular identifier labels of a set of molecular identifier labels may be less than about 40% homologous. A sequence of label regions of two or more molecular identifier labels of a set of molecular identifier labels may be less than about 20% homologous. A label region may be referred to as a cellular label region.

[00682] The kits and methods disclosed herein may further comprise one or more primers. The one or more primers may comprise a sequence that is at least partially complementary to the universal PCR region. The one or more primers may comprise a sequence that is at least about 50% complementary to the universal PCR region. The one or more primers may comprise a sequence that is at least about 80% complementary to the universal PCR region.

[00683] The kits and methods disclosed herein may further comprise one or more amplification agents. The amplification agents may comprise a fixed panel of primers. The amplification agents may comprise one or more custom primers. The amplification agents may comprise one or more control primers. The amplification agents may comprise one or more housekeeping gene primers. The amplification agents may comprise one or more PCR reagents. The one or more PCR reagents may comprise polymerases, deoxyribonucleotide triphosphates (dNTPs), buffers, or a combination thereof.

[00684] The kits and methods disclosed herein may further comprise one or more beads. The molecular barcodes may be attached to the one or more beads. The sample tags may be attached to the one or more beads. The molecular identifier labels may be attached to the one or more beads.

[00685] Further disclosed herein are methods for generating one or more sets of beads. The method may comprise: a) depositing a plurality of first nucleic acids into a plurality of wells, wherein two or more different wells of the plurality of wells may comprise two or more different nucleic acids of the plurality of nucleic acids; b) contacting one or more wells of the plurality of wells with one or fewer beads to produce a plurality of single label beads,

wherein a single label bead of the plurality of first labeled beads comprises a bead attached to a nucleic acid of the plurality of first nucleic acids; c) pooling the plurality of first labeled beads from the plurality of wells to produce a pool of first labeled beads; d) distributing the pool of first labeled beads to a subsequent plurality of wells, wherein two or more wells of the subsequent plurality of wells comprise two or more different nucleic acids of a plurality of subsequent nucleic acids; and e) attaching one or more nucleic acids of the plurality of subsequent nucleic acids to one or more first labeled beads to produce a plurality of uniquely labeled beads.

[00686] The methods and kits disclosed herein may be used to analyze a plurality of nucleic acids. The methods and kits disclosed herein may be used to analyze less than about 100,000,000 nucleic acids. The methods and kits disclosed herein may be used to analyze less than about 10,000,000 nucleic acids. The methods and kits disclosed herein may be used to analyze less than about 1,000,000 nucleic acids. Further disclosed herein are methods of analyzing a plurality of proteins. The method may comprise: a) producing a plurality of sample-tagged polypeptides by: i) contacting a first sample comprising a plurality of polypeptides with a plurality of first sample tags to produce a plurality of first sample-tagged polypeptides; and ii) contacting a second sample comprising a plurality of polypeptides with a plurality of second sample tags to produce a plurality of second sample-tagged polypeptides, wherein the plurality of first sample tags are different from the plurality of second sample tags; b) contacting the plurality of sample-tagged polypeptides with a plurality of molecular identifier labels to produce a plurality of labeled polypeptides; and c) detecting at least a portion of the labeled polypeptides, thereby determining a count of the plurality of polypeptides in the plurality of samples.

[00687] The methods of analyzing polypeptides in a plurality of samples may further comprise determining the identity of one or more labeled polypeptides. Determining the identity of the one or more labeled polypeptides may comprise mass spectrometry. The method may further comprise combining the labeled polypeptides of the first sample with the labeled polypeptides of the second sample. The labeled polypeptides may be combined prior to determining the number of different labeled polypeptides. The method may further comprise combining the first sample-tagged polypeptides and the second sample-tagged polypeptides. The first sample-tagged polypeptides and the second sample-tagged polypeptides may be combined prior to contact with the plurality of molecular identifier labels. Determining the number of different labeled polypeptides may comprise detecting at least a portion of the tagged labeled polypeptide. Detecting at least a portion of the tagged

labeled polypeptide may comprise detecting at least a portion of the sample tag, molecule-specific tag, polypeptide, or a combination thereof.

[00688] The methods disclosed herein may comprise contacting a plurality of samples with a plurality of sample tags and a plurality of molecular identifier labels. Contacting the plurality of samples with the plurality of sample tags and the plurality of molecular identifier labels may occur simultaneously. Contacting the plurality of samples with the plurality of sample tags and the plurality of molecular identifier labels may occur concurrently. Contacting the plurality of samples with the plurality of sample tags and the plurality of molecular identifier labels may occur sequentially. Contacting the plurality of samples with the plurality of sample tags may occur prior to contacting the plurality of samples with the plurality of molecular identifier labels. Contacting the plurality of samples with the plurality of sample tags may occur after contacting the plurality of samples with the plurality of molecular identifier labels.

[00689] The methods disclosed herein may comprise contacting a first sample with a first plurality of sample tags and a first plurality of molecular identifier labels. Contacting the first sample with the first plurality of sample tags and the first plurality of molecular identifier labels may occur simultaneously. Contacting the first sample with the first plurality of sample tags and the first plurality of molecular identifier labels may occur concurrently. Contacting the first sample with the first plurality of sample tags and the first plurality of molecular identifier labels may occur sequentially. Contacting the first sample with the first plurality of sample tags may occur prior to contacting the first sample with the first plurality of molecular identifier labels. Contacting the first sample with the first plurality of sample tags may occur after contacting the first sample with the first plurality of molecular identifier labels.

[00690] The methods disclosed herein may comprise contacting a second sample with a second plurality of sample tags and a second plurality of molecular identifier labels. Contacting the second sample with the second plurality of sample tags and the second plurality of molecular identifier labels may occur simultaneously. Contacting the second sample with the second plurality of sample tags and the second plurality of molecular identifier labels may occur concurrently. Contacting the second sample with the second plurality of sample tags and the second plurality of molecular identifier labels may occur sequentially. Contacting the second sample with the second plurality of sample tags may occur prior to contacting the second sample with the second plurality of molecular identifier labels. Contacting the second sample with the second plurality of sample tags may occur after contacting the second sample with the second plurality of molecular identifier labels.

[00691] The methods and kits disclosed herein may further comprise combining two or more samples. The methods and kits disclosed herein may further comprise combining the first sample and the second sample. The first and second samples may be combined prior to contact with the plurality of molecular identifier labels. The first and second samples may be combined prior to detecting the labeled nucleic acids. The two or more samples may be combined prior to stochastically labeling two or more molecules in the two or more samples. The two or more samples may be combined after stochastically labeling two or more molecules in the two or more samples. The two or more samples may be combined prior to detecting two or more molecules in the two or more samples. The two or more samples may be combined after detecting two or more molecules in the two or more samples. The two or more samples may be combined prior to analyzing two or more molecules in the two or more samples. The two or more samples may be combined after analyzing two or more molecules in the two or more samples. The two or more samples may be combined prior to conducting one or more assays on two or more molecules in the two or more samples. The two or more samples may be combined after conducting one or more assays on two or more molecules in the two or more samples.

[00692] The methods and kits disclosed herein may comprise conducting one or more assays on two or more molecules in a sample. The one or more assays may comprise one or more amplification reactions. The methods and kits disclosed herein may further comprise conducting one or more amplification reactions to produce labeled nucleic acid amplicons. The labeled nucleic acids may be amplified prior to detecting the labeled nucleic acids. The method may further comprise combining the first and second samples prior to conducting the one or more amplification reactions.

[00693] The amplification reactions may comprise amplifying at least a portion of the sample tag. The amplification reactions may comprise amplifying at least a portion of the label. The amplification reactions may comprise amplifying at least a portion of the sample tag, label, nucleic acid, or a combination thereof. The amplification reactions may comprise amplifying at least about 1%, 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 100% of the total number of nucleic acids of the plurality of nucleic acids. The amplification reactions may comprise amplifying at least about 1% of the total number of nucleic acids of the plurality of nucleic acids. The amplification reactions may comprise amplifying at least about 5% of the total number of nucleic acids of the plurality of nucleic acids. The amplification reactions may comprise amplifying at least about 1%, 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%,

60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 100% of the total number of labeled nucleic acids of the plurality of labeled nucleic acids. The amplification reactions may comprise amplifying at least about 1% of the total number of labeled nucleic acids of the plurality of labeled nucleic acids. The amplification reactions may comprise amplifying at least about 5% of the total number of labeled nucleic acids of the plurality of labeled nucleic acids. The amplification reactions may comprise amplifying at least about 10% of the total number of labeled nucleic acids of the plurality of labeled nucleic acids. The amplification reactions may comprise amplifying less than about 95%, 90%, 80%, 70%, 60% or 50% of the total number of nucleic acids of the plurality of nucleic acids. The amplification reactions may comprise amplifying less than about 50% of the total number of nucleic acids of the plurality of nucleic acids. The amplification reactions may comprise amplifying less than about 20% of the total number of nucleic acids of the plurality of nucleic acids. The amplification reactions may comprise amplifying less than about 10% of the total number of nucleic acids of the plurality of nucleic acids. The amplification reactions may comprise amplifying less than about 95%, 90%, 80%, 70%, 60% or 50% of the total number of labeled nucleic acids of the plurality of labeled nucleic acids. The amplification reactions may comprise amplifying less than about 40% of the total number of labeled nucleic acids of the plurality of labeled nucleic acids. The amplification reactions may comprise amplifying less than about 25% of the total number of labeled nucleic acids of the plurality of labeled nucleic acids. The amplification reactions may comprise amplifying less than about 10% of the total number of labeled nucleic acids of the plurality of labeled nucleic acids.

[00694] The one or more amplification reactions may result in amplification of about 1, 2, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 200, 300, 400, 500, 600, 700, 800, 900 or 1000 targeted nucleic acids in a sample. The one or more amplification reactions may result in amplification of about 2000 targeted nucleic acids in a sample. The one or more amplification reactions may result in amplification of about 1000 targeted nucleic acids in a sample. The one or more amplification reactions may result in amplification of about 2000 targeted molecules. The one or more amplification reactions may result in amplification of about 100 targeted nucleic acids in a sample.

[00695] The amplification reactions may comprise one or more polymerase chain reactions (PCRs). The one or more polymerase chain reactions may comprise multiplex PCR, nested PCR, absolute PCR, HD-PCR, Next Gen PCR, digital RTA, or any combination thereof. The one or more polymerase chain reactions may comprise multiplex PCR. The one or more polymerase chain reactions may comprise nested PCR.

[00696] Conducting the one or more amplification reactions may comprise the use of one or more primers. The one or more primers may comprise one or more oligonucleotides. The one or more oligonucleotides may comprise at least about 7-9 nucleotides. The one or more oligonucleotides may comprise less than 12-15 nucleotides. The one or more primers may anneal to at least a portion of the plurality of labeled nucleic acids. The one or more primers may anneal to the 3' end and/or 5' end of the plurality of labeled nucleic acids. The one or more primers may anneal to an internal region of the plurality of labeled nucleic acids. The internal region may be at least about 50, 100, 150, 200, 220, 230, 240, 250, 260, 270, 280, 290, 300, 310, 320, 330, 340, 350, 360, 370, 380, 390, 400, 410, 420, 430, 440, 450, 460, 470, 480, 490, 500, 510, 520, 530, 540, 550, 560, 570, 580, 590, 600, 650, 700, 750, 800, 850, 900 or 1000 nucleotides from the 3' ends the plurality of labeled nucleic acids. The internal region may be at least about 2000 nucleotides from the 3' ends the plurality of labeled nucleic acids. The one or more primers may comprise a fixed panel of primers. The one or more primers may comprise at least one or more custom primers. The one or more primers may comprise at least one or more control primers. The one or more primers may comprise at least one or more housekeeping gene primers. The one or more oligonucleotides may comprise a sequence selected from a group consisting of sequences in Table 1. The one or more primers may comprise a universal primer. The universal primer may anneal to a universal primer binding site. The universal primer may anneal to a universal PCR region. The one or more custom primers may anneal to at least a portion of a sample tag. The one or more custom primers may anneal to at least a portion of a molecular identifier label. The one or more custom primers may anneal to at least a portion of a molecular barcode. The one or more custom primers may anneal to the first sample tag, the second sample tag, the molecular identifier label, the nucleic acid or a product thereof. The one or more primers may comprise a universal primer and a custom primer. The one or more primers may comprise at least about 96 or more custom primers. The one or more primers may comprise at least about 960 or more custom primers. The one or more primers may comprise at least about 9600 or more custom primers. The one or more custom primers may anneal to two or more different labeled nucleic acids. The two or more different labeled nucleic acids may correspond to one or more genes.

[00697] Multiplex PCR reactions may comprise a nested PCR reaction. The nested PCR reaction may comprise a pair of primers comprising a first primer and a second primer. The first primer may anneal to a region of one or more nucleic acids of the plurality of nucleic acids. The region of the one or more nucleic acids may be at least about 300 to 400

nucleotides from the 3' end of the one or more nucleic acids. The second primer may anneal to a region of one or more nucleic acids of the plurality of nucleic acids. The region of the one or more nucleic acids may be at least 200 to 300 nucleotides from the 3' end of the one or more nucleic acids.

[00698] The methods and kits disclosed herein may further comprise conducting one or more cDNA synthesis reactions to produce one or more cDNA copies of the molecules or derivatives thereof (*e.g.*, labeled molecules). The one or more cDNA synthesis reactions may comprise one or more reverse transcription reactions.

[00699] The methods and kits disclosed herein may comprise one or more samples. The methods and kits disclosed herein may comprise a plurality of samples. The plurality of samples may comprise at least about 2, 3, 4, 5, 10, 20, 30, 40, 50, 60, 70, 80, 90 or 100 or more samples. The plurality of samples may comprise at least about 100, 200, 300, 400, 500, 600, 700, 800, 900 or 1000 or more samples. The plurality of samples may comprise at least about 1000, 2000, 3000, 4000, 5000, 6000, 7000, 8000 samples, 9000, or 10,000 samples, or 100,000 samples, or 1,000,000 or more samples. The plurality of samples may comprise at least about 10,000 samples. The plurality of samples may comprise at least about 2 samples. The plurality of samples may comprise at least about 5 samples. The plurality of samples may comprise at least about 10 samples. The plurality of samples may comprise at least about 50 samples. The plurality of samples may comprise at least about 100 samples.

[00700] The methods and kits disclosed herein may comprise one or more samples comprising one or more cells. The methods and kits disclosed herein may comprise two or more samples comprising one or more cells. A first sample may comprise one or more cells. A second sample may comprise one or more cells. The one or more cells of the first sample may be of the same cell type as the one or more cells of the second sample.

[00701] The methods and kits disclosed herein may comprise a plurality of samples. The plurality of samples may be from one or more subjects. The plurality of samples may be from two or more subjects. The plurality of samples may be from the same subject. The two or more subjects may be from the same species. The two or more subjects may be from different species. The plurality of samples may be from one or more sources. The plurality of samples may be from two or more sources. The plurality of samples may be from the same subject. The two or more sources may be from the same species. The two or more sources may be from different species.

[00702] The plurality of samples may be obtained concurrently. The plurality of samples may be obtained sequentially. The plurality of samples may be obtained over two or more

time periods. The two or more time periods may be one or more hours apart. The two or more time periods may be one or more days apart. The two or more time periods may be one or more weeks apart. The two or more time periods may be one or more months apart. The two or more time periods may be one or more years apart.

[00703] The plurality of samples may be from one or more bodily fluids, tissues, cells, organs, or muscles. The plurality of samples may comprise one or more blood samples.

[00704] The methods and kits disclosed herein may comprise one or more samples comprising one or more nucleic acids. Two or more samples may comprise one or more nucleic acids. Two or more samples may comprise two or more nucleic acids. The one or more nucleic acids of a first sample may be different from one or more nucleic acids of a second sample. The nucleic acids in a first sample may be at least about 50% identical to the nucleic acids in a second sample. The nucleic acids in a first sample may be at least about 70% identical to the nucleic acids in a second sample. The nucleic acids in a first sample may be at least about 80% identical to the nucleic acids in a second sample.

[00705] The plurality of nucleic acids in the one or more samples may comprise two or more identical sequences. At least about 5%, 10 %, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 100% of the total nucleic acids in the one or more samples may comprise the same sequence. The plurality of nucleic acids in one or more samples may comprise at least two different sequences. At least about 5%, 10 %, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 100% of the total nucleic acids in the one or more samples may comprise different sequences.

[00706] The plurality of nucleic acids may comprise RNA, DNA, cDNA, mRNA, genomic DNA, small RNA, non-coding RNA, or other nucleic acid contents of a cell. The plurality of nucleic acids may comprise mRNA. The plurality of nucleic acids may comprise RNA. The plurality of nucleic acids may comprise mRNA. The plurality of nucleic acids may comprise DNA.

[00707] The methods and kits disclosed herein may comprise one or more sample tags. The methods and kits disclosed herein may comprise one or more pluralities of sample tags. The sample tags may comprise a sample index region. The sample index region of a first plurality of sample tags may be different from the sample index region of a second plurality of sample tags. The sample tags may comprise one or more nucleotides.

[00708] The sample tags may comprise at least about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95 or 100 nucleotides. The sample tags may comprise at least about 5 or more nucleotides. The sample tags may comprise at least about 10 or more

nucleotides. The sample tags may comprise less than about 200 nucleotides. The sample tags may comprise less than about 100 nucleotides. The sample tags may comprise less than about 60 nucleotides.

[00709] The sample tags may further comprise a universal primer binding site. The sample tags may further comprise a universal PCR region. The sample tags may further comprise one or more adaptor regions. The sample tags may further comprise one or more target-specific regions.

[00710] The methods and kits disclosed herein may comprise one or more molecular identifier labels. The methods and kits disclosed herein may comprise one or more pluralities of molecular identifier labels. The one or more pluralities of molecular identifier labels may comprise two or more different molecular identifier labels. The one or more pluralities of molecular identifier labels may comprise 50 or more different molecular identifier labels. The one or more pluralities of molecular identifier labels may comprise 90 or more different molecular identifier labels. The one or more pluralities of molecular identifier labels may comprise 100 or more different molecular identifier labels. The one or more pluralities of molecular identifier labels may comprise 300 or more different molecular identifier labels. The one or more pluralities of molecular identifier labels may comprise 500 or more different molecular identifier labels. The one or more pluralities of molecular identifier labels may comprise 960 or more different molecular identifier labels. The one or more pluralities of molecular identifier labels may comprise multiple copies of one or more molecular identifier labels. Two or more pluralities of molecular identifier labels may comprise one or more identical molecular identifier labels. Two or more pluralities of molecular identifier labels may comprise 10 or more identical molecular identifier labels. The molecular identifier labels of a first plurality of molecular identifier labels may be at least about 30% identical to the molecular identifier labels of a second plurality of molecular identifier labels. The molecular identifier labels of a first plurality of molecular identifier labels may be at least about 50% identical to the molecular identifier labels of a second plurality of molecular identifier labels. The molecular identifier labels of a first plurality of molecular identifier labels may be at least about 80% identical to the molecular identifier labels of a second plurality of molecular identifier labels.

[00711] The molecular identifier labels may comprise a label region (*e.g.*, molecular label region, molecular index region). The label region of two or more molecular identifier labels of a first plurality of molecular identifier labels may be different. One or more pluralities of molecular identifier labels may comprise at least about 20 different label regions. One or

more pluralities of molecular identifier labels may comprise at least about 50 different label regions. One or more pluralities of molecular identifier labels may comprise at least about 96 different label regions. One or more pluralities of molecular identifier labels may comprise at least about 200 different label regions. One or more pluralities of molecular identifier labels may comprise at least about 500 different label regions. One or more pluralities of molecular identifier labels may comprise at least about 960 different label regions.

[00712] The molecular identifier labels may comprise at least about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more nucleotides. The molecular identifier labels may comprise at least about 20, 30, 40, 50 or more nucleotides. The molecular identifier labels may comprise at least about 21 nucleotides.

[00713] The molecular identifier labels may further comprise a target-specific region. The target-specific region may comprise an oligodT sequence.

[00714] The molecular identifier labels may further comprise one or more dye labels. The molecular identifier labels may further comprise a Cy3 dye. The molecular identifier labels may further comprise a Tye563 dye.

[00715] The methods and kits disclosed herein may comprise one or more labeled molecules. The one or more labeled molecules may be produced by contacting a plurality of molecules with a plurality of sample tags. The one or more labeled molecules may be produced by contacting a plurality of nucleic acids with a plurality of sample tags. Contacting the plurality of nucleic acids with the plurality of sample tags may comprise ligating one or more sample tags to one or more nucleic acids. Contacting the plurality of nucleic acids with the plurality of sample tag may comprise hybridizing one or more sample tags to one or more nucleic acids. Contacting the plurality of nucleic acids with the plurality of sample tag may comprise performing one or more nucleic acid extension reactions. The one or more nucleic acid extension reactions may comprise reverse transcription.

[00716] The methods and kits disclosed herein may further comprise attaching one or more oligonucleotide linkers to the plurality of nucleic acids. The method and kits may further comprise attaching one or more oligonucleotide linkers to the sample tagged nucleic acids. The methods and kits may further comprise attaching one or more oligonucleotide linkers to the labeled nucleic acids. The one or more linkers may comprise at least about 10 nucleotides.

[00717] The methods and kits disclosed herein may further comprise attaching one or more labeled nucleic acids to a support. The support may comprise a solid support. The support

may comprise a bead. The support may comprise an array. The support may comprise a glass slide.

[00718] Attachment of the labeled nucleic acids to the support may comprise amine-thiol crosslinking, maleimide crosslinking, *N*-hydroxysuccinimide or *N*-hydroxysulfosuccinimide, Zenon, SiteClick, or a combination thereof. Attaching the labeled nucleic acids to the support may comprise attaching biotin to the one or more labeled nucleic acids.

[00719] The support may comprise one or more beads. The one or more beads may be a coated bead. The coated bead may be coated with streptavidin.

[00720] The support may comprise an array. The array may comprise one or more probes. The labeled nucleic acids may be attached to the one or more probes. The one or more probes may comprise one or more oligonucleotides. The one or more probes may be attached to at least a portion of the labeled nucleic acids. The portion of the labeled nucleic acids attached to the one or more probes may comprise at least a portion of the sample tag, molecular identifier label, molecular barcode, nucleic acid, or a combination thereof.

[00721] The support may comprise a glass slide. The glass slide may comprise one or more wells. The one or more wells may be etched on the glass slide. The one or more wells may comprise at least 960 wells. The glass slide may comprise one or more probes. The one or more probes may be printed onto the glass slide. The one or more wells may further comprise one or more probes. The one or more probes may be printed within the one or more wells. The one or more probes may comprise 960 nucleic acids. The nucleic acids may be different. The nucleic acids may be the same.

[00722] The methods and kits disclosed herein may be used to determine a count of one or more molecules in one or more samples. Determining the count of one or more molecules may comprise determining the number of different labeled nucleic acids. Determining the number of different labeled nucleic acids may comprise detecting at least a portion of the labeled nucleic acid. Detecting at least a portion of the labeled nucleic acid may comprise detecting at least a portion of the sample tag, molecular identifier label, molecular barcode, nucleic acid, or a combination thereof.

[00723] Determining the number of different labeled nucleic acids may comprise sequencing. Sequencing may comprise MiSeq sequencing. Sequencing may comprise HiSeq sequencing. Determining the number of different labeled nucleic acids may comprise an array. Determining the number of different labeled nucleic acids may comprise contacting the labeled nucleic acids with the one or more probes.

[00724] Determining the number of different labeled nucleic acids may comprise contacting the labeled nucleic acids with an array. The array may comprise a plurality of probes.

Determining the number of different labeled nucleic acids may comprise contacting the labeled nucleic acids with a glass slide of a plurality of probes.

[00725] Determining the number of different labeled nucleic acids may comprise labeled probe hybridization, target-specific amplification, target-specific sequencing, sequencing with labeled nucleotides specific for target small nucleotide polymorphism, sequencing with labeled nucleotides specific for restriction enzyme digest patterns, sequencing with labeled nucleotides specific for mutations, or a combination thereof.

[00726] Determining the number of different labeled nucleic acids may comprise flow cytometry sorting of a sequence-specific label. Determining the number of different labeled nucleic acids may comprise detection of the labeled nucleic acids attached to the beads. Detection of the labeled nucleic acids attached to the beads may comprise fluorescence detection.

[00727] Determining the number of different labeled nucleic acids may comprise counting the plurality of labeled nucleic acids by fluorescence resonance energy transfer (FRET), between a target-specific probe and a labeled nucleic acid or a target-specific labeled probe. Determining the number of different labeled nucleic acids may comprise attaching the labeled nucleic acid to the support.

[00728] The methods and kits disclosed herein may further comprise immunoprecipitation of a target sequence with a nucleic-acid binding protein.

[00729] The methods and kits disclosed herein may further comprise distributing the plurality of samples into a plurality of wells of a microwell plate. One or more of the plurality of samples may comprise a plurality of cells. One or more of the plurality of samples may comprise a plurality of nucleic acids. The methods and kits disclosed herein may further comprise distributing one or fewer cells to the plurality of wells. The plurality of cells may be lysed in the microwell plate. The methods and kits disclosed herein may further comprise synthesizing cDNA in the microwell plate. Synthesizing cDNA may comprise reverse transcription of mRNA.

[00730] The methods and kits disclosed herein may further comprise distributing the plurality of first sample tags, the plurality of second sample tags, the plurality of molecular identifier labels, or any combination thereof into a microwell plate.

[00731] The methods and kits disclosed herein may further comprise distributing one or more beads in the microwell plate. The microwell plate may comprise a microwell plate

fabricated on PDMS by soft lithography, etched on a silicon wafer, etched on a glass slide, patterned photoresist on a glass slide, or a combination thereof. The microwell may comprise a hole on a microcapillary plate. The microwell plate may comprise a water-in-oil emulsion. The microwell plate may comprise at least one or more wells. The microwell plate may comprise at least about 6 wells, 12 wells, 48 wells, 96 wells, 384 wells, 960 wells or 1000 wells.

[00732] The methods and kits disclosed herein may further comprise a chip. The microwell plate may be attached to the chip. The chip may comprise at least about 6 wells, 12 wells, 48 wells, 96 wells, 384 wells, 960 wells, 1000 wells, 2000 wells, 3000 wells, 4000 wells, 5000 wells, 6000 wells, 7000 wells, 8000 wells, 9000 wells, 10,000 wells, 20,000 wells, 30,000 wells, 40,000 wells, 50,000 wells, 60,000 wells, 70,000 wells, 80,000 wells, 90,000 wells, 100,000 wells, 200,000 wells, 500,000 wells, or a million wells. The wells may comprise an area of at least about 300 microns², 400 microns², 500 microns², 600 microns², 700 microns², 800 microns², 900 microns², 1000 microns², 1100 microns², 1200 microns², 1300 microns², 1400 microns², 1500 microns². The methods and kits disclosed herein may further comprise distributing between about 10,000 and 30,000 samples on the chip.

[00733] The methods and kits disclosed herein may further comprise diagnosing a condition, disease, or disorder in a subject in need thereof.

[00734] The methods and kits disclosed herein may further comprise prognosing a condition, disease, or disorder in a subject in need thereof. The methods and kits disclosed herein may further comprise determining a treatment for a condition, disease, or disorder in a subject in need thereof.

[00735] The plurality of samples may comprise one or more samples from a subject suffering from a disease or condition. The plurality of samples may comprise one or more samples from a healthy subject.

[00736] Further disclosed herein is a method of forensic analysis comprising: a) stochastically labeling two or more molecules in two or more samples to produce two or more labeled molecules; and b) detecting the two or more labeled molecules.

[00737] The method of selecting the custom primer may further comprise selecting the custom primer based on one or more nucleic acids. The one or more nucleic acids may comprise mRNA transcripts, non-coding transcripts including structural RNAs, transcribed pseudogenes, model mRNA provided by a genome annotation process, sequences corresponding to a genomic contig, or any combination thereof. The one or more nucleic acids may be RNA. The one or more nucleic acids may be mRNA. The one or more nucleic

acids may comprise one or more exons. The method of selecting the custom primer may further comprise enriching for one or more subsets of nucleic acids. The one or more subsets comprise low abundance mRNAs. The method of selecting the custom primer may further comprise a computational algorithm.

[00738] The methods and kits disclosed herein may comprise the use of one or more controls. The one or more controls may be spiked in controls. The one or more controls may comprise nucleic acids. The one or more samples comprising a plurality of nucleic acids may be spiked with one or more control nucleic acids. The one or more control nucleic acids may be used to measure an efficiency of producing the labeled nucleic acid library.

[00739] The methods and kits disclosed herein may be used in the production of one or more nucleic acid libraries. The one or more nucleic acid libraries may comprise a plurality of labeled nucleic acids or derivatives thereof (*e.g.*, labeled amplicons). The method of producing the labeled nucleic acid library may comprise stochastically labeling two or more nucleic acids in two or more samples with two or more sets of molecular barcodes to produce a plurality of labeled nucleic acids. The method of producing a labeled nucleic acid library may comprise contacting two or more samples with a plurality of sample tags and a plurality of molecule specific labels to produce a plurality of labeled nucleic acids. The labeled nucleic acids may comprise a sample index region, a label region and a nucleic acid region. The sample index region may be used to confer a sample or sub-sample identity to the nucleic acid. The sample index region may be used to determine the source of the nucleic acid. The label region may be used to confer a unique identity to the nucleic acid, thereby enabling differentiation of two or more identical nucleic acids in the same sample or sub-sample.

[00740] The method of producing a nucleic acid library may further comprise amplifying one or more labeled nucleic acids to produce one or more enriched labeled nucleic acids. The method may further comprise conducting one or more pull-down assays of the one or more enriched labeled nucleic acids. The method may further comprise purifying the one or more enriched labeled nucleic acids.

[00741] The kits disclosed herein may comprise a plurality of beads, a primer and/or amplification agents. One or more kits may be used in the analysis of at least about 10, 20, 30, 40, 50, 60, 70, 80, 90, or 100 or more samples or sub-samples. One or more kits may be used in the analysis of at least about 96 samples. One or more kits may be used in the analysis of at least about 384 samples. The kit may further comprise instructions for primer design and optimization.

[00742] The kit may further comprise one or more microwell plates. The one or more microwell plates may be used for the distribution of one or more beads. The one or more microwell plates may be used for the distribution of one or more molecules or derivatives thereof (*e.g.*, labeled molecules, labeled amplicons) from one or more samples.

[00743] The kit may further comprise one or more additional containers. The one or more additional containers may comprise one or more additional pluralities of sample tags. The one or more additional pluralities of sample tags in the one or more additional containers may be different from the first plurality of sample tags in the first container. The one or more additional containers may comprise one or more additional pluralities of molecular identifier labels. The one or more additional pluralities of molecular identifier labels of the one or more additional containers may be at least about 50% identical to the one or more additional molecular identifier labels of the second container. The one or more additional pluralities of molecular identifier labels of the one or more additional containers may be at least about 80% identical to the one or more additional molecular identifier labels of the second container. The one or more additional pluralities of molecular identifier labels of the one or more additional containers may be at least about 90% identical to the one or more additional molecular identifier labels of the second container.

[00744] Further disclosed herein are methods of producing one or more sets of labeled beads. The method of producing the one or more sets of labeled beads may comprise attaching one or more nucleic acids to one or more beads, thereby producing one or more sets of labeled beads. The one or more nucleic acids may comprise one or more molecular barcodes. The one or more nucleic acids may comprise one or more sample tags. The one or more nucleic acids may comprise one or more molecular identifier labels. The one or more nucleic acids may comprise a) a primer region; b) a sample index region; and c) a linker or adaptor region. The one or more nucleic acids may comprise a) a primer region; b) a label region; and c) a linker or adaptor region. The one or more nucleic acids may comprise a) a sample index region; and b) a label region. The one or more nucleic acids may further comprise a primer region. The one or more nucleic acids may further comprise a target specific region. The one or more nucleic acids may further comprise a linker region. The one or more nucleic acids may further comprise an adaptor region. The one or more nucleic acids may further comprise a sample index region. The one or more nucleic acids may further comprise a label region.

[00745] The primer region of the nucleic acids for a set of labeled beads may be at least about 70% identical. The primer region of the nucleic acids for a set of labeled beads may be

at least about 90% identical. The primer region of the nucleic acids for a set of labeled beads may be the same.

[00746] The sample index region of the nucleic acids for a set of labeled beads may be at least about 70% identical. The sample index region of the nucleic acids for a set of labeled beads may be at least about 90% identical. The sample index region of the nucleic acids for a set of labeled beads may be the same. The sample index region of the nucleic acids for two or more sets of sample indexed beads may be less than about 40% identical. The sample index region of the nucleic acids for two or more sets of sample indexed beads may be less than about 50% identical. The sample index region of the nucleic acids for two or more sets of sample indexed beads may be less than about 60% identical. The sample index region of nucleic acids for two or more sets of sample indexed beads may be different.

[00747] The label region of the nucleic acids for two or more sets of labeled beads may be at least about 70% identical. The label region of the nucleic acids for two or more sets of labeled beads may be at least about 90% identical. The label region of the nucleic acids for two or more sets of labeled beads may be the same. The label region of the nucleic acids for a set of labeled beads may be less than about 40% identical. The label region of the nucleic acids for a set of labeled beads may be less than about 50% identical. The label region of the nucleic acids for a set of labeled beads may be less than about 60% identical. The label region of two or more nucleic acids for a set of labeled beads may be different.

[00748] The linker or adaptor region of the nucleic acids for a set of labeled beads may be at least about 70% identical. The linker or adaptor region of the nucleic acids for a set of labeled beads may be at least about 90% identical. The linker or adaptor region of the nucleic acids for a set of labeled beads may be the same.

[00749] The target specific region of the nucleic acids for two or more sets of target specified beads may be at least about 70% identical. The target specific region of the nucleic acids for two or more sets of target specified beads may be at least about 90% identical. The target specific region of the nucleic acids for two or more sets of target specified beads may be the same. The target specific region of the nucleic acids for a set of target specified beads may be less than about 40% identical. The target specific region of the nucleic acids for a set of target specified beads may be less than about 50% identical. The target specific region of the nucleic acids for a set of target specified beads may be less than about 60% identical. The target specific region of two or more nucleic acids for a set of target specified beads may be different.

[00750] The one or more sets of labeled beads may comprise one million or more labeled beads. The one or more sets of labeled beads may comprise ten million or more labeled beads.

[00751] Attaching the one or more nucleic acids to the beads may comprise covalent attachment. Attaching the one or more nucleic acids to the beads may comprise conjugation. Attaching the one or more nucleic acids to the beads may comprise ionic interactions.

[00752] The beads may be coated beads. The nucleic acids may be attached to one or more tags. The beads may be coated with streptavidin. The nucleic acids may be attached to biotin. The beads may also be coated with antibodies or nucleic acids, and the nucleic acids may be attached to the beads indirectly via such surface coated materials.

[00753] In one aspect, the disclosure provides for a composition comprising: a solid support, wherein said solid support comprises a plurality of oligonucleotides, wherein at least two of said plurality of oligonucleotides comprises a cellular label and a molecular label, wherein said cellular labels of said at least two of said plurality of oligonucleotides are the same, and wherein said molecular labels of said at least two of said plurality of oligonucleotides are different. In some embodiments, the plurality of oligonucleotide further comprises a sample label. In some embodiments, the plurality of oligonucleotides further comprises a target binding region. In some embodiments, the target binding region comprises a sequence is adapted to hybridize to a target nucleic acid. In some embodiments, the target binding region comprises a sequence selected from the group consisting of: a random multimer *e.g.*, a random dimer, trimer, tetramer, pentamer, hexamer, heptamer, octamer, nonamer, decamer, or higher multimer sequence of any length; a gene-specific primer; and oligo dT; or any combination thereof. In some embodiments, the plurality of oligonucleotides comprises a universal label. In some embodiments, the universal label comprises a binding site for a sequencing primer. In some embodiments, the plurality of oligonucleotides comprises a linker. In some embodiments, the linker comprises a functional group. In some embodiments, the linker is located 5' to said oligonucleotide. In some embodiments, the linker is selected from the group consisting of: C6, biotin, streptavidin, primary amines, aldehydes, and ketones, or any combination thereof. In some embodiments, the solid support is comprised of polystyrene. In some embodiments, the solid support is magnetic. In some embodiments, the solid support is selected from the group consisting of: a PDMS solid support, a glass solid support, a polypropylene solid support, an agarose solid support, a gelatin solid support, a magnetic solid support, and a pluronic solid support, or any combination thereof. In some embodiments, the solid support comprises a bead. In some embodiments, the solid support

comprises a diameter of about 20 microns. In some embodiments, the solid support comprises a diameter from about 5 microns to about 40 microns. In some embodiments, the solid support comprises a functional group. In some embodiments, the functional group is selected from the group consisting of: C6, biotin, streptavidin, primary amines, aldehydes, and ketones, or any combination thereof. In some embodiments, the cellular label comprises a plurality of cellular labels. In some embodiments, the plurality of cellular labels are interspersed with a plurality of linker label sequences. In some embodiments, the plurality of oligonucleotides comprises from 10,000 to 1 billion oligonucleotides.

[00754] In one aspect the disclosure provides for a solid support comprising: a first oligonucleotide comprising: a first cellular label comprising a first random sequence, a second random sequence, and a first linker label sequence, wherein said first linker label sequence connects said first random sequence and said second random sequence; and a first molecular label comprising a random sequence; and a second oligonucleotide comprising: a second cellular label comprising a third random sequence, a fourth random sequence, and a second linker label sequence, wherein said second linker label sequence connects said third random sequence and said fourth random sequence; and a second molecular label comprising a random sequence, wherein said first cellular label and said second cellular label are the same and said first molecular label and said second molecular label are different. In some embodiments, the first and second oligonucleotides further comprise identical sample index regions. In some embodiments, the sample index region comprises a random sequence. In some embodiments, the sample index region is 4-12 nucleotides in length. In some embodiments, the cellular label is directly attached to said molecular label. In some embodiments, the cellular label and said molecular label are attached through a linker label sequence. In some embodiments, the random sequence of said cellular label is from 4-12 nucleotides in length. In some embodiments, the constant sequence of said cellular label is at least 4 nucleotides in length. In some embodiments, the cellular label has a total length of at least 12 nucleotides. In some embodiments, the cellular label further comprises one or more additional random sequences. In some embodiments, the cellular label further comprises one or more additional linker label sequences. In some embodiments, the one or more additional linker label sequences connect the one or more additional random sequences. In some embodiments, the random sequence of the molecular label is 4-12 nucleotides in length.

[00755] In one aspect the disclosure provides for a composition comprising: a solid support, wherein said solid support comprises a plurality of oligonucleotides, wherein at least two of said plurality of oligonucleotides comprises: a cellular label, a molecular label; and a target

binding region; and a plurality of a target nucleic acids, wherein said cellular labels of said at least two of said plurality of oligonucleotides are the same, and wherein said molecular labels of said at least two of said plurality of oligonucleotides are different. In some embodiments, the target binding region comprises a sequence that is adapted to hybridize to at least one of said plurality of target nucleic acids. In some embodiments, the target binding region comprises a sequence selected from the group consisting of: a random a random multimer *e.g.*, a random dimer, trimer, quatramer, pentamer, hexamer, septamer, octamer, nonamer, decamer, or higher multimer sequence of any length; a gene-specific primer; and oligo dT; or any combination thereof. In some embodiments, the plurality of oligonucleotides comprises from 10,000 to 1 billion oligonucleotides. In some embodiments, the plurality of oligonucleotides comprises a number of oligonucleotides greater than the number of target nucleic acids of said plurality of target nucleic acids. In some embodiments, the plurality of target nucleic acids comprises multiple copies of a same target nucleic acid. In some embodiments, the plurality of target nucleic acids comprises multiple copies of different target nucleic acids. In some embodiments, the plurality of target nucleic acids are bound to said plurality of oligonucleotides. In some embodiments, the oligonucleotide further comprises a sample label. In some embodiments, the plurality of oligonucleotides comprises a universal label. In some embodiments, the universal label comprises a binding site for a sequencing primer. In some embodiments, the plurality of oligonucleotides comprises a linker. In some embodiments, the linker comprises a functional group. In some embodiments, the linker is located 5' to said oligonucleotide. In some embodiments, the functional group comprises an amino group. In some embodiments, the linker is selected from the group consisting of: C6, biotin, streptavidin, primary amines, aldehydes, and ketones, or any combination thereof. In some embodiments, the solid support is comprised of polystyrene. In some embodiments, the solid support is magnetic. In some embodiments, the solid support is selected from the group consisting of: a PDMS solid support, a glass solid support, a polypropylene solid support, an agarose solid support, a gelatin solid support, a magnetic solid support, and a pluronic solid support, or any combination thereof. In some embodiments, the solid support comprises a bead. In some embodiments, the solid support comprises a diameter of about 20 microns. In some embodiments, the solid support comprises a diameter from about 5 microns to about 40 microns. In some embodiments, the solid support comprises a functional group. In some embodiments, the functional group comprises a carboxy group. In some embodiments, the functional group is selected from the group consisting of: C6, biotin, streptavidin, primary amines, aldehydes, and ketones, or any

combination thereof. In some embodiments, the cellular label comprises a plurality of cellular labels. In some embodiments, the plurality of cellular labels is interspersed with a plurality of linker label sequences.

[00756] In one aspect the disclosure provides for a kit comprising: a first solid support, wherein said first solid support comprises a first plurality of oligonucleotides, wherein said first plurality of oligonucleotides comprises the same first cellular label, a second solid support, wherein said second solid support comprises a second plurality of oligonucleotides, wherein said second plurality of oligonucleotides comprises the same second cellular label, instructions for use, wherein said first cellular label and said second cellular label are different. In some embodiments, oligonucleotides form said first plurality of oligonucleotides and said second plurality of oligonucleotides comprises a molecular label. In some embodiments, the molecular labels of said oligonucleotides are different. In some embodiments, the molecular labels of said oligonucleotides are the same. In some embodiments, the molecular label of some of said oligonucleotides are different and some are the same. In some embodiments, the oligonucleotides from said first plurality of oligonucleotides and said second plurality of oligonucleotides comprise a target binding region. In some embodiments, the kit further comprises: a microwell array. In some embodiments, the kit further comprises: a buffer. In some embodiments, the buffer is selected from the group consisting of: a reconstitution buffer, a dilution buffer, and a stabilization buffer, or any combination thereof.

[00757] In one aspect the disclosure provides for a method for determining an amount of a target nucleic acid comprising: contacting a sample with a solid support, wherein said solid support comprises a plurality of oligonucleotides, wherein at least two of said plurality of oligonucleotides comprises a cellular label and a molecular label, wherein said cellular labels of said at least two of said plurality of oligonucleotides are the same, and wherein said molecular labels of said at least two of said plurality of oligonucleotides are different; and hybridizing said target nucleic acid from said sample to an oligonucleotide of said plurality of oligonucleotides. In some embodiments, the sample comprises cells. In some embodiments, the sample is lysed prior to said hybridizing. In some embodiments, the hybridizing comprising hybridizing multiple copies of a same target nucleic acid to said plurality of oligonucleotides. In some embodiments, the method further comprises: amplifying said target nucleic acid. In some embodiments, the amplifying comprises reverse transcribing said target nucleic acid. In some embodiments, the amplifying comprises amplification using a method selected from the group consisting of: PCR, quantitative PCR, real-time PCR, and digital

PCR, or any combination thereof. In some embodiments, the amplifying is performed directly on said solid support. In some embodiments, the amplifying is performed on a template transcribed from said solid support. In some embodiments, the method further comprises: sequencing said target nucleic acid. In some embodiments, the sequencing comprises sequencing said target nucleic acid and said molecular label. In some embodiments, the method further comprises: determining an amount of said target nucleic acid. In some embodiments, the determining comprises quantifying levels of said target nucleic acid. In some embodiments, the determining comprises counting the number of sequenced molecular labels for said target nucleic acid. In some embodiments, the contacting occurs in a microwell. In some embodiments, the microwell is made from a material selected from the group consisting of: hydrophilic plastic, plastic, elastomer, and hydrogel, or any combination thereof. In some embodiments, the microwell comprises agarose. In some embodiments, the microwell is one microwell of a microwell array. In some embodiments, the microwell array comprises at least 90 microwells. In some embodiments, the microwell array comprises at least 150,000 microwells. In some embodiments, the microwell comprises at least one solid support per well. In some embodiments, the microwell comprises at most two solid supports per well. In some embodiments, the microwell is of a size that accommodates at most two of said solid support. In some embodiments, the microwell is of a size that accommodates at most one solid support. In some embodiments, the microwell is at least 25 microns deep. In some embodiments, the microwell is at least 25 microns in diameter.

[00758] In one aspect the disclosure provides for a method to reduce amplification bias of a target nucleic acid comprising: contacting a sample to a solid support, wherein said solid support comprises a plurality of oligonucleotides, wherein at least two of said plurality of oligonucleotides comprises a cellular label and a molecular label, wherein said cellular labels of said at least two of said plurality of oligonucleotides are the same, and wherein said molecular labels of said at least two of said plurality of oligonucleotides are different; and hybridizing a target nucleic acid from said sample to said plurality of oligonucleotides; amplifying said target nucleic acid; sequencing said target nucleic acid, wherein said sequencing sequences said target nucleic acid and said molecular label of said oligonucleotide to which said target nucleic acid is bound; and determining an amount of said target nucleic acid. In some embodiments, the hybridizing comprising hybridizing multiple copies of a same target nucleic acid to said plurality of oligonucleotides. In some embodiments, the determining comprises counting a number of sequenced molecular labels for a same target nucleic acid. In some embodiments, the counting counts the number of

copies of said same target nucleic acid. In some embodiments, the sample comprises cells. In some embodiments, the amplifying comprises reverse transcribing said target nucleic acid. In some embodiments, the amplifying comprises amplification using a method selected from the group consisting of: PCR, quantitative PCR, real-time PCR, and digital PCR, or any combination thereof. In some embodiments, the amplifying is performed directly on said solid support. In some embodiments, the amplifying is performed on a template transcribed from said solid support.

[00759] In one aspect the disclosure provides for a composition comprising: a microwell; a cell; and a solid support, wherein said solid support comprises a plurality of oligonucleotides, wherein at least two of said plurality of oligonucleotides comprises a cellular label and a molecular label, wherein said cellular labels of said at least two of said plurality of oligonucleotides are the same, and wherein said molecular labels of said at least two of said plurality of oligonucleotides are different. In some embodiments, the at least two of said plurality of oligonucleotides further comprises a sample label. In some embodiments, the at least two of said plurality of oligonucleotides further comprises a target binding region. In some embodiments, the target binding region comprises a sequence selected from the group consisting of: a random multimer *e.g.*, a random dimer, trimer, tetramer, pentamer, hexamer, septamer, octamer, nonamer, decamer, or higher multimer sequence of any length; a gene-specific primer; and oligo dT; or any combination thereof. In some embodiments, the plurality of oligonucleotides comprises a universal label. In some embodiments, the universal label comprises a binding site for a sequencing primer. In some embodiments, the solid support is comprised of polystyrene. In some embodiments, the solid support is magnetic. In some embodiments, the solid support is selected from the group consisting of: a PDMS solid support, a glass solid support, a polypropylene solid support, an agarose solid support, a gelatin solid support, a magnetic solid support, and a pluronic solid support, or any combination thereof. In some embodiments, the solid support comprises a bead. In some embodiments, the solid support has a diameter of about 20 microns. In some embodiments, the solid support has a diameter from about 5 microns to about 40 microns. In some embodiments, the cellular label comprises a plurality of cellular labels. In some embodiments, the plurality of cellular labels is interspersed with a plurality of linker sequences. In some embodiments, the microwell is made from a material selected from the group consisting of: hydrophilic plastic, plastic, elastomer, and hydrogel, or any combination thereof. In some embodiments, the microwell comprises agarose. In some embodiments, the microwell is a microwell of a microwell array. In some embodiments, the microwell

comprises at least one solid support per well. In some embodiments, the microwell comprises at most two solid supports per well. In some embodiments, the microwell is of a size that accommodates at least one of said solid support and at least one of said cell. In some embodiments, the microwell is of a size that accommodates at most one of said solid support and at least one of said cell. In some embodiments, the microwell is at least 25 microns deep. In some embodiments, the microwell is at least 25 microns in diameter. In some embodiments, the microwell is flat.

[00760] In one aspect the disclosure provides for a device comprising: a first substrate comprising a first microwell array; wherein said first microwell array comprises a plurality of first microwells in a first pre-determined spatial arrangement configured to perform multiplexed, single cell stochastic labeling and molecular indexing assays.

[00761] In some embodiments, the device comprises a first substrate comprising at least a second microwell array, wherein said at least second microwell array comprises a plurality of at least second microwells in an at least second pre-determined spatial arrangement. In some embodiments, the first microwells and the at least second microwells are the same. In some embodiments, the first microwells and the at least second microwells are different. In some embodiments, the first pre-determined spatial arrangement and the at least second pre-determined spatial arrangement are the same. In some embodiments, the first pre-determined spatial arrangement and the at least second pre-determined spatial arrangement are different. In some embodiments, a pre-determined spatial arrangement comprises a one dimensional or two dimensional array pattern. In some embodiments, the two dimensional array pattern comprises a square grid, a rectangular grid, or a hexagonal grid. In some embodiments, the microwells comprise a cylindrical geometry, a conical geometry, a hemispherical geometry, a rectangular geometry, a polyhedral geometry, or a combination thereof. In some embodiments, a diameter of the microwells is between about 5 microns and about 50 microns. In some embodiments, a depth of the microwells is between about 10 microns and about 60 microns. In some embodiments, a center-to-center spacing between two adjacent microwells is between about 15 microns and about 75 microns. In some embodiments, a total number of microwells in a first or at least second microwell array is between about 96 and about 5,000,000. In some embodiments, the first substrate comprises silicon, fused-silica, glass, a polymer, a metal, or a combination thereof. In some embodiments, the first substrate further comprises agarose or a hydrogel. In some embodiments, a microwell array further comprises at least one surface feature, wherein said surface feature surrounds one or more

individual microwells or straddles a surface between individual microwells, and wherein said surface feature is domed, ridged, or peaked.

[00762] In one aspect the disclosure provides for a device comprising: a first substrate comprising at least a first microwell array; and a mechanical fixture comprising a top plate, a bottom plate, and a gasket; wherein when the first substrate and mechanical fixture are in assembled form, the first substrate is positioned between the gasket and the bottom plate, the gasket forms a leak-proof seal with the first substrate, and the top plate and gasket form at least a first chamber encompassing said at least first microwell array such that a cell sample and a bead-based oligonucleotide label may be dispensed into said at least first chamber to perform multiplexed, single cell stochastic labeling and a molecular indexing assays.

[00763] In some embodiments, the at least first microwell array is any described herein. In some embodiments, the gasket is fabricated from polydimethylsiloxane (PDMS) or a similar elastomeric material. In some embodiments, the top and bottom plates are fabricated from aluminum, anodized aluminum, stainless steel, teflon, polymethylmethacrylate, polycarbonate, or a similar rigid polymer material.

[00764] In one aspect the disclosure provides for a device comprising: at least one substrate further comprising at least one microwell array; and a flow cell; wherein the flow cell encloses or is attached to said at least one substrate, and includes at least one inlet port and at least one outlet port for the purpose of delivering fluids to said microwell arrays; and wherein the device is configured to perform multiplexed, single cell stochastic labeling and molecular indexing assays.

[00765] In some embodiments, said at least one substrate comprise at least one microwell array as described herein. In some embodiments, the flow cell further comprises a plurality of microarray chambers that interface with a plurality of microwell arrays such that one or more samples may be processed in parallel. In some embodiments, the flow cell further comprises a porous barrier or flow diffuser to provide more uniform delivery of cells and beads to the at least one microwell array. In some embodiments, the flow cell further comprises dividers that divide each chamber containing a microwell array into subsections that collectively cover the same total array area and provide for more uniform delivery of cells and beads to the at least one microwell array. In some embodiments, the width of fluid channels incorporated into the device is between about 50 microns and 20 mm. In some embodiments, the depth of fluid channels incorporated into the device is between about 50 microns and about 2 mm. In some embodiments, the flow cell is fabricated from a material selected from the group consisting of silicon, fused-silica, glass, polydimethylsiloxane (PDMS; elastomer),

polymethylmethacrylate (PMMA), polycarbonate (PC), polypropylene (PP), polyethylene (PE), high density polyethylene (HDPE), polyimide, cyclic olefin polymers (COP), cyclic olefin copolymers (COC), polyethylene terephthalate (PET), epoxy resin, metal, or a combination of these materials. In some embodiments, the device comprises a fixed component of an instrument system configured to perform automated multiplexed, single cell stochastic labeling and molecular indexing assays. In some embodiments, the device comprises a removable component of an instrument system configured to perform automated multiplexed, single cell stochastic labeling and molecular indexing assays.

[00766] In one aspect the disclosure provides for a cartridge comprising: at least a first substrate further comprising at least a first microwell array; at least a first flow cell or microwell array chamber; one or more sample or reagent reservoirs; and wherein the cartridge further comprises at least one inlet port and at least one outlet port for the purpose of delivering fluids to said at least first microwell array; and wherein the cartridge is configured to perform multiplexed, single cell stochastic labeling and molecular indexing assays.

[00767] In some embodiments, said at least first substrate comprises at least a first microwell array as described herein. In some embodiments, the cartridge comprises a plurality of microwell arrays and is configured to process one or more samples in parallel. In some embodiments, the at least first flow cell or microwell array chamber further comprises a porous barrier or flow diffuser to provide more uniform delivery of cells and beads to the at least first microwell arrays. In some embodiments, the at least first flow cell or microwell array chamber further comprises dividers that divide the at least first flow cell or microwell array chamber into subsections that collectively cover the same total array area and provide for more uniform delivery of cells and beads to the microwell arrays. In some embodiments, the width of fluid channels incorporated into the cartridge is between about 50 microns and 200 microns. In some embodiments, the width of the fluid channels incorporated into the cartridge is between about 200 microns and 2 mm. In some embodiments, the width of the fluid channels incorporated into the cartridge is between about 2 mm and 10 mm. In some embodiments, the width of the fluid channels incorporated into the cartridge is between about 10 mm and 20 mm. In some embodiments, the depth of fluid channels incorporated into the cartridge is between about 50 microns and about 2 mm. In some embodiments, the depth of fluid channels incorporated into the cartridge is between about 500 microns and 1 mm. In some embodiments, the depth of fluid channels incorporated into the cartridge is between about 1 mm and about 2 mm. In some embodiments, the one or more flow cells or microwell

array chambers are fabricated from a material selected from the group consisting of silicon, fused-silica, glass, polydimethylsiloxane (PDMS; elastomer), polymethylmethacrylate (PMMA), polycarbonate (PC), polypropylene (PP), polyethylene (PE), high density polyethylene (HDPE), polyimide, cyclic olefin polymers (COP), cyclic olefin copolymers (COC), polyethylene terephthalate (PET), epoxy resin, metal, or a combination of these materials. In some embodiments, the device comprises a removable, consumable component of an instrument system configured to perform automated multiplexed, single cell stochastic labeling and molecular indexing assays. In some embodiments, the cartridge further comprises bypass channels or other design features for providing self-metering of cell samples or bead suspensions dispensed or injected into the cartridge. In some embodiments, the cartridge further comprises integrated miniature pumps for controlling fluid flow through the device. In some embodiments, the cartridge further comprises integrated miniature valves for compartmentalizing pre-loaded reagents and for controlling fluid flow through the device. In some embodiments, the cartridge further comprises vents for providing an escape path for trapped air. In some embodiments, the cartridge further comprises design elements for creating physical or chemical barriers that effectively increase pathlength and prevent or minimize diffusion of molecules between microwells, wherein the design elements are selected from the group consisting of: a pattern of serpentine channels for delivery of cells and beads to the at least first microwell array, a retractable platen or deformable membrane that is pressed into contact with the surface of the at least first microwell array, or the release of an immiscible, hydrophobic fluid from a reservoir within the cartridge. In some embodiments, the cartridge further comprises integrated temperature control components or an integrated thermal interface for providing good thermal contact with an external instrument system. In some embodiments, the cartridge further comprises an optical interface or window for optical imaging of the at least first microwell array. In some embodiments, the cartridge further comprises one or more removable sample collection chambers that are configured to interface with stand-alone PCR thermal cyclers and/or sequencing instruments. In some embodiments, the cartridge itself is configured to interface directly with stand-alone PCR thermal cyclers and/or sequencing instruments.

[00768] In one aspect the disclosure provides for an instrument system comprising: at least a first flow cell or cartridge further comprising at least a first microwell array; and a flow controller; wherein the flow controller controls the delivery of cell samples, bead-based oligonucleotide labeling reagents, and other assay reagents to the at least first microwell

array, and the instrument system is configured to perform multiplexed, single cell stochastic labeling and molecular indexing assays.

[00769] In some embodiments, the at least first microwell array as described herein. In some embodiments, the at least first flow cell is a fixed component of the system. In some embodiments, the at least first flow cell is a removable, consumable component of the system. In some embodiments, the at least first cartridge is a removable, consumable component of the system. In some embodiments, cell samples and bead-based oligonucleotide reagents are dispensed or injected directly into the cartridge by the user. In some embodiments, assay reagents other than cell samples are preloaded in the cartridge. In some embodiments, the instrument system further comprises an imaging system for imaging the at least first microwell array. In some embodiments, the instrument system further comprises a cell or bead distribution system for facilitating uniform distribution of cells and beads across the at least first microwell array, wherein the mechanism underlying said distribution system is selected from the group consisting of rocking, shaking, swirling, recirculating flow, low frequency agitation, or high frequency agitation. In some embodiments, the instrument system further comprises a cell lysis system wherein the system uses a high frequency piezoelectric transducer for sonicating the cells. In some embodiments, the instrument system further comprises a temperature controller for maintaining a user-specified temperature, or for ramping temperature between two or more specified temperatures over two or more specified time intervals. In some embodiments, the instrument system further comprises a magnetic field controller for use in eluting beads from microwells. In some embodiments, the instrument system further comprises a computer or processor programmed to provide a user interface and control of system functions. In some embodiments, the instrument system further comprises program code for providing real-time image analysis capability. In some embodiments, the real-time image analysis and instrument control functions are coupled, so that cell and bead sample loading steps can be prolonged or repeated until optimal cell/bead distributions are achieved. In some embodiments, the instrument system further comprises an integrated PCR thermal cycler for amplification of oligonucleotide labels. In some embodiments, the instrument system further comprises an integrated sequencer for sequencing of oligonucleotide libraries, thereby providing sample-to-answer capability. In some embodiments, the cell samples comprise patient samples and the results of the multiplexed, single cell stochastic labeling and molecular indexing assay are used for clinical diagnostic applications. In some embodiments, the cell samples comprise patient samples and the results of the multiplexed, single cell stochastic labeling and

molecular indexing assay are used by a healthcare provider to make informed healthcare treatment decisions.

[00770] In one aspect the disclosure provides for software residing in a computer readable medium programmed to perform one or more of the following sequence data analysis functions: determining the number of reads per gene per cell, and the number of unique transcript molecules per gene per cell; principal component analysis or other statistical analysis to predict confidence intervals for determinations of the number of transcript molecules per gene per cell; alignment of gene sequence data with known reference sequences; decoding/demultiplexing of sample barcodes, cell barcodes, and molecular barcodes; and automated clustering of molecular labels to compensate for amplification or sequencing errors; wherein the sequence data is generated by performing multiplexed, single cell stochastic labeling and molecular indexing assays.

[00771] In one aspect the disclosure provides for a composition comprising: a solid support, wherein the solid support comprises a plurality of oligonucleotides, wherein at least two of the plurality of oligonucleotides comprises a cellular label and a molecular label, wherein the cellular labels of the at least two of the plurality of oligonucleotides are the same, and wherein the molecular labels of the at least two of the plurality of oligonucleotides are different.

[00772] In some embodiments, the plurality of oligonucleotide further comprises a sample label. In some embodiments, the plurality of oligonucleotides further comprises a target binding region. In some embodiments, the target binding region comprises a sequence is adapted to hybridize to a target nucleic acid. In some embodiments, the target nucleic acid comprises a plurality of target nucleic acids comprising at least 0.01%, 0.02%, 0.05%, 0.1%, 0.2%, 0.3%, 0.4%, 0.5%, 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 60%, 70%, 80%, 90%, or 100% of the transcripts of a transcriptome of an organism. In some embodiments, the target nucleic acid is DNA. In some embodiments, the target nucleic acid is RNA. In some embodiments, the target nucleic acid is mRNA. In some embodiments, the DNA is genomic DNA. In some embodiments, the genomic DNA is sheared. In some embodiments, the sheared genomic DNA comprises at least 0.01%, 0.02%, 0.05%, 0.1%, 0.2%, 0.3%, 0.4%, 0.5%, 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 60%, 70%, 80%, 90%, or 100% of the genes of a genome of an organism. In some embodiments, the target binding region comprises a sequence selected from the group consisting of: a random multimer *e.g.*, a

random dimer, trimer, quatramer, pentamer, hexamer, septamer, octamer, nonamer, decamer, or higher multimer sequence of any length; a gene-specific primer; and oligo dT; or any combination thereof. In some embodiments, the plurality of oligonucleotides comprises a universal label. In some embodiments, the universal label comprises a binding site for a sequencing primer. In some embodiments, the plurality of oligonucleotides comprises a linker. In some embodiments, the linker comprises a functional group. In some embodiments, the linker is located 5' to the oligonucleotide. In some embodiments, the linker is selected from the group consisting of: C6, biotin, streptavidin, primary amines, aldehydes, and ketones, or any combination thereof. In some embodiments, the solid support is comprised of polystyrene. In some embodiments, in solid support is magnetic. In some embodiments, the solid support is selected from the group consisting of: a PDMS solid support, a glass solid support, a polypropylene solid support, an agarose solid support, a gelatin solid support, a magnetic solid support, and a pluronic solid support, or any combination thereof. In some embodiments, the solid support comprises a bead. In some embodiments, the solid support comprises a diameter of about 20 microns. In some embodiments, the solid support comprises a diameter from about 5 microns to about 40 microns. In some embodiments, the solid support comprises a functional group. In some embodiments, the functional group is selected from the group consisting of: C6, biotin, streptavidin, primary amines, aldehydes, and ketones, or any combination thereof. In some embodiments, the cellular label comprises a plurality of cellular labels. In some embodiments, the plurality of cellular labels is interspersed with a plurality of linker label sequences. In some embodiments, the plurality of oligonucleotides comprises from 10,000 to 1 billion oligonucleotides. In some embodiments, the plurality of oligonucleotides comprises from 10,000 to 1 billion target binding regions. In some embodiments, the plurality of oligonucleotides comprises from 10,000 to 1 billion different target binding regions. In some embodiments, the plurality of oligonucleotides comprises from 10,000 to 1 billion same target binding regions. In some embodiments, the different target binding regions can hybridize to at least 0.01%, 0.02%, 0.05%, 0.1%, 0.2%, 0.3%, 0.4%, 0.5%, 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 60%, 70%, 80%, 90%, or 100% of the transcripts of a transcriptome of an organism. In some embodiments, the different target binding regions can hybridize to at least 0.01%, 0.02%, 0.05%, 0.1%, 0.2%, 0.3%, 0.4%, 0.5%, 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 60%, 70%, 80%, 90%, or 100% of the transcripts of a transcriptome of an organism.

[00773] In one aspect the disclosure provides for a composition comprising: a solid support, wherein the solid support comprises a plurality of oligonucleotides, wherein at least two of the plurality of oligonucleotides comprises a cellular label and a molecular label, wherein the cellular labels of the at least two of the plurality of oligonucleotides are the same, and wherein the molecular labels of the at least two of the plurality of oligonucleotides are different.

[00774] In some embodiments, the plurality of oligonucleotide further comprises a sample label. In some embodiments, the plurality of oligonucleotides further comprises a target binding region. In some embodiments, the target binding region comprises a sequence is adapted to hybridize to a target nucleic acid. In some embodiments, the target nucleic acid comprises a plurality of target nucleic acids comprising at least 0.01%, 0.02%, 0.05%, 0.1%, 0.2%, 0.3%, 0.4%, 0.5%, 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 60%, 70%, 80%, 90%, or 100% of the transcripts of a transcriptome of an organism. In some embodiments, the target nucleic acid comprises sheared genomic DNA wherein the wherein the sheared genomic DNA comprises at least 0.01%, 0.02%, 0.05%, 0.1%, 0.2%, 0.3%, 0.4%, 0.5%, 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 60%, 70%, 80%, 90%, or 100% of the genes of a genome of an organism. In some embodiments, the target binding region comprises an oligo dT. In some embodiments, the at least two of the plurality of oligonucleotides comprises a first oligonucleotide and a second oligonucleotide, wherein the first oligonucleotide comprises a first cellular label and a first molecular label, wherein the first cellular label comprises a first random sequence, a second random sequence, and a first linker label sequence, wherein the first linker label sequence connects the first random sequence and the second random sequence; and the first molecular label comprises a random sequence; and the second oligonucleotide comprises a second cellular label and a second molecular label, wherein the second cellular label comprises a third random sequence, a fourth random sequence, and a second linker label sequence, wherein the second linker label sequence connects the third random sequence and the fourth random sequence; and the second molecular label comprising a random sequence, and wherein the first cellular label and the second cellular label are the same and the first molecular label and the second molecular label are different.

[00775] In one aspect the disclosure provides for a kit comprising any composition described herein and instructions for use.

[00776] In one aspect the disclosure provides for a method, comprising: contacting a sample with a solid support, wherein the solid support comprises a plurality of oligonucleotides, wherein at least two of the plurality of oligonucleotides comprises a cellular label and a molecular label, wherein the cellular labels of the at least two of the plurality of oligonucleotides are the same, and wherein the molecular labels of the at least two of the plurality of oligonucleotides are different; and hybridizing the target nucleic acid from the sample to an oligonucleotide of the plurality of oligonucleotides.

[00777] In some embodiments, the sample comprises cells. In some embodiments, the sample is lysed prior to the hybridizing. In some embodiments, the hybridizing comprising hybridizing multiple copies of a same target nucleic acid to the plurality of oligonucleotides. In some embodiments, the method further comprises reverse transcribing the target nucleic acid. In some embodiments, the method further comprises performing an oligonucleotide amplification. In some embodiments, the amplifying comprises amplification using a method selected from the group consisting of: PCR, quantitative PCR, real-time PCR, and digital PCR, or any combination thereof.

[00778] In one aspect the disclosure provides for a A solid support comprising: a first oligonucleotide comprising: a first cellular label comprising a first random sequence, a second random sequence, and a first linker label sequence, wherein the first linker label sequence connects the first random sequence and the second random sequence; and a first molecular label comprising a random sequence; and a second oligonucleotide comprising: a second cellular label comprising a third random sequence, a fourth random sequence, and a second linker label sequence, wherein the second linker label sequence connects the third random sequence and the fourth random sequence; and a second molecular label comprising a random sequence, wherein the first cellular label and the second cellular label are the same and the first molecular label and the second molecular label are different. In some embodiments, the first and second oligonucleotides further comprise identical sample index regions. In some embodiments, the sample index region comprises a random sequence. In some embodiments, the sample index region is 4-12 nucleotides in length. In some embodiments, the cellular label is directly attached to the molecular label. In some embodiments, the cellular label and the molecular label are attached through a linker label sequence. In some embodiments, the random sequence of the cellular label is from 4-12 nucleotides in length. In some embodiments, the constant sequence of the cellular label is at least 4 nucleotides in length. In some embodiments, the cellular label has a total length of at least 12 nucleotides. In some embodiments, the cellular label further comprises one or more

additional random sequences. In some embodiments, the cellular label further comprises one or more additional linker label sequences. In some embodiments, the one or more additional linker label sequences connect the one or more additional random sequences. In some embodiments, the random sequence of the molecular label is 4-12 nucleotides in length.

[00779] In one aspect the disclosure provides for a composition comprising: a solid support, wherein the solid support comprises a plurality of oligonucleotides, wherein at least two of the plurality of oligonucleotides comprises: a cellular label, a molecular label; and a target binding region; and a plurality of target nucleic acids, wherein the cellular labels of the at least two of the plurality of oligonucleotides are the same, and wherein the molecular labels of the at least two of the plurality of oligonucleotides are different.

[00780] In some embodiments, the target binding region comprises a sequence that is adapted to hybridize to at least one of the plurality of target nucleic acids. In some embodiments, the target binding region comprises a sequence selected from the group consisting of: a random multimer *e.g.*, a random dimer, trimer, tetramer, pentamer, hexamer, heptamer, octamer, nonamer, decamer, or higher multimer sequence of any length; a gene-specific primer; and oligo dT; or any combination thereof. In some embodiments, the plurality of oligonucleotides comprises from 10,000 to 1 billion oligonucleotides. In some embodiments, the plurality of oligonucleotides comprises a number of oligonucleotides greater than the number of target nucleic acids of the plurality of target nucleic acids. In some embodiments, the plurality of target nucleic acids comprises multiple copies of a same target nucleic acid. In some embodiments, the plurality of target nucleic acids comprises multiple copies of different target nucleic acids. In some embodiments, the plurality of target nucleic acids are bound to the plurality of oligonucleotides. In some embodiments, the oligonucleotide further comprises a sample label. In some embodiments, the plurality of oligonucleotides comprises a universal label. In some embodiments, the universal label comprises a binding site for a sequencing primer. In some embodiments, the plurality of oligonucleotides comprises a linker. In some embodiments, the linker comprises a functional group. In some embodiments, the linker is located 5' to the oligonucleotide. In some embodiments, the functional group comprises an amino group. In some embodiments, the linker is selected from the group consisting of: C6, biotin, streptavidin, primary amines, aldehydes, and ketones, or any combination thereof. In some embodiments, the solid support is comprised of polystyrene. In some embodiments, the solid support is magnetic. In some embodiments, the solid support is selected from the group consisting of: a PDMS solid support, a glass solid support, a polypropylene solid support, an agarose solid support, a

gelatin solid support, a magnetic solid support, and a pluronic solid support, or any combination thereof. In some embodiments, the solid support comprises a bead. In some embodiments, the solid support comprises a diameter of about 20 microns. In some embodiments, the solid support comprises a diameter from about 5 microns to about 40 microns. In some embodiments, the solid support comprises a functional group. In some embodiments, the functional group comprises a carboxy group. In some embodiments, the functional group is selected from the group consisting of: C6, biotin, streptavidin, primary amines, aldehydes, and ketones, or any combination thereof. In some embodiments, the cellular label comprises a plurality of cellular labels. In some embodiments, the plurality of cellular labels is interspersed with a plurality of linker label sequences.

[00781] In one aspect the disclosure provides for a kit comprising: a first solid support, wherein the first solid support comprises a first plurality of oligonucleotides, wherein the first plurality of oligonucleotides comprises a same first cellular label, a second solid support, wherein the second solid support comprises a second plurality of oligonucleotides, wherein the second plurality of oligonucleotides comprises a same second cellular label, and instructions for use, wherein the first cellular label and the second cellular label are different.

[00782] In some embodiments, oligonucleotides from the first plurality of oligonucleotides and the second plurality of oligonucleotides comprises a molecular label. In some embodiments, the molecular label of the oligonucleotides are different. In some embodiments, the molecular label of the oligonucleotides are the same. In some embodiments, the molecular label of some of the oligonucleotides are different and some are the same. In some embodiments, oligonucleotides from the first plurality of oligonucleotides and the second plurality of oligonucleotides comprise a target binding region. In some embodiments, the kit further comprises a microwell array. In some embodiments, the kit further comprises a buffer. In some embodiments, the buffer is selected from the group consisting of: a reconstitution buffer, a dilution buffer, and a stabilization buffer, or any combination thereof.

[00783] In one aspect the disclosure provides for a method for determining an amount of a target nucleic acid comprising: contacting a sample with a solid support, wherein the solid support comprises a plurality of oligonucleotides, wherein at least two of the plurality of oligonucleotides comprises a cellular label and a molecular label, wherein the cellular labels of the at least two of the plurality of oligonucleotides are the same, and wherein the molecular labels of the at least two of the plurality of oligonucleotides are different; and

hybridizing the target nucleic acid from the sample to an oligonucleotide of the plurality of oligonucleotides.

[00784] In some embodiments, the sample comprises cells. In some embodiments, the sample is lysed prior to the hybridizing. In some embodiments, the hybridizing comprising hybridizing multiple copies of a same target nucleic acid to the plurality of oligonucleotides. In some embodiments, the method further comprises amplifying the target nucleic acid. In some embodiments, the amplifying comprises reverse transcribing the target nucleic acid. In some embodiments, the amplifying comprises amplification using a method selected from the group consisting of: PCR, quantitative PCR, real-time PCR, and digital PCR, or any combination thereof. In some embodiments, the amplifying is performed directly on the solid support. In some embodiments, the amplifying is performed on a template transcribed from the solid support. In some embodiments, the method further comprises sequencing the target nucleic acid. In some embodiments, the sequencing comprises sequencing the target nucleic acid and the molecular label. In some embodiments, the method further comprises determining an amount of the target nucleic acid. In some embodiments, the determining comprises quantifying levels of the target nucleic acid. In some embodiments, the determining comprises counting the number of sequenced molecular labels for the target nucleic acid. In some embodiments, the contacting occurs in a microwell. In some embodiments, the microwell is made from a material selected from the group consisting of: hydrophilic plastic, plastic, elastomer, and hydrogel, or any combination thereof. In some embodiments, the microwell comprises agarose. In some embodiments, the microwell is one microwell of a microwell array. In some embodiments, the microwell array comprises at least 90 microwells. In some embodiments, the microwell array comprises at least 150,000 microwells. In some embodiments, the microwell comprises at least one solid support per well. In some embodiments, the microwell comprises at most two solid supports per well. In some embodiments, the microwell is of a size that accommodates at most two of the solid support. In some embodiments, the microwell is of a size that accommodates at most one solid support. In some embodiments, the microwell is at least 25 microns deep. In some embodiments, the microwell is at least 25 microns in diameter.

[00785] In one aspect the disclosure provides for a method to reduce amplification bias of a target nucleic acid comprising: contacting a sample to a solid support, wherein the solid support comprises a plurality of oligonucleotides, wherein at least two of the plurality of oligonucleotides comprises a cellular label and a molecular label, wherein the cellular labels of the at least two of the plurality of oligonucleotides are the same, and wherein the

molecular labels of the at least two of the plurality of oligonucleotides are different; and hybridizing a target nucleic acid from the sample to the plurality of oligonucleotides; amplifying the target nucleic acid or complement thereof. sequencing the target nucleic acid or complement thereof, wherein the sequencing sequences the target nucleic acid or complement thereof and the molecular label of the oligonucleotide to which the target nucleic acid or complement thereof is bound. determining an amount of the target nucleic acid.

[00786] In some embodiments, the hybridizing comprising hybridizing multiple copies of a same target nucleic acid to the plurality of oligonucleotides. In some embodiments, the determining comprises counting a number of sequenced molecular labels for a same target nucleic acid. In some embodiments, the counting counts the number of copies of the same target nucleic acid. In some embodiments, the sample comprises cells. In some embodiments, the amplifying comprises reverse transcribing the target nucleic acid. In some embodiments, the amplifying comprises amplification using a method selected from the group consisting of: PCR, quantitative PCR, real-time PCR, and digital PCR, or any combination thereof. In some embodiments, the amplifying is performed directly on the solid support. In some embodiments, the amplifying is performed on a template transcribed from the solid support.

[00787] In one aspect the disclosure provides for a composition comprising a microwell; a cell; and a solid support, wherein the solid support comprises a plurality of oligonucleotides, wherein at least two of the plurality of oligonucleotides comprises a cellular label and a molecular label, wherein the cellular labels of the at least two of the plurality of oligonucleotides are the same, and wherein the molecular labels of the at least two of the plurality of oligonucleotides are different.

[00788] In some embodiments, the at least two of the plurality of oligonucleotides further comprises a sample label. In some embodiments, the at least two of the plurality of oligonucleotides further comprises a target binding region. In some embodiments, the target binding region comprises a sequence selected from the group consisting of: a random multimer *e.g.*, a random dimer, trimer, tetramer, pentamer, hexamer, septamer, octamer, nonamer, decamer, or higher multimer sequence of any length; a gene-specific primer; and oligo dT; or any combination thereof. In some embodiments, the plurality of oligonucleotides comprises a universal label. In some embodiments, the universal label comprises a binding site for a sequencing primer. In some embodiments, the solid support is comprised of polystyrene. In some embodiments, in solid support is magnetic. In some embodiments, the solid support is selected from the group consisting of: a PDMS solid support, a glass solid support, a polypropylene solid support, an agarose solid support, a gelatin solid support, a

magnetic solid support, and a pluronic solid support, or any combination thereof. In some embodiments, the solid support comprises a bead. In some embodiments, the solid support has a diameter of about 20 microns. In some embodiments, the solid support has a diameter from about 5 microns to about 40 microns. In some embodiments, the cellular label comprises a plurality of cellular labels. In some embodiments, the plurality of cellular labels is interspersed with a plurality of linker sequences. In some embodiments, the microwell is made from a material selected from the group consisting of: hydrophilic plastic, plastic, elastomer, and hydrogel, or any combination thereof. In some embodiments, the microwell comprises agarose. In some embodiments, the microwell is a microwell of a microwell array. In some embodiments, the microwell comprises at least one solid support per well. In some embodiments, the microwell comprises at most two solid supports per well. In some embodiments, the microwell is of a size that accommodates at least one of the solid support and at least one of the cell. In some embodiments, the microwell is of a size that accommodates at most one of the solid support and at least one of the cell. In some embodiments, the microwell is at least 25 microns deep. In some embodiments, the microwell is at least 25 microns in diameter. In some embodiments, the microwell is flat

[00789] In one aspect the disclosure provides for a device, comprising a plurality of microwells, wherein the plurality of microwells comprises at least two microwells; and wherein each microwell of the plurality of microwells has a volume ranging from about 1,000 μm^3 to about 120,000 μm^3 . In some embodiments, each microwell of the plurality of microwells has a volume of about 20,000 μm^3 . In some embodiments, the plurality of microwells comprises from about 1,000 to about 5,000,000 microwells. In some embodiments, the plurality of microwells comprises about 100,000 to about 200,000 microwells. In some embodiments, the microwells are comprised in a single layer of a material. In some embodiments, at least about 10 % of the microwells further comprise a cell. In some embodiments, at least about 10% of the microwells further comprise a solid support which comprises a plurality of oligonucleotides, wherein at least two of the plurality of oligonucleotides comprise a cellular label and a molecular label, wherein the cellular labels of the at least two of the plurality of oligonucleotides are the same, and wherein the molecular labels of the at least two of the plurality of oligonucleotides are different. In some embodiments, the solid supports are magnetized.

[00790] In one aspect the disclosure provides for an apparatus comprising any device described herein, and a liquid handler.

[00791] In some embodiments, the liquid handler delivers liquid to the plurality of microwells in about 1 second. In some embodiments, the apparatus delivers liquid to the plurality of microwells from a single input port. In some embodiments, the apparatus further comprises a magnet. In some embodiments, the apparatus further comprises at least one of: an inlet port, an outlet port, a pump, a valve, a vent, a reservoir, a sample collection chamber, a temperature control apparatus, or any combination thereof. In some embodiments, the apparatus comprises the sample collection chamber, wherein the sample collection chamber is removable from the apparatus. In some embodiments, the apparatus further comprises an optical imager. In some embodiments, the optical imager produces an output signal which is used to control the liquid handler. In some embodiments, the apparatus further comprises a thermal cycling mechanism configured to perform polymerase chain reaction (PCR) amplification of oligonucleotides.

[00792] In one aspect the disclosure provides for a method of producing a clinical diagnostic test result, comprising producing the clinical diagnostic test result with any device or apparatus described herein. In some embodiments, the clinical diagnostic test result is transmitted via a communication medium.

[00793] In one aspect the disclosure provides for a device comprising: one or more substrates further comprising one or more microwell arrays; wherein the microwell arrays are used to perform multiplexed, single cell stochastic labeling and molecular indexing assays.

[00794] In some embodiments, the microwell arrays of the substrates comprise microwells arranged in a one dimensional or two dimensional array pattern. In some embodiments, the two dimensional array pattern of microwells is selected from the group including a square grid, a rectangular grid, or a hexagonal grid.

[00795] In some embodiments, the microwells of the microwell arrays are fabricated using a well geometry selected from the group including cylindrical, conical, hemispherical, rectangular, or polyhedral. In some embodiments, the microwells of the microwell arrays are fabricated using a overall geometry that comprises two or more component geometries selected from the group including cylindrical, conical, hemispherical, rectangular, or polyhedral. In some embodiments, the diameter of the microwells in the microwell arrays is between about 5 microns and about 50 microns. In some embodiments, the depth of the microwells in the microwell arrays is between about 10 microns and about 60 microns. In some embodiments, the center-to-center spacing between microwells in the microwell arrays is between about 15 microns and about 75 microns. In some embodiments, the total number of microwells in each of the microwell arrays is between about 96 and about 5,000,000. In some

embodiments, the one or more substrates are fabricated from a material selected from the group including silicon, fused-silica, glass, a polymer, or a metal. In some embodiments, the one or more substrates are fabricated from agarose or a hydrogel. In some embodiments, the microwell arrays further comprise surface features between microwells that surround the microwells or straddle the surface between microwells, and are selected from the group including domed, ridged, or peaked surface features.

[00796] In one aspect the disclosure provides for a device comprising: a substrate further comprising one or more microwell arrays; and a mechanical fixture comprising a top plate, a bottom plate, and a gasket; wherein when assembled the substrate is positioned between the gasket and the bottom plate, the gasket forms a leak-proof seal with the substrate, and the top plate and gasket form one or more chambers encompassing the microwell arrays such that one or more cell samples and bead-based oligonucleotide labels may be dispensed into the chambers for the purpose of performing multiplexed, single cell stochastic labeling and molecular indexing assays.

[00797] In some embodiments, the substrate comprises any one or more microwell arrays as described herein. In some embodiments, the gasket is fabricated from polydimethylsiloxane (PDMS) or a similar elastomeric material. In some embodiments, the top and bottom plates are fabricated from aluminum, anodized aluminum, stainless steel, teflon, polymethylmethacrylate, polycarbonate, or a similar rigid polymer material.

[00798] In one aspect the disclosure provides for a device comprising: one or more substrates further comprising one or more microwell arrays; and one or more flow cells; wherein the one or more flow cells enclose or are attached to the one or more substrates, and include at least one inlet port and at least one outlet port for the purpose of delivering fluids to the microwell arrays; and wherein the device is used to perform multiplexed, single cell stochastic labeling and molecular indexing assays.

[00799] In some embodiments, the one or more substrates comprise any one or more microwell arrays as described herein. In some embodiments, each of the one or more flow cells further comprise a plurality of microarray chambers that interface with a plurality of microwell arrays such that one or more samples may be processed in parallel. In some embodiments, the one or more flow cells further comprise a porous barrier or flow diffuser to provide more uniform delivery of cells and beads to the microwell arrays. In some embodiments, the one or more flow cells further comprise dividers that divide chambers containing microwell arrays into subsections that collectively cover the same total array area and provide for more uniform delivery of cells and beads to the microwell arrays. In some

embodiments, the width of fluid channels incorporated into the device is between about 50 microns and 20 mm. In some embodiments, the depth of fluid channels incorporated into the device is between about 50 microns and about 2 mm. In some embodiments, the one or more flow cells are fabricated from a material selected from the group consisting of silicon, fused-silica, glass, polydimethylsiloxane (PDMS; elastomer), polymethylmethacrylate (PMMA), polycarbonate (PC), polypropylene (PP), polyethylene (PE), high density polyethylene (HDPE), polyimide, cyclic olefin polymers (COP), cyclic olefin copolymers (COC), polyethylene terephthalate (PET), epoxy resin, metal, or a combination of these materials. In some embodiments, the device comprises a fixed component of an instrument system for performing automated multiplexed, single cell stochastic labeling and molecular indexing assays. In some embodiments, the device comprises a removable component of an instrument system for performing automated multiplexed, single cell stochastic labeling and molecular indexing assays.

[00800] In one aspect the disclosure provides for a cartridge comprising: one or more substrates further comprising one or more microwell arrays; one or more flow cells or microwell array chambers; one or more sample or reagent reservoirs; and wherein the cartridge further comprises at least one inlet port and at least one outlet port for the purpose of delivering fluids to the microwell arrays; and wherein the cartridge is used to perform multiplexed, single cell stochastic labeling and molecular indexing assays.

[00801] In some embodiments, the one or more substrates comprise any one or more microwell arrays as described herein. In some embodiments, the one or more flow cells or microwell array chambers interface with a plurality of microwell arrays such that one or more samples may be processed in parallel. In some embodiments, the one or more flow cells or microwell array chambers further comprise a porous barrier or flow diffuser to provide more uniform delivery of cells and beads to the microwell arrays. In some embodiments, the one or more flow cells or microwell array chambers further comprise dividers that divide the flow cells or chambers into subsections that collectively cover the same total array area and provide for more uniform delivery of cells and beads to the microwell arrays. In some embodiments, the width of fluid channels incorporated into the cartridge is between about 50 microns and 200 microns. In some embodiments, the width of the fluid channels incorporated into the cartridge is between about 200 microns and 2 mm. In some embodiments, the width of the fluid channels incorporated into the cartridge is between about 2 mm and 10 mm. In some embodiments, the width of the fluid channels incorporated into the cartridge is between about 10 mm and 20 mm. In some embodiments, the depth of fluid channels incorporated

into the cartridge is between about 50 microns and about 10 mm. In some embodiments, the depth of fluid channels incorporated into the cartridge is between about 500 microns and 1 mm. In some embodiments, the depth of fluid channels incorporated into the cartridge is between about 1 mm and about 2 mm. In some embodiments, the one or more flow cells or microwell array chambers are fabricated from a material selected from the group consisting of silicon, fused-silica, glass, polydimethylsiloxane (PDMS; elastomer), polymethylmethacrylate (PMMA), polycarbonate (PC), polypropylene (PP), polyethylene (PE), high density polyethylene (HDPE), polyimide, cyclic olefin polymers (COP), cyclic olefin copolymers (COC), polyethylene terephthalate (PET), epoxy resin, metal, or a combination of these materials. In some embodiments, the device comprises a removable, consumable component of an instrument system for performing automated multiplexed, single cell stochastic labeling and molecular indexing assays. In some embodiments, the cartridge further comprises bypass channels or other design features for providing self-metering of cell samples or bead suspensions dispensed or injected into the cartridge. In some embodiments, the cartridge further comprises integrated miniature pumps for controlling fluid flow through the device. In some embodiments, the cartridge further comprises integrated miniature valves for compartmentalizing pre-loaded reagents and for controlling fluid flow through the device. In some embodiments, the cartridge further comprises vents for providing an escape path for trapped air. In some embodiments, the cartridge further comprises design elements for creating physical or chemical barriers that effectively increase pathlength and prevent or minimize diffusion of molecules between microwells, wherein the design elements are selected from the group consisting of: a pattern of serpentine channels for delivery of cells and beads to the microwell array, a retractable platen or deformable membrane that is pressed into contact with the surface of the microwell array, or the release of an immiscible, hydrophobic fluid from a reservoir within the cartridge. In some embodiments, the cartridge further comprises integrated temperature control components or an integrated thermal interface for providing good thermal contact with an external instrument system. In some embodiments, the cartridge further comprises an optical interface or window for optical imaging of the one or more microwell arrays. In some embodiments, the cartridge further comprises one or more removable sample collection chambers that are configured to interface with stand-alone PCR thermal cyclers and/or sequencing instruments. In some embodiments, the cartridge itself is configured to interface directly with stand-alone PCR thermal cyclers and/or sequencing instruments.

[00802] In one aspect the disclosure provides for an instrument system comprising: one or more flow cells or cartridges further comprising one or more microwell arrays; and a flow controller; wherein the flow controller controls the delivery of cell samples, bead-based oligonucleotide labeling reagents, and other assay reagents to the microwell arrays, and the instrument system is used to perform multiplexed, single cell stochastic labeling and molecular indexing assays.

[00803] In some embodiments, the one or more microwell arrays are any described herein. In some embodiments, the one or more flow cells are a fixed component of the system. In some embodiments, the one or more flow cells are a removable, consumable component of the system. In some embodiments, the one or more cartridges are removable, consumable components of the system. In some embodiments, cell samples and bead-based oligonucleotide reagents are dispensed or injected directly into the cartridge by the user. In some embodiments, assay reagents other than cell samples are preloaded in the cartridge. In some embodiments, the instrument system further comprises an imaging system for imaging the microwell arrays. In some embodiments, the instrument system further comprises a cell or bead distribution system for facilitating uniform distribution of cells and beads across the microwell arrays, wherein the mechanism underlying the distribution system is selected from the group consisting of rocking, shaking, swirling, recirculating flow, low frequency agitation, or high frequency agitation. In some embodiments, the instrument system further comprises a cell lysis system wherein the system uses a high frequency piezoelectric transducer for somicating the cells. In some embodiments, the instrument system further comprises a temperature controller for maintaining a user-specified temperature, or for ramping temperature between two or more specified temperatures over two or more specified time intervals. In some embodiments, the instrument system further comprises a magnetic field controller for use in eluting beads from microwells. In some embodiments, the instrument system further comprises a computer or processor programmed to provide a user interface and control of system functions. In some embodiments, the instrument system further comprises program code for providing real-time image analysis capability. In some embodiments, the real-time image analysis and instrument control functions are coupled, so that cell and bead sample loading steps can be prolonged or repeated until optimal cell/bead distributions are achieved. In some embodiments, the instrument system further comprises an integrated PCR thermal cycler for amplification of oligonucleotide labels. In some embodiments, the instrument system further comprises an integrated sequencer for sequencing of oligonucleotide libraries, thereby providing sample-to-answer capability. In

some embodiments, the cell samples comprise patient samples and the results of the multiplexed, single cell stochastic labeling and molecular indexing assay are used for clinical diagnostic applications. In some embodiments, the cell samples comprise patient samples and the results of the multiplexed, single cell stochastic labeling and molecular indexing assay are used by a healthcare provider to make informed healthcare treatment decisions.

[00804] In one aspect the disclosure provides for software residing in a computer readable medium programmed to perform one or more of the following sequence data analysis: determining the number of reads per gene per cell, and the number of unique transcript molecules per gene per cell; principal component analysis or other statistical analysis to predict confidence intervals for determinations of the number of transcript molecules per gene per cell; alignment of gene sequence data with known reference sequences; decoding/demultiplexing of sample barcodes, cell barcodes, and molecular barcodes; and automated clustering of molecular labels to compensate for amplification or sequencing errors; wherein the sequence data is generated by performing multiplexed, single cell stochastic labeling and molecular indexing assays.

CLAIMS**WHAT IS CLAIMED IS:**

1. A method, comprising
 - (a) obtaining a sample comprising a plurality of cells;
 - (b) labeling at least a portion of two or more polynucleotide molecules, complements thereof, or reaction products therefrom, from a first cell of the plurality and a second cell of the plurality with
 - (i) a first same cell label specific to the first cell and a second same cell label specific to the second cell; and
 - (ii) a molecular label specific to each of the two or more polynucleotide molecules, complements thereof, or reaction products therefrom, wherein each molecular label of the two or more polynucleotide molecules, complements thereof, or reaction products therefrom, from the first cell are unique with respect to each other, and wherein each molecular label of the two or more polynucleotide molecules, complements thereof, or reaction products therefrom, from the second cell are unique with respect to each other.
2. The method of claim 1, further comprising sequencing the at least a portion of two or more polynucleotide molecules, complements thereof, or reaction products therefrom.
3. The method of claim 2, further comprising analyzing sequence data from the sequencing to identify a number of individual molecules of the polynucleotides in a specific one of the cells.
4. The method of claim 1, wherein the cells are cancer cells.
5. The method of claim 1, wherein the cells are infected with viral polynucleotides.
6. The method of claim 1, wherein the cells are bacteria or fungi.
7. The method of claim 2, wherein the sequencing comprises sequencing with read lengths of at least 100 bases.
8. The method of claim 2, wherein the sequencing comprises sequencing with read lengths of at least 500 bases.

9. The method of claim 1, wherein the polynucleotide molecules are mRNAs or micro RNAs, and the complements thereof and reaction products thereof are complements of and reaction products therefrom the mRNAs or micro RNAs.
10. The method of claim 1, wherein the molecular labels are on a bead.
11. The method of claim 1, wherein the label specific to an individual cell is on a bead.
12. The method as recited in claim 1, wherein the label specific to an individual cell and the molecular labels are on beads.
13. The method of any one of claims 1-12, that is performed at least in part in an emulsion.
14. The method of any one of claims 1-12, that is performed at least in part in a well or microwell of an array.
15. The method of any one of claims 1-14, wherein the presence of a polynucleotide that is associated with a disease or condition is detected.
16. The method of any one of claim 15, wherein the disease or condition is a cancer.
17. The method of any one of claims 1-16, wherein at least a portion of a microRNA, complement thereof, or reaction product therefrom is detected.
18. The method of claim 15, wherein the disease or condition is a viral infection.
19. The method of claim 18, wherein the viral infection is from an enveloped virus.
20. The method of claim 18, wherein the viral infection is from a non-enveloped virus.
21. The method of claim 20, wherein the virus contains viral DNA that is double stranded.
22. The method of claim 20, wherein the virus contains viral DNA that is single stranded.
23. The method of claim 18, wherein the virus is selected from the group consisting of a pox virus, a herpes virus, a varicella zoster virus, a cytomegalovirus, an Epstein-Barr virus, a hepatitis virus, a papovavirus, polyomavirus, and any combination thereof.
24. The method of any one of claims 1-23, wherein the first cell is from a person not having a disease or condition and the second cell is from a person having the disease or condition.
25. The method of claim 24, wherein the persons are different.
26. The method of claim 24, wherein the persons are the same but cells are taken at different time points.

27. The method of any one of claims 1-23, wherein the first cell is from a person having the disease or condition and the second cell is from the same person.
28. The method of any one of claims 1-23, wherein the cells in the sample comprise cells from a tissue or organ.
29. The method of any one of claims 1-28, wherein the cells in the sample comprise cells from a thymus, white blood cells, red blood cells, liver cells, spleen cells, lung cells, heart cells, brain cells, skin cells, pancreas cells, stomach cells, cells from the oral cavity, cells from the nasal cavity, colon cells, small intestine cells, kidney cells, cells from a gland, brain cells, neural cells, glial cells, eye cells, reproductive organ cells, bladder cells, gamete cells, human cells, fetal cells, amniotic cells, or any combination thereof.
30. A solid support comprising a plurality of oligonucleotides each comprising a cellular label and a molecular label, wherein each cellular label of the plurality of oligonucleotides are the same, and each molecular label of the plurality of oligonucleotides are different; and wherein
- (a) the solid support is a bead,
 - (b) the cellular label is specific to the solid support,
 - (c) the solid support, when placed at the center of a three dimensional Cartesian coordinate system, has oligonucleotides extending into at least seven of eight octants,
 - or
 - (d) any combination of (a) – (c).
31. The solid support of claim 30, wherein the plurality of oligonucleotides further comprises at least one of:
- (a) a sample label;
 - (b) a universal label; and
 - (c) a target nucleic acid binding region.
32. The solid support of claim 31, wherein the solid support comprises the target nucleic acid binding region, wherein the target nucleic acid binding region comprises a sequence selected from the group consisting of a gene-specific sequence, an oligo-dT sequence, a random multimer, and any combination thereof.
33. The solid support of claim 31 or 32, further comprising a target nucleic acid or complement thereof.

34. The solid support of claim 33, comprising a plurality of target nucleic acids or complements thereof comprising
- (a) from about 0.01% to about 100% of transcripts of a transcriptome of an organism or complements thereof, or
 - (b) from about 0.01% to about 100% of genes of a genome of an organism or complements thereof.
35. The solid support of any one of claims 30-34, wherein
- (a) the cellular labels of the plurality of oligonucleotides comprise a first random sequence connected to a second random sequence by a first label linking sequence; and
 - (b) the molecular labels of the plurality of oligonucleotides comprise random sequences.
36. The solid support of any one of claims 30-35, wherein the solid support is selected from the group consisting of a polydimethylsiloxane (PDMS) solid support, a polystyrene solid support, a glass solid support, a polypropylene solid support, an agarose solid support, a gelatin solid support, a magnetic solid support, a pluronic solid support, and any combination thereof.
37. The solid support of any one of claims 30-35, wherein
- (a) the plurality of oligonucleotides comprise a linker comprising a linker functional group, and
 - (b) the solid support comprises a solid support functional group; wherein the solid support functional group and linker functional group connect to each other.
38. The solid support of claim 37, wherein the linker functional group and the solid support functional group are individually selected from the group consisting of C6, biotin, streptavidin, primary amine(s), aldehyde(s), ketone(s), and any combination thereof.
39. The solid support of any one of claims 30-38, wherein molecular labels of the plurality of oligonucleotides comprise at least 15 nucleotides.
40. A kit comprising the solid support of any one of claims 30-39, and instructions for use.
41. The kit of claim 40, further comprising a well.
42. The kit of claim 41, wherein the well is comprised in an array.
43. The kit of any one of claims 40-42, wherein the well is a microwell.

44. The kit of any one of claims 40-43, further comprising a buffer.
45. An emulsion comprising the solid support of any one of claims 30-39.
46. A composition comprising a well and the solid support of any one of claims 30-39.
47. A composition comprising a cell and the solid support of any one of claims 30-39.
48. The emulsion of claim 45 or composition of claim 46, further comprising a cell.
49. The composition or emulsion of any one of claims 47 or 48, wherein the cell is a single cell.
50. The composition of claim 46, wherein the well is a microwell.
51. The composition of claim 50, wherein the microwell has a volume ranging from about $1,000 \mu\text{m}^3$ to about $120,000 \mu\text{m}^3$.
52. A method, comprising:
 - (a) contacting a sample with the solid support of any one of claims 30-32 or 35-39,
 - (b) hybridizing a target nucleic acid from the sample to an oligonucleotide of the plurality of oligonucleotides.
53. The method of claim 52, further comprising
 - (c) amplifying the target nucleic acid or complement thereof.
54. The method of claim 53, further comprising
 - (d) sequencing the target nucleic acid or complement thereof, wherein the sequencing comprises sequencing the molecular label of the oligonucleotide to which the target nucleic acid or complement thereof is bound.
55. The method of claim 54, further comprising
 - (e) determining an amount of the target nucleic acid or complement thereof, wherein the determining comprises:
 - (i) quantifying levels of the target nucleic acid or complement thereof;
 - (ii) counting a number of sequences comprising the same molecular label;
 - (iii) or a combination of (i) and (ii).
56. The method of claim 54 or 55, wherein the method does not comprise aligning any same molecular labels or any same cellular labels.
57. The method of any one of claims 53-56, wherein the amplifying comprises reverse transcribing the target nucleic acid.

58. The method of any one of claims 53-57, wherein the amplifying employs a method selected from the group consisting of: PCR, nested PCR, quantitative PCR, real time PCR, digital PCR, and any combination thereof.
59. The method of any one of claims 53-58, wherein the amplifying is performed
- (a) directly on the solid support;
 - (b) on a template transcribed from the solid support; or
 - (c) a combination of (a) and (b).
60. The method of any one of claims 52-59, wherein the sample comprises a cell.
61. The method of claim 60, wherein the cell is a single cell.
62. The method of any one of claims 52-61, wherein the contacting occurs in a well.
63. The method of claim 62, wherein the well is a microwell and is contained in an array of microwells.
64. A device, comprising a plurality of microwells, wherein each microwell of the plurality of microwells has a volume ranging from about $1,000 \mu\text{m}^3$ to about $120,000 \mu\text{m}^3$.
65. The device of claim 64, wherein each microwell of the plurality of microwells has a volume of about $20,000 \mu\text{m}^3$.
66. The device of claim 64 or 65, wherein the plurality of microwells comprises from about 96 to about 200,000 microwells.
67. The device of any one of claims 64-66, wherein the microwells are comprised in a layer of a material.
68. The device of any one of claims 64-67, wherein at least about 10 % of the microwells further comprise a cell.
69. Then device of any one of claims 64-69, further comprising the solid support of any one of claims 30-39.
70. An apparatus comprising the device of any one of claims 64-69, and a liquid handler.
71. The apparatus of claim 70, wherein the liquid handler delivers liquid to the plurality of microwells in about one second.
72. The apparatus of claim 70 or 71, wherein the liquid handler delivers liquid to the plurality of microwells from a single input port.

73. The apparatus of any one of claims 70-72, further comprising a magnet.
74. The apparatus of any one of claims 70-73, further comprising at least one of: an inlet port, an outlet port, a pump, a valve, a vent, a reservoir, a sample collection chamber, a temperature control apparatus, or any combination thereof.
75. The apparatus of claim 74, comprising the sample collection chamber, wherein the sample collection chamber is removable from the apparatus.
76. The apparatus of any one of claims 70-75, further comprising an optical imager.
77. The apparatus of claim 76, wherein the optical imager produces an output signal which is used to control the liquid handler.
78. The apparatus of any one of claims 70-77, further comprising a thermal cycling mechanism configured to perform a polymerase chain reaction (PCR) amplification of oligonucleotides.
79. A method of producing a clinical diagnostic test result, comprising producing the clinical diagnostic test result with:
- (a) the device or apparatus of any one of claims 70-78;
 - (b) the solid support of any one of claims 30-39;
 - (c) the method of any one of claims 1-29 or 52-63; or
 - (d) any combination of (a)-(c).
80. The method of claim 79, wherein the clinical diagnostic test result is transmitted via a communication medium.
81. A method of making the solid support of any one of claims 30-39, comprising
- (a) attaching to a solid support: a first polynucleotide comprising
 - (i) a first portion of the cellular label, and
 - (ii) a first linker; and
 - (b) contacting a second polynucleotide comprising
 - (i) a second portion of the cellular label,
 - (ii) a sequence complementary to the first linker, and
 - (iii) the molecular label.
82. The method of claim 81, wherein the third polynucleotide further comprises a target nucleic acid binding region.
83. The kit of any one of claims 40-44, wherein the kit is contained in a package.

84. The kit of claim 83, wherein the package is a box.
85. The kit of claim 83 or 84, wherein the package or box has a volume of 2 cubic feet or less.
86. The kit of claim 83 or 84, wherein the package or box has a volume of 1 cubic foot or less.
87. The method of any one of claims 1-86, wherein an emulsion, microwell, or well contains only one cell.
88. The method of any one of claims 1-87, wherein from 1 to 2,000,000 emulsions, microwells, or wells each contain only one cell.
89. The method of any one of claims 1-88, comprising distributing at most one cell into each emulsion, microwell, or well.
90. The method of any one of claims 1-89, wherein a single solid support and a single cell are distributed to an emulsion, microwell, or well.
91. The method of any one of claims 1-90, wherein from 1 to 2,000,000 emulsions, microwells, or wells each have distributed thereto one cell and one solid support.
92. The method of any one of claims 1-91, comprising distributing at most one solid support per emulsion, microwell, or well.
93. The method of any one of claims 1-92, comprising distributing one solid support and one cell to each of from 1 to 2,000,000 microwells, emulsions, or wells.
94. The method of any one of claims 1-93, wherein cell distribution is random or non-random.
95. The method of any one of claims 1-94, wherein cell distribution is stochastic.
96. The method of any one of claims 1-95, wherein a cell is distributed by a cell sorter.
97. The method of any one of claims 1-96, wherein a cell is distributed by contacting one or more wells, microwells, or emulsions with a dilute solution of cells diluted so that at most one cell is distributed to the one or more wells, microwells, or emulsions.
98. The method of any one of claims 1-97, wherein the target specific regions, target specific regions of the plurality of oligonucleotides, or the target specific region of the two or more polynucleotide molecules, comprise sequences complementary to two or more targets of a target panel.
99. The method of claim 98, wherein the two or more targets of the target panel are biomarkers.

100. The method of claim 99, wherein the biomarkers are biomarkers for a disease or condition.
101. The method of claim 100, wherein the disease or condition is a cancer, an infection, a viral infection, an inflammatory disease, a neurodegenerative disease, a fungal disease, a bacterial infection, or any combination thereof.
102. The method of any one of claims 98-101, wherein the panel comprises from: 2-50,000, 2-40,000, 2-30,000, 2-20,000, 2-10,000, 2-9000, 2-8,000, 2-7,000, 2-6,000, 2-5,000, 2-1,000, 2-800, 2-700, 2-600, 2-500, 2-400, 2-300, 2-200, 2-100, 2-75, 2-50, 2-40, 2-30, 2-20, 2-10, or 2-5 biomarkers.

1/116




FIG. 1

2/116




FIG. 2A

3/116




FIG. 2B




FIG. 2C




FIG. 3




FIG. 4

6/116




FIG. 5

7/116




FIG. 6A




FIG. 6B

8/116




FIG. 6C

well volume	2.12E-14	m ³
bead volume	4.19E-15	m ³
well volume-bead volume	1.70E-14	m ³
well volume-bead volume	1.70E-11	L
#mRNA per cell	2.50E+05	molecules
concentration of mRNA	1.47E+16	molecules per L
concentration of mRNA	2.44E-08	moles per L
concentration of mRNA	24.40	nM

FIG. 7

9/116




FIG. 8A




FIG. 8B




FIG. 8C

10/116




FIG. 9

11/116



FIG. 10

12/116




FIG. 11A




FIG. 11B

13/116




FIG. 11C




FIG. 11D

14/116




FIG. 11E




FIG. 11F

15/116




FIG. 11G




FIG. 11H

16/116




FIG. 11I




FIG. 12A

17/116




FIG. 12B




FIG. 12C

18/116




FIG. 12D




FIG. 12E

19/116




FIG. 12F




FIG. 12G

20/116




FIG. 12H




FIG. 12I

21/116




FIG. 12J

22/116




FIG. 12K

23/116




FIG. 12L