

Allele specific primer designing for SNPs of ACHE gene related to oxidative stress

Rajdeep Kaur

RIMT University Mandigobindgarh, Punjab

Abstract: Pesticides are the chemicals which are used for agricultural and non-agricultural processes such as insecticide, herbicide, fungicide, termiticide, piscicide etc. Pesticides are generally known as chemical or biological substance designed to kill or retard the growth of pests that damage or interfere with the growth of crops, shrubs, trees, timber and other vegetation desired by humans. Pesticide formulations are complex mixtures which contain, besides the active ingredients, several other components, such as solvents, wetting and emulsifying agents, and additives (Hayes, 1991). Practically all chemical pesticides, pose long-term danger to the environment and humans through their persistence in nature and body tissue. Most of the pesticides are non-specific, and may kill life forms that are harmless or useful. Pesticides can also cause deadly diseases like cancer, Alzheimer, Parkinson, skin diseases etc. They are in direct contact with humans through air, food, or when sprayed on crops, fruits etc. They can enter in human body and mixed up. They can cause DNA damage and various type of mutations can also occur. Poisonous effects of pesticides include damage to DNA such as changes or losses of nucleotic bases, and double and single-strand breakage of DNA and sister chromatid; exchanges Damaged DNA can be repaired by the different types of DNA repair mechanisms like BER, NER, mis-match repair, homologous recombinant repair etc. Pesticide formulations are complex mixtures which contain, besides the active ingredients, several other components, such as solvents, wetting and emulsifying agents, and additives. Pesticides cause oxidative stress leading to the generation of free radicals and alterations in antioxidants or free oxygen radical scavenging enzyme systems. Oxidative stress can be defined most simply as the imbalance between the production of free radicals capable of causing peroxidation of the lipid layer of cells and the body's antioxidant defence. Free radicals are defined as atoms or molecules that contain one or more unpaired electrons. The toxicity of many pesticides is associated with the production of free radicals, which are not only toxic themselves, but are also implicated in the pathophysiology of many diseases. Epidemiological studies in humans long-term exposed to a mixture of pesticides (OPs, synthetic PYRs and NMC) have reported stimulated antioxidant enzymes and lipid peroxidation in erythrocytes even in the absence of a decrease in acetylcholinesterase – AChE.

IndexTerms - ACHE, Pesticides, WASP, OPs

Introduction:

Pesticides are the substances used against pests including insects, pathogens, weeds, birds, roundworms and microbes that compete with humans for food, destroy properties or are vectors for diseases. Organophosphates are more abundantly used pesticides and it is more toxic than other classes of pesticides (Al-Salon 1994). Ops that are used now the days in agricultural practices results in an environmental pollution and number of acute and chronic poisoning events. Ops are mainly absorbed through skin. Most OP exert toxicity on target or non target organs through inhibition of activity of acetylcholinesterase in nerve and muscles (Surajudiin et al 2015). The mode of exposure to ops includes the gastrointestinal and inhalatory and dermal one (Hussain 2010). Ops mainly target acetylcholinesterase (ACHE) that hydrolyze acetylcholinesterase. The main effect of Ops involves irreversible inhibition of the activity of ACHE enzyme in blood and nervous system resulting accumulation of acetylcholine and activation of muscarinic and nicotinic receptor which may ultimately lead to death (Yshanid et al 2005). Acetylcholine esterase hydrolyzes the neurotransmitter, acetylcholine at neuromuscular junction's and brain cholinergic synapse. It is mapped to chromosome 7q22 (Ehrlich et al., 1994). ACHE is present in molecular forms which possess same catalytic properties but they are different in their oligomeric assembly and mode of cell attachment. All enzymes are encoded by single ACHE gene. Due to alternative mRNA splicing and post translational associations of catalytic and structural subunits there is structural diversity. The quantity of Ach accumulation depends on the pesticide dose. In human beings there are two genes which encode acetylcholine hydrolyzing enzymes.

Web Based Allele Specific Primer (WASP):

AS PCR is also known as amplification refractory mutation system (ARMS). This technique is a quick and dependable genotyping protocol that requires minimal instruments found in most laboratories. It is based on the extension of primer only when its 3'-end is a perfect complement to the allele present in the input sample. Thus, if a single base polymorphism occurs, the genotyping results can be observed by simply comparing the length of PCR products. To address these difficulties a SNP database and intuitive graphical interface should be integrated with the AS primer design tool while primer destabilizing condition must be considered to ensure effectiveness of primer results. WASP is a web application constructed using Ruby on Rails Framework. The application is integrated into the local SNP database running MySQL database server. This database frequently collects public SNP information and related reference data from public SNP databases. SNPs can be detected using allele-specific PCR primers based on the 3' terminal nucleotide of a primer that corresponds to a specific SNP site. However, reliable discrimination between the alleles is not sufficient to achieve using this described method. To overcome this problem, allele specific primers with an additional base pair change within the three bases closest to the SNP site between alleles have been used. WASP offers a tool for designing AS primers for both SNPs and mutations. By integrating the database for known SNPs (using gene ID or rs number), this tool facilitates the awkward process of getting flanking sequences and other related information from public SNP databases. It takes into account the underlying destabilizing effect to ensure the effectiveness of designed primers. With user-friendly SVG interface, WASP intuitively presents resulting designed primers, which assist users to export or to make further adjustment to the design. This software can be freely accessed at <http://bioinfo.biotech.or.th/WASP>.

Material and methods:

Literature is studied for different SNPs of ACHE gene related to oxidative stress diseases. Allele specific primers have been designed for these SNPs using WASP software.

Results:

The study was carried out to design allele specific primers for various SNP's of ACHE gene.

SNP's of ACHE:

Total 6 SNP were found in different research papers which were related to diseases caused by oxidative stress.

s.no.	rs no.
1	rs17228581
2	rs7636
3	rs3757869
4	rs1799805
5	rs17228669
6	rs2396755

1. rs17228581

Sequence Size: 601

Primer Picking Parameters:

Primer Size Opt: 20 Min: 18 Max: 36

GC% Opt: 50.0 Min: 20.0 Max: 85.0

Tm Opt: 55.0 Min: 40.0 Max: 65.0

Max Tm Diff: 100.0

Max Self Complementarity: 8.0

Max 3' Complementarity: 3.0

Max PolyX in Primer: 3

Number Primer Return: 5

Mismatch at the penultimate primer position: 'Yes'

In-silico PCR Filtering: 'Yes' with Max Product Size: 4000

Mispriming Filtering: 'No'

Oligo 1 :

Wildtype Reverse Primer 5': GCCTCTCCCATGCCCAAT

Mutant Reverse Primer 5': GCCTCTCCCATGCCCAAC

Common Forward Primer 5': AATGTGGGTCTCCTGGAT

Product Size: 224

In-silico PCR Accepted.

2. rs7636

AS Primer Picking Result for rs7636

Sequence Size: 601

Primer Picking Parameters:

Primer Size Opt: 20 Min: 18 Max: 36

GC% Opt: 50.0 Min: 20.0 Max: 85.0

Tm Opt: 55.0 Min: 40.0 Max: 65.0

Max Tm Diff: 100.0

Max Self Complementarity: 8.0

Max 3' Complementarity: 3.0

Max PolyX in Primer: 3

Number Primer Return: 5

Mismatch at the penultimate primer position: 'Yes'

In-silico PCR Filtering: 'Yes' with Max Product Size: 4000

Mispriming Filtering: 'No'

Oligo 1

Wildtype Reverse Primer 5': AACTCGATCTCGTAGCCGTCG

Mutant Reverse Primer 5': AACTCGATCTCGTAGCCGTCA

Common Forward Primer 5': GGTCTGCATTACACAGACT

Product Size: 225

In-silico PCR Accepted

Oligo 2

Wildtype Reverse Primer 5': ACTCGATCTCGTAGCCGTCG

Mutant Reverse Primer 5': ACTCGATCTCGTAGCCGTCA

Common Forward Primer 5': ATTACACAGACTGGCTGCAT

Product Size: 216

In-silico PCR Accepted

Oligo 3

Wildtype Reverse Primer 5': CTCGATCTCGTAGCCGTCG

Mutant Reverse Primer 5': CTCGATCTCGTAGCCGTCA

Common Forward Primer 5': CTGCATTACACAGACTGGC

Product Size: 219

In-silico PCR Accepted

Oligo 4 :

Wildtype Reverse Primer 5': TCGATCTCGTAGCCGTCG

Mutant Reverse Primer 5': TCGATCTCGTAGCCGTCA
 Common Forward Primer 5': GTGGTCCTGCATTACACAG
 Product Size: 224
 In-silico PCR Accepted

3. rs3757869

AS Primer Picking Result for rs3757869
 Sequence Size: 601
 Primer Picking Parameters:
 Primer Size Opt: 20 Min: 18 Max: 36
 GC% Opt: 50.0 Min: 20.0 Max: 85.0
 Tm Opt: 55.0 Min: 40.0 Max: 65.0
 Max Tm Diff: 100.0
 Max Self Complementarity: 8.0
 Max 3' Complementarity: 3.0
 Max PolyX in Primer: 3
 Number Primer Return: 5
 Mismatch at the penultimate primer position: 'Yes'
 In-silico PCR Filtering: 'Yes' with Max Product Size: 4000
 Mispriming Filtering: 'No'

Oligo 1

Wildtype Forward Primer 5': CATGTGGGAGTGTGGCAGAAG
 Mutant Forward Primer 5': CATGTGGGAGTGTGGCAGAAT
 Common Reverse Primer 5': CCACCTCATTCCACTAGC
 Product Size: 144
 In-silico PCR Accepted:

Oligo 2

Wildtype Forward Primer 5': ATGTGGGAGTGTGGCAGAAG
 Mutant Forward Primer 5': ATGTGGGAGTGTGGCAGAAT
 Common Reverse Primer 5': ACCTCATTCCACTAGCGAT
 Product Size: 141
 In-silico PCR Accepted

Oligo 3

Wildtype Forward Primer 5': TGTGGGAGTGTGGCAGAAG
 Mutant Forward Primer 5': TGTGGGAGTGTGGCAGAAT
 Common Reverse Primer 5': ACCTCATTCCACTAGCGA
 Product Size: 140
 In-silico PCR Accepted: There is only one PCR product in the current human genome.

Oligo 4

Wildtype Forward Primer 5': GTGGGAGTGTGGCAGAAG
 Mutant Forward Primer 5': GTGGGAGTGTGGCAGAAT
 Common Reverse Primer 5': ACCTCATTCCACTAGCGA
 Product Size: 139
 In-silico PCR Accepted.

4. rs1799805

AS Primer Picking Result for rs1799805
 Sequence Size: 601
 Primer Picking Parameters:
 Primer Size Opt: 20 Min: 18 Max: 36
 GC% Opt: 50.0 Min: 20.0 Max: 85.0
 Tm Opt: 55.0 Min: 40.0 Max: 65.0
 Max Tm Diff: 100.0
 Max Self Complementarity: 8.0
 Max 3' Complementarity: 3.0
 Max PolyX in Primer: 3
 Number Primer Return: 5
 Mismatch at the penultimate primer position: 'Yes'
 In-silico PCR Filtering: 'Yes' with Max Product Size: 4000
 Mispriming Filtering: 'No'

Oligo 1 :

Wildtype Forward Primer 5': TCATCAACGCGGGAGACTTTA
 Mutant Forward Primer 5': TCATCAACGCGGGAGACTTTC
 Common Reverse Primer 5': GAGGGATGCAGAGAAAGAG
 Product Size: 237
 In-silico PCR Accepted

Oligo 2

Wildtype Forward Primer 5': CATCAACGCGGGAGACTTTA
 Mutant Forward Primer 5': CATCAACGCGGGAGACTTTC
 Common Reverse Primer 5': AGGGATGCAGAGAAAGAGA

Product Size: 235
 In-silico PCR Accepted
 Oligo 3 :
 Wildtype Forward Primer 5': ATCAACGCGGGAGACTTTA
 Mutant Forward Primer 5': ATCAACGCGGGAGACTTTC
 Common Reverse Primer 5': AGGGATGCAGAGAAAGAGAA
 Product Size: 234
 In-silico PCR Accepted

Oligo 4 :
 Wildtype Forward Primer 5': TCAACGCGGGAGACTTTA
 Mutant Forward Primer 5': TCAACGCGGGAGACTTTC
 Common Reverse Primer 5': GAGGACTTCTGGGACTTCTG
 Product Size: 151
 In-silico PCR Accepted.

5. rs17228609

AS Primer Picking Result for rs17228609

Sequence Size: 601

Primer Picking Parameters:

Primer Size Opt: 20 Min: 18 Max: 36
 GC% Opt: 50.0 Min: 20.0 Max: 85.0
 Tm Opt: 55.0 Min: 40.0 Max: 65.0

Max Tm Diff: 100.0

Max Self Complementarity: 8.0

Max 3' Complementarity: 3.0

Max PolyX in Primer: 3

Number Primer Return: 5

Mismatch at the penultimate primer position: 'Yes'

In-silico PCR Filtering: 'Yes' with Max Product Size: 4000

Mispriming Filtering: 'No'

Oligo 1 :

Wildtype Reverse Primer 5': AGGGCTGGGCTATAACACAGAAG

Mutant Reverse Primer 5': AGGGCTGGGCTATAACACAGAAA

Common Forward Primer 5': GCTGTAACAGTTTATTGGCA

Oligo 2 :

Wildtype Reverse Primer 5': GGGCTGGGCTATAACACAGAAG

Mutant Reverse Primer 5': GGGCTGGGCTATAACACAGAAA

Common Forward Primer 5': CCGTGGCTGTAACAGTTTAT

Product Size: 158

In-silico PCR Accepted

Oligo 3 :

Wildtype Reverse Primer 5': GGCTGGGCTATAACACAGAAG

Mutant Reverse Primer 5': GGCTGGGCTATAACACAGAAA

Common Forward Primer 5': CCGTGGCTGTAACAGTTTAT

Product Size: 157

In-silico PCR Accepted

Oligo 4 :

Wildtype Reverse Primer 5': GCTGGGCTATAACACAGAAG

Mutant Reverse Primer 5': GCTGGGCTATAACACAGAAA

Common Forward Primer 5': GCTGTAACAGTTTATTGGCA

Product Size: 151

In-silico PCR Accepted

Oligo 5 :

Wildtype Reverse Primer 5': CTGGGCTATAACACAGAAG

Mutant Reverse Primer 5': CTGGGCTATAACACAGAAA

Common Forward Primer 5': GCTGTAACAGTTTATTGGCA

Product Size: 150

In-silico PCR Accepted.

6. rs2396755

AS Primer Picking Result for rs239675

Sequence Size: 601

Primer Picking Parameters:

Primer Size Opt: 20 Min: 18 Max: 36
 GC% Opt: 50.0 Min: 20.0 Max: 85.0
 Tm Opt: 55.0 Min: 40.0 Max: 65.0

Max Tm Diff 100.0

Max Self Complementarity: 8.0

Max 3' Complementarity: 3.0

Max PolyX in Primer: 3
 Number Primer Return: 5
 Mismatch at the penultimate primer position: 'Yes'
 In-silico PCR Filtering: 'Yes' with Max Product Size: 4000
 Mispriming Filtering: 'No'

Oligo 1 :

Wildtype Reverse Primer 5': GGGCTGGAGGGCAAGATG
 Mutant Reverse Primer 5': GGGCTGGAGGGCAAGATA
 Common Forward Primer 5': AGGAGCTCCCACAATGCT
 Product Size: 216

Discussion:

Genotyping is the process of determining differences in the genetic make-up (*genotype*) of an individual by examining the individual's DNA sequence using biological assays and comparing it to another individual's sequence or a reference sequence. It reveals the alleles an individual has inherited from their parents. Genotyping is the process of determining which genetic variants an individual possesses. Genotyping can be performed through a variety of different methods, depending on the variants of interest and resources available. Single nucleotide polymorphisms (SNPs) are the most common source of human genetic variation. The potential use of SNPs for genetic mapping of complex traits, pharmacogenetics, and medical diagnostics has been much discussed. These are single base differences between DNA of different individuals. These are more suitable for genotyping markers compared to the conventional markers such as RFLP (Restriction fragment length polymorphism), AFLP (Amplified fragment length polymorphism) and SSR (Simple Sequence Repeats). SNPs are becoming genetic markers that are used in detection of risk-associated alleles linked to human diseases. Recently, massive parallel sequencing platforms such as GSFLX (Roche), Solexa (Illumina) and SOLID (Applied Biosystems) have significantly reduced the cost of high throughput sequencing. A large variety of techniques for high-throughput SNP genotyping have also been developed using Taqman, Amplifluor, genome re-sequencing, and SNP arrays. These techniques are expensive and time consuming and require specialized equipments. The original SNP genotyping methods—DNA sequencing and PCR-RFLP—are laborious and expensive because they require multiple steps including size separation. AS-PCR (Allele-specific PCR) is widely used for low-throughput applications in research. Our aim was to design allele specific primers for *ACHE* gene with WASP software. A total of 10 SNPs were selected for allele specific primers and were designed by the WASP software. Very less scientific literature is available regarding the use of allele specific primer designing of *ACHE* gene for genotyping purpose. Wangkumhang *et al* in 2007 designed five allele specific primers for *C4P21D6* gene and found WASP a good tool for designing AS Primers for both SNP and Mutations. No other study is available in the literature for the comparison with our study. In conclusion, WASP proved to be a good tool for designing AS Primers for both SNP and Mutations

9.Conclusion:

The allele specific primers were designed with WASP (a web based allele specific primer) design application for a total of 10 SNPs of *ACHE* gene. The tool has potential in conveniently assisting scientists in getting SNP information from local SNP database.

10. References:

- [1] Agrawal D, Sultana P, Gupta G. Oxidative damage and changes in the glutathione redox system in erythrocytes from rats treated with hexachlorocyclohexane. *Food Chem. Toxicol.* 1991;29:459–462.
- [2] Ahmed RS, Pasha ST VS, Banerjee BD. Influence of dietary ginger (*Zingiber Officinale* Rose) on oxidative stress induced by Malathion in rats. *Food and Chemical Toxicology.* 2000;38:443–450.
- [3] Almeida M, Fanini F, Davino SC, Aznar AE, Koch OR, Barros SBM. Pro and anti-oxidant parameters in rat liver after short-term exposure to hexachlorobenzene. *Hum. Exp. Toxicol.* 1997;16:257–261.
- [4] Banerjee BD, Seth V, Ahmed RS. Pesticide-induced oxidative stress: perspectives and trends. *Rev Environ Health.* 2001;16:1–40.
- [5] Barbazuk WB, Emrich SJ, Chen HD, Li L, Schnable PS: SNP discovery via 454 transcriptome sequencing. *Plant J* 2007, 51:910–918.
- [6] Bolognesi C. Genotoxicity of pesticides: a review of human biomonitoring studies *Mutation Research* 543 2003;251–272.
- [7] Butterfield DA: Amyloid beta-peptide (1–42)-induced oxidative stress and neurotoxicity: implications for neurodegeneration in Alzheimer's disease brain. A review. *Free Radic Res*, 2002;36:1307–13.
- [8] Cha RS, Zarbl H, Keohavong P, Thilly WG: Mismatch amplification mutation assay (MAMA): application to the c-H-ras gene. *PCR Methods Appl* 1992;2:14–20.
- [9] Choi YJ, Seelbach MJ, Pu H. Polychlorinated biphenyls disrupt intestinal integrity via NADPH oxidase-induced alterations of tight junction protein expression. *Environ Health Perspect.* 2010;118:976–981.
- [10] Cok, I.; Bilgili, A.; Ozdemir, M.; Ozbek, H.; Bilgili, N. & Burgaz, S. Organochlorine pesticide residues in human breast milk from agricultural regions of Turkey, 1995–1996. *Bulletin of Environmental Contamination and Toxicology.* 1997. 59., 577–582.
- [11] Costa C, Silva S, Coelho P, Torres J, Teixeira J, Mayan O Micronucleus analysis in a Portuguese population exposed to pesticides: Preliminary survey. *Int. J. Hyg. Environ.-Health* 210 (2007) 415–418.
- [12] Das P, John G. Induction of sister chromatid exchanges and chromosome aberrations in vivo in *Etioplosuratus* (Bloch) following exposure to organophosphorus pesticides. *Toxicol Lett.* 1999;104:111–116.
- [13] Dong, L.M.; Potter, J.D.; White, E.; Ulrich, C.M.; Cardon, L.R. & Peters, U. Genetic susceptibility to cancer: the role of polymorphisms in candidate genes. *Journal of the American Medical Association.* 2008. 299., 2423–2436.
- [14] Eberle MA, Ng PC, Kuhn K, Zhou L, Peiffer DA, Galver L, Viaud-Martinez KA, Lawley CT, Gunderson KL, Shen R, Murray SS: Power to detect risk alleles using genome-wide tag SNP panels. *PLoS Genet* 2007;3:1827–1837.
- [15] Feltus FA, Wan J, Schulze SR, Estill JC, Jiang N, Paterson AH: An SNP resource for rice genetics and breeding based on subspecies indica and japonica genome alignments. *Genome Res* 2004, 14:1812–1819.
- [16] Ferro R, Parvathaneni A, Patel S, Cheriya P, Pesticides and Breast Cancer. *Advances in Breast Cancer Research.* 2012;1:30–35.
- [17] Flint-Garcia SA, Thornsberry JM, Buckler ES: Structure of linkage disequilibrium in plants. *Annu Rev Plant Biol* 2003, 54:357–37
- [18] Gilden RC, Huffling K, Sattler B. Pesticides and health risks. *J Obstet Gynecol Neonatal Nurs.* 2010;39(1):103–10.

- [19] Green RM, Hodges NJ, Chipman JK, O'Donovan MR, Graham M. Reactive oxygen species from the uncoupling of human cytochrome P450 1B1 may contribute to the carcinogenicity of dioxin-like polychlorinated biphenyls. *Mutagenesis*. 2008; 23:457-463
- [20] Gut IG: Automation in genotyping of single nucleotide polymorphisms *Hum Mutat* 2001, 17:475-492.
- [21] Halliwell, B. Oxidative stress and cancer: have we moved forward? *Biochemical Journal*. 2007. 401., 1-11.
- [22] Hayes WJ, Laws ER. Dosage and other factors influencing toxicity. *Handbook of pesticides toxicology*, Academic Press, San Diego. 1991;1:39-105.
- [23] Hernández A, Lacasa M, Gil F, Barranco M, Antonio P, Guarnido O. Evaluation of pesticide-induced oxidative stress from a gene-environment interaction perspective. *Toxicology* 307 ;2005; 95- 102.
- [24] Hillier LW, Miller RD, Baird SE, Chinwalla A, Fulton LA, Koboldt DC, Waterston H: Comparison of *C. elegans* and *C. briggsae* genome sequences reveals extensive conservation of chromosome organization and synteny. *PLoS Biol* 2007, 5:e167.
- [25] Huang J, Wei W, Zhang J, Liu G, Bignell GR, Stratton MR, Futreal PA, Wooster R, Jones KW, Shaperro MH: Whole genome DNA copy number changes identified by high density oligonucleotide arrays. *Hum Genomics* 2004, 1:287-299.
- [26] Khrrer JP. Free radicals as mediator of tissue injury and disease. *Crit. Rev. Toxicol.* 1993;23:21-48.
- [27] Kolaczinski JH, Curtis CF. Chronic illness as a result of low level exposure to synthetic pyrethroid insecticides. *Food Chem Toxicol.* 2004;42:697-706.
- [28] Kwok PY: Methods for genotyping single nucleotide polymorphisms. *Annu Rev Genomics Hum Genet* 2001, 2:235-258.
- [29] Kwok S, Chang SY, Sninsky JJ, Wang A: A guide to the design and use of mismatched and degenerate primers. *PCR Methods Appl* 1994, 3:S39-S47
- [30] Li R, Li Y, Fang X, Yang H, Wang J, Kristiansen K, Wang J: SNP detection for massively parallel whole-genome resequencing. *Genome Res* 2009,19:1124-1132.
- [31] Li, Q.; Kobayashi, M. & Kawada, T. (2007). Organophosphorus pesticides induce apoptosis in human NK cells, *Toxicology*, 239., 89-95.
- [32] Lioi MB, Scarfi MR, Santoro A, Barbieri R, Zeni O, Berardino DD, Ursini MV. Genotoxicity and oxidative stress induced by pesticide exposure in bovine lymphocyte cultures in vitro. *Mutat Res.* 1988;403:13-20.
- [33] Livak KJ, Marmaro J, Todd JA: Towards fully automated genome-wide polymorphism screening. *Nat Genet* 1995, 9:341-342.
- [34] Malarvannan, G.; Kunisue, T.; Isobe, T.; Sudaryanto, A.; Takahashi, S.; Prudente, M.; Subramanian, A. & Tanabe, S. Organohalogen compounds in human breast milk from mothers living in Payatas and Malate, the Philippines: levels, accumulation kinetics and infant health risk. *Environmental Pollution*. 2009. 157., 1924-1932.
- [35] Matsuzaki H, Dong S, Loi H, Di X, Liu G, Hubbell E, Law J, Berntsen T, Chadha M, Hui H, Yang G, Kennedy GC, Webster TA, Cawley S, Walsh PS, Jones KW, Fodor SP, Mei R: Genotyping over 100,000 SNPs on a pair of oligonucleotide arrays. *Nat Methods* 2004, 1:109-111.
- [36] Mena, S.; Ortega, A. & Estrela, J.M. (2009). Oxidative stress in environmental-induced carcinogenesis. *Mutation Research*, 674., 36-44.
- [37] Mino CP, Bustamante G, Sanchez ME, Leone PE. Cytogenetic monitoring in a population occupationally exposed to pesticides in Ecuador. *Environ Health Perspect.* 2002;110:1077-1085.
- [38] Myakishev MV, Khripin Y, Hu S, Hamer DH: High-throughput SNP genotyping by allele-specific PCR with universal energy-transfer-labeled primers. *Genome Res* 2001, 11:163-169.
- [39] Newton CR, Graham A, Heptinstall LE, Powell SJ, Summers C, Kalsheker N, Smith JC, Markham AF: Analysis of any point mutation in DNA. The amplification refractory mutation system (ARMS). *Nucleic Acids Res* 1989, 17 (7): 2503-16. 10.1093/nar/17.7.2503.
- [40] Newton CR, Graham A, Heptinstall LE, Powell SJ, Summers C, Kalsheker N, Smith JC, Markham AF: Analysis of any point mutation in DNA. The amplification refractory mutation system (ARMS). *Nucleic Acids Res* 1989, 17(7):2503-16.
- [41] Perera, F.P, Rauh, V, Whyatt, R.M, Tang, D Tsai, W.Y. Bernert, J.T. Tu, Y.H.; Andrews, H.Barr, D.B.; Camann, D.E.; Diaz, D.; Dietrich, J.; Reyes, A. & Kinney, P.L. A summary of recent findings on birth outcomes and developmental effects of prenatal ETS, PAH, and pesticide exposures. *Neurotoxicology*, 2005. 26., 573-587.
- [42] Pongsakorn Wangkumhang, Kridsakorn Chaichoompu, Chumpol Ngamphiw, Uttapong Ruangrit, Juntima Chanprasert, Anuchai Assawamakin and Sissade Tongsimma *Genomics* 2007;275, 10.1186/1471-2164-8-27, 2007.
- [43] Shaperro MH, Zhang J, Loraine A, Liu W, Di X, Liu G, Jones KW: MARA: a novel approach for highly multiplexed locus-specific SNP genotyping using high-density DNA oligonucleotide arrays. *Nucleic Acids Res* 2004,32:e181.
- [44] Shen R, Fan JB, Campbell D, Chang W, Chen J, Doucet D, Yeakley J, Bibikova M, Wickham Garcia E, McBride C, Steemers F, Garcia F, Kermani BG, Gunderson K, Oliphant A: High-throughput SNP genotyping on universal bead arrays. *Mutat Res* 2005, 573:70-82.
- [45] Shendure J, Mitra RD, Varma C, Church GM: Advanced sequencing technologies: methods and goals. *Nat Rev Genet* 2004, 5:335-344.
- [46] Sherer TB, Richardson JR, Testa CM. Mechanism of toxicity of pesticides acting at complex I: relevance to environmental etiologies of Parkinson's disease. *J Neurochem.* 2007;100:1469-1479.
- [47] Stevenson, D.E.; Walborg Jr., E.F.; North, D.W.; Sielken Jr., R.L.; Ross, C.E.; Wright, A.S.; Xu, Y. Kamendulis, L.M. & Klaunig, J.E. Monograph: reassessment of human cancer risk of aldrin/dieldrin. *Toxicology Letters*. 1999. 109., 123-186.
- [48] Trick M, Long Y, Meng J, Bancroft I: Single nucleotide polymorphism (SNP) discovery in the polyploid *Brassica napus* using Solexa transcriptome sequencing. *Plant Biotechnol J* 2009, 7:334-346.
- [49] Wang C, Liu Z: Arabidopsis ribonucleotide reductases are critical for cell cycle progression, DNA damage repair, and plant development. *Plant Cell* 2006, 18:350-365.
- [50] Wang, Y. Bulky DNA lesions induced by reactive oxygen species. *Chemical Research in Toxicology*. 2008. 21., 276-281.
- [51] Yadav AS, Sehrawat G. Evaluation of Genetic Damage in Farmers Exposed to Pesticide Mixtures. *Int J Hum Genet.* 2011;11(2):105-109.
- [52] Ye S, Dhillon S, Ke X, Collins AR, Day IN: An efficient procedure for genotyping single nucleotide polymorphisms. *Nucleic Acids Res.* 2001, 29 (17): E88-8. 10.1093/nar/29.17.e88.
- [53] Ye S, Dhillon S, Ke X, Collins AR, Day IN: An efficient procedure for genotyping single nucleotide polymorphisms. *Nucleic Acids Res* 2001, 29(17):E88-8.