

potato research

Journal of the European Association for Potato Research
Zeitschrift der Europäischen Gesellschaft für Kartoffelforschung
Revue de l' Association Européenne pour la Recherche
sur la Pomme de Terre
Volume 14 Number 1 March 1971 Pages 1 - 109

Continuation of »European Potato Journal«

Potato research

Journal of the European Association for Potato Research
Zeitschrift der Europäischen Gesellschaft für Kartoffelforschung
Revue de l' Association Européenne pour la Recherche
sur la Pomme de Terre
Volume 14 Number 1 March 1971 Pages 1 - 109

 Continuation of »European Potato Journal«

Jong, H. de and Rowe, P. R. (1971): Inbreeding in cultivated diploid potatoes. *Potato Res.* 14: 74–83.

The effects of inbreeding in cultivated diploid potatoes were studied by using self-fertile hybrids from crosses between Group Phureja and Group Stenotomum and haploids of Group Tuberosum. S₁, S₂ and S₃ progenies usually exhibited decreased performance and vigour with increased inbreeding. In most families the inbreeding depression seemed to coincide with a curve calculated on the basis of the expected loss of heterozygosity in a diploid organism. The performance of some progenies deviated markedly from expectations apparently because a non-random sample of parents contributed to later generations. Many inbred genotypes were lost owing to poor seedling vigour or tuberization. This indicated the presence of many harmful recessive genes. The rapid increase in homozygosity following selfing at the diploid level provides a means for developing homozygous clones with a potential for use in future genetic investigations.

Gately, T. F. (1971): Effects of nitrogen on potato yields and on the total-N and nitrate-N content of the tops. *Potato Res.* 14: 84–90.

The effects of 4 nitrogen levels on tuber yield, and on the total-N and nitrate-N content of plant samples taken during the early growing season were examined using the variety *Kerr's Pink* grown on nineteen sites.

The mean yield of the zero N treatment was 38.7 tons/ha. Application of 45, 90 and 135 kg/ha increased yields by 4.5, 6.5 and 8.8 tons/ha, respectively.

The total-N and nitrate-N content of the plant samples increased as the amount of N fertiliser was increased. They were highest at sites where the response to N fertiliser was least.

There were no significant relationships between yields and plant analyses, but correlations between percent responses and plant analyses were significant.

Howard, H. W. (1971): The stability of L1-mutant periclinal potato chimeras. *Potato Res.* 14: 91–93.

The varieties *Red Craigs Royal* and *Red King* are periclinal chimeras with the constitution, L1 uniform red – L2 white-splashed-pink – L3 white-splashed-pink. About 1 in a 100 plants has a tuber or part of a tuber showing reversion to L1 white-splashed-pink.

Verma, S. C., Malhotra, V. P., Joshi, K. C. and Sharma, T. R. (1971): Specific gravity and dry matter content of potato. *Potato Res.* 14: 94–95.

Data obtained from 516 pairs of specific gravity and dry matter determinations were used to derive a relationship for estimation of dry matter content of potatoes grown in India.

Burton, W. G. and Meigh, D. F. (1971): The production of growth-suppressing volatile substances by stored potato tubers. *Potato Res.* 14: 96–101.

The sprout suppressing constituent(s) of the volatiles may be removed by a solution of silver sulphate in 96% H₂SO₄ or by aqueous solutions of mercuric acetate or chloride, but not by 96% H₂SO₄ alone, suggesting that they may be olefinic, ethereal and/or sulphur-containing. Ethylene may be produced in very small quantities, insufficient to be an active olefinic constituent.

(Continuation of page IV)

Potato Research

Journal of the European Association for Potato Research

Zeitschrift der Europäischen Gesellschaft für Kartoffelforschung

Revue de l'Association Européenne pour la Recherche sur la Pomme de terre

Editorial Board – Schriftleitung – Comité de rédaction:

The Council of the Association – Der Vorstand der Gesellschaft – Le Conseil de l'Association

Editors – Schriftleiter – Rédacteurs:

Dr. W. G. Burton, A.R.C., Food Research Institute, Colney Lane, Norwich, NOR 70F, England

Dr. K. Müller, Institut für Agrikulturchemie der Universität Göttingen, Nikolausbergerweg 7
34 Göttingen, Deutschland Bundesrepublik

Dr. Ir. D. E. van der Zaag, Postbus 20, Wageningen, The Netherlands

Assistant Editor – Assistent-Schriftleiter – Rédacteur-adjoint

Ir. H. P. Beukema, Wageningen, The Netherlands

Preparation for press – Fertigstellung für den Druck – Présentation:

Centre for Agricultural Publishing and Documentation, P.O. Box 4, Wageningen,
The Netherlands



Composition of the Journal Original contributions on fundamental and applied research on potatoes, subject surveys, short, provisional or preliminary communications, reports of section meetings (including abstracts of papers), reviews and Association news.

A volume of the Journal consists of four issues published in the same year. Papers are in English, German or French with a short summary in the language used and extended summaries in the other two.

Zusammenstellung der Zeitschrift Originalbeiträge über grundlegende und angewandte Forschung an der Kartoffel, Fachberichte, kurze, vorläufige oder einleitende Mitteilungen, Berichte über Sektionstagungen (einschliesslich Kurzfassungen der Vorträge), Buch- oder andere Besprechungen und Berichte der Gesellschaft.

Ein Jahrgang der Zeitschrift besteht aus vier Hefen. Die Beiträge erscheinen in English, Deutsch oder Französisch mit einer kurzen Zusammenfassung in der Originalsprache und ausführlicheren Zusammenfassungen in den beiden andern Sprachen.

Composition du Journal Contributions originales sur des recherches fondamentales et appliquées à la Pomme de terre, aperçus sur divers sujets, courtes communications provisoires ou

préliminaires, rapports de réunions de section (comprenant des résumés de communications,) revues et nouvelles de l'Association.

Un volume du journal comprend les quatre numéros publiés la même année. Les articles sont rédigés en anglais, en allemand ou en français et sont accompagnés d'un bref résumé dans la langue de la publication et d'un résumé étendu dans les deux autres langues.

Subscription to non-members 25 Dutch florins

Bezugspreis für Nichtmitglieder 25 holl. Gulden

Abonnement pour non-membres 25 florins hollandais

Back-numbers 6,25 Dutch florins each

Frühere Hefte je 6,25 holl. Gulden

Anciens numéros 6,25 florins hollandais l'exemplaire

All correspondence should normally be addressed to Postbus 20, Wageningen, The Netherlands.

Alle Korrespondenz ist in der Regel zu richten an Postbus 20, Wageningen, Niederlande.

Toute la correspondance doit être normalement adressée à Postbus 20, Wageningen, Pays-Bas.

Contents

E. Winkler Kartoffelbau in Tirol II. Photosynthesevermögen und Respiration von verschiedenen Kartoffelsorten	1
B. Mica Einfluss der Düngung auf den Gehalt an essentiellen Aminosäuren im Kartoffel- eiweiss	19
F. A. Langton The development of a laboratory test for assessing potato varietal susceptibility to gangrene caused by <i>Phoma exigua</i> var. <i>foveata</i>	29
Bärbel Schöber Physiologische Veränderungen in der Kartoffelknolle nach Verletzung und Infek- tion mit <i>Phytophthora infestans</i>	39
J. K. A. Bleasdale and R. Thompson The suitability for canning of some potato varieties	49
R. T. V. Fox, J. G. Manners and A. Myers Ultrastructure of entry and spread of <i>Erwinia carotovora</i> var. <i>atroseptica</i> into pota- to tubers	61
H. de Jong and P. R. Rowe Inbreeding in cultivated diploid potatoes	74
T. F. Gately Effects of nitrogen on potato yields and on the total-N and nitrate-N content of the tops	84
Short communications	
H. W. Howard The stability of L1-mutant periclinal potato chimeras	91
S. C. Verma, V. P. Malhotra, K. C. Joshi and T. R. Sharma Specific gravity and dry matter content of potato	94
W. G. Burton and D. F. Meigh The production of growth-suppressing volatile substances by stored potato tubers	96
Reviews and publications received	97
Association news	103

Inbreeding in cultivated diploid potatoes*

H. DE JONG¹ and P. R. ROWE²

¹ Department of Horticulture, University of Wisconsin, Madison (Wisconsin), USA

² Crops Research Division, University of Wisconsin, Madison (Wisconsin), USA

Accepted for publication: 10 September 1970

Zusammenfassung, Résumé p. 81

Summary

The effects of inbreeding in cultivated diploid potatoes were studied by using self-fertile hybrids from crosses between Group Phureja and Group Stenotomum and haploids of Group Tuberosum. S₁, S₂ and S₃ progenies usually exhibited decreased performance and vigour with increased inbreeding. In most families the inbreeding depression seemed to coincide with a curve calculated on the basis of the expected loss of heterozygosity in a diploid organism. The performance of some progenies deviated markedly from expectations apparently because a non-random sample of parents contributed to later generations. Many inbred genotypes were lost owing to poor seedling vigour or tuberization. This indicated the presence of many harmful recessive genes. The rapid increase in homozygosity following selfing at the diploid level provides a means for developing homozygous clones with a potential for use in future genetic investigations.

Introduction

The success of hybridization between inbred lines in maize has led to many similar investigations with other crops. Several workers have reported inbreeding studies in cultivated tetraploid potatoes (Engel, 1957; Feistritzer, 1952; Gowen, 1956; Krantz, 1924, 1946; Krantz and Hutchins, 1929; Mullin and Lauer, 1966; Rudorf, 1958). Such studies have not been reported in cultivated diploid potatoes, probably because most of the diploid *Solanum* species are self-incompatible (Cipar et al., 1964). Research in this area is now possible as a result of studies of haploids of the cultivated tetraploid potato (*S. tuberosum* L. Group Tuberosum). Crosses between cultivated diploid species and male-fertile Tuberosum haploids produce vigorous, highly fertile progeny. Most of these 24-chromosome hybrids are self-incompatible, but crosses involving haploids from an inbred Minnesota seedling, *Minn. 20-20-34*, (primarily *US-W4*) were found to be self-fertile (Smiley, 1963; Cipar, 1964).

This paper describes the effect of inbreeding on populations derived from diploid-haploid hybrids. An inbreeding study with a genetic base of diploid hybrids, has several advantages: 1) since the tetraploid varieties and self-incompatible diploid species may have several recessive genes not normally expressed, these hybrids should

* Cooperative investigations of the Crops Research Division, Agricultural Research Service, U.S. Department of Agriculture, and the Wisconsin Agricultural Experiment Station.

be an extremely fertile source of unexplored genetic variability (Hougas and Peloquin, 1962); 2) the rate of progress towards homozygosity following selfing is much faster with diploids than with autotetraploids. The number of generations needed to reduce heterozygosity by 50 per cent by inbreeding is 1 in diploids and 3.8 in autotetraploids (Bartlett and Haldane, 1934); 3) stocks homozygous for marker genes of potential value for future genetic investigations can be obtained much faster; 4) the segregation ratios obtained for progenies from selfed seed would be much easier to analyze with a disomic inheritance pattern. Progeny analysis based on selfing can be a powerful tool for evaluating potential parents for a crossing programme (Feistritzer, 1952; Krantz and Hutchins, 1929).

Materials and methods

The experiments were conducted at the Potato Introduction Station, Sturgeon Bay, Wisconsin. The initial genetic base consisted of the following hybrid families:

US-W 5309 (Phu PI 195191 × US-W4)

US-W 5314 (Phu PI 225708.1 × US-W4)

US-W 5315 (Phu PI 225675.1 × US-W4)

US-W 5328 (Stn PI 230513.1 × US-W4)

In each generation, plants were self-pollinated and the seed produced by self-fertile plants bulked to produce subsequent generations. Observations were made on populations grown from true seed and from tubers. Families grown from true seed were started in the greenhouse and transplanted to the field. Two replicates of seedling populations were obtained by dividing a bulked population of seed into 2 equal lots. All plants producing tubers were saved for evaluation in tuber plantings. Replicated trials were planted in the field in 2 years.


The following characteristics were measured:

1. Plant height – in cm when the plants had reached maximum height.
2. Vigour index – based on total amount of foliage. Individual plants were rated on a scale from 1 (very weak or small) to 5 (very vigorous).
3. Number of tubers per hill.
4. Tuber weight – g per hill.
5. Self-, pollen-, and ovule-fertility. The fertility data were obtained by using the decapitation technique (Peloquin and Hougas, 1959). Tests were conducted on all replicates planted from tubers in 1968 (Table 3). For each inbred clone, 3 stems with 4 buds each were collected at random from a large population of Phureja plants, and these 12 flowers were then used as females to test the pollen fertility of the inbred population. Up to 2 stems, also with 4 buds each, were cut from each flowering clone of the inbred populations. On each stem three flowers were used to test the self-fertility, and one flower was emasculated and pollinated with a mixture of pollen from the Phureja population to measure the ovule fertility. A total of 3651 pollinations were made; of these 935 were made to measure self-fertility, 2375 to measure pollen fertility and 341 to measure ovule fertility.

Results

Inbreeding depression was expressed by all measured characters. When averaged over the 4 families, the inbreeding depression generally followed the rate expected based on reduction of heterozygosity. In the case of tuber weight per hill for the populations grown from tubers in 1968, it was possible to calculate a curve which agreed very well with the observed data (Fig. 1). Considerable variation in response to inbreeding existed between families (Table 1). Family *US-W 5328* exhibited inbreeding depression at each generation of inbreeding. Family *US-W 5309* had a sharp depression from the

Fig. 1. Average tuber weight in g per hill (1968).



g per hill – g pro Staude – gr/touffe
 generation – Generation – génération
 observed – festgestellt – observé
 expected – erwartet – attendu

Abb. 1. Mittleres Knollengewicht in g/pro Staude (1968).
 Fig. 1. Poids moyens de tubercules en gr/touffe (1968).

INBREEDING IN CULTIVATED DIPLOID POTATOES

Table 1. Effect of inbreeding on four diploid potato families.

Family ¹	Number of clones ²			Plant height ³ (cm)			Vigour index ⁴			Tubers/hill ⁵			Tuber weight g/hill ⁶		
	S ₀	S ₁	S ₂	S ₀	S ₁	S ₂	S ₀	S ₁	S ₂	S ₀	S ₁	S ₂	S ₀	S ₁	S ₂
First clonal generation ⁷															
US-W 5309	12	16	16	43	25	27	3.5	2.3	2.5	21	16	15	563	248	269
US-W 5328	9	12	8	41	27	17	3.6	2.4	1.6	19	15	8	390	425	167
US-W 5314	0	12	11		27	23		2.4	2.2		13	7		287	88
US-W 5315	0	10	9		25	29		2.6	2.7		20	21		297	415
Total	21	50	44												
Weighted average ⁸				42	26	25	3.5	2.4	2.2	20	16	14	499	304	181
% of S ₀ ⁹				100	62	60	100	69	63	100	80	70	100	61	36
Second or later clonal generations ¹⁰															
US-W 5309	12	10	11	31	23	31	3.6	2.8	3.7	10	14	12	198	165	217
US-W 5328	7	11	4	37	28	14	4.0	3.3	2.0	20	13	5	370	330	13
US-W 5314	0	10	4		22	22		2.7	2.6		13	13		100	95
US-W 5315	0	9	7		20	25		2.5	3.5		13	14		202	237
Total	19	40	26												
Weighted average ⁸				33	24	25	3.7	2.9	3.2	14	13	12	260	211	175
% of S ₀ ⁹				100	73	76	100	78	86	100	93	86	100	81	67

¹ Familie – Familles

² Anzahl Klone – Nombre de clones,

³ Pflanzenhöhe – hauteur des plantes

⁴ Frohwüchsigkeit (Index) – Index de vigueur

⁵ Knollen/Staude – Tubercules/touffe

⁶ Knollengewicht g/Staude – Poids (gr) de tub. par touffe

⁷ Erste Klonegeneration – première génération clonale

⁸ Gewogenes Mittel – Poids moyen

⁹ in % von S₀ – % de S₀

¹⁰ Zweite oder spätere Klonegenerationen – Seconde ou suivantes générations clonales

Tabelle 1. Einfluss der Inzucht auf vier diploide Kartoffelfamilien.

Tableau 1. Effet de l'autofécondation sur quatre familles de Pomme de terre diploïdes.

S₀ to the S₁, but the subsequent S₂ generation was not depressed. The performance of families US-W 5314 and US-W 5315, respectively, was slightly lower and slightly higher in the S₂ than in the S₁ (Table 1).

The inbreeding depression of populations obtained by clonal reproduction was nearly always less than in the asexual parent population (Tables 1 and 2). In all cases the asexual progenies contained fewer genotypes than their parent populations, because of lack of tuberization in the parent populations. Clones planted from tubers exhibited reduced tuberization that increased with each subsequent level of inbreeding (Table 3).

There was a severe inbreeding depression from S₀ to S₁ for all characters for all

Table 2. Effect of inbreeding on potato populations grown from true seed and the first clonal generation

Character ¹	Seedling populations (1968) ²				First clonal generation (1969) ³			
	S ₀	S ₁	S ₂	S ₃	S ₀	S ₁	S ₂	S ₃
Number of plants ⁴	29	163	184	163	11	41	48	35
Plant height (cm) ⁵	25	16	16	15	46	21	20	20
% of S ₀ ⁶	100	64	64	59	100	46	43	42
Vigour index ⁷	5	2	3	3	5	3	3	3
% of S ₀ ⁶	100	51	57	57	100	60	60	60
Tubers/hill ⁸	10	6	5	5	23	16	20	16
% of S ₀ ⁶	100	60	50	50	100	70	87	70
Tuber weight (g/hill) ⁹	61	20	16	14	426	236	234	193
% of S ₀ ⁶	100	33	26	23	100	55	55	45
% tuberizing ¹⁰	69	81	80	71	97	93	94	83
% with ≥ 1 viable tubers after storage ¹¹	62	70	72	63				
% with ≥ 4 viable tubers after storage ¹²	38	33	32	25				

¹ Eigenschaft – Caractère² Sämlingspopulationen (1968) – Populations de plantules (1968)³ Erste Klongeneration (1969) – Première génération clonale (1969)⁴ Anzahl pflanzen – Nombre de plantes⁵ Pflanzenhöhe (cm) – Hauteur des plantes (cm)⁶ in % von S₀ – % de S₀⁷ Frohwüchsigkeit, Index – Index de vigueur⁸ Knollen/Staude – Tub./tousse⁹ Knollengewicht (g/Staude) – Poids des tub. (gr/touffe)¹⁰ % mit Knollenansatz – % de tubérisation¹¹ % mit ≥ 1 lebensfähiger Knolle nach der Lagerung – % avec ≥ 1 tubercule viable après conservation¹² % mit ≥ 4 lebensfähigen Knollen nach der Lagerung – % avec ≥ 4 tubercules viables après conservation

Tabelle 2. Einfluss der Inzucht auf Kartoffelpopulationen, entstanden aus Samen und der ersten Klongeneration.

Tableau 2. Effet de l'autofécondation sur des populations de pomme de terre issues de vraies semences et sur la première génération clonale.

families when grown from true seed. There was little if any depression in the subsequent S₂ and S₃ generations. These populations tuberized fairly well in the field even in the advanced generations of inbreeding. Most of these tubers were very small, however, and were not viable the following spring after standard winter storage conditions (Table 2).

The per cent of plants that flowered, the per cent of fertile flowering plants, and the level of fertility decreased very sharply from the S₀ to the S₁ but not from S₁ to S₂ (Table 4). The self-fertility was much lower than either the pollen or ovule fertility at all levels of inbreeding.

INBREEDING IN CULTIVATED DIPLOID POTATOES

Table 3. Effect of inbreeding on asexual reproduction.

Year ¹	Number of clones planted ²		
	S ₀	S ₁	S ₂
1968	21	50	44
1969	19	40	26
Per cent replanted in 1969 ³	90	80	59

¹ Jahr – Année

² Anzahl ausgepflanzter Klone – Nombre de clones plantés

³ % 1969 nachgebaute Klone – Pourcents replantés en 1969

Tabelle 3. Einfluss der Inzucht auf die ungeschlechtliche Vermehrung.

Tableau 3. Effet de l'autofécondation sur la reproduction asexuée.

Table 4. Effect of inbreeding on flowering and fertility.

Level of inbreeding ¹	Clones planted ²	Per cent stand ³	Per cent flowering ⁴	Fertility in per cent of flowering clones ⁵			Seeds/fruit for fertile plants ⁶		
				Self- ⁷	Pollen- ⁸	Ovule- ⁹	Self- ⁷	Pollen- ⁸	Ovule- ⁹
S ₀	21	95	100	47	85	86	12	54	76
S ₁	50	91	54	16	34	23	9	36	34
S ₂	44	74	54	16	41	47	6	38	30

¹ Inzuchtgrad – Degré de consanguinité

² Anzahl gepflanzter Klone – Clones plantés

³ Prozent Pflanzenentwicklung – Situation en %

⁴ Prozent Blühzustand – % de floraison

⁵ Fertilität in % der blühenden Klone – Fertilité en % des clones fleurissant

⁶ Samen/Frucht für fruchtbare Pflanzen – Semences par fruit chez les plantes fertiles

⁷ Selbst – Auto

⁸ Pollen- – Pollen-

⁹ Eizellen – Ovule

Tabelle 4. Einfluss der Inzucht auf den Blühzustand und die Fruchtbarkeit.

Tableau 4. Effet de l'autofécondation sur la floraison et la fertilité.

Discussion

Generally the average performance of the 4 families decreased with each subsequent level of inbreeding. For some characters, the inbreeding depression coincided with a curve calculated on the basis of heterozygosity in a diploid organism (Fig. 1 and Table 1). Such a response is expected of a crop that is normally highly heterozygous, cross-pollinated and vegetatively propagated. Upon inbreeding tetraploids, Krantz (1946) obtained an S₂ that yielded 82% of the S₀, whereas in this experiment the first clonal generation S₂ yielded only 36% of the S₀ (Table 1). The difference can be attributed to the more rapid increase in homozygosity at the diploid level. The 4 families responded

differently to inbreeding. The fluctuations in response were partly due to reproductive difficulties (both sexual and asexual) which increase with inbreeding. The genetic differences between the original parent populations (S_0) could also have contributed to differences in response to inbreeding. This was observed at the tetraploid level by Krantz and Hutchins (1929).

Attempts to obtain each subsequent level of inbreeding from a representative sample of parents at the preceding level were not always successful. Only the more vigorous plants flowered, and only a portion of these were self-fertile in the advanced generations of inbreeding (Table 4). A selection for vigour in the sexual cycle has also been observed upon inbreeding at the tetraploid level (Krantz, 1946; Krantz and Hutchins, 1929). The self-fertility was considerably lower than either pollen or ovule fertility for all levels of inbreeding when expressed either as per cent of flowering clones or as seeds per fruit. A reduction in seed set for self-, pollen- and ovule-fertility could be demonstrated with each subsequent level of inbreeding (Table 4). A considerably higher seed set upon selfing the S_0 of similar material has been reported (Hollenback, 1966; Smiley, 1963). In our experiment, however, special efforts were made to obtain selfed seed from the weaker clones as well as from the more vigorous ones. This procedure was necessary to obtain a sample of selfed seed as nearly random as possible, but probably contributed to a lower self-fertility level in the S_0 than reported by others. Rudolf (1958) reported similar reproductive difficulties upon inbreeding tetraploids. He could not obtain the S_3 in most of his inbred lines due to their lack of vigour, and to virus infection and reduced fertility.

Approximately 45% of the genotypes were lost between the time of planting the seed to the transplanting stage. Similar losses were reported by Engel (1957). For several genotypes we were unable, due to lack of tuberization, to obtain the second clonal generation. When the performance of the first clonal generations was compared with the asexual progenies, the depression in relation to the respective S_0 's was smaller for most progenies than for their respective asexual parental populations. (Tables 1 and 2). This can probably be attributed in part to the difficulties with asexual reproduction. Many plants that could be classified as 'tuberizing' in the seedling year, nevertheless only produced small immature tubers which were not viable the following spring after standard storage conditions. The per cent of seedlings with 4 or more viable tubers the following spring was considerably lower in the S_3 than in the S_0 , although the per cent of plants classified as 'tuberizing' was approximately the same for the S_0 and S_3 (Table 2). The problems with asexual reproduction indicate that harmful recessive genes may have been uncovered, the expression of which is delayed until after the first clonal generation, or until an unfavourable reaction between the environment and the weaker genotypes occurs. The genotypes in which lethal or semi-lethal genes are expressed are automatically excluded from contributing to the subsequent generation, which in turn affects the rate of inbreeding. In spite of these problems in reproduction, several fairly vigorous S_5 lines have now been obtained. Many of these lines exhibit a high degree of uniformity in leaf type, tuber shape and patterns of pigmentation. Other S_5 lines are still segregating for these same characters.

It appears, therefore, that it is possible to obtain homozygous clones for future genetic studies on the diploid level. Assuming no selection for heterozygosity, the diploid S_5 generation is about 97% homozygous. In comparison, the autotetraploid S_5 generation, assuming random chromosome segregation, would only be 53% homozygous (Bartlett and Haldane, 1934). The unavoidable selection for vigour during reproduction, although kept to a minimum, probably resulted to a certain extent in some selection for heterozygosity as well. However, the diploid S_5 generation probably represents the greatest degree of inbreeding ever imposed upon potatoes.

Inbreeding by selfing imposes selection for self-fertility and, although such lines have not yet been detected, lines homozygous for this trait might be expected. They would be very useful in investigating the genetics of self-fertility. Linkage relations with the gene(s) for self-fertility will have special interest. Simmonds (1966) reported a case of linkage of a harmful recessive gene to the S-locus in potato. If the gene(s) for self-fertility are linked to recessive lethals, it may be difficult to develop true breeding self-fertile clones.

The rapid decline of vigour upon inbreeding suggests that hybrid vigour may be obtained when two inbred clones with a high combining ability are crossed. Considering the success with hybrid vigour in other crops, this area certainly warrants further investigation in potatoes, especially on the diploid level where advanced inbred lines can be obtained in a relatively short time. A broader genetic base will probably be necessary to develop more diverse inbred lines that can be used as parents to produce maximum hybrid vigour.

Zusammenfassung

Inzucht bei kultivierten diploiden Kartoffeln

Inzucht bei kultivierten diploiden Kartoffeln wurde möglich, nachdem man herausgefunden hatte, dass Kreuzungen zwischen kultivierten diploiden Arten und gewissen Haploiden von *Solanum tuberosum* in hohem Masse selbstfertil sind. Die ursprüngliche genetische Basis dieses Versuches bestand aus 4 Familien, die durch Kreuzungen zwischen der Gruppe Phureja und der Gruppe Stenotomum mit der Gruppe *Tuberosum* haploid *US-W 4* erzielt wurden. Bei einer Anzahl der untersuchten Eigenschaften stimmte die durchschnittliche Inzuchtdepression der 4 Familien mit einer Kurve überein, die auf der Basis der Heterozygotie in einem diploiden Organismus (Tabellen 1, 2 und Abb. 1) berechnet wurde. Die vier Familien reagierten unterschiedlich auf die Einkreuzung (Tabelle 1). Dies kann teilweise auf genetische Unterschiede zwischen den ursprünglichen Elternpopulationen (S_0) und auf eine Auslese auf Frohwüchsigkeit sowohl im

geschlechtlichen als auch im ungeschlechtlichen Vermehrungszyklus zurückgeführt werden. Nur die kräftigeren Pflanzen blühten, und nur ein kleiner Teil davon war in den fortgeschrittenen Inzuchtgenerationen selbstfertil (Tabelle 4). Dies zeigt, dass Allele für Selbstunverträglichkeit in den Populationen noch vorhanden sind. Der niedrigere Grad der Selbstfertilität, ausgedrückt in Samen pro Frucht, weist beim Vergleich mit der Pollen- und der Eizellenfertilität darauf hin, dass rezessive lethale Gene vorhanden sind. Fortgeschrittene Inzuchtgenerationen wiesen eine abnehmende Knollenbildung auf (Tabelle 3). Beim Vergleich der Leistung der ersten Klongenerationen mit jener der ungeschlechtlichen Nachkommenschaften war die Depression in bezug auf die entsprechenden S_0 's bei den meisten Nachkommenschaften kleiner als bei ihren entsprechenden ungeschlechtlichen Elternpopulationen (Tabellen 1 und 2). Es wurden einige kräftige S_5 -Linien er-

zielt, manche von ihnen besitzen einen hohen Grad von Einheitlichkeit. Die rasche Abnahme der Frohwüchsigkeit nach der Einkreuzung deutet darauf hin, dass ein Heterosiseffekt bei Kreuzung fortgeschrittener Inzuchtklone erreicht werden könnte. Da Inzucht durch Selbstung eine

Auslese auf Selbstfertilität erfordert, könnte dieses Vorgehen einige homozygote, selbstfertile Linien hervorbringen, was für eine Untersuchung der Genetik der Selbstfertilität sehr nützlich wäre.

Résumé

L'autofécondation dans les pommes de terre cultivées diploïdes

L'autofécondation chez les pommes de terre cultivées diploïdes devint possible quand on découvrit que des croisements entre espèces cultivées diploïdes et certains haploïdes de *Solanum tuberosum* étaient hautement auto-fertiles. La base génétique initiale de la présente étude comprend 4 familles issues de croisements entre groupe Phureja et groupe Stenotomum avec le haploïde US-W4 du groupe Tuberosum. Pour un certain nombre de caractères, la dépression moyenne de consanguinité des 4 familles coïncide avec une courbe calculée sur la base de l'hétérozygotie dans un organisme diploïde (Tableaux 1, 2 et Fig. 1). Les 4 familles réagissaient différemment à l'autofécondation (Tableau 1). Ceci peut partiellement être attribué à des différences génétiques entre les populations parentales originales (S_0) et à une sélection pour la vigueur dans les cycles de reproduction à la fois sexuelle et asexuelle. Seules les plantes les plus vigoureuses fleurissaient, et seule une petite partie de celles-ci étaient auto-fertiles dans les générations successives autofécondées (Tableau 4). Ceci indique que les allèles d'autoincompatibilité sont toujours présents

dans les populations. Le niveau plus faible d'autofertilité exprimée par le nombre de graines par fruit, quand on la compare à la fertilité du pollen et de l'ovule fait supposer que des gènes létaux récessifs sont présents. Les générations autofécondées successives montrent une tubérisation décroissante (Tableau 3). Quand on compare les performances des premières générations clonales avec les descendance assexuées, la dépression, comparée aux S_0 respectifs, est plus faible chez la plupart des descendance que chez les populations parentales assexuées respectives (Tableaux 1 et 2). Plusieurs lignées S_5 bien vigoureuses ont été obtenues, plusieurs de celles-ci montrant un haut degré d'uniformité. Le déclin rapide dans la vigueur après autofécondation suggère qu'on peut obtenir des hybrides vigoureux de croisements entre clones précédemment autofécondés. Puisque la consanguinité par autofécondation impose une sélection pour l'autofertilité, on peut de la sorte produire quelques lignées homozygotes auto-fertiles qui pourraient être utiles pour une recherche sur la génétique de l'autofertilité.

References

- Bartlett, M. S. & Haldane, J. B. S., 1934. The theory of inbreeding in autotetraploids. *J. Genet.* 29: 175-180.
- Cipar, M. S., 1964. Self-compatibility in hybrids between Phureja and haploid Andigena clones of *Solanum tuberosum*. *Eur. Potato J.* 7: 152-160.
- Cipar, M. S., Peloquin, S. J., & Hougas, R. W., 1964. Inheritance of incompatibility in hybrids between *Solanum tuberosum* haploids and diploid species. *Euphytica* 13: 163-172.
- Engel, K. H., 1957. Grundlegende Fragen zu einem Schema fuer Arbeiten mit Inzuchten bei Kartoffeln. *Der Züchter* 27: 98-124.
- Feistritzer, W., 1952. Die Selbstungsanalyse, eine Voraussetzung fuer die Kreuzungszucht der Kartoffel. *Ztschr. Pflanzenzüchtung* 31: 173-195.
- Gowen, F. A., 1956. Selfing in the potato as a means of securing improved parental clones. Ph. D. Thesis. Univ. Minnesota, 104 p. Univ. Microfilms, Ann Arbor, Mich. (*Diss. Abstr.* 16: 2286).
- Hollenback, J. G., 1966. Variation in self- and pseudo-compatibility in haploid Tuberosum-Phureja

INBREEDING IN CULTIVATED DIPLOID POTATOES

- hybrid families. Ph. D. Thesis Univ. Wisconsin, 72 p. Univ. Microfilms. Ann Arbor, Mich. (*Diss. Abstr.* 28: 496-B).
- Hougas, R. W. & Peloquin, S. J., 1962. Exploitation of *Solanum* germplasm. In D. S. Correll: The potato and its wild relatives. Texas Research Foundation, Renner, Texas. p. 21–24.
- Krantz, F. A., 1924. Potato breeding methods. *Minnesota Agric. Exp. Stn. Tech. Bull.* 25. 32 p.
- Krantz, F. A., 1946. Potato breeding methods III. A suggested procedure for potato breeding. *Minnesota agric. Exp. Stn Tech. Bull.* 173. 24 p.
- Krantz, F. A. & Hutchins, A. E., 1929. Potato breeding methods II. Selection in inbred lines. *Minnesota agric. Exp. Stn Tech. Bull.* 58. 23 p.
- Mullin, R. & Lauer, F. I., 1966. Breeding behavior of F₁ and inbred potato clones. *Proc. Amer. Soc. Hort. Sci.* 89: 449–455.
- Peloquin, S. J. & Hougas, R. W., 1959. Decapitation and genetic markers as related to haploidy in *Solanum tuberosum*. *Eur. Potato J.* 2: 176–183.
- Rudorf, W., 1958. Zuchtmethoden. In: H. Kappert and W. Rudorf, Handbuch der Pflanzenzüchtung, Vol. III., p. 157. Paul Parey, Berlin and Hamburg.
- Simmonds, N. W., 1966. Linkage to the S-locus in diploid potatoes. *Heredity* 21: 473–479.
- Smiley, J. H., 1963. Behavior of self-fertility and its utilization in *Solanum* diploid species – haploid *Solanum tuberosum* hybrids. Ph. D. Thesis. Univ. Wisconsin, 33 p. Univ. Microfilms. Ann Arbor, Mich. (*Diss. Abstr.* 24: 2234).