

Under the Paperwork Reduction Act of 1995 no persons are required to respond to a collection of information unless it displays a valid OMB control number.

UTILITY PATENT APPLICATION TRANSMITTAL

(Only for new nonprovisional applications under 37 CFR 1.53(b))

Attorney Docket No.	224990/10-039C4-US/459258
First Named Inventor	Jo Ann Wilson
Title	Inventor for Preparing Composite and Prioritized Composite Con
Priority Mail Express® Label No.	

APPLICATION ELEMENTS See MPEP chapter 600 concerning utility patent application contents.	ADDRESS TO: Commissioner for Patents P.O. Box 1450 Alexandria, VA 22313-1450
☒ Fee Transmittal Form (PTO/SB/17 or equivalent) ☐ Applicant asserts small entity status. See 37 CFR 1.27 ☐ Applicant certifies micro entity status. See 37 CFR 1.29. Applicant must attach form PTO/SB/15A or B or equivalent. ☒ Specification [Total Pages <u>47</u>] Both the claims and abstract must start on a new page. (See MPEP § 602.01(a) for information on the preferred arrangement) ☐ Drawing(s) (35 U.S.C. 113) [Total Sheets _____] ☐ Inventor's Oath or Declaration [Total Pages <u>2</u>] (including substitute statements under 37 CFR 1.64 and assignments serving as an oath or declaration under 37 CFR 1.63(e)) ☐ Newly executed (original or copy) ☒ A copy from a prior application (37 CFR 1.63(d)) ☒ Application Data Sheet * See note below. See 37 CFR 1.76 (PTO/AIA/14 or equivalent) ☐ CD-ROM or CD-R in duplicate, large table, or Computer Program (Appendix) ☐ Landscape Table on CD ☐ Nucleotide and/or Amino Acid Sequence Submission (if applicable, items a. - c. are required) ☐ Computer Readable Form (CRF) ☐ Specification Sequence Listing on: ☐ CD-ROM or CD-R (2 copies); or ☐ Paper ☐ Statements verifying identity of above copies	ACCOMPANYING APPLICATION PAPERS ☐ Assignment Papers (cover sheet & document(s)) Name of Assignee _____ ☐ 37 CFR 3.73(c) Statement ☒ Power of Attorney (when there is an assignee) ☐ English Translation Document (if applicable) ☐ Information Disclosure Statement (PTO/SB/08 or PTO-1449) ☐ Copies of citations attached ☒ Preliminary Amendment ☐ Return Receipt Postcard (MPEP § 503) (Should be specifically itemized) ☐ Certified Copy of Priority Document(s) (if foreign priority is claimed) ☐ Nonpublication Request Under 35 U.S.C. 122(b)(2)(B)(i). Applicant must attach form PTO/SB/35 or equivalent. ☒ Other: <u>Cover Letter to the Commissioner for Patents;</u> <u>Certificate of Transmission under 37 C.F.R. § 1.8(a)</u>

*Note: (1) Benefit claims under 37 CFR 1.78 and foreign priority claims under 1.55 must be included in an Application Data Sheet (ADS).
 (2) For applications filed under 35 U.S.C. 111, the application must contain an ADS specifying the applicant if the applicant is an assignee, person to whom the inventor is under an obligation to assign, or person who otherwise shows sufficient proprietary interest in the matter. See 37 CFR 1.46(b).

19. CORRESPONDENCE ADDRESS

☒ The address associated with Customer Number: 91684 OR ☐ Correspondence address below

Name			
Address			
City	State	Zip Code	
Country	Telephone	Email	
Signature			Date
Name (Print/Type)	Heidi M. Berven, Ph.D.		Registration No. (Attorney/Agent)
		48,951	

This collection of information is required by 37 CFR 1.53(b). The information is required to obtain or retain a benefit by the public which is to file (and by the USPTO to process) an application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.11 and 1.14. This collection is estimated to take 12 minutes to complete, including gathering, preparing, and submitting the completed application form to the USPTO. Time will vary depending upon the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, P.O. Box 1450, Alexandria, VA 22313-1450. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.

If you need assistance in completing this form, call 1-800-PTO-9199 and select option 2.

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

Application Data Sheet 37 CFR 1.76		Attorney Docket Number	224990/10-039C4-US/459258
		Application Number	
Title of Invention	Processes for Preparing Quinoline Compounds and Pharmaceutical Compositions Containing Such Compounds		
The application data sheet is part of the provisional or nonprovisional application for which it is being submitted. The following form contains the bibliographic data arranged in a format specified by the United States Patent and Trademark Office as outlined in 37 CFR 1.76. This document may be completed electronically and submitted to the Office in electronic format using the Electronic Filing System (EFS) or the document may be printed and included in a paper filed application.			

Secrecy Order 37 CFR 5.2:

☐ Portions or all of the application associated with this Application Data Sheet may fall under a Secrecy Order pursuant to 37 CFR 5.2 (Paper filers only. Applications that fall under Secrecy Order may not be filed electronically.)

Inventor Information:

Inventor 1					Remove	
Legal Name						
Prefix	Given Name	Middle Name	Family Name	Suffix		
	Jo Ann		Wilson			
Residence Information (Select One) ☒ US Residency ☐ Non US Residency ☐ Active US Military Service						
City	San Francisco	State/Province	CA	Country of Residence	US	
Mailing Address of Inventor:						
Address 1		74 Seward Street				
Address 2						
City	San Francisco	State/Province	CA			
Postal Code	94114	Country i	US			
Inventor 2					Remove	
Legal Name						
Prefix	Given Name	Middle Name	Family Name	Suffix		
	Khalid		Shah			
Residence Information (Select One) ☒ US Residency ☐ Non US Residency ☐ Active US Military Service						
City	Half Moon Bay	State/Province	CA	Country of Residence	US	
Mailing Address of Inventor:						
Address 1		281 Eagle Trace Drive				
Address 2						
City	Half Moon Bay	State/Province	CA			
Postal Code	94019	Country i	US			
All Inventors Must Be Listed - Additional Inventor Information blocks may be generated within this form by selecting the Add button.						
Add						

Correspondence Information:

Enter either Customer Number or complete the Correspondence Information section below.
For further information see 37 CFR 1.33(a).

Application Data Sheet 37 CFR 1.76		Attorney Docket Number	224990/10-039C4-US/459258
		Application Number	
Title of Invention	Processes for Preparing Quinoline Compounds and Pharmaceutical Compositions Containing Such Compounds		

☐ An Address is being provided for the correspondence information of this application.

Customer Number	91684		
Email Address	patent@honigman.com	Add Email	Remove Email

Application Information:

Title of the Invention	Processes for Preparing Quinoline Compounds and Pharmaceutical Compositions Containing Such Compounds		
Attorney Docket Number	224990/10-039C4-US/459258	Small Entity Status Claimed	☐
Application Type	Nonprovisional		
Subject Matter	Utility		
Total Number of Drawing Sheets (if any)		Suggested Figure for Publication (if any)	

Filing By Reference:

Only complete this section when filing an application by reference under 35 U.S.C. 111(c) and 37 CFR 1.57(a). Do not complete this section if application papers including a specification and any drawings are being filed. Any domestic benefit or foreign priority information must be provided in the appropriate section(s) below (i.e., "Domestic Benefit/National Stage Information" and "Foreign Priority Information").

For the purposes of a filing date under 37 CFR 1.53(b), the description and any drawings of the present application are replaced by this reference to the previously filed application, subject to conditions and requirements of 37 CFR 1.57(a).

Application number of the previously filed application	Filing date (YYYY-MM-DD)	Intellectual Property Authority or Country

Publication Information:

☐ Request Early Publication (Fee required at time of Request 37 CFR 1.219)

Request Not to Publish. I hereby request that the attached application not be published under

☐ 35 U.S.C. 122(b) and certify that the invention disclosed in the attached application **has not and will not** be the subject of an application filed in another country, or under a multilateral international agreement, that requires publication at eighteen months after filing.

Representative Information:

Representative information should be provided for all practitioners having a power of attorney in the application. Providing this information in the Application Data Sheet does not constitute a power of attorney in the application (see 37 CFR 1.32). Either enter Customer Number or complete the Representative Name section below. If both sections are completed the customer Number will be used for the Representative Information during processing.

Please Select One:	☒ Customer Number	☐ US Patent Practitioner	☐ Limited Recognition (37 CFR 11.9)
Customer Number	91684		

Application Data Sheet 37 CFR 1.76		Attorney Docket Number	224990/10-039C4-US/459258
		Application Number	
Title of Invention	Processes for Preparing Quinoline Compounds and Pharmaceutical Compositions Containing Such Compounds		

Domestic Benefit/National Stage Information:

This section allows for the applicant to either claim benefit under 35 U.S.C. 119(e), 120, 121, 365(c), or 386(c) or indicate National Stage entry from a PCT application. Providing benefit claim information in the Application Data Sheet constitutes the specific reference required by 35 U.S.C. 119(e) or 120, and 37 CFR 1.78.

When referring to the current application, please leave the "Application Number" field blank.

Prior Application Status	Pending	Remove			
Application Number	Continuity Type	Prior Application Number	Filing or 371(c) Date (YYYY-MM-DD)		
	Division of	16151653	2018-10-04		
Prior Application Status	Patented	Remove			
Application Number	Continuity Type	Prior Application Number	Filing Date (YYYY-MM-DD)	Patent Number	Issue Date (YYYY-MM-DD)
16151653	Division of	15348716	2016-11-10	10123999	2018-11-13
Prior Application Status	Patented	Remove			
Application Number	Continuity Type	Prior Application Number	Filing Date (YYYY-MM-DD)	Patent Number	Issue Date (YYYY-MM-DD)
15348716	Division of	13984559	2014-03-20	9717720	2017-08-01
Prior Application Status	Expired	Remove			
Application Number	Continuity Type	Prior Application Number	Filing or 371(c) Date (YYYY-MM-DD)		
13984559	a 371 of international	PCT/US2012/024591	2012-02-10		
Prior Application Status	Expired	Remove			
Application Number	Continuity Type	Prior Application Number	Filing or 371(c) Date (YYYY-MM-DD)		
PCT/US2012/024591	Claims benefit of provisional	61441520	2011-02-10		
Prior Application Status	Expired	Remove			
Application Number	Continuity Type	Prior Application Number	Filing or 371(c) Date (YYYY-MM-DD)		
PCT/US2012/024591	Claims benefit of provisional	61441527	2011-02-10		
Additional Domestic Benefit/National Stage Data may be generated within this form by selecting the Add button.					

Foreign Priority Information:

Application Data Sheet 37 CFR 1.76		Attorney Docket Number	224990/10-039C4-US/459258
		Application Number	
Title of Invention	Processes for Preparing Quinoline Compounds and Pharmaceutical Compositions Containing Such Compounds		

This section allows for the applicant to claim priority to a foreign application. Providing this information in the application data sheet constitutes the claim for priority as required by 35 U.S.C. 119(b) and 37 CFR 1.55. When priority is claimed to a foreign application that is eligible for retrieval under the priority document exchange program (PDX)ⁱ the information will be used by the Office to automatically attempt retrieval pursuant to 37 CFR 1.55(i)(1) and (2). Under the PDX program, applicant bears the ultimate responsibility for ensuring that a copy of the foreign application is received by the Office from the participating foreign intellectual property office, or a certified copy of the foreign priority application is filed, within the time period specified in 37 CFR 1.55(g)(1).

			Remove
Application Number	Country ⁱ	Filing Date (YYYY-MM-DD)	Access Code ¹ (if applicable)

Additional Foreign Priority Data may be generated within this form by selecting the **Add** button.

Statement under 37 CFR 1.55 or 1.78 for AIA (First Inventor to File) Transition Applications

This application (1) claims priority to or the benefit of an application filed before March 16, 2013 and (2) also contains, or contained at any time, a claim to a claimed invention that has an effective filing date on or after March 16, 2013. ☐ NOTE: By providing this statement under 37 CFR 1.55 or 1.78, this application, with a filing date on or after March 16, 2013, will be examined under the first inventor to file provisions of the AIA.
--

Application Data Sheet 37 CFR 1.76		Attorney Docket Number	224990/10-039C4-US/459258
		Application Number	
Title of Invention	Processes for Preparing Quinoline Compounds and Pharmaceutical Compositions Containing Such Compounds		

Authorization or Opt-Out of Authorization to Permit Access:

When this Application Data Sheet is properly signed and filed with the application, applicant has provided written authority to permit a participating foreign intellectual property (IP) office access to the instant application-as-filed (see paragraph A in subsection 1 below) and the European Patent Office (EPO) access to any search results from the instant application (see paragraph B in subsection 1 below).

Should applicant choose not to provide an authorization identified in subsection 1 below, applicant **must opt-out** of the authorization by checking the corresponding box A or B or both in subsection 2 below.

NOTE: This section of the Application Data Sheet is **ONLY** reviewed and processed with the **INITIAL** filing of an application. After the initial filing of an application, an Application Data Sheet cannot be used to provide or rescind authorization for access by a foreign IP office(s). Instead, Form PTO/SB/39 or PTO/SB/69 must be used as appropriate.

1. Authorization to Permit Access by a Foreign Intellectual Property Office(s)

A. Priority Document Exchange (PDX) - Unless box A in subsection 2 (opt-out of authorization) is checked, the undersigned hereby **grants the USPTO authority** to provide the European Patent Office (EPO), the Japan Patent Office (JPO), the Korean Intellectual Property Office (KIPO), the State Intellectual Property Office of the People's Republic of China (SIPO), the World Intellectual Property Organization (WIPO), and any other foreign intellectual property office participating with the USPTO in a bilateral or multilateral priority document exchange agreement in which a foreign application claiming priority to the instant patent application is filed, access to: (1) the instant patent application-as-filed and its related bibliographic data, (2) any foreign or domestic application to which priority or benefit is claimed by the instant application and its related bibliographic data, and (3) the date of filing of this Authorization. See 37 CFR 1.14(h)(1).

B. Search Results from U.S. Application to EPO - Unless box B in subsection 2 (opt-out of authorization) is checked, the undersigned hereby **grants the USPTO authority** to provide the EPO access to the bibliographic data and search results from the instant patent application when a European patent application claiming priority to the instant patent application is filed. See 37 CFR 1.14(h)(2).

The applicant is reminded that the EPO's Rule 141(1) EPC (European Patent Convention) requires applicants to submit a copy of search results from the instant application without delay in a European patent application that claims priority to the instant application.

2. Opt-Out of Authorizations to Permit Access by a Foreign Intellectual Property Office(s)

☐ A. Applicant **DOES NOT** authorize the USPTO to permit a participating foreign IP office access to the instant application-as-filed. If this box is checked, the USPTO will not be providing a participating foreign IP office with any documents and information identified in subsection 1A above.

☐ B. Applicant **DOES NOT** authorize the USPTO to transmit to the EPO any search results from the instant patent application. If this box is checked, the USPTO will not be providing the EPO with search results from the instant application.

NOTE: Once the application has published or is otherwise publicly available, the USPTO may provide access to the application in accordance with 37 CFR 1.14.

Application Data Sheet 37 CFR 1.76		Attorney Docket Number	224990/10-039C4-US/459258
		Application Number	
Title of Invention	Processes for Preparing Quinoline Compounds and Pharmaceutical Compositions Containing Such Compounds		

Applicant Information:

Providing assignment information in this section does not substitute for compliance with any requirement of part 3 of Title 37 of CFR to have an assignment recorded by the Office.			
Applicant 1			
If the applicant is the inventor (or the remaining joint inventor or inventors under 37 CFR 1.45), this section should not be completed. The information to be provided in this section is the name and address of the legal representative who is the applicant under 37 CFR 1.43; or the name and address of the assignee, person to whom the inventor is under an obligation to assign the invention, or person who otherwise shows sufficient proprietary interest in the matter who is the applicant under 37 CFR 1.46. If the applicant is an applicant under 37 CFR 1.46 (assignee, person to whom the inventor is obligated to assign, or person who otherwise shows sufficient proprietary interest) together with one or more joint inventors, then the joint inventor or inventors who are also the applicant should be identified in this section.			
			Clear
☒ Assignee	☐ Legal Representative under 35 U.S.C. 117	☐ Joint Inventor	
☐ Person to whom the inventor is obligated to assign.	☐ Person who shows sufficient proprietary interest		
If applicant is the legal representative, indicate the authority to file the patent application, the inventor is:			
Name of the Deceased or Legally Incapacitated Inventor:			
If the Applicant is an Organization check here. ☒			
Organization Name	Exelixis, Inc.		
Mailing Address Information For Applicant:			
Address 1	1851 Harbor Bay Parkway		
Address 2			
City	Alameda	State/Province	CA
Country	US	Postal Code	94502
Phone Number		Fax Number	
Email Address			
Additional Applicant Data may be generated within this form by selecting the Add button.			

Assignee Information including Non-Applicant Assignee Information:

Providing assignment information in this section does not substitute for compliance with any requirement of part 3 of Title 37 of CFR to have an assignment recorded by the Office.

Application Data Sheet 37 CFR 1.76		Attorney Docket Number	224990/10-039C4-US/459258	
		Application Number		
Title of Invention	Processes for Preparing Quinoline Compounds and Pharmaceutical Compositions Containing Such Compounds			

Assignee 1

Complete this section if assignee information, including non-applicant assignee information, is desired to be included on the patent application publication. An assignee-applicant identified in the "Applicant Information" section will appear on the patent application publication as an applicant. For an assignee-applicant, complete this section only if identification as an assignee is also desired on the patent application publication.

If the Assignee or Non-Applicant Assignee is an Organization check here. ☐

Prefix	Given Name	Middle Name	Family Name	Suffix

Mailing Address Information For Assignee including Non-Applicant Assignee:

Address 1				
Address 2				
City		State/Province		
Country		Postal Code		
Phone Number		Fax Number		
Email Address				

Additional Assignee or Non-Applicant Assignee Data may be generated within this form by selecting the Add button.

Signature:

NOTE: This Application Data Sheet must be signed in accordance with 37 CFR 1.33(b). However, if this Application Data Sheet is submitted with the INITIAL filing of the application and either box A or B is not checked in subsection 2 of the "Authorization or Opt-Out of Authorization to Permit Access" section, then this form must also be signed in accordance with 37 CFR 1.14(c).

This Application Data Sheet must be signed by a patent practitioner if one or more of the applicants is a juristic entity (e.g., corporation or association). If the applicant is two or more joint inventors, this form must be signed by a patent practitioner, all joint inventors who are the applicant, or one or more joint inventor-applicants who have been given power of attorney (e.g., see USPTO Form PTO/AIA/81) on behalf of all joint inventor-applicants.

See 37 CFR 1.4(d) for the manner of making signatures and certifications.

Signature			Date (YYYY-MM-DD)	2019-12-06	
First Name	Heidi M.	Last Name	Berven	Registration Number	48,951

Additional Signature may be generated within this form by selecting the Add button.

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

Application Data Sheet 37 CFR 1.76		Attorney Docket Number	224990/10-039C4-US/459258
		Application Number	
Title of Invention	Processes for Preparing Quinoline Compounds and Pharmaceutical Compositions Containing Such Compounds		

This collection of information is required by 37 CFR 1.76. The information is required to obtain or retain a benefit by the public which is to file (and by the USPTO to process) an application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.14. This collection is estimated to take 23 minutes to complete, including gathering, preparing, and submitting the completed application data sheet form to the USPTO. Time will vary depending upon the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, P.O. Box 1450, Alexandria, VA 22313-1450. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. **SEND TO: Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.**



WIPO | PCT



(10) International Publication Number

WO 2012/109510 A1

(43) International Publication Date
16 August 2012 (16.08.2012)

(51) International Patent Classification:

C07D 215/227 (2006.01) A61P 35/00 (2006.01)
A61K 31/47 (2006.01)

(21) International Application Number:

PCT/US2012/024591

(22) International Filing Date:

10 February 2012 (10.02.2012)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:


61/441,520 10 February 2011 (10.02.2011) US
61/441,527 10 February 2011 (10.02.2011) US(71) Applicant (for all designated States except US): EXELIX-
IS, INC. [US/US]; 210 East Grand Avenue, South San
Francisco, CA 94080 (US).

(72) Inventor; and

(75) Inventor/Applicant (for US only): WILSON, Jo Ann
[US/US]; 74 Seward Street, San Francisco, CA 94114
(US).(74) Agent: BERVEN, Heidi, M.; Honigman, Miller Schwart
And Cohn, LLP, 350 East Michigan Avenue, Suite 300,
Kalamazoo, MI 49007-3800 (US).(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ,
CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO,
DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN,
HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR,
KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME,
MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ,
OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD,
SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR,
TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ,
UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU,
TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE,
DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU,
LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK,
SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: PROCESSES FOR PREPARING QUINOLINE COMPOUNDS AND PHARMACEUTICAL COMPOSITIONS CON-
TAINING SUCH COMPOUNDS(57) Abstract: The present invention is directed to processes for making and compositions containing quinolines such as formula I
or pharmaceutically acceptable salts thereof wherein: X1 is H, Br, Cl, or X2 is H, Br, Cl, or n1 is 1-2; and n2 is 1-2.

PROCESSES FOR PREPARING QUINOLINE COMPOUNDS AND PHARMACEUTICAL COMPOSITIONS CONTAINING SUCH COMPOUNDS

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the benefit of priority to U.S. Provisional Application No. 61/441,520, filed February 10, 2011, and 61/441,527, filed February 10, 2011, which are incorporated herein by reference

FIELD OF THE INVENTION

[0002] This disclosure relates to processes for preparing compounds useful for modulating protein kinase enzymatic activity. More specifically, this disclosure relates to processes for preparing quinolines that are useful for modulating cellular activities such as proliferation, differentiation, programmed cell death, migration, and chemo-invasion and to pharmaceutical compositions containing such compounds.

BACKGROUND OF THE INVENTION

[0003] Traditionally, dramatic improvements in the treatment of cancer are associated with identification of therapeutic agents acting through novel mechanisms. One mechanism that can be exploited in cancer treatment is the modulation of protein kinase activity because signal transduction through protein kinase activation is responsible for many of the characteristics of tumor cells. Protein kinase signal transduction is of particular relevance in, for example, renal, gastric, head and neck, lung, breast, prostate, and colorectal cancers; hepatocellular carcinoma; as well as in the growth and proliferation of brain tumor cells.

[0004] Protein kinases can be categorized as receptor type or non-receptor type. Receptor-type tyrosine kinases are comprised of a large number of transmembrane receptors with diverse biological activity. For a detailed discussion of the receptor-type tyrosine kinases, see Plowman et al., DN&P 7(6): 334-339, 1994. Since protein kinases and their ligands play critical roles in various cellular activities, deregulation of protein kinase enzymatic activity can lead to altered cellular properties, such as uncontrolled cell growth associated with cancer. In addition to oncological indications, altered kinase signaling is implicated in numerous other pathological diseases, including, for example, immunological disorders, cardiovascular diseases, inflammatory diseases, and degenerative diseases. Therefore, protein kinases are attractive targets for small molecule drug discovery. Particularly attractive targets for small-molecule modulation with respect to antiangiogenic

and antiproliferative activity include receptor type tyrosine kinases c-Met, KDR, c-Kit, Axl, flt-3, and flt-4.

[0005] The kinase c-Met is the prototypic member of a subfamily of heterodimeric receptor tyrosine kinases (RTKs) which include Met, Ron, and Sea. The endogenous ligand for c-Met is the hepatocyte growth factor (HGF), a potent inducer of angiogenesis. Binding of HGF to c-Met induces activation of the receptor via autophosphorylation, resulting in an increase of receptor dependent signaling, which promotes cell growth and invasion. Anti-HGF antibodies or HGF antagonists have been shown to inhibit tumor metastasis *in vivo* (See: Maulik et al Cytokine & Growth Factor Reviews 2002 13, 41-59). c-Met overexpression has been demonstrated on a wide variety of tumor types including breast, colon, renal, lung, squamous cell myeloid leukemia, hemangiomas, melanomas, astrocytomas, and glioblastomas. Additionally, activating mutations in the kinase domain of c-Met have been identified in hereditary and sporadic renal papilloma and squamous cell carcinoma. (See, e.g., Maulik et al., Cytokine & growth Factor reviews 2002 13, 41-59; Longati et al., Curr Drug Targets 2001, 2, 41-55; Funakoshi et al., Clinica Chimica Acta 2003 1-23).


[0006] Inhibition of epidermal growth factor (EGF), vascular endothelial growth factor (VEGF), and ephrin signal transduction will prevent cell proliferation and angiogenesis, two key cellular processes needed for tumor growth and survival (Matter A., Drug Disc. Technol. 2001 6, 1005-1024). Kinase KDR (refers to kinase insert domain receptor tyrosine kinase) and flt-4 (fms-like tyrosine kinase-4) are both VEGF receptors. Inhibition of EGF, VEGF, and ephrin signal transduction will prevent cell proliferation and angiogenesis, two key cellular processes needed for tumor growth and survival (Matter A. Drug Disc. Technol. 2001 6, 1005-1024). EGF and VEGF receptors are desirable targets for small molecule inhibition. All members of the VEGF family stimulate cellular responses by binding to tyrosine kinase receptors (the VEGFRs) on the cell surface, causing them to dimerize and become activated through transphosphorylation. The VEGF receptors have an extracellular portion with immunoglobulin-like domains, a single transmembrane spanning region, and an intracellular portion containing a split tyrosine-kinase domain. VEGF binds to VEGFR-1 and VEGFR-2. VEGFR-2 is known to mediate almost all of the known cellular responses to VEGF.

[0007] Kinase c-Kit (also called stem cell factor receptor or steel factor receptor) is a type 3 receptor tyrosine kinase (RTK) belonging to the platelet-derived growth factor receptor subfamily. Overexpression of c-Kit and c-Kit ligand has been described in variety of human

diseases, including human gastrointestinal stromal tumors, mastocytosis, germ cell tumors, acute myeloid leukemia (AML), NK lymphoma, small-cell lung cancer, neuroblastomas, gynecological tumors, and colon carcinoma. Moreover, elevated expression of c-Kit may also relate to the development of neoplasia associated with neurofibromatosis type 1 (NF-1), mesenchymal tumors GISTs, and mast cell disease, as well as other disorders associated with activated c-Kit.

[0008] Kinase Flt-3 (fms-like tyrosine kinase-3) is constitutively activated via mutation, either in the juxtamembrane region or in the activation loop of the kinase domain, in a large proportion of patients with AML (Reilly, *Leuk. Lymphoma*, 2003, 44: 1-7).

[0009] Small-molecule compounds that specifically inhibit, regulate, and/or modulate the signal transduction of kinases, such as c-Met, VEGFR2, KDR, c-Kit, Axl, flt-3, and flt-4 described above, are particularly desirable as a means to treat or prevent disease states associated with abnormal cell proliferation and angiogenesis. One such small-molecule is compound IA, which has the chemical structure:



IA

WO2005/030140 describes the synthesis of compound IA (Table 2, Compound 12, Example 48) and also discloses the therapeutic activity of this molecule to inhibit, regulate, and/or modulate the signal transduction of kinases (Assays, Table 4, entry 289), the entire contents of which is incorporated herein by reference.


[0010] Although therapeutic efficacy is the primary concern for a therapeutic agent, the pharmaceutical composition can be equally important to its development. Generally, the drug developer endeavors to discover a pharmaceutical composition that possesses desirable properties, such as satisfactory water-solubility (including rate of dissolution), storage stability, hygroscopicity, and reproducibility, all of which can impact the processability, manufacture, and/or bioavailability of the drug.

[0011] Accordingly, there is a need for the discovery of new processes for making quinolines such as compound IA that minimize the formation of undesirable process contaminants or byproducts. There is also a need for new pharmaceutical compositions containing quinolines such as compound IA that are essentially free of process byproducts.

SUMMARY OF THE INVENTION

[0012] These and other needs are met by the present disclosure, which is directed to processes for making and compositions containing quinolines or pharmaceutically acceptable salts thereof.

[0013] In one aspect, the disclosure relates to processes for preparing a compound of formula I:



or a pharmaceutically acceptable salt thereof, wherein:

X^1 is H, Br, Cl, or F;

X^2 is H, Br, Cl, or F;


n_1 is 1-2; and

n_2 is 1-2.


[0014] Intermediates useful in preparing the above compounds are also disclosed.

[0015] The compounds of formula I are useful as protein kinase modulators, and they inhibit various protein kinases including Ret and c-Met.

[0016] In another aspect, the disclosure provides a process for preparing compound IB:



from compound IA:




IA,

comprising:


- (a) heating and agitating a mixture comprising compound IA and L-malic acid, methylethyl ketone, and water;
- (b) cooling the mixture;
- (c) vacuum distilling the mixture successively; and
- (d) isolating the compound of IB by filtration.

[0017] In another aspect, the disclosure provides a process for preparing compound IB:



IB

from compound IA:



IA,


comprising:

- (a) heating and agitating a mixture comprising compound IA and L-malic acid, methylethyl ketone, and water;
- (b) cooling the mixture;
- (c) seeding the mixture with compound IB;
- (d) vacuum distilling the mixture; and

(e) isolating compound IB by filtration.

[0018] In another aspect, the disclosure provides compound I, IA, or IB admixed with

less than 100 ppm 6,7-dimethoxy-quinoline-4-ol, the structure of which is



[0019] In another aspect, the disclosure provides pharmaceutical compositions containing the compound of formula I, compound IA, or compound IB for oral administration.

[0020] In another aspect, the disclosure provides a pharmaceutical tablet composition according to Table 1.

Table 1

Ingredient	% w/w
Compound I	31.68
Microcrystalline Cellulose (MCC) (Avicel PH102)	38.85
Lactose anhydrous 60 M	19.42
Hydroxypropyl Cellulose, EXF	3.00
Croscarmellose Sodium	3.00
Total Intra-granular	95.95
Silicon dioxide, Colloidal PWD	0.30
Croscarmellose Sodium	3.00
Magnesium Stearate	0.75
Total	100.00

[0021] In another aspect, the disclosure provides a pharmaceutical tablet composition according to Table 2.

Table 2

Ingredient	(% w/w)
Compound I	25.0-33.3
Microcrystalline Cellulose, NF	q.s
Hydroxypropyl Cellulose, NF	3
Poloxamer, NF	0-3
Croscarmellose Sodium, NF	6.0
Colloidal Silicon Dioxide, NF	0.5
Magnesium Stearate, NF	0.5-1.0
Total	100

[0022] In another aspect, the disclosure provides a pharmaceutical tablet composition according to Table 2A.

Table 2A

Ingredient	% w/w
Compound IB (10 % drug load as Compound IA)	12.67
MCC	51.52
Lactose	25.76
Hydroxypropyl cellulose	3.0
Croscarmellose Sodium	6.0
Colloidal Silicon Dioxide	0.3
Magnesium Stearate	0.75
Total	100

[0023] In another aspect, the disclosure provides a pharmaceutical capsule composition according to Table 3.

Table 3

Ingredient	mg/unit dose
Compound IB (10 % drug load as Compound IA)	25
Silicified Microcrystalline Cellulose	196.75
Croscarmellose sodium	12.5
Sodium starch glycolate	12.5
Fumed Silica	0.75
Stearic acid	2.5
Total Fill Weight	250

[0024] In another aspect, the disclosure provides a pharmaceutical capsule composition according to Table 4.

Table 4

Ingredient	mg/unit dose
Compound IB (50 % drug load as Compound IA)	100
Silicified Microcrystalline Cellulose	75.40
Croscarmellose sodium	10.00
Sodium Starch Glycolate	10.00
Fumed silica	0.6
Stearic Acid	4.0
Total Fill Weight	200

[0025] In another aspect, the disclosure provides a pharmaceutical capsule composition according to Table 5, wherein the IB weight equivalents are provided.

Table 5


	mg/unit dose
Ingredient	50mg
Compound IB	63.35
Microcrystalline Cellulose	95.39
Croscarmellose sodium	9.05
Sodium starch glycolate	9.05
Fumed Silica	0.54
Stearic acid	3.62
Total Fill Weight	181.00

[0026] In another aspect, the disclosure provides a pharmaceutical capsule composition according to Table 6, wherein the IB weight equivalents are provided.

Table 6

	mg/unit dose
Ingredient	60mg
Compound IB	73.95
Microcrystalline Cellulose	114.36
Croscarmellose sodium	10.85
Sodium starch glycolate	10.85
Fumed Silica	0.65
Stearic acid	4.34
Total Fill Weight	217.00

[0027] In another aspect, the invention is directed to a pharmaceutical composition comprising compound I, IA, or IB admixed with less than 100 ppm 6,7-dimethoxy-quinoline-



4-ol, the structure of which is

[0028] There are many different aspects and embodiments of the disclosure described herein, and each aspect and each embodiment is non-limiting in regard to the scope of the disclosure. The terms "aspects" and "embodiments" are meant to be non-limiting regardless of where the terms "aspect" or "embodiment" appears in this specification. The transitional term "comprising," as used herein, which is synonymous with "including," "containing," or "characterized by," is inclusive or open-ended and does not exclude additional, unrecited elements.

DETAILED DESCRIPTION OF THE INVENTION

Definitions

[0029] As used herein, the following words and phrases are generally intended to have the meanings as set forth below, except to the extent that the context in which they are used indicates otherwise or they are expressly defined to mean something different.

[0030] The word “can” is used in a non-limiting sense and in contradistinction to the word “must.” Thus, for example, in many aspects of the invention a certain element is described as “can” having a specified identity, which is meant to convey that the subject element is permitted to have that identity according to the invention but is not required to have it.

[0031] If a group “R” is depicted as “floating” on a ring system, then unless otherwise defined, the substituent(s) “R” can reside on any atom of the ring system, assuming replacement of a depicted, implied, or expressly defined hydrogen from one of the ring atoms, so long as a stable structure is formed.

[0032] When there are more than one such depicted “floating” groups, such as where there are two groups; then, unless otherwise defined, the “floating” groups can reside on any atoms of the ring system, again assuming each replaces a depicted, implied, or expressly defined hydrogen on the ring.

[0033] “Pharmaceutically acceptable salts” include acid addition salts.

[0034] “Pharmaceutically acceptable acid addition salt” refers to those salts that retain the biological effectiveness of the free bases and that are not biologically or otherwise undesirable, formed with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, and the like, or mixtures thereof, as well as organic acids such as acetic acid, trifluoroacetic acid, propionic acid, glycolic acid, pyruvic acid, oxalic acid, maleic acid, malonic acid, succinic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, p-toluenesulfonic acid, salicylic acid, and the like, or mixtures thereof.

[0035] “Essentially free” as used in the phrase “essentially free of process byproducts or contaminants,” means that a compound or composition as disclosed here in is admixed with 200 parts per million (ppm) or less of such byproducts or contaminants.


[0036] The disclosure is further illustrated by the following examples, which are not to be construed as limiting the disclosure in scope or spirit to the specific procedures described in them. Unless specified otherwise, the starting materials and various intermediates may be

obtained from commercial sources, prepared from commercially available organic compounds, or prepared using well-known synthetic processes.

Processes

Aspect 1: Processes for Making Compounds of Formula I

[0037] Aspect (1) of the invention relates to a process of preparing a compound of formula I:



or a pharmaceutically acceptable salt thereof, wherein:

X^1 is H, Br, Cl, or F;


X^2 is H, Br, Cl, or F;

n_1 is 1-2; and

n_2 is 1-2;

the process comprising:

contacting the compound of formula g(1) with reactant z(1) to yield the compound of formula I:



[0038] The reaction is advantageously carried out under suitable reaction conditions.

Non-limiting examples of suitable reaction conditions include using basic conditions. Non-limiting examples of basic conditions that can be used in Aspect (1) include the use of inorganic bases, such as aqueous KOH, NaOH, K_2CO_3 , Na_2CO_3 , K_3PO_4 , Na_3PO_4 , K_2HPO_4 , Na_2HPO_4 , and the like, or mixtures thereof. Other non-limiting examples of suitable reaction conditions include using suitable solvents. Non-limiting examples of suitable solvents that can be used include water miscible solvents, such as THF, acetone, ethanol, and the like, or mixtures thereof. Other non-limiting examples of suitable reaction conditions include using suitable temperatures. Suitable temperatures that may be used for the reaction include a

temperature at a range from about 10 °C to about 30 °C, or alternatively, at a range from about 15 °C to about 28 °C, or alternatively, at a range from about 20 °C to about 25 °C. The product formed by the reaction is in the free base form, and this free base form may be converted into a pharmaceutically acceptable salt thereof by processes known in the art. For example, the compound of formula I can be converted to the L-malate salt by the addition of L-malic acid and a suitable solvent.

[0039] Utilities of the compound of formula I are further described in WO 2005/030140 A2, which is incorporated herein by reference.

Embodiments of Aspect (1) Part A

- [0040] In another embodiment of Aspect (1), X^1 is Cl or F.
- [0041] In another embodiment of Aspect (1), X^2 is Cl or F.
- [0042] In another embodiment of Aspect (1), X^1 is F.
- [0043] In another embodiment of Aspect (1), X^2 is F.
- [0044] In another embodiment of Aspect (1), X^1 is H.
- [0045] In another embodiment of Aspect (1), X^2 is H.
- [0046] In another embodiment of Aspect (1), n_1 is 1.
- [0047] In another embodiment of Aspect (1), n_2 is 1.
- [0048] In another embodiment of Aspect (1), n_1 is 2.
- [0049] In another embodiment of Aspect (1), n_2 is 2.
- [0050] All compounds of formula I for Aspect (1) disclosed above include any of the disclosed alternative embodiments in Part A for each of X^1 , X^2 , n_1 or n_2 , in combination with any other of the disclosed alternative embodiments in Part A for each of X^1 , X^2 , n_1 , or n_2 , as well as a pharmaceutically acceptable salt of any such combination.


Embodiments of Aspect (1) Part B

- [0051] In another embodiment of Aspect (1), n_1 and n_2 are each 1.
- [0052] In another embodiment of Aspect (1), n_1 and n_2 are each 2.
- [0053] In another embodiment of Aspect (1), n_1 is 1; and n_2 is 2.
- [0054] In another embodiment of Aspect (1), n_1 is 2 and n_2 is 1.
- [0055] In another embodiment of Aspect (1), X^1 is H; and X^2 is F.
- [0056] In another embodiment of Aspect (1), X^1 is F; and X^2 is H.
- [0057] In another embodiment of Aspect (1), X^1 and X^2 are each H.
- [0058] In another embodiment of Aspect (1), X^1 and X^2 are each F.

- [0059] In another embodiment of Aspect (1), X^1 is Cl; and X^2 is H.
 [0060] In another embodiment of Aspect (1), X^1 is H; and X^2 is Cl.
 [0061] In another embodiment of Aspect (1), X^1 and X^2 are each Cl.
 [0062] In another embodiment of Aspect (1), X^1 is Cl; and X^2 is F.
 [0063] In another embodiment of Aspect (1), X^1 is F; and X^2 is Cl.

Embodiments of Aspect (1) Part C

- [0064] In an embodiment of Aspect (1), the compound of formula g(1) can be made by reacting a compound of formula f(1) with reactant y(1) to yield the compound of g(1):



wherein LG represents a leaving group, and each of X^2 , and $n2$ are as defined in Aspect (1), or as in any of the embodiments of Aspect (1) Part A. A non-limiting example of a leaving group includes a halo group such as Cl, Br, or F. Various compounds of reactant y(1) are commercially available, such as 2-fluoro-4-aminophenol and 4-aminophenol. Also, the skilled artisan would be able to make any variation of reactant y(1) using commercially available starting materials and by using known techniques to modify these commercially available starting materials to yield various compounds within the scope of reactant y(1).

- [0065] The reaction in this embodiment is advantageously carried out under suitable reaction conditions. Non-limiting examples of suitable reaction conditions include using suitable solvents such as polar solvents. Non-limiting examples of polar solvents that can be used include tetrahydrofuran (THF), dimethylacetamide (DMA), dimethylsulfoxide (DMSO), dimethylformamide (DMF), ethyl acetate, N-methyl pyrrolidone (NMP), propylene carbonate, and the like, or mixtures thereof. In another embodiment, the polar solvent is dimethylacetamide (DMA). In another embodiment, the polar solvent is dimethylsulfoxide (DMSO). In another embodiment, the polar solvent is dimethylformamide (DMF). In another embodiment, the polar solvent is ethyl acetate. In another embodiment, the polar solvent is N-methyl pyrrolidone (NMP). In another embodiment, the polar solvent is propylene carbonate. In another embodiment, the solvent is a mixture of solvents, such as a mixture comprising THF and DMA.

[0066] The reactants f(1) and y(1) can be added together at a temperature ranging from about 10 °C to about 30 °C, or alternatively, from about 15 °C to about 28 °C, or alternatively, from about 20 °C to about 25 °C. The mixture is then heated to a temperature ranging from about 80 °C to about 125 °C, or alternatively, from about 95 °C to about 110 °C, or alternatively, from about 100 °C to about 105 °C, and the selected temperature is maintained until the reaction is complete.

[0067] Other non-limiting examples of suitable reaction conditions in this step of Aspect (1) include the use of a suitable base, such as a metal hydroxide or a non-nucleophilic base. Examples of metal hydroxides include sodium hydroxide or potassium hydroxide. Non-limiting examples of non-nucleophilic bases that can be used include lithium diisopropylamide, lithium tetramethylpiperidide, and alkali metal alkoxides such as sodium tert-butoxide, potassium tert-butoxide, sodium-pentoxide, and the like, or mixtures thereof. Preferably, the base is sodium tert-butoxide or sodium tert-pentoxide. In one embodiment, the base is sodium tert-pentoxide. Typically the sodium tert-pentoxide is commercially available as 35 weight percent solution of base in tetrahydrofuran, or as a 95 weight percent solid reagent. Preferably, the sodium tert-pentoxide is a 95 weight percent solid.

[0068] Typically, approximately 1.1 to 3.0 molar equivalents of base are used relative the moles of f(1) that are used. More preferably, 1.3 to 2.5 molar equivalents of base are used relative the moles of f(1) that are used. More preferably, 1.5 to 2.2 molar equivalents of base are used relative the moles of f(1) that are used. More preferably, 1.7 to 2.1 molar equivalents of base are used relative the moles of f(1) that are used.

[0069] Typically, the amount of molar equivalents of amino phenol that are used exceeds the molar equivalents of base that are used. In one embodiment, 1.1 to 2 molar equivalents of amino phenol are used relative to the molar equivalents of base that are used.

[0070] Once the reaction is substantially complete, the reaction mixture can be cooled to a temperature ranging from about 10 °C to about 25 °C. Precooled water can be charged at a rate to maintain a temperature that ranges from about 5 °C to about 35 °C. Alternatively, the precooled water can be charged at a rate to maintain a temperature that ranges from about 10 °C to about 25 °C. As a non-limiting example, the precooled water can be at a temperature ranging from about 0 °C to about 10 °C. As another non-limiting example, the precooled water can be at a temperature ranging from about 2 °C to about 7 °C. The precipitate can be collected by filtration under standard conditions and purified by standard purification techniques.

Embodiments of Aspect (1) Part D

[0071] In an embodiment of Aspect (1), the compound of formula f(1) can be made by converting a compound of formula e(1) to the compound of formula f(1):




wherein LG represents a leaving group. Non-limiting examples of leaving groups that can be used include halo groups (e.g., Cl, Br, or F) that can be added by halogenating agents. Non-limiting examples of halogenating agents that can be used include chlorinating agents, such as SOCl_2 , SO_2Cl_2 , COCl_2 , PCl_5 , POCl_3 , and the like.

[0072] The reaction is advantageously carried out under suitable reaction conditions. Non-limiting examples of suitable reaction conditions in Part D of Aspect (1) include the use of suitable solvents. Non-limiting example of suitable solvents that can be used during the halogenation of the compound of formula e(1) include a polar, aprotic solvent, such as CH_3CN , DMF, and the like, or mixtures thereof. In other embodiments, the chlorination can be carried out using POCl_3 in acetonitrile, COCl_2 in DMF, or SOCl_2 in DMF. The addition of the chlorination agent is advantageously carried out at a temperature ranging from about 60 °C to about 90 °C. In another embodiment, the addition of the chlorination agent can be carried out at a temperature ranging from about 70 °C to about 85 °C. In another embodiment, the addition of the chlorination agent can be carried out at a temperature ranging from about 74 °C to about 80 °C. The product can then be collected by filtration and purified using standard techniques.

Embodiments of Aspect (1) Part E

[0073] In an embodiment of Aspect (1), reactant z(1) can be made by reacting reactant z(1a) with a chlorinating agent to yield reactant z(1):




wherein X^1 is Br, Cl, or F; and n1 is 1-2. Compounds of reactant z(1a) can be made according to the process described in Example 25 of WO 2005/030140 A2, and the skilled artisan would be able to make any necessary substitutions using commercially available starting materials to come up with various compounds within the scope of reactant z(1a). Example 25 in WO 2005/030140 A2 is incorporated herein by reference.

[0074] The reaction is advantageously carried out under suitable reaction conditions. Non-limiting examples of suitable reaction conditions include using a chlorinating agent such as POCl_3 , oxalyl chloride, and the like. In another embodiment, oxalyl chloride is used as a chlorinating agent. Non-limiting examples of suitable reaction conditions include carrying out the reaction at a temperature in the range from about 0 °C to about 25 °C, or alternatively at a temperature in the range from about 5 °C to about 20 °C. Other non-limiting examples of suitable reaction conditions include carrying out the reaction in a suitable solvent. Non-limiting examples of suitable solvents that can be used include polar aprotic solvents, such as halogenated hydrocarbons (e.g., dichloromethane and chloroform), ethers (e.g., Et_2O), dioxane, tetrahydrofuran (THF) containing catalytic DMF, and the like, or mixtures thereof. The resulting solution containing reactant z(1) can be used, without further processing, to make the compound of formula I.

Other Embodiments of Aspect (1)


[0075] In another embodiment of Aspect (1), the compound of formula I is a compound of formula IA-1:




or a pharmaceutically acceptable salt thereof, wherein:

X^1 is H, Cl, Br, or F; and X^2 is H, Cl, Br, or F. Compound IA can be in the free base form or it can be converted to a pharmaceutically acceptable salt thereof. Accordingly, compound IA can be converted to its L-malate salt by the addition of L-malic acid and a suitable solvent.


[0076] In another embodiment of Part D of Aspect (1), the compound of formula e(1) is compound e(2):



and the compound of formula f(1) is compound f(2):




[0077] In another embodiment of Part C of Aspect (1), the compound of formula f(1) is compound f(2):



reactant y(1) is reactant (y)(2):




wherein X^2 is hydrogen or fluoro; and
the compound of formula g(1) is of formula g(2):




[0078] In a further embodiment, the reaction employs a non-nucleophilic base. In a further embodiment, the non-nucleophilic base is an alkali metal alkoxide; and the reaction is carried out in a polar solvent. In a further embodiment, the alkali metal alkoxide is sodium *tert*-butoxide, and the polar solvent is DMA.


[0079] In another embodiment of Part C of Aspect (1), the compound of formula f(1) is compound f(2):




reactant y(1) is reactant (y)(3):




wherein X^2 is hydrogen or fluoro; and
the compound of formula g(1) is compound g(3):




[0080] In another embodiment of Aspect (1) of this disclosure, the compound of formula g(1) is compound g(3):



reactant z(1) is reactant (z)(2):



the compound of formula I is compound IA:



IA

[0081] In a further embodiment, the reaction is carried out in the presence of an inorganic base. In a further embodiment, the inorganic base is K₂CO₃, and the solvent employed in this reaction is a combination of THF and H₂O.

[0082] In another embodiment of Aspect (1) of this disclosure, X¹ and X² for each of formula g(1) and reactant z(1) are each selected from Cl or F. In another embodiment, X¹ and X² for each of formula g(2), and reactant z(1) are both F.


[0083] The compound of formula f(2), or a pharmaceutically acceptable salt thereof, can be made by converting the compound of formula e(2) to a compound of formula f(2) with a chlorinating agent in a suitable solvent:




The compound of formula f(2) can be in its free base form or converted to a pharmaceutically acceptable salt thereof. The reaction conditions that can be used in this aspect include any of the reaction conditions disclosed in Part E of Aspect (1).

Aspect 2: Processes for Making Compounds of Formula g(2)

[0084] Aspect (2) of the disclosure relates to a process of preparing compound g(2):



or a pharmaceutically acceptable salt thereof; the process comprising reacting compound f(2) with reactant y(3) under basic conditions (e.g., using 2,6-lutidine) in an appropriate solvent to yield compound g(3):




The reaction conditions that can be used in this aspect include any of the reaction conditions disclosed in Part C of Aspect (1).


[0085] Alternative reaction conditions that can be used in this aspect include any of the reaction conditions disclosed in Parts C and D of Aspect (1).

Aspect 3: Processes for Making Compounds of IB

[0086] As indicated above, in one aspect, the invention provides a process for preparing compound IB:



from compound IA:



IA;

comprising:

- (a) heating and agitating a mixture comprising compound IA and L-malic acid, methylethyl ketone, and water;
- (b) cooling the mixture;
- (c) vacuum distilling the mixture successively; and
- (d) isolating the compound of IB by filtration.

[0087] In one embodiment of this aspect, compound IA is admixed with a sufficient amount of L-malic acid in a methylethyl ketone (MEK)/water (1:1) mixture. Alternatively, L-malic acid is added as a solution in water to a mixture of compound IA in methyl ethyl ketone. Generally the amount of L-malic is greater than 1 molar equivalent relative to compound IA. The mixture of compound IA and L-malic acid in MEK/water is heated at about 40-70 °C, and preferably at about 50-60 °C, and more preferably at about 55-60 °C with agitation, such as by stirring or the like, for about 1 to about 5 hours. At the end of the heating, the mixture is optionally clarified by filtering to give a clear solution. The resulting clear solution is then vacuum distilled from 1 to about 5 times at 150 to 200 mm Hg and a maximum jacket temperature of 55 °C to provide the desired crystalline compound of IB.

[0088] In one embodiment, L-malic acid is charged as a solution in water to compound IA. Generally the amount of L-malic is greater than 1 molar equivalent relative to compound IA. The mixture of compound IA and L-malic acid in MEK/water is heated at about 40-70 °C, and preferably at about 50-60 °C, and more preferably at about 55-60 °C with agitation, such as by stirring or the like, for about 1 to about 5 hours. At the end of the heating, the mixture is optionally clarified by filtering to give a clear solution which is at a temperature of about 30-40 °C, and more preferably at a temperature of about 33-37 °C. This clear solution is optionally seeded to facilitate crystallization. After seeding, the resulting mixture is vacuum distilled as provided above.

[0089] In one embodiment, compound IB is in the N-1 form. In another embodiment, compound IB is in the N-2 form. In another embodiment, compound IB is a mixture of the N-1 form and the N-2 form. Processes for preparing the N-1 and N-2 forms of compound IB

are disclosed in WO 2010/083414 (PCT/US2010021194), the entire contents of which are incorporated herein by reference.

[0090] In another embodiment, the disclosure relates to compound IA or IB admixed with 100 ppm or less of 6,7-dimethoxy-quinoline-4-ol. In one embodiment, the compound is admixed with 50 ppm or less of 6,7-dimethoxy-quinoline-4-ol. In another embodiment, the compound is admixed with 25 ppm or less of 6,7-dimethoxy-quinoline-4-ol. In another embodiment, the compound is admixed with 10 ppm or less of 6,7-dimethoxy-quinoline-4-ol. In another embodiment, the compound is admixed with 5 ppm or less of 6,7-dimethoxy-quinoline-4-ol. In another embodiment, the compound is admixed with 2.5 ppm or less of 6,7-dimethoxy-quinoline-4-ol.

[0091] In another embodiment, the disclosure relates to compound IA admixed with 100 ppm or less of 6,7-dimethoxy-quinoline-4-ol. In one embodiment, the compound is admixed with 50 ppm or less of 6,7-dimethoxy-quinoline-4-ol. In another embodiment, the compound is admixed with 25 ppm or less of 6,7-dimethoxy-quinoline-4-ol. In another embodiment, the compound is admixed with 10 ppm or less of 6,7-dimethoxy-quinoline-4-ol. In another embodiment, the compound is admixed with 5 ppm or less of 6,7-dimethoxy-quinoline-4-ol. In another embodiment, the compound is admixed with 2.5 ppm or less of 6,7-dimethoxy-quinoline-4-ol.

[0092] In another embodiment, the disclosure relates to compound IB admixed with 100 ppm or less of 6,7-dimethoxy-quinoline-4-ol. In one embodiment, the compound is admixed with 50 ppm or less of 6,7-dimethoxy-quinoline-4-ol. In another embodiment, the compound is admixed with 25 ppm or less of 6,7-dimethoxy-quinoline-4-ol. In another embodiment, the compound is admixed with 10 ppm or less of 6,7-dimethoxy-quinoline-4-ol. In another embodiment, the compound is admixed with 5 ppm or less of 6,7-dimethoxy-quinoline-4-ol. In another embodiment, the compound is admixed with 2.5 ppm or less of 6,7-dimethoxy-quinoline-4-ol.

[0093] In another embodiment, the disclosure relates to compound IB in the N-1 form admixed with 100 ppm or less of 6,7-dimethoxy-quinoline-4-ol. In one embodiment, the compound is admixed with 50 ppm or less of 6,7-dimethoxy-quinoline-4-ol. In another embodiment, the compound is admixed with 25 ppm or less of 6,7-dimethoxy-quinoline-4-ol. In another embodiment, the compound is admixed with 10 ppm or less of 6,7-dimethoxy-quinoline-4-ol. In another embodiment, the compound is admixed with 5 ppm or less of 6,7-dimethoxy-quinoline-4-ol. In another embodiment, the compound is admixed with 2.5 ppm or less of 6,7-dimethoxy-quinoline-4-ol.

[0094] In another embodiment, the disclosure relates to compound IB in the N-2 form admixed with 100 ppm or less of 6,7-dimethoxy-quinoline-4-ol. In one embodiment, the compound is admixed with 50 ppm or less of 6,7-dimethoxy-quinoline-4-ol. In another embodiment, the compound is admixed with 25 ppm or less of 6,7-dimethoxy-quinoline-4-ol. In another embodiment, the compound is admixed with 10 ppm or less of 6,7-dimethoxy-quinoline-4-ol. In another embodiment, the compound is admixed with 5 ppm or less of 6,7-dimethoxy-quinoline-4-ol. In another embodiment, the compound is admixed with 2.5 ppm or less of 6,7-dimethoxy-quinoline-4-ol.

[0095] In another embodiment, the disclosure relates to compound IB as a mixture of the N-1 form and the N-2 form admixed with 100 ppm or less of 6,7-dimethoxy-quinoline-4-ol. In one embodiment, the compound is admixed with 50 ppm or less of 6,7-dimethoxy-quinoline-4-ol. In another embodiment, the compound is admixed with 25 ppm or less of 6,7-dimethoxy-quinoline-4-ol. In another embodiment, the compound is admixed with 10 ppm or less of 6,7-dimethoxy-quinoline-4-ol. In another embodiment, the compound is admixed with 5 ppm or less of 6,7-dimethoxy-quinoline-4-ol. In another embodiment, the compound is admixed with 2.5 ppm or less of 6,7-dimethoxy-quinoline-4-ol.

Pharmaceutical Compositions

[0096] In another embodiment, the disclosure relates to a pharmaceutical composition comprising Compound I, IA, or IB. Various carriers used in formulating pharmaceutically acceptable compositions and known techniques for their bulk preparation and subsequent production into unit dosage forms are employed to make the pharmaceutical compositions disclosed herein and are described in Remington: The Science and Practice of Pharmacy, 21st edition, 2005, ed. D. B. Troy, Lippincott Williams & Wilkins, Philadelphia, and Encyclopedia of Pharmaceutical Technology, eds. J. Swarbrick and J. C. Boylan, 1988-1999, Marcel Dekker, New York. The amount of carriers and excipients used in a composition can be varied proportionally according to the amount of active ingredient used (that is, Compound I, IA or IB).

[0097] In one embodiment, the pharmaceutical composition is a tablet.

[0098] In another embodiment, the pharmaceutical composition is a capsule.

[0099] In another embodiment, the pharmaceutical composition comprises Compound IA.

[00100] In another embodiment, the pharmaceutical composition comprises Compound IB.

[00101] In another embodiment, the pharmaceutical composition comprises Compound IB as the N-1 polymorph.

[00102] In another embodiment, the pharmaceutical composition comprises Compound IB as the N-2 polymorph.

[00103] In another embodiment, the pharmaceutical composition comprises Compound IB as a mixture of the N-1 form and the N-2 form.

[00104] In another embodiment, the disclosure relates to a pharmaceutical composition comprising Compound IA or IB; one or more fillers; one or more disintegrants; one or more glidants; and one or more lubricants.

[00105] In this embodiment, the filler comprises microcrystalline cellulose.

[00106] In this embodiment, the disintegrant comprises croscarmellose sodium.

[00107] In this embodiment, the disintegrant comprises croscarmellose sodium and sodium starch glycolate.

[00108] In this embodiment, the glidant comprises fumed silica.

[00109] In this embodiment, the lubricant comprises stearic acid.

[00110] In another embodiment, the disclosure relates to a pharmaceutical composition comprising Compound IA or IB; microcrystalline cellulose; lactose; hydroxypropyl cellulose; croscarmellose sodium; colloidal silicon dioxide; and magnesium stearate.

[00111] In another embodiment, the disclosure relates to a pharmaceutical composition comprising Compound IA or IB; microcrystalline cellulose; hydroxypropyl cellulose; a surfactant; croscarmellose sodium; colloidal silicon dioxide; and magnesium stearate.

[00112] In another embodiment, the disclosure relates to a pharmaceutical composition comprising Compound IA or IB; microcrystalline cellulose; croscarmellose sodium; fumed silica; and stearic acid.

[00113] In another embodiment, the disclosure relates to a pharmaceutical composition comprising Compound IA or IB; microcrystalline cellulose; anhydrous lactose; hydroxypropyl cellulose; croscarmellose sodium; silicon dioxide; and magnesium stearate.

[00114] In another embodiment, the disclosure relates to a pharmaceutical composition comprising Compound IA or IB; microcrystalline cellulose; anhydrous lactose; hydroxypropyl cellulose; a surfactant; croscarmellose sodium; silicon dioxide; and magnesium stearate.

[00115] In another aspect, the disclosure provides a pharmaceutical composition according to Tables 1, 2, 2A, 3, 4, 5, and 6 as provided above. The compositions are prepared according to methods available to the skilled artisan. For example, the Tablet


formulations are prepared by combining, blending, and compacting the components of the tablet compositions. The capsule compositions are prepared by combining and blending the components and then placing the blend in a gelatin capsule.

[00116] For example, the 25 mg capsules (Table 3, 10 percent drug load formulation) are prepared as follows. The drug substance is delumped through a mill. The delumped drug substance is then co-screened with an equal volume Prosolv HD90. The excipients, except for stearic acid, are screened and charged to a blender along with the co-screened drug substance. The mixture is blended in a V-Blender. This process is repeated to manufacture a second subplot of unlubricated blend. The two sublots are then combined together in a V-blender and lubricated with stearic acid which has been co-screened with an equal volume of unlubricated blend. The final blend is then encapsulated into opaque, size 1 gelatin capsules using an automated capsule filling machine. The capsules are then weight sorted through an automatic weight sorter.

[00117] The 100-mg capsules (Table 4, 50% drug load formulation) are manufactured in two equal sublots of 5 kg of blend which are combined prior to lubricant blend. The drug substance is delumped through a mill. The excipients, except for stearic acid, are screened and charged to the mixer along with the delumped drug substance. The mixture is blended with a high shear mixer. The process is repeated to manufacture a second subplot of unlubricated blend. The final blend is then encapsulated into Swedish opaque, size 1 gelatin capsules using an automated capsule filling machine. The capsules are then weight sorted through an automatic weight sorter.

[00118] The 50 and 60 mg capsules (Tables 5 and 6) are prepared in a similar fashion as the 25 and 100 mg capsules.

[00119] In another aspect, the disclosure relates to a pharmaceutical composition comprising a compound of Formula IA or IB and a pharmaceutically acceptable carrier admixed with less than 100 ppm of 6,7-dimethoxy-quinoline-4-ol. 6,7-dimethoxy-quinoline-

4-ol, the structure of which is , can be used as reagent c(1) to make chloride f(1) and is a byproduct that may form during the synthesis of Compound IA or IB. Minimizing the concentration of contaminants or byproducts such as 6,7-dimethoxy-quinoline-4-ol in pharmaceutical compositions destined for human administration is desirable.

[00120] In one embodiment, the pharmaceutical composition as defined in any of the previous embodiments (for example, the pharmaceutical composition of Tables 1, 2, 2A, 3, 4, 5, and 6) is admixed with 100 ppm 6,7-dimethoxy-quinoline-4-ol.

[00121] In another embodiment, the pharmaceutical composition as defined in any of the previous embodiments is admixed with 50 ppm 6,7-dimethoxy-quinoline-4-ol.

[00122] In another embodiment, the pharmaceutical composition as defined in any of the previous embodiments is admixed with 25 ppm 6,7-dimethoxy-quinoline-4-ol.

[00123] In another embodiment, the pharmaceutical composition as defined in any of the previous embodiments is admixed with 15 ppm 6,7-dimethoxy-quinoline-4-ol.

[00124] In another embodiment, the pharmaceutical composition as defined in any of the previous embodiments is admixed with 10 ppm 6,7-dimethoxy-quinoline-4-ol.

[00125] In another embodiment, the pharmaceutical composition as defined in any of the previous embodiments is admixed with 5 ppm 6,7-dimethoxy-quinoline-4-ol.

[00126] In another embodiment, the pharmaceutical composition as defined in any of the previous embodiments is admixed with 2.5 ppm 6,7-dimethoxy-quinoline-4-ol.

[00127] In another embodiment, the disclosure relates to a pharmaceutical composition comprising Compound IA or IB; one or more fillers; one or more disintegrants; one or more glidants; and one or more lubricants admixed with 100 ppm or less 6,7-dimethoxy-quinoline-4-ol.

[00128] In this embodiment, the filler comprises microcrystalline cellulose.

[00129] In this embodiment, the disintegrant comprises croscarmellose sodium.

[00130] In this embodiment, the disintegrant comprises croscarmellose sodium and sodium starch glycolate.

[00131] In this embodiment, the glidant comprises fumed silica.

[00132] In this embodiment, the lubricant comprises stearic acid.

[00133] In another embodiment, the disclosure relates to a pharmaceutical composition comprising Compound IA or IB; microcrystalline cellulose; croscarmellose sodium; fumed silica; and stearic acid; admixed with 100 ppm or less of 6,7-dimethoxy-quinoline-4-ol. In one embodiment of this embodiment, the composition is admixed with 50 ppm or less of 6,7-dimethoxy-quinoline-4-ol. In another embodiment of this embodiment, the composition is admixed with 25 ppm or less of 6,7-dimethoxy-quinoline-4-ol. In another embodiment of this embodiment, the composition is admixed with 10 ppm or less of 6,7-dimethoxy-quinoline-4-ol. In another embodiment of this embodiment, the composition is admixed with

5 ppm or less of 6,7-dimethoxy-quinoline-4-ol. In another embodiment of this embodiment, the composition is admixed with 2.5 ppm or less of 6,7-dimethoxy-quinoline-4-ol.

[00134] In another embodiment, the disclosure relates to a pharmaceutical composition comprising Compound IA or IB; microcrystalline cellulose; anhydrous lactose; hydroxypropyl cellulose; a surfactant; croscarmellose sodium; silicon dioxide; and magnesium stearate; admixed with 100 ppm or less of 6,7-dimethoxy-quinoline-4-ol. In one embodiment of this embodiment, the composition is admixed with 50 ppm or less of 6,7-dimethoxy-quinoline-4-ol. In another embodiment of this embodiment, the composition is admixed with 25 ppm or less of 6,7-dimethoxy-quinoline-4-ol. In another embodiment of this embodiment, the composition is admixed with 10 ppm or less of 6,7-dimethoxy-quinoline-4-ol. In another embodiment of this embodiment, the composition is admixed with 5 ppm or less of 6,7-dimethoxy-quinoline-4-ol. In another embodiment of this embodiment, the composition is admixed with 2.5 ppm or less of 6,7-dimethoxy-quinoline-4-ol.

[00135] In another embodiment, the disclosure relates to a pharmaceutical composition comprising Compound IA or IB; microcrystalline cellulose; croscarmellose sodium; fumed silica; and stearic acid; ; admixed with 100 ppm or less of 6,7-dimethoxy-quinoline-4-ol. In one embodiment of this embodiment, the composition is admixed with 50 ppm or less of 6,7-dimethoxy-quinoline-4-ol. In another embodiment of this embodiment, the composition is admixed with 25 ppm or less of 6,7-dimethoxy-quinoline-4-ol. In another embodiment of this embodiment, the composition is admixed with 10 ppm or less of 6,7-dimethoxy-quinoline-4-ol. In another embodiment of this embodiment, the composition is admixed with 5 ppm or less of 6,7-dimethoxy-quinoline-4-ol. In another embodiment of this embodiment, the composition is admixed with 2.5 ppm or less of 6,7-dimethoxy-quinoline-4-ol.

EXAMPLES


[00136] The invention is illustrated further by the following examples in Scheme 1 and the description thereof, which are not to be construed as limiting the invention in scope or spirit to the specific procedures described in them. Those having skill in the art will recognize that the starting materials may be varied and additional steps employed to produce compounds encompassed by the invention, as demonstrated by the following examples. Those skilled in the art will also recognize that it may be necessary to utilize different solvents or reagents to achieve some of the above transformations. Unless otherwise specified, all reagents and solvents are of standard commercial grade and are used without further purification. The

appropriate atmosphere to run the reaction under, for example, air, nitrogen, hydrogen, argon, and the like, will be apparent to those skilled in the art.

Preparation of N-(4-([6,7-bis(methoxy)quinolin-4-yl]oxy)phenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide and the (L)-malate salt thereof.

[00137] A synthetic route that can be used for the preparation of N-(4-([6,7-bis(methoxy)quinolin-4-yl]oxy)phenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide and the (L)-malate salt thereof is depicted in Figure 1:

Figure 1



Preparation of 4-Chloro-6,7-dimethoxy-quinoline

[00138] A reactor was charged sequentially with 6,7-dimethoxy-quinoline-4-ol (47.0 kg) and acetonitrile (318.8 kg). The resulting mixture was heated to approximately 60 °C, and phosphorus oxychloride (POCl₃, 130.6 kg) was added. After the addition of POCl₃, the temperature of the reaction mixture was raised to approximately 77 °C. The reaction was deemed complete (approximately 13 hours) when less than 3% of the starting material remained (in-process high-performance liquid chromatography [HPLC] analysis). The reaction mixture was cooled to approximately 2 – 7 °C and then quenched into a chilled solution of dichloromethane (DCM, 482.8 kg), 26 % NH₄OH (251.3 kg), and water (900 L). The resulting mixture was warmed to approximately 20 – 25 °C, and phases were separated.

The organic phase was filtered through a bed of AW hyflo super-cel NF (Celite; 5.4 kg) and the filter bed was washed with DCM (118.9 kg). The combined organic phase was washed with brine (282.9 kg) and mixed with water (120 L). The phases were separated and the organic phase was concentrated by vacuum distillation with the removal of solvent (approximately 95 L residual volume). DCM (686.5 kg) was charged to the reactor containing organic phase and concentrated by vacuum distillation with the removal of solvent (approximately 90 L residual volume). Methyl t-butyl ether (MTBE, 226.0 kg) was then charged and the temperature of the mixture was adjusted to -20 to -25 °C and held for 2.5 hours resulting in solid precipitate which was then filtered and washed with n-heptane (92.0 kg), and dried on a filter at approximately 25 °C under nitrogen to afford the title compound. (35.6 kg).

Preparation of 4-(6,7-dimethoxy-quinoline-4-yloxy)-phenylamine

[00139] 4-Aminophenol (24.4 kg) dissolved in N,N-dimethylacetamide (DMA, 184.3 kg) was charged to a reactor containing 4-chloro-6,7-dimethoxyquinoline (35.3 kg), sodium t-butoxide (21.4 kg) and DMA (167.2 kg) at 20-25 °C. This mixture was then heated to 100-105 °C for approximately 13 hours. After the reaction was deemed complete as determined using in-process HPLC analysis (<2% starting material remaining), the reactor contents were cooled at 15 to 20 °C and water (pre-cooled, 2 to 7 °C, 587 L) charged at a rate to maintain 15 to 30 °C temperature. The resulting solid precipitate was filtered, washed with a mixture of water (47 L) and DMA (89.1 kg) and finally with water (214 L). The filter cake was then dried at approximately 25 °C on filter to yield crude 4-(6,7-dimethoxy-quinoline-4-yloxy)-phenylamine (59.4 kg wet, 41.6 kg dry calculated based on LOD). Crude 4-(6,7-dimethoxy-quinoline-4-yloxy)-phenylamine was refluxed (approximately 75 °C) in a mixture of tetrahydrofuran (THF, 211.4 kg) and DMA (108.8 kg) for approximately 1 hour and then cooled to 0-5 °C and aged for approximately 1 h after which time the solid was filtered, washed with THF (147.6 kg) and dried on a filter under vacuum at approximately 25 °C to yield 4-(6,7-dimethoxy-quinoline-4-yloxy)-phenylamine (34.0 kg).

Alternative Preparation of 4-(6,7-dimethoxy-quinoline-4-yloxy)-phenylamine

[00140] 4-chloro-6,7-dimethoxyquinoline (34.8 kg) and 4-aminophenol (30.8 kg) and sodium tert pentoxide (1.8 equivalents) 88.7 kg, 35 wt percent in THF) were charged to a reactor, followed by N,N-dimethylacetamide (DMA, 293.3 kg). This mixture was then heated to 105-115 °C for approximately 9 hours. After the reaction was deemed complete as

determined using in-process HPLC analysis (<2% starting material remaining), the reactor contents were cooled at 15 to 25 °C and water (315 kg) was added over a two hour period while maintaining the temperature between 20 and 30 °C. The reaction mixture was then agitated for an additional hour at 20 to 5 °C. The crude product was collected by filtration and washed with a mixture of 88 kg water and 82.1 kg DMA, followed by 175 kg water. The product was dried on a filter drier for 53 hours. The LOD showed less than 1% w/w.

[00141] In an alternative procedure, 1.6 equivalents of sodium tert-pentoxide were used and the reaction temperature was increased from 110-120 °C. In addition, the cool down temperature was increased to 35-40 °C and the starting temperature of the water addition was adjusted to 35-40 °C, with an allowed exotherm to 45 °C.

Preparation of 1-(4-fluoro-phenylcarbamoyl)-cyclopropanecarboxylic acid

[00142] Triethylamine (19.5 kg) was added to a cooled (approximately 5 °C) solution of cyclopropane-1,1-dicarboxylic acid (24.7 kg) in THF (89.6 kg) at a rate such that the batch temperature did not exceed 5 °C. The solution was stirred for approximately 1.3 hours, and then thionyl chloride (23.1 kg) was added, keeping the batch temperature below 10 °C. When the addition was complete, the solution was stirred for approximately 4 h keeping temperature below 10 °C. A solution of 4-fluoroaniline (18.0 kg) in THF (33.1 kg) was then added at a rate such that the batch temperature did not exceed 10 °C. The mixture was stirred for approximately 10 hours after which the reaction was deemed complete. The reaction mixture was then diluted with isopropyl acetate (218.1 kg). This solution was washed sequentially with aqueous sodium hydroxide (10.4 kg, 50 % dissolved in 119 L of water) further diluted with water (415 L), then with water (100 L) and finally with aqueous sodium chloride (20.0 kg dissolved in 100 L of water). The organic solution was concentrated by vacuum distillation (100 L residual volume) below 40 °C followed by the addition of n-heptane (171.4 kg), which resulted in the precipitation of solid. The solid was recovered by filtration and washed with n-Heptane (102.4 kg) resulting in wet crude, 1-(4-fluoro-phenylcarbamoyl)-cyclopropanecarboxylic acid (29.0 kg). The crude, 1-(4-fluoro-phenylcarbamoyl)-cyclopropanecarboxylic acid was dissolved in methanol (139.7 kg) at approximately 25 °C followed by the addition of water (320 L) resulting in slurry which was recovered by filtration, washed sequentially with water (20 L) and n-heptane (103.1 kg) and then dried on the filter at approximately 25 °C under nitrogen to afford the title compound (25.4 kg).

Preparation of 1-(4-Fluoro-phenylcarbamoyl)-cyclopropanecarbonyl chloride

[00143] Oxalyl chloride (12.6 kg) was added to a solution of 1-(4-fluoro-phenylcarbamoyl)-cyclopropanecarboxylic acid (22.8 kg) in a mixture of THF (96.1 kg) and N, N-dimethylformamide (DMF; 0.23 kg) at a rate such that the batch temperature did not exceed 25 °C. This solution was used in the next step without further processing.

Alternative Preparation of 1-(4-Fluoro-phenylcarbamoyl)-cyclopropanecarbonyl chloride

[00144] A reactor was charged with 1-(4-fluoro-phenylcarbamoyl)-cyclopropanecarboxylic acid (35 kg), 344 g DMF, and 175kg THF. The reaction mixture was adjusted to 12-17 °C and then to the reaction mixture was charged 19.9 kg of oxalyl chloride over a period of 1 hour. The reaction mixture was left stirring at 12-17 °C for 3 to 8 hours. This solution was used in the next step without further processing.

Preparation of cyclopropane-1,1-dicarboxylic acid [4-(6,7-dimethoxy-quinoline-4-yloxy)-phenyl]-amide (4-fluoro-phenyl)-amide (Compound 1A)

[00145] The solution from the previous step containing 1-(4-fluoro-phenylcarbamoyl)-cyclopropanecarbonyl chloride was added to a mixture of compound 4-(6,7-dimethoxy-quinoline-4-yloxy)-phenylamine (23.5 kg) and potassium carbonate (31.9 kg) in THF (245.7 kg) and water (116 L) at a rate such that the batch temperature did not exceed 30 °C. When the reaction was complete (in approximately 20 minutes), water (653 L) was added. The mixture was stirred at 20-25 °C for approximately 10 hours, which resulted in the precipitation of the product. The product was recovered by filtration, washed with a pre-made solution of THF (68.6 kg) and water (256 L), and dried first on a filter under nitrogen at approximately 25 °C and then at approximately 45 °C under vacuum to afford the title compound (41.0 kg, 38.1 kg, calculated based on LOD).

Alternative Preparation of cyclopropane-1,1-dicarboxylic acid [4-(6,7-dimethoxy-quinoline-4-yloxy)-phenyl]-amide (4-fluoro-phenyl)-amide

[00146] A reactor was charged with 4-(6,7-dimethoxy-quinoline-4-yloxy)-phenylamine (35.7 kg, 1 equivalent), followed by 412.9 kg THF. To the reaction mixture was charged a solution of 48.3 K₂CO₃ in 169 kg water. The acid chloride solution of described in the Alternative Preparation of 1-(4-Fluoro-phenylcarbamoyl)-cyclopropanecarbonyl chloride above was transferred to the reactor containing 4-(6,7-dimethoxy-quinoline-4-yloxy)-

phenylamine while maintaining the temperature between 20-30 °C over a minimum of two hours. The reaction mixture was stirred at 20-25 °C for a minimum of three hours. The reaction temperature was then adjusted to 30-25 °C and the mixture was agitated. The agitation was stopped and the phases of the mixture were allowed to separate. The lower aqueous phase was removed and discarded. To the remaining upper organic phase was added 804 kg water. The reaction was left stirring at 15-25 °C for a minimum of 16 hours.

[00147] The product precipitated. The product was filtered and washed with a mixture of 179 kg water and 157.9 THF in two portions. The crude product was dried under a vacuum for at least two hours. The dried product was then taken up in 285.1 kg THF. The resulting suspension was transferred to reaction vessel and agitated until the suspension became a clear (dissolved) solution, which required heating to 30-35 °C for approximately 30 minutes. 456 kg water was then added to the solution, as well as 20 kg SDAG-1 ethanol (ethanol denatured with methanol over two hours). The mixture was agitated at 15-25 °C for at least 16 hours. The product was filtered and washed with a mixture of 143 kg water and 126.7 THF in two portions. The product was dried at a maximum temperature set point of 40 °C.

[00148] In an alternative procedure, the reaction temperature during acid chloride formation was adjusted to 10-15 °C. The recrystallization temperature was changed from 15-25 °C to 45-50 °C for 1 hour and then cooled to 15-25 °C over 2 hours.

Preparation of cyclopropane-1,1-dicarboxylic acid [4-(6,7-dimethoxy-quinoline-4-yloxy)-phenyl]-amide (4-fluoro-phenyl)-amide, (L) malate salt (Compound IB)

[00149] Cyclopropane-1,1-dicarboxylic acid [4-(6,7-dimethoxy-quinoline-4-yloxy)-phenyl]-amide (4-fluoro-phenyl)-amide (1-5; 13.3 kg), L-malic acid (4.96 kg), methyl ethyl ketone (MEK; 188.6 kg) and water (37.3 kg) were charged to a reactor and the mixture was heated to reflux (approximately 74 °C) for approximately 2 hours. The reactor temperature was reduced to 50 to 55 °C and the reactor contents were filtered. These sequential steps described above were repeated two more times starting with similar amounts of 1-5 (13.3 kg), L-Malic acid (4.96 kg), MEK (198.6 kg) and water (37.2 kg). The combined filtrate was azeotropically dried at atmospheric pressure using MEK (1133.2 kg) (approximate residual volume 711 L; KF ≤ 0.5 % w/w) at approximately 74 °C. The temperature of the reactor contents was reduced to 20 to 25 °C and held for approximately 4 hours resulting in solid precipitate which was filtered, washed with MEK (448 kg) and dried under vacuum at 50 °C to afford the title compound (45.5 kg).

Alternative Preparation of cyclopropane-1,1-dicarboxylic acid [4-(6,7-dimethoxy-quinoline-4-yloxy)-phenyl]-amide (4-fluoro-phenyl)-amide, (L) malate salt

[00150] Cyclopropane-1,1-dicarboxylic acid [4-(6,7-dimethoxy-quinoline-4-yloxy)-phenyl]-amide (4-fluoro-phenyl)-amide (47.9 kg), L-malic acid (17.2), 658.2 kg methyl ethyl ketone, and 129.1 kg water (37.3 kg) were charged to a reactor and the mixture was heated 50-55 °C for approximately 1-3 hours, and then at 55-60 °C for an additional 4-5 hours. The mixture was clarified by filtration through a 1 µm cartridge. The reactor temperature was adjusted to 20-25 °C and vacuum distilled with a vacuum at 150-200 mm Hg with a maximum jacket temperature of 55 °C to the volume range of 558-731 L.

[00151] The vacuum distillation was performed two more times with the charge of 380 kg and 380.2 kg methyl ethyl ketone, respectively. After the third distillation, the volume of the batch was adjusted to 18 w/w of Cyclopropane-1,1-dicarboxylic acid [4-(6,7-dimethoxy-quinoline-4-yloxy)-phenyl]-amide (4-fluoro-phenyl)-amide by charging 159.9 kg methyl ethyl ketone to give a total volume of 880L. An additional vacuum distillation was carried out by adjusting 245.7 methyl ethyl ketone. The reaction mixture was left with moderate agitation at 20-25 °C for at least 24 hours. The product was filtered and washed with 415.1 kg methyl ethyl ketone in three portions. The product was dried under a vacuum with the jacket temperature set point at 45 °C.

[00152] In an alternative procedure, the order of addition was changed so that a solution of 17.7 kg L-malic acid dissolved in 129.9 kg water was added to Cyclopropane-1,1-dicarboxylic acid [4-(6,7-dimethoxy-quinoline-4-yloxy)-phenyl]-amide (4-fluoro-phenyl)-amide (48.7 kg) in methyl ethyl ketone (673.3 kg).

Preparation of Compound IB, Form N-1

[00153] A solution was prepared by adding tetrahydrofuran (12 mL/g-bulk-LR (limiting reagent); 1.20 L) and N-(4-{{6,7-bis(methyloxy)-quinolin-4-yl}oxy}phenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide, (100 g; 1.00 equiv; 100.00 g) and (L)-malic acid (1.2 equiv (molar); 32.08 g) to a 1 L reactor. Water (0.5317 mL/g-bulk-LR; 53.17 mL) was added and the solution was heated to 60 °C and maintained at that temperature for one hour until the solids were fully dissolved. The solution was passed through a Polish Filter.

[00154] At 60 °C, acetonitrile (12 mL/g-bulk-LR; 1.20 L) was added over a period of 8 hours. The solution was held at 60 °C for 10 hours. The solution was then cooled to 20 °C and held for 1 hour. The solids were filtered and washed with acetonitrile (12 mL/g-bulk-LR;

1.20 L). The solids were dried at 60 °C (25 mm Hg) for 6 hours to afford compound (I), Form N-1 (108 g; 0.85 equivalent; 108.00 g; 85.22% yield) as a white crystalline solid.

Alternate Preparation of Compound IB, Form N-1

[00155] A solution was prepared with 190 mL tetrahydrofuran (110 mL), methyl isobutyl ketone, and 29 mL water. Next, 20 mL of this solution were transferred into an amber bottle, and then saturated by adding N-(4-[[6,7-bis(methyloxy)-quinolin-4-yl]oxy}phenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide, (L)-malate until a thick slurry formed, and aging for at least 2 hours with stirring at room temperature. The solids were removed by filtration through a Buchner funnel, rendering a clear saturated solution.

[00156] Separately, a powder blend was made with known amounts of two batches of compound IB: (1) 300 mg of batch 1, which contained approximately 41% compound IB, Form N-1 and 59% compound IB, Form N-2 by Raman spectroscopy analysis, and (2) 200 mg of batch 2, which had a XPRD pattern similar to compound IB, Form N-2.

[00157] The compound IB, Form N-1 and compound (I), Form N-2 powder blend was added into the saturated solution, and the slurry was aged under magnetic stirring at room temperature for 25 days. The slurry was then sampled and filtered through a Buchner funnel to obtain 162 mg of wet cake. The wet cake was dried in a vacuum oven at 45 °C to afford 128 mg of crystalline compound IB in the N-1 form.

Preparation of Crystalline Compound IB, Form N-2

Preparation of Crystalline Compound IB, Form N-2 Seed Crystals

[00158] A solution was prepared by combining 20 ml of acetone and 300 mg of compound IA (N-(4-[[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide) in a 25ml screw capped vial. Next, 0.758ml of a 0.79M (L)-malic acid stock solution was added to the vial with magnetic stirring. The solution was then left stirring for 24 hours at ambient temperature. The sample was then suction filtered with 0.45µm PTFE filter cartridge and dried *in vacuo* at ambient temperature overnight.

Preparation of Crystalline Compound IB, Form N-2.

[00159] To a reactor were added N-(4-[[6,7-bis(methyloxy)-quinolin-4-yl]oxy}phenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (48 g; 1.00 equiv; 48.00 g) and tetrahydrofuran (16.5 mL/g-bulk-LR; 792.00 mL). The water content was adjusted to 1 wt% water. The solution was heated to 60 °C. Once dissolved, the solution was passed through a polish filter to provide the first solution.

[00160] In a separate reactor, (L)-malic acid (1.2 equiv (molar); 15.40 g) was dissolved into methyl isobutyl ketone (10 mL/g-bulk-LR; 480.00 mL) and tetrahydrofuran (1 mL/g-bulk-LR; 48.00 mL). Next, 50 mL of the (L)-malic acid solution was added to the first solution at 50 °C. Seed crystals were added (1%, 480 mg) and the malic acid solution was added at 50 °C dropwise via an addition funnel (1.3 mL/min over 3 hours). The slurry was held at 50 °C for 18 hours and then was cooled to 25 °C over 30 minutes. The solids were filtered, and washed with 20% tetrahydrofuran/methyl isobutyl ketone (10V, 480 mL). The solids were dried under vacuum at 60 °C for 5 hours to afford compound IB (55.7 g; 0.92 equivalent; 55.70 g; 91.56% yield) as an off-white crystalline solid.

Stability Studies of Pharmaceutical Compositions

[00161] The pharmaceutical capsule compositions of Tables 3 and 4 were prepared by mixing the ingredients according to processes known in the art.

Table 3

Ingredient	mg/unit dose
Compound IB (10 % drug load as Compound IA)	25
Microcrystalline Cellulose	196.75
Croscarmellose sodium	12.5
Sodium starch glycolate	12.5
Fumed Silica	0.75
Stearic acid	2.5
Total Fill Weight	250

Table 4

Ingredient	mg/unit dose
Compound IB (50 % drug load as Compound IA)	100
Silicified Microcrystalline Cellulose	75.40
Croscarmellose sodium	10.00
Sodium Starch Glycolate	10.00
Fumed silica	0.6
Stearic Acid	4.0
Total Fill Weight	200

[00162] The capsule compositions were subjected to stability studies to monitor the formation of 6,7-dimethoxy-quinoline-4-ol at various temperatures and relative humidities over time. The results are summarized in Tables 7A and 7B and Tables 8A and 8B.

Table 7A: Stability of 25 Mg Capsules (Table 3)

	Conditions	Bottle A1	Bottle A2	Bottle A3	Bottle A4
		PPM of 6,7-dimethoxy-quinoline-4-ol			
Initial T=0	25 °C/60% RH	2	2	3	3
1 Month	25 °C/60% RH	3	4	5	NA
	30 °C/75% RH	4	4	NA	NA
	40 °C/75% RH	9	9	10	NA
3 Months	25 °C/60% RH	5	5	7	NA
	30 °C/75% RH	7	6	NA	NA
	40 °C/75% RH	22	23	24	NA
6 Months	25 °C/60% RH	6	6	7	7 (3M in blister)
	30 °C/75% RH	9	9	NA	NA
	40 °C/75% RH	40	44	43	27 (3M in blister)
9 Months	25 °C/60% RH	7	7	9	8 (6M in blister)
	30 °C/75% RH	13	12	NA	NA
	40 °C/75% RH	NA	NA	68	60 (6M in blister)

M=Months; NA=Not Applicable; RH= Relative Humidity; PPM= Parts per Million. A portion of Bottle A4 was repackaged in a blister pack after being stored in bottles for 3 months.

Table 7B: Stability of 25 Mg Capsules (Table 3)

	Conditions	Bottle B1	Bottle B	Bottle B3	Bottle B4
		PPM of 6,7-dimethoxy-quinoline-4-ol			
Initial T=0	25 °C/60% RH	3	1	2	2
1 Month	25 °C/60% RH	<2	<2	<2	NA
	30 °C/75% RH	<2	<2	NA	NA
	40 °C/75% RH	2	<2	<2	NA
3 Months	25 °C/60% RH	2	<2	<2	NA
	30 °C/75% RH	2	<2	NA	NA
	40 °C/75% RH	3	<2	<2	NA
6 Months	25 °C/60% RH	<2	<2	<2	<2 (3M in blister)
	30 °C/75% RH	2	<2	NA	NA
	40 °C/75% RH	4	<2	3	3 (3M in blister)
9 Months	25 °C/60% RH	<2	<2	<2	<2 (6M in blister)
	30 °C/75% RH	3	<2	NA	NA
	40 °C/75% RH	NA	NA	5	4 (6M in blister)

A portion of Bottle B4 was repackaged in a blister pack after being stored in bottles for 3 months.

Table 8A: Stability of 100 Mg Capsules (Table 4)

	Conditions	Bottle A1	Blister A2	Bottle A3
		PPM of 6,7-dimethoxy-quinoline-4-ol		
Initial T=0	25 °C/60% RH	4	4	6
1 Month	25 °C/60% RH	4	4	6
	30 °C/75% RH	4	NA	6
	40 °C/75% RH	6	6	9
3 Months	25 °C/60% RH	5	5	7
	30 °C/75% RH	6	NA	7
	40 °C/75% RH	10	10	12
6 Months	25 °C/60% RH	5	5	
	30 °C/75% RH	6	NA	
	40 °C/75% RH	11	17	

M=Months; NA=Not Applicable; RH= Relative Humidity; PPM= Parts per Million.

Table 8B: Stability of 100 Mg Capsules (Table 4)

	Conditions	Bottle B1	Blister B2	Bottle B3
		PPM of 6,7-dimethoxy-quinoline-4-ol		
Initial T=0	25 °C/60% RH	1	1	2
1 Month	25 °C/60% RH	<2	<2	<2
	30 °C/75% RH	<2	<2	2
	40 °C/75% RH	<2	<2	2
3 Months	25 °C/60% RH	<2	<2	<2
	30 °C/75% RH	<2	NA	<2
	40 °C/75% RH	<2	<2	2
6 Months	25 °C/60% RH	<2	<2	
	30 °C/75% RH	<2	NA	
	40 °C/75% RH	2	2	

M=Months; NA=Not Applicable; RH= Relative Humidity; PPM= Parts per Million.


[00163] The results summarized in Tables 7A and 7B and 8A and 8B indicate that formation of 6,7-dimethoxy-quinoline-4-ol was minimized to 50 ppm or less over time in the capsule formulations.

[00164] The foregoing disclosure has been described in some detail by way of illustration and example, for purposes of clarity and understanding. The invention has been described with reference to various specific and preferred embodiments and techniques. However, it should be understood that many variations and modifications can be made while remaining within the spirit and scope of the invention. It will be obvious to one of skill in the art that

changes and modifications can be practiced within the scope of the appended claims. Therefore, it is to be understood that the above description is intended to be illustrative and not restrictive. The scope of the invention should, therefore, be determined not with reference to the above description, but should instead be determined with reference to the following appended claims, along with the full scope of equivalents to which such claims are entitled. All references cited herein are incorporated by reference in their entirety.

CLAIMS

1. A process for preparing a compound of formula I:



or a pharmaceutically acceptable salt thereof, wherein:

X^1 is H, Br, Cl, or F;


X^2 is H, Br, Cl, or F;

$n1$ is 1-2; and


$n2$ is 1-2;

the process comprising:

contacting the compound of formula g(1) with reactant z(1) to yield the compound of formula I:



2. The process according to claim 1, wherein the compound of formula g(1) is made by reacting a compound of formula f(1) with reactant y(1) to yield the compound of formula g(1):




wherein LG represents a leaving group, and each X^2 and $n2$ are as defined in claim 1.

3. The process according to claim 2, wherein the compound of formula f(1) is made by converting compound e(1) to the compound of formula f(1):




wherein LG represents a leaving group.

4. The process according to claim 1, wherein reactant z(1) is made by reacting reactant z(1a) with a chlorinating agent to yield reactant z(1):



wherein X¹ is Br, Cl, or F; and n₁ is 1-2.

5. The process according to claim 1, wherein the compound of formula I is compound IA-1:




or a pharmaceutically acceptable salt thereof, wherein:


X¹ is H, Cl, Br, or F; and

X² is H, Cl, Br, or F.


6. The process according to claim 5, wherein the compound of formula f(1) is compound f(2):




reactant y(1) is reactant (y)(2):




wherein X^2 is hydrogen or fluoro; and
the compound of formula g(1) is of formula g(2):




7. The process according to claim 6, wherein the reaction employs a non-nucleophilic base.
8. The process according to claim 7, wherein the non-nucleophilic base is an alkali metal alkoxide; and the reaction is carried out in a polar solvent.
9. The process according to claim 8, wherein the alkali metal alkoxide is sodium tert-butoxide, and the polar solvent is DMA.
10. The process according to claim 8, wherein the alkali metal alkoxide is sodium tert-pentoxide, and the polar solvent is DMA.
11. The process according to claim 1, wherein the compound of formula g(1) is of compound g(3):




reactant z(1) is reactant z(2):



the compound of formula I is compound IA:




12. The process according to claim 11, wherein the reaction is carried out in the presence of an inorganic base.
13. The process according to claim 12, wherein the inorganic base is K_2CO_3 , and the solvent employed in this reaction is a combination of THF and H_2O .
14. A process for preparing compound IB:



41

IB


from compound IA:




comprising:

- (a) heating and agitating a mixture comprising compound IA and L-malic acid, methylethyl ketone, and water;
- (b) cooling the mixture;
- (c) vacuum distilling the mixture successively; and
- (d) isolating the compound of IB by filtration.

15. A process for preparing compound IB:



from compound IA:



comprising:

- (a) heating and agitating a mixture comprising compound IA and L-malic acid, methylethyl ketone, and water;
 - (b) cooling the mixture;
 - (c) seeding the mixture with the compound of IB;
 - (d) vacuum distilling the mixture; and
 - (e) isolating the compound of IB by filtration.
16. Compound IA or IB admixed with 100 ppm or less of 6,7-dimethoxy-quinoline-4-ol.
17. Compound IB admixed with 100 ppm or less of 6,7-dimethoxy-quinoline-4-ol.
18. Compound IB in the N-1 form admixed with 100 ppm or less of 6,7-dimethoxy-quinoline-4-ol.
19. Compound IB in the N-2 form admixed with 100 ppm or less of 6,7-dimethoxy-quinoline-4-ol.
20. Compound IB as a mixture of the N-1 form and N-2 form admixed with 100 ppm or less of 6,7-dimethoxy-quinoline-4-ol.
21. A pharmaceutical composition according to Table 1.

Table 1

Ingredient	% w/w
Compound I	31.68
Microcrystalline Cellulose (Avicel PH102)	38.85
Lactose anhydrous 60 M	19.42
Hydroxypropyl Cellulose, EXF	3.00
Croscarmellose Sodium	3.00
Total Intra-granular	95.95
Silicon dioxide, Colloidal PWD	0.30
Croscarmellose Sodium	3.00
Magnesium Stearate	0.75
Total	100.00

22. A pharmaceutical composition according to Table 2.

Table 2

Ingredient	(% w/w)
Compound I	25.0-33.3
Microcrystalline Cellulose, NF	q.s
Hydroxypropyl Cellulose, NF	3
Poloxamer, NF	0-3
Croscarmellose Sodium, NF	6.0
Colloidal Silicon Dioxide, NF	0.5
Magnesium Stearate, NF	0.5-1.0
Total	100

23. A pharmaceutical composition according to Table 2A.

Table 2A

Ingredient	% w/w
Compound IB (10 % drug load as Compound IA)	12.67
MCC	51.52
Lactose	25.76
Hydroxypropyl cellulose	3.0
Croscarmellose Sodium	6.0
Colloidal Silicon Dioxide	0.3
Magnesium Stearate	0.75
Total	100

24. A pharmaceutical composition according to Table 3.

Table 3

Ingredient	mg/unit dose
Compound IB (10 % drug load as Compound IA)	25
Silicified Microcrystalline Cellulose	196.75
Croscarmellose sodium	12.5
Sodium starch glycolate	12.5
Fumed Silica	0.75
Stearic acid	2.5
Total Fill Weight	250

25. A pharmaceutical composition according to Table 4.

Table 4

Ingredient	mg/unit dose
Compound IB (50 % drug load as Compound IA)	100
Silicified Microcrystalline Cellulose	75.40
Croscarmellose sodium	10.00
Sodium Starch Glycolate	10.00
Fumed silica	0.6
Stearic Acid	4.0
Total Fill Weight	200

26. A pharmaceutical composition according to Table 5.

Table 5

Ingredient	mg/unit dose
Compound IB (10 % drug load as Compound IA)	50mg
Microcrystalline Cellulose	63.35
Croscarmellose sodium	95.39
Sodium starch glycolate	9.05
Fumed Silica	9.05
Stearic acid	0.54
Total Fill Weight	3.62

27. A pharmaceutical composition according to Table 6.

Table 6

Ingredient	mg/unit dose
Compound IB	60mg
Microcrystalline Cellulose	73.95
Croscarmellose sodium	114.36
Sodium starch glycolate	10.85
Fumed Silica	10.85
Stearic acid	0.65
Total Fill Weight	4.34

28. The pharmaceutical composition of claims 20-27 admixed with 100 ppm or less of 6,7-dimethoxy-quinoline-4-ol.

29. The pharmaceutical composition of claims 20-27 admixed with 50 ppm or less of 6,7-dimethoxy-quinoline-4-ol.

30. The pharmaceutical composition of claims 20-27 admixed with 25 ppm or less of 6,7-dimethoxy-quinoline-4-ol.
31. The pharmaceutical composition of claims 20-27 admixed with 15 ppm or less of 6,7-dimethoxy-quinoline-4-ol.
32. The pharmaceutical composition of claims 20-27 admixed with 10 ppm or less of 6,7-dimethoxy-quinoline-4-ol.
33. The pharmaceutical composition of claims 20-27 admixed with 5 ppm or less of 6,7-dimethoxy-quinoline-4-ol.
34. The pharmaceutical composition of claims 20-27 admixed with 2.5 ppm or less of 6,7-dimethoxy-quinoline-4-ol.

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2012/024591

A. CLASSIFICATION OF SUBJECT MATTER

INV. C07D215/227 A61K31/47 A61P35/00
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPD-Internal, CHEM ABS Data, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2010/083414 A1 (EXELIXIS INC [US]; ST CLAIR BROWN ADRIAN [GB]; LAMB PETER [US]; GALLAG) 22 July 2010 (2010-07-22) cited in the application Scheme 1 on page24; claims;claim 11 -----	1-34

☐ Further documents are listed in the continuation of Box C.

☒ See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"Z" document member of the same patent family

Date of the actual completion of the international search

16 May 2012

Date of mailing of the international search report

25/05/2012

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3018

Authorized officer

Wolf, Claudia

INTERNATIONAL SEARCH REPORT

Information on patent family members


International application No

PCT/US2012/024591

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2010083414	A1	22-07-2010	
		AR 075025 A1	02-03-2011
		AU 2010204619 A1	01-09-2011
		CA 2758030 A1	22-07-2010
		CN 102388024 A	21-03-2012
		EP 2387563 A1	23-11-2011
		KR 20120013301 A	14-02-2012
		SG 173014 A1	29-08-2011
		TW 201036948 A	16-10-2010
		US 2012035212 A1	09-02-2012
		WO 2010083414 A1	22-07-2010

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

**DECLARATION (37 CFR 1.63) FOR UTILITY OR DESIGN APPLICATION USING AN
APPLICATION DATA SHEET (37 CFR 1.76)**

Title of Invention	Processes for Preparing Quinoline Compounds and Pharmaceutical Compositions Containing Such Compounds
As the below named inventor, I hereby declare that: This declaration is directed to: ☐ The attached application, or ☒ United States application or PCT International application number <u>15/348,716</u> filed on <u>November 10, 2016</u>. The above-identified application was made or authorized to be made by me. I believe that I am the original inventor or an original joint inventor of a claimed invention in the application. I hereby acknowledge that any willful false statement made in this declaration is punishable under 18 U.S.C. 1001 by fine or imprisonment of not more than five (5) years, or both. WARNING: Petitioner/applicant is cautioned to avoid submitting personal information in documents filed in a patent application that may contribute to identity theft. Personal information such as social security numbers, bank account numbers, or credit card numbers (other than a check or credit card authorization form PTO-2038 submitted for payment purposes) is never required by the USPTO to support a petition or an application. If this type of personal information is included in documents submitted to the USPTO, petitioners/applicants should consider redacting such personal information from the documents before submitting them to the USPTO. Petitioner/applicant is advised that the record of a patent application is available to the public after publication of the application (unless a non-publication request in compliance with 37 CFR 1.213(a) is made in the application) or issuance of a patent. Furthermore, the record from an abandoned application may also be available to the public if the application is referenced in a published application or an issued patent (see 37 CFR 1.14). Checks and credit card authorization forms PTO-2038 submitted for payment purposes are not retained in the application file and therefore are not publicly available.	
LEGAL NAME OF INVENTOR Inventor: <u>Jo Ann Wilson</u> Date (Optional): _____ Signature: 	
Note: An application data sheet (PTO/SB/14 or equivalent), including naming the entire inventive entity, must accompany this form or must have been previously filed. Use an additional PTO/AIA/01 form for each additional inventor.	

This collection of information is required by 35 U.S.C. 115 and 37 CFR 1.63. The information is required to obtain or retain a benefit by the public which is to file (and by the USPTO to process) an application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.11 and 1.14. This collection is estimated to take 1 minute to complete, including gathering, preparing, and submitting the completed application form to the USPTO. Time will vary depending upon the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, P.O. Box 1450, Alexandria, VA 22313-1450. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.

If you need assistance in completing the form, call 1-800-PTO-9199 and select option 2.

**DECLARATION (37 CFR 1.63) FOR UTILITY OR DESIGN APPLICATION USING AN
APPLICATION DATA SHEET (37 CFR 1.76)**

Title of Invention	Processes for Preparing Quinoline Compounds and Pharmaceutical Compositions Containing Such Compounds
As the below named inventor, I hereby declare that: This declaration is directed to: ☐ The attached application, or ☒ United States application or PCT international application number <u>16/151,653</u> filed on <u>October 4, 2018</u> The above-identified application was made or authorized to be made by me. I believe that I am the original inventor or an original joint inventor of a claimed invention in the application. I hereby acknowledge that any willful false statement made in this declaration is punishable under 18 U.S.C. 1001 by fine or imprisonment of not more than five (5) years, or both. WARNING: Petitioner/applicant is cautioned to avoid submitting personal information in documents filed in a patent application that may contribute to identity theft. Personal information such as social security numbers, bank account numbers, or credit card numbers (other than a check or credit card authorization form PTO-2038 submitted for payment purposes) is never required by the USPTO to support a petition or an application. If this type of personal information is included in documents submitted to the USPTO, petitioners/applicants should consider redacting such personal information from the documents before submitting them to the USPTO. Petitioner/applicant is advised that the record of a patent application is available to the public after publication of the application (unless a non-publication request in compliance with 37 CFR 1.213(a) is made in the application) or issuance of a patent. Furthermore, the record from an abandoned application may also be available to the public if the application is referenced in a published application or an issued patent (see 37 CFR 1.14). Checks and credit card authorization forms PTO-2038 submitted for payment purposes are not retained in the application file and therefore are not publicly available.	
LEGAL NAME OF INVENTOR Inventor: <u>Khalid Shah</u> Date (Optional): <u>14 June 2019</u> Signature: <u></u>	
Note: An application data sheet (PTO/SB/14 or equivalent), including naming the entire inventive entity, must accompany this form or must have been previously filed. Use an additional PTO/AIA/01 form for each additional inventor.	

This collection of information is required by 35 U.S.C. 115 and 37 CFR 1.63. The information is required to obtain or retain a benefit by the public which is to file (and by the USPTO to process) an application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.11 and 1.14. This collection is estimated to take 1 minute to complete, including gathering, preparing, and submitting the completed application form to the USPTO. Time will vary depending upon the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, P.O. Box 1450, Alexandria, VA 22313-1450. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.

If you need assistance in completing the form, call 1-800-PTO-9199 and select option 2.

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Inventors:	Jo Ann Wilson, et al.	Art Unit:	To Be Determined
Application No.:	To Be Determined	Examiner:	To Be Determined
Filing Date:	December 6, 2019	Conf. No.:	To Be Determined
Atty. Docket No.:	224990/10-039C4-US/459258		

Title: Processes for Preparing Quinoline Compounds and Pharmaceutical Compositions Containing Such Compounds

Commissioner for Patents
United States Patent and Trademark Office
P. O. Box 1450
Alexandria, VA 22313-1450

PRELIMINARY AMENDMENT UNDER 37 C.F.R. § 1.115

Dear Sir or Madam:

Prior to examination of the application, please amend the above-referenced application as follows:

Amendments to the specification begin on page 2.

Remarks begin on page 3.

Inventors: Jo Ann Wilson, et al.
Application No.: To Be Determined
Filing Date: December 6, 2019
Atty. Docket No.: 224990/10-039C4-US/459258

Art Unit: To Be Determined
Examiner: To Be Determined
Conf. No.: To Be Determined

AMENDMENT TO THE SPECIFICATION

On page 1 of the specification, after the title and header, please replace paragraph [0001] with the following paragraph:

[0001] This application is a divisional of United States Application Serial Number 16/151,653, filed October 4, 2018, which is a divisional of United States Application Serial Number 15/348,716, filed November 10, 2016, now United States Patent Number 10,123,999, which is a divisional of United States Application Serial Number 13/984,559, filed March 20, 2014, now United States Patent Number 9,717,720, which claims priority under 35 U.S.C. § 371 to Patent Cooperation Treaty Application Number PCT/US2012/024591, filed February 10, 2012, which claims the benefit of United States Provisional Application Number 61/441,520, filed February 10, 2011, and United States Provisional Application Number 61/441,527, filed February 10, 2011, the entire contents of which are incorporated herein by reference.

Inventors: Jo Ann Wilson, et al.
Application No.: To Be Determined
Filing Date: December 6, 2019
Atty. Docket No.: 224990/10-039C4-US/459258

Art Unit: To Be Determined
Examiner: To Be Determined
Conf. No.: To Be Determined

REMARKS

Amendments to the Specification

The Applicant has amended the application to state that this application is a divisional of United States Application Serial Number 16/151,653, filed October 4, 2018, which is a divisional of United States Application Serial Number 15/348,716, filed November 10, 2016, now United States Patent Number 10,123,999, which is a divisional of United States Application Serial Number 13/984,559, filed March 20, 2014, now United States Patent Number 9,717,720, which claims priority under 35 U.S.C. § 371 to Patent Cooperation Treaty Application Number PCT/US2012/024591, filed February 10, 2012, which claims the benefit of United States Provisional Application Number 61/441,520, filed February 10, 2011, and United States Provisional Application Number 61/441,527, filed February 10, 2011, the entire contents of which are incorporated herein by reference. It is believed that no new matter is added through the entry of these amendments.

Conclusion and Required Fees

It is believed that due with this submission is the Basic Utility Application Fee of \$300.00, the Utility Search Fee of \$660.00, and the Utility Examination Fee of \$760.00. The Applicant hereby authorizes the Patent Office to charge this total of \$1,720.00 to Deposit Account No. 503145, referencing Attorney Docket Number 224990/10-039C4-US/459258.

Respectfully submitted,

HONIGMAN LLP

Dated: December 6, 2019


Heidi M. Berven, Ph.D.
Reg. No. 48,951
650 Trade Centre Way, Suite 200
Kalamazoo, MI 49002-0402
Telephone: (269) 337-7710
Facsimile: (269) 337-7711