

Oncologist 2012;17:813-822. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/22643536>.

48. Wang Z, Lai ST, Xie L, et al. Metformin is associated with reduced risk of pancreatic cancer in patients with type 2 diabetes mellitus: A systematic review and meta-analysis. Diabetes Res Clin Pract 2014;106:19-26. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/24837144>.

49. Sadeghi N, Abbruzzese JL, Yeung SC, et al. Metformin use is associated with better survival of diabetic patients with pancreatic cancer. Clin Cancer Res 2012;18:2905-2912. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/22465831>.

50. Toriola AT, Stolzenberg-Solomon R, Dalidowicz L, et al. Diabetes and pancreatic cancer survival: a prospective cohort-based study. Br J Cancer 2014;111:181-185. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/24786605>.

51. Hruban RH, Canto MI, Goggins M, et al. Update on familial pancreatic cancer. Adv Surg 2010;44:293-311. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/20919528>.

52. Humphris JL, Johns AL, Simpson SH, et al. Clinical and pathologic features of familial pancreatic cancer. Cancer 2014. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/25313458>.

53. Lynch HT, Smyrk T, Kern SE, et al. Familial pancreatic cancer: a review. Semin Oncol 1996;23:251-275. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/8623061>.

54. Wang W, Chen S, Brune KA, et al. PancPRO: risk assessment for individuals with a family history of pancreatic cancer. J Clin Oncol 2007;25:1417-1422. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/17416862>.

55. Hemminki A, Markie D, Tomlinson I, et al. A serine/threonine kinase gene defective in Peutz-Jeghers syndrome. Nature 1998;391:184-187. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9428765>.

56. Jenne DE, Reimann H, Nezu J, et al. Peutz-Jeghers syndrome is caused by mutations in a novel serine threonine kinase. Nat Genet 1998;18:38-43. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/9425897>.

57. Korsse SE, Harinck F, van Lier MG, et al. Pancreatic cancer risk in Peutz-Jeghers syndrome patients: a large cohort study and implications for surveillance. J Med Genet 2013;50:59-64. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/23240097>.

58. Giardiello FM, Brensinger JD, Tersmette AC, et al. Very high risk of cancer in familial Peutz-Jeghers syndrome. Gastroenterology 2000;119:1447-1453. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/11113065>.

59. van Lier MG, Wagner A, Mathus-Vliegen EM, et al. High cancer risk in Peutz-Jeghers syndrome: a systematic review and surveillance recommendations. Am J Gastroenterol 2010;105:1258-1264; author reply 1265. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/20051941>.

60. Su GH, Hruban RH, Bansal RK, et al. Germline and somatic mutations of the STK11/LKB1 Peutz-Jeghers gene in pancreatic and biliary cancers. Am J Pathol 1999;154:1835-1840. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/10362809>.

61. Weiss FU. Pancreatic cancer risk in hereditary pancreatitis. Front Physiol 2014;5:70. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/24600409>.

62. LaRusch J, Solomon S, Whitcomb DC. Pancreatitis Overview. In: Pagon RA, Adam MP, Ardinger HH, et al., eds. GeneReviews(R). Seattle (WA): University of Washington, Seattle; 2014.

63. Howes N, Lerch MM, Greenhalf W, et al. Clinical and genetic characteristics of hereditary pancreatitis in Europe. *Clin Gastroenterol Hepatol* 2004;2:252-261. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15017610>.
64. Lowenfels AB, Maisonneuve P, DiMaggio EP, et al. Hereditary pancreatitis and the risk of pancreatic cancer. International Hereditary Pancreatitis Study Group. *J Natl Cancer Inst* 1997;89:442-446. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9091646>.
65. Rebours V, Levy P, Ruzsiewicz P. An overview of hereditary pancreatitis. *Dig Liver Dis* 2012;44:8-15. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21907651>.
66. Whelan AJ, Bartsch D, Goodfellow PJ. Brief report: a familial syndrome of pancreatic cancer and melanoma with a mutation in the CDKN2 tumor-suppressor gene. *N Engl J Med* 1995;333:975-977. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/7666917>.
67. de Snoo FA, Bishop DT, Bergman W, et al. Increased risk of cancer other than melanoma in CDKN2A founder mutation (p16-Leiden)-positive melanoma families. *Clin Cancer Res* 2008;14:7151-7157. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18981015>.
68. Vasen HF, Gruis NA, Frants RR, et al. Risk of developing pancreatic cancer in families with familial atypical multiple mole melanoma associated with a specific 19 deletion of p16 (p16-Leiden). *Int J Cancer* 2000;87:809-811. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10956390>.
69. Lynch HT, Brand RE, Hogg D, et al. Phenotypic variation in eight extended CDKN2A germline mutation familial atypical multiple mole melanoma-pancreatic carcinoma-prone families: the familial atypical mole melanoma-pancreatic carcinoma syndrome. *Cancer* 2002;94:84-96. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11815963>.
70. Ghiorzo P, Fornarini G, Sciallero S, et al. CDKN2A is the main susceptibility gene in Italian pancreatic cancer families. *J Med Genet* 2012;49:164-170. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22368299>.
71. Aaltonen LA, Salovaara R, Kristo P, et al. Incidence of hereditary nonpolyposis colorectal cancer and the feasibility of molecular screening for the disease. *N Engl J Med* 1998;338:1481-1487. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9593786>.
72. Lindor NM, Petersen GM, Spurdle AB, et al. Pancreatic Cancer and a Novel MSH2 Germline Alteration. *Pancreas* 2011;40:1138-1140. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21926548>.
73. Lynch HT, de la Chapelle A. Hereditary colorectal cancer. *N Engl J Med* 2003;348:919-932. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12621137>.
74. Hampel H, Frankel WL, Martin E, et al. Screening for the Lynch syndrome (hereditary nonpolyposis colorectal cancer). *N Engl J Med* 2005;352:1851-1860. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15872200>.
75. Hampel H, Frankel WL, Martin E, et al. Feasibility of screening for Lynch syndrome among patients with colorectal cancer. *J Clin Oncol* 2008;26:5783-5788. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18809606>.
76. Boland CR, Goel A. Microsatellite instability in colorectal cancer. *Gastroenterology* 2010;138:2073-2087 e2073. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20420947>.
77. Kastrinos F, Mukherjee B, Tayob N, et al. Risk of pancreatic cancer in families with Lynch syndrome. *Jama* 2009;302:1790-1795. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19861671>.
78. Win AK, Young JP, Lindor NM, et al. Colorectal and other cancer risks for carriers and noncarriers from families with a DNA mismatch repair gene mutation: a prospective cohort study. *J Clin Oncol*

2012;30:958-964. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/22331944>.

79. Cancer risks in BRCA2 mutation carriers. J Natl Cancer Inst 1999;91:1310-1316. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/10433620>.

80. Al-Sukhni W, Rothenmund H, Borgida AE, et al. Germline BRCA1 mutations predispose to pancreatic adenocarcinoma. Hum Genet 2008;124:271-278. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/18762988>.

81. Ferrone CR, Levine DA, Tang LH, et al. BRCA germline mutations in Jewish patients with pancreatic adenocarcinoma. J Clin Oncol 2009;27:433-438. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/19064968>.

82. Hahn SA, Greenhalf B, Ellis I, et al. BRCA2 germline mutations in familial pancreatic carcinoma. J Natl Cancer Inst 2003;95:214-221. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12569143>.

83. Iqbal J, Ragone A, Lubinski J, et al. The incidence of pancreatic cancer in BRCA1 and BRCA2 mutation carriers. Br J Cancer 2012;107:2005-2009. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/23099806>.

84. van Asperen CJ, Brohet RM, Meijers-Heijboer EJ, et al. Cancer risks in BRCA2 families: estimates for sites other than breast and ovary. J Med Genet 2005;42:711-719. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/16141007>.

85. Liede A, Karlan BY, Narod SA. Cancer risks for male carriers of germline mutations in BRCA1 or BRCA2: a review of the literature. J Clin Oncol 2004;22:735-742. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/14966099>.

86. Couch FJ, Johnson MR, Rabe K, et al. Germ line Fanconi anemia complementation group C mutations and pancreatic cancer. Cancer

Res 2005;65:383-386. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/15695377>.

87. Slater EP, Langer P, Niemczyk E, et al. PALB2 mutations in European familial pancreatic cancer families. Clin Genet 2010;78:490-494. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20412113>.

88. van der Heijden MS, Yeo CJ, Hruban RH, Kern SE. Fanconi anemia gene mutations in young-onset pancreatic cancer. Cancer Res 2003;63:2585-2588. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/12750283>.

89. Roberts NJ, Jiao Y, Yu J, et al. ATM mutations in patients with hereditary pancreatic cancer. Cancer Discov 2012;2:41-46. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/22585167>.

90. Klein AP, Brune KA, Petersen GM, et al. Prospective risk of pancreatic cancer in familial pancreatic cancer kindreds. Cancer Res 2004;64:2634-2638. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/15059921>.

91. Clores MJ, Thosani A, Buscaglia JM. Multidisciplinary diagnostic and therapeutic approaches to pancreatic cystic lesions. J Multidiscip Healthc 2014;7:81-91. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/24520195>.

92. Farrell JJ, Fernandez-del Castillo C. Pancreatic cystic neoplasms: management and unanswered questions. Gastroenterology 2013;144:1303-1315. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/23622140>.

93. Law JK, Hruban RH, Lennon AM. Management of pancreatic cysts: a multidisciplinary approach. Curr Opin Gastroenterol 2013;29:509-516. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23872487>.

94. Tanaka M, Fernandez-del Castillo C, Adsay V, et al. International consensus guidelines 2012 for the management of IPMN and MCN of

the pancreas. *Pancreatology* 2012;12:183-197. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/22687371>.

95. Del Chiaro M, Verbeke C, Salvia R, et al. European experts consensus statement on cystic tumours of the pancreas. *Dig Liver Dis* 2013;45:703-711. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/23415799>.

96. Canto MI, Goggins M, Hruban RH, et al. Screening for early pancreatic neoplasia in high-risk individuals: a prospective controlled study. *Clin Gastroenterol Hepatol* 2006;4:766-781; quiz 665. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16682259>.

97. Canto MI, Hruban RH, Fishman EK, et al. Frequent detection of pancreatic lesions in asymptomatic high-risk individuals. *Gastroenterology* 2012;142:796-804; quiz e714-795. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22245846>.

98. Al-Sukhni W, Borgida A, Rothenmund H, et al. Screening for pancreatic cancer in a high-risk cohort: an eight-year experience. *J Gastrointest Surg* 2012;16:771-783. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22127781>.

99. Poley JW, Kluijdt I, Gouma DJ, et al. The yield of first-time endoscopic ultrasonography in screening individuals at a high risk of developing pancreatic cancer. *Am J Gastroenterol* 2009;104:2175-2181. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19491823>.

100. Langer P, Kann PH, Fendrich V, et al. Five years of prospective screening of high-risk individuals from families with familial pancreatic cancer. *Gut* 2009;58:1410-1418. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19470496>.

101. Ding Z, Wu H, Zhang J, et al. MicroRNAs as novel biomarkers for pancreatic cancer diagnosis: a meta-analysis based on 18 articles. *Tumour Biol* 2014. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24880590>.

102. Kobayashi T, Nishiumi S, Ikeda A, et al. A novel serum metabolomics-based diagnostic approach to pancreatic cancer. *Cancer Epidemiol Biomarkers Prev* 2013;22:571-579. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23542803>.

103. Mayers JR, Wu C, Clish CB, et al. Elevation of circulating branched-chain amino acids is an early event in human pancreatic adenocarcinoma development. *Nat Med* 2014. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25261994>.

104. Schultz NA, Dehlendorff C, Jensen BV, et al. MicroRNA biomarkers in whole blood for detection of pancreatic cancer. *Jama* 2014;311:392-404. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24449318>.

105. Liggett T, Melnikov A, Yi QL, et al. Differential methylation of cell-free circulating DNA among patients with pancreatic cancer versus chronic pancreatitis. *Cancer* 2010;116:1674-1680. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20143430>.

106. O'Brien DP, Sandanayake NS, Jenkinson C, et al. Serum CA19-9 is significantly up-regulated up to 2 years prior to diagnosis with pancreatic cancer: implications for early disease detection. *Clin Cancer Res* 2014. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24938522>.

107. Canto MI, Harinck F, Hruban RH, et al. International Cancer of the Pancreas Screening (CAPS) Consortium summit on the management of patients with increased risk for familial pancreatic cancer. *Gut* 2013;62:339-347. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23135763>.

108. Callery MP, Chang KJ, Fishman EK, et al. Pretreatment assessment of resectable and borderline resectable pancreatic cancer: expert consensus statement. *Ann Surg Oncol* 2009;16:1727-1733. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19396496>.

109. Al-Hawary MM, Francis IR, Chari ST, et al. Pancreatic ductal adenocarcinoma radiology reporting template: consensus statement of

the society of abdominal radiology and the american pancreatic association. *Gastroenterology* 2014;146:291-304.e291. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24355035>.

110. Edge SB, Byrd DR, Compton CC, et al., eds. *AJCC Cancer Staging Manual* (ed 7th). New York: Springer; 2010.

111. Bilimoria KY, Bentrem DJ, Ko CY, et al. Validation of the 6th edition AJCC Pancreatic Cancer Staging System: report from the National Cancer Database. *Cancer* 2007;110:738-744. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17580363>.

112. Wong JC, Lu DSK. Staging of pancreatic adenocarcinoma by imaging studies. *Clin Gastroenterol Hepatol* 2008;6:1301-1308. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18948228>.

113. Fuhrman GM, Charnsangavej C, Abbruzzese JL, et al. Thin-section contrast-enhanced computed tomography accurately predicts the resectability of malignant pancreatic neoplasms. *Am J Surg* 1994;167:104-111. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/7906097>.

114. Horton KM, Fishman EK. Adenocarcinoma of the pancreas: CT imaging. *Radiol Clin North Am* 2002;40:1263-1272. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12479710>.

115. House MG, Yeo CJ, Cameron JL, et al. Predicting resectability of periampullary cancer with three-dimensional computed tomography. *J Gastrointest Surg* 2004;8:280-288. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15019924>.

116. Klaus M, Schobinger M, Wolf I, et al. Value of three-dimensional reconstructions in pancreatic carcinoma using multidetector CT: initial results. *World J Gastroenterol* 2009;15:5827-5832. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19998504>.

117. McNulty NJ, Francis IR, Platt JF, et al. Multi-detector row helical CT of the pancreas: effect of contrast-enhanced multiphasic imaging on

enhancement of the pancreas, peripancreatic vasculature, and pancreatic adenocarcinoma. *Radiology* 2001;220:97-102. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11425979>.

118. Walters DM, Lapar DJ, de Lange EE, et al. Pancreas-protocol imaging at a high-volume center leads to improved preoperative staging of pancreatic ductal adenocarcinoma. *Ann Surg Oncol* 2011;18:2764-2771. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21484522>.

119. Schima W, Ba-Ssalamah A, Goetzinger P, et al. State-of-the-art magnetic resonance imaging of pancreatic cancer. *Top Magn Reson Imaging* 2007;18:421-429. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18303400>.

120. Vachiranubhap B, Kim YH, Balci NC, Semelka RC. Magnetic resonance imaging of adenocarcinoma of the pancreas. *Top Magn Reson Imaging* 2009;20:3-9. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19687720>.

121. Agarwal B, Abu-Hamda E, Molke KL, et al. Endoscopic ultrasound-guided fine needle aspiration and multidetector spiral CT in the diagnosis of pancreatic cancer. *Am J Gastroenterol* 2004;99:844-850. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15128348>.

122. Deerenberg EB, Poley JW, Hermans JJ, et al. Role of endoscopic ultrasonography in patients suspected of pancreatic cancer with negative helical MDCT scan. *Dig Surg* 2011;28:398-403. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22188923>.

123. Nawaz H, Fan CY, Kloke J, et al. Performance characteristics of endoscopic ultrasound in the staging of pancreatic cancer: a meta-analysis. *JOP* 2013;14:484-497. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24018593>.

124. Wang W, Shpaner A, Krishna SG, et al. Use of EUS-FNA in diagnosing pancreatic neoplasm without a definitive mass on CT. *Gastrointest Endosc* 2013;78:73-80. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23523302>.

125. Rosch T, Braig C, Gain T, et al. Staging of pancreatic and ampullary carcinoma by endoscopic ultrasonography. Comparison with conventional sonography, computed tomography, and angiography. *Gastroenterology* 1992;102:188-199. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/1727753>.

126. Varadarajulu S, Wallace MB. Applications of endoscopic ultrasonography in pancreatic cancer. *Cancer Control* 2004;11:15-22. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14749619>.

127. Buchs NC, Chilcott M, Poletti PA, et al. Vascular invasion in pancreatic cancer: Imaging modalities, preoperative diagnosis and surgical management. *World J Gastroenterol* 2010;16:818-831. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20143460>.

128. Inoue K, Ohuchida J, Ohtsuka T, et al. Severe localized stenosis and marked dilatation of the main pancreatic duct are indicators of pancreatic cancer instead of chronic pancreatitis on endoscopic retrograde balloon pancreatography. *Gastrointest Endosc* 2003;58:510-515. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14520282>.

129. Nallamotheu G, Hilden K, Adler DG. Endoscopic retrograde cholangiopancreatography for non-gastroenterologists: what you need to know. *Hosp Pract (Minneap)* 2011;39:70-80. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21576899>.

130. Pavey DA, Gress FG. The role of EUS-guided FNA for the evaluation of biliary strictures. *Gastrointest Endosc* 2006;64:334-337. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16923478>.

131. Tirkes T, Sandrasegaran K, Sanyal R, et al. Secretin-enhanced MR cholangiopancreatography: spectrum of findings. *Radiographics* 2013;33:1889-1906. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24224585>.

132. Farma JM, Santillan AA, Melis M, et al. PET/CT fusion scan enhances CT staging in patients with pancreatic neoplasms. *Ann Surg*

2008;15:2465-2471. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18551347>.

133. Rijkers AP, Valkema R, Duivenvoorden HJ, van Eijck CH. Usefulness of F-18-fluorodeoxyglucose positron emission tomography to confirm suspected pancreatic cancer: a meta-analysis. *Eur J Surg Oncol* 2014;40:794-804. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24755095>.

134. Wang Z, Chen JQ, Liu JL, et al. FDG-PET in diagnosis, staging and prognosis of pancreatic carcinoma: a meta-analysis. *World J Gastroenterol* 2013;19:4808-4817. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23922481>.

135. Ahmed SI, Bochkarev V, Oleynikov D, Sasson AR. Patients with pancreatic adenocarcinoma benefit from staging laparoscopy. *J Laparoendosc Adv Surg Tech A* 2006;16:458-463. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17004868>.

136. Allen VB, Gurusamy KS, Takwoingi Y, et al. Diagnostic accuracy of laparoscopy following computed tomography (CT) scanning for assessing the resectability with curative intent in pancreatic and periampullary cancer. *Cochrane Database Syst Rev* 2013;11:Cd009323. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24272022>.

137. Warshaw AL, Gu ZY, Wittenberg J, Waltman AC. Preoperative staging and assessment of resectability of pancreatic cancer. *Arch Surg* 1990;125:230-233. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/2154172>.

138. Andersson R, Vagianos CE, Williamson RCN. Preoperative staging and evaluation of resectability in pancreatic ductal adenocarcinoma. *HPB (Oxford)* 2004;6:5-12. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18333037>.

139. Alexakis N, Gomatos IP, Sbarounis S, et al. High serum CA 19-9 but not tumor size should select patients for staging laparoscopy in

radiological resectable pancreas head and peri-ampullary cancer. Eur J Surg Oncol 2014. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25266999>.

140. Karachristos A, Scarneas N, Hoffman JP. CA 19-9 levels predict results of staging laparoscopy in pancreatic cancer. J Gastrointest Surg 2005;9:1286-1292. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16332484>.

141. White R, Winston C, Gonen M, et al. Current utility of staging laparoscopy for pancreatic and peripancreatic neoplasms. J Am Coll Surg 2008;206:445-450. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18308214>.

142. Ferrone CR, Haas B, Tang L, et al. The influence of positive peritoneal cytology on survival in patients with pancreatic adenocarcinoma. J Gastrointest Surg 2006;10:1347-1353. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17175453>.

143. Brugge WR, De Witt J, Klapman JB, et al. Techniques for cytologic sampling of pancreatic and bile duct lesions: The Papanicolaou Society of Cytopathology Guidelines. Cytojournal 2014;11:2. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25191516>.

144. Micames C, Jowell PS, White R, et al. Lower frequency of peritoneal carcinomatosis in patients with pancreatic cancer diagnosed by EUS-guided FNA vs. percutaneous FNA. Gastrointest Endosc 2003;58:690-695. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14595302>.

145. Okasha HH, Naga MI, Esmat S, et al. Endoscopic Ultrasound-Guided Fine Needle Aspiration versus Percutaneous Ultrasound-Guided Fine Needle Aspiration in Diagnosis of Focal Pancreatic Masses. Endosc Ultrasound 2013;2:190-193. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24949394>.

146. Layfield LJ, Dodd L, Factor R, Schmidt RL. Malignancy risk associated with diagnostic categories defined by the Papanicolaou

Society of Cytopathology pancreaticobiliary guidelines. Cancer Cytopathol 2014;122:420-427. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24339321>.

147. Chen YK, Pleskow DK. SpyGlass single-operator peroral cholangiopancreatography system for the diagnosis and therapy of bile-duct disorders: a clinical feasibility study (with video). Gastrointest Endosc 2007;65:832-841. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17466202>.

148. Strasberg SM, Middleton WD, Teefey SA, et al. Management of diagnostic dilemmas of the pancreas by ultrasonographically guided laparoscopic biopsy. Surgery 1999;126:736-741; discussion 741-733. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10520923>.

149. NIH state-of-the-science statement on endoscopic retrograde cholangiopancreatography (ERCP) for diagnosis and therapy. NIH Consens State Sci Statements 2002;19:1-26. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14768653>.

150. Campisi A, Brancatelli G, Vullierme MP, et al. Are pancreatic calcifications specific for the diagnosis of chronic pancreatitis? A multidetector-row CT analysis. Clin Radiol 2009;64:903-911. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19664481>.

151. Kajiwarra M, Kojima M, Konishi M, et al. Autoimmune pancreatitis with multifocal lesions. J Hepatobiliary Pancreat Surg 2008;15:449-452. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18670850>.

152. Kalady MF, Peterson B, Baillie J, et al. Pancreatic duct strictures: identifying risk of malignancy. Ann Surg Oncol 2004;11:581-588. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15150064>.

153. Menges M, Lerch MM, Zeitz M. The double duct sign in patients with malignant and benign pancreatic lesions. Gastrointest Endosc 2000;52:74-77. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10882966>.

154. Finkelberg DL, Sahani D, Deshpande V, Brugge WR. Autoimmune pancreatitis. *N Engl J Med* 2006;355:2670-2676. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17182992>.
155. Law R, Bronner M, Vogt D, Stevens T. Autoimmune pancreatitis: a mimic of pancreatic cancer. *Cleve Clin J Med* 2009;76:607-615. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19797461>.
156. Salla C, Chatzipantelis P, Konstantinou P, et al. EUS-FNA contribution in the identification of autoimmune pancreatitis: a case report. *JOP* 2007;8:598-604. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17873466>.
157. Holmes BJ, Hruban RH, Wolfgang CL, Ali SZ. Fine needle aspirate of autoimmune pancreatitis (lymphoplasmacytic sclerosing pancreatitis): cytomorphologic characteristics and clinical correlates. *Acta Cytol* 2012;56:228-232. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22555522>.
158. Learn PA, Grossman EB, Do RK, et al. Pitfalls in avoiding operation for autoimmune pancreatitis. *Surgery* 2011;150:968-974. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21893326>.
159. Hardacre JM, Iacobuzio-Donahue CA, Sohn TA, et al. Results of pancreaticoduodenectomy for lymphoplasmacytic sclerosing pancreatitis. *Ann Surg* 2003;237:853-858; discussion 858-859. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12796582>.
160. Sah RP, Chari ST. Autoimmune pancreatitis: an update on classification, diagnosis, natural history and management. *Curr Gastroenterol Rep* 2012;14:95-105. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22350841>.
161. Hamano H, Kawa S, Horiuchi A, et al. High serum IgG4 concentrations in patients with sclerosing pancreatitis. *N Engl J Med* 2001;344:732-738. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11236777>.
162. van Heerde MJ, Buijs J, Hansen BE, et al. Serum level of Ca 19-9 increases ability of IgG4 test to distinguish patients with autoimmune pancreatitis from those with pancreatic carcinoma. *Dig Dis Sci* 2014;59:1322-1329. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24385012>.
163. Safi F, Roscher R, Bittner R, et al. High sensitivity and specificity of CA 19-9 for pancreatic carcinoma in comparison to chronic pancreatitis. Serological and immunohistochemical findings. *Pancreas* 1987;2:398-403. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/3306667>.
164. Morris-Stiff G, Taylor MA. Ca19-9 and pancreatic cancer: Is it really that good? *J Gastrointest Oncol* 2012;3:88-89. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22811875>.
165. Ballehaninna UK, Chamberlain RS. The clinical utility of serum CA 19-9 in the diagnosis, prognosis and management of pancreatic adenocarcinoma: An evidence based appraisal. *J Gastrointest Oncol* 2012;3:105-119. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22811878>.
166. Hartwig W, Strobel O, Hinz U, et al. CA19-9 in potentially resectable pancreatic cancer: perspective to adjust surgical and perioperative therapy. *Ann Surg Oncol* 2013;20:2188-2196. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23247983>.
167. Kim YC, Kim HJ, Park JH, et al. Can preoperative CA19-9 and CEA levels predict the resectability of patients with pancreatic adenocarcinoma? *J Gastroenterol Hepatol* 2009;24:1869-1875. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19686409>.
168. Kondo N, Murakami Y, Uemura K, et al. Prognostic impact of perioperative serum CA 19-9 levels in patients with resectable pancreatic cancer. *Ann Surg Oncol* 2010;17:2321-2329. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20336387>.
169. Bauer TM, El-Rayes BF, Li X, et al. Carbohydrate antigen 19-9 is a prognostic and predictive biomarker in patients with advanced

pancreatic cancer who receive gemcitabine-containing chemotherapy: a pooled analysis of 6 prospective trials. *Cancer* 2013;119:285-292. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22786786>.

170. Berger AC, Garcia M, Hoffman JP, et al. Postresection CA 19-9 predicts overall survival in patients with pancreatic cancer treated with adjuvant chemoradiation: a prospective validation by RTOG 9704. *J Clin Oncol* 2008;26:5918-5922. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19029412>.

171. Berger AC, Winter K, Hoffman JP, et al. Five Year Results of US Intergroup/RTOG 9704 With Postoperative CA 19-9 $\leq$ 90 U/mL and Comparison to the CONKO-001 Trial. *Int J Radiat Oncol Biol Phys* 2012;84:e291-297. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22682806>.

172. Ferrone CR, Finkelstein DM, Thayer SP, et al. Perioperative CA19-9 levels can predict stage and survival in patients with resectable pancreatic adenocarcinoma. *J Clin Oncol* 2006;24:2897-2902. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16782929>.

173. Humphris JL, Chang DK, Johns AL, et al. The prognostic and predictive value of serum CA19.9 in pancreatic cancer. *Ann Oncol* 2012;23:1713-1722. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22241899>.

174. Montgomery RC, Hoffman JP, Riley LB, et al. Prediction of recurrence and survival by post-resection CA 19-9 values in patients with adenocarcinoma of the pancreas. *Ann Surg Oncol* 1997;4:551-556. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9367020>.

175. Tzeng CW, Balachandran A, Ahmad M, et al. Serum carbohydrate antigen 19-9 represents a marker of response to neoadjuvant therapy in patients with borderline resectable pancreatic cancer. *HPB (Oxford)* 2014;16:430-438. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23991810>.

176. Hess V, Glimelius B, Grawe P, et al. CA 19-9 tumour-marker response to chemotherapy in patients with advanced pancreatic cancer enrolled in a randomised controlled trial. *Lancet Oncol* 2008;9:132-138. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18249033>.

177. Pelzer U, Hilbig A, Sinn M, et al. Value of carbohydrate antigen 19-9 in predicting response and therapy control in patients with metastatic pancreatic cancer undergoing first-line therapy. *Front Oncol* 2013;3:155. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23785668>.

178. Halm U, Schumann T, Schiefke I, et al. Decrease of CA 19-9 during chemotherapy with gemcitabine predicts survival time in patients with advanced pancreatic cancer. *Br J Cancer* 2000;82:1013-1016. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10737382>.

179. Ishii H, Okada S, Sato T, et al. CA 19-9 in evaluating the response to chemotherapy in advanced pancreatic cancer. *Hepatogastroenterology* 1997;44:279-283. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9058159>.

180. Ko AH, Hwang J, Venook AP, et al. Serum CA19-9 response as a surrogate for clinical outcome in patients receiving fixed-dose rate gemcitabine for advanced pancreatic cancer. *Br J Cancer* 2005;93:195-199. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15999098>.

181. Wong D, Ko AH, Hwang J, et al. Serum CA19-9 decline compared to radiographic response as a surrogate for clinical outcomes in patients with metastatic pancreatic cancer receiving chemotherapy. *Pancreas* 2008;37:269-274. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18815548>.

182. Tempero MA, Uchida E, Takasaki H, et al. Relationship of carbohydrate antigen 19-9 and Lewis antigens in pancreatic cancer. *Cancer Res* 1987;47:5501-5503. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/3308077>.

183. Mann DV, Edwards R, Ho S, et al. Elevated tumour marker CA19-9: clinical interpretation and influence of obstructive jaundice. *Eur J*

Surg Oncol 2000;26:474-479. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/11016469>.

184. Marrelli D, Caruso S, Pedrazzani C, et al. CA19-9 serum levels in obstructive jaundice: clinical value in benign and malignant conditions. Am J Surg 2009;198:333-339. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/19375064>.

185. Mackey JR, Mani RS, Selner M, et al. Functional nucleoside transporters are required for gemcitabine influx and manifestation of toxicity in cancer cell lines. Cancer Res 1998;58:4349-4357. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9766663>.

186. Farrell JJ, Elsaleh H, Garcia M, et al. Human equilibrative nucleoside transporter 1 levels predict response to gemcitabine in patients with pancreatic cancer. Gastroenterology 2009;136:187-195. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18992248>.

187. Greenhalf W, Ghaneh P, Neoptolemos JP, et al. Pancreatic cancer hENT1 expression and survival from gemcitabine in patients from the ESPAC-3 trial. J Natl Cancer Inst 2014;106:djt347. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24301456>.

188. Liu ZQ, Han YC, Zhang X, et al. Prognostic value of human equilibrative nucleoside transporter1 in pancreatic cancer receiving gemcitabine-based chemotherapy: a meta-analysis. PLoS One 2014;9:e87103. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/24475233>.

189. Marechal R, Bachet JB, Mackey JR, et al. Levels of gemcitabine transport and metabolism proteins predict survival times of patients treated with gemcitabine for pancreatic adenocarcinoma. Gastroenterology 2012;143:664-674 e661-666. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/22705007>.

190. Saif M, Lee Y, Kim R. Harnessing gemcitabine metabolism: a step towards personalized medicine for pancreatic cancer. Ther Adv Med

Oncol 2012;4:341-346. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/23118809>.

191. Zhu Y, Qi M, Lao L, et al. Human Equilibrative Nucleoside Transporter 1 Predicts Survival in Patients with Pancreatic Cancer Treated with Gemcitabine: A Meta-Analysis. Genet Test Mol Biomarkers 2014. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24625353>.

192. Ormanns S, Heinemann V, Raponi M, et al. Human equilibrative nucleoside transporter 1 is not predictive for gemcitabine efficacy in advanced pancreatic cancer: translational results from the AIO-PK0104 phase III study with the clone SP120 rabbit antibody. Eur J Cancer 2014;50:1891-1899. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/24857044>.

193. Sinn M, Sinn BV, Stieler J, et al. Hent1 expression in patients with pancreatic cancer treated with gemcitabine after curative intended resection: Results from the CONKO-001 trial [abstract]. ASCO Meeting Abstracts 2014;32:4124. Available at:
http://meeting.ascopubs.org/cgi/content/abstract/32/15_suppl/4124.

194. Poplin E, Wasan H, Rolfe L, et al. Randomized, multicenter, phase II study of CO-101 versus gemcitabine in patients with metastatic pancreatic ductal adenocarcinoma: including a prospective evaluation of the role of hENT1 in gemcitabine or CO-101 sensitivity. J Clin Oncol 2013;31:4453-4461. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/24220555>.

195. Burris HA, 3rd, Moore MJ, Andersen J, et al. Improvements in survival and clinical benefit with gemcitabine as first-line therapy for patients with advanced pancreas cancer: a randomized trial. J Clin Oncol 1997;15:2403-2413. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/9196156>.

196. Oettle H, Post S, Neuhaus P, et al. Adjuvant chemotherapy with gemcitabine vs observation in patients undergoing curative-intent resection of pancreatic cancer: a randomized controlled trial. JAMA

2007;297:267-277. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/17227978>.

197. Oettle H, Neuhaus P, Hochhaus A, et al. Adjuvant chemotherapy with gemcitabine and long-term outcomes among patients with resected pancreatic cancer: the CONKO-001 randomized trial. *JAMA* 2013;310:1473-1481. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/24104372>.

198. Grunewald R, Abbruzzese JL, Tarassoff P, Plunkett W. Saturation of 2',2'-difluorodeoxycytidine 5'-triphosphate accumulation by mononuclear cells during a phase I trial of gemcitabine. *Cancer Chemother Pharmacol* 1991;27:258-262. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/1998982>.

199. Tempero M, Plunkett W, Ruiz Van Haperen V, et al. Randomized phase II comparison of dose-intense gemcitabine: thirty-minute infusion and fixed dose rate infusion in patients with pancreatic adenocarcinoma. *J Clin Oncol* 2003;21:3402-3408. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/12885837>.

200. Poplin E, Feng Y, Berlin J, et al. Phase III, randomized study of gemcitabine and oxaliplatin versus gemcitabine (fixed-dose rate infusion) compared with gemcitabine (30-minute infusion) in patients with pancreatic carcinoma E6201: a trial of the Eastern Cooperative Oncology Group. *J Clin Oncol* 2009;27:3778-3785. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19581537>.

201. Demols A, Peeters M, Polus M, et al. Gemcitabine and oxaliplatin (GEMOX) in gemcitabine refractory advanced pancreatic adenocarcinoma: a phase II study. *Br J Cancer* 2006;94:481-485. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16434988>.

202. Fine RL, Fogelman DR, Schreiber SM, et al. The gemcitabine, docetaxel, and capecitabine (GTX) regimen for metastatic pancreatic cancer: a retrospective analysis. *Cancer Chemother Pharmacol* 2008;61:167-175. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/17440727>.

203. Ko AH, Espinoza AM, Jones KA, et al. Optimizing the administration of fixed-dose rate gemcitabine plus capecitabine using an alternating-week schedule: a dose finding and early efficacy study in advanced pancreatic and biliary carcinomas. *Am J Clin Oncol* 2012;35:411-417. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/21552099>.

204. Berlin JD, Catalano P, Thomas JP, et al. Phase III study of gemcitabine in combination with fluorouracil versus gemcitabine alone in patients with advanced pancreatic carcinoma: Eastern Cooperative Oncology Group Trial E2297. *J Clin Oncol* 2002;20:3270-3275. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/12149301>.

205. Colucci G, Giuliani F, Gebbia V, et al. Gemcitabine alone or with cisplatin for the treatment of patients with locally advanced and/or metastatic pancreatic carcinoma: a prospective, randomized phase III study of the Gruppo Oncologia dell'Italia Meridionale. *Cancer* 2002;94:902-910. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/11920457>.

206. Colucci G, Labianca R, Di Costanzo F, et al. Randomized phase III trial of gemcitabine plus cisplatin compared with single-agent gemcitabine as first-line treatment of patients with advanced pancreatic cancer: the GIP-1 study. *J Clin Oncol* 2010;28:1645-1651. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/20194854>.

207. Cunningham D, Chau I, Stocken DD, et al. Phase III randomized comparison of gemcitabine versus gemcitabine plus capecitabine in patients with advanced pancreatic cancer. *J Clin Oncol* 2009;27:5513-5518. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19858379>.

208. Heinemann V, Boeck S, Hinke A, et al. Meta-analysis of randomized trials: evaluation of benefit from gemcitabine-based combination chemotherapy applied in advanced pancreatic cancer. *BMC Cancer* 2008;8:82-82. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/18373843>.

209. Heinemann V, Quietzsch D, Gieseler F, et al. Randomized phase III trial of gemcitabine plus cisplatin compared with gemcitabine alone in advanced pancreatic cancer. *J Clin Oncol* 2006;24:3946-3952. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16921047>.

210. Heinemann V, Labianca R, Hinke A, Louvet C. Increased survival using platinum analog combined with gemcitabine as compared to single-agent gemcitabine in advanced pancreatic cancer: pooled analysis of two randomized trials, the GERCOR/GISCAD intergroup study and a German multicenter study. *Ann Oncol* 2007;18:1652-1659. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17660491>.

211. Herrmann R, Bodoky G, Ruhstaller T, et al. Gemcitabine plus capecitabine compared with gemcitabine alone in advanced pancreatic cancer: a randomized, multicenter, phase III trial of the Swiss Group for Clinical Cancer Research and the Central European Cooperative Oncology Group. *J Clin Oncol* 2007;25:2212-2217. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17538165>.

212. Louvet C, Labianca R, Hammel P, et al. Gemcitabine in combination with oxaliplatin compared with gemcitabine alone in locally advanced or metastatic pancreatic cancer: results of a GERCOR and GISCAD phase III trial. *J Clin Oncol* 2005;23:3509-3516. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15908661>.

213. Reni M, Cordio S, Milandri C, et al. Gemcitabine versus cisplatin, epirubicin, fluorouracil, and gemcitabine in advanced pancreatic cancer: a randomised controlled multicentre phase III trial. *Lancet Oncol* 2005;6:369-376. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15925814>.

214. Rocha Lima CM, Green MR, Rotche R, et al. Irinotecan plus gemcitabine results in no survival advantage compared with gemcitabine monotherapy in patients with locally advanced or metastatic pancreatic cancer despite increased tumor response rate. *J Clin Oncol* 2004;22:3776-3783. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15365074>.

215. Ciliberto D, Botta C, Correale P, et al. Role of gemcitabine-based combination therapy in the management of advanced pancreatic cancer: A meta-analysis of randomised trials. *Eur J Cancer* 2013;49:593-603. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22989511>.

216. Sun C, Ansari D, Andersson R, Wu DQ. Does gemcitabine-based combination therapy improve the prognosis of unresectable pancreatic cancer? *World J Gastroenterol* 2012;18:4944-4958. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23002368>.

217. Kulke MH, Tempero MA, Niedzwiecki D, et al. Randomized phase II study of gemcitabine administered at a fixed dose rate or in combination with cisplatin, docetaxel, or irinotecan in patients with metastatic pancreatic cancer: CALGB 89904. *J Clin Oncol* 2009;27:5506-5512. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19858396>.

218. Stathopoulos GP, Syrigos K, Aravantinos G, et al. A multicenter phase III trial comparing irinotecan-gemcitabine (IG) with gemcitabine (G) monotherapy as first-line treatment in patients with locally advanced or metastatic pancreatic cancer. *Br J Cancer* 2006;95:587-592. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16909140>.

219. Goncalves A, Gilibert M, Francois E, et al. BAYPAN study: a double-blind phase III randomized trial comparing gemcitabine plus sorafenib and gemcitabine plus placebo in patients with advanced pancreatic cancer. *Ann Oncol* 2012;23:2799-2805. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22771827>.

220. Von Hoff DD, Ramanathan RK, Borad MJ, et al. Gemcitabine Plus nab-Paclitaxel Is an Active Regimen in Patients With Advanced Pancreatic Cancer: A Phase I/II Trial. *J Clin Oncol* 2011;29:4548-4554. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21969517>.

221. Von Hoff DD, Ervin T, Arena FP, et al. Increased survival in pancreatic cancer with nab-paclitaxel plus gemcitabine. *N Engl J Med*

2013;369:1691-1703. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/24131140>.

222. Goldstein D, El-Maraghi RH, Hammel P, et al. Analyses of updated overall survival (OS) and prognostic effect of neutrophil-to-lymphocyte ratio (NLR) and CA 19-9 from the phase III MPACT study of nab-paclitaxel (nab-P) plus gemcitabine (Gem) versus Gem for patients (pts) with metastatic pancreatic cancer (PC) [abstract]. ASCO Meeting Abstracts 2014;32:4027. Available at:

http://meeting.ascopubs.org/cgi/content/abstract/32/15_suppl/4027.

223. Kindler HL, Friberg G, Singh DA, et al. Phase II trial of bevacizumab plus gemcitabine in patients with advanced pancreatic cancer. J Clin Oncol 2005;23:8033-8040. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16258101>.

224. Xiong HQ, Rosenberg A, LoBuglio A, et al. Cetuximab, a monoclonal antibody targeting the epidermal growth factor receptor, in combination with gemcitabine for advanced pancreatic cancer: a multicenter phase II Trial. J Clin Oncol 2004;22:2610-2616. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15226328>.

225. Moore MJ, Goldstein D, Hamm J, et al. Erlotinib plus gemcitabine compared with gemcitabine alone in patients with advanced pancreatic cancer: a phase III trial of the National Cancer Institute of Canada Clinical Trials Group. J Clin Oncol 2007;25:1960-1966. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/17452677>.

226. Philip PA, Benedetti J, Corless CL, et al. Phase III study comparing gemcitabine plus cetuximab versus gemcitabine in patients with advanced pancreatic adenocarcinoma: Southwest Oncology Group-directed intergroup trial S0205. J Clin Oncol 2010;28:3605-3610. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/20606093>.

227. Kindler HL, Niedzwiecki D, Hollis D, et al. Gemcitabine plus bevacizumab compared with gemcitabine plus placebo in patients with advanced pancreatic cancer: phase III trial of the Cancer and Leukemia

Group B (CALGB 80303). J Clin Oncol 2010;28:3617-3622. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20606091>.

228. Kindler HL, Ioka T, Richel DJ, et al. Axitinib plus gemcitabine versus placebo plus gemcitabine in patients with advanced pancreatic adenocarcinoma: a double-blind randomised phase 3 study. Lancet Oncol 2011;12:256-262. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/21306953>.

229. Van Cutsem E, Vervenne WL, Bennouna J, et al. Phase III trial of bevacizumab in combination with gemcitabine and erlotinib in patients with metastatic pancreatic cancer. J Clin Oncol 2009;27:2231-2237. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19307500>.

230. Rougier P, Riess H, Manges R, et al. Randomised, placebo-controlled, double-blind, parallel-group phase III study evaluating aflibercept in patients receiving first-line treatment with gemcitabine for metastatic pancreatic cancer. Eur J Cancer 2013;49:2633-2642. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/23642329>.

231. Aranda E, Manzano JL, Rivera F, et al. Phase II open-label study of erlotinib in combination with gemcitabine in unresectable and/or metastatic adenocarcinoma of the pancreas: relationship between skin rash and survival (Pantar study). Ann Oncol 2012;23:1919-1925. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/22156621>.

232. Stepanski EJ, Reyes C, Walker MS, et al. The association of rash severity with overall survival: findings from patients receiving erlotinib for pancreatic cancer in the community setting. Pancreas 2013;42:32-36. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/22699203>.

233. Golan T, Kanji ZS, Epelbaum R, et al. Overall survival and clinical characteristics of pancreatic cancer in BRCA mutation carriers. Br J Cancer 2014;111:1132-1138. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/25072261>.

234. Majdak EJ, Debniak J, Milczek T, et al. Prognostic impact of BRCA1 pathogenic and BRCA1/BRCA2 unclassified variant mutations

in patients with ovarian carcinoma. Cancer 2005;104:1004-1012. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16047333>.

235. Stefansson OA, Jonasson JG, Johannsson OT, et al. Genomic profiling of breast tumours in relation to BRCA abnormalities and phenotypes. Breast Cancer Res 2009;11:R47. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19589159>.

236. Oliver GR, Sugar E, Laheru D, Diaz LA. Family history of cancer and sensitivity to platinum chemotherapy in pancreatic adenocarcinoma [abstract]. Gastrointestinal Cancers Symposium 2010:180. Available at: <http://meetinglibrary.asco.org/content/2395-72>.

237. Lowery MA, Kelsen DP, Stadler ZK, et al. An emerging entity: pancreatic adenocarcinoma associated with a known BRCA mutation: clinical descriptors, treatment implications, and future directions. Oncologist 2011;16:1397-1402. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21934105>.

238. Li Q, Yan H, Liu W, et al. Efficacy and safety of gemcitabine-fluorouracil combination therapy in the management of advanced pancreatic cancer: a meta-analysis of randomized controlled trials. PLoS One 2014;9:e104346. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25093849>.

239. De Jesus-Acosta A, Oliver GR, Blackford A, et al. A multicenter analysis of GTX chemotherapy in patients with locally advanced and metastatic pancreatic adenocarcinoma. Cancer Chemother Pharmacol 2012;69:415-424. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21800112>.

240. Neoptolemos JP, Stocken DD, Friess H, et al. A randomized trial of chemoradiotherapy and chemotherapy after resection of pancreatic cancer. N Engl J Med 2004;350:1200-1210. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15028824>.

241. Neoptolemos JP, Stocken DD, Bassi C, et al. Adjuvant chemotherapy with fluorouracil plus folinic acid vs gemcitabine following

pancreatic cancer resection: a randomized controlled trial. JAMA 2010;304:1073-1081. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20823433>.

242. Comparison of fluorouracil with additional levamisole, higher-dose folinic acid, or both, as adjuvant chemotherapy for colorectal cancer: a randomised trial. QUASAR Collaborative Group. Lancet 2000;355:1588-1596. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10821362>.

243. Jager E, Heike M, Bernhard H, et al. Weekly high-dose leucovorin versus low-dose leucovorin combined with fluorouracil in advanced colorectal cancer: results of a randomized multicenter trial. Study Group for Palliative Treatment of Metastatic Colorectal Cancer Study Protocol 1. J Clin Oncol 1996;14:2274-2279. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8708717>.

244. O'Connell MJ. A phase III trial of 5-fluorouracil and leucovorin in the treatment of advanced colorectal cancer. A Mayo Clinic/North Central Cancer Treatment Group study. Cancer 1989;63:1026-1030. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/2465076>.

245. Ychou M, Conroy T, Seitz JF, et al. An open phase I study assessing the feasibility of the triple combination: oxaliplatin plus irinotecan plus leucovorin/ 5-fluorouracil every 2 weeks in patients with advanced solid tumors. Ann Oncol 2003;14:481-489. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12598357>.

246. Conroy T, Paillet B, Francois E, et al. Irinotecan plus oxaliplatin and leucovorin-modulated fluorouracil in advanced pancreatic cancer--a Groupe Tumeurs Digestives of the Federation Nationale des Centres de Lutte Contre le Cancer study. J Clin Oncol 2005;23:1228-1236. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15718320>.

247. Ychou M, Desseigne F, Guimbaud R, et al. Randomized phase II trial comparing folfox (5FU/leucovorin [LV], irinotecan [I] and oxaliplatin [O]) vs gemcitabine (G) as first-line treatment for metastatic pancreatic adenocarcinoma (MPA). First results of the ACCORD 11 trial [abstract]. J Clin Oncol 2007;25 (June 20 Suppl):4516. Available at:

http://meeting.ascopubs.org/cgi/content/abstract/25/18_suppl/4516?sid=8904e4d6-689f-4715-8b08-891737076e04.

248. Conroy T, Desseigne F, Ychou M, et al. FOLFIRINOX versus gemcitabine for metastatic pancreatic cancer. *N Engl J Med* 2011;364:1817-1825. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21561347>.

249. Gourgou-Bourgade S, Bascoul-Molle C, Desseigne F, et al. Impact of FOLFIRINOX compared with gemcitabine on quality of life in patients with metastatic pancreatic cancer: results from the PRODIGE 4/ACCORD 11 randomized trial. *J Clin Oncol* 2013;31:23-29. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23213101>.

250. Lowery MA, Yu KH, Adel NG, et al. Activity of front-line FOLFIRINOX (FFX) in stage III/IV pancreatic adenocarcinoma (PC) at Memorial Sloan-Kettering Cancer Center (MSKCC) [abstract]. *ASCO Meeting Abstracts* 2012;30:4057. Available at: http://meeting.ascopubs.org/cgi/content/abstract/30/15_suppl/4057.

251. Boeck S, Vehling-Kaiser U, Waldschmidt D, et al. Erlotinib 150 mg daily plus chemotherapy in advanced pancreatic cancer: an interim safety analysis of a multicenter, randomized, cross-over phase III trial of the 'Arbeitsgemeinschaft Internistische Onkologie'. *Anticancer Drugs* 2010;21:94-100. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19770635>.

252. Cartwright TH, Cohn A, Varkey JA, et al. Phase II study of oral capecitabine in patients with advanced or metastatic pancreatic cancer. *J Clin Oncol* 2002;20:160-164. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11773165>.

253. Pelzer U, Schwaner I, Stieler J, et al. Best supportive care (BSC) versus oxaliplatin, folinic acid and 5-fluorouracil (OFF) plus BSC in patients for second-line advanced pancreatic cancer: a phase III-study from the German CONKO-study group. *Eur J Cancer* 2011;47:1676-1681. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21565490>.

254. Xiong HQ, Varadhachary GR, Blais JC, et al. Phase 2 trial of oxaliplatin plus capecitabine (XELOX) as second-line therapy for patients with advanced pancreatic cancer. *Cancer* 2008;113:2046-2052. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18756532>.

255. Reni M, Cereda S, Milella M, et al. Maintenance sunitinib or observation in metastatic pancreatic adenocarcinoma: a phase II randomised trial. *Eur J Cancer* 2013;49:3609-3615. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23899530>.

256. Rahma OE, Duffy A, Liewehr DJ, et al. Second-line treatment in advanced pancreatic cancer: a comprehensive analysis of published clinical trials. *Ann Oncol* 2013;24:1972-1979. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23670093>.

257. Maisey N, Chau I, Cunningham D, et al. Multicenter randomized phase III trial comparing protracted venous infusion (PVI) fluorouracil (5-FU) with PVI 5-FU plus mitomycin in inoperable pancreatic cancer. *J Clin Oncol* 2002;20:3130-3136. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12118027>.

258. Pelzer U, Kubica K, Stieler J, et al. A randomized trial in patients with gemcitabine refractory pancreatic cancer. Final results of the CONKO 003 study [abstract]. *J Clin Oncol* 2008;26 (May 20 suppl):4508. Available at: http://meeting.ascopubs.org/cgi/content/abstract/26/15_suppl/4508?sid=31ed1378-2eda-4a39-8a2a-177afb2c9fb4.

259. Saif MW. New developments in the treatment of pancreatic cancer. Highlights from the "44th ASCO Annual Meeting". Chicago, IL, USA. May 30 - June 3, 2008. *JOP* 2008;9:391-397. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18648128>.

260. Oettle H, Riess H, Stieler JM, et al. Second-line oxaliplatin, folinic acid, and fluorouracil versus folinic acid and fluorouracil alone for gemcitabine-refractory pancreatic cancer: outcomes from the CONKO-003 trial. *J Clin Oncol* 2014;32:2423-2429. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24982456>.

261. Gill S, Ko Y-J, Cripps MC, et al. PANCREOX: A randomized phase 3 study of 5FU/LV with or without oxaliplatin for second-line advanced pancreatic cancer (APC) in patients (pts) who have received gemcitabine (GEM)-based chemotherapy (CT) [abstract]. ASCO Meeting Abstracts 2014;32:4022. Available at: http://meeting.ascopubs.org/cgi/content/abstract/32/15_suppl/4022.
262. Heinemann V, Vehling-Kaiser U, Waldschmidt D, et al. Gemcitabine plus erlotinib followed by capecitabine versus capecitabine plus erlotinib followed by gemcitabine in advanced pancreatic cancer: final results of a randomised phase 3 trial of the 'Arbeitsgemeinschaft Internistische Onkologie' (AIO-PK0104). Gut 2013;62:751-759. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22773551>.
263. Seiwert TY, Salama JK, Vokes EE. The concurrent chemoradiation paradigm--general principles. Nat Clin Pract Oncol 2007;4:86-100. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17259930>.
264. Kalser MH, Ellenberg SS. Pancreatic cancer. Adjuvant combined radiation and chemotherapy following curative resection. Arch Surg 1985;120:899-903. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/4015380>.
265. Moertel CG, Frytak S, Hahn RG, et al. Therapy of locally unresectable pancreatic carcinoma: a randomized comparison of high dose (6000 rads) radiation alone, moderate dose radiation (4000 rads + 5-fluorouracil), and high dose radiation + 5-fluorouracil: The Gastrointestinal Tumor Study Group. Cancer 1981;48:1705-1710. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/7284971>.
266. Klinkenbijl JH, Jeekel J, Sahmoud T, et al. Adjuvant radiotherapy and 5-fluorouracil after curative resection of cancer of the pancreas and periampullary region: phase III trial of the EORTC gastrointestinal tract cancer cooperative group. Ann Surg 1999;230:776-782. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10615932>.
267. Smeenk HG, van Eijck CHJ, Hop WC, et al. Long-term survival and metastatic pattern of pancreatic and periampullary cancer after adjuvant chemoradiation or observation: long-term results of EORTC trial 40891. Ann Surg 2007;246:734-740. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17968163>.
268. Regine WF, Winter KA, Abrams RA, et al. Fluorouracil vs gemcitabine chemotherapy before and after fluorouracil-based chemoradiation following resection of pancreatic adenocarcinoma: a randomized controlled trial. JAMA 2008;299:1019-1026. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18319412>.
269. Garofalo MC, Abrams RA, Regine WF. Adjuvant therapy for pancreatic cancer: no 'definite' standard. Oncology 2007;21:726-730. Available at: <http://www.cancernetwork.com/display/article/10165/61708>.
270. Regine WF, Winter KA, Abrams R, et al. Fluorouracil-based chemoradiation with either gemcitabine or fluorouracil chemotherapy after resection of pancreatic adenocarcinoma: 5-year analysis of the U.S. Intergroup/RTOG 9704 phase III trial. Ann Surg Oncol 2011;18:1319-1326. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21499862>.
271. Crane CH, Ben-Josef E, Small W. Chemotherapy for pancreatic cancer. N Engl J Med 2004;350:2713-2715. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15218575>.
272. Koshy MC, Landry JC, Cavanaugh SX, et al. A challenge to the therapeutic nihilism of ESPAC-1. Int J Radiat Oncol Biol Phys 2005;61:965-966. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15752874>.
273. Morris SL, Beasley M, Leslie M. Chemotherapy for pancreatic cancer. N Engl J Med 2004;350:2713-2715; author reply 2713-2715. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15215490>.
274. Van Laethem JL, Hammel P, Mornex F, et al. Adjuvant gemcitabine alone versus gemcitabine-based chemoradiotherapy after curative resection for pancreatic cancer: a randomized EORTC-40013-

22012/FFCD-9203/GERCOR phase II study. J Clin Oncol 2010;28:4450-4456. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20837948>.

275. Schmidt J, Abel U, Debus J, et al. Open-label, multicenter, randomized phase III trial of adjuvant chemoradiation plus interferon Alfa-2b versus fluorouracil and folinic acid for patients with resected pancreatic adenocarcinoma. J Clin Oncol 2012;30:4077-4083. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23008325>.

276. Ren F, Xu YC, Wang HX, et al. Adjuvant chemotherapy, with or without postoperative radiotherapy, for resectable advanced pancreatic adenocarcinoma: continue or stop? Pancreatology 2012;12:162-169. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22487527>.

277. Liao WC, Chien KL, Lin YL, et al. Adjuvant treatments for resected pancreatic adenocarcinoma: a systematic review and network meta-analysis. Lancet Oncol 2013;14:1095-1103. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24035532>.

278. Kooby DA, Gillespie TW, Liu Y, et al. Impact of adjuvant radiotherapy on survival after pancreatic cancer resection: an appraisal of data from the national cancer data base. Ann Surg Oncol 2013;20:3634-3642. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23771249>.

279. Morganti AG, Falconi M, van Stiphout RG, et al. Multi-institutional Pooled Analysis on Adjuvant Chemoradiation in Pancreatic Cancer. Int J Radiat Oncol Biol Phys 2014. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25220717>.

280. Neoptolemos JP, Stocken DD, Dunn JA, et al. Influence of resection margins on survival for patients with pancreatic cancer treated by adjuvant chemoradiation and/or chemotherapy in the ESPAC-1 randomized controlled trial. Ann Surg 2001;234:758-768. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11729382>.

281. Herman JM, Swartz MJ, Hsu CC, et al. Analysis of fluorouracil-based adjuvant chemotherapy and radiation after pancreaticoduodenectomy for ductal adenocarcinoma of the pancreas: results of a large, prospectively collected database at the Johns Hopkins Hospital. J Clin Oncol 2008;26:3503-3510. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18640931>.

282. Corsini MM, Miller RC, Haddock MG, et al. Adjuvant radiotherapy and chemotherapy for pancreatic carcinoma: the Mayo Clinic experience (1975-2005). J Clin Oncol 2008;26:3511-3516. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18640932>.

283. Hsu CC, Herman JM, Corsini MM, et al. Adjuvant chemoradiation for pancreatic adenocarcinoma: the Johns Hopkins Hospital-Mayo Clinic collaborative study. Ann Surg Oncol 2010;17:981-990. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20087786>.

284. Butturini G, Stocken DD, Wente MN, et al. Influence of resection margins and treatment on survival in patients with pancreatic cancer: meta-analysis of randomized controlled trials. Arch Surg 2008;143:75-83; discussion 83. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18209156>.

285. Redmond KJ, Wolfgang CL, Sugar EA, et al. Adjuvant chemoradiation therapy for adenocarcinoma of the distal pancreas. Ann Surg Oncol 2010;17:3112-3119. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20680697>.

286. Stocken DD, Buchler MW, Dervenis C, et al. Meta-analysis of randomised adjuvant therapy trials for pancreatic cancer. Br J Cancer 2005;92:1372-1381. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15812554>.

287. Kim R, Saif MW. Is there an optimal neoadjuvant therapy for locally advanced pancreatic cancer? JOP 2007;8:279-288. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17495356>.

288. Crane CH, Abbruzzese JL, Evans DB, et al. Is the therapeutic index better with gemcitabine-based chemoradiation than with 5-fluorouracil-based chemoradiation in locally advanced pancreatic cancer? *Int J Radiat Oncol Biol Phys* 2002;52:1293-1302. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11955742>.

289. Blackstock AW, Tepper JE, Niedwiecki D, et al. Cancer and leukemia group B (CALGB) 89805: phase II chemoradiation trial using gemcitabine in patients with locoregional adenocarcinoma of the pancreas. *Int J Gastrointest Cancer* 2003;34:107-116. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15361643>.

290. Girard N, Mornex F, Bossard N, et al. Estimating optimal dose of twice-weekly gemcitabine for concurrent chemoradiotherapy in unresectable pancreatic carcinoma: mature results of GEMRT-01 Phase I trial. *Int J Radiat Oncol Biol Phys* 2010;77:1426-1432. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20056351>.

291. Loehrer PJ, Powell ME, Cardenes HR, et al. A randomized phase III study of gemcitabine in combination with radiation therapy versus gemcitabine alone in patients with localized, unresectable pancreatic cancer: E4201 [abstract]. *J Clin Oncol* 2008;26 (May 20 suppl):4506. Available at: http://meeting.ascopubs.org/cgi/content/abstract/26/15_suppl/4506?sid=526086db-bb84-4435-b063-52f14bdacd3c.

292. Murphy JD, Adusumilli S, Griffith KA, et al. Full-dose gemcitabine and concurrent radiotherapy for unresectable pancreatic cancer. *Int J Radiat Oncol Biol Phys* 2007;68:801-808. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17379445>.

293. Shibuya K, Oya N, Fujii T, et al. Phase II Study of Radiation Therapy Combined With Weekly Low-Dose Gemcitabine for Locally Advanced, Unresectable Pancreatic Cancer. *Am J Clin Oncol* 2010;34:115-119. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20065850>.

294. Huang J, Robertson JM, Margolis J, et al. Long-term results of full-dose gemcitabine with radiation therapy compared to 5-fluorouracil with radiation therapy for locally advanced pancreas cancer. *Radiother Oncol* 2011;99:114-119. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21621866>.

295. Zhu CP, Shi J, Chen YX, et al. Gemcitabine in the chemoradiotherapy for locally advanced pancreatic cancer: a meta-analysis. *Radiother Oncol* 2011;99:108-113. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21571383>.

296. Mukherjee S, Hurt CN, Bridgewater J, et al. Gemcitabine-based or capecitabine-based chemoradiotherapy for locally advanced pancreatic cancer (SCALOP): a multicentre, randomised, phase 2 trial. *Lancet Oncol* 2013;14:317-326. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23474363>.

297. Chen Y, Sun XJ, Jiang TH, Mao AW. Combined radiochemotherapy in patients with locally advanced pancreatic cancer: a meta-analysis. *World J Gastroenterol* 2013;19:7461-7471. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24259979>.

298. Treatment of locally unresectable carcinoma of the pancreas: comparison of combined-modality therapy (chemotherapy plus radiotherapy) to chemotherapy alone. Gastrointestinal Tumor Study Group. *J Natl Cancer Inst* 1988;80:751-755. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/2898536>.

299. Klaassen DJ, MacIntyre JM, Catton GE, et al. Treatment of locally unresectable cancer of the stomach and pancreas: a randomized comparison of 5-fluorouracil alone with radiation plus concurrent and maintenance 5-fluorouracil—an Eastern Cooperative Oncology Group study. *J Clin Oncol* 1985;3:373-378. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/3973648>.

300. Brunner TB, Grabenbauer GG, Kastl S, et al. Preoperative Chemoradiation in Locally Advanced Pancreatic Carcinoma: A Phase II

Study. *Onkologie* 2000;23:436-442. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/11441238>.

301. Macchia G, Valentini V, Mattiucci GC, et al. Preoperative chemoradiation and intra-operative radiotherapy for pancreatic carcinoma. *Tumori* 2007;93:53-60. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/17455872>.

302. Thomas CR, Jr., Weiden PL, Traverso LW, Thompson T. Concomitant intraarterial cisplatin, intravenous 5-fluorouracil, and split-course radiation therapy for locally advanced unresectable pancreatic adenocarcinoma: a phase II study of the Puget Sound Oncology Consortium (PSOC-703). *Am J Clin Oncol* 1997;20:161-165. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9124192>.

303. Cinar P, Ko AH. Evolving treatment options for locally advanced unresectable pancreatic ductal adenocarcinoma. *J Natl Compr Canc Netw* 2014;12:167-172. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/24586078>.

304. Loehrer PJ, Sr., Feng Y, Cardenes H, et al. Gemcitabine Alone Versus Gemcitabine Plus Radiotherapy in Patients With Locally Advanced Pancreatic Cancer: An Eastern Cooperative Oncology Group Trial. *J Clin Oncol* 2011;29:4105-4112. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/21969502>.

305. Philip PA. Locally Advanced Pancreatic Cancer: Where Should We Go From Here? *J Clin Oncol* 2011;29:4066-4068. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/21969514>.

306. Chauffert B, Mornex F, Bonnetain F, et al. Phase III trial comparing intensive induction chemoradiotherapy (60 Gy, infusional 5-FU and intermittent cisplatin) followed by maintenance gemcitabine with gemcitabine alone for locally advanced unresectable pancreatic cancer. Definitive results of the 2000-01 FFCD/SFRO study. *Ann Oncol* 2008;19:1592-1599. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/18467316>.

307. Huguet F, Andre T, Hammel P, et al. Impact of chemoradiotherapy after disease control with chemotherapy in locally advanced pancreatic adenocarcinoma in GERCOR phase II and III studies. *J Clin Oncol* 2007;25:326-331. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/17235048>.

308. Huguet F, Girard N, Guerche CS-E, et al. Chemoradiotherapy in the management of locally advanced pancreatic carcinoma: a qualitative systematic review. *J Clin Oncol* 2009;27:2269-2277. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19307501>.

309. Krishnan S, Rana V, Janjan NA, et al. Induction chemotherapy selects patients with locally advanced, unresectable pancreatic cancer for optimal benefit from consolidative chemoradiation therapy. *Cancer* 2007;110:47-55. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/17538975>.

310. Hammel P, Huguet F, Van Laethem J-L, et al. Comparison of chemoradiotherapy (CRT) and chemotherapy (CT) in patients with a locally advanced pancreatic cancer (LAPC) controlled after 4 months of gemcitabine with or without erlotinib: Final results of the international phase III LAP 07 study [abstract]. *ASCO Meeting Abstracts* 2013;31:LBA4003. Available at:
http://meeting.ascopubs.org/cgi/content/abstract/31/18_suppl/LBA4003.

311. Huguet F, Hammel P, Vernerey D, et al. Impact of chemoradiotherapy (CRT) on local control and time without treatment in patients with locally advanced pancreatic cancer (LAPC) included in the international phase III LAP 07 study [abstract]. *ASCO Meeting Abstracts* 2014;32:4001. Available at:
http://meeting.ascopubs.org/cgi/content/abstract/32/15_suppl/4001.

312. Bai YR, Wu GH, Guo WJ, et al. Intensity modulated radiation therapy and chemotherapy for locally advanced pancreatic cancer: results of feasibility study. *World J Gastroenterol* 2003;9:2561-2564. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14606097>.

313. Combs SE, Habermehl D, Kessel K, et al. Intensity modulated radiotherapy as neoadjuvant chemoradiation for the treatment of patients with locally advanced pancreatic cancer. Outcome analysis and comparison with a 3D-treated patient cohort. *Strahlenther Onkol* 2013;189:738-744. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/23896630>.

314. Crane CH, Antolak JA, Rosen, II, et al. Phase I study of concomitant gemcitabine and IMRT for patients with unresectable adenocarcinoma of the pancreatic head. *Int J Gastrointest Cancer* 2001;30:123-132. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/12540024>.

315. Milano MT, Chmura SJ, Garofalo MC, et al. Intensity-modulated radiotherapy in treatment of pancreatic and bile duct malignancies: toxicity and clinical outcome. *Int J Radiat Oncol Biol Phys* 2004;59:445-453. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15145161>.

316. Spalding AC, Jee K-W, Vineberg K, et al. Potential for dose-escalation and reduction of risk in pancreatic cancer using IMRT optimization with lexicographic ordering and gEUD-based cost functions. *Med Phys* 2007;34:521-529. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/17388169>.

317. Yovino S, Poppe M, Jabbour S, et al. Intensity-modulated radiation therapy significantly improves acute gastrointestinal toxicity in pancreatic and ampullary cancers. *Int J Radiat Oncol Biol Phys* 2011;79:158-162. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/20399035>.

318. Chang DT, Schellenberg D, Shen J, et al. Stereotactic radiotherapy for unresectable adenocarcinoma of the pancreas. *Cancer* 2009;115:665-672. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19117351>.

319. Chuong MD, Springett GM, Freilich JM, et al. Stereotactic body radiation therapy for locally advanced and borderline resectable pancreatic cancer is effective and well tolerated. *Int J Radiat Oncol Biol*

Phys 2013;86:516-522. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/23562768>.

320. Herman JM, Koong AC. Stereotactic body radiation therapy: a new standard option for pancreatic cancer? *J Natl Compr Canc Netw* 2014;12:1489-1493. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/25313185>.

321. Rwigema JC, Parikh SD, Heron DE, et al. Stereotactic body radiotherapy in the treatment of advanced adenocarcinoma of the pancreas. *Am J Clin Oncol* 2011;34:63-69. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/20308870>.

322. Tozzi A, Comito T, Alongi F, et al. SBRT in unresectable advanced pancreatic cancer: preliminary results of a mono-institutional experience. *Radiat Oncol* 2013;8:148. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/23799996>.

323. Wild AT, Hiniker SM, Chang DT, et al. Re-irradiation with stereotactic body radiation therapy as a novel treatment option for isolated local recurrence of pancreatic cancer after multimodality therapy: experience from two institutions. *J Gastrointest Oncol* 2013;4:343-351. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/24294505>.

324. Gunderson LL, Martin JK, Kvols LK, et al. Intraoperative and external beam irradiation +/- 5-FU for locally advanced pancreatic cancer. *Int J Radiat Oncol Biol Phys* 1987;13:319-329. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/3104244>.

325. Gunderson LL, Martin JK, Jr., Earle JD, et al. Intraoperative and external beam irradiation with or without resection: Mayo pilot experience. *Mayo Clin Proc* 1984;59:691-699. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/6482514>.

326. Mohiuddin M, Regine WF, Stevens J, et al. Combined intraoperative radiation and perioperative chemotherapy for

unresectable cancers of the pancreas. J Clin Oncol 1995;13:2764-2768. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/7595736>.

327. Roldan GE, Gunderson LL, Nagorney DM, et al. External beam versus intraoperative and external beam irradiation for locally advanced pancreatic cancer. Cancer 1988;61:1110-1116. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/3342371>.

328. Ashman JB, Moss AA, Rule WG, et al. Preoperative chemoradiation and IOERT for unresectable or borderline resectable pancreas cancer. J Gastrointest Oncol 2013;4:352-360. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24294506>.

329. Cai S, Hong TS, Goldberg SI, et al. Updated long-term outcomes and prognostic factors for patients with unresectable locally advanced pancreatic cancer treated with intraoperative radiotherapy at the Massachusetts General Hospital, 1978 to 2010. Cancer 2013;119:4196-4204. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24006012>.

330. Jingu K, Tanabe T, Nemoto K, et al. Intraoperative radiotherapy for pancreatic cancer: 30-year experience in a single institution in Japan. Int J Radiat Oncol Biol Phys 2012;83:e507-511. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22445002>.

331. Palta M, Willett C, Czito B. The role of intraoperative radiation therapy in patients with pancreatic cancer. Semin Radiat Oncol 2014;24:126-131. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24635869>.

332. Faris JE, Blaszkowsky LS, McDermott S, et al. FOLFIRINOX in locally advanced pancreatic cancer: the Massachusetts General Hospital Cancer Center experience. Oncologist 2013;18:543-548. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23657686>.

333. Ammori JB, Colletti LM, Zalupski MM, et al. Surgical resection following radiation therapy with concurrent gemcitabine in patients with previously unresectable adenocarcinoma of the pancreas. J

Gastrointest Surg 2003;7:766-772. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/13129554>.

334. Bickenbach KA, Gonen M, Tang LH, et al. Downstaging in pancreatic cancer: a matched analysis of patients resected following systemic treatment of initially locally unresectable disease. Ann Surg Oncol 2012;19:1663-1669. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22130621>.

335. Habermehl D, Kessel K, Welzel T, et al. Neoadjuvant chemoradiation with Gemcitabine for locally advanced pancreatic cancer. Radiat Oncol 2012;7:28. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22385572>.

336. Kadera BE, Sunjaya DB, Isacoff WH, et al. Locally advanced pancreatic cancer: association between prolonged preoperative treatment and lymph-node negativity and overall survival. JAMA Surg 2014;149:145-153. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24306217>.

337. Massucco P, Capussotti L, Magnino A, et al. Pancreatic resections after chemoradiotherapy for locally advanced ductal adenocarcinoma: analysis of perioperative outcome and survival. Ann Surg Oncol 2006;13:1201-1208. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16955382>.

338. Mondo EL, Noel MS, Katz AW, et al. Unresectable locally advanced pancreatic cancer: treatment with neoadjuvant leucovorin, fluorouracil, irinotecan, and oxaliplatin and assessment of surgical resectability. J Clin Oncol 2013;31:e37-39. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23233707>.

339. Mornex F, Girard N, Delpero J-R, Partensky C. Radiochemotherapy in the management of pancreatic cancer--part I: neoadjuvant treatment. Semin Radiat Oncol 2005;15:226-234. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16183476>.

340. Quiros RM, Brown KM, Hoffman JP. Neoadjuvant therapy in pancreatic cancer. *Cancer Invest* 2007;25:267-273. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17612937>.
341. White RR, Hurwitz HI, Morse MA, et al. Neoadjuvant chemoradiation for localized adenocarcinoma of the pancreas. *Ann Surg Oncol* 2001;8:758-765. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11776488>.
342. Gillen S, Schuster T, Meyer Zum Buschenfelde C, et al. Preoperative/neoadjuvant therapy in pancreatic cancer: a systematic review and meta-analysis of response and resection percentages. *PLoS Med* 2010;7:e1000267. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20422030>.
343. Li D, Xie K, Wolff R, Abbruzzese JL. Pancreatic cancer. *Lancet* 2004;363:1049-1057. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15051286>.
344. Gudjonsson B. Cancer of the pancreas. 50 years of surgery. *Cancer* 1987;60:2284-2303. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/3326653>.
345. Crist DW, Sitzmann JV, Cameron JL. Improved hospital morbidity, mortality, and survival after the Whipple procedure. *Ann Surg* 1987;206:358-365. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/3632096>.
346. Allison DC, Piantadosi S, Hruban RH, et al. DNA content and other factors associated with ten-year survival after resection of pancreatic carcinoma. *J Surg Oncol* 1998;67:151-159. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9530884>.
347. Howard TJ, Krug JE, Yu J, et al. A margin-negative R0 resection accomplished with minimal postoperative complications is the surgeon's contribution to long-term survival in pancreatic cancer. *J Gastrointest Surg* 2006;10:1338-1345. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17175452>.
348. Sohn TA, Yeo CJ, Cameron JL, et al. Resected adenocarcinoma of the pancreas-616 patients: results, outcomes, and prognostic indicators. *J Gastrointest Surg* 2000;4:567-579. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11307091>.
349. Bilimoria KY, Talamonti MS, Sener SF, et al. Effect of hospital volume on margin status after pancreaticoduodenectomy for cancer. *J Am Coll Surg* 2008;207:510-519. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18926452>.
350. Winter JM, Cameron JL, Campbell KA, et al. 1423 pancreaticoduodenectomies for pancreatic cancer: A single-institution experience. *J Gastrointest Surg* 2006;10:1199-1210; discussion 1210-1191. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17114007>.
351. Zervos EE, Rosemurgy AS, Al-Saif O, Durkin AJ. Surgical management of early-stage pancreatic cancer. *Cancer Control* 2004;11:23-31. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14749620>.
352. Abrams RA, Lowy AM, O'Reilly EM, et al. Combined modality treatment of resectable and borderline resectable pancreas cancer: expert consensus statement. *Ann Surg Oncol* 2009;16:1751-1756. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19390900>.
353. Bockhorn M, Uzunoglu FG, Adham M, et al. Borderline resectable pancreatic cancer: a consensus statement by the International Study Group of Pancreatic Surgery (ISGPS). *Surgery* 2014;155:977-988. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24856119>.
354. Sobin LH, Gospodarowicz MK, Wittekind C, eds. *TNM Classification of Malignant Tumours* (ed 7th): John Wiley & Sons; 2009.
355. Varadhachary GR, Tamm EP, Abbruzzese JL, et al. Borderline resectable pancreatic cancer: definitions, management, and role of preoperative therapy. *Ann Surg Oncol* 2006;13:1035-1046. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16865597>.

356. Katz MH, Marsh R, Herman JM, et al. Borderline resectable pancreatic cancer: need for standardization and methods for optimal clinical trial design. *Ann Surg Oncol* 2013;20:2787-2795. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23435609>.

357. Talamonti M. Borderline resectable pancreatic cancer: a new classification for an old challenge. *Ann Surg Oncol* 2006;13:1019-1020. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16865593>.

358. Gumbs AA, Rodriguez Rivera AM, Milone L, Hoffman JP. Laparoscopic Pancreatoduodenectomy: A Review of 285 Published Cases. *Ann Surg Oncol* 2011;18:1335-1341. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21207166>.

359. Venkat R, Edil BH, Schulick RD, et al. Laparoscopic distal pancreatectomy is associated with significantly less overall morbidity compared to the open technique: a systematic review and meta-analysis. *Ann Surg* 2012;255:1048-1059. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22511003>.

360. Nakeeb A, Lillemoe KD, Grosfeld JL. Surgical techniques for pancreatic cancer. *Minerva Chir* 2004;59:151-163. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15238889>.

361. Yeo TP, Hruban RH, Leach SD, et al. Pancreatic cancer. *Curr Probl Cancer* 2002;26:176-275. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12399802>.

362. Baque P, Iannelli A, Delotte J, et al. Division of the right posterior attachments of the head of the pancreas with a linear stapler during pancreaticoduodenectomy: vascular and oncological considerations based on an anatomical cadaver-based study. *Surg Radiol Anat* 2009;31:13-17. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18712270>.

363. Evans DB, Pisters PW. Novel applications of endo GIA linear staplers during pancreaticoduodenectomy and total pancreatectomy.

Am J Surg 2003;185:606-607. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12781900>.

364. Harrison LE, Klimstra DS, Brennan MF. Isolated portal vein involvement in pancreatic adenocarcinoma. A contraindication for resection? *Ann Surg* 1996;224:342-347; discussion 347-349. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8813262>.

365. Riediger H, Makowiec F, Fischer E, et al. Postoperative morbidity and long-term survival after pancreaticoduodenectomy with superior mesenterico-portal vein resection. *J Gastrointest Surg* 2006;10:1106-1115. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16966029>.

366. Tseng JF, Raut CP, Lee JE, et al. Pancreaticoduodenectomy with vascular resection: margin status and survival duration. *J Gastrointest Surg* 2004;8:935-949. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15585381>.

367. Stitzenberg KB, Watson JC, Roberts A, et al. Survival after pancreatectomy with major arterial resection and reconstruction. *Ann Surg Oncol* 2008;15:1399-1406. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18320285>.

368. Mollberg N, Rahbari NN, Koch M, et al. Arterial resection during pancreatectomy for pancreatic cancer: a systematic review and meta-analysis. *Ann Surg* 2011;254:882-893. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22064622>.

369. Worni M, Castleberry AW, Clary BM, et al. Concomitant Vascular Reconstruction During Pancreatectomy for Malignant Disease: A Propensity Score-Adjusted, Population-Based Trend Analysis Involving 10 206 Patients. *Arch Surg* 2012;1-8. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23247767>.

370. Christein JD, Kendrick ML, Iqbal CW, et al. Distal pancreatectomy for resectable adenocarcinoma of the body and tail of the pancreas. *J Gastrointest Surg* 2005;9:922-927. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16137585>.

371. Shoup M, Conlon KC, Klimstra D, Brennan MF. Is extended resection for adenocarcinoma of the body or tail of the pancreas justified? J Gastrointest Surg 2003;7:946-952; discussion 952. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14675703>.

372. Strasberg SM, Linehan DC, Hawkins WG. Radical antegrade modular pancreateosplenectomy procedure for adenocarcinoma of the body and tail of the pancreas: ability to obtain negative tangential margins. J Am Coll Surg 2007;204:244-249. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17254928>.

373. Stauffer JA, Rosales-Velderrain A, Goldberg RF, et al. Comparison of open with laparoscopic distal pancreatectomy: a single institution's transition over a 7-year period. HPB (Oxford) 2013;15:149-155. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23297726>.

374. Pericleous S, Middleton N, McKay SC, et al. Systematic review and meta-analysis of case-matched studies comparing open and laparoscopic distal pancreatectomy: is it a safe procedure? Pancreas 2012;41:993-1000. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22836858>.

375. Tran Cao HS, Lopez N, Chang DC, et al. Improved perioperative outcomes with minimally invasive distal pancreatectomy: results from a population-based analysis. JAMA Surg 2014;149:237-243. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24402232>.

376. Fortner JG. Regional pancreatectomy for cancer of the pancreas, ampulla, and other related sites. Tumor staging and results. Ann Surg 1984;199:418-425. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/6712317>.

377. Fuhrman GM, Leach SD, Staley CA, et al. Rationale for en bloc vein resection in the treatment of pancreatic adenocarcinoma adherent to the superior mesenteric-portal vein confluence. Pancreatic Tumor Study Group. Ann Surg 1996;223:154-162. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8597509>.

378. Leach SD, Lee JE, Charnsangavej C, et al. Survival following pancreaticoduodenectomy with resection of the superior mesenteric-portal vein confluence for adenocarcinoma of the pancreatic head. Br J Surg 1998;85:611-617. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9635805>.

379. Clavien PA, Rudiger HA. A simple technique of portal vein resection and reconstruction during pancreaticoduodenectomy. J Am Coll Surg 1999;189:629-634. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10589601>.

380. Launois B, Stasik C, Bardaxoglou E, et al. Who benefits from portal vein resection during pancreaticoduodenectomy for pancreatic cancer? World J Surg 1999;23:926-929. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10449822>.

381. Taschieri AM, Elli M, Rovati M, et al. Surgical treatment of pancreatic tumors invading the spleno-mesenteric-portal vessels. An Italian Multicenter Survey. Hepatogastroenterology 1999;46:492-497. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10228849>.

382. van Geenen RC, ten Kate FJ, de Wit LT, et al. Segmental resection and wedge excision of the portal or superior mesenteric vein during pancreatoduodenectomy. Surgery 2001;129:158-163. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11174708>.

383. Yu XZ, Li J, Fu DL, et al. Benefit from synchronous portal-superior mesenteric vein resection during pancreaticoduodenectomy for cancer: a meta-analysis. Eur J Surg Oncol 2014;40:371-378. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24560302>.

384. Kelly KJ, Winslow E, Kooby D, et al. Vein involvement during pancreaticoduodenectomy: is there a need for redefinition of "borderline resectable disease"? J Gastrointest Surg 2013;17:1209-1217; discussion 1217. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23620151>.

385. Traverso LW, Longmire WP, Jr. Preservation of the pylorus in pancreaticoduodenectomy. Surg Gynecol Obstet 1978;146:959-962. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/653575>.

386. Yeo CJ, Cameron JL, Sohn TA, et al. Six hundred fifty consecutive pancreaticoduodenectomies in the 1990s: pathology, complications, and outcomes. Ann Surg 1997;226:248-257. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9339931>.

387. van Berge Henegouwen MI, van Gulik TM, DeWit LT, et al. Delayed gastric emptying after standard pancreaticoduodenectomy versus pylorus-preserving pancreaticoduodenectomy: an analysis of 200 consecutive patients. J Am Coll Surg 1997;185:373-379. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9328386>.

388. Kozuschek W, Reith HB, Waleczek H, et al. A comparison of long term results of the standard Whipple procedure and the pylorus preserving pancreatoduodenectomy. J Am Coll Surg 1994;178:443-453. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/7909485>.

389. Lin PW, Lin YJ. Prospective randomized comparison between pylorus-preserving and standard pancreaticoduodenectomy. Br J Surg 1999;86:603-607. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10361177>.

390. Matsumoto I, Shinzeki M, Asari S, et al. A prospective randomized comparison between pylorus- and subtotal stomach-preserving pancreatoduodenectomy on postoperative delayed gastric emptying occurrence and long-term nutritional status. J Surg Oncol 2014;109:690-696. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24619624>.

391. Morel P, Mathey P, Corboud H, et al. Pylorus-preserving duodenopancreatectomy: long-term complications and comparison with the Whipple procedure. World J Surg 1990;14:642-646. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/2238666>.

392. Roder JD, Stein HJ, Huttli W, Siewert JR. Pylorus-preserving versus standard pancreatico-duodenectomy: an analysis of 110 pancreatic and periampullary carcinomas. Br J Surg 1992;79:152-155. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/1348201>.

393. Seiler CA, Wagner M, Sadowski C, et al. Randomized prospective trial of pylorus-preserving vs. Classic duodenopancreatectomy (Whipple procedure): initial clinical results. J Gastrointest Surg 2000;4:443-452. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11077317>.

394. Yeo CJ, Cameron JL, Maher MM, et al. A prospective randomized trial of pancreaticogastrostomy versus pancreaticojejunostomy after pancreaticoduodenectomy. Ann Surg 1995;222:580-588. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/7574936>.

395. Topal B, Fieuws S, Aerts R, et al. Pancreaticojejunostomy versus pancreaticogastrostomy reconstruction after pancreaticoduodenectomy for pancreatic or periampullary tumours: a multicentre randomised trial. Lancet Oncol 2013;14:655-662. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23643139>.

396. Wolfgang CL, Pawlik TM. Pancreaticoduodenectomy: time to change our approach? Lancet Oncol 2013;14:573-575. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23643140>.

397. Hallet J, Zih FS, Deobald RG, et al. The impact of pancreaticojejunostomy versus pancreaticogastrostomy reconstruction on pancreatic fistula after pancreaticoduodenectomy: meta-analysis of randomized controlled trials. HPB (Oxford) 2014. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25040921>.

398. Gomez T, Palomares A, Serradilla M, Tejedor L. Reconstruction after pancreatoduodenectomy: Pancreatojejunostomy vs pancreatogastrostomy. World J Gastrointest Oncol 2014;6:369-376. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25232462>.

399. Bassi C, Falconi M, Molinari E, et al. Duct-to-mucosa versus end-to-side pancreaticojejunostomy reconstruction after

pancreaticoduodenectomy: results of a prospective randomized trial. Surgery 2003;134:766-771. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14639354>.

400. Sikora SS, Posner MC. Management of the pancreatic stump following pancreaticoduodenectomy. Br J Surg 1995;82:1590-1597. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8548218>.

401. Strasberg SM, Drebin JA, Mokadam NA, et al. Prospective trial of a blood supply-based technique of pancreaticojejunostomy: effect on anastomotic failure in the Whipple procedure. J Am Coll Surg 2002;194:746-758. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12081065>.

402. Winter JM, Cameron JL, Campbell KA, et al. Does pancreatic duct stenting decrease the rate of pancreatic fistula following pancreaticoduodenectomy? Results of a prospective randomized trial. J Gastrointest Surg 2006;10:1280-1290. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17114014>.

403. Lowy AM, Lee JE, Pisters PW, et al. Prospective, randomized trial of octreotide to prevent pancreatic fistula after pancreaticoduodenectomy for malignant disease. Ann Surg 1997;226:632-641. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9389397>.

404. Yeo CJ, Cameron JL, Lillemoe KD, et al. Does prophylactic octreotide decrease the rates of pancreatic fistula and other complications after pancreaticoduodenectomy? Results of a prospective randomized placebo-controlled trial. Ann Surg 2000;232:419-429. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10973392>.

405. Allen PJ, Gonen M, Brennan MF, et al. Pasireotide for postoperative pancreatic fistula. N Engl J Med 2014;370:2014-2022. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24849084>.

406. Lillemoe KD, Cameron JL, Kim MP, et al. Does fibrin glue sealant decrease the rate of pancreatic fistula after pancreaticoduodenectomy?

Results of a prospective randomized trial. J Gastrointest Surg 2004;8:766-772. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15531229>.

407. Cubilla AL, Fortner J, Fitzgerald PJ. Lymph node involvement in carcinoma of the head of the pancreas area. Cancer 1978;41:880-887. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/638975>.

408. Nagai H, Kuroda A, Morioka Y. Lymphatic and local spread of T1 and T2 pancreatic cancer. A study of autopsy material. Ann Surg 1986;204:65-71. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/3015059>.

409. Glanemann M, Shi B, Liang F, et al. Surgical strategies for treatment of malignant pancreatic tumors: extended, standard or local surgery? World J Surg Oncol 2008;6:123. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19014474>.

410. Pisters P, Brennan M. Regional lymph node dissection for pancreatic adenocarcinoma. In: Evans D, Pisters P, Abbruzzese J, eds., eds. Pancreatic Cancer. New York: Springer-Verlag; 2002:139-151.

411. Pedrazzoli S, DiCarlo V, Dionigi R, et al. Standard versus extended lymphadenectomy associated with pancreatoduodenectomy in the surgical treatment of adenocarcinoma of the head of the pancreas: a multicenter, prospective, randomized study. Lymphadenectomy Study Group. Ann Surg 1998;228:508-517. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9790340>.

412. Yeo CJ, Cameron JL, Sohn TA, et al. Pancreaticoduodenectomy with or without extended retroperitoneal lymphadenectomy for periampullary adenocarcinoma: comparison of morbidity and mortality and short-term outcome. Ann Surg 1999;229:613-622; discussion 622-614. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10235519>.

413. Riall TS, Cameron JL, Lillemoe KD, et al. Pancreaticoduodenectomy with or without distal gastrectomy and extended retroperitoneal lymphadenectomy for periampullary

adenocarcinoma--part 3: update on 5-year survival. J Gastrointest Surg 2005;9:1191-1204; discussion 1204-1196. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16332474>.

414. Yeo CJ, Cameron JL, Lillemoe KD, et al. Pancreaticoduodenectomy with or without distal gastrectomy and extended retroperitoneal lymphadenectomy for periampullary adenocarcinoma, part 2: randomized controlled trial evaluating survival, morbidity, and mortality. Ann Surg 2002;236:355-366; discussion 366-358. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12192322>.

415. Nimura Y, Nagino M, Takao S, et al. Standard versus extended lymphadenectomy in radical pancreatoduodenectomy for ductal adenocarcinoma of the head of the pancreas: long-term results of a Japanese multicenter randomized controlled trial. J Hepatobiliary Pancreat Sci 2012;19:230-241. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22038501>.

416. Michalski CW, Kleeff J, Wente MN, et al. Systematic review and meta-analysis of standard and extended lymphadenectomy in pancreaticoduodenectomy for pancreatic cancer. Br J Surg 2007;94:265-273. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17318801>.

417. Sun J, Yang Y, Wang X, et al. Meta-analysis of the Efficacies of Extended and Standard Pancreatoduodenectomy for Ductal Adenocarcinoma of the Head of the Pancreas. World J Surg 2014;38:2708-2715. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24912627>.

418. Tol JA, Gouma DJ, Bassi C, et al. Definition of a standard lymphadenectomy in surgery for pancreatic ductal adenocarcinoma: a consensus statement by the International Study Group on Pancreatic Surgery (ISGPS). Surgery 2014;156:591-600. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25061003>.

419. Farnell MB, Aranha GV, Nimura Y, Michelassi F. The role of extended lymphadenectomy for adenocarcinoma of the head of the

pancreas: strength of the evidence. J Gastrointest Surg 2008;12:651-656. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18085343>.

420. Shrikhande SV, Barreto SG. Extended pancreatic resections and lymphadenectomy: An appraisal of the current evidence. World J Gastrointest Surg 2010;2:39-46. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21160848>.

421. Cordera F, Arciero CA, Li T, et al. Significance of common hepatic artery lymph node metastases during pancreaticoduodenectomy for pancreatic head adenocarcinoma. Ann Surg Oncol 2007;14:2330-2336. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17492334>.

422. Shimada K, Sakamoto Y, Sano T, Kosuge T. The role of paraaortic lymph node involvement on early recurrence and survival after macroscopic curative resection with extended lymphadenectomy for pancreatic carcinoma. J Am Coll Surg 2006;203:345-352. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16931307>.

423. Bottger TC, Junginger T. Factors influencing morbidity and mortality after pancreaticoduodenectomy: critical analysis of 221 resections. World J Surg 1999;23:164-171. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9880426>.

424. Braasch JW, Gray BN. Considerations that lower pancreatoduodenectomy mortality. Am J Surg 1977;133:480-484. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/848682>.

425. Lerut JP, Gianello PR, Otte JB, Kestens PJ. Pancreaticoduodenal resection. Surgical experience and evaluation of risk factors in 103 patients. Ann Surg 1984;199:432-437. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/6712319>.

426. Gundry SR, Strodel WE, Knol JA, et al. Efficacy of preoperative biliary tract decompression in patients with obstructive jaundice. Arch Surg 1984;119:703-708. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/6428380>.

427. Hatfield AR, Tobias R, Terblanche J, et al. Preoperative external biliary drainage in obstructive jaundice. A prospective controlled clinical trial. *Lancet* 1982;2:896-899. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/6126752>.
428. Heslin MJ, Brooks AD, Hochwald SN, et al. A preoperative biliary stent is associated with increased complications after pancreatoduodenectomy. *Arch Surg* 1998;133:149-154. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9484726>.
429. Lai EC, Mok FP, Fan ST, et al. Preoperative endoscopic drainage for malignant obstructive jaundice. *Br J Surg* 1994;81:1195-1198. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/7741850>.
430. McPherson GA, Benjamin IS, Hodgson HJ, et al. Pre-operative percutaneous transhepatic biliary drainage: the results of a controlled trial. *Br J Surg* 1984;71:371-375. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/6372935>.
431. Pitt HA, Gomes AS, Lois JF, et al. Does preoperative percutaneous biliary drainage reduce operative risk or increase hospital cost? *Ann Surg* 1985;201:545-553. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/2986562>.
432. Thomas JH, Connor CS, Pierce GE, et al. Effect of biliary decompression on morbidity and mortality of pancreatoduodenectomy. *Am J Surg* 1984;148:727-731. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/6439064>.
433. Cavell LK, Allen PJ, Vinoya C, et al. Biliary self-expandable metal stents do not adversely affect pancreaticoduodenectomy. *Am J Gastroenterol* 2013;108:1168-1173. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23545711>.
434. van der Gaag NA, Rauws EA, van Eijck CH, et al. Preoperative biliary drainage for cancer of the head of the pancreas. *N Engl J Med* 2010;362:129-137. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20071702>.
435. Pisters PW, Hudec WA, Hess KR, et al. Effect of preoperative biliary decompression on pancreaticoduodenectomy-associated morbidity in 300 consecutive patients. *Ann Surg* 2001;234:47-55. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11420482>.
436. Aadam AA, Evans DB, Khan A, et al. Efficacy and safety of self-expandable metal stents for biliary decompression in patients receiving neoadjuvant therapy for pancreatic cancer: a prospective study. *Gastrointest Endosc* 2012;76:67-75. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22483859>.
437. Mullen JT, Lee JH, Gomez HF, et al. Pancreaticoduodenectomy after placement of endobiliary metal stents. *J Gastrointest Surg* 2005;9:1094-1104; discussion 1104-1095. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16269380>.
438. Varadhachary GR, Wolff RA, Crane CH, et al. Preoperative gemcitabine and cisplatin followed by gemcitabine-based chemoradiation for resectable adenocarcinoma of the pancreatic head. *J Clin Oncol* 2008;26:3487-3495. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18640929>.
439. Varadhachary GR, Wolff RA. The War on Pancreatic Cancer: Are We Gaining Ground? *Oncology* 2011;24:1335-1336. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21294479>.
440. Krokidis M, Fanelli F, Orgera G, et al. Percutaneous Palliation of Pancreatic Head Cancer: Randomized Comparison of ePTFE/FEP-Covered Versus Uncovered Nitinol Biliary Stents. *Cardiovasc Intervent Radiol* 2010;34:352-361. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20467870>.
441. Kullman E, Frozanpor F, Soderlund C, et al. Covered versus uncovered self-expandable nitinol stents in the palliative treatment of malignant distal biliary obstruction: results from a randomized, multicenter study. *Gastrointest Endosc* 2010;72:915-923. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21034892>.

442. Chun HJ, Kim ES, Hyun JJ, et al. Gastrointestinal and biliary stents. J Gastroenterol Hepatol 2010;25:234-243. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20136988>.
443. Ho H, Mahajan A, Gosain S, et al. Management of complications associated with partially covered biliary metal stents. Dig Dis Sci 2010;55:516-522. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19267200>.
444. Telford JJ, Carr-Locke DL, Baron TH, et al. A randomized trial comparing uncovered and partially covered self-expandable metal stents in the palliation of distal malignant biliary obstruction. Gastrointest Endosc 2010;72:907-914. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21034891>.
445. Lieberman MD, Kilburn H, Lindsey M, Brennan MF. Relation of perioperative deaths to hospital volume among patients undergoing pancreatic resection for malignancy. Ann Surg 1995;222:638-645. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/7487211>.
446. Gordon TA, Burleyson GP, Tielsch JM, Cameron JL. The effects of regionalization on cost and outcome for one general high-risk surgical procedure. Ann Surg 1995;221:43-49. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/7826160>.
447. Ho V, Heslin MJ. Effect of hospital volume and experience on in-hospital mortality for pancreaticoduodenectomy. Ann Surg 2003;237:509-514. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12677147>.
448. Imperato PJ, Nenner RP, Starr HA, et al. The effects of regionalization on clinical outcomes for a high risk surgical procedure: a study of the Whipple procedure in New York State. Am J Med Qual 1996;11:193-197. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8972936>.
449. Rosemurgy AS, Bloomston M, Serafini FM, et al. Frequency with which surgeons undertake pancreaticoduodenectomy determines length of stay, hospital charges, and in-hospital mortality. J Gastrointest Surg 2001;5:21-26. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11309644>.
450. Sosa JA, Bowman HM, Gordon TA, et al. Importance of hospital volume in the overall management of pancreatic cancer. Ann Surg 1998;228:429-438. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9742926>.
451. Gouma DJ, van Geenen RC, van Gulik TM, et al. Rates of complications and death after pancreaticoduodenectomy: risk factors and the impact of hospital volume. Ann Surg 2000;232:786-795. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11088073>.
452. Simunovic M, To T, Theriault M, Langer B. Relation between hospital surgical volume and outcome for pancreatic resection for neoplasm in a publicly funded health care system. CMAJ 1999;160:643-648. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10101998>.
453. van Heek NT, Kuhlmann KF, Scholten RJ, et al. Hospital volume and mortality after pancreatic resection: a systematic review and an evaluation of intervention in the Netherlands. Ann Surg 2005;242:781-788, discussion 788-790. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16327488>.
454. Birkmeyer JD, Finlayson SR, Tosteson AN, et al. Effect of hospital volume on in-hospital mortality with pancreaticoduodenectomy. Surgery 1999;125:250-256. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10076608>.
455. Birkmeyer JD, Siewers AE, Finlayson EVA, et al. Hospital volume and surgical mortality in the United States. N Engl J Med 2002;346:1128-1137. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11948273>.
456. Bilimoria KY, Bentrem DJ, Ko CY, et al. Multimodality therapy for pancreatic cancer in the U.S. : utilization, outcomes, and the effect of

hospital volume. Cancer 2007;110:1227-1234. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/17654662>.

457. La Torre M, Nigri G, Ferrari L, et al. Hospital volume, margin status, and long-term survival after pancreaticoduodenectomy for pancreatic adenocarcinoma. Am Surg 2012;78:225-229. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/22369834>.

458. Hyder O, Dodson RM, Nathan H, et al. Influence of patient, physician, and hospital factors on 30-day readmission following pancreatoduodenectomy in the United States. JAMA Surg 2013;148:1095-1102. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/24108580>.

459. Verbeke CS. Resection margins and R1 rates in pancreatic cancer—are we there yet? Histopathology 2008;52:787-796. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18081813>.

460. Tang LH, Berlin J, Branton P, et al. Protocol for the Examination of Specimens from Patients with Carcinoma of the Exocrine Pancreas. 2013. Available at:
http://www.cap.org/apps/docs/committees/cancer/cancer_protocols/2013/PancreasEndo_13protocol_3201.pdf.

461. Gebhardt C, Meyer W, Reichel M, Wunsch PH. Prognostic factors in the operative treatment of ductal pancreatic carcinoma. Langenbecks Arch Surg 2000;385:14-20. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/10664114>.

462. Mitsunaga S, Hasebe T, Iwasaki M, et al. Important prognostic histological parameters for patients with invasive ductal carcinoma of the pancreas. Cancer Sci 2005;96:858-865. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/16367904>.

463. Huebner M, Kendrick M, Reid-Lombardo KM, et al. Number of lymph nodes evaluated: prognostic value in pancreatic adenocarcinoma. J Gastrointest Surg 2012;16:920-926. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/22421988>.

464. Opfermann KJ, Wahlquist AE, Garrett-Mayer E, et al. Adjuvant Radiotherapy and Lymph Node Status for Pancreatic Cancer: Results of a Study From the Surveillance, Epidemiology, and End Results (SEER) Registry Data. Am J Clin Oncol 2014;37:112-116. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/23211221>.

465. Valsangkar NP, Bush DM, Michaelson JS, et al. N0/N1, PNL, or LNR? The Effect of Lymph Node Number on Accurate Survival Prediction in Pancreatic Ductal Adenocarcinoma. J Gastrointest Surg 2013;17:257-266. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/23229885>.

466. Ashfaq A, Pockaj BA, Gray RJ, et al. Nodal Counts and Lymph Node Ratio Impact Survival After Distal Pancreatectomy for Pancreatic Adenocarcinoma. J Gastrointest Surg 2014. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/24916590>.

467. John BJ, Naik P, Ironside A, et al. Redefining the R1 resection for pancreatic ductal adenocarcinoma: tumour lymph nodal burden and lymph node ratio are the only prognostic factors associated with survival. HPB (Oxford) 2013;15:674-680. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/23458477>.

468. Robinson SM, Rahman A, Haugk B, et al. Metastatic lymph node ratio as an important prognostic factor in pancreatic ductal adenocarcinoma. Eur J Surg Oncol 2012;38:333-339. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/22317758>.

469. Shamseddine AI, Mukherji D, Melki C, et al. Lymph node ratio is an independent prognostic factor after resection of periaampullary malignancies: data from a tertiary referral center in the middle East. Am J Clin Oncol 2014;37:13-18. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/23111358>.

470. Wentz SC, Zhao ZG, Shyr Y, et al. Lymph node ratio and preoperative CA 19-9 levels predict overall survival and recurrence-free survival in patients with resected pancreatic adenocarcinoma. World J

Gastrointest Oncol 2012;4:207-215. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/23444312>.

471. Classification of pancreatic cancer (ed 2). Tokyo: Kanehara, Japan Pancreas Society 2003.

472. Campbell F, Foulis AK, Verbeke CC. Dataset for the histopathological reporting of carcinomas of the pancreas, ampulla of Vater and common bile duct. The Royal College of Pathologists 2010. Available at:
<http://www.rcpath.org/Resources/RCPPath/Migrated%20Resources/Documents/D/datasethistopathologicalreportingcarcinomasmay10.pdf>.

473. Hruban RH, Pitman MB, Klimstra DS. Tumors of the Pancreas: Afip Atlas of Tumor Pathology; 4th Series Fascicle 6: American Registry of Pathology; Armed Forces Institutes of Pathology; 2007.

474. Konstantinidis IT, Warshaw AL, Allen JN, et al. Pancreatic ductal adenocarcinoma: is there a survival difference for R1 resections versus locally advanced unresectable tumors? What is a "true" R0 resection? Ann Surg 2013;257:731-736. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/22968073>.

475. Frampton AE, Gall TM, Krell J, et al. Is there a 'margin' for error in pancreatic cancer surgery? Future Oncol 2013;9:31-34. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/23252561>.

476. Gnerlich JL, Luka SR, Deshpande AD, et al. Microscopic margins and patterns of treatment failure in resected pancreatic adenocarcinoma. Arch Surg 2012;147:753-760. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/22911074>.

477. Delperio JR, Bachellier P, Regenet N, et al. Pancreaticoduodenectomy for pancreatic ductal adenocarcinoma: a French multicentre prospective evaluation of resection margins in 150 evaluable specimens. HPB (Oxford) 2014;16:20-33. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/23464850>.

478. Valle JW, Palmer D, Jackson R, et al. Optimal duration and timing of adjuvant chemotherapy after definitive surgery for ductal adenocarcinoma of the pancreas: ongoing lessons from the ESPAC-3 study. J Clin Oncol 2014;32:504-512. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/24419109>.

479. Reni M. Neoadjuvant treatment for resectable pancreatic cancer: time for phase III testing? World J Gastroenterol 2010;16:4883-4887. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20954273>.

480. Araujo RL, Gaujoux S, Huguet F, et al. Does pre-operative chemoradiation for initially unresectable or borderline resectable pancreatic adenocarcinoma increase post-operative morbidity? A case-matched analysis. HPB (Oxford) 2013;15:574-580. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/23458208>.

481. Lim KH, Chung E, Khan A, et al. Neoadjuvant therapy of pancreatic cancer: the emerging paradigm? Oncologist 2012;17:192-200. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22250057>.

482. Dholakia AS, Hacker-Prietz A, Wild AT, et al. Resection of borderline resectable pancreatic cancer after neoadjuvant chemoradiation does not depend on improved radiographic appearance of tumor–vessel relationships. J Radiat On 2013;2:413-425. Available at: <http://citations.springer.com/item?doi=10.1007/s13566-013-0115-6>.

483. Katz MH, Fleming JB, Bhosale P, et al. Response of borderline resectable pancreatic cancer to neoadjuvant therapy is not reflected by radiographic indicators. Cancer 2012;118:5749-5756. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/22605518>.

484. Esnaola NF, Chaudhary UB, O'Brien P, et al. Phase 2 trial of induction gemcitabine, oxaliplatin, and cetuximab followed by selective capecitabine-based chemoradiation in patients with borderline resectable or unresectable locally advanced pancreatic cancer. Int J Radiat Oncol Biol Phys 2014;88:837-844. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/24606850>.

485. Festa V, Andriulli A, Valvano MR, et al. Neoadjuvant chemo-radiotherapy for patients with borderline resectable pancreatic cancer: a meta-analytical evaluation of prospective studies. Jop 2013;14:618-625. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24216547>.
486. Kim EJ, Ben-Josef E, Herman JM, et al. A multi-institutional phase 2 study of neoadjuvant gemcitabine and oxaliplatin with radiation therapy in patients with pancreatic cancer. Cancer 2013;119:2692-2700. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23720019>.
487. Landry J, Catalano PJ, Staley C, et al. Randomized phase II study of gemcitabine plus radiotherapy versus gemcitabine, 5-fluorouracil, and cisplatin followed by radiotherapy and 5-fluorouracil for patients with locally advanced, potentially resectable pancreatic adenocarcinoma. J Surg Oncol 2010;101:587-592. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20461765>.
488. Marti JL, Hochster HS, Hiotis SP, et al. Phase I/II trial of induction chemotherapy followed by concurrent chemoradiotherapy and surgery for locoregionally advanced pancreatic cancer. Ann Surg Oncol 2008;15:3521-3531. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18830756>.
489. Van Buren G, 2nd, Ramanathan RK, Krasinskas AM, et al. Phase II study of induction fixed-dose rate gemcitabine and bevacizumab followed by 30 Gy radiotherapy as preoperative treatment for potentially resectable pancreatic adenocarcinoma. Ann Surg Oncol 2013;20:3787-3793. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23904005>.
490. McClaine RJ, Lowy AM, Sussman JJ, et al. Neoadjuvant therapy may lead to successful surgical resection and improved survival in patients with borderline resectable pancreatic cancer. HPB (Oxford) 2010;12:73-79. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20495649>.
491. Stokes JB, Nolan NJ, Stelow EB, et al. Preoperative Capecitabine and Concurrent Radiation for Borderline Resectable Pancreatic Cancer. Ann Surg Oncol 2011;18:619-627. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21213060>.
492. Laurence JM, Tran PD, Morarji K, et al. A Systematic Review and Meta-analysis of Survival and Surgical Outcomes Following Neoadjuvant Chemoradiotherapy for Pancreatic Cancer. J Gastrointest Surg 2011;15:2059-2069. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21913045>.
493. Christians KK, Tsai S, Mahmoud A, et al. Neoadjuvant FOLFIRINOX for Borderline Resectable Pancreas Cancer: A New Treatment Paradigm? Oncologist 2014;19:266-274. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24569947>.
494. Tinchon C, Hubmann E, Pichler A, et al. Safety and efficacy of neoadjuvant FOLFIRINOX treatment in a series of patients with borderline resectable pancreatic ductal adenocarcinoma. Acta Oncol 2013;52:1231-1233. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23445338>.
495. Artinyan A, Anaya DA, McKenzie S, et al. Neoadjuvant therapy is associated with improved survival in resectable pancreatic adenocarcinoma. Cancer 2011;117:2044-2049. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21523715>.
496. Breslin TM, Hess KR, Harbison DB, et al. Neoadjuvant chemoradiotherapy for adenocarcinoma of the pancreas: treatment variables and survival duration. Ann Surg Oncol 2001;8:123-132. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11258776>.
497. Evans DB, Rich TA, Byrd DR, et al. Preoperative chemoradiation and pancreaticoduodenectomy for adenocarcinoma of the pancreas. Arch Surg 1992;127:1335-1339. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/1359851>.
498. Evans DB, Varadhachary GR, Crane CH, et al. Preoperative gemcitabine-based chemoradiation for patients with resectable

adenocarcinoma of the pancreatic head. J Clin Oncol 2008;26:3496-3502. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18640930>.

499. Hoffman JP, Weese JL, Solin LJ, et al. A pilot study of preoperative chemoradiation for patients with localized adenocarcinoma of the pancreas. Am J Surg 1995;169:71-77. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/7818001>.

500. Hoffman JP, Lipsitz S, Pisansky T, et al. Phase II trial of preoperative radiation therapy and chemotherapy for patients with localized, resectable adenocarcinoma of the pancreas: an Eastern Cooperative Oncology Group Study. J Clin Oncol 1998;16:317-323. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9440759>.

501. Palmer DH, Stocken DD, Hewitt H, et al. A randomized phase 2 trial of neoadjuvant chemotherapy in resectable pancreatic cancer: gemcitabine alone versus gemcitabine combined with cisplatin. Ann Surg Oncol 2007;14:2088-2096. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17453298>.

502. Spitz FR, Abbruzzese JL, Lee JE, et al. Preoperative and postoperative chemoradiation strategies in patients treated with pancreaticoduodenectomy for adenocarcinoma of the pancreas. J Clin Oncol 1997;15:928-937. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9060530>.

503. Talamonti MS, Small W, Mulcahy MF, et al. A multi-institutional phase II trial of preoperative full-dose gemcitabine and concurrent radiation for patients with potentially resectable pancreatic carcinoma. Ann Surg Oncol 2006;13:150-158. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16418882>.

504. Abbott DE, Tzeng CW, Merkow RP, et al. The cost-effectiveness of neoadjuvant chemoradiation is superior to a surgery-first approach in the treatment of pancreatic head adenocarcinoma. Ann Surg Oncol 2013;20 Suppl 3:S500-508. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23397153>.

505. Palta M, Willett C, Czito B. Role of radiation therapy in patients with resectable pancreatic cancer. Oncology (Williston Park) 2011;25:715-721, 727. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21874833>.

506. Takahashi H, Ogawa H, Ohigashi H, et al. Preoperative chemoradiation reduces the risk of pancreatic fistula after distal pancreatectomy for pancreatic adenocarcinoma. Surgery 2011;150:547-556. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21621236>.

507. Andriulli A, Festa V, Botteri E, et al. Neoadjuvant/preoperative gemcitabine for patients with localized pancreatic cancer: a meta-analysis of prospective studies. Ann Surg Oncol 2012;19:1644-1662. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22012027>.

508. Chua TC, Saxena A. Preoperative chemoradiation followed by surgical resection for resectable pancreatic cancer: a review of current results. Surg Oncol 2011;20:e161-168. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21704510>.

509. Pingpank JF, Hoffman JP, Ross EA, et al. Effect of preoperative chemoradiotherapy on surgical margin status of resected adenocarcinoma of the head of the pancreas. J Gastrointest Surg 2001;5:121-130. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11331473>.

510. Golcher H, Brunner TB, Witzigmann H, et al. Neoadjuvant chemoradiation therapy with gemcitabine/cisplatin and surgery versus immediate surgery in resectable pancreatic cancer : Results of the first prospective randomized phase II trial. Strahlenther Onkol 2014. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25252602>.

511. Tachezy M, Gebauer F, Petersen C, et al. Sequential neoadjuvant chemoradiotherapy (CRT) followed by curative surgery vs. primary surgery alone for resectable, non-metastasized pancreatic adenocarcinoma: NEOPA- a randomized multicenter phase III study (NCT01900327, DRKS00003893, ISRCTN82191749). BMC Cancer

2014;14:411. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/24906700>.

512. Furman MJ, Lambert LA, Sullivan ME, Whalen GF. Rational follow-up after curative cancer resection. J Clin Oncol 2013;31:1130-1133. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23358986>.

513. Tzeng CW, Fleming JB, Lee JE, et al. Yield of clinical and radiographic surveillance in patients with resected pancreatic adenocarcinoma following multimodal therapy. HPB (Oxford) 2012;14:365-372. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22568412>.

514. Tzeng CW, Abbott DE, Cantor SB, et al. Frequency and intensity of postoperative surveillance after curative treatment of pancreatic cancer: a cost-effectiveness analysis. Ann Surg Oncol 2013;20:2197-2203. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23408126>.

515. Witkowski ER, Smith JK, Ragulin-Coyne E, et al. Is it worth looking? Abdominal imaging after pancreatic cancer resection: a national study. J Gastrointest Surg 2012;16:121-128. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21972054>.

516. Tempero MA, Berlin J, Ducreux M, et al. Pancreatic cancer treatment and research: an international expert panel discussion. Ann Oncol 2011;22:1500-1506. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21199884>.

517. Katz MH, Wang H, Fleming JB, et al. Long-term survival after multidisciplinary management of resected pancreatic adenocarcinoma. Ann Surg Oncol 2009;16:836-847. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19194760>.

518. Meyers MO, Meszoely IM, Hoffman JP, et al. Is reporting of recurrence data important in pancreatic cancer? Ann Surg Oncol 2004;11:304-309. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14993026>.

519. Arnaoutakis GJ, Rangachari D, Laheru DA, et al. Pulmonary resection for isolated pancreatic adenocarcinoma metastasis: an analysis of outcomes and survival. J Gastrointest Surg 2011;15:1611-1617. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21725701>.

520. House MG, Choti MA. Palliative therapy for pancreatic/biliary cancer. Surg Clin North Am 2005;85:359-371. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15833477>.

521. Soderlund C, Linder S. Covered metal versus plastic stents for malignant common bile duct stenosis: a prospective, randomized, controlled trial. Gastrointest Endosc 2006;63:986-995. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16733114>.

522. Moss AC, Morris E, Mac Mathuna P. Palliative biliary stents for obstructing pancreatic carcinoma. Cochrane Database Syst Rev 2006:CD004200. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16625598>.

523. Kitano M, Yamashita Y, Tanaka K, et al. Covered self-expandable metal stents with an anti-migration system improve patency duration without increased complications compared with uncovered stents for distal biliary obstruction caused by pancreatic carcinoma: a randomized multicenter trial. Am J Gastroenterol 2013;108:1713-1722. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24042190>.

524. Maire F, Hammel P, Ponsot P, et al. Long-term outcome of biliary and duodenal stents in palliative treatment of patients with unresectable adenocarcinoma of the head of pancreas. Am J Gastroenterol 2006;101:735-742. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16635221>.

525. Lillemoe KD, Cameron JL, Hardacre JM, et al. Is prophylactic gastrojejunostomy indicated for unresectable periampullary cancer? A prospective randomized trial. Ann Surg 1999;230:322-328. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10493479>.

526. Van Heek NT, De Castro SM, van Eijck CH, et al. The need for a prophylactic gastrojejunostomy for unresectable periampullary cancer: a prospective randomized multicenter trial with special focus on assessment of quality of life. *Ann Surg* 2003;238:894-902; discussion 902-895. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14631226>.
527. Lillemoe KD, Cameron JL, Kaufman HS, et al. Chemical splanchnicectomy in patients with unresectable pancreatic cancer. A prospective randomized trial. *Ann Surg* 1993;217:447-455; discussion 456-447. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/7683868>.
528. Wyse JM, Carone M, Paquin SC, et al. Randomized, double-blind, controlled trial of early endoscopic ultrasound-guided celiac plexus neurolysis to prevent pain progression in patients with newly diagnosed, painful, inoperable pancreatic cancer. *J Clin Oncol* 2011;29:3541-3546. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21844506>.
529. Wong GY, Schroeder DR, Carns PE, et al. Effect of neurolytic celiac plexus block on pain relief, quality of life, and survival in patients with unresectable pancreatic cancer: a randomized controlled trial. *JAMA* 2004;291:1092-1099. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14996778>.
530. Jeurnink SM, Polinder S, Steyerberg EW, et al. Cost comparison of gastrojejunostomy versus duodenal stent placement for malignant gastric outlet obstruction. *J Gastroenterol* 2010;45:537-543. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20033227>.
531. Jeurnink SM, Steyerberg EW, Hof G, et al. Gastrojejunostomy versus stent placement in patients with malignant gastric outlet obstruction: a comparison in 95 patients. *J Surg Oncol* 2007;96:389-396. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17474082>.
532. Jeurnink SM, van Eijck CH, Steyerberg EW, et al. Stent versus gastrojejunostomy for the palliation of gastric outlet obstruction: a systematic review. *BMC Gastroenterol* 2007;7:18. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17559659>.
533. Gurusamy KS, Kumar S, Davidson BR. Prophylactic gastrojejunostomy for unresectable periampullary carcinoma. *Cochrane Database Syst Rev* 2013;2:CD008533. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23450583>.
534. Gao L, Yang YJ, Xu HY, et al. A randomized clinical trial of nerve block to manage end-stage pancreatic cancerous pain. *Tumour Biol* 2014;35:2297-2301. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24163058>.
535. Zhong W, Yu Z, Zeng JX, et al. Celiac plexus block for treatment of pain associated with pancreatic cancer: a meta-analysis. *Pain Pract* 2014;14:43-51. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23682788>.
536. Dominguez-Munoz JE. Pancreatic enzyme therapy for pancreatic exocrine insufficiency. *Curr Gastroenterol Rep* 2007;9:116-122. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17418056>.
537. Keller J, Laver P. Human pancreatic exocrine response to nutrients in health and disease. *Gut* 2005;54 Suppl 6:vi1-28. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15951527>.
538. Sikkens EC, Cahen DL, Kuipers EJ, Bruno MJ. Pancreatic enzyme replacement therapy in chronic pancreatitis. *Best Pract Res Clin Gastroenterol* 2010;24:337-347. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20510833>.
539. Dominguez-Munoz JE. Pancreatic exocrine insufficiency: diagnosis and treatment. *J Gastroenterol Hepatol* 2011;26 Suppl 2:12-16. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21323992>.
540. Lemaire E, O'Toole D, Sauvanet A, et al. Functional and morphological changes in the pancreatic remnant following pancreaticoduodenectomy with pancreaticogastric anastomosis. *Br J Surg* 2000;87:434-438. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10759738>.

541. Epstein AS, O'Reilly EM. Exocrine pancreas cancer and thromboembolic events: a systematic literature review. J Natl Compr Canc Netw 2012;10:835-846. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22773799>.
542. Khorana AA, Francis CW, Culakova E, et al. Thromboembolism in hospitalized neutropenic cancer patients. J Clin Oncol 2006;24:484-490. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16421425>.
543. Lee AYY, Levine MN, Baker RI, et al. Low-molecular-weight heparin versus a coumarin for the prevention of recurrent venous thromboembolism in patients with cancer. N Engl J Med 2003;349:146-153. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12853587>.
544. Riess H, Pelzer U, Deutschinoff G, et al. A prospective, randomized trial of chemotherapy with or without the low molecular weight heparin (LMWH) enoxaparin in patients (pts) with advanced pancreatic cancer (APC): Results of the CONKO 004 trial [abstract]. J Clin Oncol 2009;27(suppl):LBA4506. Available at: <http://meeting.ascopubs.org/cgi/content/abstract/27/18S/LBA4506?sid=e598f786-51a5-42d1-82a4-08d6f1163f76>.
545. Boyd AD, Brown D, Henrickson C, et al. Screening for depression, sleep-related disturbances, and anxiety in patients with adenocarcinoma of the pancreas: a preliminary study. ScientificWorldJournal 2012;2012:650707. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22666142>.
546. Turaga KK, Malafa MP, Jacobsen PB, et al. Suicide in patients with pancreatic cancer. Cancer 2011;117:642-647. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20824626>.
547. Philip PA, Mooney M, Jaffe D, et al. Consensus report of the national cancer institute clinical trials planning meeting on pancreas cancer treatment. J Clin Oncol 2009;27:5660-5669. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19858397>.
548. Van Laethem JL, Verslype C, Iovanna JL, et al. New strategies and designs in pancreatic cancer research: consensus guidelines report from a European expert panel. Ann Oncol 2012;23:570-576. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21810728>.
549. Tempero MA, Klimstra D, Berlin J, et al. Changing the Way We Do Business: Recommendations to Accelerate Biomarker Development in Pancreatic Cancer. Clin Cancer Res 2013;19:538-540. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23344262>.
550. Ellis LM, Bernstein DS, Voest EE, et al. American society of clinical oncology perspective: raising the bar for clinical trials by defining clinically meaningful outcomes. J Clin Oncol 2014;32:1277-1280. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24638016>.
551. Philip PA, Chansky K, LeBlanc M, et al. Historical controls for metastatic pancreatic cancer: benchmarks for planning and analyzing single-arm phase II trials. Clin Cancer Res 2014;20:4176-4185. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24914040>.
552. Varadhachary GR, Evans DB. Rational study endpoint(s) for preoperative trials in pancreatic cancer: pathologic response rate, margin negative resection, overall survival or 'all of the above'? Ann Surg Oncol 2013;20:3712-3714. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23943023>.
553. Pogue-Geile KL, Chen R, Bronner MP, et al. Palladin mutation causes familial pancreatic cancer and suggests a new cancer mechanism. PLoS Med 2006;3:e516. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17194196>.
554. Wayne JD, Abdalla EK, Wolff RA, et al. Localized adenocarcinoma of the pancreas: the rationale for preoperative chemoradiation. Oncologist 2002;7:34-45. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11854545>.