



US010472427B2

(12) **United States Patent**
Desjarlais et al.(10) **Patent No.: US 10,472,427 B2**(45) **Date of Patent: *Nov. 12, 2019**(54) **HETERODIMERIC PROTEINS**(71) Applicant: **Xencor, Inc.**, Monrovia, CA (US)(72) Inventors: **John Desjarlais**, Pasadena, CA (US);
Gregory Moore, Azusa, CA (US);
Rumana Rashid, Temple City, CA (US); **Matthew Bennett**, Monrovia, CA (US)(73) Assignee: **Xencor, Inc.**, Monrovia, CA (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

This patent is subject to a terminal disclaimer.

(21) Appl. No.: **15/633,629**(22) Filed: **Jun. 26, 2017**(65) **Prior Publication Data**

US 2018/0215834 A1 Aug. 2, 2018

Related U.S. Application Data

(63) Continuation of application No. 14/155,344, filed on Jan. 14, 2014, now Pat. No. 9,701,759.

(60) Provisional application No. 61/780,310, filed on Mar. 13, 2013, provisional application No. 61/778,148, filed on Mar. 12, 2013, provisional application No. 61/764,954, filed on Feb. 14, 2013, provisional application No. 61/752,347, filed on Jan. 14, 2013.

(51) **Int. Cl.****C07K 16/00** (2006.01)**C07K 16/46** (2006.01)**C07K 16/28** (2006.01)(52) **U.S. Cl.**CPC **C07K 16/468** (2013.01); **C07K 16/2803** (2013.01); **C07K 16/2809** (2013.01); **C07K 16/2896** (2013.01)(58) **Field of Classification Search**

CPC C07K 16/00

USPC 424/136.1

See application file for complete search history.

(56) **References Cited****U.S. PATENT DOCUMENTS**3,773,919 A 11/1973 Boswell et al.
4,169,888 A 10/1979 Hanka et al.
4,179,337 A 12/1979 Davis et al.
4,256,746 A 3/1981 Miyashita et al.
4,294,757 A 10/1981 Asai
4,301,144 A 11/1981 Iwashita et al.
4,307,016 A 12/1981 Asai et al.
4,313,946 A 2/1982 Powell et al.
4,315,929 A 2/1982 Freedman et al.
4,322,348 A 3/1982 Asai et al.
4,331,598 A 5/1982 Hasegawa et al.
4,361,650 A 5/1982 Asai et al.
4,362,663 A 12/1982 Kida et al.
4,364,866 A 12/1982 Asai et al.4,364,935 A 12/1982 Kung et al.
4,371,533 A 2/1983 Akimoto et al.
4,424,219 A 1/1984 Hashimoto et al.
4,450,254 A 5/1984 Isley et al.
4,496,689 A 1/1985 Mitra
4,640,835 A 2/1987 Shimizu et al.
4,670,417 A 6/1987 Iwasaki et al.
4,791,192 A 12/1988 Nakagawa et al.
4,880,935 A 11/1989 Thorpe
4,923,990 A 5/1990 Nakano et al.
4,943,533 A 7/1990 Mendelsohn et al.
4,970,198 A 11/1990 Lee et al.
5,053,394 A 10/1991 Ellestad et al.
5,070,092 A 12/1991 Kanda et al.
5,084,468 A 1/1992 Saito et al.
5,101,038 A 3/1992 Nakano et al.
5,122,368 A 6/1992 Greenfield et al.
5,187,186 A 2/1993 Kanda et al.
5,208,020 A 5/1993 Chari et al.
5,264,586 A 11/1993 Nicolaou et al.
5,384,412 A 1/1995 Nicolaou et al.
5,416,064 A 5/1995 Chari et al.
5,475,092 A 12/1995 Chari et al.
5,500,362 A 3/1996 Robinson et al.
5,530,101 A 6/1996 Queen et al.
5,541,087 A 7/1996 Lo et al.
5,550,246 A 8/1996 Nicolaou et al.
5,558,864 A 9/1996 Bendig et al.
5,585,089 A 12/1996 Queen et al.
5,585,097 A 12/1996 Bolt et al.
5,585,499 A 12/1996 Chari et al.
5,622,929 A 4/1997 Willner et al.

(Continued)

FOREIGN PATENT DOCUMENTSEP 0425235 B1 9/1996
EP 1752471 2/2007

(Continued)

OTHER PUBLICATIONS

Muramatsu et al., Production and characterization of an active single-chain variable fragment antibody recognizing CD25., Cancer Lett. Jul. 28, 2005;225(2):225-36. Epub Jan. 23, 2005.

(Continued)

Primary Examiner — Sean E Aeder(74) *Attorney, Agent, or Firm* — Robin M. Silva; Louis T. Nguyen; Morgan, Lewis & Bockius, LLP(57) **ABSTRACT**

The invention provides heterodimeric antibodies comprising a first heavy chain comprising a first Fc domain and a single chain Fv region that binds a first antigen. The heterodimeric antibodies also comprise a second heavy chain comprising a second Fc domain, a first variable heavy chain and a first variable light chain, wherein the first and second Fc domains are different.

14 Claims, 104 Drawing Sheets**Specification includes a Sequence Listing.**

(56)

References Cited

U.S. PATENT DOCUMENTS

5,635,483 A	6/1997	Pettit et al.	2002/0076406 A1	6/2002	Leung
5,641,780 A	6/1997	Amishiro et al.	2002/0103345 A1	8/2002	Zhu
5,663,149 A	9/1997	Pettit et al.	2002/0131968 A1	9/2002	Waldmann et al.
5,677,171 A	10/1997	Hudziak et al.	2003/0003097 A1	1/2003	Reff et al.
5,693,761 A	12/1997	Queen et al.	2003/0017979 A1	1/2003	Mack et al.
5,693,762 A	12/1997	Queen et al.	2003/0091561 A1	5/2003	Van de Winkel
5,703,080 A	12/1997	Nakakura et al.	2003/0157108 A1	8/2003	Presta
5,712,374 A	1/1998	Kuntsmann et al.	2003/0223999 A1	12/2003	Lindhofer
5,714,586 A	2/1998	Kuntsmann et al.	2004/0071696 A1	4/2004	Adams et al.
5,726,044 A	3/1998	Lo et al.	2004/0162411 A1	8/2004	Lanzavecchia
5,731,168 A	3/1998	Carter et al.	2004/0170626 A1	9/2004	Schuurman
5,736,137 A	4/1998	Anderson et al.	2004/0242851 A1	12/2004	Zhu
5,739,116 A	4/1998	Hamann et al.	2005/0114037 A1	5/2005	Desjarlais et al.
5,767,237 A	6/1998	Sakakibara et al.	2005/0136050 A1	6/2005	Kufer et al.
5,767,285 A	6/1998	Hamann et al.	2005/0142133 A1	6/2005	Lazar et al.
5,770,701 A	6/1998	McGahren et al.	2005/0176028 A1	8/2005	Hofmeister et al.
5,770,710 A	6/1998	McGahren et al.	2005/0191702 A1	9/2005	Mack et al.
5,773,001 A	6/1998	Hamann et al.	2005/0238648 A1	10/2005	Jacobs
5,780,588 A	7/1998	Pettit et al.	2005/0238649 A1	10/2005	Doronina
5,807,706 A	9/1998	Carter et al.	2006/0008883 A1	1/2006	Lazar
5,821,333 A	10/1998	Carter et al.	2006/0018897 A1	1/2006	Lee et al.
5,821,337 A	10/1998	Carter et al.	2006/0024298 A1	2/2006	Lazar et al.
5,824,805 A	10/1998	King et al.	2006/0024317 A1	2/2006	Boyd
5,846,545 A	12/1998	Chari et al.	2006/0073142 A1	4/2006	Chan et al.
5,859,205 A	1/1999	Adair et al.	2006/0074008 A1	4/2006	Senter
5,877,291 A	3/1999	Mezes et al.	2006/0115481 A1	6/2006	Lindhofer et al.
5,877,296 A	3/1999	Hamann et al.	2006/0121032 A1	6/2006	Dahiyat et al.
5,891,996 A	4/1999	Mateo de Acosta del Rio et al.	2006/0134105 A1	6/2006	Lazar et al.
5,892,020 A	4/1999	Mezes et al.	2006/0235208 A1	10/2006	Lazar
5,945,311 A	8/1999	Lindhofer et al.	2007/0071675 A1	3/2007	Wu et al.
5,968,509 A	10/1999	Gorman et al.	2007/0105199 A1	5/2007	Yan et al.
6,054,297 A	4/2000	Carter et al.	2007/0123479 A1	5/2007	Kufer et al.
6,071,515 A	6/2000	Mezes et al.	2007/0148170 A1	6/2007	Desjarlais
6,124,431 A	9/2000	Sakakibara et al.	2007/0287170 A1	12/2007	Davis et al.
6,180,370 B1	1/2001	Queen et al.	2008/0044413 A1	2/2008	Hammond et al.
6,214,345 B1	4/2001	Firestone et al.	2008/0050370 A1	2/2008	Glaser et al.
6,235,883 B1	5/2001	Jakobovits et al.	2008/0138335 A1	6/2008	Takahashi et al.
6,329,507 B1	12/2001	Mezes et al.	2008/0213273 A1	9/2008	Burge
6,407,213 B1	6/2002	Carter et al.	2008/0219974 A1	9/2008	Bernett et al.
6,441,163 B1	8/2002	Chari et al.	2008/0242845 A1	10/2008	Lazar et al.
6,455,677 B1	9/2002	Park et al.	2009/0082213 A1	3/2009	Horowitz et al.
6,506,883 B2	1/2003	Meteo de Acosta del Rio et al.	2009/0136485 A1	5/2009	Chu et al.
6,602,684 B1	8/2003	Umana et al.	2009/0163699 A1	6/2009	Desjarlais
6,632,927 B2	10/2003	Adair et al.	2009/0214539 A1	8/2009	Grosmaire et al.
6,706,265 B1	3/2004	Bolt et al.	2009/0252683 A1	10/2009	Kischel et al.
6,716,410 B1	4/2004	Witztum	2009/0252729 A1	10/2009	Farrington et al.
6,723,538 B2	4/2004	Mack et al.	2009/0274692 A1	11/2009	Tan et al.
6,884,869 B2	4/2005	Senter et al.	2009/0311253 A1	12/2009	Ghayur et al.
6,989,452 B2	1/2006	Ng et al.	2009/0317869 A1	12/2009	Alley et al.
7,087,600 B2	8/2006	Ng et al.	2010/0004431 A1	1/2010	Bernett et al.
7,112,324 B1	9/2006	Dorken et al.	2010/0015133 A1	1/2010	Igawa et al.
7,129,261 B2	10/2006	Ng et al.	2010/0080814 A1	4/2010	Desjarlais et al.
7,276,497 B2	10/2007	Chari et al.	2010/0150918 A1	6/2010	Kufer et al.
7,303,749 B1	12/2007	Chari	2010/0174053 A1	7/2010	Johnson et al.
7,368,565 B2	5/2008	Chari et al.	2010/0178298 A1	7/2010	Lindhofer
7,498,302 B2	3/2009	Ng et al.	2010/0183554 A1	7/2010	Mach et al.
7,507,420 B2	3/2009	Ng et al.	2010/0226925 A1	9/2010	Dillon et al.
7,517,903 B2	4/2009	Chen et al.	2010/0239567 A1	9/2010	Esue
7,601,354 B2	10/2009	Chari	2010/0239582 A1	9/2010	Humphreys et al.
7,642,228 B2	1/2010	Carter et al.	2010/0256339 A1	10/2010	Bossenmaier et al.
7,691,962 B2	4/2010	Boyd et al.	2010/0256340 A1	10/2010	Brinkmann et al.
7,695,936 B2	4/2010	Carter et al.	2010/0298542 A1	11/2010	Igawa et al.
7,696,338 B2	4/2010	Neville, Jr. et al.	2010/0322933 A1	12/2010	Lindhofer et al.
7,728,114 B2	6/2010	Mach et al.	2010/0330089 A1	12/2010	Damle et al.
8,063,187 B2	11/2011	Chu et al.	2010/0331527 A1	12/2010	Davis et al.
8,114,967 B2	2/2012	Bhatt et al.	2011/0054151 A1	3/2011	Lazar et al.
8,216,805 B2	7/2012	Carter et al.	2011/0076275 A1	3/2011	Igawa et al.
8,236,308 B2	8/2012	Kischel et al.	2011/0177500 A1	7/2011	Winther et al.
8,309,690 B2	11/2012	Allan et al.	2011/0189178 A1	8/2011	Desjarlais et al.
8,367,805 B2	2/2013	Chamberlain et al.	2011/0189209 A1	8/2011	Neville, Jr. et al.
8,409,568 B2	4/2013	Gao et al.	2011/0201032 A1	8/2011	Zeng et al.
8,592,562 B2	11/2013	Kannan et al.	2011/0217302 A1	9/2011	Odegard et al.
8,637,641 B2	1/2014	Dahiyat et al.	2011/0262439 A1	10/2011	Kufer et al.
8,946,387 B2	2/2015	Koenig et al.	2011/0275787 A1	11/2011	Kufer et al.
2001/0035606 A1	11/2001	Schoen	2011/0293619 A1	12/2011	Kufer et al.
			2012/0028304 A1	2/2012	Dahiyat et al.
			2012/0034228 A1	2/2012	Kufer et al.
			2012/0149876 A1	6/2012	Von Kreudenstein et al.
			2012/0156207 A1	6/2012	Chu et al.

(56)	References Cited			WO	WO2002098883	12/2002	
	U.S. PATENT DOCUMENTS			WO	WO2004010957	2/2004	
				WO	WO2004043493	5/2004	
				WO	WO2004103272	12/2004	
				WO	WO2004106383	12/2004	
	2012/0251531	A1	10/2012	Baehner et al.	WO	WO2005063816	7/2005
	2012/0251541	A1	10/2012	Baurin et al.	WO	WO2005112919	A2 12/2005
	2013/0089541	A1	4/2013	D'Angelo et al.	WO	WO2005118635	12/2005
	2013/0095097	A1	4/2013	Blakenship et al.	WO	WO2006020258	2/2006
	2013/0101586	A1	4/2013	Riegler et al.	WO	WO2006034488	3/2006
	2013/0115208	A1	5/2013	Ho et al.	WO	WO2006036834	4/2006
	2013/0129723	A1	5/2013	Blakenship et al.	WO	WO2006072620	7/2006
	2013/0142793	A1	6/2013	Ledbetter et al.	WO	WO2006110476	A2 10/2006
	2013/0171095	A1	7/2013	Bernett et al.	WO	WO2006106905	12/2006
	2013/0195849	A1	8/2013	Von Kreudenstein et al.	WO	WO2007005612	1/2007
	2013/0209355	A1	8/2013	De Weers et al.	WO	WO2007018431	A2 2/2007
	2013/0267686	A1	10/2013	Brinkmann	WO	WO2007033230	3/2007
	2013/0336981	A1	12/2013	de Kruif et al.	WO	WO2007042261	4/2007
	2014/0024111	A1	1/2014	Kannan et al.	WO	WO2007046006	4/2007
	2014/0072581	A1	3/2014	Dixit et al.	WO	WO2007047829	4/2007
	2014/0086916	A1	3/2014	Zha	WO	WO2007059404	A2 5/2007
	2014/0212435	A1	7/2014	Moore et al.	WO	WO2007062037	5/2007
	2014/0212436	A1	7/2014	Moore et al.	WO	WO2007084342	7/2007
	2014/0249297	A1	9/2014	Lazar et al.	WO	WO2007089149	A2 8/2007
	2014/0288275	A1	9/2014	Moore et al.	WO	WO2007093630	8/2007
	2014/0294759	A1	10/2014	Chu et al.	WO	WO2007098934	9/2007
	2014/0294823	A1	10/2014	Moore et al.	WO	WO2007110205	10/2007
	2014/0294833	A1	10/2014	Desjarlais et al.	WO	WO2007113648	10/2007
	2014/0294835	A1	10/2014	Moore et al.	WO	WO2007147901	12/2007
	2014/0294836	A1	10/2014	Chu et al.	WO	WO20070147901	12/2007
	2014/0302064	A1	10/2014	Moore	WO	WO2008003103	1/2008
	2014/0322217	A1	10/2014	Moore et al.	WO	WO2008003115	1/2008
	2014/0356381	A1	12/2014	Moore et al.	WO	WO2008003116	1/2008
	2014/0363426	A1	12/2014	Moore et al.	WO	WO2008119096	10/2008
	2014/0370013	A1	12/2014	Desjarlais et al.	WO	WO2008119566	10/2008
	2014/0370020	A1	12/2014	Kuramochi et al.	WO	WO2008124858	10/2008
	2014/0377269	A1	12/2014	Mabry et al.	WO	WO2008145142	12/2008
	2014/0377270	A1	12/2014	Moore et al.	WO	WO2008150494	12/2008
	2015/0071948	A1	3/2015	Lazar et al.	WO	WO2009000006	12/2008
	2015/0307629	A1	10/2015	Bernett et al.	WO	WO2009017394	A1 2/2009
	2016/0060360	A1	3/2016	Moore et al.	WO	WO2009017823	2/2009
	2016/0068588	A1	3/2016	Bernett et al.	WO	WO2009030734	3/2009
	2016/0176969	A1	6/2016	Bernett et al.	WO	WO2009032782	3/2009
	2016/0215063	A1	7/2016	Bernett et al.	WO	WO2009086320	7/2009
	2016/0229924	A1	8/2016	Bernett et al.	WO	WO2009089004	7/2009
	2017/0020963	A1	1/2017	Qu et al.	WO	WO2009106096	9/2009
	FOREIGN PATENT DOCUMENTS				WO	WO2009106321	9/2009
					WO	WO2010028796	3/2010
					WO	WO2010033736	3/2010
					WO	WO2010034441	4/2010
	EP	1829895	5/2007		WO	WO2010037835	4/2010
	EP	2006381	12/2008		WO	WO2010042904	4/2010
	EP	2009101	A1 12/2008		WO	WO2010062171	A2 6/2010
	EP	2194066	6/2010		WO	WO2010085682	7/2010
	EP	2202245	A1 6/2010		WO	WO2010106180	9/2010
	EP	2522724	6/2011		WO	WO2010115551	10/2010
EP	2155788	2/2014		WO	WO2010115552	10/2010	
WO	WO8705330	9/1987		WO	WO2010115553	10/2010	
WO	WO9211018	7/1992		WO	WO2010115589	10/2010	
WO	WO9321232	10/1993		WO	WO2010119119	10/2010	
WO	WO9413804	5/1994		WO	WO20100112193	10/2010	
WO	WO9520045	1/1995		WO	WO2010151792	12/2010	
WO	WO9640210	6/1996		WO	WO2010151808	12/2010	
WO	WO96027011	9/1996		WO	WO2011005621	1/2011	
WO	WO98050431	11/1998		WO	WO2011028952	3/2011	
WO	WO199937791	7/1999		WO	WO2011036183	3/2011	
WO	WO99054440	10/1999		WO	WO2011066342	5/2011	
WO	WO99066951	12/1999		WO	WO2011063348	3/2011	
WO	WO200061739	A1 10/2000		WO	WO2011066501	6/2011	
WO	WO200124763	A2 4/2001		WO	WO2011131746	10/2011	
WO	WO200129246	A1 4/2001		WO	WO2011133886	10/2011	
WO	WO200162931	A1 8/2001		WO	WO2011143545	11/2011	
WO	WO200188138	11/2001		WO	WO2012016227	2/2012	
WO	WO2001083525	11/2001		WO	WO2012018687	2/2012	
WO	WO2001090192	11/2001		WO	WO2012032080	3/2012	
WO	WO200216368	2/2002		WO	WO2012058768	5/2012	
WO	WO200230954	A1 4/2002		WO	WO2012062596	5/2012	
WO	WO200231140	A1 4/2002		WO	WO2012107417	8/2012	
WO	WO2002088172	A2 7/2002		WO	WO2012116453	9/2012	
WO	WO2002062850	8/2002					
WO	WO2002083180	10/2002					

(56)

References Cited

FOREIGN PATENT DOCUMENTS

WO	WO2012125495	9/2012
WO	WO2012125850	9/2012
WO	WO2012131555	10/2012
WO	WO2012146394	11/2012
WO	WO2012146628	11/2012
WO	WO2012162067	11/2012
WO	WO2013006544	1/2013
WO	WO2013016714	1/2013
WO	WO2013022855	2/2013
WO	WO2013026833	2/2013
WO	WO2013033008	3/2013
WO	WO2013047748	4/2013
WO	WO2013055809	4/2013
WO	WO2013063702	5/2013
WO	WO2013096828	6/2013
WO	WO2013125667	8/2013
WO	WO2013180201	12/2013
WO	WO2014004586	1/2014
WO	WO2014012085	1/2014
WO	WO2014056783	4/2014
WO	WO2014079000	5/2014
WO	WO2014110601	7/2014
WO	WO2014113510	7/2014
WO	WO2014145806	9/2014
WO	WO2014145907	9/2014
WO	WO2014164553	10/2014
WO	WO2014209804	12/2014
WO	WO2015018528	2/2015
WO	WO2015026892	2/2015
WO	WO2015063339	5/2015
WO	WO2015103072	7/2015
WO	WO2015143079	9/2015
WO	WO2015149077	10/2015
WO	WO2015184207	12/2015
WO	WO2016014984	1/2016
WO	WO2016086186	6/2016
WO	WO2016086189	6/2016
WO	WO2016086196	6/2016
WO	WO2016105450	6/2016
WO	WO2016141387	9/2016
WO	WO2016182751	11/2016

OTHER PUBLICATIONS

Sforzini et al., Targeting of saporin to Hodgkin's lymphoma cells by anti-CD30 and anti-CD25 bispecific antibodies., *Br J Haematol.* Sep. 1998;102(4):1061-8.

U.S. Appl. No. 12/631,508, filed Dec. 4, 2009, Chari et al.

(No Author Name) "A method for making multispecific antibodies having heteromultimeric and common components", *Expert Opinion on Therapeutic Patents*, Genentech, Inc. (1999) 9(6): 785-790, pp. 785-790.

"Polythene Glycol and Derivatives for Advanced PEGylation", Catalog 2005-2006, Nektar Therapeutics.

"Xencor Provides Data Updates on XmaB Bispecific Antibody Program and Announces Presentations at Upcoming American Society of Hematology 2014 Annual Meeting", Nov. 6, 2014, XP055255549, retrieved from the Internet: http://files.shareholder.com/downloads/AMDA-2B2V8N/0x0x792404/77590b72-837a-4085-bc55-78fa500638dc/XNCR_News_2014_11_6_General_Releases.pdf.

Abbott Laboratories, Strategies and Current Approaches for Improving Drug-Like-Properties During Biologics Drug Candidate Selection, AAPS Webinar—Nov. 10, 2011.

Adams, et al., Avidity-Mediated Enhancement of in vivo Tumor Targeting by Single-Chain Fv Dimers, *Clin Cancer Res*, 2006, vol. 12(5), pp. 1599-1605, doi:10.1158/1078-0432.CCR-05-2217.

Alberola-Ila et al., Stimulation Through the TCR/CD3 Complex Up-Regulates the CD2 Surface Expression on Human T Lymphocytes, Feb. 15, 1991.

Alibaud et al., A New Monoclonal Anti-CD3? Antibody Reactive on Paraffin Sections, *Journal of Histochemistry & Cytochemistry*, 2000, vol. 48, p. 1609.

An, et al., IgG2m4, an engineered antibody isotype with reduced Fc function, *mAbs*, 2009, vol. 1, Issue 6, pp. 572-579, www.landesbioscience.com/journals/mabs/article/10185.

Aplin et al., Preparation, properties, and applications of carbohydrate conjugates of proteins and lipids, 1981, *CRC Crit. Rev. Biochem.*, pp. 259-306.

Arnett, et al., Crystal structure of a human CD3- ϵ/δ dimer in complex with a UCHT1 single-chain antibody fragment, *PNAS*, 2004, vol. 101, No. 46, pp. 16268-16273.

Asano, et al., Cytotoxic enhancement of a bispecific diabody (Db) by format conversion to tandem single-chain variable fragment (taFv): The Case of the hEx3 Diabody, *JBC Papers in Press*, 2010, <http://www.jbc.org/cgi/doi/10.1074/jbc.M110.172957>.

Asano, et al., Highly Effective Recombinant Format of a Humanized IgG-like Bispecific Antibody for Cancer Immunotherapy with Retargeting of Lymphocytes to Tumor Cells, *The Journal of Biological Chemistry*, 2007, vol. 282, No. 38, pp. 27659-27665.

Atwell, et al., Stable Heterodimers from Remodeling the Domain Interface of a Homodimer using a Phage Display Library, *J. Mol. Biol.*, 1997, vol. 270, pp. 26-35.

Baca et al., Antibody humanization using monovalent phage display, 1997, *J. Biol. Chem.* 272(16):10678-10684.

Baeuerle, et al., Response to Letter, "Correct TandAb protein," *Molecular Immunology*, 2007, vol. 44, p. 3084.

Baeuerle, et al., Review—Bispecific T-Cell Engaging Antibodies for Cancer Therapy, *Cancer Res*, 2009, vol. 69: (12), pp. 4941-4944.

Barbas, et al. In vitro evolution of a neutralizing human antibody to human immunodeficiency virus type 1 to enhance affinity and broaden strain cross-reactivity, 1994, *Proc. Nat. Acad. Sci. USA* 91:3809-3813.

Bargou et al., Tumor Regression in Cancer Patients by Very Low Doses of a T Cell-Engaging Antibody, *Science*, 2008, vol. 321, pp. 974-977.

Bhatt, Sea Lane—DDD presentation, "Surrobody™—A Novel Approach to Bispecifics . . .," Aug. 8, 2012.

Bibollet-Ruche et al., The Quality of Chimpanzee T-Cell Activation and Simian Immunodeficiency Virus/Human Immunodeficiency Virus Susceptibility Achieved via Antibody-Mediated T-Cell Receptor/CD3 Stimulation Is a Function of the Anti-CD3 Antibody Isotype, *J. Biol. Chem.*, Jul. 30, 2008.

Biochemica, Your apoptosis specialist, 1999, No. 2, pp. 34-37 (Roche Molecular Biochemicals).

Bird et al., Single-chain antigen-binding proteins, 1988, *Science* 242:423-426.

Blumel, et al., Epitope distance to the target cell membrane and antigen size determine the potency of T cell-mediated lysis by BiTE antibodies specific for a large melanoma surface antigen, *Cancer Immunol Immunother.*, 2010, vol. 59(8), pp. 1197-1209.

Borras, et al., Generic Approach for the Generation of Stable Humanized Single-chain Fv Fragments from Rabbit Monoclonal Antibodies, *The Journal of Biological Chemistry*, 2010, vol. 285, No. 12, pp. 9054-9066.

Bortoletto, Nicola et al., "Optimizing anti-CD3 affinity for effective T cell targeting against tumor cells," *Eur J Immunol. Nov.* 2002;32(11):3102-7.

Boswell et al., Effects of Charge on Antibody Tissue Distribution and Pharmacokinetics, 2010, *Bioconjugate Chem.*, 21(21):2153-2163.

Brandi, et al., Bispecific antibody fragments with CD20 3 CD28 specificity allow effective autologous and allogeneic T-cell activation against malignant cells in peripheral blood and bone marrow cultures from patients with B-cell lineage leukemia and lymphoma, *Experimental Hematology*, 1999, vol. 27, pp. 1264-1270.

Brinkmann, et al., presentation slideshow—"Roche Penzberg & Roche Glycart, Schlieren: Centers of Excellence for Recombinant Proteins".

Brinkmann, et al., A recombinant immunotoxin containing a disulfide-stabilized Fv fragment, *Proc. Natl. Acad. Sci. USA*, 1993, vol. 90, pp. 7538-7542.

Cao, et al., Oligomerization is required for the activity of recombinant soluble LOX-1, *FEBS J. Sep.* 2009;276(17):4909-20. doi: 10.1111/j.1742-4658.2009.07190.x. Epub Jul. 31, 2009.

(56)

References Cited

OTHER PUBLICATIONS

- Carpenter, et al., Non-Fc Receptor-Binding Humanized Anti-CD3 Antibodies Induce Apoptosis of Activated Human T Cells, *J. Immunol.*, 2000, vol. 165, No. 11, pp. 6205-6213.
- Carter et al., Antibody-drug conjugates for cancer therapy, 2008, *Cancer J.* 14(3):154-169.
- Carter et al., Humanization of an anti-p185HER2 antibody for human cancer therapy, 1992, *Proc Natl Acad Sci USA* 89:4285-9.
- Castoldi, et al., Molecular characterization of novel trispecific ErbB-cMet-IGF1R antibodies and their antigen-binding properties, *Protein Engineering, Design & Selection*, 2012, vol. 25, No. 10, pp. 551-559.
- Cemerski, et al., Suppression of mast cell degranulation through a dual-targeting tandem IgE-IgG Fc domain biologic engineered to bind with high affinity to FcγRIIb., *Immunol Lett.* Mar. 30, 2012;143(1):34-43. doi: 10.1016/j.imlet.2012.01.008. Epub Jan. 25, 2012.
- Chames et al., Bispecific antibodies for cancer therapy—The light at the end of the tunnel?, *mAbs*, 2009, vol. 1, Issue 6, pp. 1-9.
- Chang, et al., Monoclonal antibodies against oxidized low-density lipoprotein bind to apoptotic cells and inhibit their phagocytosis by elicited macrophages: evidence that oxidation-specific epitopes mediate macrophage recognition., *Proc Natl Acad Sci U S A.* May 25, 1999;96(11):6353-8.
- Chari et al., Immunoconjugates containing novel maytansinoids: promising anticancer drugs, 1992, *Cancer Research* 52: 127-131.
- Chatal, 1989, *Monoclonal Antibodies in Immunoscintigraphy*, CRC Press (Book Abstract).
- Chelius, et al., Structural and functional characterization of the trifunctional antibody catumaxomab, *mAbs*, 2010, vol. 2, Issue 3, pp. 309-319.
- Chothia et al., Canonical structures for the hypervariable regions of immunoglobulins, 1987, *J. Mol. Biol.* 196:901-917.
- Chothia, et al., Structural Determinants in the Sequences of Immunoglobulin Variable Domain, *J. Mol. Biol.*, 1998, vol. 278, pp. 457-479.
- Chu et al., Reduction of total IgE by targeted coengagement of IgE B-cell receptor and FcγRIIb with Fc-engineered antibody., *J Allergy Clin Immunol.* Apr. 2012;129(4):1102-15. doi: 10.1016/j.jaci.2011.11.029. Epub Jan. 16, 2012.
- Conrad, et al., TCR and CD3 Antibody Cross-Reactivity in 44 Species, *Cytometry Part A*, 2007, vol. 71A, pp. 925-933.
- Conrath, et al., Antigen Binding and Solubility Effects upon the Veneering of a Camel VHH in Framework-2 to Mimic a VH, *J. Mol. Biol.*, 2005, vol. 350, pp. 112-125.
- Counterman et al., "Volumes of Individual Amino Acid Residues in Gas-Phase Peptide Ions.", *J. Am. Chem. Soc.*, 1999,121 (16), pp. 4031-4039.
- Cuesta, et al., Multivalent antibodies: when design surpasses evolution, *Trends in Biotechnology*, 2010, vol. 28, No. 7, pp. 355-362, doi:10.1016/j.tibtech.2010.03.007.
- d'Argouges, et al., Combination of rituximab with blinatumomab (MT103/MEDI-538), a T cell-engaging CD19-/CD3-bispecific antibody, for highly efficient lysis of human B lymphoma cells, *Leukemia Research*, 2009, vol. 33, pp. 465-473.
- Davies et al., Expression of GnTIII in recombinant anti-CD20 CHO production cell line: expression of antibodies with altered glycoforms leads to an increase in ADCC through higher affinity for FcγRIII, 2001, *Biotechnol Bioeng* 74:288-294.
- Davila, et al., Efficacy and Toxicity Management of 19-28z CAR T Cell Therapy in B Cell Acute Lymphoblastic Leukemia, *Sci. Transl. Med.*, 2014, vol. 6, Issue 224, pp. 1-10, 224ra25.
- Davis, et al., SEEDbodies: fusion proteins based on strand-exchange engineered domain (SEED) C₄3 heterodimers in an Fc analogue platform for asymmetric binders or immunofusions and bispecific antibodies, *Protein Engineering, Design & Selection*, 2010, vol. 23, No. 4 pp. 195-202.
- De Groot et al., De-Immunization of Therapeutic Proteins by T-Cell Epitope Modification, 2005, *Dev. In Biologicals*, 2005, 122:171-194.
- De Pascalis et al., Grafting of "abbreviated" complementarity-determining regions containing specificity-determining residues essential for ligand contact to engineer a less immunogenic humanized monoclonal antibody, 2002, *J. Immunol.* 169:3076-3084.
- Del Nagro et al., A critical role for complement C3d and the B cell coreceptor (CD19/CD21) complex in the initiation of inflammatory arthritis., *J Immunol.* Oct. 15, 2005;175(8):5379-89.
- Demarest et al., Antibody therapeutics, antibody engineering, and the merits of protein stability, *Current Opin in Drug Discovery & Development*, 2008 11(5): 675-587, Sep. 11, 2008.
- Deyev, et al., Multivalency: the hallmark of antibodies used for optimization of tumor targeting by design, *BioEssays*, 2008, vol. 30, pp. 904-918.
- DiGiammarino et al., Ligand association rates to the inner-variable-domain of a dual-variable-domain immunoglobulin are significantly impacted by linker design, *mAbs*3:5, 1-8; Sep.-Oct.; 3(5):487-94, Landes Bioscience, Sep. 1, 2011.
- Dixon, et al., Activation of Human T Lymphocytes by Crosslinking of Anti-CD3 Monoclonal Antibodies, *Journal of Leukocyte Biology*, 1989, vol. 46, pp. 214-220.
- Dong et al., A stable IgG-like bispecific antibody targeting the epidermal growth factor receptor and the type I insulin-like growth factor receptor demonstrates superior anti-tumor activity, *mAbs* 3:3, May-Jun. 2011: 273-288, May 1, 2011.
- Doronina , Development of potent monoclonal antibody auristatin conjugates for cancer therapy, 2003, *Nat Biotechnol* 21(7):778-784.
- Dreier, et al., Extremely Potent, Rapid and Costimulation-Independent Cytotoxic T-cell Response Against Lymphoma Cells Catalyzed by a Single-Chain Bispecific Antibody, *Int. J. Cancer*, 2002, vol. 100, pp. 690-697.
- Dreier, et al., T Cell Costimulus-Independent and Very Efficacious Inhibition of Tumor Growth in Mice Bearing Subcutaneous or Leukemic Human B Cell Lymphoma Xenografts by a CD19-/CD3-Bispecific Single-Chain Antibody Construct, *The Journal of Immunology*, 2003, vol. 170, pp. 4397-4402.
- Dubowchik et al., Receptor-mediated and enzyme-dependent targeting of cytotoxic anticancer drugs, 1999, *Pharm. Therapeutics* 83:67-123.
- Ducry et al., Antibody-drug conjugates: linking cytotoxic payloads to monoclonal antibodies, 2010, *Bioconjugate Chem.* 21:5-13.
- Dudgeon, et al., General strategy for the generation of human antibody variable domains with increased aggregation resistance, *PNAS Early Edition*, 2012, pp. 10879-10884, www.pnas.org/cgi/doi/10.1073/pnas.1202866109 & Supporting Information.
- Duke, et al., Measurement of apoptosis and other forms of cell death, 2004, *Curr protocols immunol.* 3.17.1-3.17.16.
- DukSin et al., Relationship of the structure and biological activity of the natural homologues of tunicamycin, 1982, *J. Biol. Chem.* 257:3105.
- Duval, et al., A Bispecific Antibody Composed of a Nonneutralizing Antibody to the gp41 Immunodominant Region and an Anti-CD89 Antibody Directs Broad Human Immunodeficiency Virus Destruction by Neutrophils, *Journal of Virology*, 2008, pp. 4671-4674, doi:10.1128/JVI.02499-07.
- Edge et al., Deglycosylation of glycoproteins by trifluoromethanesulfonic acid, 1981, *Anal. Biochem.* 118:131.
- Elliott, et al., Antiparallel Conformation of Knob and Hole Aglycosylated Half-Antibody Homodimers is Mediated by a CH2—CH3 Hydrophobic Interaction, *Journal of Molecular Biology*, 2014, vol. 426, Issue 9, pp. 1947-1957.
- Feldmann et al., Novel Humanized and Highly Efficient Bispecific Antibodies Mediate Killing of Prostate Stem Cell Antigen-Expressing Tumor Cells by CD8+ and CD4+ T cells, Aug. 8, 2012.
- Feldmann et al., Retargeting of T Cells to Prostate Stem Cell Antigen Expressing Tumor Cells: Comparison of Different Antibody Formats, Dec. 28, 2010.
- Fernandes, et al., T Cell Receptors are Structures Capable of Initiating Signaling in the Absence of Large Conformational Rearrangements, *The Journal of Biological Chemistry*, 2012, vol. 287, No. 16, pp. 13324-13335.
- Fischer, Nicolas et al., "Bispecific antibodies: molecules that enable novel therapeutic strategies", 2007, vol. 74, pp. 3-14.

(56)

References Cited

OTHER PUBLICATIONS

- Foreman, et al., ErbB3 Inhibitory Surroboodies Inhibit Tumor Cell Proliferation In Vitro and In Vivo, *Mol Cancer Ther*, 2012, vol. 11(7), pp. 1411-1420.
- Foreman, et al., PEGS poster, "ErbB3 Inhibitory Surroboodies Inhibit Tumor Cell Proliferation In Vitro and In Vivo," 2012.
- Fraker et al., Crystal structure of peptide cyclo-(D-VAL-L-PRO-L-VAL-D-PRO)₃, 1978, *Biochem. Biophys. Res. Commun.* 80(4):849-57.
- Francois, et al., Construction of a Bispecific Antibody Reacting with the α - and β -Chains of the Human IL-2 Receptor, *The Journal of Immunology*, May 15, 1993, vol. 150, No. 10, pp. 4610-4619.
- F-star Modular Antibodies Fact Sheet, Apr. 2008, "Modular Antibody Technology" (w/ reference to Ruker WO 2006/072620 A1).
- F-star Modular Antibodies Press Release, Mar. 28, 2008, "Antibody Engineering Company F-Star Buys Back Royalty Obligations. TVM Capital Joins Investor Syndicate."
- Fudenberg, et al., Serologic Demonstration of Dual Specificity of Rabbit Bivalent Hybrid Antibody, *The Journal of Experimental Medicine*, 1964, vol. 119(1), pp. 151-166.
- Ganesan, et al., Fc γ RIIb on Liver Sinusoidal Endothelium Clears Small Immune Complexes, *The Journal of Immunology*, Nov. 15, 2012, vol. 189 No. 10, pp. 4981-4988.
- GenBank AAA38124.1, immunoglobulin heavy-chain VJ region [Mus musculus] Protein/NCBI.
- GenBank AAA39180.1, immunoglobulin light-chain VJ region [Mus musculus] Protein/NCBI.
- Ghändler et al., One of the CD3 ϵ Subunits within a T Cell Receptor Complex Lies in Close Proximity to the C β FG Loop, *J. Exp. Med.*, 1998, vol. 187, No. 9, pp. 1529-1536.
- Ghetie et al., Multiple roles for the major histocompatibility complex Class I-related receptor FcRn, 2000, *Annu Rev Immunol* 18:739-766.
- Gilliland, et al., Universal bispecific antibody for targeting tumor cells for destruction by cytotoxic T cells, *Proc. Natl. Acad. Sci. USA*, 1988, vol. 85, pp. 7719-7723.
- Gorman et al., Reshaping a therapeutic CD4 antibody, *Proc. Natl. Acad. Sci. USA* 88:4181-4185.
- Grodzki & Bernstein, "Antibody Purification: Ion-Exchange Chromatography," *Methods Mol Biol* 2010 :588:27-32.
- Gunasekaran et al., Enhancing Antibody Fc Heterodimer Formation through Electrostatic Steering Effects, *Journal of Biological Chemistry*, vol. 285, No. 25, pp. 19637-10946, Apr. 16, 2010 & Supplementary Tables.
- Haagen, et al., The Efficacy of CD3 x CD19 Bispecific Monoclonal Antibody (BsAb) in a Clonogenic Assay: The Effect of Repeated Addition of BsAb, and Interleukin-2, *Blood*, 1995, vol. 85, No. 11, pp. 3208-3212.
- Hakimuddin et al., A chemical method for the deglycosylation of proteins, 1987, *Arch. Biochem. Biophys.* 259:52.
- Hamel, et al., The Role of the V_L- and V_H- Segments in the Preferential Reassociation of Immunoglobulin Subunits, *Molecular Immunology*, 1986, vol. 23, No. 5, pp. 503-510.
- HAwkins et al., Selection of phage antibodies by binding affinity mimicking affinity maturation, 1992, *J. Mol. Biol.* 226:889-896.
- Hayden-Ledbetter, et al., CD20-Directed Small Modular Immunopharmaceutical, TRU-015, Depletes Normal and Malignant B Cells, *Clin Cancer Res*, 2009, vol. 15(8), pp. 2739-2746.
- He et al., Humanization and pharmacokinetics of a monoclonal antibody with specificity for both E- and P-selectin, 1998, *J. Immunol.* 160:1029-1035.
- Hennecke et al., "Non-repetitive single-chain Fv linkers selected by selectively infective phage (SIP) technology," *Protein Eng.* May 1998;11(5):405-10.
- Hernandez-Caselles, et al., A study of CD33 (SIGLEC-3) antigen expression and function on activated human T and NK cells: two isoforms of CD33 are generated by alternative splicing, *J. Leukoc. Biol.*, 2006, vol. 79, pp. 46-58.
- Hexham, et al., Influence of relative binding affinity on efficacy in a panel of anti-CD3 scFv immunotoxins, *Molecular Immunology*, 2001, vol. 38, pp. 397-408.
- Hinman et al., Preparation and characterization of monoclonal antibody conjugates of the calicheamicins: a novel and potent family of antitumor antibodies, 1993 *Cancer Res.* 53:3336-3342.
- Hoffmann, et al., Serial killing of tumor cells by cytotoxic T cells redirected with a CD19-/CD3-bispecific single-chain antibody construct, *Int. J. Cancer*, 2005, vol. 115, pp. 98-104.
- Holliger et al., "Diabodies": Small Bivalent and bispecific antibody fragments, 1993, *Proc. Natl. Acad. Sci. U.S.A.* 90:6444-6448.
- Holliger et al., Engineering bispecific antibodies, 1993, *Current Opinion Biotechnol.* 4:446-449.
- Houtenbos, et al., The novel bispecific diabody α CD40/ α CD28 strengthens leukaemic dendritic cell-induced T-cell reactivity, *British Journal of Haematology*, 2008, vol. 142, pp. 273-283.
- Hu et al., Minibody: A novel engineered anti-carcinoembryonic antigen antibody fragment (single-chain Fv-C_H3) which exhibits rapid, high-level targeting of xenografts, 1996, *Cancer Res.* 56:3055-3061.
- Huston et al., Protein engineering antibody binding sites: recovery of specific activity in an anti-digoxin single-chain Fv analogue produced in *Escherichia coli*, 1988, *Proc. Natl. Acad. Sci. U.S.A.* 85:5879-5883.
- Igawa et al., Reduced elimination of IgG antibodies by engineering the variable region, 2010, *PEDS.* 23(5): 385-392.
- Igawa, V_H/V_L interface engineering to promote selective expression and inhibit conformational isomerization of thrombopoietin receptor agonist single-chain diabody, *Protein Engineering, Design & Selection*, 2010, vol. 23, No. 8, pp. 667-677.
- Ishigaki et al., Impact of Plasma Oxidized Low-Density Lipoprotein Removal on Atherosclerosis., *Circulation* 118: 75-83, 2008.
- Jackson et al., In vitro antibody maturation, 1995, *J. Immunol.* 154(7):3310-9.
- Jäger, et al., The Trifunctional Antibody Ertumaxomab Destroys Tumor Cells That Express Low Levels of Human Epidermal Growth Factor Receptor 2, *Cancer Res*, 2009, vol. 69(10), pp. 4270-4276.
- Jefferis et al., Interaction sites on human IgG-Fc for Fc γ R: current models, 2002, *Immunol Lett* 82:57-65.
- Jespers, et al., Crystal Structure of HEL4, a Soluble, Refoldable Human V_H Single Domain with a Germ-line Scaffold, *J. Mol. Biol.*, 2004, vol. 337, pp. 893-903.
- Jimenez, et al., A recombinant, fully human, bispecific antibody neutralizes the biological activities mediated by both vascular endothelial growth factor receptors 2 and 3, *Mol Cancer Ther*, 2005, vol. 4(3), pp. 427-434.
- Jin et al. The Design and Engineering of IgG-Like Bispecific Antibodies., Chapter 9, *Bispecific Antibodies*, pp. 151-169.
- Jin, et al., MetMAb, the One-Armed 5D5 Anti-c-Met Antibody, Inhibits Orthotopic Pancreatic Tumor Growth and Improves Survival, *Cancer Res* 2008, vol. 68, pp. 4360-4368.
- Johnson et al., Anti-tumor activity of CC49-doxorubicin immunoconjugates, 1995, *Anticancer Res.* 15:1387-93.
- Johnson, et al., Effector Cell Recruitment with Novel Fv-based Dual-affinity Re-targeting Protein Leads to Potent Tumor Cytolysis and In Vivo B-cell Depletion, *J. Mol. Biol.*, 2010, vol. 399, pp. 436-449.
- Jones et al., Replacing the complementarity-determining regions in a human antibody with those from a mouse, 1986, *Nature* 321:522-525.
- Jordan et al., Structural understanding of stabilization patterns in engineered . . . bispecific Ig-like antibody molecules, *Proteins* 2009; 77:832-841, Jun. 19, 2009.
- Jung, et al., Design of interchain disulfide bonds in the framework region of the Fv fragment of the monoclonal antibody B3, *Proteins*, 1994, vol. 19(1), pp. 35-47.
- Jung, et al., Target Cell-restricted Triggering of the CD95 (APO-1/Fas) Death Receptor with Bispecific Antibody Fragments, *Cancer Research*, 2001, vol. 61, pp. 1846-1848.
- Jungbluth et al., A monoclonal antibody recognizing human cancers with amplification/overexpression of the human epidermal growth factor receptor, 2003, *Proc Natl Acad Sci U S A.* 100(2):639-44.

(56)

References Cited

OTHER PUBLICATIONS

- Kabat et al., 1991, Sequences of proteins of immunological interest, Department of Health and Human Services, Bethesda, vol. 1, 5th Ed.
- Kakutani et al., Accumulation of LOX-1 ligand in plasma and atherosclerotic lesions of Watanabe heritable hyperlipidemic rabbits: identification by a novel enzyme immunoassay, *Biochem Biophys Res Commun.* Mar. 23, 2001;282(1):180-5.
- Kanakaraj, et al., Simultaneous targeting of TNF and Ang2 with a novel bispecific antibody enhances efficacy in an in vivo model of arthritis, *mAbs*, 2012, vol. 4, Issue 5, pp. 600-613, <http://dx.doi.org/10.4161/mabs.21227> & Supplemental Data.
- Kettleborough et al., Humanization of a mouse monoclonal antibody by CDR-grafting: the importance of framework residues on loop conformation, 1991, *Protein Eng.* 4(7):773-83.
- Keyna, et al., Surrogate Light Chain-Dependent Selection of Ig Heavy Chain V Regions, *J. Immunol.*, 1995, vol. 155, pp. 5536-5542.
- Kiewe, et al., Phase I Trial of the Trifunctional Anti-HER2 x Anti-CD3 Antibody Ertumaxomab in Metastatic Breast Cancer, *Clin Cancer Res.*, 2006, vol. 12(10), pp. 3085-3091.
- Kim et al., "Localization of the site of murine IgG1 molecule that is involved in binding the murine intestinal Fc receptor," *Eur. J. Immunol.*, 24:2429-2434, 1994.
- Kipriyanov, et al., Bispecific CD3 x CD19 Diabody for T Cell-Mediated Lysis of Malignant Human B Cells, *Int. J. Cancer*, 1998, vol. 77, pp. 763-772.
- Kipriyanov, et al., Bispecific Tandem Diabody for Tumor Therapy with Improved Antigen Binding and Pharmacokinetics, *J. Mol. Biol.*, 1999, vol. 293, pp. 41-56.
- Kipriyanov, et al., Effect of Domain Order on the Activity of Bacterially Produced Bispecific Single-chain Fv Antibodies, *J. Mol. Biol.*, 2003, vol. 330, pp. 99-111.
- Kipriyanov, et al., Two amino acid mutations in an anti-human CD3 single chain Fv antibody fragment that affect the yield on bacterial secretion but not the affinity, *Protein Engineering*, 1997, vol. 10, No. 4, pp. 445-453.
- Klein et al., Progression of metastatic human prostate cancer to androgen independence in immunodeficient SDIC mice, 1997, *Nature Medicine* 3: 402-408.
- Klein, et al., Progress in overcoming the chain association issue in bispecific heterodimeric IgG antibodies, *mAbs*, Nov.-Dec. 2012, vol. 4, issue 6, pp. 653-663, doi: 10.4161/mabs.21379, Epub Aug. 27, 2012.
- Klinger, et al., Immunopharmacologic response of patients with B-lineage acute lymphoblastic leukemia to continuous infusion of T cell—engaging CD19/CD3-bispecific BiTE antibody blinatumomab, *Blood*, 2012, vol. 119, No. 26, pp. 6226-6233.
- Koristka, et al., Retargeting of Human Regulatory T Cells by Single-Chain Bispecific Antibodies, *The Journal of Immunology*, 2012, vol. 188, pp. 1551-1558, www.jimmunol.org/cgi/doi/10.4049/jimmunol.1101760.
- Kostelny, et al., Formation of a Bispecific Antibody by the Use of Leucine Zippers, *The Journal of Immunology* 1992, vol. 148, pp. 1547-1553.
- Krauss et al., Specificity grafting of human antibody frameworks selected from a phage display library: generation of a highly stable humanized anti-CD22 single-chain Fv fragment, 2003, *Protein Engineering* 16(10):753-759.
- Krupka, et al., CD33 target validation and sustained depletion of Aml blasts in long-term cultures by the bispecific T-cell—engaging antibody AMG 330, *Blood*, 2014, vol. 123, No. 3, pp. 356-365, Prepublished online Dec. 3, 2013; doi:10.1182/blood-2013-08-523548 & Data Supplement.
- Kung, et al., Monoclonal Antibodies Defining Distinctive Human T Cell Surface Antigens, *Science*, 1979, vol. 206, pp. 347-349.
- Kuppen, peter et al., The development and purification of a bispecific antibody for lymphokine-activated killer cell targeting against the rat colon carcinoma CC531., *Cancer Immunol Immunother.* Jun. 1993;36(6):403-8.
- Labrijn, et al., Efficient generation of stable bispecific IgG1 by controlled Fab-arm exchange, www.pnas.org/cgi/doi/10.1073/pnas.1220145110 & Supporting Information.
- Laszlo et al., Cellular determinants for preclinical activity of a novel CD33/CD3 bispecific T-cell engager (BiTE) antibody, AMG 330, against human AML, *blood* 2014 123: 554-561, Dec. 5, 2013.
- Lau et al., Conjugation of Doxorubicin to monoclonal anti-carcinoembryonic antigen antibody via novel thiol-directed cross-linking reagents, 1995, *Bioorg-Med-Chem.* 3(10):1299-1304.
- Lau et al., Novel doxorubicin-monoconal anti-carcinoembryonic antigen antibody immunoconjugate activity in vitro, 1995, *Bioorg-Med-Chem.* 3(10):1305-12.
- Lazar Declaration, Dec. 27, 2010, pp. 1-4.
- Lewis, et al., Generation of bispecific IgG antibodies by structure-based design of an orthogonal Fab interface, *Nature Biotechnology*, 2014, doi:10.1038/nbt.2797 & Supplemental Information.
- Li, et al., Construction and characterization of a humanized anti-human CD3 monoclonal antibody 12F6 with effective immunoregulation functions, *Immunology*, 2005, vol. 116, pp. 487-498.
- Lindhofer, et al., Preferential Species-Restricted Heavy/Light Chain Pairing in Rat/Mouse. Quadromas: Implications for a Single-Step Purification of Bispecific Antibodies, *The Journal of Immunology*, 1995, vol. 155, pp. 219-225.
- Ling, et al., Interspecies Scaling of Therapeutic Monoclonal Antibodies: Initial Look, *J Clin Pharmacol*, 2009, vol. 49, pp. 1382-1402, doi: 10.1177/0091270009337134.
- Link, et al., Production and Characterization of a Bispecific IgG Capable of Inducing T-Cell-Mediated Lysis of Malignant B Cells, *Blood*, 1993, vol. 81, No. 12, pp. 3343-3349.
- Linke, et al., Catumaxomab, Clinical development and future directions, *mAbs*, 2010, vol. 2, Issue 2, pp. 129-136.
- Little, et al., Letter to the Editor, "Flawed TandAb production," *Molecular Immunology*, 2007, vol. 44, p. 3083.
- Liu et al., Asymmetrical Fc Engineering Greatly Enhances Antibody-dependent Cellular Cytotoxicity (ADCC) Effector Function and Stability of the Modified Antibodies, *J. Biol. Chem.* 2014, 289: 3571-3590, Dec. 5, 2013.
- Liu et al., Eradication of large colon tumor xenografts by targeted delivery of maytansinoids, 1996 *Proc. Natl. Acad. Sci. USA* 93:8618-8623.
- Liu, et al., Crystallization of a Deglycosylated T Cell Receptor (TCR) Complexed with an Anti-TCR Fab Fragment, *The Journal of Biological Chemistry*, 1996, vol. 271, No. 52, pp. 33639-33646.
- Lode et al., Targeted therapy with a novel enediene antibiotic calicheamicins o₁ effectively suppress growth and dissemination of liver metastases in a syngeneic model of murine neuroblastoma, 1998, *Cancer Res.* 58:2928.
- Löffler, et al., A recombinant bispecific single-chain antibody, CD19 x CD3, induces rapid, and high lymphoma-directed cytotoxicity by unstimulated T lymphocytes, *Blood*, 2000, vol. 95, No. 6, pp. 2098-2103.
- Lu, et al., a Fully Human Recombinant IgG-like Bispecific Antibody to Both the Epidermal Growth Factor Receptor and the Insulin-like Growth Factor Receptor for Enhanced Antitumor Activity, *The Journal of Biological Chemistry*, 2005, vol. 280, No. 20, pp. 19665-19672.
- Lu, et al., Di-diabody: a novel tetravalent bispecific antibody molecule by design, *Journal of Immunological Methods*, 2003, vol. 279, pp. 219-232.
- Lu, et al., Fab-scFv fusion protein: an efficient approach to production of bispecific antibody fragments, *Journal of Immunological Methods*, 2002, vol. 267, pp. 213-226.
- Lu, et al., The effect of variable domain orientation and arrangement on the antigen-binding activity of a recombinant human bispecific diabody, *Biochemical and Biophysical Research Communications*, 2004, vol. 318, pp. 507-513.
- Lum, et al., The new face of bispecific antibodies: targeting cancer and much more, *Experimental Hematology*, 2006, vol. 34, pp. 1-6.
- Lutterbues, et al., AACR Poster, "Conversion of Cetuximab, Panitumumab, Trastuzumab and Omalizumab into T Cell-engaging BiTE Antibodies Creates Novel Drug Candidates of High Potency," 2008.

(56)

References Cited

OTHER PUBLICATIONS

- Lutterbuese, et al., T cell-engaging BiTE antibodies specific for EGFR potently eliminate KRAS- and BRAF-mutated colorectal cancer cells, *PNAS Early Edition*, 2010, www.pnas.org/cgi/doi/10.1073/pnas.1000976107 & Supporting Information.
- Ma, et al., Expression and Characterization of a Divalent Chimeric Anti-Human CD3 Single Chain Antibody, *Scand.J.Immunol*, 1996, vol. 43, pp. 134-139.
- Mabry, et al., A dual-targeting PDGFR β /VEGF-A molecule assembled from stable antibody fragments demonstrates anti-angiogenic activity in vitro and in vivo, *mAbs*, 2010, vol. 2, Issue 1, pp. 20-34; www.landesbioscience.com/journals/mabs/article/10498 & Supplemental Information.
- Mabry, et al., Engineering of stable bispecific antibodies targeting IL-17A and IL-23, *Protein Engineering, Design & Selection*, 2009, vol. 23, No. 3, pp. 115-127; doi:10.1093/protein/gzp073 & Supplementary Figures 1-8.
- Mack, et al., A small bispecific antibody construct expressed as a functional single-chain molecule with high tumor cell cytotoxicity, *Proc. Natl. Acad. Sci. USA*, 1995, vol. 92, pp. 7021-7025.
- Mack, et al., Biologic Properties of a Bispecific Single-Chain Antibody Directed Against 17-1A (EpCAM) and CD3—Tumor Cell-Dependent T Cell Stimulation and Cytotoxic Activity, *The Journal of Immunology*, 1997, vol. 158, pp. 3965-3970.
- MacroGenics Factsheet, Dual Affinity Re-Targeting (“DART”) Platform, 2010.
- Mandler et al., Immunoconjugates of geldanamycin and anti-HER2 Monoclonal antibodies: antiproliferative activity on human breast carcinoma cell lines, 2000, *J. Nat. Cancer Inst.* 92(19):1573-1581.
- Mandler et al., Modifications in synthesis strategy improve the yield and efficacy of geldanamycin-herceptin immunoconjugates, 2002, *Bioconjugate Chem.* 13:786-791).
- Mandler et al., Synthesis and evaluation of antiproliferative activity of a geldanamycin-herceptin™ immunoconjugates, 2000, *Bioorganic & Med. Chem. Letters* 10:1025-1028.
- Mandy, et al., Effect of Reduction of Several Disulfide Bonds on the Properties and Recombination of Univalent Fragments of Rabbit Antibody, *The Journal of Biological Chemistry*, 1963, vol. 238, No. 1, pp. 206-213.
- Mandy, et al., Recombination of Univalent Subunits Derived from Rabbit Antibody, *The Journal of Biological Chemistry*, 1961, vol. 236, No. 12, pp. 3221-3226.
- Marks et al., By-passing immunization: building high affinity human antibodies by chain shuffling, 1992, *Biotechnology* 10:779-783.
- Martin, et al., Generation of the Germline Peripheral B Cell Repertoire: VH8IX-A B Cells Are Unable to Complete All Developmental Programs, *J. Immunol.*, 1998, vol. 160, pp. 3748-3758.
- Martinez, et al., Characterization of a novel modification on IgG2 light chain: Evidence for the presence of O-linked mannosylation, *J. Chromatogr. A*, 2007, vol. 1156 pp. 183-187.
- Marvin, Bispecific antibodies for dual-modality cancer therapy: killing two signaling cascades with one stone, *Curr Opin Drug Discov Devel*, 2006, vol. 9(2), pp. 184-193.
- Marvin, et al., Recombinant approaches to IgG-like bispecific antibodies, *Acta Pharmacologica Sinica*, 2005, vol. 26 (6), pp. 649-658.
- Mateo et al, Humanization of a mouse monoclonal antibody that blocks the epidermal growth factor receptor: recovery of antagonistic activity, 1997, *Immunotechnology*, 3(1):71-81.
- McPhee, Engineering human immunodeficiency virus 1 protease heterodimers as macromolecular inhibitors of viral maturation, *Proc. Natl. Acad. Sci. USA*, 1996, vol. 93, pp. 11477-11481.
- Meijer, et al., Isolation of Human Antibody Repertoires with Preservation of the Natural Heavy and Light Chain Pairing, *J. Mol. Biol.*, 2006, vol. 358, pp. 764-772.
- Merchant, et al., An efficient route to human bispecific IgG, *Nature Biotechnology*, 1998, vol. 16, pp. 677-681.
- Mertens, Nico, “Tribodies: Fab-scFv fusion proteins as a platform to create multi-functional pharmaceuticals.”, SpringerLink 2011, 135-149.
- Metz, et al., Bispecific antibody derivatives with restricted binding functionalities that are activated by proteolytic processing, *Protein Engineering, Design & Selection*, 2012, vol. 25, No. 10, pp. 571-580.
- Metz, et al., Bispecific digoxigenin-binding antibodies for targeted payload delivery, *PNAS*, 2011, vol. 108, No. 20, pp. 8194-8199.
- Michaelson et al., Anti-tumor activity of stability-engineered IgG-like bispecific antibodies targeting TRAIL-R2 and LTbetaR, [mAbs 1:2, 128-141; Mar./Apr. 2009]; Mar. 11, 2009.
- Michalk et al., Characterization of a novel single-chain bispecific antibody for retargeting of T cells to tumor cells via the TCR co-receptor CD8., *PLoS One*. Apr. 21, 2014;9(4):e95517. doi: 10.1371/journal.pone.0095517.
- Miller et al., Stability engineering of scFvs for the development of bispecific and multivalent antibodies, *PEDS*, 2010, vol. 23, No. 7, pp. 549-557 & Supplementary Data.
- Miller, biogen idec Stability Engineering and Production of IgG-like Bispecific Antibodies, AAPS National Biotechnology Conference, Jun. 24 to Jun. 27, 2007.
- Milutinovic, et al., Sanford Burnham Medical Research Institute / AACR Poster, #4318, “Development of a novel dual agonist Surboby™ that simultaneously activates both death receptors DR4 and DR5 and induces cancer cell death with high potency”.
- Mimoto et al., Engineered antibody Fc variant with selectively enhanced FcγRIIb binding over both FcγRIIa(R131) and FcγRIIa(H131)., *Protein Eng Des Sel*. Oct. 2013;26(10):589-98. doi: 10.1093/protein/gzt022. Epub Jun. 5, 2013.
- Mimoto, et al., Novel asymmetrically engineered antibody Fc variant with superior FcγR binding affinity and specificity compared with afucosylated Fc variant, *mAbs*, 2013, vol. 5, Issue 2, pp. 229-236.
- Modjtahedi et al, Phase I trial and tumour localization of the anti-EGFR monoclonal antibody ICR62 in head and neck or lung cancer, 1996, *Br J Cancer*, 73(2):228-35.
- Modjtahedi et al, Targeting of cells expressing wild-type EGFR and type-III mutant EGFR (EGFRVIII) by anti-EGFR MaB ICR62: a two-pronged attack for tumor therapy, 2003, *Int J Cancer*, 105(2):273-80.
- Modjtahedi et al., Antitumor activity of combinations of antibodies directed against different epitopes on the extracellular domain of the human EGF receptor, 1993, *J. Cell Biophys.* 1993, 22(1-3):129-46.
- Modjtahedi et al., The human EGF receptor as a target for cancer therapy: six new rat mAbs against the receptor on the breast carcinoma MDA-MB 468, 1993, *Br J Cancer*. 1993, 67(2):247-53.
- Mølhøj, et al., CD19-/CD3-bispecific antibody of the BiTE class is far superior to tandem diabody with respect to redirected tumor cell lysis, *Molecular Immunology* 2007, vol. 44 , pp. 1935-1943.
- Moore et al., Tuning T Cell Affinity Improves Efficacy and Safety of Anti-CD38 x Anti-CD3 Bispecific Antibodies in Monkeys—a Potential Therapy for Multiple Myeloma., 57th ASH Annual Meeting and Exposition (Dec. 5-8, 2015), American Society of Hematology, Orlando, Florida.
- Moore, et al., A novel bispecific antibody format enables simultaneous bivalent and monovalent co-engagement of distinct target antigens., *MABs*. Nov.-Dec. 2011; 3(6): 546-557; Published online Nov. 1, 2011. doi: 10.4161/mabs.3.6.18123.
- Moore, et al., Application of dual affinity retargeting molecules to achieve optimal redirected T-cell killing of B-cell lymphoma, *Blood*, 2011, vol. 117, No. 17, pp. 4542-4551.
- Moretti et al., BEAT® the bispecific challenge: a novel and efficient platform for the expression of bispecific IgGs. *BMC Proceedings* 2013 7(Suppl 6):09.
- Morrison, et al., News and Views: Two heads are better than one, *Nature Biotechnology*, 2007, vol. 25, No. 11, pp. 1233-1234.
- Mosmann, 1983, Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays, *J. Immunol. Methods* 65:55-63.
- Muda, et al., Therapeutic assessment of SEED: a new engineered antibody platform designed to generate mono and bispecific antibodies, *Protein Engineering, Design & Selection*, 2011, vol. 24, No. 5, pp. 447-454.

(56)

References Cited

OTHER PUBLICATIONS

- Murthy et al., Binding of an antagonistic monoclonal antibody to an intact and fragmented EGF-receptor polypeptide, 1987, *Arch Biochem Biophys.* 252(2):549-60.
- Nagorsen, et al., Blinatumomab: A historical perspective, *Pharmacology & Therapeutics*, 2012, vol. 136, pp. 334-342, <http://dx.doi.org/10.1016/j.pharmthera.2012.07.013>.
- Nelson, et al., Point of View: Antibody fragments—Hope and hype, *mAbs*, 2010, vol. 2, Issue 1, pp. 77-83.
- Neville et al., Enhancement of immunotoxin efficacy by acid-cleavable cross-linking agents utilizing diphtheria toxin and toxin mutants, 1989, *Biol. Chem.* 264:14653-14661.
- Nielsen, et al., Human T cells resistant to complement lysis by bivalent antibody can be efficiently lysed by dimers of monovalent antibody, *Blood*, 2002, vol. 100, No. 12, pp. 4067-4073.
- Nisonoff, et al., Letters to the Editors: Recombination of a Mixture of Univalent Antibody Fragments of Different Specificity, *Arch. Biochem. Biophys.*, 1961, pp. 460-462.
- Nisonoff, et al., Quantitative Estimation of the Hybridization of Rabbit Antibodies, *Nature*, 1962, vol. 194, No. 4826, pp. 355-359.
- North, et al., A New Clustering of Antibody CDR Loop Conformations, *J. Mol. Biol.*, 2011, vol. 406, pp. 228-256, doi:10.1016/j.jmb.2010.10.030.
- O'Connor et al., Humanization of an antibody against human protein C and calcium-dependence involving framework residues, 1998, *Protein Eng.* 11:321-8.
- Olafsen, et al., Covalent disulfide-linked anti-CEA diabody allows site-specific conjugation and radiolabeling for tumor targeting applications, *Protein Engineering, Design & Selection*, 2004, vol. 17, No. 1, pp. 21-27.
- Page et al., 1993, *Intermational. Journal of Oncology* 3:473-476.
- Panke, et al., Quantification of cell surface proteins with bispecific antibodies, *Protein Engineering, Design & Selection*, 2013, vol. 26, No. 10, pp. 645-654.
- Pessano, et al., The T3/T cell receptor complex: antigenic distinction between the two 20-kd T3 (T3- δ and T3- ϵ) subunits, *The EMBO Journal*, 1985, vol. 4, No. 2, pp. 337-344.
- Pettit et al., Antineoplastic agents 365. Dolastatin 10 SAR probes, 1998, *Anti-Cancer Drug Design* 13:243-277.
- Pettit et al., Dolastatins 24. Synthesis of (–)-dolastatin 10. I X-ray molecular structure of N,N-dimethylvalyl-valyl-dolaisoleuine tert-butyl ester, 1996, *J. Chem. Soc. Perkin Trans. 1* 5:859-863.
- Pettit et al., Specific activities of dolastatin 10 and peptide derivatives against *Cryptococcus neoformans*, 1998, *Antimicrob. Agents Chemother.* 42(11):2961-2965.
- Pettit et al., Structure-activity studies with chiral isomers and with segments of the antimitotic marine peptide dolastatin 10, 1989, *J. Am. Chem. Soc.* 111:5463-5465.
- Pettit, et al., The dolastatins; 18: Sterospecific synthesis of dolaproine1, 1996, *Synthesis* 719-725.
- Pichler et al., Differences of T-Cell Activation by the Anti-CD3 Antibodies Leu4 and BMA030, Mar. 30, 1987.
- Potapov et al., Protein—Protein Recognition: Juxtaposition of Domain and Interface Cores in Immunoglobulins and Other Sandwich-like Proteins, *J. Mol. Biol.*, 2004, vol. 342, pp. 665-679.
- Presta et al., Humanization of an anti-vascular endothelial growth factor monoclonal antibody for the therapy of solid tumors and other disorders, 1997, *Cancer Res.* 57(20):4593-9.
- Queen et al., A humanized antibody that binds to the interleukin 2 receptor, 1989, *Proc Natl Acad Sci, USA* 86:10029-33.
- Rader et al., A phage display approach for rapid antibody humanization: designed combinatorial V gene libraries, 1998, *Proc. Natl. Acad. Sci. USA* 95: 8910-8915.
- Raghavan et al., Fc receptors and their interactions with immunoglobulins, 1996, *Annu Rev Cell Dev Biol* 12:181-220.
- Rattel, et al., AACR Poster, "Validation of *Cynomolgus* Monkeys as Relevant Species for Safety Assessment of a Novel Human BiTE Antibody Platform for Cancer Therapy," 2010.
- Reddy et al., Elimination of Fc receptor-dependent effector functions of a modified IgG4 monoclonal antibody to human CD4., *J Immunol.* Feb. 15, 2000;164(4):1925-33.
- Reiter et al., Disulfide stabilization of antibody Fv: computer predictions and experimental evaluation, *Protein Eng.*, 1995, vol. 8(12), pp. 1323-31.
- Reiter et al., Engineering interchain disulfide bonds into conserved framework regions of Fv fragments: improved biochemical characteristics of recombinant immunotoxins containing disulfide-stabilized Fv, *Protein Eng.*, 1994, vol. 7(5), pp. 697-704.
- Remington's *Pharmaceutical Sciences* 16th edition, Osol, a. Ed. (1980) (Book Abstract).
- Repp, et al., Combined Fc-protein- and Fc-glyco-engineering of scFv-Fc fusion proteins synergistically enhances CD16a binding but does not further enhance NK-cell mediated ADCC, *Journal of Immunological Methods*, 2011, vol. 373, Issues 1-2, pp. 67-78.
- Ridgway, et al., 'Knobs-into-holes' engineering of antibody CH3 domains for heavy chain heterodimerization, *Protein Engineering*, 1996, vol. 9, No. 7, pp. 617-621.
- Riechmann et al., Reshaping human antibodies for therapy, 1988, *Nature* 332:323-329.
- Riethmuller, Symmetry breaking: bispecific antibodies, the beginnings, and 50 years on, *Cancer Immunity*, 2012, vol. 12, p. 12, pp. 1-7.
- Rodeck et al., Interactions between growth factor receptors and corresponding monoclonal antibodies in human tumors, 1987, *J Cell Biochem.* 35(4):315-20.
- Roguska et al., Humanization of murine monoclonal antibodies through variable domain resurfacing, 1994, *Proc. Natl. Acad. Sci. USA* 91:969-973.
- Roosnek, et al., Triggering T Cells by Otherwise Inert Hybrid Anti-CD3/Antitumor Antibodies Requires Encounter with the Specific Target Cell, *J. Exp. Med.*, 1989, vol. 170, pp. 297-302.
- Roque et al., Antibodies and genetically engineered related molecules: production and purification, 2004, *Biotechnol. Prog.* 20:639-654.
- Rose, et al., Mutation of Y407 in the CH3 domain dramatically alters glycosylation and structure of human IgG, *mAbs*, 2013, vol. 5, Issue 2, pp. 219-228.
- Rose, et al., Quantitative Analysis of the Interaction Strength and Dynamics of Human IgG4 Half Molecules by Native Mass Spectrometry, *Structure*, 2011, vol. 19, pp. 1274-1282.
- Rosok et al., A combinatorial library strategy for the rapid humanization of anticarcinoma BR 96 Fab, 1996, *J. Biol. Chem.* 271(37): 22611-22618.
- Rossi, et al., A new class of bispecific antibodies to redirect T cells for cancer immunotherapy, *mAbs* 2014, vol. 6, Issue 2, pp. 381-391.
- Roux, et al., Structural analysis of the nurse shark (new) antigen receptor (NAR): Molecular convergence of NAR and unusual mammalian immunoglobulins, *Proc. Natl. Acad. Sci. USA*, 1998, vol. 95, pp. 11804-11809.
- Rudnick, et al., Affinity and Avidity in Antibody-Based Tumor Targeting, *Cancer Biotherapy and Radiopharmaceuticals*, 2009, vol. 24, No. 2, pp. 155-161, doi: 10.1089/cbr.2009.0627.
- Röthlisberger, et al., Domain Interactions in the Fab Fragment: A Comparative Evaluation of the Single-chain Fv and Fab Format Engineered with Variable Domains of Different Stability, *J. Mol. Biol.*, 2005, vol. 347, pp. 773-789.
- Salmeron et al., A conformational epitope expressed upon association of CD3-epsilon with either CD3-delta or CD3-gamma is the main target for recognition by anti-CD3 monoclonal antibodies, Nov. 1, 1991.
- Sancho et al., CD3- Surface Expression is Required for CD4-p56lck-mediated Up-regulation of T Cell Antigen Receptor-CD3 Signaling in T Cells, Apr. 16, 1992.
- Schaefer, et al., A Two-in-One Antibody against HER3 and EGFR Has Superior Inhibitory Activity Compared with Monospecific Antibodies, *Cancer Cell*, 2011, vol. 20, pp. 472-486 & Supplemental Information, pp. 1-21.
- Schaefer, et al., Immunoglobulin domain crossover as a generic approach for the production of bispecific IgG antibodies, *PNAS*, 2011, vol. 108, No. 27, pp. 11187-11192.

(56)

References Cited

OTHER PUBLICATIONS

- Schlapsch, et al., Functional humanization of an anti-CD16 Fab fragment: obstacles of switching from murine Λ to human Λ or κ light chains, *Protein Engineering, Design & Selection*, 2009, vol. 22, No. 3, pp. 175-188, doi:10.1093/protein/gzn066.
- Schlereth, et al., Eradication of Tumors from a Human Colon Cancer Cell Line and from Ovarian Cancer Metastases in Immunodeficient Mice by a Single-Chain Ep-CAM-/CD3-Bispecific Antibody Construct, *Cancer Res* 2005, vol. 65(7), pp. 2882-2889.
- Schlereth, et al., T-cell activation and B-cell depletion in chimpanzees treated with a bispecific anti-CD19/anti-CD3 single-chain antibody construct, *Cancer Immunol Immunother*, 2006, vol. 55, pp. 503-514, doi:10.1007/s00262-005-0001-1.
- Schoonjans, et al., Fab Chains as an Efficient Heterodimerization Scaffold for the Production of Recombinant Bispecific and Trispecific Antibody Derivatives, *The Journal of Immunology*, 2000, vol. 165, pp. 7050-7057.
- Schroder et al., *The Peptides*, vol. pp. 76-136, 1965, Academic Press.
- Senter et al, *Proceedings of the American Association for Cancer Research*, 2004, vol. 45, Abstract No. 623.
- Senter, Potent antibody drug conjugates for cancer therapy, 2009, *Current Opin. Chem. Biol.* 13:235.
- Shalaby, et al., Development of Humanized Bispecific Antibodies Reactive with Cytotoxic Lymphocytes and Tumor Cells Overexpressing the HER2 Protooncogene, *J.Exp.Med.*, 1992, vol. 175, pp. 217-225.
- Shan, et al., Characterization of scFv-Ig Constructs Generated from the Anti-CD20 mAb 1F5 Using Linker Peptides of Varying Lengths, *J Immunol*, 1999, vol. 162, pp. 6589-6595.
- Shearman, et al., Construction, Expression and Characterization of Humanized Antibodies Directed Against the Human $\alpha\beta$ T Cell Receptor, *The Journal of Immunology*, 1991, vol. 147, No. 12, pp. 4366-4373.
- Shen, et al., Catumaxomab, a rat/murine hybrid trifunctional bispecific monoclonal antibody for the treatment of cancer, *Curr Opin Mol Ther*, 2008, vol. 10(3), pp. 273-284.
- Shen, et al., Single Variable Domain-IgG Fusion: A Novel Recombinant Approach to Fc Domain-Containing Bispecific Antibodies, *The Journal of Biological Chemistry*, 2006, vol. 281, No. 16, pp. 10706-10714.
- Shields et al., Lack of fucose on human IgG1 N-linked oligosaccharide improves binding to human Fc γ R11 and antibody-dependent cellular toxicity, 2002, *J Biol Chem* 277:26733-26740.
- Shier et al., Identification of functional and structural amino-acid residues by parsimonious mutagenesis, 1995, *Gene* 169:147-155.
- Shinkawa et al., The absence of fucose but not the presence of galactose or bisecting N-acetylglucosamine of human IgG1 complex-type oligosaccharides shows the critical role of enhancing antibody-dependent cellular cytotoxicity, 2003, *J Biol Chem* 278:3466-3473.
- Skehan et al., Identification of functional and structural amino-acid residues by parsimonious mutagenesis, 1990, *J. Natl. Cancer Inst.* 82(13):1107-12.
- Soumyarani et al, Oxidatively modified high density lipoprotein promotes inflammatory response in human monocytes-macrophages by enhanced production of ROS, TNF- α , MMP-9, and MMP-2., *Mol Cell Biochem. Jol.* 2012;366(1-2):277-85. doi: 10.1007/s11010-012-1306-y. Epub Apr. 17, 2012.
- Spies et al., Alternative molecular formats and therapeutic applications for bispecific antibodies., *Mol Immunol.* Jan. 27, 2015. pii: S0161-5890(15)00005X. doi: 10.1016/j.molimm.2015.01.003.
- Spiess, et al., Bispecific antibodies with natural architecture produced by co-culture of bacteria expressing two distinct half-antibodies, *Nature Biotechnology*, 2013, doi:10.1038/nbt.2621 & Supplemental Information.
- Stamova, Unexpected recombinations in single chain bispecific anti-CD3-anti-CD33 antibodies can be avoided by a novel linker module, Oct. 29, 2011.
- Stanfield, et al., Maturation of Shark Single-domain (IgNAR) Antibodies: Evidence for Induced-fit Binding, *J. Mol. Biol.*, 2007, vol. 367, pp. 358-372.
- Stewart, et al., Recombinant CD36 inhibits oxLDL-induced ICAM-1-dependent monocyte adhesion., *Mol Immunol.* Feb. 2006;43(3):255-67.
- Strop, P. et al., Generating Bispecific Human IgG1 and IgG2 Antibodies from Any Antibody Pair, *J. Mol. Biol.*, 2012, doi:10.1016/j.jmb.2012.04.020.
- Tan et al., "Superhumanized" antibodies: reduction of immunogenic potential by complementarity-determining region grafting with human germline sequences: application to an anti-CD28, 2002, *J. Immunol.* 169:1119-1125.
- Tan, Philip, Presentation at PepTalk, Jan. 25, 2013, "Bi-specific ADAPTIR Molecule Targeting CD86 and Delivering Monomeric IL10 to Inhibit Antigen Presenting Cells".
- Tarcsa et al, Chapter 10 Dual-Variable Domain Immunoglobulin (DVD-IgTM) Technology: A Versatile, Novel Format for the Next Generation of Dual-Targeting Biologics, *Bispecific Antibodies* 2011, pp. 171-185, 2011.
- Teachey, et al., Cytokine release syndrome after blinatumomab treatment related to abnormal macrophage activation and ameliorated with cytokine-directed therapy, *Blood*, 2013, vol. 121, No. 26, pp. 5154-5157.
- Tedgui, et al., Cytokines in atherosclerosis: pathogenic and regulatory pathways., *Physiol Rev.* Apr. 2006;86(2):515-81.
- Thompson, et al., An Anti-CD3 Single-chain Immunotoxin with a Truncated Diphtheria Toxin Avoids Inhibition by Pre-existing Antibodies in Human Blood, *J.Biol.Chem.*, 1995, vol. 270, No. 47, pp. 28037-28041.
- Thompson, et al., Improved binding of a bivalent single-chain immunotoxin results in increased efficacy for in vivo T-cell depletion, *Protein Engineering*, 2001, vol. 14, No. 12, pp. 1035-1041.
- Thorne, et al., CD36 is a receptor for oxidized high density lipoprotein: implications for the development of atherosclerosis., *FEBS Lett.* Mar. 20, 2007;581(6):1227-32. Epub Feb. 28, 2007.
- Thorpe et al., New coupling agents for the synthesis of immunotoxins containing a hindered disulfide bond with improved stability in Vivo, 1987, *Cancer Res.* 47:5924-5931.
- Thotakura et al., Enzymatic deglycosylating of glycoproteins, 1987, *Meth. Enzymol.* 138:350.
- Thurman et al., Detection of complement activation using monoclonal antibodies against C3d., *J Clin Invest.* May 2013;123(5):2218-30. doi: 10.1172/JCI65861. Epub Apr. 24, 2013.
- Tomlinson et al., Methods for generating multivalent and bispecific antibody fragments, 2000, *Methods Enzymol.* 326:461-479.
- Topp, et al., Targeted Therapy With the T-Cell—Engaging Antibody Blinatumomab of Chemotherapy-Refractory Minimal Residual Disease in B-Lineage Acute Lymphoblastic Leukemia Patients Results in High Response Rate and Prolonged Leukemia-Free Survival, *J Clin Oncol* vol. 29, No. 18, pp. 2493-2498.
- Trautnecker, et al., Bispecific single chain molecules (Janusins) target cytotoxic lymphocytes on Hiv infected cells, *The EMBO Journal*, 1991, vol. 1, No. 12, pp. 3655-3659.
- Tsurushita et al., Humanization of monoclonal antibodies, 2004, *Molecular Biology of B Cells* 533-545.
- Umaña et al., Engineered glycoforms of an antineuro-blastoma IgG1 with optimized antibody-dependent cellular cytotoxic activity, 1999, *Nat Biotechnol* 17:176-180.
- Valliere-Douglass, et al., O-Fucosylation of an antibody light chain: Characterization of a modification occurring on an IgG1 molecule, *Glycobiology*, 2009, vol. 19, No. 2, pp. 144-152, doi:10.1093/glycob/cwn116.
- van Boxel, et al., Some lessons from the systematic production and structural analysis of soluble $\alpha\beta$ T-cell receptors, *Journal of Immunological Methods*, 2009, vol. 350, pp. 14-21.
- van Wauwe, et al., OKT3: A Monoclonal Anti-Human T Lymphocyte Antibody with Potent Mitogenic Properties, *The Journal of Immunology*, 1980, vol. 124, No. 6, pp. 2708-2713.
- Verdier, et al., Determination of lymphocyte subsets and cytokine levels in Cynomolgus monkeys, *Toxicology*, 1995, vol. 105, pp. 81-90.
- Verhoeven et al., Reshaping human antibodies: grafting an antilysozyme activity, 1988, *Science*, 239:1534-1536.

(56)

References Cited

OTHER PUBLICATIONS

- Veri, et al., Therapeutic Control of B Cell Activation via Recruitment of Fcγ Receptor IIb (CD32B) Inhibitory Function With a Novel Bispecific Antibody Scaffold, *Arthritis & Rheumatism*, 2010, vol. 62, No. 7, pp. 1933-1943.
- Vettermann, et al., Powered by pairing: The surrogate light chain amplifies immunoglobulin heavy chain signaling and pre-selects the antibody repertoire, *Seminars in Immunology* 18,2006, pp. 44-55.
- Von Kreudenstein, et al., Improving biophysical properties of a bispecific antibody scaffold to aid developability: Quality by molecular design, *mAbs*, 2013, vol. 5, Issue 5, pp. 1-9, <http://dx.doi.org/10.4161/mabs.25632> & Supplemental Material.
- Wang et al., Conserved amino acid networks involved in antibody variable domain interactions, *Proteins*, 2009, vol. 76, pp. 99-114.
- Wang et al., Expression and characterization of recombinant soluble monkey CD3 molecules: mapping the FN18 polymorphic epitope, *Molecular Immunology*, 2004, vol. 40, pp. 1179-1188.
- Wang, et al., A block in both early T lymphocyte and natural killer cell development in transgenic mice with high-copy numbers of the human CD3E gene, *Proc. Natl. Acad. Sci. USA*, 1994, vol. 91, pp. 9402-9406.
- Ward, et al., Protein Engineering of Homodimeric Tyrosyl-tRNA Synthetase to Produce Active Heterodimers, *The Journal of Biological Chemistry*, 1986, vol. 261, No. 21, pp. 9576-9578.
- Wawrzyniczak et al., Methods for preparing immunotoxins: Effect of the linkage on activity and stability. In *Immunoconjugates. Antibody Conjugates in Radio imaging and Therapy of Cancer*. (C.-W. Vogel, editor). New York, Oxford University Press, pp. 28-55.
- Weatherill, et al., Towards a universal disulphide stabilised single chain Fv format: importance of interchain disulphide bond location and vL-vH orientation, *Protein Engineering, Design & Selection*, 2012, vol. 25, No. 7, pp. 321-329.
- Weiner, et al., The Role of T Cell Activation Bispecific Antibody Therapy in Anti-CD3 X Antitumor, *Journal of Immunology*, 1994, vol. 152, pp. 2385-2392.
- Wesolowski, et al., Single domain antibodies: promising experimental and therapeutic tools in infection and immunity, *Med Microbiol Immunol*, 2009, vol. 198, pp. 157-174.
- Whitlow, et al., An improved linker for single-chain Fv with reduced aggregation and enhanced proteolytic stability, *Protein Engineering*, 1993, vol. 6, No. 8, pp. 989-995.
- Wong, et al., The Mechanism of Anti-CD3 Monoclonal Antibodies, *Transplantation*, 1990, vol. 50, No. 4, pp. 683-689.
- Woods, et al., LC-MS characterization and purity assessment of a prototype bispecific antibody, *mAbs*, 2013, vol. 5, Issue 5, pp. 711-722, <http://dx.doi.org/10.4161/mabs.25488>.
- Woyke et al., In vitro activities and postantifungal effects of the potent dolastatin 10 derivative auristatin PHE, 2001, *Antimicrob. Agents and Chemother.* 45(12):3580-3584.
- Wu et al., Molecular construction and optimization of anti-human IL-11a/β variable domain immunoglobulin (DVD-IgTM) molecules, [*mAbs* 1:4, 339-347; Jul./Aug. 2009]; Landes Bioscience, Apr. 10, 2009.
- Wu et al., Simultaneous targeting of multiple disease mediators by a dual-variable-domain immunoglobulin, (DVD-IgTM) molecules, *Jul.-Aug. 2009*; 339-347, Oct. 14, 2007.
- Wu et al., Humanization of a murine monoclonal antibody by simultaneous optimization of framework and CDR residues, 1999, *J. Mol. Biol.* 294:151-162.
- Wu, et al., Multimerization of a chimeric anti-CD20 single-chain Fv-Fc fusion protein is mediated through variable domain exchange, *Protein Engineering*, 2001, vol. 14, No. 12, pp. 1025-1033.
- Wucherpfennig, et al., Structural Biology of the T-cell Receptor: Insights into Receptor Assembly, Ligand Recognition, and Initiation of Signaling, *Cold Spring Harb Perspect Biol* 2010;2:a005140.
- Xie, et al., A new format of bispecific antibody: highly efficient heterodimerization, expression and tumor cell lysis, *Journal of Immunological Methods*, 2005, vol. 296, pp. 95-101, doi:10.1016/j.jim.2004.11.005.
- Xu, et al., Combinatorial surrobody libraries, *PNAS*, 2008, vol. 105, No. 31, pp. 10756-10761.
- Xu, et al., Rapid optimization and prototyping for therapeutic antibody-like molecules, *mAbs*, 2013, vol. 5, Issue 2, pp. 237-254.
- Xu, et al., Surrobody with Functional Tails, *J. Mol. Biol.*, 2010, vol. 397, pp. 352-360.
- Yang et al., Differential in vitro activation of CD8-CD4+ and CD4-CD8+ T lymphocytes by combinations of anti-CD2 and anti-CD3 antibodies, *Apr. 1, 1988*.
- Yelton et al., Affinity maturation of the BR96 anti-carcinoma antibody by codon-based mutagenesis, 1995, *J. Immunol.* 155:1994-2004.
- Yeung, et al., Engineering human IgG1 affinity to human neonatal Fc receptor: impact of affinity improvement on pharmacokinetics in primates, *J Immunol.* Jun. 15, 2009;182(12):7663-71. doi: 10.4049/jimmunol.0804182.
- Yoshino et al., Upgrading of flow cytometric analysis for absolute counts, cytokines and other antigenic molecules of cynomolgus monkeys (*Macaca fascicularis*) by using anti-human cross-reactive antibodies, *Exp. Anim.*, 2000, vol. 49(2), pp. 97-100.
- Yu et al., The biosynthetic gene cluster of the maytansinoids antitumor agent ansamitocin from *actinosynnema pretiosum*, 2002, *PNAS* 99:7968-7973.
- Zamyatin AA., *Amino acid, peptide, and protein volume in solution.*, *Annu Rev Biophys Bioeng.* 1984;13:145-65.
- Zeidler, et al., The Fc-region of a new class of intact bispecific antibody mediates activation of accessory cells and NK cells and induces direct phagocytosis of tumour cells, *Br J Cancer*, 2000, vol. 83(2), pp. 261-266.
- Ziebig et al., Effect of the oxLDL Binding Protein Fc-CD68 on Plaque Extension and Vulnerability in Atherosclerosis., *Circulation Research* 108: 695-703, 2011.
- Zhu, et al., Identification of Heavy Chain Residues in a Humanized Anti-CD3 Antibody Important for Efficient Antigen Binding and T Cell Activation, *The Journal of Immunology*, 1995, vol. 155, pp. 1903-1910.
- Zhu, et al., Remodeling domain interfaces to enhance heterodimer formation, *Protein Science*, 1997, vol. 6, pp. 781-788.
- Zuo, et al., An efficient route to the production of an IgG-like bispecific antibody, *Protein Engineering*, 2000, vol. 13, No. 5, pp. 361-367.
- Szymkowski et al., Creating the next generation of protein therapeutics through rational drug design, *Current opinion in drug discovery & development*, Sep. 1, 2005, p. 590, XP055354917, England.
- Tang et al., Selection of linkers for a catalytic single-chain antibody using phage display technology., *Journal of Biological Chemistry*, American Society for Biochemistry and Molecular Biology, US, vol. 271, No. 26, Jan. 1, 1996, pp. 15682-19258.
- Chu et al., Inhibition of B cell receptor-mediated activation of primary human B cells by coengagement of CD19 and FcγRIIb with Fc-engineered antibodies., *Mol Immunol.* Sep. 2008;45(15):3926-33. doi: 10.1016/j.molimm.2008.06.027. Epub Aug. 8, 2008.
- Wigginton et al., An immunoglobulin E-reactive chimeric human immunoglobulin G1 anti-idiotypic inhibits basophil degranulation through cross-linking of FcεR1 with FcγRIIb., *Clinical & Experimental Allergy*, 38: 313-319.
- Zalevsky et al., "Enhanced antibody half-life improves in vivo activity." *Nature Biotechnology*, vol. 28, No. 2, Feb. 1, 2010, pp. 157-159.
- Kim et al., Mutational approaches to improve the biophysical properties of human single-domain antibodies., *Biochim Biophys Acta.* Nov. 2014;1844(11):1983-2001. doi: 10.1016/j.bbapap.2014.07.008. Epub Jul. 24, 2014.
- Krah et al., "Single-domain antibodies for biomedical applications.", *Immunopharmacol Immunotoxicol.* 2016;38(1):21-8. doi: 10.3109/08923973.2015.1102934. Epub Nov. 9, 2015.
- Chichili et al., A CD3xCD123 bispecific DART for redirecting host T cells to myelogenous leukemia: preclinical activity and safety in nonhuman primates., *Sci Transl Med.* May 27, 2015;7(289):289ra82. doi: 10.1126/scitranslmed.aaa5693.

(56)

References Cited**OTHER PUBLICATIONS**

Chu et al., Immunotherapy with Long-Lived Anti-CD123 × Anti-CD3 Bispecific Antibodies Stimulates Potent T Cell Mediated Killing of Human AML Cell Lines and of CD123+ Cells in Monkeys: A Potential Therapy for Acute Myelogenous Leukemia, *Blood* 2014, 124:2316.

Terry M., "FDA Places Clinical Hold on AML Drug Co-Developed by Johnson & Johnson (JNJ) and Genmab A/S (Gen Co.)", Biospace 2016, Retrieved from the internet: <https://www.biospace.com/article/fda-places-clinical-hold-on-aml-drug-co-developed-by-johnson-and-johnson-and-genmab-a-s/>.

Kharmate et al., Inhibition of tumor promoting signals by activation of SSTR2 and opioid receptors in human breast cancer cells., *Cancer Cell Int. Sep. 23, 2013;13(1):93*. doi: 10.1186/1475-2867-13-93.

Ott et al., CTLA-4 and PD-1/PD-L1 blockade: new immunotherapeutic modalities with durable clinical benefit in melanoma patients., *Clin Cancer Res. Oct. 1, 2013;19(19):5300-9*. doi: 10.1158/1078-0432.CCR-13-0143.

Spranger et al., Mechanism of tumor rejection with doublets of CTLA-4, PD-1/PD-L1, or IDO blockade involves restored IL-2 production and proliferation of CD8(+) T cells directly within the tumor microenvironment., *J Immunother Cancer. Feb. 18, 2014;2:3*. doi: 10.1186/2051-1426-2-3. eCollection 2014.

Bernett et al., Multiple Bispecific Checkpoint Combinations Promote T cell activation., Nov. 11, 2016, retrieved from the internet: http://files.shareholder.com/downloads/AMDA-2B2V8N/0x0x916283/67AE1A8B-40E8-4316-9F79-384D06B2C395/XNCR_SITC_2016_PD1xCTLA4_Poster126_12Nov2016.pdf.

Hedvat et al., Dual Blockade of PD-1 and CTLA-4 with Bispecific Antibodies Promotes Human T cell Activation and Proliferation., Nov. 11, 2016, retrieved from the internet: http://files.shareholder.com/downloads/AMDA-2B2V8N/0x0x916284/D8084990-61EC-4DFE-81376-60CF58B8C06F/CPI_bispecifics.pdf.

Chichili et al., Co-targeting of PD-1 and CTLA-4 Inhibitory Pathways with Bispecific DART® and TRIDENT™ Molecules., Apr. 4, 2017, retrieved from the internet: http://files.shareholder.com/downloads/AMDA-278VRP/0x0x935572/8CC86417-40BA-41C0-935D-EF1B7DBOB5BB/AACR_2017_-Co-targeting_PD-1_and_CTLA-4_Inhibitory_Pathways_with_DART_and_TRIDENT_Molecules.pdf.

Smith et al., Mouse model recapitulating human Fcγ receptor structural and functional diversity., *Proc Natl Acad Sci U S A. Apr. 17, 2012;109(16):6181-6*. doi: 10.1073/pnas.1203954109. Epub Apr. 2, 2012.

Tabrizi et al., Biodistribution mechanisms of therapeutic monoclonal antibodies in health and disease., *AAPS J. Mar. 2010;12(1):33-43*. doi: 10.1208/s12248-009-9157-5. Epub Nov. 19, 2009.

DiGiandomenico et al., A multifunctional bispecific antibody protects against *Pseudomonas aeruginosa*., *Sci Transl Med. Nov. 12, 2014;6(262):262ra155*. doi: 10.1126/scitranslmed.3009655.




Figure 1C

Figure 1B

Figure 1A




Figure 2E

Figure 2D

Figure 2C

Figure 2B

Figure 2A





Figure 4A

Preferred steric variants that favor Fc heterodimerization.

Monomer 1	Monomer 2
F405A	T394F
S364D	Y349K
S364E	L368K
S364E	Y349K
S364F	K370G
S364H	Y349K
S364H	Y349T
S364Y	K370G
T411K	K370E
V397S/F405A	T394F
K370R/T411K	K370E/T411E
L351E/S364D	Y349K/L351K
L351E/S364E	Y349K/L351K
L351E/T366D	L351K/T366K
P395T/V397S/F405A	T394F
S364D/K370G	S364Y/K370R
S364D/T394F	Y349K/F405A
S364E/F405A	Y349K/T394F
S364E/F405S	Y349K/T394Y
S364E/T411E	Y349K/D401K
S364H/D401K	Y349T/T411E
S364H/F405A	Y349T/T394F
S364H/T394F	Y349T/F405A
Y349C/S364E	Y349K/S354C
L351E/S364D/F405A	Y349K/L351K/T394F
L351K/S364H/D401K	Y349T/L351E/T411E
S364E/T411E/F405A	Y349K/T394F/D401K
S364H/D401K/F405A	Y349T/T394F/T411E
S364H/F405A/T411E	Y349T/T394F/D401K
Y349T	S364H
T394F	F405A
Y349T/T394F	S364H/F405A
K370E	T411K
K370E/T411D	T411K
K370E/T411E	K370R/T411K
L368E/K409E	L368K
Y349T/T411E	S364H/D401K
Y349T/T394F/S354C	S364H/F405A/Y349C
T411E	D401K
T411E	D401R/T411R
Q347E/K360E	Q347R
L368E	S364K
L368E/K370S	S364K
L368E/K370T	S364K
L368E/D401R	S364K
L368E/D401N	S364K
L368E	E357S/S364K
L368E	S364K/K409E
L368E	S364K/K409V
L368D	S364K

Figure 4B

Specifically preferred steric variants that favor Fc heterodimerization.

Variant 1	Variant 2
F405A	T394F
S364D	Y349K
S364E	Y349K
S364H	Y349T
L351K	L351E
D401K	T411E
S364D/T394F	Y349K/F405A
S364E/F405A	Y349K/T394F
S364H/D401K	Y349T/T411E
S364H/F405A	Y349T/T394F
S364H/T394F	Y349T/F405A
L351K/S364H/D401K	Y349T/L351E/T411E
S364H/D401K/F405A	Y349T/T394F/T411E
S364H/F405A/T411E	Y349T/T394F/D401K
Y349T	S364H
T394F	F405A
Y349T/T394F	S364H/F405A
K370E	T411K
K370E/T411D	T411K
K370E/T411E	K370R/T411K
L368E/K409E	L368K
Y349T/T411E	S364H/D401K
Y349T/T394F/S354C	S364H/F405A/Y349C
T411E	D401K
T411E	D401R/T411R
Q347E/K360E	Q347R
L368E	S364K
L368E/K370S	S364K
L368E/K370T	S364K
L368E/D401R	S364K
L368E/D401N	S364K
L368E	E357S/S364K
L368E	S364K/K409E
L368E	S364K/K409V
L368D	S364K

Figure 5A

PREFERRED HETERODIMERIZATION VARIANTS

Fc monomer 1 substitutions	Fc monomer 2 substitutions
Y407T	T366Y
F405A	T394W
T366Y/F405A	T394W/Y407T
Y407A	T366W
T366S/L368A/Y407V	T366W
T366S/L368A/Y407V/Y349C	T366W/S354C
K392D/K409D	E356K/D399K
K370D/K392D/K409D	E356K/E357K/D399K
Y349T	S364H
T394F	F405A
Y349T/T394F	S364H/F405A
K370E	T411K
K370E/T411D	T411K
K370E/T411E	K370R/T411K
L368E/K409E	L368K
Y349T/T411E	S364H/D401K
Y349T/T394F/S354C	S364H/F405A/Y349C
T411E	D401K
T411E	D401R/T411R
Q347E/K360E	Q347R
L368E	S364K
I199T/N203D/K247Q/R355Q/N384S/ K392N/V397M/Q419E/K447_ (deletion of K447)	Q196K/I199T/P217R/P228R/N276K
I199T/N203D/K247Q/R355Q/N384S/K3 92N/V397M/Q419E/K447_	Q196K/I199T/N276K
N384S/K392N/V397M/Q419E	N276K
D221E/P228E/L368E	D221R/P228R/K409R
C220E/P228E/L368E	C220R/E224R/P228R/K409R
F405L	K409R
T366I/K392M/T394W	F405A/Y407V
T366V/K409F	L351Y/Y407A
T366A/K392E/K409F/T411E	D399R/S400R/Y407A
Y349T	S364H
T394F	F405A
Y349T/T394F	S364H/F405A
K370E	T411K
K370E/T411D	T411K
K370E/T411E	K370R/T411K
L368E/K409E	L368K
Y349T/T411E	S364H/D401K
Y349T/T394F/S354C	S364H/F405A/Y349C
T411E	D401K
T411E	D401R/T411R

Figure 5B

Q347E/K360E	Q347R
L368E	S364K
L368E/K370S	S364K
L368E/K370T	S364K
L368E/D401R	S364K
L368E/D401N	S364K
L368E	E357S/S364K
L368E	S364K/K409E
L368E	S364K/K409V
L368D	S364K

Figure 6

Fc monomer 1 substitutions	Fc monomer 2 substitutions
I199T/N203D/K247Q/R355Q/N384S/K392N/V397M/Q419E/K447_	Q196K/I199T/P217R/P228R/N276K
I199T/N203D/K247Q/R355Q/N384S/K392N/V397M/Q419E/K447_	Q196K/I199T/N276K
N203D/K247Q/R355Q/N384S/K392N/V397M/Q419E/K447_	Q196K/P217R/P228R/N276K
N203D/K247Q/R355Q/N384S/K392N/V397M/Q419E/K447_	Q196K/N276K
N203D/K247Q/R355Q/N384S/K392N/Q419E/K447_	Q196K/P217R/P228R/N276K
N203D/K247Q/R355Q/N384S/K392N/Q419E/K447_	Q196K/N276K

Figure 7A

XENP11874 HC-Fab SEQ ID NO: 441

EVQLVESGGGLVKPGGSLKLSAASGYTFTSYVMHWVRQAPGKGLEWIGYINPYNDGTYNEKFQGRVTISS
DKSISTAYMELSSLRSEDTAMYYCARGTYYYGTRVFDYWGGQGLTVTVSSASTKGPSVFPLAPSSKSTSGGTA
ALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVD
KKVEPKSCDKTHTCPPCPAPELLRGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEV
HNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKARPAIEKTISKAKGQPREPQVYTLPPSRE
EMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFPSCSV
MHEALHNHYTQKSLSLSPGK

XENP11874 HC-scFv SEQ ID NO: 442

EVQLVESGGGLVQPGGSLRLSAAAGYTFNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRFTI
SRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLTVTVSSGGGGSGGGSGGGGSQAVV
TQEPSTLVSPGGTPTLTGSSSTGAVTTSNYANWVQQKPGQAPRGLIGGTNKRAPGVPARFSGSLLGGKAALT
LSGAQPEDEAEYYCALWYSNLWVFGGGTKLTVLERKSSDKTHTCPCPAPELLRGPSVFLFPPKPKDTLMIS
RTPEVTCVVVDVSHEDPEVKFKWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNK
ARPAIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLD
SDGSFFLYSKLTVDKSRWQQGNVFPSCSVMHEALHNHYTQKSLSLSPGK

XENP11874 LC SEQ ID NO: 443

DIVMTQSPATLSLSPGERATLSCRSSKSLQNVNGNTYLYWFQQKPGQSPQLLIYRMSNLNSGVFDRFSGSGS
GTEFTLTISLLEPEDFAVYYCMQHLEYPIITFGAGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNF
YPREAKVQWKVDNALQSGNSQESVTEQDSKDSTYSLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNR
GEC

XENP11924 HC-Fab SEQ ID NO: 444

EVQLVESGGGLVKPGGSLKLSAASGYTFTSYVMHWVRQAPGKGLEWIGYINPYNDGTYNEKFQGRVTISS
DKSISTAYMELSSLRSEDTAMYYCARGTYYYGTRVFDYWGGQGLTVTVSSASTKGPSVFPLAPSSKSTSGGTA
ALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYTCNVNDRKPSNTKVD
KKVEPKSCDKTHTCPPCPAPELLRGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVQFNWYVDGVEV
HNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKARPAIEKTISKAKGQPREPQVYTLPPSQE
EMTKNQVSLTCLVKGFYPSDIAVEWESSGQPENNYNTTTPMLDSDGSFFLYSKLTVDKSRWQQEGNVFPSCSV
MHEALHNHYTQKSLSLSPG

Figure 7B

XENP11924 HC-scFv SEQ ID NO: 445

EVQLVESGGGLVQPGGSLRLSCAASGFTFNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRFTI
SRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGT¹LVTVSSGGGGSGGGSGGGGSQAVV
TQEP²SLTVSPGGTVTLTCGSSTGAVTTSNYANWVQOKPGQAPRGLIGGTNRRAPGVPA³RFSGSLLGGKAALT
LSGAQPEDEAEYYCALWYSNLWVFGGGTKLT⁴VLEKSSDKTHTCPRCPAPELLRGPSVFLFPPKPKDTLMIS
RTPEVTCVVDVSHEDPEVKFKWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNK
ARPAPIEKTI⁵SKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTT⁶PFVL
DSDGSFFLYSKLTVDKSRWQQGNV⁷FSCSVMH⁸EALHNHYTQKSLSLSPGK

XENP11924 LC SEQ ID NO: 446

DIVMTQSPATLSLSPGERATLSCRSSKSLQNVNGNTYLYWFQQKPGQSPQLLIYRMSNLNSGV¹PD²RFSGSGS
GTEFTLT³ISSLEPEDFAVYYCMQHLEYPITFGAGTKLEIKRTVAAPS⁴VFI⁵PPSDEQLKSGTASV⁶VCLLN⁷F
YPREAKVQWKVDNALQSGNSQESVTEQDSK⁸STYSL⁹STLTLSKADY¹⁰EKKH¹¹VYACEVTHQGLSS¹²PFVTK¹³SFNR
GEC

Figure 8

Humanized Anti-CD3 VH with Kabat CDRs underlined SEQ ID NO: 447

EVQLVESGGGLVQPGGSLRLSCAASGFTFNTYAMNWVRQAPGKGLEWVGRIIRSKYNNYATYYADSVKGRFTI
SRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFNGNSYVSWFAYWGQGLTVTVSS

Humanized Anti-CD3 VL with Kabat CDRs underlined SEQ ID NO: 448

QAVVTQEPSTLVSPGGTVTLTCGSSTGAVTTSNYANWVQQKPGQAPRGLIGGTNKRAPGVPARFSGSLLGK
AALTLSGAQPEDEAEYYCALWYSNLWVFGGGTKLTVL




Figure 10




Figure 11A




Figure 11B





Figure 11C

Figure 12




Figure 13

XENP11925 HC-Fab SEQ ID NO: 449

EVQLVESGGGLVQPGGSLRLSCAASGFD~~F~~SR~~S~~WMNWVRQAPGKGLEWVSEINPDSSTINYATSVKGRFTISR
DNSKNTLYLQMNSLRAEDTAVYYCARYGNWFFPYWGQGT~~L~~TVSSASTKGPSVFPLAPSSKSTSGGTAALGCL
VKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPS~~S~~SLGTQTYTCNVD~~H~~KRPSNTKVDKKVEP
KSCDKHTCPCPCPAPELLRGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVQFNWYVDGVEVHNAKT
KPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKARPAIEKTISKAKGQPREPQVYTLPPS~~Q~~EEMTKN
QVSLTCLVKGFYPSDIAVEWESSGQ~~P~~ENNYNTTPPMLDS~~D~~GSFFLYSKLTVDKSRWQEGNVFSCSVMHEALH
NHYTQKSLSLSPG

XENP11925 HC-scFv SEQ ID NO: 450

EVQLVESGGGLVQPGGSLRLSCAASGFTFNTYAMNWVRQAPGKGLEWVGRI~~R~~SKYNNYATYYADSVKGRFTI
SRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGT~~L~~TVSSGGGGSGGGGSGGGGSQAVV
TQEPSTLVSPGGTVTLTCCSSTGAVTTSNYANWVQKPGQAPRGLIGGTINKRAPGVPARFSGSLLGGKAALT
LSGAQPEDEAEYYCALWYSNLWVFGGGTKLTVLERKSSDKTHTCPRCPAPELLRGPSVFLFPPKPKDTLMIS
RTPEVTCVVVDVSHEDPEVKFKWYVDGVEVHNAKT~~K~~PREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNK
ARPAIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQ~~P~~ENNYKTTPPV
LDS~~D~~GSFFLYSKLTVDKSRWQQGNV~~F~~SCSVMHEALH~~N~~HYTQKSLSLSPGK

XENP11925 LC SEQ ID NO: 451

DIVMTQSPSSLSASVGDRVTITCRASQNVDTNVAWYQKPGQSPKALIIYSASYRYS~~G~~VPDRFTGSGSGTDFT
LTIS~~S~~LQPEDFATYFCQQYDSYPLTFGGG~~T~~KLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREA
KVQWKVDNALQSGNSQESVTEQDSKDS~~T~~YSLSSLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC




Figure 14C

Figure 15

Fluorescent LDH RTCC assay with
10k RPMI8226 cells (P12) and 300k T cell (24h)
XENP11925, purified IEX fractions




Figure 16A

XENP11924 Heavy Chain HC-Fab SEQ ID NO: 452

EVQLVESGGGLVQPGGSLKLSCAASGYTFTSYVMHWVRQAPGKGLEWIGYINPYNDGTYNEKFQGRVTI
SSDKSISTAYMELSSLRSEDTAMYYCARGTYYYGTRVFDYWGQGTTLVTVSSASTKGPSVFPLAPSSKSTS
GGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYTCNVDRKP
SNTKVDKKVEPKSCDKTHTCPPCPAPELLRGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVQFN
WYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKARPAIEKTISKAKGQPREP
QVYTLPPSQEEMTKNQVSLTCLVKGFYPSDIAVEWESSGQPENNYNTTPEMLDSDGSFFLYSKLTVDKSR
WQEGNVFSCSVMEALHNHYTQKSLSLSPG

XENP11924 Heavy Chain HC-scFv SEQ ID NO: 453

EVQLVESGGGLVQPGGSLRLSCAASGFTTFNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGTTLVTVSSGGGGSGGGGSGGGGS
QAVVTQEPSTLTVSPGGTVTLTCCSSTGAVTTSNYANWVQKPGQAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTLSGAQPEDEAEYYCALWYSNLWVFGGCTKLTVLERKSSDKTHTCPRCPAPELLRGPSVFLFPPK
PKDTLMISRTPEVTCVVVDVSHEDPEVKFKWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNG
KEYKCKVSNKARPAIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQ
PENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPGK

XENP12152 Heavy Chain HC-Fab (T411E) SEQ ID NO: 454

EVQLVESGGGLVQPGGSLKLSCAASGYTFTSYVMHWVRQAPGKGLEWIGYINPYNDGTYNEKFQGRVTI
SSDKSISTAYMELSSLRSEDTAMYYCARGTYYYGTRVFDYWGQGTTLVTVSSASTKGPSVFPLAPSSKSTS
GGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYTCNVDRKP
SNTKVDKKVEPKSCDKTHTCPPCPAPELLRGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVQFN
WYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKARPAIEKTISKAKGQPREP
QVYTLPPSQEEMTKNQVSLTCLVKGFYPSDIAVEWESSGQPENNYNTTPEMLDSDGSFFLYSKLEVDKSR
WQEGNVFSCSVMEALHNHYTQKSLSLSPG

XENP12152 Heavy Chain HC-scFv (D401K) SEQ ID NO: 455

EVQLVESGGGLVQPGGSLRLSCAASGFTTFNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGTTLVTVSSGGGGSGGGGSGGGGS
QAVVTQEPSTLTVSPGGTVTLTCCSSTGAVTTSNYANWVQKPGQAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTLSGAQPEDEAEYYCALWYSNLWVFGGCTKLTVLERKSSDKTHTCPRCPAPELLRGPSVFLFPPK
PKDTLMISRTPEVTCVVVDVSHEDPEVKFKWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNG
KEYKCKVSNKARPAIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQ
PENNYKTTTPVLDSKGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPGK

Figure 16B

XENP12155 Heavy Chain HC-Fab (L368E) SEQ ID NO: 456

EVQLVESGGGLVQPGGSLRLSCAASGFTFTSYVMHWVRQAPGKGLEWIGYINPYNDGTTYNEKFQGRVTI
SSDKSISTAYMELSSLRSEDTAMYYCARGTYYYGTRVFDYWGGQTLTVTVSSASTKGPSVFPLAPSSKSTS
GGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVFSSSLGTQTYTCNVDPKPK
SNTKVDKKVEPKSCDKTHTCTPPCPAPELLRGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVQFN
WYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKARPAIEKTISKAKGQPREP
QVYTLPPSQEEMTKNQVSLTCEVKGFYPSDIAVEWESSGQPENNYNTTPPMLDSDGSEFFLYSKLTVDKSR
WQEGNVFSCSVMEALHNHYTQKSLSLSPG

XENP12155 Heavy Chain HC-scFv (S364K) SEQ ID NO: 457

EVQLVESGGGLVQPGGSLRLSCAASGFTFTSYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGTLLTVTVSSGGGGSGGGGSGGGGS
QAVVTQEPSTLVSPGGTVTLTCGSSGTGAVTTSNYANWVQKPGQAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTLSGAQPEDEAEYYCALWYSLWVFGGGTKLTVLERKSSDKTHTCTPRCPAPELLRGPSVFLFPPK
PKDTLMISRTPEVTCVVVDVSHEDPEVKFKWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNG
KEYKCKVSNKARPAIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQ
PENNYKTTTPVLDSGSEFFLYSKLTVDKSRWQQGNVSCSVMEALHNHYTQKSLSLSPGK

XENP11924, XENP12152, XENP12155 Light Chain SEQ ID NO: 458

DIVMTQSPATLSLSPGERATLSCRSSKSLQNVNGNTYLYWFQQKPGQSPQLLIYRMSNLSGVDPDRFSGS
GSGTEFTLTISSELEPEDFAVYYCMQHLEYPITFGAGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCL
LNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSSTYSLSSTLTLSKADYEKHKVYACEVTHQGLSSPV
TKSFNRGEC

Figure 17




Figure 18




Figure 19

AMINO ACID	pI
Alanine Ala A	6.00
Arginine Arg R	11.15
Asparagine Asn N	5.41
Aspartic acid Asp D	2.77
Cysteine Cys C	5.02
Glutamic acid Glu E	3.22
Glutamine Gln Q	5.65
Glycine Gly G	5.97
Histidine His H	7.47
Isoleucine Ile I	5.94
Leucine Leu L	5.98
Lysine Lys K	9.59
Methionine Met M	5.74
Phenylalanine Phe F	5.48
Proline Pro P	6.30
Serine Ser S	5.68
Threonine Thr T	5.64
Tryptophan Trp W	5.89
Tyrosine Tyr Y	5.66
Valine Val V	5.96

Figure 20A

XenP#	Name (HC)	SEQ ID NO (HC)	Name (LC)	SEQ ID NO (LC)	Calc. pI	# KR	Delta KR (vs. WT)	# DE	Delta DE (vs. WT)	Charge State	# HC Mutations vs IgG1	# LC Mutations vs IgG1	Total # of Mutations
XENP004547	IgG1-WT	2	Ck-WT	112	8.10	122	0	116	0	6			
XENP006384	IgG2-WT	3	Ck-WT	113	7.31	118	-4	118	2	0	27	0	27
NA	IgG3-WT	4	Ck-WT	NA							22		22
NA	IgG4-WT	5	Ck-WT	NA							28		28
XENP007349	IgG1/2-HC	6	Ck-WT	114	8.11	120	-2	114	-2	6	11	0	11
XENP005653	IgG1-434S	50	Ck-WT	115	8.1	122	0	116	0	6	1	0	1
XENP006389	IgG2-434S	51	Ck-WT	116	7.31	118	-4	118	2	0	28	0	28
XENP009491	IgG1-Ch1-pI(6)	7	Ck-WT	1	6.21	116	-6	118	12	-18	6	0	6
XENP009492	IgG1-WT	2	Ck-pI(6)	8	6.21	116	-6	118	12	-18	0	6	6
XENP009493	IgG1-Ch1-pI(6)	192	Ck-pI(6)	117	5.49	110	-12	140	24	-30	6	6	12
XENP009492	IgG1-Ch1-pI(6)-434S	52	Ck-pI(6)	117	5.49	110	-12	140	24	-30	7	6	13
XENP009993	IgG1-Ch1-pI(6)-428L/434S	53	Ck-pI(6)	117	5.49	110	-12	140	24	-30	8	6	14
XENP010088	IgG1-WT	2	Ck-pI(3)	120	6.58	118	-8	122	6	-6	0	3	3
XENP010089	IgG1-WT	2	Ck-pI(6-DEDE)	121	5.85	118	-8	136	20	-30	0	10	10
XENP010090	IgG2-WT	3	Ck-pI(3)	122	6.16	112	-10	124	8	-12	27	3	30
XENP010091	IgG2-WT	3	Ck-pI(6)	123	5.88	112	-10	130	14	-18	27	6	33
XENP010092	IgG2-WT	3	Ck-pI(6-DEDE)	124	5.58	112	-10	138	22	-26	27	10	37
XENP010093	pI-Iso1	13	Ck-WT	125	6.20	110	-12	122	6	-12	13	0	13
XENP010094	pI-Iso1(NF)	14	Ck-WT	126	6.20	110	-12	122	6	-12	15	0	15
XENP010095	pI-Iso1(NF-VE)	15	Ck-WT	127	6.16	110	-12	122	6	-12	19	0	19
XENP010096	pI-Iso1(NF-VE)	15	Ck-pI(3)	128	5.63	104	-16	128	12	-24	19	3	22
XENP010101	pI-Iso1(NF-VE)	15	Ck-pI(6)	129	5.43	104	-16	134	16	-30	19	6	25
XENP010102	pI-Iso1(NF-VE)	15	Ck-pI(6-DEDE)	130	5.23	104	-16	142	26	-38	19	10	29
XENP010103	pI-Iso1(NF-VE-DEDE)	16	Ck-WT	131	5.79	110	-12	130	14	-20	22	0	22

Figure 20B

XenP#	Name (HC)	SEQ ID NO (HC)	Name (LC)	SEQ ID NO (LC)	Calc. pI	# KR	Delta KR (vs. WT)	# DE	Delta DE (vs. WT)	Charge State	# HC Mutations vs IgG1	# LC Mutations vs IgG1	Total # of Mutations
XENP010104	pI-Iso1(NF-VE-DEDE)	16	CK-pl(3)	132	5.37	104	-18	136	20	-32	22	3	25
XENP010105	pI-Iso1(NF-VE-DEDE)	16	CK-pl(6)	133	5.22	104	-18	142	28	-38	22	6	28
XENP010106	pI-Iso1(NF-VE-DEDE)	16	CK-pl(6-DEDE)	134	5.07	104	-18	150	34	-46	22	10	32
XENP010107	IgG1-pl(7)	25	CK-pl(4)	135	5.31	100	-18	136	20	-38	7	4	11
XENP010108	IgG1-pl(11)	26	CK-pl(4)	136	4.98	92	-32	144	28	-50	11	4	15
XENP010109	IgG1/2-pl(7)	27	CK-pl(4)	137	5.36	100	-30	134	18	-48	17	4	21
XENP010110	IgG1/2-pl(11)	28	CK-pl(4)	145	5.01	92	-32	142	28	-48	21	4	25
XENP010017	IgG1-pl(6)-Neutral-to-DE	56	CK-N152D S156E	138	6.59	122	0	128	12	-6	3	3	6
XENP010018	IgG1-pl(6)-KR to-Neutral	57	CK-K126Q K145Q	139	6.58	110	-12	116	0	-6	3	3	6
XENP010019	IgG1-pl(6)-KR to-DE	58	CK-K126E K145E	140	5.92	110	-12	128	12	-18	3	3	6
XENP010178	IgG-pl-Iso2	19	CK-WT	141	6.27	110	-12	120	4	-10	28	0	28
XENP010179	IgG-pl-Iso3	22	CK-WT	142	6.20	110	-12	122	6	-12	19	0	19
XENP010180	IgG-pl-Iso2-4-34S	62	CK-WT	143	6.27	110	-12	130	4	-10	29	0	29
XENP010181	IgG-pl-Iso3-4-34S	63	CK-WT	144	6.20	110	-12	122	6	-12	20	0	20
XENP010182	IgG-pl-Iso2	19	CK-pl(4)	145	5.55	102	-30	128	12	-28	28	4	32
XENP010183	IgG-pl-Iso3	22	CK-pl(4)	145	5.54	102	-30	130	14	-28	19	4	23
XENP010184	IgG-pl-Iso2-4-34S	62	CK-pl(4)	145	5.55	102	-30	128	12	-28	29	4	33
XENP010185	IgG-pl-Iso3-4-34S	63	CK-pl(4)	145	5.54	102	-30	130	14	-28	20	4	24
XENP010265	IgG-pl-Iso3-2-22K	70	any light chain	170	6.31	112	-10	122	6	-10	18	0	18
XENP010266	IgG-pl-Iso3-2-74K	71	any light chain	171	6.31	112	-10	122	6	-10	18	0	18
XENP010267	IgG-pl-Iso3-2-96Y	72	any light chain	172	6.20	110	-12	122	6	-12	18	0	18
XENP010268	IgG-pl-Iso3-3-02Y	73	any light chain	173	6.20	110	-12	122	6	-12	18	0	18
XENP010269	IgG-pl-Iso3-3-09L	74	any light chain	174	6.20	110	-12	122	6	-12	18	0	18

Figure 20C

XenP#	Name (HC)	SEQ ID NO (HC)	Name (LC)	SEQ ID NO (LC)	Calc. pI	# KR	Delta KR (vs. WT)	# DE	Delta DE (vs. WT)	Charge State	# HC Mutations vs IgG1	# LC Mutations vs IgG1	Total # of Mutations
XENP010270	IgG-pH-Iso3-339A	75	any light chain	175	6.20	110	-12	122	6	-12	18	0	18
XENP010271	IgG-pH-Iso3-355Q	76	any light chain	176	6.31	112	-10	122	8	-10	18	0	18
XENP010272	IgG-pH-Iso3-384N	77	any light chain	177	6.20	110	-12	122	6	-12	18	0	18
XENP010273	IgG-pH-Iso3-392K	78	any light chain	178	6.31	112	-10	122	6	-10	18	0	18
XENP010274	IgG-pH-Iso3-397V	79	any light chain	179	6.20	110	-12	122	6	-12	18	0	18
XENP010275	IgG-pH-Iso3-413Q	80	any light chain	180	6.31	110	-12	120	4	-10	18	0	18
XENP010276	IgG-pH-Iso3-296Y/300Y	81	any light chain	181			-12		6	-12	17	0	17
XENP010277	IgG-pH-Iso3-384N/392K/397V	82	any light chain	182	6.20	110	-10	122	6	-10	16	0	16
XENP010278	IgG-pH-Iso3-137G	83	any light chain	183	6.31	112	-12	120	4	-10	18	0	18
XENP010279	IgG-pH-Iso3-138G	84	any light chain	184	6.20	110	-12	122	8	-12	18	0	18
XENP010280	IgG-pH-Iso3-192S	85	any light chain	185	6.20	110	-12	122	6	-12	18	0	18
XENP010281	IgG-pH-Iso3-193L	86	any light chain	186	6.20	110	-12	122	6	-12	18	0	18
XENP010282	IgG-pH-Iso3-199H	87	any light chain	187	6.20	110	-12	122	6	-12	18	0	18
XENP010283	IgG-pH-Iso3-203N	88	any light chain	188	6.31	110	-12	120	4	-10	18	0	18
XENP010284	IgG-pH-Iso3-214K	89	any light chain	189	6.31	112	-10	122	6	-10	18	0	18
XENP010285	IgG-pH-Iso3-137G/138G	90	any light chain	146			-12		4	-10	17	0	17
XENP010286	IgG-pH-Iso3-51	23	Ck-WT	194	6.20	110	-12	122	6	-12	17	0	17
XENP010287	IgG-pH-Iso3-199H/203N	91	any light chain	195			-12		4	-10	17	0	17
XENP010288	IgG-pH-Iso3-214K/222K	92	any light chain	196	6.31	110	-8	120	6	-8	17	0	17
XENP010289	IgG-pH-Iso3-138G/192S/193L	93	any light chain	197	6.44	114	-12	122	8	-12	16	0	16
XENP010290	IgG-PH-Iso3-137G/138G/192S/	94	any light chain	198	6.20	110	-12	122	4	-10	15	0	15

Figure 20D

XenP#	Name (HC)	SEQ ID NO (HC)	Name (LC)	SEQ ID NO (LC)	Calc. pI	# KR	Delta KR (vs. WT)	# DE	Delta DE Charge (vs. WT)	Charge State	# HC Mutations vs IgG1	# LC Mutations vs IgG1	Total # of Mutations
XENP010324	IgG-pl-Iso3	22	CK-Iso(3)	148	5.92	106	-16	124	8	-18	19	3	22
XENP010325	IgG-pl-Iso3	22	CK-Iso(4)	149	5.83	104	-18	124	8	-20	19	4	23
XENP010326	IgG-pl-Iso3	22	CK-Iso(5)	150	5.75	104	-18	126	10	-22	19	5	24
XENP010327	IgG-pl-Iso3	22	CK-Iso(6)	151	5.68	104	-18	128	12	-24	19	6	25
XENP010425	H3L0_IgG2_CH1_I	95	any light chain	199	7.30	120	-2	120	4	0	16	0	16
XENP010426	IgG-pl-Iso3-charges-only	24	Cx-WT	152	6.20	110	-12	122	6	-12	9	0	9
XENP010427	IgG-pl-Iso2-charges-only	21	Cx-WT	153	6.27	110	-12	120	4	-10	18	0	18
XENP010428	H3L0_IgG2_CH1_I	161	any light chain	200	7.67	122	-12	120	4	2	9	0	9
XENP010456	IgG-pl-Iso3-SL-4345	98	Cx-WT	152	6.20	110	-12	122	6	-12	18	0	18
XENP010467	IgG-pl-Iso3-SL-4384/4345	97	Cx-WT	152	6.20	110	-12	122	6	-12	19	0	19
XENP010468	H3L0_pl_Iso3_S13	162	any light chain	201	6.20	110	-12	122	4	-12	17	0	17
XENP010469	7G/S138G/N192S/	163	any light chain	202	6.31	110	-12	120	4	-10	16	0	16
XENP010470	IgG-pl-Iso2-SL-4345	98	Cx-WT	152	6.27	110	-12	120	4	-10	26	0	26
XENP010471	IgG-pl-Iso2-SL-4345	98	Cx-WT	152	6.27	110	-12	120	4	-10	27	0	27

Figure 20E

XenP#	Name (HC)	SEQ ID NO (HC)	Name (LC)	SEQ ID NO (LC)	Calc. pI	# KR	Delta KR (vs. WT)	# DE	Delta DE Charge (vs. WT) State	# HC Mutations vs IgG1	# LC Mutations vs IgG1	Total # of Mutations
XENP010472	lgG-pl-Iso3- charges-only-434S	191	Ck-WT	152	6.20	110	-12	122	6 -12	10	0	10
XENP010473	lgG-pl-Iso2- charges-only-434S	192	Ck-WT	152	6.27	110	-12	130	4 -10	19	0	19
XENP010474	H0L0_IgG2_CH1_I gG1_CH2_CH3_N4	164	any light chain	203	7.30	120	-2	130	4 0	17	0	17
XENP010475	H0L0_IgG2_CH1_I gG1_Hinge_CH2_C	165	any light chain	204	7.67	122	-22	120	20 -38	8	4	12
XENP010476	lgG-pl-Iso3- charges-only-434S	25	Ck-pl(4)	135	5.31	100	-22	136	18 -34	18	4	22
XENP010477	lgG-pl-Iso3- charges-only-434S	166	Ck-pl(4)	135	5.36	100	-12	134	4 -10	23	0	23
XENP010478	H0L0_pl_Iso1/2_c harges_only	167	any light chain	205	6.27	110	-16	130	8 -18	17	3	20
XENP010511	lgG-pl-Iso3-SL	23	Ck-Iso(3)	148	5.92	106	-18	124	8 -20	17	4	21
XENP010512	lgG-pl-Iso3-SL	23	Ck-Iso(4)	149	5.83	104	-18	124	10 -22	17	5	22
XENP010513	lgG-pl-Iso3-SL	23	Ck-Iso(5)	150	5.75	104	-18	126	8 -18	18	3	21
XENP010517	lgG-pl-Iso3-SL- 434S	98	Ck-Iso(3)	148	5.92	106	-18	124	8 -20	18	4	22
XENP010518	lgG-pl-Iso3-SL- 434S	98	Ck-Iso(4)	149	5.83	104	-18	124	10 -22	18	5	23
XENP010519	lgG-pl-Iso3-SL- 434S	98	Ck-Iso(5)	150	5.75	104	-18	126				

Figure 20F

XenP#	Name (HC)	SEQ ID NO (HC)	Name (LC)	SEQ ID NO (LC)	Calc. pI	# KR	Delta KR (vs. WT)	# DE	Delta DE (vs. WT)	Charge State	# HC Mutations vs IgG1	# LC Mutations vs IgG1	Total # of Mutations
XENP010520	IgG-pl-Iso3-SL-428L/434S	97	CK-Iso(3)	148	5.92	106	-18	124	8	-18	19	3	22
XENP010521	IgG-pl-Iso3-SL-428L/434S	97	CK-Iso(4)	149	5.83	104	-18	124	8	-30	19	4	23
XENP010522	IgG-pl-Iso3-SL-428L/434S	97	CK-Iso(5)	150	5.75	104	-18	126	10	-22	19	5	24
XENP010525	IgG-pl-Iso3-SL-434S	98	CK-pl(4)	149	5.54	102	-20	130	14	-38	18	4	22
XENP010526	IgG-pl-Iso3-434S	99	CK-Iso(5)	150	5.75	104	-18	126	10	-22	20	5	25
XENP010527	IgG-pl-Iso2-SL-434S	86	CK-Iso(5)	150	5.78	104	-18	124	8	-20	27	5	32

Figure 21A

Variant	# of sub(s)	pl / pl	pl / WT	WT / WT	avg delta pl
G137E/N203D/K274Q/R355Q/K392N/Q419E/K447_	7	6.43	7.14	8.02	-0.79
G137E/N203D/K274Q/R355Q/K392N/Q419E	6	6.58	7.30	8.02	-0.72
G137E/N203D/K274Q/R355Q/K392N/K447_	6	6.58	7.30	8.02	-0.72
G137E/N203D/K274Q/R355Q/Q419E/K447_	6	6.58	7.30	8.02	-0.72
G137E/N203D/K274Q/K392N/Q419E/K447_	6	6.58	7.30	8.02	-0.72
G137E/N203D/R355Q/K392N/Q419E/K447_	6	6.58	7.30	8.02	-0.72
G137E/K274Q/R355Q/K392N/Q419E/K447_	6	6.58	7.30	8.02	-0.72
N203D/K274Q/R355Q/K392N/Q419E/K447_	6	6.58	7.30	8.02	-0.72
G137E/N203D/K274Q/R355Q/K392N	5	6.76	7.46	8.02	-0.63
G137E/N203D/K274Q/R355Q/Q419E	5	6.76	7.46	8.02	-0.63
G137E/N203D/K274Q/R355Q/K447_	5	6.76	7.46	8.02	-0.63
G137E/N203D/K274Q/K392N/Q419E	5	6.76	7.46	8.02	-0.63
G137E/N203D/K274Q/K392N/K447_	5	6.76	7.46	8.02	-0.63
G137E/N203D/K274Q/Q419E/K447_	5	6.76	7.46	8.02	-0.63
G137E/N203D/R355Q/K392N/Q419E	5	6.76	7.46	8.02	-0.63
G137E/N203D/R355Q/K392N/K447_	5	6.76	7.46	8.02	-0.63
G137E/N203D/R355Q/Q419E/K447_	5	6.76	7.46	8.02	-0.63
G137E/N203D/K392N/Q419E/K447_	5	6.76	7.46	8.02	-0.63
G137E/K274Q/R355Q/K392N/Q419E	5	6.76	7.46	8.02	-0.63
G137E/K274Q/R355Q/K392N/K447_	5	6.76	7.46	8.02	-0.63
G137E/K274Q/R355Q/Q419E/K447_	5	6.76	7.46	8.02	-0.63
G137E/K274Q/K392N/Q419E/K447_	5	6.76	7.46	8.02	-0.63
G137E/R355Q/K392N/Q419E/K447_	5	6.76	7.46	8.02	-0.63
N203D/K274Q/R355Q/K392N/Q419E	5	6.76	7.46	8.02	-0.63
N203D/K274Q/R355Q/K392N/K447_	5	6.76	7.46	8.02	-0.63
N203D/K274Q/R355Q/Q419E/K447_	5	6.76	7.46	8.02	-0.63
N203D/K274Q/K392N/Q419E/K447_	5	6.76	7.46	8.02	-0.63
N203D/R355Q/K392N/Q419E/K447_	5	6.76	7.46	8.02	-0.63
K274Q/R355Q/K392N/Q419E/K447_	5	6.76	7.46	8.02	-0.63
G137E/N203D/K274Q/R355Q	4	7.00	7.61	8.02	-0.51
G137E/N203D/K274Q/K392N	4	7.00	7.61	8.02	-0.51
G137E/N203D/K274Q/Q419E	4	7.00	7.61	8.02	-0.51
G137E/N203D/K274Q/K447_	4	7.00	7.61	8.02	-0.51
G137E/N203D/R355Q/K392N	4	7.00	7.61	8.02	-0.51
G137E/N203D/R355Q/Q419E	4	7.00	7.61	8.02	-0.51
G137E/N203D/R355Q/K447_	4	7.00	7.61	8.02	-0.51
G137E/N203D/K392N/Q419E	4	7.00	7.61	8.02	-0.51
G137E/N203D/K392N/K447_	4	7.00	7.61	8.02	-0.51
G137E/N203D/Q419E/K447_	4	7.00	7.61	8.02	-0.51
G137E/K274Q/R355Q/K392N	4	7.00	7.61	8.02	-0.51
G137E/K274Q/R355Q/Q419E	4	7.00	7.61	8.02	-0.51
G137E/K274Q/R355Q/K447_	4	7.00	7.61	8.02	-0.51
G137E/K274Q/K392N/Q419E	4	7.00	7.61	8.02	-0.51
G137E/K274Q/K392N/K447_	4	7.00	7.61	8.02	-0.51
G137E/K274Q/Q419E/K447_	4	7.00	7.61	8.02	-0.51
G137E/R355Q/K392N/Q419E	4	7.00	7.61	8.02	-0.51
G137E/R355Q/K392N/K447_	4	7.00	7.61	8.02	-0.51
G137E/R355Q/Q419E/K447_	4	7.00	7.61	8.02	-0.51
G137E/K392N/Q419E/K447_	4	7.00	7.61	8.02	-0.51
N203D/K274Q/R355Q/K392N	4	7.00	7.61	8.02	-0.51
N203D/K274Q/R355Q/Q419E	4	7.00	7.61	8.02	-0.51
N203D/K274Q/R355Q/K447_	4	7.00	7.61	8.02	-0.51
N203D/K274Q/K392N/Q419E	4	7.00	7.61	8.02	-0.51

Figure 21B

N203D/K274Q/K392N/K447_	4	7.00	7.61	8.02	-0.51
N203D/K274Q/Q419E/K447_	4	7.00	7.61	8.02	-0.51
N203D/R355Q/K392N/Q419E	4	7.00	7.61	8.02	-0.51
N203D/R355Q/K392N/K447_	4	7.00	7.61	8.02	-0.51
N203D/R355Q/Q419E/K447_	4	7.00	7.61	8.02	-0.51
N203D/K392N/Q419E/K447_	4	7.00	7.61	8.02	-0.51
K274Q/R355Q/K392N/Q419E	4	7.00	7.61	8.02	-0.51
K274Q/R355Q/K392N/K447_	4	7.00	7.61	8.02	-0.51
K274Q/R355Q/Q419E/K447_	4	7.00	7.61	8.02	-0.51
K274Q/K392N/Q419E/K447_	4	7.00	7.61	8.02	-0.51
R355Q/K392N/Q419E/K447_	4	7.00	7.61	8.02	-0.51
G137E/N203D/K274Q	3	7.30	7.74	8.02	-0.36
G137E/N203D/R355Q	3	7.30	7.74	8.02	-0.36
G137E/N203D/K392N	3	7.30	7.74	8.02	-0.36
G137E/N203D/Q419E	3	7.30	7.74	8.02	-0.36
G137E/N203D/K447_	3	7.30	7.74	8.02	-0.36
G137E/K274Q/R355Q	3	7.30	7.74	8.02	-0.36
G137E/K274Q/K392N	3	7.30	7.74	8.02	-0.36
G137E/K274Q/Q419E	3	7.30	7.74	8.02	-0.36
G137E/K274Q/K447_	3	7.30	7.74	8.02	-0.36
G137E/R355Q/K392N	3	7.30	7.74	8.02	-0.36
G137E/R355Q/Q419E	3	7.30	7.74	8.02	-0.36
G137E/R355Q/K447_	3	7.30	7.74	8.02	-0.36
G137E/K392N/Q419E	3	7.30	7.74	8.02	-0.36
G137E/K392N/K447_	3	7.30	7.74	8.02	-0.36
G137E/Q419E/K447_	3	7.30	7.74	8.02	-0.36
N203D/K274Q/R355Q	3	7.30	7.74	8.02	-0.36
N203D/K274Q/K392N	3	7.30	7.74	8.02	-0.36
N203D/K274Q/Q419E	3	7.30	7.74	8.02	-0.36
N203D/K274Q/K447_	3	7.30	7.74	8.02	-0.36
N203D/R355Q/K392N	3	7.30	7.74	8.02	-0.36
N203D/R355Q/Q419E	3	7.30	7.74	8.02	-0.36
N203D/R355Q/K447_	3	7.30	7.74	8.02	-0.36
N203D/K392N/Q419E	3	7.30	7.74	8.02	-0.36
N203D/K392N/K447_	3	7.30	7.74	8.02	-0.36
N203D/Q419E/K447_	3	7.30	7.74	8.02	-0.36
K274Q/R355Q/K392N	3	7.30	7.74	8.02	-0.36
K274Q/R355Q/Q419E	3	7.30	7.74	8.02	-0.36
K274Q/R355Q/K447_	3	7.30	7.74	8.02	-0.36
K274Q/K392N/Q419E	3	7.30	7.74	8.02	-0.36
K274Q/K392N/K447_	3	7.30	7.74	8.02	-0.36
K274Q/Q419E/K447_	3	7.30	7.74	8.02	-0.36
R355Q/K392N/Q419E	3	7.30	7.74	8.02	-0.36
R355Q/K392N/K447_	3	7.30	7.74	8.02	-0.36
R355Q/Q419E/K447_	3	7.30	7.74	8.02	-0.36
K392N/Q419E/K447_	3	7.30	7.74	8.02	-0.36
G137E/N203D	2	7.61	7.85	8.02	-0.21
G137E/K274Q	2	7.61	7.85	8.02	-0.21
G137E/R355Q	2	7.61	7.85	8.02	-0.21
G137E/K392N	2	7.61	7.85	8.02	-0.21
G137E/Q419E	2	7.61	7.85	8.02	-0.21
G137E/K447_	2	7.61	7.85	8.02	-0.21
N203D/K274Q	2	7.61	7.85	8.02	-0.21
N203D/R355Q	2	7.61	7.85	8.02	-0.21
N203D/K392N	2	7.61	7.85	8.02	-0.21
N203D/Q419E	2	7.61	7.85	8.02	-0.21
N203D/K447_	2	7.61	7.85	8.02	-0.21

Figure 21C

K274Q/R355Q	2	7.61	7.85	8.02	-0.21
K274Q/K392N	2	7.61	7.85	8.02	-0.21
K274Q/Q419E	2	7.61	7.85	8.02	-0.21
K274Q/K447_	2	7.61	7.85	8.02	-0.21
R355Q/K392N	2	7.61	7.85	8.02	-0.21
R355Q/Q419E	2	7.61	7.85	8.02	-0.21
R355Q/K447_	2	7.61	7.85	8.02	-0.21
K392N/Q419E	2	7.61	7.85	8.02	-0.21
K392N/K447_	2	7.61	7.85	8.02	-0.21
Q419E/K447_	2	7.61	7.85	8.02	-0.21
G137E	1	7.85	7.94	8.02	-0.09
N203D	1	7.85	7.94	8.02	-0.09
K274Q	1	7.85	7.94	8.02	-0.09
R355Q	1	7.85	7.94	8.02	-0.09
K392N	1	7.85	7.94	8.02	-0.09
Q419E	1	7.85	7.94	8.02	-0.09
K447_	1	7.85	7.94	8.02	-0.09

Figure 22

Variant	# of sub(s)	pl / pl	pl / WT	WT / WT	avg delta pl
Q196K/P217R/P228R/N276K/H435R	5	8.53	8.32	8.02	0.25
Q196K/P217R/P228R/N276K	4	8.45	8.27	8.02	0.22
Q196K/P217R/P228R/H435R	4	8.46	8.28	8.02	0.22
Q196K/P217R/N276K/H435R	4	8.45	8.27	8.02	0.22
Q196K/P228R/N276K/H435R	4	8.45	8.27	8.02	0.22
P217R/P228R/N276K/H435R	4	8.46	8.28	8.02	0.22
Q196K/P217R/P228R	3	8.37	8.22	8.02	0.17
Q196K/P217R/N276K	3	8.37	8.22	8.02	0.17
Q196K/P217R/H435R	3	8.37	8.22	8.02	0.17
Q196K/P228R/N276K	3	8.37	8.22	8.02	0.17
Q196K/P228R/H435R	3	8.37	8.22	8.02	0.17
Q196K/N276K/H435R	3	8.37	8.22	8.02	0.17
P217R/P228R/N276K	3	8.37	8.22	8.02	0.17
P217R/P228R/H435R	3	8.37	8.22	8.02	0.18
P217R/N276K/H435R	3	8.37	8.22	8.02	0.17
P228R/N276K/H435R	3	8.37	8.22	8.02	0.17
Q196K/P217R	2	8.27	8.16	8.02	0.13
Q196K/P228R	2	8.27	8.16	8.02	0.13
Q196K/N276K	2	8.27	8.16	8.02	0.13
Q196K/H435R	2	8.27	8.16	8.02	0.13
P217R/P228R	2	8.28	8.16	8.02	0.13
P217R/N276K	2	8.27	8.16	8.02	0.13
P217R/H435R	2	8.28	8.16	8.02	0.13
P228R/N276K	2	8.27	8.16	8.02	0.13
P228R/H435R	2	8.28	8.16	8.02	0.13
N276K/H435R	2	8.27	8.16	8.02	0.13
Q196K	1	8.16	8.10	8.02	0.07
P217R	1	8.16	8.10	8.02	0.07
P228R	1	8.16	8.10	8.02	0.07
N276K	1	8.16	8.10	8.02	0.07
H435R	1	8.16	8.10	8.02	0.07

Figure 23A

HETERODIMERIZATION VARIANTS

	IgG1	IgG2	IgG3	IgG4	Variants
196	Q	Q	Q	K	K
199	I	T	T	T	T
203	N	D	N	D	D
217	P	R	L	S	R
220	C	C	P	G	E,R (IgG2)
221	D	—	LGD	—	E,R (IgG1)
222	K	V	T	—	E,R (IgG2)
223	T	—	T	—	D,E,R,K
225	T	—	T	P	D,E,R,K
228	P	P	RCPEPK SCDTPP PCPRCP EPKSCD TPPPCP RCPEPK SCDTPP PCPR	S	D,E,R,K
247	P	P	P	P	Q
276	N	N	K	N	K
340	K	K	K	K	E,Q
345	E	E	E	E	K
	Q	Q	Q	Q	E,K,R
349	Y	Y	Y	Y	A,C,D,E,I,K,S,T,W
350	T	T	T	T	I
351	L	L	L	L	E,K,V,Y
354	S	S	S	S	C
355	R	R	R	Q	E,Q
356	D	E	E	E	K,L,R
357	E	E	E	E	K,R,Q,T
	K	K	K	K	D,E
362	Q	Q	Q	Q	E,K
	S	S	S	S	C,D,E,F,G,H,K,R,T,Y
366	T	T	T	T	A,D,I,K,L,M,S,V,W,Y
	L	L	L	L	A,D,E,K,S all but C,P
370	K	K	K	K	C,D,E,G,R,S,T,V all but C,P
371	G	G	G	G	D
384	N	N	S	N	S
390	N	N	N	N	D,E,K,R
392	K	K	N	K	C,D,E,F,L,M,N
394	T	T	T	T	F,S,V,W,Y

Figure 23B

	IgG1	IgG2	IgG3	IgG4	Variants
395	P	P	P	P	T,V
396	P	P	P	P	T,V
397	V	M	M	V	M,S,T
399	D	D	D	D	all but C,P C,K,R
400	S	S	S	S	A,D,E,K,R
401	D	D	D	D	K,N,R
405	F	F	F	F	L, all but C,P A,F,L,M,S,T,V
407	Y	Y	Y	Y	T,V all but C,P A,L,M,V
409	K	K	K	R	R ,all but C,P D,E,F,K,L,M,V,W
	T	T	T	T	D,E,K,L,N,R,S
419	Q	Q	Q	E	E
439	K	K	K	K	D,E

FIGURE 24

COMBINATIONS OF VARIANTS INTO HETERODIMERIZATION FORMATS

Heterodimerization format	FcRn variants Monomer 1 and/or Monomer 2	Fc variants Monomer 1 and/or Monomer 2	pl variants	Steric variants (including charge pairs)	Combinations (See Legend E)
"bottle opener" scFv-Fab-Fc (Figure 1B)	See Legend A	See Legend B	See Legend C	See Legend D	1), 2), 3), 4), 5), 6), 7), 8), 9) and 10)
Trimeric bottle opener (Figure 5A)	See Legend A	See Legend B	See Legend C	See Legend D	1), 2), 3), 4), 5), 6), 7), 8), 9) and 10)
Trimeric bottle opener (scFv- Fab-Fv-scFv; Figure 5A)	See Legend A	See Legend B	See Legend C	See Legend D	1), 2), 3), 4), 5), 6), 7), 8), 9) and 10)
Trimeric bottle opener (scFv- Fab-Fv-scFv; Figure 5B)	See Legend A	See Legend B	See Legend C	See Legend D	1), 2), 3), 4), 5), 6), 7), 8), 9) and 10)
Trimeric bottle opener (scFv- Fab-Fv-scFv; Figure 5C)	See Legend A	See Legend B	See Legend C	See Legend D	1), 2), 3), 4), 5), 6), 7), 8), 9) and 10)
Trimeric bottle opener (scFv- Fab-Fv-Fab; Figure 5D)	See Legend A	See Legend B	See Legend C	See Legend D	1), 2), 3), 4), 5), 6), 7), 8), 9) and 10)
Trimeric bottle opener (scFv- Fab-Fv-Fab; Figure 5E)	See Legend A	See Legend B	See Legend C	See Legend D	1), 2), 3), 4), 5), 6), 7), 8), 9) and 10)

Figure 25A

XENP	VH ID	VL ID	VH Substitutions	VL Substitutions
11502	H1	L1.4		
11794	H1.3	L1.4	L11D	
11795	H1.4	L1.4	L11K	
11796	H1.5	L1.4	L11N	
11801	H1.6	L1.4	L11S	
11802	H1.7	L1.4	R19K	
11803	H1.8	L1.4	N30S	
11804	H1.9	L1.4	N35H	
11805	H1.10	L1.4	N35S	
11806	H1.11	L1.4	P41S	
11807	H1.12	L1.4	Y52CA	
11808	H1.13	L1.4	Y52CA/N54S	
11809	H1.14	L1.4	Y58A	
11810	H1.15	L1.4	D61A	
11811	H1.16	L1.4	D73N	
11812	H1.17	L1.4	D73T	
11813	H1.18	L1.4	L78A	
11814	H1.19	L1.4	A84D	
11815	H1.20	L1.4	A84K	
11816	H1.21	L1.4	A84S	
11817	H1.22	L1.4	V93A	
11818	H1.23	L1.4	V93T	
11819	H1.24	L1.4	N100D	
11820	H1.25	L1.4	N100Q	
11821	H1.26	L1.4	S100AT	
11822	H1.27	L1.4	A101D	
11823	H1.28	L1.4	L108T	
11824	H1.29	L1.30	G44S	G100S
11825	H1	L1.10		E7P
11826	H1	L1.11		T12S
11827	H1	L1.12		G17A
11828	H1	L1.13		G17Q
11829	H1	L1.14		T29S
11830	H1	L1.15		T29S/S30G/N31H
11831	H1	L1.16		G49Y/G50D
11832	H1	L1.17		K53N
11833	H1	L1.18		P56S
11834	H1	L1.19		N52S/K53N/R54K/A55H/P56S/G57W/V58T
11835	H1	L1.20		A60D
11836	H1	L1.21		L66K/L67S
11837	H1	L1.22		L66S/L67S

Figure 25B

XENP	VH ID	VL ID	VH Substitutions	VL Substitutions
11838	H1	L1.23		L75I
11839	H1	L1.24		P80A
11840	H1	L1.25		P80S
11841	H1	L1.26		E85D
11842	H1	L1.27		A89L
11843	H1	L1.28		L95H
11844	H1	L1.29		T105E/V106I/L106AK
11920	H1	L3.1		
11921	H1	L5.1		
11922	H1	L1.31		L75D/E85D/L95H
11923	H1	L1.32		Q38E
12107	H1.38	L1.4	V89T	
12108	H1	L1.33		Q42K
12109	H1	L1.34		Q42T
12110	H1	L1.35		Q42S/A43P
12111	H1	L1.36		A43P
12112	H1	L1.37		A43G
12113	H1	L1.38		A43S
12114	H1	L1.39		A43D
12131	H1.30	L1.4	N30S/N100D	
12132	H1.31	L1.4	N30S/N35S/N100D	
12133	H1.32	L1.4	N30S/Y52CA/N100D	
12134	H1.33	L1.4	N30S/N100D/A101D	
12135	H1.34	L1.4	N30S/N35S/Y52CA/N100D/A101D	
12136	H1.35	L1.4	G10E/L11V/V12K/Q13K/G16E	
12137	H1.36	L1.4	K52BN/Y52C_/N53_/N54G/Y55G/A56S	
12138	H1.37	L1.4	Q39K	
12139	H1.37	L1.32	Q39K	Q38E
12149	H1	L1.40		L75I/E85D/L95H
12213	H1.30	L1.40	N30S/N100D	L75I/E85D/L95H
12214	H1.8	L1.40	N30S	L75I/E85D/L95H
12215	H1.39	L1.40	N30S/Q39K	L75I/E85D/L95H
12216	H1.40	L1.40	N30S/Q39K/N100D	L75I/E85D/L95H
12217	H1.30	L1.41	N30S/N100D	L75I/L95H
12218	H1.8	L1.41	N30S	L75I/L95H
12219	H1.39	L1.41	N30S/Q39K	L75I/L95H
12220	H1.40	L1.41	N30S/Q39K/N100D	L75I/L95H

Figure 25C

XENP	VH ID	VL ID	VH Substitutions	VL Substitutions
12221	H1.30	L1.42	N30S/N100D	L75I/Q42K/L95H
12222	H1.8	L1.42	N30S	L75I/Q42K/L95H
12223	H1.39	L1.42	N30S/Q39K	L75I/Q42K/L95H
12224	H1.40	L1.42	N30S/Q39K/N100D	L75I/Q42K/L95H
12225	H1.30	L1.43	N30S/N100D	L75I/A43S/L95H
12226	H1.8	L1.43	N30S	L75I/A43S/L95H
12227	H1.39	L1.43	N30S/Q39K	L75I/A43S/L95H
12228	H1.40	L1.43	N30S/Q39K/N100D	L75I/A43S/L95H
12229	H1.30	L1.44	N30S/N100D	L75I/Q42K/A43S/L95H
12230	H1.8	L1.44	N30S	L75I/Q42K/A43S/L95H
12231	H1.39	L1.44	N30S/Q39K	L75I/Q42K/A43S/L95H
12232	H1.40	L1.44	N30S/Q39K/N100D	L75I/Q42K/A43S/L95H
12233	H1.30	L1.45	N30S/N100D	L75I/Q42K/E85D/L95H
12234	H1.8	L1.45	N30S	L75I/Q42K/E85D/L95H
12235	H1.39	L1.45	N30S/Q39K	L75I/Q42K/E85D/L95H
12236	H1.40	L1.45	N30S/Q39K/N100D	L75I/Q42K/E85D/L95H
12237	H1.30	L1.46	N30S/N100D	L75I/A43S/E85D/L95H
12238	H1.8	L1.46	N30S	L75I/A43S/E85D/L95H
12239	H1.39	L1.46	N30S/Q39K	L75I/A43S/E85D/L95H
12240	H1.40	L1.46	N30S/Q39K/N100D	L75I/A43S/E85D/L95H
12241	H1.30	L1.47	N30S/N100D	L75I/Q42K/A43S/E85D/L95H
12242	H1.8	L1.47	N30S	L75I/Q42K/A43S/E85D/L95H
12243	H1.39	L1.47	N30S/Q39K	L75I/Q42K/A43S/E85D/L95H
12244	H1.40	L1.47	N30S/Q39K/N100D	L75I/Q42K/A43S/E85D/L95H

Figure 26B

SEQ ID NO. 9.

EVQLVESGGGDVQPGGSLRLSCAASGFTFNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLTVTVSS

SEQ ID NO. 10.

QAVVTQEPSTLVSPGGTVTLTCSSTGAVTTSNYANWVQKPGQAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTLSGAQPEDEAEYYCALWYSNLWVFGGGTKLTVL

XENP11795 H1.4_L1.4

SEQ ID NO. 11.

EVQLVESGGGKVQPGGSLRLSCAASGFTFNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLTVTVSS
QAVVTQEPSTLVSPGGTVTLTCSSTGAVTTSNYANWVQKPGQAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTLSGAQPEDEAEYYCALWYSNLWVFGGGTKLTVLGSHHHHHH

SEQ ID NO. 12.

EVQLVESGGGKVQPGGSLRLSCAASGFTFNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLTVTVSS
QAVVTQEPSTLVSPGGTVTLTCSSTGAVTTSNYANWVQKPGQAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTLSGAQPEDEAEYYCALWYSNLWVFGGGTKLTVL

SEQ ID NO. 13.

EVQLVESGGGKVQPGGSLRLSCAASGFTFNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLTVTVSS

SEQ ID NO. 14.

QAVVTQEPSTLVSPGGTVTLTCSSTGAVTTSNYANWVQKPGQAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTLSGAQPEDEAEYYCALWYSNLWVFGGGTKLTVL

XENP11796 H1.5_L1.4

SEQ ID NO. 15.

EVQLVESGGGNVQPGGSLRLSCAASGFTFNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLTVTVSS
QAVVTQEPSTLVSPGGTVTLTCSSTGAVTTSNYANWVQKPGQAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTLSGAQPEDEAEYYCALWYSNLWVFGGGTKLTVLGSHHHHHH

XENP11796 H1.5_L1.4, cont.

SEQ ID NO. 16.

EVQLVESGGGNVQPGGSLRLSCAASGFTFNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLTVTVSS
QAVVTQEPSTLVSPGGTVTLTCSSTGAVTTSNYANWVQKPGQAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTLSGAQPEDEAEYYCALWYSNLWVFGGGTKLTVL

SEQ ID NO. 17.

EVQLVESGGGNVQPGGSLRLSCAASGFTFNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLTVTVSS

Figure 26C

SEQ ID NO. 18.

QAVVTQEP~~SLTV~~SPGGT~~VT~~LT~~CG~~SSTGAVTTSNYANWVQ~~QKPGQ~~APRGLIGGTNKRAPGV~~PAR~~FSGSLLG
GKAALTLSGAQPEDEAEYYCALWYSNLWVFGGGTKLTVL

XENP11801 H1.6_L1.4

SEQ ID NO. 19.

EVQLVESGGGSVQPGGSLRLSCAASGFTFNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFNGNSYVSWFAYWGQGT~~LVTVSS~~
QAVVTQEP~~SLTV~~SPGGT~~VT~~LT~~CG~~SSTGAVTTSNYANWVQ~~QKPGQ~~APRGLIGGTNKRAPGV~~PAR~~FSGSLLG
GKAALTLSGAQPEDEAEYYCALWYSNLWVFGGGTKLTVLGSHHHHHH

SEQ ID NO. 20.

EVQLVESGGGSVQPGGSLRLSCAASGFTFNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFNGNSYVSWFAYWGQGT~~LVTVSS~~
QAVVTQEP~~SLTV~~SPGGT~~VT~~LT~~CG~~SSTGAVTTSNYANWVQ~~QKPGQ~~APRGLIGGTNKRAPGV~~PAR~~FSGSLLG
GKAALTLSGAQPEDEAEYYCALWYSNLWVFGGGTKLTVL

SEQ ID NO. 21.

EVQLVESGGGSVQPGGSLRLSCAASGFTFNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFNGNSYVSWFAYWGQGT~~LVTVSS~~

SEQ ID NO. 22.

QAVVTQEP~~SLTV~~SPGGT~~VT~~LT~~CG~~SSTGAVTTSNYANWVQ~~QKPGQ~~APRGLIGGTNKRAPGV~~PAR~~FSGSLLG
GKAALTLSGAQPEDEAEYYCALWYSNLWVFGGGTKLTVL

XENP11802 H1.7_L1.4

SEQ ID NO. 23.

EVQLVESGGGLVQPGGSLKLSCAASGFTFNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFNGNSYVSWFAYWGQGT~~LVTVSS~~
QAVVTQEP~~SLTV~~SPGGT~~VT~~LT~~CG~~SSTGAVTTSNYANWVQ~~QKPGQ~~APRGLIGGTNKRAPGV~~PAR~~FSGSLLG
GKAALTLSGAQPEDEAEYYCALWYSNLWVFGGGTKLTVLGSHHHHHH

SEQ ID NO. 24.

EVQLVESGGGLVQPGGSLKLSCAASGFTFNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFNGNSYVSWFAYWGQGT~~LVTVSS~~
QAVVTQEP~~SLTV~~SPGGT~~VT~~LT~~CG~~SSTGAVTTSNYANWVQ~~QKPGQ~~APRGLIGGTNKRAPGV~~PAR~~FSGSLLG
GKAALTLSGAQPEDEAEYYCALWYSNLWVFGGGTKLTVL

SEQ ID NO. 25.

EVQLVESGGGLVQPGGSLKLSCAASGFTFNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFNGNSYVSWFAYWGQGT~~LVTVSS~~

SEQ ID NO. 26.

QAVVTQEP~~SLTV~~SPGGT~~VT~~LT~~CG~~SSTGAVTTSNYANWVQ~~QKPGQ~~APRGLIGGTNKRAPGV~~PAR~~FSGSLLG
GKAALTLSGAQPEDEAEYYCALWYSNLWVFGGGTKLTVL

Figure 26H

XENP11811 H1.16_L1.4

SEQ ID NO. 59.

EVQLVESGGGLVQPFGGSLRLSCAASGFTFTNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDN SKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLVTVSSGGGGGGGGGGGGGGGGGG
QAVVTQEP SLTVSPGGTVTLTCGSSTGAVTTSNYANWVQQKPGQAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTLSGAQPEDEAEYYCALWYSNLWVFGGGTKLTVLGSHHHHHH

SEQ ID NO. 60.

EVQLVESGGGLVQPFGGSLRLSCAASGFTFTNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDN SKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLVTVSSGGGGGGGGGGGGGGGGGG
QAVVTQEP SLTVSPGGTVTLTCGSSTGAVTTSNYANWVQQKPGQAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTLSGAQPEDEAEYYCALWYSNLWVFGGGTKLTVL

SEQ ID NO. 61.

EVQLVESGGGLVQPFGGSLRLSCAASGFTFTNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDN SKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLVTVSS

SEQ ID NO. 62.

QAVVTQEP SLTVSPGGTVTLTCGSSTGAVTTSNYANWVQQKPGQAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTLSGAQPEDEAEYYCALWYSNLWVFGGGTKLTVL

XENP11812 H1.17_L1.4

SEQ ID NO. 63.

EVQLVESGGGLVQPFGGSLRLSCAASGFTFTNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDT SKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLVTVSSGGGGGGGGGGGGGGGGGG
QAVVTQEP SLTVSPGGTVTLTCGSSTGAVTTSNYANWVQQKPGQAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTLSGAQPEDEAEYYCALWYSNLWVFGGGTKLTVLGSHHHHHH

SEQ ID NO. 64.

EVQLVESGGGLVQPFGGSLRLSCAASGFTFTNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDT SKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLVTVSSGGGGGGGGGGGGGGGGGG
QAVVTQEP SLTVSPGGTVTLTCGSSTGAVTTSNYANWVQQKPGQAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTLSGAQPEDEAEYYCALWYSNLWVFGGGTKLTVL

SEQ ID NO. 65.

EVQLVESGGGLVQPFGGSLRLSCAASGFTFTNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDT SKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLVTV

SEQ ID NO. 66.

QAVVTQEP SLTVSPGGTVTLTCGSSTGAVTTSNYANWVQQKPGQAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTLSGAQPEDEAEYYCALWYSNLWVFGGGTKLTVL

Figure 26N

XENP11823 H1.28_L1.4

SEQ ID NO. 107.

EVQLVESGGGLVQPGGSLRLSCAASGFTFTNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFNGNSYVSWFAYWGQGT~~TVTVSS~~
QAVVTQEPSLTVSPGGTVTLTCSSTGAVTTSNYANWVQQKPGQAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTLSGAQPEDEAEYYCALWYSNLWVFGGGTKLTVLGSHHHHHH

SEQ ID NO. 108.

EVQLVESGGGLVQPGGSLRLSCAASGFTFTNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFNGNSYVSWFAYWGQGT~~TVTVSS~~
QAVVTQEPSLTVSPGGTVTLTCSSTGAVTTSNYANWVQQKPGQAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTLSGAQPEDEAEYYCALWYSNLWVFGGGTKLTVL

SEQ ID NO. 109.

EVQLVESGGGLVQPGGSLRLSCAASGFTFTNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFNGNSYVSWFAYWGQGT~~TVTVSS~~

SEQ ID NO. 110.

QAVVTQEPSLTVSPGGTVTLTCSSTGAVTTSNYANWVQQKPGQAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTLSGAQPEDEAEYYCALWYSNLWVFGGGTKLTVL

XENP11824 H1.29_L1.30

SEQ ID NO. 111.

EVQLVESGGGLVQPGGSLRLSCAASGFTFTNTYAMNWVRQAPGKSLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFNGNSYVSWFAYWGQGT~~TVTVSS~~
QAVVTQEPSLTVSPGGTVTLTCSSTGAVTTSNYANWVQQKPGQAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTLSGAQPEDEAEYYCALWYSNLWVFGGGTKLTVLGSHHHHHH

SEQ ID NO. 112.

EVQLVESGGGLVQPGGSLRLSCAASGFTFTNTYAMNWVRQAPGKSLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFNGNSYVSWFAYWGQGT~~TVTVSS~~
QAVVTQEPSLTVSPGGTVTLTCSSTGAVTTSNYANWVQQKPGQAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTLSGAQPEDEAEYYCALWYSNLWVFGGGTKLTVL

SEQ ID NO. 113.

EVQLVESGGGLVQPGGSLRLSCAASGFTFTNTYAMNWVRQAPGKSLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFNGNSYVSWFAYWGQGT~~TVTVSS~~

SEQ ID NO. 114.

QAVVTQEPSLTVSPGGTVTLTCSSTGAVTTSNYANWVQQKPGQAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTLSGAQPEDEAEYYCALWYSNLWVFGGGTKLTVL

Figure 26T

XENP11835 H1_L1.20

SEQ ID NO. 155.

EVQLVESGGGLVQPGGSLRLSCAASGFTFNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDS KNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLVTVSSGGGGSGGGSGGGG
QAVVTQEP SLTVSPGGTVTLTCGSSTGAVTTSNYANWVQKPGQAPRGLIGGTNKRAPGV PDRFSGSLLG
GKAALTL SGAQPEDEAEYYCALWYSNLWVFGGGTKLTVLGSHHHHHH

SEQ ID NO. 156.

EVQLVESGGGLVQPGGSLRLSCAASGFTFNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDS KNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLVTVSSGGGGSGGGSGGGG
QAVVTQEP SLTVSPGGTVTLTCGSSTGAVTTSNYANWVQKPGQAPRGLIGGTNKRAPGV PDRFSGSLLG
GKAALTL SGAQPEDEAEYYCALWYSNLWVFGGGTKLTVL

SEQ ID NO. 157.

EVQLVESGGGLVQPGGSLRLSCAASGFTFNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDS KNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLVTVSS

SEQ ID NO. 158.

QAVVTQEP SLTVSPGGTVTLTCGSSTGAVTTSNYANWVQKPGQAPRGLIGGTNKRAPGV PDRFSGSLLG
GKAALTL SGAQPEDEAEYYCALWYSNLWVFGGGTKLTVL

XENP11836 H1_L1.21

SEQ ID NO. 159.

EVQLVESGGGLVQPGGSLRLSCAASGFTFNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDS KNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLVTVSSGGGGSGGGSGGGG
QAVVTQEP SLTVSPGGTVTLTCGSSTGAVTTSNYANWVQKPGQAPRGLIGGTNKRAPGV PARFSGSKSG
GKAALTL SGAQPEDEAEYYCALWYSNLWVFGGGTKLTVLGSHHHHHH

SEQ ID NO. 160.

EVQLVESGGGLVQPGGSLRLSCAASGFTFNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDS KNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLVTVSSGGGGSGGGSGGGG
QAVVTQEP SLTVSPGGTVTLTCGSSTGAVTTSNYANWVQKPGQAPRGLIGGTNKRAPGV PARFSGSKSG
GKAALTL SGAQPEDEAEYYCALWYSNLWVFGGGTKLTVL

SEQ ID NO. 161.

EVQLVESGGGLVQPGGSLRLSCAASGFTFNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDS KNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLVTVSS

SEQ ID NO. 162.

QAVVTQEP SLTVSPGGTVTLTCGSSTGAVTTSNYANWVQKPGQAPRGLIGGTNKRAPGV PARFSGSKSG
GKAALTL SGAQPEDEAEYYCALWYSNLWVFGGGTKLTVL

Figure 26U

XENP11837 H1_L1.22

SEQ ID NO. 163.

EVQLVESGGGLVQPGGSLRLSCAASGFTFTNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLVTVSSGGGSGGGSGGGG
QAVVTQEPSTLVSPGGTVTLTCSSTGAVTTSNYANWVQKPGQAPRGLIGGTNKRAPGVPARFSGSSSG
GKAALTLSGAQPEDEAEYYCALWYSNLWVFGGGTKLTVLGSHHHHHH

SEQ ID NO. 164.

EVQLVESGGGLVQPGGSLRLSCAASGFTFTNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLVTVSSGGGSGGGSGGGG
QAVVTQEPSTLVSPGGTVTLTCSSTGAVTTSNYANWVQKPGQAPRGLIGGTNKRAPGVPARFSGSSSG
GKAALTLSGAQPEDEAEYYCALWYSNLWVFGGGTKLTVL

SEQ ID NO. 165.

EVQLVESGGGLVQPGGSLRLSCAASGFTFTNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLVTVSS

SEQ ID NO. 166.

QAVVTQEPSTLVSPGGTVTLTCSSTGAVTTSNYANWVQKPGQAPRGLIGGTNKRAPGVPARFSGSSSG
GKAALTLSGAQPEDEAEYYCALWYSNLWVFGGGTKLTVL

XENP11838 H1_L1.23

SEQ ID NO. 167.

EVQLVESGGGLVQPGGSLRLSCAASGFTFTNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLVTVSSGGGSGGGSGGGG
QAVVTQEPSTLVSPGGTVTLTCSSTGAVTTSNYANWVQKPGQAPRGLIGGTNKRAPGVPARFSGSLG
GKAALTLSGAQPEDEAEYYCALWYSNLWVFGGGTKLTVLGSHHHHHH

SEQ ID NO. 168.

EVQLVESGGGLVQPGGSLRLSCAASGFTFTNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLVTVSSGGGSGGGSGGGG
QAVVTQEPSTLVSPGGTVTLTCSSTGAVTTSNYANWVQKPGQAPRGLIGGTNKRAPGVPARFSGSLG
GKAALTLSGAQPEDEAEYYCALWYSNLWVFGGGTKLTVL

SEQ ID NO. 169.

EVQLVESGGGLVQPGGSLRLSCAASGFTFTNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLVTVSS

SEQ ID NO. 170.

QAVVTQEPSTLVSPGGTVTLTCSSTGAVTTSNYANWVQKPGQAPRGLIGGTNKRAPGVPARFSGSLG
GKAALTLSGAQPEDEAEYYCALWYSNLWVFGGGTKLTVL

Figure 26X

XENP11843 H1_L1.28

SEQ ID NO. 187.

EVQLVESGGGLVQPGGSLRLSCAASGFTFTNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLVTVSSGGGGSGGGGGGGGGG
QAVVTQEPSTLVSPGGTVTLTCSSTGAVTTSNYANWVQKPGQAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTLSGAQPEDEAEYYCALWYSNHWVFGGGTKLTVLGSHHHHHH

SEQ ID NO. 188.

EVQLVESGGGLVQPGGSLRLSCAASGFTFTNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLVTVSSGGGGSGGGGGGGGGG
QAVVTQEPSTLVSPGGTVTLTCSSTGAVTTSNYANWVQKPGQAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTLSGAQPEDEAEYYCALWYSNHWVFGGGTKLTVL

SEQ ID NO. 189.

EVQLVESGGGLVQPGGSLRLSCAASGFTFTNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLVTVSS

SEQ ID NO. 190.

QAVVTQEPSTLVSPGGTVTLTCSSTGAVTTSNYANWVQKPGQAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTLSGAQPEDEAEYYCALWYSNHWVFGGGTKLTVL

XENP11844 H1_L1.29

SEQ ID NO. 191.

EVQLVESGGGLVQPGGSLRLSCAASGFTFTNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLVTVSSGGGGSGGGGGGGGGG
QAVVTQEPSTLVSPGGTVTLTCSSTGAVTTSNYANWVQKPGQAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTLSGAQPEDEAEYYCALWYSNLWVFGGGTKLEIKGSHHHHHH

SEQ ID NO. 192.

EVQLVESGGGLVQPGGSLRLSCAASGFTFTNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLVTVSSGGGGSGGGGGGGGGG
QAVVTQEPSTLVSPGGTVTLTCSSTGAVTTSNYANWVQKPGQAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTLSGAQPEDEAEYYCALWYSNLWVFGGGTKLEIKGS

SEQ ID NO. 193.

EVQLVESGGGLVQPGGSLRLSCAASGFTFTNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLVTVSS

SEQ ID NO. 194.

QAVVTQEPSTLVSPGGTVTLTCSSTGAVTTSNYANWVQKPGQAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTLSGAQPEDEAEYYCALWYSNLWVFGGGTKLEIKGS

Figure 26Z

XENP11922 H1_L1.31

SEQ ID NO. 203.

EVQLVESGGGLVQPGGSLRLSCAASGFTFNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLVTVSS
QAVVTQEPSTLVSPGGTVTLTCGSSTGAVTTSNYANWVQKPGQAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTDSGAQPEDEADYYCALWYSNHWFVGGGTKLTVLGSHHHHHH

SEQ ID NO. 204.

EVQLVESGGGLVQPGGSLRLSCAASGFTFNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLVTVSS
QAVVTQEPSTLVSPGGTVTLTCGSSTGAVTTSNYANWVQKPGQAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTDSGAQPEDEADYYCALWYSNHWFVGGGTKLTVL

SEQ ID NO. 205.

EVQLVESGGGLVQPGGSLRLSCAASGFTFNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLVTVSS

SEQ ID NO. 206.

QAVVTQEPSTLVSPGGTVTLTCGSSTGAVTTSNYANWVQKPGQAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTDSGAQPEDEADYYCALWYSNHWFVGGGTKLTVL

XENP11923 H1_L1.32

SEQ ID NO. 207.

EVQLVESGGGLVQPGGSLRLSCAASGFTFNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLVTVSS
QAVVTQEPSTLVSPGGTVTLTCGSSTGAVTTSNYANWVQKPGQAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTLGAQPEDEAEYYCALWYSNLWVFGGKTKLTVLGSHHHHHH

SEQ ID NO. 208.

EVQLVESGGGLVQPGGSLRLSCAASGFTFNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLVTVSS
QAVVTQEPSTLVSPGGTVTLTCGSSTGAVTTSNYANWVQKPGQAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTLGAQPEDEAEYYCALWYSNLWVFGGKTKLTVL

SEQ ID NO. 209.

EVQLVESGGGLVQPGGSLRLSCAASGFTFNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLVTVSS

SEQ ID NO. 210.

QAVVTQEPSTLVSPGGTVTLTCGSSTGAVTTSNYANWVQKPGQAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTLGAQPEDEAEYYCALWYSNLWVFGGKTKLTVL

Figure 26BB

XENP12109 H1_L1.34

SEQ ID NO. 219.

EVQLVESGGGLVQPGGSLRLSCAASGFTFTNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGT~~LVTVSS~~
QAVVTQEP~~SLTVSPGGTVTLTCGSSTGAVTTSNYANWVQKPGTAPRGLIGGT~~NKRAPGV~~PARFSGSLLG~~
GKAALTLSGAQPEDEAEYYCALWY~~SNLWVFGGGTKLTVLG~~SHHHHHH

SEQ ID NO. 220.

EVQLVESGGGLVQPGGSLRLSCAASGFTFTNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGT~~LVTVSS~~
QAVVTQEP~~SLTVSPGGTVTLTCGSSTGAVTTSNYANWVQKPGTAPRGLIGGT~~NKRAPGV~~PARFSGSLLG~~
GKAALTLSGAQPEDEAEYYCALWY~~SNLWVFGGGTKLTVL~~

SEQ ID NO. 221.

EVQLVESGGGLVQPGGSLRLSCAASGFTFTNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGT~~LVTVSS~~

SEQ ID NO. 222.

QAVVTQEP~~SLTVSPGGTVTLTCGSSTGAVTTSNYANWVQKPGTAPRGLIGGT~~NKRAPGV~~PARFSGSLLG~~
GKAALTLSGAQPEDEAEYYCALWY~~SNLWVFGGGTKLTVL~~

XENP12110 H1_L1.35

SEQ ID NO. 223.

EVQLVESGGGLVQPGGSLRLSCAASGFTFTNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGT~~LVTVSS~~
QAVVTQEP~~SLTVSPGGTVTLTCGSSTGAVTTSNYANWVQKPGSPPRGLIGGT~~NKRAPGV~~PARFSGSLLG~~
GKAALTLSGAQPEDEAEYYCALWY~~SNLWVFGGGTKLTVLG~~SHHHHHH

SEQ ID NO. 224.

EVQLVESGGGLVQPGGSLRLSCAASGFTFTNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGT~~LVTVSS~~
QAVVTQEP~~SLTVSPGGTVTLTCGSSTGAVTTSNYANWVQKPGSPPRGLIGGT~~NKRAPGV~~PARFSGSLLG~~
GKAALTLSGAQPEDEAEYYCALWY~~SNLWVFGGGTKLTVL~~

SEQ ID NO. 225.

EVQLVESGGGLVQPGGSLRLSCAASGFTFTNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGT~~LVTVSS~~

SEQ ID NO. 226.

QAVVTQEP~~SLTVSPGGTVTLTCGSSTGAVTTSNYANWVQKPGSPPRGLIGGT~~NKRAPGV~~PARFSGSLLG~~
GKAALTLSGAQPEDEAEYYCALWY~~SNLWVFGGGTKLTVL~~

Figure 26DD

XENP12113 H1_L1.38

SEQ ID NO. 235.

EVQLVESGGGLVQPGGSLRLSCAASGFTFNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLVTVSS
QAVVTQEPSLTVSPGGTVTLTCGSSTGAVTTSNYANWVQKPGQSPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTLSGAQPEDEAEYYCALWYSNLWVFGGGTKLTVLGSHHHHHH

SEQ ID NO. 236.

EVQLVESGGGLVQPGGSLRLSCAASGFTFNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLVTVSS
QAVVTQEPSLTVSPGGTVTLTCGSSTGAVTTSNYANWVQKPGQSPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTLSGAQPEDEAEYYCALWYSNLWVFGGGTKLTVL

SEQ ID NO. 237.

EVQLVESGGGLVQPGGSLRLSCAASGFTFNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLVTVSS

SEQ ID NO. 238.

QAVVTQEPSLTVSPGGTVTLTCGSSTGAVTTSNYANWVQKPGQSPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTLSGAQPEDEAEYYCALWYSNLWVFGGGTKLTVL

XENP12114 H1_L1.39

SEQ ID NO. 239.

EVQLVESGGGLVQPGGSLRLSCAASGFTFNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLVTVSS
QAVVTQEPSLTVSPGGTVTLTCGSSTGAVTTSNYANWVQKPGQDPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTLSGAQPEDEAEYYCALWYSNLWVFGGGTKLTVLGSHHHHHH

SEQ ID NO. 240.

EVQLVESGGGLVQPGGSLRLSCAASGFTFNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLVTVSS
QAVVTQEPSLTVSPGGTVTLTCGSSTGAVTTSNYANWVQKPGQDPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTLSGAQPEDEAEYYCALWYSNLWVFGGGTKLTVL

SEQ ID NO. 241.

EVQLVESGGGLVQPGGSLRLSCAASGFTFNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLVTVSS

SEQ ID NO. 242.

QAVVTQEPSLTVSPGGTVTLTCGSSTGAVTTSNYANWVQKPGQDPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTLSGAQPEDEAEYYCALWYSNLWVFGGGTKLTVL

Figure 26HH

XENP12137 H1.36_L1.4

SEQ ID NO. 267.

EVQLVESGGGLVQPGGSLRLSCAASGFTFNTYAMNWVRQAPGKGLEWVGRIRSNGGSTYYADSVKGRFTI
SRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLVTVSSGGGGGGGGGGGGQA
VVTQEPSLTVSPGGTVTLTCSSTGAVTTSNYANWVQKPGQAPRGLIGGTNKRAPGVPARFSGSLLGK
AALTLSGAQPEDEAEYYCALWYSNLWVFGGGTKLTVLGSHHHHHH

SEQ ID NO. 268.

EVQLVESGGGLVQPGGSLRLSCAASGFTFNTYAMNWVRQAPGKGLEWVGRIRSNGGSTYYADSVKGRFTI
SRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLVTVSSGGGGGGGGGGGGQA
VVTQEPSLTVSPGGTVTLTCSSTGAVTTSNYANWVQKPGQAPRGLIGGTNKRAPGVPARFSGSLLGK
AALTLSGAQPEDEAEYYCALWYSNLWVFGGGTKLTVL

SEQ ID NO. 269.

EVQLVESGGGLVQPGGSLRLSCAASGFTFNTYAMNWVRQAPGKGLEWVGRIRSNGGSTYYADSVKGRFTI
SRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLVTVSS

SEQ ID NO. 270.

QAVVTQEPSLTVSPGGTVTLTCSSTGAVTTSNYANWVQKPGQAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTLSGAQPEDEAEYYCALWYSNLWVFGGGTKLTVL

XENP12138 H1.37_L1.4

SEQ ID NO. 271.

EVQLVESGGGLVQPGGSLRLSCAASGFTFNTYAMNWVRKAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLVTVSSGGGGGGGGGGGG
QAVVTQEPSLTVSPGGTVTLTCSSTGAVTTSNYANWVQKPGQAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTLSGAQPEDEAEYYCALWYSNLWVFGGGTKLTVLGSHHHHHH

SEQ ID NO. 272.

EVQLVESGGGLVQPGGSLRLSCAASGFTFNTYAMNWVRKAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLVTVSSGGGGGGGGGGGG
QAVVTQEPSLTVSPGGTVTLTCSSTGAVTTSNYANWVQKPGQAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTLSGAQPEDEAEYYCALWYSNLWVFGGGTKLTVL

SEQ ID NO. 273.

EVQLVESGGGLVQPGGSLRLSCAASGFTFNTYAMNWVRKAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLVTVSS

SEQ ID NO. 274.

QAVVTQEPSLTVSPGGTVTLTCSSTGAVTTSNYANWVQKPGQAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTLSGAQPEDEAEYYCALWYSNLWVFGGGTKLTVL

Figure 26II

XENP12139 H1.37 L1.32

SEQ ID NO. 275.

EVQLVESGGGLVQPGGSLRLSCAASGFTFTNTYAMNWVRKAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGTILTVTSSS
QAVVTQEPSLTVSPGGTIVLTCSSTGAVTTSNYANWVQKEPGQAPRGLIGGTNKRAPGVPARFSGLLG
GKAALTLSGAOPEDEAEYYCALWYSNLWVFGGGTKITVLGSHHHHHH

SEO ID NO. 276.

[illegible]

SEO ID NO. 277.

EVQLVESGGGLVQPGGSLRLSCAASGFTFTNTYAMNWVRKAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLOMNSLRAEDTAVYYCVRHGNEFGNSYVSWFAYWGOGTLTVSS

SEQ ID NO. 278.

QAVVTQEPSLTVSPGGTVTLLTCGSSTGAVTTSNYANWVQEKPGQAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTLSGAOPDEAEYYCALWYSNLWVFGGGTKLTVL

XENP12149 H1 L1.40

SEQ ID NO. 279.

EVQLVESGGGLVQPGGSLKLSKAASGFTFTNTYAMNWRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAWYWGQGLTVTVSS
QAVVTQEPSLTVSPGGT~~TVTL~~TCGSSTGAVTTSNYANWVQKQPGQAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTISGAOPEDEADYYCALWYSNHWFVGGG~~TKLTVL~~GSHHHHHH

SEO ID NO. 280.

[illegible]

SEQ ID NO. 281.

EVQLVESGGGLVQPGGSLRLSCAASGFTFNTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLOMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGOGTLTVSS

SEQ ID NO. 282.

QAVVTQEPSLTVSPGGTVTLLTCGSSTGAVTTSNYANWVQKPGQAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTISGAOPEDADYYCALWYSNHHVFGGGTKLTVL

Figure 26NN

XENP12221 H1.30_L1.42

SEQ ID NO. 315.

EVQLVESGGGLVQPGGSLRLSCAASGFTTFSTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGDSYVSWFAYWGQGLTVTVSSGGGGSGGGSGGGGSGGGG
QAVVTQEPSLTVSPGGTVTLTCGSSTGAVTTSNYANWVQQKPGKAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTISGAQPEDEAEYYCALWYSNHWVFVGGGTKLTVLGSHHHHHH

SEQ ID NO. 316.

EVQLVESGGGLVQPGGSLRLSCAASGFTTFSTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGDSYVSWFAYWGQGLTVTVSSGGGGSGGGSGGGGSGGGG
QAVVTQEPSLTVSPGGTVTLTCGSSTGAVTTSNYANWVQQKPGKAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTISGAQPEDEAEYYCALWYSNHWVFVGGGTKLTVL

SEQ ID NO. 317.

EVQLVESGGGLVQPGGSLRLSCAASGFTTFSTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGDSYVSWFAYWGQGLTVTVSS

SEQ ID NO. 318.

QAVVTQEPSLTVSPGGTVTLTCGSSTGAVTTSNYANWVQQKPGKAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTISGAQPEDEAEYYCALWYSNHWVFVGGGTKLTVL

XENP12222 H1.8_L1.42

SEQ ID NO. 319.

EVQLVESGGGLVQPGGSLRLSCAASGFTTFSTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLTVTVSSGGGGSGGGSGGGGSGGGG
QAVVTQEPSLTVSPGGTVTLTCGSSTGAVTTSNYANWVQQKPGKAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTISGAQPEDEAEYYCALWYSNHWVFVGGGTKLTVLGSHHHHHH

SEQ ID NO. 320.

EVQLVESGGGLVQPGGSLRLSCAASGFTTFSTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLTVTVSSGGGGSGGGSGGGGSGGGG
QAVVTQEPSLTVSPGGTVTLTCGSSTGAVTTSNYANWVQQKPGKAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTISGAQPEDEAEYYCALWYSNHWVFVGGGTKLTVL

SEQ ID NO. 321.

EVQLVESGGGLVQPGGSLRLSCAASGFTTFSTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLTVTVSS

SEQ ID NO. 322.

QAVVTQEPSLTVSPGGTVTLTCGSSTGAVTTSNYANWVQQKPGKAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTISGAQPEDEAEYYCALWYSNHWVFVGGGTKLTVL

Figure 2600

XENP12223 H1.39_L1.42

SEQ ID NO. 323.

EVQLVESGGGLVQPGGSLRLSCAASGFTFSTYAMNWVRKAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFNGNSYVSWFAYWGQGLVTVSSGGGGSGGGGSGGGG
QAVVTQEPSLTVSPGGTVTLTCSSTGAVTTSNYANWVQQKPGKAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTISGAQPEDEAEYYCALWYSNHWFVGGGTKLTVLGSHHHHHH

SEQ ID NO. 324.

EVQLVESGGGLVQPGGSLRLSCAASGFTFSTYAMNWVRKAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFNGNSYVSWFAYWGQGLVTVSSGGGGSGGGGSGGGG
QAVVTQEPSLTVSPGGTVTLTCSSTGAVTTSNYANWVQQKPGKAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTISGAQPEDEAEYYCALWYSNHWFVGGGTKLTVL

SEQ ID NO. 325.

EVQLVESGGGLVQPGGSLRLSCAASGFTFSTYAMNWVRKAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFNGNSYVSWFAYWGQGLVTVSS

SEQ ID NO. 326.

QAVVTQEPSLTVSPGGTVTLTCSSTGAVTTSNYANWVQQKPGKAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTISGAQPEDEAEYYCALWYSNHWFVGGGTKLTVL

XENP12224 H1.40_L1.42

SEQ ID NO. 327.

EVQLVESGGGLVQPGGSLRLSCAASGFTFSTYAMNWVRKAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGDSYVSWFAYWGQGLVTVSSGGGGSGGGGSGGGG
QAVVTQEPSLTVSPGGTVTLTCSSTGAVTTSNYANWVQQKPGKAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTISGAQPEDEAEYYCALWYSNHWFVGGGTKLTVLGSHHHHHH

SEQ ID NO. 328.

EVQLVESGGGLVQPGGSLRLSCAASGFTFSTYAMNWVRKAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGDSYVSWFAYWGQGLVTVSSGGGGSGGGGSGGGG
QAVVTQEPSLTVSPGGTVTLTCSSTGAVTTSNYANWVQQKPGKAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTISGAQPEDEAEYYCALWYSNHWFVGGGTKLTVL

SEQ ID NO. 329.

EVQLVESGGGLVQPGGSLRLSCAASGFTFSTYAMNWVRKAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGDSYVSWFAYWGQGLVTVSS

SEQ ID NO. 330.

QAVVTQEPSLTVSPGGTVTLTCSSTGAVTTSNYANWVQQKPGKAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTISGAQPEDEAEYYCALWYSNHWFVGGGTKLTVL

Figure 26UU

XENP12235 H1.39_L1.45

SEQ ID NO. 371.

EVQLVESGGGLVQPGGSLRLSCAASGFTTFSTYAMNWVRKAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLTVTVSSGGGGSGGGSGGGG
QAVVTQEPSLTVSPGGTVTLTCGSSTGAVTTSNYANWVQQKPGKAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTISGAQPEDEADYYCALWYSNHWVFEGGGTKLTVLGSHHHHHH

SEQ ID NO. 372.

EVQLVESGGGLVQPGGSLRLSCAASGFTTFSTYAMNWVRKAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLTVTVSSGGGGSGGGSGGGG
QAVVTQEPSLTVSPGGTVTLTCGSSTGAVTTSNYANWVQQKPGKAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTISGAQPEDEADYYCALWYSNHWVFEGGGTKLTVL

SEQ ID NO. 373.

EVQLVESGGGLVQPGGSLRLSCAASGFTTFSTYAMNWVRKAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLTVTVSS

SEQ ID NO. 374.

QAVVTQEPSLTVSPGGTVTLTCGSSTGAVTTSNYANWVQQKPGKAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTISGAQPEDEADYYCALWYSNHWVFEGGGTKLTVL

XENP12236 H1.40_L1.45

SEQ ID NO. 375.

EVQLVESGGGLVQPGGSLRLSCAASGFTTFSTYAMNWVRKAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGDSYVSWFAYWGQGLTVTVSSGGGGSGGGSGGGG
QAVVTQEPSLTVSPGGTVTLTCGSSTGAVTTSNYANWVQQKPGKAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTISGAQPEDEADYYCALWYSNHWVFEGGGTKLTVLGSHHHHHH

SEQ ID NO. 376.

EVQLVESGGGLVQPGGSLRLSCAASGFTTFSTYAMNWVRKAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGDSYVSWFAYWGQGLTVTVSSGGGGSGGGSGGGG
QAVVTQEPSLTVSPGGTVTLTCGSSTGAVTTSNYANWVQQKPGKAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTISGAQPEDEADYYCALWYSNHWVFEGGGTKLTVL

SEQ ID NO. 377.

EVQLVESGGGLVQPGGSLRLSCAASGFTTFSTYAMNWVRKAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGDSYVSWFAYWGQGLTVTVSS

SEQ ID NO. 378.

QAVVTQEPSLTVSPGGTVTLTCGSSTGAVTTSNYANWVQQKPGKAPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTISGAQPEDEADYYCALWYSNHWVFEGGGTKLTVL

Figure 26VV

XENP12237 H1.30_L1.46

SEQ ID NO. 379.

EVQLVESGGGLVQPGGSLRLSCAASGFTTFSTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDS KNTLYLQMNSLRAEDTAVYYCVRHGNFGDSYVSWFAYWGQGLTVTVSS
QAVVTQEP SLTVSPGGTVTLT CGSSTGAVTTSNYANWVQKPGQSPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTISGAQPEDEADYYCALWYSNHWFVGGGTKLTVLGSHHHHHH

SEQ ID NO. 380.

EVQLVESGGGLVQPGGSLRLSCAASGFTTFSTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDS KNTLYLQMNSLRAEDTAVYYCVRHGNFGDSYVSWFAYWGQGLTVTVSS
QAVVTQEP SLTVSPGGTVTLT CGSSTGAVTTSNYANWVQKPGQSPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTISGAQPEDEADYYCALWYSNHWFVGGGTKLTVL

SEQ ID NO. 381.

EVQLVESGGGLVQPGGSLRLSCAASGFTTFSTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDS KNTLYLQMNSLRAEDTAVYYCVRHGNFGDSYVSWFAYWGQGLTVTVSS

SEQ ID NO. 382.

QAVVTQEP SLTVSPGGTVTLT CGSSTGAVTTSNYANWVQKPGQSPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTISGAQPEDEADYYCALWYSNHWFVGGGTKLTVL

XENP12238 H1.8_L1.46

SEQ ID NO. 383.

EVQLVESGGGLVQPGGSLRLSCAASGFTTFSTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDS KNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLTVTVSS
QAVVTQEP SLTVSPGGTVTLT CGSSTGAVTTSNYANWVQKPGQSPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTISGAQPEDEADYYCALWYSNHWFVGGGTKLTVLGSHHHHHH

SEQ ID NO. 384.

EVQLVESGGGLVQPGGSLRLSCAASGFTTFSTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDS KNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLTVTVSS
QAVVTQEP SLTVSPGGTVTLT CGSSTGAVTTSNYANWVQKPGQSPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTISGAQPEDEADYYCALWYSNHWFVGGGTKLTVL

SEQ ID NO. 385.

EVQLVESGGGLVQPGGSLRLSCAASGFTTFSTYAMNWVRQAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDS KNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLTVTVSS

SEQ ID NO. 386.

QAVVTQEP SLTVSPGGTVTLT CGSSTGAVTTSNYANWVQKPGQSPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTISGAQPEDEADYYCALWYSNHWFVGGGTKLTVL

Figure 26WW

XENP12239 H1.39_L1.46

SEQ ID NO. 387.

EVQLVESGGGLVQPGGSLRLSCAASGFTFSTYAMNWVRKAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFNGNSYVSWFAYWGQGLVTVSS
QAVVTQEPSLTVSPGGTVTLTCSSTGAVTTSNYANWVQKPGQSPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTISGAQPEDEADYYCALWYSNHWVFGGGTKLTVLGSHHHHH

SEQ ID NO. 388.

EVQLVESGGGLVQPGGSLRLSCAASGFTFSTYAMNWVRKAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFNGNSYVSWFAYWGQGLVTVSS
QAVVTQEPSLTVSPGGTVTLTCSSTGAVTTSNYANWVQKPGQSPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTISGAQPEDEADYYCALWYSNHWVFGGGTKLTVL

SEQ ID NO. 389.

EVQLVESGGGLVQPGGSLRLSCAASGFTFSTYAMNWVRKAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFNGNSYVSWFAYWGQGLVTVSS

SEQ ID NO. 390.

QAVVTQEPSLTVSPGGTVTLTCSSTGAVTTSNYANWVQKPGQSPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTISGAQPEDEADYYCALWYSNHWVFGGGTKLTVL

XENP12240 H1.40_L1.46

SEQ ID NO. 391.

EVQLVESGGGLVQPGGSLRLSCAASGFTFSTYAMNWVRKAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGDSYVSWFAYWGQGLVTVSS
QAVVTQEPSLTVSPGGTVTLTCSSTGAVTTSNYANWVQKPGQSPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTISGAQPEDEADYYCALWYSNHWVFGGGTKLTVLGSHHHHH

SEQ ID NO. 392.

EVQLVESGGGLVQPGGSLRLSCAASGFTFSTYAMNWVRKAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGDSYVSWFAYWGQGLVTVSS
QAVVTQEPSLTVSPGGTVTLTCSSTGAVTTSNYANWVQKPGQSPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTISGAQPEDEADYYCALWYSNHWVFGGGTKLTVL

SEQ ID NO. 393.

EVQLVESGGGLVQPGGSLRLSCAASGFTFSTYAMNWVRKAPGKGLEWVGRIRSKYNNYATYYADSVKGRF
TISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGDSYVSWFAYWGQGLVTVSS

SEQ ID NO. 394.

QAVVTQEPSLTVSPGGTVTLTCSSTGAVTTSNYANWVQKPGQSPRGLIGGTNKRAPGVPARFSGSLLG
GKAALTISGAQPEDEADYYCALWYSNHWVFGGGTKLTVL

Figure 27A

XENP	VH ID	VL ID	Yield (mg/L)	T _m (°C)
11502	H1	L1.4	13.39	62.7
11794	H1.3	L1.4	4.00	62
11795	H1.4	L1.4	18.01	62.5
11796	H1.5	L1.4	0.15	61.5
11801	H1.6	L1.4	2.13	62.5
11802	H1.7	L1.4	14.66	62.25
11803	H1.8	L1.4	1.85	64.125
11804	H1.9	L1.4	20.57	65
11805	H1.10	L1.4	29.45	68.5
11806	H1.11	L1.4	1.71	59.75
11807	H1.12	L1.4	6.27	63.375
11808	H1.13	L1.4	9.70	62.375
11809	H1.14	L1.4	3.22	63
11810	H1.15	L1.4	15.38	62
11811	H1.16	L1.4	3.98	58
11812	H1.17	L1.4	0.16	60
11813	H1.18	L1.4	9.09	61
11814	H1.19	L1.4	2.53	62
11815	H1.20	L1.4	5.29	62.5
11816	H1.21	L1.4	19.89	62
11817	H1.22	L1.4	7.38	59
11818	H1.23	L1.4	7.90	62
11819	H1.24	L1.4	3.10	63.125
11820	H1.25	L1.4	4.57	61.875
11821	H1.26	L1.4	57.12	61.5
11822	H1.27	L1.4	25.53	65.5
11823	H1.28	L1.4	6.66	62.5
11824	H1.29	L1.30	30.48	61
11825	H1	L1.10	18.02	62
11826	H1	L1.11	6.36	62.5
11827	H1	L1.12	19.82	61.75
11828	H1	L1.13	2.22	62
11829	H1	L1.14	20.47	62.5
11830	H1	L1.15	20.63	62
11831	H1	L1.16	34.29	61.75
11832	H1	L1.17	17.89	62.25
11833	H1	L1.18	3.85	60
11834	H1	L1.19	6.02	60.75
11835	H1	L1.20	22.46	62.5
11836	H1	L1.21	8.16	55
11837	H1	L1.22	3.51	56.75
11838	H1	L1.23	15.90	63.5

Figure 27B

XENP	VH ID	VL ID	Yield (mg/L)	T _m (°C)
11839	H1	L1.24	15.94	61.75
11840	H1	L1.25	15.11	61.5
11841	H1	L1.26	7.30	64
11842	H1	L1.27	4.39	57.5
11843	H1	L1.28	15.61	62.875
11844	H1	L1.29	12.74	61.5
11920	H1	L3.1	0.72	52
11921	H1	L5.1	1.88	55.5
11922	H1	L1.31	8.89	58.25
11923	H1	L1.32	5.02	58.25
12107	H1.38	L1.4	19.28	62.5
12108	H1	L1.33	14.77	63
12109	H1	L1.34	10.77	62
12110	H1	L1.35	11.61	62.5
12111	H1	L1.36	17.51	62.5
12112	H1	L1.37	13.42	61
12113	H1	L1.38	13.03	63
12114	H1	L1.39	9.78	55.5
12131	H1.30	L1.4	12.73	65
12132	H1.31	L1.4	31.88	71
12133	H1.32	L1.4	24.00	65.5
12134	H1.33	L1.4	26.84	67.5
12135	H1.34	L1.4	40.40	72.5
12136	H1.35	L1.4	8.34	60.5
12137	H1.36	L1.4	4.30	60
12138	H1.37	L1.4	14.19	64.75
12139	H1.37	L1.32	6.75	63.5
12149	H1	L1.40	58.86	65.5

Figure 28A

XENP	% Improvement from H1_L1.4
11796	125.1
11803	118.0
11804	12.2
11805	16.3
11807	31.6
11808	54.4
11809	40.2
11502	129.3
11819	69.2
11820	98.8
11822	7.6
11838	169.8
11841	95.1
11843	152.5
11794	112.1
11795	241.8
11801	77.1
11802	232.6
11806	93.8
11810	263.9
11811	42.4
11812	5.1
11813	107.4
11814	83.8
11815	173.6
11816	230.7
11817	46.4
11818	33.3
11823	127.3
11825	227.4
11826	137.3
11827	197.1
11828	82.3
11829	225.4
11830	206.7
11832	79.9
11833	35.7
11834	66.9
11835	194.7
11836	191.6

Figure 28B

XENP	% Improvement from H1_L1.4
11837	73.9
11839	212.5
11840	204.0
11842	3.4
11844	112.3
12131	29.0
12132	0.1
12133	10.4
12134	1.0
12136	64.4
12137	0.1
12138	108.9
12108	145.2
12113	112.9
12149	128.7

Figure 29

HUMAN T-cell surface glycoprotein CD3 epsilon chain SEQ ID NO: 459

MQSGTHWRVLGLCLLSVGWVGQDGNEEMGGITQTPYKVSISGTTVILTCPQYPG
SEILWQHNDKNIGGDEDDKNIGSDEDHLSLKEFSELEQSGYYVCYPRGSK
PEDANFYLYLRARVCENCMEMDVMSVATIVIVDICTGGLLLVYYWSK
NRKAKAKPVTRGAGAGGRQRGQNKERPPVPNPDYEPKRGQRDLYSGL
NQRRI

Figure 30A

Optimized scFv CD3 variable regions

Xen #	vhCDR1	vhCDR2	vhCDR3	vICDR1	vICDR2	vICDR3
XENP11502 H1_L1.4 SEQ ID NO: 4	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVKG SEQ ID NO: 413	HGNFGNSVVSWEFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLMV SEQ ID NO: 430
XENP11794 H1.3_L1.4 SEQ ID NO: 8	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVKG SEQ ID NO: 413	HGNFGNSVVSWEFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLMV SEQ ID NO: 430
XENP11795 H1.4_L1.4 SEQ ID NO: 12	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVKG SEQ ID NO: 413	HGNFGNSVVSWEFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLMV SEQ ID NO: 430
XENP11796 H1.5_L1.4 SEQ ID NO: 16	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVKG SEQ ID NO: 413	HGNFGNSVVSWEFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLMV SEQ ID NO: 430
XENP11801 H1.6_L1.4 SEQ ID NO: 20	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVKG SEQ ID NO: 413	HGNFGNSVVSWEFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLMV SEQ ID NO: 430
XENP11802 H1.7_L1.4 SEQ ID NO: 24	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVKG SEQ ID NO: 413	HGNFGNSVVSWEFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLMV SEQ ID NO: 430
XENP11803 H1.8_L1.4 SEQ ID NO: 28	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVKG SEQ ID NO: 413	HGNFGNSVVSWEFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLMV SEQ ID NO: 430
XENP11804 H1.9_L1.4 SEQ ID NO: 32	TYAMH SEQ ID NO: 432	RIRSKYNNYATYYADSVKG SEQ ID NO: 413	HGNFGNSVVSWEFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLMV SEQ ID NO: 430
XENP11805 H1.10_L1.4 SEQ ID NO: 36	TYAMS SEQ ID NO: 412	RIRSKYNNYATYYADSVKG SEQ ID NO: 413	HGNFGNSVVSWEFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLMV SEQ ID NO: 430
XENP11806 H1.11_L1.4 SEQ ID NO: 40	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVKG SEQ ID NO: 413	HGNFGNSVVSWEFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLMV SEQ ID NO: 430
XENP11807 H1.12_L1.4 SEQ ID NO: 44	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVKG SEQ ID NO: 414	HGNFGNSVVSWEFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLMV SEQ ID NO: 430

Figure 30B

Xen #	vhCDR1	vhCDR2	vhCDR3	viCDR1	viCDR2	viCDR3
XENP11808 H1.13.L1.4 SEQ ID NO: 48	TYAMN SEQ ID NO: 411	RIRSKANSYATYYADSVRG SEQ ID NO: 434	HGNFGNSVSWFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLWV SEQ ID NO: 430
XENP11809 H1.14.L1.4 SEQ ID NO: 52	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVRG SEQ ID NO: 429	HGNFGNSVSWFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLWV SEQ ID NO: 430
XENP11810 H1.15.L1.4 SEQ ID NO: 56	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYAASVRG SEQ ID NO: 431	HGNFGNSVSWFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLWV SEQ ID NO: 430
XENP11811 H1.16.L1.4 SEQ ID NO: 60	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVRG SEQ ID NO: 413	HGNFGNSVSWFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLWV SEQ ID NO: 430
XENP11812 H1.17.L1.4 SEQ ID NO: 64	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVRG SEQ ID NO: 413	HGNFGNSVSWFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLWV SEQ ID NO: 430
XENP11813 H1.18.L1.4 SEQ ID NO: 68	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVRG SEQ ID NO: 413	HGNFGNSVSWFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLWV SEQ ID NO: 430
XENP11814 H1.19.L1.4 SEQ ID NO: 72	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVRG SEQ ID NO: 413	HGNFGNSVSWFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLWV SEQ ID NO: 430
XENP11815 H1.20.L1.4 SEQ ID NO: 76	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVRG SEQ ID NO: 413	HGNFGNSVSWFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLWV SEQ ID NO: 430
XENP11816 H1.21.L1.4 SEQ ID NO: 80	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVRG SEQ ID NO: 413	HGNFGNSVSWFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLWV SEQ ID NO: 430
XENP11817 H1.22.L1.4 SEQ ID NO: 84	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVRG SEQ ID NO: 413	HGNFGNSVSWFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLWV SEQ ID NO: 430
XENP11818 H1.23.L1.4 SEQ ID NO: 88	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVRG SEQ ID NO: 413	HGNFGNSVSWFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLWV SEQ ID NO: 430
XENP11819 H1.24.L1.4 SEQ ID NO: 92	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVRG SEQ ID NO: 413	HGNFGNSVSWFAY SEQ ID NO: 417	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLWV SEQ ID NO: 430

Figure 30C

Xen #	vhCDR1	vhCDR2	vhCDR3	viCDR1	viCDR2	viCDR3
XENP11820 H1.25_L1.4 SEQ ID NO: 96	TYAMN SEQ ID NO: 411	RIRSKYNNIATYYADSVKG SEQ ID NO: 413	HGNFGNSYVSWFAY SEQ ID NO: 418	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLWV SEQ ID NO: 430
XENP11821 H1.26_L1.4 SEQ ID NO: 100	TYAMN SEQ ID NO: 411	RIRSKYNNIATYYADSVKG SEQ ID NO: 413	HGNFGNTYVSWFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLWV SEQ ID NO: 430
XENP11822 H1.27_L1.4 SEQ ID NO: 104	TYAMN SEQ ID NO: 411	RIRSKYNNIATYYADSVKG SEQ ID NO: 413	HGNFGNSYVSWFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLWV SEQ ID NO: 430
XENP11823 H1.28_L1.4 SEQ ID NO: 108	TYAMN SEQ ID NO: 411	RIRSKYNNIATYYADSVKG SEQ ID NO: 413	HGNFGNSYVSWFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLWV SEQ ID NO: 430
XENP11824 H1.29_L1.30 SEQ ID NO: 112	TYAMN SEQ ID NO: 411	RIRSKYNNIATYYADSVKG SEQ ID NO: 413	HGNFGNSYVSWFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLWV SEQ ID NO: 430
XENP11825 H1.L1.10 SEQ ID NO: 116	TYAMN SEQ ID NO: 411	RIRSKYNNIATYYADSVKG SEQ ID NO: 413	HGNFGNSYVSWFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLWV SEQ ID NO: 430
XENP11826 H1.L1.11 SEQ ID NO: 120	TYAMN SEQ ID NO: 411	RIRSKYNNIATYYADSVKG SEQ ID NO: 413	HGNFGNSYVSWFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLWV SEQ ID NO: 430
XENP11827 H1.L1.12 SEQ ID NO: 124	TYAMN SEQ ID NO: 411	RIRSKYNNIATYYADSVKG SEQ ID NO: 413	HGNFGNSYVSWFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLWV SEQ ID NO: 430
XENP11828 H1.L1.13 SEQ ID NO: 128	TYAMN SEQ ID NO: 411	RIRSKYNNIATYYADSVKG SEQ ID NO: 413	HGNFGNSYVSWFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLWV SEQ ID NO: 430
XENP11829 H1.L1.14 SEQ ID NO: 132	TYAMN SEQ ID NO: 411	RIRSKYNNIATYYADSVKG SEQ ID NO: 413	HGNFGNSYVSWFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 421	GTNKRAP SEQ ID NO: 425	ALWYSNLWV SEQ ID NO: 430
XENP11830 H1.L1.15 SEQ ID NO: 136	TYAMN SEQ ID NO: 411	RIRSKYNNIATYYADSVKG SEQ ID NO: 413	HGNFGNSYVSWFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 422	GTNKRAP SEQ ID NO: 425	ALWYSNLWV SEQ ID NO: 430
XENP11831 H1.L1.16 SEQ ID NO: 140	TYAMN SEQ ID NO: 411	RIRSKYNNIATYYADSVKG SEQ ID NO: 413	HGNFGNSYVSWFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	DTNKRAP SEQ ID NO: 426	ALWYSNLWV SEQ ID NO: 430

Figure 30D

Xen #	vhCDR1	vhCDR2	vhCDR3	vICDR1	vICDR2	vICDR3
XENP11832 H1_L1.17 SEQ ID NO: 144	TYAMN SEQ ID NO: 411	RIRSKYNNVATYYADSVKRG SEQ ID NO: 413	HGNFGNSYVSWFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 427	ALWYSNLWV SEQ ID NO: 430
XENP11833 H1_L1.18 SEQ ID NO: 148	TYAMN SEQ ID NO: 411	RIRSKYNNVATYYADSVKRG SEQ ID NO: 413	HGNFGNSYVSWFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAS SEQ ID NO: 428	ALWYSNLWV SEQ ID NO: 430
XENP11834 H1_L1.19 SEQ ID NO: 152	TYAMN SEQ ID NO: 411	RIRSKYNNVATYYADSVKRG SEQ ID NO: 413	HGNFGNSYVSWFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTSPKHS SEQ ID NO: 462	ALWYSNLWV SEQ ID NO: 430
XENP11835 H1_L1.20 SEQ ID NO: 156	TYAMN SEQ ID NO: 411	RIRSKYNNVATYYADSVKRG SEQ ID NO: 413	HGNFGNSYVSWFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLWV SEQ ID NO: 430
XENP11836 H1_L1.21 SEQ ID NO: 160	TYAMN SEQ ID NO: 411	RIRSKYNNVATYYADSVKRG SEQ ID NO: 413	HGNFGNSYVSWFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLWV SEQ ID NO: 430
XENP11837 H1_L1.22 SEQ ID NO: 164	TYAMN SEQ ID NO: 411	RIRSKYNNVATYYADSVKRG SEQ ID NO: 413	HGNFGNSYVSWFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLWV SEQ ID NO: 430
XENP11838 H1_L1.23 SEQ ID NO: 168	TYAMN SEQ ID NO: 411	RIRSKYNNVATYYADSVKRG SEQ ID NO: 413	HGNFGNSYVSWFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLWV SEQ ID NO: 430
XENP11839 H1_L1.24 SEQ ID NO: 172	TYAMN SEQ ID NO: 411	RIRSKYNNVATYYADSVKRG SEQ ID NO: 413	HGNFGNSYVSWFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLWV SEQ ID NO: 430
XENP11840 H1_L1.25 SEQ ID NO: 176	TYAMN SEQ ID NO: 411	RIRSKYNNVATYYADSVKRG SEQ ID NO: 413	HGNFGNSYVSWFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLWV SEQ ID NO: 430
XENP11841 H1_L1.26 SEQ ID NO: 180	TYAMN SEQ ID NO: 411	RIRSKYNNVATYYADSVKRG SEQ ID NO: 413	HGNFGNSYVSWFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLWV SEQ ID NO: 430
XENP11842 H1_L1.27 SEQ ID NO: 184	TYAMN SEQ ID NO: 411	RIRSKYNNVATYYADSVKRG SEQ ID NO: 413	HGNFGNSYVSWFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	LLWYSNLWV SEQ ID NO: 417
XENP11843 H1_L1.28 SEQ ID NO: 188	TYAMN SEQ ID NO: 411	RIRSKYNNVATYYADSVKRG SEQ ID NO: 413	HGNFGNSYVSWFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNHWV SEQ ID NO: 433

Figure 30E

Xen #	vHCDR1	vHCDR2	vHCDR3	vICDR1	vICDR2	vICDR3
XENP11844 H1_L1.29 SEQ ID NO: 192	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVKG SEQ ID NO: 413	HGNFGNSYVSWFAY SEQ ID NO: 416	GSSTGAVTTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLWV SEQ ID NO: 430
XENP11920 H1_L3.1 SEQ ID NO: 196	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVKG SEQ ID NO: 413	HGNFGNSYVSWFAY SEQ ID NO: 416	RSSTGAVTTTSNYAN SEQ ID NO: 423	GTNKRAP SEQ ID NO: 425	ALWYSNLWV SEQ ID NO: 430
XENP11921 H1_L5.1 SEQ ID NO: 200	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVKG SEQ ID NO: 413	HGNFGNSYVSWFAY SEQ ID NO: 416	KSSTGAVTTTSNYAN SEQ ID NO: 424	GTNKRAP SEQ ID NO: 425	ALWYSNLWV SEQ ID NO: 430
XENP11922 H1_L1.31 SEQ ID NO: 204	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVKG SEQ ID NO: 413	HGNFGNSYVSWFAY SEQ ID NO: 416	GSSTGAVTTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLWV SEQ ID NO: 430
XENP11923 H1_L1.32 SEQ ID NO: 208	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVKG SEQ ID NO: 413	HGNFGNSYVSWFAY SEQ ID NO: 416	GSSTGAVTTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLWV SEQ ID NO: 430
XENP12107 H1.38_L1.4 SEQ ID NO: 212	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVKG SEQ ID NO: 413	HGNFGNSYVSWFAY SEQ ID NO: 416	GSSTGAVTTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLWV SEQ ID NO: 430
XENP12108 H1_L1.33 SEQ ID NO: 216	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVKG SEQ ID NO: 413	HGNFGNSYVSWFAY SEQ ID NO: 416	GSSTGAVTTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLWV SEQ ID NO: 430
XENP12109 H1_L1.34 SEQ ID NO: 220	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVKG SEQ ID NO: 413	HGNFGNSYVSWFAY SEQ ID NO: 416	GSSTGAVTTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLWV SEQ ID NO: 430
XENP12110 H1_L1.35 SEQ ID NO: 224	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVKG SEQ ID NO: 413	HGNFGNSYVSWFAY SEQ ID NO: 416	GSSTGAVTTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLWV SEQ ID NO: 430
XENP12111 H1_L1.36 SEQ ID NO: 228	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVKG SEQ ID NO: 413	HGNFGNSYVSWFAY SEQ ID NO: 416	GSSTGAVTTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLWV SEQ ID NO: 430
XENP12112 H1_L1.37 SEQ ID NO: 232	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVKG SEQ ID NO: 413	HGNFGNSYVSWFAY SEQ ID NO: 416	GSSTGAVTTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLWV SEQ ID NO: 430
XENP12113 H1_L1.38 SEQ ID NO: 236	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVKG SEQ ID NO: 413	HGNFGNSYVSWFAY SEQ ID NO: 416	GSSTGAVTTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLWV SEQ ID NO: 430

Figure 30F

Xen #	vhCDR1	vhCDR2	vhCDR3	vICDR1	vICDR2	vICDR3
XENP12114 H1_L1.39 SEQ ID NO: 240	TYAMN SEQ ID NO: 411	RIRSKYNNVATYYADSVKG SEQ ID NO: 413	HGNFGNSYVSWFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLMV SEQ ID NO: 430
XENP12131 H1.30_L1.4 SEQ ID NO: 244	TYAMN SEQ ID NO: 411	RIRSKYNNVATYYADSVKG SEQ ID NO: 413	HGNFGDSYVSWFAY SEQ ID NO: 417	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLMV SEQ ID NO: 430
XENP12132 H1.31_L1.4 SEQ ID NO: 248	TYAMS SEQ ID NO: 412	RIRSKYNNVATYYADSVKG SEQ ID NO: 413	HGNFGDSYVSWFAY SEQ ID NO: 417	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLMV SEQ ID NO: 430
XENP12133 H1.32_L1.4 SEQ ID NO: 252	TYAMN SEQ ID NO: 411	RIRSKANNVATYYADSVKG SEQ ID NO: 414	HGNFGDSYVSWFAY SEQ ID NO: 417	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLMV SEQ ID NO: 430
XENP12134 H1.33_L1.4 SEQ ID NO: 256	TYAMN SEQ ID NO: 411	RIRSKYNNVATYYADSVKG SEQ ID NO: 413	HGNFGDSYVSWFDY SEQ ID NO: 419	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLMV SEQ ID NO: 430
XENP12135 H1.34_L1.4 SEQ ID NO: 260	TYAMS SEQ ID NO: 412	RIRSKANNVATYYADSVKG SEQ ID NO: 414	HGNFGDSYVSWFDY SEQ ID NO: 419	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLMV SEQ ID NO: 430
XENP12136 H1.35_L1.4 SEQ ID NO: 264	TYAMN SEQ ID NO: 411	RIRSKYNNVATYYADSVKG SEQ ID NO: 413	HGNFGNSYVSWFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLMV SEQ ID NO: 430
XENP12137 H1.36_L1.4 SEQ ID NO: 268	TYAMN SEQ ID NO: 411	RIRSGSGSTYYADSVKG SEQ ID NO: 415	HGNFGNSYVSWFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLMV SEQ ID NO: 430
XENP12138 H1.37_L1.4 SEQ ID NO: 272	TYAMN SEQ ID NO: 411	RIRSKYNNVATYYADSVKG SEQ ID NO: 413	HGNFGNSYVSWFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLMV SEQ ID NO: 430
XENP12139 H1.37_L1.32 SEQ ID NO: 276	TYAMN SEQ ID NO: 411	RIRSKYNNVATYYADSVKG SEQ ID NO: 413	HGNFGNSYVSWFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLMV SEQ ID NO: 430
XENP12149 H1_L1.40 SEQ ID NO: 280	TYAMN SEQ ID NO: 411	RIRSKYNNVATYYADSVKG SEQ ID NO: 413	HGNFGNSYVSWFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLMV SEQ ID NO: 430
XENP12213 H1.30_L1.40 SEQ ID NO: 284	TYAMN SEQ ID NO: 411	RIRSKYNNVATYYADSVKG SEQ ID NO: 413	HGNFGDSYVSWFAY SEQ ID NO: 417	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNLMV SEQ ID NO: 430

Figure 30G

Xen #	VhCDR1	VhCDR2	VhCDR3	ViCDR1	ViCDR2	ViCDR3
XENP12214 H1.8 L1.40 SEQ ID NO: 288	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVKG SEQ ID NO: 413	HGNFGDSYVSWFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNHWV SEQ ID NO: 433
XENP12215 H1.39 L1.40 SEQ ID NO: 292	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVKG SEQ ID NO: 413	HGNFGDSYVSWFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNHWV SEQ ID NO: 433
XENP12216 H1.40 L1.40 SEQ ID NO: 296	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVKG SEQ ID NO: 413	HGNFGDSYVSWFAY SEQ ID NO: 417	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNHWV SEQ ID NO: 433
XENP12217 H1.30 L1.41 SEQ ID NO: 300	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVKG SEQ ID NO: 413	HGNFGDSYVSWFAY SEQ ID NO: 417	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNHWV SEQ ID NO: 433
XENP12218 H1.8 L1.41 SEQ ID NO: 304	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVKG SEQ ID NO: 413	HGNFGDSYVSWFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNHWV SEQ ID NO: 433
XENP12219 H1.39 L1.41 SEQ ID NO: 308	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVKG SEQ ID NO: 413	HGNFGDSYVSWFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNHWV SEQ ID NO: 433
XENP12220 H1.40 L1.41 SEQ ID NO: 312	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVKG SEQ ID NO: 413	HGNFGDSYVSWFAY SEQ ID NO: 417	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNHWV SEQ ID NO: 433
XENP12221 H1.30 L1.42 SEQ ID NO: 316	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVKG SEQ ID NO: 413	HGNFGDSYVSWFAY SEQ ID NO: 417	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNHWV SEQ ID NO: 433
XENP12222 H1.8 L1.42 SEQ ID NO: 320	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVKG SEQ ID NO: 413	HGNFGDSYVSWFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNHWV SEQ ID NO: 433
XENP12223 H1.39 L1.42 SEQ ID NO: 324	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVKG SEQ ID NO: 413	HGNFGDSYVSWFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNHWV SEQ ID NO: 433
XENP12224 H1.40 L1.42 SEQ ID NO: 328	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVKG SEQ ID NO: 413	HGNFGDSYVSWFAY SEQ ID NO: 417	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNHWV SEQ ID NO: 433
XENP12225 H1.30 L1.43 SEQ ID NO: 332	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVKG SEQ ID NO: 413	HGNFGDSYVSWFAY SEQ ID NO: 417	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNHWV SEQ ID NO: 433

Figure 30H

Xen #	vHCDR1	vHCDR2	vHCDR3	vICDR1	vICDR2	vICDR3
XENP12226 H1.8 L1.43 SEQ ID NO: 336	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVKVG SEQ ID NO: 413	HGNFGDSYVSWFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNHHV SEQ ID NO: 433
XENP12227 H1.39 L1.43 SEQ ID NO: 340	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVKVG SEQ ID NO: 413	HGNFGDSYVSWFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNHHV SEQ ID NO: 433
XENP12228 H1.40 L1.43 SEQ ID NO: 344	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVKVG SEQ ID NO: 413	HGNFGDSYVSWFAY SEQ ID NO: 417	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNHHV SEQ ID NO: 433
XENP12229 H1.30 L1.44 SEQ ID NO: 348	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVKVG SEQ ID NO: 413	HGNFGDSYVSWFAY SEQ ID NO: 417	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNHHV SEQ ID NO: 433
XENP12230 H1.8 L1.44 SEQ ID NO: 352	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVKVG SEQ ID NO: 413	HGNFGDSYVSWFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNHHV SEQ ID NO: 433
XENP12231 H1.39 L1.44 SEQ ID NO: 356	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVKVG SEQ ID NO: 413	HGNFGDSYVSWFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNHHV SEQ ID NO: 433
XENP12232 H1.40 L1.44 SEQ ID NO: 360	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVKVG SEQ ID NO: 413	HGNFGDSYVSWFAY SEQ ID NO: 417	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNHHV SEQ ID NO: 433
XENP12233 H1.30 L1.45 SEQ ID NO: 364	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVKVG SEQ ID NO: 413	HGNFGDSYVSWFAY SEQ ID NO: 417	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNHHV SEQ ID NO: 433
XENP12234 H1.8 L1.45 SEQ ID NO: 368	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVKVG SEQ ID NO: 413	HGNFGDSYVSWFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNHHV SEQ ID NO: 433
XENP12235 H1.39 L1.45 SEQ ID NO: 372	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVKVG SEQ ID NO: 413	HGNFGDSYVSWFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNHHV SEQ ID NO: 433
XENP12236 H1.40 L1.45 SEQ ID NO: 376	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVKVG SEQ ID NO: 413	HGNFGDSYVSWFAY SEQ ID NO: 417	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNHHV SEQ ID NO: 433
XENP12237 H1.30 L1.46 SEQ ID NO: 380	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVKVG SEQ ID NO: 413	HGNFGDSYVSWFAY SEQ ID NO: 417	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNHHV SEQ ID NO: 433

Figure 301

Xen #	vhCDR1	vhCDR2	vhCDR3	vICDR1	vICDR2	vICDR3
XENP12238 H1.8_L1.46 SEQ ID NO: 384	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVKRG SEQ ID NO: 413	HGNFGNSYVSWEFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNHHV SEQ ID NO: 433
XENP12239 H1.39_L1.46 SEQ ID NO: 388	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVKRG SEQ ID NO: 413	HGNFGNSYVSWEFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNHHV SEQ ID NO: 433
XENP12240 H1.40_L1.46 SEQ ID NO: 392	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVKRG SEQ ID NO: 413	HGNFGDSYVSWEFAY SEQ ID NO: 417	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNHHV SEQ ID NO: 433
XENP12241 H1.30_L1.47 SEQ ID NO: 396	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVKRG SEQ ID NO: 413	HGNFGDSYVSWEFAY SEQ ID NO: 417	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNHHV SEQ ID NO: 433
XENP12242 H1.8_L1.47 SEQ ID NO: 400	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVKRG SEQ ID NO: 413	HGNFGNSYVSWEFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNHHV SEQ ID NO: 433
XENP12243 H1.39_L1.47 SEQ ID NO: 404	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVKRG SEQ ID NO: 413	HGNFGNSYVSWEFAY SEQ ID NO: 416	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNHHV SEQ ID NO: 433
XENP12244 H1.40_L1.47 SEQ ID NO: 408	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVKRG SEQ ID NO: 413	HGNFGDSYVSWEFAY SEQ ID NO: 417	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNHHV SEQ ID NO: 433
XENP ??	TYAMN SEQ ID NO: 411	RIRSKYNNYATYYADSVKRG SEQ ID NO: 413	HGNFGDSYVSWEFAY SEQ ID NO: 417	GSSTGAVTTSNYAN SEQ ID NO: 420	GTNKRAP SEQ ID NO: 425	ALWYSNHHV SEQ ID NO: 433
Consensus sequences	TYAMXaa1, wherein Xaa1 is N, S or H (SEQ ID NO:435)	R-I-K-S-K-Xaa1-N- Xaa2-Y-A-T-Xaa3-Y-Y- A-Xaa4-S-V-K-G, wherein Xaa1 is Y or A, Xaa2 is N or S, Xaa3 is Y or A and Xaa4 is D or A (SEQ ID NO:436)	H-Q-N-F-G- Xaa1-S-Y-V-S- W-F-Xaa2-Y, wherein Xaa1 is N, D or Q and Xaa2 is A or D (SEQ ID NO:437)	Xaa1-S-S-T-G-A- V-T-Xaa2-Xaa3- Xaa4-Y-A-N, wherein Xaa1 is G, R or K, Xaa2 is T or S, Xaa3 is S or G and Xaa4 is N or H, (SEQ ID NO:438)	Xaa1-T-N- Xaa2-R-A- Xaa3, wherein Xaa1 is G or D, Xaa2 is K or N, and Xaa3 is P or S (SEQ ID NO:439)	Xaa1-L-W-Y- S-N-Xaa2-W- V, wherein Xaa1 is A or L and Xaa2 is L or H (SEQ ID NO:440)

1

HETERODIMERIC PROTEINS**PRIORITY**

This application is a continuation of U.S. patent application Ser. No. 14/155,344, filed Jan. 14, 2014 which claims priority to U.S. Provisional Patent Application Nos. 61/752,348 filed on Jan. 14, 2013, No. 61/780,310 filed on Mar. 13, 2013, No. 61/778,148 filed Mar. 12, 2013, and No. 61/764,954 filed Feb. 14, 2013, the entire contents of which are incorporated herein for all purposes by this reference.

SEQUENCE LISTING

The instant application contains a Sequence Listing which has been submitted electronically in ASCII format and is hereby incorporated by reference in its entirety. Said ASCII copy, created on Jun. 26, 2017, is named 067461-5161-US06_ST25.txt and is 763,759 bytes in size.

INCORPORATION OF RELATED APPLICATIONS

The following applications are incorporated by reference in their entirety, U.S. Ser. Nos. 61/302,707, 61/732,813, 61/598,686, 61/441,552, 61/311,472, 61/450,457, 61/545,498, 61/368,969, 61/391,509, 61/391,515, 61/439,263, 61/368,962 and 61/593,846, 61/515,745, 61/752,349, 13/194,904, 13/568,028, 13/648,951, 12/875,015, as well as US Publication Nos. 2009/0163,699 and 2006/0024298;

TECHNICAL FIELD

The present invention describes novel immunoglobulin compositions that simultaneously co-engage antigens, where both of the antigens are bound monovalently. The novel immunoglobulins described preferably utilize heterodimeric Fc regions. Methods of using the novel immunoglobulin compositions, particularly for therapeutic purposes, are also described herein.

BACKGROUND OF THE INVENTION

Antibody-based therapeutics have been used successfully to treat a variety of diseases, including cancer and autoimmune/inflammatory disorders. Yet improvements to this class of drugs are still needed, particularly with respect to enhancing their clinical efficacy. One avenue being explored is the engineering of additional and novel antigen binding sites into antibody-based drugs such that a single immunoglobulin molecule co-engages two different antigens. Such non-native or alternate antibody formats that engage two different antigens are often referred to as bispecifics. Because the considerable diversity of the antibody variable region (Fv) makes it possible to produce an Fv that recognizes virtually any molecule, the typical approach to bispecific generation is the introduction of new variable regions into the antibody.

A number of alternate antibody formats have been explored for bispecific targeting (Chames & Baty, 2009, mAbs 1[6]:1-9; Holliger & Hudson, 2005, Nature Biotechnology 23[9]:1126-1136; and Kontermann, 2012 MAb 4(2):182, all of which are expressly incorporated herein by reference). Initially, bispecific antibodies were made by fusing two cell lines that each produced a single monoclonal antibody (Milstein et al., 1983, Nature 305:537-540). Although the resulting hybrid hybridoma or quadroma did

2

produce bispecific antibodies, they were only a minor population, and extensive purification was required to isolate the desired antibody. An engineering solution to this was the use of antibody fragments to make bispecifics. Because such fragments lack the complex quaternary structure of a full length antibody, variable light and heavy chains can be linked in single genetic constructs. Antibody fragments of many different forms have been generated, including diabodies, single chain diabodies, tandem scFv's, and Fab2 bispecifics (Chames & Baty, 2009, mAbs 1[6]:1-9; Holliger & Hudson, 2005, Nature Biotechnology 23[9]:1126-1136; expressly incorporated herein by reference). While these formats can be expressed at high levels in bacteria and may have favorable penetration benefits due to their small size, they clear rapidly in vivo and can present manufacturing obstacles related to their production and stability. A principal cause of these drawbacks is that antibody fragments typically lack the constant region of the antibody with its associated functional properties, including larger size, high stability, and binding to various Fc receptors and ligands that maintain long half-life in serum (i.e., the neonatal Fc receptor FcRn) or serve as binding sites for purification (i.e., protein A and protein G).

More recent work has attempted to address the shortcomings of fragment-based bispecifics by engineering dual binding into full length antibody-like formats (Wu et al., 2007, Nature Biotechnology 25[11]:1290-1297; U.S. Ser. No. 12/477,711; Michaelson et al., 2009, mAbs 1[2]:128-141; PCT/US2008/074693; Zuo et al., 2000, Protein Engineering 13[5]:361-367; U.S. Ser. No. 09/865,198; Shen et al., 2006, J Biol Chem 281[16]:10706-10714; Lu et al., 2005, J Biol Chem 280[20]:19665-19672; PCT/US2005/025472; and Kontermann, 2012 MAb 4(2):182, all of which are expressly incorporated herein by reference). These formats overcome some of the obstacles of the antibody fragment bispecifics, principally because they contain an Fc region. One significant drawback of these formats is that, because they build new antigen binding sites on top of the homodimeric constant chains, binding to the new antigen is always bivalent.

For many antigens that are attractive as co-targets in a therapeutic bispecific format, the desired binding is monovalent rather than bivalent. For many immune receptors, cellular activation is accomplished by cross-linking of a monovalent binding interaction. The mechanism of cross-linking is typically mediated by antibody/antigen immune complexes, or via effector cell to target cell engagement. For example, the low affinity Fc gamma receptors (FcγRs) such as FcγRIIa, FcγRIIb, and FcγRIIIa bind monovalently to the antibody Fc region. Monovalent binding does not activate cells expressing these FcγRs; however, upon immune complexation or cell-to-cell contact, receptors are cross-linked and clustered on the cell surface, leading to activation. For receptors responsible for mediating cellular killing, for example FcγRIIIa on natural killer (NK) cells, receptor cross-linking and cellular activation occurs when the effector cell engages the target cell in a highly avid format (Bowles & Weiner, 2005, J Immunol Methods 304:88-99, expressly incorporated by reference). Similarly, on B cells the inhibitory receptor FcγRIIb downregulates B cell activation only when it engages into an immune complex with the cell surface B-cell receptor (BCR), a mechanism that is mediated by immune complexation of soluble IgG's with the same antigen that is recognized by the BCR (Heyman 2003, Immunol Lett 88[2]:157-161; Smith and Clatworthy, 2010, Nature Reviews Immunology 10:328-343; expressly incorporated by reference). As another example, CD3 activation

3

of T-cells occurs only when its associated T-cell receptor (TCR) engages antigen-loaded MHC on antigen presenting cells in a highly avid cell-to-cell synapse (Kuhns et al., 2006, *Immunity* 24:133-139). Indeed nonspecific bivalent cross-linking of CD3 using an anti-CD3 antibody elicits a cytokine storm and toxicity (Perruche et al., 2009, *J Immunol* 183 [2]:953-61; Chatenoud & Bluestone, 2007, *Nature Reviews Immunology* 7:622-632; expressly incorporated by reference). Thus for practical clinical use, the preferred mode of CD3 co-engagement for redirected killing of targets cells is monovalent binding that results in activation only upon engagement with the co-engaged target.

Thus while bispecifics generated from antibody fragments suffer biophysical and pharmacokinetic hurdles, a drawback of those built with full length antibody-like formats is that they engage co-target antigens multivalently in the absence of the primary target antigen, leading to nonspecific activation and potentially toxicity. The present invention solves this problem by introducing a novel bispecific format that enables the co-engagement of distinct target antigens.

SUMMARY OF THE INVENTION

Accordingly, in one aspect, the invention provides heterodimeric antibodies comprising a first heavy chain comprising a first Fc domain and a single chain Fv region that binds a first antigen. The heterodimeric antibodies also comprise a second heavy chain comprising a second Fc domain, a first variable heavy chain and a first variable light chain, wherein the first and second Fc domains are different.

In a further aspect, the pIs of the first heavy chain and the second heavy chain are at least 0.5 logs apart.

In an additional aspect, the heterodimeric antibodies of the invention further comprise one or two additional antigen binding domains that can be either or both of an scFv and a Fab.

In a further aspect, the heterodimeric antibodies of the invention comprise heterodimerization variants according to FIGS. 4, 5, 6, 20, 21 and 22.

In an additional aspect, the heterodimeric antibodies comprise pI variants in each monomer, with monomer one are ISO(-): I199T/N203D/K274Q/R355Q/N384S/K392N/V397M/Q419E/DEL44 and the pI substitutions in monomer 2 are ISO(+RR): Q196K/I199T/P217R/P228R/N276K.

In additional aspects, the present invention provides heterodimeric antibodies wherein one of the heavy chains comprises a variant selected from the group consisting of S119E, K133E, K133Q, R133E (in case of IgG2-4), R133Q (in case of IgG2-4), T164E, K205E, K205Q, N208D, K210E, K210Q, K274E, K320E, K322E, K326E, K334E, R355E, K392E, Deletion of K447, adding peptide DEDE at the c-terminus, G137E, N203D, K274Q, R355Q, K392N and Q419E, 349A, 349C, 349E, 349I, 349K, 349S, 349T, 349W, 351E, 351K, 354C, 356K, 357K, 364C, 364D, 364E, 364F, 364G, 364H, 364R, 364T, 364Y, 366D, 366K, 366S, 366W, 366Y, 368A, 368E, 368K, 368S, 370C, 370D, 370E, 370G, 370R, 370S, 370V, 392D, 392E, 394F, 394S, 394W, 394Y, 395T, 395V, 396T, 397E, 397S, 397T, 399K, 401K, 405A, 405S, 407T, 407V, 409D, 409E, 411D, 411E, 411K, 439D, 349C/364E, 349K/351K, 349K/351K/394F, 349K/354C, 349K/394F, 349K/394F/401K, 349K/394Y, 349K/401K, 349K/405A, 349T/351E/411E, 349T/394F, 349T/394F/401K, 349T/394F/411E, 349T/405A, 349T/411E, 351E/364D, 351E/364D/405A, 351E/364E, 351E/366D, 351K/364H/401K, 351K/366K, 364D/370G, 364D/394F, 364E/405A, 364E/405S, 364E/411E, 364E/411E/405A, 364H/394F, 364H/401K, 364H/401K/405A, 364H/405A,

4

364H/405A/411E, 364Y/370R, 370E/411E, 370R/411K, 395T/397S/405A and 397S/405A.

The present invention describes methods for generating the novel compositions of the invention. The present invention describes purification methods for the immunoglobulins herein, particularly methods for separating heterodimeric and homodimeric protein species. Also described are methods of testing the immunoglobulins herein, including in vitro and in vitro experiments.

The present invention provides isolated nucleic acids encoding the novel immunoglobulin compositions described herein. The present invention provides vectors comprising said nucleic acids, optionally, operably linked to control sequences. The present invention provides host cells containing the vectors, and methods for producing and optionally recovering the immunoglobulin compositions.

The present invention provides compositions comprising immunoglobulin polypeptides described herein, and a physiologically or pharmaceutically acceptable carrier or diluent.

The present invention contemplates therapeutic and diagnostic uses for the immunoglobulin polypeptides disclosed herein.

BRIEF DESCRIPTION OF THE DRAWINGS

Figures for Bottle Cap

FIG. 1A-1C. Illustration of the "triple F" format for bispecific immunoglobulins. FIG. 1A shows a scFv-Fc format. FIG. 1C depicts a more standard bispecific format, also utilizing the pI variants of the invention (and optionally and independently the other heterodimerization variants). FIG. 1B shows the "triple F" format (sometimes also referred to as the "bottle-opener" configuration; (and optionally and independently the other heterodimerization variants).

FIGS. 2A-2E depict a variety of "higher multispecificity" embodiments of the present invention. FIG. 2A shows a "triple F" configuration with another scFv attached to the Fab monomer (this one, along with FIG. 2D, has a greater molecular weight differential as well). FIG. 2B depicts a "triple F" with another scFv attached to the scFv monomer. FIG. 2C depicts a "three scFv" format. FIG. 2D depicts an additional Fab attached to the Fab monomer. FIG. 2E depicts a Fab hooked to one of the scFv monomers.

FIGS. 3A-3E show additional varieties of "higher multispecificity" embodiments of the "triple F" format, all with one monomer comprising an scFv and all of which have molecular weight differentials which can be exploited for purification of the heterodimers). FIG. 3A shows a "Fab-Fv" format with binding to two different antigens, with FIG. 3B depicting the "Fab-Fv" format with binding to a single antigen (e.g. bivalent binding to antigen 1). FIGS. 3C and 3D depicts a "Fv-Fab" format with similar bivalent or monovalent additional antigen binding. FIG. 3E depicts one monomer with a CH1-CL attached to the second scFv.

FIGS. 4A and 4B show novel steric variants. As will be understood by those in the art, the first column of each table represents "corresponding" monomer pairs or "sets": that is, monomer 1 has 405A and the corresponding steric variant is 394F. Again, these can be optionally and independently combined, as long as some "strandedness" is maintained.

FIG. 5A-5B depicts heterodimerization variants that find particular use in the present invention.

FIG. 6 depicts novel pI heterodimerization variants of particular use in the creation of heterodimeric proteins. As will be understood by those in the art, the first row of each table represents "corresponding" monomer pairs or sets. Note that the I199T and V397M variants are "neutral" (e.g.

5

not for the purposes of altering pI of one of the monomers). Rather, they contribute to the reduction of immunogenicity by helping preserve a close epitope; e.g. I199T follows pI variants N203D or Q196K, and V397M follows K392N. Thus these two variants can optionally and independently be removed from any of the sequences above.

FIG. 7A-7B. Amino acid sequences for an anti-CD19 Fab x anti-CD3 scFv “triple F” embodiment, showing the HC-Fab sequence, the HC-scFv sequence, and the light chain. Amino acid substitutions made to allow for efficient purification of the desired “triple F” heterodimer over the undesired dual scFv-Fc and mAb homodimers are underlined.

FIG. 8. Amino acid sequences for the humanized anti-CD3 variable regions finding particular use in the invention (with CDRs underlined).

FIG. 9A-9C. Production of XENP11874, a “triple F” bispecific with an anti-CD19 Fab and anti-CD3 scFv. In FIG. 9A, the ion exchange purification of the desired “triple F” heterodimer from the undesired dual scFv-Fc and mAb homodimers is shown. The purity of the “triple F” fraction was checked by IEF gel, shown in FIG. 9B. Finally, SEC was used to confirm the homogenous size of the “triple F” product (see FIG. 9C).

FIG. 10. XENP11874, anti-CD19 Fab x anti-CD3 scFv “triple F” bispecific, potently recruits T cells for B cell depletion.

FIG. 11A-11C. The production of XENP11924, a “triple F” bispecific with an anti-CD19 Fab and anti-CD3 scFv. In FIG. 11A, the ion exchange purification of the desired “triple F” heterodimer from the undesired dual scFv-Fc and mAb homodimers is shown. The purity of the “triple F” fraction was checked by IEF gel, shown in FIG. 11B. Finally, SEC was used to confirm the homogenous size of the “triple F” product (see FIG. 11C).

FIG. 12. XENP11924, anti-CD19 Fab x anti-CD3 scFv “triple F” bispecific, potently recruits T cells for the killing of the Raji tumor cell line.

FIG. 13. Amino acid sequences for the anti-CD38 Fab x anti-CD3 scFv “triple F” embodiment, showing the HC-Fab sequence, the HC-scFv sequence, and the light chain. Amino acid substitutions made to allow for efficient purification of the desired “triple F” heterodimer over the undesired dual scFv-Fc and mAb homodimers are underlined.

FIG. 14A-14C. The production of XENP11925, a “triple F” bispecific with an anti-CD38 Fab and anti-CD3 scFv. In FIG. 14A, the ion exchange purification of the desired “triple F” heterodimer from the undesired dual scFv-Fc and mAb homodimers is shown. The purity of the “triple F” fraction was checked by IEF gel, shown in FIG. 14B. Finally, SEC was used to confirm the homogenous size of the “triple F” product (see FIG. 14C).

FIG. 15. XENP11925, anti-CD38 Fab x anti-CD3 scFv “triple F” bispecific, potently recruits T cells for the killing of the RPMI8226 tumor cell line.

FIG. 16A-16B. Amino acid sequences for embodiments of the anti-CD19 Fab x anti-CD3 scFv “triple F” bispecific, showing the HC-Fab sequence, the HC-scFv sequence, and the light chain for each bispecific. Amino acid substitutions made to “skew” the formation toward desired “triple F” heterodimer over undesired dual scFv-Fc and mAb homodimers are underlined.

FIG. 17. Isoelectric focusing (IEF) gel providing analysis of various “triple F” bispecifics. The desired “triple F” heterodimer has an apparent pI of 8.4 (middle band). Note the increase in heterodimer yield for XENP12152 and XENP12155.

6

FIG. 18. Analytical ion exchange chromatography providing analysis of various “triple F” bispecifics. Yield of the desired “triple F” heterodimer is boxed. Note the increase in heterodimer yield for XENP12155.

FIG. 19 shows the literature pIs of the 20 amino acids. It should be noted that the listed pIs are calculated as free amino acids; the actual pI of any side chain in the context of a protein is different, and thus this list is used to show pI trends and not absolute numbers for the purposes of the invention.

FIG. 20A-20F. Data table of exemplary pI-engineered variants listing:

15	XenP#	the internal reference number
	Name (HC)	heavy chain sequence designation
	SEQ ID NO (HC)	corresponding SEQ ID NO of the heavy chain sequence
	Name (LC)	light chain sequence designation
20	SEQ ID NO (LC)	corresponding SEQ ID NO of the light chain sequence
	Calc. pI	calculated pI value for the entire antibody sequence, including heavy and light chain Fv + constant domains, with the Fv of bevacizumab and the constant domains as defined in the table
	#KR	number of Lys or Arg residues in IgG1 with the Fv of bevacizumab and the constant domains as defined in the table
25	Delta KR (vs. WT)	change in the number of Lys or Arg residues relative to IgG1 wild-type sequence of bevacizumab
	#DE	number of Asp or Glu residues in IgG1 with the Fv of bevacizumab and the constant domains as defined in the table
30	Delta DE (vs. WT)	change in the number of Asp or Glu acid residues relative to IgG1 wild-type sequence of bevacizumab
	Charge state	derived from the total number of Lys and Arg minus the total number of Asp and Glu residues, assuming a pH of 7
35	# HC Mutations vs IgG1	number of mutations in the heavy chain constant domain as compared to IgG1
	# LC Mutations vs IgG1	number of mutations in the light chain constant domain as compared to IgG1
	Total # of Mutations	total number of mutations in the heavy chain and light chain constant domains as compared to IgG1

It should be noted that FIG. 20A-20F has SEQ ID NO:s that are associated with the sequence listing filed in U.S. Ser. No. 13/648,951, and are hereby expressly incorporated by reference.

FIG. 21A-21C. depicts a number of pI variants that can additionally find use in the present invention, which is a list of all possible reduced pI variants created from isotypic substitutions of IgG1-4. Shown are the pI values for the three expected species as well as the average delta pI between the heterodimer and the two homodimer species present when the variant heavy chain is transfected with IgG1-WT heavy chain.

FIG. 22 List of all possible increased pI variants created from isotypic substitutions of IgG1-4. Shown are the pI values for the three expected species as well as the average delta pI between the heterodimer and the two homodimer species present when the variant heavy chain is transfected with IgG1-WT heavy chain.

FIG. 23A-23B is an additional list of potential heterodimerization variants for use in the present invention.

FIG. 24 depicts a matrix of possible combinations of heterodimerization formats, heterodimerization variants (separated into pI variants and steric variants (which includes charge pair variants), Fc variants, FcRn variants and combinations. Legend A are suitable FcRn variants: 434A, 434S, 428L, 308F, 259I, 428L/434S, 259I/308F, 436I/428L, 436I or V/434S, 436V/428L, 252Y, 252Y/254T/

256E and 259I/308F/428L. That is, the Triple F format of FIG. 1B can have any of these FcRn variants. For clarity, as each heavy chain is different, FcRn variants (as well as the Fc variants) can reside on one or both monomers. Legend B are suitable Fc variants: 236A, 239D, 239E, 332E, 332D, 239D/332E, 267D, 267E, 328F, 267E/328F, 236A/332E, 239D/332E/330Y, 239D, 332E/330L, 236R, 328R, 236R/328R, 236N/267E, 243L, 298A and 299T. (Note, additional suitable Fc variants are found in FIG. 41 of US 2006/0024298, the figure and legend of which are hereby incorporated by reference in their entirety). Legend C are suitable pI variants, and these, for brevity are imported from FIG. 23A-23B, again with the understanding that there is a "strandedness" to pI variants. Legend D are suitable steric variants (including charge pair variants); again, for brevity are imported from FIG. 21A-21C, again with the understanding that there is a "strandedness" to steric variants. Legend E reflects the following possible combinations, again, with each variant being independently and optionally combined from the appropriate source Legend: 1) pI variants plus FcRn variants; 2) pI variants plus Fc variants; 3) pI variants plus FcRn variants plus Fc variants; 4) steric variants plus FcRn variants; 5) steric variants plus Fc variants; 6) steric variants plus FcRn variants plus Fc variants; 7) pI variants plus steric variants plus FcRn variants; 8) pI variants plus steric variants plus Fc variants; 9) pI variants plus steric variants plus FcRn variants plus Fc variants; and 10) pI variants plus steric variants.

FIGS. 25A-25C depict stability-optimized, humanized anti-CD3 variant scFvs. Substitutions are given relative to the H1_L1.4 scFv sequence. Amino acid numbering is Kabat numbering.

FIGS. 26A-26YY depict the amino acid sequences of stability-optimized, humanized anti-CD3 variant scFvs, variable heavy and variable light sequences. (Note also that the first sequence is the histidine tagged version for ease of purification). CDRs are underlined. It should be understood that the increased stability of the optimized variable and optimized light chains (as well as the scFv chains) can be attributed to framework regions as well as the CDRs. Thus, it should be understood that the disclosure of the entire variable region includes the disclosure of the framework regions, although they are not separately numbered. In addition, the scFv linkers are shown in grey. Each scFv linker can be replaced with a charged scFv linker as depicted in FIG. 7A-7B. That is, any charged scFv linker, whether positive or negative, including those depicted in FIG. 7A-7B, can be substituted for the highlighted region in FIGS. 2A to 2YY.

FIGS. 27A-27B show yields after Ni-NTA purification and melting temperatures (T_m) as determined by DSF (Differential Scanning Fluorimetry) of stability-optimized, humanized anti-CD3 variant scFvs. Similar DSF stability measurements for the I2C anti-CD3 variable region in a comparable scFv format gave a T_m of 59° C.

FIGS. 28A-28B show fold improvement (relative to H1_L1.4 scFv) in cell surface binding affinities (IC₅₀) of stability-optimized, humanized anti-CD3 variant scFvs as determined in a competitive binding experiment using purified human T cells from PBMCs.

FIG. 29. Figure depicts the sequence of human CD3.

FIGS. 30A-30I depict a collation of all the CD3 vHCDR1-3 and vLCDR1-3 sequences useful in the present invention.

DETAILED DESCRIPTION OF THE INVENTION

FIGS. 1-76 of U.S. Ser. No. 61/593,846 and the associated legends and discussion in the specification are hereby incorporated by reference.

FIGS. 2 and 17 of U.S. Ser. No. 61/778,157, inclusive of all the sequences including the optimized CD3 sequences are expressly incorporated herein by reference.

1. Overview

The present invention is directed to novel constructs to provide bispecific antibodies (or, as discussed below, trispecific or tetraspecific antibodies can also be made). An ongoing problem in antibody technologies is the desire for "bispecific" (and/or multispecific) antibodies that bind to two (or more) different antigens simultaneously, in general thus allowing the different antigens to be brought into proximity and resulting in new functionalities and new therapies. In general, these antibodies are made by including genes for each heavy and light chain into the host cells. This generally results in the formation of the desired heterodimer (A-B), as well as the two homodimers (A-A and B-B). However, a major obstacle in the formation of multispecific antibodies is the difficulty in purifying the heterodimeric antibodies away from the homodimeric antibodies and/or biasing the formation of the heterodimer over the formation of the homodimers.

The present invention is generally directed to the creation of heterodimeric proteins such as antibodies that can co-engage antigens in several ways, relying on amino acid variants in the constant regions that are different on each chain to promote heterodimeric formation and/or allow for ease of purification of heterodimers over the homodimers.

Thus, the present invention is directed to novel immunoglobulin compositions that co-engage at least a first and a second antigen. First and second antigens of the invention are herein referred to as antigen-1 and antigen-2 respectively. One heavy chain of the antibody contains a single chain Fv ("scFv", as defined below) and the other heavy chain is a "regular" FAb format, comprising a variable heavy chain and a light chain. This structure is sometimes referred to herein as "triple F" format (scFv-FAb-Fc) or the "bottle-opener" format, due to a rough visual similarity to a bottle-opener (see FIG. 1A-1C). The two chains are brought together by the use of amino acid variants in the constant regions (e.g. the Fc domain and/or the hinge region) that promote the formation of heterodimeric antibodies as is described more fully below.

There are several distinct advantages to the present "triple F" format. As is known in the art, antibody analogs relying on two scFv constructs often have stability and aggregation problems, which can be alleviated in the present invention by the addition of a "regular" heavy and light chain pairing. In addition, as opposed to formats that rely on two heavy chains and two light chains, there is no issue with the incorrect pairing of heavy and light chains (e.g. heavy 1 pairing with light 2, etc.)

There are a number of mechanisms that can be used to generate the heterodimers of the present invention. In addition, as will be appreciated by those in the art and described more fully below, these mechanisms can be combined to ensure high heterodimerization.

One mechanism is generally referred to in the art as "knobs and holes" ("KIH"), referring to amino acid engineering that creates steric influences to favor heterodimeric formation and disfavor homodimeric formation can also optionally be used; this is sometimes referred to as "knobs

and holes”, as described in U.S. Ser. No. 61/596,846, Ridgway et al., Protein Engineering 9(7):617 (1996); Atwell et al., J. Mol. Biol. 1997 270:26; U.S. Pat. No. 8,216,805, all of which are hereby incorporated by reference in their entirety. FIGS. 4A and 4B, further described below, identifies a number of “monomer A-monomer B” pairs that rely on “knobs and holes”. In addition, as described in Merchant et al., Nature Biotech. 16:677 (1998), these “knobs and hole” mutations can be combined with disulfide bonds to skew formation to heterodimerization. Some of these variants are shown in FIGS. 16, 20, 21 and 22.

An additional mechanism that finds use in the generation of heterodimers is sometimes referred to as “electrostatic steering” as described in Gunasekaran et al., J. Biol. Chem. 285(25):19637 (2010), hereby incorporated by reference in its entirety. This is sometimes referred to herein as “charge pairs”. In this embodiment, electrostatics are used to skew the formation towards heterodimerization. As those in the art will appreciate, these may also have an effect on pI, and thus on purification, and thus could in some cases also be considered pI variants. However, as these were generated to force heterodimerization and were not used as purification tools, they are classified as “steric variants”. These include, but are not limited to, D221E/P228E/L368E paired with D221R/P228R/K409R (e.g. these are “monomer corresponding sets) and C220E/P228E/368E paired with C220R/E224R/P228R/K409R.

In the present invention, there are two basic mechanisms that can lead to ease of purifying heterodimeric proteins; one relies on the use of pI variants, such that each monomer has a different pI, thus allowing the isoelectric purification of A-A, A-B and B-B dimeric proteins. Alternatively, the “triple F” format also allows separation on the basis of size. As is further outlined below, it is also possible to “skew” the formation of heterodimers over homodimers, as is generally outlined below.

In the present invention that utilizes pI as a separation mechanism to create the heterodimeric Triple F format, amino acid variants can be introduced into one or both of the monomer polypeptides; that is, the pI of one of the monomers (referred to herein for simplicity as “monomer A”) can be engineered away from monomer B, or both monomer A and B change be changed, with the pI of monomer A increasing and the pI of monomer B decreasing. As is outlined more fully below, the pI changes of either or both monomers can be done by removing or adding a charged residue (e.g. a neutral amino acid is replaced by a positively or negatively charged amino acid residue, e.g. glycine to glutamic acid), changing a charged residue from positive or negative to the opposite charge (aspartic acid to lysine) or changing a charged residue to a neutral residue (e.g. loss of a charge; lysine to serine). A number of these variants are shown in FIGS. 6, 20 and 21.

Accordingly, in this embodiment of the present invention provides for creating a sufficient change in pI in at least one of the monomers such that heterodimers can be separated from homodimers. As will be appreciated by those in the art, and as discussed further below, this can be done by using a “wild type” heavy chain constant region and a variant region that has been engineered to either increase or decrease its pI (wt A⁺B or wt A⁻B⁻), or by increasing one region and decreasing the other region (A⁺B⁻ or A⁻B⁺). It should be noted that in this discussion it does not matter which monomer comprises the scFv and which the Fab.

Thus, in general, a component of the present invention are amino acid variants in the constant regions of antibodies that are directed to altering the isoelectric point (pI) of at least

one, if not both, of the monomers of a dimeric protein to form “pI heterodimers” (when the protein is an antibody, these are referred to as “pI antibodies”) by incorporating amino acid substitutions (“pI variants” or “pI substitutions”) into one or both of the monomers. As shown herein, the separation of the heterodimers from the two homodimers can be accomplished if the pIs of the two monomers differ by as little as 0.1 pH unit, with 0.2, 0.3, 0.4 and 0.5 or greater all finding use in the present invention.

Furthermore, as will be appreciated by those in the art and outlined herein, heterodimers can be separated from homodimers on the basis of size. For example, as shown in FIG. 1A-1B, heterodimers with two scFvs (FIG. 1A) can be separated by those of the “triple F” format (FIG. 1B) and a bispecific mAb (FIG. 1C). This can be further exploited in higher valency with additional antigen binding sites being utilized. For example, in as shown in FIG. 2D, one monomer will have two Fab fragments and the other will have one scFv, resulting in a differential in size and thus molecular weight. Other examples are shown in FIGS. 2 and 3.

In addition, as will be appreciated by those in the art and outlined herein, the format outlined herein can be expanded to provide trispecific and tetraspecific antibodies as well. In this embodiment, some variations of which are depicted in FIGS. 2 and 3, it will be recognized that it is possible that some antigens are bound divalently (e.g. two antigen binding sites to a single antigen; for example, A and B could be part of a typical bivalent association and C and D can be optionally present and optionally the same or different). As will be appreciated, any combination of Fab and scFvs can be utilized to achieve the desired result and combinations.

In the case where pI variants are used to achieve heterodimerization, by using the constant region(s) of the heavy chain(s), a more modular approach to designing and purifying multispecific proteins, including antibodies, is provided. In addition, in some embodiments, the possibility of immunogenicity resulting from the pI variants is significantly reduced by importing pI variants from different IgG isotypes such that pI is changed without introducing significant immunogenicity. Thus, an additional problem to be solved is the elucidation of low pI constant domains with high human sequence content, e.g. the minimization or avoidance of non-human residues at any particular position.

In one embodiment, the heterodimeric antibody provides for monovalent engagement of one antigen using a scFv and monovalent engagement of the other antigen using a Fab. As outlined below, this format can also be varied; in some embodiments, there is monovalent engagement of three antigens, divalent engagement of one antigen and monovalent engagement of a second antigen (e.g. A and C are to the same antigen and B is to a different antigen), etc.

A side benefit that can occur with this pI engineering is also the extension of serum half-life and increased FcRn binding. That is, as described in U.S. Ser. No. 13/194,904 (incorporated by reference in its entirety), lowering the pI of antibody constant domains (including those found in antibodies and Fc fusions) can lead to longer serum retention in vivo. These pI variants for increased serum half life also facilitate pI changes for purification.

In addition to all or part of a variant heavy constant domain, one or both of the monomers may contain one or two fusion partners, such that the heterodimers form multivalent proteins. As is generally depicted in the Figures, the fusion partners are depicted as A, B, C and D, with all combinations possible. In general, A, B, C and D are selected such that the heterodimer is at least bispecific or bivalent in its ability to interact with additional proteins. In

11

the context of the present “triple F” format, generally A and B are an scFv and a Fv (as will be appreciated, either monomer can contain the scFv and the other the Fv/Fab) and then optionally one or two additional fusion partners.

Accordingly, the present invention provides novel constructs to produce multivalent antibodies.

Furthermore, as outlined herein, additional amino acid variants may be introduced into the bispecific antibodies of the invention, to add additional functionalities. For example, amino acid changes within the Fc region can be added (either to one monomer or both) to facilitate increased ADCC or CDC (e.g. altered binding to Fcγ receptors); to allow or increase yield of the addition of toxins and drugs (e.g. for ADC), as well as to increase binding to FcRn and/or increase serum half-life of the resulting molecules. As is further described herein and as will be appreciated by those in the art, any and all of the variants outlined herein can be optionally and independently combined with other variants.

II. Definitions

In order that the application may be more completely understood, several definitions are set forth below. Such definitions are meant to encompass grammatical equivalents.

By “ablation” herein is meant a decrease or removal of activity. Thus for example, “ablating FcγR binding” means the Fc region amino acid variant has less than 50% starting binding as compared to an Fc region not containing the specific variant, with less than 70-80-90-95-98% loss of activity being preferred, and in general, with the activity being below the level of detectable binding in a Biacore assay. Of particular use in the ablation of FcγR binding is the double variant 236R/328R.

By “ADCC” or “antibody dependent cell-mediated cytotoxicity” as used herein is meant the cell-mediated reaction wherein nonspecific cytotoxic cells that express FcγRs recognize bound antibody on a target cell and subsequently cause lysis of the target cell. ADCC is correlated with binding to FcγRIIIa; increased binding to FcγRIIIa leads to an increase in ADCC activity.

By “ADCP” or antibody dependent cell-mediated phagocytosis as used herein is meant the cell-mediated reaction wherein nonspecific cytotoxic cells that express FcγRs recognize bound antibody on a target cell and subsequently cause phagocytosis of the target cell.

By “modification” herein is meant an amino acid substitution, insertion, and/or deletion in a polypeptide sequence or an alteration to a moiety chemically linked to a protein. For example, a modification may be an altered carbohydrate or PEG structure attached to a protein. By “amino acid modification” herein is meant an amino acid substitution, insertion, and/or deletion in a polypeptide sequence. For clarity, unless otherwise noted, the amino acid modification is always to an amino acid coded for by DNA, e.g. the 20 amino acids that have codons in DNA and RNA.

By “amino acid substitution” or “substitution” herein is meant the replacement of an amino acid at a particular position in a parent polypeptide sequence with a different amino acid. In particular, in some embodiments, the substitution is to an amino acid that is not naturally occurring at the particular position, either not naturally occurring within the organism or in any organism. For example, the substitution E272Y refers to a variant polypeptide, in this case an Fc variant, in which the glutamic acid at position 272 is replaced with tyrosine. For clarity, a protein which has been engineered to change the nucleic acid coding sequence but not change the starting amino acid (for example exchanging CGG (encoding arginine) to CGA (still encoding arginine) to

12

increase host organism expression levels) is not an “amino acid substitution”; that is, despite the creation of a new gene encoding the same protein, if the protein has the same amino acid at the particular position that it started with, it is not an amino acid substitution.

By “amino acid insertion” or “insertion” as used herein is meant the addition of an amino acid sequence at a particular position in a parent polypeptide sequence. For example, -233E or 233E designates an insertion of glutamic acid after position 233 and before position 234. Additionally, -233ADE or A233ADE designates an insertion of AlaAsp-Glu after position 233 and before position 234.

By “amino acid deletion” or “deletion” as used herein is meant the removal of an amino acid sequence at a particular position in a parent polypeptide sequence. For example, E233- or E233# designates a deletion of glutamic acid at position 233. Additionally, EDA233- or EDA233# designates a deletion of the sequence GluAspAla that begins at position 233.

By “variant protein” or “protein variant”, or “variant” as used herein is meant a protein that differs from that of a parent protein by virtue of at least one amino acid modification. Protein variant may refer to the protein itself, a composition comprising the protein, or the amino sequence that encodes it. Preferably, the protein variant has at least one amino acid modification compared to the parent protein, e.g. from about one to about seventy amino acid modifications, and preferably from about one to about five amino acid modifications compared to the parent. As described below, in some embodiments the parent polypeptide, for example an Fc parent polypeptide, is a human wild type sequence, such as the Fc region from IgG1, IgG2, IgG3 or IgG4, although human sequences with variants can also serve as “parent polypeptides”. The protein variant sequence herein will preferably possess at least about 80% identity with a parent protein sequence, and most preferably at least about 90% identity, more preferably at least about 95-98-99% identity. Variant protein can refer to the variant protein itself, compositions comprising the protein variant, or the DNA sequence that encodes it. Accordingly, by “antibody variant” or “variant antibody” as used herein is meant an antibody that differs from a parent antibody by virtue of at least one amino acid modification. “IgG variant” or “variant IgG” as used herein is meant an antibody that differs from a parent IgG (again, in many cases, from a human IgG sequence) by virtue of at least one amino acid modification, and “immunoglobulin variant” or “variant immunoglobulin” as used herein is meant an immunoglobulin sequence that differs from that of a parent immunoglobulin sequence by virtue of at least one amino acid modification. “Fc variant” or “variant Fc” as used herein is meant a protein comprising an amino acid modification in an Fc domain. The Fc variants of the present invention are defined according to the amino acid modifications that compose them. Thus, for example, N434S or 434S is an Fc variant with the substitution serine at position 434 relative to the parent Fc polypeptide, wherein the numbering is according to the EU index. Likewise, M428L/N434S defines an Fc variant with the substitutions M428L and N434S relative to the parent Fc polypeptide. The identity of the WT amino acid may be unspecified, in which case the aforementioned variant is referred to as 428L/434S. It is noted that the order in which substitutions are provided is arbitrary, that is to say that, for example, 428L/434S is the same Fc variant as M428L/N434S, and so on. For all positions discussed in the present invention that relate to antibodies, unless otherwise noted, amino acid position numbering is according to the EU index. The EU index or

EU index as in Kabat or EU numbering scheme refers to the numbering of the EU antibody (Edelman et al., 1969, *Proc Natl Acad Sci USA* 63:78-85, hereby entirely incorporated by reference.) The modification can be an addition, deletion, or substitution. Substitutions can include naturally occurring amino acids and, in some cases, synthetic amino acids. Examples include U.S. Pat. No. 6,586,207; WO 98/48032; WO 03/073238; US2004-0214988A1; WO 05/35727A2; WO 05/74524A2; J. W. Chin et al., (2002), *Journal of the American Chemical Society* 124:9026-9027; J. W. Chin, & P. G. Schultz, (2002), *ChemBioChem* 11:1135-1137; J. W. Chin, et al., (2002), *PICAS United States of America* 99:11020-11024; and, L. Wang, & P. G. Schultz, (2002), *Chem.* 1-10, all entirely incorporated by reference.

As used herein, "protein" herein is meant at least two covalently attached amino acids, which includes proteins, polypeptides, oligopeptides and peptides. The peptidyl group may comprise naturally occurring amino acids and peptide bonds, or synthetic peptidomimetic structures, i.e. "analogs", such as peptoids (see Simon et al., *PNAS USA* 89(20):9367 (1992), entirely incorporated by reference). The amino acids may either be naturally occurring or synthetic (e.g. not an amino acid that is coded for by DNA); as will be appreciated by those in the art. For example, homophenylalanine, citrulline, ornithine and noreleucine are considered synthetic amino acids for the purposes of the invention, and both D- and L-(R or S) configured amino acids may be utilized. The variants of the present invention may comprise modifications that include the use of synthetic amino acids incorporated using, for example, the technologies developed by Schultz and colleagues, including but not limited to methods described by Cropp & Shultz, 2004, *Trends Genet.* 20(12):625-30, Anderson et al., 2004, *Proc Natl Acad Sci USA* 101 (2):7566-71, Zhang et al., 2003, 303(5656):371-3, and Chin et al., 2003, *Science* 301(5635): 964-7, all entirely incorporated by reference. In addition, polypeptides may include synthetic derivatization of one or more side chains or termini, glycosylation, PEGylation, circular permutation, cyclization, linkers to other molecules, fusion to proteins or protein domains, and addition of peptide tags or labels.

By "residue" as used herein is meant a position in a protein and its associated amino acid identity. For example, Asparagine 297 (also referred to as Asn297 or N297) is a residue at position 297 in the human antibody IgG1.

By "Fab" or "Fab region" as used herein is meant the polypeptide that comprises the VH, CHL VL, and CL immunoglobulin domains. Fab may refer to this region in isolation, or this region in the context of a full length antibody, antibody fragment or Fab fusion protein. By "Fv" or "Fv fragment" or "Fv region" as used herein is meant a polypeptide that comprises the VL and VH domains of a single antibody.

By "IgG subclass modification" or "isotype modification" as used herein is meant an amino acid modification that converts one amino acid of one IgG isotype to the corresponding amino acid in a different, aligned IgG isotype. For example, because IgG1 comprises a tyrosine and IgG2 a phenylalanine at EU position 296, a F296Y substitution in IgG2 is considered an IgG subclass modification.

By "non-naturally occurring modification" as used herein is meant an amino acid modification that is not isotypic. For example, because none of the IgGs comprise a serine at position 434, the substitution 434S in IgG1, IgG2, IgG3, or IgG4 (or hybrids thereof) is considered a non-naturally occurring modification.

By "amino acid" and "amino acid identity" as used herein is meant one of the 20 naturally occurring amino acids that are coded for by DNA and RNA.

By "effector function" as used herein is meant a biochemical event that results from the interaction of an antibody Fc region with an Fc receptor or ligand. Effector functions include but are not limited to ADCC, ADCP, and CDC.

By "IgG Fc ligand" as used herein is meant a molecule, preferably a polypeptide, from any organism that binds to the Fc region of an IgG antibody to form an Fc/Fc ligand complex. Fc ligands include but are not limited to FcγRI, FcγRII, FcγRIII, FcRn, C1q, C3, mannan binding lectin, mannose receptor, staphylococcal protein A, streptococcal protein G, and viral FcγR. Fc ligands also include Fc receptor homologs (FcRH), which are a family of Fc receptors that are homologous to the FcγRs (Davis et al., 2002, *Immunological Reviews* 190:123-136, entirely incorporated by reference). Fc ligands may include undiscovered molecules that bind Fc. Particular IgG Fc ligands are FcRn and Fc gamma receptors. By "Fc ligand" as used herein is meant a molecule, preferably a polypeptide, from any organism that binds to the Fc region of an antibody to form an Fc/Fc ligand complex.

By "Fc gamma receptor", "FcγR" or "FcγammaR" as used herein is meant any member of the family of proteins that bind the IgG antibody Fc region and is encoded by an FcγR gene. In humans this family includes but is not limited to FcγRI (CD64), including isoforms FcγRIa, FcγRIb, and FcγRIc; FcγRII (CD32), including isoforms FcγRIIa (including allotypes H131 and R131), FcγRIIb (including FcγRIIb-1 and FcγRIIb-2), and FcγRIIc; and FcγRIII (CD16), including isoforms FcγRIIIa (including allotypes V158 and F158) and FcγRIIIb (including allotypes FcγRIIb-NA1 and FcγRIIb-NA2) (Jefferis et al., 2002, *Immunol Lett* 82:57-65, entirely incorporated by reference), as well as any undiscovered human FcγRs or FcγR isoforms or allotypes. An FcγR may be from any organism, including but not limited to humans, mice, rats, rabbits, and monkeys. Mouse FcγRs include but are not limited to FcγRI (CD64), FcγRII (CD32), FcγRIII (CD16), and FcγRIII-2 (CD16-2), as well as any undiscovered mouse FcγRs or FcγR isoforms or allotypes.

By "FcRn" or "neonatal Fc Receptor" as used herein is meant a protein that binds the IgG antibody Fc region and is encoded at least in part by an FcRn gene. The FcRn may be from any organism, including but not limited to humans, mice, rats, rabbits, and monkeys. As is known in the art, the functional FcRn protein comprises two polypeptides, often referred to as the heavy chain and light chain. The light chain is beta-2-microglobulin and the heavy chain is encoded by the FcRn gene. Unless otherwise noted herein, FcRn or an FcRn protein refers to the complex of FcRn heavy chain with beta-2-microglobulin. A variety of FcRn variants used to increase binding to the FcRn receptor, and in some cases, to increase serum half-life, are discussed in Legend A of FIG. 24.

By "parent polypeptide" as used herein is meant a starting polypeptide that is subsequently modified to generate a variant. The parent polypeptide may be a naturally occurring polypeptide, or a variant or engineered version of a naturally occurring polypeptide. Parent polypeptide may refer to the polypeptide itself, compositions that comprise the parent polypeptide, or the amino acid sequence that encodes it. Accordingly, by "parent immunoglobulin" as used herein is meant an unmodified immunoglobulin polypeptide that is modified to generate a variant, and by "parent antibody" as used herein is meant an unmodified antibody that is modified

15

to generate a variant antibody. It should be noted that "parent antibody" includes known commercial, recombinantly produced antibodies as outlined below.

By "Fc fusion protein" or "immunoadhesin" herein is meant a protein comprising an Fc region, generally linked (optionally through a linker moiety, as described herein) to a different protein, such as a binding moiety to a target protein, as described herein).

By "position" as used herein is meant a location in the sequence of a protein. Positions may be numbered sequentially, or according to an established format, for example the EU index for antibody numbering.

By "strandedness" in the context of the monomers of the heterodimeric proteins of the invention herein is meant that, similar to the two strands of DNA that "match", heterodimerization variants are incorporated into each monomer so as to preserve the ability to "match" to form heterodimers. For example, if some pI variants are engineered into monomer A (e.g. making the pI higher) then steric variants that are "charge pairs" that can be utilized as well do not interfere with the pI variants, e.g. the charge variants that make a pI higher are put on the same "strand" or "monomer" to preserve both functionalities.

By "target antigen" as used herein is meant the molecule that is bound specifically by the variable region of a given antibody. A target antigen may be a protein, carbohydrate, lipid, or other chemical compound. A wide number of suitable target antigens are described below.

By "target cell" as used herein is meant a cell that expresses a target antigen.

By "variable region" as used herein is meant the region of an immunoglobulin that comprises one or more Ig domains substantially encoded by any of the V.kappa., V.lamda., and/or VH genes that make up the kappa, lambda, and heavy chain immunoglobulin genetic loci respectively.

By "wild type or WT" herein is meant an amino acid sequence or a nucleotide sequence that is found in nature, including allelic variations. A WT protein has an amino acid sequence or a nucleotide sequence that has not been intentionally modified.

Heterodimeric Proteins

The present invention is directed to the generation of multispecific, particularly bispecific binding proteins, and in particular, multispecific antibodies that have one monomer comprising an scFv and the other an Fv.

Antibody

The present invention relates to the generation of multispecific antibodies, generally therapeutic antibodies. As is discussed below, the term "antibody" is used generally. Antibodies that find use in the present invention can take on a number of formats as described herein, including traditional antibodies as well as antibody derivatives, fragments and mimetics, described below. In general, the term "antibody" includes any polypeptide that includes at least one constant domain, including, but not limited to, CH1, CH2, CH3 and CL.

Traditional antibody structural units typically comprise a tetramer. Each tetramer is typically composed of two identical pairs of polypeptide chains, each pair having one "light" (typically having a molecular weight of about 25 kDa) and one "heavy" chain (typically having a molecular weight of about 50-70 kDa). Human light chains are classified as kappa and lambda light chains. The present invention is directed to the IgG class, which has several subclasses, including, but not limited to IgG1, IgG2, IgG3, and IgG4. Thus, "isotype" as used herein is meant any of the subclasses of immunoglobulins defined by the chemical and

16

antigenic characteristics of their constant regions. It should be understood that therapeutic antibodies can also comprise hybrids of isotypes and/or subclasses. For example, as shown in US Publication 2009/0163699, incorporated by reference, the present invention covers pI engineering of IgG1/G2 hybrids.

The amino-terminal portion of each chain includes a variable region of about 100 to 110 or more amino acids primarily responsible for antigen recognition, generally referred to in the art and herein as the "Fv domain" or "Fv region". In the variable region, three loops are gathered for each of the V domains of the heavy chain and light chain to form an antigen-binding site. Each of the loops is referred to as a complementarity-determining region (hereinafter referred to as a "CDR"), in which the variation in the amino acid sequence is most significant. "Variable" refers to the fact that certain segments of the variable region differ extensively in sequence among antibodies. Variability within the variable region is not evenly distributed. Instead, the V regions consist of relatively invariant stretches called framework regions (FRs) of 15-30 amino acids separated by shorter regions of extreme variability called "hypervariable regions" that are each 9-15 amino acids long or longer.

Each VH and VL is composed of three hypervariable regions ("complementary determining regions," "CDRs") and four FRs, arranged from amino-terminus to carboxy-terminus in the following order: FR1-CDR1-FR2-CDR2-FR3-CDR3-FR4.

The hypervariable region generally encompasses amino acid residues from about amino acid residues 24-34 (LCDR1; "L" denotes light chain), 50-56 (LCDR2) and 89-97 (LCDR3) in the light chain variable region and around about 31-35B (HCDR1; "H" denotes heavy chain), 50-65 (HCDR2), and 95-102 (HCDR3) in the heavy chain variable region; Kabat et al., SEQUENCES OF PROTEINS OF IMMUNOLOGICAL INTEREST, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, Md. (1991) and/or those residues forming a hypervariable loop (e.g. residues 26-32 (LCDR1), 50-52 (LCDR2) and 91-96 (LCDR3) in the light chain variable region and 26-32 (HCDR1), 53-55 (HCDR2) and 96-101 (HCDR3) in the heavy chain variable region; Chothia and Lesk (1987) J. Mol. Biol. 196:901-917. Specific CDRs of the invention are described below.

Throughout the present specification, the Kabat numbering system is generally used when referring to a residue in the variable domain (approximately, residues 1-107 of the light chain variable region and residues 1-113 of the heavy chain variable region) (e.g. Kabat et al., supra (1991)).

The CDRs contribute to the formation of the antigen-binding, or more specifically, epitope binding site of antibodies. "Epitope" refers to a determinant that interacts with a specific antigen binding site in the variable region of an antibody molecule known as a paratope. Epitopes are groupings of molecules such as amino acids or sugar side chains and usually have specific structural characteristics, as well as specific charge characteristics. A single antigen may have more than one epitope.

The epitope may comprise amino acid residues directly involved in the binding (also called immunodominant component of the epitope) and other amino acid residues, which are not directly involved in the binding, such as amino acid residues which are effectively blocked by the specifically antigen binding peptide; in other words, the amino acid residue is within the footprint of the specifically antigen binding peptide.

17

Epitopes may be either conformational or linear. A conformational epitope is produced by spatially juxtaposed amino acids from different segments of the linear polypeptide chain. A linear epitope is one produced by adjacent amino acid residues in a polypeptide chain. Conformational and nonconformational epitopes may be distinguished in that the binding to the former but not the latter is lost in the presence of denaturing solvents.

An epitope typically includes at least 3, and more usually, at least 5 or 8-10 amino acids in a unique spatial conformation. Antibodies that recognize the same epitope can be verified in a simple immunoassay showing the ability of one antibody to block the binding of another antibody to a target antigen, for example "binning."

The carboxy-terminal portion of each chain defines a constant region primarily responsible for effector function. Kabat et al. collected numerous primary sequences of the variable regions of heavy chains and light chains. Based on the degree of conservation of the sequences, they classified individual primary sequences into the CDR and the framework and made a list thereof (see SEQUENCES OF IMMUNOLOGICAL INTEREST, 5th edition, NIH publication, No. 91-3242, E. A. Kabat et al., entirely incorporated by reference).

In the IgG subclass of immunoglobulins, there are several immunoglobulin domains in the heavy chain. By "immunoglobulin (Ig) domain" herein is meant a region of an immunoglobulin having a distinct tertiary structure. Of interest in the present invention are the heavy chain domains, including, the constant heavy (CH) domains and the hinge domains. In the context of IgG antibodies, the IgG isotypes each have three CH regions. Accordingly, "CH" domains in the context of IgG are as follows: "CH1" refers to positions 118-220 according to the EU index as in Kabat. "CH2" refers to positions 237-340 according to the EU index as in Kabat, and "CH3" refers to positions 341-447 according to the EU index as in Kabat. As shown herein and described below, the pI variants can be in one or more of the CH regions, as well as the hinge region, discussed below.

It should be noted that the sequences depicted herein start at the CH1 region, position 118; the variable regions are not included except as noted. For example, the first amino acid of SEQ ID NO: 2, while designated as position "1" in the sequence listing, corresponds to position 118 of the CH1 region, according to EU numbering.

Another type of Ig domain of the heavy chain is the hinge region. By "hinge" or "hinge region" or "antibody hinge region" or "immunoglobulin hinge region" herein is meant the flexible polypeptide comprising the amino acids between the first and second constant domains of an antibody. Structurally, the IgG CH1 domain ends at EU position 220, and the IgG CH2 domain begins at residue EU position 237. Thus for IgG the antibody hinge is herein defined to include positions 221 (D221 in IgG1) to 236 (G236 in IgG1), wherein the numbering is according to the EU index as in Kabat. In some embodiments, for example in the context of an Fc region, the lower hinge is included, with the "lower hinge" generally referring to positions 226 or 230. As noted herein, pI variants can be made in the hinge region as well.

The light chain generally comprises two domains, the variable light domain (containing the light chain CDRs and together with the variable heavy domains forming the Fv region), and a constant light chain region (often referred to as CL or Ck).

Another region of interest for additional substitutions, outlined below, is the Fc region. By "Fc" or "Fc region" or "Fc domain" as used herein is meant the polypeptide com-

18

prising the constant region of an antibody excluding the first constant region immunoglobulin domain and in some cases, part of the hinge. Thus Fc refers to the last two constant region immunoglobulin domains of IgA, IgD, and IgG, the last three constant region immunoglobulin domains of IgE and IgM, and the flexible hinge N-terminal to these domains. For IgA and IgM, Fc may include the J chain. For IgG, the Fc domain comprises immunoglobulin domains C γ 2 and C γ 3 (C γ 2 and C γ 3) and the lower hinge region between C γ 1 (C γ 1) and C γ 2 (C γ 2). Although the boundaries of the Fc region may vary, the human IgG heavy chain Fc region is usually defined to include residues C226 or P230 to its carboxyl-terminus, wherein the numbering is according to the EU index as in Kabat. In some embodiments, as is more fully described below, amino acid modifications are made to the Fc region, for example to alter binding to one or more Fc γ R receptors or to the Fc γ Rn receptor.

In some embodiments, the antibodies are full length. By "full length antibody" herein is meant the structure that constitutes the natural biological form of an antibody, including variable and constant regions, including one or more modifications as outlined herein.

Alternatively, the antibodies can include a variety of structures, including, but not limited to, antibody fragments, monoclonal antibodies, bispecific antibodies, minibodies, domain antibodies, synthetic antibodies (sometimes referred to herein as "antibody mimetics"), chimeric antibodies, humanized antibodies, antibody fusions (sometimes referred to as "antibody conjugates"), and fragments of each, respectively.

In one embodiment, the antibody is an antibody fragment, as long as it contains at least one constant domain which can be engineered to produce heterodimers, such as pI engineering. Other antibody fragments that can be used include fragments that contain one or more of the CH1, CH2, CH3, hinge and CL domains of the invention that have been pI engineered. For example, Fc fusions are fusions of the Fc region (CH2 and CH3, optionally with the hinge region) fused to another protein. A number of Fc fusions are known the art and can be improved by the addition of the heterodimerization variants of the invention. In the present case, antibody fusions can be made comprising CH1; CH1, CH2 and CH3; CH2; CH3; CH2 and CH3; CH1 and CH3, any or all of which can be made optionally with the hinge region, utilizing any combination of heterodimerization variants described herein.

In some embodiments of the present invention, one monomer comprises a heavy chain comprises a scFv linked to an Fc domain, and the other monomer comprises a heavy chain comprising a Fab linked to an Fc domain, e.g. a "typical" heavy chain, and a light chain. By "Fab" or "Fab region" as used herein is meant the polypeptide that comprises the VH, CH1, VL, and CL immunoglobulin domains. Fab may refer to this region in isolation, or this region in the context of a full length antibody, antibody fragment or Fab fusion protein. By "Fv" or "Fv fragment" or "Fv region" as used herein is meant a polypeptide that comprises the VL and VH domains of a single antibody.

Chimeric and Humanized Antibodies

In some embodiments, the antibody can be a mixture from different species, e.g. a chimeric antibody and/or a humanized antibody. In general, both "chimeric antibodies" and "humanized antibodies" refer to antibodies that combine regions from more than one species. For example, "chimeric antibodies" traditionally comprise variable region(s) from a mouse (or rat, in some cases) and the constant region(s) from a human. "Humanized antibodies" generally refer to non-

19

human antibodies that have had the variable-domain framework regions swapped for sequences found in human antibodies. Generally, in a humanized antibody, the entire antibody, except the CDRs, is encoded by a polynucleotide of human origin or is identical to such an antibody except within its CDRs. The CDRs, some or all of which are encoded by nucleic acids originating in a non-human organism, are grafted into the beta-sheet framework of a human antibody variable region to create an antibody, the specificity of which is determined by the engrafted CDRs. The creation of such antibodies is described in, e.g., WO 92/11018, Jones, 1986, *Nature* 321:522-525, Verhoeyen et al., 1988, *Science* 239:1534-1536, all entirely incorporated by reference. "Backmutation" of selected acceptor framework residues to the corresponding donor residues is often required to regain affinity that is lost in the initial grafted construct (U.S. Pat. Nos. 5,530,101; 5,585,089; 5,693,761; 5,693,762; 6,180,370; 5,859,205; 5,821,337; 6,054,297; 6,407,213, all entirely incorporated by reference). The humanized antibody optimally also will comprise at least a portion of an immunoglobulin constant region, typically that of a human immunoglobulin, and thus will typically comprise a human Fc region. Humanized antibodies can also be generated using mice with a genetically engineered immune system. Roque et al., 2004, *Biotechnol. Prog.* 20:639-654, entirely incorporated by reference. A variety of techniques and methods for humanizing and reshaping non-human antibodies are well known in the art (See Tsurushita & Vasquez, 2004, *Humanization of Monoclonal Antibodies*, Molecular Biology of B Cells, 533-545, Elsevier Science (USA), and references cited therein, all entirely incorporated by reference). Humanization methods include but are not limited to methods described in Jones et al., 1986, *Nature* 321:522-525; Riechmann et al., 1988, *Nature* 332:323-329; Verhoeyen et al., 1988, *Science*, 239:1534-1536; Queen et al., 1989, *Proc Natl Acad Sci, USA* 86:10029-33; He et al., 1998, *J. Immunol.* 160: 1029-1035; Carter et al., 1992, *Proc Natl Acad Sci USA* 89:4285-9; Presta et al., 1997, *Cancer Res.* 57(20):4593-9; Gorman et al., 1991, *Proc. Natl. Acad. Sci. USA* 88:4181-4185; O'Connor et al., 1998, *Protein Eng* 11:321-8, all entirely incorporated by reference. Humanization or other methods of reducing the immunogenicity of nonhuman antibody variable regions may include resurfacing methods, as described for example in Roguska et al., 1994, *Proc. Natl. Acad. Sci. USA* 91:969-973, entirely incorporated by reference. In one embodiment, the parent antibody has been affinity matured, as is known in the art. Structure-based methods may be employed for humanization and affinity maturation, for example as described in U.S. Ser. No. 11/004,590. Selection based methods may be employed to humanize and/or affinity mature antibody variable regions, including but not limited to methods described in Wu et al., 1999, *J. Mol. Biol.* 294:151-162; Baca et al., 1997, *J. Biol. Chem.* 272(16):10678-10684; Rosok et al., 1996, *J. Biol. Chem.* 271(37): 22611-22618; Rader et al., 1998, *Proc. Natl. Acad. Sci. USA* 95: 8910-8915; Krauss et al., 2003, *Protein Engineering* 16(10):753-759, all entirely incorporated by reference. Other humanization methods may involve the grafting of only parts of the CDRs, including but not limited to methods described in U.S. Ser. No. 09/810,510; Tan et al., 2002, *J. Immunol.* 169:1119-1125; De Pascalis et al., 2002, *J. Immunol.* 169:3076-3084, all entirely incorporated by reference.

Multispecific Antibody Constructs

As will be appreciated by those in the art and discussed more fully below, the heterodimeric fusion proteins of the

20

present invention take on a number variety of configurations, with a preferred embodiment shown in FIG. 1B as a "triple F" construct.

Heterodimeric Heavy Chain Constant Regions

Accordingly, the present invention provides heterodimeric proteins based on the use of monomers containing variant heavy chain constant regions as a first domain. By "monomer" herein is meant one half of the heterodimeric protein. It should be noted that traditional antibodies are actually tetrameric (two heavy chains and two light chains). In the context of the present invention, one pair of heavy-light chains (if applicable, e.g. if the monomer comprises an Fab) is considered a "monomer". Similarly, a heavy chain region comprising the scFv is considered a monomer. In the case where an Fv region is one fusion partner (e.g. heavy and light chain) and a non-antibody protein is another fusion partner, each "half" is considered a monomer. Essentially, each monomer comprises sufficient heavy chain constant region to allow heterodimerization engineering, whether that be all the constant region, e.g. CH1-hinge-CH2-CH3, the Fc region (CH2-CH3), or just the CH3 domain.

The variant heavy chain constant regions can comprise all or part of the heavy chain constant region, including the full length construct, CH1-hinge-CH2-CH3, or portions thereof, including for example CH2-CH3 or CH3 alone. In addition, the heavy chain region of each monomer can be the same backbone (CH1-hinge-CH2-CH3 or CH2-CH3) or different. N- and C-terminal truncations and additions are also included within the definition; for example, some pl variants include the addition of charged amino acids to the C-terminus of the heavy chain domain.

Thus, in general, one monomer of the present "triple F" construct is a scFv region-hinge-Fc domain and the other is (VH-CH1-hinge-CH2-CH3 plus associated light chain), with heterodimerization variants, including steric and pl variants, Fc and FcRn variants, and additional antigen binding domains (with optional linkers) included in these regions.

In addition to the heterodimerization variants (e.g. steric and pl variants) outlined herein, the heavy chain regions may also contain additional amino acid substitutions, including changes for altering FcγR and FcRn binding as discussed below.

In addition, some monomers can utilize linkers between the variant heavy chain constant region and the fusion partner. For the scFv portion of the "bottle-opener", standard linkers as are known in the art can be used. In the case where additional fusion partners are made (e.g. FIGS. 2 and 3), traditional peptide linkers can be used, including flexible linkers of glycine and serine. In some cases, the linkers for use as components of the monomer are different from those defined below for the ADC constructs, and are in many embodiments not cleavable linkers (such as those susceptible to proteases), although cleavable linkers may find use in some embodiments.

The heterodimerization variants include a number of different types of variants, including, but not limited to, steric variants (including charge variants) and pl variants, that can be optionally and independently combined with any other variants. In these embodiments, it is important to match "monomer A" with "monomer B"; that is, if a heterodimeric protein relies on both steric variants and pl variants, these need to be correctly matched to each monomer: e.g. the set of steric variants that work (1 set on monomer A, 1 set on monomer B) is combined with pl

21

variant sets (1 set on monomer A, 1 set on monomer B), such that the variants on each monomer are designed to achieve the desired function.

It is important to note that the heterodimerization variants outlined herein (for example, including but not limited to those variants shown in FIGS. 4, 5 and 6), can be optionally and independently combined with any other variants, and on any other monomer. Thus, for example, pI variants for monomer 1 from FIG. 6 can be added to other heterodimerization variants for monomer 1 in FIGS. 5A-5B and 6 or from monomer 2. That is, what is important for the heterodimerization is that there are "sets" of variants, one set for one monomer and one set for the other. Whether these are combined from the FIGS. 1 to 1 (e.g. monomer 1 listings can go together) or switched (monomer 1 pI variants with monomer 2 steric variants) is irrelevant. However, as noted herein, "strandedness" should be preserved when combinations are made as outlined above. Furthermore, for the additional Fc variants (such as for FcγR binding, FcRn binding, etc.), either monomer, or both monomers, can include any of the listed variants, independently and optionally. In some cases, both monomers have the additional variants and in some only one monomer has the additional variants, or they can be combined.

Heterodimerization Variants

The present invention provides multispecific antibody formats, on a "triple F" or "bottle opener" scaffold as depicted in FIG. 1A-3E.

Steric Variants

In some embodiments, the formation of heterodimers can be facilitated by the addition of steric variants. That is, by changing amino acids in each heavy chain, different heavy chains are more likely to associate to form the heterodimeric structure than to form homodimers with the same Fc amino acid sequences. Suitable steric variants are shown in FIGS. 4 and 5.

One mechanism is generally referred to in the art as "knobs and holes", referring to amino acid engineering that creates steric influences to favor heterodimeric formation and disfavor homodimeric formation can also optionally be used; this is sometimes referred to as "knobs and holes", as described in U.S. Ser. No. 61/596,846, Ridgway et al., Protein Engineering 9(7):617 (1996); Atwell et al., J. Mol. Biol. 1997 270:26; U.S. Pat. No. 8,216,805, all of which are hereby incorporated by reference in their entirety. FIGS. 4 and 5, further described below, identifies a number of "monomer A-monomer B" pairs that rely on "knobs and holes". In addition, as described in Merchant et al., Nature Biotech. 16:677 (1998), these "knobs and hole" mutations can be combined with disulfide bonds to skew formation to heterodimerization.

An additional mechanism that finds use in the generation of heterodimers is sometimes referred to as "electrostatic steering" as described in Gunasekaran et al., J. Biol. Chem. 285(25):19637 (2010), hereby incorporated by reference in its entirety. This is sometimes referred to herein as "charge pairs". In this embodiment, electrostatics are used to skew the formation towards heterodimerization. As those in the art will appreciate, these may also have an effect on pI, and thus on purification, and thus could in some cases also be considered pI variants. However, as these were generated to force heterodimerization and were not used as purification tools, they are classified as "steric variants". These include, but are not limited to, D221E/P228E/L368E paired with D221R/P228R/K409R (e.g. these are "monomer corresponding sets") and C220E/P228E/368E paired with C220R/E224R/P228R/K409R.

22

Additional monomer A and monomer B variants that can be combined with other variants, optionally and independently in any amount, such as pI variants outlined herein or other steric variants that are shown in FIG. 37 of US 2012/0149876, the figure and legend of which are incorporated expressly by reference herein.

In some embodiments, the steric variants outlined herein can be optionally and independently incorporated with any pI variant (or other variants such as Fc variants, FcRn variants, etc.) into one or both monomers.

pI (Isoelectric Point) Variants for Heterodimers

In general, as will be appreciated by those in the art, there are two general categories of pI variants: those that increase the pI of the protein (basic changes) and those that decrease the pI of the protein (acidic changes). As described herein, all combinations of these variants can be done: one monomer may be wild type, or a variant that does not display a significantly different pI from wild-type, and the other can be either more basic or more acidic. Alternatively, each monomer is changed, one to more basic and one to more acidic.

Preferred combinations of pI variants are shown in FIG. 6, with additional pI variants disclosed in FIGS. 20, 21 and 22.

Heavy Chain Acidic pI Changes

Accordingly, when one monomer comprising a variant heavy chain constant domain is to be made more positive (e.g. lower the pI), one or more of the following substitutions can be made: S119E, K133E, K133Q, T164E, K205E, K205Q, N208D, K210E, K210Q, K274E, K320E, K322E, K326E, K334E, R355E, K392E, a deletion of K447, adding peptide DEDE at the c-terminus, G137E, N203D, K274Q, R355Q, K392N and Q419E. As outlined herein and shown in the figures, these changes are shown relative to IgG1, but all isotypes can be altered this way, as well as isotype hybrids.

In the case where the heavy chain constant domain is from IgG2-4, R133E and R133Q can also be used.

Basic pI Changes

Accordingly, when one monomer comprising a variant heavy chain constant domain is to be made more negative (e.g. increase the pI), one or more of the following substitutions can be made: Q196K, P217R, P228R, N276K and H435R. As outlined herein and shown in the figures, these changes are shown relative to IgG1, but all isotypes can be altered this way, as well as isotype hybrids.

Antibody Heterodimers Light Chain Variants

In the case of antibody based heterodimers, e.g. where at least one of the monomers comprises a light chain in addition to the heavy chain domain, pI variants can also be made in the light chain. Amino acid substitutions for lowering the pI of the light chain include, but are not limited to, K126E, K126Q, K145E, K145Q, N152D, S156E, K169E, S202E, K207E and adding peptide DEDE at the c-terminus of the light chain. Changes in this category based on the constant lambda light chain include one or more substitutions at R108Q, Q124E, K126Q, N138D, K145T and Q199E. In addition, increasing the pI of the light chains can also be done.

Isotypic Variants

In addition, many embodiments of the invention rely on the "importation" of pI amino acids at particular positions from one IgG isotype into another, thus reducing or eliminating the possibility of unwanted immunogenicity being introduced into the variants. That is, IgG1 is a common isotype for therapeutic antibodies for a variety of reasons, including high effector function. However, the heavy con-

23

stant region of IgG1 has a higher pI than that of IgG2 (8.10 versus 7.31). By introducing IgG2 residues at particular positions into the IgG1 backbone, the pI of the resulting monomer is lowered (or increased) and additionally exhibits longer serum half-life. For example, IgG1 has a glycine (pI 5.97) at position 137, and IgG2 has a glutamic acid (pI 3.22); importing the glutamic acid will affect the pI of the resulting protein. As is described below, a number of amino acid substitutions are generally required to significantly affect the pI of the variant antibody. However, it should be noted as discussed below that even changes in IgG2 molecules allow for increased serum half-life.

In other embodiments, non-isotypic amino acid changes are made, either to reduce the overall charge state of the resulting protein (e.g. by changing a higher pI amino acid to a lower pI amino acid), or to allow accommodations in structure for stability, etc. as is more further described below.

In addition, by pI engineering both the heavy and light constant domains, significant changes in each monomer of the heterodimer can be seen. As discussed herein, having the pIs of the two monomers differ by at least 0.5 can allow separation by ion exchange chromatography or isoelectric focusing, or other methods sensitive to isoelectric point. Calculating pI

The pI of each monomer can depend on the pI of the variant heavy chain constant domain and the pI of the total monomer, including the variant heavy chain constant domain and the fusion partner. Thus, in some embodiments, the change in pI is calculated on the basis of the variant heavy chain constant domain, using the chart in the Figures. Alternatively, the pI of each monomer can be compared.

pI Variants that Also Confer Better FcRn In Vivo Binding

In the case where the pI variant decreases the pI of the monomer, they can have the added benefit of improving serum retention in vivo.

Although still under examination, Fc regions are believed to have longer half-lives in vivo, because binding to FcRn at pH 6 in an endosome sequesters the Fc (Ghetie and Ward, 1997 Immunol Today. 18(12): 592-598, entirely incorporated by reference). The endosomal compartment then recycles the Fc to the cell surface. Once the compartment opens to the extracellular space, the higher pH, ~7.4, induces the release of Fc back into the blood. In mice, Dall'Acqua et al. showed that Fc mutants with increased FcRn binding at pH 6 and pH 7.4 actually had reduced serum concentrations and the same half life as wild-type Fc (Dall'Acqua et al. 2002, J. Immunol. 169:5171-5180, entirely incorporated by reference). The increased affinity of Fc for FcRn at pH 7.4 is thought to forbid the release of the Fc back into the blood. Therefore, the Fc mutations that will increase Fc's half-life in vivo will ideally increase FcRn binding at the lower pH while still allowing release of Fc at higher pH. The amino acid histidine changes its charge state in the pH range of 6.0 to 7.4. Therefore, it is not surprising to find His residues at important positions in the Fc/FcRn complex.

Recently it has been suggested that antibodies with variable regions that have lower isoelectric points may also have longer serum half-lives (Igawa et al., 2010 PEDS. 23(5): 385-392, entirely incorporated by reference). However, the mechanism of this is still poorly understood. Moreover, variable regions differ from antibody to antibody. Constant region variants with reduced pI and extended half-life would provide a more modular approach to improving the pharmacokinetic properties of antibodies, as described herein.

24

pI variants that find use in this embodiment, as well as their use for purification optimization, are disclosed in FIG. 20A-20F.

Combination of Heterodimeric Variants

As will be appreciated by those in the art, all of the recited heterodimerization variants can be optionally and independently combined in any way, as long as they retain their "strandedness" or "monomer partition". In addition, all of these variants can be combined into any of the heterodimerization formats. See FIG. 24.

In the case of pI variants, while embodiments finding particular use are shown in the Figures, other combinations can be generated, following the basic rule of altering the pI difference between two monomers to facilitate purification.

Target Antigens

The immunoglobulins of the invention may target virtually any antigens. The "triple F" format is particularly beneficial for targeting two (or more) distinct antigens. (As outlined herein, this targeting can be any combination of monovalent and divalent binding, depending on the format). Thus the immunoglobulins herein preferably co-engage two target antigens, although in some cases, three or four antigens can be monovalently engaged. Each monomer's specificity can be selected from the lists below. While the triple F immunoglobulins described herein are particularly beneficial for targeting distinct antigens, in some cases it may be beneficial to target only one antigen. That is, each monomer may have specificity for the same antigen.

Particular suitable applications of the immunoglobulins herein are co-target pairs for which it is beneficial or critical to engage each target antigen monovalently. Such antigens may be, for example, immune receptors that are activated upon immune complexation. Cellular activation of many immune receptors occurs only by cross-linking, achieved typically by antibody/antigen immune complexes, or via effector cell to target cell engagement. For some immune receptors, for example the CD3 signaling receptor on T cells, activation only upon engagement with co-engaged target is critical, as nonspecific cross-linking in a clinical setting can elicit a cytokine storm and toxicity. Therapeutically, by engaging such antigens monovalently rather than multivalently, using the immunoglobulins herein, such activation occurs only in response to cross-linking only in the microenvironment of the primary target antigen. The ability to target two different antigens with different valencies is a novel and useful aspect of the present invention. Examples of target antigens for which it may be therapeutically beneficial or necessary to co-engage monovalently include but are not limited to immune activating receptors such as CD3, FcγRs, toll-like receptors (TLRs) such as TLR4 and TLR9, cytokine, chemokine, cytokine receptors, and chemokine receptors. In many embodiments, one of the antigen binding sites binds to CD3, and in some embodiments it is the scFv-containing monomer.

Virtually any antigen may be targeted by the immunoglobulins herein, including but not limited to proteins, subunits, domains, motifs, and/or epitopes belonging to the following list of target antigens, which includes both soluble factors such as cytokines and membrane-bound factors, including transmembrane receptors: 17-1A, 4-1BB, 4Dc, 6-keto-PGF1a, 8-iso-PGF2a, 8-oxo-dG, A1 Adenosine Receptor, A33, ACE, ACE-2, Activin, Activin A, Activin AB, Activin B, Activin C, Activin RIA, Activin RIA ALK-2, Activin RIB ALK-4, Activin RIIA, Activin RIIIB, ADAM, ADAM10, ADAM12, ADAM15, ADAM17/TACE, ADAM8, ADAM9, ADAMTS, ADAMTS4, ADAMTS5, Addressins, aFGF, ALCAM, ALK, ALK-1, ALK-7, alpha-

25

1-antitrypsin, alpha-V/beta-1 antagonist, ANG, Ang, APAF-1, APE, API, APP, APRIL, AR, ARC, ART, Artemin, anti-Id, ASPARTIC, Atrial natriuretic factor, av/b3 integrin, Axl, b2M, B7-1, B7-2, B7-H, B-lymphocyte Stimulator (BlyS), BACE, BACE-1, Bad, BAFF, BAFF-R, Bag-1, BAK, Bax, BCA-1, BCAM, Bcl, BCMA, BDNF, b-ECGF, bFGF, BID, Bik, BIM, BLC, BL-CAM, BLK, BMP, BMP-2 BMP-2a, BMP-3 Osteogenin, BMP-4 BMP-2b, BMP-5, BMP-6 Vgr-1, BMP-7 (OP-1), BMP-8 (BMP-8a, OP-2), BMPR, BMPR-1A (ALK-3), BMPR-1B (ALK-6), BRK-2, RPK-1, BMPR-II (BRK-3), BMPs, b-NGF, BOK, Bombesin, Bone-derived neurotrophic factor, BPDE, BPDE-DNA, BTC, complement factor 3 (C3), C3a, C4, C5, C5a, C10, CA125, CAD-8, Calcitonin, cAMP, carcinoembryonic antigen (CEA), carcinoma-associated antigen, Cathepsin A, Cathepsin B, Cathepsin C/DPPI, Cathepsin D, Cathepsin E, Cathepsin H, Cathepsin L, Cathepsin O, Cathepsin S, Cathepsin V, Cathepsin X/Z/P, CBL, CCI, CCK2, CCL, CCL1, CCL11, CCL12, CCL13, CCL14, CCL15, CCL16, CCL17, CCL18, CCL19, CCL2, CCL20, CCL21, CCL22, CCL23, CCL24, CCL25, CCL26, CCL27, CCL28, CCL3, CCL4, CCL5, CCL6, CCL7, CCL8, CCL9/10, CCR, CCR1, CCR10, CCR10, CCR2, CCR3, CCR4, CCR5, CCR6, CCR7, CCR8, CCR9, CD1, CD2, CD3, CD3E, CD4, CD5, CD6, CD7, CD8, CD10, CD11a, CD11b, CD11c, CD13, CD14, CD15, CD16, CD18, CD19, CD20, CD21, CD22, CD23, CD25, CD27L, CD28, CD29, CD30, CD30L, CD32, CD33 (p67 proteins), CD34, CD38, CD40, CD40L, CD44, CD45, CD46, CD49a, CD52, CD54, CD55, CD56, CD61, CD64, CD66e, CD74, CD80 (B7-1), CD89, CD95, CD123, CD137, CD138, CD140a, CD146, CD147, CD148, CD152, CD164, CEACAM5, CFTR, cGMP, CINC, *Clostridium botulinum* toxin, *Clostridium perfringens* toxin, CKb8-1, CLC, CMV, CMV UL, CNTF, CNTN-1, COX, C-Ret, CRG-2, CT-1, CTACK, CTGF, CTLA-4, CX3CL1, CX3CR1, CXCL, CXCL1, CXCL2, CXCL3, CXCL4, CXCL5, CXCL6, CXCL7, CXCL8, CXCL9, CXCL10, CXCL11, CXCL12, CXCL13, CXCL14, CXCL15, CXCL16, CXCR, CXCR1, CXCR2, CXCR3, CXCR4, CXCR5, CXCR6, cytokeratin tumor-associated antigen, DAN, DCC, DcR3, DC-SIGN, Decay accelerating factor, des(1-3)-IGF-I (brain IGF-1), Dhh, digoxin, DNAM-1, Dnase, Dpp, DPPIV/CD26, Dtk, ECAD, EDA, EDA-A1, EDA-A2, EDAR, EGF, EGFR (ErbB-1), EMA, EMMPRIN, ENA, endothelin receptor, Enkephalinase, eNOS, Eot, eotaxinl, EpCAM, Ephrin B2/EphB4, EPO, ERCC, E-selectin, ET-1, Factor IIa, Factor VII, Factor VIIIc, Factor IX, fibroblast activation protein (FAP), Fas, FcR1, FEN-1, Ferritin, FGF, FGF-19, FGF-2, FGF3, FGF-8, FGFR, FGFR-3, Fibrin, FL, FLIP, Flt-3, Flt-4, Follicle stimulating hormone, Fractalkine, FZD1, FZD2, FZD3, FZD4, FZD5, FZD6, FZD7, FZD8, FZD9, FZD10, G250, Gas 6, GCP-2, GCSF, GD2, GD3, GDF, GDF-1, GDF-3 (Vgr-2), GDF-5 (BMP-14, CDMP-1), GDF-6 (BMP-13, CDMP-2), GDF-7 (BMP-12, CDMP-3), GDF-8 (Myostatin), GDF-9, GDF-15 (MIC-1), GDNF, GDNF, GFAP, GFRa-1, GFR-alpha, GFR-alpha2, GFR-alpha3, GTR, Glucagon, Glut 4, glycoprotein Hb/IIIa (GP Hb/IIIa), GM-CSF, gp130, gp72, GRO, Growth hormone releasing factor, Hapten (NP-cap or NIP-cap), HB-EGF, HCC, HCMV gB envelope glycoprotein, HCMV gH envelope glycoprotein, HCMV UL, Hemopoietic growth factor (HGF), Hep B gp120, heparanase, Her2, Her2/neu (ErbB-2), Her3 (ErbB-3), Her4 (ErbB-4), herpes simplex virus (HSV) gB glycoprotein, HSV gD glycoprotein, HGFA, High molecular weight melanoma-associated antigen (HMW-MAA), HIV gp120, HIV IIIB gp 120 V3 loop, HLA, HLA-DR, HM1.24, HMFG PEM, HRG, Hrk, human car-

26

diac myosin, human cytomegalovirus (HCMV), human growth hormone (HGH), HVEM, 1-309, IAP, ICAM, ICAM-1, ICAM-3, ICE, ICOS, IFNg, Ig, IgA receptor, IGF, IGF binding proteins, IGF-1R, IGFBP, IGF-I, IGF-II, IL, IL-1, IL-1R, IL-2, IL-2R, IL-4, IL-4R, IL-5, IL-5R, IL-6, IL-6R, IL-8, IL-9, IL-10, IL-12, IL-13, IL-15, IL-18, IL-18R, IL-23, interferon (INF)-alpha, INF-beta, INF-gamma, Inhibin, iNOS, Insulin A-chain, Insulin B-chain, Insulin-like growth factor 1, integrin alpha2, integrin alpha3, integrin alpha4, integrin alpha4/beta1, integrin alpha4/beta7, integrin alpha5 (alphaV), integrin alpha5/beta1, integrin alpha5/beta3, integrin alpha6, integrin beta1, integrin beta2, interferon gamma, IP-10, I-TAC, JE, Kallikrein 2, Kallikrein 5, Kallikrein 6, Kallikrein 11, Kallikrein 12, Kallikrein 14, Kallikrein 15, Kallikrein L1, Kallikrein L2, Kallikrein L3, Kallikrein L4, KC, KDR, Keratinocyte Growth Factor (KGF), laminin 5, LAMP, LAP, LAP (TGF-1), Latent TGF-1, Latent TGF-1 bpl, LBP, LDGF, LECT2, Lefty, Lewis-Y antigen, Lewis-Y related antigen, LFA-1, LFA-3, Lfo, LIF, LIGHT, lipoproteins, LIX, LKN, Lptn, L-Selectin, LT-a, LT-b, LTB4, LTBP-1, Lung surfactant, Luteinizing hormone, Lymphotoxin Beta Receptor, Mac-1, MadCAM, MAG, MAP2, MARC, MCAM, MCAM, MCK-2, MCP, M-CSF, MDC, Mer, METALLOPROTEASES, MGDF receptor, MGMT, MHC (HLA-DR), MIF, MIG, MIP, MIP-1-alpha, MK, MMAC1, MMP, MMP-1, MMP-10, MMP-11, MMP-12, MMP-13, MMP-14, MMP-15, MMP-2, MMP-24, MMP-3, MMP-7, MMP-8, MMP-9, MPIF, Mpo, MSK, MSP, mucin (Muc1), MUC18, Muellerian-inhibitin substance, Mug, MuSK, NAIP, NAP, NCAD, N-Cadherin, NCA 90, NCAM, NCAM, Neprilysin, Neurotrophin-3, -4, or -6, Neurturin, Neuronal growth factor (NGF), NGFR, NGF-beta, nNOS, NO, NOS, Npn, NRG-3, NT, NTN, OB, OGG1, OPG, OPN, OSM, OX40L, OX40R, p150, p95, PADPr, Parathyroid hormone, PARC, PARP, PBR, PBSF, PCAD, P-Cadherin, PCNA, PDGF, PDGF, PDK-1, PECAM, PEM, PF4, PGE, PGE, PG12, PGJ2, PIN, PLA2, placental alkaline phosphatase (PLAP), PIGF, PLP, PP14, Proinsulin, Prorelaxin, Protein C, PS, PSA, PSCA, prostate specific membrane antigen (PSMA), PTEN, PTHrp, Ptk, PTN, R51, RANK, RANKL, RANTES, RANTES, Relaxin A-chain, Relaxin B-chain, renin, respiratory syncytial virus (RSV) F, RSV Fgp, Ret, Rheumatoid factors, RLIP76, RPA2, RSK, S100, SCF/KL, SDF-1, SERINE, Serum albumin, sFRP-3, Shh, SIGIRR, SK-1, SLAM, SLPI, SMAC, SMDF, SMOH, SOD, SPARC, Stat, STEAP, STEAP-II, TACE, TACI, TAG-72 (tumor-associated glycoprotein-72), TARC, TCA-3, T-cell receptors (e.g., T-cell receptor alpha/beta), TdT, TECK, TEM1, TEMS, TEM7, TEM8, TERT, testicular PLAP-like alkaline phosphatase, Tfr, TGF, TGF-alpha, TGF-beta, TGF-beta Pan Specific, TGF-beta RI (ALK-5), TGF-beta RII, TGF-beta RIIB, TGF-beta RIII, TGF-beta1, TGF-beta2, TGF-beta3, TGF-beta4, TGF-beta5, Thrombin, Thymus Ck-1, Thyroid stimulating hormone, Tie, TIMP, TIQ, Tissue Factor, TMEFF2, Tmpo, TMPRSS2, TNF, TNF-alpha, TNF-alpha beta, TNF-beta2, TNF-c, TNF-RI, TNF-RII, TNFRSF10A (TRAIL R1 Apo-2, DR4), TNFRSF10B (TRAIL R2 DRS, KILLER, TRICK-2A, TRICK-B), TNFRSF10C (TRAIL R3 DcR1, LIT, TRID), TNFRSF10D (TRAIL R4 DcR2, TRUND), TNFRSF11A (RANK ODF R, TRANCE R), TNFRSF11B (OPG OCIF, TR1), TNFRSF12 (TWEAK R FN14), TNFRSF13B (TACI), TNFRSF13C (BAFF R), TNFRSF14 (HVEM ATAR, HveA, LIGHT R, TR2), TNFRSF16 (NGFR p75NTR), TNFRSF17 (BCMA), TNFRSF18 (GITR AITR), TNFRSF19 (TROY TAJ, TRADE), TNFRSF19L (RELT), TNFRSF1A (TNF RI CD120a, p55-60), TNFRSF1B (TNF

RII CD120b, p75-80), TNFRSF26 (TNFRH3), TNFRSF3 (LTbR TNF RIII, TNFC R), TNFRSF4 (OX40 ACT35, TXGP1 R), TNFRSF5 (CD40 p50), TNFRSF6 (Fas Apo-1, APT1, CD95), TNFRSF6B (DcR3 M68, TR6), TNFRSF7 (CD27), TNFRSF8 (CD30), TNFRSF9 (4-1BB CD137, ILA), TNFRSF21 (DR6), TNFRSF22 (DcTRAIL R2 TNFRH2), TNFRST23 (DcTRAIL R1 TNFRH1), TNFRSF25 (DR3 Apo-3, LARD, TR-3, TRAMP, WSL-1), TNFSF10 (TRAIL Apo-2 Ligand, TL2), TNFSF11 (TRANCE/RANK Ligand, OPG Ligand), TNFSF12 (TWEAK Apo-3 Ligand, DR3 Ligand), TNFSF13 (APRIL TALL2), TNFSF13B (BAFF BLYS, TALL1, THANK, TNFSF20), TNFSF14 (LIGHT HVEM Ligand, LTg), TNFSF15 (TL1A/VEGI), TNFSF18 (GITR Ligand AITR Ligand, TL6), TNFSF1A (TNF-a Conectin, DIF, TNFSF2), TNFSF1B (TNF-b LTa, TNFSF1), TNFSF3 (LTb TNFC, p33), TNFSF4 (OX40 Ligand gp34, TXGP1), TNFSFS (CD40 Ligand CD154, gp39, HIGM1, IMD3, TRAP), TNFSF6 (Fas Ligand Apo-1 Ligand, APT1 Ligand), TNFSF7 (CD27 Ligand CD70), TNFSF8 (CD30 Ligand CD153), TNFSF9 (4-1BB Ligand CD137 Ligand), TP-1, t-PA, Tpo, TRAIL, TRAIL R, TRAIL-R1, TRAIL-R2, TRANCE, transferring receptor, TRF, Trk, TROP-2, TSG, TSLP, tumor-associated antigen CA 125, tumor-associated antigen expressing Lewis Y related carbohydrate, TWEAK, TXB2, Ung, uPAR, uPAR-1, Urokinase, VCAM, VCAM-1, VECAD, VE-Cadherin, VE-cadherin-2, VEGFR-1 (flt-1), VEGF, VEGFR, VEGFR-3 (flt-4), VEGI, VIM, Viral antigens, VLA, VLA-1, VLA-4, VNR integrin, von Willebrands factor, WIF-1, WNT1, WNT2, WNT2B/13, WNT3, WNT3A, WNT4, WNT5A, WNT5B, WNT6, WNT7A, WNT7B, WNT8A, WNT8B, WNT9A, WNT9A, WNT9B, WNT10A, WNT10B, WNT11, WNT16, XCL1, XCL2, XCR1, XCR1, XEDAR, XIAP, XPD, and receptors for hormones and growth factors. To form the bispecific or trispecific antibodies of the invention, antibodies to any combination of these antigens can be made; that is, each of these antigens can be optionally and independently included or excluded from a multispecific antibody according to the present invention.

Exemplary antigens that may be targeted specifically by the immunoglobulins of the invention include but are not limited to: CD20, CD19, Her2, EGFR, EpCAM, CD3, FcγRIIIa (CD16), FcγRIIa (CD32a), FcγRIIb (CD32b), FcγRI (CD64), Toll-like receptors (TLRs) such as TLR4 and TLR9, cytokines such as IL-2, IL-5, IL-13, IL-12, IL-23, and TNFα, cytokine receptors such as IL-2R, chemokines, chemokine receptors, growth factors such as VEGF and HGF, and the like. To form the multispecific antibodies of the invention, antibodies to any combination of these antigens can be made; that is, each of these antigens can be optionally and independently included or excluded from a multispecific antibody according to the present invention.

Particularly preferred combinations for bispecific antibodies are an antigen-binding domain to CD3 and an antigen binding domain to CD19; an antigen-binding domain to CD3 and an antigen binding domain to CD33; an antigen-binding domain to CD3 and an antigen binding domain to CD 38. Again, in many embodiments, the CD3 binding domain is the scFv, having an exemplary sequence as depicted in the Figures and/or CD3 CDRs as outlined.

The choice of suitable target antigens and co-targets depends on the desired therapeutic application. Some targets that have proven especially amenable to antibody therapy are those with signaling functions. Other therapeutic antibodies exert their effects by blocking signaling of the receptor by inhibiting the binding between a receptor and its

cognate ligand. Another mechanism of action of therapeutic antibodies is to cause receptor down regulation. Other antibodies do not work by signaling through their target antigen. The choice of co-targets will depend on the detailed biology underlying the pathology of the indication that is being treated.

Monoclonal antibody therapy has emerged as an important therapeutic modality for cancer (Weiner et al., 2010, *Nature Reviews Immunology* 10:317-327; Reichert et al., 2005, *Nature Biotechnology* 23[9]:1073-1078; herein expressly incorporated by reference). For anti-cancer treatment it may be desirable to target one antigen (antigen-1) whose expression is restricted to the cancerous cells while co-targeting a second antigen (antigen-2) that mediates some immunological killing activity. For other treatments it may be beneficial to co-target two antigens, for example two angiogenic factors or two growth factors, that are each known to play some role in proliferation of the tumor. Exemplary co-targets for oncology include but are not limited to HGF and VEGF, IGF-1R and VEGF, Her2 and VEGF, CD19 and CD3, CD20 and CD3, Her2 and CD3, CD19 and FcγRIIIa, CD20 and FcγRIIIa, Her2 and FcγRIIIa. An immunoglobulin of the invention may be capable of binding VEGF and phosphatidylserine; VEGF and ErbB3; VEGF and PLGF; VEGF and ROBO4; VEGF and BSG2; VEGF and CDCP1; VEGF and ANPEP; VEGF and c-MET; HER-2 and ERB3; HER-2 and BSG2; HER-2 and CDCP1; HER-2 and ANPEP; EGFR and CD64; EGFR and BSG2; EGFR and CDCP1; EGFR and ANPEP; IGF1R and PDGFR; IGF1R and VEGF; IGF1R and CD20; CD20 and CD74; CD20 and CD30; CD20 and DR4; CD20 and VEGFR2; CD20 and CD52; CD20 and CD4; HGF and c-MET; HGF and NRP1; HGF and phosphatidylserine; ErbB3 and IGF1R; ErbB3 and IGF1,2; c-Met and Her-2; c-Met and NRP1; c-Met and IGF1R; IGF1,2 and PDGFR; IGF1,2 and CD20; IGF1,2 and IGF1R; IGF2 and EGFR; IGF2 and HER2; IGF2 and CD20; IGF2 and VEGF; IGF2 and IGF1R; IGF1 and IGF2; PDGFRa and VEGFR2; PDGFRa and PLGF; PDGFRa and VEGF; PDGFRa and c-Met; PDGFRa and EGFR; PDGFRb and VEGFR2; PDGFRb and c-Met; PDGFRb and EGFR; RON and c-Met; RON and MTSP1; RON and MSP; RON and CDCP1; VGFR1 and PLGF; VGFR1 and RON; VGFR1 and EGFR; VEGFR2 and PLGF; VEGFR2 and NRP1; VEGFR2 and RON; VEGFR2 and DLL4; VEGFR2 and EGFR; VEGFR2 and ROBO4; VEGFR2 and CD55; LPA and SIP; EPHB2 and RON; CTLA4 and VEGF; CD3 and EPCAM; CD40 and IL6; CD40 and IGF; CD40 and CD56; CD40 and CD70; CD40 and VEGFR1; CD40 and DRS; CD40 and DR4; CD40 and APRIL; CD40 and BCMA; CD40 and RANKL; CD28 and MAPG; CD80 and CD40; CD80 and CD30; CD80 and CD33; CD80 and CD74; CD80 and CD2; CD80 and CD3; CD80 and CD19; CD80 and CD4; CD80 and CD52; CD80 and VEGF; CD80 and DRS; CD80 and VEGFR2; CD22 and CD20; CD22 and CD80; CD22 and CD40; CD22 and CD23; CD22 and CD33; CD22 and CD74; CD22 and CD19; CD22 and DRS; CD22 and DR4; CD22 and VEGF; CD22 and CD52; CD30 and CD20; CD30 and CD22; CD30 and CD23; CD30 and CD40; CD30 and VEGF; CD30 and CD74; CD30 and CD19; CD30 and DR5; CD30 and DR4; CD30 and VEGFR2; CD30 and CD52; CD30 and CD4; CD138 and RANKL; CD33 and FTL3; CD33 and VEGF; CD33 and VEGFR2; CD33 and CD44; CD33 and DR4; CD33 and DR5; DR4 and CD137; DR4 and IGF1,2; DR4 and IGF1R; DR4 and DR5; DR5 and CD40; DR5 and CD137; DR5 and CD20; DR5 and EGFR; DR5 and IGF1,2; DR5 and IGF1R,

29

DR5 and HER-2, and EGFR and DLL4. Other target combinations include one or more members of the EGF/erb-2/erb-3 family.

Other targets (one or more) involved in oncological diseases that the immunoglobulins herein may bind include, but are not limited to those selected from the group consisting of: CD52, CD20, CD19, CD3, CD4, CD8, BMP6, IL12A, IL1A, IL1B, IL2, IL24, INHA, TNF, TNFSF10, BMP6, EGF, FGF1, FGF10, FGF11, FGF12, FGF13, FGF14, FGF16, FGF17, FGF18, FGF19, FGF2, FGF20, FGF21, FGF22, FGF23, FGF3, FGF4, FGF5, FGF6, FGF7, FGF8, FGF9, GRP, IGF1, IGF2, IL12A, IL1A, IL1B, IL2, INHA, TGFA, TGFB1, TGFB2, TGFB3, VEGF, CDK2, FGF10, FGF18, FGF2, FGF4, FGF7, IGF1R, IL2, BCL2, CD164, CDKN1A, CDKN1B, CDKN1C, CDKN2A, CDKN2B, CDKN2C, CDKN3, GNRH1, IGFBP6, IL1A, IL1B, ODZ1, PAWR, PLG, TGFB11, AR, BRCA1, CDK3, CDK4, CDK5, CDK6, CDK7, CDK9, E2F1, EGFR, ENO1, ERBB2, ESR1, ESR2, IGFBP3, IGFBP6, IL2, INSL4, MYC, NOX5, NR6A1, PAP, PCNA, PRKCQ, PRKD1, PRL, TP53, FGF22, FGF23, FGF9, IGFBP3, IL2, INHA, KLK6, TP53, CHGB, GNRH1, IGF1, IGF2, INHA, INSL3, INSL4, PRL, KLK6, SHBG, NR1D1, NR1H3, NR1I3, NR2F6, NR4A3, ESR1, ESR2, NROB1, NROB2, NR1D2, NR1H2, NR1H4, NR112, NR2C1, NR2C2, NR2E1, NR2E3, NR2F1, NR2F2, NR3C1, NR3C2, NR4A1, NR4A2, NR5A1, NR5A2, NR6 PGR, RARB, FGF1, FGF2, FGF6, KLK3, KRT1, APOC1, BRCA1, CHGA, CHGB, CLU, COL1A1, COL6A1, EGF, ERBB2, ERK8, FGF1, FGF10, FGF11, FGF13, FGF14, FGF16, FGF17, FGF18, FGF2, FGF20, FGF21, FGF22, FGF23, FGF3, FGF4, FGF5, FGF6, FGF7, FGF8, FGF9, GNRH1, IGF1, IGF2, IGFBP3, IGFBP6, IL12A, IL1A, IL1B, IL2, IL24, INHA, INSL3, INSL4, KLK10, KLK12, KLK13, KLK14, KLK15, KLK3, KLK4, KLK5, KLK6, KLK9, MMP2, MMP9, MSMB, NTN4, ODZ1, PAP, PLAUI, PRL, PSAP, SERPINA3, SHBG, TGFA, TIMP3, CD44, CDH1, CDH10, CDH19, CDH20, CDH7, CDH9, CDH1, CDH10, CDH13, CDH18, CDH19, CDH20, CDH7, CDH8, CDH9, ROBO2, CD44, ILK, ITGA1, APC, CD164, COL6A1, MTSS1, PAP, TGFB11, AGR2, AIG1, AKAP1, AKAP2, CANT1, CAV1, CDH12, CLDN3, CLN3, CYB5, CYC1, DAB2IP, DES, DNCL1, ELAC2, ENO2, ENO3, FASN, FLJ12584, FLJ25530, GAGEB1, GAGEC1, GGT1, GSTP1, HIP1, HUMCYT2A, IL29, K6HF, KAI1, KRT2A, MIB1, PART1, PATE, PCA3, PIAS2, PIK3CG, PPID, PRI, PSCA, SLC2A2, SLC33 μ l, SLC43 μ l, STEAP, STEAP2, TPM1, TPM2, TRPC6, ANGPT1, ANGPT2, ANPEP, ECGF1, EREG, FGF1, FGF2, FIGF, FLT1, JAG1, KDR, LAMA5, NRP1, NRP2, PGF, PLXDC1, STAB 1, VEGF, VEGFC, ANGPTL3, BAI1, COL4A3, IL8, LAMA5, NRP1, NRP2, STAB 1, ANGPTL4, PECAM1, PF4, PROK2, SERPINF1, TNFAIP2, CCL11, CCL2, CXCL1, CXCL10, CXCL3, CXCL5, CXCL6, CXCL9, IFNA1, IFNB1, IFNG, IL1B, IL6, MDK, EDG1, EFNA1, EFNA3, EFNB2, EGF, EPHB4, FGFR3, HGF, IGF1, ITGB3, PDGFA, TEK, TGFA, TGFB1, TGFB2, TGFB1, CCL2, CDH5, COL1A1, EDG1, ENG, ITGAV, ITGB3, THBS1, THBS2, BAD, BAG1, BCL2, CCNA1, CCNA2, CCND1, CCNE1, CCNE2, CDH1 (E-cadherin), CDKN1B (p27Kip1), CDKN2A (p16INK4a), COL6A1, CTNNB1 (b-catenin), CTSB (cathepsin B), ERBB2 (Her-2), ESR1, ESR2, F3 (TF), FOSL1 (FRA-1), GATA3, GSN (Gelsolin), IGFBP2, IL2RA, IL6, IL6R, IL6ST (glycoprotein 130), ITGA6 (a6 integrin), JUN, KLK5, KRT19, MAP2K7 (c-Jun), MKI67 (Ki-67), NGFB (GF), NGFR, NME1 (M23A), PGR, PLAUI (uPA), PTEN, SERPINB5 (maspin), SERPINE1 (PAI-1), TGFA, THBS1

30

(thrombospondin-1), TIE (Tie-1), TNFRSF6 (Fas), TNFSF6 (FasL), TOP2A (topoisomerase IIA), TP53, AZGP1 (zinc-glycoprotein), BPAG1 (plectin), CDKN1A (p21Wap/Cip1), CLDN7 (claudin-7), CLU (clusterin), ERBB2 (Her-2), FGF1, FLRT1 (fibronectin), GABRP (GABAa), GNAS1, ID2, ITGA6 (a6 integrin), ITGB4 (b4 integrin), KLF5 (GC Box BP), KRT19 (Keratin 19), KRTHB6 (hair-specific type II keratin), MACMARCKS, MT3 (metallothionein-III), MUC1 (mucin), PTGS2 (COX-2), RAC2 (p21Rac2), S100A2, SCGB1D2 (lipophilin B), SCGB2A1 (mammaglobin 2), SCGB2A2 (mammaglobin 1), SPRR1B (Sprl), THBS1, THBS2, THBS4, and TNFAIP2 (B94), RON, c-Met, CD64, DLL4, PLGF, CTLA4, phosphatidylserine, ROBO4, CD80, CD22, CD40, CD23, CD28, CD80, CD55, CD38, CD70, CD74, CD30, CD138, CD56, CD33, CD2, CD137, DR4, DR5, RANKL, VEGFR2, PDGFR, VEGFR1, MTSP1, MSP, EPHB2, EPHA1, EPHA2, EpCAM, PGE2, NKG2D, LPA, SIP, APRIL, BCMA, MAPG, FLT3, PDGFR alpha, PDGFR beta, ROR1, PSMA, PSCA, SCD1, and CD59. To form the bispecific or trispecific antibodies of the invention, antibodies to any combination of these antigens can be made; that is, each of these antigens can be optionally and independently included or excluded from a multispecific antibody according to the present invention.

Monoclonal antibody therapy has become an important therapeutic modality for treating autoimmune and inflammatory disorders (Chan & Carter, 2010, *Nature Reviews Immunology* 10:301-316; Reichert et al., 2005, *Nature Biotechnology* 23[9]:1073-1078; herein expressly incorporated by reference). Many proteins have been implicated in general autoimmune and inflammatory responses, and thus may be targeted by the immunoglobulins of the invention. Autoimmune and inflammatory targets include but are not limited to C5, CCL1 (1-309), CCL11 (eotaxin), CCL13 (mcp-4), CCL15 (MIP-1d), CCL16 (HCC-4), CCL17 (TARC), CCL18 (PARC), CCL19, CCL2 (mcp-1), CCL20 (MIP-3a), CCL21 (MIP-2), CCL23 (MIP-1), CCL24 (MIP-2/eotaxin-2), CCL25 (TECK), CCL26, CCL3 (MIP-1a), CCL4 (MIP-1b), CCL5 (RANTES), CCL7 (mcp-3), CCL8 (mcp-2), CXCL1, CXCL10 (IP-10), CXCL11 (1-TAC/IP-9), CXCL12 (SDF1), CXCL13, CXCL14, CXCL2, CXCL3, CXCL5 (ENA-78/LIX), CXCL6 (GCP-2), CXCL8, IL13, IL8, CCL13 (mcp-4), CCR1, CCR2, CCR3, CCR4, CCR5, CCR6, CCR7, CCR8, CCR9, CX3CR1, IL8RA, XCR1 (CX3CR1), IFNA2, IL10, IL13, IL17C, IL1A, IL1B, IL1F10, IL1F5, IL1F6, IL1F7, IL1F8, IL1F9, IL22, IL5, IL8, IL9, LTA, LTB, MIF, SCYE1 (endothelial Monocyte-activating cytokine), SPP1, TNF, TNFSF5, IFNA2, IL10RA, IL10RB, IL13, IL13RA1, IL5RA, IL9, IL9R, ABCF1, BCL6, C3, C4A, CEBPB, CRP, ICEBERG, IL1R1, IL1RN, IL8RB, LTB4R, TOLLIP, FADD, IRAK1, IRAK2, MYD88, NCK2, TNFAIP3, TRADD, TRAF1, TRAF2, TRAF3, TRAF4, TRAF5, TRAF6, ACVR1, ACVR1B, ACVR2, ACVR2B, ACVRL1, CD28, CD3E, CD3G, CD3Z, CD69, CD80, CD86, CNR1, CTLA4, CYSLTR1, FCER1A, FCER2, FCGR3A, GPR44, HAVCR2, OPRD1, P2RX7, TLR2, TLR3, TLR4, TLR5, TLR6, TLR7, TLR8, TLR9, TLR10, BLR1, CCL1, CCL2, CCL3, CCL4, CCL5, CCL7, CCL8, CCL11, CCL13, CCL15, CCL16, CCL17, CCL18, CCL19, CCL20, CCL21, CCL22, CCL23, CCL24, CCL25, CCR1, CCR2, CCR3, CCR4, CCR5, CCR6, CCR7, CCR8, CCR9, CX3CL1, CX3CR1, CXCL1, CXCL2, CXCL3, CXCL5, CXCL6, CXCL10, CXCL11, CXCL12, CXCL13, CXCR4, GPR2, SCYE1, SDF2, XCL1, XCL2, XCR1, AMH, AMHR2, BMPR1A, BMPR1B, BMPR2, C19orf10 (IL27w), CER1, CSF1, CSF2, CSF3, DKFZp451J0118, FGF2, GFII, IFNA1, IFNB1, IFNG, IGF1, IL1A, IL1B,

31

IL1R1, IL1R2, IL2, IL2RA, IL2RB, IL2RG, IL3, IL4, IL4R, IL5, IL5RA, IL6, IL6R, IL6ST, IL7, IL8, IL8RA, IL8RB, IL9, IL9R, IL10, IL10RA, IL10RB, IL11, IL12RA, IL12A, IL12B, IL12RB1, IL12RB2, IL13, IL13RA1, IL13RA2, IL15, IL15RA, IL16, IL17, IL17R, IL18, IL18R1, IL19, IL20, KITLG, LEP, LTA, LTB, LTB4R, LTB4R2, LTBR, MIF, NPPB, PDGFB, TBX21, TDGF1, TGFA, TGFB1, TGFB1I1, TGFB2, TGFB3, TGFB1, TGFB1R1, TGFB2R2, TGFB3R3, TH1L, TNF, TNFRSF1A, TNFRSF1B, TNFRSF7, TNFRSF8, TNFRSF9, TNFRSF11A, TNFRSF21, TNFSF4, TNFSF5, TNFSF6, TNFSF11, VEGF, ZFPM2, and RNF110 (ZNF144). To form the bispecific or trispecific antibodies of the invention, antibodies to any combination of these antigens can be made; that is, each of these antigens can be optionally and independently included or excluded from a multispecific antibody according to the present invention.

Exemplary co-targets for autoimmune and inflammatory disorders include but are not limited to IL-1 and TNF α , IL-6 and TNF α , IL-6 and IL-1, IgE and IL-13, IL-1 and IL-13, IL-4 and IL-13, IL-5 and IL-13, IL-9 and IL-13, CD19 and Fc γ RIIb, and CD79 and Fc γ RIIb.

Immunoglobulins of the invention with specificity for the following pairs of targets to treat inflammatory disease are contemplated: TNF and IL-17A; TNF and RANKL; TNF and VEGF; TNF and SOST; TNF and DKK; TNF and α V β 3; TNF and NGF; TNF and IL-23p19; TNF and IL-6; TNF and SOST; TNF and IL-6R; TNF and CD-20; IgE and IL-13; IL-13 and IL23p19; IgE and IL-4; IgE and IL-9; IgE and IL-9; IgE and IL-13; IL-13 and IL-9; IL-13 and IL-4; IL-13 and IL-9; IL-13 and IL-9; IL-13 and IL-4; IL-13 and IL-23p19; IL-13 and IL-9; IL-6R and VEGF; IL-6R and IL-17A; IL-6R and RANKL; IL-17A and IL-1 β ; IL-1 β and RANKL; IL-1 β and VEGF; RANKL and CD-20; IL-l α and IL-1 β ; IL-l α and IL-1 β .

Pairs of targets that the immunoglobulins described herein can bind and be useful to treat asthma may be determined. In an embodiment, such targets include, but are not limited to, IL-13 and IL-1 β , since IL-1 β is also implicated in inflammatory response in asthma; IL-13 and cytokines and chemokines that are involved in inflammation, such as IL-13 and IL-9; IL-13 and IL-4; IL-13 and IL-5; IL-13 and IL-25; IL-13 and TARC; IL-13 and MDC; IL-13 and MIF; IL-13 and TGF- β ; IL-13 and LHR agonist; IL-13 and CCL25; IL-13 and SPRR2a; IL-13 and SPRR2b; and IL-13 and ADAMS. The immunoglobulins herein may have specificity for one or more targets involved in asthma selected from the group consisting of CSF1 (M-CSF), CSF2 (GM-CSF), CSF3 (G-CSF), FGF2, IFN α 1, IFN β 1, IFN γ , histamine and histamine receptors, IL1A, IL1B, IL2, IL3, IL4, IL5, IL6, IL7, IL8, IL9, IL10, IL11, IL12A, IL12B, IL13, IL14, IL15, IL16, IL17, IL18, IL19, KITLG, PDGFB, IL2RA, IL4R, IL5RA, IL8RA, IL8RB, IL12RB1, IL12RB2, IL13RA1, IL13RA2, IL18R1, TSLP, CCL1, CCL2, CCL3, CCL4, CCL5, CCL7, CCL8, CCL13, CCL17, CCL18, CCL19, CCL20, CCL22, CCL24, CX3CL1, CXCL1, CXCL2, CXCL3, XCL1, CCR2, CCR3, CCR4, CCR5, CCR6, CCR7, CCR8, CX3CR1, GPR2, XCR1, FOS, GATA3, JAK1, JAK3, STAT6, TBX21, TGFB1, TNF, TNFSF6, YY1, CYS-LTR1, FCER1A, FCER2, LTB4R, TB4R2, LTBR, and Chitinase. To form the bispecific or trispecific antibodies of the invention, antibodies to any combination of these antigens can be made; that is, each of these antigens can be optionally and independently included or excluded from a multispecific antibody according to the present invention.

Pairs of targets involved in rheumatoid arthritis (RA) may be co-targeted by the invention, including but not limited to

32

TNF and IL-18; TNF and IL-12; TNF and IL-23; TNF and IL-1 β ; TNF and MIF; TNF and IL-17; and TNF and IL-15.

Antigens that may be targeted in order to treat systemic lupus erythematosus (SLE) by the immunoglobulins herein include but are not limited to CD-20, CD-22, CD-19, CD28, CD4, CD80, HLA-DRA, IL10, IL2, IL4, TNFRSF5, TNFRSF6, TNFSF5, TNFSF6, BLR1, HDAC4, HDAC5, HDAC7A, HDAC9, ICOSL, IGBP1, MS4A1, RGS1, SLA2, CD81, IFN β 1, IL10, TNFRSF5, TNFRSF7, TNFSF5, AICDA, BLNK, GALNAC4S-6ST, HDAC4, HDAC5, HDAC7A, HDAC9, IL10, IL11, IL4, INHA, INHBA, KLF6, TNFRSF7, CD28, CD38, CD69, CD80, CD83, CD86, DPP4, FCER2, IL2RA, TNFRSF8, TNFSF7, CD24, CD37, CD40, CD72, CD74, CD79A, CD79B, CR2, ILIR2, ITGA2, ITGA3, MS4A1, ST6GALI, CDIC, CHST10, HLA-A, HLA-DRA, and NT5E.; CTLA4, B7.1, B7.2, BlyS, BAFF, C5, IL-4, IL-6, IL-10, IFN- α , and TNF- α . To form the bispecific or trispecific antibodies of the invention, antibodies to any combination of these antigens can be made; that is, each of these antigens can be optionally and independently included or excluded from a multispecific antibody according to the present invention.

The immunoglobulins herein may target antigens for the treatment of multiple sclerosis (MS), including but not limited to IL-12, TWEAK, IL-23, CXCL13, CD40, CD40L, IL-18, VEGF, VLA-4, TNF, CD45RB, CD200, IFN γ , GM-CSF, FGF, C5, CD52, and CCR2. An embodiment includes co-engagement of anti-IL-12 and TWEAK for the treatment of MS.

One aspect of the invention pertains to immunoglobulins capable of binding one or more targets involved in sepsis, in an embodiment two targets, selected from the group consisting of TNF, IL-1, MIF, IL-6, IL-8, IL-18, IL-12, IL-23, FasL, LPS, Toll-like receptors, TLR-4, tissue factor, MIP-2, ADORA2A, CASP1, CASP4, IL-10, IL-1B, NF κ B1, PROC, TNFRSF1A, CSF3, CCR3, ILIRN, MIF, NF κ B1, PTAFR, TLR2, TLR4, GPR44, HMOX1, midkine, IRAK1, NF κ B2, SERPINA1, SERPINE1, and TREM1. To form the bispecific or trispecific antibodies of the invention, antibodies to any combination of these antigens can be made; that is, each of these antigens can be optionally and independently included or excluded from a multispecific antibody according to the present invention.

In some cases, immunoglobulins herein may be directed against antigens for the treatment of infectious diseases.

The antibodies of the present invention are generally isolated or recombinant. "Isolated," when used to describe the various polypeptides disclosed herein, means a polypeptide that has been identified and separated and/or recovered from a cell or cell culture from which it was expressed. Ordinarily, an isolated polypeptide will be prepared by at least one purification step. An "isolated antibody," refers to an antibody which is substantially free of other antibodies having different antigenic specificities.

"Specific binding" or "specifically binds to" or is "specific for" a particular antigen or an epitope means binding that is measurably different from a non-specific interaction. Specific binding can be measured, for example, by determining binding of a molecule compared to binding of a control molecule, which generally is a molecule of similar structure that does not have binding activity. For example, specific binding can be determined by competition with a control molecule that is similar to the target.

Specific binding for a particular antigen or an epitope can be exhibited, for example, by an antibody having a KD for an antigen or epitope of at least about 10⁻⁴ M, at least about

33

10-5 M, at least about 10-6 M, at least about 10-7 M, at least about 10-8 M, at least about 10-9 M, alternatively at least about 10-10 M, at least about 10-11 M, at least about 10-12 M, or greater, where KD refers to a dissociation rate of a particular antibody-antigen interaction. Typically, an antibody that specifically binds an antigen will have a KD that is 20-, 50-, 100-, 500-, 1000-, 5,000-, 10,000- or more times greater for a control molecule relative to the antigen or epitope.

Also, specific binding for a particular antigen or an epitope can be exhibited, for example, by an antibody having a KA or Ka for an antigen or epitope of at least 20-, 50-, 100-, 500-, 1000-, 5,000-, 10,000- or more times greater for the epitope relative to a control, where KA or Ka refers to an association rate of a particular antibody-antigen interaction. Modified Antibodies

In addition to the modifications outlined above, other modifications can be made. For example, the molecules may be stabilized by the incorporation of disulphide bridges linking the VH and VL domains (Reiter et al., 1996, Nature Biotech. 14:1239-1245, entirely incorporated by reference). In addition, there are a variety of covalent modifications of antibodies that can be made as outlined below.

Covalent modifications of antibodies are included within the scope of this invention, and are generally, but not always, done post-translationally. For example, several types of covalent modifications of the antibody are introduced into the molecule by reacting specific amino acid residues of the antibody with an organic derivatizing agent that is capable of reacting with selected side chains or the N- or C-terminal residues.

Cysteiny residues most commonly are reacted with a-haloacetates (and corresponding amines), such as chloroacetic acid or chloroacetamide, to give carboxymethyl or carboxyamidomethyl derivatives. Cysteiny residues may also be derivatized by reaction with bromotrifluoroacetone, α -bromo- β -(5-imidazolyl)propionic acid, chloroacetyl phosphate, N-alkylmaleimides, 3-nitro-2-pyridyl disulfide, methyl 2-pyridyl disulfide, p-chloromercuribenzoate, 2-chloromercuri-4-nitrophenol, or chloro-7-nitrobenzo-2-oxa-1,3-diazole and the like.

In addition, modifications at cysteines are particularly useful in antibody-drug conjugate (ADC) applications, further described below. In some embodiments, the constant region of the antibodies can be engineered to contain one or more cysteines that are particularly "thiol reactive", so as to allow more specific and controlled placement of the drug moiety. See for example U.S. Pat. No. 7,521,541, incorporated by reference in its entirety herein.

Histidyl residues are derivatized by reaction with diethylpyrocarbonate at pH 5.5-7.0 because this agent is relatively specific for the histidyl side chain. Para-bromophenacyl bromide also is useful; the reaction is preferably performed in 0.1M sodium cacodylate at pH 6.0.

Lysiny residues and amino terminal residues are reacted with succinic or other carboxylic acid anhydrides. Derivatization with these agents has the effect of reversing the charge of the lysiny residues. Other suitable reagents for derivatizing alpha-amino-containing residues include imidoesters such as methyl picolinimide; pyridoxal phosphate; pyridoxal; chloroborohydride; trinitrobenzenesulfonic acid; O-methylisourea; 2,4-pentanedione; and transaminase-catalyzed reaction with glyoxylate.

Arginy residues are modified by reaction with one or several conventional reagents, among them phenylglyoxal, 2,3-butanedione, 1,2-cyclohexanedione, and ninhydrin. Derivatization of arginine residues requires that the reaction

34

be performed in alkaline conditions because of the high pKa of the guanidine functional group. Furthermore, these reagents may react with the groups of lysine as well as the arginine epsilon-amino group.

The specific modification of tyrosyl residues may be made, with particular interest in introducing spectral labels into tyrosyl residues by reaction with aromatic diazonium compounds or tetranitromethane. Most commonly, N-acetylimidazole and tetranitromethane are used to form O-acetyl tyrosyl species and 3-nitro derivatives, respectively. Tyrosyl residues are iodinated using 125I or 131I to prepare labeled proteins for use in radioimmunoassay, the chloramine T method described above being suitable.

Carboxyl side groups (aspartyl or glutamyl) are selectively modified by reaction with carbodiimides ($R'-N=C=N-R'$), where R and R' are optionally different alkyl groups, such as 1-cyclohexyl-3-(2-morpholinyl-4-ethyl) carbodiimide or 1-ethyl-3-(4-azonia-4,4-dimethylpentyl) carbodiimide. Furthermore, aspartyl and glutamyl residues are converted to asparaginy and glutaminy residues by reaction with ammonium ions.

Derivatization with bifunctional agents is useful for cross-linking antibodies to a water-insoluble support matrix or surface for use in a variety of methods, in addition to methods described below. Commonly used crosslinking agents include, e.g., 1,1-bis(diazoacetyl)-2-phenylethane, glutaraldehyde, N-hydroxysuccinimide esters, for example, esters with 4-azidosalicylic acid, homobifunctional imidoesters, including disuccinimidyl esters such as 3,3'-dithiobis (succinimidylpropionate), and bifunctional maleimides such as bis-N-maleimido-1,8-octane. Derivatizing agents such as methyl-3-[(p-azidophenyl)dithio]propionimide yield photoactivatable intermediates that are capable of forming crosslinks in the presence of light. Alternatively, reactive water-insoluble matrices such as cynomolgusogen bromide-activated carbohydrates and the reactive substrates described in U.S. Pat. Nos. 3,969,287; 3,691,016; 4,195,128; 4,247,642; 4,229,537; and 4,330,440, all entirely incorporated by reference, are employed for protein immobilization.

Glutaminy and asparaginy residues are frequently deamidated to the corresponding glutamyl and aspartyl residues, respectively. Alternatively, these residues are deamidated under mildly acidic conditions. Either form of these residues falls within the scope of this invention.

Other modifications include hydroxylation of proline and lysine, phosphorylation of hydroxyl groups of seryl or threony residues, methylation of the α -amino groups of lysine, arginine, and histidine side chains (T. E. Creighton, Proteins: Structure and Molecular Properties, W. H. Freeman & Co., San Francisco, pp. 79-86 [1983], entirely incorporated by reference), acetylation of the N-terminal amine, and amidation of any C-terminal carboxyl group.

In addition, as will be appreciated by those in the art, labels (including fluorescent, enzymatic, magnetic, radioactive, etc. can all be added to the antibodies (as well as the other compositions of the invention).

Glycosylation

Another type of covalent modification is alterations in glycosylation. In another embodiment, the antibodies disclosed herein can be modified to include one or more engineered glycoforms. By "engineered glycoform" as used herein is meant a carbohydrate composition that is covalently attached to the antibody, wherein said carbohydrate composition differs chemically from that of a parent antibody. Engineered glycoforms may be useful for a variety of purposes, including but not limited to enhancing or reducing

35

effector function. A preferred form of engineered glycoform is afucosylation, which has been shown to be correlated to an increase in ADCC function, presumably through tighter binding to the FcγRIIIa receptor. In this context, "afucosylation" means that the majority of the antibody produced in the host cells is substantially devoid of fucose, e.g. 90-95-98% of the generated antibodies do not have appreciable fucose as a component of the carbohydrate moiety of the antibody (generally attached at N297 in the Fc region). Defined functionally, afucosylated antibodies generally exhibit at least a 50% or higher affinity to the FcγRIIIa receptor.

Engineered glycoforms may be generated by a variety of methods known in the art (Umaria et al., 1999, *Nat Biotechnol* 17:176-180; Davies et al., 2001, *Biotechnol Bioeng* 74:288-294; Shields et al., 2002, *J Biol Chem* 277:26733-26740; Shinkawa et al., 2003, *J Biol Chem* 278:3466-3473; U.S. Pat. No. 6,602,684; U.S. Ser. No. 10/277,370; U.S. Ser. No. 10/113,929; PCT WO 00/61739A1; PCT WO 01/29246A1; PCT WO 02/31140A1; PCT WO 02/30954A1, all entirely incorporated by reference; (Potelligent® technology [Biowa, Inc., Princeton, N.J.]; GlycoMAb® glycosylation engineering technology [Glycart Biotechnology AG, Zürich, Switzerland]). Many of these techniques are based on controlling the level of fucosylated and/or bisecting oligosaccharides that are covalently attached to the Fc region, for example by expressing an IgG in various organisms or cell lines, engineered or otherwise (for example Lec-13 CHO cells or rat hybridoma YB2/0 cells, by regulating enzymes involved in the glycosylation pathway (for example FUT8 [α 1,6-fucosyltransferase] and/or β 1-4-N-acetylglucosaminyltransferase III [GnTIII]), or by modifying carbohydrate(s) after the IgG has been expressed. For example, the "sugar engineered antibody" or "SEA technology" of Seattle Genetics functions by adding modified saccharides that inhibit fucosylation during production; see for example 20090317869, hereby incorporated by reference in its entirety. Engineered glycoform typically refers to the different carbohydrate or oligosaccharide; thus an antibody can include an engineered glycoform.

Alternatively, engineered glycoform may refer to the IgG variant that comprises the different carbohydrate or oligosaccharide. As is known in the art, glycosylation patterns can depend on both the sequence of the protein (e.g., the presence or absence of particular glycosylation amino acid residues, discussed below), or the host cell or organism in which the protein is produced. Particular expression systems are discussed below.

Glycosylation of polypeptides is typically either N-linked or O-linked. N-linked refers to the attachment of the carbohydrate moiety to the side chain of an asparagine residue. The tri-peptide sequences asparagine-X-serine and asparagine-X-threonine, where X is any amino acid except proline, are the recognition sequences for enzymatic attachment of the carbohydrate moiety to the asparagine side chain. Thus, the presence of either of these tri-peptide sequences in a polypeptide creates a potential glycosylation site. O-linked glycosylation refers to the attachment of one of the sugars N-acetylgalactosamine, galactose, or xylose, to a hydroxyamino acid, most commonly serine or threonine, although 5-hydroxyproline or 5-hydroxylysine may also be used.

Addition of glycosylation sites to the antibody is conveniently accomplished by altering the amino acid sequence such that it contains one or more of the above-described tri-peptide sequences (for N-linked glycosylation sites). The alteration may also be made by the addition of, or substi-

36

tution by, one or more serine or threonine residues to the starting sequence (for O-linked glycosylation sites). For ease, the antibody amino acid sequence is preferably altered through changes at the DNA level, particularly by mutating the DNA encoding the target polypeptide at preselected bases such that codons are generated that will translate into the desired amino acids.

Another means of increasing the number of carbohydrate moieties on the antibody is by chemical or enzymatic coupling of glycosides to the protein. These procedures are advantageous in that they do not require production of the protein in a host cell that has glycosylation capabilities for N- and O-linked glycosylation. Depending on the coupling mode used, the sugar(s) may be attached to (a) arginine and histidine, (b) free carboxyl groups, (c) free sulfhydryl groups such as those of cysteine, (d) free hydroxyl groups such as those of serine, threonine, or hydroxyproline, (e) aromatic residues such as those of phenylalanine, tyrosine, or tryptophan, or (f) the amide group of glutamine. These methods are described in WO 87/05330 and in Aplin and Wriston, 1981, *CRC Crit. Rev. Biochem.*, pp. 259-306, both entirely incorporated by reference.

Removal of carbohydrate moieties present on the starting antibody (e.g. post-translationally) may be accomplished chemically or enzymatically. Chemical deglycosylation requires exposure of the protein to the compound trifluoromethanesulfonic acid, or an equivalent compound. This treatment results in the cleavage of most or all sugars except the linking sugar (N-acetylglucosamine or N-acetylgalactosamine), while leaving the polypeptide intact. Chemical deglycosylation is described by Hakimuddin et al., 1987, *Arch. Biochem. Biophys.* 259:52 and by Edge et al., 1981, *Anal. Biochem.* 118:131, both entirely incorporated by reference. Enzymatic cleavage of carbohydrate moieties on polypeptides can be achieved by the use of a variety of endo- and exo-glycosidases as described by Thotakura et al., 1987, *Meth. Enzymol.* 138:350, entirely incorporated by reference. Glycosylation at potential glycosylation sites may be prevented by the use of the compound tunicamycin as described by Duskin et al., 1982, *J. Biol. Chem.* 257:3105, entirely incorporated by reference. Tunicamycin blocks the formation of protein-N-glycoside linkages.

Another type of covalent modification of the antibody comprises linking the antibody to various nonproteinaceous polymers, including, but not limited to, various polyols such as polyethylene glycol, polypropylene glycol or polyoxyalkylenes, in the manner set forth in, for example, 2005-2006 PEG Catalog from Nektar Therapeutics (available at the Nektar website) U.S. Pat. Nos. 4,640,835; 4,496,689; 4,301,144; 4,670,417; 4,791,192 or 4,179,337, all entirely incorporated by reference. In addition, as is known in the art, amino acid substitutions may be made in various positions within the antibody to facilitate the addition of polymers such as PEG. See for example, U.S. Publication No. 2005/0114037A1, entirely incorporated by reference.

Additional Fc Variants for Additional Functionality

In addition to pI amino acid variants, there are a number of useful Fc amino acid modification that can be made for a variety of reasons, including, but not limited to, altering binding to one or more FcγR receptors, altered binding to FcRn receptors, etc.

Accordingly, the proteins of the invention can include amino acid modifications, including the heterodimerization variants outlined herein, which includes the pI variants FcγR Variants

Accordingly, there are a number of useful Fc substitutions that can be made to alter binding to one or more of the FcγR

receptors. Substitutions that result in increased binding as well as decreased binding can be useful. For example, it is known that increased binding to Fc γ RIIIa generally results in increased ADCC (antibody dependent cell-mediated cytotoxicity; the cell-mediated reaction wherein nonspecific cytotoxic cells that express Fc γ Rs recognize bound antibody on a target cell and subsequently cause lysis of the target cell). Similarly, decreased binding to Fc γ RIIb (an inhibitory receptor) can be beneficial as well in some circumstances. Amino acid substitutions that find use in the present invention include those listed in U.S. Ser. No. 11/124,620 (particularly FIG. 41), Ser. Nos. 11/174,287, 11/396,495, 11/538,406, all of which are expressly incorporated herein by reference in their entirety and specifically for the variants disclosed therein. Particular variants that find use include, but are not limited to, 236A, 239D, 239E, 332E, 332D, 239D/332E, 267D, 267E, 328F, 267E/328F, 236A/332E, 239D/332E/330Y, 239D, 332E/330L and 299T.

In addition, there are additional Fc substitutions that find use in increased binding to the FcRn receptor and increased serum half life, as specifically disclosed in U.S. Ser. No. 12/341,769, hereby incorporated by reference in its entirety, including, but not limited to, 434S, 428L, 308F, 259I, 428L/434S, 259I/308F, 436I/428L, 436I or V/434S, 436V/428L and 259I/308F/428L.

Linkers

The present invention optionally provides linkers as needed, for example in the addition of additional antigen binding sites, as depicted for example in FIG. 2A-2E, where "the other end" of the molecule contains additional antigen binding components. In addition, as outlined below, linkers are optionally also used in antibody drug conjugate (ADC) systems. When used to join the components of the central mAb-Fv constructs, the linker is generally a polypeptide comprising two or more amino acid residues joined by peptide bonds and are used to link one or more of the components of the present invention. Such linker polypeptides are well known in the art (see e.g., Holliger, P., et al. (1993) *Proc. Natl. Acad. Sci. USA* 90:6444-6448; Poljak, R. J., et al. (1994) *Structure* 2:1121-1123). A variety of linkers may find use in some embodiments described herein. As will be appreciated by those in the art, there are at least three different linker types used in the present invention.

"Linker" herein is also referred to as "linker sequence", "spacer", "tethering sequence" or grammatical equivalents thereof. Homo- or hetero-bifunctional linkers are well known (see, 1994 Pierce Chemical Company catalog, technical section on cross-linkers, pages 155-200, incorporated entirely by reference). A number of strategies may be used to covalently link molecules together. These include, but are not limited to polypeptide linkages between N- and C-termini of proteins or protein domains, linkage via disulfide bonds, and linkage via chemical cross-linking reagents. In one aspect of this embodiment, the linker is a peptide bond, generated by recombinant techniques or peptide synthesis. The linker peptide may predominantly include the following amino acid residues: Gly, Ser, Ala, or Thr. The linker peptide should have a length that is adequate to link two molecules in such a way that they assume the correct conformation relative to one another so that they retain the desired activity. In one embodiment, the linker is from about 1 to 50 amino acids in length, preferably about 1 to 30 amino acids in length. In one embodiment, linkers of 1 to 20 amino acids in length may be used. Useful linkers include glycine-serine polymers, including for example (GS)_n, (GSGGS)_n, (GGGGS)_n, and (GGGS)_n, where n is an integer of at least one, glycine-alanine polymers, alanine-serine polymers, and

other flexible linkers. Alternatively, a variety of nonproteinaceous polymers, including but not limited to polyethylene glycol (PEG), polypropylene glycol, polyoxyalkylenes, or copolymers of polyethylene glycol and polypropylene glycol, may find use as linkers, that is may find use as linkers.

Other linker sequences may include any sequence of any length of CL/CH1 domain but not all residues of CL/CH1 domain; for example the first 5-12 amino acid residues of the CL/CH1 domains. Linkers can be derived from immunoglobulin light chain, for example C κ or C λ . Linkers can be derived from immunoglobulin heavy chains of any isotype, including for example C γ 1, C γ 2, C γ 3, C γ 4, C α 1, C α 2, C δ , C ϵ , and C μ . Linker sequences may also be derived from other proteins such as Ig-like proteins (e.g. TCR, FcR, KIR), hinge region-derived sequences, and other natural sequences from other proteins.

Antibody-Drug Conjugates

In some embodiments, the multispecific antibodies of the invention are conjugated with drugs to form antibody-drug conjugates (ADCs). In general, ADCs are used in oncology applications, where the use of antibody-drug conjugates for the local delivery of cytotoxic or cytostatic agents allows for the targeted delivery of the drug moiety to tumors, which can allow higher efficacy, lower toxicity, etc. An overview of this technology is provided in Ducry et al., *Bioconjugate Chem.*, 21:5-13 (2010), Carter et al., *Cancer J.* 14(3):154 (2008) and Senter, *Current Opin. Chem. Biol.* 13:235-244 (2009), all of which are hereby incorporated by reference in their entirety.

Thus the invention provides multispecific antibodies conjugated to drugs. Generally, conjugation is done by covalent attachment to the antibody, as further described below, and generally relies on a linker, often a peptide linkage (which, as described below, may be designed to be sensitive to cleavage by proteases at the target site or not). In addition, as described above, linkage of the linker-drug unit (LU-D) can be done by attachment to cysteines within the antibody. As will be appreciated by those in the art, the number of drug moieties per antibody can change, depending on the conditions of the reaction, and can vary from 1:1 to 10:1 drug:antibody. As will be appreciated by those in the art, the actual number is an average.

Thus the invention provides multispecific antibodies conjugated to drugs. As described below, the drug of the ADC can be any number of agents, including but not limited to cytotoxic agents such as chemotherapeutic agents, growth inhibitory agents, toxins (for example, an enzymatically active toxin of bacterial, fungal, plant, or animal origin, or fragments thereof), or a radioactive isotope (that is, a radioconjugate) are provided. In other embodiments, the invention further provides methods of using the ADCs.

Drugs for use in the present invention include cytotoxic drugs, particularly those which are used for cancer therapy. Such drugs include, in general, DNA damaging agents, anti-metabolites, natural products and their analogs. Exemplary classes of cytotoxic agents include the enzyme inhibitors such as dihydrofolate reductase inhibitors, and thymidylate synthase inhibitors, DNA intercalators, DNA cleavers, topoisomerase inhibitors, the anthracycline family of drugs, the *vinca* drugs, the mitomycins, the bleomycins, the cytotoxic nucleosides, the pteridine family of drugs, diynes, the podophyllotoxins, dolastatins, maytansinoids, differentiation inducers, and taxols.

Members of these classes include, for example, methotrexate, methopterin, dichloromethotrexate, 5-fluorouracil, 6-mercaptopurine, cytosine arabinoside, melphalan, leurosine, leurosideine, actinomycin, daunorubicin, doxorubicin, mitomycin C, mitomycin A, caminomycin, aminopterin,

tallysomycin, podophyllotoxin and podophyllotoxin derivatives such as etoposide or etoposide phosphate, vinblastine, vincristine, vindesine, taxanes including taxol, taxotere retinoic acid, butyric acid, N8-acetyl spermidine, camptothecin, calicheamicin, esperamicin, ene-diyne, duocarmycin A, duocarmycin SA, calicheamicin, camptothecin, maytansinoids (including DM1), monomethylauristatin E (MMAE), monomethylauristatin F (MMAF), and maytansinoids (DM4) and their analogues.

Toxins may be used as antibody-toxin conjugates and include bacterial toxins such as diphtheria toxin, plant toxins such as ricin, small molecule toxins such as geldanamycin (Mandler et al (2000) J. Nat. Cancer Inst. 92(19):1573-1581; Mandler et al (2000) Bioorganic & Med. Chem. Letters 10:1025-1028; Mandler et al (2002) Bioconjugate Chem. 13:786-791), maytansinoids (EP 1391213; Liu et al., (1996) Proc. Natl. Acad. Sci. USA 93:8618-8623), and calicheamicin (Lode et al (1998) Cancer Res. 58:2928; Hinman et al (1993) Cancer Res. 53:3336-3342). Toxins may exert their cytotoxic and cytostatic effects by mechanisms including tubulin binding, DNA binding, or topoisomerase inhibition.

Conjugates of a multispecific antibody and one or more small molecule toxins, such as a maytansinoids, dolastatins, auristatins, a trichothecene, calicheamicin, and CC1065, and the derivatives of these toxins that have toxin activity, are contemplated.

Maytansinoids

Maytansine compounds suitable for use as maytansinoid drug moieties are well known in the art, and can be isolated from natural sources according to known methods, produced using genetic engineering techniques (see Yu et al (2002) PNAS 99:7968-7973), or maytansinol and maytansinol analogues prepared synthetically according to known methods. As described below, drugs may be modified by the incorporation of a functionally active group such as a thiol or amine group for conjugation to the antibody.

Exemplary maytansinoid drug moieties include those having a modified aromatic ring, such as: C-19-dechloro (U.S. Pat. No. 4,256,746) (prepared by lithium aluminum hydride reduction of ansamycin P2); C-20-hydroxy (or C-20-demethyl)+/-C-19-dechloro (U.S. Pat. Nos. 4,361,650 and 4,307,016) (prepared by demethylation using *Streptomyces* or *Actinomyces* or dechlorination using LAH); and C-20-demethoxy, C-20-acyloxy (—OCOR), +/-dechloro (U.S. Pat. No. 4,294,757) (prepared by acylation using acyl chlorides) and those having modifications at other positions.

Exemplary maytansinoid drug moieties also include those having modifications such as: C-9-SH (U.S. Pat. No. 4,424,219) (prepared by the reaction of maytansinol with H₂S or P₂S₅); C-14-alkoxymethyl(demethoxy/CH₂OR) (U.S. Pat. No. 4,331,598); C-14-hydroxymethyl or acyloxymethyl (CH₂OH or CH₂OAc) (U.S. Pat. No. 4,450,254) (prepared from *Nocardia*); C-15-hydroxy/acyloxy (U.S. Pat. No. 4,364,866) (prepared by the conversion of maytansinol by *Streptomyces*); C-15-methoxy (U.S. Pat. Nos. 4,313,946 and 4,315,929) (isolated from *Trewia nudiflora*); C-18-N-demethyl (U.S. Pat. Nos. 4,362,663 and 4,322,348) (prepared by the demethylation of maytansinol by *Streptomyces*); and 4,5-deoxy (U.S. Pat. No. 4,371,533) (prepared by the titanium trichloride/LAH reduction of maytansinol).

Of particular use are DM1 (disclosed in U.S. Pat. No. 5,208,020, incorporated by reference) and DM4 (disclosed in U.S. Pat. No. 7,276,497, incorporated by reference). See also a number of additional maytansinoid derivatives and methods in 5,416,064, WO/01/24763, 7,303,749, 7,601,354, U.S. Ser. No. 12/631,508, WO02/098883, 6,441,163, 7,368,

565, WO02/16368 and WO04/1033272, all of which are expressly incorporated by reference in their entirety.

ADCs containing maytansinoids, methods of making same, and their therapeutic use are disclosed, for example, in U.S. Pat. Nos. 5,208,020; 5,416,064; 6,441,163 and European Patent EP 0 425 235 B1, the disclosures of which are hereby expressly incorporated by reference. Liu et al., Proc. Natl. Acad. Sci. USA 93:8618-8623 (1996) described ADCs comprising a maytansinoid designated DM1 linked to the monoclonal antibody C242 directed against human colorectal cancer. The conjugate was found to be highly cytotoxic towards cultured colon cancer cells, and showed antitumor activity in an in vivo tumor growth assay.

Chari et al., Cancer Research 52:127-131 (1992) describe ADCs in which a maytansinoid was conjugated via a disulfide linker to the murine antibody A7 binding to an antigen on human colon cancer cell lines, or to another murine monoclonal antibody TA.1 that binds the HER-2/neu oncogene. The cytotoxicity of the TA.1-maytansinoid conjugate was tested in vitro on the human breast cancer cell line SK-BR-3, which expresses 3×10⁵ HER-2 surface antigens per cell. The drug conjugate achieved a degree of cytotoxicity similar to the free maytansinoid drug, which could be increased by increasing the number of maytansinoid molecules per antibody molecule. The A7-maytansinoid conjugate showed low systemic cytotoxicity in mice.

Auristatins and Dolastatins

In some embodiments, the ADC comprises a multispecific antibody conjugated to dolastatins or dolastatin peptidic analogs and derivatives, the auristatins (U.S. Pat. Nos. 5,635,483; 5,780,588). Dolastatins and auristatins have been shown to interfere with microtubule dynamics, GTP hydrolysis, and nuclear and cellular division (Woyke et al (2001) Antimicrob. Agents and Chemother. 45(12):3580-3584) and have anticancer (U.S. Pat. No. 5,663,149) and antifungal activity (Pettit et al (1998) Antimicrob. Agents Chemother. 42:2961-2965). The dolastatin or auristatin drug moiety may be attached to the antibody through the N (amino) terminus or the C (carboxyl) terminus of the peptidic drug moiety (WO 02/088172).

Exemplary auristatin embodiments include the N-terminus linked monomethylauristatin drug moieties DE and DF, disclosed in "Senter et al, Proceedings of the American Association for Cancer Research, Volume 45, Abstract Number 623, presented Mar. 28, 2004 and described in United States Patent Publication No. 2005/0238648, the disclosure of which is expressly incorporated by reference in its entirety.

An exemplary auristatin embodiment is MMAE (see U.S. Pat. No. 6,884,869 expressly incorporated by reference in its entirety).

Another exemplary auristatin embodiment is MMAF (see US 2005/0238649, 5,767,237 and 6,124,431, expressly incorporated by reference in their entirety).

Additional exemplary embodiments comprising MMAE or MMAF and various linker components (described further herein) have the following structures and abbreviations (wherein Ab means antibody and p is 1 to about 8):

Typically, peptide-based drug moieties can be prepared by forming a peptide bond between two or more amino acids and/or peptide fragments. Such peptide bonds can be prepared, for example, according to the liquid phase synthesis method (see E. Schroder and K. Lubke, "The Peptides", volume 1, pp 76-136, 1965, Academic Press) that is well known in the field of peptide chemistry. The auristatin/dolastatin drug moieties may be prepared according to the methods of: U.S. Pat. Nos. 5,635,483; 5,780,588; Pettit et al

41

(1989) J. Am. Chem. Soc. 111:5463-5465; Pettit et al (1998) Anti-Cancer Drug Design 13:243-277; Pettit, G. R., et al. Synthesis, 1996, 719-725; Pettit et al (1996) J. Chem. Soc. Perkin Trans. 1 5:859-863; and Doronina (2003) Nat Biotechnol 21(7):778-784. Calicheamicin

In other embodiments, the ADC comprises an antibody of the invention conjugated to one or more calicheamicin molecules. For example, Mylotarg is the first commercial ADC drug and utilizes calicheamicin γ 1 as the payload (see U.S. Pat. No. 4,970,198, incorporated by reference in its entirety). Additional calicheamicin derivatives are described in U.S. Pat. Nos. 5,264,586, 5,384,412, 5,550,246, 5,739,116, 5,773,001, 5,767,285 and 5,877,296, all expressly incorporated by reference. The calicheamicin family of antibiotics are capable of producing double-stranded DNA breaks at sub-picomolar concentrations. For the preparation of conjugates of the calicheamicin family, see U.S. Pat. Nos. 5,712,374, 5,714,586, 5,739,116, 5,767,285, 5,770,701, 5,770,710, 5,773,001, 5,877,296 (all to American Cyanamid Company). Structural analogues of calicheamicin which may be used include, but are not limited to, γ 1I, α 2I, α 2I, N-acetyl- γ 1I, PSAG and 01I (Hinman et al., Cancer Research 53:3336-3342 (1993), Lode et al., Cancer Research 58:2925-2928 (1998) and the aforementioned U.S. patents to American Cyanamid). Another anti-tumor drug that the antibody can be conjugated is QFA which is an antifolate. Both calicheamicin and QFA have intracellular sites of action and do not readily cross the plasma membrane. Therefore, cellular uptake of these agents through antibody mediated internalization greatly enhances their cytotoxic effects.

Duocarmycins

CC-1065 (see 4,169,888, incorporated by reference) and duocarmycins are members of a family of antitumor antibiotics utilized in ADCs. These antibiotics appear to work through sequence-selectively alkylating DNA at the N3 of adenine in the minor groove, which initiates a cascade of events that result in apoptosis.

Important members of the duocarmycins include duocarmycin A (U.S. Pat. No. 4,923,990, incorporated by reference) and duocarmycin SA (U.S. Pat. No. 5,101,038, incorporated by reference), and a large number of analogues as described in U.S. Pat. Nos. 7,517,903, 7,691,962, 5,101,038; 5,641,780; 5,187,186; 5,070,092; 5,070,092; 5,641,780; 5,101,038; 5,084,468, 5,475,092, 5,585,499, 5,846,545, WO2007/089149, WO2009/017394A1, 5,703,080, 6,989,452, 7,087,600, 7,129,261, 7,498,302, and 7,507,420, all of which are expressly incorporated by reference.

Other Cytotoxic Agents

Other antitumor agents that can be conjugated to the antibodies of the invention include BCNU, streptozocin, vincristine and 5-fluorouracil, the family of agents known collectively LL-E33288 complex described in U.S. Pat. Nos. 5,053,394, 5,770,710, as well as esperamicins (U.S. Pat. No. 5,877,296).

Enzymatically active toxins and fragments thereof which can be used include diphtheria A chain, nonbinding active fragments of diphtheria toxin, exotoxin A chain (from *Pseudomonas aeruginosa*), ricin A chain, abrin A chain, modeccin A chain, alpha-sarcin, *Aleurites fordii* proteins, dianthin proteins, *Phytolacca americana* proteins (PAPI, PAPII, and PAP-S), *momordica charantia* inhibitor, curcumin, crocin, *sapaonararia officinalis* inhibitor, gelonin, mitogellin, restrictocin, phenomycin, enomycin and the tricothecenes. See, for example, WO 93/21232 published Oct. 28, 1993.

The present invention further contemplates an ADC formed between an antibody and a compound with nucle-

42

olytic activity (e.g., a ribonuclease or a DNA endonuclease such as a deoxyribonuclease; DNase).

For selective destruction of the tumor, the antibody may comprise a highly radioactive atom. A variety of radioactive isotopes are available for the production of radioconjugated antibodies. Examples include At211, I131, I125, Y90, Re186, Re188, Sm153, Bi212, P32, Pb212 and radioactive isotopes of Lu.

The radio- or other labels may be incorporated in the conjugate in known ways. For example, the peptide may be biosynthesized or may be synthesized by chemical amino acid synthesis using suitable amino acid precursors involving, for example, fluorine-19 in place of hydrogen. Labels such as Tc99m or I123, Re186, Re188 and In111 can be attached via a cysteine residue in the peptide. Yttrium-90 can be attached via a lysine residue. The IODOGEN method (Fraker et al (1978) Biochem. Biophys. Res. Commun. 80: 49-57 can be used to incorporate Iodine-123. "Monoclonal Antibodies in Immunoscintigraphy" (Chatal, CRC Press 1989) describes other methods in detail.

For compositions comprising a plurality of antibodies, the drug loading is represented by p, the average number of drug molecules per Antibody. Drug loading may range from 1 to 20 drugs (D) per Antibody. The average number of drugs per antibody in preparation of conjugation reactions may be characterized by conventional means such as mass spectroscopy, ELISA assay, and HPLC. The quantitative distribution of Antibody-Drug-Conjugates in terms of p may also be determined.

In some instances, separation, purification, and characterization of homogeneous Antibody-Drug-conjugates where p is a certain value from Antibody-Drug-Conjugates with other drug loadings may be achieved by means such as reverse phase HPLC or electrophoresis. In exemplary embodiments, p is 2, 3, 4, 5, 6, 7, or 8 or a fraction thereof.

The generation of Antibody-drug conjugate compounds can be accomplished by any technique known to the skilled artisan. Briefly, the Antibody-drug conjugate compounds can include a multispecific antibody as the Antibody unit, a drug, and optionally a linker that joins the drug and the binding agent.

A number of different reactions are available for covalent attachment of drugs and/or linkers to binding agents. This is can be accomplished by reaction of the amino acid residues of the binding agent, for example, antibody molecule, including the amine groups of lysine, the free carboxylic acid groups of glutamic and aspartic acid, the sulfhydryl groups of cysteine and the various moieties of the aromatic amino acids. A commonly used non-specific methods of covalent attachment is the carbodiimide reaction to link a carboxy (or amino) group of a compound to amino (or carboxy) groups of the antibody. Additionally, bifunctional agents such as dialdehydes or imidoesters have been used to link the amino group of a compound to amino groups of an antibody molecule.

Also available for attachment of drugs to binding agents is the Schiff base reaction. This method involves the periodate oxidation of a drug that contains glycol or hydroxy groups, thus forming an aldehyde which is then reacted with the binding agent. Attachment occurs via formation of a Schiff base with amino groups of the binding agent. Isothiocyanates can also be used as coupling agents for covalently attaching drugs to binding agents. Other techniques are known to the skilled artisan and within the scope of the present invention.

In some embodiments, an intermediate, which is the precursor of the linker, is reacted with the drug under

appropriate conditions. In other embodiments, reactive groups are used on the drug and/or the intermediate. The product of the reaction between the drug and the intermediate, or the derivatized drug, is subsequently reacted with an multispecific antibody of the invention under appropriate conditions.

It will be understood that chemical modifications may also be made to the desired compound in order to make reactions of that compound more convenient for purposes of preparing conjugates of the invention. For example a functional group e.g. amine, hydroxyl, or sulfhydryl, may be appended to the drug at a position which has minimal or an acceptable effect on the activity or other properties of the drug.

Linker Units

Typically, the antibody-drug conjugate compounds comprise a Linker unit between the drug unit and the antibody unit. In some embodiments, the linker is cleavable under intracellular or extracellular conditions, such that cleavage of the linker releases the drug unit from the antibody in the appropriate environment. For example, solid tumors that secrete certain proteases may serve as the target of the cleavable linker; in other embodiments, it is the intracellular proteases that are utilized. In yet other embodiments, the linker unit is not cleavable and the drug is released, for example, by antibody degradation in lysosomes.

In some embodiments, the linker is cleavable by a cleaving agent that is present in the intracellular environment (for example, within a lysosome or endosome or caveolea). The linker can be, for example, a peptidyl linker that is cleaved by an intracellular peptidase or protease enzyme, including, but not limited to, a lysosomal or endosomal protease. In some embodiments, the peptidyl linker is at least two amino acids long or at least three amino acids long or more.

Cleaving agents can include, without limitation, cathepsins B and D and plasmin, all of which are known to hydrolyze dipeptide drug derivatives resulting in the release of active drug inside target cells (see, e.g., Dubowchik and Walker, 1999, *Pharm. Therapeutics* 83:67-123). Peptidyl linkers that are cleavable by enzymes that are present in CD38-expressing cells. For example, a peptidyl linker that is cleavable by the thiol-dependent protease cathepsin-B, which is highly expressed in cancerous tissue, can be used (e.g., a Phe-Leu or a Gly-Phe-Leu-Gly linker (SEQ ID NO: 460)). Other examples of such linkers are described, e.g., in U.S. Pat. No. 6,214,345, incorporated herein by reference in its entirety and for all purposes.

In some embodiments, the peptidyl linker cleavable by an intracellular protease is a Val-Cit linker or a Phe-Lys linker (see, e.g., U.S. Pat. No. 6,214,345, which describes the synthesis of doxorubicin with the val-cit linker).

In other embodiments, the cleavable linker is pH-sensitive, that is, sensitive to hydrolysis at certain pH values. Typically, the pH-sensitive linker hydrolyzable under acidic conditions. For example, an acid-labile linker that is hydrolyzable in the lysosome (for example, a hydrazone, semicarbazone, thiosemicarbazone, cis-aconitic amide, orthoester, acetal, ketal, or the like) may be used. (See, e.g., U.S. Pat. Nos. 5,122,368; 5,824,805; 5,622,929; Dubowchik and Walker, 1999, *Pharm. Therapeutics* 83:67-123; Neville et al., 1989, *Biol. Chem.* 264:14653-14661.) Such linkers are relatively stable under neutral pH conditions, such as those in the blood, but are unstable at below pH 5.5 or 5.0, the approximate pH of the lysosome. In certain embodiments, the hydrolyzable linker is a thioether linker (such as, e.g., a thioether attached to the therapeutic agent via an acylhydrazone bond (see, e.g., U.S. Pat. No. 5,622,929).

In yet other embodiments, the linker is cleavable under reducing conditions (for example, a disulfide linker). A variety of disulfide linkers are known in the art, including, for example, those that can be formed using SATA (N-succinimidyl-5-acetylthioacetate), SPDP (N-succinimidyl-3-(2-pyridyldithio)propionate), SPDB (N-succinimidyl-3-(2-pyridyldithio)butyrate) and SMPT (N-succinimidyl-oxy carbonyl- α -methyl- α -(2-pyridyl-dithio) toluene)-, SPDB and SMPT. (See, e.g., Thorpe et al., 1987, *Cancer Res.* 47:5924-5931; Wawrzynczak et al., In *Immunconjugates: Antibody Conjugates in Radioimaging and Therapy of Cancer* (C. W. Vogel ed., Oxford U. Press, 1987. See also U.S. Pat. No. 4,880,935.)

In other embodiments, the linker is a malonate linker (Johnson et al., 1995, *Anticancer Res.* 15:1387-93), a maleimidobenzoyl linker (Lau et al., 1995, *Bioorg-Med-Chem.* 3(10):1299-1304), or a 3'-N-amide analog (Lau et al., 1995, *Bioorg-Med-Chem.* 3(10):1305-12).

In yet other embodiments, the linker unit is not cleavable and the drug is released by antibody degradation. (See U.S. Publication No. 2005/0238649 incorporated by reference herein in its entirety and for all purposes).

In many embodiments, the linker is self-immolative. As used herein, the term "self-immolative Spacer" refers to a bifunctional chemical moiety that is capable of covalently linking together two spaced chemical moieties into a stable tripartite molecule. It will spontaneously separate from the second chemical moiety if its bond to the first moiety is cleaved. See for example, WO 2007059404A2, WO06110476A2, WO05112919A2, WO2010/062171, WO09/017394, WO07/089149, WO 07/018431, WO04/043493 and WO02/083180, which are directed to drug-cleavable substrate conjugates where the drug and cleavable substrate are optionally linked through a self-immolative linker and which are all expressly incorporated by reference.

Often the linker is not substantially sensitive to the extracellular environment. As used herein, "not substantially sensitive to the extracellular environment," in the context of a linker, means that no more than about 20%, 15%, 10%, 5%, 3%, or no more than about 1% of the linkers, in a sample of antibody-drug conjugate compound, are cleaved when the antibody-drug conjugate compound presents in an extracellular environment (for example, in plasma).

Whether a linker is not substantially sensitive to the extracellular environment can be determined, for example, by incubating with plasma the antibody-drug conjugate compound for a predetermined time period (for example, 2, 4, 8, 16, or 24 hours) and then quantitating the amount of free drug present in the plasma.

In other, non-mutually exclusive embodiments, the linker promotes cellular internalization. In certain embodiments, the linker promotes cellular internalization when conjugated to the therapeutic agent (that is, in the milieu of the linker-therapeutic agent moiety of the antibody-drug conjugate compound as described herein). In yet other embodiments, the linker promotes cellular internalization when conjugated to both the auristatin compound and the multispecific antibodies of the invention.

A variety of exemplary linkers that can be used with the present compositions and methods are described in WO 2004-010957, U.S. Publication No. 2006/0074008, U.S. Publication No. 20050238649, and U.S. Publication No. 2006/0024317 (each of which is incorporated by reference herein in its entirety and for all purposes).

Drug Loading

Drug loading is represented by p and is the average number of Drug moieties per antibody in a molecule. Drug

loading ("p") may be 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 or more moieties (D) per antibody, although frequently the average number is a fraction or a decimal. Generally, drug loading of from 1 to 4 is frequently useful, and from 1 to 2 is also useful. ADCs of the invention include collections of antibodies conjugated with a range of drug moieties, from 1 to 20. The average number of drug moieties per antibody in preparations of ADC from conjugation reactions may be characterized by conventional means such as mass spectroscopy and, ELISA assay.

The quantitative distribution of ADC in terms of p may also be determined. In some instances, separation, purification, and characterization of homogeneous ADC where p is a certain value from ADC with other drug loadings may be achieved by means such as electrophoresis.

For some antibody-drug conjugates, p may be limited by the number of attachment sites on the antibody. For example, where the attachment is a cysteine thiol, as in the exemplary embodiments above, an antibody may have only one or several cysteine thiol groups, or may have only one or several sufficiently reactive thiol groups through which a linker may be attached. In certain embodiments, higher drug loading, e.g. $p > 5$, may cause aggregation, insolubility, toxicity, or loss of cellular permeability of certain antibody-drug conjugates. In certain embodiments, the drug loading for an ADC of the invention ranges from 1 to about 8; from about 2 to about 6; from about 3 to about 5; from about 3 to about 4; from about 3.1 to about 3.9; from about 3.2 to about 3.8; from about 3.2 to about 3.7; from about 3.2 to about 3.6; from about 3.3 to about 3.8; or from about 3.3 to about 3.7. Indeed, it has been shown that for certain ADCs, the optimal ratio of drug moieties per antibody may be less than 8, and may be about 2 to about 5. See US 2005-0238649 A1 (herein incorporated by reference in its entirety).

In certain embodiments, fewer than the theoretical maximum of drug moieties are conjugated to an antibody during a conjugation reaction. An antibody may contain, for example, lysine residues that do not react with the drug-linker intermediate or linker reagent, as discussed below. Generally, antibodies do not contain many free and reactive cysteine thiol groups which may be linked to a drug moiety; indeed most cysteine thiol residues in antibodies exist as disulfide bridges. In certain embodiments, an antibody may be reduced with a reducing agent such as dithiothreitol (DTT) or tricarboylethylphosphine (TCEP), under partial or total reducing conditions, to generate reactive cysteine thiol groups. In certain embodiments, an antibody is subjected to denaturing conditions to reveal reactive nucleophilic groups such as lysine or cysteine.

The loading (drug/antibody ratio) of an ADC may be controlled in different ways, e.g., by: (i) limiting the molar excess of drug-linker intermediate or linker reagent relative to antibody, (ii) limiting the conjugation reaction time or temperature, (iii) partial or limiting reductive conditions for cysteine thiol modification, (iv) engineering by recombinant techniques the amino acid sequence of the antibody such that the number and position of cysteine residues is modified for control of the number and/or position of linker-drug attachments (such as thioMab or thioFab prepared as disclosed herein and in WO2006/034488 (herein incorporated by reference in its entirety)).

It is to be understood that where more than one nucleophilic group reacts with a drug-linker intermediate or linker reagent followed by drug moiety reagent, then the resulting product is a mixture of ADC compounds with a distribution of one or more drug moieties attached to an antibody. The average number of drugs per antibody may be calculated

from the mixture by a dual ELISA antibody assay, which is specific for antibody and specific for the drug. Individual ADC molecules may be identified in the mixture by mass spectroscopy and separated by HPLC, e.g. hydrophobic interaction chromatography.

In some embodiments, a homogeneous ADC with a single loading value may be isolated from the conjugation mixture by electrophoresis or chromatography.

Methods of Determining Cytotoxic Effect of ADCs

Methods of determining whether a Drug or Antibody-Drug conjugate exerts a cytostatic and/or cytotoxic effect on a cell are known. Generally, the cytotoxic or cytostatic activity of an Antibody Drug conjugate can be measured by: exposing mammalian cells expressing a target protein of the Antibody Drug conjugate in a cell culture medium; culturing the cells for a period from about 6 hours to about 5 days; and measuring cell viability. Cell-based in vitro assays can be used to measure viability (proliferation), cytotoxicity, and induction of apoptosis (caspase activation) of the Antibody Drug conjugate.

For determining whether an Antibody Drug conjugate exerts a cytostatic effect, a thymidine incorporation assay may be used. For example, cancer cells expressing a target antigen at a density of 5,000 cells/well of a 96-well plated can be cultured for a 72-hour period and exposed to 0.5 μ Ci of 3H-thymidine during the final 8 hours of the 72-hour period. The incorporation of 3H-thymidine into cells of the culture is measured in the presence and absence of the Antibody Drug conjugate.

For determining cytotoxicity, necrosis or apoptosis (programmed cell death) can be measured. Necrosis is typically accompanied by increased permeability of the plasma membrane; swelling of the cell, and rupture of the plasma membrane. Apoptosis is typically characterized by membrane blebbing, condensation of cytoplasm, and the activation of endogenous endonucleases. Determination of any of these effects on cancer cells indicates that an Antibody Drug conjugate is useful in the treatment of cancers.

Cell viability can be measured by determining in a cell the uptake of a dye such as neutral red, trypan blue, or ALA-MAR™ blue (see, e.g., Page et al., 1993, Intl. J. Oncology 3:473-476). In such an assay, the cells are incubated in media containing the dye, the cells are washed, and the remaining dye, reflecting cellular uptake of the dye, is measured spectrophotometrically. The protein-binding dye sulforhodamine B (SRB) can also be used to measure cytotoxicity (Skehan et al., 1990, J. Natl. Cancer Inst. 82:1107-12).

Alternatively, a tetrazolium salt, such as MTT, is used in a quantitative colorimetric assay for mammalian cell survival and proliferation by detecting living, but not dead, cells (see, e.g., Mosmann, 1983, J. Immunol. Methods 65:55-63).

Apoptosis can be quantitated by measuring, for example, DNA fragmentation. Commercial photometric methods for the quantitative in vitro determination of DNA fragmentation are available. Examples of such assays, including TUNEL (which detects incorporation of labeled nucleotides in fragmented DNA) and ELISA-based assays, are described in Biochemica, 1999, no. 2, pp. 34-37 (Roche Molecular Biochemicals).

Apoptosis can also be determined by measuring morphological changes in a cell. For example, as with necrosis, loss of plasma membrane integrity can be determined by measuring uptake of certain dyes (e.g., a fluorescent dye such as, for example, acridine orange or ethidium bromide). A method for measuring apoptotic cell number has been

described by Duke and Cohen, *Current Protocols in Immunology* (Coligan et al. eds., 1992, pp. 3.17.1-3.17.16). Cells also can be labeled with a DNA dye (e.g., acridine orange, ethidium bromide, or propidium iodide) and the cells observed for chromatin condensation and margination along the inner nuclear membrane. Other morphological changes that can be measured to determine apoptosis include, e.g., cytoplasmic condensation, increased membrane blebbing, and cellular shrinkage.

The presence of apoptotic cells can be measured in both the attached and "floating" compartments of the cultures. For example, both compartments can be collected by removing the supernatant, trypsinizing the attached cells, combining the preparations following a centrifugation wash step (e.g., 10 minutes at 2000 rpm), and detecting apoptosis (e.g., by measuring DNA fragmentation). (See, e.g., Piazza et al., 1995, *Cancer Research* 55:3110-16).

In vivo, the effect of a therapeutic composition of the multispecific antibody of the invention can be evaluated in a suitable animal model. For example, xenogenic cancer models can be used, wherein cancer explants or passaged xenograft tissues are introduced into immune compromised animals, such as nude or SCID mice (Klein et al., 1997, *Nature Medicine* 3: 402-408). Efficacy can be measured using assays that measure inhibition of tumor formation, tumor regression or metastasis, and the like.

The therapeutic compositions used in the practice of the foregoing methods can be formulated into pharmaceutical compositions comprising a carrier suitable for the desired delivery method. Suitable carriers include any material that when combined with the therapeutic composition retains the anti-tumor function of the therapeutic composition and is generally non-reactive with the patient's immune system. Examples include, but are not limited to, any of a number of standard pharmaceutical carriers such as sterile phosphate buffered saline solutions, bacteriostatic water, and the like (see, generally, *Remington's Pharmaceutical Sciences* 16th Edition, A. Osal., Ed., 1980).

Antibody Compositions for In Vivo Administration

Formulations of the antibodies used in accordance with the present invention are prepared for storage by mixing an antibody having the desired degree of purity with optional pharmaceutically acceptable carriers, excipients or stabilizers (*Remington's Pharmaceutical Sciences* 16th edition, Osal, A. Ed. [1980]), in the form of lyophilized formulations or aqueous solutions. Acceptable carriers, excipients, or stabilizers are nontoxic to recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate, and other organic acids; antioxidants including ascorbic acid and methionine; preservatives (such as octadecyldimethylbenzyl ammonium chloride; hexamethonium chloride; benzalkonium chloride, benzethonium chloride; phenol, butyl or benzyl alcohol; alkyl parabens such as methyl or propyl paraben; catechol; resorcinol; cyclohexanol; 3-pentanol; and m-cresol); low molecular weight (less than about 10 residues) polypeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, histidine, arginine, or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugars such as sucrose, mannitol, trehalose or sorbitol; salt-forming counter-ions such as sodium; metal complexes (e.g. Zn-protein complexes); and/or non-ionic surfactants such as TWEEN™, PLURONICS™ or polyethylene glycol (PEG).

The formulation herein may also contain more than one active compound as necessary for the particular indication being treated, preferably those with complementary activities that do not adversely affect each other. For example, it may be desirable to provide antibodies with other specificities. Alternatively, or in addition, the composition may comprise a cytotoxic agent, cytokine, growth inhibitory agent and/or small molecule antagonist. Such molecules are suitably present in combination in amounts that are effective for the purpose intended.

The active ingredients may also be entrapped in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization, for example, hydroxymethylcellulose or gelatin-microcapsules and poly-(methylmethacrylate) microcapsules, respectively, in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nano-particles and nanocapsules) or in macroemulsions. Such techniques are disclosed in *Remington's Pharmaceutical Sciences* 16th edition, Osal, A. Ed. (1980).

The formulations to be used for in vivo administration should be sterile, or nearly so. This is readily accomplished by filtration through sterile filtration membranes.

Sustained-release preparations may be prepared. Suitable examples of sustained-release preparations include semipermeable matrices of solid hydrophobic polymers containing the antibody, which matrices are in the form of shaped articles, e.g. films, or microcapsules. Examples of sustained-release matrices include polyesters, hydrogels (for example, poly(2-hydroxyethyl-methacrylate), or poly(vinylalcohol)), polylactides (U.S. Pat. No. 3,773,919), copolymers of L-glutamic acid and .gamma. ethyl-L-glutamate, non-degradable ethylene-vinyl acetate, degradable lactic acid-glycolic acid copolymers such as the LUPRON DEPOT™ (injectable microspheres composed of lactic acid-glycolic acid copolymer and leuprolide acetate), and poly-D-(-)-3-hydroxybutyric acid. While polymers such as ethylene-vinyl acetate and lactic acid-glycolic acid enable release of molecules for over 100 days, certain hydrogels release proteins for shorter time periods.

When encapsulated antibodies remain in the body for a long time, they may denature or aggregate as a result of exposure to moisture at 37° C., resulting in a loss of biological activity and possible changes in immunogenicity. Rational strategies can be devised for stabilization depending on the mechanism involved. For example, if the aggregation mechanism is discovered to be intermolecular S—S bond formation through thio-disulfide interchange, stabilization may be achieved by modifying sulfhydryl residues, lyophilizing from acidic solutions, controlling moisture content, using appropriate additives, and developing specific polymer matrix compositions.

Administrative Modalities

The antibodies and chemotherapeutic agents of the invention are administered to a subject, in accord with known methods, such as intravenous administration as a bolus or by continuous infusion over a period of time, by intramuscular, intraperitoneal, intracerebrospinal, subcutaneous, intra-articular, intrasynovial, intrathecal, oral, topical, or inhalation routes. Intravenous or subcutaneous administration of the antibody is preferred.

Treatment Modalities

In the methods of the invention, therapy is used to provide a positive therapeutic response with respect to a disease or condition. By "positive therapeutic response" is intended an improvement in the disease or condition, and/or an improvement in the symptoms associated with the disease or con-

dition. For example, a positive therapeutic response would refer to one or more of the following improvements in the disease: (1) a reduction in the number of neoplastic cells; (2) an increase in neoplastic cell death; (3) inhibition of neoplastic cell survival; (5) inhibition (i.e., slowing to some extent, preferably halting) of tumor growth; (6) an increased patient survival rate; and (7) some relief from one or more symptoms associated with the disease or condition.

Positive therapeutic responses in any given disease or condition can be determined by standardized response criteria specific to that disease or condition. Tumor response can be assessed for changes in tumor morphology (i.e., overall tumor burden, tumor size, and the like) using screening techniques such as magnetic resonance imaging (MRI) scan, x-radiographic imaging, computed tomographic (CT) scan, bone scan imaging, endoscopy, and tumor biopsy sampling including bone marrow aspiration (BMA) and counting of tumor cells in the circulation.

In addition to these positive therapeutic responses, the subject undergoing therapy may experience the beneficial effect of an improvement in the symptoms associated with the disease.

Thus for B cell tumors, the subject may experience a decrease in the so-called B symptoms, i.e., night sweats, fever, weight loss, and/or urticaria. For pre-malignant conditions, therapy with an multispecific therapeutic agent may block and/or prolong the time before development of a related malignant condition, for example, development of multiple myeloma in subjects suffering from monoclonal gammopathy of undetermined significance (MGUS).

An improvement in the disease may be characterized as a complete response. By "complete response" is intended an absence of clinically detectable disease with normalization of any previously abnormal radiographic studies, bone marrow, and cerebrospinal fluid (CSF) or abnormal monoclonal protein in the case of myeloma.

Such a response may persist for at least 4 to 8 weeks, or sometimes 6 to 8 weeks, following treatment according to the methods of the invention. Alternatively, an improvement in the disease may be categorized as being a partial response. By "partial response" is intended at least about a 50% decrease in all measurable tumor burden (i.e., the number of malignant cells present in the subject, or the measured bulk of tumor masses or the quantity of abnormal monoclonal protein) in the absence of new lesions, which may persist for 4 to 8 weeks, or 6 to 8 weeks.

Treatment according to the present invention includes a "therapeutically effective amount" of the medicaments used. A "therapeutically effective amount" refers to an amount effective, at dosages and for periods of time necessary, to achieve a desired therapeutic result.

A therapeutically effective amount may vary according to factors such as the disease state, age, sex, and weight of the individual, and the ability of the medicaments to elicit a desired response in the individual. A therapeutically effective amount is also one in which any toxic or detrimental effects of the antibody or antibody portion are outweighed by the therapeutically beneficial effects.

A "therapeutically effective amount" for tumor therapy may also be measured by its ability to stabilize the progression of disease. The ability of a compound to inhibit cancer may be evaluated in an animal model system predictive of efficacy in human tumors.

Alternatively, this property of a composition may be evaluated by examining the ability of the compound to inhibit cell growth or to induce apoptosis by *in vitro* assays known to the skilled practitioner. A therapeutically effective

amount of a therapeutic compound may decrease tumor size, or otherwise ameliorate symptoms in a subject. One of ordinary skill in the art would be able to determine such amounts based on such factors as the subject's size, the severity of the subject's symptoms, and the particular composition or route of administration selected.

Dosage regimens are adjusted to provide the optimum desired response (e.g., a therapeutic response). For example, a single bolus may be administered, several divided doses may be administered over time or the dose may be proportionally reduced or increased as indicated by the exigencies of the therapeutic situation. Parenteral compositions may be formulated in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the subjects to be treated; each unit contains a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier.

The specification for the dosage unit forms of the present invention are dictated by and directly dependent on (a) the unique characteristics of the active compound and the particular therapeutic effect to be achieved, and (b) the limitations inherent in the art of compounding such an active compound for the treatment of sensitivity in individuals.

The efficient dosages and the dosage regimens for the multispecific antibodies used in the present invention depend on the disease or condition to be treated and may be determined by the persons skilled in the art.

An exemplary, non-limiting range for a therapeutically effective amount of an multispecific antibody used in the present invention is about 0.1-100 mg/kg, such as about 0.1-50 mg/kg, for example about 0.1-20 mg/kg, such as about 0.1-10 mg/kg, for instance about 0.5, about such as 0.3, about 1, or about 3 mg/kg. In another embodiment, the antibody is administered in a dose of 1 mg/kg or more, such as a dose of from 1 to 20 mg/kg, e.g. a dose of from 5 to 20 mg/kg, e.g. a dose of 8 mg/kg.

A medical professional having ordinary skill in the art may readily determine and prescribe the effective amount of the pharmaceutical composition required. For example, a physician or a veterinarian could start doses of the medication employed in the pharmaceutical composition at levels lower than that required in order to achieve the desired therapeutic effect and gradually increase the dosage until the desired effect is achieved.

In one embodiment, the multispecific antibody is administered by infusion in a weekly dosage of from 10 to 500 mg/kg such as of from 200 to 400 mg/kg. Such administration may be repeated, e.g., 1 to 8 times, such as 3 to 5 times. The administration may be performed by continuous infusion over a period of from 2 to 24 hours, such as of from 2 to 12 hours.

In one embodiment, the multispecific antibody is administered by slow continuous infusion over a long period, such as more than 24 hours, if required to reduce side effects including toxicity.

In one embodiment the multispecific antibody is administered in a weekly dosage of from 250 mg to 2000 mg, such as for example 300 mg, 500 mg, 700 mg, 1000 mg, 1500 mg or 2000 mg, for up to 8 times, such as from 4 to 6 times. The administration may be performed by continuous infusion over a period of from 2 to 24 hours, such as of from 2 to 12 hours. Such regimen may be repeated one or more times as necessary, for example, after 6 months or 12 months. The dosage may be determined or adjusted by measuring the amount of compound of the present invention in the blood

51

upon administration by for instance taking out a biological sample and using anti-idiotypic antibodies which target the antigen binding region of the multispecific antibody.

In a further embodiment, the multispecific antibody is administered once weekly for 2 to 12 weeks, such as for 3 to 10 weeks, such as for 4 to 8 weeks.

In one embodiment, the multispecific antibody is administered by maintenance therapy, such as, e.g., once a week for a period of 6 months or more.

In one embodiment, the multispecific antibody is administered by a regimen including one infusion of an multispecific antibody followed by an infusion of an multispecific antibody conjugated to a radioisotope. The regimen may be repeated, e.g., 7 to 9 days later.

As non-limiting examples, treatment according to the present invention may be provided as a daily dosage of an antibody in an amount of about 0.1-100 mg/kg, such as 0.5, 0.9, 1.0, 1.1, 1.5, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 40, 45, 50, 60, 70, 80, 90 or 100 mg/kg, per day, on at least one of day 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, or 40, or alternatively, at least one of week 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20 after initiation of treatment, or any combination thereof, using single or divided doses of every 24, 12, 8, 6, 4, or 2 hours, or any combination thereof.

In some embodiments the multispecific antibody molecule thereof is used in combination with one or more additional therapeutic agents, e.g. a chemotherapeutic agent. Non-limiting examples of DNA damaging chemotherapeutic agents include topoisomerase I inhibitors (e.g., irinotecan, topotecan, camptothecin and analogs or metabolites thereof, and doxorubicin); topoisomerase II inhibitors (e.g., etoposide, teniposide, and daunorubicin); alkylating agents (e.g., melphalan, chlorambucil, busulfan, thiotepa, ifosfamide, carmustine, lomustine, semustine, streptozocin, decarbazine, methotrexate, mitomycin C, and cyclophosphamide); DNA intercalators (e.g., cisplatin, oxaliplatin, and carboplatin); DNA intercalators and free radical generators such as bleomycin; and nucleoside mimetics (e.g., 5-fluorouracil, capecitabine, gemcitabine, fludarabine, cytarabine, mercaptopurine, thioguanine, pentostatin, and hydroxyurea).

Chemotherapeutic agents that disrupt cell replication include: paclitaxel, docetaxel, and related analogs; vincristine, vinblastin, and related analogs; thalidomide, lenalidomide, and related analogs (e.g., CC-5013 and CC-4047); protein tyrosine kinase inhibitors (e.g., imatinib mesylate and gefitinib); proteasome inhibitors (e.g., bortezomib); NF- κ B inhibitors, including inhibitors of I κ B kinase; antibodies which bind to proteins overexpressed in cancers and thereby downregulate cell replication (e.g., trastuzumab, rituximab, cetuximab, and bevacizumab); and other inhibitors of proteins or enzymes known to be upregulated, over-expressed or activated in cancers, the inhibition of which downregulates cell replication.

In some embodiments, the antibodies of the invention can be used prior to, concurrent with, or after treatment with Velcade® (bortezomib).

All cited references are herein expressly incorporated by reference in their entirety.

Whereas particular embodiments of the invention have been described above for purposes of illustration, it will be appreciated by those skilled in the art that numerous varia-

52

tions of the details may be made without departing from the invention as described in the appended claims.

EXAMPLES

Examples are provided below to illustrate the present invention. These examples are not meant to constrain the present invention to any particular application or theory of operation. For all constant region positions discussed in the present invention, numbering is according to the EU index as in Kabat (Kabat et al., 1991, Sequences of Proteins of Immunological Interest, 5th Ed., United States Public Health Service, National Institutes of Health, Bethesda, entirely incorporated by reference). Those skilled in the art of antibodies will appreciate that this convention consists of nonsequential numbering in specific regions of an immunoglobulin sequence, enabling a normalized reference to conserved positions in immunoglobulin families. Accordingly, the positions of any given immunoglobulin as defined by the EU index will not necessarily correspond to its sequential sequence.

Example 1

Prototype "Triple F" Bispecific Antibody

The present invention describes novel immunoglobulin compositions that co-engage a first and second antigen. One heavy chain of the antibody contains a single chain Fv ("scFv", as defined below) and the other heavy chain is a "regular" Fab format, comprising a variable heavy chain and a light chain (see FIG. 1). This structure is sometimes referred to herein as "triple F" format (scFv-Fab-Fc). The two chains are brought together by the dimeric Fc region (see FIG. 2). The Fc region can be modified by amino acid substitution to allow for efficient purification of the "triple F" heterodimer. Further, the Fc region can be modified by amino acid substitution to promote the formation of the "triple F" heterodimer. Examples of Fc substitutions are described more fully below.

Fc substitutions can be included in the "triple F" format to allow for efficient purification of the desired "triple F" heterodimer over the undesired dual scFv-Fc and mAb homodimers. An example of this is in the inclusion of Fc substitutions that alter the isoelectric point (pI) of each monomer so that each monomer has a different pI. In this case the desired "triple F" heterodimer will have a different pI than that of the undesired dual scFv-Fc and mAb homodimers, thus facilitating isoelectric purification of the "triple F" heterodimer (e.g., anionic exchange columns, cationic exchange columns). These substitutions also aid in the determination and monitoring of any contaminating dual scFv-Fc and mAb homodimers post-purification (e.g., IEF gels, cIEF, and analytical IEX columns). See FIG. 3 for a list of substitutions that can be made in Fc monomer 1 and Fc monomer 2 to allow for efficient purification of the desired "triple F" heterodimer.

Fc substitutions can be included in the "triple F" format to "skew" the formation toward the desired "triple F" heterodimer over the undesired dual scFv-Fc and mAb homodimers. For example, see FIG. 4 for a list of substitutions that can be made in Fc monomer 1 and Fc monomer 2 to "skew" production toward the "triple F" heterodimer. Amino acid substitutions listed in FIG. 3 and FIG. 4 can be combined, leading to an increased yield of "triple F" heterodimer that can be easily purified away from any contaminating dual scFv-Fc and mAb homodimers.

53

After optimization of an scFv domain for inclusion in the “triple F” format, an optimized scFv domain can be coupled with a variety of standard antibody heavy chains in a convenient fashion. For example, an anti-CD3 scFv for recruiting T cell cytotoxicity can be coupled with a variety of anti-tumor antigen antibody heavy chains (e.g., those binding CD5, CD20, CD30, CD40, CD33, CD38, EGFR, EpCAM, Her2, HM1.24, or other tumor antigen). Further examples of optimized scFv domains that can be conveniently coupled with standard antibody heavy chains include anti-CD16 scFv for natural killer cell cytotoxicity; anti-CD32b scFv for inhibitory activity (here the coupled antibody heavy chain would bind, e.g., CD19, CD40, CD79a, CD79b, or other immune receptors); and anti-transferrin receptor scFv, anti-insulin receptor, or anti-LRP1 for transport across the blood-brain barrier.

Example 2

Multi-specific Antibodies Derived from the “Triple F” Format

Multi-specific antibodies can be constructed by attaching additional scFv or Fab domains that bind a third antigen to the C-terminus of one of the “triple F” heavy chains. See FIG. 5 for examples. Alternatively, the C-terminal scFv or Fab may bind the first or second antigen, thus conferring bivalency and an increase in overall binding affinity for that antigen.

Multi-specific antibodies can also be constructed by coupling the scFv-Fc heavy chain of the “triple F” format may with rearranged antibody heavy chains as depicted in FIG. 6. Such rearranged heavy chains may include an additional Fv region that binds a third antigen or an additional Fv region that binds the first antigen or second antigen, thus conferring bivalency and an increase in overall binding affinity for that antigen.

Example 3

Anti-CD19 Fab x Anti-CD3 scFv “Triple F” Bispecific

Amino acid sequences for anti-CD19 Fab x anti-CD3 scFv “triple F” bispecifics are listed in FIG. 7. Amino acid substitutions made to allow for efficient purification of the desired “triple F” heterodimer over the undesired dual scFv-Fc and mAb homodimers are underlined. Amino acid sequences for preferred humanized anti-CD3 variable regions are listed in FIG. 8 (with CDRs underlined). Some

54

examples of expression and purification of the desired “triple F” species and its bioactivity are given below.

The production of XENP11874, a “triple F” bispecific with an anti-CD19 Fab and anti-CD3 scFv, is outlined in FIG. 9. In FIG. 9A, the ion exchange purification of the desired “triple F” heterodimer from the undesired dual scFv-Fc and mAb homodimers is shown. The purity of the “triple F” fraction was checked by IEF gel, shown in FIG. 9B. Finally, SEC was used to confirm the homogenous size of the “triple F” product (see FIG. 9C).

XENP11874, anti-CD19 Fab x anti-CD3 scFv “triple F” bispecific, was shown to have potent bioactivity. The ability of XENP11874 to potentially recruit T cells for B cell depletion is shown in FIG. 10.

The production of XENP11924, a “triple F” bispecific with an anti-CD19 Fab and anti-CD3 scFv, is outlined in FIG. 11. In FIG. 11A, the ion exchange purification of the desired “triple F” heterodimer from the undesired dual scFv-Fc and mAb homodimers is shown. The purity of the “triple F” fraction was checked by IEF gel, shown in FIG. 11B. Finally, SEC was used to confirm the homogenous size of the “triple F” product (see FIG. 11C).

XENP11924, anti-CD19 Fab x anti-CD3 scFv “triple F” bispecific, was shown to have potent bioactivity. The ability of XENP11924 to potentially recruit T cells for the killing of the Raji tumor cell line is shown in FIG. 12.

Example 4

Anti-CD38 Fab x Anti-CD3 scFv “Triple F” Bispecific

Amino acid sequences for anti-CD38 Fab x anti-CD3 scFv “triple F” bispecifics are listed in FIG. 13. Amino acid substitutions made to allow for efficient purification of the desired “triple F” heterodimer over the undesired dual scFv-Fc and mAb homodimers are underlined. Some examples of expression and purification of the desired “triple F” species and its bioactivity are given below.

The production of XENP11925, a “triple F” bispecific with an anti-CD38 Fab and anti-CD3 scFv, is outlined in FIG. 14. In FIG. 14A, the ion exchange purification of the desired “triple F” heterodimer from the undesired dual scFv-Fc and mAb homodimers is shown. The purity of the “triple F” fraction was checked by IEF gel, shown in FIG. 14B. Finally, SEC was used to confirm the homogenous size of the “triple F” product (see FIG. 14C).

XENP11925, anti-CD38 Fab x anti-CD3 scFv “triple F” bispecific, was shown to have potent bioactivity. The ability of XENP11925 to potentially recruit T cells for the killing of the RPMI8226 tumor cell line is shown in FIG. 15.

SEQUENCE LISTING

The patent contains a lengthy “Sequence Listing” section. A copy of the “Sequence Listing” is available in electronic form from the USPTO web site (<http://seqdata.uspto.gov/?pageRequest=docDetail&DocID=US10472427B2>). An electronic copy of the “Sequence Listing” will also be available from the USPTO upon request and payment of the fee set forth in 37 CFR 1.19(b)(3).

We claim:

1. A heterodimeric protein comprising:

- a) a first monomer comprising a first variant Fc domain, and
- b) a second monomer comprising a second variant Fc domain,

wherein said first and second variant Fc domains comprise amino acid variant set L368D and S364K, according to the EU index as in Kabat.

2. The heterodimeric protein according to claim 1, wherein:

55

- a) said first monomer further comprises a single chain Fv (scFv) region that binds a first antigen;
 - b) said second monomer further comprises a first variable heavy chain;
- and wherein said heterodimeric protein further comprises: 5
- c) a third monomer comprising:
 - i) a first variable light chain; and
 - ii) a first constant light chain.
3. A heterodimeric protein comprising:
- a) a first monomer comprising: 10
 - i) a first variant Fc domain; and
 - ii) a single chain Fv (scFv) region that binds a first antigen; and
 - b) a second monomer comprising a VH-CH1-hinge-CH2-CH3 region, wherein said CH2-CH3 is a second variant Fc domain; and 15
 - c) third monomer comprising:
 - i) a variable light chain; and
 - ii) a constant light chain;
- wherein said first and second variant Fc domains comprise amino acid variant set L368D and S364K, according to the EU index as in Kabat. 20
4. The heterodimeric protein according to claim 1, wherein said first variant Fc domain comprises amino acid substitution L368D and said second variant Fc domain comprises amino acid substitution S364K. 25
5. The heterodimeric protein according to claim 1, wherein said first variant Fc domain comprises amino acid substitution S364K and said second variant Fc domain comprises amino acid substitution L368D. 30
6. The heterodimeric protein according to claim 1 or 2, wherein said first and second variant Fc domains are selected from the group consisting of human IgG1, IgG2 and IgG4 variant Fc domains.
7. A nucleic acid composition comprising: 35
- a) a first nucleic acid encoding the first monomer of claim 1; and

56

- b) a second nucleic acid encoding the second monomer of claim 1.
8. A nucleic acid composition comprising:
- a) a first nucleic acid encoding the first monomer of claim 2;
 - b) a second nucleic acid encoding the second monomer of claim 2; and
 - c) a third nucleic acid encoding the third monomer of claim 2.
9. A nucleic acid composition comprising:
- a) a first nucleic acid encoding the first heavy chain of claim 3;
 - b) a second nucleic acid encoding the second heavy chain of claim 3; and
 - c) a third nucleic acid encoding the third monomer of claim 3.
10. An expression vector comprising the nucleic acid composition of any one of claims 7 to 9.
11. A host cell comprising the expression vector of claim 10.
12. A method of making a heterodimeric protein according to any one of claims 1 to 3, the method comprising culturing a host cell according to claim 11 under conditions whereby said heterodimeric protein is produced.
13. A method of purifying a heterodimeric protein according to any one of claim 1, 2 or 3 comprising:
- a) providing a composition comprising the heterodimeric protein;
 - b) loading said composition onto an ion exchange column; and
 - c) collecting a fraction containing said heterodimeric protein.
14. A composition comprising a heterodimeric protein according to any one of claim 1, 2 or 3 and a pharmaceutical carrier.

* * * * *