



US 20210085608A1

(19) **United States**

(12) **Patent Application Publication** (10) **Pub. No.: US 2021/0085608 A1**
(43) **Pub. Date: Mar. 25, 2021**

(54) **LYOPHILIZED COMPOSITIONS OF
PHENOBARBITAL SODIUM SALT**

Publication Classification

(71) Applicant: **Nivagen Pharmaceuticals, Inc.,**
Sacramento, CA (US)

(51) **Int. Cl.**
A61K 9/19 (2006.01)
A61K 9/00 (2006.01)
A61K 31/515 (2006.01)

(72) Inventors: **Bala Tripura Sundari Chodavarapu,**
Davis, CA (US); **Thirupathi Mangali,**
Sacramento, CA (US); **Jay Shukla,**
Sacramento, CA (US); **Anand Shukla,**
Denver, CO (US); **Dasaradhi**
Lakkaraju, Sacramento, CA (US)

(52) **U.S. Cl.**
CPC *A61K 9/19* (2013.01); *A61K 31/515*
(2013.01); *A61K 9/0019* (2013.01)

(73) Assignee: **Nivagen Pharmaceuticals, Inc.,**
Sacramento, CA (US)

(57) **ABSTRACT**

(21) Appl. No.: **17/025,881**

(22) Filed: **Sep. 18, 2020**

Related U.S. Application Data

(60) Provisional application No. 62/903,511, filed on Sep.
20, 2019.

A composition of, method for producing, and use of an amorphous lyophilized Phenobarbital Sodium having high purity are presented. The amorphous lyophilized Phenobarbital Sodium is storage-stable being essentially void of impurities (e.g., phenylethylacetylurea (PEAU), 2-ethyl-2-phenylmalonamide (2EPMM), and/or alpha-phenylbutyrylguanidine (PBG)) upon reconstitution in water.



Fig. 1A




Fig. 1B




Fig. 1C




Fig. 1D




Fig. 1E




Fig. 1F




Fig. 2A






Fig. 2B




	Peak Name	RT	Area	s/n	USP Plate Count	USP Tailing
1	Phenobarbital	4.597	9769	1054.848570	7861.1	1.1

Fig. 2C




	PeakName	RT	Area	Amount	s/n	USP Plate Count	USP Tailing
1	Phenobarbital	4.597	38614	0.001	2906.846635	7746.3	1.2

Fig. 2D



	PeakName	RT	Area	% Area	sh	USP Plate Count	USP Tailing	Purity/ Angle	Purity/ Threshold
1	2EPMM	2.119	18787	100.00	1472	10573.6	1.2	0.674	1.833

Fig. 2E



	PeakName	RT	Area	%Area	sn	USP Plate Count	USP Tailing	Purity/ Angle	Purity/ Threshold
1	PBG	3.350	41776	100.00	2090	88019	15	21.648	62.982

Fig. 2F




Fig. 2G






Fig. 2H




	PeakName	RT	Area	Amount	s/n	USP Plate Count	USP Tailing
1	Phenobarbital	4.582	897147	0.476	50940.058185	7493.0	1.2

Fig. 3A




	Peak Name	RT	Area	Amount	USP Resolution	USP Plate Count	USP Tailing	Assay
1	Phenobarbital	10.130	2396458	60.823	17.9	17534.1	1.0	112.1

Fig. 3B




	Peak Name	RT	Area	% Area	Amount	Units	Assay	USP Resolution	USP Tailing
1	Benzyl Alcohol	3.391	87091	10.96	14.994	mg/mL	105.2	23.7	1.7
2	Blank	1.356	828	0.10	0.068	mg/mL			1.0
3	Phenobarbital	4.537	697184	87.71	57.200	mg/mL	96.3	7.2	1.5
4	RRT_1.82	8.319	9745	1.23	0.800	mg/mL		15.0	1.3

Fig. 3C



	Peak Name	RT	Area	% Area	Amount	Units	Assay	USP Resolution	USP Tailing
1	Benzyl Alcohol	3.392	84435	11.25	14.537	mg/mL	102.0	23.7	1.7
2	Blank	1.357	798	0.11	0.065	mg/mL			1.1
3	Phenobarbital	4.536	627538	83.64	51.486	mg/mL	86.7	7.2	1.5
4	RRT_1.82	8.321	37484	5.00	3.075	mg/mL		14.6	1.2


Fig. 3D



SampleName RLD_058407_65 mg/mL; Channel Description 2998 (205-400)nm; Result Id 3537


	PeakName	RT	Area	% Area	USP Resolution	sn	USP Plate Count	USP Tailing
1	Benzylalcohol	3.855	4623585	21.07	17.1	390.1	12312.3	1.5
2	Pheno_impurity-A	1.829	2803	0.01	5.2	0.3	7608.2	1.6
3	Phenobarbital	5.238	17188084	78.32	8.0	1018.9	11238.0	1.4
4	RRT_1.967_Phenol degradation product	9.781	92937	0.42	16.5	3.5	13191.8	1.2

Fig. 3E




	Peak Name	RT	Area	% Area	Amount	Units	USP Resolution	USP Tailing
1	Benzyl Alcohol	3.391	3972837	21.25	16.272	mg/mL	13.8	1.7
2	Phenobarbital	4.537	14515521	77.64	56.435	mg/mL	7.1	1.5
3	RRT_1.82	8.321	200110	1.07	0.778	mg/mL	14.6	1.3

Fig. 3F




	Peak Name	RT	Area	% Area	Amount	Units	USP Resolution	USP Tailing
1	Benzyl Alcohol	3.392	3833550	21.50	15.701	mg/mL	12.8	1.7
2	Phenobarbital	4.536	13183528	73.94	51.257	mg/mL	7.1	1.5
3	RRT_1.82	8.321	762558	4.28	2.965	mg/mL	14.6	1.3
4	RRT_1.949	9.000	42716	0.24	0.168	mg/mL	1.9	1.3

Fig. 3G




	PeakName	RT	Area	Amount	USP Resolution	USP Plate Count	USP Tailing	Assay
1	Phenobarbital	10.131	2358542	119.721	17.9	17563.1	1.0	110.3

Fig. 3H




	Peak Name	RT	Area	% Area	Amount	Units	Assay	USP Resolution	USP Tailing
1	Benzyl Alcohol	3.391	43232	5.84	14.886	mg/mL	104.5		1.7
2	Phenobarbital	4.535	888747	93.10	113.015	mg/mL	95.2	7.2	1.5
3	RRT_1.82	8.322	7837	1.06	1.286	mg/mL		14.5	1.2

Fig. 3I




	Peak Name	RT	Area	% Area	Amount	Units	Assay	USP Resolution	USP Tailing
1	Benzyl Alcohol	3.391	43440	5.94	14.958	mg/mL	105.0		1.7
2	Phenobarbital	4.537	857548	89.99	107.895	mg/mL	90.9	7.2	1.5
3	RRT_1.82	8.323	29742	4.07	4.880	mg/mL		14.6	1.2

Fig. 3J




	PeakName	RT	Area	% Area	USP Resolution	s/n	USP Plate Count	USP Tailing
1	Benzylalcohol	3853	2394223	11.87	20.8	2239	12438.0	15
2	Phenobarbital	5235	17661984	87.59	8.0	11444	11214.3	14
3	RRT_1.667_Phenodegrinproduct	9776	83640	0.41	16.6	35	13612.7	11

Fig. 3K




	Peak Name	RT	Area	% Area	Amount	Units	USP Resolution	USP Tailing
1	Benzyl Alcohol	3.391	1963213	11.91	16.081	mg/mL	13.4	1.7
2	Phenobarbital	4.535	14347574	87.06	111.565	mg/mL	7.1	1.5
3	RRT_1.82	8.320	162620	0.99	1.265	mg/mL	14.6	1.3

Fig. 3L



	Peak Name	RT	Area	% Area	Amount	Units	USP Resolution	USP Tailing
1	Benzyl Alcohol	3.391	1984788	12.10	16.258	mg/mL	12.8	1.7
2	Phenobarbital	4.536	13768702	83.93	107.064	mg/mL	7.1	1.5
3	RRT_1.82	8.322	594916	3.63	4.626	mg/mL	14.6	1.2
4	RRT_1.949	8.984	48220	0.29	0.375	mg/mL	1.8	1.3


Fig. 4A



SampleName #NPH1907_Lyo_65mg/mL_Initial; Channel Description W2489 ChB 254nm; Result Id 13991

	PeakName	RT	Area	Amount	Units	Assay	USP Resolution	s/n	USP Plate Count	USP Tailing
1	Phenobarbital	4.706	887698	58.858	mg/mL	99.1	19.3	52343.5	7952.5	1.2
2	Sodium	1.153	3626					318.4	943.9	1.1


Fig. 4B



SampleName NPH1907_65mg_RT_6M; Channel Description W2489 ChB 254nm; Result Id 31317

	Peak Name	RT	Area	Amount	Units	Assay	USP Resolution	s/n	USP Plate Count	USP Tailing
1	Phenobarbital	4.556	903872	59.388	mg/mL	100.0	20.8	44822.6	8704.0	1.1
2	Sodium/Blank	1.129	10057	0.661	mg/mL			616.7	1292.0	1.3


Fig. 4C



SampleName NPH1907_65mg_ACC_6M; Channel Description W2489 ChB 254nm; Result Id 31320

	PeakName	RT	Area	Amount	Units	Assay	USP Resolution	s/n	USP Plate Count	USP Tailing
1	Phenobarbital	4.552	896990	58.936	mg/mL	99.3	211	48295.1	8573.3	1.1
2	Sodium/Blank	1.122	5564	0.366	mg/mL			364.9	1275.7	1.3


Fig. 4D



SampleName #NPH1907_Lyo_65mg/mL_Initial; Channel Description W2489 ChA 210nm; Result Id 14043


	PeakName	RT	Area	% Area	Amount	Units	USP Resolution	s/n	USP Plate Count	USP Tailing
1	Phenobarbital	4.715	18143119	99.88	58.872	mg/mL	17.3	1873101.3	7804.1	1.2
2	RRT_0.37	1.688	7983	0.04	0.026	mg/mL	37	772.4	2802.6	1.2
3	Sodium	1.158	13483	0.07	0.044	mg/mL		1197.5	1009.2	1.2

Fig. 4E



	PeakName	RT	Area	% Area	Amount	Units	USP Resolution	s/n	USP Plate Count	USP Tailing
1	Blank2	1.364	4013	0.02			19	626.1	2241.7	1.1
2	PEAU	8.276	4512	0.02	0.013	mg/mL	14.8	278.0	12299.1	1.1
3	Phenobarbital	4.565	18206837	99.56	53.971	mg/mL	15.2	1616675.4	8239.9	1.1
4	RRT_0.37	1.644	12808	0.07	0.038	mg/mL	19	1270.5	1640.7	1.1
5	SodiumBlank	1.135	58253	0.32	0.173	mg/mL		8406.1	1363.0	1.1


Fig. 4F



SampleName NPH1907_65mg_ACC_6M; ChannelDescription W2489 ChA 210nm; ResultId 31632

	Peak Name	RT	Area	% Area	Amount	Units	USP Resolution	s/n	USP Plate Count	USP Tailing
1	Blank2	1.361	5648	0.03			1.9	899.0	2143.8	1.0
2	PEAU	8.275	6741	0.04	0.020	mg/mL	14.6	427.7	11918.8	1.0
3	Phenobarbital	4.561	18093753	99.70	53.638	mg/mL	15.2	1838135.6	8445.9	1.1
4	RRT_0.37	1.638	10466	0.06	0.031	mg/mL	1.9	1064.2	1482.5	1.1
5	SodiumBlank	1.131	32159	0.18	0.095	mg/mL		4942.0	1362.2	1.1


Fig. 4G



SampleName #NPH1912_Lyo_130mg/mL_Initial; Channel Description W2489 ChB 254nm; Result Id 13996

	Peak Name	RT	Area	Amount	Units	Assay	USP Resolution	s/n	USP Plate Count	USP Tailing
1	Phenobarbital	4.704	880753	116.796	mg/mL	98.3	19.3	40192.1	7866.8	1.2
2	Sodium	1.151	2171					148.3	959.0	1.0


Fig. 4H



SampleName NPH1912_130mg_RT_6M; Channel Description W2489 ChB 254nm; Result Id 31325

	PeakName	RT	Area	Amount	Units	Assay	USP Resolution	sn	USP Plate Count	USP Tailing
1	Phenobarbital	4.564	905459	118.985	mg/mL	100.2	20.7	55858.2	8639.9	1.1
2	Sodium/Blank	1.133	4686	0.616	mg/mL			358.4	1316.2	1.4


Fig. 4I



SampleName NPH1912_130mg_ACC_6M ; Channel Description W2489 ChB 254nm; Result Id 31328

	Peak Name	RT	Area	Amount	Units	Assay	s/n	USP Plate Count	USP Tailing
1	Phenobarbital	4.562	8698191	118.000	mg/mL	99.4	55375.6	8698.8	1.1


Fig. 4J



SampleName #NPH1912_Lyo_130mg/mL_Initial, Channel Description W2489 ChA 210nm, Result Id 14048

	PeakName	RT	Area	% Area	Amount	Units	USP Resolution	s/n	USP Plate Count	USP Tailing
1	Phenobarbital	4.713	18027682	99.93	116.995	mg/mL	17.0	1001026.6	7649.3	1.2
2	RRT_0.37	1.682	4819	0.03	0.031	mg/mL	36	433.7	2401.3	1.2
3	Sodium	1.156	7967	0.04	0.052	mg/mL		649.1	941.5	1.1


Fig. 4K



— SampleName NPH1912_130mg_RT_6M; Channel Description W2489 ChA 210nm; Result Id 31637

	PeakName	RT	Area	% Area	Amount	Units	USP Resolution	s/n	USP Plate Count	USP Tailing
1	Blank2	1.386	1069	0.01			19	143.5	2247.2	1.1
2	PEAU	8.291	4082	0.02	0.024	mg/mL	14.8	213.9	12463.8	1.0
3	Phenobarbital	4.572	18248178	99.78	106.887	mg/mL	15.9	1373532.2	8426.5	1.1
4	RRT_0.37	1.650	9642	0.05	0.057	mg/mL	20	832.7	1618.4	1.1
5	SodiumBlank	1.139	28215	0.15	0.167	mg/mL		3438.2	1382.8	1.1

Fig. 4L



SampleName NPH1912_130mg_ACC_6M; Channel Description W2489 ChA 210nm; Result Id 31640

	Peak Name	RT	Area	% Area	Amount	Units	USP Resolution	s/n	USP Plate Count	USP Tailing
1	Blank2	1.365	5481	0.03			1.9	687.2	2149.1	1.1
2	PEAU	8.292	5068	0.03	0.030	mg/mL	14.6	256.0	12320.6	1.0
3	Phenobarbital	4.570	18122566	99.73	107.442	mg/mL	15.1	132658.2	6330.4	1.1
4	RRT_0.37	1.646	9209	0.05	0.065	mg/mL	1.8	725.3	1481.0	1.1
5	SodiumBlank	1.135	28641	0.16	0.171	mg/mL		3440.5	1362.6	1.1

Fig. 5A




Fig. 5B




Fig. 5C





Fig. 5D



LYOPHILIZED COMPOSITIONS OF PHENOBARBITAL SODIUM SALT

RELATED APPLICATIONS

[0001] This application claims the benefit of U.S. Provisional Patent Application No. 62/903,511, filed on Sep. 20, 2019, which is incorporated by reference in its entirety.

FIELD OF THE INVENTION

[0002] The present disclosure relates to compositions of storage-stable lyophilized Phenobarbital Sodium having high purity that are essentially void of impurities upon reconstitution in water.

BACKGROUND OF THE INVENTION

[0003] The background description includes information that may be useful in understanding the present disclosure. It is not an admission that any of the information provided herein is prior art or relevant to the presently claimed invention, or that any publication specifically or implicitly referenced is prior art.

[0004] All publications and patent applications herein are incorporated by reference to the same extent as if each individual publication or patent application were specifically and individually indicated to be incorporated by reference. Where a definition or use of a term in an incorporated reference is inconsistent or contrary to the definition of that term provided herein, the definition of that term provided herein applies and the definition of that term in the reference does not apply.

[0005] Phenobarbital Sodium is a well-known barbiturate that is most commonly administered as an anti-convulsant for the treatment of seizures and may also be administered as a sedative, a sleep aide, or a pre-anesthetic. Phenobarbital Sodium is typically a liquid formulation administered intramuscularly (IM) or intravenously (IV) with crystalline granules or crystals dissolved in an aqueous solvent. For example, WO 2017/085687 to Parker et al. discloses a liquid formulation of Phenobarbital Sodium dissolved in water along with a C1-C4 alcohol and a glycol.

[0006] The stability of solubilized Phenobarbital Sodium has been a recurring problem because it readily hydrolyzes in water forming various degradation products. In particular, in water alone or in water with an increase in pH, Phenobarbital Sodium forms the degradation products phenylethylacetylurea (PEAU), 2-ethyl-2-phenylmalonamide (2EPM), and alpha-phenylbutyrylguanidine (PBG). While these degradation impurities may not directly compromise the effectiveness of the Phenobarbital Sodium for its therapeutic activity, these degradation impurities are toxic and cause severe side effects. Specifically, PEAU was previously administered as Pheneturide for severely affected epileptics. However, the Food and Drug Administration (FDA) withdrew approval of Pheneturide due to its toxic effects. In particular, Pheneturide is reported to be a potent liver enzyme inducer that disrupts both calcium and folate metabolism. (Nayyar et al., 2012, *J. of Appl. Pharm Sci.*, 2:230-235.) Prior to its removal as an FDA-approved drug, patients treated with Pheneturide showed low levels of serum calcium and folate.

[0007] Over the years of medical administration, efforts have been made to decrease the formation of the toxic degradation products in the Phenobarbital Sodium liquid

formulation. For example, Gupta, V. D., (*J. Pharm Sci.*, 1984, 73:1661-1662) describes the effects of ethanol, glycerol, and propylene glycol on the stability of Phenobarbital Sodium in aqueous solutions. Specifically, Gupta discloses that ethanol has the best effect followed by propylene glycol and glycerol relative to water alone. Years after Gupta, Parker et al., (WO 2017/085687) focused on minimizing the total amount of water to reduce the hydrolysis and consequently the formation of the degradation products. Parker addresses the hydrolysis issue by limiting the amount of water (e.g., to no more than 50 mg/mL) in the liquid formulation. Parker's minimized water formulations also include a C1-C4 alcohol such as ethanol and a glycol such as propylene glycol. However, while the amount of impurities in Parker's liquid formulation may have decreased compared to prior formulations, impurities such as PEAU are still present in undesirable amounts, particularly after extended storage.

[0008] In addition to the degradation impurities, the current marketed liquid formulation of Phenobarbital Sodium (e.g., Phenobarbital Sodium for Injection USP, 65 mg/mL or 130 mg/mL), includes organic solvents such as alcohol, which are not desirable in general and indeed not suitable for administration to newborns as described in the art, e.g., Williams et al., 1998, "Evaluating Toxic Alcohol Poisoning in the Emergency Setting," *Laboratory Medicine*, Vol. 29, No. 2; Wood et al., 2007, *J. of Perinatology*, 27:183-185; Kumar, 1985, "Adverse Drug Reactions in the Newborn," *Annals of Clin. And Lab. Sci.*, Vol. 15, No. 3; and Cuzzolin, 2018, *J. Pediatric and Neonatal Individualized Med.*, doi: 10.7363/070112.

[0009] Accordingly, there is still a need for a stable Phenobarbital Sodium formulation in a formulation that is storage stable and void or essentially void of degradation impurities upon solubilization in water.

SUMMARY

[0010] Disclosed herein are compositions of, methods of producing, and use of a lyophilized Phenobarbital Sodium formulation. The contemplated composition is an amorphous lyophilized Phenobarbital Sodium with high stability and purity. The lyophilized composition may include no less than 98% Phenobarbital Sodium. The amorphous composition is a storage-stable form of lyophilized Phenobarbital Sodium.

[0011] The lyophilized amorphous formulation that was produced resulted in surprising, unexpected and desirable results. The quantity of undesirable and/or toxic degradation impurities that are produced from this formulation are less than those in current commercial products. And the stability of this formulation is superior to that of current commercial products. These results are particularly surprising and unexpected in view of the fact that conventional wisdom teaches that crystalline material is more stable than amorphous material (see e.g., C. Ahlneck and G. Zografi "The Molecular Basis for Moisture Effects on the Physical and Chemical Stability of Drugs in the Solid-State," *Int. J. Pharm.* 1990, 62, 87-95). Thus, one of ordinary skill in the art would want to avoid using amorphous material in pharmaceutical formulations.

[0012] Preferably the percentage of the lyophilized formulation that is amorphous is at least 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95%. Alternatively, the percentage of the lyo-

philized formulation that is amorphous can be 100%. In addition, the percentage of the lyophilized formulation that is amorphous can be within any range between (and including) 20% to 100% (e.g., 20%-100%, 25%-95%, etc.).

[0013] Methods for producing the storage-stable form of lyophilized Phenobarbital Sodium composition result in a lyophilized powder in which the impurity phenylethylacetylurea (PEAU) is below detection limit and is never more than 0.2% of the specification limit even after storage of the lyophilized Phenobarbital Sodium for up to 6 months under accelerated stability conditions (40° C., 75% RH). In some embodiments, the method for producing the storage-stable lyophilized Phenobarbital Sodium includes adding Phenobarbital Sodium to water to form a Phenobarbital Sodium solution, having a pH between 9.2 to 10.2, and lyophilizing the bulk solution. In some embodiments, the water is Water for Injection.

[0014] In preferred embodiments, the Water for Injection is sparged with nitrogen prior to adding the Phenobarbital Sodium. Typically, the Water for Injection is sparged with nitrogen for at least 30 minutes prior to adding the Phenobarbital Sodium. In additional embodiments, the water (e.g., Water for Injection) is cooled to or between 2 to 8° C. prior to adding the Phenobarbital Sodium.

[0015] For adjusting the pH of the Phenobarbital Sodium solution, the method for producing the storage-stable form of lyophilized Phenobarbital Sodium composition may also include measuring the pH of the Phenobarbital Sodium solution, wherein if the pH is higher than 10.2, the adjusting includes adding hydrochloric acid (HCl).

[0016] Typically, the resulting lyophilized Phenobarbital Sodium forms no more than 0.1% PEAU when reconstituted in an aqueous solution after storage of the lyophilized Phenobarbital Sodium for up to 6 months both at RT and accelerated stability conditions. More typically, the resulting lyophilized Phenobarbital Sodium forms no more than 0.05% PEAU when reconstituted in an aqueous solution after storage of the lyophilized Phenobarbital Sodium for up to 6 months at room temperature (RT) and under accelerated stability conditions (40° C.). Additionally, or alternatively, the resulting lyophilized Phenobarbital Sodium forms either an amount below the quantitative limit (BQL, 0.05%) or no detectable amount of PEAU, 2-ethyl-2-phenylmalonamide (2EPMM), or alpha-phenylbutyrylguanidine (PBG) when reconstituted in the aqueous solution after storage of the lyophilized Phenobarbital Sodium for up to 6 months at RT and accelerated stability conditions.

[0017] According to contemplated embodiments as disclosed herein, the amorphous lyophilized Phenobarbital Sodium composition is stable up to at least 6 months under accelerated stability conditions which is equivalent to 24 months at RT. In this respect, in some embodiments, the stability of the lyophilized Phenobarbital Sodium is determined by reconstituting the lyophilized Phenobarbital Sodium in an aqueous solution and measuring the amount of or the presence of phenylethylacetylurea (PEAU), 2-ethyl-2-phenylmalonamide (2EPMM), and/or alpha-phenylbutyrylguanidine (PBG).

[0018] Additional embodiments include the method for producing the storage-stable form of lyophilized Phenobarbital Sodium composition as disclosed in which the Phenobarbital Sodium solution is filtered prior to lyophilization.

[0019] Further embodiments include the method for producing the storage-stable form of lyophilized Phenobarbital

Sodium composition as disclosed in which the Phenobarbital Sodium solution is aliquoted into a vial prior to lyophilization, and after lyophilization, a closure is applied to the vial under vacuum.

[0020] Aspects of the present disclosure also include a method of reducing formation of degradation products of Phenobarbital Sodium reconstituted in an aqueous solution, in which the method includes adding Phenobarbital Sodium to water to form a Phenobarbital Sodium solution, adjusting the pH of the Phenobarbital Sodium solution to or between 9.2 to 10.2, and lyophilizing the Phenobarbital Sodium solution to form stable white lyophilized powder of Phenobarbital Sodium.

[0021] In preferred embodiments, the Water for Injection is sparged with nitrogen prior to adding the Phenobarbital Sodium. Typically, the Water for Injection is sparged with nitrogen for at least 30 minutes prior to adding the Phenobarbital Sodium. In additional embodiments, the water (e.g., Water for Injection) is cooled to or between 2 to 8° C. prior to adding the Phenobarbital Sodium.

[0022] For adjusting the pH of the Phenobarbital Sodium solution, the method of reducing formation of degradation products of Phenobarbital Sodium reconstituted in an aqueous solution may also include measuring the pH of the Phenobarbital Sodium solution, wherein if the pH is higher than 10.2, the adjusting includes adding Hydrochloric acid (HCl).

[0023] In typical embodiments, the degradation products of Phenobarbital Sodium reconstituted in an aqueous solution include phenylethylacetylurea (PEAU), 2-ethyl-2-phenylmalonamide (2EPMM), and/or alpha-phenylbutyrylguanidine (PBG), and analysis of the formation of the degradation products includes reconstituting the stable lyophilized Phenobarbital Sodium in an aqueous solution to form a reconstituted solution, and quantitating the amount of or presence of PEAU, 2 EPMM, and/or PBG in the reconstituted solution.

[0024] In typical embodiments, the method of reducing formation of degradation products of Phenobarbital Sodium reconstituted in an aqueous solution results in the reconstituted solution having no less than 96% Phenobarbital Sodium. In more typical embodiments, the reconstituted solution has no less than 97% Phenobarbital Sodium. In some preferred embodiments, the reconstituted solution has no less than 98% Phenobarbital Sodium, and in other preferred embodiments between 95% and 105% Phenobarbital Sodium. In additional or alternative embodiments, the reconstituted solution has no more than 0.2% PEAU, 2 EPMM, and/or PBG, and preferably has 0.05% or less PEAU and no detectable amount of 2 EPMM or PBG.

[0025] Further aspects of the present disclosure include methods of treating an individual in need of Phenobarbital Sodium. These methods include adding saline or dextrose to the lyophilized powder of Phenobarbital Sodium as disclosed herein immediately prior to administration to form a Phenobarbital Sodium reconstituted solution and administering the Phenobarbital Sodium reconstituted solution to the individual. In some embodiments, the individual in need of Phenobarbital Sodium suffers from epilepsy. In particular embodiments, the individual in need of Phenobarbital Sodium is a newborn suffering from neonatal epilepsy. In other embodiments, the lyophilized amorphous Phenobarbital Sodium is a powder dose of 65 mg, 130 mg, 100 mg or 200 mg. In typical embodiments, the administering of the

Phenobarbital Sodium reconstituted solution to the individual includes intramuscular injection or intravenous injection.

[0026] Additional aspects of the present disclosure include a pharmaceutical product including a vial containing the white lyophilized powder of Phenobarbital Sodium for reconstitution prior to administration, in which the composition was lyophilized in the vial and the vial includes a closure positioned in order to seal the vial under vacuum after Lyophilization.

[0027] Therefore, the inventors also contemplate lyophilized Phenobarbital Sodium.

[0028] Various objects, features, aspects, and advantages will become more apparent from the following detailed description of preferred embodiments, along with the accompanying drawings.

BRIEF DESCRIPTION OF DRAWINGS

[0029] FIG. 1A is a polarized light photomicrograph of the presently disclosed Phenobarbital Sodium in water during Lyophilization microscopy at -50° C., as indicated.

[0030] FIG. 1B is a polarized light photomicrograph of the presently disclosed Phenobarbital Sodium in water during Lyophilization microscopy at -45.5° C. as indicated.

[0031] FIG. 1C is a polarized light photomicrograph of the presently disclosed Phenobarbital Sodium in water during Lyophilization microscopy at -40.0° C. as indicated.

[0032] FIG. 1D is a polarized light photomicrograph of the presently disclosed Phenobarbital Sodium in water during Lyophilization microscopy at -39.6° C. as indicated.

[0033] FIG. 1E is a polarized light photomicrograph of the presently disclosed Phenobarbital Sodium in water during Lyophilization microscopy at -39.2° C. as indicated.

[0034] FIG. 1F is a polarized light photomicrograph of the presently disclosed Phenobarbital Sodium in water during Lyophilization microscopy at -38.8° C. as indicated.

[0035] FIG. 2A is a chromatogram from an injection of a "blank solution" for assay by high performance liquid chromatography (HPLC) at a detection wavelength of 210 nanometers (nm), as described herein.

[0036] FIG. 2B is a chromatogram from an injection of the "Sensitivity solution" of Phenobarbital for related substances by HPLC at a detection wavelength of 210 nm, as described herein.

[0037] FIG. 2C is a chromatogram from an injection of a "related substance" (RS) working standard at 0.2% impurity level) for assay of related substances by HPLC at a detection wavelength of 210 nm, as described herein.

[0038] FIG. 2D is a chromatogram from an injection of 2-ethyl-2-phenylmalonamide (2EPM) for assay of related substances by HPLC at a detection wavelength of 210 nm, as described herein.

[0039] FIG. 2E is a chromatogram from an injection of alpha-phenylbutyrylguanidine (PBG) for assay of related substances by HPLC at a detection wavelength of 210 nm, as described herein.

[0040] FIG. 2F is a chromatogram from an injection of phenylethylacetylurea (PEAU) for assay of related substances by HPLC at a detection wavelength of 210 nm, as described herein.

[0041] FIG. 2G is a chromatogram from an injection of a "blank solution" for assay by HPLC at a detection wavelength of 254 nm, as described herein.

[0042] FIG. 2H is a chromatogram from an injection of an "assay working standard" for assay by HPLC at a detection wavelength of 254 nm, as described herein.

[0043] FIG. 3A is a chromatogram from an injection of a marketed liquid formulation of Phenobarbital Sodium at 65 mg/mL at initial time point for assay by HPLC at a detection wavelength of 254 nm, as described herein.

[0044] FIG. 3B is a chromatogram from an injection of a marketed liquid formulation of Phenobarbital Sodium at 65 mg/mL at 6 months from initial time point at room temperature (RT) for assay by HPLC at a detection wavelength of 254 nm, as described herein.

[0045] FIG. 3C is a chromatogram from an injection of a marketed liquid formulation of Phenobarbital Sodium at 65 mg/mL at 6 months from initial time point at accelerated stability conditions (ACC) for assay by HPLC at a detection wavelength of 254 nm, as described herein.

[0046] FIG. 3D is a chromatogram from an injection of a marketed liquid formulation of Phenobarbital Sodium at 65 mg/mL at initial time point for related substances by HPLC at a detection wavelength of 210 nm, as described herein.

[0047] FIG. 3E is a chromatogram from an injection of a marketed liquid formulation of Phenobarbital Sodium at 65 mg/mL at 6 months from initial time point at room temperature (RT) for related substances by HPLC at a detection wavelength of 210 nm, as described herein.

[0048] FIG. 3F is a chromatogram from an injection of a marketed liquid formulation of Phenobarbital Sodium at 65 mg/mL at 6 months from initial time point at accelerated stability conditions (ACC) for related substances by HPLC at a detection wavelength of 210 nm, as described herein.

[0049] FIG. 3G is a chromatogram from an injection of a marketed liquid formulation of Phenobarbital Sodium at 130 mg/mL at initial time point for assay by HPLC at a detection wavelength of 254 nm, as described herein.

[0050] FIG. 3H is a chromatogram from an injection of a marketed liquid formulation of Phenobarbital Sodium at 130 mg/mL at 6 months from initial time point at room temperature (RT) for assay by HPLC at a detection wavelength of 254 nm, as described herein.

[0051] FIG. 3I is a chromatogram from an injection of a marketed liquid formulation of Phenobarbital Sodium at 130 mg/mL at 6 months from initial time point at accelerated stability conditions (ACC) for assay by HPLC at a detection wavelength of 254 nm, as described herein.

[0052] FIG. 3J is a chromatogram from an injection of a marketed liquid formulation of Phenobarbital Sodium at 130 mg/mL at initial time point for related substances by HPLC at a detection wavelength of 210 nm, as described herein.

[0053] FIG. 3K is a chromatogram from an injection of a marketed liquid formulation of Phenobarbital Sodium at 130 mg/mL at 6 months from initial time point at room temperature (RT) for related substances by HPLC at a detection wavelength of 210 nm, as described herein.

[0054] FIG. 3L is a chromatogram from an injection of a marketed liquid formulation of Phenobarbital Sodium at 130 mg/mL at 6 months from initial time point at accelerated stability conditions (ACC) for related substances by HPLC at a detection wavelength of 210 nm, as described herein.

[0055] FIG. 4A is a chromatogram from an injection of the disclosed reconstituted lyophilized formulation of Phenobarbital Sodium at 65 mg/mL at initial time point for assay by HPLC at a detection wavelength of 254 nm, according to embodiments of the present invention.

[0056] FIG. 4B is a chromatogram from an injection of the disclosed reconstituted lyophilized formulation of Phenobarbital Sodium at 65 mg/mL at 6 months from initial time point at room temperature (RT) for assay by HPLC at a detection wavelength of 254 nm, according to embodiments of the present invention.

[0057] FIG. 4C is a chromatogram from an injection of the disclosed reconstituted lyophilized formulation of Phenobarbital Sodium at 65 mg/mL at 6 months from initial time point at accelerated stability conditions (ACC) for assay by HPLC at a detection wavelength of 254 nm, according to embodiments of the present invention.

[0058] FIG. 4D is a chromatogram from an injection of the disclosed reconstituted lyophilized formulation of Phenobarbital Sodium at 65 mg/mL at initial time point for related substances by HPLC at a detection wavelength of 210 nm, according to embodiments of the present invention.

[0059] FIG. 4E is a chromatogram from an injection of the disclosed reconstituted lyophilized formulation of Phenobarbital Sodium at 65 mg/mL at 6 months from initial time point at room temperature (RT) for related substances by HPLC at a detection wavelength of 210 nm, according to embodiments of the present invention.

[0060] FIG. 4F is a chromatogram from an injection of the disclosed reconstituted lyophilized formulation of Phenobarbital Sodium at 65 mg/mL at 6 months from initial time point at accelerated stability conditions (ACC) for related substances by HPLC at a detection wavelength of 210 nm, according to embodiments of the present invention.

[0061] FIG. 4G is a chromatogram from an injection of the disclosed reconstituted lyophilized formulation of Phenobarbital Sodium at 130 mg/mL at initial time point for assay by HPLC at a detection wavelength of 254 nm, according to embodiments of the present invention.

[0062] FIG. 4H is a chromatogram from an injection of the disclosed reconstituted lyophilized formulation of Phenobarbital Sodium at 130 mg/mL at 6 months from initial time point at room temperature (RT) for assay by HPLC at a detection wavelength of 254 nm, according to embodiments of the present invention.

[0063] FIG. 4I is a chromatogram from an injection of the disclosed reconstituted lyophilized formulation of Phenobarbital Sodium at 130 mg/mL at 6 months from initial time point at accelerated stability conditions (ACC) for assay by HPLC at a detection wavelength of 254 nm, according to embodiments of the present invention.

[0064] FIG. 4J is a chromatogram from an injection of the disclosed reconstituted lyophilized formulation of Phenobarbital Sodium at 130 mg/mL at initial time point for related substances by HPLC at a detection wavelength of 210 nm, according to embodiments of the present invention.

[0065] FIG. 4K is a chromatogram from an injection of the disclosed reconstituted lyophilized formulation of Phenobarbital Sodium at 130 mg/mL at 6 months from initial time point at room temperature (RT) for related substances by HPLC at a detection wavelength of 210 nm, according to embodiments of the present invention.

[0066] FIG. 4L is a chromatogram from an injection of the disclosed reconstituted lyophilized formulation of Phenobarbital Sodium at 130 mg/mL at 6 months from initial time point at accelerated stability conditions (ACC) for related substances by HPLC at a detection wavelength of 210 nm, according to embodiments of the present invention.

[0067] FIG. 5A is an X-ray powder diffraction (XRPD) pattern of Phenobarbital Sodium Grade P

[0068] FIG. 5B is an XRPD pattern of Phenobarbital Sodium for Injection, USP, 65 mg/vial.

[0069] FIG. 5C is an XRPD pattern of Phenobarbital Sodium for Injection, USP, 130 mg/vial.

[0070] FIG. 5D is a stack plot of the XRPD patterns of FIGS. 5A, 5B, and 5C.

DETAILED DESCRIPTION

[0071] The inventors have surprisingly discovered a method for making a storage-stable and highly purified Phenobarbital Sodium composition that is void of any detectable impurities in its lyophilized powder form and is essentially void of impurities upon reconstitution in water. The contemplated Phenobarbital Sodium composition is a lyophilized powder that is no less than 98% Phenobarbital Sodium and is stable (25° C., 65% RH and 40° C., 75% RH) for up to at least 6 months under accelerated conditions which is equivalent to 24 months at RT. Upon reconstitution (e.g. solubilization/resuspension) in water, the lyophilized Phenobarbital Sodium forms no more than 0.05% of 2-ethyl-2-phenylmalonamide (2EPMM) or alpha-phenylbutyrylguanidine (PBG), and phenylethylacetylurea (PEAU).

[0072] The Phenobarbital Sodium composition as described herein refers to Phenobarbital Sodium having an IUPAC (International Union of Pure and Applied Chemistry) name of sodium 5-ethyl-4,6-dioxo-5-phenyl-1H-pyrimidin-2-olate, a Chemical Abstract Service (CAS) No. 57-30-7, and a molecular formula of $C_{12}H_{11}N_2NaO_3$.

[0073] In order to avoid hydrolysis of Phenobarbital Sodium and therefore the formation of its toxic degradation products, the inventors have unexpectedly discovered that a lyophilized Phenobarbital Sodium formed by the methods disclosed herein results in a storage-stable lyophilized powder of Phenobarbital Sodium. In particular, with reference to the polarized light photomicrographs of FIGS. 1A-1F and the XRPD analysis shown in FIG. 5A-5D, the storage-stable lyophilized powder of Phenobarbital Sodium has an amorphous structure. Without wishing to be bound by any theory or hypothesis, the lyophilized Phenobarbital Sodium produced following the methods according to embodiments of the present disclosure produces an amorphous structure that contributes to the observed stability of this composition upon reconstitution in water. Irrespective of the mechanism of action, the contemplated lyophilized Phenobarbital Sodium composition is stable and upon reconstitution in water, does not form significant (e.g., not measurable or not detectable) amounts of the toxic degradation products—namely PEAU, 2EPMM, or PBG.

[0074] The contemplated method for producing a storage-stable lyophilized Phenobarbital Sodium composition includes adding purified Phenobarbital Sodium to water. Preferably the purified Phenobarbital Sodium solid is purchased from a chemical supplier (e.g., Siegfried, USA) that adheres to the United States Pharmacopeia (USP) manufacturing and synthesis guidelines. The USP grade of the Phenobarbital Sodium is preferably USP “purified” or “practical” (P) grade indicating that it is a good quality chemical. As understood in the art, the USP grade (P) active pharmaceutical ingredient (API) of Phenobarbital Sodium is not suitable for administration because it is not sterile, is hygroscopic, and as such, maintaining anhydrous conditions of this grade of API is not feasible. Structurally, the analyzed

USP grade P API (Siegfried, USA) is polymorphous—containing both crystalline and non-crystalline material. With reference to FIG. 5A, an X-ray powder diffraction (XRPD) pattern is shown of this compound.

[0075] Considering the need to avoid hydrolysis of the Phenobarbital Sodium, the Phenobarbital Sodium is preferably added to Water for Injection and later lyophilized to obtain a stable lyophilized finished product. In order to remove dissolved oxygen (O₂) gas from the water (e.g., Water for Injection), it is also preferred that the water is sparged with nitrogen (N₂) or other inert gas prior to the addition of the Phenobarbital Sodium solid. For effective sparging, a volume of about 100-200 ml water is sparged with nitrogen gas for at least 30 minutes. In exemplary protocols for producing either 65 mg, 130 mg of lyophilized Phenobarbital Sodium, 100 milliliters (ml) of Water for Injection is nitrogen sparged for 30 minutes. In additional embodiments, the water may also be cooled to approximately 2 to 8° C. In preferred embodiments, the water is cooled to approximately 2 to 8° C.

[0076] The addition of the Phenobarbital Sodium solid to the water (e.g., deionized nitrogen sparged water cooled to 2 to 8° C.) may be carried out using any suitable mixing. Preferably, the water (e.g., Water for Injection) is first added to a glass/stainless steel container for nitrogen sparging and cooling, and this prepped water remains in the glass/stainless steel container or a measured amount is transferred to a preferably weighed glass container/stainless steel container and the Phenobarbital Sodium solid is added thereto. For mixing, any suitable mixing may be used. For example, a magnetic stir bar may be added to the glass/stainless steel container. Upon addition of the Phenobarbital Sodium solid to the water, the mixture is stirred.

[0077] An additional consideration for stability of the Phenobarbital Sodium is maintaining a pH between 9.2 and 10.2 in the bulk solution of Phenobarbital Sodium. After dissolution of the Phenobarbital Sodium solid in the water, the pH is adjusted to within this range of 9.2 and 10.2. Accordingly, the pH of the water solubilized Phenobarbital Sodium is first measured upon dissolution. If the pH is higher than 10.2 or lower than 9.2, the pH is adjusted. For example, if necessary, the pH may be lowered to or between 9.2 and 10.2 using Hydrochloric acid (HCl) and the pH may be increased to or between 9.2 and 10.2 using Sodium Hydroxide (NaOH).

[0078] Following the pH measurement and any adjustment, the Phenobarbital Sodium solution is preferably filtered to remove any foreign particles. For example, the Phenobarbital Sodium solution may be filtered using a 0.2 µm (or 0.22 µm) membrane filter. For example, the membrane filter may be polyvinylidene difluoride (PVDF) or any other compatible filter membranes such as Polyethersulfone (PES) filter.

[0079] According to aspects of the contemplated method, the Phenobarbital Sodium solution dissolved in the water (e.g., nitrogen sparged Water for Injection cooled to 2 to 8° C.) having a pH of or between 9.2 and 10.2 and preferably filtered is ready for Lyophilization. Depending on the desired amount of the lyophilized Phenobarbital Sodium product, the Phenobarbital Sodium solution may be lyophilized in any amount or the solution may be aliquoted into sterile containers (e.g., vials). For example, in order to obtain a measured dose of lyophilized Phenobarbital Sodium in a container, the glass container, water, and Phenobarbital

Sodium solid are weighed prior to mixing in order to determine an actual weight of the Phenobarbital Sodium after pH measurement and adjustment. For example, to produce 65 mg or 130 mg doses of lyophilized Phenobarbital Sodium, a batch may be dissolved, mixed, pH adjusted if necessary, and filtered and then aliquoted into single use dose vials for lyophilization. In a specific example, a 50 mL batch of Phenobarbital Sodium solution may be prepared where for 65 mg doses, 3.25 grams of Phenobarbital Sodium solid (e.g., USP Grade P) is added to 40 ml of water (e.g., nitrogen sparged Water for Injection cooled to 5 to 8° C.) and for 130 mg doses, 6.5 grams of the Phenobarbital Sodium is added to the 40 ml of water. Following dissolution, pH adjustment if necessary, adjusting final volume to 50.0 mL and filtering, 1 ml aliquots of this Phenobarbital Sodium solution are added to open (e.g., unstoppered) vials and then loosely stoppered (e.g., with lyophilization rubber stoppers) prior to lyophilization. Following lyophilization, each vial contains 65 mg or 130 mg of lyophilized Phenobarbital Sodium.

[0080] In further embodiments, following lyophilization, the vials may be completely closed (e.g., stoppered) under vacuum.

[0081] Another embodiment is directed to a pharmaceutical composition that consists of lyophilized amorphous Phenobarbital Sodium.

[0082] Another embodiment is directed to a method of producing a storage-stable form of lyophilized Phenobarbital sodium that forms no more than 0.2% phenylethylacetylurea (PEAU) when reconstituted in an aqueous solution. The method comprising: includes adding Phenobarbital Sodium to water to form a Phenobarbital Sodium solution, if necessary, adjusting the pH of the Phenobarbital Sodium solution to or between 9.2 to 10.2; and lyophilizing the Phenobarbital Sodium solution.

[0083] In one embodiment, the water is Water for Injection.

[0084] In another embodiment the Water for Injection is sparged with nitrogen prior to adding the Phenobarbital Sodium.

[0085] In another embodiment, the Water for Injection is sparged with nitrogen for at least 30 minutes prior to adding the Phenobarbital Sodium.

[0086] In another embodiment, the water is cooled to or between 2 to 8° C. prior to adding the Phenobarbital Sodium.

[0087] In another embodiment, the adjusting includes measuring the pH of the Phenobarbital Sodium solution, and if the pH is higher than 10.2, the adjusting includes adding Hydrochloric acid (HCl).

[0088] In another embodiment, the lyophilized Phenobarbital Sodium forms no more than 0.1% phenylethylacetylurea (PEAU) when reconstituted in an aqueous solution after storage of the lyophilized Phenobarbital Sodium for up to 6 months at room temperature (RT) and accelerated stability conditions (ACC).

[0089] In another embodiment, the lyophilized Phenobarbital Sodium forms no more than 0.05% phenylethylacetylurea (PEAU) when reconstituted in an aqueous solution after storage of the lyophilized Phenobarbital Sodium for up to 6 months at room temperature (RT) and accelerated stability conditions (ACC).

[0090] In another embodiment, the lyophilized Phenobarbital Sodium forms no detectable amount of 2-ethyl-2-phenylmalonamide (2EPMM) or alpha-phenylbutyrylguan-

dine (PBG) when reconstituted in an aqueous solution after storage of the lyophilized Phenobarbital Sodium for up to 6 months at room temperature (RT) and accelerated stability conditions (ACC).

[0091] In another embodiment, the lyophilized Phenobarbital Sodium is stable up to at least 6 months at accelerated stability conditions (ACC) which is equivalent to 24 months at room temperature (RT).

[0092] In another embodiment, the stability of the lyophilized Phenobarbital Sodium is determined by adding the lyophilized Phenobarbital Sodium to an aqueous solution to form a reconstituted lyophilized Phenobarbital Sodium and analyzing to quantitate the amount of phenylethylacetylurea (PEAU), 2-ethyl-2-phenylmalonamide (2EPMM), and/or alpha-phenylbutyrylguanidine (PBG) in the reconstituted lyophilized Phenobarbital Sodium.

[0093] In another embodiment, the Phenobarbital Sodium solution is at a concentration of 65 mg/mL, 100 mg/mL, 130 mg/mL or 200 mg/mL.

[0094] In another embodiment, the method further includes filtering the Phenobarbital Sodium solution prior to lyophilizing.

[0095] In another embodiment, the method further includes aliquoting the Phenobarbital Sodium solution into a vial prior to lyophilization; and after lyophilization, applying a closure to the vial under vacuum.

[0096] Another embodiment is directed to a method of treating an individual in need of Phenobarbital Sodium. The method includes adding saline or dextrose to a composition containing lyophilized amorphous Phenobarbital Sodium immediately prior to administration to form a Phenobarbital Sodium solution; and administering the Phenobarbital Sodium solution to the individual.

[0097] In another embodiment, the individual suffers from epilepsy.

[0098] In another embodiment, the individual is a newborn suffering from neonatal epilepsy.

[0099] In another embodiment, the composition is a powder dose of 65 mg, 100, 130 mg or 200 mg.

[0100] In another embodiment, the administering includes intramuscular injection or intravenous injection.

[0101] Surprisingly, the inventors have determined that the presently disclosed method produces a storage-stable lyophilized Phenobarbital Sodium that maintains its stability up to at least 6 months under accelerated stability conditions which is equivalent to 24 months at RT. With reference to FIGS. 5B-5C, the lyophilized Phenobarbital Sodium is amorphous. In particular, the lyophilized Phenobarbital Sodium is non-crystalline unlike the USP grade PAPI shown comparatively in FIG. 5D. The lyophilized Phenobarbital Sodium is stable and ready for reconstitution in either water for injection, 0.9% Saline or 5% Dextrose for intravenous (IV) or intramuscular (IM) administration. The presently disclosed formulation does not contain any organic solvents or preservatives which is specifically advantageous for administration to newborns and neonates. Moreover, the resulting lyophilized Phenobarbital Sodium forms below the quantitation limit of 0.05% phenylethylacetylurea (PEAU), and no detectable amount of 2-ethyl-2-phenylmalonamide (2EPMM) or alpha-phenylbutyrylguanidine (PBG) when reconstituted in water after storage of the lyophilized Phenobarbital Sodium for up to 6 months under accelerated stability conditions which is equivalent to 24 months at RT.

[0102] In a comparison analysis, the presently disclosed lyophilized powder of Phenobarbital Sodium at 65 mg or 130 mg doses was reconstituted in water and assayed for impurities and stability along with marketed liquid formulation of Phenobarbital Sodium for Injection USP, 65 mg/mL or 130 mg/mL (Westward Pharmaceuticals). The formulation details for this comparative analysis are presented in Table 1. Representative chromatograms from HPLC analysis are shown in FIGS. 2A-2H, 3A-3L, and 4A-4L.

TABLE 1

Formulation Details			
Dosage form		Liquid	Lyophilized
Components	Chemical name	mg/mL	mg/vial
Drug Substance Inactive ingredients	Phenobarbital Sodium	65 or 130	65 or 130
	Propylene Glycol	705	NA
	Benzyl alcohol	15.6	NA
	Ethanol	78.9	NA
	HCL	For pH adjustment	NA
	Water for Injection	QS	NA
		(Quantity Supplied)	

Lyophilized Phenobarbital Sodium for Injection USP, 65 mg/vial and 130 mg/vial
Liquid Phenobarbital Sodium for Injection USP, 65 mg/mL or 130 mg/mL

[0103] With reference to Tables 2 and 3, the presently disclosed 65 mg lyophilized Phenobarbital Sodium composition was reconstituted in water and compared to the 65 mg Liquid Phenobarbital Sodium for Injection USP (65 mg/mL marketed Liquid Formulation) (Table 2) (FIGS. 3A-3F, FIGS. 4A-4F), and the presently disclosed 130 mg lyophilized Phenobarbital Sodium for Injection USP (130 mg/vial) composition was reconstituted in water and compared to the 130 mg liquid Phenobarbital Sodium for Injection USP (130 mg/mL marketed Liquid Formulation) (Table 3) (FIGS. 3G-3L, FIGS. 4G-4L). In addition to the noted characteristics of each product, the products were assayed for the degradation products—phenylethylacetylurea (PEAU), 2-ethyl-2-phenylmalonamide (2EPMM), and alpha-phenylbutyrylguanidine (PBG) by HPLC analysis, as further described in the Examples. As indicated in both Tables 2 and 3, the initial amount of PEAU (phenylethylacetylurea) assayed in the 65 mg/mL and 130 mg/mL liquid formulations was 0.5% and 0.5%, respectively. Comparatively, no initial amount of PEAU was detected in either the 65 mg or 130 mg lyophilized Phenobarbital Sodium compositions. Furthermore, after 3 and 6 months from the date of manufacture, both the liquid and lyophilized formulations at 65 mg/mL and 130 mg/mL stored at room temperature (RT) (25° C.) and under accelerated stability conditions (ACC) at 40° C. were analyzed by HPLC for assay and related substances. Under these conditions, the amount of PEAU assayed in the 65 mg/mL liquid formulations at 3 months was 1.0% (RT) and 3.7% (ACC) and at 6 months was 1.4% (RT) and 5.8% (ACC) (Table 2). Similarly, the amount of PEAU assayed in 130 mg/mL liquid formulations at 3 months was 0.9% (RT) and 2.8% (ACC) and at 6 months was 1.1% (RT) and 4.3% (ACC) (Table 3). In contrast, for the presently disclosed lyophilized compositions, the amount of PEAU assayed at 3 and 6 months for the 65 mg and the 130 mg lyophilized Phenobarbital Sodium compositions was below quantitation limit (BQL) (0.05%) both at RT and ACC stability conditions (Tables 2-3).

TABLE 2

65 mg/ml Liquid Formulation and 65 mg/vial Lyophilized Powder											
Product Name											
Phenobarbital Sodium for Injection USP, 65 mg/mL (Marketed product)						Phenobarbital Sodium for Injection USP, 65 mg/Vial (Nivagen formulation- NPH1907)					
Formulation Type											
Liquid						Lyophilized powder					
Stability Data											
Test	Initial	3 Months		6 Months		3 Months		6 Months		15 Months	
		RT	ACC	RT	ACC	RT	ACC	RT	ACC	RT**	
		25° C.,	40° C.,	25° C.,	40° C.,	25° C.,	40° C.,	25° C.,	40° C.,	25° C.,	
		60%	75%	60%	75%	60%	75%	60%	75%	60%	
		RH	RH	RH	RH	Initial	RH	RH	RH	RH	RH
Description	CCVS	CCVS	CCVS	CCVS	CCVS	WLP	WLP	WLP	WLP	WLP	WLP
Appearance after reconstitution	NA	NA	NA	NA	NA	CCS	CCS	CCS	CCS	CCS	CCS
Identification	\$	\$	\$	\$	\$	\$	\$	\$	\$	\$	\$
pH*	9.3	9.8	9.7	9.6	9.6	9.8	9.6	9.6	9.7	9.7	9.8
% Assay of Phenobarbital Sodium	102.4	97.6	103.1	96.3	86.7	99.1	98.7	97.4	100.0	99.3	96.0
% degradation (Area percent)	ND	ND	BQL	BQL	BQL	ND	ND	ND	ND	ND	ND
2EPMM											
PBG	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
PEAU	0.5	1.0	3.7	1.4	5.8	ND	BQL	BQL	BQL	BQL	BQL
Any unspecified degradation product	BQL	BQL	0.1	BQL	0.3	BQL	BQL	BQL	0.07	0.06	0.04
Total degradation products	0.5	1.0	3.8	1.4	6.1	BQL	BQL	BQL	0.07	0.06	0.04

CCVS: Clear Colorless Viscous Solution;

CCS: Clear Colorless Solution;

WLP: White Lyophilized powder

\$: In Assay analysis, the retention time of Phenobarbital in standard and sample solutions should be same.

BQL: Below Quantitation Limit; (QL-0.05%);

ND: Not Detected

2EPMM: 2-Ethyl-2-Phenylmalonamide, monohydrate;

PBG: α -phenylbutyrylguanidine,

PEAU: phenylethylacetylurea or Pheneturide

*pH of both liquid and lyophilized formulations are measured at 65 mg/mL concentration

**The samples were stored at 30 $\pm$ 2° C., 65% RH $\pm$ 5

Notably, the Lyophilized powder was stable even after 15 months, and no significant increase in the impurities was

observed, rendering such formulation superior to known liquid formulations and enable easy administration.

TABLE 3

130 mg/mL Liquid Formulation and 130 mg/vial Lyophilized formulation											
Test	Product Name										
	Phenobarbital Sodium for Injection USP, 130 mg/mL (Marketed product)						Phenobarbital Sodium for Injection USP, 130 mg/vial (Nivagen formulation-NPH1912)				
	Formulation Type										
	Liquid					Lyophilized powder					
	Stability Data										
		3 Months		6 Months			3 Months		6 Months		15 Months
		RT 25° C., 60% RH	ACC 40° C., 75% RH	RT 25° C., 60% RH	ACC 40° C., 75% RH		RT 25° C., 60% RH	ACC 40° C., 75% RH	RT 25° C., 60% RH	ACC 40° C., 75% RH	RT** 25° C., 60% RH
Description	CCVS	CCVS	CCVS	CCVS	CCVS	WLP	WLP	WLP	WLP	WLP	WLP
Appearance after reconstitution	NA	NA	NA	NA	NA	CCS	CCS	CCS	CCS	CCS	CCS
Identification	\$	\$	\$	\$	\$	\$	\$	\$	\$	\$	\$
pH*	9.5	9.9	9.9	9.8	9.7	9.8	9.8	9.8	9.8	9.8	10.0
% Assay of Phenobarbital Sodium	100.8	95.5	92.9	95.2	90.9	98.3	100.1	97.9	100.2	99.4	98.3
Related Substances (% Area)	ND	ND	BQL	BQL	BQL	ND	ND	ND	ND	ND	ND
2EPMM											
PBG	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
PEAU	0.5	0.9	2.8	1.1	4.3	ND	BQL	BQL	BQL	BQL	BQL
Any unspecified degradation product	ND	BQL	0.1	BQL	0.4	BQL	BQL	BQL	0.05	0.05	0.04
Total degradation products	0.5	0.9	2.9	1.1	4.7	BQL	BQL	BQL	0.05	0.05	0.04

CCVS: Clear Colorless Viscous Solution;

CCS: Clear Colorless Solution;

WLP: White Lyophilized powder

\$: In Assay analysis, the retention time of Phenobarbital in standard and sample solutions should be same.

BQL: Below Quantitation Limit; (QL<0.05%);

ND: Not Detected

2EPMM: 2-Ethyl-2-Phenylmalonamide, monohydrate;

PBG: α -phenylbutyrylguanidine,

PEAU: phenylethylacetylurea or Pheneturide

*pH for liquid formulation is measured at 130 mg/mL and for lyophilized formulation at 100 mg/mL dilution

**The samples were stored at 30 $\pm$ 2° C., 65% RH $\pm$ 5

[0104] In typical embodiments, the method of reducing formation of degradation products of a lyophilized Phenobarbital Sodium composition reconstituted in water results in the reconstituted solution having no less than 96% Phenobarbital Sodium. In more typical embodiments, the reconstituted solution has no less than 97% Phenobarbital Sodium. In more preferred embodiments, the reconstituted solution has no less than 98% Phenobarbital Sodium. In additional or alternative embodiments, the reconstituted solution has no quantitative amount (e.g., 0.05% or less) of PEAU, 2 EPMM, or PBG.

[0105] According to contemplated embodiments, assaying the stability of the lyophilized Phenobarbital Sodium composition includes reconstituting the composition in water at a concentration of 65 mg/mL, 100 mg/mL, 130 mg/mL or 200 mg/mL.

[0106] Further aspects of the present disclosure include methods of treating an individual in need of Phenobarbital

Sodium. For administration to an individual (e.g., a human patient) the lyophilized powder of Phenobarbital Sodium powder may be reconstituted in saline or dextrose as disclosed herein immediately prior to administration to form a Phenobarbital Sodium reconstituted solution. In particular embodiments, the individual in need of Phenobarbital Sodium suffers from epilepsy. In preferred embodiments, the individual in need of Phenobarbital Sodium is a newborn suffering from neonatal epilepsy. In other embodiments, the lyophilized Phenobarbital Sodium is a powder dose of 65 mg, 100 mg, 130 mg or 200 mg. In typical embodiments, the administering of the Phenobarbital Sodium reconstituted solution to the individual includes intramuscular injection or intravenous injection.

Examples

[0107] Lyophilization Microscopy. Lyophilization microscopy (FIGS. 1A-1F) was performed using a Leica DM LP

microscope equipped with a Linkam FDCS196 stage, TMS93 controller, vacuum pump, LNP unit, and a Spot Insight color camera. A 20x, 040 N.A. objective was used with crossed polarizers and a first order red compensator to view sample. Images were acquired using Spot Advanced software (v.4.5.9).

[0108] Lyophilization parameters for the lyophilization microscopy including a start temperature of -50°C . with heating at $1^{\circ}\text{C}/\text{minute}$ to -40°C . followed by heating at $0.5^{\circ}\text{C}/\text{minute}$ to -32.0°C .

[0109] HPLC Assays. High Performance Liquid Chromatography (HPLC) was used to assay Phenobarbital Sodium standards and sample solutions as disclosed herein (e.g., Tables 2, 3, FIGS. 3A-3L, and 4A-4L). Both Ultra High-Performance Liquid Chromatographic system (UHPLC) and High-Performance Liquid Chromatographic system (HPLC) (ThermoFisher/Waters) were used together with Data Acquisition Software (Empower 3), equipped with UV and Photodiode Array Detector (UV and PDA), column thermostat and auto sampler compartments or equivalent HPLC system and software.

[0110] An exemplary HPLC assay protocol is provided in the following:

[0111] Instruments/Apparatus: HPLC with UV detector/PDA Detector, Analytical balance, Sonicator, Volumetric flasks, Beakers, Pipettes.

[0112] Reagents, Solvents, Standards: Phenobarbital Reference Standard, Deionized water/Water for Injection, Potassium Monobasic Phosphate, Acetonitrile.

[0113] Preparation of Mobile Phase Solutions: Potassium phosphate monobasic (7 mM) and Acetonitrile (70:30) pH 4.5-6.5; Diluent: Mobile phase as a diluent.

[0114] Wash Solvent: Water: Acetonitrile (70:30)

[0115] CHROMATOGRAPHIC CONDITIONS: The liquid chromatography equipped with a UV and PDA detector, an injector and a data processor is operated as follows:

Parameters	Conditions
Column	L7 packaging column, 150×4.6 mm, Particle Size: $5\ \mu\text{m}$ or equivalent
Flow rate	$0.9\text{--}1.2$ mL/min.
Detection Wavelength	254 ± 2 nm for assay and 210 ± 2 nm for Related Substances
Injection volume	$10\ \mu\text{L}$

-continued

Parameters	Conditions
Run time	15 Minutes
Column Oven Temperature	30°C .
Sampler Cooler Temperature	25°C .
Retention Time	About 5 minutes for Phenobarbital
Pump Mode	Isocratic

[0116] Phenobarbital Assay Standard Solution: $0.2\ \text{mg/mL}$ to $0.8\ \text{mg/mL}$ in diluent; Phenobarbital Related Substances Standard Solution: $0.26\ \mu\text{g/mL}$ to $15.6\ \mu\text{g/mL}$ in diluent; Phenobarbital Related Substances resolution standard solution: $0.475\ \text{mg/mL}$ Phenobarbital and about $1\ \mu\text{g/mL}$ for related substances (2-Ethyl-2-Phenyl malonamide, monohydrate (2EPMM), Phenylbutyrylguanidine (PBG) and Phenylethylacetylurea (PEAU) in diluent; Phenobarbital Sodium Sample Solution for Assay: $0.26\ \text{mg/mL}$ to $0.78\ \text{mg/mL}$ in diluent

[0117] Phenobarbital Sodium related Substances sample solution: $0.52\ \text{mg/mL}$ of Phenobarbital Sodium and related substances 2-Ethyl-2-Phenylmalonamide, monohydrate (2EPMM), Phenyl butyrylguanidine (PBG) and Phenylethylacetylurea (PEAU) and any unspecified degradation products at $0.26\ \mu\text{g/mL}$ to $13.2\ \mu\text{g/mL}$

[0118] SYSTEM SUITABILITY REQUIREMENTS: The relative standard deviation of Phenobarbital peak from Six replication injections of related substances standard should not be more than 6.0%. The related standard deviation of Phenobarbital peak from five replicate injections in assay standard solution should not be more than 2.0%. Tailing factor for the Phenobarbital is not more than 2.0. USP Plate count for the Phenobarbital is not less than 2500. The S/N Ratio of the Sensitivity Standard Solution is NLT 10 for related substance test. The resolution between Phenobarbital peak and nearest impurity peak is NLT 1.5

[0119] Two alternative formulations are disclosed in Table 6. One of these formulations is $200\ \text{mg/vial}$, Phenobarbital Sodium reconstituted in $10\ \text{mL}$ of Water for Injection or 5% Dextrose to obtain a concentration of $20\ \text{mg/mL}$. A second formulation is $200\ \text{mg/vial}$ of Phenobarbital Sodium along with $90\ \text{mg}$ Sodium Chloride reconstituted in $10\ \text{mL}$ of Water for Injection to obtain a concentration of $20\ \text{mg/mL}$ Phenobarbital Sodium for Injection and $9\ \text{mg/mL}$ or 0.9% Sodium Chloride.

TABLE 6

Phenobarbital Sodium for Injection, USP, 200 mg/vial in 10 mL vial. Formulation Details.					
		NPH2031B (Bulk Solution @ 100 mg/mL Phenobarbital Sodium and 45 mg/mL Sodium Chloride) Compounding at $2\text{--}8^{\circ}\text{C}$. Fill Volume: 2.1 mL			
		NPH2031A (Bulk Solution @ 100 mg/mL Phenobarbital Sodium) Compounding at $2\text{--}8^{\circ}\text{C}$.			
Ingredients	Manufacturer	mg/mL (after reconstitution)	mg/vial	mg/mL (after reconstitution)	mg/vial
Phenobarbital Sodium	Sieglfried USA, LLC	20	200	20	200

TABLE 6-continued

Phenobarbital Sodium for Injection, USP, 200 mg/vial in 10 mL vial. Formulation Details.					
		NPH2031B (Bulk Solution @ 100 mg/mL Phenobarbital Sodium and 45 mg/mL Sodium Chloride) Compounding at 2-8° C. Fill Volume: 2.1 mL			
		NPH2031A (Bulk Solution @ 100 mg/mL Phenobarbital Sodium) Compounding at 2-8° C.			
Ingredients	Manufacturer	mg/mL (after reconstitution)	mg/vial	mg/mL (after reconstitution)	mg/vial
Sodium Chloride	JT Baker	NA	NA	9	90
D1 Water	In house	qs	qs	qs	qs
100% Nitrogen	Airgas	NA	NA	NA	NA
10 mL Tubular Serum Bottle clear		Wheaton Part#223686			
20 mm Igloo Bromobutyl stopper		Dartwyler			
Seal 20 mm Filp-Cap, Red		Wheaton Part# 224203			

Reconstitution Volume 10 mL; Diluents that can be used are Water for Injection, 0.9% Saline or 5% Dextrose.

[0120] These formulations along with the previously disclosed formulations were developed to enable physicians and medical staff at hospitals to easily reconstitute to a desired concentration (e.g., 20 mg/mL) and administer the medicine to patients in need thereof. The reconstituted solution is stable at 2-8° C. for 24 hrs and 3 to 4 hrs when stored at RT (see e.g., Tables 7, 8 and 9).

TABLE 7

Reconstituted Solution Stability of Phenobarbital Sodium for Injection USP, 200 mg/vial (NPH2031A Phenobarbital Sodium @ 200 mg/vial) (Reconstitution in 10 mL Water)				
Test	Spec.	Initial DOA:	Reconstituted Solution Stability after 20 hrs DOA: Jun. 6, 2020	
		Jun. 5, 2020	2-8° C.	RT
Description	WLP	WLP	WLP	WLP
Appearance	CCS	CCS	CCS	CCS
pH (20 mg/mL)	9.2-10.2	9.9	9.7	9.8
% Assay	90.0-105.0	100.1	100.1	100.1
Phenobarbital Sodium				
Related Substances (% Area)	NMT 0.2%	BQL	ND	ND
2EPMM				
PBG	NMT 0.2%	ND	ND	ND
PEAU	NMT 0.2%	BQL	0.03	0.03
Any unspecified degradation product	NMT 0.2%	BQL	0.06	0.24
Total degradation products	NMT 1.5%	BQL	0.09	0.27
Water Content by Karl Fischer	NMT 3.0%	0.84	NA	NA

TABLE 7-continued

Reconstituted Solution Stability of Phenobarbital Sodium for Injection USP, 200 mg/vial (NPH2031A Phenobarbital Sodium @ 200 mg/vial) (Reconstitution in 10 mL Water)				
Test	Spec.	Initial DOA:	Reconstituted Solution Stability after 20 hrs DOA: Jun. 6, 2020	
		Jun. 5, 2020	2-8° C.	RT
Note Book Reference		192045-163 to 189	192045-163-183	

DOA: Date of Analysis;
WLP: White Lyophilized Powder;
CCS: Clear Colorless Solution, 1 minute after reconstitution;
ND: Not Detected;
NMT: Not More Than;
BQL: Below Quantitation Limit (0.05%)
2EPMM: 2-Ethyl-2-Phenylmalonamide, monohydrate;
PBG: α -Phenylbutyrylguanidine;
PEAU: Phenylethylacetylurea

TABLE 8

Reconstituted Solution Stability of Phenobarbital Sodium for Injection USP, 200 mg/vial in Sodium Chloride (NPH2031B Phenobarbital Sodium @ 200 mg/vial and Sodium Chloride @ 90 mg/vial) (Reconstitution in 10 mL Water)				
Test	Specification	Reconstituted Solution Stability after 20 hrs		
		Initial	2-8° C.	RT
Description	WLP	WLP	WLP	WLP
Appearance	CCS	CCS	CCS	CCS
pH (20 mg/mL)	9.2-10.2	9.8	9.7	9.8
% Assay	90.0-105.0	103.3	103.4	102.8
Phenobarbital Sodium				
Related Substances (% Area)	NMT 0.2%	BQL	ND	BQL
2EPMM				
PBG	NMT 0.2%	ND	ND	ND
PEAU	NMT 0.2%	BQL	0.03	0.03
Any unspecified	NMT 0.2%	BQL	0.06	0.27

TABLE 8-continued

Reconstituted Solution Stability of Phenobarbital Sodium for Injection USP, 200 mg/vial in Sodium Chloride (NPH2031B Phenobarbital Sodium @ 200 mg/vial and Sodium Chloride @ 90 mg/vial) (Reconstitution in 10 mL Water)				
Test	Specification	Reconstituted Solution Stability after 20 hrs		
		Initial	2-8° C.	RT
degradation product				
Total degradation products	NMT 1.5%	BQL	0.09	0.3
Water Content by Karl Fischer	NMT 3.0%	0.84	NA	NA
WLP: White Lyophilized Powder; CCS: Clear Colorless Solution, 1 minute after reconstitution; ND: Not Detected; NMT: Not More Than; BQL: Below Quantitation Limit (0.05%) 2EPMM: 2-Ethyl-2-Phenylmalonamide, monohydrate; PBG: α -Phenylbutyrylguanidine; PEAU: Phenylethylacetylurea				

TABLE 9

Reconstituted Solution Stability of Phenobarbital Sodium for injection USP, 130 mg/vial (NPH2030) (20 mg/mL dilution In Water)								
Tests	Specifications	Time						
		Shelf life	2 hrs		4 hrs		24 hrs	
			Initial	2-8 C.	RT	2-8 C.	RT	2-8 C.
Appearance	CCS	CCS	CCS	CCS	CCS	CCS	CCS	CCS
pH (20 mg/mL dilution)	9.2-10.2	9.5	9.5	9.6	9.6	9.5	9.6	9.5
Assay of Phenobarbital Sodium	90.0%-105.0%	99.8	102.1	100.8	101.7	101.2	100.4	100.3
Related Substances (% Area)	NMT 0.2%	ND	ND	ND	ND	ND	ND	BQL
2EPMM								
PBG	NMT 0.2%	ND	ND	ND	ND	ND	ND	ND
PEAU	NMT 0.2%	BQL	BQL	BQL	BQL	BQL	BQL	BQL

TABLE 9-continued

Reconstituted Solution Stability of Phenobarbital Sodium for injection USP, 130 mg/vial (NPH2030) (20 mg/mL dilution In Water)								
Tests	Specifications	Time						
		Initial	2 hrs		4 hrs		24 hrs	
			2-8 C.	RT	2-8 C.	RT	2-8 C.	RT
Any unspecified degradation product @ RRT 0.37	NMT 0.2%	0.015	0.02	0.04	0.02	0.06	0.04	0.18
Total degradation products	NMT 1.5%	0.015	0.02	0.04	0.02	0.06	0.04	0.18
Osmolality mOsmol/Kg	To be reported	158	NP	NP	NP	NP	159	158

CCS: Clear colourless solution;
 ND: Not detected;
 BQL: Below Quantitation limit,
 NP: Not performed;
 2EPMM: 2-ethylphenylmalonamide monohydrate;
 PBG: Phenylbutyrylguanidine;
 PEAU: Phenylethylacetylurea

[0121] Six months stability data for Phenobarbital Sodium for Injection, USP, 65 mg/vial is included in Table 10. Six months stability data for Phenobarbital Sodium for Injection, USP, 130 mg/vial is included in Table 11. Three months stability data for Phenobarbital Sodium for Injection, USP,

200 mg/vial is included in Table 12. Three months stability data for Phenobarbital Sodium for Injection, USP (Phenobarbital Sodium and 9% Saline), 200 mg/vial is included in Table 13.

TABLE 10

Product Name	Phenobarbital Sodium for Injection USP, 65 mg/vial				Batch#NPH2020 Fill Volume: 1.05 mL		Date Manufactured: Jan. 20, 2020	
Vial Make	2 mL Fiolax Clear Blow Back Vial (Schott, Article#1590892);				Date Initiated: Feb. 14, 2020			
API Mfg.	Phenobarbital Sodium, USP Grade P, C-IV: Siegfried USA, LLC Lot#1845X007;				Re-test Date: Nov. 16, 2023			
Stopper Used	Aptar stelmi 13 mm Grey Bromobutyl 2 leg Lyophilization Rubber stopper (C1621 6730GC 6 TP3)				Flip Off Seals: Wheaton, 13 mm Red Plastic top flip off with Aluminum Seal, Lot#224202			
Configuration	Inverted		Stability		25° C., 60% RH (RT) & 40° C., 75% RH (ACC)			
		Initial DOA:	1 Month DOA: Mar. 29, 2020		3 Months DOA: May 20, 2020		6 Months DOA: Sep. 4, 2020	
Test	Spec.	Feb. 18, 2020	RT	ACC	RT	ACC	RT	ACC
Description	WLP	WLP	WLP	WLP	WLP	WLP	WLP	WLP
Appearance	CCS	CCS	CCS	CCS	CCS	CCS	CCS	CCS
pH (65 mg/mL)	9.2-10.2	9.9	9.9	9.9	9.9	9.9	9.5	9.5
% Assay	90.0%-105.0%	98.6	301.6	101.5	101.3	100.5	100.0	101.2
Phenobarbital Sodium Related Substances (% Area) 2EPMM	NMT 0.2%	BQL	ND	ND	ND	ND	ND	ND
PBG	NMT 0.2%	ND	ND	ND	ND	ND	ND	ND
PEAU	NMT 0.2%	BQL	BQL	BQL	BQL	BQL	BQL	BQL
Any unspecified degradation product	NMT 0.2%	BQL	BQL	BQL	BQL	BQL	BQL	BQL
Total degradation products	NMT 1.5%	BQL	BQL	BQL	BQL	BQL	BQL	BQL

TABLE 10-continued

Water Content	NMT 3.0%	0.6	0.9	1.3	1.4	1.5	1.5	2.1
by Karl Fischer								
Note Book Reference		192039-150 to 158	192046-23 to 45	192045-135 to 153	192051-84 to 122			

DOA: Date of Analysis;
WLP: White Lyophilized Powder;
CCS: Clear Colorless Solution, 1 minute after reconstitution;
ND: Not Detected;
NMT: Not More Than;
BQL: Below Quantitation Limit (0.05%)
2EPMM: 2-Ethyl-2-Phenylmalonamide, monohydrate;
PBG: α -Phenylbutyrylguanidine;
PEAU: Phenylethylacetylurea

TABLE 11

Product Name	Phenobarbital Sodium for Injection USP,	Batch#NPH2020	Date Manufactured:
	65 mg/vial	Fill Volume: 1.05 mL	Jan. 20, 2020
Vial Make	2 mL Fiolax Clear Blow Back Vial (Schott, Article#1590892);		Date Initiated:
			Feb. 14, 2020
API Mfg.	Phenobarbital Sodium, USP Grade P, C-IV: Siegfried USA, LLC Lot#1845X007; Re-test Date: Nov. 16, 2023		
Stopper Used	Aptar stelmi 13 mm Grey Bromobutyl 2 leg Lyophilization	Flip Off Seals: Wheaton, 13 mm	
	Rubber stopper (C1621 6730GC 6 TP3)	Red Plastic top flip off with	
Configuration	Inverted	Aluminum Seal, Lot#224202	
	Stability	25° C., 60% RH (RT) & 40° C.,	
		75% RH (ACC)	

Test	Spec.	Initial DOA: Feb. 18, 2020	1 Month DOA: Mar. 23, 2020		3 Months DOA: May 20, 2020		6 Months DOA: Sep. 4, 2020	
			RT	ACC	RT	ACC	RT	ACC
Description	WLP	WLP	WLP	WLP	WLP	WLP	WLP	WLP
Appearance	CCS	CCS	CCS	CCS	CCS	CCS	CCS	CCS
pH (65 mg/mL)	9.2-10.2	9.9	9.9	9.9	9.9	9.9	9.5	9.5
% Assay	90.0%-105.0%	98.6	101.6	101.5	101.3	100.5	100.0	101.2
Phenobarbital Sodium Related Substances (% Area)								
2EPMM								
PBG	NMT 0.2%	ND	ND	ND	ND	ND	ND	ND
PEAU	NMT 0.2%	BQL	BQL	BQL	BQL	BQL	BQL	BQL
Any unspecified degradation product	NMT 0.2%	BQL	BQL	BQL	BQL	BQL	BQL	BQL
Total degradation products	NMT 1.5%	BQL	BQL	BQL	BQL	BQL	BQL	BQL
Water Content	NMT 3.0%	0.6	0.9	1.3	1.4	1.5	1.5	2.1
by Karl Fischer								
Note Book Reference		192039-150 to 158	192045-23 to 45	192045-135 to 153	192051-84 to 122			

DOA: Date of Analysis;
WLP: White Lyophilized Powder;
CCS: Clear Colorless Solution, 1 minute after reconstitution;
ND: Not Detected;
NMT: Not More Than;
BQL: Below Quantitation Limit (0.05%)
2EPMM: 2-Ethyl-2-Phenylmalonamide, monohydrate;
PBG: α -Phenylbutyrylguanidine;
PEAU: Phenylethylacetylurea

TABLE 12

Product Name	Phenobarbital Sodium for Injection USP, 200 mg/vial	Batch#NPH2031A	Date Manufactured:
Vial	10 mL Tubular Serum Bottle Clear	Fill Volume: 2.1 mL	Jun. 3, 2020
Make	Wheaton, Part# 223586; Batch#5192875);		Date Initiated:
			Jun. 9, 2020

TABLE 12-continued

API Mfg.	Phenobarbital Sodium, USP Grade P, C-IV: Siegfried USA, LLC Lot#1924X009; Re-test Date: Jul. 20, 2024			
Stopper Used	Datwyler 20 mm Igloo Bromobutyl Lyophilization Rubber stopper			
Configuration	Inverted	Stability	Flip Off Seals: Wheaton, 20 mm Red Plastic top flip off with Aluminum Seal, Lot#224203 25° C., 60% RH (RT) & 40° C., 75% RH (ACC)	
Test	Spec.	Initial DOA:	3 Months DOA: Sep. 4, 2020	
		Jun. 5, 2020	RT	ACC
Description	WLP	WLP	WLP	WLP
Appearance	CCS	CCS	CCS	CCS
pH (20 mg/mL)	9.2-10.2	9.9	9.6	9.7
% Assay	90.0-105.0	100.1	98.1	101.0
Phenobarbital Sodium Related Substances (% Area) 2EPM	NMT 0.2%	ND	ND	ND
PBG	NMT 0.2%	ND	ND	ND
PEAU	NMT 0.2%	BQL	0.03	0.04
Any unspecified degradation product	NMT 0.2%	BQL	0.06	0.05
Total degradation products	NMT 1.5%	BQL	0.09	0.09
Water Content by Karl Fischer	NMT 3.0%	1.1	1.0	1.3
Note Book Reference		192045-163 to 189	192051-84 to 122	

DOA: Date of Analysis;
WLP: White Lyophilized Powder;
CCS: Clear Colorless Solution, 1 minute after reconstitution;
ND: Not Detected;
NMT: Not More Than;
BQL: Below Quantitation Limit (0.05%)
2EPM: 2-Ethyl-2-Phenylmalonamide, monohydrate;
PBG: α -Phenylbutyrylguanidine;
PEAU: Phenylethylacetylurea

TABLE 13

Product Name	Phenobarbital Sodium for Injection USP, 200 mg/vial	Batch#NPH2031A	Date Manufactured:	
Vial	10 mL Tubular Serum Bottle Clear	Fill Volume: 2.1 mL	Jun. 3, 2020	
Make	Wheaton, Part# 223586; Batch#5192875);		Date Initiated:	
API Mfg.	Phenobarbital Sodium, USP Grade P, C-IV: Siegfried USA, LLC Lot#1924X009; Re-test Date: Jul. 20, 2024		Jun. 9, 2020	
Stopper Used	Datwyler 20 mm Igloo Bromobutyl Lyophilization Rubber stopper		Flip Off Seals: Wheaton, 20 mm Red Plastic top flip off with Aluminum Seal, Lot#224203	
Configuration	Inverted	Stability	25° C., 60% RH (RT) & 40° C., 75% RH (ACC)	
		Initial DOA:	3 Months DOA: Sep. 4, 2020	
Test	Spec.	Jun. 5, 2020	RT	ACC
Description	WLP	WLP	WLP	WLP
Appearance	CCS	CCS	CCS	CCS
pH (20 mg/mL)	9.2-10.2	9.8	9.5	9.5
% Assay	90.0-105.0	103.3	102.8	103.6
Phenobarbital Sodium Related Substances (% Area) 2EPM	NMT 0.2%	BQL	ND	ND

TABLE 13-continued

PBG	NMT 0.2%	ND	ND	ND
PEAU	NMT 0.2%	BQL	0.03	0.04
Any unspecified degradation product	NMT 0.2%	BQL	0.06	0.05
Total degradation products	NMT 1.5%	BQL	0.1	0.08
Water Content by Karl Fischer	NMT 3.0%	0.8	0.9	1.1
Note Book Reference		192045-163 to 189	192051-84 to 122	

DOA: Date of Analysis;

WLP: White Lyophilized Powder;

CCS: Clear Colorless Solution, 1 minute after reconstitution;

ND: Not Detected;

NMT: Not More Than;

BQL: Below Quantitation Limit (0.05%);

2EPMM: 2-Ethyl-2-Phenylmalonamide, monohydrate;

PBG: α -Phenylbutyrylguanidine;

PEAU: Phenylethylacetylurea

[0122] Data for Osmolality, Particulate matter and assay for Sodium Chloride is disclosed in Table 14.

TABLE 14

Phenobarbital Sodium for Injection, USP, 200 mg/vial in Water or Sodium Chloride - Osmolality, Particulate Matter and Assay for Sodium Chloride.				
Batch No	Osmolality* (mOsmol/Kg) (Specification: To be	Particulate Matter (By Light Obscuration method) Ref: USP<788>		Assay of Sodium Chloride (titrimetric; Ref USP monograph for Sodium
	reported)	$\geq 10 \mu\text{M}$ Specification: (NMT 6000 particles/vial)	$\geq 25 \mu\text{M}$ Specification: (NMT 600 particles/vial)	Chloride Injection Specification: (95.0%-105.0%)
	Reconstitution in 10 mL (20 mg/mL)			
NPH2031A (200 mg/vial)	148	80.0	0.67	NA
NPH2031B (200 mg/vial)	372	171	1.33	99.99
(Phenobarbital Sodium 200 mg, 9% Sodium Chloride)				
Water-Blank	0	NA	NA	NA
0.9% Sodium Chloride	270	NA	NA	NA

[0123] Lyophilization parameters used for the 65 mg/vial, 100 mg/vial, 130 mg/vial and 200 mg/vial are disclosed in Table 15.

TABLE 15

Lyophilization Cycle parameters.								
Section Phase		Lyophilization Cycle 1*			Lyophilization Cycle 2**			
		Condenser set point C.	Vacuum [mTorr]	Temperature ° C.	Hold Time [mm]	Condenser set point ° C.	Vacuum [mTorr]	Temperature ° C.
1	Loading		off	5	n.a.		off	-5
2	Freezing		off	-5	5		off	-5
3	Freezing		off	-50	60		off	-52
								120 (Ramp)

TABLE 15-continued

Lyophilization Cycle parameters.									
Section	Phase	Lyophilization Cycle 1*				Lyophilization Cycle 2**			
		Condenser set point ° C.	Vacuum [mTorr]	Temperature ° C.	Hold Time [mm]	Condenser set point ° C.	Vacuum [mTorr]	Temperature ° C.	Hold Time [mm]
4	Freezing	-50	off	-10	30	-60	off	-52	180 (Hold)
5	Freezing		off	-50	120		n.a.	n.a.	n.a.
6	Evacuation		75	-50	30		50	-45	30
7	Primary drying		75	-50	400 (Hold)		50	-42	100 (Ramp)
8	Primary drying		75	-45	350 (Hold)		50	-42	1120 (Hold)
9	Primary drying		75	-43	300 (Hold)		n.a.	n.a.	n.a.
10	Primary drying		75	-35	250 (Hold)		n.a.	n.a.	n.a.
11	Primary drying		75	-20	60 (Hold)		n.a.	n.a.	n.a.
12	Primary drying		75	0	120 (Hold)		n.a.	n.a.	n.a.
13	secondary drying		75	50	120 (Hold)		50	35	100 (Ramp)
14	secondary drying	75	25	120 (Hold)	50	35	420 (Hold)		

Lyophilization Cycle 1*: 65 mg/vial, 130 mg/vial and 100 mg/vial samples disclosed in Tables 1, 2, 3, 16 and 17

Lyophilization Cycle 2**: NPH2031A (200 mg/vial Phenobarbital Sodium) and NPH2031B (200 mg/vial in Phenobarbital Sodium and 90 mg/vial Sodium Chloride, NPH2020 (65 mg/vial) and NPH2021 (130 mg/vial)

[0124] Six months stability data for Phenobarbital Sodium for Injection USP, 100 mg/vial is included in Table 16.

TABLE 16

Product Name	Phenobarbital Sodium for Injection USP, 100 mg/vial	Batch#NPH1980				Date Manufactured: Sep. 25, 2019			
Vial Make	5 mL Clear Untreated Article#9621139054					Date Initiated: Sep. 30, 2019			
API Mfg.	Phenobarbital Sodium, USP Grade P, C-IV: Siegfried USA, LLC Lot#1845X006; Re-test Date: Nov. 16, 2023								
Stopper Used	Aptar stelmi 13 mm Grey Bromobutyl 2 leg Lyophilization Rubber stopper, Lot#G702/91884					Flip Off Seals: Wheaton, 13 mm Red Plastic top flip off with Aluminum Seal, Lot# NA			
Configuration	Inverted	Stability				25° C., 60% RH (RT) & 40° C., 75% RH (ACC)			

Test	Spec.	1 Month								3 Months				6 Months			
		Initial		RT		ACC		RT		ACC		RT		ACC		RT	
		UT	T	UT	T	UT	T	UT	T	UT	T	UT	T	UT	T	UT	T
Description	WLP	WLP	WLP	WLP	WLP	WLP	WLP	WLP	WLP	WLP	WLP	WLP	WLP	WLP	WLP	WLP	WLP
Appearance	CCS	CCS	CCS	CCS	CCS	CCS	CCS	CCS	CCS	CCS	CCS	CCS	CCS	CCS	CCS	CCS	CCS
pH	9.2-10.2	9.7	9.7	9.6	9.7	9.6	9.5	9.7	9.7	9.7	9.7	9.7	9.8	9.9	9.8	9.8	9.9
% Assay	90.0%-105.0%	98.4	97.8	98.6	99.0	98.0	100.4	99.1	98.7	97.0	96.9	97.7	98.3	97.2	97.6		
Phenobarbital Sodium Related Substances (% Area) 2EPM	NMT 0.2%	ND	ND	ND	ND	ND	ND	BQL	BQL	BQL	BQL	ND	ND	ND	ND		
PBG	NMT 0.2%	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND		
PEAU	NMT 0.2%	BQL	BQL	BQL	BQL	BQL	BQL	BQL	BQL	BQL	BQL	BQL	BQL	BQL	BQL		
Any unspecified degradation product	NMT 0.2%	0.05	0.06	0.05	0.05	0.05	0.05	BQL	BQL	BQL	BQL	BQL	BQL	BQL	BQL		
Total degradation products	NMT 1.5%	0.06	0.06	0.05	0.05	0.05	0.05	BQL	BQL	BQL	BQL	BQL	BQL	BQL	BQL		

TABLE 16-continued

Water Content	NMT 3.0%	0.7	0.6	1.0	1.0	1.1	1.3	1.2	1.1	1.6	1.5	1.6	1.2	1.5	2.0
by Karl Fischer															
Note Book Reference		192031-91 to 101		192031-169 to 183				192039-44 to 52				192045-52 to 85			

WLP: White Lyophilized powder;

CCS: Clear Colorless Solution, 1 minute after reconstitution;

UT: Untreated;

T: Treated

ND: Not Detected;

NMT: Not More Than;

2EPMM: 2-Ethyl-2-Phenylmalonamide, monohydrate;

PBG: α -Phenylbutyrylguanidine;

TABLE 17

Phenobarbital Sodium for Injection, USP, 100 mg/vial Formulation Details.				
Ingredients	Manufacturer/Lot number	Concentration (mg/mL)	Concentration (mg/vial)	Batch Quantity (100 mL)
Phenobarbital Sodium, USP Grade P C-IV	Siegfried USA, LLC Mat No: 300040 Lot# 1845X006 Retest date: Nov. 16, 2023	101.13 mg (~100 mg Phenobarbital Sodium after LOD correction	100 mg	10.113 g
Water for Injection	In house	Q.S. to 1 mL	NA	Q.S. to 100 mL or 103.3 g
100% Nitrogen 5 mL, untreated	Airgas Gerresheimer Article#9621139054	NA	NA	NA
5 mL, treated	Gerresheimer Article#9621130486	NA	NA	NA
13 mm Bromobutyl Lyophilization stoppers	Aptar-Stelmi C16216720GC6TP3	NA	NA	NA
Seal: 13 mm Seal, Flip-Cap, Red	Wheaton Unlined	NA	NA	NA

Diluents that can be used: Water for Injection, 0.9% Saline or 5% Dextrose
Proposed Reconstitution Volume: 5 mL

[0125] Representative chromatograms and/or corresponding calculations are shown in FIGS. 2A-211, 3A-3L, 4A-4L, and Tables 2 and 3.

[0126] X-Ray Powder Diffraction. With reference to FIGS. 5A-5D, XRPD was performed using the Rigaku Smart-Lab X-ray diffraction system configured for reflection Bragg-Brentano geometry using a line source X-ray beam. The x-ray source is a Cu Long Fine Focus tube that was operated at 40 kV and 44 mA. That source provides an incident beam profile at the sample that changes from a narrow line at high angles to a broad rectangle at low angles. Beam conditioning slits were used on the line X-ray source to ensure that the maximum beam size is less than 10 mm both along the line and normal to the line. The Bragg-Brentano geometry is a para-focusing geometry controlled by passive divergence and receiving slits with the sample itself acting as the focusing component for the optics. The inherent resolution of Bragg-Brentano geometry is governed in part by the diffractometer radius and the width of the receiving slit used. Typically, the Rigaku Smart-Lab is operated to give peak widths of $0.1^\circ 2\theta$ or less. The axial divergence of the X-ray beam is controlled by 5.0-degree Soller slits in both the incident and diffracted beam paths. The samples were placed in a low-background, silicon holder using light manual pressure to keep the sample surfaces flat and level with the reference surface of the holder. The samples were analyzed from 2 to $40^\circ 2\theta$ using a continuous scan of $6^\circ 2\theta$ per minute with an effective step size of $0.02^\circ 2\theta$.

[0127] The recitation of ranges of values herein is merely intended to serve as a shorthand method of referring individually to each separate value falling within the range. Unless otherwise indicated herein, each individual value is incorporated into the specification as if it were individually recited herein. All methods described herein can be performed in any suitable order unless otherwise indicated herein or otherwise clearly contradicted by context. The use of any and all examples, or exemplary language (e.g., "such as") provided with respect to certain embodiments herein is intended merely to better illuminate the full scope of the present disclosure, and does not pose a limitation on the scope of the invention otherwise claimed. No language in the specification should be construed as indicating any non-claimed element essential to the practice of the claimed invention.

[0128] It should be apparent to those skilled in the art that many more modifications besides those already described are possible without departing from the full scope of the concepts disclosed herein. The disclosed subject matter, therefore, is not to be restricted except in the scope of the appended claims. Moreover, in interpreting both the specification and the claims, all terms should be interpreted in the broadest possible manner consistent with the context. In particular, the terms "comprises" and "comprising" should be interpreted as referring to elements, components, or steps in a non-exclusive manner, indicating that the referenced elements, components, or steps may be present, or utilized,

or combined with other elements, components, or steps that are not expressly referenced. Where the specification claims refers to at least one of something selected from the group consisting of A, B, C . . . and N, the text should be interpreted as requiring only one element from the group, not A plus N, or B plus N, etc.

What is claimed is:

1. A pharmaceutical composition comprising lyophilized amorphous Phenobarbital Sodium.

2. A method of producing a storage-stable form of lyophilized Phenobarbital sodium that forms no more than 0.2% phenylethylacetylurea (PEAU) when reconstituted in an aqueous solution, the method comprising:

adding Phenobarbital Sodium to water to form a Phenobarbital Sodium solution;

if necessary, adjusting the pH of the Phenobarbital Sodium solution to or between 9.2 to 10.2; and
lyophilizing the Phenobarbital Sodium solution.

3. The method of claim 2, wherein the water is Water for Injection.

4. The method of claim 3, wherein the Water for Injection is sparged with nitrogen prior to adding the Phenobarbital Sodium.

5. The method of claim 4, wherein the Water for Injection is sparged with nitrogen for at least 30 minutes prior to adding the Phenobarbital Sodium.

6. The method of claim 2, wherein the water is cooled to or between 2 to 8° C. prior to adding the Phenobarbital Sodium.

7. The method of claim 2, wherein the adjusting comprises measuring the pH of the Phenobarbital Sodium solution, and wherein if the pH is higher than 10.2, the adjusting comprises adding Hydrochloric acid (HCl).

8. The method of claim 2, wherein the lyophilized Phenobarbital Sodium forms no more than 0.1% phenylethylacetylurea (PEAU) when reconstituted in an aqueous solution after storage of the lyophilized Phenobarbital Sodium for up to 6 months at room temperature (RT) and accelerated stability conditions (ACC).

9. The method of claim 2, wherein the lyophilized Phenobarbital Sodium forms no more than 0.05% phenylethylacetylurea (PEAU) when reconstituted in an aqueous solution after storage of the lyophilized Phenobarbital Sodium for up to 6 months at room temperature (RT) and accelerated stability conditions (ACC).

10. The method of claim 2, wherein the lyophilized Phenobarbital Sodium forms no detectable amount of 2-ethyl-2-phenylmalonamide (2EPM) or alpha-phenylbutyrylguanidine (PBG) when reconstituted in an aqueous solution after storage of the lyophilized Phenobarbital Sodium for up to 6 months at room temperature (RT) and accelerated stability conditions (ACC).

11. The method of claim 2, wherein the lyophilized Phenobarbital Sodium is stable up to at least 6 months at accelerated stability conditions (ACC) which is equivalent to 24 months at room temperature (RT).

12. The method of claim 11, wherein the stability of the lyophilized Phenobarbital Sodium is determined by adding the lyophilized Phenobarbital Sodium to an aqueous solution to form a reconstituted lyophilized Phenobarbital Sodium and analyzing to quantitate the amount of phenylethylacetylurea (PEAU), 2-ethyl-2-phenylmalonamide (2EPM), and/or alpha-phenylbutyrylguanidine (PBG) in the reconstituted lyophilized Phenobarbital Sodium.

13. The method of claim 2, wherein the Phenobarbital Sodium solution is at a concentration of 65 mg/mL, 100 mg/mL, 130 mg/mL or 200 mg/mL.

14. The method of claim 2, further comprising filtering the Phenobarbital Sodium solution prior to lyophilizing.

15. The method of claim 2, further comprising:

aliquoting the Phenobarbital Sodium solution into a vial prior to lyophilization; and

after lyophilization, applying a closure to the vial under vacuum.

16. A method of treating an individual in need of Phenobarbital Sodium, the method comprising:

adding saline or dextrose to a composition comprising lyophilized amorphous Phenobarbital Sodium immediately prior to administration to form a Phenobarbital Sodium solution; and

administering the Phenobarbital Sodium solution to the individual.

17. The method of claim 16, wherein the individual suffers from epilepsy.

18. The method of claim 17, wherein the individual is a newborn suffering from neonatal epilepsy.

19. The method of claim 16, wherein the composition is a powder dose of 65 mg, 100, 130 mg or 200 mg.

20. The method of claim 16, wherein the administering comprises intramuscular injection or intravenous injection.

* * * * *