0163-769X/86/0701-0044$02.00/0
Endocrine Reviews
Copyright © 1986 by The Endocrine Society
Vol. 7, No. 1
Printed in U.S.A.

Gonadotropin-Releasing Hormone Analog Design. Structure-Function Studies Toward the Development of Agonists and Antagonists: Rationale and Perspective

MARVIN J. KARTEN AND JEAN E. RIVIER

Contraceptive Development Branch (M.J.K.), Center for Population Research, National Institute of Child Health and Human Development, National Institutes of Health, Bethesda, Maryland 20892; and The Clayton Foundation Laboratories for Peptide Biology (J.E.R.), The Salk Institute, La Jolla, California 92037

Downloaded from https://academic.oup.com/edrv/article/7/1/44/2548841 by University of California, Los Angeles user on 19 April 2025

Introduction

ON JUNE 24, 1971, Andrew V. Schally announced the determination of the primary structure of porcine GnRH at The Endocrine Society Meeting in San Francisco. This announcement was followed by publications by Matsuo *et al.* (1, 2) and Baba *et al.* (3) on the proposed amino acid sequence for porcine GnRH and its synthesis and by Burgus *et al.* (4) who characterized ovine GnRH and found the sequence to be identical with that of porcine GnRH. The physiological and therapeutic importance attributed to the discovery of the new substance was greatly increased by the prospect of the design of potent and long acting GnRH agonists and antagonists. Since that time more than 2000 analogs of GnRH have been synthesized. The impact of research of GnRH and its analogs on clinical medicine recently led Ziporyn (5) to note, "There's almost no subspecialty of medicine that will be left untouched by the [research] advances associated with LHRH or its analogs." It is the intent of this article to provide a historical review of the major, and some minor, aspects of the chemical development of GnRH agonists and antagonists up to the present state of development (July 1, 1985). The synthetic chemical efforts have been devoted largely to increasing the affinity of the peptides to the GnRH receptor and their resistance to degradation or elimination in *in vivo* systems, characteristics which, for the GnRH analogs, are generally interrelated.

An annual compilation and review of structure-activity relationships of GnRH analogs is available in the Specialist Periodical Reports which cover the literature published during 1971–1980 (volumes 4–13) (6–15). At irregular intervals, Annual Reports in Medicinal Chemistry furnish brief updates of the studies of structure-activity relationships of GnRH analogs (16–21). These reports, in conjunction with two recently published comprehensive monographs should provide the reader with a balanced account of the development of GnRH analogs (22–29).

Synthesis, Purification, and Characterization of GnRH Analogs

It is important to note that the rapid development of GnRH analogs was made possible through the extensive use of solid phase peptide synthesis (SPPS) introduced by Merrifield (30). As one of the codevelopers of the method (31), Stewart (32) has pointed out that the use of automated equipment for SPPS, benzhydrylamine-like resins for peptide amide synthesis (33, 34), and adequate methods for the purification of peptides, particularly reverse phase HPLC (RP-HPLC) in recent years (35), have made the synthesis of mammalian GnRH, <Glu–His–Trp–Ser–Tyr–Gly–Leu–Arg–Pro–Gly–NH_2, and its analogs, a relatively simple task. Although classical (solution) methods have been employed for the synthesis of GnRH (see Ref. 36 and references therein) and its analogs (37), it is quite clear that the use of SPPS and RP-HPLC were essential for the rapid exploration of structure-activity relationships as well as providing investigators with relatively large amounts of these substances for pharmacological, toxicological, and clinical studies.

While the purity of the agonists synthesized either by SPPS or classical (solution) methods was always of concern in terms of the concomitant biological activity of potential racemization products, it was a critical factor in the biological evaluation of the antagonists. This was particularly true in the early stages of development when the GnRH inhibitory activities were very low and could be masked by a small amount of racemized material

Address requests for reprints to: Dr. Marvin J. Karten, Contraceptive Development Branch, Center for Population Research, National Institute of Child Health and Human Development, National Institutes of Health, Bethesda, Maryland 20892.

EMPOWER - 1021

present as a contaminant and acting as an agonist. For example, the high GnRH potency of [D-Ala^6]GnRH (350–450% of GnRH) would barely be affected by a 10% racemization contaminant of [L-Ala^6]GnRH (4% of GnRH) (38). However, a very weak antagonist, *e.g.* [D-Phe^2]GnRH, (39) could have its activity masked by the presence of small amounts of [L-Phe^2]GnRH (40) with only 2–4% of the potency of GnRH but with full intrinsic activity *in vitro*. The separation of the possible diastereomers, *e.g.* [D-Ala^6]GnRH and [L-Ala^6]GnRH, which could result from racemization, was eventually made feasible through the use of HPLC, thus eliminating one element of uncertainty from the interpretation of the biological results (35). Similarly, a preparation of [D-His^2]GnRH exhibiting 10% GnRH-like potency (39) was subsequently shown, when purified by HPLC, to be inactive either as an agonist or an antagonist (41). Racemization of histidine during peptide synthesis is well documented and difficult to prevent irrespective of whether classical methods of synthesis or SPPS are used. The widespread use of HPLC resulted from the recognition that classical methods of purification were inferior to HPLC as a tool for the separation of these diastereomeric peptides. Most GnRH analogs reported in the literature have been characterized by amino acid analysis only, often without a quantitative determination of the unnatural amino acids. Investigators have relied on the presumed authenticity of the starting protected amino acid and on high coupling efficiency during the assembling of the peptides on the resins rather than pursuing rigorous methods of characterization of the peptides. Nuclear magnetic resonance (NMR) and mass spectrometry were employed in those cases where definite proof of structure was required. Optical rotations were generally measured, and TLC and HPLC, in several systems, were used for proof of homogeniety.

Development of GnRH Agonists

The original incentive for the development of more potent GnRH agonists was the expectation that the knowledge of the LH-releasing and ovulation-inducing effects of GnRH observed in laboratory animals could be applied to the treatment of male and female infertility (42). However, the half-life of GnRH is very short (43, 44) and more potent and longer acting analogs were thought to be necessary for practical clinical utility, regardless of any anticipated or unanticipated therapeutic applications. Potent agonists, referred to as superagonists, were rapidly produced and were subsequently discovered, along with GnRH, to have, ironically, antireproductive effects. They were available for reproductive pharmacological evaluation (42, 45, 46) within 3 yr of the structure elucidation of GnRH; this accounts for their rapid clinical exploration exemplified by the first demonstration (in 1978) of inhibition of reproductive function in women by a superagonist (47). Once superagonists had been synthesized and their potential therapeutic value was recognized, further incentive, after 1976, to seek structurally novel and longer acting agonists was provided by promising commercial considerations.

The various biological assays and animal models that have been utilized for the testing of GnRH agonists have been recently reviewed (42, 48). The most widely used *in vitro* assays have been, initially, the dispersed pituitary cell for the measurement of LH and FSH secretions and, more recently, the receptor binding assay using purified pituitary membrane fractions for the estimation of the potencies of the analogs (49, 50). *In vivo* biological assays, which have been utilized to determine the potencies of GnRH agonists, include induction of ovulation, disruption of the estrus cycle, stimulation of uterine growth, inhibition of pregnancy, stimulation of LH release in ovariectomized rats, and stimulation of LH/FSH release in immature rats using an infusion technique (42, 48).

Pro^9-ethylamide (NEt) modifications

The first important structural modification of GnRH leading to increased potency was discovered by Fujino and co-workers (37), who examined the effect of structural modifications at the C terminus of GnRH. Although the des-amide of GnRH (GnRH free acid) exhibited very low GnRH potency in ovariectomized rats (51) and Pro^9-GnRH showed only 10% of the potency of GnRH *in vitro* (34), replacement of the glycine amide terminus with alkyl amines produced nonapeptide alkyl amides with significantly greater ovulation-inducing potency than GnRH itself (37). Thus, [Pro^9-ethylamide (NEt)]GnRH, the most potent analog of the series, was 5 times more potent than GnRH and more potent than either the Pro^9-methylamide (NHMe) or Pro^9-propylamide (NHPr) modifications. The fluorinated ethylamide analog, [Pro^9-NHCH$_2$CF$_3$]GnRH was subsequently reported by Coy *et al.* (52) to be twice as potent as [Pro^9-NEt]GnRH in releasing LH when administered to immature male rats. The data of Fujino *et al.* (37), it was noted, suggested that the terminal glycine amide was not essential for high potency and that the total chain length of the peptide played an important role in the binding affinity of the analog for the pituitary receptor. It was also suggested that the introduction of this Pro-alkylamide moiety may also increase the duration of action of these analogs by virtue of their greater resistance to enzymatic degradation (53). These two desired properties, greater binding affinity and enzymatic resistance to proteolysis, were to become the basis for the rational

Downloaded from https://academic.oup.com/edrv/article/7/1/44/2548841 by University of California, Los Angeles user on 19 April 2025

design and for the explanation of activity, or lack thereof, of all the GnRH analogs regardless of the site of structural modification. The concept that protection from renal elimination would also lead to prolonged action was eventually incorporated into the design of GnRH analogs.

D-Xaa^6 modifications

The second important structural modification of GnRH, discovered by Monahan *et al.* (38), was the replacement of the Gly^6 residue with D-alanine yielding [D-Ala^6]GnRH with a potency of approximately 350–450% that of GnRH both *in vitro* and in ovariectomized rats. The corresponding [L-Ala^6]GnRH had only 4% of the potency of GnRH, and it was suggested that since the potencies determined *in vivo* were in agreement with the *in vitro* results, it was unlikely that the differences in the biological activities could be solely explained by differences in clearance rates. Instead, the increased biological potency was attributed to the conformational stabilizing effect of the D-alanine which was favorable for binding (and activity) at the receptor. This study is also noteworthy for its suggestion that GnRH may conformationally contain a β-II type bend (involving Ser–Tyr–Gly–Leu) which is stabilized in [D-Ala^6]GnRH and preferable for binding at the receptor site. This point will be discussed below when the contribution of conformational studies to the design of GnRH analogs, particularly antagonists, is reviewed. Also to be deferred for later discussion is our knowledge of the enzymatic degradation of peptides and its contribution to the design (if any) of GnRH analogs. However, it should be noted that, regardless of whether the Gly^6–Leu^7 bond or the Tyr^5-Gly^6 bond is considered to be a major site of proteolytic cleavage, substitution of glycine by D-amino acids is likely to render either linkage more resistant to enzymatic degradation (26).

Additive effects (or lack thereof) of Pro^9-NEt and D-Xaa^6 modifications

It is often assumed that the biological effect of combining several structural changes in one molecule will be additive (or, more correctly, multiplicative). Thus, according to this additivity rule (54), if one structural modification leads to a relative potency of *a* and a second modification to a relative potency of *b* then the combination of both structural modifications in a single molecule would be expected to yield an analog with a biological potency of *a* multiplied by *b*. The potential additivity of biological potency of the Pro^9-NEt and D-Ala^6 modifications was immediately tested and, in rapid succession, two reports [Coy *et al.* (55) and Fujino *et al.* (56)] on this important combination appeared. Infusion experiments with immature male rats showed that the three analogs, [D-Ala^6,Pro^9-NEt]GnRH, [D-Ala^6]GnRH, and [Pro^9-NEt]GnRH, had LH/FSH releasing potency ratios of 12–16:7–8:2.5, respectively, compared with GnRH (55). Similar ovulation-inducing potency ratios were observed (56) among the three analogs, but the potencies relative to GnRH were much higher. These *in vivo* results correlated with the *in vitro* results of Vale *et al.* (57) who, using the stimulation of LH secretion from rat pituitary cells in culture by GnRH agonists as an index of potency, noted that the combination of the two structural modifications yielded an analog with a potency approximately equal to the product of the potencies of the individual modifications. The *in vitro* results of Fujino *et al.* (56) did not agree with these findings. Although it is self-evident that comparisons of biological data by different groups are valid only if the same bioassays are employed in precisely the same manner (48), nevertheless, many of the apparent disagreements in the data reported, regarding comparisons of potencies of superagonists, can be attributed to a disregard of this axiom. Thus, the validity of the extension of the additivity principle to the combination of a D-aromatic amino acid in position 6 and the Pro^9-NEt modification became a focal point of interest with the publication of apparently conflicting reports on this subject (57–59). The observation was made that agonists with D-aromatic amino acids such as [D-Trp^6]GnRH and [D-Trp^6,Pro^9-NEt]GnRH are much more potent (36 times and 144 times the *in vitro* potency of GnRH, respectively) than the corresponding substitutions with aliphatic amino acids, such as [D-Ala^6]LHRH and [D-Ala^6,Pro^9-NEt]GnRH, which are approximately 4 times and 14 times, respectively, the potency of GnRH (57). The subsequent binding studies of Perrin *et al.* (50) showed increased binding potencies for the Pro^9-NEt modifications as compared with [D-Ala^6]GnRH or [D-Trp^6]GnRH; however, when compared with the biological potencies, in stimulating LH secretion *in vitro*, the increases were far less dramatic. The data for the D-Trp^6 analogs are in accord with the subsequent binding studies of Barron *et al.* (60) but are not in accord with the earlier *in vitro* data reported by Coy *et al.* (58). *In vivo* measurements in immature rats comparing [D-Leu^6]GnRH and [D-Leu^6,Pro^9-NEt]GnRH (Leuprolide), using integrated levels of LH over a 6-h period after injection, showed a similar additive effect (61), but the additive effect was reported not to hold true for the corresponding D-aromatic amino acid modifications in this same assay system (59). Thus, in male rats, the Pro^9-NEt modification was reported to decrease the potency of [D-Phe^6]GnRH and [D-Trp^6]GnRH by a factor of nearly 2 (59). However, it has been more recently reported that, using estrus suppression (62) as an index of agonist activity, [D-Trp^6]GnRH and [D-Trp^6,Pro^9-NEt]

Downloaded from https://academic.oup.com/edrv/article/7/1/44/2548841 by University of California, Los Angeles user on 19 April 2025

GnRH were equipotent. Postcoital comparisons in rats (Naqvi, R. and M. Lindberg, unpublished observations) also indicated that the two analogs were equipotent. In women, it was reported that the two analogs were equipotent with respect to the sc dose required for maximal LH release (63). Barron *et al.* (60) showed that MCRs in pregnant women were similar for both analogs. They concluded that since the NEt residue, which is reported to protect the peptide from postproline-cleaving enzyme activity, did not lead to a prolonged survival time in pregnant women, degradation by this enzyme in human tissues contributes minimally to GnRH clearance. Support for this conclusion was found in reports that both analogs were equipotent *in vivo* (42, 64). Thus, the overwhelming evidence points to *in vivo* equipotency for the two D-Trp^6 analogs irrespective of the *in vitro* results and binding studies supporting the additive effects on the biological potency of the D-Trp^6 and Pro^9-NEt modifications.

Hydrophobic modifications at position 6

The trend toward seeking more potent agonists with increasing hydrophobic character resulted in the addition of two more superagonists to the growing list of analogs available for clinical exploration. [N^τ-Bzl-D-His^6,Pro^9-NEt]GnRH (Histerelin) was designed by Rivier *et al.* (22) to have the characteristics of high water solubility at acidic pH and greater lipophilic character *in vivo*, while retaining high biological potency. A correlation was noted between the *in vitro* potencies of certain position 6 superagonists and their HPLC retention times at physiological pH. These correlations included [N$^\tau$-Bzl-D-His^6,Pro^9-NEt]GnRH and [D-Trp^6,Pro^9-NEt]GnRH, with the former being slightly more potent than the latter, *in vitro. In vivo* results showed a similar trend (48).

It had previously been observed that the incorporation of D-amino acids, with larger and more lipophilic side chains than in [D-Leu^6]GnRH, such as [D-Trp^6]GnRH, resulted in more potent agonists (57, 59). Nadasdi and Medzihradszky (65) proposed a quantitative correlation to exist between the potency of position 6 substituted GnRH analogs and the calculated hydrophobicity of the amino acid side chain. It is accepted that increased lipophilicity of drugs is generally associated with greater retention of the drug in the body and, therefore, prolonged duration of action (66). The retention may be the result of enhanced renal reabsorbtion or fat storage of nonionized fat-soluble compounds. Protection of the drug from renal excretion, through plasma protein binding, will also affect its duration of action. Plasma protein binding generally increases, in a given series of analogs, with increasing hydrophobicity (66). With this in mind, Nestor *et al.* (62, 67) postulated that analogs with greater hydrophobicity could have an extended biological half-life resulting from a whole body depot effect. They would attribute this effect to a decreased rate of clearance of the analog from the circulation and the increased binding capacity of the analog for hydrophobic plasma carrier proteins (67). The results of the study on a wide range of hydrophobic analogs showed that the most potent ones were found in a hydrophobicity range, as measured by their retention time on RP-HPLC (68), greater than that of the D-Trp^6 analogs. As an example, [D-Nal(2)^6]GnRH (Nafarelin acetate), the most potent of this series, was reported to be 200 times more potent than GnRH in suppressing estrus in rats. It was pointed out, however, that analogs with greater hydrophobicity than Nafarelin acetate were less potent and this includes the analog incorporating the Pro^9-NEt modification into Nafarelin acetate (62). Interestingly, [D-Nal(2)^6]GnRH and [D-Nal(1)^6]GnRH were isolipophilic but the latter was 4-fold less potent. Other examples (68, 69) also bear witness to the inadequacy of using hydrophobicity alone as a prediction of agonist potency (28).

Nafarelin acetate, which was reported to be twice as potent as [D-Trp^6]GnRH, [D-Trp^6,Pro^9-NEt]GnRH, or [N^τ-Bzl-D-His^6,Pro^9-NEt]GnRH in estrus suppression comparisons (26), became the last superagonist to be made available for clinical exploration. The improvement of the pharmacokinetics with the more hydrophobic agonist may appear to be supported by the comparisons of the reported half-lives of GnRH [$t_{1/2}$ = 8 min, constant infusion, (44)], [D-Trp^6]GnRH [$t_{1/2} \simeq$ 30 min, constant infusion, (44)], and [D-Nal(2)^6]GnRH [$t_{1/2}$ = 2.4 h, sc administration, (67)]; however, comparison of the three peptides under identical conditions is not available. The considerably longer plasma elimination half-lives reported for Nafarelin acetate in rats, monkeys, and humans than reported for GnRH or [D-Leu^6,Pro^9-NEt]GnRH (Leuprolide) were attributed, at least in part, to the more extensive plasma binding of Nafarelin acetate (70).

N^ω,$N^{\omega'}$-dialkyl-D-homoarginines were incorporated into position 6 of GnRH agonists (67, 71) as a result of successful GnRH antagonist investigations with this unnatural amino acid. The most potent, [N^ω,$N^{\omega'}$-diethyl-D-Har^6,Pro^9-NEt]GnRH, was only slightly less potent than [D-Nal(2)^6]GnRH in the rat estrus suppression assay.

Other C-terminally modified analogs

Another structural modification which has generally led to increases in potency, in combination with D-amino acids in position 6, is the α-aza-Gly^{10}(-NHNHCO-) substitution. A series of α-aza analogs of GnRH were syn-

Downloaded from https://academic.oup.com/edrv/article/7/1/44/2548841 by University of California, Los Angeles user on 19 April 2025

thesized by Dutta *et al.* (72, 73) with the expectation that the presence of an α-aza residue might be conformationally favorable, leading to higher binding affinity, and be more resistant to enzymatic degradation. Replacement of amino acids in position 6, 9, and 10 of GnRH by α-aza amino acids alone did not confer any potency advantage but when the α-aza-Gly^{10} modification was combined with, for example, the D-Ser(Bu^t)^6 substitution, the resulting analog, [D-Ser(Bu^t)^6, α-aza-Gly^{10}]GnRH, currently undergoing clinical development, was considered to be at least 5 times more potent than [D-Ser(Bu^t)^6,Pro^9-NEt]GnRH, Buserelin, (74), using induction of ovulation as a measurement of potency (72, 73). (Buserelin has been, clinically, the most extensively studied GnRH analog.) It was unclear, to Dutta *et al.*, which individual factor was primarily responsible for this enhancement of biological potency. [2-D-Nal^6, α-aza-Gly^{10}] GnRH was reported to be slightly more potent than Nafarelin acetate and approximately 2.5 times more potent than the corresponding Pro^9-NEt modification in the estrus suppression assay (26). However, Nestor (26) noted that if the α-aza-Gly^{10} substitution conferred high potency by virtue of its enhanced resistance to the post-proline-cleaving enzyme in rat plasma, then the relevance of this substitution to human therapy was less clear since the amount of postproline-cleaving enzyme present in human plasma was reported to be 2–5 times less than in rat plasma (75, 76). Does the statement of Nestor also imply that any C-terminal amide modifications generally would not confer any advantage over the parent Gly^{10}-NH$_2$ function in humans? On the basis of the human data available on [D-Trp^6]GnRH and [D-Trp^6,Pro^9-NEt]GnRH (60, 63, 64), the answer would appear to be yes, although systematic comparisons would have to be made.

Position 7 modifications

[*N*-Me-Leu^7]GnRH was found to be equipotent with GnRH and [D-Ala^6,*N*-Me-Leu^7]GnRH was found to be at least as active as [D-Ala^6]GnRH *in vitro* (77). In fact, the *N*-Me-Leu^7 modification has been incorporated into [D-Trp^6,Pro^9-NEt]GnRH yielding [D-Trp^6,*N*-Me-Leu^7,Pro^9-NEt]GnRH, an analog currently undergoing clinical development (63). Generally, the effect of the *N*-Me-Leu^7 modification in enhancing the potency of the parent peptide depends on the D-amino acid at position 6 and the bioassay used to compare their potencies (45). The introduction of the bulky alkyl side chain, the tert-butyl group, as an ether into serine, which proved to be successful in significantly enhancing potency when incorporated into the 6-position (*e.g.* Buserelin), also enhanced potency at the 7-position, of nonapeptide NEt analogs of GnRH (74). The combination of two D-Ser(Bu^t) groups at position 6 and 7 not only failed to enhance, but actually decreased, the ovulation-inducing potency of Buserelin.

Conformationally constrained and backbone modifications

With the recognition that the biological activity of [D-Ala^6, *N*-Me-Leu^7]GnRH was consistent with that of a β-turn conformation for residues 5–8 of GnRH, Freidinger *et al.* (78) introduced a γ-lactam as a conformational constraint into the 6,7 position of GnRH and found that the resulting analog was 9 times more potent than GnRH *in vitro* and, by iv injection in ovariectomized rats, 2.4 times more potent than GnRH. Further exploration of the γ-lactam modification has not been made with agonists, *per se*, but this modification has been tried with GnRH antagonists. Various attempts to obtain a conformationally restricted agonist through cyclic analog design have yielded inactive analogs (79) or agonists with low biological potency (22, 80). Spatola (81) has reviewed the effect of peptide backbone modification on structure-activity relationships. Backbone modifications, as new approaches to GnRH agonist design, resulted in relatively little *in vitro* potency in the cases of peptide bond reversals at the 5–6 or 5–6 and 6–7 position (retro-inverso analogs) (82) or in the cases of substitution of a thiomethylene (-CH$_2$S-) group for the peptide linkage at the 5–6, 6–7, or 9–10 position of GnRH. The latter substitution at 9–10 had 10% of the *in vitro* potency compared to that of GnRH, indicating the necessity of more precise conformational requirements for residues 5–8 than for residues 9–10 (23).

Before closing the discussion on the current stage of development of the GnRH agonists, it is necessary to comment upon efforts to increase the potency of the agonists by modifying other amino acid residues.

Miscellaneous modifications

The 1-L-Nal^3 substitution (83) in GnRH is the only modification in position 3 which resulted in a peptide with greater potency than GnRH (twice as potent). Interestingly, [2-L-Nal^3]GnRH (84) was only half as potent as GnRH in stimulating LH release in ovariectomized rats. Although substitutions at the other remaining positions, 1, 2, 4, 5, 8, and 9, by naturally or unnaturally occurring amino acids, have resulted in decreased potency relative to GnRH, none of these residues are fundamentally required for GnRH activity. Each of these residues can be substituted to give active analogs, albeit with reduced potency. Fragments of GnRH and truncated (deletion) analogs of GnRH possessed very low GnRH potency. A series of peptides successively shortened from the C-terminus of GnRH were essentially inactive, except for Pro^9-GnRH which had 10% of the

Downloaded from https://academic.oup.com/edrv/article/7/1/44/2548841 by University of California, Los Angeles user on 19 April 2025

potency of GnRH *in vitro* (34). A concise review of these very early explorations is given by Coy *et al.* (85), and an extensive compilation is offered in the earlier volumes of the Specialist Periodical Reports cited above.

Modifications of naturally occurring vertebrate GnRH

The sequences of salmon GnRH with Trp^7,Leu^8 substitutions (86), chicken I GnRH with a Gln^8 substitution (87–89), and chicken II GnRH with His^5,Trp^7,Tyr^8 substitutions (90), relative to mammalian GnRH, have been recently discovered. Salmon GnRH and chicken I GnRH were isolated using RIAs based on antibodies raised against mammalian GnRH. Chicken II GnRH, with three drastic changes in a portion of the molecule considered important for receptor binding/activation, was isolated on the basis of its ability to release LH in an *in vitro* dispersed cell culture assay. The high relative potency of chicken II GnRH (30% that of mammalian GnRH) was unexpected in view of the results obtained with salmon and chicken I GnRH, which were only 1–2% as potent as mammalian GnRH in this same assay. The lead brought about by this observation, that [His^5,Trp^7,Tyr^8]GnRH (or chicken II GnRH) can still result in an analog with high *in vitro* relative potency, has been, so far, largely unexplored. Preliminary studies including substitution by D-amino acids at position 6 of some of the naturally occurring nonmammalian GnRH peptides have yielded analogs that are more potent than the parent peptides (91, 92), suggesting a secondary structure similar to that of GnRH. The Pro^9-NEt modification has also been introduced in conjunction with a D-amino acid at position 6 into nonmammalian GnRH (92). On the basis of preliminary biological studies conducted in goldfish, however, Peter *et al.* (92) concluded that the structural modifications that determine superactivity of mammalian, chicken, and salmon GnRHs in goldfish differ from what is known for mammals. Finally, a frog brain GnRH was partially characterized, and its structure was proposed to be identical with that of GnRH on the basis of the HPLC behavior of the isolated material and its immunological characteristics in several RIAs (93, 94).

Enzyme Degradation Studies

The literature on enzymatic degradation of GnRH and its analogs has been recently reviewed by Flouret *et al.* (95). The recognition of the enzymatic instability of GnRH as the reason for its relatively short lived biological effect prompted early half-life studies with GnRH. However, the discovery of the stabilizing effect of the position 6 modification (whether enzymatic, or conformational and resulting in greater binding affinity) preceded any attempts to experimentally determine the enzymatic cleavage sites of GnRH or its analogs. Since the high potency of the [D-Xaa^6]GnRH agonists had been rapidly realized, there was little practical incentive to design more potent agonists based on subsequent metabolic studies. It may be for this reason only that Flouret *et al.* (95) can legitimately claim: "The design of active LHRH analogs has not been guided by metabolic studies, but rather metabolic studies have been used to rationalize the high biological activity of some analogs." Certainly knowledge of the actual enzymatic cleavage sites was used to design some early antagonists (96). However, if this statement of Flouret *et al.* is historically valid, then it is also true that the retrospective rationalization has been highly speculative and, not necessarily, physiologically relevant. There is sufficient disparity between the *in vivo* and *in vitro* enzymatic results so as to lead Flouret *et al.* (95) to also conclude: "Attempts to correlate analog potency with resistance to degradation by tissue homogenates or by purified tissue peptidases must be cautiously evaluated, as these enzymes probably are compartmentalized and might not come in contact with LHRH or its analogs under physiological conditions."

It was originally reported by Koch *et al.* (97), using rat hypothalamic extracts, that Gly^6-Leu^7 was a major cleavage site of GnRH. A study by Marks and Stern (98), using rat brain homogenates, implicated Tyr^5-Gly^6 and Pro^9-Gly^{10}, in addition to Gly^6-Leu^7, as cleavage sites. Koch *et al.* (99) correlated the increased resistance of [D-Ala^6]GnRH and [D-Trp^6]GnRH to rat hypothalamus and anterior pituitary GnRH degrading endopeptidases with increased biological potency. Bauer and co-workers found <Glu^1-His^2 (100), Tyr^5-Gly^6 (and His^2-Trp^3, from the resulting N-terminal pentapeptide) (101), and Pro^9-Gly^{10} (102) as major cleavage sites of GnRH but not Gly^6-Leu^7, using purified enzymes isolated from bovine brain and pituitary homogenates. Thus, *in vitro*, three enzymes were identified: pyroglutamate aminopeptidase, endopeptidase, and postproline-cleaving enzyme. However, the relevance of these *in vitro* findings in contrast to the *in vivo* results, where <Glu^1-His^2 (43), His^2-Trp^3, Trp^3-Ser^4, and Ser^4-Tyr^5 (95) have been identified as cleavage sites of GnRH, is unclear.

Horsthemke *et al.* (103) concluded that even though superagonists modified at positions 6 and 10 are more resistant than GnRH to enzymatic degradation, there was no strict colinearity between their enhanced agonist activity and their resistance to degradation. Therefore, it was noted, factors other than resistance to degradation, such as increased receptor binding or stimulation, must be considered as being responsible for the potency of the superagonists. They also obtained evidence that substituents at positions remote from the enzymatic site of attack affect the cleavage rates. More recently, Flouret

Downloaded from https://academic.oup.com/edrv/article/7/1/44/2548841 by University of California, Los Angeles user on 19 April 2025

et al. (104) have conducted renal tissue studies with [D-Ser^4]GnRH, [D-Trp^6]GnRH, and [D-Ser^4,D-Trp^6]GnRH. It was found that [D-Ser^4]GnRH blocked enzymatic cleavage at position 4 but then Gly^6-Leu^7 was revealed as an ancillary scissile site. [D-Trp^6]GnRH, similarly, was not cleaved at position 6, nor, surprisingly, at the Ser^4-Tyr^5 bond, but was cleaved at the His^2-Trp^3 and Trp^3-Ser^4 bonds. Finally, [D-Ser^4,D-Trp^6]GnRH was degraded *in vivo* by proximal tubules at the His^2-Trp^3 bond. Flouret *et al.* (104) concluded that their data demonstrate important inhibitory effects of D-amino acid substituents remote from the scissile peptide bond. In fact, however, their data also indicate the general vulnerability of peptidic linkages, unprotected by multiple D-amino acids, to enzymatic degradation by various tissue peptidases.

Clinical Explorations

Some of the superagonists currently undergoing clinical trials or development have been identified above. The following is a summary: [D-Leu^6,Pro^9-NEt]GnRH (Abbott Laboratories, North Chicago, IL); [D-Trp^6,Pro^9-NEt]GnRH (Salk Institute, La Jolla, CA); [D-Trp^6] GnRH (Tulane University, New Orleans, LA); [D-Trp^6,*N*-Me-Leu^7, Pro^9-NEt]GnRH (Wyeth Laboratories, Philadelphia, PA); [D-Ser(Bu^t)^6,Pro^9-NEt]GnRH (Hoechst AG, Frankfurt, West Germany); [D-Ser(Bu^t)^6, α-aza-Gly^{10}]GnRH (Imperial Chemical Industries, Macclesfield, Cheshire, England); [N^τ-Bzl-His^6,Pro^9-NEt] GnRH (Salk Institute); [D-Nal(2)^6]GnRH (Syntex Laboratories, Inc., Palo Alto, CA). All of these superagonists have D-amino acids at position 6. Some are also modified at position 10, incorporating the Pro^9-NEt modification, and one has the α-aza-Gly^{10} replacement. One analog incorporates a *N*-Me-Leu^7 substitution into the combined modifications at positions 6 and 10. Leupron (Leuprolide) is now available in the United States for use in the treatment of prostatic cancer. All other analogs are in various stages of clinical exploration and/or development both in academia and industry.

Rationale for Development of GnRH Antagonists

The original impetus for the development of the superagonists was the treatment of infertility and, only subsequently, were the antireproductive effects discovered. However, the rationale for the development of the antagonists was derived from the immediate recognition that a competitive antagonist of GnRH had the potential of being a nonsteroidal contraceptive agent (105). In fact, a specific and generally confined physiological action was expected of the antagonists (106) which were intended primarily for female contraception (42). These analogs were expected to be free of the liabilities and/or toxicities associated with either of the components of the estrogen-progestagen combination pill. The greater safety anticipated for a metabolically and rapidly degradable peptidic contraceptive was another attractive feature of GnRH analogs even though it was recognized that if the analogs were to act as competitive antagonists to endogenous GnRH they would have to be continuously present at the receptor site in order to be efficacious. One approach would have been to generate irreversible inhibitors of GnRH; such inhibitors, however, were rejected as possible candidates due to their anticipated toxic and receptor-altering properties.

Animal Models/Biological Assay Correlations

The various biological assays employed for the testing of both agonists and antagonists have recently been reviewed (42, 48). With the discovery of the relatively potent disubstituted GnRH antagonists and the successful demonstration, for the first time, that an antagonist can suppress the preovulatory proestrus surge in the normal (4-day) cycling rat and thereby inhibit ovulation, a simple animal model became available for *in vivo* testing of GnRH antagonists (106). Other previously employed rat (107) and hamster models (108) were eventually abandoned for the routine testing of antagonists.

Corbin and Beattie (106) discussed the various biological models employed (pituitary cells; ovariectomized, steroid-blocked rats, *etc.*) and concluded that data translated from those models to the normal cycling female had to be employed with caution. Yardley *et al.* (109) noted that the *in vitro* and *in vivo* potencies of the analog pair, (des-His^2[D-Ala^6]GnRH and its Pro^9-NEt modification), were nonparallel and concluded that, by itself, antagonism of GnRH-induced LH secretion in cell cultures is inadequate in predicting antiovulatory activity. In contradistinction, Beattie *et al.* (110) demonstrated that potent antiovulatory analogs suppressed the proestrus, preovulatory LH surge, thereby clearly suggesting that this represented the basis for their antiovulatory activity. Later, with the availability of more potent antagonists, there was no longer a question of whether *in vitro* activity was generally predictive (110), qualitatively, of antiovulatory activity, but rather the question of whether there was a quantitative correlation between *in vitro* and antiovulatory activity in the rat. Bowers *et al.* (25) had concluded that for most of the antagonists, the *in vitro* and *in vivo* potencies were closely parallel. The most potent inhibitors of ovulation were always very potent *in vitro*, although there were instances of other antagonists possessing *in vitro* potencies identical with those of the potent ovulation inhibitors, but with little or no antiovulatory activity even at much higher doses (111). It was also concluded that quantitative and consistent estimates of antagonist potency could be obtained

Downloaded from https://academic.oup.com/edrv/article/7/1/44/2548841 by University of California, Los Angeles user on 19 April 2025

in vitro using either intact pituitaries of 21-day-old immature female rats (111) or dispersed pituitary cell cultures (112); however, potencies of antagonists are consistently much greater when they are assayed by the dispersed cell method than by the intact pituitary method with respect to the antagonist-GnRH dosage ratio as an index of antagonist potency.

It is accepted that the initial action of GnRH is to bind to specific receptors on the surface of its target cells. From a comparison of labeled GnRH agonists or antagonists, Perrin *et al.* (113) concluded that (1) similar kinetic and equilibrium behavior were displayed by agonists and antagonists and (2) the relative binding affinities of agonists and antagonists were not significantly different from one another irrespective of which radioligand was used in the radioreceptor assays. They further concluded that antagonists bound competitively to the same receptor sites available to agonists. They cited additional support for this latter conclusion from a report by Conn *et al.* (114), who showed that an antagonist can exhibit agonist properties when the antagonist is capable of causing receptor microaggregation. Photoaffinity studies with an antagonist led Hazum and Keinan (115) to the same conclusion: GnRH agonists and antagonists bind to the same receptor. This was consistent with their previous binding studies (116). Clayton and Catt (49) and Perrin *et al.* (50) noted that there was a general positive correlation between receptor binding affinity and relative *in vitro* antagonist activity.

More relevant, however, is the observation that the potency rankings of the antagonists depend upon the type of bioassay used (117). Receptor affinity represents only one parameter, and pharmacokinetic factors such as absorption, distribution, resistance to degradation, and elimination (excretion and/or metabolism of intact drug from plasma) also influence the *in vivo* efficacy of the analogs. Ultimately, the following question must be asked: "Is the rat antiovulatory assay, which has been the standard *in vivo* screening assay used, a predictive model for suppression of gonadotropins and for antifertility activity in the human female?" A similar question may be asked of the male rat as a predictor of efficacy in the human male. These questions will no doubt be discussed, in part, by other contributors to this volume. Nevertheless, is is important to note that antagonists, possessing a wide spectrum of potencies in rats, have indeed shown varying degrees of gonadotropin suppression in nonhuman primates as well as humans (see Table 2 under *Clinical explorations*). Finally, one anticipated advantage of the antagonists is the lack of an initial stimulation of gonadotropin release inherent in the GnRH agonists which may be undesirable in the use of agonists as contraceptive agents. Although little is certain concerning the possible uses of potent and long acting antagonists of GnRH, one can anticipate that they will be tested in all the clinical situations presently employed for GnRH superagonists.

Although the antiovulatory effects of the GnRH antagonists will be emphasized in this review, other antireproductive effects in female animals have been reported and recently reviewed (42, 118–120). Similarly, in the male, the gonadotropin suppression properties of the antagonists have been described (121) and recently reviewed (120, 122–124). Species differences have also been reported (125).

Development of GnRH Antagonists

Before it was even established that an antagonist could be developed, some doubts were expressed that the GnRH antagonist program of the Center for Population Research would have any more success than prior, partially successful efforts to develop other peptide antagonists (*e.g.* angiotensin, vasopressin, glucagon). The dramatic and rapid potency increases that were observed during the development of the GnRH agonists were not to be realized with the antagonists even when some of the structural features of the superagonists were incorporated early into the design of antagonists. The development of GnRH antagonists, with its requirement of high affinity for the GnRH receptor (without intrinsic activity) and resistance to enzymatic degradation, was slow and notable for the small, incremental increases in potency and by potency plateaus, as represented by a family of equipotent GnRH antagonists, which periodically appeared.

Early modifications: positions 2; 2,6; and 2,3,6

The first competitive antagonist of GnRH to be reported was des-His^2-GnRH by Vale *et al.* (126) using the dispersed pituitary cell culture assay. While [Gly^2]GnRH had partial agonist activity, des-His^2-GnRH showed no agonist activity at the doses tested, and both analogs reduced the secretion rate of LH in dispersed rat pituitary cells, as stimulated by GnRH, at molar ratios of 1,000 to 10,000 times that of GnRH. Explanations suggesting that the presence of His^2 was required, for recognition by the receptor or for GnRH activity, were clearly inadequate since [Phe^2]GnRH had 2–4% of the potency of GnRH, and [Trp^2]GnRH was 40% as potent *in vitro* (40). des-His^2-GnRH acts as a weak competitive inhibitor of GnRH because it lacks any functionality at position 2, which is required for some intrinsic activity, but still retains a structure with sufficient topological features remaining for recognition by the receptor. In this regard, both des-<Glu^1-GnRH and des-(<Glu^1-His^2)-GnRH showed very low, if any, GnRH agonist activity in rat pituitary tissue cultures (127). Other dele-

Downloaded from https://academic.oup.com/edrv/article/7/1/44/2548841 by University of California, Los Angeles user on 19 April 2025

tion analogs also showed very low agonist potency with the exception of Pro^9-GnRH which showed an *in vitro* potency of 10% that of GnRH (128). Thus, des-His^2-GnRH appears to be the only deletion analog reported to have antagonist activity in the absence of any other structural changes. Claims of lack of *in vivo* antagonist activity (129) for des-His^2-GnRH were attributed to its low intrinsic GnRH activity. This claim was refuted by a counter-demonstration of antagonist activity for des-His^2-GnRH in normal male rats (130) as well as equipotency with des-His^2[Pro^9-NEt]GnRH. The latter had been claimed by Coy *et al.* (129) to be the first inhibitor of GnRH found to be active *in vivo* (in ovariectomized and in estrogen- and progesterone-treated rats); subsequently, this inhibitor was reported to block ovulation induced by GnRH (131). Monahan *et al.* (38), utilizing the observations of *in vitro* antagonist activity for des-His^2-GnRH and high agonist activity for [D-Ala^6]GnRH, synthesized des-His^2[D-Ala^6]GnRH, which had 3 times the antagonist potency of des-His^2-GnRH. Monahan *et al.* (132) were the first to demonstrate that a D-amino acid substitution, *e.g.* [D-Ala^2]GnRH, would lead to a GnRH partial antagonist. However, a breakthrough (reported by Yardley *et al.* and Corbin and Beattie) resulted from combining the structural features of the weakly active antagonist, [D-Phe^2]GnRH (39), with those of [D-Ala^6]GnRH, yielding [D-Phe^2,D-Ala^6]GnRH, which was the first antagonist to inhibit ovulation (at 6 × 1 mg doses in a corn oil vehicle injected sc) in the normal cycling rat (106, 109). Other modifications at position 6, such as [D-Phe^2,D-Phe^6]GnRH, did not improve antiovulatory potency, but [4-F-D-Phe^2,D-Ala^6]GnRH (110) was the first example of the use of a halogenated phenylalanine residue in position 2, a substitution which would later play a very important role in greatly increasing antagonist potency. Incorporating the Pro^9-NEt modification into [D-Phe^2,D-Ala^6]GnRH yielded an antagonist which had *in vitro* activity (109), but lacked antiovulatory activity even at 10 times the dose of the parent, [D-Phe^2,D-Ala^6]GnRH. des-His^2[D-Ala^6,Pro^9-NEt]GnRH was also found to be less potent than des-His^2[D-Ala^6]GnRH *in vivo* and this was attributed to the higher inherent GnRH agonistic activity of the former (133). The Pro^9-NEt modification, even for the later, very potent antagonists, generally reduced rather than improved potency, suggesting that improved *in vitro* potency in the agonist series may be due to a stabilized secondary structure of the active conformer involving a dipole-dipole interaction of the NEt functionality with the imidazole ring of histidine. This folding would be analogous to that suggested in the case of TRH by Donzel *et al.* (134). The only reported all L- disubstituted antagonist, [Leu^2,Leu^3]GnRH, had very weak *in vitro* potency (135). The D-Ala^6 modification, [Leu^2,Leu^3,D-Ala^6]GnRH also possessed weak *in vitro* potency (136). However, by incorporating an aromatic L-amino acid into position 3 of [D-Phe^2,D-Phe^6]GnRH, Coy *et al.* (137) and de la Cruz *et al.* (138) found that the resulting analog, [D-Phe^2,Phe^3,D-Phe^6]GnRH, improved potency 2-fold, exhibiting complete antiovulatory activity at 3 × 1 mg doses. Similarly, Humphries *et al.* (139) reported that incorporation of L-proline into position 3 of [D-Phe^2,D-Trp^6]GnRH yielded [D-Phe^2,Pro^3,D-Trp^6]GnRH, which completely inhibited ovulation at a single dose of 750 μg. The apparent discrepancy in the antiovulatory and *in vitro* potencies of di- and trisubstituted analogs, *e.g.* [D-Phe^2,D-Phe^6]GnRH [inhibitory dose ratio ((IDR_{50}) = 23/1)] and [D-Phe^2,Pro^3,D-Trp^6]GnRH (IDR_{50} = 48/1), was subsequently attributed by Rivier and Vale and coworkers (140, 141) to greater resistance to degradation or to a decrease in residual intrinsic activity resulting from the presence of the position 3 modification. [D-Phe^2,D-Trp^3,D-Phe^6]GnRH, with three D-amino acid residues, showed 83% inhibition of ovulation with a single dose of 1 mg while N^{ϵ}-pentapeptidyl-Lys^6 derivatives of [D-Phe^2,D-Trp^3,D-Lys^6]GnRH were, on a molar basis, slightly more potent than the parent D-Phe^6 analog (142). Dutta *et al.* (72) combined the α-aza-Gly^{10} modification, previously used to increase agonist potency (72, 73), with [D-Phe^2,D-Phe^6]GnRH to give [D-Phe^2,D-Phe^6, α-aza-Gly^{10}]GnRH, which showed an 8-fold increase in antiovulatory potency, compared with the parent antagonist, using inhibition of GnRH-induced ovulation as an index of potency. No comparisons were reported using the normal cycling rat. The α-aza-Gly^{10} moiety was utilized occasionally in subsequent, potent antagonists, but its success in improving potency was limited and usually erratic (143).

Conformation studies: position 1 modifications

The octapeptide, des-His^2[D-<Glu^1,Pro^9-NEt]GnRH was prepared specifically with the anticipation that an N-terminal D-amino acid would decrease the rate of degradation of the enzyme (96), presumably, pyroglutamate aminopeptidase (100, 144). The analog had some antagonist activity in ovariectomized rats, but the successful utilization of a D-<Glu^1 substitution was not realized until the next major breakthrough occurred.

Until this time, progress in increasing the potency of the GnRH analogs had been based largely on the utilization of classical functional group modifications of the side chains of GnRH coupled with changes in chain length of the peptide. The concept of conformational stabilization of a β-II type bend involving Ser-Tyr-Gly-Leu residues of GnRH had been invoked to explain the high potency of [D-Ala^6]GnRH relative to GnRH itself (38). The first attempt to introduce a conformational

Downloaded from https://academic.oup.com/edrv/article/7/1/44/2548841 by University of California, Los Angeles user on 19 April 2025

constraint into a GnRH agonist, based on the premise of Monahan *et al.* (38) and a test of the use of conformation-activity relationships for the rational design of GnRH analogs, was reported by Donzel *et al.* (79) who synthesized [Glu^4,D-Ala^6,Orn^7]GnRH. The biological inactivity of both the cyclic and linear analog was attributed to the differences in the side chains of Glu^4 and Orn^7 as compared with those of Ser^4 and Leu^7 (in GnRH). Molecular models were used to represent peptide conformations such as that of GnRH (145, 146), but, because of the inherent flexibility in peptide bonds, the use of such models is unenlightening in the absence of any other structural information (38). Spectroscopic studies employing NMR (147, 148) and optical rotary dispersion/circular dichroism (149, 150) led to the conclusion that GnRH behaves as a random coil in water and is devoid of any intrachain residue interactions (150). Marche *et al.* (149) noted a trend toward ordered structures upon increasing the temperature at pH 7.4 while Mabrey and Klotz (150) observed that GnRH exhibited a conformational transition in trifluoroethanol with the formation of a β-structure which may be relevant to biological activity at the receptor site. Momany (151–153) was the first to employ semiempirical conformational energy calculations on GnRH and its analogs. Since it was not feasible to examine all of the possible conformations of GnRH, Momany (151) restricted his calculations by utilizing 1) the observation concerning the conformational stabilization resulting from the introduction of D-Ala^6 into GnRH (38), 2) a ^{13}C-NMR study showing that the proline peptide bond, in GnRH, was completely trans (154) and 3) the "computationally derived low energy 'dipeptide' conformations of the residues involved." He concluded, by energy minimization techniques, that there were three low energy conformations of GnRH, one of which appeared to explain the observed analog data: the so-called CC conformer with the N- and C-terminus in close proximity, as speculated by Grant and Vale (146). He also noted that the low energy structures had a modified type II bend involving Tyr-Gly-Leu-Arg rather than Ser-Tyr-Gly-Leu as proposed by Monahan *et al.* (38). These observations, supported by the subsequent work of Freidinger *et al.* (78), would now explain why [D-Ala^6,*N*-Me-Leu^7]GnRH had enhanced and not reduced biological potency as originally predicted (77). Leu^7 is not involved with Ser^4 in hydrogen bonding in this modified β-II type bend, and, therefore, *N*-Me-Leu^7 would not disrupt the modified β-II type bend. In spite of the limitations of Momany's study (the use of semiempirical calculations to determine preferred conformations and the inability to identify a single global minimum amongst the low energy conformers), a working model with which to test further structure-activity relationships of GnRH analogs was provided.

Rivier and Vale (140) proceeded to use, as a rationale for introducing D-<Glu^1 in position 1, Momany's observation that the analogs containing L-<Glu^1 and D-amino acids at positions 2, 6, and/or 3 had low energy conformers with a common configurational property. The *cis* peptide bond of the L-<Glu ring in these antagonists had changed orientation with respect to that of the L-<Glu ring in the CC conformer of GnRH. Reversing this situation, by replacing L-<Glu^1 with D-<Glu^1 in the antagonists, would reorient the *cis* peptide bond of the D-<Glu ring in the antagonists to coincide with the orientation of the L-<Glu ring found in the CC conformer of GnRH. Although the analog suggested by Momany [D-<Glu^1,Phe^2,D-Xaa^6]GnRH, was not synthesized, the corresponding D-Phe^2 trisubstituted analog [D-<Glu^1,D-Phe^2,D-Phe^6]GnRH (155) was prepared. This analog lacked antiovulatory activity at the dose tested (750 μg) but did exhibit *in vitro* antagonist activity. However, when D-<Glu^1 was combined with [D-Phe^2,D-Trp^3,D-Trp^6]GnRH by Rivier and Vale (140), the resulting antagonist was found to have complete antiovulatory activity at 250 μg/rat and to exhibit *in vitro* potency in a molar ratio of 3:1 with GnRH. This was a 5-fold increase in antagonist potency with respect to the parent analog *in vitro*. Other D-<Glu^1 analogs were less potent or showed no advantage over the prototype, *in vitro* or *in vivo* (140). [D-<Glu^1,D-Phe^2,D-Trp^3,D-Trp^6]GnRH also demonstrated prolonged activity in the antiovulatory assay, possibly due to greater enzymatic resistance *in vivo* or to delayed absorption when suspended in corn oil. It was the first antagonist reported to totally block ovulation when administered as a single injection at 0900 h on the day of proestrus. The N^{ϵ}-isophthaloyl dimer of [D-<Glu^1,D-Phe^2,D-Trp^3,D-Lys^6]GnRH (156) completely inhibited ovulation at 250 μg; it had 6 times the potency of its monomeric parent. The use of cyclopentane carboxylic acid in position 1, as suggested by Momany (153), was unsuccessful, *in vivo* (157). It should also be noted that, even with the evidence for the modified β-II type bend, the conformation of GnRH or its analogs was still largely unknown. It was proposed that it was more likely that GnRH and its analogs exist as an ensemble of conformers in solution rather than a single conformer (158). Subsequent NMR studies (159) supported the modified β-turn involving Tyr-Gly-Leu-Arg. Some empirical evidence for the occurrence of the CC conformation in solution was offered by NMR and circular dichroism measurements (160), but under no circumstances could this evidence be considered as proof. However, the accumulated evidence supports the β-turn, or the folding involving Tyr-Gly-Leu-Arg, in GnRH, as the bioactive conformation in this region (78, 159). X-ray studies on GnRH and its analogs have been hampered

Downloaded from https://academic.oup.com/edrv/article/7/1/44/2548841 by University of California, Los Angeles user on 19 April 2025

by the difficulty encountered by crystallographers in growing appropriate crystals necessary for such studies.

4-Cl(and F)-D-Phe² modifications: increasing hydrophobicity

The pace of discovery of new potent antagonists quickened. Humphries *et al.* (161) showed that Ac-Pro¹ could replace D-<Glu¹ with comparable antiovulatory potency for [Ac-Pro¹,D-Phe²,D-Trp³,⁶]GnRH, emphasizing that a D-amino acid in position 1 was not essential and, in fact, the corresponding Ac-D-Pro¹ analog was less potent than either the D-<Glu¹ or the Ac-Pro¹ modification. Channabasavaiah and Stewart (162) extended the range of position 1 acetyl-D-amino acids that could be accommodated in the antagonist structure to Ac-D-Ala and Ac-D-aromatic amino acids, in general, with antiovulatory activity in the 100–250 μg range. The necessity of the acetyl group for high potency in this case was realized when it was subsequently shown (156) that [D-Phe¹,D-Phe²,D-Trp³,D-Phe⁶]GnRH was inactive at 1 mg/rat. However, with the additional discovery by Coy *et al.* (156) that the use of 4-Cl-D-Phe in position 2 dramatically increased the antiovulatory potency of this peptide (82% inhibition at 250 μg), the exact electronic requirements (carbonyl *vs.* aromaticity) at position 1 were unclear. Nevertheless, combining 4-Cl-D-Phe² with Ac-D-Phe¹ yielded [Ac-D-Phe¹,4-Cl-D-Phe²,D-Trp³,⁶]GnRH which exhibited complete antiovulatory activity at 62 μg in a propylene glycol-saline vehicle and significant inhibition of ovulation at 15 μg in a corn oil vehicle (possibly due to prolongation of release of the antagonist from the corn oil medium). With the increasing hydrophobic alignment of aromatic D-amino acids at positions 2,3, and 6, Spatola *et al.* (163) found that even the nonbulky Ac-Gly¹ provided a potent antagonist, [Ac-Gly¹,4-Cl-D-Phe²,D-Trp³,⁶]GnRH, with 100% antiovulatory activity at 25 μg in a corn oil vehicle. Exploration by Rivier *et al.* (22, 68) of the parent analog [D-Phe²,D-Trp³,⁶]GnRH led to the conclusion that [Ac-Δ³-Pro¹,D-Phe²,D-Trp³,⁶]GnRH, containing the acetyl-3,4-dehydroproline¹ residue, was significantly more potent than other position 1 substituents. When this modification was then combined with 4-Cl-D-Phe², the resulting analog, [Ac-Δ³-Pro¹,4-Cl-D-Phe²,D-Trp³,⁶]GnRH, showed complete inhibition of ovulation at 7.5 μg in corn oil (22, 68).

Predictive and deductive approaches

A more systematic attempt was then made to improve potency by applying the manual method of Topliss (164) to the Hansch approach of drug design (22, 68). Since the apparent correlation of biological potency of certain GnRH superagonists with their overall hydrophobic character, as measured by HPLC, did not hold for a series of GnRH antagonists, it was concluded that the receptor requirements for recognition and binding may involve other factors in addition to hydrophobicity, such as localized electronic density and steric effects of substituents. The Topliss method offered an opportunity to identify these essential parameters, found in Hansch-type correlations, for loci which contain aromatic side chains. This method was applied to a series of [Ac-Δ³-Pro¹,Xaa²,D-Trp³,⁶]GnRH analogs. Although one analog was discovered (4-Br-D-Phe²) with higher *in vitro* potency, the classically identified 4-Cl-D-Phe² and the closely related 4-F-D-Phe² analogs were the most potent *in vivo*. Using the Topliss approach, the investigators were unable to identify a unique set of operative parameters, as many of the antagonists turned out to have similar potencies. It is likely that this approach, which has seen considerable success in small, rigid systems, is inapplicable to the much more flexible and larger peptide systems where biological potency resulting from electronic and steric changes in the molecule may be influenced by other parameters including conformational effects (Gierasch, L. M., private communication). Still further *in vivo*, but not *in vitro*, improvements were made by incorporating β-(2-naphthyl)-D-alanine into positions 3 and 6, which resulted in analogs such as [Ac-Δ³-Pro¹,4-F-D-Phe²,2-D-Nal³,⁶]GnRH. This analog completely inhibited ovulation at a dose of 2.5 μg (22). Previously, this D-amino acid had been used to obtain a potent agonist (62). Investigation of a series of hydrophobic antagonists, using 2-D-Nal in positions 3 and/or 6, led to the conclusion that Ac-Pro¹ could be substituted for Ac-Δ³-Pro¹, resulting in analogs with equipotency in an antiovulatory assay as measured by an ED_{50} rather than an ED_{100} (28). The *N*-Me-Leu⁷ modification did not significantly improve the potency of the Ac-Δ³-Pro¹ modification (22) nor did additional forays into modifications of position 5 using Topliss' manual method (27).

D-Ala¹⁰ modification

The use of D-Ala¹⁰ by Erchegyi *et al.* (165) resulted in the potent analog, [Ac-4-Cl-D-Phe¹,4-Cl-D-Phe²,D-Trp³,D-Phe⁶,D-Ala¹⁰]GnRH, which blocked ovulation in seven of eight rats at 7.5 μg. This analog is of interest since it contains five D-amino acid residues, which may confer enzyme resistance to peptide bonds throughout much of the sequence, except at the Ser⁴-Tyr⁵ and Leu⁷-Arg⁸-Pro⁹ portions of the peptide. The introduction of sterically larger groups, such as D-Ser¹⁰ and D-Leu¹⁰, resulted in less potent analogs, implying, perhaps, that side chain branching at position 10 hinders receptor binding. This D-Ala¹⁰ analog was the most potent analog found in a classical structure-activity study dealing particularly with positions 1 and 2 (165, 166). Administra-

Downloaded from https://academic.oup.com/edrv/article/7/1/44/2548841 by University of California, Los Angeles user on 19 April 2025

tion of analogs, including the early disubstituted ones, to rats before diestrus (106) or to immature male rats (167) allowed for the determination of duration of action of the antagonists *in vivo*. With the observation of prolonged activity with the potent tetrasubstituted analog (140), increased attention was turned to evaluating analogs for prolonged activity. Several Ac-Δ^3-Pro^1 analogs, examined by Rivier *et al.* (118) as well as the above D-Ala^{10} analog, examined by Coy *et al.* (24), showed extended duration of action in the animal models employed.

D-Arg^6 modification: efforts to increase hydrophilicity of antagonists

The rapid development of the GnRH antagonists was, by now, being accompanied by numerous residue changes, and it was necessary to frequently reevaluate individual amino acid substitutions relative to the rest of the molecule in order to optimize potency. With this in mind, and, in order to improve water solubility by the introduction of more hydrophilic amino acids, position 6 was reexamined in relation to the Ac-4-Cl-D-Phe^1 and D-Ala^{10} modification by Coy *et al.* (168). With their introduction of basic D-amino acids, such as D-Arg^6, the next breakthrough was realized. The most potent analog reported was [Ac-4-Cl-D-Phe^1,4-Cl-D-Phe^2,D-Trp^3,D-Arg^6,D-Ala^{10}]GnRH, with 100% inhibition of ovulation in the 1.5- to 3-μg range in corn oil and somewhat less potent in propylene glycol-saline. The observed difference in potency in the two vehicles was suggested (168) to be due to delayed and/or prolonged absorption of this D-Arg^6 antagonist. The corresponding tetrasubstituted analog, lacking D-Ala^{10}, was reported to be 10-fold less potent. The corresponding D-Lys^6,D-Ala^{10} analog was less potent than the D-Arg^6,D-Ala^{10} modification. Extension of this observation by Horvath *et al.* (169) showed that the Ac-2-D-Nal^1 pentasubstituted derivative, [Ac-2-D-Nal^1,4-Cl-D-Phe^2,D-Trp^3,D-Arg^6,D-Ala^{10}]GnRH, had approximately 3 times the potency of the corresponding Ac-4-Cl-D-Phe^1 analog. This study, and the need to frequently reevaluate other positions each time a significant single positional improvement is discovered in order to optimize potency, strongly indicates that there is still a large empirical component to GnRH analog design.

The hydrophilic tetrasubstituted antagonist, [Ac-Δ^3-Pro^1,4-F-D-Phe^2,D-Trp^3,D-Lys^6]GnRH, exhibited very high binding affinity (113), but no antiovulatory data were reported for this analog. However, the corresponding D-Arg^6 analog with comparable binding affinity showed unexpectedly low antiovulatory potency: 60% inhibition of ovulation at 20 μg in corn oil (27). This finding was similar to that of Coy *et al.* (168) who had previously synthesized and tested [Ac-Δ^3-Pro^1,4-Cl-D-Phe^2,D-Trp^3,D-Arg^6]GnRH and found no antiovulatory activity at 7.5 μg in propylene glycol-saline. However, high *in vivo* potency could be restored to the tetrasubstituted D-Arg^6 analogs (without D-Ala^{10}) by the introduction of the very hydrophobic residue, 2-D-Nal, at position 1. Thus, [Ac-2-D-Nal^1,4-F-D-Phe^2,D-Trp^3,D-Arg^6]GnRH (27) was reported to have complete antiovulatory activity at 1 μg in corn oil. Rivier *et al.* (27) concluded that, generally, the GnRH receptor will tolerate either a hydrophobic or a hydrophilic residue at position 6, provided that it is concomitantly paired with a hydrophilic (Ac-Δ^3-Pro) or hydrophobic (Ac-2-D-Nal) residue, respectively, at position 1. Coy and Nekola (170) arrived at the same conclusion on the basis of independent data. Commenting on the hydrophilic-hydrophobic surface model, proposed by one of us (M.J.K.), Coy and Nekola (170) observed that the modifications to the hydrophilic part of the peptide chain (or looped structure stabilized by the position 6 D-amino acid) often must be accompanied by suitable alterations in the complimentary hydrophobic region in order to maintain or increase antagonist potency. It is the alteration of the global properties of these peptides, sometimes in a very subtle manner, that has enabled investigators to achieve antiovulatory potency in the nanogram range. It should be noted that while the referenced hydrophilic-hydrophobic surface model of the antagonist is consonant with many of the past structure-activity studies, the model is no doubt an oversimplification which has already been contradicted by a subsequent study of Hocart *et al.* (171). One would have predicted, on the basis of this model, that position 7 could accommodate hydrophilic amino acids. However, the introduction of Lys or Thr, at position 7, was detrimental to potency, while Phe^7, a hydrophobic residue, improved potency (171). Thus, the model is not compatible with these biological results.

[Ac-2-D-Nal^1,4-F-D-Phe^2,D-Trp^3,D-Arg^6]GnRH was tested for duration of action and required 100 μg when given on diestrus II, as compared with 1 μg when given on proestrus, for complete inhibition of ovulation (27). Nekola and Coy (172) had shown that a 100 μg dose of [Ac-2-D-Nal^1,4-Cl-D-Phe^2,D-Trp^3,D-Arg^6,D-Ala^{10}]GnRH suppressed LH levels for more than 30 h in ovariectomized rats. [Ac-2-D-Nal^1,4-F-D-Phe^2,D-Trp^3,D-Arg^6]GnRH also had 100% oral antiovulatory potency at 2.5 mg (27). Nekola *et al.* (173) had previously shown that [Ac-4-Cl-D-Phe^1,4-Cl-D-Phe^2,D-Trp^3,D-Arg^6,D-Ala^{10}]GnRH was completely effective, at 2.0 mg, orally, in blocking ovulation. The low oral antiovulatory potency (after gavage) of these analogs as compared with their sc potency was attributed, by both groups of investigators, to poor oral absorption (<1% of the absorption rate by injection), enzymatic degradation, or elimination. Structure-activity relationships (based on potencies obtained by oral *vs.* sc administration) do not necessarily correlate

Downloaded from https://academic.oup.com/edrv/article/7/1/44/2548841 by University of California, Los Angeles user on 19 April 2025

(172), but it has been generally observed that the oral-sc potency ratio is about 1000:1 under the conditions tested.

Alterations at the C terminus, either by use of the Pro^9-NEt modification (27) or by α-aza-Gly^{10} (143), in conjunction with the D-Arg^6 or the 2-D-Nal^6 antagonist series, were usually, but not always (28, 173a), detrimental to improving biological potency. The *in vitro* or *in vivo* enhancements of potency that were seen in the GnRH agonist series by use of Pro^9-NEt or α-aza-Gly^{10} were rarely realized during the development of the antagonists.

The idea of combining basicity, aromaticity, and hydrophilicity into a single amino acid led Folkers *et al.* to introduce the heterocyclic amino acids, β-(3-pyridyl)-alanine (174) and β-(3-quinolyl)-alanine (175), into a series of GnRH antagonists at positions 3 and/or 6. These amino acids are less basic than arginine but have the aromatic properties of tryptophane and naphthylalanine. The most potent analog, [Ac-2-D-Nal^1,4-Cl-D-Phe^2,3-D-Pal^3,D-Arg^6,D-Ala^{10}]GnRH, was reported to completely inhibit ovulation at 500 ng, suggesting greater potency than the corresponding D-Trp^3 analog (169). Simultaneous use of 3-D-Pal at position 3 and 6 decreased potency with respect to the parent D-Trp^3,D-Arg^6 analogs. β-(3-Quinolyl)-D-alanine offered a comparable replacement for the D-Trp^6 antagonists but not for the D-Arg^6 antagonists in terms of biological potency.

In order to further test the predictions of conformation based on minimum energy calculations (153) Roeske and Anantharamaiah (176) imposed conformational constraints at the N terminus with the introduction of α-methyl-D-amino acids at positions 2 and/or 3. The biological results supported the proposed helical type III bend for position 2 substitutions, but the use of α-methyl-D-amino acids at position 3 caused considerable loss of biological potency compared with the unmethylated parent analogs. Rivier *et al.* (22) had previously found that introduction of α-Me-4-Cl-D-Phe^2 was only marginally detrimental to biological potency when compared with the corresponding unmethylated analog. Roeske and Anantharamaiah extended the use of α-methyl amino acids at position 2 to the more potent 3-D-Pal^3,D-Arg^6 antagonists. The resulting analog [Ac-2-D-Nal^1,α-Me-4-Cl-D-Phe^2,3-D-Pal^3,D-Arg^6,D-Ala^{10}]GnRH, was completely active in inhibiting ovulation at 500 ng (Roeske, R., N. Chaturvedi, and T. Hrinyo, unpublished observations).

N^ω,$N^{\omega\prime}$-Dialkyl-D-Har^6 modifications

Nestor *et al.* (62, 67) applied their concept of the importance of the depot effect to GnRH antagonists (28, 67, 177). A prolonged biological half-life is more critical for antagonists than agonists since effective competition with endogenous GnRH pulses demands the continuous presence of the antagonist at the pituitary receptor site. They noted that [Ac-4-Cl-D-Phe^1,4-Cl-D-Phe^2,D-Trp^3,D-Arg^6,D-Ala^{10}]GnRH (168), which contains the hydrophilic D-Arg^6 moiety, had a prolonged duration of action compared with [Ac-Pro^1,4-F-D-Phe^2,2-D-Nal^{3,6}]GnRH. They suggested that the prolonged activity of the D-Arg^6 analog may be due to a depot effect involving an electrostatic interaction between the negatively charged phosphate group of the phospholipid membrane and the positively charged guanidine group of D-Arg. [It may be appropriate to cite a model proposed by Schwyzer *et al.* (178) wherein the lipid phase of the target cell membrane initially interacts with the peptide and induces secondary structures and topological arrangements, which then facilitate receptor-peptide interaction.] To further stabilize the hypothesized interactions, a series of N^ω,$N^{\omega\prime}$-dialkyl-D-arginine and homoarginine residues were incorporated into GnRH antagonists. These residues, it was suggested, would have the potential for both electrostatic and hydrophobic interaction with the phospholipid bilayer. The most potent pentasubstituted analog in this series [Ac-2-D-Nal^1,4-Cl-D-Phe^2,D-Trp^3,N^ω,$N^{\omega\prime}$-diethyl-D-Har^6,D-Ala^{10}]GnRH, was compared, biologically, with some literature standards but not with the more closely related Ac-2-D-Nal^1 standard [Ac-2-D-Nal^1,4-Cl-D-Phe^2,D-Trp^3,D-Arg^6,D-Ala^{10}]GnRH (169, 172), which contains, as the only change, D-Arg^6 rather than N^ω,$N^{\omega\prime}$-diethyl-Har^6. In the only two structurally analogous comparisons with a D-Arg^6 analog, an Ac-4-Cl-D-Phe^1,N^ω,$N^{\omega\prime}$-diethyl-D-Har^6 analog was slightly more potent, in an antiovulatory assay, when the analog was administered on diestrus II (rather than on proestrus), than the corresponding D-Arg^6 analog and [Ac-2-D-Nal^1,4-F-D-Phe^2,D-Trp^3,N^ω,$N^{\omega\prime}$-diethyl-D-Har^6]GnRH was approximately 2.5 times more potent on diestrus II than the corresponding D-Arg analog (67). These results suggest, in the two comparisons cited, that the N^ω,$N^{\omega\prime}$-diethyl-D-homoarginine substitution for D-Arg in position 6 does not significantly increase acute biological potency but may, according to the investigators, substantially increase potency in terms of duration of action. However, it should be noted that there was a greater difference in potency, in terms of duration of action, between [Ac-2-D-Nal^1,4-Cl-D-Phe^2,D-Trp^3,N^ω,$N^{\omega\prime}$-diethyl-Har^6]GnRH (ED_{50} = 2.5 μg on diestrus II) and the corresponding Ac-4-Cl-D-Phe^1 analog (ED_{50} = 16 μg on diestrus II) than there was between the former analog and [Ac-2-D-Nal^1,4-F-D-Phe^2,D-Trp^3,D-Arg^6]GnRH (ED_{50} = 9 μg on diestrus II), which lacks the N^ω,$N^{\omega\prime}$-diethyl-Har^6 modification.

Position 7 modifications

Folkers *et al.* (179) used, as a rationale for further substitutions into the more potent antagonist, the ob-

Downloaded from https://academic.oup.com/edrv/article/7/1/44/2548841 by University of California, Los Angeles user on 19 April 2025

served differences between mammalian GnRH, salmon GnRH (86), chicken I GnRH (87–89), and chicken II GnRH (90). The four vertebrate GnRH structures were postulated (179) to have similar receptors, with minor structural changes enabling the different GnRH structures to bind to the different parent receptors. Thus, it was suggested (179) that the mammalian GnRH receptor might be able to bind an antagonist with changes in the 5, 8, and particularly in the 7 position, the attractive feature of the position 7 modification being that chicken II and salmon GnRH have Trp^7 substitutions. Milton *et al.* (180), however, had noted that their results suggested that there was a difference between the chicken receptor and the mammalian GnRH receptor in the recognition of GnRH analogs substituted in position 8. Nevertheless, the most potent analog of the series [Ac-2-D-Nal^1,4-Cl-D-Phe^2,3-D-Pal^3,D-Arg^6,Trp^7,D-Ala^{10}]GnRH was reported to show complete inhibition of ovulation at 500 ng and 90% inhibition (2/20 rats ovulated) at 250 ng, whereas the parent antagonist (without Trp^7) showed complete inhibition at 500 ng but only partial inhibition (3/7 rats ovulated) at 250 ng (179). A variety of other hydrophobic and aromatic basic amino acid substitutions at position 7 proved less potent. The combined substitution of Trp^7 with Trp^5, Phe^5, or 3-Pal^5 similarly showed no improvement in antagonist potency. Rivier *et al.* (181) and Hocart *et al.* (171) synthesized some position 7 modifications and obtained somewhat different biological results. Rivier *et al.* (181) reported that [Ac-2-D-Nal^1,4-Cl-D-Phe^2,D-Trp^3,D-Arg^6,Trp^7,D-Ala^{10}] GnRH, which contains the naturally occurring Trp^7 of fish GnRH (86), was equipotent (ED_{100} = 2.5 μg) with the corresponding Met^7 and Phe^7 in the antiovulatory assay using corn oil as a vehicle. Hocart *et al.* (171) approached position 7 modifications from another point of view, specifically, as a test of the hydrophilic-hydrophobic surface model (170). When hydrophilic substitutions at position 7 showed at least 2-fold less potency, and Phe^7 showed greater potency, than the parent analog in antiovulatory assays using 40% propylene glycol-saline as a vehicle, a variety of aromatic amino acid substitutions were made. The most potent, [Ac-2-D-Nal^1,4-Cl-D-Phe^2,D-Trp^3,D-Arg^6,Phe^7,D-Ala^{10}]GnRH (identical to that of Rivier *et al.* above) was reported to be approximately twice as potent (64% inhibition of ovulation at 0.5 μg) as the Tyr^7 antagonist and more potent than either the corresponding Trp^7 or 2-Nal^7 analogs. Hocart *et al.* concluded that the results indicate a preference for hydrophobic residues at position 7, but that beyond a certain point, increasing hydrophobicity has little effect on antagonist potency.

Transposition modifications involving positions 5 and 6

A transposition involving the introduction of a basic amino acid into position 5 was reported by Roeske *et al.* (182). They had considered interchanging the Arg^5,D-Tyr^6 residues for the Tyr^5,D-Arg^6 analog, based upon the Momany model of GnRH wherein residues 5, 6, and 7 have their side chains oriented along the outside of the modified β-II type bend. Similar to previous suggestions (67, 178), was the proposal that these side chains may not be in contact with specific groups in the receptor, but possibly interact nonspecifically with components of the cell membrane. [Ac-2-D-Nal^1,4-Cl-D-Phe^2,D-Trp^3,Arg^5,D-Tyr^6,D-Ala^{10}]GnRH completely inhibited ovulation at 2.5 μg or at approximately twice the dose of the parent Tyr^5,D-Arg^6 analog. The $N^{\omega},N^{\omega'}$-diethyl-Har^5,D-Tyr^6 and the N^{ϵ}-isopropyl-Lys^5,D-Tyr^6 analogs were somewhat more potent, showing almost complete inhibition (1/10 rats ovulated) at 1.0 μg. However, the corresponding Glu^5,D-Arg^6 analog was impotent at 10 μg. These results were interpreted as evidence in support of the view that the position 5 side chain interacts with the cell membrane.

Chain length and backbone modifications

References to shorter chain GnRH antagonists such as the Pro^9-NEt modifications and their general inapplicability to improvements in potency have already been given. One longer chain analog, an undecapeptide, endo-Pro^{1a}[D-Phe^2,D-Trp^{3,6}]GnRH (183), was equipotent with its parent Ac-Pro^1 analog, but this example is not necessarily a satisfactory test of the effect of chain length on antiovulatory potency, since the corresponding endo-Gly^{1a} and the Gly^1,endo-Pro^{1a} analogs were less potent. A number of pseudodipeptide backbone modifications of antagonists were made by Spatola *et al.* (23) at position 1–2, 6–7, and 9–10 using —CH_2S— (a methylene sulfide bond) to replace the peptide bond. As was previously found for the agonists (23), the least potent analogs *in vitro* had replacements at the 6–7 position, and it was concluded that such replacements lead to undue flexibility, whereas currently acceptable conformational models demand stabilization of a β-turn involving residues 6 and 7. Replacements at the 1–2 and 9–10 positions resulted in analogs which were equipotent with their parent peptides *in vitro*, but were impotent in antiovulatory assays at the doses tested. This lack of *in vivo* potency was attributed to impaired absorption in cases wherein the analog structures were beyond optimal solubility in the Hansch formulation (184). The sulfoxide replacement, —CH_2SO—, although more polar than —CH_2S—, did not remedy the situation. The methyl-substituted, chiral bond amide replacements, —$CHCH_3S$—, proved to be the most successful of the sulfur-based amide bond replacements, with *in vitro* potencies equivalent to the parent peptide bond analogs though they were still considerably less potent *in vivo* than anticipated.

Downloaded from https://academic.oup.com/edrv/article/7/1/44/2548841 by University of California, Los Angeles user on 19 April 2025

Cyclic antagonists: computer-assisted analog design

Early attempts to impose cyclic constraints on GnRH analogs led to agonists with relatively low potency or no activity at the doses tested (22, 79, 80). Nevertheless, the results of Seprodi *et al.* (80) were interesting in view of the fact that the N- to C-terminal cyclic analogs, cyclo[-β-Ala^1,D-Ala^6,Gly^{10}-]GnRH and cyclo[-6-aminohexanoic acid^1,D-Ala^6,Gly^{10}-]GnRH, having 1.2% and 0.65% of the potency of GnRH, respectively, *in vivo*, were more potent than the corresponding linear analogs. The corresponding 2-D-Nal^6 analogs exhibited 60% and 20% intrinsic activity, respectively, when tested *in vitro* and exhibited high binding affinity, [dissociation constant (K_d) = 5 and 0.5 mM as compared to K_d = 5 nM for GnRH (185)]. This partial antagonist activity could not have been uncovered *in vivo*, thus explaining the apparent discrepancies between these results and those of Seprodi *et al.* (80). In conclusion, Rivier *et al.* (185) proposed that GnRH interacts with its receptor in a folded conformation that can be locked in by covalent bonding of the C and N termini. Most GnRH antagonists have been characterized by the lack of L-His^2 or by the substitution of L-His^2 by a D-amino acid, which resulted in selectively impairing the transducing ability of the peptide. These cyclic analogs demonstrate that at least partial agonism can also be achieved through the introduction of conformational constraints.

Based upon the fact that GnRH antagonists with drastic changes in positions 1, 2, and 3 still exhibited high binding affinity for the GnRH receptor, a variety of cyclic antagonists, particularly D- and L-Cys^1,Cys^{10} residues, were examined by Rivier *et al.* (22). A number of Cys-containing antagonists exhibited *in vitro* potencies 3–20 times lower than [D-<Glu^1,D-Phe^2,D-Trp^{3,6}]GnRH, but had no antiovulatory activity at the doses and under the conditions tested. The most potent, and interesting, however, was cyclo[-Δ^3-Pro^1,4-Cl-D-Phe^2,D-Trp^{3,6},*N*-Me-Leu^7,β-Ala^{10}-]GnRH with an *in vitro* potency equivalent to that exhibited by [D-<Glu^1,D-Phe^2,D-Trp^{3,6}] GnRH. Struthers *et al.* (186, 187) compared this constrained cyclic antagonist with GnRH (using the theoretical techniques of molecular dynamics, valence force field energy minimization, and template forcing) and derived a computer simulation of their conformation, energetics, and dynamics. It was concluded that since the cyclic antagonist was extremely rigid and had essentially a single backbone conformation (but with differing side chain conformations), this particular backbone conformation must be responsible for the ability of the antagonist to bind to the receptor. Template forcing was then used to force GnRH along a minimum energy path into the cyclic antagonist conformation. At a moderate loss of strain energy, it was found that residues 4–9 of GnRH could adopt the conformation exhibited by residues 4–9 of the cyclic antagonist. Up to this point the best cyclic antagonist showed complete antiovulatory activity in the milligram range. However, when the information derived from the molecular dynamics studies and previously reported data on Cys$^{4-9}$ and Cys$^{4-10}$ GnRH agonists (22) were combined, several 4–10 cyclic antagonists were synthesized. One of those, *e.g.* [Ac-2-D-Nal^1,4-Cl-D-Phe^2,3-D-Pal^3,Dpr^4,Arg^5,3-D-Pal^6,Asp^{10}] GnRH, with its side chains optimized for water solubility, was the most potent cyclic antagonist yet reported (185). This analog, and the corresponding Cys$^{4-10}$ derivative, showed complete inhibition of ovulation at 10–25 μg/rat using a saline vehicle and K_d values of 4.5 and 0.85, respectively. These results further support the conclusion that GnRH does, indeed, interact with its receptor in a folded conformation that can be locked in by covalent bonding of the C to N termini or through side chains of amino acids in positions 4 and 10. The results of two-dimensional proton NMR spectroscopy studies (188) on several cyclic antagonists support the proposed conformation resulting from the molecular dynamics study. A smaller cyclic hexapeptide, cyclo(-Tyr-D-Trp-Leu-Arg-Trp-Pro-), was designed by Freidinger *et al.* (189) as an antagonist based on conformational considerations (*i.e.* the suggestion of the presence of a Tyr-Gly-Leu-Arg modified Type II β-turn in GnRH), the fact that the N-terminal region which precedes the β-turn is largely hydrophobic in the most potent antagonist, and the fact that enhancement of potency is observed with hydrophobic D-amino acids in place of Gly^6. This small cyclic peptide exhibited weak *in vitro* antagonist potency approximately 3 times greater than that of the early linear antagonist, [D-Phe^2]GnRH.

The use of the γ-lactam as a conformational constraint, which was successful with GnRH (78), resulted in high *in vivo* potency when applied to an antagonist. [Ac-2-D-Nal^1,α-Me-4-Cl-D-Phe^2,D-Trp^3,2-(3′-amino-2′-oxo-1′-pyrrolidino)-4-methyl-pentanoic acid^{6,7},D-Ala^{10}] GnRH completely inhibited ovulation at 10 μg (corn oil vehicle), demonstrating, again, that a β-bend at the 6,7-position of GnRH analogs is compatible with high potency (Roeske, R., N. Chaturvedi, T. Hrinyo, and K. Kopple, unpublished observations).

Optimization of antagonist potency

The trend toward optimization of GnRH antagonist antiovulatory potency with increasing substitution is noted in Table 1. The analogs represented therein are those exhibiting the maximum potency for the given number of amino acid substitutions. While it appears that there is a relationship between increasing substitution and increased antiovulatory potency, the relation-

Downloaded from https://academic.oup.com/edrv/article/7/1/44/2548841 by University of California, Los Angeles user on 19 April 2025

TABLE 1. Maximum antiovulatory potencies of GnRH antagonists for a given number of amino acid substitutions

1 2 3 4 5 6 7 8 9 10 <Glu-His-Trp-Ser-Tyr-Gly-Leu-Arg-Pro-Gly-NH_2 GnRH antagonists		Total Antiovulatory dose (μg) ED_{100}
No. of substitutions	Positions(s) of substitution(s)	
1	2	—[a]
2	2, 6	6000[b]
3	2, 3, 6	750[c]
4	1, 2, 3, 6	1[d]
5	1, 2, 3, 6, 10	0.5[e]
6	1, 2, 3, 6, 7, 10	0.5[f]

[a] [D-Phe^2]GnRH (39).
[b] [4-F-D-Phe^2,D-Ala^6]GnRH (110).
[c] [D-Phe^2,Pro^3,D-Trp^6]GnRH (139).
[d] [Ac-2-D-Nal^1,4-F-D-Phe^2,D-Trp^3,D-Arg^6]GnRH (27).
[e] [Ac-2-D-Nal^1,α-Me-4-Cl-D-Phe^2,D-Trp^3,D-Arg^6,D-Ala^{10}]GnRH (Roeske, R., N. Chaturvedi, and T. Hrinyo, unpublished observations).
[f] [Ac-2-D-Nal^1,4-Cl-D-Phe^2,3-D-Pal^3,D-Arg^6,Trp^7,D-Ala^{10}]GnRH (179).

ship is, in fact, more complex. Merely increasing the number of amino acid substitutions does not necessarily lead to higher potency. The antagonists require precise topological features for high binding affinity to the receptor. The simple inversion of the D- and L-amino acid residues of a potent antagonist resulted in an analog [Ac-4-Cl-Phe^1,4-Cl-Phe^2,Trp^3,D-Ser^4,D-Tyr^5,Arg^6,D-Tyr^7,D-Arg^8,D-Pro^9,Ala^{10}]GnRH which had a relative binding affinity approximately 3000 times lower than that of its apparent mirror image [Ac-4-Cl-D-Phe^1,4-Cl-D-Phe^2,D-Trp^3,D-Arg^6,D-Ala^{10}]GnRH (Coy, D. H., M. V. Nekola, and M. J. Karten, unpublished observations). Folkers *et al.* (190), using the results of empirical energy calculations, synthesized an analog containing six D-amino acid substitutions: [Ac-Thr^1,D-Phe^2,D-Trp^3,D-Ser^4,D-Tyr^5,D-Trp^6,D-Arg^8]GnRH. Compared with the model tetrasubstituted analog [Ac-Thr^1,D-Phe^2,D-Trp^{3,6}]GnRH, it appeared to be equipotent *in vitro* and somewhat more potent *in vivo* with partial inhibition of ovulation at 25 μg, but, nevertheless, considerably less potent than the most potent tetrasubstituted analogs. Moreover, very potent superagonists with high binding affinity have been designed with only one or two substitutions, and these superagonists have K_d values equal to those of the superantagonists. Thus, the relationship between increased biological (*in vivo*) potency of antagonists and increased substitution are unclear, but conformationally related receptor binding effects and pharmacokinetic factors must be considered.

Peptide-induced histamine release

This completes the discussion of the development of GnRH antagonists with regard to attempts to design ever more potent analogs. Unfortunately, with the introduction of D-Arg and other basic side chains into position 6, came the unexpected finding by Schmidt *et al.* that a potent antagonist [Ac-2-D-Nal^1,4-F-D-Phe^2,D-Trp^3,D-Arg^6]GnRH produced transient edema of the face and extremities when administered sc in rats (191) at 1.25 mg/kg or 50–100 times the effective antiovulatory dose. This edematogenic effect was attributed to a change in vascular permeability, but it was not observed sc in mice, rabbits, or monkeys at the doses tested. Additionally, this analog was subsequently (192) shown to induce a cutaneous anaphylactoid-like reaction in rats, causing a dose-related whealing response at 10 μg/rat. At the same dose, 1.25 mg/kg, neither [Ac-Δ^3-Pro^1,4-F-D-Phe^2,2-D-Nal^{3,6}]GnRH nor the corresponding D-Trp^{3,6} analog, exhibited the edema effect (191). Similarly, a potent agonist, [N^r-Bzl-D-His^6,Pro^9-NEt]GnRH, did not exhibit this effect nor did it block the edema effects of the D-Arg^6 antagonist (191). Other potent D-Arg^6 (or D-Lys^6) antagonists showed similar edema effects (Naqvi, R. and M. Lindberg, unpublished observations). In a structure-function study of histamine release (from rat mast cells) triggered by GnRH analogs, Hook *et al.* (193) reported that [Ac-2-D-Nal^1,4-F-D-Phe^2,D-Trp^3,D-Arg^6]GnRH was 3000 times more potent than GnRH, while [D-Trp^6,Pro^9-NEt]GnRH was only 10 times more potent than GnRH, in triggering the release of histamine. By themselves, two basic side chains, in close proximity, are insufficient to impart high histamine-releasing activity to GnRH analogs. This is demonstrated by the fact that [D-Arg^6]GnRH and [Arg^6]GnRH are approximately 200 times less potent than [Ac-2-D-Nal^1,4-F-D-Phe^2,D-Trp^3,D-Arg^6]GnRH in triggering the release of histamine (Hook, W., M. J. Karten, and R. Siraganian, unpublished observations). The most potent analogs in triggering histamine release had a structural combination of a basic D-amino acid side chain at position 6 (in close proximity to the Arg^8) and a cluster of hydrophobic aromatic amino acids at the N terminus. This observation was confirmed for [Ac-4-Cl-D-Phe^{1,2},D-Trp^3,D-Arg^6,D-Ala^{10}]GnRH and for the corresponding Ac-2-D-Nal^1 analog; it was also observed for the corresponding Phe^7 modifications (194). These peptides also induced skin lesions similar to those caused by the histamine releaser, compound 48/80 (195), when tested in a cutaneous anaphylaxis test (194). Qualitatively consistent with the observation that [Ac-Δ^3-Pro^1,4-F-D-Phe^2,D-Trp^{3,6}]GnRH did not exhibit the edema effect at 1.25 mg/kg (191) or at 5 mg/kg (Naqvi, R., and M. Lindberg, unpublished observations) was the observation that this antagonist was 230 times less potent than the corresponding Ac-2-D-Nal^1-D-Arg^6 antagonist, in triggering the release of histamine (193). The transposition antagonist, [Ac-2-D-Nal^1,α-Me-4-Cl-D-Phe^2,D-Trp^3,Arg^5,D-Tyr^6,D-Ala^{10}]GnRH (182), was ap-

Downloaded from https://academic.oup.com/edrv/article/7/1/44/2548841 by University of California, Los Angeles user on 19 April 2025

TABLE 2. Nonhuman primate and/or clinical studies with GnRH antagonists

GnRH analog	References
[D-Phe^2,Phe^3,D-Phe^6]	(200)
[D-Phe^2,D-Trp^3,D-Phe^6]	(201, 202[a], 203[a], 204[a])
[D-Phe^2,Pro^3,D-Phe^6]	(25, 205)
[D-<Glu^1,D-Phe^2,D-Trp^{3,6}]	(25, 206[a])
[D-<Glu^1,D-Phe^2,D-Trp^3,D-Lys^6](N^{ϵ}-isophthaloyl-Lys^6 dimer)	(207[a])
Endo-Pro^{1a}[D-Phe^2,D-Trp^{3,6}]	(25, 205)
[Ac-Pro^1,D-Phe^2,D-Trp^{3,6}]	(25)
[Ac-Δ^3-Pro^1,4-Cl-D-Phe^2,D-Trp^{3,6}]	(208)
[Ac-Δ^3-Pro^1,4-F-D-Phe^2,D-Trp^{3,6}]	(209, 210, 211[a])
[Ac-Δ^3-Pro^1,4-Cl-D-Phe^2,2-D-Nal^{3,6}]	(212)
[Ac-Δ^3-Pro^1,4-F-D-Phe^2,2-D-Nal^{3,6}]	(119)
[Ac-Δ^3-Pro^1,4-Cl-D-Phe^2,D-Trp^{3,6},N-Me-Leu^7]	(208)
[Ac-D-Phe^1,4-Cl-D-Phe^2,D-Trp^{3,6}]	(213[a])
[Ac-D-Trp^1,4-Cl-D-Phe^2,D-Trp^3,D-Phe^6,D-Ala^{10}]	(214)
[Ac-4-Cl-D-Phe^{1,2},D-Trp^3,D-Phe^6,D-Ala^{10}]	(215[a])
[Ac-4-Cl-D-Phe^{1,2},D-Trp^3,D-Arg^6,D-Ala^{10}]	(216–221)
[Ac-2-D-Nal^1,4-F-D-Phe^2,D-Trp^3,D-Arg^6]	(222), ([b])[a]
[Ac-2-D-Nal^1,4-Cl-D-Phe^2,D-Trp^3,N^{ω},$N^{\omega'}$-diethyl-D-Har^6,D-Ala^{10}]	(223–225), ([c])[a]

[a] Denotes references to clinical studies.
[b] Crowley, W. F., unpublished results.
[c] Henzl, M., unpublished results.

proximately 20–40 times less potent than the D-Arg^6 antagonists in triggering histamine release; hence separation of the two arginine residues partially reduces the histamine release potential of these antagonists. The diethyl-homoarginine^6 analog, [Ac-2-D-Nal^1,4-F-D-Phe^2,D-Trp^3,N^{ω},$N^{\omega'}$-diethyl-Har^6,D-Ala^{10}]GnRH was half as potent as [Ac-2-D-Nal^1,4-F-D-Phe^2,D-Trp^3,D-Arg^6]GnRH in triggering the release of histamine (Hook, W., M. J. Karten, and R. Siraganian, unpublished observations). Thus, it appears that GnRH agonists and antagonists, as well as a variety of other peptides such as substance P, somatostatin, vasoactive intestinal peptide, gastrin, *etc.* (195–199), trigger the release of histamine, apparently as a function of certain structural parameters that are independent of other inherent biological activities. Current structural modifications of GnRH antagonists are being undertaken to drastically reduce the histamine release potential while maintaining and/or increasing the GnRH antagonist potency.

Clinical Explorations

The relatively few completed and current clinical trials as well as the more extensive nonhuman primate studies have been cited in Table 2. The earliest trisubstituted GnRH antagonists to be tested in nonhuman primates or humans contained D-Phe in positions 2 and 6 and an L- or a D-amino acid in position 3. The next generation of antagonists to be tested in primates, the tetrasubstituted analogs, all contained hydrophobic D-amino acids in positions 2, 3, and 6 (D-Phe^2 or 4-Cl (or F)-D-Phe^2 and D-Trp^{3,6} or 2-D-Nal^{3,6}) and an L- or a D-amino acid in position 1. The current generation of tetra- and pentasubstituted antagonists being explored clinically, or in nonhuman primates, all contain hydrophobic D-amino acids in positions 1, 2, and 3 [Ac-4-Cl-D-Phe^1 or Ac-2-D-Nal^1,4-Cl(or F)-D-Phe^2 and D-Trp^3] and a basic D-amino acid in position 6 (D-Arg^6 or N^{ω},$N^{\omega'}$-diethyl-Har^6). The pentasubstituted analogs also contain D-Ala in position 10.

Acknowledgments

We wish to acknowledge the contributions made by Drs. Rehan Naqvi and Majorie Lindberg of the EG & G Mason Research Institute (Worcester, MA) to the Center for Population Research's program on the testing of some of the GnRH analogs; and by Dr. Pemmaraju N. Rao of the Southwest Foundation for Biomedical Research (San Antonio, TX) who provided a constant supply of some commercially unavailable D-amino acids to the Center's contractors involved in the synthesis of some of the GnRH analogs.

References

1. Matsuo H, Baba Y, Nair RM, Arimura A, Schally AV 1971 Structure of the porcine LH- and FSH-releasing hormone. I. The proposed amino acid sequence. Biochem Biophys Res Commun 43:1334
2. Matsuo H, Arimura A, Nair RMG, Schally AV 1971 Synthesis of the porcine LH- and FSH-releasing hormone by the solid-phase method. Biochem Biophys Res Commun 45:822
3. Baba Y, Matsuo H, Schally AV 1971 Structure of the porcine LH- and FSH-releasing hormone. II. Confirmation of the proposed structure by conventional sequential analyses. Biochem Biophys Res Commun 44:459
4. Burgus R, Butcher M, Amoss M, Ling N, Monahan MW, Rivier J, Fellows R, Blackwell R, Vale W, Guillemin R 1972 Primary structure of the ovine hypothalamic luteinizing hormone-releasing factor (LRF). Proc Natl Acad Sci USA 69:278
5. Ziporyn T 1985 LHRH: Clinical applications growing. JAMA 253:469
6. Jones JH, Ridge B 1972 Peptide synthesis. In: Young GT (ed) Amino-acids, Peptides, and Proteins (Specialist Periodical Reports). Chemical Society, London, vol 4:309
7. Law HD 1974 Chemical structure and biological activity. In: Sheppard RC (ed) Amino-acids, Peptides, and Proteins (Specialist Periodical Reports). Chemical Society, London, vol 5:384
8. Camble R, Dutta AS 1975 Chemical structure and biological activity. In: Sheppard RC (ed) Amino-acids, Peptides, and Proteins (Specialist Periodical Reports). Chemical Society, London, vol 6:419
9. Schafer DJ 1976 Peptide synthesis. In: Sheppard RC (ed) Amino-acids, Peptides, and Proteins (Specialist Periodical Reports). Chemical Society, London, vol 7:247
10. Schafer DJ, Szelke M 1976 Chemical structure and biological activity. In: Sheppard RC (ed) Amino-acids, Peptides and Proteins (Specialist Periodical Reports). Chemical Society, London, vol 8:339
11. Ludescher U, Studer RO, Gillessen D 1978 Chemical structure and biological activity of hormones and related compounds. In: Sheppard RC (ed) Amino-acids, Peptides, and Proteins (Specialist Periodical Reports). Chemical Society, London, vol 9:427
12. Hallett A, Stachulski AV 1979 Peptide synthesis. In: Sheppard RC (ed) Amino-acids, Peptides, and Proteins (Specialist Periodical Reports). Chemical Society, London, vol 10:310
13. Galpin MFJ 1981 Peptide synthesis. In: Sheppard RC (ed) Amino-

Downloaded from https://academic.oup.com/edrv/article/7/1/44/2548841 by University of California, Los Angeles user on 19 April 2025

acids, Peptides, and Proteins (Specialist Periodical Reports). Royal Society of Chemistry, London, vol 11:320
14. Coy DH 1981 Chemical structure and biological activity of hormones and related compounds. In: Sheppard RC (ed) Aminoacids, Peptides, and Proteins (Specialist Periodical Reports). Royal Society of Chemistry, London, vol 12:464
15. Coy DH 1982 Chemical structure and biological activity of hormones and related compounds. In: Sheppard RC (ed) Aminoacids, Peptides, and Proteins (Specialist Periodical Reports). Royal Society of Chemistry, London, vol 13:325
16. Burgus R 1972 Peptide hormones of the hypothalamus and pituitary. In: Heinzelman RV (ed) Annual Reports in Medicinal Chemistry. Academic Press, New York, vol 7:194
17. White WF 1973 Peptide hormones of the hypothalamus and pituitary. In: Heinzelman RV (ed) Annual Reports in Medicinal Chemistry. Academic Press, New York, vol 8:204
18. Meienhofer J 1975 Peptide hormones of the hypothalamus and pituitary. In: Heinzelman RV (ed) Annual Reports in Medicinal Chemistry. Academic Press, New York, vol 10:202
19. Bell MR, Christiansen RG, Schane Jr HP 1979 Chemical control of fertility. In: Hess HJ (ed) Annual Reports in Medicinal Chemistry. Academic Press, New York, vol 14:168
20. Spatola AF 1981 Peptides of the hypothalamus. In: Hess HJ (ed) Annual Reports in Medicinal Chemistry. Academic Press, New York, vol 16:199
21. Hruby VJ, Krstenausky JL, Cody WL 1984 Recent progress in the rational design of peptide hormones and neurotransmitters. In: Hess HJ (ed) Annual Reports in Medicinal Chemistry. Academic Press, Orlando, FL, vol 19:303
22. Rivier J, Rivier C, Perrin M, Porter J, Vale WW 1981 GnRH analogs: structure activity relationships. In: Zatuchni GI, Shelton JD, Sciarra JJ (eds) LHRH Peptides as Female and Male Contraceptives. Harper & Row, Philadelphia, p 13
23. Spatola AF, Bettag AL, Agarwal NS, Saneii H, Vale WW, Bowers CY 1981 The role of peptide backbone modifications in increasing biological stability of LHRH analogs. In: Zatuchni GI, Shelton JD, Sciarra JJ (eds) LHRH Peptides as Female and Male Contraceptives. Harper & Row, Philadelphia, p 24
24. Coy DH, Nekola MV, Erchegyi J, Coy EJ, Schally AV 1981 Contraceptive effects of potent LHRH antagonist analogs. In: Zatuchni GI, Shelton JD, Sciarra JJ (eds) LHRH Peptides as Female and Male Contraceptives. Harper & Row, Philadelphia, p 37
25. Bowers CY, Folkers K, Friebel K, Lutz W, Reynolds GA, Momany F 1981 A critique on analog antagonists of LHRH. In: Zatuchni GI, Shelton JD, Sciarra JJ (eds) LHRH Peptides as Female and Male Contraceptives. Harper & Row, Philadelphia, p 46
26. Nestor Jr JJ 1984 Development of agonistic LHRH analogs. In: Vickery BH, Nestor Jr JJ, Hafez ESE (eds) LHRH and Its Analogs. MTP Press, Lancaster, UK, p 3
27. Rivier J, Rivier C, Perrin M, Porter J, Vale WW 1984 LHRH analogs as antiovulatory agents. In: Vickery BH, Nestor Jr JJ, Hafez ESE (eds) LHRH and Its Analogs. MTP Press, Lancaster, UK, p 11
28. Nestor Jr JJ, Ho TL, Tahilramani R, Horner BL, Simpson RA, Jones GH, McRae GI, Vickery BH 1984 LHRH agonists and antagonists containing very hydrophobic amino acids. In: Vickery BH, Nestor Jr JJ, Hafez ESE (eds) LHRH and Its Analogs. MTP Press, Lancaster, UK, p 23
29. Nekola MV, Coy DH 1984 LHRH antagonists in females. In: Vickery BH, Nestor Jr JJ, Hafez ESE (eds) LHRH and its Analogs. MTP Press, Lancaster, UK, p 125
30. Merrifield RB 1963 Solid phase peptide synthesis. I. The synthesis of a tetrapeptide. J Am Chem Soc 85:2149
31. Merrifield RB, Stewart JM, Jernberg N 1966 Instrument for automated synthesis of peptides. Anal Chem 38:1905
32. Stewart JM 1981 Pharmacology of LHRH and analogs. In: Zatuchni GI, Shelton JD, Sciarra JJ (eds) LHRH Peptides as Females and Male Contraceptives. Harper & Row, Philadelphia, p 2
33. Pietta PG, Marshall GR 1970 Amide protein and amide supports in solid phase peptide synthesis. J Chem Soc Chem Commun, p 650
34. Rivier J, Monahan M, Vale W, Grant G, Amoss M, Blackwell R, Guillemin R, Burguss R 1972 Solid phase peptide synthesis on a benzhydrylamine resin of LRF (luteinizing hormone releasing factor) and analogues including antagonists. Chimia 26:300
35. Burgus R, Rivier J 1976 Use of high pressure liquid chromatography in the purification of peptides. In: Loffet A (ed) Peptides 1976: Proceedings of the Fourteenth European Peptide Symposium. Editions de l'Universite dè Bruxelles, Bruxelles, Belgium, p 85
36. Immer H, Nelson VR, Revesz C, Sestanj K, Gotz M 1974 Luteinizing hormone-releasing hormone and analogs. Synthesis and biological activity. J Med Chem 17:1059
37. Fujino M, Kobayashi S, Obayashi M, Shinagawa S, Fukuda T, Kitada C, Nakayaga R, Yamazaki I, White WF, Rippel RH 1972 Structure-activity relationships in the C-terminal part of luteinizing hormone releasing hormone (LH-RH). Biochem Biophys Res Commun 49:863
38. Monahan MW, Amoss MS, Anderson HA, Vale W 1973 Synthetic analogs of the hypothalamic luteinizing hormone releasing factor with increased agonist or antagonist properties. Biochemistry 12:4616
39. Rees WAR, Foell TJ, Chai SY, Grant N 1974 Synthesis and biological activities of analogs of the luteinizing hormone-releasing hormone (LH-RH) modified in position 2. J Med Chem 17:1016
40. Grant GF, Vale W 1974 Hypothalamic control of anterior pituitary hormone secretion-characterized hypothalamic-hypophysiotropic peptides. In: James VHT, Martini L (eds) Current Topics in Experimental Endocrinology. Academic Press, New York, vol 2:38,62
41. Rivier JE, Brown MR 1978 Bombesin, bombesin analogues, and related peptides: effects on thermoregulation. Biochemistry 17:1766
42. Schally AV, Arimura A, Coy DH 1980 Recent approaches to fertility control based on derivatives of LH-RH. In: Munson PL, Diczfalusy E, Glover J, Olson RE (eds) Vitamins and Hormones. Academic Press, New York, vol 38:257
43. Redding TW, Kastin AJ, Gonzalez-Barcena D, Coy DH, Coy EJ, Schalch DS, Schally AV 1973 The half-life, metabolism and excretion of tritiated luteinizing hormone (LHRH) in man. J Clin Endocrinol Metab 37:626
44. Barron JL, Millar RP, Searle D 1982 Metabolic clearance and plasma half-disappearance time of D-Trp^6 and exogenous LHRH. J Clin Endocrinol Metab 54:1169
45. Corbin A, Bex FJ 1981 Physiology and contraceptive effects of LHRH and agonistic analogs in female animals. In: Zatuchni GI, Shelton JD, Sciarra JJ (eds) LHRH Peptides as Female and Male Contraceptives. Harper & Row, Philadelphia, p 68
46. Vale W, Rivier C, Brown M 1977 Regulatory peptides of the hypothalamus. Annu Rev Physiol 39:473
47. Nillius SJ, Bergquist C, Wide L 1978 Inhibition of ovulation in women by chronic treatment with a stimulatory LRH analogue—a new approach to birth control? Contraception 17:537
48. Hahn DW, McGuire JL, Vale W, Rivier J 1984 Biological assays utilized to characterize LHRH and its analogs. In: Vickery BH, Nestor Jr JJ, Hafez ESE (eds) LHRH and Its Analogs. MTP Press, Lancaster, UK, p 49
49. Clayton RN, Catt KJ 1980 Receptor binding affinity of gonadotropin-releasing hormone analogs: analysis by radioligand receptor assay. Endocrinology 106:1154
50. Perrin MH, Rivier JE, Vale WW 1980 Radioligand assay for gonadotropin releasing hormone: relative potencies of agonists and antagonists. Endocrinology 106:1289
51. Sievertsson H, Chang JK, Bogentoft C, Currie BL, Folkers K, Bowers CY 1971 Synthesis of the luteinizing releasing hormone of the hypothalamus and its hormonal activity. Biochem Biophys Res Commun 44:1566
52. Coy DH, Vilchez-Martinez JA, Coy EJ, Nishi N, Arimura A, Schally AV 1975 Perfluoroalkylamine derivatives of luteinizing hormone-releasing hormone. Biochemistry 14:1848

Downloaded from https://academic.oup.com/edrv/article/7/1/44/2548841 by University of California, Los Angeles user on 19 April 2025

53. Fujino M, Shinagawa S, Obayashi M, Kobagashi S, Fukuda T, Yamazaki I, Nakayama R, White WF, Rippel R 1973 Further studies on the structure-activity relationships in the C-terminal part of the luteinizing hormone-releasing hormone. J Med Chem 16:1144
54. Rudinger J 1971 The design of peptide hormone analogs. In: Ariens EJ (ed) Drug Design. Academic Press, New York, vol II:319
55. Coy DH, Coy EJ, Schally AV, Vilchez-Martinez J, Hirotsu Y, Arimura A 1974 Synthesis and biological properties of [D-Ala-6,des-GlyNH_2-10]LH-RH ethylamide, a peptide with greatly enhanced LH- and FSH-releasing activity. Biochem Biophys Res Commun 57:335
56. Fujino M, Yamazaki I, Kobayashi S, Fukuda T, Shinagawa S, Nakayama R, White WF, Rippel RH 1974 Some analogs of luteinizing hormone releasing hormone (LH-RH) having intense ovulation-inducing activity. Biochem Biophys Res Commun 57:1248
57. Vale W, Rivier C, Brown M, Ling N, Monahan M, Rivier J 1976 Pharmacology of hypothalamic regulatory peptides. Clin Endocrinol (Oxf) 5 [Suppl]: 261s
58. Coy DH, Labrie F, Savory M, Coy EJ, Schally AV 1975 LH-Releasing activity of potent LH-RH analogs *in vitro*. Biochem Biophys Res Commun 67:576
59. Coy DH, Vilchez-Martinez JA, Coy EJ, Schally AV 1976 Analogs of luteinizing hormone-releasing hormone with increased biological activity produced by D-amino acid substitutions in position 6. J Med Chem 19:423
60. Barron J, Griffiths E, Tsalacopoulos G, Millar RP 1984 Metabolism of [D-Trp^6]LHRH. In: Vickery BH, Nestor Jr JJ, Hafez ESE (eds) LHRH and Its Analogues. MTP Press, Lancaster, UK, p 411
61. Vilchez-Martinez JA, Coy DH, Arimura A, Coy EJ, Hirotsu Y, Schally AV 1974 Synthesis and biological properties of [Leu-6]-LH-RH and [D-Leu-6,des-Gly-NH_2^{10}]-LH-RH ethylamide. Biochem Biophys Res Commun 59:1226
62. Nestor Jr JJ, Ho TL, Simpson RA, Horner BL, Jones GH, McRae GI, Vickery BH 1982 Synthesis and biological activity of some very hydrophobic superagonist analogues of luteinizing hormone releasing hormone. J Med Chem 25:795
63. Corbin A 1982 From contraception to cancer: a review of the therapeutic applications of LHRH analogues as antitumor agents. Yale J Biol Med 55:27
64. Wass JA, Besser GM, Gomez-Pan A, Scanlon MF, Hall R, Kastin AJ, Coy DH, Schally AV 1979 Comparison of long-acting analogs of LHRH in man. Clin Endocrinol (Oxf) 10:419
65. Nadasdi L, Medzihradszky K 1981 A study of the applicability of QSAR calculation for peptide hormones. Biochem Biophys Res Commun 99:451
66. Ariens EJ 1971 Modulation of pharmacokinetics by molecular manipulation. In: Ariens EJ (ed) Drug Design. Academic Press, New York, vol II:1
67. Nestor Jr JJ, Ho TL, Tahilramani R, McRae GI, Vickery BH 1984 Long-acting LHRH agonists and antagonists. In: Labrie F, Belanger A, Dupont A (eds) LHRH and Its Analogues. Elsevier, Amsterdam, p 24
68. Rivier J, Rivier C, Perrin M, Porter J, Vale W 1981 Applications of the Hansch approach to design GnRH analogs using Topliss' manual method. In: Brunfeldt K (ed) Peptide 1980: Proceedings of the Sixteenth European Peptide Symposium. Scriptor, Copenhagen, p 566
69. Zeelen FJ 1983 The strategy of the development of peptide drugs. Chemtech, p 419
70. Chan RL, Chaplin MD 1985 Plasma binding of LHRH and nafarelin acetate, a highly potent LHRH agonist. Biochem Biophys Res Commun 127:673
71. Nestor Jr JJ, Tahilramani R, Ho TL, McRae GI, Vickery BH 1985 Potent LHRH agonists containing N^G, $N^{G'}$-dialkyl-D-homoarginines. In: Deber CM, Hruby VJ, Kopple KB (eds) Peptides: Structure and Function. Proceedings of the Ninth American Peptide Symposium. Pierce Chemical Co, Rockford, IL, p 557
72. Dutta AS, Furr BJA, Giles MB, Valcaccia B, Walpole AL 1978 Potent agonist and antagonist analogues of luliberin containing an azaglycine residue in position 10. Biochem Biophys Res Commun 81:382
73. Dutta AS, Furr BJA, Giles MB, Valcaccia B 1978 Synthesis and biological activity of highly active α-aza analogues of luliberin. J Med Chem 21:1018
74. König W, Sandow J, Geiger R 1975 Structure-function relationships of LH-RH/FSH-RH. In: Walter R, Meienhofer J (eds) Peptides: Chemistry, Structure and Biology. Proceedings of the Fourth American Peptide Symposium. Ann Arbor Science, Ann Arbor, MI, p 883
75. Benuk N, Marks N 1976 Differences in the degradation of hypothalamic releasing factors by rat and human serum. Life Sci 19:1271
76. Walter R, Neidle A, Marks N 1975 Significant differences in the degradation of Pro-Leu-Gly-NH_2 by human serum and that of other species. Proc Soc Exp Biol Med 148:98
77. Ling N, Vale W 1975 Analogs of luteinizing hormone releasing factor (LRF): Synthesis and biological activity of [(N^α-Me)Leu^7] LRF and [D-Ala^6,N^α-Me)-Leu^7]LRF. Biochem Biophys Res Commun 63:801
78. Freidinger RM, Veber DF, Perlow DS, Brooks JR, Sapperstein R 1980 Bioactive conformation of luteinizing hormone-releasing hormone: evidence from a conformationally constrained analog. Science 210:656
79. Donzel B, Rivier J, Goodman M 1977 Synthesis of a cyclic analog of the luteinizing hormone releasing factor: [Glu^4,D-Ala^6,Orn^7] LRF. Biopolymers 16:2587
80. Seprodi J, Coy DH, Vilchez-Martinez JA, Pedroza E, Huang WY, Schally AV 1978 Cyclic analogues of luteinizing hormone-releasing hormone with significant biological activities. J Med Chem 21:993
81. Spatola AF 1983 Peptide backbone modifications: a structure-activity analysis of peptides containing amide bond surrogates, conformational constraints, and related backbone replacements. In: Weinstein B (ed) Chemistry and Biochemistry of Amino Acids, Peptides, and Proteins. Marcel Dekker, New York, vol 7:267
82. Chorev M, Chaturvedi N, Hoeprich P, Pallai P, Goodman M 1979 Novel partially modified retro-inverso analogs of biologically active peptides. In: Gross E, Meienhofer J (eds) Peptides: Structure and Biological Function. Proceedings of the Sixth American Peptide Symposium. Pierce Chemical Co, Rockford, IL, p 455
83. Yabe Y, Miura C, Horikoshi H, Baba Y 1976 Analogues of luteinizing hormone-releasing hormone with modification in position 3. Chem Pharm Bull (Tokyo) 24:3149
84. Prasad KU, Roeske RW, Weitl FL, Vilchez-Martinez JA, Schally AV 1975 Structure-activity relationships in luteinizing hormone-releasing hormone. In: Walter R, Meienhofer J (eds) Peptides: Chemistry, Structure and Biology. Proceedings of the Fourth American Peptide Symposium. Ann Arbor Science, Ann Arbor, MI, p 871
85. Coy DH, Coy EJ, Schally AV 1975 Structure-activity relationship of LH and FSH releasing hormone. In: Marks N, Rodnight R (eds) Research Methods in Neurochemistry. Plenum Publishing, New York, vol 3:393
86. Sherwood N, Eiden L, Brownstein M, Spiess J, Rivier J, Vale W 1983 Characterization of a teleost gonadotropin-releasing hormone. Proc Natl Acad Sci USA 80:2794
87. Miyamoto K, Hasegawa Y, Minegishi T, Nomura M, Takahashi Y, Igarashi M, Kanagawa K, Matsuo H 1982 Isolation and characterization of chicken hypothalamic luteinizing hormone-releasing hormone. Biochem Biophys Res Commun 107:820
88. King JA, Millar RP 1982 Structure of chicken hypothalamic and luteinizing hormone-releasing hormone. II. Isolation and characterization. J Biol Chem 257:10729
89. King JA, Tobler CJ, Roeske RW, Day WA, Rivier JE, Millar RP 1983 A radioimmunoassay specific for [Gln^8]LH-RH: application in the confirmation of the structure of chicken hypothalamic luteinizing hormone-releasing hormone. Peptides 4:883
90. Miyamoto K, Hasegawa Y, Nomura M, Igarashi M, Kangawa K, Matsuo H 1984 Identification of the second gonadotropin-releasing hormone in chicken hypothalamus: evidence that gonadotro-

Downloaded from https://academic.oup.com/edrv/article/7/1/44/2548841 by University of California, Los Angeles user on 19 April 2025

pin secretion is probably controlled by two distinct gonadotropin-releasing hormones in avian species. Proc Natl Acad Sci USA 81:3874
91. Millar RP, King JA 1983 Synthesis and biological activity of [D-Trp^6] chicken luteinizing hormone-releasing hormone. Peptides 4:425
92. Peter RE, Nahorniak CS, Sokolowska M, Chang JP, Rivier JE, Vale WW, King JA, Millar RP 1985 Structure-activity relationships of mammalian, chicken and salmon gonadotropin releasing hormones *in vivo* in goldfish. Gen Comp Endocrinol 58:231
93. King JA, Millar RP 1979 Heterogeneity of vertebrate luteinizing hormone releasing hormone. Science 206:67
94. Rivier J, Rivier C, Branton D, Millar R, Spiess J, Vale W 1981 HPLC purification of ovine CRF, rat extra hypothalamic brain somatostatin and frog brain GnRH. In: Rich DH, Gross E (eds) Peptides: Synthesis-Structure-Function. Proceedings of the Seventh American Peptide Symposium. Pierce Chemical Co, Rockford, IL, p 771
95. Flouret G, Stetler-Stevenson MA, Carone FA, Peterson DR 1984 Enzymatic degradation of LHRH and analogs. In: Vickery BH, Nestor Jr JJ, Hafez ESE (eds) LHRH and Its Analogs. MTP Press, Lancaster, UK, p 397
96. Vilchez-Martinez JA, Coy DH, Coy EJ, Schally AV, Arimura A 1975 Antiluteinizing hormone (LH)-releasing activity of several analogues of LH-releasing hormone. Fertil Steril 26:554
97. Koch Y, Baram T, Chobsieng P, Fridkin M 1974 Enzymatic degradation of luteinizing hormone-releasing hormone (LHRH) by hypothalamic tissue. Biochem Biophys Res Commun 61:95
98. Marks N, Stern F 1974 Enzymatic mechanisms for the inactivation of luteinizing hormone-releasing hormone (LHRH). Biochem Biophys Res Commun 61:1458
99. Koch Y, Baram T, Hazum E, Fridkin M 1977 Resistance to enzymatic degradation of LHRH analogues possessing increased biological activity. Biochem Biophys Res Commun 74:488
100. Bauer K, Horsthemke B, Knisatschek H, Norvak P, Kleinhauf H 1979 Degradation of luliberin (LH-RF) by brain and pituitary tissue enzymes. Z Physiol Chem 360:229
101. Horsthemke B, Bauer K 1980 Characterization of a nonchymotrypsin-like endopeptidase from anterior pituitary that hydrolyzes luteinizing hormone-releasing hormone at the tyrosyl-glycine and histidyl-tryptophan bonds. Biochemistry 19:2867
102. Knisatschek H, Bauer K 1979 Characterization of "thyroliberin-deamidating enzyme" as a post-proline-cleaving enzyme. J Biol Chem 254:10936
103. Horsthemke B, Knisatschek H, Rivier J, Sandow J, Bauer K 1981 Degradation of luteinizing hormone-releasing hormone and analogs by adenohypophyseal peptidases. Biochem Biophys Res Commun 100:753
104. Flouret G, Carone FA, Peterson DR, Ferronyalka E 1985 Effects of D-amino acid substituents on sites of LHRH analog cleavage by renal tissue. In: Deber CM, Hruby VJ, Kopple KB (eds) Peptides: Structure and Function. Proceedings of the Ninth American Peptide Symposium. Pierce Chemical Co, Rockford, IL, p 553
105. Schally AV, Kastin AJ 1971 Stimulation and inhibition of fertility through hypothalamic agents. Drug Ther Bull 1:29
106. Corbin A, Beattie CW 1975 Inhibition of the pre-ovulatory proestrous gonadotropin surge, ovulation and pregnancy with a peptide analogue of luteinizing hormone releasing hormone. Endocr Res Commun 2:1
107. Vilchez-Martinez JA, Coy DH, Coy EJ, Arimura A, Schally AV 1976 Comparison of the anti-LH/FSH-RH and anti-ovulatory activities of [D-Phe^2,D-Leu^6]-LHRH and [D-Phe^2,D-Ala^6]-LH-RH. Endocr Res Commun 3:231
108. De la Cruz A, Coy DH, Schally AV, Coy EJ, de la Cruz KG, Arimura A 1975 Blockade of the pre-ovulatory LH surge in hamsters by an inhibitory analog of LHRH (38855). Proc Soc Exp Biol Med 149:576
109. Yardley JP, Foell TJ, Beattie CW, Grant NH 1975 Antagonism of luteinizing hormone release and of ovulation by an analog of the luteinizing hormone-releasing hormone. J Med Chem 18:1244
110. Beattie CW, Corbin A, Foell TJ, Garsky V, McKinley WA, Rees RWA, Sarantakis D, Yardley JP 1975 Luteinizing hormone-releasing hormone. Antiovulation activity of analogs substituted in position 2 and 6. J Med Chem 18:1247
111. Bowers CY, Humphries J, Wasiak T, Folkers K, Reynolds GA, Reichert Jr LE 1980 On the inhibitory effects of luteinizing hormone-releasing hormone analogs. Endocrinology 106:674
112. Vale W, Grant G, Amoss M, Blackwell R, Guillemin R 1972 Culture of enzymatically dispersed anterior pituitary cells: functional validation of a method. Endocrinology 91:562
113. Perrin MH, Haas Y, Rivier JE, Vale WW 1983 Gonadotropin-releasing hormone binding to rat anterior pituitary membrane homogenates. Comparison of antagonists and agonists using radiolabeled antagonist and agonist. Mol Pharmacol 23:44
114. Conn PM, Rogers DC, Stewart JM, Niedel J, Sheffield T 1982 Conversion of a GnRH antagonist to an agonist: implication for a receptor microaggregate as the functional unit for signal transduction. Nature 296:653
115. Hazum E, Keinan D 1983 Gonadotropin releasing-hormone receptors: photoaffinity labeling with an antagonist. Biochem Biophys Res Commun 110:116
116. Hazum E 1981 Some characteristics of GnRH receptors in rat pituitary membranes: differences between an agonist and an antagonist. Mol Cell Endocrinol 23:275
117. Rivier C, Rivier J, Perrin M, Vale W 1983 Comparison of the effect of several gonadotropin releasing hormone antagonists on luteinizing hormone secretion, receptor binding and ovulation. Biol Reprod 29:374
118. Rivier C, Rivier J, Vale WW 1981 GnRH antagonists: physiologic and contraceptive applications in the female rat. In: Zatuchni GI, Shelton JD, Sciarra JJ, (eds) LHRH Peptides as Female and Male Contraceptives. Harper & Row, Philadelphia, p 140
119. McRae GI, Vickery BH, Nestor Jr JJ, Bremner WJ, Badger TM 1984 Biological activity of a highly potent LH-RH antagonist. In: Vickery BH, Nestor Jr JJ, Hafez ESE (eds) LH-RH and Its Analogs. MTP Press, Lancaster, UK, p 137
120. Rivier C, Vale W, Rivier J 1983 Effects of gonadotropin releasing hormone agonists and antagonists on reproductive functions. J Med Chem 26:1545
121. Rivier C, Rivier J, Vale W 1980 Anti-reproductive effects of a potent GnRH antagonist in the male rat. Science 210:93
122. Heber D, Swerdloff RS 1984 LHRH antagonists for male contraception. In: Vickery BH, Nestor Jr JJ, Hafez ESE (eds) LHRH and Its Analogs. MTP Press, Lancaster, UK, p 153
123. Heber D, Dodson R, Peterson M, Channabasavaiah KC, Stewart JM, Swerdloff RS 1984 Counteractive effects of agonistic and antagonistic gonadotropin-releasing hormone analogs on spermatogenesis: sites of action. Fertil Steril 41:309
124. Debeljuk L, Maines VM, Coy DH, Schally AV 1983 Effects of a powerful antagonist of LH-RH on testicular function in prepubertal male rats. Arch Androl 11:89
125. Sundaram K, Schmidt F, Thau RB, Rivier J, Vale W, Bardin CW 1984 Species differences in the sensitivity to the antitesticular effects of [Ac-D-Nal(2)^1,D-pF-Phe^2,D-Trp^3,D-Arg^6]LHRH, a potent LHRH antagonist. Contraception 29:271
126. Vale W, Grant G, Rivier J, Monahan M, Amoss M, Blackwell R, Burgus R, Guillemin R 1972 Synthetic polypeptide antagonists of the hypothalamic luteinizing hormone releasing factor. Science 176:933
127. Yanaihara N, Yanaihara C, Sakagami M, Tsuji K, Hashimoto T, Kaneko T, Oka H, Schally AV, Arimura A, Redding T 1973 Synthesis and biological evaluation of LH and FSH releasing hormone and its analogs. J Med Chem 16:373
128. Rivier J, Amoss MS, Rivier C, Vale W 1974 Synthetic luteinizing hormone releasing factor. Short chain analogs. J Med Chem 17:230
129. Coy DH, Vilchez-Martinez JA, Coy EJ, Arimura A, Schally AV 1973 A peptide inhibitor of luteinizing hormone-releasing hormone (LH-RH). J Clin Endocrinol Metab 37:331
130. Amoss Jr MS, Monahan M, Rivier J, Guillemin R, Inhibition of LRF induced LH release *in vivo* by LRF analogues. Program of the 56th Annual Meeting of the Endocrine Society, Atlanta, GA, 1974, p A-78 (Abstract)

Downloaded from https://academic.oup.com/edrv/article/7/1/44/2548841 by University of California, Los Angeles user on 19 April 2025

131. Vilchez-Martinez JA, Schally AV, Coy DH, Coy EJ, Debeljuk L, Arimura A 1974 *In vivo* inhibition of LH release by a synthetic antagonist of LH-releasing hormone (LH-RH). Endocrinology 95:213
132. Monahan M, Vale W, Rivier C, Grant G, Guillemin R, Analogues of LRF with inhibitory action or greater potency than the natural deca-peptide hormone. Program of the 55th Annual Meeting of the Endocrine Society, Chicago, IL, 1973, p A-145 (Abstract)
133. Vilchez-Martinez JA, Schally AV, Coy DH, Coy EJ, Miller CM, Arimura A 1975 An *in vivo* assay for anti-LH-RH and anti-FSH-RH activity of inhibitory analogues of LH-RH. Endocrinology 96:1130
134. Donzel B, Rivier J, Goodman M 1974 Conformational studies on the hypothalamic releasing factor and related compounds by ^{1}H nuclear magnetic resonance spectroscopy. Biopolymers 13:2631
135. Humphries J, Fisher G, Wan YP, Folkers K, Bowers CY 1974 Analogs of the luteinizing hormone-releasing hormone to study conformational aspects of the aromatic amino acid moieties and inhibition. J Med Chem 17:569
136. Wan YP, Humphries J, Fisher G, Folkers K, Bowers CY 1976 Inhibitors of the luteinizing hormone-releasing hormone based upon modifications in the 2, 3 and 6 positions. J Med Chem 19:199
137. Coy DH, Nikolics K, Seprodi JA, Teplan I, Coy EJ, Schally AV 1975 Synthesis of D-Phe^{2},Phe^{3},D-Phe^{6}-LHRH, a potent inhibitor of LH-RH. Acta Chim (Budapest) 87:423
138. De la Cruz A, Coy DH, Vilchez-Martinez JA, Arimura A, Schally AV 1976 Blockade of ovulation in rats by inhibitory analogs of luteinizing hormone-releasing hormone. Science 191:195
139. Humphries J, Wan YP, Folkers K, Bowers CY 1976 Presence of proline in position 3 for potent inhibition of the activity of the luteinizing hormone releasing hormone and of ovulation. Biochem Biophys Res Commun 72:939
140. Rivier JE, Vale WW 1978 [D-p-Glu^{1},D-Phe^{2},D-Trp^{3,6}]-LRF. A potent luteinizing hormone releasing factor antagonist *in vitro* and inhibitor of ovulation in the rat. Life Sci 23:869
141. Vale W, Rivier C, Brown M, Rivier J 1977 Pharmacology of thyrotropin releasing factor (TRF), luteinizing hormone releasing factor (LRF) and somatostatin. In: Porter JC (ed) Hypothalamic Peptide Hormones and Pituitary Regulation. Advances in Experimental Medicine and Biology. Plenum Press, New York, p 123
142. Seprodi J, Coy DH, Vilchez-Martinez JA, Pedroza E, Schally AV 1978 Branched-chain analogues of luteinizing hormone-releasing hormone. J Med Chem 21:276
143. Folkers K, Bowers CY, Stepinski J, Plucinski T, Sakagami M, Kubiak T, 1984 Analogs of the luteinizing hormone releasing hormone having the azagly^{10} moiety with antiovulatory activity. Z Naturforsch [B] 39:528
144. Sandow JK, Jerabek-Sandow G, Krauss B, Stoll W 1981 Metabolic and dispositional studies with LHRH analogs. In: Zatuchni GI, Shelton JD, Sciarra JJ (eds) LHRH Peptides as Female and Male Contraceptives. Harper & Row, Philadelphia, p 338
145. Chang JK, Williams RH, Humphries AJ, Johansson NG, Folkers K, Bowers CY 1972 Luteinizing releasing hormone, synthesis and Arg^{8}-analogs, and conformation-sequence-activity relationships. Biochem Biophys Res Commun 47:727
146. Grant G, Vale W 1972 Speculations on structural relationships between the hypothalamic releasing factors of pituitary hormones. Nature 237:182
147. Deslauriers R, Levy GC, McGregor WH, Sarantakis D, Smith ICP 1975 Conformational flexibility of luteinizing hormone releasing hormone (LH-RH) in aqueous solution. A carbon-13 spin-lattice relaxation time study. Biochemistry 14:4335
148. Wessels PL, Feeney J, Gregory H, Gormley JJ 1973 High resolution nuclear magnetic resonance studies of the conformation of luteinizing hormone releasing hormone (LH-RH) and its component peptides. J Chem Soc [Perkin II], p 1691
149. Marche P, Morgat JL, Fromageot P 1973 Solvent effects on luteinizing and follicle-stimulating-hormone releasing factor polymorphism studied by circular dichroism. Eur J Biochem 40:513
150. Mabrey S, Klotz IM 1976 Conformation of gonadotropin releasing hormone. Biochemistry 15:234
151. Momany FA 1976 Conformational energy analysis of the molecule, luteinizing hormone-releasing hormone. 1. Native decapeptide. J Am Chem Soc 98:2990
152. Momany FA 1976 Conformational energy analysis of the molecule, luteinizing hormone-releasing hormone. 2. Tetrapeptide and decapeptide analogues. J Am Chem Soc 98:2996
153. Momany FA 1978 Conformational analysis of the molecule luteinizing hormone-releasing hormone. 3. Analog inhibitors and antagonists. J Med Chem 21:63
154. Deslauriers R, Walter R, Smith ICP 1973 ^{13}C nuclear magnetic resonance studies of the conformation of the X—Pro bond in the oligopeptide hormones, thyrotropin-releasing hormone, luteinizing hormone-release factor, angiotensin and melanocyte-stimulating hormone release-inhibiting factor. Biochem Biophys Res Commun 53:244
155. Humphries J, Wan YP, Wasiak T, Folkers K 1979 Structural requirements in positions 1, 2, 3, and 6 of the luteinizing hormone-releasing hormone (LH-RH) for antiovulatory activity. J Med Chem 22:774
156. Coy DH, Mezo I, Pedroza E, Nekola MV, Vilchez J, Piyachaturawat P, Schally AV, Seprodi J, Teplan I 1979 LH-RH antagonists with potent antiovulatory activity. In: Gross E, Meinhofer J (eds) Peptides: Structure and Biological Function. Sixth American Peptide Symposium. Pierce Chemical Co, Rockford, IL, p 775
157. Humphries J, Wan YP, Folkers K, Bowers CY 1977 On the importance of position one of ovulation inhibitors, as based on studies on [D-Phe^{2},Pro^{3},D-Phe^{6}]-LHRH. Biochem Biophys Res Commun 78:506
158. Cann JR, Channabasavaiah K, Stewart JM 1979 Circular dichroism study of the solution conformation of luteinizing hormone releasing hormone. Biochemistry 18:5776
159. Kopple KD 1981 Peptide backbone folding in LHRH and analogs. In: Rich DH, Gross E (eds) Peptides: Synthesis-Structure-Function. Proceedings of the Seventh American Peptide Symposium. Pierce Chemical Co, Rockford, IL, p 295
160. Rapp J, Kerling KET 1981 Studies on polypeptides. XXXIV. Replacement of histidine-2 in luliberin (LH-RH) by L-homohistidine. Recl Trav Chim Pays-Bas 100:62
161. Humphries J, Wasiak T, Wan YP, Folkers K, Bowers CY 1978 An antiovulatory decapeptide of higher potency which has an L-amino acid (Ac-Pro) in position 1. Biochem Biophys Res Commun 85:709
162. Channabasavaiah K, Stewart JM 1979 New analogs of luliberin which inhibit ovulation in the rat. Biochem Biophys Res Commun 86:1266
163. Spatola AF, Agarwal NS 1980 A highly potent antiovulatory LH-RH analogue with no 1-position side chain. Biochem Biophys Res Commun 97:1571
164. Topliss JG 1977 A manual method for applying the Hansch approach to drug design. J Med Chem 20:463
165. Erchegyi J, Coy DH, Nekola MV, Coy EJ, Schally AV, Mezo I, Teplan I 1981 Luteinizing hormone-releasing hormone analogs with increased antiovulatory activity. Biochem Biophys Res Commun 100:915
166. Erchegyi J, Coy DH, Nekola MV, Pedroza E, Coy EJ, Mezo I, Schally AV 1981 LH-RH antagonists: further analogs with ring-substituted aromatic residues. Peptides 2:251
167. Vilchez-Martinez JA, Coy DH, Coy EJ, Arimura A, Schally AV 1976 Prolonged anti-luteinizing hormone/follicle-stimulating hormone-releasing activities of some synthetic antagonists of luteinizing hormone-releasing hormone. Fertil Steril 27:628
168. Coy DH, Horvath A, Nekola MV, Coy EJ, Erchegyi J, Schally AV 1982 Peptide antagonists of LH-RH: large increases in antiovulatory activities produced by basic D-amino acids in the six position. Endocrinology 110:1445
169. Horvath A, Coy DH, Nekola MV, Coy EJ, Schally AV, Teplan I 1982 Synthesis and biological activity of LH-RH antagonists modified in position 1. Peptides 3:969
170. Coy DH, Nekola MV 1984 Structure-function of LH-RH analogs and design applications to other peptide systems. In: McKerns KW, Naor Z (eds) Hormonal Control of the Hypothalamo-Pituitary-Gonadal Axis. Plenum Press, New York, p 521
171. Hocart SJ, Nekola MV, Coy DH 1985 Improved antagonists of

Downloaded from https://academic.oup.com/edrv/article/7/1/44/2548841 by University of California, Los Angeles user on 19 April 2025

luteinizing hormone-releasing hormone modified in position 7. J Med Chem 28:967
172. Nekola MV, Coy DH 1984 LHRH antagonists in females. In: Vickery BH, Nestor Jr JJ, Hafez ESE (eds) LHRH and Its Analogs. MTP Press, Lancaster, UK, p 125
173. Nekola MV, Horvath A, Ge LJ, Coy DH, Schally AV 1982 Suppression of ovulation in the rat by an orally active antagonist of luteinizing hormone-releasing hormone. Science 218:160
173a. Nestor Jr JJ, Tahilramani R, Ho TL, McRae GI, Vickery BH 1984 Luteinizing hormone-releasing hormone antagonists containing very hydrophobic amino acids. J Med Chem 27:1170
174. Folkers K, Bowers CY, Kubiak T, Stepinski J 1983 Antagonists of the luteinizing hormone releasing hormone with pyridyl-alanines which completely inhibit ovulation at nanogram dosage. Biochem Biophys Res Commun 111:1089
175. Folkers K, Bowers CY, Kubiak T, Stepinski J 1983 Synthesis and antiovulatory activities in rats of analogs of the luteinizing hormone-releasing hormone having a moiety of β-(3-quinolyl)-D-α-alanine in positions 3 and 6. Z Naturforsch [B] 38:1253
176. Roeske RW, Anantharamaiah GM 1983 LHRH antagonists with restricted conformation: α-methyl residues and disulfides. In: Hruby VJ, Rich DH (eds) Peptides: Structure and Function. Proceedings of the Eighth American Peptide Symposium. Pierce Chemical Co, Rockford, IL, p 333
177. Nestor Jr JJ, Tahilramani R, Ho TL, McRae GI, Vickery BH 1983 New luteinizing hormone-releasing factor antagonists. In: Hruby VJ, Rich DH (eds) Peptides: Structure and Function. Proceedings of the Eighth American Peptide Symposium. Pierce Chemical Co, Rockford, IL, p 861
178. Schwyzer R, Gremlich HU, Gysin B, Sargent DF, Fringeli UP 1983 Specific interactions between peptide hormones and artificial lipid membranes. In: Hruby VJ, Rich DH (eds) Peptides: Structure and Function. Proceedings of the Eighth American Peptide Symposium. Pierce Chemical Co, Rockford, IL, p 657
179. Folkers K, Bowers CY, Shieh HM, Lui YZ, Xiao SB, Tang PFL, Chu JY 1984 Antagonists of the luteinizing hormone releasing hormone (LHRH) with emphasis on the Trp^7 of the salmon and chicken II LHRH's. Biochem Biophys Res Commun 123:1221
180. Milton RCL, King JA, Badminton MN, Tobler CJ, Lindsey GG, Fridkin M, Millar RP 1983 Comparative structure-activity studies on mammalian [Arg^8]LH-RH and chicken [Gln^8]LH-RH by fluorimetric titration. Biochem Biophys Res Commun 111:1082
181. Rivier J, Varga J, Porter J, Perrin M, Rivier C, Vale W, Struthers S, Hagler A 1984 Design of cyclic GnRH antagonists. In: Labrie F, Belanger A, DuPont A (eds) LHRH and Its Analogues. Elsevier, Amsterdam, p 19
182. Roeske RW, Chaturvedi N, Rivier J, Vale W, Porter J, Perrin M 1985 Substitution of Arg^5 for Tyr^5 in GnRH antagonists. In: Deber CM, Hruby VJ, Kopple KD (eds) Peptides: Structure and Function. Proceedings of the Ninth American Peptide Symposium. Pierce Chemical Co, Rockford, IL, p 561
183. Wasiak T, Humphries J, Folkers K, Bowers CY 1979 A new category of ovulation inhibitors. Linear LH-RH analogs having more than ten residues. Biochem Biophys Res Commun 86:843
184. Hansch C 1971 Quantitative structure activity relationships in drug design. In: Ariens EJ (ed) Drug Design. Academic Press, New York, vol I:271
185. Rivier J, Varga J, Porter J, Perrin M, Haas Y, Corrigan A, Rivier C, Struthers S, Hagler A, Vale W 1985 Potent conformationally constrained analogs of GnRH. In: Deber CM, Hruby VJ, Kopple KD (eds) Peptides: Structure and Function. Proceedings of the Ninth American Peptide Symposium. Pierce Chemical Co, Rockford, IL, p 541
186. Struthers RS, Rivier J, Hagler AT 1985 Molecular dynamics and minimum energy conformations of GnRH and analogs: a methodology for computer-aided drug design. Ann New York Acad Sci 439:81
187. Struthers RS, Rivier J, Hagler AT 1984 Theoretical simulation of conformation, energetics, and dynamics in the design of peptide analogs. In: Vida JA, Gordon M (eds) Conformationally Directed Drug Design: Peptides and Nucleic Acids as Templates or Targets. American Chemical Society, Washington, DC, p 239
188. Baniak EL, Gierasch LM, Hagler AT, Rivier J, Solmajer T, Struthers RS, Conformational analysis of a cyclic LH-RH antagonist by two-dimensional very high field proton NMR. Program of the Ninth American Peptide Symposium, Toronto, Canada, 1985, p 125 (Abstract)
189. Freidinger RM, Colton CD, Saperstein R, Brady EJ, Randall WC, Veber DF 1985 A cyclic hexapeptide LH-RH antagonist. In: Deber CM, Hruby VJ, Kopple KD (eds) Peptides: Structure and Function. Proceedings of the Ninth American Peptide Symposium. Pierce Chemical Co, Rockford, IL, p 549
190. Folkers K, Bowers CY, Momany F, Friebel KJ, Kubiak T, Maher J 1982 Antiovulatory potency and conformation of an antagonist of the lutenizing hormone-releasing hormone having six D-amino acids. Z Naturforsch [B] 37:872
191. Schmidt F, Sundaram K, Thau RB, Bardin CW 1984 [Ac-D-$Nal(2)^1$,4FD-Phe^2,D-Trp^3,D-Arg^6]-LHRH, a potent antagonist of LHRH, produces transient edema and behavioral changes in rats. Contraception 29:283
192. Hahn DW, McGuire JL, Vale WW, Rivier J 1985 Reproductive/endocrine and anaphylactoid properties of an LHRH antagonist, ORF 18260 [Ac-D-$Nal(2)^1$,4FDPhe^2,D-Trp^3,D-Arg^6]-GnRH. Life Sci 37:505
193. Hook WA, Karten M, Siraganian RP 1985 Histamine release by structural analogs of LHRH. Fed Proc 44:1323 (Abstract)
194. Nekola MV, O'Neil C, Morgan J, Coy DH 1984 Antagonists of luteinizing hormone releasing hormone (LHRH): potent releasers of histamine in rats. Clin Res 32:865A (Abstract)
195. Theoharides TC, Douglas WW 1981 Mast cell histamine secretion in response to somatostatin analogues: structural considerations. Eur J Pharmacol 73:131
196. Foreman J, Jordon C 1983 Histamine release and vascular changes induced by neuropeptides. Agents Actions 13:105
197. Irman-Florjanc T, Erjavec F 1983 Compound 48/80 and substance P induced release of histamine and serotonin from rat peritoneal mast cells. Agents Actions 13:138
198. Foreman JC, Piotrowski W 1984 Peptides and histamine release. J Allergy Clin Immunol 74:127
199. Tharp MD, Thirlby R, Sullivan TJ 1984 Gastrin induces histamine release from human cutaneous mast cells. J Allergy Clin Immunol 74:159
200. Hagino N, Coy DH, Schally AV, Arimura A 1977 Inhibition of release in the baboon by inhibitory analogs of luteinizing hormone releasing hormone. Hormone Metab Res 9:247
201. Gosselin RE, Fuller BG, Coy DH, Schally AV, Hobson WC 1979 Inhibition of gonadotropin release in chimpanzees by the LH-RH antagonist (D-Phe^2,D-Trp^3,D-Phe^6)-LH-RH. Proc Soc Exp Biol Med 161:21
202. Gonzalez-Barcena D, Kastin AJ, Coy DH, Nikolics K, Schally AV 1977 Suppression of gonadotropin release in man by an inhibitory analogue of luteinizing hormone-releasing hormone. Lancet 2:997
203. Canales ES, Monvelinsky H, Zarate A, Kastin AJ, Coy DH, Schally AV 1980 Suppressive effect of an inhibitory LH-RH analog on the gonadotropin response to LH-RH in normal women. Int J Fertil 25:190
204. Gonzalez-Barcena D, Kastin AJ, Schally AV, Coy DH, Vilchez-Martinez JA, Pedroza-Garcia E, Nikolics K, Seprodi J 1978 Inhibition of luteinizing hormone (LH)-releasing hormone (LH-RH)-induced LH and follicle-stimulating hormone release in man by synthetic competitive antagonistic LH-RH analogs. Fertil Steril 29:246
205. Wilks JW, Folkers K, Smith CW 1981 Activities of LHRH antagonists in rhesus monkeys during the periovulatory interval of the menstrual cycle and early pregnancy. In: Zatuchni GI, Shelton JD, Sciarra JJ (eds) LH-RH Peptides as Female and Male Contraceptives. Harper & Row, Philadelphia, p 151
206. Casper RF, Sheehan K, Erickson G, Yen SSC 1980 Neuropeptides and fertility control in the female. In: Zatuchni GI, Labbok MH, Sciarra JJ (eds) Research Frontiers in Fertility Regulation. Harper & Row, Hagerstown, MD, p 409
207. Schally AV, Coy DH 1983 Stimulatory and inhibitory analogs of LH-releasing hormone: basic and clinical studies. In: McCann

Downloaded from https://academic.oup.com/edrv/article/7/1/44/2548841 by University of California, Los Angeles user on 19 April 2025

SM, Dhindsa DS (eds) Role of Peptides and Proteins in Control of Reproduction. Elsevier Biomedical, New York, p 89
208. Graham CE, Gosselin RE, Vale WW, Rivier J, Hobson WC 1981 Antigonadotropic analysis of GnRH antagonists in higher primates in relation to ovulation inhibition. In: Zatuchni GI, Shelton JD, Sciarra JJ (eds) LH-RH Peptides as Female and Male Contraceptives. Harper & Row, Philadelphia, p 168
209. Pineda JL, Lee BC, Spiliotis BE, Rivier J, Brown TJ, Bercu BB 1983 Effect of GnRH antagonist, [Ac-Δ^3Pro^1,pFDPhe^2,D-Trp^{3,6}] GnRH, on pulsatile gonadotropin secretion in the castrate male primate. J Clin Endocrinol Metab 56:420
210. Bercu BB, Spiliotis BE, Lee BC, Brown TJ, Vale W, Rivier J, Nixon WE, Reid R 1984 Effects of a potent gonadotropin releasing hormone antagonist on pulsatile testosterone and gonadotropin secretion in the male nonhuman primate. Life Sci 35:381
211. Cetel NS, Rivier J, Vale W, Yen SS 1983 The dynamics of gonadotropin inhibition in women induced by an antagonistic analog of gonadotropin-releasing hormone. J Clin Endocrinol Metab 57:62
212. Siler-Khodr TM, Kuehl TJ, Vickery BH 1984 Effects of a gonadotropin-releasing hormone antagonist on hormonal levels in the pregnant baboon and on fetal outcome. Fertil Steril 41:448
213. Zarate A, Canales ES, Schally AV, Coy DH, Coumaru-Schally AM 1981 The use of LHRH agonists and antagonists as antifertility agents in the human female. In: Zatuchni GI, Shelton JD, Sciarra JJ (eds) LHRH Peptides as Female and Male Contraceptives. Harper & Row, Philadelphia, p 227
214. Asch RH, Balmaceda JP, Borghi M, 1984 LH-RH antagonists in rhesus and cynomolgus monkeys. In: Vickery BH, Nestor Jr JJ, Hafesz ESE (eds) LH-RH and Its Analogs. MTP Press, Lancaster, UK, p 107
215. Zarate A, Canales ES, Sthory I, Coy DH, Coumaru-Schally AM, Schally AV 1981 Antiovulatory effect of a LHRH antagonist in women. Contraception 24:315
216. Akhtar FB, Weinbauer GF, Nieschlag E 1985 Acute and chronic effects of a gonadotropin-releasing hormone antagonist on pituitary and testicular function in monkeys. J Endocrinol 104:345
217. Hodgen GD, Werlin LB, Kenigsberg D, Littman BA, Platia MP, Schenken RS 1985 Reversible pharmacologic castration of the female: implications for endometriosis, contraception and ovulation induction. In: Schmidt-Gollwitzer M, Schley R (eds) LHRH and Its Analogues—Fertility and Antifertility Aspects. Walter de Gruyter, Berlin, p 93
218. Kenigsberg D, Littman BA, Hodgen GD 1984 Medical hypophysectomy. I. Dose-response using a gonadotropin releasing hormone antagonist. Fertil Steril 42:112
219. Kenigsberg D, Littman BA, Williams RF, Hodgen GD 1984 Medical hypophysectomy. II. Variability of ovarian response to gonadotropin therapy. Fertil Steril 42:116
220. Burgos-Briceno LA, Schally AV, Bartke A, Asch RH 1984 Inhibition of serum luteinizing hormone and testosterone with an inhibitory analog of luteinizing hormone-releasing hormone in adult male rhesus monkeys. J Clin Endocrinol Metab 59:601
221. Asch RH, Balmaceda JP, Borghi MR, Niesvisky R, Coy DH, Schally AV 1983 Suppression of the positive feedback of estradiol benzoate on gonadotropin secretion by an inhibitory analog of luteinizing hormone-releasing hormone (LRH) in oophorectomized rhesus monkeys: evidence for a necessary synergism between LRH and estrogens. J Clin Endocrinol Metab 57:367
222. Mann DR, Gould KG, Patonay NR, Collins DC 1985 Effect of a GnRH antagonist on fertility in male monkeys. Biol Reprod [Suppl 1] 32:101 (Abstract)
223. Adams LA, Bremner WJ, Steiner RA 1985 Primate studies with a new, potent gonadotropin releasing hormone (GnRH) antagonist. Clin Res 33:24A (Abstract)
224. Fraser HM, Baird DT, McRae GI, Nestor JJ, Vickery BH 1985 Suppression of luteal progesterone secretion in the stumptailed macaque by an antagonist analogue of luteinizing hormone releasing hormone. J Endocrinol 104:R1
225. Weinbauer GF, Surmann FJ, Akhtar FB, Shah GV, Vickery BH, Nieschlag E 1984 Reversible inhibition of testicular function by a gonadotropin hormone-releasing hormone antagonist in monkeys (*Macaca fascicularis*). Fertil Steril 42:906

Downloaded from https://academic.oup.com/edrv/article/7/1/44/2548841 by University of California, Los Angeles user on 19 April 2025