

EUROPEAN RESPIRATORY JOURNAL

December 9, 2004

HEALTH SCIENCES
LIBRARIES

OFFICIAL
JOURNAL OF
THE EUROPEAN
RESPIRATORY
SOCIETY

On-line
www.ersnet.org

EDITORIAL

Peripheral lung remodelling in asthma and chronic obstructive pulmonary disease

ORIGINALS

Value of imprint cytology for ultrasound-guided transthoracic core biopsy

Predictive value of BAL cell differentials in the diagnosis of interstitial lung diseases

Comparison between exhaled and sputum oxidative stress bio markers in chronic airway inflammation

REVIEW

Diagnosis and pharmacotherapy of severe acute respiratory syndrome: what have we learnt?

SERIES

On the nature of *Mycobacterium tuberculosis*-latent bacilli

On-line submissions: <http://mc.manuscriptcentral.com/erj>

VOLUME 24 • NUMBER 6

DECEMBER 2004



EUROPEAN RESPIRATORY JOURNAL

CHIEF EDITORS

Peter J. Sterk and Klaus F. Rabe
Dept Pulmonology, C3-P, Leiden University Medical Center,
Albinusdreef 2, P.O. Box 9600, Leiden, NL-2300 RC,
The Netherlands
Tel: 31 71 5262950 Fax: 31 71 5266927 e-mail: erj@lumc.nl

SECRETARY TO THE CHIEF EDITORS

Mrs A. van der Mey
Tel: 31 71 5262950 Fax: 31 71 5266927 e-mail: erj@lumc.nl

PUBLICATIONS OFFICE

Linda Arnold
Publications Manager, Suite 2.4, Hutton's Building,
146 West Street, Sheffield, S1 4ES, UK
Tel: 44 114 278 0498 Fax: 44 114 278 0501
e-mail: Linda.Arnold@ersj.org.uk

SECTION EDITORS

J. Gerritsen
Groningen, The Netherlands

A.T. Dinh-Xuan
Paris, France

D. Nowak
Munich, Germany

A.M. Vignola
Palermo, Italy

C. Vogelmeier
Marburg, Germany

M. Woodhead
Manchester, UK

ASSOCIATE EDITORS

N. Ambrosino
Pisa, Italy

M. Belvisi
London, UK

C. Brambilla
Grenoble, France

P.M. Calverley
Liverpool, UK

P. Chanez
Montpellier, France

R. Djukanovic
Southampton, UK

S.G. von Essen
Omaha, USA

S. Ewig
Bochum, Germany

R. Farré
Barcelona, Spain

Y. Fukuchi
Tokyo, Japan

P.G. Gibson
Newcastle, Australia

J. Hedner
Gothenburg, Sweden

M. Kraft
Denver, USA

P.N. Le Souëf
Perth, Australia

C.E. Mapp
Ferrara, Italy

G. Massard
Strasbourg, France

J. Müller-Quernheim
Freiburg, Germany

M.I. Polkey
London, UK

G. Raghu
Seattle, USA

I. Rahman
Rochester, USA

S. Rennard
Omaha, USA

R. Rodriguez-Roisin
Barcelona, Spain

M. Saetta
Padova, Italy

T. Sandström
Umea, Sweden

A.M.W.J. Schols
Maastricht, The Netherlands

F. Sériès
Quebec City, Canada

E.K. Silverman
Boston, USA

S.G. Spiro
London, UK

J. Sunyer
Barcelona, Spain

A. Torres
Barcelona, Spain

C.S. Ulrik
Copenhagen, Denmark

J. Vestbo
Copenhagen, Denmark

U. Wahn
Berlin, Germany

K. Webb
Manchester, UK

S.G. Zakynthinos
Athens, Greece

A. Bankier
Vienna, Austria
Radiological Advisor

S. Chinn
London, UK
Statistical Advisor

C. Melot
Brussels, Belgium
Statistical Advisor

E. Berglund
Gothenburg, Sweden
Case for Diagnosis Advisor

R. Rodriguez-Roisin
Barcelona, Spain
Supplement Editor

Past Chief Editors: J-C. Yernault, Brussels, Belgium;
E. Berglund, Gothenburg, Sweden; P. Vermeire, Antwerp,
Belgium; U. Costabel, Essen, Germany; M. Decramer,
Leuven, Belgium.



CONTENTS

EDITORIALS

Volume 24, Number 6 December 2004

- 893** Peripheral lung remodelling in asthma and chronic obstructive pulmonary disease
J. Hogg
- 895** Controversial topics in tuberculosis
J.A. Caminero, A. Torres
- 896** Antibiotics in exacerbations of COPD: lessons from the past
M. Miravittles, A. Torres

CLINICAL FORUM: ONCOLOGY

- 898** Lung cancer: clinical presentation and specialist referral time
G. Bucchieri, D. Ferrigno
- 905** Value of imprint cytology for ultrasound-guided transthoracic core biopsy
W.-Y. Liao, J.-S. Jerng, K.-Y. Chen, Y.-L. Chang, P.-C. Yang, S.-H. Kuo

ORIGINAL ARTICLES

Bench to Bedside: Airway Remodelling

- 910** Airway remodelling assessed by sputum and high-resolution computed tomography in asthma and COPD
A.M. Vignola, F. Paganin, L. Capieu, N. Scichilone, M. Bellia, L. Maakel, V. Bellia, P. Godard, J. Bousquet, P. Chanez

Tissue Repair and Fibrosis

- 918** Thrombin induces collagen gel contraction partially through PAR1 activation and PKC- ϵ
Q. Fang, X. Liu, S. Abe, T. Kobayashi, X.Q. Wang, T. Kohyama, M. Hashimoto, T. Wyatt, S.I. Rennard

Paediatric Asthma

- 925** Influenza vaccination in asthmatic children: effects on quality of life and symptoms
H.J. Bueving, J.C. van der Wouden, H. Raat, R.M.D. Bernsen, J.C. de Jongste, L.W.A. van Suijlekom-Smit, A.D.M.E. Osterhaus, G.F. Rimmelzwaan, M. Rutten-van Molken, S. Thomas

- 932** Treatment of exercise-induced asthma with beclomethasone dipropionate in children with asthma
R. Petersen, L. Agertoft, S. Pedersen

Epidemiology

- 938** Ecological associations between asthma prevalence and potential exposure to farming
L. Elliott, K. Yeatts, D. Loomis
- 942** A prospective study of asthma incidence and its predictors: the RHINE study
K. Torén, T. Gislason, E. Omenaas, R. Jögi, B. Forsberg, L. Nyström, A.-C. Olin, C. Svanes, C. Janson, on behalf of the RHINE study group

ORIGINAL ARTICLES (CONT.)

COPD

- 947** Levofloxacin versus clarithromycin in COPD exacerbation: focus on exacerbation-free interval
H. Lode, J. Eller, A. Linnhoff, M. Ioanas, and the Evaluation of Therapy-Free Interval in COPD Patients Study Group
- 954** Assessments for oxygen therapy in COPD: are we under correcting arterial oxygen tensions?
K. Dheda, K. Lim, B. Ollivere, J. Leftley, F.C. Lampe, A. Salisbury, J.P. Dilworth, R.K. Rajakulasingum
- 958** Vasoactive intestinal peptide receptors in the airways of smokers with chronic bronchitis
D. Miotto, P. Boschetto, I. Bononi, E. Zeni, G. Cavallresco, L.M. Fabbri, C.E. Mapp

Surfactant and Emphysema

- 964** Effects of ageing and smoking on SP-A and SP-D levels in bronchoalveolar lavage fluid
T. Betsuyaku, Y. Kuroki, K. Nagai, Y. Nasuhara, M. Nishimura

Respiratory Muscles

- 971** Adaptation of the diaphragm and the vastus lateralis in mild-to-moderate COPD
M. Doucet, R. Debigaré, D.R. Joannisse, C. Côté, P. LeBlanc, J. Grégoire, J. Deslauriers, R. Vaillancourt, F. Maltais

Sleep-Disordered Breathing

- 980** Frontal brain lobe impairment in obstructive sleep apnoea: a proton MR spectroscopy study
M. Alchanatis, N. Deligiorgis, N. Zias, A. Amfilochiou, E. Gotsis, A. Karakatsani, A. Papadimitriou
- 987** Night-to-night variation of the oxygen desaturation index in sleep apnoea syndrome
I. Fietze, K. Dingli, K. Diefenbach, N.J. Douglas, M. Glos, M. Tallafuss, W. Terhalle, C. Witt

ARDS

- 994** Alveolar antioxidant status in patients with acute respiratory distress syndrome
R. Schmidt, T. Luboeinski, P. Markart, C. Ruppert, C. Daum, F. Grimminger, W. Seeger, A. Günther

Interstitial Lung Disease

- 1000** Predictive value of BAL cell differentials in the diagnosis of interstitial lung diseases
L. Welker, R.A. Jörres, U. Costabel, H. Magnussen

Pulmonary Hypertension

- 1007** Combination therapy with bosentan and sildenafil in idiopathic pulmonary arterial hypertension
M.M. Hoepfer, C. Faulenbach, H. Golpon, J. Winkler, T. Welte, J. Niedermeyer

Sputum and Exhaled Air

- 1011** Comparison between exhaled and sputum oxidative stress biomarkers in chronic airway inflammation
M. Corradi, P. Pignatti, P. Manini, R. Andreoli, M. Goldoni, M. Poppa, G. Moscato, B. Balbi, A. Mutti
- 1018** Granulocyte markers in hypertonic and isotonic saline-induced sputum of asthmatic subjects
S. Cianchetti, E. Bacci, L. Ruocco, M.L. Bartoli, M. Ricci, T. Pavia, F.L. Dente, A. Di Franco, B. Vagaggini, P.L. Paggiaro

EUROPEAN RESPIRATORY JOURNAL

AIMS & SCOPE

The *European Respiratory Journal*, the official Journal of The European Respiratory Society, publishes, in the English language, clinical and experimental work dealing with all aspects of respiratory medicine. In addition to original material, the Journal will print Editorials and Reviews as well as Technical notes, Case studies and Letters to the Editor.

This Journal is abstracted/indexed in: Ad Referendum, ADONIS, Biological Abstracts/BIOSIS, Chemical Abstracts, Current Contents: Life Sciences; Current Contents: Clinical Medicine, Elsevier BIOBASE/Current Awareness in Biological Sciences, EMBASE/Excerpta Medica, Index Medicus/MEDLINE, Medical Documentation Service, Research Alert, Science Citation Index, SciSearch and SIIC Data Base.

SUBSCRIPTIONS

European Respiratory Journal is published monthly by the European Respiratory Society (ERS). Twelve issues are published annually in two volumes of six issues each. The volumes for 2004 are Volumes 23 and 24. Subscribers will receive any supplements to the Journal free of charge.

Subscription rates for institutions (Volumes 23 & 24, 2004): £443.00; US\$757.00 in North America. Subscription rates include air priority postage. Subscriptions are payable in advance.

Subscribers within the European Union must provide their VAT (Value Added Tax) number or add VAT to the subscription rate at the rate prevailing in their own country.

Subscriptions are managed on behalf of the ERS by: Maney Publishing, Hudson Road, Leeds LS9 7DL, UK (fax +44 (0)113 248 6983; E-mail: subscriptions@maney.co.uk) or to: Maney Publishing North America, 875 Massachusetts Avenue, 7th Floor, Cambridge, MA 02139, USA (tel: (toll-free) 866 297 5154; fax: 617 354 6875; E-mail: maney@maneyusa.com). All orders, payments, and claims for missing issues should be sent to Maney Publishing or to any subscription agent.

ON-LINE ACCESS

All subscribers have on-line access to the full text of each issue as part of their subscription. The on-line version is available to members via www.ersnet.org. Institutional subscribers can access on-line through the Ingenta Journals Service at www.ingentaselect.com.

Members of the ERS receive the Journal as part of their membership.

Business matters (enquiries, advertisement bookings and individual copies) should be addressed to: *European Respiratory Journals Publications Office, Suite 2.4, Hutton's Building, 146 West Street, Sheffield S1 4ES, UK. Fax: +44 (0) 1142 780501. E-mail: info@ersj.org.uk*

EUROPEAN RESPIRATORY JOURNAL

TECHNICAL EDITORS

Carol Wilson
Fiona Shepherd
Pippa Powell
Fiona Marks
Catherine Biles
Mark Livingston
Elin Reeves

For all enquires concerning advertising or marketing please
e-mail marketing@ersj.org.uk

© 2004 ERS. Authorization to photocopy items for internal or personal use, or the internal or personal use of specific clients, is granted for libraries and other users registered with the Copyright Clearance Center (CCC) Transactional Reporting Service, provided that the base fee of USD 12.00 per copy is paid directly to CCC, 27 Congress Street, Salem, MA 01970. 0903-1936/03/USD 12.00. **Academic Permissions Services.** Prior to photocopying items for educational classroom use, please contact the Copyright Clearance Center, Customer Service, 222 Rosewood Drive, Danvers MA 01923, USA. (978)750-8400. All other rights, including microfilm, electronic and optical storing and/or copying reserved.

(USPS # 015-990) Periodicals postage paid at Rahway, N.J. and additional mailing offices. USA Postmaster for North American subscribers: send address changes to the European Respiratory Journal, c/o Mercury Airfreight International Ltd, 365 Blair Road, Avenel, NJ 07001. Printed in UK by H. Charlesworth & Co. Ltd.

"The advertiser shall be liable for the content of any of its advertisements published according to its instructions. The advertiser shall indemnify and hold harmless the European Respiratory Society against and from any and all claims, damages, liabilities, costs and expenses whatsoever, including reasonable counsel fees, arising from the content of any of its advertisements."

ERJ on-line at www.ersnet.org

REVIEWS

- 1025** Diagnosis and pharmacotherapy of severe acute respiratory syndrome: what have we learnt?
K.W. Tsang, G.C. Ooi, P.L. Ho
- 1033** The immune response to resistive breathing
T. Vassilakopoulos, C. Roussos, S. Zakynthinos

SERIES

- SERIES "CONTROVERSIAL ISSUES IN TUBERCULOSIS"**
Edited by A. Torres and J. Caminero
Number 1 in this Series
- 1044** On the nature of *Mycobacterium tuberculosis*-latent bacilli
P-J. Cardona, J. Ruiz-Manzano
- SERIES "RESPIRATORY MONITORING: REVISITING CLASSICAL PHYSIOLOGICAL PRINCIPLES WITH NEW TOOLS"**
Edited by M.I. Polkey, R. Farré, A.T. Dinh-Xuan
Number 2 in this Series
- 1052** Noninvasive monitoring of respiratory mechanics during sleep
R. Farré, J.M. Montserrat, D. Navajas

CASE FOR DIAGNOSIS

- 1062** Relapse of respiratory insufficiency one year after organising pneumonia
J-M. Naccache, M. Kambouchner, F. Girard, M. Antoine, A. Parrot, J. Piquet, M. Brauner, D. Valeyre
- 1066** A painful and swollen right breast in a young male
G.E. Kapotsis, Z. Daniil, S. Klimopoulos, K. Malagari, C. Roussos, D. Rontogianni, S.A. Papiris

CORRESPONDENCE

- 1069** Radicals in the church
T.M.C.M. de Kok, J.G.F. Hogervorst, J.C.S. Kleinjans, J.J. Briedé
- 1070** Combination therapy with maintenance budesonide and formoterol in COPD
A.R.L. Medford, P.M.A. Calverley
- 1071** Progressive damage on high-resolution computed tomography
D. Rawlings, D. Tennant, J. Furness, P.A. de Jong, M.H. Lequin, J.R. Mayo, P.D. Paré, H.A.W.M. Tiddens
- 1072** Variable phenotype associated with SP-C gene mutations: fatal case with the 173T mutation
S. Percopo, H.S. Cameron, L.M. Nogee, G. Pettinato, S. Montella, F. Santamaria
- 1074** ERRATUM
- 1075** AUTHORS' CORRECTION
- 1076** BOOK REVIEWS

d,f,h,i,p MEETING CALENDAR

SHORT REPORT

Combination therapy with bosentan and sildenafil in idiopathic pulmonary arterial hypertension

M.M. Hoeper*, C. Faulenbach*, H. Golpon*, J. Winkler[#], T. Welte*, J. Niedermeyer*

Combination therapy with bosentan and sildenafil in idiopathic pulmonary arterial hypertension. M.M. Hoeper, C. Faulenbach, H. Golpon, J. Winkler, T. Welte, J. Niedermeyer. ©ERS Journals Ltd 2004.

ABSTRACT: It has been proposed that targeted treatments should be combined for patients with idiopathic pulmonary arterial hypertension (IPAH) responding insufficiently to monotherapy.

This study followed the clinical course of nine patients with severe IPAH, in whom the endothelin receptor antagonist bosentan caused transient clinical improvement, eventually followed by a decline in exercise tolerance, who received adjunct treatment with the phosphodiesterase-5-inhibitor sildenafil. Measurements included the 6-min walk distance (6MWD) and cardiopulmonary exercise testing (CPET).

The 6MWD at baseline was 346 ± 66 m and improved to 403 ± 80 m 3 months after introduction of bosentan treatment. However, this effect was not sustained and, after an interval of 11 ± 5 months, the walk distance had declined to 277 ± 80 m. At this point, sildenafil was added to bosentan. Three months later, the 6MWD had increased to 392 ± 61 m and the patients remained stable throughout the median follow-up of 9 months (range 6–12). Measurement of the maximum oxygen uptake during CPET confirmed these results. The combination of bosentan and sildenafil was well tolerated by all patients.

These preliminary data suggest that combining bosentan and sildenafil may be safe and effective in patients with idiopathic pulmonary arterial hypertension.

Eur Respir J 2004; 24: 1007–1010.

*Dept of Respiratory Medicine, Hannover Medical School, Hannover, and [#]Dept of Cardiopulmonary Medicine, University of Leipzig, Leipzig, Germany.

Correspondence: M.M. Hoeper
Hannover Medical School
Dept of Respiratory Medicine
Carl-Neuberg-Str. 1
30625 Hannover
Germany
Fax: 49 5115328536
E-mail: hoeper.marius@mh-hannover.de

Keywords: Bosentan
endothelin receptor antagonists
hypertension
phosphodiesterase inhibitors
pulmonary
sildenafil

Received: April 29 2004
Accepted after revision: July 7 2004

Idiopathic pulmonary arterial hypertension (IPAH), formerly called primary pulmonary hypertension (PPH), is a disease of unknown aetiology, characterised by progressive pulmonary vascular remodelling, ultimately leading to right heart failure and death [1, 2]. Without targeted treatment, IPAH carries a grave prognosis, with a median survival of 2.8 yrs after diagnosis [3].

In the last few years, a number of novel treatments for this devastating disease have been developed and are now in use. Intravenous epoprostenol has been shown in several studies to improve haemodynamics, exercise tolerance and survival [4–6]. To circumvent the risks, inconveniences and costs associated with continuous intravenous epoprostenol, stable prostacyclin analogues have been introduced that can be administered orally, subcutaneously or by inhalation [7]. In patients with IPAH, all these substances have been shown to improve exercise tolerance and haemodynamics in relatively short-term clinical studies [8–10], but the effects on long-term survival have not been thoroughly investigated.

Besides prostaglandins, novel substances with different mechanisms of action have been introduced in recent years. The dual endothelin receptor antagonist bosentan has been approved throughout the western world for treatment of pulmonary arterial hypertension (PAH) [11]. In addition, several case series and one small controlled study have suggested that phosphodiesterase-5-inhibitors, such as sildenafil, may have beneficial effects in PAH [12, 13].

Despite the fact that all of these substances have positive effects in IPAH, they do not provide a cure and, in many

patients, the disease will eventually progress. The optimal management for patients who exhibit clinical deterioration despite targeted treatment is unclear. The introduction of intravenous epoprostenol is an option for patients in whom treatment with endothelin receptor antagonists, nonintravenous prostanoids or sildenafil fails. Another option is combination treatment [14]. A few uncontrolled clinical studies have suggested that combinations of aerosolised iloprost and sildenafil, as well as aerosolised iloprost and bosentan, are safe and effective in selected patients with progressive disease despite prostanoid treatment [15, 16]. Combining bosentan and sildenafil may also be an attractive choice, since both substances are orally available, act *via* different mechanisms and are usually well tolerated. This paper describes the clinical course of nine patients with severe and progressive IPAH who were treated with a combination of bosentan and sildenafil.

Patients and methods

Between January 2002 and December 2003, 58 patients received bosentan for treatment of IPAH at Hannover Medical School (Hannover, Germany). Bosentan was used either as first-line therapy or in addition to a prostanoid (*i.e.* inhaled iloprost, oral beraprost or intravenous iloprost). Treatment efficacy was monitored primarily by the 6-min walk distance (6MWD) and cardiopulmonary exercise testing (CPET), and was targeted to reach pre-defined goals

based on prognostically relevant variables: a 6MWD >380 m [6] and a peak oxygen uptake measured during CPET >10.4 mL·min⁻¹·kg⁻¹ [17]. Patients who did not reach these goals on two consecutive visits were offered additional treatment with sildenafil. There was no formal study protocol. All patients were informed about the fact that this approach was investigational, that long-term data on safety and efficacy of sildenafil in PAH were not available, that there were no experiences with the combination of bosentan and sildenafil, and about the alternative option to receive intravenous prostacyclin treatment.

Treatments and follow-up

All patients had an extensive diagnostic work-up to define the aetiology of pulmonary hypertension, including, but not limited to, ventilation-perfusion scanning, computed tomography of the chest, right heart catheterisation and pulmonary angiography, when indicated. Bosentan was considered first-line treatment for patients newly diagnosed with IPAH. After initiation of bosentan treatment, the patients were seen in the outpatient clinic in 3–4 monthly intervals. Follow-up examinations included regular 6MWD [18] and CPET [16]. Repeated right heart catheterisations were not part of the routine follow-up programme.

Bosentan treatment was started at a dose of 2×62.5 mg and the dose was increased to 2×125 mg after 4 weeks in all patients. Liver enzymes were monitored every 4 weeks. The dose of bosentan was not changed throughout the observation period.

Sildenafil was started at a dose of 3×25 mg (or 4×25 mg) and was increased after 4–12 weeks to 3×50 mg when the response to the initial dose was not sufficient (*i.e.* when the treatment goals as defined above were not reached).

Analysis

All values are given as mean±SD. Two baselines were set: the first one before bosentan was started and the second one before sildenafil was started. The treatment effects after 3 months were compared with the baseline values using paired *t*-tests (two-sided). A *p*-value <0.05 was considered statistically significant.

Results

Within the observation period, 2002–2003, nine out of 58 patients with IPAH, initially treated with bosentan or a combination of bosentan and a prostanoid, fulfilled the predefined criteria of clinical worsening as defined above. These patients suffered from severe and progressive pulmonary hypertension, and were in New York Heart Association (NYHA) functional class III or IV before treatment was started. All patients were nonresponders to acute vasodilator challenge. The baseline characteristics and haemodynamics of these patients are shown in table 1.

Functional classification

At baseline, eight patients were functional class (FC) III and one patient was FC IV. Three months after bosentan was started, two patients had improved to FC II and the patient who formerly was FC IV had improved to FC III. However, after a follow-up of 11±5 months, deterioration occurred in all patients and, before sildenafil was added to bosentan, seven patients were FC III and two patients were FC IV. After 3 months of treatment with bosentan and sildenafil, six patients were FC III and three patients were FC II. During the follow-up period of 6–12 months, two further patients improved from FC III to FC II (table 2).

Table 1. – Patients' baseline characteristics and haemodynamics

Age yrs	39±9
Sex female/male	7/2
NYHA class III/IV	8/1
6MWD m	337±73
Mean right atrial pressure mmHg	9±5
Mean pulmonary arterial pressure mmHg	62±12
Cardiac output L·min ⁻¹	3.1±0.7
Cardiac index L·min ⁻¹ ·m ⁻²	1.6±0.3
Pulmonary vascular resistance dynes·s·cm ⁻⁵	1549±440
Mixed venous oxygen saturation %	53±10

NYHA: New York Heart Association; 6MWD: 6-min walk distance. 1 mmHg=0.133 kPa.

Table 2. – Individual functional classes (New York Heart Association (NYHA)) and 6-min walk distance (6-MWD)

Patient no.	Baseline 1 before bosentan		3 months after introduction of bosentan		Baseline 2 before sildenafil		3 months after addition of sildenafil		6–12 months after addition of sildenafil	
	NYHA	6-MWD	NYHA	6-MWD	NYHA	6-MWD	NYHA	6-MWD	NYHA	6-MWD
1	III	393	II	478	III	297	III	363	III	373
2 [#]	III	336	III	428	III	226	III	354	II	410
3	III	367	III	406	III	354	II	450	II	462
4	III	337	III	403	IV	132	II	414	II	382
5	III	420	II	497	III	371	III	421	II	481
6	III	328	III	406	III	353	III	425	III	425
7 [‡]	III	330	III	343	III	303	III	335	III	345
8	IV	198	III	232	III	252	III	290	III	328
9	III	406	III	434	IV	203	II	480	II	450

[#]: receiving inhaled iloprost since 15 months before bosentan was started and iloprost was continued throughout the observation period; [‡]: bosentan replaced continuous intravenous iloprost that had been given for 2 yrs but had to be withdrawn after several episodes of life-threatening line sepsis, sildenafil was added 11 months after introduction of bosentan.

Six-min walk distance

As shown in figure 1 and table 2, the 6MWD at baseline was 346 ± 66 m. After introduction of bosentan treatment, the 6MWD improved by 57 m to 403 ± 80 m ($p=0.0003$). However, this effect was not sustained and, after an interval of 11 ± 5 months, the walk distance had declined to 277 ± 80 m. At this point, sildenafil was added to bosentan. Three months later, the 6MWD had increased to 392 ± 61 m ($p=0.007$) and the patients remained stable throughout the median follow-up of 9 months (range 6–12).

Cardiopulmonary exercise testing

Complete sets of data from CPET at all time points (baseline 1, three months after introduction of bosentan, baseline 2 and 3 months after the addition of sildenafil) were available from six patients. In these patients, the peak oxygen uptake ($\dot{V}O_{2,\text{peak}}$) was 9.9 ± 1.9 mL $\cdot$ min $^{-1}\cdot$ kg $^{-1}$ at baseline and increased to 12.3 ± 1.9 mL $\cdot$ min $^{-1}\cdot$ kg $^{-1}$ 3 months after introduction of bosentan ($p=0.02$). As with the walking distance, this effect was not sustained and, before introduction of sildenafil, $\dot{V}O_{2,\text{peak}}$ had decreased to 10.4 ± 2.3 mL $\cdot$ min $^{-1}\cdot$ kg $^{-1}$. Three months after sildenafil was added to bosentan, $\dot{V}O_{2,\text{peak}}$ had increased to 13.8 ± 1.5 mL $\cdot$ min $^{-1}\cdot$ kg $^{-1}$ ($p=0.006$).

Side-effects

None of the patients in this case series died or experienced any drug-related serious adverse event. There were no abnormalities of liver enzymes before or after the introduction of bosentan, or after addition of sildenafil. All patients reported minor headache and flushing when sildenafil was added to bosentan, but these problems resolved within a few days without dose adjustments. The same was true with heartburn, reported by one patient after sildenafil was started. No patient described symptoms attributable to hypotension and there were no episodes of syncope during the observation period.




Fig. 1.—Six-min walk distance (6MWD) of nine patients with idiopathic pulmonary arterial hypertension at baseline 1 (before treatment with bosentan), 3 months after introduction of bosentan, baseline 2 (11 ± 5 months later before addition of sildenafil), 3 months later and 6–12 months later. Open arrow: bosentan started; closed arrow sildenafil started.

Discussion

This case series may serve as a pilot trial showing that combining bosentan and sildenafil might be feasible in patients with IPAH. This combination was well tolerated by all patients and proved to be highly efficient. Of note, this patient population consisted of a "negatively" selected subgroup of young patients with very severe and rapidly progressive IPAH. Before targeted treatment was initiated, all patients were in functional class NYHA III or IV, and haemodynamic assessment confirmed the presence of advanced pulmonary hypertension with a mean pulmonary artery pressure of 62 ± 12 mmHg, a cardiac index of 1.6 ± 0.3 L $\cdot$ min $^{-1}\cdot$ m $^{-2}$ and a pulmonary vascular resistance of $1,549 \pm 440$ dynes $\cdot$ s $\cdot$ cm $^{-5}$. In accordance with current recommendations [7, 14], bosentan was started as first-line treatment, resulting in clinical improvement and increased exercise tolerance as determined by 6MWD and CPET. However, in contrast to the majority of patients receiving treatment with bosentan, this effect was not sustained. After an average period of close to 1 yr, the patients showed clinical deterioration and a marked decline in 6MWD, as well as in $\dot{V}O_{2,\text{peak}}$. It is the unfortunate clinical reality that, even with the novel targeted treatments available for PAH, a certain proportion of patients are going to deteriorate after initial improvement. Those patients are very likely to die from their underlying disease and they do require aggressive treatment. The most recent world conference on pulmonary hypertension recommended that prostacyclin treatment or, alternatively, combination treatment should be initiated in patients not responding sufficiently to treatment with endothelin receptor antagonists or nonintravenous prostanoids [14]. Since bosentan clearly had a beneficial effect in all of the patients in the current study, the authors decided to initiate combination treatment, reserving intravenous prostacyclin for treatment failures. For reasons discussed below, sildenafil was chosen as an add-on treatment. The combination of bosentan and sildenafil was surprisingly well tolerated by all patients and resulted in marked clinical improvement. The pre-defined treatment goals (6MWD >380 m and $\dot{V}O_{2,\text{peak}} >10.4$ mL $\cdot$ min $^{-1}\cdot$ kg $^{-1}$) were reached in six out of nine patients. The remaining patients were judged as sufficiently stable and showed a tendency towards further improvement during follow-up, so that introduction of intravenous prostacyclin was not deemed immediately necessary. It is noteworthy that none of the patients died during the observation period of 21 ± 3 months, even though the expected 1-yr and 2-yr survival probabilities of these patients based on their baseline haemodynamics according to the formula of the National Institutes of Health were 0.64 and 0.51, respectively [3].

Obviously, this observational report has several inherent limitations: there was no control group, the patient population was small and the overall observation time was rather limited. In addition, treatment with bosentan and sildenafil was not part of a formal study protocol, and serial right heart catheter examinations were not performed and, therefore, haemodynamic follow-up data were not available. Although these data would be of value, it is the authors' practice to base most therapeutic decisions on the results of patient's self reporting (functional classification), clinical examination, 6MWT and CPET, and to reserve right heart catheterisation for open questions or for certain therapeutic decisions, such as listing for highly urgent lung transplantation.

Bosentan and sildenafil were combined rather than bosentan and a prostanoid for several reasons: at the time when combination treatment was initiated in the patients, bosentan was the only drug approved for treatment of IPAH

in Germany (sildenafil, beraprost and treprostinil have not been approved in Germany, and aerosolised iloprost was officially introduced in January 2004); beraprost or treprostinil were not considered as first choices for combination treatment because the former drug has not been shown to exhibit long-term clinical efficacy [19] and the latter one commonly causes substantial infusion site pain [10]; compared to inhaled iloprost, sildenafil has the practical advantage of being orally administered rather than requiring six to nine inhalations daily; and in addition, although it has been shown that inhaled iloprost and sildenafil have synergistic effects [15, 20, 21], it was hypothesised that the combination of substances with different intracellular modes of action might be advantageous in the long run, although comparative studies are required to resolve this issue.

Economical considerations also contributed to the decision to combine bosentan and sildenafil. In Germany, the annual treatment costs for a single patient are ~€45,000 for bosentan, €70,000 for inhaled iloprost and €5,000–10,000 (depending on the dose) for sildenafil. Thus, the annual costs for combining bosentan and sildenafil are €50,000–55,000, compared to €75,000–80,000 for inhaled iloprost plus sildenafil and €115,000 for inhaled iloprost plus bosentan. In comparison, in Germany, the annual treatment costs for intravenous epoprostenol are €230,000 (at a dose of 25 ng·kg⁻¹·min⁻¹, which is a typical average dose in patients receiving epoprostenol for >1 yr [5, 6]) and those for intravenous iloprost are €180,000 (at a dose of 2 ng·kg⁻¹·min⁻¹, a typical average dose for intravenous iloprost [16]). Thus, when the clinical situation mandates combination treatment or the introduction of intravenous prostacyclin, the annual savings with the combination bosentan and sildenafil are >€100,000 for every patient, meaning that the approach in this study resulted in savings of >€1,000,000 just based on the nine patients described.

Despite these considerations, many open questions remain: it is unknown whether combination treatment is truly more efficient than single treatment (although theoretical reasoning clearly favours combination treatment, switching to sildenafil may have been equally effective as combining bosentan and sildenafil); it is unknown which combination regimens exert favourable long-term results (this will be difficult to answer as the number of potentially useful combinations is exponentially increasing with the introduction of novel treatments for PAH); and finally, so far, the combination of bosentan and sildenafil has only been used in patients with IPAHA, and it is unknown whether these observations also apply to other forms of pulmonary hypertension.

In conclusion, the data presented here provide preliminary evidence that the combination of bosentan and sildenafil may be safe and effective in selected patients with idiopathic pulmonary arterial hypertension.

References

- Rubin LJ. Primary pulmonary hypertension. *N Engl J Med* 1997; 336: 111–117.
- Gaine S. Pulmonary hypertension. *JAMA* 2000; 284: 3160–3168.
- D'Alonzo GE, Barst RJ, Ayres SM, *et al.* Survival in patients with primary pulmonary hypertension. Results from a national prospective registry. *Ann Intern Med* 1991; 115: 343–349.
- Barst RJ, Rubin LJ, Long WA, *et al.* A comparison of continuous intravenous epoprostenol (prostacyclin) with conventional therapy for primary pulmonary hypertension. The Primary Pulmonary Hypertension Study Group. *N Engl J Med* 1996; 334: 296–302.
- McLaughlin VV, Shillington A, Rich S. Survival in primary pulmonary hypertension: the impact of epoprostenol therapy. *Circulation* 2002; 106: 1477–1482.
- Sitbon O, Humbert M, Nunes H, *et al.* Long-term intravenous epoprostenol infusion in primary pulmonary hypertension. Prognostic factors and survival. *J Am Coll Cardiol* 2002; 40: 780–788.
- Hoepfer MM, Galie N, Simonneau G, Rubin LJ. New treatments for pulmonary arterial hypertension. *Am J Respir Crit Care Med* 2002; 165: 1209–1216.
- Galie N, Humbert M, Vachiery JL, *et al.* Effects of beraprost sodium, an oral prostacyclin analogue, in patients with pulmonary arterial hypertension: a randomized, double-blind, placebo-controlled trial. *J Am Coll Cardiol* 2002; 39: 1496–1502.
- Olschewski H, Simonneau G, Galie N, *et al.* Inhaled iloprost for severe pulmonary hypertension. *N Engl J Med* 2002; 347: 322–329.
- Simonneau G, Barst RJ, Galie N, *et al.* Continuous subcutaneous infusion of treprostinil, a prostacyclin analogue, in patients with pulmonary arterial hypertension. A double-blind, randomized, placebo-controlled trial. *Am J Respir Crit Care Med* 2002; 165: 800–804.
- Rubin LJ, Badesch DB, Barst RJ, *et al.* Bosentan therapy for pulmonary arterial hypertension. *N Engl J Med* 2002; 346: 896–903.
- Michelakis ED, Tymchak W, Noga M, *et al.* Long-term treatment with oral sildenafil is safe and improves functional capacity and hemodynamics in patients with pulmonary arterial hypertension. *Circulation* 2003; 20: 20.
- Sastry BK, Narasimhan C, Reddy NK, Raju BS. Clinical efficacy of sildenafil in primary pulmonary hypertension: a randomized, placebo-controlled, double-blind, crossover study. *J Am Coll Cardiol* 2004; 43: 1149–1153.
- Galie N, Seeger W, Naeije R, Simonneau G, Rubin LJ. Comparative analysis of clinical trials and evidence-based treatment algorithm in pulmonary arterial hypertension. *J Am Coll Cardiol* 2004; 43: S81–S88.
- Ghofrani HA, Rose F, Schermuly RT, *et al.* Oral sildenafil as long-term adjunct therapy to inhaled iloprost in severe pulmonary arterial hypertension. *J Am Coll Cardiol* 2003; 42: 158–164.
- Hoepfer MM, Taha N, Bekjarova A, Gatzke R, Spiekerkoetter E. Bosentan treatment in patients with primary pulmonary hypertension receiving nonparenteral prostanoids. *Eur Respir J* 2003; 22: 330–334.
- Wensel R, Opitz C, Anker SD, *et al.* Assessment of survival in patients with primary pulmonary hypertension: importance of cardiopulmonary exercise testing. *Circulation* 2002; 106: 319–324.
- ATS statement: guidelines for the six-minute walk test. *Am J Respir Crit Care Med* 2002; 166: 111–117.
- Barst RJ, McGoon M, McLaughlin V, *et al.* Beraprost therapy for pulmonary arterial hypertension. *J Am Coll Cardiol* 2003; 41: 2119–2125.
- Wilkins H, Guth A, Koenig J, *et al.* Effect of inhaled iloprost plus oral sildenafil in patients with primary pulmonary hypertension. *Circulation* 2001; 104: 1218–1222.
- Ghofrani HA, Wiedemann R, Rose F, *et al.* Combination therapy with oral sildenafil and inhaled iloprost for severe pulmonary hypertension. *Ann Intern Med* 2002; 136: 515–522.