

Converting Peptides into Drug Leads by Lipidation

L. Zhang* and G. Bulaj

Department of Medicinal Chemistry, University of Utah, Salt Lake City, Utah, USA

Abstract: Lipidation is a posttranslational modification of proteins that has also found its use in designing peptide drugs. The presence of a lipid group in peptides modulates their hydrophobicity, secondary structures and self-assembling propensities while retaining their abilities to bind to target receptors. Lipidation improves peptides' metabolic stability, membrane permeability, bioavailability, and changes peptides' pharmacokinetic and pharmacodynamic properties. Herein, we review the applications of various lipidation strategies in peptide drug design, the effects of the chain length and anchor position of fatty acids in peptide lipidation, the physicochemical and biological properties of selected lipidated peptides and the synthesis strategies for peptide lipidation.

Keywords: ADME, lipidation, lipidated peptide, lipoamino acids, peptide, peptide SAR, peptide secondary structure, peptide self-assembling, peptide serum stability, peptide synthesis.

INTRODUCTION

Lipidation is a posttranslational modification of peptides and proteins. Some nonribosomally synthesized bacterial antibiotics [1] and ghrelin (octanoyl-Ser³, Fig. 1a) [2, 3] are naturally-occurring lipidated peptides. Protein lipidations, such as prenylation, S-acylation, *N*-myristoylation, S- and *N*-palmitoylation play important roles as posttranslational modifications of proteins to maintain and modulate their functions [4, 5]. For example, lipidation is necessary for RAS superfamily proteins to maintain their cellular functions [6]. Thus, due to the significant impacts on the properties of peptides following lipidation, this chemical modification has proven to be a useful tool in peptide drug design.

Peptide drugs offer many desirable drug properties such as potent bioactivity, high receptor binding affinity and specificity, and relatively low toxicity. However, most peptides are quickly degraded in serum and exhibit rapid clearance *in vivo*. Many strategies have been employed in an attempt to overcome these apparent disadvantages including lipidation, glycosylation, cyclization, backbone modification, unnatural or D-amino acid conjugation and PEGylation. Among these chemical modification methods, lipidation has proven to be one of the most robust strategies for the generation of new therapeutic peptide leads as discussed in this review paper. Lipidation can dramatically change peptides' physicochemical properties and pharmacological properties [7]. Additionally, lipidation increases peptides' plasma stability and decreases peptides' kidney clearance.

Four lipidized peptide drugs are currently approved for use in humans: two long-acting anti-diabetics human glucagon-like peptide-1 (GLP-1) analogue liraglutide (Victoza®, NN2211, Fig. 1b) and insulin detemir (Levemir®, (N^εB²⁹-tetradecanoyl)-des-(B30)-insulin, Fig. 1c), and two antibiotics daptomycin and polymyxin B. A number of lipidated peptides, lipoproteins and lipidated peptide vaccines such as Lymerix™ (against Lyme disease) are at different phases of clinical trials. As indicated in Table 1, lipidation was extensively exploited in the drug developments of angiotensin II, desmopressin, enkephalin, galanin, neuropeptide Y (NPY), octreotide, thyrotropin releasing hormone (TRH), luteinizing hormone releasing hormone (LHRH), calcitonin, gastric inhibitory polypeptide (GIP), ghrelin, GLP-1, growth hormone-releasing hormone (GHRH), tetragastrin and insulin. In addition, lipidized peptides have also been used as protein function probes, gene delivery vectors and vaccine adjuvants [8, 9]; palmitoyl pentapeptide analogues were used in cosmetics [10].

Lipidation can be classified into three types based on lipid bond formation methods with peptide chain or side chains: amidation,

esterification (S- or O-) and S-bond (ether or disulfide) formation. Amidation and O-esterification are strong and irreversible covalent bonds, whereas S-esterification and disulfide bond formation are weak and reversible covalent bonds. Non-natural lipoamino acids (LAA) (Fig. 2a) were also extensively exploited in the development of peptide drugs and vaccines by the Toth group [59-62, 69-73]. Lipidation methods, alkyl chains, lipid anchor position and the spacer between lipids and peptides have significant impacts on the physicochemical properties, receptor binding affinities and bioactivities of lipidated peptides.

Several related review papers on protein lipidation [74-76], vaccine lipidation [77, 78], lipid core peptide delivery system [79], lipid-based delivery system [80, 81], anti-infective lipidated peptide drugs [82, 83], amphiphilic lipidated peptide self-assembling system [84], and lipopeptide molecular probes [85-87] were published. Herein, we review the peptide lipidation methods and discuss the structure-activity relationship (SAR) of peptide lipidation with particular focus on peptide fatty acid lipidation.

1. PHYSICOCHEMICAL AND PHARMACOLOGICAL PROPERTIES OF LIPIDATED PEPTIDES

1.1. Lipophilicity

Most naturally-occurring peptides are hydrophilic peptides. Lipidation provides a simple way to modulate peptide lipophilicity. Since lipophilicity affects the absorption, distribution, metabolism and excretion (ADME) and bioavailability of drugs, lipidation is an attractive strategy to convert peptides into drug leads. Partition between the aqueous phase and the membrane phase of drugs can be simulated through partition of drugs between two immiscible solvents [88]. The *n*-octanol-water partition coefficient (logD) is commonly used to evaluate the permeability of drugs across biological membranes. Many research groups have evaluated the hydrophobicity of lipidated peptides by determining their logD [8, 12, 14, 15, 18, 31, 55, 89, 90]. Lipidation of TRH with lauroyl (*N*-lauroyl-TRH, Fig. 2b) increased the logD of TRH from 0.068 to 1.91 [38]. Lipidation of NPY-BBB1(K3) with palmitoyl (NPY-BBB1, Table 2) changed the logD of NPY-BBB1(K3) from -1.38 to 2.82 [15]. All the palmitoyl lipidated galanin and neuropeptide W (NPW) analogues showed similar logD values, ~1.2 and ~1.6, respectively (Table 2) [14, 31]. The logD was not affected by the anchor positions of the palmitoyl group on the KKKK-palmitoyl scaffold. More specifically, the logD was linearly related to the chain length of the fatty acids of the galanin lipidated Gal-B2 analogues from Gal-B2-C₈ to Gal-B2-C₁₈ (Fig. 2c; C_n, n is the carbon number of linear fatty acids) [18].

1.2. Secondary Structures

Lipidation can change the secondary structures of peptides through the hydrophobic interactions of the fatty acid chain with the

*Address correspondence to this author at the 421 Wakara Way, Suite 360, Salt Lake City, Utah 84108, USA; Tel: (801)581-4371; Fax: 801-585-6354; E-mail: liuyin.zhang@pharm.utah.edu




Fig. (1). Structures of ghrelin, GLP-1 (liraglutide) and insulin detemir.

Table 1. Lipidation Methods Applied in Peptide Modifications

Type			Peptides investigated
Fatty acid	linear	Amide	Enkephalin [11-13], Galanin [14-18], GIP [7, 19], GLP-1 (liraglutide) [20], hPP [21], IGF1R JM inhibitor [22], Insulin [23-29]; Leuteinizing hormone releasing hormone (LHRH) [30], NPW [31], NPY [15], NT [15], RC-160 [32, 33], STAT3 [34], Tetragastrin [35-37], Thyrotropin releasing hormone (TRH) [28, 30, 38, 39]
		Ester	Angiotensin II [40, 41], BBI [42-44], Desmopressin [45], Ghrelin [2, 46-48], Salmon calcitonin (sCT) [49-51], TRH [52], Octreotide [53]
	branched		Liraglutide [20]
Amino fatty acid			Conotoxin SIIIA [54], Ghrelin [46], Liraglutide [20], Met-Enkephalin [55], TAN-1511A [56], Tenecin-1 [57]
Acidic fatty acid (dicarboxylic acid)			Growth hormone-releasing hormone antagonists (GHRH) [58]
Halo-fatty acid			Ghrelin [46]
Lipoamino acid			Core peptide [59, 60], Thymopentin TP5 [61], Peptide A2 [62], Polycationic lipophilic core dendron excipient [63, 64]
Alkyl ether (O-, S-)			Angiotensin [41]
Alkyl amine			Compound 3 [65]
Adamantylation			Ghrelin [46]
Cholesterol			C34 [66], Insulin [67,68]




Fig. (2). Structures of lipoamino acids (LAA), N-lauroyl-TRH and Gal-B2 analogues.

peptide backbone or side chains [91, 92]. The lipidation analogues of random-coiled salmon calcitonin (sCT), REAL-sCT (Fig. 3a) and Mal-sCT (Fig. 3b) adopted α -helix secondary structures [49]. The octanoylation analogues on Lys¹⁶ or Lys²⁸ of random-coiled human β -amyloid (1-40) (Fig. 3c) adopted β -sheet structures [93]. Palmitoyl-kallidin (Fig. 3d) adopted a non-ideal type I β -turn in the presence of dodecylphosphocholine (DPC) micelles while palmitoyl-Gly-kallidin adopted a γ -turn structure [94].

Long-chain fatty acid ($>\text{C}_8$) lipidations have strong effects on the secondary structures of peptides when compared to short-chain fatty acid lipidations, likely due to their strong hydrophobic interactions with peptides. For example, a random-coiled peptide **16r** acquired an α -helix structure after N-terminal lipidation with

fatty acids (Fig. 3e); the α -helical content of C_{16} -**16r** was higher than that of C_6 -**16r** in water or 50% trifluoroethanol (TFE) [95]. The melting temperatures (T_m) of lipidated collagen derived peptides (Fig. 3e) from triple helix state to random coil state were also correlated with the chain length of fatty acids [95]. The lipidated analogues of **16r**-Tyr-SPARC₁₁₉₋₁₂₂ (Fig. 3e) showed similar increasing trends of α -helical contents and T_m with the elongation of fatty acid chain [96]. A de novo α -helix peptide **1a** (Fig. 3f) acquired β -sheet conformations after N-terminal lipidation with long-chain fatty acids (from C_{12} to C_{16}), while the lipidated analogues of **1a** with short-chain fatty acids (from C_2 to C_{10}) resulted in random-coiled structures [97]. The analogues of a de novo α -helix dimer **2a** (Fig. 3f) lipidated with long-chain fatty acids (from C_{12} to C_{16}) at its two N-termini maintained α -helix

Table 2. The SAR Study Analogues of Neuropeptide Galanin, NPW, NT and NPY (Lys-Pal, *N*^c-Palmitoyl-Lysine; Ava, 5-Amino-Valeric Acid)

Analogue	Sequence
Gal-LAA [14]	(Sar)WTLNSAGYLLGPHA(Lys-Pal)-NH ₂
Gal-(K)4	(Sar)WTLNSAGYLLGPKKKK-NH ₂
Gal-B1	(Sar)WTLNSAGYLLGPKKK(Lys-Pal)-NH ₂
Gal-B2	(Sar)WTLNSAGYLLGPKK(Lys-Pal)K-NH ₂
Gal-B3	(Sar)WTLNSAGYLLGPK(Lys-Pal)KK-NH ₂
Gal-B4	(Sar)WTLNSAGYLLGP(Lys-Pal)KKK-NH ₂
Gal-B11	(Sar)WTLNSAGYLLGKK(Lys-Pal)K-NH ₂
Gal-B12	(Sar)WTLNSAGYLLKK(Lys-Pal)K-NH ₂
NPW-B2 [31]	Ac-WYK(Lys-Pal)KKK(Ava)GRAAGLLMGL-NH ₂
NPW-B3	Ac-WYKK(Lys-Pal)KK(Ava)GRAAGLLMGL-NH ₂
NPW-B4	Ac-WYKKK(Lys-Pal)K(Ava)GRAAGLLMGL-NH ₂
NPW-B5	Ac-WYKKKK(Lys-Pal)(Ava)GRAAGLLMGL-NH ₂
NT-BBB1(K4) [15]	Ac-KKKKPYIL-OH
NT-BBB1	Ac-K(Lys-Pal)KKPYIL-OH
NPY-BBB1(K3) [15]	Ac-YKKK(Ahx)RHYINLITRQRY-NH ₂
NPY-BBB1	Ac-YKK(Lys-Pal)(Ahx)RHYINLITRQRY-NH ₂
NPY-BBB2(K3)	Ac-YKKK(Ahx)ARHYINLITRQRY-NH ₂
NPY-BBB2	Ac-YKK(Lys-Pal)(Ahx)ARHYINLITRQRY-NH ₂

structures while the analogues of **2α** lipidated with short-chain fatty acids (from C₄ to C₁₀) induced β -sheet structures [97].

Lipid position was found to be important in stabilizing the conformations of lipidated peptides. The *N*-terminal lipidation analogue of α -conotoxin MII, LAA₁₀-MII (Fig. 3g; LAA₁₀, 2-amino-D,L-dodecanoic acid), exhibited consistent helical tertiary structure [69]. However, the analogue [Asn⁵]LAA₁₀-MII (Fig. 3g) lipidated in the middle of peptide chain did not adopt the bioactive conformation [69]. Lipidation of β -amyloid near one β -sheet region (Lys¹⁶) or the bend region of the two β -sheets (Lys²⁸) yielded two analogues (C₈-Lys¹⁶)- β -amyloid(1-40) and (C₈-Lys²⁸)- β -amyloid(1-40) (Fig. 3c) with enhanced secondary conformations as compared to that of the native β -amyloid(1-40) [93]. Lipidation at the unstructured *N*-terminal region of β -amyloid, such as *N*-C₈- β -amyloid(1-40), did not affect the secondary conformations [93].

Unsaturated fatty acids and hydroxyl fatty acids exhibit less strong hydrophobic interactions with peptide chains when compared to the corresponding saturated fatty acids, resulting in less influences on the secondary structures of lipidated peptides. Oleoyl-**16r**-Tyr-SPARC₁₁₉₋₁₂₂ (Fig. 3e) showed a lower helical content and *T_m* than that of C₁₈-**16r**-Tyr-SPARC₁₁₉₋₁₂₂ [96]. The helical content of ricinoleoyl-**16r**-Tyr-SPARC₁₁₉₋₁₂₂ is higher than that of oleoyl-**16r**-Tyr-SPARC₁₁₉₋₁₂₂ due to the compensation of the hydrogen bond of the hydroxyl group but is not strong than that of C₁₈-**16r**-Tyr-SPARC₁₁₉₋₁₂₂ [96].

The interactions of lipidated peptides with membranes significantly influence peptide secondary structures. SMOi2-9 (Ac-LTYAWHTSFKAL-NH₂) showed random conformations in aqueous solutions; lipidated peptide SMOi2-43 (Fig. 3h) adopted a β -type conformation in aqueous solution and an α -helix conformation in DPC micelles [22]. *N*-Octanoyl-mastoparan-X (Fig. 3i) and *N*-propanoyl-mastoparan-X adopted β -sheet conformations in HEPES solution. However, both analogues adopted α -helix conformations in 1-palmitoyl-2-oleoyl-phosphatidylcholine (POPC) or 1-palmitoyl-2-oleoyl-phosphatidylglycerol (POPG) solutions with 50-60% α -helical contents [98]. The lipidated analogue of human lactoferrin C₁₂-LF11 (Fig. 3j) adopted an α -helix conformation in DPC micelles but formed a loop in sodium dodecyl

sulphate (SDS) micelles [99]. Lipidated peptides with specific conformation properties also appeared as useful molecular probes to study protein folding [100].

1.3. Oligomers and Self-Assembling Macromolecules

Lipidated peptides can form stable oligomers, self-assembling structures (such as micelles, tubules, vesicles, mono- or bilayers, nanofibers) and aggregates. *N*-Terminal lipidated analogues of dimer **2α** (from C₁₀- to C₁₆-, Fig. 3f) formed stable oligomers [97]. Compound 3 (Fig. 4a) formed hollow tubules or helical ribbons spontaneously; these self-assembled structures protected compound 3 from trypsin degradation [65]. Several reviews describing self-assembly mechanisms, compositions and applications of lipidated peptides were available [84, 101-104]. Herein, we briefly describe the effects of lipid anchor position and the chain length of fatty acids on the self-assembly and aggregation properties of lipidated peptide.

In lipidated peptides, long-chain fatty acid conjugations provide stronger hydrophobic driving force for aggregation formation as compared to short-chain fatty acid conjugations. *N*-Terminal lipidation of PA4 with hexanoic acid (C₆-) (Fig. 4b) formed random size and shape precipitates at the concentrations as low as 1%, while *N*-terminal lipidation of PA4 with long-chain fatty acids (C₁₀-, C₁₆- or C₂₂-) formed different morphological nanofibers at the concentration of 0.25% [105]. In some cases long-chain fatty acids facilitated the formation of large particles regardless of the peptide sequences [95]. Lipidation strategies also have impacts on the particle formation of lipidated peptides. REAL-sCT (Fig. 3a) and Mal-sCT (Fig. 3b) formed particles with 30- to 40-fold differences in mean particle radii, 3.1 - 3.6 nm (11.2 - 1000 μ M) and 105 - 134.3 nm (2.5 - 224 μ M), respectively [49]. Lipidated analogues of the GBV-C/HGV NS3 protein fragments 505-514 (SAELSMQRRG) and 513-522 (RGRTGRGRSG) demonstrated different aggregation properties [106]. Palmitoyl-SAELSMQRRG formed fibrous aggregates in aqueous solution, whereas palmitoyl-RGRTGRGRSG failed to form any fibrous aggregates under the same conditions [106].




Fig. (3). Structures of the lipitated analogues of sCT (Real-sCT and Mal-sCT), β -amyloid, kallidin, 16r, collagen, 1a/2a, LAA-MII, SMOi2-43, mastoparan-X and C₁₂-LF11.

Lipid anchor positions have certain impacts on the critical micelle concentration (CMC) of lipitated peptides. For example, the CMCs of (*N*^ε-C₈-Lys¹⁶)- β -amyloid-(1-40) and (*N*^ε-C₈-Lys²⁸)- β -amyloid-(1-40) (Fig. 3c) were 10 μ M and 8 μ M, respectively; whereas the CMC of *N*-C₈- β -amyloid-(1-40) was equal to the CMC of the native β -amyloid (1-40), ~50 μ M [93].

1.4. Membrane Binding

Lipidation affects peptide-membrane interactions [88]. The interactions may have tremendous effects on the physical properties of membranes, such as the fluidity, free volume, intrinsic curvature and morphologies of the membrane [106]. Lipitated analogues of

the pentapeptide YIGSR-NH₂ (Fig. 5a) showed weak disturbances in dipalmitoylphosphatidylcholine (DPPC) liposome bilayers due to the similarity of peptide lipid chains with DPPC lipid chains [107]. The interactions also affect the membrane trafficking, localization and signal transduction of lipitated peptides, as indicated in Ras proteins [108, 109].

The chain lengths of fatty acids have important impacts on peptide-membrane interactions. The fatty acid moieties of the lipitated analogues of tripeptide GHK (with C₆, C₁₀ or C₁₄, Fig. 5b) were inserted into the external bilayer of DPPC liposomes while the peptide chain protruded from the liposome surface [110]. The membrane binding energies of *N*-terminal acylated glycine analogues (*N*-acyl-Gly) were increased with the elongation of the




Fig. (4). Structure of compound 3 and PA4 analogues.

fatty acid chains [111]. The effective charges and the distance between membranes and peptide chains of lipidated peptides were decreased due to the strong hydrophobicity of lipidated peptides [98]. Lipidated analogues of mastoparan-X (Fig. 3i) showed reduced free energy upon insertion in neutral POPC and negatively charged POPC/POPG (1:3) membranes as compared to that of the native mastoparan-X. Despite of the less hydrophobicity of propanoyl group compare to octanoyl group, *N*-propanoyl-mastoparan-X showed reduced free energy upon insertion of water into the membrane compared to *N*-octanoyl-mastoparan-X [98]. Lipidation may have negative effects on the membrane binding properties of lipidated peptides due to the increased aggregation propensity and the reduced effective monomer concentration [112].

1.5. Receptor Binding

The anchor positions and the chain lengths of fatty acids have great influence on the receptor binding ability of lipidated peptides. Lipidation on a noncritical terminus is an important strategy to maintain peptides' receptor binding ability. The C-terminus of endomorphin-1 is important to maintain its bioactivity; endomorphin-1-LAA_n (n = 8, 12; Fig. 6a) showed decreased binding affinities to opioid receptors due to the blocking of the C-terminus whereas the affinities of LAA₈-endomorphin-1 were not affected [113]. DiC₁₀-Leu-enkephalin (Fig. 6b) exhibited a remarkable decrease of the opioid receptor binding affinity due to the lipidation on the N-terminus which is critical for maintaining its bioactivity [13]. The strategy of lipidation on a noncritical terminus was applied in the modifications of galanin, NPY and neurotensin (NT) and yielded novel anticonvulsant neuropeptides without losing their receptor binding affinities [14, 15]. The lipidated galanin analogues Gal-B1 to Gal-B4 (Table 2) showed a decreasing binding trend to the human GalR1 receptor (hGalR1) and GalR2 receptor (hGalR2) with the increasing distance between the palmitoyl moiety and the pharmacophore of galanin (pharmacophore: Sar-WTLNSAGYLLGP) [14].

Long-chain fatty acid lipidation can cause considerable decreases in peptide receptor binding affinity as compared to short-chain fatty acid lipidation. The hGalR1 receptor binding affinity of lipidated Gal-(K)4 analogues (from C₈ to C₁₈, Fig. 2c) demonstrated a decreasing tendency with the elongation of the chain length of fatty acids; the hGalR1 receptor binding ability of C₁₈-Gal-B2 (*K*_i 4.0 ± 2 nM) was ten-fold less than that of Gal-(K)4 (*K*_i 0.4 ± 0.1 nM) [18]. Similarly, the somatostatin receptor affinities of C₄-RC-160 and C₁₈-RC-160 (Fig. 6c) were decreased 10- and 30-fold, respectively, as compared to that of the native RC-160 [33].

Lipidation may have different receptor binding effects toward different receptor subtypes. [K¹³(E-Pamitoyl), Q³⁴]hPP (Fig. 6d), a lipidated analogue of hormone pancreatic polypeptide (hPP), showed a similar binding ability toward human neuropeptide Y2 and Y5 receptors and a modest decrease of binding ability toward human neuropeptide Y4 receptor as compared to that of the parent peptide [K¹³, Q³⁴]hPP [21]. The binding affinities of Gal-B2 toward hGalR1 and hGalR2 receptors were decreased modestly as compared to Gal-(K)4, 10- and 2-fold, respectively [14].

Despite the common negative effects, some positive effects of lipidation on peptides' receptor binding affinity were also reported in the literature. LAA₁₀-MII (Fig. 3g) exhibited equal potency as the native MII in the inhibition of the neuronal nicotinic acetylcholine receptor subtype α3β2 [69]. The lipidated analogues of RC-160 (with C₈-, C₁₄-, C₁₆-, Fig. 6c) showed similar or modestly increased binding affinities toward somatostatin receptor as compared to that of the native RC-160 [32]. *N*-terminal lipidated GHRH antagonist analogues of JV-1-102 (from C₈- to C₁₆-, Fig. 6e) exhibited 30- to 100-fold increases of binding abilities to the membrane receptors on rat anterior pituitary cells [58].

2. ADME PROPERTIES OF LIPIDATED PEPTIDES

2.1. Absorption

Lipidation has strong and diverse effects on the membrane permeabilities of lipidated peptides. The three lipidated tetragastrin (TG) analogues, C₂-TG, C₆-TG and C₁₂-TG (Fig. 7a), have equal gastric acid output activities following *i.v.* administration [114]. However, the total gastric acid output amounts of C₂-TG and C₆-TG following large intestine loop administration were increased up to 133- and 74-fold, respectively, as compared to that of the native TG, whereas C₁₂-TG showed similar activity as the native TG [114]. The permeability across the upper small intestine of *N*-lauroyl-TRH (Fig. 2b) was increased, but the permeabilities across the other portions of small intestine and large intestine were not affected [39].

Lipidation improves the *in vitro* and *in vivo* uptakes of lipidated peptides [115]. The transportations of tyrosine kinase inhibitors ABL-myristoyl (Fig. 7b) and ABL-ss-myristoyl (Fig. 7c) across B lymphocyte cell line BA/F3 were increased 12- and 22-fold, respectively, as compared to that of the native ABL [116]. The Caco-2 cellular uptake of [¹²⁵I]DiC₁₀-Leu-enkephalin (Fig. 6b) was 131-fold higher than that of Leu-enkephalin [13]. C₂-TG and C₆-TG, exhibited increased penetration across the Caco-2 monolayers, the jejuna and colon tissues as compared to that of the native TG




Fig. (5). Structures of the lipidated analogues of YIGSR-NH₂ and GHK-OH.




Fig. (6). Structures of the lipidated analogues of endomorphin-1, DiC₁₀-Leu-enkephalin, RC-160, hPP, hGHRH antagonists.




Fig. (7). Structures of the lipidated analogues of tetragastrin, ABL-myristoyl, ABL-ss-myristoyl, insulin, DADLE, Phe-Gly, and DiC₁₆-desmopressin. (in figure 7b and 7c; 5-FAM, 5'-isomer of fluorescein).

[35-37]. (*N*^{B1}-palmitoyl)-insulin and (*N*^{B1}-palmitoyl)-(*N*^{B29}-palmitoyl)-insulin (Fig. 7d) showed remarkably increased large

intestinal absorptions and bioavailabilities when compared to that of the native insulin [25, 26, 28].




Fig. (8). Structures of the lipidated analogues of GIP, angiotensin II, REAL-octreotide and SDF1 α .

Lipidation with long-chain fatty acids may limit the membrane transportation of lipidated peptides in some scenarios. The C₁₂-TG cannot pass through the jejunal and colonic mucosas, while C₂-TG and C₆-TG showed increased penetration across the jejunal and colonic mucosas [35-37]. “Bell-shape” profiles were found in the intestinal permeabilities of lipidated TG and DADLE analogues (C₂- to C₈-, Fig. 7e) [12, 37]. Due to strong membrane binding, C₈-Phe-Gly (Fig. 7f) was retained in the skin stratum corneum and its diffusion across intestinal epithelia was reduced as compared to that of C₄-Phe-Gly and C₆-Phe-Gly [89, 115]. Similarly, N-terminal lipidated TG analogues (Fig. 7a) showed decreased transdermal permeabilities with the elongation of the fatty acid chain, C₂-TG > C₄-TG > C₆-TG [36]. Prolonged absorption phases were found in the pharmacokinetic study of DiC₁₆-desmopressin (Fig. 7g, *i.v.* tail vein injection) [45], insulin detemir (subcutaneous injection, (*s.c.*)) [117], Mal-sCT and REAL-sCT (*s.c.*) [49].

2.2. Metabolic Stability

Lipidated peptides bind to serum albumin; this phenomenon affects serum metabolism of peptides. N^{6,B29}-Palmitoyl-human insulin (Fig. 7d) was co-eluted with albumin in size exclusion chromatography [29]. The insulin-secretion stimulating activity of the lipidated GIP analogue (Lys³⁷-Palmitoyl)GIP (Fig. 8a) was correlated with the concentrations of albumin [19]. The uptake of lipidated Bowman-Birk protease inhibitor (BBI) analogue BBIsPal into the Caco-2 cell monolayers was increased up to 140-fold more than that of the native BBI in serum-free medium, but this uptake was reduced dramatically after adding fetal bovine serum [42]. The radioactivity of the supernatant of the mixture of lipidated hPP analogue ⁶⁸Ga-DOTA-[K¹³(E-Pamitoyl),Q³⁴]hPP and plasma after precipitation was dramatically decreased compared to that of the supernatant of the mixture of non-lipidated hPP analogue ⁶⁸Ga-DOTA-[K¹³,Q³⁴]hPP and plasma (Fig. 6d), indicating the strong binding of plasma protein with the lipopeptide [21].

The molecular biological mechanism of the interactions of lipidated peptides with serum albumin has not been fully studied. While the mechanism of the interactions of fatty acids with serum albumin shed a light on this unknown mechanism [118-121]. Seven common fatty acid binding sites among the identified eleven binding sites on human serum albumin were classified into three high-affinity binding sites (site V, IV and II) and four low-affinity binding sites (site I, III, VI and VII) [122-124]. Fatty acids have diverse binding abilities on those binding sites. Medium- and long-chain fatty acids prefer binding on sub-domains IIA and IIIA, respectively [122]. The binding of insulin detemir at IIA and IIIA domains can be displaced by long-chain fatty acids [27, 125].


Many lipidated peptides showed increased plasma stability, such as lipidated analogues of TG, RC-160, sCT, hPP, galanin, NT, NPY, NPW and insulin. The half-life of C₆-TG in rat plasma (10 mg protein per mL) was 300-fold higher than the half-life of the native TG, 32 h and 6.48 min, respectively [37]. The half-life of C₁₂-RC-160 in mouse serum (90%) was increased up to 12-fold compared to the half-life of the native RC-160, 24 h and 2 h, respectively [32]. The half-lives of lipidated analogues of galanin (17 mer, in 25% rat serum), salmon calcitonin (32 mer, mice, *s.c.*) and liraglutide (31 mer, pig, *s.c.*) were also increased but shorter than 12 h [18, 20, 51]. The half-life of ⁶⁸Ga-DOTA-[K¹³(E-Palmitoyl),Q³⁴]hPP in rat after tail vein injection was 25 min, while the half-life of the non-lipidized analogue ⁶⁸Ga-DOTA-[K¹³,Q³⁴]hPP was 3.6 min [21].

The chain length of fatty acids plays an important role in modulating the stability of lipidated peptides. Systemic study of the effects of the chain length of fatty acids on plasma stability of modified galanin analogues resulted in a bell-shape profile with the greatest stability achieved at myristoylation analogue (C₁₄-galanin), likely due to its albumin binding properties [18]. The stability of

Lipidation also has significant impacts on the stability of lipidated peptides in tissues and organs. Lipidated peptides [Tyr³]REAL-octreotide (Fig. 8c) [53], Mal-sCT [50] and [K⁶⁸(E-Pam)]SDF1 α (Fig. 8d; SDF1 α , stromal cell-derived factor-1 α) [126] were more stable (2 to 12-fold) than their parent peptides in liver slices, liver fluid and liver homogenates, respectively. Lipidated analogues of Phe-Gly with C₄-, C₆- and C₈- (Fig. 7f) showed similar stabilities in rat skin homogenates [89]. The stability trends of lipidated TG analogues (Fig. 7a) were different in jejunum and colon homogenates (C₁₂-TG >> C₂-TG > C₆-TG >> TG) [37] and in Caco-2 homogenates (C₄-TG > C₂-TG > C₆-TG >> TG) [35]. The organ stabilities of C₆-TG were in rectum > liver > colon > jejunum [37]. The half-lives of diC₁₀-Leu-enkephalin in small intestinal mucosal homogenate, liver homogenate and plasma were ~12 min, ~80 min and ~193 h, respectively [13].

The distribution and elimination of lipidated peptides were protracted due to their serum albumin binding properties. DiC₁₀-Leu-enkephalin, DiC₁₆-desmopressin (Fig. 7g), REAL-octreotide and ⁶⁸Ga-DOTA-[K¹³(E-Palmitoyl),Q³⁴]hPP were accumulated in the liver [13, 21, 45, 53]. Hedgehog antagonist SMOi2-17 (Ac-AKFSTHWAYTL(N⁶-Pal-K), all-D) was accumulated in the liver and the lung after 30 min following *i.v.* administration [22]. The brain concentration of this lipidated peptide reached up to 10 nM; the blood concentration of this peptide was 400 nM [22]. DiC₁₀-Leu-enkephalin was eliminated into the small intestine with bile following *i.v.* administration, while the native Leu-enkephalin was quickly degraded in blood [13]. DiC₁₆-desmopressin showed a similar anti-diuretic linear dose-response relationship as the native desmopressin but at 250-fold less dosage [45].

Side chain lipidation on N^ϵ -lysine was extensively applied in many peptides. Lipidation of N^ϵ -^{B29}-Lys of insulin (Fig. 7d) with fatty acids (from C₁₀- to C₁₆-) increased the stability, the duration of action and the glucose-lowering effects of insulin [23-25, 27]. Insulin detemir, (N^ϵ -^{B29}-Myristoyl)-des-30B-L-Thr-human insulin (Fig. 1c), demonstrated a great serum albumin binding affinity and a blood glucose-lowering effect. This long-acting (up to 24 hours)



anti-diabetic drug was launched in 2004 with a brand name Levemir®. (N^{B1} -Caproyl)-($N^{E,B29}$ -caproyl)-insulin (Fig. 7d) showed significant absorptions in small intestine and large intestine *in vitro*; it also showed equal hypoglycemic activity when compared to (N^{B1} -caproyl)-insulin and the native insulin following *i.v.* injections in rats [23, 24]. The hypoglycemic activity was decreased slightly for the (N^{B1})- or (N^{B1} , $N^{E,B29}$)-myristoyl and palmitoyl lipidated insulin analogues [24].

GIP is a promising glucose-dependent anti-diabetic drug candidate, but GIP is degraded rapidly by dipeptidyl peptidase IV (DPP-IV). Moreover, the metabolite GIP(3-42) is an antagonist to the GIP receptor. Palmitoylation on N^E -Lys¹⁶ or N^E -Lys³⁷ of GIP and N -pGluGIP (Fig. 8a) dramatically improved the metabolic stability and the antihyperglycemic and insulinotropic activities of GIP and N -pGluGIP following *i.p.* injections in mice [7, 19]. (N^E -Palmitoyl-Lys¹⁶) N -pGluGIP and (N^E -Palmitoyl-Lys³⁷) N -pGluGIP were fully intact even after a 24 h incubation with DPP-IV, while the native GIP was completely degraded by DPP-IV in 8 h [7]. Similarly, the GIP receptor antagonist analogues, (Pro³)GIP and (Hyp³)GIP lipidated at N^E -Lys¹⁶ with palmitoyl, showed significantly improved stabilities toward DPP-IV and enhanced inhibitions of GIP-stimulated cAMP production and cellular insulin secretions [129, 130].

GLP-1 is a promising anti-diabetic drug candidate lacking hypoglycemic side effects, but GLP-1 is degraded rapidly by DPP-IV. The half-life of GLP-1 is only 2 min (*i.v.*) or 1 h (*s.c.*) [131-133]. Lipidations of GLP-1 led to the discovery of the recently launched once-daily (*s.c.*) anti-diabetic drug liraglutide (Arg³⁴-Lys²⁶-(N^E -(γ -Glu(N -palmitoyl))-GLP-1(7-37), Fig. 1b). Liraglutide has a prolonged half-life (10~12 h in humans, *s.c.*) and equipotent receptor binding ability toward GLP-1 receptor as the native GLP-1 [134-136]. The lipidation analogues of GLP-1 (from C₁₀- to C₁₈-, Fig. 1b) via the spacer γ -glutamic acid insertion have no discernible effects on GLP-1 receptor binding potency with the exception of the C₁₈-analogue [20]. The half-lives of lipidated GLP-1 analogues following *s.c.* administration in pigs were protracted with the


elongation of the chain length of fatty acids. A phase I clinical trial for an oral variant of liraglutide, NN9924, was initiated recently. Inspired by the success of liraglutide, human pancreatic polypeptide (hPP) was modified with palmitoyl lipidation via the spacer γ -glutamic acid insertion at Lys¹³; this lipidated hPP analogue (Fig. 6d) demonstrated a prolonged duration of anorectic effect [21].

Lipidation analogue of cecropin A-melittin hybrid peptide CA(1-7)M(2-9) at N^E -Lys⁷ with C₈-, C₁₂-, C₁₄- and C₁₆- (Fig. 9c) showed impaired secondary structures and decreased bactericidal activities, whereas the analogues that were lipidated at terminal positions (N^{1a} - or N^{1e} -) exhibited 7-fold increases in the inhibition of promastigote or amastigote proliferation as compared to the native CA(1-7)M(2-9) [137].

The half-lives of lipidated human calcitonin analogues N^E -acetyl-Lys¹⁸-human calcitonin and N^E -caproyl-Lys¹⁸-human calcitonin (Fig. 9d) were increased up to 3 to 10-folds compared to the native human calcitonin in small intestine and large intestine mucosal homogenates [138]. Both lipidated analogues showed improved hypocalcemic effects following administration into small intestinal loop, while the native human calcitonin was not active [138].


Our research group developed a peptide delivery method into the central nervous system (CNS) via the conjugation of a lipidated oligolysine motif (so called "cationization-lipidization motif") [14-18, 31, 139]. This strategy was successfully applied to four neuropeptides: galanin (Fig. 2c) [14, 17, 18], NT (Fig. 9e) [15], NPY (Fig. 9f) [15] and NPW (Fig. 9g) [31]. Neuropeptides modified with this strategy showed increased BBB penetration, improved serum stability, preserved peptide receptor binding properties and enhanced anticonvulsant activities.

Galanin is an endogenous neuropeptide exhibiting an anticonvulsant activity. Gal-LAA, lipidated at the C-terminus of galanin (1-16) (Table 2), did not demonstrate any anticonvulsant activity following systemic administration [14]. Surprisingly, a combination of lipidation and oligolysine "cationization" yielded




R = acetyl, methoxycarbonyl

(10a) Dermorphine



(10b) Peptide A2



(10c) TP5 analogues



(10d) Core peptide

Fig. (10). Structures of the lipidated analogues of dermorphine, peptide A2, TP5 and core peptide.

potent, systematically-active anticonvulsant compounds [14]. Interestingly, the anchor position of the palmitoyl moiety and the presence of cationic amino acids (Lys, Arg and/or His) significantly affected the bioactivities of the lipidated galanin analogues [14]. The SAR study of lipidated analogues of Gal-B2 (Fig. 2c) revealed that the anticonvulsant activities were increased with the elongation of fatty acid chains from C₆ to C₁₈ but were saturated at Gal-B2-C₁₆ and Gal-B2-C₁₈ in mice [18]. Several galanin analogues that exhibit promising activities in suppressing seizures in several animal models of epilepsy are under preclinical evaluations [17].

Anticonvulsant neuropeptides NPY and NT are not active *in vivo* after systemic delivery. The two neuropeptides conjugated with cationization-lipidation motif, NPY-BBB2 and NT-BBB1, exhibited potent anticonvulsant activities (less than 1 mg/kg, *i.p.*) in the 6 Hz epilepsy model [15]. Central insertion of the cationization-lipidation motif into NPW also yielded active anticonvulsant compounds [31]. Taken together, the cationization-lipidation conjugation provides an attractive modification method to improve peptides' CNS activity and to generate new drug leads.

3.3. Esterification with Fatty Acids

Esterification is often regarded as a reversible lipidation method due to its hydrolysis propensity *via* enzyme-mediated degradation resulting in the release of intact peptides. Reversible posttranslational S-acylation of proteins plays many important roles in protein membrane association, stability and cellular signaling [6]. The inactive analogue [Ac-(O-Palmitoyl)Ser¹-Val⁵]Ang (Fig. 8b) recovered its innate bioactivity after being hydrolyzed to [Ac-Ser¹-Val⁵]Ang by mild saponification or lipolysis in an aorta strip contraction assay [41].

Ghrelin (Fig. 1a) is the only known endogenous hormone which is naturally lipidized with octanoyl at Ser³; decanoylation or decenoylation analogues were also exist [2, 47]. The hydrophobic interaction of the octanoyl group of ghrelin with human growth hormone secretagogue receptor type 1A (hGHSR1a) is essential for its anti-obesity bioactivity. Lipidations of [Ser³]ghrelin with other fatty acids were also tolerable, such as octyl, 3-octenoyl, 2,4,6-octatrienoyl, 4-methylpentanoyl, 2-propylpentanoyl-, undecanoyl, 3-phenylpropionyl, adamantaneacetyl and other long-chain fatty acid lipidations [46, 48]. Site-specific activities were exhibited in the ghrelin analogues lipidated at different positions of serine [46]. [(O-Octanoyl)-Ser²]ghrelin preserved equipotent activity as the native ghrelin in aequorin bioluminescence functional assay but [(O-octanoyl)-Ser⁶]ghrelin and [(O-octanoyl)-Ser¹⁸]ghrelin were not active [46].

Acylation on the hydroxy group of the tyrosine of dermorphin analogue with various fatty acids (Fig. 10a) increased the oral bioavailabilities of the peptides; acetylation and methoxycarbonylation analogues showed similar antinociceptive activities as the parent peptide [140].

3.4. The REAL Technology and Cysteine Lipidation

The reversible aqueous lipidation technology (REAL) aims to lipidize peptides in aqueous solution by the disulfide bond formation of (*N*-palmitoylcysteinyl)-2-pyridyl-disulfide (Pal-CPD) with thiol groups of either Cys or a coupled thiopropionyl moiety on peptide chain [42]. The REAL technology increases peptides' plasma stability, blood circulation, gastrointestinal absorption and accumulation in the liver. The REAL technology was applied successfully in the modification of desmopressin (Fig. 7g) [45], salmon calcitonin (Fig. 3a) [51, 53], octreotide (Fig. 8c) [51, 53] and soy Bowman-Birk protease inhibitor (BBI) [42-44].

DiC₁₆-desmopressin was synthesized by the conjugation of Pal-CPD with thiol groups of Cys⁵ and *N*-thioacetyl of desmopressin [45]. DiC₁₆-desmopressin showed 4-fold duration of

anti-diuretic activity without body weight loss in rats when compared to desmopressin (4 days vs. 1 day) [45]. REAL-sCT (Fig. 3a) showed similar plasma calcium-lowering ability as the native sCT following *i.v.* administration [51]. Oral administration of REAL-sCT at a 10-fold higher dosage demonstrated similar effects as *s.c.* administration in stabilizing urinary deoxypyridinoline level. REAL-sCT was mainly retained in the blood and liver. REAL-octreotide was more stable in plasma and liver slices and more potent in the inhibition of growth hormone release following *s.c.* administration in rats as compared to that of the native octreotide [53].

BBIssPal was synthesized with REAL technology *via* the conjugation of Pal-CPD with 3-thiopropionyl-Lysine groups of BBI (8 KDa) [42, 43]. The blood concentration following *i.v.* tail injection in mice and the uptake across the Caco-2 cell monolayers of BBIssPal was increased up to 10-fold and 140-fold, respectively, when compared to the native BBI. The potency of BBI in preventing the carcinogen-induced transformation of C3H10T1/2 cells was also maintained in BBIssPal. BBIssPal was accumulated predominantly in the liver. The other lipidated analogue of BBI modified with REAL technology on cysteines, BBI-Pal, exhibited a long retention time in the stomach following oral administration [44].

S-Alkylation of sCT with *N*^ε-maleimido-L-Lys(Palmitoyl)-OH in aqueous medium yielded Mal-sCT (Fig. 3b) [49, 50]. Mal-sCT showed enhanced stabilities in the homogenates of intestine and liver, increased uptake across the Caco-2 cell monolayers, and extended hypocalcemic activity following oral administration as compared to that of the native sCT.

3.5. Lipoamino Acids (LAAs)

The application of unnatural lipoamino acids (LAAs, Fig. 2a) were developed by the Toth group [73]. Lipidation of ryanodine receptor agonist [D-Arg¹⁸]peptide A2 (Fig. 10b) with LAA maintained peptides' α -helical structures, increased peptides' Ca²⁺ release activities and permeabilities across the Caco-2 cell monolayers as compared to that of the native peptide [62]. LAA conjugations enhanced the serum stability of thymopentin TP5 (RKDVY, Fig. 10c) [61, 141]. The plasma half-lives of [(TP5)₄-lys]₂-lys-LAA₁₄-LAA₁₄ and TP5-LAA₁₄ were increased up to 7 and 18 min, respectively; while the half-life of the native TP5 was only ~1 min [61]. The C-terminal lipidated analogues of Glu-Trp-NH₂ with LAAs showed increased stabilities in Caco-2 cell homogenates and permeabilities across the Caco-2 monolayers [142]. Lipidated analogue of core peptide (CP, a transmembrane sequence of T-cell antigen receptor α -chain) LAA₁₀-CP (Fig. 10d) showed increased permeability across Caco-2 cell monolayers and anti-arthritis activity in rats as compared to that of the native CP [59, 60]. The AUC and C_{max} of low molecular weight heparin were improved following co-administration of polylysine-(LAA₁₀)₂ per oral [63]. Of note, the enantiomeric diversity of LAAs should be considered in LAA lipidation [30, 69, 70, 143].

3.6. Lipidation with Amino Fatty Acids

Lipidation with amino fatty acids produced mixed results on the modification of several peptides. The serum stability and the anti-bacterial activity of the pharmacophore of tenecin-1 were increased following *N*-terminal lipidation with 6-aminoheptanoic acid (Ahx) (Fig. 11a) but *N*-terminal acetylation or benzylation analogues showed decreased antibacterial activity due to lacking the *N*-terminal amino group [57]. Replacing the palmitoyl group of liraglutide (Fig. 1b) with 11-aminoundecanoyl dramatically decreased its stability in pigs (half-lives 16 h vs 1.1 h, respectively) [20]. Replacing the octanoyl group of ghrelin (Fig. 1a) with 7-aminoheptanoyl (Ahp) decreased its binding affinity and agonist activity on hGHSR1a [46].

Amino fatty acids were also exploited as backbone spacers to replace noncritical amino acids in peptide chain. Replacing Pyroglu¹-Asn² and Gly⁶-Gly⁷ of conotoxin SIIIA with Ahx yielded Ahx-SIIIA (Fig. 11b). Ahx-SIIIA showed improved inhibitions of TTX-sensitive sodium current in dorsal root ganglia neurons and compound action potentials in the sciatic nerve and enhanced analgesic activity in the formalin pain model [54]. Replacing the non-essential Gly²-Gly³ of Met-enkephalin with aminobutanoyl (Abt), aminopentanoyl (Apn), Ahx or Ahp (Fig. 11c) yielded lipidated Met-enkephalin analogues with reduced affinities and agonist activities on opioid receptors [55]. However, these analogues showed analgesic and anticonvulsant activities comparable to the native Met-enkephalin. Replacing Gly-Gly-Gly moiety of TAN-1511A with 4-aminobenzoyl and 4-aminomethylbenzoyl (Fig. 11d) improved the peptide's proliferative activities in bone marrow cells, but this improvement was not found in the lipidated analogues substituted with Ahx and aminooctanoyl (Aot) [56]. Replacing non-critical amino acids of NPY with Ahx yielded lipidated NPY analogues (Table 2) with similar receptor binding abilities [144].

3.7. Dicarboxylic Acid Modification

Dicarboxylic acid acylation transforms the positively charged cationic amine group into the negatively charged anionic acid group in the lipidated peptides. The most successful application of this strategy is the discovery of ultra-long-acting basal insulin Degludec (*N*^{ε29B}-[*N*-(15-carboxypentadecanoyl)-L-γ-glutamyl]-des-30B-L-threonine-human insulin, NN1250, Fig. 11e), which is currently being evaluated in a phase III clinical trial [145]. The *N*-terminal lipidation of growth hormone-releasing hormone antagonist GHRH(1-29)NH₂ (Fig. 6e) with dicarboxylic acids (ranging from heptane diacid to hexadecane diacid) and fatty acids were systemically studied [58]. Decane diacid-GHRH(1-29)NH₂ showed similar binding affinity and inhibitory ability as compared to that of decanoyl-GHRH(1-29)NH₂. All analogues had significant anti-

proliferative activities against MiaPaCa-2 human pancreatic cancer cell line [58].

3.8. Miscellaneous Lipidation Strategies

The *C*-terminus esterification analogues of DADLE *via* hydrazine spacer with C₂-, C₄-, C₆- or C₈- (Fig. 7e) showed increased permeabilities across intestine and stabilities *in vitro* as compared with the native DADLE [12]. The *C*-terminal esterification of dermorphin (Fig. 10a) with various aliphatic alcohols increased its oral bioavailability and maintained its antinociceptive activity [140]. Modification of core peptide (CP, Fig. 10d) with *C*-terminal multi-palmitoyl groups *via* spacer Gly-Tris improved the binding ability to the lipid membrane; mono- and tri-palmitoyl-Tris-Gly-CP analogues inhibited IL-2 production more effectively than CP [59]. Lipidation on the imidazole group of histidine was reported in the chemical modification of thyrotropin-releasing hormone (TRH), but this strategy was not continued due to the unstable bond of acyl-imidazole [146, 147].

A novel isostere of palmitoyl cysteine, (S)-2-Amino-4-(2-pentadecyl-1,3-dioxolan-2-yl)-butanoic acid, was reported to substitute *S*-palmitoyl moiety of H-Ras protein [148]. Fmoc-Met-enkephalin exhibited increased BBB permeability, enhanced analgesic activity, and controlled release following spontaneous chemical hydrolysis *in vivo* [149]. Replacing the octanoyl group of human ghrelin with 1-adamantaneacetyl or 8-bromooctanoyl maintained its binding affinity to receptor hGHSR1a [46]. Conjugation on the C34 fragment of HIV fusion inhibitor gp41 with cholesterol moiety Gly-Ser-Gly-Cys(cholesterol) dramatically improved its antiviral potency [66]. Conjugation on *N*^{B29}-insulin with one or two deoxycholic acids maintained its binding to human hepatoma cell line HepG2; these analogues produced similar blood glucose nadir as the native insulin but with a longer duration following *i.v.* administration [67, 68].




Fig. (11). Structures of the lipidated analogues of tenecin-1, Ahx-SIIIA, Met-enkephalin, TAN-1511A and Degludec.

Glycosylation compensates the low aqueous solubility of lipidated peptides and increases their membrane transportation through glucose transporters. Liposaccharide, the combination of LAA lipidation and glycosylation, is applied in the development of small-molecule drugs, peptides, vaccines and antibiotics [59, 72, 73, 80, 150-154]. Hemolytic potency of liposaccharides was correlated with the chain length of LAA [72]. Two naturally-occurring ghrelin-like peptides lipidated with octanoyl at Ser³ and glycosylated with mucin-type sugar chain (*N*-acetyl-hexosamine)₃-(hexose)₂ at Thr¹¹ or Ser¹⁰-Thr¹¹ were also found in red stingray *Dasyatis akajei* [155].

It is noteworthy that stapled peptides can also be considered as "lipidated" peptides with non-natural olefin amino acids inserted into the peptide backbone. The helical contents of the analogues of BH3 segment of BID stapled with alkenes 4-octene (i+4 span) or 4-tetradecene (i+7 span) were increased up to 5-fold [156]. Properties of the stapled peptides are reviewed in [157].

Insertion of glycine, glutamic acid, amino fatty acids, PEG or hydrazine between fatty acids and peptides is a useful strategy to minimize the disturbance of fatty acids on the bioactivities of lipidated peptides. For example, the spacer γ -Glu conjugated between Lys²⁶ of Arg³⁴-GLP-1(7-37) and fatty acids was important in maintaining the receptor binding potency [20].

Reports on unfavorable results of lipidated peptides can provide useful information on designing peptide drug leads. The CNS activity of *N*-lauroyl-TRH were decreased in pentobarbital sleeping assay and the endocrine activity in thyroid-stimulating hormone releasing activity assay [38]. Replacing the palmitoyl moiety of liraglutide with 2-hexyldecanoyl, 10-cyclohexane-decanoyl, 10-benzyldecanoyl or 4-pentylbenzenesulfonyl abolished the peptides' receptor binding affinity [20, 41]. Alkyl ether [Sar¹,Tyr⁴(O-C₁₈H₃₇)]Ang showed a very low binding affinity on bovine adrenocortical membranes as compared to human angiotensin analogues [Sar¹]Ang and [Sar¹,Tyr⁴(O-Pal)]Ang [41].

Hemolysis was found in long-chain fatty acid lipidated peptides due to their strong membrane binding affinity [98, 127]. Cytotoxicity and aggregation propensity were related to the position of the octanoyl group on β -amyloid [93]. The analogues of β -amyloid lipidated at Lys¹⁶ and Lys²⁸ with octanoylation showed more toxic than the native β -amyloid and *N*-octanoyl- β -amyloid toward Neuro2A cells [93]. Lipidation of 2a (Fig. 3f) with 1-adamantanecarbonyl transformed the structure of 2a from α -helix into β -sheet and formed amyloid fibrils in aqueous solution [158].

4. CHEMICAL SYNTHESIS METHODS

Automated peptide synthesis on solid support is the most commonly used method for the syntheses of lipidated peptides. Building blocks such as *N*^ε-acyl-Lys and LAA can be chemically synthesized and directly coupled on resin by the Fmoc solid phase peptide synthesis (SPPS) method [18, 159-161]. *S*-Acylated cysteine is not recommended due to possible aminolysis by the α -amino group [162].

The on-resin lipidation is an applicable strategy for the syntheses of long-chain peptides *via* orthogonal protecting strategy. Fmoc-orthogonal protecting groups for the side chain of lysine can

be selected from Dde-, Nosyl-, Alloc-, Mtt- and Mmt [163]. Dde group can be deprotected with hydrazine or NH₂OH·HCl/imidazole in NMP/CH₂Cl₂ [161, 164, 165]. Nosyl group can be deprotected with *p*-methoxythiophenol/K₂CO₃ in CH₃CN/ DMSO [166, 167]. Alloc can be deprotected with Pd⁰ catalysis [161]. Trityl protecting groups Mmt, Mtt and PhFI can be removed by 1%-3% TFA in CH₂Cl₂ with scavenger 2% TES [168, 169]. Mmt protecting strategy is also applicable in the side chain protection of Ser/Thr and Cys residues [170-172]. Meanwhile, Cys(Trt) needs 50% TFA to deprotect Trityl group [172]. 2-ClTrt protecting group on the side chain of Tyr is also sensitive to 2% TFA [173]. Deprotecting tert-Butylsulfanyl group of Cys(StBu) with PBu₃ may cause β -elimination [162].

Boc- group can be deprotected with trimethylsilyl trifluoromethanesulfonate (TMSOTf) with lutidine; this deprotecting method is orthogonal with Fmoc and unsaturated double bond [174]. Unsaturated fatty acid lipidations such as prenylation are sensitive to strong acids; special linkers such as trityl, oxime, hydrazino-benzoyl, (2-phenyl-2-trimethylsilyl)ethyl (PTMSEL) and ellman sulfonamide should be considered [6, 74, 169, 175].

Solution lipidation methods such as REAL technology are attractive strategies for peptides and proteins [13, 45, 49, 51]. Palmitoyl esterification of Ser or Cys can be accomplished by conjugating palmitoyl chloride to peptides in TFA solvent [176]. The TFA condition prevents acylation at amino groups. Rearrangement of *S*-acyl to *N*-acyl in terminal Cys lipidation can occur in the Fmoc deprotecting process with piperidine, so 1% DBU should be used to remove Fmoc [175]. On the other hand, the rearrangement can be exploited as a peptide lipidation method. Conjugating hydrazinoacetyl peptides with lipidated α -oxo-aldehyde is a convenient *in situ* lipidation method and can be applied in parallel synthesis (Fig. 12) [177-180].

5. SUMMARY AND OUTLOOK

The modification of peptides and proteins *via* lipidation has been continuously growing since 1970's; the marketing of the lipidated peptide drugs is a considerable validation that lipidation is a successful strategy to develop more peptide drug leads in the future. As discussed herein, lipidation improved pharmacological and pharmaceutical properties of many peptides. It becomes apparent that even subtle changes in lipidation strategy, including the type of fatty acids, lipid anchor position or spacer insertion can produce profound impacts on the physicochemical properties and bioactivities of lipidated peptides.

Extensive SAR studies are necessary to take full advantage of lipidation to convert peptides into drugs, since mixed results on ADME properties of lipidated peptides emphasize the complexity of this modification strategy. With systematic and judicious approaches, lipidation will increase peptides' hydrophobicity, secondary structure, self-assembly, albumin binding, metabolic stability, bioavailability and ADME properties while maintaining peptides' superior bioactivity including potency and specificity for target receptors. With increased interest of exploring peptides as therapeutics, lipidation will continue to be one of the leading chemical modifications to produce clinical drug candidates.




Fig. (12). In situ hydrazine lipidation strategy.

ACKNOWLEDGEMENTS

This work was supported by NIH grant U01 NS066991. We thank Dan McDougale and Massiell L. German for critical reading of the manuscript. Conflict of Interest: G.B. is a scientific founder of NeuroAdjuvants, Inc.

ABBREVIATIONS

Abt	=	aminobutanoyl
ADME	=	absorption, distribution, metabolism and excretion
Ahp	=	aminoheptanoyl
Ahx	=	aminohexanoic acid
β -Alanine	=	3-aminopropanoic acid
Alloc	=	allyloxycarbonyl
Aot	=	aminooctanoyl
Apn	=	aminopentanoyl
AUC	=	area under the curve
Ava	=	5-amino-valeric acid
BBB	=	blood-brain barrier
BBi	=	Bowman-Birk protease inhibitor
C ₂	=	acetyl
C ₃	=	propanoyl
C ₆	=	hexanoyl or caproyl
C ₈	=	octanoyl
C ₁₀	=	decanoyl
C ₁₂	=	lauroyl
C ₁₄	=	myristoyl
C ₁₆	=	palmitoyl
C ₁₈	=	stearoyl
CD	=	circular dichroism
2-ClTrt	=	2-chlorotrityl
CMC	=	critical micelle concentration
CNS	=	central nervous system
CP	=	core peptide
Dde	=	4,4-dimethyl-2,6-dioxocyclohex-1-ylidene)ethyl
DBU	=	1,8-diazabicyclo[5.4.0]undec-7-ene
DIC	=	<i>N,N'</i> -diisopropylcarbodiimide
DPC	=	dodecylphosphocholine
DPP-IV	=	dipeptidyl peptidase IV
DPPC	=	dipalmitoylphosphatidylcholine
5-FAM	=	5'-isomer of fluorescein
Fmoc	=	<i>N</i> -(9-fluorenyl)methoxycarbonyl
GalR1	=	galanin receptors subtype 1
GalR2	=	galanin receptors subtype 2
GHRH	=	growth hormone-releasing hormone
GIP	=	gastric inhibitory polypeptide or glucose-dependent insulinotropic polypeptide
GLP-1	=	glucagon-like peptide-1
GPCRs	=	G-protein coupled receptors
Har	=	homoarginine

hGHSR1a	=	human growth hormone secretagogue receptor type 1A
HOBt	=	1-hydroxybenzotriazole
hPP	=	hormone pancreatic polypeptide
IGF1R JM	=	insulin-like growth factor 1 receptor juxtamembrane segment
<i>i.p.</i>	=	intraperitoneally
<i>i.v.</i>	=	intravenous
LAA	=	lipoamino acid
LAA ₈	=	2-amino-D,L-octanoic acid
LAA ₁₀	=	2-amino-D,L-dodecanoic acid
LAA ₁₂	=	2-amino-D,L-lauric acid
Mmt	=	4-methoxytrityl
Mtt	=	4-methyltrityl
Nle	=	norleucine
Nosyl	=	2-nitro-benzenesulfonyl
NPY	=	neuropeptide Y
NT	=	neurotensin
Pal-CPD	=	(<i>N</i> -palmitoylcysteinyl)-2-pyridyl-disulfide
PBu ₃	=	tributylphosphine
Phe(<i>p</i> Cl)	=	<i>p</i> -chlorophenylalanine
pGlu	=	paroglutamyl
PhFI	=	9-phenylfluoren-9-yl
POPC	=	1-palmitoyl-2-oleoyl-phosphatidylcholine
POPG	=	1-palmitoyl-2-oleoyl-phosphatidylglycerol
PTMSEL	=	(2-phenyl-2-trimethylsilyl)ethyl
REAL	=	reversible aqueous lipidation
Sar	=	sarcosine
SAR	=	structure-activity relationship
<i>s.c.</i>	=	subcutaneous injection
sCT	=	salmon calcitonin
SDF1 α	=	stromal cell-derived factor-1 α
SDS	=	sodium dodecylsulphate
SPPS	=	solid phase peptide synthesis
STAT	=	signal transducer and activator of transcription protein
TES	=	triethylsilane
TFA	=	trifluoroacetic acid
TFE	=	2,2,2-trifluoroethanol
TG	=	tetragastrin
<i>T_m</i>	=	melting temperature
TMSOTf	=	trimethylsilyl trifluoromethanesulfonate
TRH	=	thyrotropin-releasing hormone.

REFERENCES

- [1] Strieker, M.; Marahiel, M.A. The structural diversity of acidic lipopeptide antibiotics. *ChemBioChem*, **2009**, 10 (4), 607-616.
- [2] Hosoda, H.; Kojima, M.; Mizushima, T.; Shimizu, S.; Kangawa, K. Structural divergence of human Ghrelin. *J. Biol. Chem.*, **2003**, 278 (1), 64-70.
- [3] Kojima, M.; Hosoda, H.; Date, Y.; Nakazato, M.; Matsuo, H.; Kangawa, K. Ghrelin is a growth-hormone-releasing acylated peptide from stomach. *Nature*, **1999**, 402 (6762), 656-660.

- [4] Linder, M.E.; Deschenes, R.J. Palmitoylation: policing protein stability and traffic. *Nat. Rev. Mol. Cell Biol.*, **2007**, *8* (1), 74-84.
- [5] Walsh, C.T.; Garneau-Tsodikova, S.; Gatto, G.J. Protein posttranslational modifications: the chemistry of proteome diversifications. *Angew. Chem. Int. Ed.*, **2005**, *44* (45), 7342-7372.
- [6] Brunsfeld, L.; Kuhlmann, J.; Waldmann, H. Synthesis of palmitoylated Ras-peptides and -proteins. *Methods*, **2006**, *40* (2), 151-165.
- [7] Irwin, N.; Green, B.D.; Gault, V.A.; Greer, B.; Harriott, P.; Bailey, C.J.; Flatt, P.R.; O'Harte, F.P. Degradation, insulin secretion, and antihyperglycemic actions of two palmitate-derivitized N-terminal pyroglutamyl analogues of glucose-dependent insulinotropic polypeptide. *J. Med. Chem.*, **2005**, *48* (4), 1244-1250.
- [8] Laczkó, I.; Váró, G.; Bottka, S.; Bálint, Z.; Illyés, E.; Vass, E.; Bertrand, J.-R.; Malvy, C.; Hollósi, M. N-terminal acylation of the SV40 nuclear localization signal peptide enhances its oligonucleotide binding and membrane translocation efficiencies. *Arch. Biochem. Biophys.*, **2006**, *454* (2), 146-154.
- [9] Tarwadi; Jazayeri, J.A.; Prankerd, R.J.; Pouton, C.W. Preparation and *in vitro* evaluation of novel lipopeptide transfection agents for efficient gene delivery. *Bioconjugate Chem.*, **2008**, *19* (4), 940-950.
- [10] Robinson, L.R.; Fitzgerald, N.C.; Doughty, D.G.; Dawes, N.C.; Berge, C.A.; Bissett, D.L. Topical palmitoyl pentapeptide provides improvement in photaged human facial skin. *Int. J. Cosmet Sci.*, **2005**, *27* (3), 155-160.
- [11] Kellam, B.; Drouillard, B.; Dekany, G.; Starr, M.S.; Toth, I. Synthesis and *in vitro* evaluation of lipooamino acid and carbohydrate-modified enkephalins as potential antinociceptive agents. *Int. J. Pharm.*, **1998**, *161* (1), 55-64.
- [12] Uchiyama, T.; Kotani, A.; Tatsumi, H.; Kishida, T.; Okamoto, A.; Okada, N.; Murakami, M.; Fujita, T.; Fujiwara, Y.; Kiso, Y.; Muranishi, S.; Yamamoto, A. Development of novel lipophilic derivatives of DADLE (leucine enkephalin analogue): intestinal permeability characteristics of DADLE derivatives in rats. *Pharm. Res.*, **2000**, *17* (12), 1461-1467.
- [13] Wang, J.; Hogenkamp, D.J.; Tran, M.; Li, W.Y.; Yoshimura, R.F.; Johnstone, T.B.; Shen, W.C.; Gee, K.W. Reversible lipidization for the oral delivery of leu-enkephalin. *J. Drug Target.*, **2006**, *14* (3), 127-136.
- [14] Bulaj, G.; Green, B.R.; Lee, H.K.; Robertson, C.R.; White, K.; Zhang, L.; Sochanska, M.; Flynn, S.P.; Scholl, E.A.; Pruess, T.H.; Smith, M.D.; White, H.S. Design, synthesis, and characterization of high-affinity, systemically-active galanin analogues with potent anticonvulsant activities. *J. Med. Chem.*, **2008**, *51* (24), 8038-8047.
- [15] Green, B.R.; White, K.L.; McDougle, D.R.; Zhang, L.; Klein, B.; Scholl, E.A.; Pruess, T.H.; White, H.S.; Bulaj, G. Introduction of lipidization-cationization motifs affords systemically bioavailable neuropeptide Y and neurotensin analogs with anticonvulsant activities. *J. Pept. Sci.*, **2010**, *16* (9), 486-495.
- [16] Robertson, C.R.; Scholl, E.A.; Pruess, T.H.; Green, B.R.; White, H.S.; Bulaj, G. Engineering galanin analogues that discriminate between GalR1 and GalR2 receptor subtypes and exhibit anticonvulsant activity following systemic delivery. *J. Med. Chem.*, **2010**, *53* (4), 1871-1875.
- [17] White, H.S.; Scholl, E.A.; Klein, B.D.; Flynn, S.P.; Pruess, T.H.; Green, B.R.; Zhang, L.; Bulaj, G. Developing novel antiepileptic drugs: characterization of NAX 5055, a systemically-active galanin analog, in epilepsy models. *Neurotherapeutics*, **2009**, *6* (2), 372-380.
- [18] Zhang, L.; Robertson, C.R.; Green, B.R.; Pruess, T.H.; White, H.S.; Bulaj, G. Structural requirements for a lipooamino acid in modulating the anticonvulsant activities of systemically active galanin analogues. *J. Med. Chem.*, **2009**, *52* (5), 1310-1316.
- [19] Irwin, N.; O'Harte, F.P.M.; Gault, V.A.; Green, B.D.; Greer, B.; Harriott, P.; Bailey, C.J.; Flatt, P.R. GIP(Lys¹⁶PAL) and GIP(Lys²⁷PAL): novel long-acting acylated analogues of glucose-dependent insulinotropic polypeptide with improved antidiabetic potential. *J. Med. Chem.*, **2006**, *49* (3), 1047-1054.
- [20] Madsen, K.; Knudsen, L.B.; Agersoe, H.; Nielsen, P.F.; Thøgersen, H.; Wilken, M.; Johansen, N.L. Structure-activity and protraction relationship of long-acting Glucagon-like Peptide-1 derivatives: importance of fatty acid length, polarity, and bulkiness. *J. Med. Chem.*, **2007**, *50* (24), 6126-6132.
- [21] Bellmann-Sickert, K.; Elling, C.E.; Madsen, A.N.; Little, P.B.; Lundgren, K.; Gerlach, L.-O.; Bergmann, R.; Holst, B.; Schwartz, T.W.; Beck-Sickinger, A.G. Long-acting lipidated analogue of human pancreatic polypeptide is slowly released into circulation. *J. Med. Chem.*, **2011**, *54* (8), 2658-2667.
- [22] Johannessen, L.; Remsberg, J.; Gaponenko, V.; Adams, K.M.; Barchi, J.J.; Tarasov, S.G.; Jiang, S.; Tarasova, N.I. Peptide structure stabilization by membrane anchoring and its general applicability to the development of potent cell-permeable inhibitors. *ChemBioChem*, **2011**, *12* (6), 914-921.
- [23] Asada, H.; Douen, T.; Mizokoshi, Y.; Fujita, T.; Murakami, M.; Yamamoto, A.; Muranishi, S. Stability of acyl derivatives of Insulin in the small intestine: relative importance of Insulin association characteristics in aqueous solution. *Pharm. Res.*, **1994**, *11* (8), 1115-1120.
- [24] Asada, H.; Douen, T.; Waki, M.; Adachi, S.; Fujita, T.; Yamamoto, A.; Muranishi, S. Absorption characteristics of chemically modified-insulin derivatives with various fatty acids in the small and large intestine. *J. Pharm. Sci.*, **1995**, *84* (6), 682-687.
- [25] Hashimoto, M.; Takada, K.; Kiso, Y.; Muranishi, S. Synthesis of palmitoyl derivatives of insulin and their biological activities. *Pharm. Res.*, **1989**, *6* (2), 171-176.
- [26] Hashizume, M.; Douen, T.; Murakami, M.; Yamamoto, A.; Takada, K.; Muranishi, S. Improvement of large intestinal absorption of insulin by chemical modification with palmitic acid in rats. *J. Pharm. Pharmacol.*, **1992**, *44* (7), 555-559.
- [27] Kurtzhals, P.; Havelund, S.; Jonassen, I.; Kiehr, B.; Larsen, U.D.; Ribøl, U.; Markussen, J. Albumin binding of insulins acylated with fatty acids: characterization of the ligand-protein interaction and correlation between binding affinity and timing of the insulin effect *in vivo*. *Biochem. J.*, **1995**, *312* (3), 725-731.
- [28] Muranishi, S.; Murakami, M.; Hashizume, M.; Yamada, K.; Tajima, S.; Kiso, Y. Trials of lipid modification of peptide hormones for intestinal delivery. *J. Controlled Release*, **1992**, *19* (1-3), 179-188.
- [29] Myers, S.R.; Yakubu-Madus, F.E.; Johnson, W.T.; Baker, J.E.; Cusick, T.S.; Williams, V.K.; Tinsley, F.C.; Kriaucinas, A.; Manetta, J.; Chen, V.J. Acylation of human insulin with palmitic acid extends the time action of human insulin in diabetic dogs. *Diabetes*, **1997**, *46* (4), 637-642.
- [30] Toth, I.; Flinn, N.; Hillery, A.; Gibbons, W.A.; Artursson, P. Lipidic conjugates of luteinizing hormone releasing hormone (LHRH)⁺ and thyrotropin releasing hormone (TRH)⁺ that release and protect the native hormones in homogenates of human intestinal epithelial (Caco-2) cells. *Int. J. Pharm.*, **1994**, *105* (3), 241-247.
- [31] Green, B.R.; Smith, M.; White, K.L.; White, H.S.; Bulaj, G. Analgesic neuropeptide W suppresses seizures in the brain revealed by rational repositioning and peptide engineering. *ACS Chem. Neurosci.*, **2011**, *2* (1), 55-56.
- [32] Dasgupta, P.; Mukherjee, R. Lipophilization of somatostatin analog RC-160 with long chain fatty acid improves its antiproliferative and antiangiogenic activity *in vitro*. *Br. J. Pharmacol.*, **2000**, *129* (1), 101-109.
- [33] Dasgupta, P.; Singh, A.T.; Mukherjee, R. Lipophilization of somatostatin analog RC-160 improves its bioactivity and stability. *Pharm. Res.*, **1999**, *16* (7), 1047-1053.
- [34] Timofeeva, O.A.; Gaponenko, V.; Lockett, S.J.; Tarasov, S.G.; Jiang, S.; Michejda, C.J.; Perantoni, A.O.; Tarasova, N.I. Rationally designed inhibitors identify STAT3 N-domain as a promising anticancer drug target. *ACS Chem. Biol.*, **2007**, *2* (12), 799-809.
- [35] Fujita, T.; Kawahara, I.; Quan, Y.-s.; Hattori, K.; Takenaka, K.; Muranishi, S.; Yamamoto, A. Permeability characteristics of Tetragastrins across intestinal membranes using the Caco-2 monolayer system: comparison between acylation and application of protease inhibitors. *Pharm. Res.*, **1998**, *15* (9), 1387-1392.
- [36] Setoh, K.; Murakami, M.; Araki, N.; Fujita, T.; Yamamoto, A.; Muranishi, S. Improvement of transdermal delivery of tetragastrin by lipophilic modification with fatty acids. *J. Pharm. Pharmacol.*, **1995**, *47* (10), 808-811.
- [37] Yodoya, E.; Uemura, K.; Tenma, T.; Fujita, T.; Murakami, M.; Yamamoto, A.; Muranishi, S. Enhanced permeability of tetragastrin across the rat intestinal membrane and its reduced degradation by acylation with various fatty acids. *J. Pharmacol. Exp. Ther.*, **1994**, *271* (3), 1509-1513.
- [38] Muranishi, S.; Sakai, A.; Yamada, K.; Murakami, M.; Takada, K.; Kiso, Y. Lipophilic peptides: synthesis of Lauroyl Thyrotropin-Releasing Hormone and its biological activity. *Pharm. Res.*, **1991**, *8* (5), 649-652.
- [39] Tanaka, K.; Fujita, T.; Yamamoto, Y.; Murakami, M.; Yamamoto, A.; Muranishi, S. Enhancement of intestinal transport of thyrotropin-releasing hormone *via* a carrier-mediated transport system by chemical modification with lauric acid. *Biochim. Biophys. Acta - Biomembranes*, **1996**, *1283* (1), 119-126.
- [40] Maletinská, L.; Neugebauer, W.; Parê, M.-C.; Jacqueline, P.; Pham, D.; Escher, E. Lipid masking and reactivation of Angiotensin analogues. *Helv. Chim. Acta*, **1996**, *79* (7), 2023 - 2034.
- [41] Maletinská, L.; Neugebauer, W.; Jacqueline, P.; Lefebvre, M.; Escher, E. Angiotensin analogues palmitoylated in positions 1 and 4. *J. Med. Chem.*, **1997**, *40* (20), 3271-3279.
- [42] Ekrami, H.M.; Kennedy, A.R.; Shen, W.-C. Water-soluble fatty acid derivatives as acylating agents for reversible lipidization of polypeptides. *FEBS Lett.*, **1995**, *371* (3), 283-286.
- [43] Honeycutt, L.; Wang, J.; Ekrami, H.; Shen, W.C. Comparison of pharmacokinetic parameters of a polypeptide, the Bowman-Birk protease inhibitor (BBI), and its palmitic acid conjugate. *Pharm. Res.*, **1996**, *13* (9), 1373-1377.
- [44] Wang, J.; Shen, W.C. Gastric retention and stability of lipidized Bowman-Birk protease inhibitor in mice. *Int. J. Pharm.*, **2000**, *204* (1-2), 111-116.
- [45] Wang, J.; Shen, D.; Shen, W.-C. Preparation, purification, and characterization of a reversibly lipidized desmopressin with potentiated anti-diuretic activity. *Pharm. Res.*, **1999**, *16* (11), 1674-1679.
- [46] Bednarek, M.A.; Feighner, S.D.; Pong, S.S.; McKee, K.K.; Hreniuk, D.L.; Silva, M.V.; Warren, V.A.; Howard, A.D.; Van Der Ploeg, L.H.; Heck, J.V. Structure-function studies on the new growth hormone-releasing peptide, ghrelin: minimal sequence of ghrelin necessary for activation of growth hormone secretagogue receptor 1a. *J. Med. Chem.*, **2000**, *43* (23), 4370-4376.
- [47] Chollet, C.; Meyer, K.; Beck-Sickinger, A.G. Ghrelin-a novel generation of anti-obesity drug: design, pharmacomodulation and biological activity of ghrelin analogues. *J. Pept. Sci.*, **2009**, *15* (11), 711-730.
- [48] Matsumoto, M.; Hosoda, H.; Kitajima, Y.; Morozumi, N.; Minamitake, Y.; Tanaka, S.; Matsuo, H.; Kojima, M.; Hayashi, Y.; Kangawa, K. Structure-activity relationship of ghrelin: pharmacological study of ghrelin peptides. *Biochem. Biophys. Res. Commun.*, **2001**, *287* (1), 142-146.

- [49] Cheng, W.; Lim, L.Y. Comparison of reversible and nonreversible aqueous-soluble lipidized conjugates of Salmon Calcitonin. *Mol. Pharmaceutics*, **2008**, *5* (4), 610-621.
- [50] Cheng, W.; Satyanarayanajois, S.; Lim, L.Y. Aqueous-soluble, non-reversible lipid conjugate of salmon calcitonin: synthesis, characterization and *in vivo* activity. *Pharm. Res.*, **2007**, *24* (1), 99-110.
- [51] Wang, J.; Chow, D.; Heiati, H.; Shen, W.C. Reversible lipidization for the oral delivery of salmon calcitonin. *J. Controlled Release*, **2003**, *88* (3), 369-380.
- [52] Lundin, S.; Møss, J.; Bundgaard, H.; Artursson, P. Absorption of thyrotropin-releasing hormone (TRH) and a TRH prodrug in a human intestinal cell line (Caco-2). *Int. J. Pharm.*, **1991**, *76* (3), R1-R4.
- [53] Yuan, L.; Wang, J.; Shen, W.C. Reversible lipidization prolongs the pharmacological effect, plasma duration, and liver retention of octreotide. *Pharm. Res.*, **2005**, *22* (2), 220-227.
- [54] Green, B.R.; Catlin, P.; Zhang, M.M.; Fiedler, B.; Bayudan, W.; Morrison, A.; Norton, R.S.; Smith, B.J.; Yoshikami, D.; Olivera, B.M.; Bulaj, G. Conotoxins containing nonnatural backbone spacers: cladistic-based design, chemical synthesis, and improved analgesic activity. *Chem. Biol.*, **2007**, *14* (4), 399-407.
- [55] Lee, H.K.; Smith, M.D.; Smith, B.J.; Grussendorf, J.; Xu, L.; Gillies, R.J.; White, H.S.; Bulaj, G. Anticonvulsant Met-enkephalin analogues containing backbone spacers reveal alternative non-opioid signaling in the brain. *ACS Chem. Biol.*, **2009**, *4* (8), 659-671.
- [56] Itoh, F.; Nishikimi, Y.; Hasuoka, A.; Yoshioka, Y.; Yukishige, K.; Tanida, S.; Aono, T. Synthesis and structure-activity relationships of TAN-1511 analogues as potent hematopoietic agents. *Chem. Pharm. Bull. (Tokyo)*, **1998**, *46* (2), 255-273.
- [57] Oh, H.S.; Kim, S.; Cho, H.; Lee, K.H. Development of novel lipid-peptide hybrid compounds with antibacterial activity from natural cationic antibacterial peptides. *Bioorg. Med. Chem. Lett.*, **2004**, *14* (5), 1109-1113.
- [58] Zarandi, M.; Varga, J.L.; Schally, A.V.; Horvath, J.E.; Toller, G.L.; Kovacs, M.; Letsch, M.; Groot, K.; Armatis, P.; Halmos, G. Lipopeptide antagonists of growth hormone-releasing hormone with improved antitumor activities. *Proc. Natl. Acad. Sci. U. S. A.*, **2006**, *103* (12), 4610-4615.
- [59] Amon, M.A.; Ali, M.; Bender, V.; Chan, Y.-N.; Toth, I.; Manolios, N. Lipidation and glycosylation of a T cell antigen receptor (TCR) transmembrane hydrophobic peptide dramatically enhances *in vitro* and *in vivo* function. *Biochim. Biophys. Acta*, **2006**, *1763* (8), 879-888.
- [60] Chan, E.; Amon, M.; Marano, R.J.; Wimmer, N.; Kearns, P.S.; Manolios, N.; Rakoczy, P.E.; Toth, I. Novel cationic lipophilic peptides for oligodeoxynucleotide delivery. *Bioorg. Med. Chem.*, **2007**, *15* (12), 4091-4097.
- [61] Pignatello, R.; Pecora, T.M. Conjugation of thymopentin (TP5) with lipophilic amino acid residues increases the hydrolytic stability and preserves the biological activity. *Pharmazie*, **2007**, *62* (9), 663-667.
- [62] Dulhunty, A.F.; Cengia, L.; Young, J.; Pace, S.M.; Harvey, P.J.; Lamb, G.D.; Chan, Y.N.; Wimmer, N.; Toth, I.; Casarotto, M.G. Functional implications of modifying RyR-activating peptides for membrane permeability. *Br. J. Pharmacol.*, **2005**, *144* (6), 743-754.
- [63] Hayes, P.Y.; Ross, B.P.; Thomas, B.G.; Toth, I. Polycationic lipophilic-core dendrons as penetration enhancers for the oral administration of low molecular weight heparin. *Bioorg. Med. Chem.*, **2006**, *14* (1), 143-152.
- [64] Ross, B.P.; Toth, I. Gastrointestinal absorption of heparin by lipidization or coadministration with penetration enhancers. *Curr. Drug Deliv.*, **2005**, *2* (3), 277-287.
- [65] Lee, K.C.; Carlson, P.A.; Goldstein, A.S.; Yager, P.; Gelb, M.H. Protection of a decapeptide from proteolytic cleavage by lipidation and self-assembly into high-axial-ratio microstructures: a kinetic and structural study. *Langmuir*, **1999**, *15* (17), 5500-5508.
- [66] Ingallinella, P.; Bianchi, E.; Ladwa, N.A.; Wang, Y.-J.; Hrin, R.; Veneziano, M.; Bonelli, F.; Ketas, T.J.; Moore, J.P.; Miller, M.D.; Pessi, A. Addition of a cholesterol group to an HIV-1 peptide fusion inhibitor dramatically increases its antiviral potency. *Proc. Natl. Acad. Sci. U. S. A.*, **2009**, *106* (14), 5801-5806.
- [67] Lee, S.; Kim, K.; Kumar, T.S.; Lee, J.; Kim, S.K.; Lee, D.Y.; Lee, Y.-k.; Byun, Y. Synthesis and biological properties of Insulin-deoxycholic acid chemical conjugates. *Bioconjugate Chem.*, **2005**, *16* (3), 615-620.
- [68] Lee, S.; Lee, J.; Lee, D.; Kim, S.; Lee, Y.; Byun, Y. A new drug carrier, N'-deoxycholyl-L-lysyl-methylester, for enhancing insulin absorption in the intestine. *Diabetologia*, **2005**, *48* (3), 405-411.
- [69] Blanchfield, J.T.; Dutton, J.L.; Hogg, R.C.; Gallagher, O.P.; Craik, D.J.; Jones, A.; Adams, D.J.; Lewis, R.J.; Alewood, P.F.; Toth, I. Synthesis, structure elucidation, *in vitro* biological activity, toxicity, and Caco-2 cell permeability of lipophilic analogues of alpha-conotoxin MII. *J. Med. Chem.*, **2003**, *46* (7), 1266-1272.
- [70] Pignatello, R.; Noce, C.; Campisi, A.; Acquaviva, R.; Bucolo, C.; Puglisi, G.; Toth, I. Evaluation of cell tolerability of a series of lipophilic acids using biological membranes and a biomembrane model. *Curr. Drug Deliv.*, **2007**, *4* (2), 109-121.
- [71] Ross, B.P.; Braddy, A.C.; McGeary, R.P.; Blanchfield, J.T.; Prokai, L.; Toth, I. Micellar aggregation and membrane partitioning of bile salts, fatty acids, sodium dodecyl sulfate, and sugar-conjugated fatty acids: correlation with hemolytic potency and implications for drug delivery. *Mol. Pharm.*, **2004**, *1* (3), 233-245.
- [72] Ross, B.P.; DeCruz, S.E.; Lynch, T.B.; Davis-Goff, K.; Toth, I. Design, synthesis, and evaluation of a liposaccharide drug delivery agent: application to the gastrointestinal absorption of gentamicin. *J. Med. Chem.*, **2004**, *47* (5), 1251-1258.
- [73] Toth, I. A novel chemical approach to drug delivery: lipidic amino acid conjugates. *J. Drug Target.*, **1994**, *2* (3), 217-239.
- [74] Görmer, K.; Waldmann, H.; Brunsfeld, L. In: *Comprehensive Natural Products II*, Mander, L., Lui, H.-W., Ed.; Elsevier: Oxford, **2010**; pp. 531-585.
- [75] Magee, T.; Seabra, M.C. Fatty acylation and prenylation of proteins: what's hot in fat. *Curr. Opin. Cell Biol.*, **2005**, *17* (2), 190-196.
- [76] Resh, M.D. Regulation of cellular signalling by fatty acid acylation and prenylation of signal transduction proteins. *Cell. Signal.*, **1996**, *8* (6), 403-412.
- [77] BenMohamed, L.; Wechsler, S.L.; Nesburn, A.B. Lipopeptide vaccines--yesterday, today, and tomorrow. *Lancet Infect Dis.*, **2002**, *2* (7), 425-431.
- [78] Raulin, J. Development in lipid drugs. *Mini. Rev. Med. Chem.*, **2005**, *5* (5), 489-498.
- [79] Zhong, W.; Skwarczynski, M.; Toth, I. Lipid Core Peptide System for gene, drug, and vaccine delivery. *Aust. J. Chem.*, **2009**, *62* (9), 956-967.
- [80] Wong, A.; Toth, I. Lipid, sugar and liposaccharide based delivery systems. *Curr. Med. Chem.*, **2001**, *8* (9), 1123-1136.
- [81] Yagi, N.; Yano, Y.; Hatanaka, K.; Yokoyama, Y.; Okuno, H. Synthesis and evaluation of a novel lipid-peptide conjugate for functionalized liposome. *Bioorg. Med. Chem. Lett.*, **2007**, *17* (9), 2590-2593.
- [82] Jerala, R. Synthetic lipopeptides: a novel class of anti-infectives. *Expert Opin. Invest. Drugs*, **2007**, *16* (8), 1159-1169.
- [83] Toniolo, C.; Crisma, M.; Formaggio, F.; Peggion, C.; Epand, R.F.; Epand, R.M. Lipopeptidols, a novel family of membrane active, antimicrobial peptides. *Cell. Mol. Life Sci.*, **2001**, *58* (9), 1179-1188.
- [84] Cavalli, S.; Kros, A. Scope and applications of amphiphilic alkyl- and lipopeptides. *Adv. Mater.*, **2008**, *20* (3), 627-631.
- [85] Charron, G.; Wilson, J.; Hang, H.C. Chemical tools for understanding protein lipidation in eukaryotes. *Curr. Opin. Chem. Biol.*, **2009**, *13* (4), 382-391.
- [86] Peters, C.; Wagner, M.; Volkert, M.; Waldmann, H. Bridging the gap between cell biology and organic chemistry: chemical synthesis and biological application of lipidated peptides and proteins. *Naturwissenschaften*, **2002**, *89* (9), 381-390.
- [87] Silvius, J.R. Lipidated peptides as tools for understanding the membrane interactions of lipid-modified proteins. *Curr. Top. Membr.*, **2002**, *52*, 371-395.
- [88] Epand, R.M. Biophysical studies of lipopeptide-membrane interactions. *Biopolymers*, **1997**, *43* (1), 15-24.
- [89] Yamamoto, A.; Morishita, Y.; Sugishita, S.; Hayami, T.; Okada, N.; Fujita, T.; Muranishi, S. Enhanced permeability of phenylalanyl-glycine (Phe-Gly) across the intestinal membranes by chemical modification with various fatty acids. *Drug Metab. Pharmacokinet.*, **2003**, *18* (1), 23-32.
- [90] Yamamoto, A.; Setoh, K.; Murakami, M.; Shironoshita, M.; Kobayashi, T.; Fujimoto, K.; Okada, N.; Fujita, T.; Muranishi, S. Enhanced transdermal delivery of phenylalanyl-glycine by chemical modification with various fatty acids. *Int. J. Pharm.*, **2003**, *250* (1), 119-128.
- [91] Macquaire, F.; Baleux, F.; Giacobbi, E.; Huynh-Dinh, T.; Neumann, J.M.; Sanson, A. Peptide secondary structure induced by a micellar phospholipid interface: proton NMR conformational study of a lipopeptide. *Biochemistry*, **1992**, *31* (9), 2576-2582.
- [92] Pakalns, T.; Haverstick, K.L.; Fields, G.B.; McCarthy, J.B.; Mooradian, D.L.; Tirrell, M. Cellular recognition of synthetic peptide amphiphiles in self-assembled monolayer films. *Biomaterials*, **1999**, *20* (23-24), 2265-2279.
- [93] Qahwash, I.M.; Boire, A.; Lanning, J.; Krausz, T.; Pytel, P.; Meredith, S.C. Site-specific effects of peptide lipidation on beta-amyloid aggregation and cytotoxicity. *J. Biol. Chem.*, **2007**, *282* (51), 36987-36997.
- [94] Giragossian, C.; Nardi, E.; Savary, C.; Pellegrini, M.; Meini, S.; Maggi, C.A.; Papini, A.M.; Mierke, D.F. Structural characterization of lipopeptide agonists for the bradykinin B2 receptor. *Biopolymers*, **2001**, *58* (5), 511-520.
- [95] Forns, P.; Lauer-Fields, J.L.; Gao, S.; Fields, G.B. Induction of protein-like molecular architecture by monoalkyl hydrocarbon chains. *Biopolymers*, **2000**, *54* (7), 531-546.
- [96] Malkar, N.B.; Lauer-Fields, J.L.; Juska, D.; Fields, G.B. Characterization of peptide-amphiphiles possessing cellular activation sequences. *Biomacromolecules*, **2003**, *4* (3), 518-528.
- [97] Takahashi, Y.; Ueno, A.; Mihara, H. Optimization of hydrophobic domains in peptides that undergo transformation from alpha-helix to beta-fibril. *Bioorg. Med. Chem.*, **1999**, *7* (1), 177-185.
- [98] Etzerodt, T.; Henriksen, J.R.; Rasmussen, P.; Clausen, M.H.; Andresen, T.L. Selective acylation enhances membrane charge sensitivity of the antimicrobial peptide Mastoparan-X. *Biophys. J.*, **2011**, *100* (2), 399-409.
- [99] Japelj, B.; Zorko, M.; Majerle, A.; Pristovsek, P.; Sanchez-Gomez, S.; Martinez de Tejada, G.; Moriyon, I.; Blondelle, S.E.; Brandenburg, K.; Andra, J.; Lohner, K.; Jerala, R. The acyl group as the central element of the structural organization of antimicrobial lipopeptide. *J. Am. Chem. Soc.*, **2007**, *129* (5), 1022-1023.
- [100] Yu, Y.-C.; Berndt, P.; Tirrell, M.; Fields, G.B. Self-assembling amphiphiles for construction of protein molecular architecture. *J. Am. Chem. Soc.*, **1996**, *118* (50), 12515-12520.

- [101] Cui, H.; Webber, M.J.; Stupp, S.I. Self-assembly of peptide amphiphiles: From molecules to nanostructures to biomaterials. *Peptide Sci.*, **2010**, *94* (1), 1-18.
- [102] Gazit, E. Self-assembled peptide nanostructures: the design of molecular building blocks and their technological utilization. *Chem. Soc. Rev.*, **2007**, *36*, 1263-1269.
- [103] Kokkoli, E.; Mardilovich, A.; Wedekind, A.; Rexeis, E.L.; Garg, A.; Craig, J.A. Self-assembly and applications of biomimetic and bioactive peptide-amphiphiles. *Soft Matter*, **2006**, *2* (12), 1015-1024.
- [104] Scanlon, S.; Aggeli, A. Self-assembling peptide nanotubes. *Nano Today*, **2008**, *3* (3-4), 22-30.
- [105] Hartgerink, J.D.; Beniash, E.; Stupp, S.I. Peptide-amphiphile nanofibers: a versatile scaffold for the preparation of self-assembling materials. *Proc. Natl. Acad. Sci. U. S. A.*, **2002**, *99* (8), 5133-5138.
- [106] Weroniski, K.J.; Díez-Pérez, I.; Busquets, M.A.; López-Iglesias, C.; Girona, V.; Prat, J. Interaction of lipidated GBV-C/HGV NS3 (513-522) and (505-514) peptides with phospholipids monolayer. An AFM study. *Colloids Surf., B Biointerfaces*, **2010**, *75* (1), 25-33.
- [107] Reig, F.; Haro, I.; Polo, D.; Egea, M.A.; Alsina, M.A. Interfacial interactions of hydrophobic peptides with lipid bilayers. *J. Colloid Interface Sci.*, **2002**, *246* (1), 60-69.
- [108] Brunsfeld, L.; Waldmann, H.; Huster, D. Membrane binding of lipidated Ras peptides and proteins - the structural point of view. *Biochim. Biophys. Acta - Biomembranes*, **2009**, *1788* (1), 273-288.
- [109] Eisenberg, S.; Henis, Y.I. Interactions of Ras proteins with the plasma membrane and their roles in signaling. *Cell. Signal.*, **2008**, *20* (1), 31-39.
- [110] Almiñana, N.; Alsina, M.A.; Reig, F. New GHK hydrophobic derivatives: Interaction with phospholipid bilayers. *Colloids Surf., B Biointerfaces*, **2007**, *57* (2), 243-249.
- [111] Peitzsch, R.M.; McLaughlin, S. Binding of acylated peptides and fatty acids to phospholipid vesicles: Pertinence to myristoylated proteins. *Biochemistry*, **1993**, *32* (39), 10436-10443.
- [112] Gatto, E.; Mazzuca, C.; Stella, L.; Venanzi, M.; Toniolo, C.; Pispisa, B. Effect of peptide lipidation on membrane perturbing activity: a comparative study on two trichogin analogues. *J. Phys. Chem. B*, **2006**, *110* (45), 22813-22818.
- [113] Koda, Y.; Del Borgo, M.; Wessling, S.T.; Lazarus, L.H.; Okada, Y.; Toth, I.; Blanchfield, J.T. Synthesis and *in vitro* evaluation of a library of modified endomorphin 1 peptides. *Bioorg. Med. Chem.*, **2008**, *16* (11), 6286-6296.
- [114] Tenma, T.; Yodoya, E.; Tashima, S.; Fujita, T.; Murakami, M.; Yamamoto, A.; Muranishi, S. Development of new lipophilic derivatives of tetragastrin: physicochemical characteristics and intestinal absorption of acyl-tetragastrin derivatives in rats. *Pharm. Res.*, **1993**, *10* (10), 1488-1492.
- [115] Yamamoto, A. Improvement of intestinal absorption of peptide and protein drugs by chemical modification with fatty acids. *Nippon Rinsho*, **1998**, *56* (3), 601-607.
- [116] Nelson, A.R.; Borland, L.; Allbritton, N.L.; Sims, C.E. Myristoyl-based transport of peptides into living cells. *Biochemistry*, **2007**, *46* (51), 14771-14781.
- [117] Heinemann, L.; Sinha, K.; Weyer, C.; Loftager, M.; Hirschberger, S.; Heise, T. Time-action profile of the soluble, fatty acid acylated, long-acting insulin analogue NN304. *Diabet Med.*, **1999**, *16* (4), 332-338.
- [118] Choi, J.-K.; Ho, J.; Curry, S.; Qin, D.; Bittman, R.; Hamilton, J.A. Interactions of very long-chain saturated fatty acids with serum albumin. *J. Lipid Res.*, **2002**, *43* (7), 1000-1010.
- [119] Chuang, V.T.G.; Otagiri, M. Stereoselective binding of human serum albumin. *Chirality*, **2006**, *18* (3), 159-166.
- [120] Hamilton, J.A. Fatty acid interactions with proteins: what X-ray crystal and NMR solution structures tell us. *Prog. Lipid Res.*, **2004**, *43* (3), 177-199.
- [121] Spector, A.A. Fatty acid binding to plasma albumin. *J. Lipid Res.*, **1975**, *16* (3), 165-179.
- [122] Bhattacharya, A.A.; Grüne, T.; Curry, S. Crystallographic analysis reveals common modes of binding of medium and long-chain fatty acids to human serum albumin. *J. Mol. Biol.*, **2000**, *303* (5), 721-732.
- [123] Fujiwara, S.; Amisaki, T. Identification of high affinity fatty acid binding sites on human serum albumin by MM-PBSA method. *Biophys. J.*, **2008**, *94* (1), 95-103.
- [124] Petitpas, I.; Grüne, T.; Bhattacharya, A.A.; Curry, S. Crystal structures of human serum albumin complexed with monounsaturated and polyunsaturated fatty acids. *J. Mol. Biol.*, **2001**, *314* (5), 955-960.
- [125] Kurtzhals, P.; Havelund, S.; Jonassen, I.; Markussen, J. Effect of fatty acids and selected drugs on the albumin binding of a long-acting, acylated insulin analogue. *J. Pharm. Sci.*, **1997**, *86* (12), 1365-1368.
- [126] Bellmann-Sickert, K.; Beck-Sickinger, A.G. Palmitoylated SDF1 α shows increased resistance against proteolytic degradation in liver homogenates. *ChemMedChem*, **2011**, *6* (1), 193-200.
- [127] Rivett, D.E.; Hewish, D.; Kirkpatrick, A.; Werkmeister, J. Inhibition of membrane-active peptides by fatty acid-peptide hybrids. *J. Protein Chem.*, **1999**, *18* (3), 291-295.
- [128] Yang, P.; Abe, S.; Sato, Y.; Yamashita, T.; Matsuda, F.; Hamayasu, T.; Imai, K.; Suzuki, K. A palmitoyl conjugate of insect pentapeptide Yamamarin arrests cell proliferation and respiration. *J. Insect Biotech. Seric.*, **2007**, *76*, 263-269.
- [129] Gault, V.A.; Hunter, K.; Irwin, N.; Greer, B.; Green, B.D.; Harriott, P.; O'Harte, F.P.; Flatt, P.R. Characterisation and glucoregulatory actions of a novel acylated form of the (Pro3)GIP receptor antagonist in type 2 diabetes. *Biol. Chem.*, **2007**, *388* (2), 173-179.
- [130] O'Harte, F.P.; Hunter, K.; Gault, V.A.; Irwin, N.; Green, B.D.; Greer, B.; Harriott, P.; Bailey, C.J.; Flatt, P.R. Antagonistic effects of two novel GIP analogs, (Hyp3)GIP and (Hyp3)GIPLys16PAL, on the biological actions of GIP and longer-term effects in diabetic ob/ob mice. *Am. J. Physiol-Endo. M.*, **2007**, *292* (6), E1674-1682.
- [131] Gutniak, M.K.; Linde, B.; Holst, J.J.; Efendic, S. Subcutaneous injection of the incretin hormone glucagon-like peptide 1 abolishes postprandial glycemia in NIDDM. *Diabetes Care*, **1994**, *17* (9), 1039-1044.
- [132] Kieffer, T.J.; McIntosh, C.H.; Pederson, R.A. Degradation of glucose-dependent insulinotropic polypeptide and truncated glucagon-like peptide 1 *in vitro* and *in vivo* by dipeptidyl peptidase IV. *Endocrinology*, **1995**, *136* (8), 3585-3596.
- [133] Vilsboll, T.; Agerso, H.; Krarup, T.; Holst, J.J. Similar elimination rates of glucagon-like peptide-1 in obese type 2 diabetic patients and healthy subjects. *J. Clin. Endocrinol Metab.*, **2003**, *88* (1), 220-224.
- [134] Agersø, H.; Jensen, L.B.; Elbrønd, B.; Rolan, P.; Zdravkovic, M. The pharmacokinetics, pharmacodynamics, safety and tolerability of NN2211, a new long-acting GLP-1 derivative, in healthy men. *Diabetologia*, **2002**, *45* (2), 195-202.
- [135] Knudsen, L.B.; Nielsen, P.F.; Huusfeldt, P.O.; Johansen, N.L.; Madsen, K.; Pedersen, F.Z.; Thøgersen, H.; Wilken, M.; Agerso, H. Potent derivatives of glucagon-like peptide-1 with pharmacokinetic properties suitable for once daily administration. *J. Med. Chem.*, **2000**, *43* (9), 1664-1669.
- [136] Ribel, U.; Larsen, M.O.; Rolin, B.; Carr, R.D.; Wilken, M.; Sturis, J.; Westergaard, L.; Deacon, C.F.; Knudsen, L.B. NN2211: a long-acting glucagon-like peptide-1 derivative with anti-diabetic effects in glucose-intolerant pigs. *Eur. J. Pharmacol.*, **2002**, *451* (2), 217-225.
- [137] Chicharro, C.; Granata, C.; Lozano, R.; Andreu, D.; Rivas, L. N-terminal fatty acid substitution increases the leishmanicidal activity of CA(1-7)M(2-9), a cecropin-melittin hybrid peptide. *Antimicrob. Agents Chemother.*, **2001**, *45* (9), 2441-2449.
- [138] Fujita, T.; Fujita, T.; Morikawa, K.; Tanaka, H.; Iemura, O.; Yamamoto, A.; Muranishi, S. Improvement of intestinal absorption of human calcitonin by chemical modification with fatty acids: Synergistic effects of acylation and absorption enhancers. *Int. J. Pharm.*, **1996**, *134* (1-2), 47-57.
- [139] Robertson, C.R.; Flynn, S.P.; White, H.S.; Bulaj, G. Anticonvulsant neuropeptides as drug leads for neurological diseases. *Nat. Prod. Rep.*, **2011**, *28* (4), 741-762.
- [140] Ogawa, T.; Araki, M.; Miyamae, T.; Okayama, T.; Hagiwara, M.; Sakurada, S.; Morikawa, T. Synthesis and antinociceptive activity of orally active opioid peptides: improvement of oral bioavailability by esterification. *Chem. Pharm. Bull. (Tokyo)*, **2003**, *51* (7), 759-771.
- [141] Heavner, G.A.; Kroon, D.J.; Audhya, T.; Goldstein, G. Biologically active analogs of thympentin with enhanced enzymatic stability. *Peptides*, **1986**, *7* (6), 1015-1019.
- [142] Bergeon, J.A.; Toth, I. Enhancement of oral drug absorption--effect of lipid conjugation on the enzymatic stability and intestinal permeability of L-Glu-L-Trp-NH₂. *Bioorg. Med. Chem.*, **2007**, *15* (22), 7048-7057.
- [143] Blanchfield, J.T.; Lew, R.A.; Smith, A.I.; Toth, I. The stability of lipidic analogues of GnRH in plasma and kidney preparations: the stereoselective release of the parent peptide. *Bioorg. Med. Chem. Lett.*, **2005**, *15* (6), 1609-1612.
- [144] Rist, B.; Wieland, H.A.; Willim, K.D.; Beck-Sickinger, A.G. A rational approach for the development of reduced-size analogues of neuropeptide Y with high affinity to the Y1 receptor. *J. Pept. Sci.*, **1995**, *1* (5), 341-348.
- [145] Heise, T.; Tack, C.J.; Cuddihy, R.; Davidson, J.; Gouet, D.; Liebl, A.; Romero, E.; Mersebach, H.; Dykiel, P.; Jorde, R. A new-generation ultra-long-acting basal insulin with a bolus boost compared with Insulin Glargine in Insulin-naïve people with type 2 diabetes. *Diabetes Care*, **2011**, *34* (3), 669-674.
- [146] Bundgaard, H.; Moss, J. Prodrugs of peptides. 6. Bioreversible derivatives of thyrotropin-releasing hormone (TRH) with increased lipophilicity and resistance to cleavage by the TRH-specific serum enzyme. *Pharm. Res.*, **1990**, *7* (9), 885-892.
- [147] Moss, J.; Buur, A.; Bundgaard, H. Prodrugs of peptides. 8. *In vitro* study of intestinal metabolism and penetration of thyrotropin-releasing hormone (TRH) and its prodrugs. *Int. J. Pharm.*, **1990**, *66* (1-3), 183-191.
- [148] Cini, E.; Lampariello, L.R.; Rodriguez, M.; Taddei, M. Synthesis and application in SPPS of a stable amino acid isostere of palmitoyl cysteine. *Tetrahedron*, **2009**, *65* (4), 844-848.
- [149] Shechter, Y.; Heldman, E.; Sasson, K.; Bachar, T.; Popov, M.; Fridkin, M. Delivery of neuropeptides from the periphery to the brain: studies with Enkephalin. *ACS Chem. Neurosci.*, **2010**, *1* (5), 399-406.
- [150] Blanchfield, J.; Toth, I. Lipid, sugar and liposaccharide based delivery systems 2. *Curr. Med. Chem.*, **2004**, *11* (17), 2375-2382.
- [151] Namane, A.; Gouyette, C.; Fillion, M.P.; Fillion, G.; Huynh Dinh, T. Improved brain delivery of AZT using a glycosyl phosphotriester prodrug. *J. Med. Chem.*, **1992**, *35* (16), 3039-3044.
- [152] Simerska, P.; Abdel-Aal, A.-B.M.; Fujita, Y.; Moyle, P.M.; McGeary, R.P.; Batzloff, M.R.; Olive, C.; Good, M.F.; Toth, I. Development of a liposaccharide-based delivery system and its application to the design of group A streptococcal vaccines. *J. Med. Chem.*, **2008**, *51* (5), 1447-1452.

- [153] Toth, I.; Malkinson, J.P.; Flinn, N.S.; Drouillat, B.; Horvath, A.; Erchegyi, J.; Idei, M.; Venetianer, A.; Artursson, P.; Lazorova, L.; Szende, B.; Keri, G. Novel lipoamino acid- and liposaccharide-based system for peptide delivery: application for oral administration of tumor-selective somatostatin analogues. *J. Med. Chem.*, **1999**, 42 (19), 4010-4013.
- [154] Wu, S.; Campbell, C.; Koda, Y.; Blanchfield, J.T.; Toth, I. Investigation of the route of absorption of lipid and sugar modified leu-enkephalin analogues and their enzymatic stability using the caco-2 cell monolayer system. *Med. Chem.*, **2006**, 2 (2), 203-211.
- [155] Kaiya, H.; Kodama, S.; Ishiguro, K.; Matsuda, K.; Uchiyama, M.; Miyazato, M.; Kangawa, K. Ghrelin-like peptide with fatty acid modification and O-glycosylation in the red stingray, *Dasyatis akajei*. *BMC Biochem.*, **2009**, 10, 30.
- [156] Walensky, L.D.; Kung, A.L.; Escher, I.; Malia, T.J.; Barbuto, S.; Wright, R.D.; Wagner, G.; Verdine, G.L.; Korsmeyer, S.J. Activation of apoptosis *in vivo* by a hydrocarbon-stapled BH3 helix. *Science*, **2004**, 305 (5689), 1466-1470.
- [157] Wolfson, W. Aileron staples peptides. *Chemistry & biology*, **2009**, 16 (9), 910-912.
- [158] Takahashi, Y.; Ueno, A.; Mihara, H. Design of a peptide undergoing α - β structural transition and Amyloid fibrillogenesis by the introduction of a hydrophobic defect. *Chem. Eur. J.*, **1998**, 4 (12), 2475-2484.
- [159] Blanchfield, J.T.; Toth, I. Modification of peptides and other drugs using lipoamino acids and sugars. *Methods Mol. Biol.*, **2005**, 298, 45-61.
- [160] Lee, H.K.; Zhang, L.; Smith, M.D.; White, H.S.; Bulaj, G. Glycosylated neurotensin analogues exhibit sub-picomolar anticonvulsant potency in a pharmacoresistant model of epilepsy. *ChemMedChem*, **2009**, 4 (3), 400-405.
- [161] Zhang, L.; Lee, H.K.; Pruess, T.H.; White, H.S.; Bulaj, G. Synthesis and applications of polyamine amino acid residues: improving the bioactivity of an analgesic neuropeptide, neurotensin. *J. Med. Chem.*, **2009**, 52 (6), 1514-1517.
- [162] Rijkers, D.T.S.; Kruijtz, J.A.W.; Killian, J.A.; Liskamp, R.M.J. A convenient solid phase synthesis of S-palmitoyl transmembrane peptides. *Tetrahedron Lett.*, **2005**, 46 (19), 3341-3345.
- [163] Isidro-Llobet, A.; Alvarez, M.; Albericio, F. Amino acid-protecting groups. *Chem. Rev.*, **2009**, 109 (6), 2455-2504.
- [164] Bycroft, B.W.; Chan, W.C.; Chhabra, S.R.; Hone, N.D. A novel lysine-protecting procedure for continuous flow solid phase synthesis of branched peptides. *J. Chem. Soc., Chem. Commun.*, **1993** (9), 778-779.
- [165] Diaz-Mochon, J.J.; Bialy, L.; Bradley, M. Full orthogonality between Dde and Fmoc: the direct synthesis of PNA-Peptide Conjugates. *Org. Lett.*, **2004**, 6 (7), 1127-1129.
- [166] Kan, T.; Fukuyama, T. New strategies: a highly versatile synthetic method for amines. *Chem. Commun.*, **2004** (4), 353-359.
- [167] Narayan, R.S.; VanNieuwenhze, M.S. Versatile and stereoselective syntheses of orthogonally protected beta-methylcysteine and beta-methylanthionine. *Org. Lett.*, **2005**, 7 (13), 2655-2658.
- [168] Kadereit, D.; Deck, P.; Heinemann, I.; Waldmann, H. Acid-labile protecting groups for the synthesis of lipidated peptides. *Chemistry*, **2001**, 7 (6), 1184-1193.
- [169] Palomo, J.M.; Lumbierres, M.; Waldmann, H. Efficient solid-phase lipopeptide synthesis employing the ellman sulfonamide linker. *Angew. Chem. Int. Ed.*, **2006**, 45 (3), 477-481.
- [170] Carpino, L.A.; Krause, E.; Sferdean, C.D.; Schümann, M.; Fabian, H.; Bienert, M.; Beyermann, M. Synthesis of 'difficult' peptide sequences: application of a depsipeptide technique to the Jung-Redemann 10- and 26-mers and the amyloid peptide A[β](1-42). *Tetrahedron Lett.*, **2004**, 45 (40), 7519-7523.
- [171] Cruz, L.J.; Cuevas, C.; Canedo, L.M.; Giral, E.; Albericio, F. Total solid-phase synthesis of marine cyclodepsipeptide IB-01212. *J. Org. Chem.*, **2006**, 71 (9), 3339-3344.
- [172] Ludolph, B.; Eisele, F.; Waldmann, H. Solid-phase synthesis of lipidated peptides. *J. Am. Chem. Soc.*, **2002**, 124 (21), 5954-5955.
- [173] Ponsati I Obiols, B.; Canas I Poblet, M.; Jodas I Farres, G.; Clemente Rodriguez, J.; Barcadit I Cabado, J. EP 1179537. July 30, 2008.
- [174] Zhang, A.J.; Russell, D.H.; Zhu, J.; Burgess, K. A method for removal of N-BOC protecting groups from substrates on TFA-sensitive resins. *Tetrahedron Lett.*, **1998**, 39 (41), 7439-7442.
- [175] Lumbierres, M.; Palomo, J.M.; Kragol, G.; Waldmann, H. Solid-phase synthesis of palmitoylated and farnesylated lipopeptides employing the fluoride-labile PTMSEL linker. *Tetrahedron Lett.*, **2006**, 47 (16), 2671-2674.
- [176] Yousefi-Salakdeh, E.; Johansson, J.; Stromberg, R. A method for S- and O-palmitoylation of peptides: synthesis of pulmonary surfactant protein-C models. *Biochem. J.*, **1999**, 343 Pt 3, 557-562.
- [177] Bonnet, D.; Bourel, L.; Gras-Masse, H.; Melnyk, O. A novel lipophilic glyoxylic acid derivative for the lipidation of peptides using salt-free hydrazine ligation. *Tetrahedron Lett.*, **2000**, 41 (51), 10003-10007.
- [178] Bourel-Bonnet, L.; Bonnet, D.; Malingue, F.; Gras-Masse, H.; Melnyk, O. Simultaneous lipidation of a characterized peptide mixture by chemoselective ligation. *Bioconjugate Chem.*, **2003**, 14 (2), 494-499.
- [179] Bourel-Bonnet, L.; Gras-Masse, H.; Melnyk, O. A novel family of amphiphilic α -oxo aldehydes for the site-specific modification of peptides by two palmitoyl groups in solution or in liposome suspensions. *Tetrahedron Lett.*, **2001**, 42 (39), 6851-6853.
- [180] Dubs, P.; Bourel-Bonnet, L.; Subra, G.; Blanpain, A.; Melnyk, O.; Pinel, A.M.; Gras-Masse, H.; Martinez, J. Parallel synthesis of a lipopeptide library by hydrazone-based chemical ligation. *J. Comb. Chem.*, **2007**, 9 (6), 973-981.