



US011840730B1

(12) **United States Patent**
Porreca et al.

(10) **Patent No.: US 11,840,730 B1**
(45) **Date of Patent: Dec. 12, 2023**

(54) **METHODS AND COMPOSITIONS FOR EVALUATING GENETIC MARKERS**

(71) Applicant: **Molecular Loop Biosciences, Inc.**,
Cambridge, MA (US)

(72) Inventors: **Gregory Porreca**, Cambridge, MA
(US); **Uri Laserson**, Boston, MA (US)

(73) Assignee: **MOLECULAR LOOP BIOSCIENCES, INC.**, Cambridge,
MA (US)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 516 days.

(21) Appl. No.: **16/952,764**

(22) Filed: **Nov. 19, 2020**

Related U.S. Application Data

(63) Continuation of application No. 15/231,687, filed on
Aug. 8, 2016, now abandoned, which is a
continuation of application No. 13/266,862, filed as
application No. PCT/US2010/001293 on Apr. 30,
2010.

(60) Provisional application No. 61/182,089, filed on May
28, 2009, provisional application No. 61/179,358,
filed on May 18, 2009, provisional application No.
61/178,923, filed on May 15, 2009, provisional
application No. 61/174,470, filed on Apr. 30, 2009.

(51) **Int. Cl.**
C12Q 1/6874 (2018.01)
C12Q 1/6827 (2018.01)
C12Q 1/6883 (2018.01)

(52) **U.S. Cl.**
CPC **C12Q 1/6874** (2013.01); **C12Q 1/6827**
(2013.01); **C12Q 1/6883** (2013.01)

(58) **Field of Classification Search**
CPC ... C12Q 1/6874; C12Q 1/6827; C12Q 1/6883
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

4,683,195 A 7/1987 Mullis et al.
4,683,202 A 7/1987 Mullis
4,988,617 A 1/1991 Landegren et al.
5,060,980 A 10/1991 Johnson et al.
5,210,015 A 5/1993 Gelfand et al.
5,234,809 A 8/1993 Boom et al.
5,242,794 A 9/1993 Whiteley et al.
5,348,853 A 9/1994 Wang et al.
5,434,049 A 7/1995 Okano et al.
5,459,307 A 10/1995 Klotz, Jr.
5,486,686 A 1/1996 Zdybel, Jr. et al.
5,491,224 A 2/1996 Bittner et al.
5,494,810 A 2/1996 Barany et al.
5,567,583 A 10/1996 Wang et al.
5,583,024 A 12/1996 McElroy et al.
5,604,097 A 2/1997 Brenner
5,636,400 A 6/1997 Young
5,674,713 A 10/1997 McElroy et al.

5,695,934 A 12/1997 Brenner
5,700,673 A 12/1997 McElroy et al.
5,701,256 A 12/1997 Marr et al.
5,830,064 A 11/1998 Bradish et al.
5,846,719 A 12/1998 Brenner et al.
5,863,722 A 1/1999 Brenner
5,866,337 A 2/1999 Schon
5,869,252 A 2/1999 Bouma et al.
5,869,717 A 2/1999 Frame et al.
5,871,921 A 2/1999 Landegren et al.
5,888,788 A 3/1999 De Miniac
5,942,391 A 8/1999 Zhang et al.
5,971,921 A 10/1999 Timbel
5,993,611 A 11/1999 Moroney et al.
5,994,056 A 11/1999 Higuchi
6,020,127 A 2/2000 MacKenzie et al.
6,033,854 A 3/2000 Kurnit et al.
6,033,872 A 3/2000 Bergsma et al.
6,100,099 A 8/2000 Gordon et al.
6,138,077 A 10/2000 Brenner
6,150,516 A 11/2000 Brenner et al.
6,171,785 B1 1/2001 Higuchi
6,172,214 B1 1/2001 Brenner
6,172,218 B1 1/2001 Brenner
6,197,508 B1 3/2001 Stanley
6,197,574 B1 3/2001 Miyamoto et al.
6,210,891 B1 4/2001 Nyren et al.
6,223,128 B1 4/2001 Allex et al.
6,235,472 B1 5/2001 Landegren et al.
6,235,475 B1 5/2001 Brenner et al.
6,235,501 B1 5/2001 Gautsch et al.
6,235,502 B1 5/2001 Weissman et al.
6,258,568 B1 7/2001 Nyren
6,274,320 B1 8/2001 Rothberg et al.
6,306,597 B1 10/2001 Macevicz
6,352,828 B1 3/2002 Brenner
6,360,235 B1 3/2002 Tilt et al.
6,361,940 B1 3/2002 Van Ness et al.
6,403,320 B1 6/2002 Read et al.
6,462,254 B1 10/2002 Vernachio et al.
6,489,105 B1 12/2002 Matlashewski et al.
6,558,928 B1 5/2003 Landegren
6,569,920 B1 5/2003 Wen et al.
6,582,938 B1 6/2003 Su et al.
6,585,938 B1 7/2003 Machida et al.
6,613,516 B1 9/2003 Christians et al.

(Continued)

FOREIGN PATENT DOCUMENTS

EP 1 321 477 A1 6/2003
EP 1 564 306 A2 8/2005

(Continued)

OTHER PUBLICATIONS

Porecca(Nature methods (2007) vol. 4, pp. 931-936).*

(Continued)

Primary Examiner — Steven Pohnert

(74) *Attorney, Agent, or Firm* — Withers Bergman LLP;
Thomas C. Meyers

(57) **ABSTRACT**

Aspects of the invention relates to methods and composi-
tions that are useful to reduce bias and increase the repro-
ducibility of multiplex analysis of genetic loci. In some
configurations, predetermined preparative steps and/or
nucleic acid sequence analysis techniques are used in mul-
tiplex analyses for a plurality of genetic loci in a plurality of
samples.

9 Claims, 16 Drawing Sheets

Specification includes a Sequence Listing.

GUARDANT HEALTH EX.1017.001

(56)

References Cited

U.S. PATENT DOCUMENTS

6,714,874 B1	3/2004	Myers et al.	9,292,527 B2	3/2016	Kennedy et al.
6,716,580 B2	4/2004	Gold et al.	9,535,920 B2	1/2017	Kennedy et al.
6,719,449 B1	4/2004	Laugharn, Jr. et al.	9,567,639 B2	2/2017	Oliphant et al.
6,818,395 B1	11/2004	Quake et al.	9,677,124 B2	6/2017	Umbarger
6,828,100 B1	12/2004	Ronaghi	10,066,259 B2	9/2018	Gore et al.
6,833,246 B2	12/2004	Balasubramanian	10,202,637 B2	2/2019	Umbarger
6,858,412 B2	2/2005	Willis et al.	10,227,635 B2	3/2019	Umbarger et al.
6,911,345 B2	6/2005	Quake et al.	10,604,799 B2	3/2020	Porreca et al.
6,913,879 B1	7/2005	Schena	10,683,533 B2	6/2020	Umbarger et al.
6,927,024 B2	8/2005	Dodge et al.	2001/0007742 A1	7/2001	Landergren
6,941,317 B1	9/2005	Chamberlin et al.	2001/0046673 A1	11/2001	French et al.
6,948,843 B2	9/2005	Laugharn, Jr. et al.	2002/0001800 A1	1/2002	Lapidus
7,034,143 B1	4/2006	Preparata et al.	2002/0040216 A1	4/2002	Dumont et al.
7,041,481 B2	5/2006	Anderson et al.	2002/0042052 A1	4/2002	Nilsen et al.
7,049,077 B2	5/2006	Yang	2002/0091666 A1	7/2002	Rice et al.
7,057,026 B2	6/2006	Barnes et al.	2002/0164629 A1	11/2002	Quake et al.
7,071,324 B2	7/2006	Preparata et al.	2002/0172954 A1	11/2002	Mao et al.
7,074,564 B2	7/2006	Landegren	2002/0182609 A1	12/2002	Arcot
7,074,586 B1	7/2006	Cheronis et al.	2002/0187496 A1	12/2002	Andersson et al.
7,115,400 B1	10/2006	Adessi et al.	2002/0190663 A1	12/2002	Rasmussen
7,169,560 B2	1/2007	Lapidus et al.	2003/0166057 A1	9/2003	Hildebrand et al.
7,211,390 B2	5/2007	Rothberg et al.	2003/0175709 A1	9/2003	Murphy et al.
7,232,656 B2	6/2007	Balasubramanian et al.	2003/0177105 A1	9/2003	Xiao et al.
7,244,559 B2	7/2007	Rothberg et al.	2003/0203370 A1	10/2003	Yakhini et al.
RE39,793 E	8/2007	Brenner	2003/0208454 A1	11/2003	Rienhoff et al.
7,264,929 B2	9/2007	Rothberg et al.	2003/0224384 A1	12/2003	Sayood et al.
7,282,337 B1	10/2007	Harris	2004/0029264 A1	2/2004	Robbins
7,297,518 B2	11/2007	Quake et al.	2004/0053275 A1	3/2004	Shafer
7,320,860 B2	1/2008	Landegren et al.	2004/0106112 A1	6/2004	Nilsson et al.
7,323,305 B2	1/2008	Leamon et al.	2004/0121373 A1	6/2004	Friedlander et al.
7,335,762 B2	2/2008	Rothberg et al.	2004/0142325 A1	7/2004	Mintz et al.
7,351,528 B2	4/2008	Landegren	2004/0152108 A1	8/2004	Keith et al.
7,393,665 B2	7/2008	Brenner	2004/0161773 A1	8/2004	Rogan et al.
7,510,829 B2	3/2009	Faham et al.	2004/0170965 A1	9/2004	Scholl et al.
7,523,117 B2	4/2009	Zhang et al.	2004/0171051 A1	9/2004	Holloway
7,537,889 B2	5/2009	Sinha et al.	2004/0197813 A1	10/2004	Hoffman et al.
7,537,897 B2	5/2009	Brenner et al.	2004/0209299 A1	10/2004	Pinter et al.
7,544,473 B2	6/2009	Brenner	2005/0003369 A1	1/2005	Christians et al.
7,582,431 B2	9/2009	Drmanac et al.	2005/0026204 A1	2/2005	Landegren
7,598,035 B2	10/2009	Macevicz	2005/0032095 A1	2/2005	Wigler et al.
7,629,151 B2	12/2009	Gold et al.	2005/0048505 A1	3/2005	Fredrick et al.
7,642,056 B2	1/2010	Ahn et al.	2005/0059048 A1	3/2005	Gunderson et al.
7,666,593 B2	2/2010	Lapidus	2005/0100900 A1	5/2005	Kawashima et al.
7,700,323 B2	4/2010	Willis et al.	2005/0112590 A1	5/2005	Boom et al.
7,774,962 B1	8/2010	Ladd	2005/0186589 A1	8/2005	Kowalik et al.
7,776,616 B2	8/2010	Heath et al.	2005/0214811 A1	9/2005	Margulies et al.
RE41,780 E	9/2010	Anderson et al.	2005/0244879 A1	11/2005	Schumm et al.
7,790,388 B2	9/2010	Landegren et al.	2005/0250147 A1	11/2005	Macevicz
7,809,509 B2	10/2010	Milosavljevic	2005/0272065 A1	12/2005	Lakey et al.
7,835,871 B2	11/2010	Kain et al.	2006/0008824 A1	1/2006	Ronaghi et al.
7,862,999 B2	1/2011	Zheng et al.	2006/0019304 A1	1/2006	Hardenbol et al.
7,865,534 B2	1/2011	Genstruct	2006/0024681 A1	2/2006	Smith et al.
7,883,849 B1	2/2011	Dahl	2006/0030536 A1	2/2006	Yu et al.
7,957,913 B2	6/2011	Chinitz et al.	2006/0078894 A1	4/2006	Winkler et al.
7,960,120 B2	6/2011	Rigatti et al.	2006/0149047 A1	7/2006	Nanduri et al.
7,985,716 B2	7/2011	Yershov et al.	2006/0177837 A1	8/2006	Borozen et al.
7,993,880 B2	8/2011	Willis et al.	2006/0183132 A1	8/2006	Fu et al.
8,024,128 B2	9/2011	Rabinowitz et al.	2006/0192047 A1	8/2006	Goossen
8,114,027 B2	2/2012	Triva	2006/0195269 A1	8/2006	Yeatman et al.
8,165,821 B2	4/2012	Zhang	2006/0246500 A1	11/2006	Browne
8,209,130 B1	6/2012	Kennedy et al.	2006/0263789 A1	11/2006	Kincaid
8,283,116 B1	10/2012	Bhattacharyya et al.	2006/0281098 A1	12/2006	Miao et al.
8,462,161 B1	6/2013	Barber	2006/0286577 A1	12/2006	Jia
8,463,895 B2	6/2013	Arora et al.	2006/0292585 A1	12/2006	Nautiyal et al.
8,474,228 B2	7/2013	Adair et al.	2006/0292611 A1	12/2006	Berka et al.
8,496,166 B2	7/2013	Burns et al.	2007/0009925 A1	1/2007	Fang et al.
8,529,744 B2	9/2013	Marziali et al.	2007/0020640 A1	1/2007	McCloskey et al.
8,738,300 B2	5/2014	Porreca et al.	2007/0042369 A1	2/2007	Reese et al.
8,778,609 B1	7/2014	Umbarger	2007/0092883 A1	4/2007	Schouten et al.
8,812,422 B2	8/2014	Nizzari et al.	2007/0114362 A1	5/2007	Feng et al.
8,847,799 B1	9/2014	Kennedy et al.	2007/0128624 A1	6/2007	Gormley et al.
8,976,049 B2	3/2015	Kennedy et al.	2007/0161013 A1	7/2007	Hantash
9,074,244 B2	7/2015	Sparks et al.	2007/0162983 A1	7/2007	Hesterkamp et al.
9,115,387 B2	8/2015	Umbarger	2007/0166705 A1	7/2007	Milton et al.
9,228,233 B2	1/2016	Kennedy et al.	2007/0212704 A1	9/2007	Dong et al.
			2007/0225487 A1	9/2007	Nilsson et al.
			2007/0238122 A1	10/2007	Allbritton et al.
			2007/0244675 A1	10/2007	Shai et al.
			2007/0264653 A1	11/2007	Berlin et al.

(56)

References Cited

U.S. PATENT DOCUMENTS

2008/0003142 A1 1/2008 Link et al.
 2008/0014589 A1 1/2008 Link et al.
 2008/0076118 A1 3/2008 Tooke et al.
 2008/0081330 A1 4/2008 Kahvejian
 2008/0085836 A1 4/2008 Kearns et al.
 2008/0090239 A1 4/2008 Shoemaker et al.
 2008/0125324 A1 5/2008 Petersdorf et al.
 2008/0176209 A1 7/2008 Muller et al.
 2008/0269068 A1 10/2008 Church et al.
 2008/0280955 A1 11/2008 McCamish
 2008/0293589 A1 11/2008 Shapero
 2009/0009904 A1 1/2009 Yasuna et al.
 2009/0019156 A1 1/2009 Mo et al.
 2009/0026082 A1 1/2009 Rothberg et al.
 2009/0029385 A1 1/2009 Christians et al.
 2009/0035777 A1 2/2009 Kokoris et al.
 2009/0042206 A1 2/2009 Schneider et al.
 2009/0098551 A1 4/2009 Landers et al.
 2009/0099041 A1 4/2009 Church et al.
 2009/0105081 A1 4/2009 Rodesch et al.
 2009/0119313 A1 5/2009 Pearce
 2009/0127589 A1 5/2009 Rothberg et al.
 2009/0129647 A1 5/2009 Dimitrova et al.
 2009/0156412 A1 6/2009 Boyce, IV et al.
 2009/0163366 A1 6/2009 Nickerson et al.
 2009/0181389 A1 7/2009 Li et al.
 2009/0191565 A1 7/2009 Lapidus et al.
 2009/0192047 A1 7/2009 Parr et al.
 2009/0202984 A1 8/2009 Cantor
 2009/0203014 A1 8/2009 Wu et al.
 2009/0220955 A1 9/2009 Verrant
 2009/0226975 A1 9/2009 Sabot et al.
 2009/0233814 A1 9/2009 Bashkirov et al.
 2009/0298064 A1 12/2009 Batzoglu et al.
 2009/0301382 A1 12/2009 Patel
 2009/0318310 A1 12/2009 Liu et al.
 2010/0035243 A1 2/2010 Muller et al.
 2010/0035252 A1 2/2010 Rothberg et al.
 2010/0063742 A1 3/2010 Hart et al.
 2010/0069263 A1 3/2010 Shendure et al.
 2010/0076185 A1 3/2010 Adey et al.
 2010/0086914 A1 4/2010 Bentley et al.
 2010/0086926 A1 4/2010 Craig et al.
 2010/0105107 A1 4/2010 Hildebrand et al.
 2010/0137143 A1 6/2010 Rothberg et al.
 2010/0137163 A1 6/2010 Link et al.
 2010/0143908 A1 6/2010 Gillevet
 2010/0159440 A1 6/2010 Messier et al.
 2010/0188073 A1 7/2010 Rothberg et al.
 2010/0196911 A1 8/2010 Hoffman et al.
 2010/0197507 A1 8/2010 Rothberg et al.
 2010/0216151 A1 8/2010 Lapidus et al.
 2010/0216153 A1 8/2010 Lapidus et al.
 2010/0248984 A1 9/2010 Shaffer et al.
 2010/0282617 A1 11/2010 Rothberg et al.
 2010/0285578 A1 11/2010 Selden et al.
 2010/0297626 A1 11/2010 McKernan et al.
 2010/0300559 A1 12/2010 Schultz et al.
 2010/0300895 A1 12/2010 Nobile et al.
 2010/0301042 A1 12/2010 Kahlert
 2010/0301398 A1 12/2010 Rothberg et al.
 2010/0304982 A1 12/2010 Hinz et al.
 2010/0311061 A1 12/2010 Korlach et al.
 2010/0330619 A1 12/2010 Willis et al.
 2011/0004413 A1 1/2011 Carnevali et al.
 2011/0009278 A1 1/2011 Kain et al.
 2011/0015863 A1 1/2011 Pevzner et al.
 2011/0021366 A1 1/2011 Chinitz et al.
 2011/0034342 A1 2/2011 Fox
 2011/0092375 A1 4/2011 Zamore et al.
 2011/0098193 A1 4/2011 Kingsmore et al.
 2011/0117544 A1 5/2011 Lexow
 2011/0159499 A1 6/2011 Hindson et al.
 2011/0166029 A1 7/2011 Margulies et al.
 2011/0224105 A1 9/2011 Kum et al.

2011/0230365 A1 9/2011 Rohlf et al.
 2011/0257889 A1 10/2011 Klammer et al.
 2011/0288780 A1 11/2011 Rabinowitz et al.
 2011/0301042 A1 12/2011 Steinmann et al.
 2012/0015050 A1 1/2012 Abkevich et al.
 2012/0021930 A1 1/2012 Schoen et al.
 2012/0046877 A1 2/2012 Hyland et al.
 2012/0059594 A1 3/2012 Hatchwell et al.
 2012/0074925 A1 3/2012 Oliver
 2012/0079980 A1 4/2012 Taylor et al.
 2012/0115736 A1 5/2012 Bjornson et al.
 2012/0164630 A1 6/2012 Porreca et al.
 2012/0165202 A1 6/2012 Porreca et al.
 2012/0179384 A1 7/2012 Kuramitsu et al.
 2012/0214678 A1 8/2012 Rava et al.
 2012/0216151 A1 8/2012 Sarkar et al.
 2012/0236861 A1 9/2012 Ganeshalingam et al.
 2012/0245041 A1 9/2012 Brenner et al.
 2012/0252020 A1 10/2012 Shuber
 2012/0252684 A1 10/2012 Selifonov et al.
 2012/0258461 A1 10/2012 Weisbart
 2012/0270212 A1 10/2012 Rabinowitz et al.
 2012/0270739 A1 10/2012 Rava et al.
 2013/0129755 A1 5/2013 Song
 2013/0130921 A1 5/2013 Gao et al.
 2013/0178378 A1 7/2013 Hatch et al.
 2013/0183672 A1 7/2013 de Laat et al.
 2013/0222388 A1 8/2013 McDonald
 2013/0268206 A1 10/2013 Porreca et al.
 2013/0268474 A1 10/2013 Nizzari et al.
 2013/0274146 A1 10/2013 Umbarger et al.
 2013/0275103 A1 10/2013 Struble et al.
 2013/0288242 A1 10/2013 Stoughton et al.
 2013/0323730 A1 12/2013 Curry et al.
 2013/0332081 A1 12/2013 Reese et al.
 2013/0337447 A1 12/2013 Porreca et al.
 2013/0344096 A1 12/2013 Chiang et al.
 2014/0129201 A1 5/2014 Kennedy et al.
 2014/0136120 A1 5/2014 Colwell et al.
 2014/0206552 A1 7/2014 Rabinowitz et al.
 2014/0222349 A1 8/2014 Higgins et al.
 2014/0228226 A1 8/2014 Yin et al.
 2014/0255931 A1 9/2014 Porreca et al.
 2014/0308667 A1 10/2014 Umbarger
 2014/0318274 A1 10/2014 Zimmerman et al.
 2014/0342354 A1 11/2014 Evans et al.
 2014/0361022 A1 12/2014 Finneran
 2015/0051085 A1 2/2015 Vogelstein et al.
 2015/0056613 A1 2/2015 Kural
 2015/0111208 A1 4/2015 Umbarger et al.
 2015/0178445 A1 6/2015 Cibulskis et al.
 2015/0258170 A1 9/2015 McCabe et al.
 2015/0299767 A1 10/2015 Armour et al.
 2015/0310163 A1 10/2015 Kingsmore et al.
 2015/0354003 A1 12/2015 Umbarger
 2016/0003812 A1 1/2016 Porreca et al.
 2016/0034638 A1 2/2016 Spence et al.
 2016/0068889 A1 3/2016 Gole et al.
 2016/0210486 A1 7/2016 Porreca et al.
 2016/0251719 A1 9/2016 Umbarger
 2017/0044610 A1 2/2017 Johnson
 2017/0129964 A1 5/2017 Cheung
 2017/0183731 A1 6/2017 Mann et al.
 2017/0275676 A1 9/2017 Umbarger
 2018/0371533 A1 12/2018 Gore et al.
 2019/0233881 A1 8/2019 Umbarger et al.
 2020/0181696 A1 6/2020 Porreca et al.

FOREIGN PATENT DOCUMENTS

EP 10770071.8 11/2010
 EP 2425240 A2 3/2012
 EP 2 437 191 A2 4/2012
 EP 2716766 A1 4/2014
 WO 95/011995 A1 5/1995
 WO 1996/019586 A1 6/1996
 WO 98/014275 A1 4/1998
 WO 98/044151 A1 10/1998
 WO 00/018957 A1 4/2000

(56)

References Cited**FOREIGN PATENT DOCUMENTS**

WO	02/093453	A2	11/2002
WO	2004/018497	A2	3/2004
WO	2004/083819	A2	9/2004
WO	2005/003304	A2	1/2005
WO	2006/084132	A2	8/2006
WO	2007/010251	A2	1/2007
WO	2007/061284	A1	5/2007
WO	2007/107717	A1	9/2007
WO	2007/123744	A2	11/2007
WO	2007/135368	A2	11/2007
WO	2008067551	A2	6/2008
WO	2009/036525	A2	3/2009
WO	2009/076238	A2	6/2009
WO	2010/024894	A1	3/2010
WO	2010/115154	A1	10/2010
WO	2010/126614	A2	11/2010
WO	2011/006020	A1	1/2011
WO	2011066476	A1	6/2011
WO	2011067378	A1	6/2011
WO	2011/102998	A2	8/2011
WO	2011/155833	A2	12/2011
WO	2012/006291	A2	1/2012
WO	2012040387	A1	3/2012
WO	2012/051208	A2	4/2012
WO	2012/087736	A1	6/2012
WO	2012/109500	A2	8/2012
WO	2012/134884	A1	10/2012
WO	2012/149171	A1	11/2012
WO	2012/170725	A2	12/2012
WO	2013/058907	A1	4/2013
WO	2013/148496	A1	10/2013
WO	2013/177086	A1	11/2013
WO	2013/191775	A2	12/2013
WO	2014/052909	A2	4/2014
WO	2014/074246	A1	5/2014
WO	2015/119941	A2	8/2015

OTHER PUBLICATIONS

Wang (Genome Biology (2007) vol. 8, R246.1-246.14).*

Schiffman (Journal of Clinical Oncology (2007) vol. 25, p. 530).*

Tan, 2014, Clinical outcome of preimplantation genetic diagnosis and screening using next generation sequencing, *GigaScience* 3(30):1-9.

Thauvin-Robinet, 2009, The very low penetrance of cystic fibrosis for the R117H mutation: a reappraisal for genetic counseling and newborn screening, *J Med Genet* 46:752-758.

Thiyagarajan, 2006, PathogenMIPer: a tool for the design of molecular inversion probes to detect multiple pathogens, *BMC Bioinformatics* 7:500.

Thompson, 1994, Clustal W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice, *Nuc Acids Res* 22:4673-80.

Thompson, 2011, The properties and applications of single-molecule DNA sequencing, *Genome Biol* 12(2):217.

Thorntenson, 1998, An Automated Hydrodynamic Process for Controlled, Unbiased DNA Shearing, *Genome Res* 8(8): 848-855.

Thorvaldsdottir, 2012, Integrative Genomics Viewer (IGV): high-performance genomics data visualization and exploration, *Brief Bioinform* 24(2):178-92.

Tkachuk, 1990, Detection of bcr-abl Fusion in Chronic Myelogenous Leukemia by in Situ Hybridization, *Science* 250:559.

Tobler, 2005, The SNPlex Genotyping System: A Flexible and Scalable Platform for SNP Genotyping, *J Biomol Tech* 16(4):398.

Tokino, 1996, Characterization of the human p57 KIP2 gene: alternative splicing, insertion/deletion polymorphisms in VNTR sequences in the coding region, and mutational analysis, *Human Genetics* 96:625-31.

Treangen, 2011, Repetitive DNA and next-generation sequencing: computational challenges and solutions, *Nat Rev Gen* 13(1):36-46.

Turner, 2009, Massively parallel exon capture and library-free resequencing across 16 genomes, *Nat Meth* 6:315-316.

Turner, 2009, Methods for genomic partitioning, *Ann Rev Hum Gen* 10:263-284.

Umbarger, 2013, Detecting contamination in Next Generation DNA sequencing libraries, American Society of Human Genetics 63rd Annual Meeting, Abstract, Oct. 22, 2013.

Umbarger, 2014, Next-generation carrier screening, *Gen Med* 16(2):132-140.

Veeneman, 2012, Oculus: faster sequence alignment by streaming read compression, *BMC Bioinformatics* 13:297.

Wahl, 1979, Efficient transfer of large DNA fragments from agarose gels to diazobenzylxymethyl-paper and rapid hybridization by using dextran sulfate, *PNAS* 76:3683-3687.

Wallace, 1979, Hybridization of synthetic oligodeoxyribonucleotides to dp x 174DNA: the effect of single base pair mismatch, *Nucl Acids Res* 6:3543-3557.

Wallace, 1987, Oligonucleotide probes for the screening of recombinant DNA libraries, *Meth Enz* 152:432-442.

Wang, 2005, Allele quantification using molecular inversion probes (MIP), *Nucleic Acids Res* 33(21):e183.

Warner, 1996, A general method for the detection of large CAG repeat expansions by fluorescent PCR, *J Med Genet* 33(12):1022-6.

Warren, 2007, Assembling millions of short DNA sequences using SSAKE, *Bioinformatics*, 23:500-501.

Waszak, 2010, Systematic inference of copy-number genotypes from personal genome sequencing data reveals extensive olfactory gene content diversity, *PLoS Comp Biol* 6(11):e1000988.

Watson, 2004, Cystic fibrosis population carrier screening: 2004 revision of American College of Medical Genetics mutation panel, *Genetics in Medicine* 6(5):387-391.

Williams, 2003, Restriction endonucleases classification, properties, and applications, *Mol Biotechnol* 23(3):225-43.

Wirth, 1999, Quantitative analysis of survival motor neuron copies, *Am J Hum Genet* 64:1340-1356.

Wittung, 1997, Extended DNA-Recognition Repertoire of Peptide Nucleic Acid (PNA): PNA-dsDNA Triplex Formed with Cytosine-Rich Homopyrimidine PNA, *Biochemistry* 36:7973.

Wu, 1998, Sequencing regular and labeled oligonucleotides using enzymatic digestion and ionspray mass spectrometry, *Anal Biochem* 263:129-138.

Wu, 2001, Improved oligonucleotide sequencing by alkaline phosphatase and exonuclease digestions with mass spectrometry, *Anal Biochem* 290:347-352.

Xu, 2012, FastUniq: A fast de novo duplicates removal tool for paired short reads, *PLoS One* 7(12):e52249.

Yau, 1996, Accurate diagnosis of carriers of deletions and duplications in Duchenne/Becker muscular dystrophy by fluorescent dosage analysis, *J Med Gen* 33(7):550-8.

Ye, 2009, Pindel: a pattern growth approach to detect break points of large deletions and medium size insertions from paired-end short reads, *Bioinformatics* 25(21):2865-2871.

Yershov, 1996, DNA analysis and diagnostics on oligonucleotide microchips, *PNAS* 93:4913-4918.

Yoo, 2009, Applications of DNA microarray in disease diagnostics, *J Microbiol Biotech* 19(7):635-46.

Yoon, 2014, MicroDuMIP: target-enrichment technique for microarray-based duplex molecular inversion probes, *Nucl Ac Res* 43(5):e28.

Yoshida, 2004, Role of BRCA1 and BRCA2 as regulators of DNA repair, transcription, and cell cycle in response to DNA damage, *Cancer Science* 95(11):866-71.

Yu, 2007, A novel set of DNA methylation markers in urine sediments for sensitive/specific detection of bladder cancer, *Clin Cancer Res* 13(24):7296-7304.

Yuan, 1981, Structure and mechanism of multifunctional restriction endonucleases, *Ann Rev Biochem* 50:285-319.

Zerbino, 2008, Velvet: Algorithms for de novo short read assembly using de Bruijn graphs, *Genome Research* 18 (5):821-829.

Zhang, 2011, Is Mitochondrial tRNA^{Phe} Variant m.593T.Ca Synergistically Pathogenic Mutation in Chinese LHON Families with m.11778G.A? *PLoS ONE* 6(10) e26511.

Zhao, 2009, PGA4genomics for comparative genome assembly based on genetic algorithm optimization, *Genomics* 94 (4):284-6.

(56)

References Cited**OTHER PUBLICATIONS**

- Zheng, 2011, iAssembler: a package for de novo assembly of Roche-454/Sanger transcriptome sequences, *BMC Bioinformatics* 12:453.
- Zhou, 2014, Bias from removing read duplication in ultra-deep sequencing experiments, *Bioinformatics* 30 (8):1073-1080.
- Zhulidov, 2004, Simple cDNA normalization using kamchatka crab duplex-specific nuclease, *Nucl Acids Res* 32(3):e37.
- Zimmerman, 2010, A novel custom resequencing array for dilated cardiomyopathy, *Gen Med* 12(5):268-78.
- Zimran, 1990, A glucocerebrosidase fusion gene in Gaucher disease, *J Clin Invest* 85:219-222.
- Zuckerman, 1987, Efficient methods for attachment of thiol specific probes to the 3'-ends of synthetic oligodeoxyribonucleotides, *Nucl Acid Res* 15(13):5305-5321.
- Craig, 1997, Removal of repetitive sequences from FISH probes, *Hum Genet* 100:472.
- Cremers, 1998, Autosomal Recessive Retinitis Pigmentosa and Cone-Rod Dystrophy Caused by Splice Site Mutations in the Stargardt's Disease Gene ABCR, *Hum Mol Gen* 7(3):355.
- Cronin, 1996, Cystic Fibrosis Mutation Detection by Hybridization to Light-Generated DNA Probe Arrays *Human Mutation* 7:244.
- Dahl, 2005, Multiplex amplification enabled by selective circularization of large sets of genomic DNA fragments, *Nucleic Acids Res* 33(8):e71.
- Danecek, 2011, The variant call format and VCFtools, *Bioinformatics* 27(15):2156-2158.
- De la Bastide, 2007, Assembling genome DNA sequences with PHRAP, *Current Protocols in Bioinformatics* 17:11.4.1-11.4.15.
- Delcher, 1999, Alignment of whole genomes, *Nuc Acids Res* 27(11):2369-2376.
- Den Dunnen, 2003, Mutation Nomenclature, *Curr Prot Hum Genet* 7.13.1-7.13.8.
- Deng et al., 2012, Supplementary Material, *Nature Biotechnology*, S1-I-S1-I 1, Retrieved from the Internet on Oct. 24, 2012.
- Deng, 2009, targeted bisulfite sequencing reveals changes in DNA methylation, *Nat Biotech* 27(4):353-360.
- Deorowicz, 2013, Data compression for sequencing data, *Alg for Mole Bio* 8:25.
- Diep, 2012, Library-free methylation sequencing with bisulfite padlock probes, *Nature Methods* 9:270-272 (and supplemental information).
- DiGiustini, 2009, De novo sequence assembly of a filamentous fungus using Sanger, 454 and Illumina sequence data, *Genome Biology*, 10:R94.
- Dolinsek, 2013, Depletion of unwanted nucleic acid templates by selection cleavage: LNAzymes, catalytically active oligonucleotides containing locked nucleic acids, open a new window for detecting rare microbial community members, *App Env Microbiol* 79(5):1534-1544.
- Dong, 2011, Mutation surveyor: An in silico tool for sequencing analysis, *Methods Mol Biol* 760:223-37.
- Drmanac, 1992, Sequencing by hybridization: towards an automated sequencing of one million M13 clones arrayed on membranes, *Electrophoresis* 13:566-573.
- Dudley, 2009, A quick guide for developing effective bioinformatics programming skills, *PLoS Comp Biol* 5(12): e1000589.
- Ericsson, 2008, A dual-tag microarray platform for high-performance nucleic acid and protein analyses, *Nucl Acids Res* 36:e45.
- Fares, 2008, Carrier frequency of autosomal-recessive disorders in the Ashkenazi Jewish population: should the rationale for mutation choice for screening be reevaluated?, *Prenatal Diagnosis* 28:236-41.
- Faulstich, 1997, A sequencing method for RNA oligonucleotides based on mass spectrometry, *Anal Chem* 69:4349-4353.
- Faust, 2014, SAMBLASTER: fast duplicate marking and structural variant read extraction, *Bioinformatics* published online May 7, 2014.
- Fitch, 1970, Distinguishing homologs from analogous proteins, *Syst Biol* 19(2):99-113.
- Flaschker, 2007, Description of the mutations in 15 subjects with variant forms of maple syrup urine disease, *J Inherit Metab Dis* 30:903-909.
- Frey, 2006, *Statistics Hacks* 108-115.
- Friedenson, 2005, BRCA1 and BRCA2 Pathways and the Risk of Cancers Other Than Breast or Ovarian, *Medscape General Medicine* 7(2):60.
- Furtado, 2011, Characterization of large genomic deletions in the FBN1 gene using multiplex ligation-dependent probe amplification, *BMC Med Gen* 12:119-125.
- Garber, 2008, Fixing the front end, *Nat Biotech* 26(10):1101-1104.
- Gemayel, 2010, Variable tandem repeats accelerate evolution of coding and regulatory sequences, *Ann Rev Genet* 44:445-77.
- Giusti, 1993, Synthesis and Characterization of 5'-Fluorescent-dye-labeled Oligonucleotides, *PCR Meth Appl* 2:223-227.
- Glover, 1995, Sequencing of oligonucleotides using high performance liquid chromatography and electrospray mass spectrometry, *Rapid Com Mass Spec* 9:897-901.
- Gnirke, 2009, Solution hybrid selection with ultra-long oligonucleotides for massively parallel targeted sequencing, *nature biotechnology* 27:182-9.
- Goto, 1994, A Study on Development of a Deductive Object-Oriented Database and Its Application to Genome Analysis, PhD Thesis, Kyushu University, Kyushu, Japan (106 pages).
- Goto, 2010, BioRuby: bioinformatics software for the Ruby programming language, *Bioinformatics* 26(20):2617-2619.
- Green, 2005, Suicide polymerase endonuclease restriction, a novel technique for enhancing PCR amplification of minor DNA template, *Appl Env Microbiol* 71(8):4721-4727.
- Guerrero-Fernandez, 2013, FQbin: a compatible and optimize dformat for storing and managing sequence data, *IWBBIO Proceedings*, Granada 337-344.
- Gupta, 1991, A general method for the synthesis of 3'-sulfhydryl and phosphate group containing oligonucleotides, *Nucl Acids Res* 19(11):3019-3025.
- Gupta, 2014, Expanding the genetic toolkit: ZFNs, TALENs, and CRISPR-Cas9, *J Clin Invest* 124(10):4154.
- Gustincich, 1991, A fast method for high-quality genomic DNA extraction from whole human blood, *BioTechniques* 11 (3):298-302.
- Gut, 1995, A procedure for selective DNA alkylation and detection by mass spectrometry, *Nucl Acids Res* 23 (8):1367-1373.
- Hallam, 2014, Validation for Clinical Use of, and Initial Clinical Experience with, a Novel Approach to Population-Based Carrier Screening using High-Throughput Next-Generation DNA Sequencing, *J Mol Diagn* 16:180-9.
- Hammond, 1996, Extraction of DNA from preserved animal specimens for use in randomly amplified polymorphic DNA analysis, *Anal Biochem* 240:298-300.
- Hardenbol, 2003, Multiplexed genotyping with sequence-tagged molecular inversion probes, *Nat Biotech* 21:673-8.
- Hardenbol, 2005, Highly multiplexed molecular inversion probe genotyping: over 10,000 targeted SNPs genotyped in a single tube assay, *Genome Res* 15:269-75.
- Harris, 2006, Defects can increase the melting temperature of DNA-nanoparticle assemblies, *J Phys Chem B* 110 (33):16393-6.
- Harris, 2008, Helicos True Single Molecule Sequencing (tSMS) *Science* 320:106-109.
- Harris, 2008, Single-molecule DNA sequencing of a viral genome, *Science* 320(5872):106-9.
- Heger, 2006, Protonation of Cresol Red in Acidic Aqueous Solutions Caused by Freezing, *J Phys Chem B* 110 (3):1277-1287.
- Heid, 1996, Real time quantitative PCR, *Genome Res* 6:986-994.
- Hiatt, 2013, Single molecule molecular inversion probes for targeted, high-accuracy detection of low-frequency variation, *Genome Res* 23:843-54.
- Hodges, 2007, Genome-wide in situ exon capture for selective resequencing, *Nat Genet* 39(12):1522-7.
- Abravaya, 1995, Detection of point mutations with a modified ligase chain reaction (GAP-LCR), *Nucleic Acids Research*, 23(4): 675-682.

(56)

References Cited**OTHER PUBLICATIONS**

- Adey, 2010, Rapid, low-input, low-bias construction of shotgun fragment libraries by high-density in vitro transposition, *Genome Biol* 11:R119.
- Agno, 1969, The alkaline denaturation of DNA, *Biophys J* 9(11):1281-1311.
- Agrawal, 1990, Site-specific functionalization of oligodeoxynucleotides for non-radioactive labelling, *Tetrahedron Let* 31:1543-1546.
- Akhras, 2007, Connector inversion probe technology: A powerful one-primer multiplex DNA amplification system for numerous scientific applications, *PLoSOne* 9:e915.
- Akhras, 2007, PathogenMip Assay: a multiplex pathogen detection assay, *PLOS One* 2:e2230.
- Alazard, 2002, Sequencing of production-scale synthetic oligonucleotides by enriching for coupling failures using matrix-assisted laser desorption/ionization time-of-flight mass spectrometry, *Anal Biochem* 301:57-64.
- Alazard, 2006, Sequencing oligonucleotides by enrichment of coupling failures using matrix-assisted laser desorption/ionization time-of-flight mass spectrometry, *Curr Protoc Nucleic Acid Chem*, Chapter 10, Unit 10:1-7.
- Albert, 2007, Direct selection of human genomic loci by microarray hybridization, *Nature Methods* 4(11):903-5.
- Aljanabi, 1997, Universal and rapid salt-extraction of high quality genomic DNA for PCR-based techniques, *Nucl. Acids Res* 25:4692-4693.
- Antonarakis and the Nomenclature Working Group, 1998, Recommendations for a nomenclature system for human gene mutations, *Human Mutation* 11:1-3.
- Archer, 2014, Selective and flexible depletion of problematic sequences from RNA-seq libraries at the cDNA stage, *BMC Genomics* 15(1):401.
- Ball, 2009, Targeted and genome-scale strategies reveal gene-body methylation signatures in human cells, *Nat Biotech* 27:361-8.
- Balzer, 2013, Filtering duplicate reads from 454 pyrosequencing data, *Bioinformatics* 29(7):830-836.
- Barany, 1991, Genetic disease detection and DNA amplification using cloned thermostable ligase, *PNAS* 88:189-193.
- Barany, 1991, The Ligase Chain Reaction in a PCR World, *Genome Research* 1:5-16.
- Bau, 2008, Targeted next-generation sequencing by specific capture of multiple genomic loci using low-volume microfluidic DNA arrays, *Analytical and Bioanal Chem* 393(1):171-5.
- Beer, 1962, Determination of base sequence in nucleic acids with the electron microscope: visibility of a marker, *PNAS* 48(3):409-416.
- Bell, 2011, Carrier testing for severe childhood recessive diseases by next-generation sequencing, *Sci Trans Med* 3 (65ra4).
- Benner, 2001, Evolution, language and analogy in functional genomics, *Trends Genet* 17:414-8.
- Bentzley, 1996, Oligonucleotide sequence and composition determined by matrix-assisted laser desorption/ionization, *Anal Chem* 68:2141-2146.
- Bentzley, 1998, Base specificity of oligonucleotide digestion by calf spleen phosphodiesterase with matrix-assisted laser desorption ionization analysis, *Anal Biochem* 258:31-37.
- Bhangale, 2006, Automating resequencing-based detection of insertion-deletion polymorphisms, *Nature Genetics* 38:1457-1462.
- Bickle, 1993, Biology of DNA Restriction, *Microbiol Rev* 57(2):434-50.
- Bonfield, 2013, Compression of FASTQ and SAM format sequencing data, *PLoS One* 8(3):e59190.
- Bose, 2012, BIND—An algorithm for loss-less compression of nucleotide sequence data, *J Biosci* 37(4):785-789.
- Boyden, 2013, High-throughput screening for SMN1 copy number loss by next-generation sequencing, American Society of Human Genetics 63rd Annual Meeting, Abstract, Oct. 22, 2013.
- Boyer, 1971, DNA restriction and modification mechanisms in bacteria, *Ann Rev Microbiol* 25:153-76.
- Braasch, 2001, Locked nucleic acid (LNA): fine-tuning the recognition of DNA and RNA, *Chemistry & Biology* 8(1):1-7.
- Braslavsky, 2003, Sequence information can be obtained from single DNA molecules, *PNAS* 100:3960-4.
- Brinkman, 2004, Splice Variants as Cancer Biomarkers, *Clin Biochem* 37:584.
- Brison, 1982, General method for cloning amplified DNA by differential screening, *Mol Cell Biol* 2(5):578-587.
- Brown, 1979, Chemical synthesis and cloning of a tyrosine tRNA gene, *Methods Enzymol* 68:109-51.
- Browne, 2002, Metal ion-catalyzed nucleic Acid alkylation and fragmentation, *J Am Chem Soc* 124(27):7950-7962.
- Brownstein, 2014, An international effort towards developing standards for best practices in analysis, interpretation and reporting of clinical genome sequencing results in the CLARITY Challenge, *Genome Biol* 15:R53.
- Bunyan, 2004, Dosage analysis of cancer predisposition genes by multiplex ligation-dependent probe amplification, *British Journal of Cancer*, 91(6):1155-59.
- Burrow, 1994, A block-sorting lossless data compression algorithm, Technical Report 124, Digital Equipment Corporation, CA. (24 pages).
- Carpenter, 2013, Pulling out the 1%: whole-genome capture for the targeted enrichment of ancient DNA sequencing libraries, *Am J Hum Genet* 93(5):852-864.
- Caruthers, 1985, Gene synthesis machines: DNA chemistry and its uses, *Science* 230:281-285.
- Castellani, 2008, Consensus on the use of and interpretation of cystic fibrosis mutation analysis in clinical practice, *J Cyst Fib* 7:179-196.
- Challis, 2012, An integrative variant analysis suite for whole exome next-generation sequencing data, *BMC Informatics* 13(8):1-12.
- Chan, 2011, Natural and engineered nicking endonucleases—from cleavage mechanism to engineering of strand-specificity, *Nucl Acids Res* 39(1):1-18.
- Chen, 2010, Identification of racehorse and sample contamination by novel 24-plex STR system, *Forensic Sci Int: Genetics* 4:158-167.
- Chennagiri, 2013, A generalized scalable database model for storing and exploring genetic variations detected using sequencing data, American Society of Human Genetics 63rd Annual Meeting, Abstract, Oct. 22, 2013.
- Chevreux, 1999, Genome sequence assembly using trace signals and additional sequence information, *Proc GCB* 99:45-56.
- Chirgwin, 1979, Isolation of biologically active ribonucleic acid from sources enriched in ribonuclease, *Biochemistry*, 18:5294-99.
- Choe, 2010, Novel CFTR Mutations in a Korean Infant with Cystic Fibrosis and Pancreatic Insufficiency, *J Korean Med Sci* 25:163-5.
- Ciotti, 2004, Triplet repeat primed PCR (TP PCR) in molecular diagnostic testing for Friedrich ataxia, *J Mol Diag* 6 (4):285-9.
- Cock, 2010, The Sanger FASTQ file format for sequences with quality scores, and the Solexa/Illumina FASTQ variants, *Nucleic Acids Res* 38(6):1767-1771.
- Collins, 2004, Finishing the euchromatic sequence of the human genome, *Nature* 431(7011):931-45.
- Maddalena, 2005, Technical standards and guidelines: molecular genetic testing for ultra-rare disorders, *Genet Med* 7:571-83.
- Malewicz, 2010, Pregel: a system for large-scale graph processing, *Proc. ACM SIGMOD Int Conf Mgmt Data* 135-46.
- Mamanova, 2010, Target-enrichment strategies for next-generation sequencing, *Nat Meth* 7(2):111-118.
- Margulies, 2005, Genome sequencing in micro-fabricated high-density picotiter reactors, *Nature*, 437:376-380.
- Marras, 1999, Multiplex detection of single-nucleotide variations using molecular beacons, *Genetic Analysis: Biomolecular Engineering* 14:151.
- Maxam, 1977, A new method for sequencing DNA, *PNAS*, 74:560-564.
- May 1988, How Many Species Are There on Earth?, *Science* 241(4872):1441-9.
- McDonnell, 2007, Antisepsis, disinfection, and sterilization: types, action, and resistance, p. 239.
- McKenna, 2010, The Genome Analysis Toolkit: A MapReduce framework for analyzing next-generation DNA sequencing data, *Genome Research* 20:1297-1303.

(56)

References Cited**OTHER PUBLICATIONS**

- Messiaen, 1999, Exon 10b of the NF1 gene represents a mutational hotspot and harbors a recurrent missense mutation Y489C associated with aberrant splicing, *Genetics in Medicine*, 1(6):248-253.
- Meyer, 2007, Targeted high-throughput sequencing of tagged nucleic acid samples, *Nucleic Acids Research* 35(15): e97 (5 pages).
- Meyer, 2008, Parallel tagged sequencing on the 454 platform, *Nat Protocol* 3(2):267-278.
- Miesenbock, 1998, Visualizing secretion and synaptic transmission with pH-sensitive green fluorescent proteins, *Nature* 394(6689):192-95.
- Miller, 2010, Assembly algorithms for next-generation sequencing data, *Genomics* 95:315-327.
- Mills, 2010, Mapping copy number variation by population-scale genome sequencing, *Nature* 470(7332):59-65.
- Miner, 2004, Molecular barcodes detect redundancy and contamination in hairpin-bisulfite PCR, *Nucl Acids Res* 32 (17):e135.
- Minton, 2011, Mutation Surveyor: software for DNA sequence analysis, *Meth Mol Biol* 688:143-53.
- Miyake, 2009, PIK3CA gene mutations and amplification in uterine cancers, *Canc Lett* 261:120-126.
- Miyazaki, 2009, Characterization of deletion breakpoints in patients with dystrophinopathy carrying a deletion of exons 15-55 of the Duchenne muscular dystrophy (DMD) gene, *J Hum Gen* 54:127-30.
- Mockler, 2005, Applications of DNA tiling arrays for whole-genome analysis, *Genomics* 85(1):1-15.
- Mohammed, 2012, Delimitate—a fast and efficient methods for loss-less compression of genomic sequences, *Bioinformatics* 28(19):2527-2529.
- Moudrianakis, 1965, Base Sequence Determination in Nucleic Acids with the Electron Microscope, III. Chemistry and Microscopy of Guanine-Labeled DNA, *PNAS*, 53:564-71.
- Mullan, 2002, Multiple sequence alignment—the gateway to further analysis, *Brief Bioinform* 3(3):303-5.
- Munne, 2012, Preimplantation genetic diagnosis for aneuploidy and translocations using array comparative genomic hybridization, *Curr Genomics* 13(6):463-470.
- Nan, 2006, A novel CFTR mutation found in a Chinese patient with cystic fibrosis, *Chinese Med J* 119(2):103-9.
- Narang, 1979, Improved phosphotriester method for the synthesis of gene fragments, *Meth Enz* 68:90-98.
- Nelson, 1989, Bifunctional oligonucleotide probes synthesized using a novel CPG support are able to detect single base pair mutations, *Nucl Acids Res* 17(18):7187-7194.
- Ng, 2009, Targeted capture and massively parallel sequencing of 12 human exomes, *Nature* 461(7261):272-6.
- Nicholas, 2002, Strategies for multiple sequence alignment, *Biotechniques* 32:572-91.
- Nickerson, 1990, Automated DNA diagnostics using an ELISA-based oligonucleotide ligation assay, *PNAS* 87:8923-7.
- Nielsen, 1999, *Peptide Nucleic Acids, Protocols and Applications* (Norfolk: Horizon Scientific Press, 1-19).
- Nilsson, 2006, Analyzing genes using closing and replicating circles, *Trends in Biotechnology* 24:83-8.
- Ning, 2001, SSAHA: a fast search method for large DNA databases, *Genome Res* 11(10):1725-9.
- Nordhoff, 1993, Ion stability of nucleic acids in infrared matrix-assisted laser desorption/ionization mass spectrometry, *Nucl Acid Res* 21(15):3347-57.
- Nuttall, 2013, Rapid and accurate large-scale genotyping of duplicated genes and discovery of interlocus gene conversions, *Nat Meth* 10(9):903-909.
- Nuttall, 2014, Resolving genomic disorder-associated breakpoints within segmental DNA duplications using massively parallel sequencing, *Nat Prot* 9(6):1496-1513.
- O'Roak, 2012, Multiplex targeted sequencing identifies recurrently mutated genes in autism spectrum disorders, *Science* 338(6114):1619-1622.
- Oefner, 1996, Efficient random sub-cloning of DNA sheared in a recirculating point-sink flow system, *Nucleic Acids Res* 24(20):3879-3886.
- Oka, 2006, Detection of loss of heterozygosity in the p53 gene in renal cell carcinoma and bladder cancer using the polymerase chain reaction, *Mol Carcinogenesis* 4(1):10-13.
- Okoniewski, 2013, Precise breakpoint localization of large genomic deletions using PacBio and Illumina next-generation sequencers, *Biotechniques* 54(2):98-100.
- Okou, 2007, Microarray-based genomic selection for high-throughput resequencing, *Nat Meth* 4(11):907-909.
- Oliphant, 2002, BeadArray technology: enabling an accurate, cost-effective approach to high-throughput genotyping, *Biotechniques Suppl*:56-8, 60-1.
- Ordahl, 1976, Sheared DNA fragment sizing: comparison of techniques, *Nucleic Acids Res* 3:2985-2999.
- Ostrer, 2001, A genetic profile of contemporary Jewish populations, *Nat Rev Genet* 2(11):891-8.
- Owens, 1998, Aspects of oligonucleotide and peptide sequencing with MALDI and electrospray mass spectrometry, *Bioorg Med Chem* 6:1547-1554.
- Parameswaran, 2007, A pyrosequencing-tailored nucleotide barcode design unveils opportunities for large-scale sample multiplexing, *Nucl Acids Res* 35:e130.
- Parkinson, 2012, Preparation of high-quality next-generation sequencing libraries from picogram quantities of target DNA, *Genome Res* 22:125-133.
- Pastor, 2010, Conceptual modeling of human genome mutations: a dichotomy between what we have and what we should have, 2010 *Proc BIOSTEC Bioinformatics*, pp. 160-166.
- Paton, 2000, Conceptual modelling of genomic information, *Bioinformatics* 16(6):548-57.
- Pearson, 1988, Improved tools for biological sequence comparison, *PNAS* 85(8):2444-8.
- Holland, 2008, BioJava: an open-source framework for bioinformatics, *Bioinformatics* 24(18):2096-2097.
- Homer, 2008, Resolving Individuals Contributing Trace Amounts of DNA to Highly Complex Mixtures Using High-Density SNP Genotyping Microarrays, *PLoS One* 4(8):e1000167.
- Homer, 2009, BFAST: An alignment tool for large scale genome resequencing, *PLoS ONE* 4(11):e7767.
- Housley, 2009, SNP discovery and haplotype analysis in the segmentally duplicated DRD5 coding region, *Ann Hum Genet* 73(3):274-282.
- Huang, 2008, Comparative analysis of common CFTR polymorphisms poly-T, TG repeats and M470V in a healthy Chinese population, *World J Gastroenterol* 14(12):1925-30.
- Husemann, 2009, *Phylogenetic Comparative Assembly, Algorithms in Bioinformatics: 9th International Workshop*, pp. 145-156, Salzberg & Warnow, Eds. Springer-Verlag, Berlin, Heidelberg.
- Illumina, 2010, *De Novo assembly using Illumina reads*, Technical Note (8 pages).
- International Human Genome Sequencing Consortium, 2004, Finishing the euchromatic sequence of the human genome, *Nature* 431:931-945.
- Iqbal, 2012, De novo assembly and genotyping of variants using colored de Bruijn graphs, *Nature Genetics* 44:226-232.
- Isosomppi, 2009, Disease-causing mutations in the CLRN1 gene alter normal CLRN1 protein trafficking to the plasma membrane, *Mol Vis* 15:1806-1818.
- Jaijo, 2010, Microarray-based mutation analysis of 183 Spanish families with Usher syndrome, *Invest Ophthalmol Vis Sci* 51(3):1311-7.
- Jensen, 2001, Orthologs and paralogs—we need to get it right, *Genome Biol* 2(8):1002-1002.3.
- Jones, 2008, Core signaling pathways in human pancreatic cancers revealed by global genomic analyses, *Science* 321(5897):1801-1806.
- Kambara, 1988, Optimization of Parameters in a DNA Sequenator Using Fluorescence Detection, *Nature Biotechnology* 6:816-821.
- Kennedy, 2013, Accessing more human genetic variation with short sequencing reads, *American Society of Human Genetics 63rd Annual Meeting, Abstract*, Oct. 22, 2013.

(56)

References Cited**OTHER PUBLICATIONS**

- Kent, 2002, BLAT—The BLAST-like alignment tool, *Genome Res* 12(4): 656-664.
- Kerem, 1989, Identification of the cystic fibrosis gene: genetic analysis, *Science* 245:1073-1080.
- Kinde, 2012, FAST-SeqS: a simple and effective method for detection of aneuploidy by massively parallel sequencing, *PLoS One* 7(7):e41162.
- Kircher, 2010, High-throughput DNA sequencing—concepts and limitations, *Bioassays* 32:524-36.
- Kirpekar, 1994, Matrix assisted laser desorption/ionization mass spectrometry of enzymatically synthesized RNA up to 150 kDa, *Nucl Acids Res* 22:3866-3870.
- Klein, 2011, LOCAS—A low coverage sequence assembly tool for re-sequencing projects, *PLoS One* 6(8):e23455.
- Kneen, 1998, Green fluorescent protein as a noninvasive intracellular pH indicator, *Biophys J* 74(3):1591-99.
- Koboldt, 2009, VarScan: variant detection in massively parallel sequencing of individual and pooled samples, *Bioinformatics* 25:2283-85.
- Krawitz, 2010, Microindel detection in short-read sequence data, *Bioinformatics* 26(6):722-729.
- Kreindler, 2010, Cystic fibrosis: exploiting its genetic basis in the hunt for new therapies, *Pharmacol Ther* 125 (2):219-229.
- Krishnakumar, 2008, A comprehensive assay for targeted multiplex amplification of human DNA sequences, *PNAS* 105:9296-301.
- Kumar, 2010, Comparing de novo assemblers for 454 transcriptome data, *Genomics* 11:571.
- Kurtz, 2004, Versatile and open software for comparing large genomes, *Genome Biol* 5:R12.
- Lam, 2008, Compressed indexing and local alignment of DNA, *Bioinformatics* 24(6):791-97.
- Langmead, 2009, Ultrafast and memory-efficient alignment of short DNA sequences to the human genome, *Genome Biol* 10:R25.
- Larkin, 2007, Clustal W and Clustal X version 2.0, *Bioinformatics*, 23(21):2947-2948.
- Lecompte, 2001, Multiple alignment of complete sequences (MACS) in the post-genomic era, *Gene* 270(1-2):17-30.
- Li, 2003, DNA binding and cleavage by the periplasmic nuclease Vyn: a novel structure with a known active site, *EMBO J* 22(15):4014-4025.
- Li, 2008, SOAP: short oligonucleotide alignment program, *Bioinformatics* 24(5):713-14.
- Li, 2009, Fast and accurate short read alignment with Burrows-Wheeler transform, *Bioinformatics*, 25 (14):1754-60.
- Li, 2009, SOAP2: an improved ultrafast tool for short read alignment, *Bioinformatics* 25(15):1666-67.
- Li, 2009, The Sequence Alignment/Map format and SAMtools, *Bioinformatics* 25(16):2078-9.
- Li, 2010, Fast and accurate long-read alignment with Burrows-Wheeler transform, *Bioinformatics* 26(5):589-95.
- Li, 2011, Improving SNP discovery by base alignment quality, *Bioinformatics* 27:1157.
- Li, 2011, Single nucleotide polymorphism genotyping and point mutation detection by ligation on microarrays, *J Nanosci Nanotechnol* 11(2):994-1003.
- Li, 2012, A new approach to detecting low-level mutations in next-generation sequence data, *Genome Biol* 13:1-15.
- Li, 2014, HUGO: Hierarchical mUlti-reference Genome compression for aligned reads, *JAMIA* 21:363-373.
- Lin, 2008, ZOOM! Zillions of Oligos Mapped, *Bioinformatics*, 24:2431.
- Lin, 2010, A molecular inversion probe assay for detecting alternative splicing, *BMC Genomics* 11(712):1-14.
- Lin, 2012, Development and evaluation of a reverse dot blot assay for the simultaneous detection of common alpha and beta thalassemia in Chinese, *Blood Cells Molecules, and Diseases* 48(2):86-90.
- Lipman, 1985, Rapid and sensitive protein similarity searches, *Science* 227(4693):1435-41.
- Liu, 2012, Comparison of next-generation sequencing systems, *J Biomed Biotech* 2012:251364.
- Llopis, 1998, Measurement of cytosolic, mitochondrial, and Golgi pH in single living cells with green fluorescent proteins, *PNAS* 95(12):6803-08.
- Ma, 2006, Application of real-time polymerase chain reaction (RT-PCR), *J Am Soc* 1-15.
- MacArthur, 2014, Guidelines for investigating causality of sequence variants in human disease, *Nature* 508:469-76.
- Pertea, 2003, TIGR gene indices clustering tools (TGICL), *Bioinformatics* 19(5):651-52.
- Pieles, 1993, Matrix-assisted laser desorption ionization time-of-flight mass spectrometry: A powerful tool for the mass and sequence analysis of natural and modified oligonucleotides, *Nucleic Acids Res* 21:3191-3196.
- Pinho, 2013, MFCompress: a compression tool for FASTA and multi-FASTA data, *Bioinformatics* 30(1):117-8.
- Porreca, 2007, Multiplex amplification of large sets of human exons, *Nat Meth* 4(11):931-936.
- Porreca, 2013, Analytical performance of a Next-Generation DNA sequencing-based clinical workflow for genetic carrier screening, *American Society of Human Genetics 63rd Annual Meeting, Abstract*, Oct. 22, 2013.
- Pourmand, 2006, PathogenMIPer: a tool for the design of molecular inversion probes, *BMC informatics* 7:500.
- Procter, 2006, Molecular diagnosis of Prader-Willi and Angelman syndromes by methylation-specific melting analysis and methylation-specific multiplex ligation-dependent probe amplification, *Clin Chem* 52(7):1276-83.
- Qiagen, 2011, *Gentra Puregene handbook*, 3d Ed. (72 pages).
- Quail, 2010, DNA: Mechanical Breakage, In *Encyclopedia of Life Sciences*, John Wiley & Sons Ltd, Chichester (5 pages).
- Rambaut, 1997, Seq-Gen: an application for the Monte Carlo simulation of DNA sequence evolution along phylogenetic trees, *Bioinformatics* 13:235-38.
- Richards, 2008, ACMG recommendations for standards for interpretation and reporting of sequence variations: Revisions, *Genet Med* 10(4):294-300.
- Richter, 2008, MetaSim—A Sequencing Simulator for Genomics and Metagenomics, *PLoS ONE* 3:e3373.
- Roberts, 1980, Restriction and modification enzymes and their recognition sequences, *Nucleic Acids Res* 8(1):r63-r80.
- Rodriguez, 2010, Constructions from Dots and Lines, *Bull Am Soc Inf Sci Tech* 36(6):35-41.
- Rosendahl, 2013, CFTR, SPINK1, CTFR and PRSS1 variants in chronic pancreatitis: is the role of mutated CFTR overestimated?, *Gut* 62:582-592.
- Rothberg, 2011, An integrated semiconductor device enabling non-optical genome sequencing, *Nature* 475:348-352.
- Rowntree, 2003, The phenotypic consequences of CFTR mutations, *Ann Hum Gen* 67:471-485.
- Baihan, 2009, Update on Usher syndrome, *Curr Op Neurology* 22(1):19-24.
- Sanger, 1977, DNA Sequencing with chain-terminating inhibitors, *PNAS* 74(12):5463-5467.
- Santa Lucia, 1998, A unified view of polymer, dumbbell, and oligonucleotide DNA nearest-neighbor thermodynamics, *PNAS* 95(4):1460-5.
- Sargent, 1987, Isolation of differentially expressed genes, *Meth Enzym* 152:423-432.
- Sauro, 2004, How Do You Calculate a Z-Score/ Sigma Level?, <https://www.measuringusability.com/zcalc.htm> (online publication).
- Sauro, 2004, What's a Z-score and Why Use It in Usability Testing?, <https://www.measuringusability.com/z.htm> (online publication).
- Schadt, 2010, A window into third-generation sequencing, *Human Mol Genet* 19(R2):R227-40.
- Schatz, 2010, Assembly of large genomes using second-generation sequencing, *Genome Res.*, 20:1165-1173.
- Schiffman, 2009, Molecular inversion probes reveal patterns of 9p21 deletion and copy number aberrations in childhood leukemia, *Cancer Genetics and Cytogenetics* 193:9-18.

(56)

References Cited**OTHER PUBLICATIONS**

- Schneeberger, 2011, Reference-guided assembly of four diverse *Arabidopsis thaliana* genomes, PNAS 108 (25):10249-10254.
- Schouten, 2002, Relative Quantification of 40 Nucleic Acid Sequences by Multiplex Ligation-Dependent Probe Amplification, Nucleic Acids Res 30 (12):257.
- Schrijver, 2005, Diagnostic testing by CFTR gene mutation analysis in a large group of Hispanics, J Mol Diag 7 (2):289-299.
- Schuette, 1995, Sequence analysis of phosphorothioate oligonucleotides via matrix-assisted laser desorption ionization time-of-flight mass spectrometry, J Pharm Biomed Anal 13:1195-1203.
- Schwartz, 2009, Identification of cystic fibrosis variants by polymerase chain reaction/oligonucleotide ligation assay, J Mol Diag 11(3):211-15.
- Schwartz, 2011, Clinical utility of single nucleotide polymorphism arrays, Clin Lab Med 31(4):581-94.
- Sequeira, 1997, Implementing generic, object-oriented models in biology, Ecological Modeling 94.1:17-31.
- Shagin, 2002, A novel method for SNP detection, Genome Res 12:1935.
- Shen, 2011, High quality DNA sequence capture of 524 disease candidate genes, PNAS 108(16):6549-6554.
- Shen, 2013, Multiplex capture with double-stranded DNA probes, Genome Medicine 5(50):1-8.
- Sievers, 2011, Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega, Mol Syst Biol 7:539.
- Simpson, 2009, ABySS: A parallel assembler for short read sequence data, Genome Res., 19(6):1117-23.
- Slater, 2005, Automated generation of heuristics for biological sequence comparison, BMC Bioinformatics 6:31.
- Smirnov, 1996, Sequencing oligonucleotides by exonuclease digestion and delayed extraction matrix-assisted laser desorption ionization time-of-flight mass spectrometry, Anal Biochem 238:19-25.
- Smith, 1985, The synthesis of oligonucleotides containing an aliphatic amino group at the 5' terminus: synthesis of fluorescent DNA primers for use in DNA sequence analysis, Nucl Acid Res 13:2399-2412.
- Smith, 2010, Highly-multiplexed barcode sequencing: an efficient method for parallel analysis of pooled samples, Nucleic Acids Research 38(13):e142 (8 pages).
- Soni, 2007, Progress toward ultrafast DNA sequencing using solid-state nanopores, Clin Chem 53(11):1996-2001.
- Spanu, 2010, Genome expansion and gene loss in powdery mildew fungi reveal tradeoffs in extreme parasitism, Science 330(6010):1543-46.
- Sproat, 1987, The synthesis of protected 5'-mercapto-2',5'-dideoxyribonucleoside-3'-O-phosphoramidites; uses of 5'-mercapto-oligodeoxyribonucleotides, Nucl Acid Res 15:4837-4848.
- Streit, 2003, CFTR gene: molecular analysis in patients from South Brazil, Molecular Genetics and Metabolism 78:259-264.
- Strom, 2005, Mutation detection, interpretation, and applications in the clinical laboratory setting, Mutat Res 673:160-67.
- Summerer, 2009, Enabling technologies of genomic-scale sequence enrichment for targeted high-throughput sequencing, Genomics 94(6):363-8.
- Summerer, 2010, Targeted High Throughput Sequencing of a Cancer-Related Exome Subset by Specific Sequence Capture With a Fully Automated Microarray Platform, Genomics 95(4):241-246.
- Sunnucks, 1996, Microsatellite and chromosome evolution of parthenogenetic sitobion aphids in Australia, Genetics 144:747-756.

* cited by examiner

FIG. 1




FIG. 2




FIG. 3




FIG. 4A




FIG. 4B




FIG. 4C





FIG. 5


FIG. 6


FIG. 7


FIG. 8

FIG. 9




FIG. 10





FIG. 11A


FIG. 11B


FIG. 11C


FIG. 12



FIG. 13

FIG. 14



1

METHODS AND COMPOSITIONS FOR EVALUATING GENETIC MARKERS

RELATED APPLICATIONS

This application is a continuation of U.S. application Ser. No. 15/231,687, filed Aug. 8, 2016, which is a continuation of U.S. application Ser. No. 13/266,862, filed Mar. 13, 2012, which is a national phase application that claims the benefit of, and priority to, international PCT Application No. PCT/US2010/001293, filed Apr. 30, 2010, which claims the benefit under 35 U.S.C. § 119(e) of U.S. Provisional Application No. 61/174,470, filed Apr. 30, 2009, U.S. Provisional Application No. 61/178,923, filed May 15, 2009, U.S. Provisional Application No. 61/179,358, filed May 18, 2009, and U.S. Provisional Application No. 61/182,089, filed May 28, 2009, the entire contents of each of which are incorporated herein by reference.

FIELD OF INVENTION

The invention relates to methods and compositions for determining genotypes in patient samples.

BACKGROUND OF THE INVENTION

Information about the genotype of a subject is becoming more important and relevant for a range of healthcare decisions as the genetic basis for many diseases, disorders, and physiological characteristics is further elucidated. Medical advice is increasingly personalized, with individual decisions and recommendations being based on specific genetic information. Information about the type and number of alleles at one or more genetic loci impacts disease risk, prognosis, therapeutic options, and genetic counseling amongst other healthcare considerations.

For cost-effective and reliable medical and reproductive counseling on a large scale, it is important to be able correctly and unambiguously identify the allelic status for many different genetic loci in many subjects.

Numerous technologies have been developed for detecting and analyzing nucleic acid sequences from biological samples. These technologies can be used to genotype subjects and determine the allelic status of any locus of interest. However, they are not sufficiently robust and cost-effective to be scaled up for reliable high throughput analysis of many genetic loci in large numbers of patients. The frequency of incorrect or ambiguous calls is too high for current technology to manage large numbers of patient samples without involving expensive and time-consuming steps to resolve uncertainties and provide confidence in the information output.

SUMMARY OF THE INVENTION

Aspects of the invention relate to preparative and analytical methods and compositions for evaluating genotypes, and in particular, for determining the allelic identity (or identities in a diploid organism) of one or more genetic loci in a subject.

Aspects of the invention are based, in part, on the identification of different sources of ambiguity and error in genetic analyses, and, in part, on the identification of one or more approaches to avoid, reduce, recognize, and/or resolve these errors and ambiguities at different stages in a genetic analysis.

2

According to aspects of the invention, certain types of genetic information can be under-represented or over-represented in a genetic analysis due to a combination of stochastic variation and systematic bias in any of the preparative stages (e.g., capture, amplification, etc.), determining stages (e.g., allele-specific detection, sequencing, etc.), data interpretation stages (e.g., determining whether the assay information is sufficient to identify a subject as homozygous or heterozygous), and/or other stages.

According to aspects of the invention, error or ambiguity may be apparent in a genetic analysis, but not readily resolved without running additional samples or more expensive assays (e.g., array-based assays may report no-calls due to noisy/low signal). According to further aspects of the invention, error or ambiguity may not be accounted for in a genetic analysis and incorrect base calls may be made even when the evidence for them is limited and/or not statistically significant (e.g., next-generation sequencing technologies may report base calls even if the evidence for them is not statistically significant). According to further aspects of the invention error or ambiguity may be problematic for a multi-step genetic analysis because it is apparent but not readily resolved in one or more steps of the analysis and not apparent or accounted for in other steps of the analysis.

In some embodiments, sources of error and ambiguity in one or more steps can be addressed by capturing and/or interrogating each target locus of interest with one or more sets of overlapping probes that are designed to overcome any systematic bias or stochastic effects that may impact the complexity and/or fidelity of the genetic information that is generated.

In some embodiments, sources of error and ambiguity in one or more steps can be addressed by capturing and/or interrogating each target locus of interest with at least one set of probes, wherein different probes are labeled with different identifiers that can be used to track the assay reactions and determine whether certain types of genetic information are under-represented or over-represented in the information that is generated.

In some embodiments, errors and ambiguities associated with the analysis of regions containing large numbers of sequence repeats are addressed by systematically analyzing frequencies of certain nucleic acids at particular stages in an assay (e.g., at a capture, sequencing, or detection stage). It should be appreciated that such techniques may be particularly useful in the context of a standardized protocol that is designed to allow many different loci to be evaluated in parallel without requiring different assay procedures for each locus. In some embodiments, the use of a single detection modality (e.g., sequencing) to assay multiple types of genetic lesions (e.g., point mutations, insertions/deletions, length polymorphisms) is advantageous in the clinical setting. In some embodiments of the invention, methods are provided that facilitate the use of multiple sample preparation steps in parallel, coupled with multiple analytical processes following sequence detection. Thus, in some embodiments of the invention, an improved workflow is provided that reduces error and uncertainty when simultaneously assaying different types of genetic lesions across multiple loci in multiple patients.

In some embodiments, aspects of the invention provide methods for overcoming preparative and/or analytical bias by combining two or more techniques, each having a different bias (e.g., a known bias towards under-representation or over-representation of one or more types of sequences), and using the resulting data to determine a genetic call for a subject with greater confidence.

It should be appreciated that in some embodiments, aspects of the invention relate to multiplex diagnostic methods. In some embodiments, multiplex diagnostic methods comprise capturing a plurality of genetic loci in parallel (e.g., one or more genetic loci from Table 1). In some embodiments, the genetic loci possess one or more polymorphisms (e.g., one or more polymorphisms from Table 2) the genotypes of which correspond to disease causing alleles. Accordingly, in some embodiments, the disclosure provides methods for assessing multiple heritable disorders in parallel. In some embodiments, methods are provided for diagnosing multiple heritable disorders in parallel at a pre-implantation, prenatal, perinatal, or postnatal stage. In some embodiments, the disclosure provides methods for analyzing multiple genetic loci (e.g., a plurality of target nucleic acids selected from Table 1) from a patient sample, such as a blood, pre-implantation embryo, chorionic villus or amniotic fluid sample, or other sample (e.g., other biological fluid or tissue sample such as a biopsy sample) as aspects of the invention are not limited in this respect.

Other samples may include tumor tissue or circulating tumor cells. In some embodiments, a patient sample (e.g., a tumor tissue or cell sample) is mosaic for one or more mutations of interest, and thus, may require higher sensitivity than is needed for a germline mutation analysis. In some embodiments, a sample comprises cells from a non-host organism (e.g., bacterial or viral infections in a human subject) or a sample for environmental monitoring (e.g., bacterial, viral, fungal composition of a soil, water, or air sample).

Accordingly, in some embodiments, aspects of the methods disclosed herein relate to genotyping a polymorphism of a target nucleic acid. In some embodiments, the genotyping may comprise determining that one or more alleles of the target nucleic acid are heterozygous or homozygous. In further embodiments, the genotyping may comprise determining the sequence of a polymorphism and comparing that sequence to a control sequence that is indicative of a disease risk. In some embodiments, the polymorphism is selected from a locus in Table 1 or Table 2. However, it should be appreciated that any locus associated with a disease or condition of interest may be used.

In some embodiments, a diagnosis, prognosis, or disease risk assessment is provided to a subject based on a genotype determined for that subject at one or more genetic loci (e.g., based on the analysis of a biological sample obtained from that subject). In some embodiments, an assessment is provided to a couple, based on their respective genotypes at one or more genetic loci, of the risk of their having one or more children having a genotype associated with a disease or condition (e.g., a homozygous or heterozygous genotype associated with a disease or condition). In some embodiments, a subject or a couple may seek genetic or reproductive counseling in connection with a genotype determined according to embodiments of the invention. In some embodiments, genetic information from a tumor or circulating tumor cells is used to determine prognosis and guide selection of appropriate drugs/treatments.

It should be appreciated that any of the methods or compositions described herein may be used in combination with any of the medical evaluations associated with one or more genetic loci as described herein.

In some embodiments, aspects of the invention provide effective methods for overcoming challenges associated with systematic errors (bias) and/or stochastic effects in multiplex genomic capture and/or analysis (including sequencing analysis). In some embodiments, aspects of the

invention are useful to avoid, reduce and/or account for variability in one or more sampling and/or analytical steps. For example, in some embodiments, variability in target nucleic acid representation and unequal sampling of heterozygous alleles in pools of captured target nucleic acids can be overcome.

Accordingly, in some embodiments, the disclosure provides methods that reduce variability in the detection of target nucleic acids in multiplex capture methods. In other embodiments, methods improve allelic representation in a capture pool and, thus, improve variant detection outcomes. In certain embodiments, the disclosure provides preparative methods for capturing target nucleic acids (e.g., genetic loci) that involve the use of different sets of multiple probes (e.g., molecular inversion probes MIPs) that capture overlapping regions of a target nucleic acid to achieve a more uniform representation of the target nucleic acids in a capture pool compared with methods of the prior art. In other embodiments, methods reduce bias, or the risk of bias, associated with large scale parallel capture of genetic loci, e.g., for diagnostic purposes. In other embodiments, methods are provided for increasing reproducibility (e.g., by reducing the effect of polymorphisms on target nucleic acid capture) in the detection of a plurality of genetic loci in parallel. In further embodiments, methods are provided for reducing the effect of probe synthesis and/or probe amplification variability on the analysis of a plurality of genetic loci in parallel.

According to some aspects, methods of analyzing a plurality of genetic loci are provided. In some embodiments, the methods comprise contacting each of a plurality of target nucleic acids with a probe set, wherein each probe set comprises a plurality of different probes, each probe having a central region flanked by a 5' region and a 3' region that are complementary to nucleic acids flanking the same strand of one of a plurality of subregions of the target nucleic acid, wherein the subregions of the target nucleic acid are different, and wherein each subregion overlaps with at least one other subregion, isolating a plurality of nucleic acids each having a nucleic acid sequence of a different subregion for each of the plurality of target nucleic acids, and analyzing the isolated nucleic acids.

In other embodiments, methods comprise contacting each of a plurality of target nucleic acids with a probe set, wherein each probe set comprises a plurality of different probes, each probe having a central region flanked by a 5' region and a 3' region that are complementary to nucleic acids flanking the same strand of one of a plurality of subregions of the target nucleic acid, wherein the subregions of the target nucleic acid are different, and wherein a portion of the 5' region and a portion of the 3' region of a probe have, respectively, the sequence of the 5' region and the sequence of the 3' region of a different probe, isolating a plurality of nucleic acids each having a nucleic acid sequence of a different subregion for each of the plurality of target nucleic acids, and analyzing the isolated nucleic acids.

Aspects of the disclosure are based, in part, on the discovery of methods for overcoming problems associated with systematic and random errors (bias) in genome capture, amplification and sequencing methods, namely high variability in the capture and amplification of nucleic acids and disproportionate representation of heterozygous alleles in sequencing libraries. Accordingly, in some embodiments, the disclosure provides methods that reduce errors associated with the variability in the capture and amplification of nucleic acids. In other embodiments, the methods improve allelic representation in sequencing libraries and, thus,

5

improve variant detection outcomes. In certain embodiments, the disclosure provides preparative methods for capturing target nucleic acids (e.g., genetic loci) that involve the use of differentiator tag sequences to uniquely tag individual nucleic acid molecules. In some embodiments, the differentiator tag sequence permit the detection of bias based on the occurrence of combinations of differentiator tag and target sequences observed in a sequencing reaction. In other embodiments, the methods reduce errors caused by bias, or the risk of bias, associated with the capture, amplification and sequencing of genetic loci, e.g., for diagnostic purposes.

Aspects of the invention relate to providing sequence tags (referred to as differentiator tags) that are useful to determine whether target nucleic acid sequences identified in an assay are from independently isolated target nucleic acids or from multiple copies of the same target nucleic acid molecule (e.g., due to bias in a preparative step, for example, amplification). This information can be used to help analyze a threshold number of independently isolated target nucleic acids from a biological sample in order to obtain sequence information that is reliable and can be used to make a genotype conclusion (e.g., call) with a desired degree of confidence. This information also can be used to detect bias in one or more nucleic acid preparative steps.

In some embodiments, the methods disclosed herein are useful for any application where reduction of bias, e.g., associated with genomic isolation, amplification, sequencing, is important. For example, detection of cancer mutations in a heterogeneous tissue sample, detection of mutations in maternally-circulating fetal DNA, and detection of mutations in cells isolated during a preimplantation genetic diagnostic procedure.

Accordingly, in some aspects, methods of genotyping a subject are provided. In some embodiments, the methods comprise determining the sequence of at least a threshold number of independently isolated nucleic acids, wherein the sequence of each isolated nucleic acid comprises a target nucleic acid sequence and a differentiator tag sequence, wherein the threshold number is a number of unique combinations of target nucleic acid and differentiator tag sequences, wherein the isolated nucleic acids are identified as independently isolated if they comprise unique combinations of target nucleic acid and differentiator tag sequences, and wherein the target nucleic acid sequence is the sequence of a genomic locus of a subject.

In some embodiments, the isolated nucleic acids are products of a circularization selection-based preparative method, e.g., molecular inversion probe capture products. In other embodiments, the isolated nucleic acids are products of an amplification-based preparative methods. In other embodiments, the isolated nucleic acids are products of hybridization-based preparative methods.

Circularization selection-based preparative methods selectively convert regions of interest (target nucleic acids) into a covalently-closed circular molecule which is then isolated typically by removal (usually enzymatic, e.g. with exonuclease) of any non-circularized linear nucleic acid. Oligonucleotide probes (e.g., molecular inversion probes) are designed which have ends that flank the region of interest (target nucleic acid) and, optionally, primer sites, e.g., sequencing primer sites. The probes are allowed to hybridize to the genomic target, and enzymes are used to first (optionally) fill in any gap between probe ends and second ligate the probe closed. Following circularization, any remaining (non-target) linear nucleic acid is typically removed, resulting in isolation (capture) of target nucleic acid. Circularization selection-based preparative methods include molecular

6

inversion probe capture reactions and 'selector' capture reactions. In some embodiments, molecular inversion probe capture of a target nucleic acid is indicative of the presence of a polymorphism in the target nucleic acid.

In amplification-based (e.g., PCR-based or LCR-based, etc.) preparative methods, genomic loci (target nucleic acids) are isolated directly by means of a polymerase chain reaction or ligase chain reaction (or other amplification method) that selectively amplifies each locus using one or more oligonucleotide primers. It is to be understood that primers will be sufficiently complementary to the target sequence to hybridize with and prime amplification of the target nucleic acid. Any one of a variety of art known methods may be utilized for primer design and synthesis. One or more of the primers may be perfectly complementary to the target sequence. Degenerate primers may also be used. Primers may also include additional nucleic acids that are not complementary to target sequences but that facilitate downstream applications, including for example restriction sites and differentiator tag sequences. Amplification-based methods include amplification of a single target nucleic acid and multiplex amplification (amplification of multiple target nucleic acids in parallel).

Hybridization-based preparative methods involve selectively immobilizing target nucleic acids for further manipulation. It is to be understood that one or more oligonucleotides (immobilization oligonucleotides), which comprise differentiator tag sequences, and which may be from 15 to 170 nucleotides in length, are used which hybridize along the length of a target region of a genetic locus to immobilize it. In some embodiments, immobilization oligonucleotides, are either immobilized before hybridization is performed (e.g., Roche/Nimblegen 'sequence capture'), or are prepared such that they include a moiety (e.g. biotin) which can be used to selectively immobilize the target nucleic acid after hybridization by binding to e.g., streptavidin-coated microbeads (e.g. Agilent 'SureSelect').

It should be appreciated that any of the circularization, amplification, and/or hybridization based methods described herein may be used in connection with one or more of the tiling/staggering, tagging, size-detection, and/or sensitivity enhancing algorithms described herein.

In some embodiments, the methods disclosed herein comprise determining the sequence of molecular inversion probe capture products, each comprising a molecular inversion probe and a target nucleic acid, wherein the sequence of the molecular inversion probe comprises a differentiator tag sequence and, optionally, a primer sequence, and wherein the target nucleic acid is a captured genomic locus of a subject, and genotyping the subject at the captured genomic locus based on the sequence of at least a threshold number of unique combinations of target nucleic acid and differentiator tag sequences of molecular inversion probe capture products.

In some embodiments, the methods disclosed herein comprise obtaining molecular inversion probe capture products, each comprising a molecular inversion probe and a target nucleic acid, wherein the sequence of the molecular inversion probe comprises a differentiator tag sequence and, optionally, a primer sequence, wherein the target nucleic acid is a captured genomic locus of the subject, amplifying the molecular inversion probe capture products, and genotyping the subject by determining, for each target nucleic acid, the sequence of at least a threshold number of unique combinations of target nucleic acid and differentiator tag sequence of molecular inversion probe capture products. In certain embodiments, obtaining comprises capturing target

nucleic acids from a genomic sample of the subject with molecular inversion probes, each comprising a unique differentiator tag sequence. In specific embodiments, capturing is performed under conditions wherein the likelihood of obtaining two or more molecular inversion probe capture products with identical combinations of target and differentiator tag sequences is equal to or less than a predetermined value, optionally wherein the predetermined value is about 0.05.

In one embodiment, the threshold number for a specific target nucleic acid sequence is selected based on a desired statistical confidence for the genotype. In some embodiments, the methods further comprising determining a statistical confidence for the genotype based on the number of unique combinations of target nucleic acid and differentiator tag sequences.

According to some aspects, methods of analyzing a plurality of genetic loci are provided. In some embodiments, the methods comprise obtaining a plurality of molecular inversion probe capture products each comprising a molecular inversion probe and a target nucleic acid, wherein the sequence of the molecular inversion probe comprises a differentiator tag sequence and, optionally, a primer sequence (e.g., a sequence that is complementary to the sequence of a nucleic acid that is used as a primer for sequencing or other extension reaction), amplifying the plurality of molecular inversion probe capture products, determining numbers of occurrence of combinations of target nucleic acid and differentiator tag sequence of molecular inversion probe capture products in the amplified plurality, and if the number of occurrence of a specific combination of target nucleic acid sequence and differentiator tag sequence exceeds a predetermined value, detecting bias in the amplification of the molecular inversion probe comprising the specific combination. In some embodiments, the methods further comprise genotyping target sequences in the plurality, wherein the genotyping comprises correcting for bias, if detected.

In some embodiments, the target nucleic acid is a gene (or portion thereof) selected from Table 1. In some embodiments, the genotyping comprises determining the sequence of a target nucleic acid (e.g., a polymorphic sequence) at one or more (both) alleles of a genome (a diploid genome) of a subject. In certain embodiments, the genotyping comprises determining the sequence of a target nucleic acid at both alleles of a diploid genome of a subject, wherein the target nucleic acid comprises, or consists of, a sequence of Table 1, Table 2, or other locus of interest.

In some embodiments, aspects of the invention provide methods and compositions for identifying nucleic acid insertions or deletions in genomic regions of interest without determining the nucleotide sequences of these regions. Aspects of the invention are particularly useful for detecting nucleic acid insertions or deletions in genomic regions containing nucleic acid sequence repeats (e.g., di- or trinucleotide repeats). However, the invention is not limited to analyzing nucleic acid repeats and may be used to detect insertions or deletions in any target nucleic acid of interest. Aspects of the invention are particularly useful for analyzing multiple loci in a multiplex assay.

In some embodiments, aspects of the invention relate to determining whether an amount of target nucleic acid that is captured in a genomic capture assay is higher or lower than expected. In some embodiments, a statistically significant deviation from an expected amount (e.g., higher or lower) is indicative of the presence of a nucleic acid insertion or deletion in the genomic region of interest. In some embodi-

ments, the amount is a number of nucleic acid molecules that are captured. In some embodiments, the amount is a number of independently captured nucleic acid molecules in a sample. It should be appreciated that the captured nucleic acids may be literally captured from a sample, or their sequences may be captured without actually capturing the original nucleic acids in the sample. For example, nucleic acid sequences may be captured in an assay that involves a template-based extension of nucleic acids having the region of interest, in the sample.

Aspects of the invention are based on the recognition that the efficiency of certain capture techniques is affected by the length of the nucleic acid being captured.

Accordingly, an increase or decrease in the length of a target nucleic acid (e.g., due to an insertion or deletion of a repeated sequence) can alter the capture efficiency of that nucleic acid. In some embodiments, a difference in the capture efficiency (e.g., a statistically significant difference in the capture efficiency) of a target nucleic acid is indicative of an insertion or deletion in the target nucleic acid. It should be appreciated that the capture efficiency for a target nucleic acid may be evaluated based on an amount of captured nucleic acid (e.g., number of captured nucleic acid molecules) relative to a control amount (e.g., based on an amount of control nucleic acid that is captured). However, the invention is not limited in this respect and other techniques for evaluating capture efficiency also may be used.

According to aspects of the invention, evaluating the capture efficiency as opposed to determining the sequence of the entire repeat region reduces errors associated with sequencing through repeat regions. Repeat sequences often give rise to stutters or skips in sequencing reactions that make it very difficult to accurately determine the number of repeats in a target region without running multiple sequencing reactions under different conditions and carefully analyzing the results. Such procedures are cumbersome and not readily scalable in a manner that is consistent with high throughput analyses of target nucleic acids. In some embodiments, repeat regions may be longer than the length of the individual sequence read, making length determination on the basis of a single read impossible. For example, when using next-generation sequencing the repeat regions may be longer than the length of the individual sequence read, making length determination on the basis of a single read impossible. Accordingly, aspects of the invention are useful to increase the sensitivity of detecting insertions or deletions in target regions, particularly target regions containing repeated sequences.

In some embodiments, aspects of the invention relate to capturing genomic nucleic acid sequences using a molecular inversion probe (e.g., MIP or Padlock probe) technique, and determining whether the amount (e.g., number) of captured sequences is higher or lower than expected. In some embodiments, the amount (e.g., number) of captured sequences is compared to an amount (e.g., number) of sequences captured in a control assay. The control assay may involve analyzing a control sample that contains a nucleic acid from the same genetic locus having a known sequence length (e.g., a known number of nucleic acid repeats). However, a control may involve analyzing a second (e.g., different) genetic locus that is not expected to contain any insertions or deletions. The second genetic locus may be analyzed in the same sample as the locus being interrogated or in a different sample where its length has been previously determined. The second genetic locus may be a locus that is not char-

acterized by the presence of nucleic acid repeats (and thus not expected to contain insertions or deletions of the repeat sequence).

In some embodiments, a target nucleic acid region that is being evaluated may be determined by the identity of the targeting arms of a probe that is designed to capture the target region (or sequence thereof). For example, the targeting arms of a MIP probe may be designed to be complementary (e.g., sufficiently complementary for selective hybridization and/or polymerase extension and/or ligation) to genomic regions flanking a target region suspected of containing an insertion or deletion. It should be appreciated that two targeting arms may be designed to be complementary (e.g., sufficiently complementary for selective hybridization and/or polymerase extension and/or ligation) to the two flanking regions that are immediately adjacent (e.g., immediately 5' and 3', respectively) to a region of a sequence repeat on one strand of a genomic nucleic acid. However, one or both targeting arms may be designed to hybridize several bases (e.g., 1-5, 5-10, 10-25, 25-50, or more) upstream or downstream from the repeat region in such a way that the captured sequence includes a region of unique genomic sequence that on one or both sides of the repeat region. This unique region can then be used to identify the captured target (e.g., based on sequence or hybridization information).

In some embodiments, two or more (e.g., 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) different loci may be interrogated in parallel in a single assay (e.g., in a multiplex assay). In some embodiments, the ratio of captured nucleic acids for each locus may be used to determine whether a nucleic acid insertion or deletion is present in one locus relative to the other. For example, the ratio may be compared to a control ratio that is representative of the two loci when neither one has an insertion or deletion relative to control sequences (e.g., sequences that are normal or known to be associated with healthy phenotypes for those loci). However, the amount of captured nucleic acids may be compared to any suitable control as discussed herein.

The locus of a captured sequence may be identified by determining a portion of unique sequence 5' and/or 3' to the repeat region in the target nucleic acid suspected of containing a deletion or insertion. This does not require sequencing the captured repeat region itself. However, some or all of the repeat region also could be sequenced as aspects of the invention are not limited in this respect.

Aspects of the invention may be combined with one or more sequence-based assays (e.g., SNP detection assays), for example in a multiplex format, to determine the genotype of one or more regions of a subject.

In some embodiments, methods of detecting a polymorphism in a nucleic acid in a biological sample are provided. In some embodiments, the methods comprise evaluating the efficiency of capture at one or more loci and determining whether one or both alleles at that locus contain an insertion or deletion relative to a control locus (e.g., a locus indicative of a length of repeat sequence that is associated with a healthy phenotype).

Accordingly, aspects of the invention relate to methods for determining whether a target nucleic acid has an abnormal length by evaluating the capture efficiency of a target nucleic acid in a biological sample from a subject, wherein a capture efficiency that is different from a reference capture efficiency is indicative of the presence, in the biological sample, of a target nucleic acid having an abnormal length. It should be appreciated that the term "abnormal" is a relative term based on a comparison to a "normal" length. In

some embodiments, a normal length is a length that is associated with a normal (e.g., healthy or non-carrier phenotype). Accordingly, an abnormal length is a length that is either shorter or longer than the normal length. In some embodiments, the presence of an abnormal length is indicative of an increased risk that the locus is associated with a disease or a disease carrier phenotype. In some embodiments, the abnormal length is indicative that the subject is either has a disease or condition or is a carrier of a disease or condition (e.g., associated with the locus). However, it should be appreciated that the description of embodiments relating to detecting the presence of an abnormal length also support detecting the presence of a length that is different from an expected or control length.

In some embodiments, aspects of the invention relate to estimating the length of a target nucleic acid (e.g., of a sub-target region within a target nucleic acid). In some embodiments, aspects of the invention relate to methods for estimating the length of a target nucleic acid by contacting the target nucleic acid with a plurality of detection probes under conditions that permit hybridization of the detection probes to the target nucleic acid, wherein each detection probe is a polynucleotide that comprises a first arm that hybridizes to a first region of the target nucleic acid and a second arm that hybridizes to a second region of the target nucleic acid, wherein the first and second regions are on a common strand of the target nucleic acid, and wherein the nucleotide sequence of the target between the 5' end of the first region and the 3' end of the second region is the nucleotide sequence of a sub-target nucleic acid; and capturing a plurality of sub-target nucleic acids that are hybridized with the plurality of detection probes; and measuring the frequency of occurrence of a sub-target nucleic acid in the plurality of sub-target nucleic acids, wherein the frequency of occurrence of the sub-target nucleic acid in the plurality of sub-target nucleic acids is indicative of the length of the sub-target nucleic acid. It should be appreciated that methods for estimating a nucleic acid length may involve comparing a capture efficiency for a target nucleic acid region to two or more reference efficiencies for known nucleic acid lengths in order to determine whether the target nucleic acid region is smaller, intermediate, or larger in size than the known control lengths. In some embodiments, a series of nucleic acids of known different lengths may be used to provide a calibration curve for evaluating the length of a target nucleic acid region of interest.

In some embodiments, the capture efficiency of a target region suspected of having a deletion or insertion is determined by comparing the capture efficiency to a reference indicative of a normal capture efficiency. In some embodiments, the capture efficiency is lower than the reference capture efficiency. In some embodiments, the subject is identified as having an insertion in the target region. In some embodiments, the capture efficiency is higher than the reference capture efficiency. In some embodiments, the subject is identified as having a deletion in the target region. In some embodiments, the subject is identified as being heterozygous for the insertion. In some embodiments, the subject is identified as being heterozygous for the deletion.

In some embodiments of any of the methods described herein (e.g., tiling/staggering, tagging, size-detection, and/or sensitivity enhancement) aspects of the invention relate to capturing a sub-target nucleic acid (or a sequence of a sub-target nucleic acid). In some embodiments, a molecular inversion probe technique is used. In some embodiments, a molecular inversion probe is a single linear strand of nucleic acid that comprises a first targeting arm at its 5' end and a

second targeting arm at its 3' end, wherein the first targeting arm is capable of specifically hybridizing to a first region flanking one end of the sub-target nucleic acid, and wherein the second targeting arm is capable of specifically hybridizing to a second region flanking the other end of the sub target nucleic acid on the same strand of the target nucleic acid. In some embodiments, the first and second targeting arms are between about 10 and about 100 nucleotides long. In some embodiments, the first and second targeting arms are about 10-20, 20-30, 30-40, or 40-50 nucleotides long. In some embodiments, the first and second targeting arms are about 20 nucleotides long. In some embodiments, the first and second targeting arms have the same length. In some embodiments, the first and second targeting arms have different lengths. In some embodiments, each pair of first and second targeting arms in a set of probes has the same length. Accordingly, if one of the targeting arms is longer, the other one is correspondingly shorter. This allows for a quality control step in some embodiments to confirm that all captured probe/target sequence products have the same length after a multiplexed plurality of capture reactions. In some embodiments, a set of probes may be designed to have the same length if the intervening region is varied to accommodate any differences in the length of either one or both of the first and second targeting arms.

In some embodiments, the hybridization Tms of the first and second targeting arms are similar. In some embodiments, the hybridization Tms of the first and second targeting arms are within 2-5° C. of each other. In some embodiments, the hybridization Tms of the first and second targeting arms are identical. In some embodiments, the hybridization Tms of the first and second targeting arms are close to empirically-determined optima but not necessarily identical.

In some embodiments, the first and second targeting arms of a molecular inversion probe have different Tms. For example, the Tm of the first targeting arm (at the 5' end of the molecular inversion probe) may be higher than the Tm of the second targeting arm (at the 3' end of the molecular inversion probe). According to aspects of the invention, and without wishing to be bound by theory, a relatively high Tm for the first targeting arm may help avoid or prevent the first targeting arm from being displaced after hybridization by the extension product of the 3' end of the second targeting arm. It should be appreciated that a reference to the Tm of a targeting arm as used herein relates to the Tm of hybridization of the targeting arm to a nucleic acid having the complementary sequence (e.g., the region of the target nucleic acid that has a sequence that is complementary to the sequence of the targeting arm). It also should be appreciated that the Tms of the targeting arms described herein may be calculated using any appropriate method. For example, in some embodiments an experimental method (e.g., a gel shift assay, a hybridization assay, a melting curve analysis, for example in a PCR machine with a SYBR dye by stepping through a temperature ramp while monitoring signal level from an intercalating dye, for example, bound to a double-stranded DNA, etc.) may be used to determine one or more Tms empirically. In some embodiments, an optimal Tm may be determined by evaluating the number of products formed (e.g., for each of a plurality of MIP probes), and determining the optimal Tm as the center point in a histogram of Tm for all targeting arms. In some embodiments, a predictive algorithm may be used to determine a Tm theoretically. In some embodiments, a relatively simple predictive algorithm may be used based on the number of G/C and A/T base pairs when the sequence is hybridized to its target and/or the

length of the hybridized product (e.g., for example, $64.9 + 41 * ([G+C] - 16.4) / (A+T+G+C)$, see for example, Wallace, R. B., Shaffer, J., Murphy, R. F., Bonner, J., Hirose, T., and Itakura, K. (1979) *Nucleic Acids Res* 6:3543-3557). In some embodiments, a more complex algorithm may be used to account for the effects of base stacking entropy and enthalpy, ion concentration, and primer concentration (see, for example, SantaLucia J (1998), *Proc Natl Acad Sci USA*, 95:1460-5). In some embodiments an algorithm may use modified parameters (e.g., nearest-neighbor parameters for basepair entropy/enthalpy values). It should be appreciated that any suitable algorithm may be used as aspects of the invention are not limited in this respect. However, it also should be appreciated that different methodologies may results in different calculated or predicted Tms for the same sequences. Accordingly, in some embodiments, the same empirical and/or theoretical method is used to determine the Tms of different sequences for a set of probes to avoid a negative impact of any systematic difference in the Tm determination or prediction when designing a set of probes with predetermined similarities or differences for different Tms.

In some embodiments, the Tm of the first targeting arm may be about 1° C., about 2° C., about 3° C., about 4° C., about 5° C., or more than about 5° C. higher than the Tm of the second targeting arm. In some embodiments, each probe in a plurality of probes (e.g., each probe in a set of 5-10, each probe in a set of at least 10, each probe in a set of 10-50, each probe in a set of 50-100, each probe in a set of 100-500, each probe in a set of 500-1,000, each probe in a set of 1,000-1,500, each probe in a set of 1,500-2,000, each probe in a set of 2,000-3,000, 3,000-5,000, 5,000-10,000 or each probe in a set of at least 5,000 different probes) has a unique first targeting arm (e.g., they all have different sequences) and a unique second targeting arm (e.g., they all have different sequences). In some embodiments, for at least 10% of the probes (e.g., at least 25%, 25%-50%, 50%-75%, 75%-90%, 90%-95% or over 95%, or all of the probes) the first targeting arm has a Tm for its complementary sequence that is higher (e.g., about 1° C., about 2° C., about 3° C., about 4° C., about 5° C., or more than about 5° C. higher) than the Tm of the second targeting arm for its complementary sequence. In some embodiments, each of the first targeting arms have similar or identical Tms for their respective complementary sequences and each of the second targeting arms have similar or identical Tms for their respective complementary sequences (and the first targeting arms have higher Tms than the second targeting arms). For example, in some embodiments, the Tm of the first arm(s) may be about 58° C. and the Tm of the second arm(s) may be about 56° C. In some embodiments, the Tm of the first arm(s) may be about 68° C., and the Tm of the second arm(s) may be about 65° C. It should be appreciated that in some embodiments the similarity (e.g., within a range of 1° C., 2° C., 3° C., 4° C., 5° C.) or identity of the Tms for the different targeting arms should be based either on empirical data for each arm or based on the same predictive algorithm for each arm (e.g., Wallace, R. B., Shaffer, J., Murphy, R. F., Bonner, J., Hirose, T., and Itakura, K. (1979) *Nucleic Acids Res* 6:3543-3557, SantaLucia J (1998), *Proc Natl Acad Sci USA*, 95:1460-5, or other algorithm).

In some embodiments, the Tm of the first targeting arm of a molecular inversion probe (at the 5' end of the molecular inversion probe) is selected to be sufficiently stable to prevent displacement of the first targeting arm from its complementary sequence on a target nucleic acid. In some embodiments, the Tm of the first targeting arm is 50-55° C.,

13

at least 55° C., 55-60° C., at least 60° C., 60-65° C., at least 65° C., at least 70° C., at least 75° C., or at least 80° C. As discussed above, it should be appreciated that the for a particular targeting arm may be determined empirically or theoretically. Different theoretical models may be used to determine a T_m and it should be appreciated that the predicted T_m for a particular sequence may be different depending on the algorithm used for the prediction. In some embodiments, each probe in a plurality of probes (e.g., each probe in a set of 5-10, each probe in a set of at least 10, each probe in a set of 10-50, each probe in a set of 50-100, each probe in a set of 100-500, or each probe in a set of at least 500 different probes) has a different first targeting arm (e.g., different sequences) but each different first targeting arm has a similar or identical T_m for its complementary sequence on a target nucleic acid. It should be appreciated that in some embodiments the similarity (e.g., within a range of 1 C, 2 C, 3 C, 4 C, 5 C) or identity of the T_ms for the different targeting arms should be based either on empirical data for each arm or based on the same predictive algorithm for each arm (e.g., Wallace, R. B., Shaffer, J., Murphy, R. F., Bonner, J., Hirose, T., and Itakura, K. (1979) *Nucleic Acids Res* 6:3543-3557, SantaLucia J (1998), *Proc Natl Acad Sci USA*, 95:1460-5, or other algorithm).

In some embodiments, the sub-target nucleic acid contains a nucleic acid repeat. In some embodiments, the nucleic acid repeat is a dinucleotide or trinucleotide repeat. In some embodiments, the sub-target nucleic acid contains 10-100 copies of the nucleic acid repeat in the absence of an abnormal increase or decrease in nucleic acid repeats. In some embodiments, the sub-target nucleic acid is a region of the Fragile-X locus that contains a nucleic acid repeat. In some embodiments, one or both targeting arms hybridize to a region on the target nucleic acid that is immediately adjacent to a region of nucleic acid repeats. In some embodiments, one or both targeting arms hybridize to a region on the target nucleic acid that is separated from a region of nucleic acid repeats by a region that does not contain any nucleic acid repeats. In some embodiments, the molecular inversion probe further comprises a primer-binding region that can be used to sequence the captured sub-target nucleic acid and optionally the first and/or second targeting arm.

In some embodiments, aspects of the invention relate to evaluating the length of a plurality of different target nucleic acids in a biological sample. In some embodiments, the plurality of target nucleic acids are analyzed using a plurality of different molecular inversion probes. In some embodiments, each different molecular inversion probe comprises a different pair of first and second targeting arms at each of the 3' and 5' ends. In some embodiments, each different molecular inversion probe comprises the same primer-binding sequence.

In some embodiments, aspects of the invention relate to analyzing nucleic acid from a biological sample obtained from a subject. In some embodiments, the biological sample is a blood sample. In some embodiments, the biological sample is a tissue sample, specific cell population, tumor sample, circulating tumor cells, or environmental sample. In some embodiments, the biological sample is a single cell. In some embodiments, nucleic acids are analyzed in biological samples obtained from a plurality of different subjects. In some embodiments, nucleic acids from a biological sample are analyzed in multiplex reactions. It should be appreciated that a biological sample contains a plurality of copies of a genome derived from a plurality of cells in the sample. Accordingly, a sample may contain a plurality of indepen-

14

dent copies of a target nucleic acid region of interest, the capture efficiency of which can be used to evaluate its size as described herein.

In some embodiments, aspects of the invention relate to evaluating a nucleic acid capture efficiency by determining an amount of target nucleic acid that is captured (e.g., an amount of sub-target nucleic acid sequences that are captured). In some embodiments, the amount of target nucleic acid that is captured is determined by determining a number of independently captured target nucleic acid molecules (e.g., the amount of independently captured molecules that have the sequence of the sub-target region). In some embodiments, the amount of target nucleic acid that is captured is compared to a reference amount of captured nucleic acid. In some embodiments, the reference amount is determined by determining a number of independently captured molecules of a reference nucleic acid. In some embodiments, the reference nucleic acid is a nucleic acid of a different locus in the biological sample that is not suspected of containing a deletion or insertion. In some embodiments, the reference nucleic acid is a nucleic acid of known size and amount that is added to the capture reaction. As described herein, a number of independently captured nucleic acid sequences can be determined by contacting a nucleic acid sample with a preparation of a probe (e.g., a MIP probe as described herein). It should be appreciated that the preparation may comprise a plurality of copies of the same probe and accordingly a plurality of independent copies of the target region may be captured by different probe molecules. The number of probe molecules that actually capture a sequence can be evaluated by determining an amount or number of captured molecules using any suitable technique. This number is a reflection of both the number of target molecules in the sample and the efficiency of capture of those target molecules, which in turn is related to the size of the target molecules as described herein. Accordingly, the capture efficiency can be evaluated by controlling for the abundance of the target nucleic acid, for example by comparing the number or amount of captured target molecules to an appropriate control (e.g., a known size and amount of control nucleic acid, or a different locus that should be present in the same amount in the biological sample and is not expected to contain any insertions or deletions). It should be appreciated that other factors may affect the capture efficiency of a particular target nucleic acid region (e.g., the sequence of the region, the GC content, the presence of secondary structures, etc.). However, these factors also can be accounted for by using appropriate controls (e.g., known sequences having similar properties, the same sequences, other genomic sequences expected to be present in the biological sample at the same frequency, etc., or any combination thereof).

In some embodiments, aspects of the invention relate to identifying a subject as having an insertion or deletion in one or more alleles of a genetic locus if the capture efficiency for that genetic locus is statistically significantly different than a reference capture efficiency.

It should be appreciated that hybridization conditions used for any of the capture techniques described herein (e.g., MIP capture techniques) can be based on known hybridization buffers and conditions.

In some embodiments, the methods disclosed herein are useful for any application where the detection of deletions or insertions is important.

In some embodiments, aspects of the invention relate to basing a nucleic acid sequence analysis on results from two or more different nucleic acid preparatory techniques that

15

have different systematic biases in the types of nucleic acids that they sample. According to the invention, different techniques have different sequence biases that are systematic and not simply due to stochastic effects during nucleic acid capture or amplification. Accordingly, the degree of over-sampling required to overcome variations in nucleic acid preparation needs to be sufficient to overcome the biases (e.g., an oversampling of 2-5 fold, 5-10 fold, 5-15 fold, 15-20 fold, 20-30 fold, 30-50 fold, or intermediate to higher fold).

According to some embodiments, different techniques have different characteristic or systematic biases. For example, one technique may bias a sample analysis towards one particular allele at a genetic locus of interest, whereas a different technique would bias the sample analysis towards a different allele at the same locus. Accordingly, the same sample may be identified as being different depending on the type of technique that is used to prepare nucleic acid for sequence analysis. This effectively represents a sensitivity limitation, because each technique has different relative sensitivities for polymorphic sequences of interest.

According to aspects of the invention, the sensitivity of a nucleic acid analysis can be increased by combining the sequences from different nucleic acid preparative steps and using the combined sequence information for a diagnostic assay (e.g., for a making a call as to whether a subject is homozygous or heterozygous at a genetic locus of interest).

In some embodiments, the invention provides a method of increasing the sensitivity of a nucleic acid detection assay by obtaining a first preparation of a target nucleic acid using a first preparative method on a biological sample, obtaining a second preparation of a target nucleic acid using a second preparative method on the biological sample, assaying the sequences obtained in both first and second nucleic acid preparations, and using the sequence information from both first and second nucleic acid preparations to determine the genotype of the target nucleic acid in the biological sample, wherein the first and second preparative methods have different systematic sequence biases. In some embodiments, the first and second nucleic acid preparations are combined prior to performing a sequence assay. In some embodiments, separate sequence assays are performed on the first and second nucleic acid preparations and the sequence information from both assays are combined to determine the genotype of the target nucleic acid in the biological sample. In some embodiments, the first preparative method is an amplification-based, a hybridization-based, or a circular probe-based preparative method. In some embodiments, the second method is an amplification-based, a hybridization-based, or a circular probe-based preparative method. In some embodiments, the first and second methods are of different types (e.g., only one of them is an amplification-based, a hybridization-based, or a circular probe-based preparative method, and the other one is one or the other two types of method). Accordingly, in some embodiments the second preparative method is an amplification-based, a hybridization-based, or a circular probe-based preparative method, provided that the second method is different from the first method. However, in some embodiments, both methods may be of the same type, provided they are different methods (e.g., both are amplification based or hybridization-based, but are different types of amplification or hybridization methods, e.g., with different relative biases).

In amplification-based (e.g., PCR-based or LCR-based, etc.) preparative methods, genomic loci (target nucleic acids) are isolated directly by means of a polymerase chain reaction or ligase chain reaction (or other amplification

16

method) that selectively amplifies each locus using a pair of oligonucleotide primers. It is to be understood that primers will be sufficiently complementary to the target sequence to hybridize with and prime amplification of the target nucleic acid. Any one of a variety of art known methods may be utilized for primer design and synthesis. One or both of the primers may be perfectly complementary to the target sequence. Degenerate primers may also be used. Primers may also include additional nucleic acids that are not complementary to target sequences but that facilitate downstream applications, including for example restriction sites and identifier sequences (e.g., source sequences). PCR based methods may include amplification of a single target nucleic acid and multiplex amplification (amplification of multiple target nucleic acids in parallel).

Hybridization-based preparative methods may involve selectively immobilizing target nucleic acids for further manipulation. It is to be understood that one or more oligonucleotides (immobilization oligonucleotides), which in some embodiments may be from 10 to 200 nucleotides in length, are used which hybridize along the length of a target region of a genetic locus to immobilize it. In some embodiments, immobilization oligonucleotides are either immobilized before hybridization is performed (e.g., Roche/Nimblegen 'sequence capture'), or are prepared such that they include a moiety (e.g., biotin) which can be used to selectively immobilize the target nucleic acid after hybridization by binding to e.g., streptavidin-coated microbeads (e.g., Agilent 'SureSelect').

Circularization selection-based preparative methods selectively convert each region of interest into a covalently-closed circular molecule which is then isolated by removal (usually enzymatic, e.g., with exonuclease) of any non-circularized linear nucleic acid. Oligonucleotide probes are designed which have ends that flank the region of interest. The probes are allowed to hybridize to the genomic target, and enzymes are used to first (optionally) fill in any gap between probe ends and second ligate the probe closed. In some embodiments, following circularization, any remaining (non-target) linear nucleic acid can be removed, resulting in isolation (capture) of target nucleic acid. Circularization selection-based preparative methods include molecular inversion probe capture reactions and 'selector' capture reactions. However, other techniques may be used as aspects of the invention are not limited in this respect. In some embodiments, molecular inversion probe capture of a target nucleic acid is indicative of the presence of a polymorphism in the target nucleic acid.

A variety of methods may be used to evaluate and compare bias profiles of each preparative technique. Next-generation sequencing may be used to quantitatively measure the abundance of each isolated target nucleic acid obtained from a certain preparative method. This abundance may be compared to a control abundance value (e.g., a known starting abundance of the target nucleic acid) and/or with an abundance determined through the use of an alternative preparative method. For example, a set of target nucleic acids may be isolated by one or more of the three preparative methods; the target nucleic acid may be observed x times using the amplification technique, y times using the hybridization enrichment technique, and z times using the circularization selection technique. A pairwise correlation coefficient may be computed between each abundance value (e.g., x and y, x and z, and y and z) to assess bias in nucleic acid isolation between pairs of preparative methods. Since the mechanisms of isolation are different in each

approach, the abundances will usually be different and largely uncorrelated with each other.

In some embodiments, the invention provides a method of obtaining a nucleic acid preparation that is representative of a target nucleic acid in a biological sample by obtaining a first preparation of a target nucleic acid using a first preparative method on a biological sample, obtaining a second preparation of a target nucleic acid using a second preparative method on the biological sample, and combining the first and second nucleic acid preparations to obtain a combined preparation that is representative of the target nucleic acid in the biological sample.

In some embodiments of any of the methods described herein, a third preparation of the target nucleic acid is obtained using a third preparative method that is different from the first and second preparative methods, wherein the first, second, and third preparative methods all have different systematic sequence biases. In some embodiments of any of the methods described herein, the different preparative methods are used for a plurality of different loci in the biological sample to increase the sensitivity of a multiplex nucleic acid analysis. In some embodiments, the target nucleic acid has a sequence of a gene selected from Table 1.

However, it should be appreciated that a genotyping method of the invention may include several steps, each of which independently may involve one or more different preparative techniques described herein. In some embodiments, a nucleic acid preparation may be obtained using one or more (e.g., 2, 3, 4, 5, or more) different techniques described herein (e.g., amplification, hybridization capture, circular probe capture, etc., or any combination thereof) and the nucleic acid preparation may be analyzed using one or more different techniques (e.g., amplification, hybridization capture, circular probe capture, etc., or any combination thereof) that are selected independently of the techniques used for the initial preparation.

In some embodiments, aspects of the invention also provide compositions, kits, devices, and analytical methods for increasing the sensitivity of nucleic acid assays. Aspects of the invention are particularly useful for increasing the confidence level of genotyping analyses. However, aspects of the invention may be used in the context of any suitable nucleic acid analysis, for example, but not limited to, a nucleic acid analysis that is designed to determine whether more than one sequence variant is present in a sample.

In some embodiments, aspects of the invention relate to a plurality of nucleic acid probes (e.g., 10-50, 50-100, 100-250, 250-500, 500-1,000, 1,000-2,000, 2,000-5,000, 5,000-7,500, 7,500-10,000, or lower, higher, or intermediate number of different probes).

In some embodiments, each probe or each of a subset of probes (e.g., 10-25%, 25-50%, 50-75%, 75-90%, or 90-99%) has a different first targeting arm. In some embodiments, each probe or each probe of a subset of probes (e.g., 10-25%, 25-50%, 50-75%, 75-90%, or 90-99%) has a different second targeting arm. In some embodiments, the first and second targeting arms are separated by the same intervening sequence. In some embodiments, the first and second targeting arms are complementary to target nucleic acid sequences that are separated by the same or a similar length (e.g., number of nucleic acids, for example, 0-25, 25-50, 50-100, 100-250, 250-500, 500-1,000, 1,000-2,500 or longer or intermediate number of nucleotides) on their respective target nucleic acids (e.g., genomic loci). In some embodiments, each probe or a subset of probes (e.g., 10-30 25%, 25-50%, 50-75%, 75-90%, or 90-99%) includes a first primer binding sequence. In some embodiments, the primer

binding sequence is the same (e.g., it can be used to prime sequencing or other extension reaction). In some embodiments, each probe or a subset of probes (e.g., 10-25%, 25-50%, 50-75%, 75-90%, or 90-99%) includes a unique identifier sequence tag (e.g., that is predetermined and can be used to distinguish each probe).

In some embodiments, the methods disclosed herein are useful for any application where sensitivity is important. For example, detection of cancer mutations in a heterogeneous tissue sample, detection of mutations in maternally-circulating fetal DNA, and detection of mutations in cells isolated during a preimplantation genetic diagnostic procedure.

According to some aspects of the invention, methods of detecting a polymorphism in a nucleic acid in a biological sample are provided. In some embodiments, the methods comprise obtaining a nucleic acid preparation using a preparative method (e.g., any of the preparative methods disclosed herein) on a biological sample, and performing a molecular inversion probe capture reaction on the nucleic acid preparation, wherein a molecular inversion probe capture (e.g., using a mutation-detection MIP) of a target nucleic acid of the nucleic acid preparation is indicative of the presence of a mutation (polymorphism) in the target nucleic acid, optionally wherein the polymorphism is selected from Table 2.

According to some aspects of the invention, methods of genotyping a nucleic acid in a biological sample are provided. In some embodiments, the methods comprise obtaining a nucleic acid preparation using a preparative method on a biological sample, sequencing a target nucleic acid of the nucleic acid preparation, and performing a molecular inversion probe capture reaction on the biological sample, wherein a molecular inversion probe capture of the target nucleic acid in the biological sample is indicative of the presence of a polymorphism in the target nucleic acid, genotyping the target nucleic acid based on the results of the sequencing and the capture reaction.

In some embodiments of the methods disclosed herein, the target nucleic acid has a sequence of a gene selected from Table 1.

It should be appreciated that any one or more embodiments described herein may be used for evaluating multiple genetic markers in parallel. Accordingly, in some embodiments, aspects of the invention relate to determining the presence of one or more markers (e.g., one or more alleles) at multiple different genetic loci in parallel.

Accordingly, the risk or presence of multiple heritable disorders may be evaluated in parallel. In some embodiments, the risk of having offspring with one or more heritable disorders may be evaluated. In some embodiments, an evaluation may be performed on a biological sample of a parent or a child (e.g., at a pre-implantation, prenatal, perinatal, or postnatal stage).

In some embodiments, the disclosure provides methods for analyzing multiple genetic loci (e.g., a plurality of target nucleic acids selected from Table 1 or 2) from a patient sample, such as a blood, pre-implantation embryo, chorionic villus or amniotic fluid sample. A patient or subject may be a human. However, aspects of the invention are not limited to humans and may be applied to other species (e.g., mammals, birds, reptiles, other vertebrates or invertebrates) as aspects of the invention are not limited in this respect. A subject or patient may be male or female. In some embodiments, in connection with reproductive genetic counseling, samples from a male and female member of a couple may be analyzed. In some embodiments, for example, in connection with an animal breeding program, samples from a plurality

of male and female subjects may be analyzed to determine compatible or optimal breeding partners or strategies for particular traits or to avoid one or more diseases or conditions. Accordingly, reproductive risks may be determined and/or reproductive recommendations may be provided based on information derived from one or more embodiments of the invention.

However, it should be appreciated that aspects of the invention may be used in connection with any medical evaluation where the presence of one or more alleles at a genetic locus of interest is relevant to a medical determination (e.g., risk or detection of disease, disease prognosis, therapy selection, therapy monitoring, etc.). Further aspects of the invention may be used in connection with detection, in tumor tissue or circulating tumor cells, of mutations in cellular pathways that cause cancer or predict efficacy of treatment regimens, or with detection and identification of pathogenic organisms in the environment or a sample obtained from a subject, e.g., a human subject.

These and other aspects of the invention are described in more detail in the following description and non-limiting examples and drawings.

BRIEF DESCRIPTION OF DRAWINGS

FIG. 1 illustrates a non-limiting embodiment of a tiled probe layout;

FIG. 2 illustrates a non-limiting embodiment of a staggered probe layout;

FIG. 3 illustrates a non-limiting embodiment of an alternating staggered probe layout;

FIGS. 4A-C depict various non-limiting methods for combining differentiator tag sequence and target sequences (NNNN depicts a differentiator tag sequence);

FIG. 5 depicts a non-limiting method for genotyping based on target and 5 differentiator tag sequences;

FIG. 6 depicts non-limiting results of a simulation of a MIP capture reaction;

FIG. 7 depicts a non-limiting graph of sequencing coverage;

FIG. 8 illustrates that shorter sequences are captured with higher efficiency than longer sequences using MIPs;

FIG. 9 illustrates a non-limiting scheme of padlock (MIP) capture of a region that includes both repetitive regions (thick wavy line) and the adjacent unique sequence (thick straight line);

FIG. 10 illustrates a non-limiting hypothetical relationship between target gap size and the relative number of reads of the repetitive region;

FIG. 11A depicts MIP capture of FMR1 repeat regions from a diploid genome;

FIG. 11B depicts preparative methods for biallelic resolution of FMR1 repeat region lengths in a diploid genome using MIP capture probes and unique differentiator tags;

FIG. 11C depicts an analysis of FMR1 repeat region lengths in a diploid genome;

FIG. 12 is a schematic of an embodiment of an algorithm of the invention;

FIG. 13 illustrates a non-limiting example of a graph of per-target abundance with MIP capture; and,

FIG. 14 shows a non-limiting a graph of correlation between two MIP capture reactions.

DETAILED DESCRIPTION

Aspects of the invention relate to preparative and analytical methods and compositions for evaluating genotypes,

and in particular, for determining the allelic identity (or identities in a diploid organism) of one or more genetic loci in a subject.

Aspects of the invention are based, in part, on the identification of different sources of ambiguity and error in genetic analyses, and, in part, on the identification of one or more approaches to avoid, reduce, recognize, and/or resolve these errors and ambiguities at different stages in a genetic analysis. Aspects of the invention relate to methods and compositions for addressing bias and/or stochastic variation associated with one or more preparative and/or analytical steps of a nucleic acid evaluation technology. In some embodiments, preparative methods can be adapted to avoid or reduce the risk of bias skewing the results of a genetic analysis. In some embodiments, analytical methods can be adapted to recognize and correct for data variations that may give rise to misinterpretation (e.g., incorrect calls such as homozygous when the subject is actually heterozygous or heterozygous when the subject is actually homozygous). Methods of the invention may be used for any type of mutation, for example a single base change (e.g., insertion, deletion, transversion or transition, etc.), a multiple base insertion, deletion, duplication, inversion, and/or any other change or combination thereof.

In some embodiments, additional or alternative techniques may be used to address loci characterized by multiple repeats of a core sequence where the length of the repeat is longer than a typical sequencing read thereby making it difficult to determine whether a deletion or duplication of one or more core sequence units has occurred based solely on a sequence read.

In some embodiments, increased confidence in an assay result may be obtained by i) selecting two or more different preparative and/or analytical techniques that have different biases (e.g., known to have different biases), ii) evaluating a patient sample using the two or more different techniques, iii) comparing the results from the two or more different techniques, and/or iv) determining whether the results are consistent for the two or more different techniques. In some embodiments, if determining in step (iv) indicates that the results are consistent (e.g., the same) then increased confidence in the assay result is obtained. In other embodiments, if determining in step (iv) indicates that the results are inconsistent (e.g., that the results are ambiguous) then one or more additional preparative and/or analytical techniques, which have a different bias (e.g., known to have a different bias) compared with the two or more different preparative and/or analytical techniques selected in step (i), are used to evaluate the patient sample, and the results of the one or more additional preparative and/or analytical techniques are compared with the results from step (ii) to resolve the inconsistency.

In some embodiments, two or more independent samples may be obtained from a subject and independently analyzed. In some embodiments, two or more independent samples are obtained at approximately the same time point. In some embodiments, two or more independent samples are obtained at multiple different time points. In some embodiments, the use of two or more independent sample facilitates the elimination, normalization, and/or quantification of stochastic measurement noise. It is to be appreciated that two or more independent samples may be obtained in connection with any of the methods disclosed herein, including, for example, methods for pathogen profiling in a human or other animal subjects, monitoring tumor progression/regression,

analyzing circulating tumor cells, analyzing fetal cells in maternal circulation, and analyzing/monitoring/profiling of environmental pathogens.

In some embodiments, one or more of the techniques described herein may be combined in a single assay protocol for evaluating multiple patient samples in parallel.

It should be appreciated that aspects of the invention may be useful for high throughput, cost-effective, yet reliable, genotyping of multiple patient samples (e.g., in parallel, for example in multiplex reactions). In some embodiments, aspects of the invention are useful to reduce the error frequency in a multiplex analysis. Certain embodiments may be particularly useful where multiple reactions (e.g., multiple loci and/or multiple patient samples) are being processed. For example, 10-25, 25-50, 50-75, 75-100 or more loci may be evaluated for each subject out of any number of subject samples that may be processed in parallel (e.g., 1-25, 25-50, 50-100, 100-500, 500-1,000, 1,000-2,500, 2,500-5,000 or more or intermediate numbers of patient samples). It should be appreciated that different embodiments of the invention may involve conducting two or more target capture reactions and/or two or more patient sample analyses in parallel in a single multiplex reaction. For example, in some embodiments a plurality of capture reactions (e.g., using different capture probes for different target loci) may be performed in a single multiplex reaction on a single patient sample. In some embodiments, a plurality of captured nucleic acids from each one of a plurality of patient samples may be combined in a single multiplex analysis reaction. In some embodiments, samples from different subjects are tagged with subject-specific (e.g., patient-specific) tags (e.g., unique sequence tags) so that the information from each product can be assigned to an identified subject. In some embodiments, each of the different capture probes used for each patient sample have a common patient-specific tag. In some embodiments, the capture probes do not have patient-specific tags, but the captured products from each subject may be amplified using one or a pair of amplification primers that are labeled with a patient-specific tag. Other techniques for associating a patient-specific tag with the captured product from a single patient sample may be used as aspects of the invention are not limited in this respect. It should be appreciated that patient-specific tags as used herein may refer to unique tags that are assigned to identified patients in a particular assay. The same tags may be used in a separate multiplex analysis with a different set of patient samples (e.g., from different patients) each of which is assigned one of the tags. In some embodiments, different sets of unique tags may be used in sequential (e.g., alternating) multiplex reactions in order to reduce the risk of contamination from one assay to the next and allow contamination to be detected on the basis of the presence of tags that are not expected to be present in a particular assay.

Embodiments of the invention may be used for any of a number of different settings: reproductive settings, disease screening, identifying subjects having cancer, identifying subjects having increased risk for a disease, stratifying a population of subjects according to one or more of a number of factors, for example responsiveness to a particular drug, lack or not of an adverse reaction (or risk therefore) to a particular drug, and/or providing information for medical records (e.g., homozygosity, heterozygosity at one or more loci). It should be appreciated that the invention is not limited to genomic analysis of patient samples. For example, aspects of the invention may be useful for high throughput genetic analysis of environment samples to detect pathogens.

In some embodiments, the methods disclosed herein are useful for diagnosis of one or more heritable disorders. In some embodiments, a heritable disorder that may be diagnosed with the methods disclosed herein is a genetic disorder that is prevalent in the Ashkenazi Jewish population. In some embodiments, the heritable disorders are selected from: 21-Hydroxylase-Deficient Congenital Adrenal Hyperplasia; ABCC8-Related Hyperinsulinism; Alpha-Thalassemia, includes Constant Spring, & MR associated; Arylsulfatase A Deficiency-Metyachromatic Leukodystrophy; Biotinidase Deficiency Holocarboxylase Synthetase Deficiency; Bloom's Syndrome; Canavan Disease; CFTR-Related Disorders-cystic fibrosis; Citrullinemia Type 1; Combined MMA & Homocystinuria-db1C; Dystrophinopathies (DMD & BMD); Familial Dysautonomia; Fanconi Anemia-FANCC; Galactosemia-Classical: Galactokinase Deficiency & Galactose Epimerase Deficiency; Gaucher Disease; GJB2-Related DFNB 1 Nonsyndromic Hearing Loss and Deafness; Glutaric acidemia Type 1; Hemoglobinopathies beta-chain disorders; Glycogen Storage Disease Type 1A; Maple Syrup Urine Disease; Types 1A, 1B, 2, 3; Medium Chain Acyl-Coenzyme A; Dehydrogenase Deficiency-MCADD; Methylmalonic Acidemia; Mucopolidosis IV; Nema-line Myopathy; Nieman-Pick Type A-Acid Sphingomyelinase Deficiency; Non-Ketotic Hyperglycinemia-Glycine Encephalopathy; Ornithine Transcarbamylase Deficiency; PKU Phenylalanine Hydroxylase Deficiency; Propionic Acidemia; Short Chain Acyl-CoA Dehydrogenase Deficiency-SCADD; Smith-Lemli-Opitz Syndrome; Spinal Muscular Atrophy (SMNI)-SMA; Tay Sachs-HexA Deficiency; Usher Syndrome-Type I (Type IB, Type IC, Type ID, Type IF, Type IG); X-Linked Mental Retardation ARX-Related Disorders; X-Linked Mental Retardation with Cerebellar Cyplopasia and distinctive Facial Appearance; X-Linked Mental Retardation; includes 9, 21, 30, 46, 58, 63, 88, 89; X linked mental retardation: FM1-Related Disorders—FRXA, Fragile X MR; X-linked SMR; Renpenning Syndrome 1; Zellweger Spectrum disorders—Peroxisomal Bifunctional Enzyme Deficiencies including Zellweger, NALD, and/or infantile Refsums. However, all of these, subsets of these, other genes, or combinations thereof may be used.

According to some aspects, the disclosure relates to multiplex diagnostic methods. In some embodiments, multiplex diagnostic methods comprise capturing a plurality of genetic loci in parallel (e.g., a genetic locus of Table 1). In some embodiments, genetic loci possess one or more polymorphisms (e.g., a polymorphism of Table 2) the genotypes of which correspond to disease causing alleles. Accordingly, in some embodiments, the disclosure provides methods for assessing multiple heritable disorders in parallel.

In some embodiments, methods are provided for diagnosing multiple heritable disorders in parallel at a pre-implantation, prenatal, perinatal, or postnatal stage. In some embodiments, the disclosure provides methods for analyzing multiple genetic loci (e.g., a plurality of target nucleic acids selected from Table 1) from a patient sample, such as a blood, pre-implantation embryo, chorionic villus or amniotic fluid sample. A patient or subject may be a human. However, aspects of the invention are not limited to humans and may be applied to other species (e.g., mammals, birds, reptiles, other vertebrates or invertebrates) as aspects of the invention are not limited in this respect. A subject or patient may be male or female. In some embodiments, in connection with reproductive genetic counseling, samples from a male and female member of a couple may be analyzed. In some embodiments, for example, in connection with an animal

breeding program, samples from a plurality of male and female subjects may be analyzed to determine compatible or optimal breeding partners or strategies for particular traits or to avoid one or more diseases or conditions.

However, it should be appreciated that any other diseases may be studied and/or risk factors for diseases or disorders including, but not limited to allergies, responsiveness to treatment, cancer tumor profiling for treatment and prognosis, monitoring and identification of patient infections, and monitoring of environmental pathogens.

1. Reducing Representational Bias in Multiplex Amplification Reactions:

In some embodiments, aspects of the invention relate to methods that reduce bias and increase reproducibility in multiplex detection of genetic loci, e.g., for diagnostic purposes.

Molecular inversion probe technology is used to detect or amplify particular nucleic acid sequences in potentially complex mixtures. Use of molecular inversion probes has been demonstrated for detection of single nucleotide polymorphisms (Hardenbol et al. 2005 *Genome Res* 15:269-75) and for preparative amplification of large sets of exons (Porreca et al. 2007 *Nat Methods* 4:931-6, Krishnakumar et al. 2008 *Proc Natl Acad Sci USA* 105:9296-301). One of the main benefits of the method is in its capacity for a high degree of multiplexing, because generally thousands of targets may be captured in a single reaction containing thousands of probes. However, challenges associated with, for example, amplification efficiency (See, e.g., Turner E H, et al., *Nat Methods*. 2009 Apr. 6:1-2.) have limited the practical utility of the method in research and diagnostic settings.

Aspects of the disclosure are based, in part, on the discovery of effective methods for overcoming challenges associated with systematic errors (bias) in multiplex genomic capture and sequencing methods, namely high variability in target nucleic acid representation and unequal sampling of heterozygous alleles in pools of captured target nucleic acids (e.g., isolated from a biological sample). Accordingly, in some embodiments, the disclosure provides methods that reduce variability in the detection of target nucleic acids in multiplex capture methods. In other embodiments, methods improve allelic representation in a capture pool and, thus, improve variant detection outcomes. In certain embodiments, the disclosure provides preparative methods for capturing target nucleic acids (e.g., genetic loci) that involve the use of different sets of multiple probes (e.g., molecular inversion probes MIPs) that capture overlapping regions of a target nucleic acid to achieve a more uniform representation of the target nucleic acids in a capture pool compared with methods of the prior art. In other embodiments, methods reduce bias, or the risk of bias, associated with large scale parallel capture of genetic loci, e.g., for diagnostic purposes. In other embodiments, methods are provided for increasing reproducibility (e.g., by reducing the effect of polymorphisms on target nucleic acid capture) in the detection of a plurality of genetic loci in parallel. In further embodiments, methods are provided for reducing the effect of probe synthesis and/or probe amplification variability on the analysis of a plurality of genetic loci in parallel.

In some aspects, the disclosure provides probe sets that comprise a plurality of different probes. As used herein, a 'probe' is a nucleic acid having a central region flanked by a 5' region and a 3' region that are complementary to nucleic acids flanking the same strand of a target nucleic acid or

subregion thereof. An exemplary probe is a molecular inversion probe (MIP). A 'target nucleic acid' may be a genetic locus.

Exemplary genetic loci are disclosed herein in Table 1 (RefSeqGene Column).

While probes have been typically designed to meet certain constraints (e.g. melting temperature, G/C content, etc.) known to partially affect capture/amplification efficiency (Ball et al (2009) *Nat Biotech* 27:361-8 AND Deng et al (2009) *Nat Biotech* 27:353-60), a set of constraints which is sufficient to ensure either largely uniform or highly reproducible capture/amplification efficiency has not previously been achieved.

As disclosed herein, uniformity and reproducibility can be increased by designing multiple probes per target, such that each base in the target is captured by more than one probe. In some embodiments, the disclosure provides multiple MIPs per target to be captured, where each MIP in a set designed for a given target nucleic acid has a central region and a 5' region and 3' region ('targeting arms') which hybridize to (at least partially) different nucleic acids in the target nucleic acid (immediately flanking a subregion of the target nucleic acid). Thus, differences in efficiency between different targeting arms and fill-in sequences may be averaged across multiple MIPs for a single target, which results in more uniform and reproducible capture efficiency.

In some embodiments, the methods involve designing a single probe for each target (a target can be as small as a single base or as large as a kilobase or more of contiguous sequence).

It may be preferable, in some cases, to design probes to capture molecules (e.g., target nucleic acids or subregions thereof) having lengths in the range of 1-200 bp (as used herein, a bp refers to a base pair on a double-stranded nucleic acid—however, where lengths are indicated in bps, it should be appreciated that single-stranded nucleic acids having the same number of bases, as opposed to base pairs, in length also are contemplated by the invention). However, probe design is not so limited. For example, probes can be designed to capture targets having lengths in the range of up to 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500, 1000, or more bps, in some cases.

It is to be appreciated that the length of a capture molecule (e.g., a target nucleic acid or subregion thereof) is selected based upon multiple considerations. For example, where analysis of a target involves sequencing, e.g., with a next-generation sequencer, the target length should typically match the sequencing read-length so that shotgun library construction is not necessary. However, it should be appreciated that captured nucleic acids may be sequenced using any suitable sequencing technique as aspects of the invention are not limited in this respect.

It is also to be appreciated that some target nucleic acids are too large to be captured with one probe. Consequently, it may be necessary to capture multiple subregions of a target nucleic acid in order to analyze the full target.

In some embodiments, a subregion of a target nucleic acid is at least 1 bp. In other embodiments, a subregion of a target nucleic acid is at least 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000 bp or more. In other embodiments, a subregion of a target nucleic acid has a length that is up to 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, or more percent of a target nucleic acid length.

The skilled artisan will also appreciate that consideration is made, in the design of MIPs, for the relationship between probe length and target length. In some embodiments, MIPs

are designed such that they are several hundred basepairs (e.g., up to 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000 bp or more) longer than corresponding target (e.g., subregion of a target nucleic acid, target nucleic acid).

In some embodiments, lengths of subregions of a target nucleic acid may differ.

For example, if a target nucleic acid contains regions for which probe hybridization is not possible or inefficient, it may be necessary to use probes that capture subregions of one or more different lengths in order to avoid hybridization with problematic nucleic acids and capture nucleic acids that encompass a complete target nucleic acid.

Aspects of the invention involve using multiple probes, e.g., MIPs, to amplify each target nucleic acid. In some embodiments, the set of probes for a given target can be designed to 'tile' across the target, capturing the target as a series of shorter sub targets. In some embodiments, where a set of probes for a given target is designed to 'tile' across the target, some probes in the set capture flanking non-target sequence). Alternately, the set can be designed to 'stagger' the exact positions of the hybridization regions flanking the target, capturing the full target (and in some cases capturing flanking non-target sequence) with multiple probes having different targeting arms, obviating the need for tiling. The particular approach chosen will depend on the nature of the target set. For example, if small regions are to be captured, a staggered-end approach might be appropriate, whereas if longer regions are desired, tiling might be chosen. In all cases, the amount of bias-tolerance for probes targeting pathological loci can be adjusted ('dialed in') by changing the number of different MIPs used to capture a given molecule.

In some embodiments, the 'coverage factor', or number of probes used to capture a basepair in a molecule, is an important parameter to specify. Different numbers of probes per target are indicated depending on whether one is using the tiling approach (see, e.g., FIG. 1) or one of the staggered approaches (see, e.g., FIG. 2 or 3).

FIG. 1 illustrates a non-limiting embodiment of a tiled probe layout showing ten captured sub-targets tiled across a single target. Each position in the target is covered by three sub-targets such that MIP performance per base pair is averaged across three probes.

FIG. 2 illustrates a non-limiting embodiment of a staggered probe layout showing the targets captured by a set of three MIPs. Each MIP captures the full target, shown in black, plus (in some cases) additional extra-target sequence, shown in gray, such that the targeting arms of each MIP fall on different sequence. Each position in the target is covered by three sub-targets such that MIP performance per basepair is averaged across three probes. Targeting arms land immediately adjacent to the black or gray regions shown. It should be appreciated that in some embodiments, the targeting arms (not shown) can be designed so that they do not overlap with each other.

FIG. 3 illustrates a non-limiting embodiment of an alternating staggered probe layout showing the targets captured by a set of three MIPs. Each MIP captures the full target, shown in black, plus (in some cases) additional extra-target sequence, shown in gray, such that the targeting arms of each MIP fall on different sequence. Each position in the target is covered by three sub-targets such that MIP performance per basepair is averaged across three probes. Targeting arms land immediately adjacent to the black or gray regions shown.

It should be appreciated that for any of the layouts, the targeting arms on adjacent tiled or staggered probes may be designed to either overlap, not overlap, or overlap for only a subset of the probes.

In certain embodiments for any of the layouts, a coverage factor of about 3 to about 10 is used. However, the methods are not so limited and coverage factors of up to 2, 3, 4, 5, 6, 7, 8, 9, 10, 20 or more may be used. It is to be appreciated that the coverage factor selected may depend the probe layout being employed. For example, in the tiling approach, for a desired coverage factor, the number of probes per target is typically a function of target length, sub-target length, and spacing between adjacent sub-target start locations (step size). For example, for a desired coverage factor of 3, a 200 bp target with a start-site separation of 20 bp and sub-target length of 60 bp may be encompassed with 12 MIPs (FIG. 1). Thus, a specific coverage factor may be achieved by varying the number of probes per target nucleic acid and the length of the molecules captured. In the staggered approach, a fixed-length target nucleic acid is captured as several sub-regions or as 'super-targets', which are molecules comprising the target nucleic acid and additional flanking nucleic acids, which may be of varying lengths. For example, a target of 50 bp can be captured at a coverage factor of 3 with 3 probes in either a 'staggered' (FIG. 2) or 'alternating staggered' configuration (FIG. 3).

The coverage factor will be driven by the extent to which detection bias is tolerable. In some cases, where the bias tolerance is small, it may be desirable to target more subregions of target nucleic acid with, perhaps, higher coverage factors. In some embodiments, the coverage factor is up to 2, 3, 4, 5, 6, 7, 8, 9, 10 or more.

In some embodiments, when a tiled probe layout is used, when the target length is greater than 1 bp and when a step size (distance between the 5'-end of a target and the 5' end of its adjacent target) is less than the length of a target or subregion thereof, it is possible to compute probe number for a particular target based on target length (T), sub target length (S), and coverage factor (C), such that probe number = $T/(S/C) + (C-1)$.

In some aspects, the disclosure provides methods to increase the uniformity of amplification efficiency when multiple molecules are amplified in parallel; methods to increase the reproducibility of amplification efficiency; methods to reduce the contribution of targeting probe variability to amplification efficiency; methods to reduce the effect on a given target nucleic acid of polymorphisms in probe hybridization regions; and/or methods to simplify downstream workflows when multiplex amplification by MIPs is used as a preparative step for analysis by nucleic acid sequencing.

Polymorphisms in the target nucleic acid under the regions flanking a target can interfere with hybridization, polymerase fill-in, and/or ligation. Furthermore, this may occur for only one allele, resulting in allelic drop-out, which ultimately decreases downstream sequencing accuracy. In some embodiments, using a set of MIPs having multiple hybridization sites for the capture of any given target, the probability of loss from polymorphism is substantially decreased because not all targeting arms in the set of MIPs will cover the location of the mutation.

Probes for MIP capture reactions may be synthesized on programmable microarrays because of the large number of sequences required. Because of the low synthesis yields of these methods, a subsequent amplification step is required to produce sufficient probe for the MIP amplification reaction. The combination of multiplex oligonucleotide synthesis and

pooled amplification results in uneven synthesis error rates and representational biases. By synthesizing multiple probes for each target, variation from these sources may be averaged out because not all probes for a given target will have the same error rates and biases.

Multiplex amplification strategies disclosed herein may be used analytically, as in detection of SNPs, or preparatively, often for next-generation sequencing or other sequencing techniques. In the preparative setting, the output of an amplification reaction is generally the input to a shotgun library protocol, which then becomes the input to the sequencing platform. The shotgun library is necessary in part because next-generation sequencing yields reads significantly shorter than amplicons such as exons. In addition to the bias-reduction afforded by the multi-tiled approach described here, tiling also obviates the need for shotgun library preparation. Since the length of the capture molecule can be specified when the probes, e.g., MIPs, are designed, it can be chosen to match the readlength of the sequencer. In this way, reads can ‘walk’ across an exon by virtue of the start position of each capture molecule in the probe set for that exon. Reducing analytical errors associated with bias in nucleic acid preparations: In some embodiments, aspects of the invention relate to preparative steps in DNA sequencing-related technologies that reduce bias and increase the reliability and accuracy of downstream quantitative applications.

There are currently many genomics assays that utilize next-generation (polony-based) sequencing to generate data, including genome resequencing, RNA-seq for gene expression, bisulphite sequencing for methylation, and Immune-seq, among others. In order to make quantitative measurements (including genotype calling), these methods utilize the counts of sequencing reads of a given genomic locus as a proxy for the representation of that sequence in the original sample of nucleic acids. The majority of these techniques require a preparative step to construct a high-complexity library of DNA molecules that is representative of a sample of interest. This may include chemical or biochemical treatment of the DNA (e.g., bisulphite treatment), capture of a specific subset of the genome (e.g., padlock probe capture, solution hybridization), and a variety of amplification techniques (e.g., polymerase chain reaction, whole genome amplification, rolling circle amplification).

Systematic and random errors are common problems associated with genome amplification and sequencing library construction techniques. For example, genomic sequencing library may contain an over- or under-representation of particular sequences from a source genome as a result of errors (bias) in the library construction process. Such bias can be particularly problematic when it results in target sequences from a genome being absent or undetectable in the sequencing libraries. For example, an under-representation of particular allelic sequences (e.g., heterozygotic alleles) from a genome in a sequencing library can result in an apparent homozygous representation in a sequencing library. As most downstream sequencing library quantification techniques depend on stochastic counting processes, these problems have typically been addressed by sampling enough (over-sampling) to obtain a minimum number of observations necessary to make statistically significant decisions. However, the strategy of oversampling is generally limited to elimination of low-count Poisson noise, and the approach wastes resources and increases the expense required to perform such experiments. Moreover, oversampling can result in a reduced statistical confidence in certain conclusions (e.g., diagnostic calls) based on the data.

Accordingly, new approaches are needed for overcoming bias in sequencing library preparatory methods.

Aspects of the disclosure are based, in part, on the discovery of methods for overcoming problems associated with systematic and random errors (bias) in genome capture, amplification and sequencing methods, namely high variability in the capture and amplification of nucleic acids and disproportionate representation of heterozygous alleles in sequencing libraries. Accordingly, in some embodiments, the disclosure provides methods that reduce variability in the capture and amplification of nucleic acids. In other embodiments, the methods improve allelic representation in sequencing libraries and, thus, improve variant detection outcomes. In certain embodiments, the disclosure provides preparative methods for capturing target nucleic acids (e.g., genetic loci) that involve the use of differentiator tag sequences to uniquely tag individual nucleic acid molecules. In some embodiments, the differentiator tag sequence permits the detection of bias based on the frequency with which pairs of differentiator tag and target sequences are observed in a sequencing reaction. In other embodiments, the methods reduce errors caused by bias, or the risk of bias, associated with the capture, amplification and sequencing of genetic loci, e.g., for diagnostic purposes.

Aspects of the invention relate to associating unique sequence tags (referred to as differentiator tag sequences) with individual target molecules that are independently captured and/or analyzed (e.g., prior to amplification or other process that may introduce bias). These tags are useful to distinguish independent target molecules from each other thereby allowing an analysis to be based on a known number of individual target molecules. For example, if each of a plurality of target molecule sequences obtained in an assay is associated with a different differentiator tag, then the target sequences can be considered to be independent of each other and a genotype likelihood can be determined based on this information. In contrast, if each of the plurality of target molecule sequences obtained in the assay is associated with the same differentiator tag, then they probably all originated from the same target molecule due to over-representation (e.g., due to biased amplification) of this target molecule in the assay. This provides less information than the situation where each nucleic acid was associated with a different differentiator tag. In some embodiments, a threshold number of independently isolated molecules (e.g., unique combinations of differentiator tag and target sequences) is analyzed to determine the genotype of a subject.

In some embodiments, the invention relates to compositions comprising pools (libraries) of preparative nucleic acids that each comprise “differentiator tag sequences” for detecting and reducing the effects of bias, and for genotyping target nucleic acid sequences. As used herein, a “differentiator tag sequence” is a sequence of a nucleic acid (a preparative nucleic acid), which in the context of a plurality of different isolated nucleic acids, identifies a unique, independently isolated nucleic acid. Typically, differentiator tag sequences are used to identify the origin of a target nucleic acid at one or more stages of a nucleic acid preparative method. For example, in the context of a multiplex nucleic acid capture reaction, differentiator tag sequences provide a basis for differentiating between multiple independent, target nucleic acid capture events. Also, in the context of a multiplex nucleic acid amplification reaction, differentiator tag sequences provide a basis for differentiating between multiple independent, primary amplicons of a target nucleic acid, for example. Thus, combinations of target nucleic acid

and differentiator tag sequence (target:differentiator tag sequences) of an isolated nucleic acid of a preparative method provide a basis for identifying unique, independently isolated target nucleic acids. FIG. 4A-C depict various non-limiting examples of methods for combining differentiator tag sequence and target sequences.

It will be apparent to the skilled artisan that differentiator tags may be synthesized using any one of a number of different methods known in the art. For example, differentiator tags may be synthesized by random nucleotide addition.

Differentiator tag sequences are typically of a predefined length, which is selected to control the likelihood of producing unique target:differentiator tag sequences in a preparative reaction (e.g., amplification-based reaction, a circularization selection-based reaction, e.g., a MIP reaction). Differentiator tag sequences may be, up to 5, up to 6, up to 7 up to 8, up to 9, up to 10, up to 11, up to 12, up to 13, up to 14, up to 15, up to 16, up to 17, up to 18, up to 19, up to 20, up to 21, up to 22, up to 23, up to 24, up to 25, or more nucleotides in length. For purposes of genotyping, isolated nucleic acids are identified as independently isolated if they comprise unique combinations of target nucleic acid and differentiator tag sequences, and observance of threshold numbers of unique combinations of target nucleic acid and differentiator tag sequences provide a certain statistical confidence in the genotype.

During a library preparation process, each nucleic acid molecule may be tagged with a unique differentiator tag sequence in a configuration that permits the differentiator tag sequence to be sequenced along with the target nucleic acid sequence of interest (the nucleic acid sequence for which the library is being prepared, e.g., a polymorphic sequence). The incorporation of the nucleic acid comprising a differentiator tag sequence at a particular step allows the detection and correction of biases in subsequent steps of the protocol.

A large library of unique differentiator tag sequences may be created by using degenerate, random-sequence polynucleotides of defined length. The differentiator tag sequences of the polynucleotides may be read at the final stage of the sequencing. The observations of the differentiator tag sequences may be used to detect and correct biases in the final sequencing read-out of the library. For example, the total possible number of differentiator tag sequences, which may be produced, e.g., randomly, is 4^N , where N is the length of the differentiator tag sequence. Thus, it is to be understood that the length of the differentiator tag sequence may be adjusted such that the size of the population of MIPs having unique differentiator tag sequences is sufficient to produce a library of MIP capture products in which identical independent combinations of target nucleic acid and differentiator tag sequence are rare. As used herein combinations of target nucleic acid and differentiator tag sequences, may also be referred to as "target:differentiator tag sequences".

In the final readout of a sequencing process, each read may have an additional unique differentiator tag sequence. In some embodiments, when differentiator tag sequences are distributed randomly in a library, all the unique differentiator tag sequences will be observed about an equal number of times. Accordingly, the number of occurrences of a differentiator tag sequence may follow a Poisson distribution.

In some embodiments, overrepresentation of target:differentiator tag sequences in a pool of preparative nucleic acids (e.g., amplified MIP capture products) is indicative of bias in the preparative process (e.g., bias in the amplification process). For example, target:differentiator tag sequence combinations that are statistically overrepresented are

indicative of bias in the protocol at one or more steps between the incorporation of the differentiator tag sequences into MIPs and the actual sequencing of the MIP capture products.

The number of reads of a given target:differentiator tag sequence may be indicative (may serve as a proxy) of the amount of that target sequence present in the originating sample. In some embodiments, the numbers of occurrence of sequences in the originating sample is the quantity of interest. For example, using the methods disclosed herein, the occurrence of differentiator tag sequences in a pool of MIPs may be predetermined (e.g., may be the same for all differentiator tag sequences). Accordingly, changes in the occurrence of differentiator tag sequences after amplification and sequencing may be indicative of bias in the protocol. Bias may be corrected to provide an accurate representation of the composition of the original MIP pool, e.g., for diagnostic purposes.

According to some aspects, a library of preparative nucleic acid molecules (e.g., MIPs, each nucleic acid in the library having a unique differentiator tag sequence, may be constructed such that the number of nucleic acid molecules in the library is significantly larger than the number prospective target nucleic acid molecules to be captured using the library. This ensures that products of the preparative methods include only unique target:differentiator tag sequence; e.g., in a MIP reaction the capture step would undersample the total population of unique differentiator tag sequences in the MIP library. For example, an experiment utilizing 1 ug of genomic DNA will contain about ~150,000 copies of a diploid genome. For a MIP library, each MIP in the library comprising a randomly produced 12-mer differentiator tag sequence (~1.6 million possible unique differentiator tag sequences), there would be more than 100 unique differentiator tag sequences per genomic copy. For a MIP library, each MIP in the library comprising a randomly produced 15-mer differentiator tag sequence (~1 billion possible unique differentiator tag sequences), there would be more than 7000 unique differentiator tag sequences per genomic copy. Therefore, the probability of the same differentiator tag sequence being incorporated multiple times is incredibly small. Thus, it is to be appreciated that the length of the differentiator tag sequence is to be selected based on the amount of target sequence in a MIP capture reaction and the desired probability for having multiple, independent occurrences of target:differentiator tag sequence combinations.

FIG. 5 depicts a non-limiting method for genotyping based on target and differentiator tag sequences. Sequencing reads of target and differentiator tags sequences are collapsed to make diploid genotype calls. FIG. 6 depicts non-limiting results of a simulation of a MIP capture reaction in which MIP probes, each having a differentiator tag sequence of 15 nucleotides, are combined with 10000 target sequence copies (e.g., genome equivalents). In this simulated reaction, the probability of capturing one or more copies of a target sequence having the same differentiator tag sequence is 0.05. The Y axis reflects the number of observations. The X axis reflects the number of independent occurrences of target:differentiator tag combinations. FIG. 7 depicts a non-limiting graph of sequencing coverage, which can help ensure that alleles are sampled to sufficient depth (e.g., either 10x or 20x minimum sampling per allele, assuming 1000 targets). In this non-limiting example, the X axis is total per-target coverage required, and the Y axis is the probability that a given total coverage will result in at least 10x or 20x coverage for each allele.

31

The skilled artisan will appreciate that as part of a MIP library preparation process, adapters may be ligated onto the ends of the molecules of interest. Adapters often contain PCR primer sites (for amplification or emulsion PCR) and/or sequencing primer sites. In addition, barcodes may be included, for example, to uniquely identify individual samples (e.g., patient samples) that may be mixed together. (See, e.g., USPTO Publication Number US 2007/0020640 A1 (McCloskey et al.))

The actual incorporation of the random differentiator tag sequences can be performed through various methods known in the art. For example, nucleic acids comprising differentiator tag sequences may be incorporated by ligation. This is a flexible method, because molecules having differentiator tag sequence can be ligated to any blunt-ended nucleic acids. The sequencing primers must be incorporated subsequently such that they sequence both the differentiator tag sequence and the target sequence. Alternatively, the sequencing adaptors can be synthesized with the random differentiator tag sequences at their 3' end (as degenerate bases), so that only one ligation must be performed. Another method is to incorporate the differentiator tag sequence into a PCR primer, such that the primer structure is arranged with the common adaptor sequence followed by the random differentiator tag sequence followed by the PCR priming sequence (in 5' to 3' order). A differentiator tag sequence and adaptor sequence (which may contain the sequencing primer site) are incorporated as tags. Another method to incorporate the differentiator tag sequences is to synthesize them into a padlock probe prior to performing a gene capture reaction. The differentiator tag sequence is incorporated 3' to the targeting arm but 5' to the amplification primer that will be used downstream in the protocol. Another method to incorporate the differentiator tag sequences is as a tag on a gene-specific or poly-dT reverse-transcription primer. This allows the differentiator tag sequence to be incorporated directly at the cDNA level.

In some embodiments, at the incorporation step, the distribution of differentiator tag sequences can be assumed to be uniform. In this case, bias in any part of the protocol would change the uniformity of this distribution, which can be observed after sequencing. This allows the differentiator tag sequence to be used in any preparative process where the ultimate output is sequencing of many molecules in parallel.

Differentiator tag sequences may be incorporated into probes (e.g., MIPs) of a plurality when they are synthesized on-chip in parallel, such that degeneracy of the incorporated nucleotides is sufficient to ensure near-uniform distribution in the plurality of probes. It is to be appreciated that

32

amplification of a pool of unique differentiator tag sequences may itself introduce bias in the initial pool. However, in most practical cases, the scale of synthesis (e.g., by column synthesis, chip based synthesis, etc.) is large enough that amplification of an initial pool of differentiator tag sequences is not necessary. By avoiding amplification or selection steps on the pool of unique differentiator tag sequences, potential bias may be minimized. One example of the use of the differentiator tag sequences is in genome re-sequencing.

Considering that the raw accuracy of most next-generation sequencing instruments is relatively low, it is crucial to oversample the genomic loci of interest.

Furthermore, since there are two alleles at every locus, it is important to sample enough to ensure that both alleles have been observed a sufficient number of times to determine with a sufficient degree of statistical confidence whether the sample is homozygous or heterozygous. Indeed, the sequencing is performed to sample the composition of molecules in the originating sample. However, after multiple reads have been collected for a given locus, it is possible that due to bias (e.g., caused by PCR amplification steps), a large fraction of the reads are derived from a single originating molecule. This would skew the population of target sequences observed, and would affect the outcome of the genotype call. For example, it is possible that a locus that is heterozygous is called as homozygous, because there are only a few observations of the second allele out of many observations of that locus. However, if information is available on differentiator tag sequences, this situation could be averted, because the over-represented allele would be seen to also have an over-represented differentiator tag sequence (i.e., the sequences with the overrepresented differentiator tag sequence all originated from the same single molecule). Therefore, the sequences and corresponding distribution of differentiator tag sequences can be used as an additional input to the genotype-calling algorithm to significantly improve the accuracy and confidence of the genotype calls.

In some aspects, the disclosure provides methods for analyzing a plurality of target sequences which are genetic loci or portions of genetic loci (e.g., a genetic locus of Table 1). The genetic loci may be analyzed by sequencing to obtain a genotype at one or more polymorphisms (e.g., SNPs). Exemplary polymorphisms are disclosed in Table 2. The skilled artisan will appreciate that other polymorphisms are known in the art and may be identified, for example, by querying the Entrez Single Nucleotide Polymorphism database, for example, by searching with a GeneID from Table 1.

TABLE 1

Target Nucleic Acids						
Gene name	Gene ID	Description	Gene aliases	OMIM	RefSeqGene	Chromosome map position
CYP21 A2	1589	cytochrome P450, family 21, subfamily A, polypeptide 2	CAH1; CPS1; CA21H; CYP21; CYP21B; P450c21B; MGC150536; MGC150537; CYP21 A2	201910	NG_008337.1	6p21.3
ABCC8	6833	ATP-binding cassette, sub-family C (CFTR/MRP), member 8	HI; SUR; HHF1; MRP8; PHH1; SUR1; ABC36; HRINS; TNDM2; ABCC8	600509	NG_008867.1	11p15.1

TABLE 1-continued

Target Nucleic Acids						
Gene name	Gene ID	Description	Gene aliases	OMIM	RefSeqGene	Chromosome map position
ATRX	546	alpha thalassemia/mental retardation syndrome X-linked (RAD54 homolog, <i>S. cerevisiae</i>)	SHS; XH2; XNP; ATR2; SFM1; RAD54; MRXHF1; RAD54L; ZNF-HX; MGC2094; ATRX	300032	NG_008838.1	Xq13.1-q21.1
ARSA	410	arylsulfatase A	MLD; ARSA	607574	NG_009260.1	22q13.31-qter; 22q13.33
PSAP	5660	Prosaposin	GLBA; SAPI; FLJ00245; MGC110993; PSAP	176801	NG_008835.1	10q21-q22
BTD	686	Biotinidase	BTD	609019	NG_008019.1	3p25
HLCS	3141	holocarboxylase synthetase (biotin-(propionyl-Coenzyme A-carboxylase (ATP-hydrolysing)) ligase)	HCS; HLCS	609018	NC_000021.7	21q22.1; 21q22.13
BLM	641	Bloom syndrome, RecQ helicase-like	BS; RECQ2; RECQL2; RECQL3; MGC126616; MGC131618; MGC131620; BLM	604610	NG_007272.1	15q26.1
ASPA	443	aspartoacylase (Canavan disease)	ASP; ACY2; ASPA	608034	NG_008399.1	17pter-P13
CFTR	1080	cystic fibrosis transmembrane conductance regulator (ATP-binding cassette sub-family C, member 7)	CF; MRP7; ABC35; ABCC7; CFTR/MRP; TNR-CFTR; dJ760C5.1; CFTR	602421	NC_000007.12	7q31.2
ASS1	445	argininosuccinate synthetase 1	ASS; CTLN1; ASS1	603470	NG_011542.1	9q34.1
MMACHC	25974	methylmalonic aciduria (cobalamin deficiency) cb1C type, with homocystinuria	cb1C; FLJ25671; DKFZp564I122; RP11-291L19.3; MMACWC	609831	NC_000001.9	1p34.1
IKBKAP	8518	inhibitor of kappa light polypeptide gene enhancer in B-cells, kinase complex-associated protein	FD; DYS; ELP1; IKAP; IKI3; TOT1; FLJ12497; DKFZp781H1425; IKBKAP	603722	NG_008788.1	9q31
FANCC	2176	Fanconi anemia, complementation group C	FA3; FAC; FACC; FLJ14675; FANCC	227645	NG_011707.1	9q22.3
GALK1	2584	galactokinase 1	GK1; GALK; GALK1	604313	NG_008079.1	17q24
GALT	2592	galactose-1-phosphate uridylyltransferase	GALT	606999	NC_000009.10	9p13
GALE	2582	UDP-galactose-4-epimerase	SDR1E1; FLJ95174; FLJ97302; GALE	606953	NG_007068.1	1p36-p35
GBA	2629	glucosidase, beta; acid (includes glucosylceramidase)	GCB; GBA1; GLUC; GBA	606463	NG_009783.1	1q21
GJB2	2706	gap junction protein, beta 2, 26 kDa	HID; KID; PPK; CX26; DFNA3; DFNB1; NSRD1; DFNA3A; DFNB1A; GJB2	121011	NG_008358.1	13q11-q12

TABLE 1-continued

Target Nucleic Acids						
Gene name	Gene ID	Description	Gene aliases	OMIM	RefSeqGene	Chromosome map position
GCDH	2639	glutaryl-Coenzyme A dehydrogenase	GCD; ACAD5; GCDH	608801	NG_009292.1	19p13.2
G6PC	2538	glucose-6-phosphatase, catalytic subunit	G6PT; GSD1; GSD1a; MGCI63350; G6PC	232200	NG_011808.1	17q21
HBB	3043	hemoglobin, beta	CD113t-C; beta-globin; HBB	141900	NG_000007.3	11p15.5
BCKDHA	593	branched chain keto acid dehydrogenase E1, alpha polypeptide	MSU; MSUD1; OVD1A; BCKDE1A; FLJ45695; BCKDHA	608348	NC_000019.8	19q13.1-q13.2
BCKDHB	594	branched chain keto acid dehydrogenase E1, beta polypeptide	E1B; FLJ17880; dJ279A18.1; BCKDHB	248611	NG_009775.1	6q13-q15
DBT	1629	dihydrolipoamide branched chain transacylase E2	E2; E2B; BCATE2; MGC9061; DBT	248610	NG_011852.1	1p31
DLD	1738	dihydrolipoamide dehydrogenase	E3; LAD; DLDH; GCSL; PHE3; DLD	238331	NG_008045.1	7q31-q32
ACADM	34	acyl-Coenzyme A dehydrogenase, C-4 to C-12 straight chain	MCAD; ACAD1; MCADH; FLJ18227; FLJ93013; FLJ99884; ACADM	607008	NG_007045.1	1p31
MMAA	166785	methylmalonic aciduria (cobalamin deficiency) cblA type	cblA; MGC120010; MGC120011; MGC120012; MGC120013; MMAA	607481	NG_007536.1	4q31.22
MMAB	326625	methylmalonic aciduria (cobalamin deficiency) cblB type	ATR; cblB; MGC20496; MMAB	607568	NG_007096.1	12q24
MUT	4594	methylmalonyl Coenzyme A mutase	MCM; MUT	609058	NG_007100.1	6p12.3
MCOLN1	57192	mucolipin 1	ML4; MLIV; MST080; TRPML1; MSTP080; TRP-ML1; TRPM-L1; MCOLN1	605248	NC_000019.8	19p13.3-pl3.2
ACTA1	58	actin, alpha 1, skeletal muscle	ACTA; ASMA; CFTD; MPFD; NEM1; NEM2; NEM3; CFTD1; CFTDM; ACTA1	102610	NG_006672.1	1q42.13
TPM3	7170	tropomyosin 3	TM3; TRK; NEM1; TM-5; TM30; TM30 nm; TPMsk3; hscp30; MGC3261; FLJ41118; MGC14582; MGC72094; OK/SW-c1.5; TPM3	191030	NG_008621.1	1q21.2
TNNT1	7138	troponin T type 1 (skeletal, slow)	ANM; TNT; STNT; TNTS; FLJ98147; MGC104241; TNNT1	191041	NG_011829.1	19q13.4
NEB	4703	nebulin	NEM2; NEB177D; FLJ11505; FLJ36536; FLJ39568; FLJ39584; DKFZp686C1456; NEB	161650	NG_009382.1	2q22
SMPD1	6609	sphingomyelin phosphodiesterase 1, acid lysosomal	ASM; NPD; SMPD1	607608	NG_011780.1	11p15.4-pl5.1
GLDC	2731	glycine dehydrogenase (decarboxylating)	GCE; NKH; GCSP; HYGN1; MGC138198; MGC138200; GLDC	238300	NC_000009.10	9p22

TABLE 1-continued

Target Nucleic Acids						
Gene name	Gene ID	Description	Gene aliases	OMIM	RefSeqGene	Chromosome map position
GCSH	2653	glycine cleavage system protein H (aminomethyl carrier)	GCE; NKH; GCSH	238330	NC_000016.8	16q23.2
AMT	275	aminomethyltransferase	GCE; NKH; GCST; AMT	238310	NC_000003.10	3p21.2-p21.1
OTC	5009	ornithine carbamoyltransferase	OCTD; MGC129967; MGC129968; MGC138856; OTC	300461	NG_008471.1	Xp21.1
PAH	5053	phenylalanine hydroxylase	PH; PKU; PKU1; PAH	612349	NG_008690.1	12g22-q24.2
DHPR	5860	quinoid dihydropteridine reductase	DHPR; PKU2; SDR33C1; FLJ42391; QDPR	612676	NG_008763.1	4p15.31
PTS	5805	6-pyruvoyltetrahydropterin synthase	PTPS; FLJ97081; PTS	261640	NG_008743.1	11q22.3-q23.3
PCCA	5095	propionyl Coenzyme A carboxylase, alpha polypeptide	PCCA	232000	NG_008768.1	13q32
PCCB	5096	propionyl Coenzyme A carboxylase, beta polypeptide	DKFZp451E113; PCCB	232050	NG_008939.1	3q21-q22
ACADS	35	acyl-Coenzyme A dehydrogenase, C-2 to C-3 short chain	SCAD; ACAD3; ACADS	606885	NG_007991.1	12g22-qter
DHCR7	1717	7-dehydrocholesterol reductase	SLOS; DHCR7	602858	NC_000011.8	11q13.2-q13.5
SMNT	6606	survival of motor neuron 1, telomeric	SMA; SMN; SMA1; SMA2; SMA3; SMA4; SMA@; SMNT; BCD541; T-BCD541; SMN1	600354	NG_008691.1	5q13
HEXA	3073	hexosaminidase A (alpha polypeptide)	TSD; MGC99608; HEXA	606869	NG_009017.1	15g23-q24
MYO7A	4647	myosin VIIA	DFNB2; MYU7A; NSRD2; USH1B; DFNA11; MYOVIIA; MYO7A	276903	NG_009086.1	11q13.5
USH1C	10083	Usher syndrome 1 C (autosomal recessive, severe)	PDZ73; AIE-75; DFNB18; PDZ-45; PDZ-73; NY-CO-37; NY-CO-38; ush1cpst; PDZ-73/NY-CO-38; USH1C	605242	NC_000011.8	11p15.1-p14
CDH23	64072	cadherin-like 23	USH1D; DFNB12; FLJ00233; FLJ36499; KIAA1774; KIAA1812; MGC102761; DKFZp434P2350; CDH23	605516	NG_008835.1	10g21-q22
PCDH15	65217	protocadherin 15	USH1F; DFNB23; DKFZp667A1711; PCDH15	605514	NG_009191.1	10q21.1
SANS	124590	Usher syndrome 1G (autosomal recessive)	SANS; ANKS4A; FLJ33924; USH1G	607696	NG_007882.1	17q25.1

TABLE 1-continued

Target Nucleic Acids						
Gene name	Gene ID	Description	Gene aliases	OMIM	RefSeqGene	Chromosome map position
ARX	170302	aristaless related homeobox	ISSX; PRTS; MRX29; MRX32; MRX33; MRX36; MRX38; MRX43; MRX54; MRX76; MRX87; MRXS1; ARX	300382	NG_008281.1	Xp21
OPHN1	4983	oligophrenin 1	OPN1; MRX60; OPHN1	300127	NG_008960.1	Xq12
JARID1C	8242	lysine (K)-specific demethylase 5C	MRXJ; SMCX; MRXSJ; XE169; JARID1C; DXS1272E; KDM5C	314690	NG_008085.1	Xp11.22-p11.21
FTSJ1	24140	FtsJ homolog 1 (<i>E. coli</i>)	JM23; MRX9; SPB1; TRM7; CDLIV; MRX44; FTSJ1	300499	NG_008879.1	Xp11.23
SLC6A8	6535	solute carrier family 6 (neurotransmitter transporter, creatine), member 8	CRT; CT1; CRTR; MGC87396; SLC6A8	300036	NC_000023.9	Xq28
DLG3	1741	discs, large homolog 3 (<i>Drosophila</i>)	MRX; MRX90; NEDLG; NE-Dlg; SAP102; SAP-102; KIAA1232; DLG3	300189	NC_000023.9	Xq13.1
TM4SF2	7102	letraspanin 7	A15; MXS1; CD231; MRX58; CCG-B7; TM4SF2; TALLA-1; TM4SF2b; DXS1692E; TSPAN7	300096	NG_009160.1	Xp11.4
ZNF41	7592	zinc finger protein 41	MRX89; MGC8941; ZNF41	314995	NG_008238.1	Xp11.23
FACL4	2182	acyl-CoA synthetase long-chain family member 4	ACS4; FACL4; LACS4; MRX63; MRX68; ACSL4	300157	NG_008053.1	Xq22.3-q23
PQBP1	10084	polyglutamine binding protein 1	SHS; MRX55; MRXS3; MRXS8; NPW38; RENS1; PQBP1	300463	NC_000023.9	Xp11.23
PEX1	5189	peroxisomal biogenesis factor 1	ZWS1; PEX1	602136	NG_008341.1	7q21.2
PXMP3	5828	peroxisomal membrane protein 3, 35 kDa	PAF1; PEX2; PMP3; PAF-1; PMP35; RNF72; PXMP3	170993	NG_008371.1	8q21.1
PEX6	5190	peroxisomal biogenesis factor 6	PAF2; PAF-2; PXAAA1; PEX6	601498	NG_008370.1	6p21.1
PEX10	5192	peroxisomal biogenesis factor 10	NALD; RNF69; MGC1998; PEX10	602859	NG_008342.1	1p36.32
PEX12	5193	peroxisomal biogenesis factor 12	PAF-3; PEX12	601758	NG_008447.1	17q12

TABLE 1-continued

Target Nucleic Acids						
Gene name	Gene ID	Description	Gene aliases	OMIM	RefSeqGene	Chromosome map position
PEX5	5830	peroxisomal biogenesis factor 5	PXR1; PTS1R; PTS1-BP; FLJ50634; FLJ50721; FLJ51948; PEX5	600414	NG_008448.1	12p13.31
PEX26	55670	peroxisomal biogenesis factor 26	FLJ20695; PEX26M1T; Pex26pM1T; PEX26	608666	NG_008339.1	22q11.21

The mutations listed in Table 2 are documented polymorphisms in several disease-associated genes (CFTR is mutated in cystic fibrosis, GBA is mutated in Gaucher disease, ASPA is mutated in Canavan disease, HEXA is mutated in Tay Sachs disease). The polymorphisms are of several types: insertion/deletion polymorphisms which will

cause frameshifts (and thus generally interrupt protein function) unless the insertion/deletion length is a multiple of 3 bp, and substitutions which can alter the amino acid sequence of the protein and in some cases cause complete inactivation by introduction of a stop codon.

TABLE 2

Non-limiting examples of polymorphisms					
Gene name	GeneID	SNP ID	Mutation	SEQ ID NO:	
CFTR	1080	rs63500661	TCACATCACCAAGTTAAAAA[A/G]G GGGCGGGGGCAGAATGAAAATT	1	
CFTR	1080	rs63107760	AAACAAGGATGAATTAAGTTTTTTT[-/T] AAAAAAGAACATTTGGTAAGGGGA	2	
CFTR	1080	rs62469443	ATCACCAAGTTAAAAAAGGG[A/G]C GGGGGGCAGAATGAAAATTGCAT	3	
CFTR	1080	rs62469442	CTATTGAACCAGAACCAACAGGAAT[A/G]C CATAGCATTTTGTAACTAACTG	4	
CFTR	1080	rs62469441	CAGGAGTTCAAGACCAGCCTACTAAA[A/C]C ACACACACACACACACACAC	5	
CFTR	1080	rs62469439	GATTAAATAATAGTGTATTATGTACCC[C/G]GC TTATAGGAGAAGGGGTGTGTGT	6	
CFTR	1080	rs62469438	ATTGTTATCTTTTCATATAAGGTAAC[A/T]GA GGCCAGAGAGATTAAATAACAT	7	
CFTR	1080	rs62469437	TAATTTTAATTAAAGTAAATTTAATTG[A/G]TA GATAAATAAGTAGATAAAAAATA	8	
CFTR	1080	rs62469436	GTATAAAAAAAAAAAAAAAGTT[A/T]G AATGTTTCTTGCAATTCAGAGCCT	9	
CFTR	1080	rs62469435	ATACTAAAAATTAAAGTTCTCTTGC[A/G]AT ATATTTTCTTAATATCTTACATC	10	
CFTR	1080	rs62469434	TGCTGGGATTACAGGCGTGAGCCACC[A/G]C GCCTGGCCTGATGGACATATTTT	11	
CFTR	1080	rs62469433	CTACAATATAAGTATAGTATTGCAAA[A/C]CC ATCAGGAAGGGTGAATTAATTTT	12	
CFTR	1080	rs61763210	GTTGTCTCCAACTTTTTTCAGGTG[-/AGA] AGGTGGCCAACCGAGCTTCGGAAG	13	
CFTR	1080	rs61720488	TTTTTTCATAAAGATTATATAAAGG[A/C]TA TTGCTTTGAATCACAAACACTA	14	
CFTR	1080	rs61481156	ATCTAGTGAGCAGTCAGGAAAGAGAA[C/T]T TCCAGATCTGGAAATCAGGGTTA	15	
CFTR	1080	rs61443875	TAGAGTATAAAAAAAAAAAAAA[-/A] GTTTGAATGTTTCTTGCAATTCAGA	16	
CFTR	1080	rs61312222	TGCAATGCCAATATCAAGATATT[C/G]GA GTATACTGTCAATAAACTTCATA	17	

TABLE 2-continued

Non-limiting examples of polymorphisms				
Gene name	GeneID	SNP ID	Mutation	SEQ ID NO:
CFTR	1080	rs61159372	TCCTCAACAGTTAGAAACAATATTTT[C/G]AG TGATTTCCTCATGCCAACTTTACT	18
CFTR	1080	rs61094145	TTTTTGGTATTGTTGTTAAATAAGTG[A/G]GA ATTCAATACAGTATAATGTCTGT	19
CFTR	1080	rs61086387	CTTGAAATCGGATATATATATATATA[-/T GTATATATATATATATATATATATATAT ACATATATATATATA]GTATTATCCCTGTTTTTC ACAGTTT	20
CFTR	1080	rs60996744	AGAGGGGCTGTGAAGGACACCAAGGA[A/G]G AGACTAAGAGCCAGGAGGGGAAAC	21
CFTR	1080	rs60960860	TAGAGTTTATTAGCTTTTACTACTCT[A/G]CTT AGTTACTTTGTGTACAGAATA	22
CFTR	1080	rs60923902	ACTAGTGATGATGAGCTTCTTTTCAT[-/AT] GTTTGTGGCTGCATAAATGTCTTC	23
CFTR	1080	rs60912824	GCAGAGAAAAGAGGGGCTGTGAAGGA[C/G]A CCAAGGAGGAGACTAAGAGCCAGG	24
CFTR	1080	rs60887846	TTCAGAGGTCTACCACTGGTGCATAC[G/T]CT AATCACAGTGTGAAATTTTAC	25
CFTR	1080	rs60793174	AAGAAAGAGCAAAAGAGGGCAAACCTT[C/T]T CATACATTTTGTATGTCGAAACCA	26
CFTR	1080	rs60788575	CCTAAAGTTTAAAAAGAAAAAAAAA[-/A] GGAAGAAGGAATTAAAAATCCAAAG	27
CFTR	1080	rs60760741	GTGTGTGTGTGTATATATATATATAT[A/T]TA TATATTTTTTTTTTCTGAGCCA	28
CFTR	1080	rs60456599	AACTGTTGATGTTTTTCATTATTTA[C/G]ATC ATTGGAAAACCTTAGATTCTAG	29
CFTR	1080	rs60363249	TTTATCCATTCTTAACGAGACAGAC[A/G]TT TTTTAGAGCTGGTCCAGGAAA	30
CFTR	1080	rs60355115	TTGAAATCGGATATATATATATATAT[A/G]TA TATATATATATATATATATATAT	31
CFTR	1080	rs60308689	TAGTTTTTTATTTCTCATATTATTT[-/T] CAGTGGCTTTTTCTCCACATCTT	32
CFTR	1080	rs60271242	ACATAGTTCTCAGTGGTACAACTACA[A/G]GT GATTTCTTTTCTTATTTCTGG	33
CFTR	1080	rs60010318	AGAGCAATGGCATCCCTTGTCTTGTG[C/T]TA TACAGGATGCAGCAATTATAGG	34
CFTR	1080	rs59961323	TTCTGTCTACATAAGATGTCATACTA[A/G]AT TATCTTTCCAGCATGCATTCAG	35
CFTR	1080	rs59961270	CAGGGTGGCATGTTAGGAGTGCTTA[A/G]A ATAAATGAGTTGGTTATACAAGTA	36
CFTR	1080	rs59837506	AGGACACACACACACACACACACA[-/CA] TGACACACATTTAAATAGATGCAT	37
CFTR	1080	rs59572090	TAAAAAATTGGTATAATGAAATTGCA[C/T]TT GTAGTCTTTGGACATTAAATCC	38
CFTR	1080	rs59548252	TTCAATACTTAAGAGGTACGCAGAG[A/G]A AAGAGGGGCTGTGAAGGACCCAA	39
CFTR	1080	rs59519859	CAGCAATGAATATTTTGAGGCTGAGG[C/T]GC TGAGGGGTAAATTCAGCCTGG	40
CFTR	1080	rs59509837	TTATGGTTTATATTTTGTGTCTTCT[-/C TTT]AACACATCTTTCTAGCAGAATTCA	41

TABLE 2-continued

Non-limiting examples of polymorphisms				
Gene name	GeneID	SNP ID	Mutation	SEQ ID NO.
GBA	2629	rs12752133	CTCTTCCGAGGTTCCACCCTGAACAC[C/T]TT CCTGCTCCCTCGTGGTGTAGAGT	67
GBA	2629	rs12747811	TTCTGACTGGCAACCAGCCCCACTCT[C/T]TG GGAGCCCTCAGGAATGAAGTTC	68
GBA	2629	rs12743554	gctcagcctcccaggtgagtgag[A/T]ggcgcgatc tcggctcaccgcaacc	69
GBA	2629	rs12041778	CATGAACCACATCAAATGAGATTAG[C/T]GG GAGTGGCACACACAGTCATGACC	70
GBA	2629	rs12034326	AAGCAGCCCTGGGGAGTCGGGGCGGG[A/G]C CTGGATTGAAAAGAGACGGTCA	71
GBA	2629	rs11558184	CTCCAAGTTCTGGGAGCAGAGTGTGC[A/G]G CTAGGCTCCTGGGATCGAGGGATG	72
GBA	2629	rs11430678	GTTCTCCAGTAAtttttttttttt[-/G/T] gttttgagacagagtcttgccctgt	73
GBA	2629	rs11264345	CTAGTACCTTACTTCCCTCAAGTTCA[A/T]TC ATCTCACAGATATTTCCTGAGCA	74
GBA	2629	rs10908459	aattagccgtgctgtggcggtgc[C/T]tgtaatccc acgtacttgggaggct	75
GBA	2629	rs10796940	CCATGGCCAGCCGGGGAGGGACGGG[A/C]A CACACAGACCCACACAGAGACTCA	76
GBA	2629	rs10668496	agcgagagagagagagagagagagagagagagag[-/AG] gagCCGGCGGAGAACTACGCATGC	77
GBA	2629	rs7416991	CGTAGCAGTTAGCAGATGATAGGCGG[C/G/T] GAAATCTTATTTCACAGGGCATTAA	78
GBA	2629	rs4024049	CTGGCCCTGGTGACAGTGGGGCTGTG[C/T]GT GGGGCCAGAGCCTTCTCAGAGGT	79
GBA	2629	rs4024048	CAGATACTGGCCCTGGTGACAGTGGG[A/G]C TGTGCGTGGGGCCAGAGCCTTCTC	80
GBA	2629	rs4024047	GACAGATACTGGCCCTGGTGACAGTG[G/T]G GCTGTGCGTGGGGCCAGAGCCTTCTC	81
GBA	2629	rs3841430	GGCTCctctctctctctctctctctct[-/TC] gctcgctctctcgctctctcgctct	82
GBA	2629	rs3754485	GTTTCAGACCAGCCTGGCCAACATAG[C/T]GA AACCCATCTCTACTAAAAATAA	83
GBA	2629	rs3205619	AGTGGGCGATTGGATGGAGCTGAGTA[C/T]G GGGCCATCCAGGCTAATCACACC	84
GBA	2629	rs2990227	CCGGGCTCCGTGAATGTTTGTACAT[C/G]TC TGAAGAACGTATGAATTACATAA	85
GBA	2629	rs2990226	GAATCCCAACCCGACGCTCGTCGCC[C/G]G GCTCCGTGAATGTTTGTACATGT	86
GBA	2629	rs2990225	GCGAATCCCAACCCGACGCTCGTCG[C/T]CG GGCTCCGTGAATGTTTGTACAT	87
GBA	2629	rs2990224	TGGGCAGAAAGTCAGGGTCCAAAGAAA[G/T]G GCAAAGAAAAGTGTcagtggtca	88
ASPA	443	rs63751297	TAAGAAAGACGTTTTTGATTTTTTTC[A/G]GA CTTCTCTGGCTCCACTACCCTGC	89
ASPA	443	rs62071301	CTGATTCCTGGCCAGGAGCGGTGGCT[C/T]AC GCCTGTAATCCAGCGCTTTGGG	90
ASPA	443	rs62071300	TAAAAATGCTGATTCCTGGCCAGGAG[C/T]GG TGGCTCACGCCGTGAATCCAGC	91

TABLE 2-continued

Non-limiting examples of polymorphisms				
Gene name	GeneID	SNP ID	Mutation	SEQ ID NO.
ASPA	443	rs62071299	TTTAAAAATGCTGATTCCTGGCCAGG[A/C]GC GGTGGCTCAGCCTGTAATCCCA	92
ASPA	443	rs62071297	CAAGACCTGTCAAAGATCTGAGAAAT[A/T]TT ACCCGACTTACAAGCTAACCATT	93
ASPA	443	rs61697033	ACTGTAATAAGTGTGTAAAAGAAAT[A/G]C ACAAAATAATATAGCAGAGGGTAT	94
ASPA	443	rs60743592	CTTGAGGTCAGGAGTTCAAGACCAGT[C/T]TG GGCAACATGGGAAAACCTTGTC	95
ASPA	443	rs60666840	AGGTTGCAGTGAGCCGAGATCATGCC[A/G]TT GCACTCCAGCCGGGCAACAAAA	96
ASPA	443	rs60147514	ACAAGTGTCTTGAAATTATCTGTGAT[C/T]TG CTATAGAGCAATACTTTTGTAAT	97
ASPA	443	rs59930743	GTGGGTATATGCAGCTCTATGCACTA[C/T]CT GCTCATTTATTTGGTAAATCTAA	98
ASPA	443	rs59690349	TGTGTGTGTGTGCGTGTGTGTGTGTG[-/T GTGTGTG]ATCATAAGAGTGGCTGCAGCAA ACT	99
ASPA	443	rs59676360	AGTCTGGAGTGCAATGGTGCAATCTC[A/G]GC TCACTGCAGCCTCCACCTCCGGG	100
ASPA	443	rs59335404	CTCCTAATGGATATTTCCCTAAATTTT[G/T]CTG AACAGAATTTAACTTGAGCTGG	101
ASPA	443	rs58879097	ATTTAAAAATGGATTTCTAGAAAAAC[A/G]AT CACATACTTGAATATTTTAGCAA	102
ASPA	443	rs58686774	CTATAAATGGGTAGCATGAGGGATTC[A/G]A GGAGGTGGCTGAAAGAAGCACGTA	103
ASPA	443	rs57511162	AAGAAACCAAGCATAGTAGAGTGTTA[A/G]A AAACCAAGCAACTAACCACTGT	104
ASPA	443	rs55859596	CGGGGCTCAGAACTTGTACAGAAAA[A/T]T AAAATATACTCCACTCAAGGGAAT	105
ASPA	443	rs55742972	TACTACACTTCACGGTACTGTACTT[-/G TACTT]TTTTTCCAAATTGAAGGTTTTTGGC	106
ASPA	443	rs55640436	TTGTTTTTGTGTTTTTGTGTTTTT[-/G TTTTTGTGTTTT]TGAGATGGAGTCTCGCTCT GTCGCC	107
ASPA	443	rs36225687	TTGCTTACTACACTTCACGGATAC[-/T GTACT]TGACTTTTTTTTCCAAATTGAAGGT	108
ASPA	443	rs36051310	GAGGTGGCTGAAAGAAGCACGTATCC[-/C] TGATGGCATGGTTGCGGTTATATG	109
ASPA	443	rs36034906	GAGAAAAGCAGTTCCTGGAACACCCC[-/C] ACCCCTTAACCCCTTATCTCTGCTT	110
ASPA	443	rs36033666	TTACATATGTATACATGTGCCATGTT[-/T] GGTGTGCCGCACCCATTAACTCGTC	111
ASPA	443	rs35730123	CTTTTTCCAGATTTTTTTTTTTTTTTT[-/T] GAGACAGAGTTTCACTCTGTTGCC	112
ASPA	443	rs35629100	TTTGGAAATCTTAAGCTTTTATTGTTG[-/G] TGTCACAGAGAAACAGGATCTGTAT	113
ASPA	443	rs35614631	TACTTTAAGTTTTAGGGTACATGTGC[-/A] CCATGTGCAGGTTGTTACATATGT	114

TABLE 2-continued

Non-limiting examples of polymorphisms				
Gene name	GeneID	SNP ID	Mutation	SEQ ID NO.
ASPA	443	rs35225782	ATTTCATGACCAGCCACATAAAATGCAC[-/A] GTATTACTTCGCAAGCATGCCAATG	115
ASPA	443	rs35178659	GTGCACTAGAATTAGCTAAAGTGGGG[-/G] AAAAAAGATGCATTTGATGGTCTA	116
ASPA	443	rs35095578	AACCTCCACCTCCCAGGTTCAAGAGA[-/A] TTCTCTGCCTCAGCCTCCCAAGTA	117
ASPA	443	rs35002210	CCTCCCTGTGATCCGAAGTAGCAGAC[A/G]TA CTTAACCTTCCATGGTGGATTGTT	118
ASPA	443	rs34744839	AAAACATTATTATATCTAGAAAAAAA[-/A] TGTATCTTAACCATTTGTTGGGAAGTG	119
ASPA	443	rs34680506	TTGAAGGTAAAATCATAGGGAGTTGG[-/G] AGCTGTCCTCTTGCCTGAATCAGT	120
ASPA	443	rs34365618	ACTTGTGGCCTTTTGGAGAGGTTAG[-/CA] ACTCTGAAAACCTGTCCCTGGACC	121
ASPA	443	rs34275920	GAAGGAGAAAAAGAGAGGAAATAAGT[-/T] AAAATAATAAACACAATTAATAAAG	122
ASPA	443	rs34109510	TGTATACATGTGCCATGTTGGTGTGC[C/T]GC ACCCATTAACCTCGTCATTTAGCA	123
ASPA	443	rs34054576	TCACCTGTCACCTCCTATAGAACTTT[-/C] CCCTGACCCCTCCTCTATAGCATTA	124
ASPA	443	rs34015272	ATAAATGATCATCATTCACAGTAGGG[-/G] TTTTGTTTTGTTTTTTTTCTGGAAA	125
ASPA	443	rs34002091	ACAGACATATCTACAAACACACTTTT[-/T] CACATATTTGTGTAAGTCATTTATG	126
ASPA	443	rs28940574	AAAGACAATAAACTAACGCTCAATG[A/C]A AAAAGTATTGCTGCTGTTTACAT	127
ASPA	443	rs28940279	TACCGTGTACCCCGTGTTTGTGAATG[A/C]GG CCGCATATTACGAAAAGAAAGAA	128
ASPA	443	rs17850703	CAGGGCTGGAGGTAAAACCATTTATT[A/G]CT AACCCAGAGCAGTGAAGAAGTG	129
ASPA	443	rs17222495	TTCTTCATTGCCTATTGAAGAGAGAG[C/T]GG AATGCTTTGGTTGCCAGATATGG	130
ASPA	443	rs17175228	CACAAGATCTCATTACTCAGGAGCTG[C/T]CC AAGTGTCTAATGTACTTAGTTAA	131
ASPA	443	rs16953074	TTCTGTGTAACATTTCAATTAAGCAA[A/G]GG ATTTCGGCAAATCAAAAATTGTCA	132
ASPA	443	rs16953070	TAAAACGTATTGAAGGTATTATTGAC[G/T]CT GTTGAAGCAAAGAGAACAAAACA	133
HEXA	3073	rs62022858	ATCTGCTCTTCCAGTTGGATGACAAG[C/T]CT TGCTGTCTAACACCTGCTGCAGA	134
HEXA	3073	rs62022857	CCATTTTTTGTGTATTTTTTTTTTC[C/T]TGAA TACTTTTTATCGCAGTTGGTT	135
HEXA	3073	rs62017872	CCCTGTCTCTAAAAGAAAAAAAAAAAA[A/G]A AAAAAAAAAGAAAACAAAACCCAA	136
HEXA	3073	rs62017871	AGTGGCTCCAAAAGGTCATGGAACC[C/T]CT TGAGGATGATGCAAATTGACTCT	137
HEXA	3073	rs61662730	TAAAGTTACTTTTCTTTTATGACTT[C/T]CCC TTATTTTTTAACCTTATGCTTT	138
HEXA	3073	rs61329913	CAGAGTTAAAAAAAAAAAAAAAAAA[-/A] GGAAGTAGCAGCAACAGCTTGGAAA	139

TABLE 2-continued

Non-limiting examples of polymorphisms				
Gene name	GeneID	SNP ID	Mutation	SEQ ID NO:
HEXA	3073	rs60920713	GTTGCCCGGGTTGAGTGCAGAGGCA[C/T]AT TTGGCTCACAGCAACCTCTGCC	140
HEXA	3073	rs60783213	AAGGCTTTTTTTTTTTTTTTTTTTT[-/T TTT]GAGACAGAGTCTTGCTGTGTCAACC	141
HEXA	3073	rs60644867	GCCTACATTCTGCAAAGAGGAGGGAA[C/G]A TTCACAGCTCCATACTTGAACCT	142
HEXA	3073	rs60288568	CCAAAGGAGAATAGCTCTAGGGGAGG[C/G]A GGTGGATGAGTATGCATGGGGGAG	143
HEXA	3073	rs59888548	GACTCCATCTCAAAAAAAAAAAAAA[-/A] TGCAGTCTAATGGCAGAATTAGACT	144
HEXA	3073	rs59733856	TTATTTATTTATTTATTTATTTTGA[A/G]ACA GGTCTCTGTGTCCAGGCTGG	145
HEXA	3073	rs59427837	TTTGAGGCAGGGTCTCACTCTGTTG[C/T]CC AGGGTTGAGTGCAGAGGCACATC	146
HEXA	3073	rs59171976	CGCCTTGCGAAGGCCCCACAGCTTGC[C/T]TG TGACAAACGTTTCATAGGCAATG	147
HEXA	3073	rs58706602	GGAGGTCTGTACAAAGCACCACTAC[C/T]TC ATGGGTCAGTTCCACAGCAGAA	148
HEXA	3073	rs58696963	GAATCTTATAATCACTGTGTACCTC[-/C CTC]TGTTTCATATTTTCGCAATTGAACT	149
HEXA	3073	rs58610850	AACATAGTATCTAATATAGCTTTTACA[C/T]CC AAAGCCAAAATATGAATACACTG	150
HEXA	3073	rs58016062	TTGTTTTGTTTTGTTTTGGGGGGGGG[-/G] TTGTTTTTCTGAGAGGGAGTCTTGC	151
HEXA	3073	rs57733983	CATACCAAAGGGCAGCTGGAGGGATAC[C/T]A GACGGAAGTCATGTGGAGAGTGAA	152
HEXA	3073	rs57476645	CAGGTGTGAGCCACCACGACCACCAA[A/T]T TAGCTCTTTTACTCCTTCCTTC	153
HEXA	3073	rs56870003	AGTGGTAGCTGATTTTGCTTCTGGAT[A/C]CT TTGCCACCTTCCCACTCTTTAAT	154
HEXA	3073	rs56338339	AAAGACCTGTTTCTTAAAAAAAAAAAA[-/A GAAAAAAAAAAAAA]GAAAGAAAAGAAAAG AAAAAACAG	155
HEXA	3073	rs55995352	TAAAAAATCTTTCAATGAGGAGATGT[C/T]CC CAGAGCAAGACAGCTGTAGGATG	156
HEXA	3073	rs55860138	AAAAGAAAAAAAAAAAAAAAAAAAAA[-/A] GAAACAAAACCCAAACCCATAAG	157
HEXA	3073	rs55743646	CCTGTCTCTAAAAGAAAAAAAAAAAAA[A/G]A AAAAAAAAAGAAAACAAACCCAAA	158
HEXA	3073	rs55665666	GTTATCATAGAAAAATATCACTCT[-/GT] CTGTATCCCACTTCCAGAACTGT	159
HEXA	3073	rs36106892	CAGGAGCTCATAGAATTACATACAAT[-/C] TTTTTTTTTTTTTTTGGAGACAGCG	160
HEXA	3073	rs36091525	TTGAGAATCTTATAATCACTGTGTA[-/C CTC]CCTCTGTTTCATATTTTCGCAATTG	161
HEXA	3073	rs35949555	CCACTACCACAGTGCCTAGAGAACA[C/T]A TGTGTTAATAATATTAAATAAT	162
HEXA	3073	rs35827424	CCCTGTCTCTAAAAGAAAAAAAAAAAAA[-/A] AAAAAAAAAGAAAACAAACCCCAA	163
HEXA	3073	rs35729578	CCATTATATCATTCATTCCCACTCA[-/T] TTCTTCATTCCAACCAAGATATAT	164

TABLE 2-continued

Non-limiting examples of polymorphisms				
Gene name	GeneID	SNP ID	Mutation	SEQ ID NO.
HEXA	3073	rs35649102	TCCGTCTCAAAAAAAAAAAAAAG[-/A] GAAAGGAATTATTCTCATGTATACA	165
HEXA	3073	rs35118677	CTGGGGCAGTTAAAAAGAAAAACAA[-/C] CCCTGGTCCCTGCCCTTGAGGAGAT	166
HEXA	3073	rs35005352	CTCCAGGGTCCCATTCCAGGACCACA[-/C] GCCTGCTACCTCTGCAGCTCACTCA	167
HEXA	3073	rs34736306	GGATTGACATATACCAGTTAGACGGA[-/T] TTTTTTTTTCCATAAACCAAGGCTCA	168
HEXA	3073	rs34607939	ACAAATAATTACTACATATCTACAAC[A/G]TT CCAGATACAGAAGAAATGGCCAA	169
HEXA	3073	rs34496117	TAAACACACTTGAAACATCATATAAA[-/A TG]ATATTACTACAAGACTTAACCGTAA	170
HEXA	3073	rs34300017	ACACAGGTAATCCATGTTTATTATAG[-/A] AAAATGCCACATTACTCTTTATTGA	171
HEXA	3073	rs34206496	AGTTATCATAGAAAAATATCACACTC[-/TG] TCTGTATCCCACTTCCAGAACTG	172
HEXA	3073	rs34110830	AATGAACCTTACAGGAAGGTAATATAT[-/G] GGAAATAAACATCTTATTGAATTTA	173
HEXA	3073	rs34093438	GGACCCCTGAAAGGCACAAGACACCC[-/T] TTCAGGTTCACTTCTCGAAAGCT	174
HEXA	3073	rs34085965	CCACCAATCACCAGAGCCTTCTGCTC[A/G]GG GGTACCTGAGGAAAACAAGCAA	175
HEXA	3073	rs34004907	AAAGACTGAAAAAACATTCTATACTA[-/T] TTTTCTTGTATCCTCGAAATGTC	176
HEXA	3073	rs28942072	TATCTTCATCTTGGAGGAGATGAGGT[C/T]GA TTTCACCTGCTGGAAGTCCAACC	177
HEXA	3073	rs28942071	TTGCCTATGAACGTTTGTACACTTC[C/T]GCT GTGAGTTGCTGAGCGAGGTGT	178
HEXA	3073	rs28941771	GCTTGCTGTTGGATACATCTCGCCAT[C/T]AC CTGCCACTCTCTAGCATCTTGA	179
HEXA	3073	rs28941770	CCGGGGCTTGCTGTTGGATACATCTC[G/T]CC ATTACCTGCCACTCTCTAGCATC	180

3. Nucleic Acid Target Length Evaluation:

In some embodiments, aspects of the invention relate to methods for detecting nucleic acid deletions or insertions in regions containing nucleic acid sequence repeats.

Genomic regions that contain nucleic acid sequence repeats are often the site of genetic instability due to the amplification or contraction of the number of sequence repeats (e.g., the insertion or deletion of one or more units of the repeated sequence). Instability in the length of genomic regions that contain high numbers of repeat sequences has been associated with a number of hereditary and non hereditary diseases and conditions.

For example, "Fragile X syndrome, or Martin-Bell syndrome, is a genetic syndrome which results in a spectrum of characteristic physical, intellectual, emotional and behavioral features which range from severe to mild in manifestation. The syndrome is associated with the expansion of a single trinucleotide gene sequence (CGG) on the X chromosome, and results in a failure to express the FMR-1 protein which is required for normal neural development.

There are four generally accepted forms of Fragile X syndrome which relate to the length of the repeated CGG sequence; Normal (29-31 CGG repeats), Premutation (55-200 CGG repeats), Full Mutation (more than 200 CGG repeats), and Intermediate or Gray Zone Alleles (40-60 repeats)."

Other examples include cancer, which has been associated with microsatellite instability (MSI) involving an increase or decrease in the genomic copy number of nucleic acid repeats at one or more microsatellite loci (e.g., BAT-25 and/or BAT-26). There are currently many sequencing-based assays for determining the number of nucleic acid sequence repeats at a particular locus and identifying the presence of nucleic acid insertions or deletions. However, such techniques are not useful in a high throughput multiplex analysis where the entire length of a region may not be sequenced.

In contrast, in some embodiments, aspects of the invention relate to detecting the presence of an insertion or deletion at a genomic locus without requiring the locus to be sequenced (or without requiring the entire locus to be

sequenced). Aspects of the invention are particularly useful for detecting an insertion or deletion in a nucleic acid region that contains high levels of sequence repeats. The presence of sequence repeats at a genetic locus is often associated with relatively high levels of polymorphism in a population due to insertions or deletions of one or more of the sequence repeats at the locus. The polymorphisms can be associated with diseases or predisposition to diseases (e.g., certain polymorphic alleles are recessive alleles associated with a disease or condition). However, the presence of sequence repeats often complicates the analysis of a genetic locus and increases the risk of errors when using sequencing techniques to determine the precise sequence and number of repeats at that locus.

In some embodiments, aspects of the invention relate to determining the size of a genetic locus by evaluating the capture frequency of a portion of that locus suspected of containing an insertion or deletion (e.g., due to the presence of sequence repeats) using a nucleic acid capture technique (e.g., a nucleic acid sequence capture technique based on molecular inversion probe technology). According to aspects of the invention, a statistically significant difference in capture efficiency for a genetic locus of interest in different biological samples (e.g., from different subjects) is indicative of different relative lengths in those samples. It should be appreciated that the length differences may be at one or both alleles of the genetic locus. Accordingly, aspects of the invention may be used to identify polymorphisms regardless of whether biological samples being interrogated at heterozygous or homozygous for the polymorphisms. According to aspects of the invention, subjects that contain one or more loci with an insertion or deletion can be identified by analyzing capture efficiencies for nucleic acids obtained from one or more biological samples using appropriate controls (e.g., capture efficiencies for known nucleic acid sizes, capture efficiencies for other regions that are not suspected of containing an insertion or deletion in the biological sample(s), or predetermined reference capture efficiencies, or any combination thereof. However, it should be appreciated that aspects of the invention are not limited by the nature or presence of the control. In some embodiments, if a statistically significant variation in capture efficiency is detected, a subject may be identified as being at risk for a disease or condition associated with insertions or deletions at that genetic locus. In some embodiments, the subject may be analyzed in greater detail in order to determine the precise nature of the insertion or deletion and whether the subject is heterozygous or homozygous for one or more insertions or deletions. For example, gel electrophoresis of an amplification (e.g., PCR) product of the locus, or Southern blotting, or any combination thereof can be used as an orthogonal approach to verify the length of the locus. In some embodiments, a more exhaustive and detailed sequence analysis of the locus can be performed to identify the number and types of insertions and deletions. However, other techniques may be used to further analyze a locus identified as having an abnormal length according to aspects of the invention.

Accordingly, aspects of the invention relate to detecting abnormal nucleic acid lengths in genomic regions of interest. In some embodiments, the invention aims to estimate the size of genomic regions that are hard to be accessed, such as repetitive elements. However, it should be appreciated that methods of the invention do not require that the precise length be estimated. In some embodiments, it is sufficient to determine that one or more alleles with abnormal lengths are

present at a locus of interest (e.g., based on the detection of abnormal capture efficiencies).

In a non-limiting example, fragile X can be used to illustrate aspects of the invention where the size of trinucleotide repeats (genotype) is linked to a symptom (phenotype). However, it should be appreciated that fragile X is a non-limiting example and similar analyses may be performed for other genetic loci (e.g., independently or simultaneously in multiplex analyses).

Use of molecular inversion probes (MIPs) has been demonstrated for detection of single nucleotide polymorphisms (Hardenbol et al. 2005 *Genome Res* 15:269-75) and for preparative amplification of large sets of exons (Porreca et al. 2007 *Nat Methods* 4:931-6, Krishnakumar et al. 2008 *Proc Natl Acad Sci USA* 105:9296-301). In both cases, oligonucleotide probes are designed which have ends ('targeting arms') that hybridize up-stream and down-stream of the locus that is to be amplified.

In some embodiments, aspects of the invention are based on the recognition that the effect of length on probe capturing efficiency can be used in the context of an assay (e.g., a high throughput and/or multiplex assay) to allow the length of sequences to be determined without requiring sequencing of the entire region being evaluated. This is particularly useful for repeat regions that are prone to changes in size. As illustrated in FIG. 8, which is reproduced from Deng et al., *Nature Biotech.* 27:353-60, (see Supplemental FIG. 1G of Deng et al.) illustrates that shorter sequences are captured with higher efficiency than longer sequences using MIPs. The statistical package R and its effects module were used for this analysis. A linear model was used, and each individual factor was assumed to be independent. The dashed lines represent a 95% confidence interval. Shorter target sequences were captured with higher efficiency than long target sequences ($p < 2 \times 10^{-16}$). However, the use of this differential capture efficiency for systematic sequence length analysis was not previously recognized.

In some embodiments, following probe hybridization, polymerase fill-in and ligation reactions are performed to convert the hybridized probe to a covalently-closed, circular molecule containing the desired target. PCR or rolling circle amplification plus exonuclease digestion of non-circularized material is performed to isolate and amplify the circular targets from the starting nucleic acid pool. Since one of the main benefits of the method is the potential for a high degree of multiplexing, generally thousands of targets are captured in a single reaction containing thousands of probes.

According to aspects of the invention, repetitive regions are surrounded by non repetitive unique sequences, which can be used to amplify the repeat-containing regions using, for example, PCR or padlock (MIP)-based method.

In addition to the repetitive regions, a probe (e.g., a MIP or padlock probe) can be designed to include at least a sequence that is sufficient to be uniquely identified in the genome (or target pool). After the probe is circularized and amplified, the amplicon can be end-sequenced so that the unique sequence can be identified and served as the "representative" of the repetitive region as illustrated in FIG. 9. FIG. 9 illustrates a non-limiting scheme of padlock (MIP) capture of a region that includes both repetitive regions (thick wavy line) and the adjacent unique sequence (thick straight line). The regions of the probe are indicated with the targeting arms shown as regions "1" and "3." An intervening region that may be, or include, a sequencing primer binding site is shown as "2." After the padlock is circularized and

amplified, it can be end-sequenced to obtain the sequence of the unique sequence, which represents the repetitive region of interest.

Although capturing efficiency is overall negatively correlated with target length, different probe sequences may have unique features. Therefore, multiple probes could be designed and tested so that an optimal one is chosen to be sensitive enough to differentiate repetitive sizes of roughly 0-150 bp, 150-600 bp, and beyond, which represent normal, premutation and full mutation of fragile X syndrome, respectively. However, it should be appreciated that other probe sizes and sequences can be designed, and optionally optimized, to distinguish a range of repeat region size differences (e.g., length differences of about 3-30 bases, about 30-60 bases, about 60-90 bases, about 90-120 bases, about 120-150 bases, about 150-300 bases, about 300-600 bases, about 600-900 bases, or any intermediate or longer length difference). It should be appreciated that a length difference may be an increase in size or a decrease in size.

In some embodiments, an initial determination of an unexpected capture frequency is indicative of the presence of size difference. In some embodiments, an increase in capture frequency is indicative of a deletion. In some embodiments, a decrease in capture frequency is indicative of an insertion. However, it should be appreciated that depending on specific sequence parameters and the relative sizes of the capture probes, the target region, and the deletions or insertions, a change in capture frequency can be associated with either an increase or decrease in target region length. In some embodiments, the precise nature of the change can be determined using one or more additional techniques as described herein.

Accordingly, in some aspects a MIP probe includes a linear nucleic acid strand that contains two hybridization sequences or targeting arms, one at each end of the linear probe, wherein each of the hybridization sequences is complementary to a separate sequence on a the same strand of a target nucleic acid, and wherein these sequences on the target nucleic acid flank the two ends of the target nucleic acid sequence of interest. It should be appreciated that upon hybridization, the two ends of the probe are inverted with respect to each other in the sense that both 5' and 3' ends of the probe hybridize to the same strand to separate regions flanking the target region (as illustrated in FIG. 9 for example).

In some embodiments, the hybridization sequences are between about 10-100 nucleotides long, for example between about 10-30, about 30-60, about 60-90, or about 20, about 30, about 40, or about 50 nucleotides long. However, other lengths may be used depending on the application. In some embodiments, the hybridization Tms of both targeting arms of a probe are designed or selected to be similar. In some embodiments, the hybridization Tms of the targeting arms of a plurality of probes designed to capture different target regions are selected or designed to be similar so that they can be used together in a multiplex reaction. Accordingly, a typical size of a MIP probe prior to fill in is about 60-80 nucleotides long. However, other sizes can be used depending on the sizes of the targeting arms and any other sequences (e.g., primer binding or tag sequences) that are present in the MIP probe. In some embodiments, MIP probes are designed to avoid sequence-dependent secondary structures. In some embodiments, MIP probes are designed such that the targeting arms do not overlap with known polymorphic regions. In some embodiments, targeting arms that can be used for capturing the repeat region of the Fragile X locus

can have the following sequences or complementary to these sequences depending on the strand that is captured.

left: CTCUITYCGGTITCACTTC (SEQ ID NO: 181)

right: ATCTTCTCTTCAGCCCTGCT (SEQ ID NO: 182)

The typical captured size using these targeting arms is about 100 nucleotides in length (e.g., about 30 repeats of a tri-nucleotide repeat).

In some embodiments, the number of reads obtained for the “representative” of the repetitive region is not informative to estimate the target length because it is dependent on the total number of reads obtained. To overcome this, it is useful to include one or more probes that target other “control” regions where no or minimal polymorphism exists among populations. Because of the systematic consistency of capturing efficiency (see, e.g., FIG. 9), the ratio of reads obtained for the repetitive “representative” to reads obtained for the control region(s) will be tuned using DNA with defined numbers of repeats. Ultimately, the ratio can serve as a measure of the repeat length as illustrated in FIG. 10. FIG. 10 illustrates a non-limiting hypothetical relationship between target gap size and the relative number of reads of the repetitive region, which is measured by the ratio of the repeat “representative” reads vs. the “control” region reads. The unit of y-axis is arbitrary.

In some embodiments, to better tell targets with similar size range apart, the whole repetitive region can be sequenced by making a shotgun library (e.g., by making a shotgun library from a captured sequence, for example a sequence captured using a MIP probe). The longer the repeat is, the more short reads of repeats will be obtained. Therefore, the target length will contribute twice to the relative number of “repetitive” reads, which will gain better resolution of differentiating targets. In some embodiments, the expectation is that the number of reads from any given repeat will be a direct function of the number of repeats present. However, in some embodiments, a Poisson sampling induced spread may need to be considered and in some embodiments may be sufficiently large to limit the resolution.

When a precise measurement of the length of both alleles from a diploid sample is desired, further manipulations may be required. This is because the capture efficiency measured will actually be the average efficiency of the two alleles. To effectively achieve separate measurements for each allele, barcodes (e.g., sequence tags) can be used that allow the efficiency of individual capture events (from individual genomic loci) to be followed. FIG. 11A-C shows the approach. For a given locus, MIPs are synthesized to contain one of a large number differentiator tags in their backbone such that the probability of any two MIPs in a reaction having the same differentiator tag sequence is low. MIP capture is performed on the sample; the reaction will be biased for shorter target lengths, and therefore the reaction product will be comprised of more ‘short’ circles than ‘long’ circles. Each circle should bear a unique differentiator tag sequence. Then, linear RCA (1RCA) is performed on the circles. In the 1RCA reaction, circles are converted into long, linear concatemers of themselves. The 1RCA reaction for a given circle stops when the concatemer has reached a ‘fixed’ length (based on the processivity/error rate of the polymerase). Concatemers derived from smaller circles will therefore contain more copies of the differentiator tag, and concatemers derived from larger circles will contain fewer copies of the differentiator tag. The number of each differentiator tag sequence is counted, for example, by next-generation sequencing.

61

When number of occurrences is plotted against differentiator tag ID, the data will naturally cluster into two groups reflecting the lengths of the two alleles in the diploid sample. The allele lengths can therefore be read directly off this graph, after absolute length calibration using known standards. In some embodiments, a sequencing technique (e.g., a next-generation sequencing technique) is used to sequence part of one or more captured targets (e.g., or amplicons thereof) and the sequences are used to count the number of different barcodes that are present. Accordingly, in some 5 10
embodiments, aspects of the invention relate to a highly-multiplexed qPCR reaction.

62

Other non-limiting examples of loci at which insertions or deletions or repeat sequences may be associated with a disease or condition are provided in Tables 3 and 4. It should be appreciated that the presence of an abnormal length at any one or more of these loci may be evaluated according to aspects of the invention. In some embodiments, two or more of these loci or other loci may be evaluated in a single multiplex reaction using different probes designed to hybridize under the same reaction conditions to different target nucleic acid in a biological sample.

TABLE 3

Polyglutamine (PolyQ) Diseases				
Type	Gene	Normal/ wildtype	Pathogenic	
DRPLA (Dentatorubropallidoluyian atrophy)	ATN1 or DRPLA	6-35	49-88	
HD (Huntington's disease)	HTT (Huntingtin)	10-35	35+	
SBMA (Spinobulbar muscular atrophy or Kennedy disease)	Androgen receptor on the X chromosome.	9-36	38-62	
SCA1 (Spinocerebellar ataxia Type 1)	ATXN1	6-35	49-88	
SCA2 (Spinocerebellar ataxia Type 2)	ATXN2	14-32	33-77	
SCA3 (Spinocerebellar ataxia Type 3 or Machado-Joseph disease)	ATXN3	12-40	55-86	
SCA6 (Spinocerebellar ataxia Type 6)	CACNA1A	4-18	21-30	
SCA7 (Spinocerebellar ataxia Type 7)	ATXN7	7-17	38-120	
SCA17 (Spinocerebellar ataxia Type 17)	TBP	25-42	47-63	

TABLE 4

Non-Polyglutamine Diseases				
Type	Gene	Codon	Normal/ wildtype	Pathogenic
FRAXA (Fragile X syndrome)	FMR1, on the X- chromosome	CGG	6-53	230+
FXTAS (Fragile X-associated tremor/ataxia syndrome)	FMR1, on the X- chromosome	CGG	6-53	55-200
FRAXE (Fragile XE mental retardation)	AFF2 or FMR2, on the X- chromosome	GCC	6-35	200+
FRDA (Friedreich's ataxia)	FXN or X25, (frataxin)	GAA	7-34	100+

TABLE 4-continued

Non-Polyglutamine Diseases				
Type	Gene	Codon	Normal/ wildtype	Pathogenic
DM (Myotonic dystrophy)	DMPK	CTG	5-37	50+
SCA8 (Spinocerebellar ataxia Type 8)	OSCA or SCA8	CTG	16-37	110-250
SCA12 (Spinocerebellar ataxia Type 12)	PPP2R2B or SCA12	CAG On 5' end	7-28	66-78

The following examples illustrate aspects and embodiments of the invention and are not intended to be limiting or restrictive. Many variations of the invention will become apparent to those skilled in the art upon review of this specification. The full scope of the invention should be determined by reference to the claims, along with their full scope of equivalents, and the specification, along with such variations.

4. Increasing Detection Sensitivity:

In some embodiments, aspects of the invention relate to methods for increasing the sensitivity of nucleic acid detection assays.

There are currently many genomic assays that utilize next-generation (e.g., polony-based) sequencing to generate data, including genome resequencing, RNA-seq for gene expression, bisulphite sequencing for methylation, and Immune-seq, among others. In order to make quantitative measurements (including genotype calling), these methods utilize the counts of sequencing reads of a given genomic locus as a proxy for the representation of that sequence in the original sample of nucleic acids. The majority of these techniques require a preparative step to construct a high-complexity library of DNA molecules that is representative of a sample of interest. Current assays use one of several alternative nucleic acid preparative techniques (e.g., amplification, for example PCR-based amplification; sequence-specific capture, for example, using immobilized capture probes; or target capture into a circularized probe followed by a sequence analysis step. In order to reduce errors associated with the unpredictability (stochastic nature) of nucleic acid isolation and sequence analysis techniques, current methods involve oversampling a target nucleic acid preparation in order to increase the likelihood that all sequences that are present in the original nucleic acid sample will be represented in the final sequence data. For example, a genomic sequencing library may contain an over- or under-representation of particular sequences from a source nucleic acid sample (e.g., genome preparation) as a result of stochastic variations in the library construction process. Such variations can be particularly problematic when they result in target sequences from a genome being absent or undetectable in a sequencing library. For example, an under-representation of particular allelic sequences (e.g., heterozygotic alleles) from a genome in a sequencing library can result in an apparent homozygous representation in a sequencing library.

In contrast, aspects of the invention relate to basing a nucleic acid sequence analysis on results from two or more different nucleic acid preparatory techniques that have different systematic biases in the types of nucleic acids that

they sample rather than simply oversampling the target nucleic acid. According to some embodiments, different techniques have different sequence biases that are systematic and not simply due to stochastic effects during nucleic acid capture or amplification. Accordingly, in some embodiments, the degree of oversampling required to overcome variations in nucleic acid preparation needs to be sufficient to overcome the biases. In some embodiments, the invention provides methods that reduce the need for oversampling by combining nucleic acid and/or sequence results obtained from two or more different nucleic acid preparative techniques that have different biases.

According to the invention, different techniques have different characteristic or systematic biases. For example, one technique may bias a sample analysis towards one particular allele at a genetic locus of interest, whereas a different technique would bias the sample analysis towards a different allele at the same locus. Accordingly, the same sample may be identified as being different depending on the type of technique that is used to prepare nucleic acid for sequence analysis. This effectively represents a sensitivity issue, because each technique has a different relative sensitivities for polymorphic sequences of interest.

According to aspects of the invention, the sensitivity of a nucleic acid analysis can be increased by combining the sequences from different nucleic acid preparative steps and using the combined sequence information for a diagnostic assay (e.g., for a making a call as to whether a subject is homozygous or heterozygous at a genetic locus of interest).

Currently, the ability of DNA sequencing to detect mutations is limited by the ability of the upstream sample isolation (e.g., by amplification, immobilization enrichment, circularization capture, etc.) methods to reliably isolate the locus of interest.

If one wishes to make heterozygote base-calls for a diploid genome (e.g. a human sample presented for molecular diagnostic sequencing), it is important in some embodiments that the isolation method produces near- or perfectly-uniform amounts of the two alleles to be sequenced (at least sufficiently uniform to be "called" unambiguously as a heterozygote or a homozygote for a locus of interest).

Sample preparative methods may fall into three classes: 1) single- or several target amplification (e.g., uniplex PCR, 'multiplex' PCR), 2) multi-target hybridization enrichment (e.g., Agilent SureSelect 'hybrid capture' [Gnirke et al 2009, *Nature methods* 27:182-9], Roche/Nimblegen 'sequence capture' [Hodges et al 2007, *Nature genetics* 39:1522-7], and 3) multi-target circularization selection (e.g. molecular inversion probes or padlock probes, [Porreca et al 2007, *Nature methods* 4:931-6, Turner et al 2009, *Nature methods*

6:315-6], 'selectors' [Dahl et al 2005, *Nucleic acids research* 33:e71]). Each of these methods can result in a pool of isolated product that does not adequately represent the input abundance distribution. For example, the two alleles at a heterozygous position can become skewed far from their input 50:50 ratio to something that results in a missed basecall during downstream sequencing. For example, if the ratio was skewed from 50:50 to 10:90, and the sample was sequenced to 10x average coverage, there is a high probability that one of the two alleles would not be observed once in the ten sequencing reads. This would reduce the sensitivity of the sequencing method by converting a heterozygous position to homozygous (where potentially the 'mutant' allele was the one not observed). In some embodiments, a skewed ratio is a particular issue that decreases the sensitivity of detecting mutations present in a heterogeneous tumor tissue. For example, if only 10% of the cells analyzed in a heterogeneous sample harbored a heterozygous mutation, the mutation would be expected to be present in 5% of sequence reads, not 50%. In this scenario, the need for robust, sensitive detection may be even more acute.

The methods disclosed herein are based, in part, on the discovery that certain classes of isolation methods have different modes of bias. The disclosure provides methods for increasing the sensitivity of the downstream sequencing by using a combination of multiple isolation methods (e.g., one or more from at least two of the classes disclosed herein) for a sample. This is particularly important in molecular diagnostics where high sensitivity is required to minimize the chances of 'missing' a disease-associated mutation. For example, given a nominal false-negative error rate of 1×10^{-3} for sequencing following circularization selection, and a false-negative error rate of 1×10^{-3} for sequencing following hybridization enrichment, one can achieve a final false-negative rate of 1×10^{-6} by performing both techniques on the sample (assuming failures in each method are fully independent). For a recessive disease with carrier frequency of 0.1, caused by a single fully-penetrant mutant allele, the number of missed carrier diagnoses would decrease from 1000 per million patients tested to 1 per million patients tested. Furthermore, if the testing was used in the context of prenatal carrier screening, the number of affected children born as a result of missing the carrier call in one parent would decrease from 25 per million to 25 per billion born.

Additionally, the disclosure provides combinations of preparative methods to effectively increase sequencing coverage in regions containing disease-associated alleles. Since heterozygote error rate is largely tied to both deviations from 50:50 allele representation, and in the case of next-generation DNA sequencing deviations from average abundance (such that less abundant isolated targets are more likely to be undersampled at one or both alleles), selectively increasing coverage in these regions will also selectively increase sensitivity. Furthermore, MIPs that detect presence or absence of specific known disease-associated mutations can be used to increase sensitivity selectively. In some embodiments, these MIPs would have a targeting arm whose 3'-most region is complementary to the expected mutation, and has a fill-in length of 0 or more bp. Thus, the MIP will form only if the mutation is present, and its presence will be detected by sequencing.

Additionally, algorithms disclosed herein may be used to determine base identity with varying levels of stringency depending on whether the given position has any known disease-associated alleles. Stringency can be reduced in such positions by decreasing the minimum number of observed mutant reads necessary to make a consensus base-call. This

will effectively increase sensitivity for mutant allele detection at the cost of decreased specificity.

An embodiment of the invention combines MIPs plus hybridization enrichment, plus optionally extra MIPs targeted to specific known, common disease-associated loci, e.g., to detect the presence of a polymorphism in a target nucleic acid. A non-limiting example is illustrated in FIG. 12 that illustrates a schematic using MIPs plus hybridization enrichment, plus optionally extra MIPs targeted to specific known, common disease-associated loci, e.g., to detect the presence of a polymorphism in a target nucleic acid.

FIGS. 13 and 14 illustrate different capture efficiencies for MIP-based captures. FIG. 13 shows a graph of per-target abundance with MIP capture. In this graph, bias largely drives the heterozygote error rate, since targets which are less abundant here are less likely to be covered in sufficient depth during sequencing to adequately sample both alleles. This is from Turner et al 2009, *Nature methods* 6:315-6. Hybridization enrichment results in a qualitatively similar abundance distribution, but the abundance of a given target is likely not correlated between the two methods. FIG. 14 shows a graph of correlation between two MIP capture reactions from Ball et al 2009, *Nature biotechnology* 27:361-8. Each point represents the target abundance in replicate 1 and replicate 2. Pearson correlation $r=0.956$. This indicates that MIP capture reproducibly biases targets to specific abundances. Hybridization enrichment is similarly correlated from one capture to the next.

According to aspects of the invention, such biases can be detected or overcome by systematically combining different capture and/or analytical techniques in an assay that interrogates a plurality of loci in a plurality of subject samples.

Accordingly, it should be appreciated that in any of the embodiments described herein (e.g., tiling/staggering, tagging, size-detection, sensitivity enhancing algorithms, or any combination thereof), aspects of the invention involve preparing genomic nucleic acid and/or contacting them with one or more different probes (e.g., capture probes, hybridization probes, MIPs, others etc.). In some embodiments, the amount of genomic nucleic acid used per subject ranges from 1 ng to 10 micrograms (e.g., 500 ng to 5 micrograms). However, higher or lower amounts (e.g., less than 1 ng, more than 10 micrograms, 10-50 micrograms, 50-100 micrograms or more) may be used. In some embodiments, for each locus of interest, the amount of probe used per assay may be optimized for a particular application. In some embodiments, the ratio (molar ratio, for example measured as a concentration ratio) of probe to genome equivalent (e.g., haploid or diploid genome equivalent, for example for each allele or for both alleles of a nucleic acid target or locus of interest) ranges from 1/100, 1/10, 1/1, 10/1, 100/1, 1000/1. However, lower, higher, or intermediate ratios may be used.

In some embodiments, the amount of target nucleic acid and probe used for each reaction is normalized to avoid any observed differences being caused by differences in concentrations or ratios. In some embodiments, in order to normalize genomic DNA and probe, the genomic DNA concentration is read using a standard spectrophotometer or by fluorescence (e.g., using a fluorescent intercalating dye). The probe concentration may be determined experimentally or using information specified by the probe manufacturer.

Similarly, once a locus has been captured (e.g., on a MIP or other probe or in another form), it may be amplified and/or sequenced in a reaction involving one or more primers. The amount of primer added for each reaction can range from 0.1 pmol to 1 nmol, 0.15 pmol to 1.5 nmol (for

example around 1.5 pmol). However, other amounts (e.g., lower, higher, or intermediate amounts) may be used.

In some embodiments, it should be appreciated that one or more intervening sequences (e.g., sequence between the first and second targeting arms on a MIP capture probe), identifier or tag sequences, or other probe sequences that are not designed to hybridize to a target sequence (e.g., a genomic target sequence) should be designed to avoid excessive complementarity (to avoid cross-hybridization) to target sequences or other sequences (e.g., other genomic sequences) that may be in a biological sample. For example, these sequences may be designed have a sufficient number of mismatches with any genomic sequence (e.g., at least 5, 10, 15, or more mismatches out of 30 bases) or as having a T_m (e.g., a mismatch T_m) that is lower (e.g., at least 5, 10, 15, 20, or more degrees C. lower) than the hybridization reaction temperature.

It should be appreciated that a targeting arm as used herein may be designed to hybridize (e.g., be complementary) to either strand of a genetic locus of interest if the nucleic acid being analyzed is DNA (e.g., genomic DNA). However, in the context of MIP probes, whichever strand is selected for one targeting arm will be used for the other one. However, in the context of RNA analysis, it should be appreciated that a targeting arm should be designed to hybridize to the transcribed RNA. It also should be appreciated that MIP probes referred to herein as “capturing” a target sequence are actually capturing it by template-based synthesis rather than by capturing the actual target molecule (other than for example in the initial stage when the arms hybridize to it or in the sense that the target molecule can remain bound to the extended MIP product until it is denatured or otherwise removed).

It should be appreciated that in some embodiments a targeting arm may include a sequence that is complementary to one allele or mutation (e.g., a SNP or other polymorphism, a mutation, etc.) so that the probe will preferentially hybridize (and capture) target nucleic acids having that allele or mutation. However, in many embodiments, each targeting arm is designed to hybridize (e.g., be complementary) to a sequence that is not polymorphic in the subjects of a population that is being evaluated. This allows target sequences to be captured and/or sequenced for all alleles and then the differences between subjects (e.g., calls of heterozygous or homozygous for one or more loci) can be based on the sequence information and/or the frequency as described herein.

It should be appreciated that sequence tags (also referred to as barcodes) may be designed to be unique in that they do not appear at other positions within a probe or a family of probes and they also do not appear within the sequences being targeted. Thus they can be used to uniquely identify (e.g., by sequencing or hybridization properties) particular probes having other characteristics (e.g., for particular subjects and/or for particular loci).

It also should be appreciated that in some embodiments probes or regions of probes or other nucleic acids are described herein as comprising or including certain sequences or sequence characteristics (e.g., length, other properties, etc.). However, it should be appreciated that in some embodiments, any of the probes or regions of probes or other nucleic acids consist of those regions (e.g., arms, central regions, tags, primer sites, etc., or any combination thereof) of consist of those sequences or have sequences with characteristics that consist of one or more characteristics (e.g., length, or other properties, etc.) as described herein in the context of any of the embodiments (e.g., for

tilled or staggered probes, tagged probes, length detection, sensitivity enhancing algorithms or any combination thereof).

It should be appreciated that probes, primers, and other nucleic acids designed or used herein may be synthetic, natural, or a combination thereof. Accordingly, as used herein, the term “nucleic acid” refers to multiple linked nucleotides (i.e., molecules comprising a sugar (e.g., ribose or deoxyribose) linked to an exchangeable organic base, which is either a pyrimidine (e.g., cytosine (C), thymidine (T) or uracil (U)) or a purine (e.g., adenine (A) or guanine (G)). “Nucleic acid” and “nucleic acid molecule” may be used interchangeably and refer to oligoribonucleotides as well as oligodeoxyribonucleotides. The terms shall also include polynucleosides (i.e., a polynucleotide minus a phosphate) and any other organic base containing nucleic acid.

The organic bases include adenine, uracil, guanine, thymine, cytosine and inosine. Unless otherwise stated, nucleic acids may be single or double stranded. The nucleic acid may be naturally or non-naturally occurring. Nucleic acids can be obtained from natural sources, or can be synthesized using a nucleic acid synthesizer (i.e., synthetic).

Harvest and isolation of nucleic acids are routinely performed in the art and suitable methods can be found in standard molecular biology textbooks. (See, for example, Maniatis’ Handbook of Molecular Biology.) The nucleic acid may be DNA or RNA, such as genomic DNA, mitochondrial DNA, mRNA, cDNA, rRNA, miRNA, or a combination thereof. Non-naturally occurring nucleic acids such as bacterial artificial chromosomes (BACs) and yeast artificial chromosomes (YACs) can also be used.

The invention also contemplates the use of nucleic acid derivatives. As will be described herein, the use of certain nucleic acid derivatives may increase the stability of the nucleic acids of the invention by preventing their digestion, particularly when they are exposed to biological samples that may contain nucleases. As used herein, a nucleic acid derivative is a non-naturally occurring nucleic acid or a unit thereof. Nucleic acid derivatives may contain non-naturally occurring elements such as non-naturally occurring nucleotides and non-naturally occurring backbone linkages.

Nucleic acid derivatives may contain backbone modifications such as but not limited to phosphorothioate linkages, phosphodiester modified nucleic acids, phosphorothiolate modifications, combinations of phosphodiester and phosphorothioate nucleic acid, methylphosphonate, alkylphosphonates, phosphate esters, alkylphosphonothioates, phosphoramidates, carbamates, carbonates, phosphate triesters, acetamidates, carboxymethyl esters, methylphosphorothioate, phosphorodithioate, p-ethoxy, and combinations thereof. The backbone composition of the nucleic acids may be homogeneous or heterogeneous.

Nucleic acid derivatives may contain substitutions or modifications in the sugars and/or bases. For example, they include nucleic acids having backbone sugars which are covalently attached to low molecular weight organic groups other than a hydroxyl group at the 3' position and other than a phosphate group at the 5' position (e.g., an 2'-O-alkylated ribose group). Nucleic acid derivatives may include non-ribose sugars such as arabinose. Nucleic acid derivatives may contain substituted purines and pyrimidines such as C-5 propyne modified bases, 5-methylcytosine, 2-aminopurine, 2-amino-6-chloropurine, 2,6-diaminopurine, hypoxanthine, 2-thiouracil and pseudoisocytosine. In some embodiments, substitution(s) may include one or more substitutions/modifications in the sugars/bases, groups attached to the base,

including biotin, fluorescent groups (fluorescein, cyanine, rhodamine, etc), chemically-reactive groups including carboxyl, NHS, thiol, etc., or any combination thereof.

A nucleic acid may be a peptide nucleic acid (PNA), locked nucleic acid (LNA), DNA, RNA, or co-nucleic acids of the same such as DNA-LNA co-nucleic acids. PNA are DNA analogs having their phosphate backbone replaced with 2-aminoethyl glycine residues linked to nucleotide bases through glycine amino nitrogen and methylenecarbonyl linkers. PNA can bind to both DNA and RNA targets by Watson-Crick base pairing, and in so doing form stronger hybrids than would be possible with DNA or RNA based oligonucleotides in some cases.

PNA are synthesized from monomers connected by a peptide bond (Nielsen, P. E. et al. *Peptide Nucleic Acids, Protocols and Applications*, Norfolk: Horizon Scientific Press, p. 1-19 (1999)). They can be built with standard solid phase peptide synthesis technology. PNA chemistry and synthesis allows for inclusion of amino acids and polypeptide sequences in the PNA design. For example, lysine residues can be used to introduce positive charges in the PNA backbone. All chemical approaches available for the modifications of amino acid side chains are directly applicable to PNA. Several types of PNA designs exist, and these include single strand PNA (ssPNA), bisPNA and pseudo-complementary PNA (pcPNA).

The structure of PNA/DNA complex depends on the particular PNA and its sequence. ssPNA binds to single stranded DNA (ssDNA) preferably in antiparallel orientation (i.e., with the N-terminus of the ssPNA aligned with the 3' terminus of the ssDNA) and with a Watson-Crick pairing. PNA also can bind to DNA with a Hoogsteen base pairing, and thereby forms triplexes with double stranded DNA (dsDNA) (Wittung, P. et al., *Biochemistry* 36:7973 (1997)).

A locked nucleic acid (LNA) is a modified RNA nucleotide. An LNA form hybrids with DNA, which are at least as stable as PNA/DNA hybrids (Braasch, D. A. et al., *Chem & Biol.* 8(1):1-7(2001)). Therefore, LNA can be used just as PNA molecules would be. LNA binding efficiency can be increased in some embodiments by adding positive charges to it. LNAs have been reported to have increased binding affinity inherently.

Commercial nucleic acid synthesizers and standard phosphoramidite chemistry are used to make LNAs. Therefore, production of mixed LNA/DNA sequences is as simple as that of mixed PNA/peptide sequences. The stabilization effect of LNA monomers is not an additive effect. The monomer influences conformation of sugar rings of neighboring deoxynucleotides shifting them to more stable configurations (Nielsen, P. E. et al. *Peptide Nucleic Acids, Protocols and Applications*, Norfolk: Horizon Scientific Press, p. 1-19 (1999)). Also, lesser number of LNA residues in the sequence dramatically improves accuracy of the synthesis. Most of biochemical approaches for nucleic acid conjugations are applicable to LNA/DNA constructs.

These and other aspects of the invention are illustrated by the following non-limiting examples.

EXAMPLES

The following examples illustrate non-limiting embodiments of the invention.

Example 1: Design a Set of Capture Probes for a Human Target Exon

All targets are captured as a set of partially-overlapping subtargets. For example, in the tiling approach, a 200 bp

target exon might be captured as a set of 12 subtargets, each 60 bp in length (FIG. 1). Each subtarget is chosen such that it partially overlaps two or three other targets.

In some embodiments, all probes are composed of three regions: 1) a 20 bp 'targeting arm' comprised of sequence which hybridizes immediately upstream from the sub-target, 2) a 30 bp 'constant region' comprised of sequence used as a pair of amplification priming sites, and 3) a second 20 bp 'targeting arm' comprised of sequence which hybridizes immediately downstream from the sub-target. Targeting arm sequences will be different for each capture probe in a set, while constant region sequence will be the same for all probes in the set, allowing all captured targets to be amplified with a single set of primers. Targeting arm sequences should be designed such that any given pair of 20 bp sequences is unique in the target genome (to prevent spurious capture of undesired sites). Additionally, melting temperatures should be matched for all probes in the set such that hybridization efficiency is uniform for all probes at a constant temperature (e.g., 60 C). Targeting arm sequences should be computationally screened to ensure they do not form strong secondary structure that would impair their ability to basepair with the genomic target.

Hybridize Capture Probes to Human Genomic Sample Assemble Hybridization Reaction:

1.0 ul capture probe mix (~2.5 μ mol)
2.0 ul 10 $\times$ Ampligase buffer (Epicentre)
6.0 ul 500 ng/ul human genomic DNA (~16.7 fmol)
11 ul dH₂O

In a thermal cycler, heat reaction to 95 C for 5 min to denature genomic DNA, then cool to 60 C. Allow to incubate at 60 C for 40 hours.

Convert Hybridized Probes into Covalently-Closed Circular Products Containing Subtargets

Prepare Fill-In/Ligation Reaction Mixture:

0.25 ul 2 mM dNTP mix (Invitrogen)
2.5 ul 10 $\times$ Ampligase buffer (Epicentre)
5.0 ul 5 U/ul Taq Stoffel fragment (Applied Biosystems)
12.5 ul 5 U/ul Ampligase (Epicentre)
4.75 ul dH₂O

Add 1.0 ul of this mix to the hybridized probe reaction, and incubate at 60 C for 10 hours.

Purify Circularized Probe/Subtarget Products from Un-Reacted Probes and Genomic DNA

Prepare Exonuclease Reaction Mixture:

21 ul fill-in/ligation reaction product
2.0 ul 10 $\times$ exonuclease I buffer (New England Biolabs)
2.0 ul 20 U/ul exonuclease I (New England Biolabs)
2.0 ul 100 U/ul exonuclease III (New England Biolabs)
Incubate at 37 C for 60 min, then heat-inactivate by incubating at 80 C for 15 min. Immediately cool to 4 C for storage.

Amplify Circular Material by PCR Using Primers Specific to the 'Constant Region' of the Probes

Prepare PCR Mixture:

5.0 ul 10 $\times$ Accuprime reaction buffer (Invitrogen)
1.5 ul 10 uM CP-2-FA (5'-GCAC-GATCCGACGGTAGTGT-3') (SEQ ID NO:183)
1.5 ul 10 uM CP-2-RA (5'-CCGTAATCGGGAAGCT-GAAG-3') (SEQ ID NO:184)
0.4 ul 25 mM dNTP mix (Invitrogen)
2.0 ul heat-inactivated exonuclease reaction mix
1.5 ul 10 $\times$ SybrGreen (Invitrogen)
0.4 ul 2.5 U/ul Accuprime Pfx polymerase (Invitrogen)
37.7 ul dH₂O

71

Thermal cycle in real-time thermal cycler according to the following protocol, but stop cycling before amplification yield plateaus (generally 8-12 cycles):

1. 95 C for 5 min
2. 95 C for 30 sec
3. 58 C for 60 sec
4. 72 C for 60 sec
5. goto 2, N more times

Prepare a Shotgun Next-Generation Sequencing Library for Analysis

Purify desired amplicon population from non-specific amplification products by gel extraction.

Concatemerize amplicons into high-molecular weight products suitable for shearing

Mechanically shear, using either a nebulizer, BioRuptor, Hydroshear, Covaris, or similar instrument. DNA should be sheared into fragments several hundred base-pairs in length.

Ligate adapters required for amplification by the sequencing platform used. If necessary, purify ligated product from unligated product and adapters.

Example 2: Use of Differentiator Tag Sequences to Detect and Correct Bias in a MIP-Capture Reaction of a Set of Exon Targets

The first step in performing the detection/correction is to determine how many differentiator tag sequences are necessary for the given sample. In this example, 1000 genomic targets corresponding to 1000 exons were captured. Since the differentiator tag sequence is part of the probe, it will measure/report biases that occur from the earliest protocol steps. Also, being located in the backbone, the differentiator tag sequence can easily be sequenced from a separate priming site, and therefore not impact the total achievable read-length for the target sequence.

MIP probes are synthesized using standard column-based oligonucleotide synthesis by any number of vendors (e.g. IDT), and differentiator tag sequences are introduced as 'degenerate' positions in the backbone. Each degenerate position increases the total number of differentiator tag sequences synthesized by a factor of 4, so a 10 nt degenerate region implies a differentiator tag sequence complexity of ~1e6 species.

Hybridize Capture Probes to Human Genomic Sample Assemble Hybridization Reaction:

- 1.0 ul capture probe mix (~2.5 pmol)
- 2.0 ul 10xAmpligase buffer (Epicentre)
- 6.0 ul 500 ng/ul human genomic DNA (~16.7 fmol)
- 11 ul dH2O

In a thermal cycler, heat reaction to 95 C for 5 min to denature genomic DNA, then cool to 60 C. Allow to incubate at 60 C for 40 hours.

Convert Hybridized Probes into Covalently-Closed Circular Products Containing Subtargets

Prepare Fill-In/Ligation Reaction Mixture:

- 0.25 ul 2 mM dNTP mix (Invitrogen)
- 2.5 ul 10xAmpligase buffer (Epicentre)
- 5.0 ul 5 U/ul Taq Stoffel fragment (Applied Biosystems)
- 12.5 ul 5 U/ul Ampligase (Epicentre)
- 4.75 ul dH2O

Add 1.0 ul of this mix to the hybridized probe reaction, and incubate at 60 C for 10 hours.

72

Purify Circularized Probe/Subtarget Products from Un-Reacted Probes and Genomic DNA

Prepare Exonuclease Reaction Mixture:

- 21 ul fill-in/ligation reaction product
- 2.0 ul 10xexonuclease I buffer (New England Biolabs)
- 2.0 ul 20 U/ul exonuclease I (New England Biolabs)
- 2.0 ul 100 U/ul exonuclease III (New England Biolabs)
- Incubate at 37 C for 60 min, then heat-inactivate by incubating at 80 C for 15 min. Immediately cool to 4 C for storage.

Amplify Circular Material by PCR Using Primers Specific to the 'Constant Region' of the Probes

Prepare PCR Mixture:

- 5.0 ul 10xAccuprime reaction buffer (Invitrogen)
- 1.5 ul 10 uM CP-2-FA (5'-GCAC-GATCCGACGGTAGTGT-3') (SEQ ID NO: 183)
- 1.5 ul 10 uM CP-2-RA (5'-CCGTAATCGGGAAGCT-GAAG-3') (SEQ ID NO: 184)
- 0.4 ul 25 mM dNTP mix (Invitrogen)
- 2.0 ul heat-inactivated exonuclease reaction mix
- 1.5 ul 10xSybrGreen (Invitrogen)
- 0.4 ul 2.5 U/ul Accuprime Pfx polymerase (Invitrogen)
- 37.7 ul dH2O

Thermal cycle in real-time thermal cycler according to the following protocol, but stop cycling before amplification yield plateaus (generally 8-12 cycles):

6. 95 C for 5 min
7. 95 C for 30 sec
8. 58 C for 60 sec
9. 72 C for 60 sec
10. goto 2, N more times

Prepare a Shotgun Next-Generation Sequencing Library for Analysis

Purify desired amplicon population from non-specific amplification products by gel extraction.

Concatemerize amplicons into high-molecular weight products suitable for shearing

Mechanically shear, using either a nebulizer, BioRuptor, Hydroshear,

Covaris, or similar instrument. DNA should be sheared into fragments several hundred basepairs in length.

Ligate adapters required for amplification by the sequencing platform used. If necessary, purify ligated product from unligated product and adapters.

Perform Sequencing of Library According to Manufacturer's Directions (e.g. Illumina, ABI, Etc), Reading Both the Target Sequence and the Differentiator Tag Sequence.

Analyze Data by Correcting for any Biases Detected by Quantitation of Differentiator Tag Sequence Abundance

Construct a table of target: differentiator tag abundances from the read data, e.g.:

Target ID	Differentiator Tag sequence ID	Count
1	3547	1
2	4762	1
1	9637	1
1	1078	5
3	4762	1
1	2984	1

All 'count' entries should be '1', since any particular target: differentiator tag mapping will not occur more than once by chance, and therefore will only be observed if bias was present somewhere in the sample preparation process. For any target: differentiator tag combination observed more

73

than once, all such reads are 'collapsed' into a single read before consensus basecalls are determined. This will cancel the effect of bias on consensus basecall accuracy. FIG. 5 depicts a method for making diploid genotype calls in which repeat target: differentiator tag combination are collapsed.

Example 3: Differentiator Tag Sequence Design for MIP Capture Reactions

For a set of targets, the number of differentiator tag sequences necessary to be confident (within some statistical bounds) that a certain differentiator tag sequence will not be observed more than once by chance in combination with a certain target sequence was determined. The total number of unique differentiator tag sequences for a certain differentiator tag sequence length is determined as $4^{(\text{Length in nucleotides of the differentiator tag sequence})}$. For a molecular inversion probe capture reaction that uses MIP probes having differentiator tag sequences, the probability of performing the capture reaction and capturing one or more copies of a target sequence having the same differentiator tag sequence is calculated as: $p = 1 - [N! / (N-M)!] / [N^M]$, wherein N is the total number of possible unique differentiator tag sequences and M is the number of target sequence copies in the capture reaction. Thus, by varying the differentiator tag sequence length it is possible to perform a MIP capture reaction in which the probability of capturing one or more copies of a target sequence having the same differentiator tag sequence is set at a predetermined probability value.

For example, for a differentiator tag sequence of 15 nucleotides in length, there are 1,073,741,824 possible differentiator tag sequences. A MIP capture reaction in which MIP probes, each having a differentiator tag sequence of 15 nucleotides, are combined with 10000 target sequence copies (e.g., genome equivalents), the probability of capturing one or more copies of a target sequence having the same differentiator tag sequence is 0.05. In this example, the MIP reaction will produce very few (usually 0, but occasionally 1 or more) targets where multiple copies are tagged with the same differentiator tag sequence. FIG. 6 depicts results of a simulation for 100000 capture reactions having 15 nucleotide differentiator tag sequences and 10000 target sequences.

Example 4: Assessment of the Probability for Obtaining Enough Sequencing Reads to Make Accurate Base-Calls at Multiple Independent Loci, as a Function of Sequencing Coverage

Monte Carlo simulations were performed to determine sequencing coverage requirements. The simulations assume 10000 genomic copies of a given locus (target) half mom alleles and half dad alleles. The simulations further assume 1% efficiency of capture for the MIP reaction. The simulation samples from a capture mix 100 times without replacement to create a set of 100 capture products. The simulation then samples from the set of 100 capture products with replacement (assuming unbiased amplification) to generate 'reads' from either mom or dad. The number of reads sampled depends on the coverage. The number of independent reads from both mom and dad necessary to make a high-quality base-call (assumed to be 10 or 20 reads) were then determined. The process was repeated 1000 times for each coverage level, and the fraction of times that enough reads from both parents were successfully obtained was determined. This fraction was raised to the power 1000, assuming we have 1000 independent loci that must obtain

74

successful base-calls, plotted (See FIG. 7). Result show that roughly 50x coverage is required to capture each allele $\geq 10x$ with >0.95 probability.

Example 5: MIP Capture of 'Target' Locus and 'Control' Loci

In some embodiments, to accurately quantify the efficiency of target locus capture, at least three sets of control loci are captured in parallel that have a priori been shown to serve as proxies for various lengths of target locus. For example, if the target locus is expected to have a length between 50 and 1000 bp, then sets of control loci having lengths of 50, 250, and 1000 bp could be captured (e.g. 20 loci per set should provide adequate protection from outliers), and their abundance digitally measured by sequencing. These loci should be chosen such that minimal variation in efficiency between samples and on multiple runs of the same sample is observed (and are therefore 'efficiency invariant'). These will serve as 'reference' points that define the shape of the curve of abundance-vs-length. Determining the length of the target is then simply a matter of 'reading' the length from the appropriate point on the calibration curve.

In some embodiments, the statistical confidence one has in the estimate of target length from this method is driven largely by three factors: 1) reproducibility/variation of the abundance data used to generate the calibration curve; 2) goodness of fit of the regression to the 'control' datapoints; 3) reproducibility of abundance data for the target locus being measured. Statistical bounds on 1) and 2) will be known in advance, having been measured during development of the assay. Additionally, statistical bounds on 3) will be known in general in advance, since assay development should include adequate population sampling and measure of technical reproducibility. Standard statistical methods should be used to combine these three measures into a single P value for any given experimental measure of target abundance.

In some embodiments, given the set of calibration observations, and a linear regression fit to that data, the regression can be used to predict the length value for n observations of the target locus whose length is unknown. First, choose an acceptable range for the confidence interval of the length estimate. For example, in the case of distinguishing "normal" (87-93 bp) from "premutation" (165-600 bp) potential cases of Fragile X, the goal is to measure length to sufficient precision to distinguish 93 bp from 165 bp. The predicted response value, computed when n observations is substituted into the equation for the regressed line, will have arbitrary precision. However, if for example a 95% confidence level is desired, that 95% confidence interval must be sufficiently short that it does not overlap both the "normal" and "premutation" length ranges. Continuing the example, if one calculates a length of 190 from n=400 MIP observations, and based on the regression from calibration data, the 95% confidence interval is 190 ± 20 bp, one can conclude the sample represents a "premutation" length with 95% certainty. Conversely, if the calibration data were less robust, error estimates of the regression would be higher, leading to larger confidence intervals on the predicted response value. In some embodiments, if the 95% CI were calculated as 190 ± 100 bp from n=400, one could not determine whether the predicted response value corresponds to a "normal" or "premutation" length.

In some embodiments, the confidence interval for a predicted response is calculated as:

75

The estimate for the response $\hat{y}$ is identical to the estimate for the mean of the response: $\hat{y} = b_0 + b_1 x^*$. The confidence interval for the predicted value is given by $\hat{y} \pm t^* s_{\hat{y}}$, where $\hat{y}$ is the fitted value corresponding to x^* . The value t^* is the upper $(1-C)/2$ critical value for the $t(n-2)$ distribution.

In some embodiments, a technique for analyzing a locus of interest can involve the following steps.

Convert Hybridized Probes into Covalently-Closed Circular Products Containing Subtargets

Prepare Fill-In/Ligation Reaction Mixture:

0.25 ul 2 mM dNTP mix (Invitrogen)
2.5 ul 10×Ampligase buffer (Epicentre)
5.0 ul 5 U/ul Taq Stoffel fragment (Applied Biosystems)
12.5 ul 5 U/ul Ampligase (Epicentre)
4.75 ul dH2O

Add 1.0 ul of this mix to the hybridized probe reaction, and incubate at 60 C for 10 hours.

Purify Circularized Probe/Subtarget Products from Un-Reacted Probes and 30 Genomic DNA

Prepare Exonuclease Reaction Mixture:

21 ul fill-in/ligation reaction product
2.0 ul 10×exonuclease I buffer (New England Biolabs)
2.0 ul 20 U/ul exonuclease I (New England Biolabs)
2.0 ul 100 U/ul exonuclease III (New England Biolabs)

Incubate at 37 C for 60 min, then heat-inactivate by incubating at 80 C for 15 min. Immediately cool to 4 C for storage.

Amplify Circular Material by PCR Using Primers Specific to the 'Constant Region' of the Probes

Prepare PCR Mixture:

5.0 ul 10×Accuprime reaction buffer (Invitrogen)
1.5 ul 10 uM CP-2-FA-Ilmn (platform-specific amplification sequence plus 'circle constant region'-specific sequence)
1.5 ul 10 uM CP-2-RA-Ilmn (platform-specific amplification sequence plus 'circle constant region'-specific sequence)
0.4 ul 25 mM dNTP mix (Invitrogen)
2.0 ul heat-inactivated exonuclease reaction mix
1.5 ul 10×SybrGreen (Invitrogen)
0.4 ul 2.5 U/ul Accuprime Pfx polymerase (Invitrogen)
37.7 ul dH2O

Thermal cycle in real-time thermal cycler according to the following protocol, but stop cycling before amplification yield plateaus (generally 8-12 cycles):

11. 95 C for 5 min
12. 95 C for 30 sec
13. 58 C for 60 sec
14. 72 C for 60 sec
15. goto 2, N more times

Perform Sequencing (e.g., Next-Generation Sequencing) on Sample for Digital Quantitation According to Manufacturer's Instructions (e.g., Illumina, ABI)

Example 6: MIP-capture reaction of a set of exon target nucleic acids

MIP probes are synthesized using standard column-based oligonucleotide synthesis by any number of vendors (e.g. IDT).

Hybridize Capture Probes to Human Genomic Sample Assemble Hybridization Reaction:

1.0 ul capture probe mix (~2.5 pmol)
2.0 ul 10×Ampligase buffer (Epicentre)
6.0 ul 500 ng/ul human genomic DNA (~16.7 fmol)
11 ul dH2O

76

In a thermal cycler, heat reaction to 95 C for 5 min to denature genomic DNA, then cool to 60 C. Allow to incubate at 60 C for 40 hours.

Convert Hybridized Probes into Covalently-Closed Circular Products Containing Target Nucleic Acids

Prepare Fill-In/Ligation Reaction Mixture:

0.25 ul 2 mM dNTP mix (Invitrogen)
2.5 ul 10×Ampligase buffer (Epicentre)
5.0 ul 5 U/ul Taq Stoffel fragment (Applied Biosystems)
12.5 ul 5 U/ul Ampligase (Epicentre)
4.75 ul dH2O

Add 1.0 ul of this mix to the hybridized probe reaction, and incubate at 60 C for 10 hours.

Purify, Circularized Probe/Target Nucleic Acid Products from Un-Reacted Probes and Genomic DNA

Prepare Exonuclease Reaction Mixture:

21 ul fill-in/ligation reaction product
2.0 ul 10×exonuclease I buffer (New England Biolabs)
2.0 ul 20 U/ul exonuclease I (New England Biolabs)
2.0 ul 100 U/ul exonuclease III (New England Biolabs)

Incubate at 37 C for 60 min, then heat-inactivate by incubating at 80 C for 15 min. Immediately cool to 4 C for storage.

Amplify Circular Material by PCR Using Primers Specific to the 'Constant Region' Of the Probes

Prepare PCR Mixture:

5.0 ul 10×Accuprime reaction buffer (Invitrogen)
1.5 ul 10 uM CP-2-FA (5'-GCAC-GATCCGACGGTAGTGT-3') (SEQ ID NO: 183)
1.5 ul 10 uM CP-2-RA (5'-CCGTAATCGGGAAGCT-GAAG-3') (SEQ ID NO: 184)
0.4 ul 25 mM dNTP mix (Invitrogen)
2.0 ul heat-inactivated exonuclease reaction mix
1.5 ul 10×SybrGreen (Invitrogen)
0.4 ul 2.5 U/ul Accuprime Pfx polymerase (Invitrogen)
37.7 ul dH2O

Thermal cycle in real-time thermal cycler according to the following protocol, but stop cycling before amplification yield plateaus (generally 8-12 cycles):

16. 95 C for 5 min
17. 95 C for 30 sec
18. 58 C for 60 sec
19. 72 C for 60 sec
20. goto 2, N more times

Prepare a Shotgun Next-Generation Sequencing Library for Analysis

Purify desired amplicon population from non-specific amplification products by gel extraction.

Concatemerize amplicons into high-molecular weight products suitable for shearing

Mechanically shear, using either a nebulizer, BioRuptor, Hydroshear, Covaris, or similar instrument. DNA should be sheared into fragments several hundred base-pairs in length.

Ligate adapters required for amplification by the sequencing platform used. If necessary, purify ligated product from unligated product and adapters.

Perform Sequencing of Library According to Manufacturer's Directions (e.g. Illumina, ABI, Etc), Reading the Target Sequence to Determine Abundance of the Target Nucleic Acid.

Example 7: Use of MIPs, Hybridization, and Mutation-Detection MIPs to Genotype a Set of 1000 Targets

MIPs, hybridization, and mutation-detection MIPs are used to genotype a set of 1000 targets. The protocol permits detection of any of 50 specific known point mutations

First, separate MIP, hybridization, and mutation-detection MIP reactions are performed on a biological sample. A MIP capture reaction is performed essentially as described in Turner et al 2009, Nature methods 6:315-6. A set of MIPs is designed such that each probe in the set flanks one of the 1000 targets. Separately, a hybridization enrichment reaction is performed using the Agilent SureSelect procedure. Prior to selection, the genomic DNA to be enriched is converted into a shotgun sequencing library using Illumina's 'Fragment Library' kit and protocol. Agilent's web interface is used to design a set of probes which will hybridize to the target nucleic acids. Separately, a set of probes are designed (mutation-detection MIPs) which will form MIPs only if mutations (e.g., specific polymorphisms) are present. Each mutation-detection MIP has a 3'-most base identity that is specific for a single known mutation. A reaction with this set of mutation-detection MIPs is performed to selectively detect the presence of any mutant alleles.

Once all three reactions have been performed, the two MIP reactions are combined (e.g., at potentially non-equimolar ratios to further increase sensitivity of mutation detection) into a single tube, and run as one sample on the next-generation DNA sequencing instrument. The hybridization-enriched reaction is run as a separate sample on the

next-generation DNA sequencing instrument. Reads from each 'sample' are combined by a software algorithm which forms a consensus diploid genotype at each position in the target set by evaluating the total coverage at each position, the origin of each read in that total coverage, the quality score of each individual read, and the presence (or absence) of any reads derived from mutation-specific MIPs overlapping the region.

It should be appreciated that the preceding examples are non-limiting and aspects of the invention may be implemented as described herein using alternative techniques and/or protocols that are available to one of ordinary skill in the art.

It will be clear that the methods may be practiced other than as particularly described in the foregoing description and examples. Numerous modifications and variations of the present disclosure are possible in light of the above teachings and, therefore, are within the scope of the claims. Preferred features of each aspect of the Disclosure are as for each of the other aspects mutatis mutandis. The documents including patents, patent applications, journal articles, or other disclosures mentioned herein are hereby incorporated by reference in their entirety. In the event of conflict, the disclosure of present application controls, other than in the event of clear error.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 184

<210> SEQ ID NO 1

<211> LENGTH: 52

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 1

tcacatcacc aagttaaaaa aaaaaarggg gcgggggggc agaataaaaa tt 52

<210> SEQ ID NO 2

<211> LENGTH: 52

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (27) .. (27)

<223> OTHER INFORMATION: May or may not be present

<400> SEQUENCE: 2

aaacaaggat gaattaagtt tttttttaa aaagaaacat ttggttaagg ga 52

<210> SEQ ID NO 3

<211> LENGTH: 52

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 3

atcaccaagt taaaaaaaa aaaggggrcgg gggggcagaa tgaaaattgc at 52

<210> SEQ ID NO 4

<211> LENGTH: 52

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 4

ctattgaacc agaaccaaac aggaatrcca tagcattttg taaactaaac tg 52

-continued

<210> SEQ ID NO 5
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 5

caggagttca agaccagcct actaaamcac acacacacac acacacacac ac 52

<210> SEQ ID NO 6
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 6

gattaaataa tagtgtttat gtacccsgct tataggagaa gaggggtgtgt gt 52

<210> SEQ ID NO 7
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 7

attgttatct tttcatataa ggtaacwgag gccagagag attaaataac at 52

<210> SEQ ID NO 8
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 8

taattttaat taagtaaatt taattgrtag ataaataagt agataaaaaa ta 52

<210> SEQ ID NO 9
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 9

gtataaaaaa aaaaaaaaaa aaagttwgaa tgttttcttg cattcagagc ct 52

<210> SEQ ID NO 10
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 10

atactaaaaa tttaaagttc tcttgcrata tattttctta atatettaca tc 52

<210> SEQ ID NO 11
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 11

tgctgggatt acaggcgtga gccaccrcgc ctggcctgat ggacatatt tt 52

<210> SEQ ID NO 12
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 12

ctacaatata agtatagtat tgcaamcca tcaggaaggg tgtaactat tt 52

-continued

<210> SEQ ID NO 13
 <211> LENGTH: 54
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (27)..(29)
 <223> OTHER INFORMATION: This region may or may not be present in its entirety

 <400> SEQUENCE: 13

 gttgtctcca aacttttttt caggtgagaa ggtggccaac cgagcttcgg aaag 54

<210> SEQ ID NO 14
 <211> LENGTH: 52
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

 <400> SEQUENCE: 14

 ttttttcata aaagattata taaaggmtat tgcttttgaa tcacaaacac ta 52

<210> SEQ ID NO 15
 <211> LENGTH: 52
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

 <400> SEQUENCE: 15

 atctagtgag cagtcaggaa agagaayttc cagatcctgg aaatcagggt ta 52

<210> SEQ ID NO 16
 <211> LENGTH: 52
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (27)..(27)
 <223> OTHER INFORMATION: May or may not be present

 <400> SEQUENCE: 16

 tagagtataa aaaaaaaaaa aaaaaaagtt tgaatgtttt cttgcattca ga 52

<210> SEQ ID NO 17
 <211> LENGTH: 52
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

 <400> SEQUENCE: 17

 tgcaaatgcc aactatcaaa gatattsgag tatactgtca ataaacttca ta 52

<210> SEQ ID NO 18
 <211> LENGTH: 52
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

 <400> SEQUENCE: 18

 tcctcaacag ttagaaacaa tattttsagt gatttcccat gccaaactta ct 52

<210> SEQ ID NO 19
 <211> LENGTH: 52
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

 <400> SEQUENCE: 19

 tttttggtat tgtgttataa taagtgrgaa ttcaatacag tataatgtct gt 52

-continued

```

<210> SEQ ID NO 20
<211> LENGTH: 87
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (27)..(72)
<223> OTHER INFORMATION: This region may or may not be present in its
        entirety

<400> SEQUENCE: 20

cttgaaatcg gatatatata tatatatgta tatatatata tatatatata tatatatata      60
tagtattatc cctgttttca cagttttt                                         87

<210> SEQ ID NO 21
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 21

agaggggctg tgaaggacac caaggargag actaagagcc aggagggaaa ac           52

<210> SEQ ID NO 22
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 22

tagagtttat tagctttttac tactctrcct agttactttg tggtacagaa ta          52

<210> SEQ ID NO 23
<211> LENGTH: 53
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (27)..(28)
<223> OTHER INFORMATION: This region may or may not be present in its
        entirety

<400> SEQUENCE: 23

actagtgatg atgagcttct tttcatatgt ttgttggtg cataaatgtc ttc           53

<210> SEQ ID NO 24
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 24

gcagagaaaa gaggggctgt gaaggasacc aaggaggaga ctaagagcca gg           52

<210> SEQ ID NO 25
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 25

ttcagaggtc taccactggt gcatackcta atcacagtgt cgaaaatttt ac           52

<210> SEQ ID NO 26
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 26

```

-continued

aagaaagagc aaaagagggc aaacttytca tacatttttg atgtcgaaac ca 52

<210> SEQ ID NO 27
 <211> LENGTH: 52
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (27)..(27)
 <223> OTHER INFORMATION: May or may not be present

<400> SEQUENCE: 27

cctaaagtgtt aaaaagaaaa aaaaaaagga agaaggaatt aaaaatccaa ag 52

<210> SEQ ID NO 28
 <211> LENGTH: 52
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 28

gtgtgtgtgt gtatatatat atatatttat atattttttt tttcctgagc ca 52

<210> SEQ ID NO 29
 <211> LENGTH: 52
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 29

aaactgttga tgttttcatt tatttasatc attggaaaac ttagattct ag 52

<210> SEQ ID NO 30
 <211> LENGTH: 52
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 30

tttatccatt cttaaccaga acagacrttt tttcagagct ggtccaggaa aa 52

<210> SEQ ID NO 31
 <211> LENGTH: 52
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 31

ttgaaatcgg atatatatat atatatrta atatatatat atatatatat at 52

<210> SEQ ID NO 32
 <211> LENGTH: 52
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (27)..(27)
 <223> OTHER INFORMATION: May or may not be present

<400> SEQUENCE: 32

tagtttttta tttcctcata ttattttcag tggcttttcc ttccacatct tt 52

<210> SEQ ID NO 33
 <211> LENGTH: 52
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 33

-continued

 acatagttct cagtgtgtaca actacargtg atttctcttt tcttatttct gg 52

<210> SEQ ID NO 34
 <211> LENGTH: 52
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 34

agagcaatgg catcccttgt cttgtgytat acaggatgca gcaatttata gg 52

<210> SEQ ID NO 35
 <211> LENGTH: 52
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 35

ttctgtctac ataagatgtc atactaratt atcttttcca gcatgcattc ag 52

<210> SEQ ID NO 36
 <211> LENGTH: 52
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 36

cagggtggca tgtaggcag tgcttaraat aaatgagttg gttatacaag ta 52

<210> SEQ ID NO 37
 <211> LENGTH: 53
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (27)..(28)
 <223> OTHER INFORMATION: This region may or may not be present in its entirety

<400> SEQUENCE: 37

aggacacaca cacacacaca cacacacatg cacacacatt taaatagatg cat 53

<210> SEQ ID NO 38
 <211> LENGTH: 52
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 38

taaaaaattg gtataatgaa attgcayttg tagtctttgg acatttaaatt cc 52

<210> SEQ ID NO 39
 <211> LENGTH: 52
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 39

tttaataact taagaggtac gcagagraaa gaggggctgt gaaggacacc aa 52

<210> SEQ ID NO 40
 <211> LENGTH: 52
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 40

cagcaatgaa tattttgagg ctgaggygct gaggggtaaa attgcagcct gg 52

<210> SEQ ID NO 41

-continued

<211> LENGTH: 55
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (27)..(30)
<223> OTHER INFORMATION: This region may or may not be present in its entirety

<400> SEQUENCE: 41

ttatgggttta tatttttgtg tcttctcttt aacacatctt ttctagcaga attca 55

<210> SEQ ID NO 42
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (27)..(27)
<223> OTHER INFORMATION: May or may not be present

<400> SEQUENCE: 42

gtatttttagt tttttttttt tgtttgtttt gttttgtttt gttttgtttt tg 52

<210> SEQ ID NO 43
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 43

tgggtgactc catttttact tttagtytgg tctgttgagg cctcgtgaga ga 52

<210> SEQ ID NO 44
<211> LENGTH: 55
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (27)..(30)
<223> OTHER INFORMATION: This region may or may not be present in its entirety

<400> SEQUENCE: 44

tattttcatg tatttttagtt tttttttttt gtttgttttg ttttgttttg ttttg 55

<210> SEQ ID NO 45
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 45

gtgtgtgtgt atatatatat atatatwtat attttttttt tcctgagcca aa 52

<210> SEQ ID NO 46
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (27)..(27)
<223> OTHER INFORMATION: May or may not be present

<400> SEQUENCE: 46

aacctattag catgtctggc agaaaaatag atacttaata aatttcttaa at 52

<210> SEQ ID NO 47
<211> LENGTH: 52

-continued

<212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

 <400> SEQUENCE: 47

 gaggccttaga cagtttaagt aactcargca tggttacaca actagctagg gc 52

 <210> SEQ ID NO 48
 <211> LENGTH: 52
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

 <400> SEQUENCE: 48

 gtgtgagtat tatgagacca tatgttrgga gattttattt ggtattgagg at 52

 <210> SEQ ID NO 49
 <211> LENGTH: 52
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

 <400> SEQUENCE: 49

 gaaacccac cccttetata gttttcycct taatatattac aatggaacca tt 52

 <210> SEQ ID NO 50
 <211> LENGTH: 52
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

 <400> SEQUENCE: 50

 catatatata tagtgtgtgt gtgtgtrtat atatatatat atatatttt tt 52

 <210> SEQ ID NO 51
 <211> LENGTH: 53
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

 <400> SEQUENCE: 51

 cgagcgagag agagagagag agagagagga gccggcgcca gaactacgca tgc 53

 <210> SEQ ID NO 52
 <211> LENGTH: 52
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

 <400> SEQUENCE: 52

 ggcaggtaat atctagtacc ttacttwtat ttcctgagca cattctacat tt 52

 <210> SEQ ID NO 53
 <211> LENGTH: 52
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

 <400> SEQUENCE: 53

 ggccaggaat gggagtgcct aggtgcragag gtggcactgt tcccgcagct gc 52

 <210> SEQ ID NO 54
 <211> LENGTH: 52
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

 <400> SEQUENCE: 54

 tgaaaactcc atcccctcag ggatcatyaga tgaagagaag accacagggg tt 52

 <210> SEQ ID NO 55

-continued

```

<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 55

tgtaggtaag ggtcacatgt gggagasgca gctgtgggta ggtcagccct gt          52

<210> SEQ ID NO 56
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (27)..(27)
<223> OTHER INFORMATION: May or may not be present

<400> SEQUENCE: 56

ccaagaaggc gccattacac tccagcctgg gcgacagggc gagactccct ca          52

<210> SEQ ID NO 57
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (27)..(27)
<223> OTHER INFORMATION: May or may not be present

<400> SEQUENCE: 57

tgccacaccc agctaatttg tgtgtgggtat gtgtgtgtat gtatgtgtgt gt          52

<210> SEQ ID NO 58
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (27)..(27)
<223> OTHER INFORMATION: May or may not be present

<400> SEQUENCE: 58

gttcctccag taattttttt ttttttttgt tttgagacag agtcttgccc tg          52

<210> SEQ ID NO 59
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (27)..(27)
<223> OTHER INFORMATION: May or may not be present

<400> SEQUENCE: 59

atcatgccca gataattttt tttttttgta ttttagtaga cacagggttt ca          52

<210> SEQ ID NO 60
<211> LENGTH: 53
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (27)..(28)
<223> OTHER INFORMATION: This region may or may not be present in its
entirety

<400> SEQUENCE: 60

cgagcgagag agagagagag agagagagga gccggcgcgga gaactacgca tgc          53

```

-continued

```

<210> SEQ ID NO 61
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (27)..(27)
<223> OTHER INFORMATION: May or may not be present

```

```

<400> SEQUENCE: 61

cctgtgaggg gcacattcct tagtagctaa ggagttgggg gtgtgaagat cc          52

```

```

<210> SEQ ID NO 62
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (27)..(27)
<223> OTHER INFORMATION: May or may not be present

```

```

<400> SEQUENCE: 62

acaggctact ggctggggccc aggcaaaggg ggccttgcca ggaaaagtcc ct          52

```

```

<210> SEQ ID NO 63
<211> LENGTH: 53
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (27)..(28)
<223> OTHER INFORMATION: This region may or may not be present in its
entirety

```

```

<400> SEQUENCE: 63

gcgagagaga gagagagaga gagaggagag cggcgcgag aactacgcat gcg          53

```

```

<210> SEQ ID NO 64
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<400> SEQUENCE: 64

aagaagaaaa ataaaaagaa agtgggycag accgagagaa caggaagcct ga          52

```

```

<210> SEQ ID NO 65
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<400> SEQUENCE: 65

aaggacaaag gcaaagagac aaaggckcaa cactgggggt cccagagag tg          52

```

```

<210> SEQ ID NO 66
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<400> SEQUENCE: 66

tacctagtca cttcctgect ccatggygca aaaggggatg ggtgtgcctc tt          52

```

```

<210> SEQ ID NO 67
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

```

-continued

<400> SEQUENCE: 67

ctcttcgag gttccacct gaacacyttc ctgctccctc gtggtgtaga gt 52

<210> SEQ ID NO 68

<211> LENGTH: 52

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 68

ttctgactgg caaccagccc cactctygtg gagccctcag gaatgaactt gc 52

<210> SEQ ID NO 69

<211> LENGTH: 52

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 69

gtcagcctc ccaggetgga gtgcagwggc gcatctcgg ctcaccgcaa cc 52

<210> SEQ ID NO 70

<211> LENGTH: 52

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 70

catgaaccac atcaaatgag atttagyggg agtggcacac acagtcatga cc 52

<210> SEQ ID NO 71

<211> LENGTH: 52

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 71

aagcagccct ggggagtcgg ggcgggrcct ggattggaaa agagacggtc ac 52

<210> SEQ ID NO 72

<211> LENGTH: 52

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 72

ctccaagttc tgggagcaga gtgtgcrctt aggtcctcgg gatcgaggga tg 52

<210> SEQ ID NO 73

<211> LENGTH: 52

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (27)..(27)

<223> OTHER INFORMATION: May or may not be present

<400> SEQUENCE: 73

gttcctccag taattttttt tttttktgtt ttgagacaga gtcttgccct gt 52

<210> SEQ ID NO 74

<211> LENGTH: 52

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 74

ctagtacett acttccctca agttcawtca tctcacagat atttcttgag ca 52

-continued

```

<210> SEQ ID NO 75
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 75

aattagccgt gcgtggtggc gggtgcytgt aatccacgt acttgggagg ct      52


<210> SEQ ID NO 76
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 76

ccatggccag cgggggaggg gacgggmaca cacagaccca cacagagact ca      52


<210> SEQ ID NO 77
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (77)..(77)
<223> OTHER INFORMATION: May or may not be present

<400> SEQUENCE: 77

agcgagagag agagagagag agagagrgag cggcgcgag aactacgcat gc      52


<210> SEQ ID NO 78
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 78

cgtagcagtt agcagatgat agcggbgaa atcttatttc acagggcatt aa      52


<210> SEQ ID NO 79
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 79

ctggccctgg tgacagtggg gctgtgygtg gggccagagc cttctcagag gt      52


<210> SEQ ID NO 80
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 80

cagatactgg ccctggtgac agtgggrctg tgcgtggggc cagagccttc tc      52


<210> SEQ ID NO 81
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 81

gacagatact ggccctggtg acagtgkggc tgtgcgtggg gccagagcct tc      52


<210> SEQ ID NO 82
<211> LENGTH: 53
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:

```

-continued

```

<221> NAME/KEY: misc_feature
<222> LOCATION: (27)..(28)
<223> OTHER INFORMATION: This region may or may not be present in its
        entirety

<400> SEQUENCE: 82

ggctctctctc tctctctctc tctctctctc tegetctctc gtctctctc tet          53

<210> SEQ ID NO 83
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 83

gtttcagacc agcctggcca acatagygaa accccatctc tactaaaaat aa          52

<210> SEQ ID NO 84
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 84

agtgggcgat tggatggagc tgagtayggg gcccatccag gctaatacaca cc          52

<210> SEQ ID NO 85
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 85

ccgggctccg tgaatgtttg tcacatstct gaagaacgta tgaattacat aa          52

<210> SEQ ID NO 86
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 86

gaatcccaac cccgacgctc gtcgcscggc tccgtgaatg tttgtcacat gt          52

<210> SEQ ID NO 87
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 87

gcgaatccca accccgacgc tcgtcgycgg gtcctgtgaa tgtttgtcac at          52

<210> SEQ ID NO 88
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 88

tgggcagaag tcagggtcca aagaaakggc aaagaaaagt gtcagtggct ca          52

<210> SEQ ID NO 89
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 89

taagaaagac gtttttgatt ttttctrgac ttctctggct ccactaccct gc          52

```

-continued

<210> SEQ ID NO 90
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 90
ctgattcctg gccaggagcg gtggctyacg cctgtaatcc cagcgctttg gg 52

<210> SEQ ID NO 91
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 91
taaaaatgct gattcctggc caggagyggg ggctcacgcc tgtaatccca gc 52

<210> SEQ ID NO 92
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 92
tttaaaaatg ctgattcctg gccaggmgcg gtggctcagc cctgtaatcc ca 52

<210> SEQ ID NO 93
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 93
caagacctgt caaagatctg agaaatwta cccgacttac aagctaacca tt 52

<210> SEQ ID NO 94
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 94
actgtaataa gtgctgtaaa agaaatrcac aaaataatat agcagagggt at 52

<210> SEQ ID NO 95
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 95
cttgaggtea ggagttcaag accagtytgg gcaacatggg gaaaaccttg tc 52

<210> SEQ ID NO 96
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 96
aggttgacgt gagccgagat catgcrrttg cactccagcc ggggcaacaa aa 52

<210> SEQ ID NO 97
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 97
acaagtgtct tgaaattatc tgtgatytc tatagagcaa tacttttgta aa 52

-continued

<210> SEQ ID NO 98
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 98

gtgggtatat gcagctctat gcactayctg ctcatttatt tggtaaactt aa 52

<210> SEQ ID NO 99
<211> LENGTH: 59
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (27)..(34)
<223> OTHER INFORMATION: This region may or may not be present in its entirety

<400> SEQUENCE: 99

tgtgtgtgtg tgcgtgtgtg tgtgtgtgtg tgtgatcata agagtggctg cagcaaaact 59

<210> SEQ ID NO 100
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 100

agtctggagt gcaatgggtgc aatctcrgct cactgcagcc tccacctccg gg 52

<210> SEQ ID NO 101
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 101

ctcctaattg atatttccta aattttkctg aacagaattt aacttgagct gg 52

<210> SEQ ID NO 102
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 102

atttaaaaat ggatttctag aaaaacratc acatacttga atatttttag aa 52

<210> SEQ ID NO 103
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 103

ctataaatgg gtagcatgag ggattcragg aggtggctga aagaagcacg ta 52

<210> SEQ ID NO 104
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 104

aagaaaccaa gcatagtaga gtgttaraaa accaaagcaa ctaaacaact gt 52

<210> SEQ ID NO 105
<211> LENGTH: 52

-continued

<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 105

cggggctcag aacttgtaac agaaaawtaa aatatactcc actcaaggga at 52

<210> SEQ ID NO 106
<211> LENGTH: 57
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (27)..(32)
<223> OTHER INFORMATION: This region may or may not be present in its
entirety

<400> SEQUENCE: 106

tactacactt cacggatact gtacttgtag tttttttcca aattgaaggt ttttggc 57

<210> SEQ ID NO 107
<211> LENGTH: 63
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (27)..(38)
<223> OTHER INFORMATION: This region may or may not be present in its
entirety

<400> SEQUENCE: 107

ttgtttttgt tttgtttttt gtttttgttt ttgttttttg agatggagtc tcgctctgtc 60
gcc 63

<210> SEQ ID NO 108
<211> LENGTH: 57
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (27)..(32)
<223> OTHER INFORMATION: This region may or may not be present in its
entirety

<400> SEQUENCE: 108

tttgccctac tacacttcac ggatactgta cttgtacttt ttttccaaat tgaaggt 57

<210> SEQ ID NO 109
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (27)..(27)
<223> OTHER INFORMATION: May or may not be present

<400> SEQUENCE: 109

gagggtggctg aaagaagcac gtatccctga tggcatggtt gcgggttata tg 52

<210> SEQ ID NO 110
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (27)..(27)
<223> OTHER INFORMATION: May or may not be present

<400> SEQUENCE: 110

-continued

gagaaaagca gttctctggaa caccgccacc ccttaacccc ttatctctgc tt 52

<210> SEQ ID NO 111
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (27)..(27)
<223> OTHER INFORMATION: May or may not be present

<400> SEQUENCE: 111

ttacatatgt atacatgtgc catggttggt gtgccgcacc cattaactcg tc 52

<210> SEQ ID NO 112
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (27)..(27)
<223> OTHER INFORMATION: May or may not be present

<400> SEQUENCE: 112

ctttttccag attttttttt tttttttgag acagagtttc actcttggtg cc 52

<210> SEQ ID NO 113
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (27)..(27)
<223> OTHER INFORMATION: May or may not be present

<400> SEQUENCE: 113

tttggaaatc ttaagctttt atttgggtgt cacagagaaa caggatctgt at 52

<210> SEQ ID NO 114
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (27)..(27)
<223> OTHER INFORMATION: May or may not be present

<400> SEQUENCE: 114

tactttaagt tttagggtac atgtgcacca tgtgcaggtt tgttacatat gt 52

<210> SEQ ID NO 115
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (27)..(27)
<223> OTHER INFORMATION: May or may not be present

<400> SEQUENCE: 115

attcatgacc agccacataa atgcacagta ttacttcgca agcatgccaa tg 52

<210> SEQ ID NO 116
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:

-continued

<221> NAME/KEY: misc_feature
<222> LOCATION: (27)..(27)
<223> OTHER INFORMATION: May or may not be present

<400> SEQUENCE: 116

gtgcactaga attagctaaa gtgggggaaa aaaagatgca ttgatgggc ta 52

<210> SEQ ID NO 117
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (27)..(27)
<223> OTHER INFORMATION: May or may not be present

<400> SEQUENCE: 117

aacctccacc tcccaggttc aagagaattc tctgcctca gcctccaag ta 52

<210> SEQ ID NO 118
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 118

cctccctgtg atccgaagta gcagacrtac ttaacttcca tgggtgattg tt 52

<210> SEQ ID NO 119
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (27)..(27)
<223> OTHER INFORMATION: May or may not be present

<400> SEQUENCE: 119

aaaacattat tatactaga aaaaaaatgt atcttaacca ttgtgggaag tg 52

<210> SEQ ID NO 120
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (27)..(27)
<223> OTHER INFORMATION: May or may not be present

<400> SEQUENCE: 120

ttgaaggtaa aatcataggg agttgggagc tgcctcttg cgctgaatca gt 52

<210> SEQ ID NO 121
<211> LENGTH: 53
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (27)..(28)
<223> OTHER INFORMATION: This region may or may not be present in its entirety

<400> SEQUENCE: 121

acttgaggcc tttttggaga ggtagcaac tctgaaaact ctgtccctgg acc 53

<210> SEQ ID NO 122
<211> LENGTH: 52
<212> TYPE: DNA

-continued

```

<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (27)..(27)
<223> OTHER INFORMATION: May or may not be present

<400> SEQUENCE: 122
gaaggagaaa aagagaggaa ataagttaaa ataataaaca caattaataa ag          52

<210> SEQ ID NO 123
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 123
tgtatacatg tgccatgttg gtgtgcygca cccattaact cgtcatttag ca          52

<210> SEQ ID NO 124
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (27)..(27)
<223> OTHER INFORMATION: May or may not be present

<400> SEQUENCE: 124
tcacctgtca cctcctatag aactttcccc tgaccctcct ctatagcatt aa          52

<210> SEQ ID NO 125
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (27)..(27)
<223> OTHER INFORMATION: May or may not be present

<400> SEQUENCE: 125
ataaatgata atcattcaca gtagggggtt tgttttgttt tttttctgga aa          52

<210> SEQ ID NO 126
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (27)..(27)
<223> OTHER INFORMATION: May or may not be present

<400> SEQUENCE: 126
acagacatat ctacaaacac actttttcac atatttgtgt aagtcattta tg          52

<210> SEQ ID NO 127
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 127
aaagacaact aaactaacgc tcaatgmaaa aagtattcgc tgctgtttac at          52

<210> SEQ ID NO 128
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 128

```

-continued

taccgtgtac cccgtgtttg tgaatgmggc cgcattattac gaaaagaaag aa 52

<210> SEQ ID NO 129
 <211> LENGTH: 52
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 129

cagggctgga ggtaaaacca tttatttcta accccagagc agtgaagaag tg 52

<210> SEQ ID NO 130
 <211> LENGTH: 52
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 130

ttcttcattg cctattgaag agagagyggga atgctttggt tgccagatat gg 52

<210> SEQ ID NO 131
 <211> LENGTH: 52
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 131

cacaagatct cattaactcag gagctgycca agtgtctaata gtacttagtt aa 52

<210> SEQ ID NO 132
 <211> LENGTH: 52
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 132

ttctgtgtaa catttcattt aagcaargga ttcggcaaat caaaaattgt ca 52

<210> SEQ ID NO 133
 <211> LENGTH: 52
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 133

taaaacgtat tgaaggtatt attgackctg ttgaagcaaa gagaacaaaa ca 52

<210> SEQ ID NO 134
 <211> LENGTH: 52
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 134

atctgctctt ccagttggat gacaagyctt gctgtctaac acctgctgca ga 52

<210> SEQ ID NO 135
 <211> LENGTH: 52
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 135

ccattttttg ttgtattttt tttttcytga atacttttta tcgcagttgg tt 52

<210> SEQ ID NO 136
 <211> LENGTH: 52
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

-continued

<400> SEQUENCE: 136

ccctgtctct aaaagaaaa aaaaaaraaa aaaaaaagaa aacaaaaccc aa 52

<210> SEQ ID NO 137

<211> LENGTH: 52

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 137

agtggctcca aaaaggtcat ggaaccyctt gaggatgatg caaattgact ct 52

<210> SEQ ID NO 138

<211> LENGTH: 52

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 138

taaagttact tttcttttat tgactttccc ttatttttta acettatgct tt 52

<210> SEQ ID NO 139

<211> LENGTH: 52

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (27)..(27)

<223> OTHER INFORMATION: May or may not be present

<400> SEQUENCE: 139

cagagttaaa aaaaaaaaa aaaaaaagga agtagcagca acagcttgga aa 52

<210> SEQ ID NO 140

<211> LENGTH: 52

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 140

gttgcccagg gttgagtga gaggcayac ttggctcaca gcaacctctg cc 52

<210> SEQ ID NO 141

<211> LENGTH: 55

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (27)..(30)

<223> OTHER INFORMATION: This region may or may not be present in its entirety

<400> SEQUENCE: 141

aaggettttt ttttttttt ttttttttt gagacagagt cttgctgtgt ccccc 55

<210> SEQ ID NO 142

<211> LENGTH: 52

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 142

gcctacattc tgcaaagagg agggaasatt cacagctcca tacttgaacc ct 52

<210> SEQ ID NO 143

<211> LENGTH: 52

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

-continued

<400> SEQUENCE: 143

ccaaaggaga atagctctag gggaggsagg tggatgagta tgcattggggg ag 52

<210> SEQ ID NO 144

<211> LENGTH: 52

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (27)..(27)

<223> OTHER INFORMATION: May or may not be present

<400> SEQUENCE: 144

gactccatct caaaaaaaaa aaaaaaatgc agtctaattgg cagaattaga ct 52

<210> SEQ ID NO 145

<211> LENGTH: 52

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 145

ttattttattt attttattat ttttgaraca gggctctctgt tgtccaggct gg 52

<210> SEQ ID NO 146

<211> LENGTH: 52

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 146

ttttgaggca gggctctcact ctgttgycca gggttgagtg cagaggcaca tc 52

<210> SEQ ID NO 147

<211> LENGTH: 52

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 147

cgcttgcga agggcccccaca gcttgcytgt gacaaacggt cataggcaaa tg 52

<210> SEQ ID NO 148

<211> LENGTH: 52

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 148

ggaggtctgt acaaagcacc acctacytca tgggtcagtt tccacagcag aa 52

<210> SEQ ID NO 149

<211> LENGTH: 55

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (27)..(30)

<223> OTHER INFORMATION: This region may or may not be present in its entirety

<400> SEQUENCE: 149

gaatcttata attcactgtg tacctccctc tgtttcatat tttcgcaatt gaact 55

<210> SEQ ID NO 150

<211> LENGTH: 52

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

-continued

<400> SEQUENCE: 150

aacatagtat ctaatatagc ttacaycca aagccaaaat atgaatacac tg	52
--	----

<210> SEQ ID NO 151

<211> LENGTH: 52

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (27)..(27)

<223> OTHER INFORMATION: May or may not be present

<400> SEQUENCE: 151

ttgttttgtt ttgtttgggg gggggggttg tttttctgag agggagtctt gc	52
---	----

<210> SEQ ID NO 152

<211> LENGTH: 52

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 152

cataccaaag ggcagctgga gggatayaga cggaagtcac gtggagagtg aa	52
---	----

<210> SEQ ID NO 153

<211> LENGTH: 52

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 153

caggtgtgag ccaccacgac caccaawtta gctcttttta ctccttcctt tc	52
---	----

<210> SEQ ID NO 154

<211> LENGTH: 52

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 154

agtggtagct gattttgctt ctggatmctt tgccaccttc ccaactcttta at	52
--	----

<210> SEQ ID NO 155

<211> LENGTH: 64

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (27)..(39)

<223> OTHER INFORMATION: This region may or may not be present in its entirety

<400> SEQUENCE: 155

aaagacctgt ttcttaaaaa aaaaaaagaa aaaaaaaaag aaagaaaaga aaagaaaaaa	60
---	----

acag	64
------	----

<210> SEQ ID NO 156

<211> LENGTH: 52

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 156

taaaaaatct ttcaatgagg agatgtyccc agagcaagac agctgtagga tg	52
---	----

<210> SEQ ID NO 157

<211> LENGTH: 52

<212> TYPE: DNA

-continued

```

<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (27)..(27)
<223> OTHER INFORMATION: May or may not be present

<400> SEQUENCE: 157

aaaagaaaaa aaaaaaaaaa aaaaaaagaa aacaaaaccc aaacccataa ag          52

<210> SEQ ID NO 158
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 158

cctgtctcta aaagaaaaaa aaaaaaaraaa aaaaaagaaa acaaaaccca aa          52

<210> SEQ ID NO 159
<211> LENGTH: 53
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (27)..(28)
<223> OTHER INFORMATION: This region may or may not be present in its
entirety

<400> SEQUENCE: 159

gttatcatag aaaaatatca cactctgtct gtatccccac ttccagaaac tgt          53

<210> SEQ ID NO 160
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (27)..(27)
<223> OTHER INFORMATION: May or may not be present

<400> SEQUENCE: 160

caggagctca tagaattaca tacaatcttt tttttttttt ttgagacag cg          52

<210> SEQ ID NO 161
<211> LENGTH: 55
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (27)..(30)
<223> OTHER INFORMATION: This region may or may not be present in its
entirety

<400> SEQUENCE: 161

ttgagaatct tataattcac tgtgtacctc cctctgtttc atattttcgc aattg          55

<210> SEQ ID NO 162
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 162

ccactaccac agtgcctaga gaacaayatg tgtttaataa tatttaaata at          52

<210> SEQ ID NO 163
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

```

-continued

<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (27)..(27)
<223> OTHER INFORMATION: May or may not be present

<400> SEQUENCE: 163

ccctgtctct aaaagaaaaa aaaaaaaaaa aaaaaaagaa aacaaaaccc aa 52

<210> SEQ ID NO 164
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (27)..(27)
<223> OTHER INFORMATION: May or may not be present

<400> SEQUENCE: 164

ccattatatc attcatttcc cactcatttt cttcattcca accaagatat at 52

<210> SEQ ID NO 165
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (27)..(27)
<223> OTHER INFORMATION: May or may not be present

<400> SEQUENCE: 165

tccgtctcaa aaaaaaaaaa aaaaagagaa aggaattatt ctcatgtata ca 52

<210> SEQ ID NO 166
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (27)..(27)
<223> OTHER INFORMATION: May or may not be present

<400> SEQUENCE: 166

ctggggcagt taaaaagaaa aacaaaacccc tgggccctgc ccttgaggag at 52

<210> SEQ ID NO 167
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (27)..(27)
<223> OTHER INFORMATION: May or may not be present

<400> SEQUENCE: 167

ctccagggtc ccattccagg accacacgcc tgctacctct gcagctcact ca 52

<210> SEQ ID NO 168
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (27)..(27)
<223> OTHER INFORMATION: May or may not be present

<400> SEQUENCE: 168

ggattgacat ataccagtta gacggatttt ttttttccat aaaccaggct ca 52

-continued

<210> SEQ ID NO 169
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 169

acaaataatt actacatatt tacaacrttc cagatacaga agaaatggcc aa 52

<210> SEQ ID NO 170
<211> LENGTH: 54
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (27)..(29)
<223> OTHER INFORMATION: This region may or may not be present in its entirety

<400> SEQUENCE: 170

taaacacact tgaacatca tataaatga tattactaca agacttaacc gtaa 54

<210> SEQ ID NO 171
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (27)..(27)
<223> OTHER INFORMATION: May or may not be present

<400> SEQUENCE: 171

acacaggtaa tccatgttta ttatagaaaa atgccacatt actctttatt ga 52

<210> SEQ ID NO 172
<211> LENGTH: 53
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (27)..(28)
<223> OTHER INFORMATION: This region may or may not be present in its entirety

<400> SEQUENCE: 172

agttatcata gaaaaatatt acactctgtc tgtatcccca cttccagaaa ctg 53

<210> SEQ ID NO 173
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (27)..(27)
<223> OTHER INFORMATION: May or may not be present

<400> SEQUENCE: 173

aatgaactta caggaaggta atatattgga aataaacatt ttattgaatt ta 52

<210> SEQ ID NO 174
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (27)..(27)
<223> OTHER INFORMATION: May or may not be present

<400> SEQUENCE: 174

-continued

ggacccctga aaggcacaag acacccttcc aggttcacac ttctgaaag ct 52

<210> SEQ ID NO 175
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 175

ccaccaatca ccagagcctt ctgctcrggg gtacctgagg gaaaacaagc aa 52

<210> SEQ ID NO 176
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (27)..(27)
<223> OTHER INFORMATION: May or may not be present

<400> SEQUENCE: 176

aaagactgaa aaaacattca taactatattt tcttggtatc ctcggaatg tc 52

<210> SEQ ID NO 177
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 177

tatcttcac ttgaggaga tgaggtygat ttcacctgct ggaagtccaa cc 52

<210> SEQ ID NO 178
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 178

ttgcctatga acgtttgtca cacttcygc gtgagttgct gaggcgaggt gt 52

<210> SEQ ID NO 179
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 179

gcttgctgtt ggatacatct cgccatyacc tgccactctc tagcatcctg ga 52

<210> SEQ ID NO 180
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 180

ccggggcttg ctgttgata catctckcca ttacctgcca ctctctagca tc 52

<210> SEQ ID NO 181
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 181

ctccgtttcg gtttcacttc 20

<210> SEQ ID NO 182

-continued

<211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 182

atcttctctt cagccctgt

20

<210> SEQ ID NO 183
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 183

gcacgatccg acggtagtgt

20

<210> SEQ ID NO 184
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 184

cgtaatcgg gaagctgaag

20

What is claimed is:

1. A method for correcting for errors or bias introduced during nucleic acid analysis workflow, the method comprising the steps of:

obtaining a biological sample comprising a plurality of target nucleic acid molecules from more than one locus of origin;

introducing a set of differentiator tags, wherein members of said set of differentiator tags are associated with members of said plurality, such that one or more of said loci of origin are associated with more than one differentiator tag;

amplifying each of the plurality of tagged target nucleic acid molecules to generate amplicons;

sequencing the amplicons obtained in said amplifying step to obtain sequence reads of each of the amplicons, wherein each of the sequence reads comprises a target nucleic acid molecule sequence and a differentiator tag sequence; and

correcting for error or bias introduced during said workflow by collapsing target: differentiator tag combinations observed more than once into a single count.

2. The method of claim 1, wherein said correcting step further comprises collapsing all sequence reads comprising a same differentiator tag into a single read before determining a consensus base call for the target nucleic acid molecule sequences.

3. The method of claim 2, further comprising the step of determining the presence of a sequencing or amplification error based upon a number of said differentiator tag sequences.

4. The method of claim 1, wherein said biological sample is selected from the group consisting of blood and biopsy tissue.

5. The method of claim 1, wherein said plurality of target nucleic acid molecules comprises bacterial or viral nucleic acids isolated from said biological sample.

6. The method of claim 1, wherein said plurality of target nucleic acid molecules are circulating tumor nucleic acid molecules.

7. The method of claim 1, further comprising the step of determining the copy number of a genomic region or transcript in a patient from whom the sample is obtained.

8. The method of claim 1, wherein said plurality of target nucleic acid molecules are maternally-circulating fetal or placental nucleic acid molecules.

9. A method for correcting for errors or bias introduced during nucleic acid analysis workflow, the method comprising the steps of:

obtaining a biological sample comprising a plurality of target nucleic acid molecules from more than one locus of origin;

introducing a set of unique differentiator tags, wherein members of said set of unique differentiator tags are randomly associated with members of said plurality of target nucleic acid molecules, such that any given tag is uniquely associated with said locus of origin;

amplifying each of the plurality of tagged target nucleic acid molecules to generate amplicons;

sequencing the amplicons obtained in said amplifying step to obtain sequence reads of each of the amplicons, wherein each of the sequence reads comprises a target nucleic acid molecule sequence and a differentiator tag sequence; and

correcting for error or bias introduced during said workflow by recognizing target/ differentiator tags having the same sequence as being derived from the same input molecule.

* * * * *